

$W^{2,\delta}$ ESTIMATES FOR SOLUTION SETS OF FULLY NONLINEAR ELLIPTIC INEQUALITIES ON $C^{1,\alpha}$ DOMAINS

DONGSHENG LI AND XUEMEI LI

ABSTRACT. In this paper, we establish boundary $W^{2,\delta}$ estimates for $u \in S(\lambda, \Lambda, f)$ on $C^{1,\alpha}$ domains with $f \in L^p$ as $n < p < \infty$ and $C^{1,\alpha}$ boundary values. Instead of straightening out the boundary, our main idea is to obtain boundary $W^{2,\delta}$ estimates from interior W^{2,δ_0} estimates and Whitney decomposition for some $\delta \leq \delta_0$.

1. INTRODUCTION

In this paper, we establish boundary $W^{2,\delta}$ estimates for $u \in S(\lambda, \Lambda, f)$ on $C^{1,\alpha}$ domains. Some notions and notations concerning $S(\lambda, \Lambda, f)$ are listed as follows. We refer to [2] and [3] for more details.

Definition 1.1. Let $0 < \lambda \leq \Lambda < \infty$. For $M \in \mathcal{S}(n)$ (the space of real $n \times n$ symmetric matrices), the Pucci's extremal operators \mathcal{M}^- and \mathcal{M}^+ are defined as follows:

$$\begin{aligned}\mathcal{M}^-(M, \lambda, \Lambda) &= \lambda \sum_{e_i > 0} e_i + \Lambda \sum_{e_i < 0} e_i, \\ \mathcal{M}^+(M, \lambda, \Lambda) &= \Lambda \sum_{e_i > 0} e_i + \lambda \sum_{e_i < 0} e_i,\end{aligned}$$

where $e_i = e_i(M)$ are the eigenvalues of M .

Definition 1.2. Let $p > n/2$ and $f \in L^p_{loc}(\Omega)$. A function $u \in C(\Omega)$ satisfies $\mathcal{M}^-(D^2u, \lambda, \Lambda) \leq f(x)$ (resp. $\mathcal{M}^+(D^2u, \lambda, \Lambda) \geq f(x)$) in Ω in the viscosity sense if $x_0 \in \Omega$, $\varphi \in W^{2,p}_{loc}(\Omega)$ and $u - \varphi$ has a local minimum (resp. maximum) at x_0 imply

$$\begin{aligned}\text{ess lim inf}_{x \rightarrow x_0} \{ \mathcal{M}^-(D^2\varphi, \lambda, \Lambda) - f(x) \} &\leq 0, \\ (\text{resp. } \text{ess lim sup}_{x \rightarrow x_0} \{ \mathcal{M}^+(D^2\varphi, \lambda, \Lambda) - f(x) \}) &\geq 0.\end{aligned}$$

Definition 1.3. Let $p > n/2$ and $f \in L^p_{loc}(\Omega)$. We denote by $S(\lambda, \Lambda, f)$ the space of continuous functions u in Ω such that

$$\mathcal{M}^-(D^2u, \lambda, \Lambda) \leq f(x) \leq \mathcal{M}^+(D^2u, \lambda, \Lambda)$$

in the viscosity sense in Ω .

Our main result is as follows.

Key words and phrases. Fully Nonlinear Elliptic Equations, $W^{2,\delta}$ Estimates, Whitney decomposition, $C^{1,\alpha}$ Domains.

Theorem 1.4. *Let $0 < \alpha < 1$ and $\Omega \subset \mathbb{R}^n$ be a bounded domain with $C^{1,\alpha}$ boundary portion $T \subset \partial\Omega$. Suppose that u satisfies*

$$\begin{cases} u \in S(\lambda, \Lambda, f) & \text{in } \Omega; \\ u = g & \text{on } T, \end{cases} \quad (1.1)$$

where

$$f \in L^p(\Omega) \text{ for } n < p < \infty \text{ and } g \in C^{1,\alpha}(T).$$

Then there exists a positive constant δ depending on $n, \lambda, \Lambda, \alpha$ and p such that for any domain $\Omega' \subset \subset \Omega \cup T$, we have

$$\|u\|_{W^{2,\delta}(\Omega')} \leq C (\|u\|_{L^\infty(\Omega)} + \|f\|_{L^p(\Omega)} + \|g\|_{C^{1,\alpha}(T)}), \quad (1.2)$$

where C depends on $n, \lambda, \Lambda, \alpha, \delta, p, T, \Omega'$ and Ω .

Remark 1.5. (i) When $T = \partial\Omega$ in Theorem 1.4, we may obtain a global $W^{2,\delta}(\Omega)$ estimate on the bounded $C^{1,\alpha}$ domain.

(ii) Interior W^{2,δ_0} estimates and boundary Hölder gradient estimates are used in the proof of Theorem 1.4, where δ_0 depends on n, λ and Λ . Interior W^{2,δ_0} estimates need $p > n - \epsilon$ with ϵ depending on n, λ and Λ . (See [5] for more details.) Boundary Hölder gradient estimates need $p > n$. (See the following Corollary 3.4 for more details.)

$W^{2,p}$ regularity for fully nonlinear elliptic equations has been studied extensively during recent decades. For $u \in S(\lambda, \Lambda, f)$ and $f \in L^n$, Caffarelli [2] proved interior W^{2,δ_0} estimates with $\delta_0 > 0$ depending on n, λ and Λ , which was first discovered by Lin [12] for linear elliptic equations in nondivergence form. Then, interior $W^{2,p}$ estimates for fully nonlinear elliptic equations $F(D^2u, x) = f(x)$ with $f \in L^p$ and $n < p < \infty$ were established under the following two assumptions: The homogeneous equations with constant coefficients $F(D^2w, x_0)$ have interior $C^{1,1}$ estimates for any $x_0 \in B_1$ and the oscillation of $F(M, x)$ in x is small in L^n sense. (See [2].) Escoriaza [5] extended the above estimates to the optimal range of exponents $n - \epsilon < p < \infty$ with ϵ depending on n, λ and Λ . In [9], Li and Zhang relaxed the Caffarelli's assumptions of F to: $F(D^2w, x_0)$ have interior W^{2,p_1} estimates for any $x_0 \in B_1$ and the oscillation of $F(M, x)$ in x is small in L^{p_2} sense, where $n - \epsilon < p_0 < p < p_1 \leq \infty$ and $1/p_1 + 1/p_2 = 1/p_0$. Winter [14] extended Caffarelli's interior $W^{2,p}$ regularity to boundary and established boundary $W^{2,p}$ estimates on flat domains and then on $C^{1,1}$ domains alongside a flattening argument. For oblique boundary value problem for $F(D^2u, x) = f(x)$, based on pointwise boundary $C^{2,\alpha}$ regularity in [11], Byun and Han [1] derived the boundary $W^{2,p}$ estimates on flat domains and then on $C^{2,\alpha}$ domains by straightening out boundary.

The technique of straightening out boundary is widely applied to deal with boundary regularity. However, it is no longer applicable for the $C^{1,\alpha}$ domains. The reason is as follows: If the technique of straightening out boundary is used, then for any $x_0 \in \partial\Omega$, one should find a neighborhood \mathcal{N} of x_0 and a diffeomorphism ψ from \mathcal{N} onto B_1 such that $\psi(\mathcal{N} \cap \Omega) = B_1^+$ and $\psi(\mathcal{N} \cap \partial\Omega) = B_1 \cap \{x^n = 0\}$. Writing $y = \psi(x)$, $\tilde{u}(y) = u(x)$ and $\tilde{f}(y) = f(x)$ for $x \in \mathcal{N} \cap \Omega$ and $y \in B_1^+$, we calculate

$$D^2u = D\psi^T(D^2\tilde{u} \circ \psi)D\psi + ((D\tilde{u} \circ \psi)D_{ij}\psi)_{1 \leq i, j \leq n}.$$

Then ψ need to be $C^{1,1}$ in view of the second term.

In this paper, instead of straightening out the boundary, we establish boundary $W^{2,\delta}$ estimates for $u \in S(\lambda, \Lambda, f)$ on $C^{1,\alpha}$ domains from interior W^{2,δ_0} estimates and Whitney decomposition. Whitney decomposition is an effective tool for obtaining boundary estimates from interior estimates. For instance, Cao, Li and Wang [4] utilized it to prove the optimal weighted $W^{2,p}$ estimates for elliptic equations with non-compatible conditions; Li, Li and Zhang [8] utilized it to prove boundary $W^{2,p}$ estimates for linear elliptic equations on $C^{1,\alpha}$ domains.

We illustrate our idea as follows. Let $\{Q_k\}_{k=1}^\infty$ be Whitney decomposition of Ω_1 (Suppose $0 \in \partial\Omega$ and denote $\Omega_r = \Omega \cap B_r$) and $\tilde{Q}_k = \frac{6}{5}Q_k$ the $\frac{6}{5}$ -dilation of Q_k with respect to its center. Let u satisfy (1.1). Deduce from Hölder's inequality and interior W^{2,δ_0} estimates that for any $\delta \leq \delta_0$,

$$\begin{aligned} \|D^2 u\|_{L^\delta(Q_k)}^\delta &\leq C d_k^{n(1-\frac{\delta}{\delta_0})} \|D^2(u-l)\|_{L^{\delta_0}(Q_k)}^\delta \\ &\leq C \left(d_k^{n-2\delta} \|u-l\|_{L^\infty(\tilde{Q}_k)}^\delta + d_k^{n-\frac{\delta n}{p}} \|f\|_{L^p(\tilde{Q}_k)}^\delta \right) \end{aligned}$$

for some universal constant C and any affine function l , where d_k denotes the diameter of Q_k . By C^{1,α_0} estimates up to the boundary, we can take l such that

$$|u(x) - l(x)| \leq C \text{dist}(x, \partial\Omega)^{1+\alpha_0}, \quad \forall x \in \Omega_{1/2}$$

for some positive constants C and proper $0 < \alpha_0 < 1$. It then follows from Whitney decomposition that

$$\|u-l\|_{L^\infty(\tilde{Q}_k)}^\delta \leq C d_k^{(1+\alpha_0)\delta}.$$

Combining the above estimates, we obtain

$$\|D^2 u\|_{L^\delta(Q_k)}^\delta \leq C \left(d_k^{-\delta+\alpha_0\delta+n} + d_k^{n-\frac{\delta n}{p}} \|f\|_{L^p(\Omega_1)}^\delta \right).$$

By taking sum on both sides with respect to k , we obtain the desired boundary $W^{2,\delta}$ estimate for any $\delta < \min\{1/(1-\alpha_0), p/n\}$ that guarantees $\sum_k (d_k^{-\delta+\alpha_0\delta+n} + d_k^{n-\frac{\delta n}{p}})$ is convergent.

The paper is organized as follows. In Section 2, Whitney decomposition and its relevant properties are concluded. In Section 3, we demonstrate some basic estimates for $u \in S(\lambda, \Lambda, f)$ concerning interior W^{2,δ_0} estimates and pointwise boundary Hölder gradient estimates. In Section 4, we prove Theorem 1.4.

We end this section by introducing some notations.

Notation.

1. $e_i = (0, \dots, 0, 1, \dots, 0) = i^{th}$ standard coordinate vector.
2. $x' = (x^1, x^2, \dots, x^{n-1})$ and $x = (x', x^n)$.
3. $\mathbb{R}_+^n = \{x \in \mathbb{R}^n : x^n > 0\}$.
4. $B_r(x_0) = \{x \in \mathbb{R}^n : |x - x_0| < r\}$ and $B_r^+(x_0) = B_r(x_0) \cap \mathbb{R}_+^n$.
5. $B'_r = \{x' \in \mathbb{R}^{n-1} : |x'| < r\}$ and $T_r = \{(x', 0) : x' \in B'_r\}$.
6. $\Omega_r(x_0) = \Omega \cap B_r(x_0)$ and $(\partial\Omega)_r(x_0) = \partial\Omega \cap B_r(x_0)$. We omit x_0 when $x_0 = 0$.
7. $\text{diam} E = \sup\{|x - y|, x, y \in E\}$, $\forall E \subset \mathbb{R}^n$.
8. $\text{dist}(E, F) = \inf\{|x - y|, x \in E, y \in F\}$, $\forall E, F \subset \mathbb{R}^n$.

2. WHITNEY DECOMPOSITION

In what follows, by a cube we mean a closed cube in \mathbb{R}^n , with sides parallel to the axes. We say two such cubes are disjoint if their interiors are disjoint.

Lemma 2.1. (*Whitney decomposition*) *Let Ω be a non-empty open set in \mathbb{R}^n . Then there exist two sequences of cubes Q_k (called the Whitney cubes of Ω) and $\tilde{Q}_k = \frac{6}{5}Q_k$ ($\frac{6}{5}$ -dilation of Q_k with respect to center of Q_k) such that*

- (i) $\Omega = \bigcup_{k=1}^{\infty} Q_k = \bigcup_{k=1}^{\infty} \tilde{Q}_k$;
- (ii) The Q_k are mutually disjoint;
- (iii) $d_k \leq \text{dist}(Q_k, \partial\Omega) \leq 4d_k$, where $d_k = \text{diam}Q_k$;
- (iv) Each point of Ω is contained in at most 12^n of the cubes \tilde{Q}_k .

For the proof of the above lemma, we refer to Theorem 1 and Proposition 3 in Section VI.1 in [13].

From now on, we make the following assumption on Ω :

(A) $0 \in \partial\Omega$ and there exists a continuous function $\varphi : B'_1 \rightarrow \mathbb{R}$ such that

$$\Omega_1 = \{x^n > \varphi(x'), |x| < 1\} \quad \text{and} \quad (\partial\Omega)_1 = \{x^n = \varphi(x'), |x| < 1\}.$$

Recall that $\Omega_1 = \Omega \cap B_1$ and $(\partial\Omega)_1 = \partial\Omega \cap B_1$.

Let $\{Q_k\}_{k=1}^{\infty}$ be Whitney decomposition of Ω_1 , $d_k = \text{diam}Q_k$ and $\tilde{Q}_k = \frac{6}{5}Q_k$. The following lemmas were first proved in [8] and we give their proofs here for completeness.

Lemma 2.2. *Suppose that Ω satisfies Assumption (A). Then*

$$\Omega_{1/12} \subset \bigcup_{\tilde{Q}_k \subset \Omega_{1/4}} Q_k. \quad (2.1)$$

Proof. If not, there exist a point $x \in \Omega_{1/12}$ and a cube Q_k such that $x \in Q_k$ but $\tilde{Q}_k \not\subset \Omega_{1/4}$. It follows that there exists a point $y \in \tilde{Q}_k$ with $|y| \geq 1/4$. Then we deduce from Lemma 2.1 (iii) that

$$\text{dist}(Q_k, \partial\Omega_1) \geq \text{diam}Q_k = 5\text{diam}\tilde{Q}_k/6 \geq 5(|y| - |x|)/6 \geq 5/36 > 1/12.$$

On the other hand, since $x \in Q_k \cap \Omega_{1/12}$ and $0 \in \partial\Omega$,

$$\text{dist}(Q_k, \partial\Omega_1) \leq |x| \leq 1/12.$$

Thus we get a contradiction. \square

Lemma 2.3. *Suppose that Ω satisfies Assumption (A) with $\varphi \in C^{0,1}(B'_1)$. If $q > n - 1$, then*

$$\sum_{\tilde{Q}_k \subset \Omega_{1/4}} d_k^q \leq C, \quad (2.2)$$

where C depends on n, q and $\|\varphi\|_{C^{0,1}(B'_1)}$.

Proof. Set

$$\mathcal{F}^s = \bigcup_k \{Q_k : 2^{-s-1} < d_k \leq 2^{-s}, \tilde{Q}_k \subset \Omega_{1/4}\}, \quad s = 1, 2, \dots \quad (2.3)$$

For any $Q_k \in \mathcal{F}^s$, since $\mathcal{F}^s \subset \Omega_{1/4}$, there exists $y_k \in (\partial\Omega)_{1/2}$ such that

$$\text{dist}(Q_k, y_k) = \text{dist}(Q_k, (\partial\Omega)_1) = \text{dist}(Q_k, \partial\Omega_1) \leq 4d_k \leq 2^{-s+2},$$

where we have used Lemma 2.1 (iii). (Recall that $(\partial\Omega)_r = \partial\Omega \cap B_r$ for $r > 0$.) It follows that

$$\text{dist}(x, (\partial\Omega)_1) \leq d_k + \text{dist}(Q_k, y_k) \leq 2^{-s} + 2^{-s+2} \leq 2^{-s+3}$$

for any $x \in Q_k$ and $Q_k \in \mathcal{F}^s$, which implies

$$\mathcal{F}^s \subset \Omega_{1/4} \cap \{\text{dist}(x, (\partial\Omega)_1) \leq 2^{-s+3}\}. \quad (2.4)$$

By Assumption (A), we have

$$\begin{aligned} & \Omega_{1/4} \cap \{\text{dist}(x, (\partial\Omega)_1) \leq 2^{-s+3}\} \\ & \subset \{|x'| \leq 1/4, \varphi(x') \leq x^n \leq \varphi(x') + (\|\varphi\|_{C^{0,1}(B'_1)} + 1)2^{-s+3}\}. \end{aligned}$$

Since

$$|\{|x'| \leq 1/4, \varphi(x') \leq x^n \leq \varphi(x') + (\|\varphi\|_{C^{0,1}(B'_1)} + 1)2^{-s+3}\}| \leq C2^{-s},$$

we have

$$|\Omega_{1/4} \cap \{\text{dist}(x, (\partial\Omega)_1) \leq 2^{-s+3}\}| \leq C2^{-s},$$

where C depends on n and $\|\varphi\|_{C^{0,1}(B'_1)}$. Therefore, by (2.4),

$$|\mathcal{F}^s| \leq C2^{-s}. \quad (2.5)$$

Observe that

$$\bigcup_{\tilde{Q}_k \subset \Omega_{1/4}} Q_k = \bigcup_{s=1}^{\infty} \bigcup_{Q_k \in \mathcal{F}^s} Q_k.$$

If $q > n - 1$, we derive from (2.3) and (2.5) that

$$\begin{aligned} \sum_{\tilde{Q}_k \subset \Omega_{1/4}} d_k^q & \leq \sum_{s=1}^{\infty} \left\{ \sum_{Q_k \in \mathcal{F}^s} (d_k^{q-n} \cdot d_k^n) \right\} \leq C \sum_{s=1}^{\infty} \left\{ 2^{-s(q-n)} \cdot \sum_{Q_k \in \mathcal{F}^s} d_k^n \right\} \\ & \leq C \sum_{s=1}^{\infty} 2^{-s(q-n)} |\mathcal{F}^s| \leq C \sum_{s=1}^{\infty} 2^{-s(q-n+1)} \leq C, \end{aligned}$$

where C depends on n, q and $\|\varphi\|_{C^{0,1}(B'_1)}$. \square

Remark 2.4. Lemma 2.3 is obvious as $q \geq n$ since

$$\sum_{\tilde{Q}_k \subset \Omega_{1/4}} d_k^q \leq \sum_{\tilde{Q}_k \subset \Omega_{1/4}} d_k^n \leq C_n |\Omega_{1/4}|$$

for some dimensional constant C_n .

3. PRELIMINARY ESTIMATES

The following lemma concerns interior W^{2,δ_0} regularity for $u \in S(\lambda, \Lambda, f)$ and $f \in L^n$. We refer to [2], [3] and [12] for its proof.

Theorem 3.1. *There exists a positive constant δ_0 depending on n , λ and Λ such that if*

$$u \in S(\lambda, \Lambda, f) \text{ in } B_1 \text{ and } f \in L^n(B_1),$$

then $u \in W^{2,\delta_0}(B_{1/2})$ and

$$\|u\|_{W^{2,\delta_0}(B_{1/2})} \leq C (\|u\|_{L^\infty(B_1)} + \|f\|_{L^n(B_1)}), \quad (3.1)$$

where C depends on n , λ and Λ .

The following lemma concerns pointwise boundary $C^{1,\bar{\alpha}}$ estimates for $u \in S(\lambda, \Lambda, 0)$ on flat boundaries with zero boundary values. We owe it to Krylov [7] and see more details in Lian and Zhang [11].

Lemma 3.2. *Let u satisfy*

$$\begin{cases} u \in S(\lambda, \Lambda, 0) & \text{in } B_1^+; \\ u = 0 & \text{on } T_1. \end{cases}$$

Then there exists a constant $0 < \bar{\alpha} < 1$ depending on n, λ and Λ such that u is $C^{1,\bar{\alpha}}$ at 0, i.e., there exists a constant a such that

$$|u(x) - ax^n| \leq C|x|^{1+\bar{\alpha}}\|u\|_{L^\infty(B_1^+)}, \quad \forall x \in B_{1/2}^+$$

and

$$|Du(0)| = |a| \leq C\|u\|_{L^\infty(B_1^+)},$$

where C depends on n , λ and Λ .

Based on Lemma 3.2 and compactness method, Lian and Zhang [11] proved the following lemma concerning pointwise boundary $C^{1,\alpha}$ estimates for $u \in S(\lambda, \Lambda, f)$ on $C^{1,\alpha}$ domains with $0 < \alpha < \bar{\alpha}$.

Lemma 3.3. *Let $\bar{\alpha}$ be as in Lemma 3.2 and $0 < \alpha < \bar{\alpha}$. Suppose that Ω satisfies Assumption (A) with $\varphi \in C^{1,\alpha}(B'_1)$ and u satisfies*

$$\begin{cases} u \in S(\lambda, \Lambda, f) & \text{in } \Omega_1; \\ u = g & \text{on } (\partial\Omega)_1, \end{cases} \quad (3.2)$$

where $g \in C^{1,\alpha}((\partial\Omega)_1)$ and $f \in L^n(\Omega_1)$ satisfying for some constant K_f ,

$$\|f\|_{L^n(\Omega_r)} \leq K_f r^\alpha \text{ for any } r \in (0, 1]. \quad (3.3)$$

Then u is $C^{1,\alpha}$ at 0, i.e., there exists an affine function l such that

$$|u(x) - l(x)| \leq C|x|^{1+\alpha}(\|u\|_{L^\infty(\Omega_1)} + K_f + \|g\|_{C^{1,\alpha}((\partial\Omega)_1)}), \quad \forall x \in \Omega_{r_0}$$

and

$$|Du(0)| = |Dl| \leq C(\|u\|_{L^\infty(\Omega_1)} + K_f + \|g\|_{C^{1,\alpha}((\partial\Omega)_1)}),$$

where C and r_0 depends on n , λ , Λ , α , $\bar{\alpha}$ and $\|\varphi\|_{C^{1,\alpha}(B'_1)}$.

If $f \in L^p(\Omega_1)$ for $n < p < \infty$, by Hölder's inequality, there exists a constant C_n depending only on n such that

$$\|f\|_{L^n(\Omega_r)} \leq C_n r^{1-n/p} \|f\|_{L^p(\Omega_r)} \quad \text{for any } r \in (0, 1],$$

which implies f satisfies (3.3) with $\alpha = 1 - n/p$. Then we have the following corollary:

Corollary 3.4. *Under the hypotheses of Lemma 3.3 with (3.3) replaced by $f \in L^p(\Omega_1)$ for $n < p < \infty$. Set $\alpha_0 = \min\{\alpha, 1 - n/p\}$. Then u is C^{1,α_0} at ∂ , i.e., there exists an affine function l_{x_0} such that*

$$|u(x) - l(x)| \leq C|x|^{1+\alpha_0} (\|u\|_{L^\infty(\Omega_1)} + \|f\|_{L^p(\Omega_1)} + \|g\|_{C^{1,\alpha}((\partial\Omega)_1)}), \quad \forall x \in \Omega_{r_0}$$

and

$$|Du(0)| = |Dl| \leq C(\|u\|_{L^\infty(\Omega_1)} + \|f\|_{L^p(\Omega_1)} + \|g\|_{C^{1,\alpha}((\partial\Omega)_1)}),$$

where C and r_0 depends on $n, \lambda, \Lambda, \alpha, \bar{\alpha}$ and $\|\varphi\|_{C^{1,\alpha}(B'_1)}$.

4. PROOF OF THEOREM 1.4

Theorem 1.4 follows easily from the following Theorem 4.1 which is proved by interior W^{2,δ_0} estimates (Theorem 3.1), boundary C^{1,α_0} estimates (Corollary 3.4) and Whitney decomposition (Lemma 2.1, 2.2 and 2.3).

Theorem 4.1. *Let $\bar{\alpha}$ be as in Lemma 3.2 and $0 < \alpha < \bar{\alpha}$. Suppose that Ω satisfies Assumption (A) with $\varphi \in C^{1,\alpha}(B'_1)$ and u satisfies*

$$\begin{cases} u \in S(\lambda, \Lambda, f) & \text{in } \Omega_1; \\ u = g & \text{on } (\partial\Omega)_1, \end{cases}$$

where

$$f \in L^p(\Omega_1) \quad \text{for } n < p < \infty \quad \text{and} \quad g \in C^{1,\alpha}((\partial\Omega)_1).$$

Let δ_0 be as in Theorem 3.1 and $\alpha_0 = \min\{\alpha, 1 - n/p\}$. Then for any

$$\delta \leq \delta_0 \quad \text{and} \quad \delta < 1/(1 - \alpha_0), \quad (4.1)$$

we have

$$\|u\|_{W^{2,\delta}(\Omega_{1/12})} \leq C(\|u\|_{L^\infty(\Omega_1)} + \|f\|_{L^p(\Omega_1)} + \|g\|_{C^{1,\alpha}((\partial\Omega)_1)}), \quad (4.2)$$

where C depends on $n, \lambda, \Lambda, \alpha, \bar{\alpha}, \delta, p$ and $\|\varphi\|_{C^{1,\alpha}(B'_1)}$.

Proof. In the following, we write for simplicity

$$\mathcal{H} = \|u\|_{L^\infty(\Omega_1)} + \|f\|_{L^p(\Omega_1)} + \|g\|_{C^{1,\alpha}((\partial\Omega)_1)}.$$

Let $\{Q_k\}_{k=1}^\infty$ be Whitney decomposition of Ω_1 , $d_k = \text{diam} Q_k$ and $\tilde{Q}_k = \frac{6}{5}Q_k$. For any $\tilde{Q}_k \subset \Omega_{1/4}$, let $y_k \in (\partial\Omega)_{1/2}$ and $\tilde{x}_k \in \partial\tilde{Q}_k$ such that

$$|\tilde{x}_k - y_k| = \text{dist}(\tilde{Q}_k, \partial\Omega_1) < \text{dist}(Q_k, \partial\Omega_1) \leq 4d_k,$$

where Lemma 2.1 (iii) is used in the last inequality. Consequently, we see that

$$|x - y_k| \leq |x - \tilde{x}_k| + |\tilde{x}_k - y_k| \leq 5d_k, \quad \forall x \in \tilde{Q}_k.$$

By Corollary 3.4, u is C^{1,α_0} at y_k and then there exists an affine function l_{y_k} (written by l for simplicity in the following) such that

$$|u(x) - l(x)| \leq C|x - y_k|^{1+\alpha_0} \mathcal{H} \leq Cd_k^{1+\alpha_0} \mathcal{H}, \quad \forall x \in \tilde{Q}_k, \quad (4.3)$$

where C depends on $n, \lambda, \Lambda, \alpha, \bar{\alpha}$ and $\|\varphi\|_{C^{1,\alpha}(B'_1)}$.

Since $u - l \in S(\lambda, \Lambda, f)$ in \tilde{Q}_k , we deduce from interior W^{2,δ_0} estimate, (4.3) and Hölder's inequality that

$$\begin{aligned} \int_{Q_k} |D^2(u-l)|^{\delta_0} dx &\leq C \left\{ d_k^{n-2\delta_0} \|u-l\|_{L^\infty(\tilde{Q}_k)}^{\delta_0} + d_k^{n-\delta_0} \left(\int_{\tilde{Q}_k} |f|^n dx \right)^{\delta_0/n} \right\} \\ &\leq C \left\{ d_k^{n-(1-\alpha_0)\delta_0} \mathcal{H}^{\delta_0} + d_k^{n-\delta_0 n/p} \left(\int_{\tilde{Q}_k} |f|^p dx \right)^{\delta_0/p} \right\}, \end{aligned}$$

where C depends on $n, \lambda, \Lambda, \alpha, \bar{\alpha}, p$ and $\|\varphi\|_{C^{1,\alpha}(B'_1)}$.

For any $\delta \leq \delta_0$, we deduce from Hölder's inequality and the above estimate that

$$\begin{aligned} \int_{Q_k} |D^2 u|^\delta dx &\leq C d_k^{n(1-\delta/\delta_0)} \left(\int_{Q_k} |D^2(u-l)|^{\delta_0} dx \right)^{\delta/\delta_0} \\ &\leq C \left\{ d_k^{n-(1-\alpha_0)\delta} \mathcal{H}^\delta + d_k^{n-\delta n/p} \left(\int_{\tilde{Q}_k} |f|^p dx \right)^{\delta/p} \right\}. \end{aligned}$$

Taking sum on both sides with respect to k , we obtain

$$\sum_{\tilde{Q}_k \subset \Omega_{1/4}} \int_{Q_k} |D^2 u|^\delta dx \leq C \left\{ \mathcal{H}^\delta \sum_{\tilde{Q}_k \subset \Omega_{1/4}} d_k^{n-(1-\alpha_0)\delta} + \|f\|_{L^p(\Omega_1)}^\delta \sum_{\tilde{Q}_k \subset \Omega_{1/4}} d_k^{n-\delta n/p} \right\}.$$

Since $\alpha_0 = \min\{\alpha, 1 - n/p\}$ and $\delta < 1/(1 - \alpha_0)$, we have

$$n - \delta n/p \geq n - (1 - \alpha_0)\delta > n - 1.$$

Consequently, by Lemma 2.2 and Lemma 2.3, for any $\delta \leq \delta_0$ and $\delta < 1/(1 - \alpha_0)$,

$$\begin{aligned} \int_{\Omega_{1/12}} |D^2 u|^\delta dx &\leq \sum_{\tilde{Q}_k \subset \Omega_{1/4}} \int_{Q_k} |D^2 u|^\delta dx \\ &\leq C \left(\|u\|_{L^\infty(\Omega_1)}^\delta + \|f\|_{L^p(\Omega_1)}^\delta + \|g\|_{C^{1,\alpha}(\partial\Omega_1)}^\delta \right), \end{aligned}$$

where C depends on $n, \lambda, \Lambda, \alpha, \bar{\alpha}, \delta, p$ and $\|\varphi\|_{C^{1,\alpha}(B'_1)}$. The desired estimate (4.2) then follows easily from the above estimate. \square

Theorem 1.4 then follows from Theorem 4.1.

Proof of Theorem 1.4.

Since $T \subset \partial\Omega$ is of $C^{1,\alpha}$, for any $x \in T$, there exist $r_x > 0$ and $\varphi_x \in C^{1,\alpha}(B'_{r_x}(x'))$ such that

$$\Omega_{r_x}(x) = \{x^n > \varphi_x(x')\} \cap B_{r_x}(x) \quad \text{and} \quad (\partial\Omega)_{r_x}(x) = \{x^n = \varphi_x(x')\} \cap B_{r_x}(x).$$

Since $\bigcup_{x \in T} B_{r_x/12}(x)$ cover $T \cap \bar{\Omega}'$ and $T \cap \bar{\Omega}'$ is compact, we choose a finite subset $B_{r_i/12}(x_i)$, $i = 1, 2, \dots, N$, of $\{B_{r_x/12}(x) : x \in T\}$ that still covers $T \cap \bar{\Omega}'$.

Let δ_0 be as in Theorem 3.1, $\bar{\alpha}$ be as in Lemma 3.2 and $\alpha_0 = \min\{\bar{\alpha}, \alpha, 1 - n/p\}$. Utilizing the scaled version of Theorem 4.1 in $\Omega_{r_i}(x_i) = \Omega \cap B_{r_i}(x_i)$, we obtain that for any $\delta \leq \delta_0$ and $\delta < 1/(1 - \alpha_0)$,

$$\begin{aligned} r_i^{-n/\delta-2} \|u\|_{L^\delta(\Omega_{r_i/12}(x_i))} + r_i^{-n/\delta-1} \|Du\|_{L^\delta(\Omega_{r_i/12}(x_i))} + r_i^{-n/\delta} \|D^2 u\|_{L^\delta(\Omega_{r_i/12}(x_i))} \\ \leq C_i \left(r_i^{-2} \|u\|_{L^\infty(\Omega_{r_i}(x_i))} + r_i^{-n/p} \|f\|_{L^p(\Omega_{r_i}(x_i))} + r_i^{-2} \|g\|_{C^{1,\alpha}(\partial\Omega_{r_i}(x_i))} \right) \end{aligned}$$

for $i = 1, 2, \dots, N$, where C_i depends on $n, \lambda, \Lambda, \alpha, \bar{\alpha}, \delta, p$ and $\|\varphi_{x_i}\|_{C^{1,\alpha}(B'_{r_i}(x'_i))}$.

Finally, we sum the above inequalities along with the interior W^{2,δ_0} estimate, to find $u \in W^{2,\delta}(\Omega')$ for $\Omega' \subset \subset \Omega \cup T$, with the desired estimate (1.2). \square

ACKNOWLEDGEMENT

This work is supported by NSFC 12071365.

REFERENCES

- [1] Byun, S.; Han, J. $W^{2,p}$ -estimates for fully nonlinear elliptic equations with oblique boundary conditions. *J. Differential Equations*. 268 (2020), 2125-2150.
- [2] Caffarelli, L. A.; Cabré, X. *Fully nonlinear elliptic equations*, Colloquium Publications, 43. American Mathematical Society, Providence, R.I., 1995.
- [3] Caffarelli, L. A.; Crandall, M.G.; Kocan, A. Świech, M. *On viscosity solutions of fully nonlinear equations with measurable ingredients*. *Commun. Pure Appl. Math.* 49 (1996), 365-397.
- [4] Cao, Y.; Li, D. S.; Wang, L. H. *The optimal weighted $W^{2,p}$ estimates of elliptic equation with non-compatible conditions*. *Commun. Pure Appl. Anal.* 10 (2011), 561-570.
- [5] Escauriaza, L. Z. $W^{2,n}$ a priori Estimates for Solutions to Fully Nonlinear Equations, Indiana Univ. Math. J. 42 (1993), 413-423.
- [6] Gilbarg, D.; Trudinger, N. S. *Elliptic partial differential equations of second order*. Reprint of the 1998 edition. Classics in Mathematics, Springer-Verlag, Berlin, 2001.
- [7] Krylov, N. V. *Boundedly inhomogeneous elliptic and parabolic equations in a domain*. (Russian) *Izv. Akad. Nauk SSSR Ser. Mat.* 47 (1983), 75-108.
- [8] Li, D. S.; Li X. M.; Zhang, K. $W^{2,p}$ estimates for elliptic equations on $C^{1,\alpha}$ domains. Preprint.
- [9] Li, D. S.; Zhang, K. $W^{2,p}$ interior estimates of fully nonlinear elliptic equations. *Bull. Lond. Math. Soc.* 47 (2015), 301-314.
- [10] Li, D. S.; Zhang, K. *Regularity for fully nonlinear elliptic equations with oblique boundary conditions*. *Arch. Ration. Mech. Anal.* 228 (2018), 923-967.
- [11] Lian, Y. Y.; Zhang, K. *Boundary pointwise $C^{1,\alpha}$ and $C^{2,\alpha}$ regularity for fully nonlinear elliptic equations*. *J. Differ. Equ.* 269 (2020), 1172-1191.
- [12] Lin, F. H. *Second derivative L^p -estimates for elliptic equations of nondivergent type*. *Proc. Amer. Math. Soc.* 96 (1986), 447-451.
- [13] Stein, E. M. *Singular Integrals and Differentiability Properties of Functions*. Princeton Math. Ser., vol. 30, Princeton University Press, Princeton, N.J., 1970.
- [14] Winter, N. $W^{2,p}$ and $W^{1,p}$ -estimates at the boundary for solutions of fully nonlinear, uniformly elliptic equations. *Z. Anal. Anwend.* 28 (2009), 129-164.

SCHOOL OF MATHEMATICS AND STATISTICS, XI'AN JIAOTONG UNIVERSITY, XI'AN, P.R.CHINA 710049.

SCHOOL OF MATHEMATICS AND STATISTICS, XI'AN JIAOTONG UNIVERSITY, XI'AN, P.R.CHINA 710049.

Email address: lidsh@mail.xjtu.edu.cn

Email address: pwflxm@stu.xjtu.edu.cn