

# Effects of squared four-fermion operators of the Standard Model Effective Field Theory on meson mixing

Luiz Vale Silva\*

Instituto de Física Corpuscular, Universitat de València – Consejo Superior de  
Investigaciones Científicas, Parc Científic, E-46980 Paterna, Valencia, Spain

## Abstract

The Standard Model Effective Field Theory (SMEFT) is a universal way of parametrizing New Physics (NP) manifesting as new heavy particle interactions with the Standard Model (SM) degrees of freedom, that respect the SM gauged symmetries. Higher order terms in the NP interactions possibly lead to sizable effects, mandatory for meaningful phenomenological studies, such as contributions to neutral meson mixing, which typically pushes the scale of NP to energy scales much beyond the reach of direct searches in colliders. I discuss the leading-order renormalization of double-insertions of dimension-6 four-fermion operators that change quark flavor by one unit (i.e.,  $|\Delta F| = 1$ ,  $F = \text{strange-}, \text{charm-}, \text{or bottom-flavor}$ ) by dimension-8 operators relevant to meson mixing (i.e.,  $|\Delta F| = 2$ ) in SMEFT, and consider the phenomenological implications of contributions proportional to large Yukawas. Given the underlying interest of SMEFT to encode full-fledged models at low energies, this work stresses the need to consider dimension-8 operators in phenomenological applications of dimension-6 operators of SMEFT.

## 1 Introduction

One strategy in searching for signs of New Physics (NP) – namely, phenomena that cannot be accommodated within the Standard Model (SM) – is the study of observables that are predicted by the SM to be suppressed, as for instance in the case of flavour changing neutral currents (FCNCs) due to the Glashow-Iliopoulos-Maiani (GIM) mechanism [1]. A different strategy consists in looking for deviations in observables that are precisely predicted, such as the observables that contribute to the extraction of the elements of the Cabibbo-Kobayashi-Maskawa (CKM) matrix [2, 3] in the SM, among which meson mixing observables, which also fall into the previous category, play an important role [4, 5].

In the SM, since the latter observables must involve, compared to the initial and final states and external momenta, the exchange of much heavier degrees of freedom (e.g.,  $W, Z$  bosons), an Effective Field Theory (EFT) provides at low energies a simpler picture of the underlying high-energy dynamics, in which Wilson coefficients and higher-dimensional operators carry the fingerprints of such heavy particles, a prominent example being Fermi’s contact interaction. Similarly, EFTs can be used to investigate the effects of non-SM new heavy degrees of freedom (e.g.,  $W', Z'$ ), that lead to contact interactions among the SM particles at low enough energies. The latter EFTs consist of higher-dimensional operators suppressed by some new large scale  $\Lambda_{\text{NP}}$ , typical of the NP extension, that encode in particular the flavour aspects of the new heavy sector, and their manifestation in observables that are suppressed in the SM or in observables that are precisely predicted can provide clear hints towards the discovery of NP.

The Standard Model Effective Field Theory (SMEFT) consists of the whole set of higher dimensional contact interactions that are consistent with Lorentz and the local symmetries of the SM, and is particularly useful when a new weak interacting sector is considered, in which case observational effects are dominated by the first terms in the power series in  $1/\Lambda_{\text{NP}}$ . In the case of operators of dimension-6, the so-called Warsaw basis [7] is divided into eight categories, among which we have four-fermions,  $\psi^4$ , that will play a central role in the discussion below, see Tab. 1.

---

\*luizva@ific.uv.es

$$(\bar{L}L)(\bar{L}L)$$

---


$$\begin{aligned}
Q_{\ell\ell} &= (\bar{\ell}\gamma_\mu\ell)(\bar{\ell}\gamma^\mu\ell) \\
Q_{qq}^{(1)} &= (\bar{q}\gamma_\mu q)(\bar{q}\gamma^\mu q) \\
Q_{qq}^{(3)} &= (\bar{q}\gamma_\mu\tau^I q)(\bar{q}\gamma^\mu\tau^I q) \\
Q_{\ell q}^{(1)} &= (\bar{\ell}\gamma_\mu\ell)(\bar{q}\gamma^\mu q) \\
Q_{\ell q}^{(3)} &= (\bar{\ell}\gamma_\mu\tau^I\ell)(\bar{q}\gamma^\mu\tau^I q)
\end{aligned}$$

$$(\bar{R}R)(\bar{R}R)$$

---


$$\begin{aligned}
Q_{ee} &= (\bar{e}\gamma_\mu e)(\bar{e}\gamma^\mu e) \\
Q_{xx} &= (\bar{x}\gamma_\mu x)(\bar{x}\gamma^\mu x) \\
Q_{ex} &= (\bar{e}\gamma_\mu e)(\bar{x}\gamma^\mu x) \\
Q_{ud}^{(1)} &= (\bar{u}\gamma_\mu u)(\bar{d}\gamma^\mu d) \\
Q_{ud}^{(8)} &= (\bar{u}\gamma_\mu T^A u)(\bar{d}\gamma^\mu T^A d)
\end{aligned}$$

$$(\bar{L}L)(\bar{R}R)$$

---


$$\begin{aligned}
Q_{\ell e} &= (\bar{\ell}\gamma_\mu\ell)(\bar{e}\gamma^\mu e) \\
Q_{\ell x} &= (\bar{\ell}\gamma_\mu\ell)(\bar{x}\gamma^\mu x) \\
Q_{qe} &= (\bar{q}\gamma_\mu q)(\bar{e}\gamma^\mu e) \\
Q_{qx}^{(1)} &= (\bar{q}\gamma_\mu q)(\bar{x}\gamma^\mu x) \\
Q_{qx}^{(8)} &= (\bar{q}\gamma_\mu T^A q)(\bar{x}\gamma^\mu T^A x)
\end{aligned}$$

$$(\bar{L}R)(\bar{R}L) + \text{h.c.}$$

---


$$Q_{\ell edq} = (\bar{\ell}^m e)(\bar{d}q_m)$$

$$(\bar{L}R)(\bar{L}R) + \text{h.c.}$$

---


$$\begin{aligned}
Q_{quqd}^{(1)} &= (\bar{q}^m u)\epsilon_{mn}(\bar{q}^n d) \\
Q_{quqd}^{(8)} &= (\bar{q}^m T^A u)\epsilon_{mn}(\bar{q}^n T^A d) \\
Q_{lequ}^{(1)} &= (\bar{\ell}^m e)\epsilon_{mn}(\bar{q}^n u) \\
Q_{lequ}^{(3)} &= (\bar{\ell}^m \sigma_{\mu\nu} e)\epsilon_{mn}(\bar{q}^n \sigma^{\mu\nu} u)
\end{aligned}$$

Table 1: Four-fermion operators of the so-called Warsaw basis, where  $x = u, d$ ;  $q (\ell)$  are weak-isospin doublet quarks (leptons), and  $u, d (e)$  are weak-isospin singlet quarks (leptons). Flavour or generation indices are omitted; when indicated in the text (e.g., as in the Wilson coefficient  $C_{\ell edq; fijk}$ ) they correspond to the fields above in that same ordering (i.e.,  $(\bar{\ell}_f^m e_i)(\bar{d}_j q_{m;k})$ ).

$$\begin{array}{l}
\frac{(\bar{L}L)(\bar{L}L)H^2}{Q_{\ell^4 H^2}^{(1)} = (\bar{\ell}\gamma_\mu\ell)(\bar{\ell}\gamma^\mu\ell)(H^\dagger H)} \\
Q_{\ell^4 H^2}^{(2)} = (\bar{\ell}\gamma_\mu\ell)(\bar{\ell}\gamma^\mu\tau^I\ell)(H^\dagger\tau^I H) \\
Q_{q^4 H^2}^{(1)} = (\bar{q}\gamma_\mu q)(\bar{q}\gamma^\mu q)(H^\dagger H) \\
Q_{q^4 H^2}^{(2)} = (\bar{q}\gamma_\mu q)(\bar{q}\gamma^\mu\tau^I q)(H^\dagger\tau^I H) \\
\\
\frac{(\bar{R}R)(\bar{R}R)H^2}{Q_{e^4 H^2} = (\bar{e}\gamma_\mu e)(\bar{e}\gamma^\mu e)(H^\dagger H)} \\
Q_{x^4 H^2} = (\bar{x}\gamma_\mu x)(\bar{x}\gamma^\mu x)(H^\dagger H) \\
\\
\frac{(\bar{L}R)(\bar{L}R)H^2}{Q_{\ell^2 e^2 H^2}^{(3)} = (\bar{\ell}eH)(\bar{\ell}eH)} \\
Q_{q^2 u^2 H^2}^{(5)} = (\bar{q}u\tilde{H})(\bar{q}u\tilde{H}) \\
Q_{q^2 u^2 H^2}^{(6)} = (\bar{q}T^A u\tilde{H})(\bar{q}T^A u\tilde{H}) \\
Q_{q^2 d^2 H^2}^{(5)} = (\bar{q}dH)(\bar{q}dH) \\
Q_{q^2 d^2 H^2}^{(6)} = (\bar{q}T^A dH)(\bar{q}T^A dH)
\end{array}$$

Table 2: Dimension-8 operators relevant to our discussion, where  $x = u, d$ . A complete and minimal basis is found in Ref. [6]. Flavour or generation indices are omitted, see caption of Tab. 1.

Recently, explicit bases for dimension-8 operators have been built [6, 8], among which one identifies operators involving four fermions, also central to our discussion, see Tab. 2 for a subset of them.

Operators of dimension-8 may have important phenomenological effects, and started to be discussed more systematically in various contexts: EW precision tests and Higgs measurements [9, 10, 11, 12, 13, 14, 15, 16, 17], Drell-Yan [18, 19], lepton flavour violation [20], different consequences of causality, analyticity and unitarity [21, 22, 23, 24], triple neutral gauge couplings [25, 26, 23].

In the case of NP effective operators involving fermion fields, effects that change flavour by one unit naturally lead to NP effects that change flavour by two units, which in the quark sector are efficiently probed by meson mixing. This is going to be the main interest here, namely, the leading effect of double-insertions of dimension-6 encoded in dimension-8 operators. To spell out, the focus is on the renormalization of such double-insertions. Focusing on the leading order, we will thus overlook a series of issues relevant at higher orders. Although here we focus on meson mixing, a similar discussion would hold for rare decays.

Double-insertions of higher-dimensional operators have been discussed in the literature in relation to other problems. Double-insertion of operators of dimension five have been discussed for instance in [27], that considers the lepton number violating dimension-5 Weinberg operator in SMEFT, and [28], in the case of a low-energy EFT respecting EM and QCD local symmetries. The renormalization of double-insertions of operators of dimension-5 and -6 by operators of dimension up to 7 in SMEFT has been discussed in [29]. The renormalization of double-insertions of bosonic operators of dimension-6 has been discussed in [30].

The leading-order calculation discussed here gives a first quantitative assessment of the size of contributions to meson mixing of double-insertions of dimension-6 operators in SMEFT, and higher-order effects are delegated to future work. In this respect, the phenomenological importance of meson mixing observables has triggered higher-order calculations in some specific extensions of the SM due to potentially large perturbative QCD corrections, including: two-Higgs-doublet model [31], supersymmetry [32], left-right model [33], leptoquark model [34].

This paper is organized as follows: in Sec. 2 we briefly discuss the basis of operators needed in SM calculations of meson mixing at the leading order, and extend the discussion to SMEFT; in Sec. 3 we specify our discussion in SMEFT to cases proportional to Yukawa couplings relatively large compared to the Yukawas of external fermion fields; in Sec. 4 we discuss phenomenological implications when tops in loops are considered, and then conclude. Apps. A and B contain respectively the expressions of the RG equations, and a discussion of the sensitivity of meson mixing observables to NP.




Figure 1: 1PI diagrams involving double-insertions and four external fermion legs, represented by solid lines. The Higgs scalar is represented by a dashed line. Gauge bosons can be attached to the internal loop in all possible ways.

## 2 Effective operators in meson mixing at the Leading-Log approximation

### 2.1 Standard Model

At low enough energies, heavy degrees of freedom are integrated out and their dynamics is encoded in the Wilson coefficients of higher-dimensional operators. One illustration of the use of an EFT is provided by meson mixing in the kaon sector in the SM [35, 36, 37], which proceeds via box diagrams at the leading order. Different internal flavours of the same type (here, up-type) can be combined as a result of the GIM mechanism, which suppresses SM contributions to meson mixing. There are three sets of contributions that are qualitatively very different, and quantitatively important, according to the elements of the CKM matrix: boxes involving (I) top- and up-,<sup>1</sup> (II) charm- and up-, (III) charm-, top- and up-quarks.<sup>2</sup> At the matching scale  $\mu_{EW}$  (where  $W, Z, H, t$  particles are integrated out and the first EFT is built from the full SM), case (I) is reproduced in the EFT by dimension-6 operators that change flavour by two units ( $|\Delta F| = 2$ ), and at the leading order cases (II) and (III) by dimension-6 operators that change flavour by one unit ( $|\Delta F| = 1$ ). Higher orders from strong interactions [36, 37, 39, 40, 38, 41] introduce new operators; the basis has also to be extended to account for  $1/m_c^2$  corrections, see [42] for a recent reference. A further suppression due to the GIM mechanism is the elimination of logarithmic contributions in case (II); at the same time, it makes double-insertions of dimension-6 operators finite in this case, i.e., the latter do not require renormalization by dimension-8 operators. GIM does not operate in the same way in case (III), for which the main contribution is given by a large logarithm; simultaneously, GIM does not eliminate the divergence in double-insertions of dimension-6 operators, and case (III) requires renormalization by dimension-8 operators.

To describe the resulting mixing of operators quantitatively, one must determine the anomalous dimension tensor  $\gamma_{ij,n}$ : given a set of Green's functions with two insertions of dimension-6 operators (indexed by  $i, j$ ) one calculates the counter-terms proportional to dimension-8 operators (indexed by  $n$ ), needed to renormalize the divergences resulting from the double-insertions. Large logarithms are resummed via the Renormalization Group (RG) evolution, see App. A:

$$\mu \frac{d}{d\mu} C_n^{(8)}(\mu) = \Sigma_{i,j} C_i^{(6)}(\mu) C_j^{(6)}(\mu) \gamma_{ij,n} + \Sigma_m C_m^{(8)}(\mu) \tilde{\gamma}_{mn}, \quad (1)$$

where the superscripts of the Wilson coefficients give the dimension of the corresponding operator. Solving these RG equations, the term proportional to two dimension-6 Wilson coefficients carries the logarithm  $\log(\mu_{low}/\mu_{EW})$  for some  $\mu_{low} \ll \mu_{EW}$ , consistently reproducing the logarithmic enhancement of case (III) above. The values of the dimension-8 Wilson coefficients  $C_n^{(8)}(\mu_{EW})$  are sub-leading, and the calculation of the anomalous dimension matrix  $\tilde{\gamma}_{mn}$  is not required at the leading order.

<sup>1</sup>Case (I) is the contribution that is largely dominant in neutral- $B_{(s)}$  meson mixing in the SM.

<sup>2</sup>The same qualitative discussion holds for the different gathering of contributions considered in [38].




Figure 2: 1PI diagrams involving single-insertions and two external fermion legs, represented by solid lines. The Higgs scalar is represented by a dashed line. Gauge bosons can be attached to the internal loop in all possible ways.

## 2.2 Beyond the Standard Model

An analogous picture can be drawn in SMEFT. We consider a case analog to case (III) above. First of all, we consider that the underlying NP sector does not generate tree or one-loop contributions to dimension-6  $|\Delta F| = 2$  operators, and tree contributions to  $|\Delta F| = 2$  operators of higher dimension, or at least that these contributions are highly suppressed.<sup>3</sup> Then, we consider that a possible GIM-like mechanism in the NP sector does not eliminate the need to renormalize double-insertions of  $|\Delta F| = 1$  dimension-6 operators (whose Wilson coefficients are taken to be non-zero), i.e., large logarithms are present, and assumed to be dominant. Under these assumptions, the leading contribution to meson mixing is captured by double-insertions of dimension-6 operators, that requires renormalization by dimension-8 operators. Solving an equation analogous to Eq. (1) valid for SMEFT results then, when  $\Lambda_{\text{NP}} \gg \mu_{EW}$ , in the large logarithm  $\log(\mu_{EW}/\Lambda_{\text{NP}})$ .

For simplicity, we focus on double-insertions of the same operator, with the same flavour content. Concrete extensions of the SM will typically involve more than one effective operator and a richer flavour structure, and the leading-log contributions therein could be evaluated via a similar calculation as described below. Furthermore, we focus on double-insertions of the full set of dimension-6 four-fermion operators of SMEFT, displayed in Tab. 1. Our starting point is the Warsaw basis of operators of dimension-6, and we will not discuss the matching of a particular model of renormalizable interactions onto that basis.

## 3 Contributions proportional to large Yukawa couplings

For phenomenological reasons, we focus on cases leading to contributions proportional to relatively large Yukawa couplings compared to the Yukawas of the external fields (which will lead to sizable contributions as we will see below). For instance, we focus on the contributions to kaon-meson mixing that can involve the Yukawas of charm-, bottom- and top-quarks, as well as of tau-leptons.

Fig. 1 shows 1PI Feynman diagrams for double-insertions of four-fermion operators that can lead to contributions to meson mixing. Operators of the schematic structure  $\psi^4 H^2$  are an obvious candidate to renormalize the divergences of the diagrams in Fig. 1. To full generality, other dimension-8 operators also show up in the renormalization programme, schematically:  $\psi^4 H D$ ,  $\psi^4 D^2$ , and  $\psi^4 X$ , see the basis in [6]. However, their contributions to meson mixing are proportional to one or two powers of the external fermion masses (of the same order of the external momentum scale), and we will thus neglect their contributions.

The top two diagrams in Fig. 1 introduce in general the need for counter-terms of the structure  $\psi^4 H^2$ , as seen from the equations of motion (EOMs) of fermion fields resulting from the SM Lagrangian.<sup>4</sup> However, these are also suppressed by the external fermion masses. Therefore, only the bottom two diagrams in Fig. 1 can lead to contributions proportional to large Yukawa couplings.

In principle, 1PI single-insertions of four-fermion operators lead to new contributions, see Fig. 2. The reason is that higher-dimensional operators change the EOMs of the SM fields [47].<sup>5</sup> However, EOMs of the scalar field and field strength tensors cannot change in presence of dimension-6 four-fermion operators at tree level, and the EOMs of fermion fields lead to contributions to meson mixing suppressed by the external fermion masses, as in the previous two paragraphs.

<sup>3</sup>For instance, in left-right models the masses of the extended (neutral) scalar sector suppress their tree-level contributions to meson mixing, see e.g. [33, 43].

<sup>4</sup>See [44, 45, 46] and references therein for a discussion in terms of field redefinitions. The use of EOMs is sufficient at the leading order considered here.

<sup>5</sup>See footnote 4.

WC	flavours	$B$ ( $f = 3, i = 1$ )	$B_s$ ( $f = 3, i = 2$ )	$K$ ( $f = 2, i = 1$ )
$C_{quqd}^{(1)}$	$33fi, (33if)^*$	$ C^2  < (7 \text{ TeV})^{-4}$	$ C^2  < (4 \text{ TeV})^{-4}$	$ \text{Im}\{C^2\}  < (70 \text{ TeV})^{-4}$
	$f33i, (i33f)^*$	$ C^2  < (5 \text{ TeV})^{-4}$	$ C^2  < (3 \text{ TeV})^{-4}$	$ \text{Im}\{C^2\}  < (50 \text{ TeV})^{-4}$
$C_{quqd}^{(8)}$	$33fi, (33if)^*$	$ C^2  < (3 \text{ TeV})^{-4}$	$ C^2  < (2 \text{ TeV})^{-4}$	$ \text{Im}\{C^2\}  < (30 \text{ TeV})^{-4}$
	$f33i, (i33f)^*$	$ C^2  < (3 \text{ TeV})^{-4}$	$ C^2  < (2 \text{ TeV})^{-4}$	$ \text{Im}\{C^2\}  < (30 \text{ TeV})^{-4}$

Table 3: Estimated bounds on the Wilson coefficients of the operators in Eq. (2). The first column indicates the Wilson Coefficient (WC)  $C$  being probed at the scale  $\Lambda_{\text{NP}}$ , together with its flavour indices (bounds correspond to the combination of the two cases provided, with no interference being present). The three remaining columns give estimated bounds accordingly to the meson system.

Basic expressions and explicit anomalous dimensions describing the mixing of double-insertions into operators of the structure  $\psi^4 H^2$  are collected in App. A.

## 4 Phenomenology for tops in loops

We focus our phenomenological discussion on contributions to meson mixing in which both internal fermions running inside the loop are top-quarks. The resulting effect is then proportional to two powers of the Yukawa of the top-quark. The scope of SMEFT operators is the following, see Tab. 1:

$$Q_{quqd}^{(1,8)}, \quad Q_{ud}^{(1,8)}, \quad Q_{qu}^{(1,8)}, \quad Q_{qd}^{(1,8)}, \quad Q_{qq}^{(1,3)}. \quad (2)$$

Since here the internal flavour is a top, below the EW scale the relevant operators are:

$$\begin{aligned}
O_1^{q_1 q_2} &= (\bar{q}_1^\alpha \gamma^\mu L q_2^\alpha) (\bar{q}_1^\beta \gamma_\mu L q_2^\beta), \\
O_2^{q_1 q_2} &= (\bar{q}_1^\alpha L q_2^\alpha) (\bar{q}_1^\beta L q_2^\beta), \\
O_3^{q_1 q_2} &= (\bar{q}_1^\alpha L q_2^\beta) (\bar{q}_1^\beta L q_2^\alpha), \\
O_4^{q_1 q_2} &= (\bar{q}_1^\alpha L q_2^\alpha) (\bar{q}_1^\beta R q_2^\beta), \\
O_5^{q_1 q_2} &= (\bar{q}_1^\alpha L q_2^\beta) (\bar{q}_1^\beta R q_2^\alpha), \\
\tilde{O}_1^{q_1 q_2} &= (\bar{q}_1^\alpha \gamma^\mu R q_2^\alpha) (\bar{q}_1^\beta \gamma_\mu R q_2^\beta), \\
\tilde{O}_2^{q_1 q_2} &= (\bar{q}_1^\alpha R q_2^\alpha) (\bar{q}_1^\beta R q_2^\beta), \\
\tilde{O}_3^{q_1 q_2} &= (\bar{q}_1^\alpha R q_2^\beta) (\bar{q}_1^\beta R q_2^\alpha),
\end{aligned} \quad (3)$$

where  $L$  ( $R$ ) are left- (right-) handed projectors,  $\alpha, \beta$  are color indices, and flavours are  $q_1, q_2 = b, c, s, d, u$ . Their hadronic matrix elements are calculated for instance in [48, 49], which are chirally enhanced in the case of kaons.

Finally, one sets constraints on the NP Wilson coefficients of the operators in Eq. (2) based on their contributions to different meson mixing observables, namely: indirect CP violation in the system of kaons, and mass differences in the systems of  $B_{(s)}$ -mesons. Although the relevant RGEs in SMEFT are provided in App. A, we do not discuss the mass difference in the system of kaons nor charm-meson mixing, due to the underlying difficulty in having precise SM predictions in these cases in consequence of long-distance effects. Similarly, although our focus is on meson mixing, we also provide in App. A the RGEs relevant for muonium oscillation,  $\mu^+ e^- \leftrightarrow \mu^- e^+$  (for a recent reference on this topic see [50]). We do not include the latter in our discussion due to the limited sensitivity of dimension-8 operators to NP effects. We exploit the bounds provided in [51, 52] on generic NP contributions, see App. B. Despite being an effect coming from a dimension-8 operator and being generated at one-loop, given the precision with which these observables are known, one reaches a sensitivity to multi-TeV NP effects, see Tabs. 3 and 4; we work in the basis in which down-type flavours are mass eigenstates. A certain number of comments are in order:

- Sub-leading effects above the EW scale may be numerically relevant if the leading logarithmic term is not largely dominant. Their determination, however, is beyond the scope of this work. Among the possible effects showing up at higher order, we mention single-insertions of  $|\Delta F| = 2$  dimension-8 operators and their renormalization by  $|\Delta F| = 2$  dimension-6 operators.

WC	flavours	$K$ ( $f = 2, i = 1$ )
$C_{ud}^{(1)}$	$33fi$	$ \text{Im}\{C^2\}  < (30 \text{ TeV})^{-4}$
$C_{ud}^{(8)}$	$33fi$	$ \text{Im}\{C^2\}  < (10 \text{ TeV})^{-4}$
$C_{qd}^{(1)}$	$33fi$	$ \text{Im}\{C^2\}  < (30 \text{ TeV})^{-4}$
$C_{qd}^{(8)}$	$33fi$	$ \text{Im}\{C^2\}  < (10 \text{ TeV})^{-4}$
$C_{qu}^{(1)}$	$fi33$	$ \text{Im}\{C^2\}  < (30 \text{ TeV})^{-4}$
$C_{qu}^{(8)}$	$fi33$	$ \text{Im}\{C^2\}  < (10 \text{ TeV})^{-4}$
$C_{qq}^{(1)}$	$fi33 = 33fi$	$ \text{Im}\{C^2\}  < (30 \text{ TeV})^{-4}$
$C_{qq}^{(3)}$	$fi33 = 33fi$	$ \text{Im}\{C^2\}  < (30 \text{ TeV})^{-4}$
	$f33i = 3if3$	$ \text{Im}\{C^2\}  < (30 \text{ TeV})^{-4}$

Table 4: See caption of Tab. 3 for comments. Additionally, the contribution of  $Q_{qq;f33i}^{(1)}$  to double-insertions proportional to the Yukawa of the top-quark vanishes at this order.

- The fit [51, 52] is done under the assumption that NP is only present in contact interactions that change flavour by two units, while here we analyse the combined effects of  $|\Delta F| = 1$  NP operators. In presence of  $|\Delta B| = 1$  operators  $Q_{qq}^{(1,3)}$ ,  $Q_{ud}^{(1,8)}$ ,  $Q_{qu}^{(1,8)}$ ,  $Q_{qd}^{(1,8)}$  there is NP affecting the interpretation of the extracted values of the unitary triangle angles  $\beta, \beta_s$  (beyond mixing-induced CP violation, that is already taken into account in that fit), namely, contributions similar to the gluonic penguin generated in the SM. These observables play a central role in setting constraints on the allowed size of NP in  $|\Delta B| = 2$  [53, 54], and for this reason  $|\Delta B| = 1$  operators are not included in Tab. 4. See also [55] for a discussion of the effects of dimension-6 operators in the extraction of the elements of the CKM matrix. A re-analysis of the global fit taking into account  $|\Delta B| = 1$  operators involving top-quarks will be the subject of future work. On the other hand, we provide bounds on NP in the kaon sector in Tab. 4, which result in the global fit of [51] from  $|\epsilon_K|$ .
- In the case of the operators shown in Tab. 4, single-insertions [43, 56, 57, 58] at one-loop can lead to contributions to meson mixing, setting bounds on  $|\text{Im}\{V_{td}V_{ts}^*C\}|$  instead of  $|\text{Im}\{C^2\}|$ , see [43, 57] where finite terms are discussed. Although these single-insertions are suppressed by off-diagonal elements of the CKM matrix, chiral enhancements in cases  $Q_{ud}^{(1,8)}$  and  $Q_{qd}^{(1,8)}$  lead to a much better sensitivity to the NP energy scale compared to Tab. 4. In cases  $Q_{qu}^{(1,8)}$ , the sensitivity is similar to the one shown in Tab. 4. The operators  $Q_{qq}^{(1,3)}$  can contribute to charm-meson mixing [56]. In the case of the operators shown in Tab. 3, contributions from finite terms are suppressed by the mass scale of external fields, and thus negligible.
- Other observables can also constrain the same NP effective couplings; e.g., many other operators are radiatively generated through single-insertions, whose anomalous dimensions at one-loop are found in [59, 60, 61]. These radiative effects result in contributions to rare semi-leptonic transitions, for instance. However, in particular,  $K \rightarrow \pi \bar{\nu} \nu$  rates are presently not known to an experimental accuracy much better than 100% [62], and the main sensitivity to the NP scale is still achieved by meson mixing.

Beyond tops in loops, one can also have other internal heavy flavours in the case of kaon-meson mixing, namely, charm- and bottom-quarks, and tau-leptons, which are much heavier than kaons. In these cases double-insertions have also to be considered below the EW scale, and require thus calculations beyond SMEFT.

## 5 Conclusions

I have discussed effects of a generic heavy NP sector that are encoded in higher-dimensional operators. More exactly, I have calculated the renormalization by dimension-8 operators of double-insertions of dimension-6 operators, where the latter changes flavour number by one unit and the

former by two units. At energy scales much below the characteristic scale of NP, the effects of double-insertions are constrained by meson mixing observables, which receive suppressed contributions from the SM, due to the GIM mechanism, that are precisely predicted in the case of many observables. The calculation here discussed provides the leading-order contribution to meson mixing in SMEFT when  $|\Delta F| = 2$  tree-level effects and dimension-6  $|\Delta F| = 2$  operators generated at one-loop are absent or suppressed, and no GIM-like mechanism in the NP sector operates to eliminate the need for renormalization.

The scope of SMEFT operators considered here extends to four-fermions of different chiralities, and to semi-leptonic operators. I have focused the phenomenological discussion on tops as the internal flavour in fermion loops, resulting in contributions proportional to the Yukawa coupling of the top. Given the level of experimental accuracy reached for meson mixing observables, even loop-suppressed double-insertions lead to meaningful and powerful bounds on NP, probing energy scales much beyond the reach of direct searches in colliders.

## Acknowledgements

I am indebted to Antonio Pich for early involvement in this project and for careful reading of the manuscript. This work has been supported in part by the Spanish Government and ERDF funds from the EU Commission [Grant FPA2017-84445-P] and the Generalitat Valenciana [Grant Prome-teo/2017/053]. This project has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement No 101031558.

## A Renormalization and RG equations

We consider on-shell renormalization with dimensional regularization ( $D = 4 - 2\epsilon$ ) and (modified) minimal subtraction; Ref. [63] is used to identify UV divergences; we also employ BMHV scheme to deal with  $\gamma_5$ .

Next, we follow closely the discussion of [37]. It will be left implicit that the operators being discussed are the ones of the main text. Up to  $\mathcal{O}(\Lambda_{\text{NP}}^{-6})$  terms, the non-renormalizable part of the Lagrangian is given by:

$$\begin{aligned} \mathcal{L}_{\text{eff; non-ren.}} = & -\frac{1}{\Lambda_{\text{NP}}^2} \sum_{i,j} \left( \mathcal{C}_i^{(6)} Z_{ij}^{-1} + \frac{m_H^2}{\Lambda_{\text{NP}}^2} \mathcal{C}_i^{(8)} \hat{Z}_{ij}^{-1} \right) \mu^{\Delta_j^{(6)}} Q_j^{(6),\text{bare}} \\ & -\frac{1}{\Lambda_{\text{NP}}^4} \sum_n \left( \sum_{i,j} \mathcal{C}_i^{(6)} \mathcal{C}_j^{(6)} Z_{ij,n}^{-1} + \sum_m \mathcal{C}_m^{(8)} \tilde{Z}_{mn}^{-1} \right) \mu^{\Delta_n^{(8)}} Q_n^{(8),\text{bare}} + \dots, \end{aligned} \quad (4)$$

where “(6)” and “(8)” are the dimensions of the operators involved, except when otherwise indicated (namely, the dimension-8 operators  $Q_{q^2 x^2 H^2}^{(6)}$ ,  $x = u, d$ , and their Wilson coefficients). The Wilson coefficients  $\mathcal{C}^{(6)}$  and  $\mathcal{C}^{(8)}$  are dimensionless (we reserve the typesetting  $C^{(6)}$  and  $C^{(8)}$  for the dimensionful ones), and the expressions multiplying the bare operators are the bare Wilson coefficients. The ellipses denote counter-terms proportional to unphysical operators, that will be omitted hereafter. Note that on-shell matrix elements of double-insertions of EOM-vanishing operators are in principle non-zero, see e.g. [64]. However, we consider as our starting point in the main text dimension-6 operators of the Warsaw basis, which we do not further reduce by the use of EOMs. Other than its non-renormalizable part,  $\mathcal{L}_{\text{eff; non-ren.}}$ , the full Lagrangian  $\mathcal{L}_{\text{eff}}$  also includes  $\mathcal{L}^{\text{“SM”}}$ , which is the SM Lagrangian together with counter-terms proportional to SM operators due to the presence of NP interactions.

Insertions of higher-dimensional operators leading to Green’s functions relevant for meson mixing are indicated as (the “SM” part of the Lagrangian is not being explicitly shown):

$$\langle \mathbf{T} \exp \left[ i \int d^D x \mathcal{L}_{\text{eff; non-ren.}}(x) \right] \rangle_{|\Delta F|=2}^{\text{“SM”}} = -i \langle a^{(6)} + a^{(8)} \rangle_{|\Delta F|=2}^{\text{“SM”}} \quad (5)$$

up to terms  $\mathcal{O}(\Lambda_{\text{NP}}^{-6})$ , where

$$a^{(6)}(x) = \frac{1}{\Lambda_{\text{NP}}^2} \sum_{i,j} \mathcal{C}_i^{(6)} Z_{ij}^{-1} \mu^{\Delta_j^{(6)}} Q_j^{(6),\text{bare}}(x) \quad (6)$$

and



$$\begin{aligned}
a^{(8)}(x) &= \frac{m_H^2}{\Lambda_{\text{NP}}^4} \Sigma_{i,j} C_i^{(8)} \hat{Z}_{ij}^{-1} \mu^{\Delta_j^{(6)}} Q_j^{(6),\text{bare}}(x) \\
&\quad + \frac{1}{\Lambda_{\text{NP}}^4} \Sigma_{m,n} C_m^{(8)} \tilde{Z}_{mn}^{-1} \mu^{\Delta_n^{(8)}} Q_n^{(8),\text{bare}}(x) + \frac{1}{\Lambda_{\text{NP}}^4} \Sigma_{i,j} C_i^{(6)} C_j^{(6)} \mathcal{R}_{ij}(x),
\end{aligned} \tag{7}$$

with

$$\mathcal{R}_{ij}(x) = \left( \Sigma_{i',j'} Z_{i'i'}^{-1} Z_{j'j}^{-1} \mu^{\Delta_{i'}^{(6)} + \Delta_{j'}^{(6)}} \mathcal{R}_{i'j'}^{\text{bare}}(x) + \Sigma_n Z_{ij,n}^{-1} \mu^{\Delta_n^{(8)}} Q_n^{(8),\text{bare}}(x) \right) \tag{8}$$

and

$$\mathcal{R}_{i'j'}^{\text{bare}}(x) = \frac{-i}{2} \int d^D y \mathbf{T} \left( Q_{i'}^{(6),\text{bare}}(x) Q_{j'}^{(6),\text{bare}}(y) + Q_{j'}^{(6),\text{bare}}(x) Q_{i'}^{(6),\text{bare}}(y) \right). \tag{9}$$

In powers of the Yukawa couplings  $y$  (possibly different Yukawa couplings are represented by the same letter  $y$ ), the renormalization factors are expanded as ( $\delta$  is the Kronecker symbol, with indices omitted):

$$Z^{-1} = \delta + \frac{y^2}{(4\pi)^2} Z^{-1,(1)} + \dots, \quad Z^{-1,(n)} = \Sigma_{r=0}^n \frac{1}{\epsilon^r} Z_r^{-1,(n)} + \mathcal{O}(\epsilon), \tag{10}$$

and similarly for  $\tilde{Z}^{-1}$ , while  $\hat{Z}^{-1}$  and the tensor  $Z_{ij,n}^{-1}$  have perturbative expansions starting at  $\mathcal{O}(y^2)$ .

From the scale independence of the bare coefficients, one gets (cf. Eq. (1), given in the context of the SM)

$$\mu \frac{d}{d\mu} C_n^{(8)}(\mu) = \Sigma_{i,j} C_i^{(6)}(\mu) C_j^{(6)}(\mu) \gamma_{ij,n} + \Sigma_m C_m^{(8)}(\mu) \tilde{\gamma}_{mn}, \tag{11}$$

with in particular

$$\gamma_{ij,n} = \frac{y^2}{(4\pi)^2} \gamma_{ij,n}^{(0)} + \dots, \quad \gamma_{ij,n}^{(0)} = \frac{\left( 2\Delta_y + \Delta_i^{(6)} + \Delta_j^{(6)} - \Delta_n^{(8)} \right)}{\epsilon} [Z_1^{-1,(1)}]_{ij,n} + \mathcal{O}(\epsilon), \tag{12}$$

where  $\Delta_y = \epsilon$  is the mass dimension of the coupling  $y$ , which satisfies the renormalization group equation  $\mu dy(\mu)/d\mu = -\Delta_y y(\mu) + \mathcal{O}(y^2)$ . Similarly,  $\Delta^{(6)} = 2\epsilon$  for four-fermion operators  $\psi^4$ , and  $\Delta^{(8)} = 4\epsilon$  for operators of the schematic form  $\psi^4 H^2$ . To achieve the simple form of Eq. (11), we have neglected terms  $\propto m_H^2$  indicated in the right-hand side of Eq. (4), which is justified in the leading order being considered here, where  $C^{(8)}(\Lambda_{\text{NP}})$  is sub-leading, see the main text.

The anomalous dimensions  $\gamma_{ij,n}^{(0)}$  are then calculated from the finiteness of the Green's functions introduced above,  $\langle \mathcal{R}_{ij} \rangle^{(0)}$  in particular (“0” indicates the leading order, shown in Figs. 1 and 2, and “[div.]” the divergent parts):

$$\frac{(y^{\text{bare}})^2}{(4\pi)^2} \Sigma_n [Z^{-1,(1)}]_{ij,n} \langle Q_n^{(8),\text{bare}} \rangle_{|\Delta F|=2}^{(0),\text{SM}} [\text{div.}] = -\langle \mathcal{R}_{ij}^{\text{bare}} \rangle_{|\Delta F|=2}^{(0),\text{SM}} [\text{div.}]. \tag{13}$$

Since we focus on the special case in which  $i = j$ ,  $\gamma^{(0)}$  is a matrix.

Finally, at the leading-log approximation:

$$C_n^{(8)}(\mu_{EW}) = \frac{y^2(\mu_{EW})}{(4\pi)^2} \ln \left( \frac{\mu_{EW}}{\Lambda_{\text{NP}}} \right) \Sigma_{i,j} C_i^{(6)}(\Lambda_{\text{NP}}) C_j^{(6)}(\Lambda_{\text{NP}}) \gamma_{ij,n}^{(0)}. \tag{14}$$

The scale  $\Lambda_{\text{NP}}$  appearing in the logarithm is set to 1 TeV in the numerical applications in the main text.

## A.1 Explicit expressions for the RG equations

The RG equations are given in the following (the sum over  $j, k$  indices is left implicit):

$$\begin{aligned}
(4\pi)^2 \mu \frac{d}{d\mu} C_{\ell^2 e^2 H^2; f i f i}^{(3)} &= G_a^{\ell e d q}[\text{lepton}] \times S_a \times \mathcal{G}_a^{\ell e d q} \times (Y_d^\dagger)_{kj}^2 \times (C_{\ell e d q; f i j k})^2 \\
&\quad + G_a^{\ell e q u(1)}[\text{lepton}] \times S_a \times \mathcal{G}_a^{\ell e q u(1)} \times (Y_u)_{kj}^2 \times (C_{\ell e q u; f i j k}^{(1)})^2 \\
&\quad + G_b^{\ell e} \times S_b \times \mathcal{G}_b^{\ell e} \times (Y_e^\dagger)_{kj}^2 \times (C_{\ell e; f k j i})^2
\end{aligned} \tag{15}$$

$$\begin{aligned}
(4\pi)^2 \mu \frac{d}{d\mu} C_{q^2 d^2 H^2; f_i f_i}^{(5)} &= G_a^{\ell e d q} [quark] \times S_a \times \mathcal{G}_a^{\ell e d q} \times (Y_e^\dagger)_{kj}^2 \times (C_{\ell e d q; k j i}^*)^2 \\
&+ G_a^{quqd(1)} \times S_a \times \mathcal{G}_a^{quqd(1)} \times (Y_u)_{kj}^2 \times (C_{quqd; j k f i}^{(1)})^2 \\
&+ G_b^{quqd(1)} \times S_b \times \mathcal{G}_b^{quqd(1)} \times (Y_u)_{kj}^2 \times (C_{quqd; f k j i}^{(1)})^2 \\
&+ G_b^{quqd(8)} [singlet] \times S_b \times \mathcal{G}_b^{quqd(8)} \times (Y_u)_{kj}^2 \times (C_{quqd; f k j i}^{(8)})^2 \\
&+ G_b^{qd(1)} [singlet] \times S_b \times \mathcal{G}_b^{qd(1)} \times (Y_d^\dagger)_{kj}^2 \times (C_{qd; f k j i}^{(1)})^2 \\
&+ G_b^{qd(8)} [singlet] \times S_b \times \mathcal{G}_b^{qd(8)} \times (Y_d^\dagger)_{kj}^2 \times (C_{qd; f k j i}^{(8)})^2
\end{aligned} \tag{16}$$

$$\begin{aligned}
(4\pi)^2 \mu \frac{d}{d\mu} C_{q^2 d^2 H^2; f_i f_i}^{(6)} &= G_a^{quqd(8)} \times S_a \times \mathcal{G}_a^{quqd(8)} \times (Y_u)_{kj}^2 \times (C_{quqd; j k f i}^{(8)})^2 \\
&+ G_b^{quqd(8)} [octet] \times S_b \times \mathcal{G}_b^{quqd(8)} \times (Y_u)_{kj}^2 \times (C_{quqd; f k j i}^{(8)})^2 \\
&+ G_b^{qd(1)} [octet] \times S_b \times \mathcal{G}_b^{qd(1)} \times (Y_d^\dagger)_{kj}^2 \times (C_{qd; f k j i}^{(1)})^2 \\
&+ G_b^{qd(8)} [octet] \times S_b \times \mathcal{G}_b^{qd(8)} \times (Y_d^\dagger)_{kj}^2 \times (C_{qd; f k j i}^{(8)})^2
\end{aligned} \tag{17}$$

$$\begin{aligned}
(4\pi)^2 \mu \frac{d}{d\mu} C_{q^2 u^2 H^2; f_i f_i}^{(5)} &= G_a^{\ell e q u(1)} [quark] \times S_a \times \mathcal{G}_a^{\ell e q u(1)} \times (Y_e)_{kj}^2 \times (C_{\ell e q u; j k f i}^{(1)})^2 \\
&+ G_a^{quqd(1)} \times S_a \times \mathcal{G}_a^{quqd(1)} \times (Y_d)_{kj}^2 \times (C_{quqd; f i j k}^{(1)})^2 \\
&+ G_b^{quqd(1)} \times S_b \times \mathcal{G}_b^{quqd(1)} \times (Y_d)_{kj}^2 \times (C_{quqd; j i f k}^{(1)})^2 \\
&+ G_b^{quqd(8)} [singlet] \times S_b \times \mathcal{G}_b^{quqd(8)} \times (Y_d)_{kj}^2 \times (C_{quqd; j i f k}^{(8)})^2 \\
&+ G_b^{qu(1)} [singlet] \times S_b \times \mathcal{G}_b^{qu(1)} \times (Y_u^\dagger)_{kj}^2 \times (C_{qu; f k j i}^{(1)})^2 \\
&+ G_b^{qu(8)} [singlet] \times S_b \times \mathcal{G}_b^{qu(8)} \times (Y_u^\dagger)_{kj}^2 \times (C_{qu; f k j i}^{(8)})^2
\end{aligned} \tag{18}$$

$$\begin{aligned}
(4\pi)^2 \mu \frac{d}{d\mu} C_{q^2 u^2 H^2; f_i f_i}^{(6)} &= G_a^{quqd(8)} \times S_a \times \mathcal{G}_a^{quqd(8)} \times (Y_d)_{kj}^2 \times (C_{quqd; f i j k}^{(8)})^2 \\
&+ G_b^{quqd(8)} [octet] \times S_b \times \mathcal{G}_b^{quqd(8)} \times (Y_d)_{kj}^2 \times (C_{quqd; j i f k}^{(8)})^2 \\
&+ G_b^{qu(1)} [octet] \times S_b \times \mathcal{G}_b^{qu(1)} \times (Y_u^\dagger)_{kj}^2 \times (C_{qu; f k j i}^{(1)})^2 \\
&+ G_b^{qu(8)} [octet] \times S_b \times \mathcal{G}_b^{qu(8)} \times (Y_u^\dagger)_{kj}^2 \times (C_{qu; f k j i}^{(8)})^2
\end{aligned} \tag{19}$$

$$\begin{aligned}
(4\pi)^2 \mu \frac{d}{d\mu} C_{\ell^4 H^2; f_i f_i}^{(1)} &= G_c^{\ell e} [\ell\ell] \times S_c \times \mathcal{G}_c^{\ell e} \times [Y_e Y_e^\dagger]_{jj} \times (C_{\ell e; f i j j})^2 \\
&+ G_c^{\ell u} [lepton] \times S_c \times \mathcal{G}_c^{\ell u} \times [Y_u Y_u^\dagger]_{jj} \times (C_{\ell u; f i j j})^2 \\
&+ G_c^{\ell d} [lepton] \times S_c \times \mathcal{G}_c^{\ell d} \times [Y_d Y_d^\dagger]_{jj} \times (C_{\ell d; f i j j})^2 \\
&+ G_c^{\ell q(1)} [lepton] \times S_c \times \mathcal{G}_c^{\ell q(1)} \times \left( [Y_u^\dagger Y_u]_{jj} + [Y_d^\dagger Y_d]_{jj} \right) \times (C_{\ell q; f i j j}^{(1)})^2 \\
&+ G_c^{\ell q(3)} [lepton] \times S_c \times \mathcal{G}_c^{\ell q(3)} \times \left( [Y_u^\dagger Y_u]_{jj} + [Y_d^\dagger Y_d]_{jj} \right) \times (C_{\ell q; f i j j}^{(3)})^2 \\
&+ G_c^{\ell\ell} \times S_c \times \mathcal{G}_c^{\ell\ell} \times [Y_e^\dagger Y_e]_{jj} \times (C_{\ell\ell; f i j j})^2 \\
&+ G_d^{\ell\ell} [singlet] \times S_d \times \mathcal{G}_d^{\ell\ell} \times [Y_e^\dagger Y_e]_{jj} \times (C_{\ell\ell; f j j i})^2
\end{aligned} \tag{20}$$

$$(4\pi)^2 \mu \frac{d}{d\mu} C_{\ell^4 H^2; f_i f_i}^{(2)} = G_d^{\ell\ell} [triplet(2)] \times S_d \times \mathcal{G}_d^{\ell\ell} \times [Y_e^\dagger Y_e]_{jj} \times (C_{\ell\ell; f j j i})^2 \tag{21}$$

$$\begin{aligned}
(4\pi)^2 \mu \frac{d}{d\mu} C_{q^4 H^2; f i f i}^{(1)} &= G_c^{qe} [quark] \times S_c \times \mathcal{G}_c^{qe} \times [Y_e Y_e^\dagger]_{jj} \times (C_{qe; f i j j})^2 & (22) \\
&+ G_c^{qu(1)} [qq] \times S_c \times \mathcal{G}_c^{qu(1)} \times [Y_u Y_u^\dagger]_{jj} \times (C_{qu; f i j j}^{(1)})^2 \\
&+ G_c^{qd(1)} [qq] \times S_c \times \mathcal{G}_c^{qd(1)} \times [Y_d Y_d^\dagger]_{jj} \times (C_{qd; f i j j}^{(1)})^2 \\
&+ G_c^{qu(8)} [qq] \times S_c \times \mathcal{G}_c^{qu(8)} \times [Y_u Y_u^\dagger]_{jj} \times (C_{qu; f i j j}^{(8)})^2 \\
&+ G_c^{qd(8)} [qq] \times S_c \times \mathcal{G}_c^{qd(8)} \times [Y_d Y_d^\dagger]_{jj} \times (C_{qd; f i j j}^{(8)})^2 \\
&+ G_c^{\ell q(1)} [quark] \times S_c \times \mathcal{G}_c^{\ell q(1)} \times [Y_e^\dagger Y_e]_{jj} \times (C_{\ell q; j j f i}^{(1)})^2 \\
&+ G_c^{\ell q(3)} [quark] \times S_c \times \mathcal{G}_c^{\ell q(3)} \times [Y_e^\dagger Y_e]_{jj} \times (C_{\ell q; j j f i}^{(3)})^2 \\
&+ G_c^{qq(1)} \times S_c \times \mathcal{G}_c^{qq(1)} \times ([Y_u^\dagger Y_u]_{jj} + [Y_d^\dagger Y_d]_{jj}) \times (C_{qq; f i j j}^{(1)})^2 \\
&+ G_c^{qq(3)} \times S_c \times \mathcal{G}_c^{qq(3)} \times ([Y_u^\dagger Y_u]_{jj} + [Y_d^\dagger Y_d]_{jj}) \times (C_{qq; f i j j}^{(3)})^2 \\
&+ G_d^{qq(1)} [singlet] \times S_d \times \mathcal{G}_d^{qq(1)} \times ([Y_u^\dagger Y_u]_{jj} + [Y_d^\dagger Y_d]_{jj}) \times (C_{qq; f j j i}^{(1)})^2 \\
&+ G_d^{qq(3)} [singlet] \times S_d \times \mathcal{G}_d^{qq(3)} \times ([Y_u^\dagger Y_u]_{jj} + [Y_d^\dagger Y_d]_{jj}) \times (C_{qq; f j j i}^{(3)})^2
\end{aligned}$$

$$\begin{aligned}
(4\pi)^2 \mu \frac{d}{d\mu} C_{q^4 H^2; f i f i}^{(2)} &= G_d^{qq(1)} [triplet(2)] \times S_d \times \mathcal{G}_d^{qq(1)} \times ([Y_d^\dagger Y_d]_{jj} - [Y_u^\dagger Y_u]_{jj}) \times (C_{qq; f j j i}^{(1)})^2 \\
&+ G_d^{qq(3)} [triplet(2)] \times S_d \times \mathcal{G}_d^{qq(3)} \times ([Y_u^\dagger Y_u]_{jj} - [Y_d^\dagger Y_d]_{jj}) \times (C_{qq; f j j i}^{(3)})^2 & (23)
\end{aligned}$$

$$\begin{aligned}
(4\pi)^2 \mu \frac{d}{d\mu} C_{e^4 H^2; f i f i} &= G_c^{\ell e} [ee] \times S_c \times \mathcal{G}_c^{\ell e} \times [Y_e^\dagger Y_e]_{jj} \times (C_{\ell e; j j f i})^2 & (24) \\
&+ G_c^{qe} [lepton] \times S_c \times \mathcal{G}_c^{qe} \times ([Y_u^\dagger Y_u]_{jj} + [Y_d^\dagger Y_d]_{jj}) \times (C_{qe; j j f i})^2 \\
&+ G_c^{eu} [lepton] \times S_c \times \mathcal{G}_c^{eu} \times [Y_u Y_u^\dagger]_{jj} \times (C_{eu; f i j j})^2 \\
&+ G_c^{ed} [lepton] \times S_c \times \mathcal{G}_c^{ed} \times [Y_d Y_d^\dagger]_{jj} \times (C_{ed; f i j j})^2 \\
&+ G_c^{ee} \times S_c \times \mathcal{G}_c^{ee} \times [Y_e Y_e^\dagger]_{jj} \times (C_{ee; f i j j})^2 \\
&+ G_d^{ee} \times S_d \times \mathcal{G}_d^{ee} \times [Y_e Y_e^\dagger]_{jj} \times (C_{ee; f j j i})^2
\end{aligned}$$

$$\begin{aligned}
(4\pi)^2 \mu \frac{d}{d\mu} C_{u^4 H^2; f i f i} &= G_c^{\ell u} [quark] \times S_c \times \mathcal{G}_c^{\ell u} \times [Y_e^\dagger Y_e]_{jj} \times (C_{\ell u; j j f i})^2 & (25) \\
&+ G_c^{qu(1)} [uu] \times S_c \times \mathcal{G}_c^{qu(1)} \times ([Y_u^\dagger Y_u]_{jj} + [Y_d^\dagger Y_d]_{jj}) \times (C_{qu; j j f i}^{(1)})^2 \\
&+ G_c^{qu(8)} [uu] \times S_c \times \mathcal{G}_c^{qu(8)} \times ([Y_u^\dagger Y_u]_{jj} + [Y_d^\dagger Y_d]_{jj}) \times (C_{qu; j j f i}^{(8)})^2 \\
&+ G_c^{eu} [quark] \times S_c \times \mathcal{G}_c^{eu} \times [Y_e Y_e^\dagger]_{jj} \times (C_{eu; j j f i})^2 \\
&+ G_c^{ud(1)} \times S_c \times \mathcal{G}_c^{ud(1)} \times [Y_d Y_d^\dagger]_{jj} \times (C_{ud; f i j j}^{(1)})^2 \\
&+ G_c^{ud(8)} \times S_c \times \mathcal{G}_c^{ud(8)} \times [Y_d Y_d^\dagger]_{jj} \times (C_{ud; f i j j}^{(8)})^2 \\
&+ G_c^{uu} \times S_c \times \mathcal{G}_c^{uu} \times [Y_u Y_u^\dagger]_{jj} \times (C_{uu; f i j j})^2 \\
&+ G_d^{uu} \times S_d \times \mathcal{G}_d^{uu} \times [Y_u Y_u^\dagger]_{jj} \times (C_{uu; f j j i})^2
\end{aligned}$$

$$\begin{aligned}
(4\pi)^2 \mu \frac{d}{d\mu} C_{d^4 H^2; f_i f_i} &= G_c^{\ell d} [quark] \times S_c \times \mathcal{G}_c^{\ell d} \times [Y_e^\dagger Y_e]_{jj} \times (C_{\ell d; j j f_i})^2 \\
&+ G_c^{qd(1)} [dd] \times S_c \times \mathcal{G}_c^{qd(1)} \times ([Y_u^\dagger Y_u]_{jj} + [Y_d^\dagger Y_d]_{jj}) \times (C_{qd; j j f_i}^{(1)})^2 \\
&+ G_c^{qd(8)} [dd] \times S_c \times \mathcal{G}_c^{qd(8)} \times ([Y_u^\dagger Y_u]_{jj} + [Y_d^\dagger Y_d]_{jj}) \times (C_{qd; j j f_i}^{(8)})^2 \\
&+ G_c^{ed} [quark] \times S_c \times \mathcal{G}_c^{ed} \times [Y_e Y_e^\dagger]_{jj} \times (C_{ed; j j f_i})^2 \\
&+ G_c^{ud(1)} \times S_c \times \mathcal{G}_c^{ud(1)} \times [Y_u Y_u^\dagger]_{jj} \times (C_{ud; j j f_i}^{(1)})^2 \\
&+ G_c^{ud(8)} \times S_c \times \mathcal{G}_c^{ud(8)} \times [Y_u Y_u^\dagger]_{jj} \times (C_{ud; j j f_i}^{(8)})^2 \\
&+ G_c^{dd} \times S_c \times \mathcal{G}_c^{dd} \times [Y_d Y_d^\dagger]_{jj} \times (C_{dd; f j j i})^2 \\
&+ G_d^{dd} \times S_d \times \mathcal{G}_d^{dd} \times [Y_d Y_d^\dagger]_{jj} \times (C_{dd; f j j i})^2
\end{aligned} \tag{26}$$

Note that double-insertions of  $Q_{\ell e q u}^{(3)}$  turn out to be finite. (This is the only case found for which the divergence in double-insertions at intermediate stages contains a term proportional to the Levi-Civita symbol.)

In the RG equations above:  $\mathcal{G}$  correspond to the residue of the  $\epsilon$ -pole extracted from the calculation indicated in Eq. (13) times the overall factor seen in Eq. (12) (without symmetry and group factors),  $S$  designate symmetry factors, and  $G$  group factors (subscripts  $a, b, c, d$  designate different topologies). They are given as follows:

$$\begin{aligned}
\mathcal{G}_a^{\ell e d q} &= \mathcal{G}_a^{\ell e q u(1)} = \mathcal{G}_a^{quqd(1)} = \mathcal{G}_a^{quqd(8)} = -4, \\
\mathcal{G}_b^{quqd(1)} &= \mathcal{G}_b^{quqd(8)} = 2, \\
\mathcal{G}_c^{\ell e} &= \mathcal{G}_c^{qx(1)} = \mathcal{G}_c^{qx(8)} = \mathcal{G}_c^{ee} = \mathcal{G}_c^{xx} = \mathcal{G}_c^{\ell\ell} = \mathcal{G}_c^{qq(1)} = \mathcal{G}_c^{qq(3)} \\
&= \mathcal{G}_c^{\ell q(1)} = \mathcal{G}_c^{\ell q(3)} = \mathcal{G}_c^{ex} = \mathcal{G}_c^{\ell x} = \mathcal{G}_c^{qe} = \mathcal{G}_c^{ud(1)} = \mathcal{G}_c^{ud(8)} = 2, \\
\mathcal{G}_b^{\ell e} &= \mathcal{G}_b^{qx(1)} = \mathcal{G}_b^{qx(8)} = -16, \\
\mathcal{G}_d^{\ell\ell} &= \mathcal{G}_d^{qq(1)} = \mathcal{G}_d^{qq(3)} = \mathcal{G}_d^{ee} = \mathcal{G}_d^{xx} = 2,
\end{aligned} \tag{27}$$

$$S_b = \frac{1}{2}, \quad S_a = \frac{1}{2}, \quad S_d = 1, \quad S_c = 1, \tag{28}$$

$$\begin{aligned}
G_d^{\ell\ell} [singlet] &= \frac{1}{2}, \quad G_d^{\ell\ell} [triplet(2)] = \frac{1}{2}, \quad G_c^{\ell\ell} = 1, \\
G_d^{qq(1)} [singlet] &= \frac{1}{2}, \quad G_d^{qq(1)} [triplet(2)] = \frac{1}{2}, \quad G_c^{qq(1)} = N_c, \\
G_d^{qq(3)} [singlet] &= \frac{5}{2}, \quad G_d^{qq(3)} [triplet(2)] = \frac{3}{2}, \quad G_c^{qq(3)} = N_c, \\
G_c^{\ell q(1)} [lepton] &= N_c, \quad G_c^{\ell q(1)} [quark] = 1, \\
G_c^{\ell q(3)} [lepton] &= N_c, \quad G_c^{\ell q(3)} [quark] = 1, \\
G_d^{ee} &= 1, \quad G_c^{ee} = 1, \\
G_d^{xx} &= 1, \quad G_c^{xx} = N_c, \\
G_c^{ex} [lepton] &= N_c, \quad G_c^{ex} [quark] = 1, \\
G_c^{ud(1)} &= N_c, \\
G_c^{ud(8)} &= \frac{1}{4} \left( 1 - \frac{1}{N_c} \right), \\
G_c^{\ell e} [\ell\ell] &= 1, \quad G_c^{\ell e} [ee] = 1, \\
G_c^{\ell x} [lepton] &= N_c, \quad G_c^{\ell x} [quark] = 1, \\
G_c^{qe} [lepton] &= N_c, \quad G_c^{qe} [quark] = 1, \\
G_c^{qx(1)} [qq] &= N_c, \quad G_c^{qx(1)} [xx] = N_c, \\
G_c^{qx(8)} [qq] &= \frac{1}{4} \left( 1 - \frac{1}{N_c} \right), \quad G_c^{qx(8)} [xx] = \frac{1}{4} \left( 1 - \frac{1}{N_c} \right),
\end{aligned} \tag{29}$$

and

$$\begin{aligned}
G_b^{\ell e} &= 1, \\
G_b^{qx(1)}[singlet] &= \frac{1}{N_c}, \quad G_b^{qx(1)}[octet] = 2, \\
G_b^{qx(8)}[singlet] &= \frac{1}{4} \left( N_c - \frac{2}{N_c} + \frac{1}{N_c^3} \right), \quad G_b^{qx(8)}[octet] = \frac{1}{2} \frac{1}{N_c^2}, \\
G_a^{\ell edq}[lepton] &= N_c, \quad G_a^{\ell edq}[quark] = 1, \\
G_b^{quqd(1)} &= 1, \quad G_a^{quqd(1)} = N_c, \\
G_b^{quqd(8)}[singlet] &= \frac{1}{4} \left( 1 - \frac{1}{N_c^2} \right), \quad G_b^{quqd(8)}[octet] = \frac{1}{2} \left( N_c - \frac{2}{N_c} \right), \quad G_a^{quqd(8)} = \frac{1}{2}, \\
G_a^{\ell equ(1)}[lepton] &= N_c, \quad G_a^{\ell equ(1)}[quark] = 1.
\end{aligned}$$

Everywhere in this paper  $T^A$  are the  $SU(3)$  generators, normalized such that  $\text{tr}\{T^A T^B\} = \frac{1}{2} \delta^{AB}$  (a different normalization was taken in [65]). Also,  $\tau^I$  are the Pauli matrices, for which  $\text{tr}\{\tau^I \tau^J\} = 2\delta^{IJ}$ .

As one cross-check, we verify the SM [36, 37] at the leading order.

## B Sensitivity of meson mixing to New Physics effects

Bounds on the size of NP in the neutral meson systems  $K$  and  $B_{(s)}$  are discussed in [51, 52] (see also [66, 55]). There, since these observables are used in the global extraction of the elements of the CKM matrix in the SM (the mass differences  $\Delta m_{d,s}$  playing presently a more relevant role in the fit than the indirect CP violating quantity  $|\epsilon_K|$ ), the extraction of the elements of the CKM matrix is redone allowing for NP contamination in the fit: NP is parametrized under the form

$$M_{12}^i = (M_{12}^i)_{\text{SM}} \times (1 + h_i \times e^{2i\sigma_i}), \quad (30)$$

and combined bounds on  $h_i$  and  $\sigma_i$  are extracted, where the index  $i$  refers to the different neutral meson systems. Ref. [51] combines contributions from NP in the kaon system with the top-up (case (I) discussed in Sect. 2) set of contributions from the SM, keeping cases (II) and (III) unmodified, which is consistent with our discussion in Sect. 4. The extracted ranges are  $h_d < 0.26$ ,  $h_s < 0.12$  [52], and  $|h_K \times \sin(2\sigma_K + 2\text{Arg}(V_{td}V_{ts}^*))| \lesssim 0.6$  [51]; all bounds are 95% Confidence Level intervals. These bounds constrain NP at the energy scale relevant for the different observables, namely,  $\sim M_{B_{(s)}}$  for the  $B_{(s)}$  mass differences, and 2 GeV for the indirect CP violating quantity in the kaon sector. To constrain different kinds of NP at the EW scale, one introduces the short-distance QCD corrections collected in [67], that provide the running and mixing of  $|\Delta F| = 2$  four-fermion contact interactions below the EW scale at the Next-to-Leading Order (NLO). One also needs the relevant bag parameters, which are taken from [48] in the  $K$  system, and [49] for the  $B_{(s)}$  systems (both for  $N_f = 2 + 1 + 1$ ). Light quark masses are taken from [68] ( $N_f = 2 + 1 + 1$ ), see also references therein. Put together, bounds on NP at the scale  $\mu_t = m_t(m_t) = 166$  GeV follow the following pattern for the Wilson coefficients  $C_1(i)$ ,  $C_2(i)$ ,  $C_3(i)$ ,  $C_4(i)$ ,  $C_5(i)$  introduced in the main text, see Eq. (3):

$$\begin{aligned}
\Delta m_{d,s} &: \quad 1 : 2 : 0.4 : 5 : 2, \\
|\epsilon_K| &: \quad 1 : 40 : 10 : 100 : 30
\end{aligned} \quad (31)$$

with respect to bounds on the Wilson coefficient  $C_1(i)$  (for instance, the bound on  $|C_4(B_{(s)})|$  is about 5 times stronger than the bound on  $|C_1(B_{(s)})|$ ); bounds on the latter are

$$\begin{aligned}
|C_1(B)| &< (1 \times 10^3 \text{ TeV})^{-2}, \\
|C_1(B_s)| &< (3 \times 10^2 \text{ TeV})^{-2}, \\
|\text{Im}\{C_1(K)\}| &< (2 \times 10^4 \text{ TeV})^{-2}.
\end{aligned} \quad (32)$$

The running effects between the NP scale  $\Lambda_{\text{NP}}$  and the EW scale  $\mu_{\text{EW}} = \mu_t$  are the concern of the main text. Although short-distance QCD effects below the EW scale are being included for NP, we are not including QCD effects above the EW scale, which are suppressed by a relatively small strong coupling.

## References

- [1] S. L. Glashow, J. Iliopoulos, and L. Maiani. Weak Interactions with Lepton-Hadron Symmetry. *Phys. Rev. D*, 2:1285–1292, 1970. doi:[10.1103/PhysRevD.2.1285](https://doi.org/10.1103/PhysRevD.2.1285).
- [2] Nicola Cabibbo. Unitary Symmetry and Leptonic Decays. *Phys. Rev. Lett.*, 10:531–533, 1963. doi:[10.1103/PhysRevLett.10.531](https://doi.org/10.1103/PhysRevLett.10.531).
- [3] Makoto Kobayashi and Toshihide Maskawa. CP Violation in the Renormalizable Theory of Weak Interaction. *Prog. Theor. Phys.*, 49:652–657, 1973. doi:[10.1143/PTP.49.652](https://doi.org/10.1143/PTP.49.652).
- [4] J. Charles et al. Current status of the Standard Model CKM fit and constraints on  $\Delta F = 2$  New Physics. *Phys. Rev. D*, 91(7):073007, 2015. arXiv:[1501.05013](https://arxiv.org/abs/1501.05013), doi:[10.1103/PhysRevD.91.073007](https://doi.org/10.1103/PhysRevD.91.073007).
- [5] M. Bona et al. The Unitarity Triangle Fit in the Standard Model and Hadronic Parameters from Lattice QCD: A Reappraisal after the Measurements of Delta m(s) and BR(B  $\rightarrow$  tau nu(tau)). *JHEP*, 10:081, 2006. arXiv:[hep-ph/0606167](https://arxiv.org/abs/hep-ph/0606167), doi:[10.1088/1126-6708/2006/10/081](https://doi.org/10.1088/1126-6708/2006/10/081).
- [6] Christopher W. Murphy. Dimension-8 operators in the Standard Model Effective Field Theory. *JHEP*, 10:174, 2020. arXiv:[2005.00059](https://arxiv.org/abs/2005.00059), doi:[10.1007/JHEP10\(2020\)174](https://doi.org/10.1007/JHEP10(2020)174).
- [7] B. Grzadkowski, M. Iskrzynski, M. Misiak, and J. Rosiek. Dimension-Six Terms in the Standard Model Lagrangian. *JHEP*, 10:085, 2010. arXiv:[1008.4884](https://arxiv.org/abs/1008.4884), doi:[10.1007/JHEP10\(2010\)085](https://doi.org/10.1007/JHEP10(2010)085).
- [8] Hao-Lin Li, Zhe Ren, Jing Shu, Ming-Lei Xiao, Jiang-Hao Yu, and Yu-Hui Zheng. Complete set of dimension-eight operators in the standard model effective field theory. *Phys. Rev. D*, 104(1):015026, 2021. arXiv:[2005.00008](https://arxiv.org/abs/2005.00008), doi:[10.1103/PhysRevD.104.015026](https://doi.org/10.1103/PhysRevD.104.015026).
- [9] Mikael Chala, Claudius Krause, and Germano Nardini. Signals of the electroweak phase transition at colliders and gravitational wave observatories. *JHEP*, 07:062, 2018. arXiv:[1802.02168](https://arxiv.org/abs/1802.02168), doi:[10.1007/JHEP07\(2018\)062](https://doi.org/10.1007/JHEP07(2018)062).
- [10] Chris Hays, Adam Martin, Verónica Sanz, and Jack Setford. On the impact of dimension-eight SMEFT operators on Higgs measurements. *JHEP*, 02:123, 2019. arXiv:[1808.00442](https://arxiv.org/abs/1808.00442), doi:[10.1007/JHEP02\(2019\)123](https://doi.org/10.1007/JHEP02(2019)123).
- [11] Chris Hays, Andreas Helset, Adam Martin, and Michael Trott. Exact SMEFT formulation and expansion to  $\mathcal{O}(v^4/\Lambda^4)$ . *JHEP*, 11:087, 2020. arXiv:[2007.00565](https://arxiv.org/abs/2007.00565), doi:[10.1007/JHEP11\(2020\)087](https://doi.org/10.1007/JHEP11(2020)087).
- [12] Tyler Corbett. The Feynman rules for the SMEFT in the background field gauge. *JHEP*, 03:001, 2021. arXiv:[2010.15852](https://arxiv.org/abs/2010.15852), doi:[10.1007/JHEP03\(2021\)001](https://doi.org/10.1007/JHEP03(2021)001).
- [13] Tyler Corbett, Andreas Helset, Adam Martin, and Michael Trott. EWPD in the SMEFT to dimension eight. *JHEP*, 06:076, 2021. arXiv:[2102.02819](https://arxiv.org/abs/2102.02819), doi:[10.1007/JHEP06\(2021\)076](https://doi.org/10.1007/JHEP06(2021)076).
- [14] Michael Trott. Methodology for theory uncertainties in the standard model effective field theory. *Phys. Rev. D*, 104(9):095023, 2021. arXiv:[2106.13794](https://arxiv.org/abs/2106.13794), doi:[10.1103/PhysRevD.104.095023](https://doi.org/10.1103/PhysRevD.104.095023).
- [15] Tyler Corbett, Adam Martin, and Michael Trott. Consistent higher order  $\sigma(\mathcal{G}\mathcal{G} \rightarrow h)$ ,  $\Gamma(h \rightarrow \mathcal{G}\mathcal{G})$  and  $\Gamma(h \rightarrow \gamma\gamma)$  in geoSMEFT. 7 2021. arXiv:[2107.07470](https://arxiv.org/abs/2107.07470).
- [16] Adam Martin and Michael Trott. The  $ggh$  variations. 9 2021. arXiv:[2109.05595](https://arxiv.org/abs/2109.05595).
- [17] Sally Dawson, Samuel Homiller, and Matthew Sullivan. Impact of dimension-eight SMEFT contributions: A case study. *Phys. Rev. D*, 104(11):115013, 2021. arXiv:[2110.06929](https://arxiv.org/abs/2110.06929), doi:[10.1103/PhysRevD.104.115013](https://doi.org/10.1103/PhysRevD.104.115013).
- [18] Simone Alioli, Radja Boughezal, Emanuele Mereghetti, and Frank Petriello. Novel angular dependence in Drell-Yan lepton production via dimension-8 operators. *Phys. Lett. B*, 809:135703, 2020. arXiv:[2003.11615](https://arxiv.org/abs/2003.11615), doi:[10.1016/j.physletb.2020.135703](https://doi.org/10.1016/j.physletb.2020.135703).
- [19] Radja Boughezal, Emanuele Mereghetti, and Frank Petriello. Dilepton production in the SMEFT at  $\mathcal{O}(1/\Lambda^4)$ . *Phys. Rev. D*, 104(9):095022, 2021. arXiv:[2106.05337](https://arxiv.org/abs/2106.05337), doi:[10.1103/PhysRevD.104.095022](https://doi.org/10.1103/PhysRevD.104.095022).
- [20] Marco Ardu and Sacha Davidson. What is Leading Order for LFV in SMEFT? *JHEP*, 08:002, 2021. arXiv:[2103.07212](https://arxiv.org/abs/2103.07212), doi:[10.1007/JHEP08\(2021\)002](https://doi.org/10.1007/JHEP08(2021)002).
- [21] Grant N. Remmen and Nicholas L. Rodd. Consistency of the Standard Model Effective Field Theory. *JHEP*, 12:032, 2019. arXiv:[1908.09845](https://arxiv.org/abs/1908.09845), doi:[10.1007/JHEP12\(2019\)032](https://doi.org/10.1007/JHEP12(2019)032).

- [22] Grant N. Remmen and Nicholas L. Rodd. Flavor Constraints from Unitarity and Analyticity. *Phys. Rev. Lett.*, 125(8):081601, 2020. [Erratum: *Phys.Rev.Lett.* 127, 149901 (2021)]. [arXiv:2004.02885](#), [doi:10.1103/PhysRevLett.127.149901](#).
- [23] Jiayin Gu, Lian-Tao Wang, and Cen Zhang. An unambiguous test of positivity at lepton colliders. 11 2020. [arXiv:2011.03055](#).
- [24] Quentin Bonnefoy, Emanuele Gendy, and Christophe Grojean. Positivity bounds on Minimal Flavor Violation. *JHEP*, 04:115, 2021. [arXiv:2011.12855](#), [doi:10.1007/JHEP04\(2021\)115](#).
- [25] John Ellis, Shao-Feng Ge, Hong-Jian He, and Rui-Qing Xiao. Probing the scale of new physics in the  $ZZ\gamma$  coupling at  $e^+e^-$  colliders. *Chin. Phys. C*, 44(6):063106, 2020. [arXiv:1902.06631](#), [doi:10.1088/1674-1137/44/6/063106](#).
- [26] John Ellis, Hong-Jian He, and Rui-Qing Xiao. Probing new physics in dimension-8 neutral gauge couplings at  $e^+e^-$  colliders. *Sci. China Phys. Mech. Astron.*, 64(2):221062, 2021. [arXiv:2008.04298](#), [doi:10.1007/s11433-020-1617-3](#).
- [27] Sacha Davidson, Martin Gorbahn, and Matthew Leak. Majorana neutrino masses in the renormalization group equations for lepton flavor violation. *Phys. Rev.*, D98(9):095014, 2018. [arXiv:1807.04283](#), [doi:10.1103/PhysRevD.98.095014](#).
- [28] Elizabeth E. Jenkins, Aneesh V. Manohar, and Peter Stoffer. Low-Energy Effective Field Theory below the Electroweak Scale: Anomalous Dimensions. *JHEP*, 01:084, 2018. [arXiv:1711.05270](#), [doi:10.1007/JHEP01\(2018\)084](#).
- [29] Mikael Chala and Arsenii Titov. Neutrino masses in the Standard Model effective field theory. *Phys. Rev. D*, 104(3):035002, 2021. [arXiv:2104.08248](#), [doi:10.1103/PhysRevD.104.035002](#).
- [30] Mikael Chala, Guilherme Guedes, Maria Ramos, and Jose Santiago. Towards the renormalisation of the Standard Model effective field theory to dimension eight: Bosonic interactions I. *SciPost Phys.*, 11:065, 2021. [arXiv:2106.05291](#), [doi:10.21468/SciPostPhys.11.3.065](#).
- [31] J. Urban, F. Krauss, U. Jentschura, and G. Soff. Next-to-leading order QCD corrections for the  $B_0$  anti- $B_0$  mixing with an extended Higgs sector. *Nucl. Phys. B*, 523:40–58, 1998. [arXiv:hep-ph/9710245](#), [doi:10.1016/S0550-3213\(98\)00043-1](#).
- [32] Marco Ciuchini et al. Delta M(K) and epsilon(K) in SUSY at the next-to-leading order. *JHEP*, 10:008, 1998. [arXiv:hep-ph/9808328](#), [doi:10.1088/1126-6708/1998/10/008](#).
- [33] Véronique Bernard, Sébastien Descotes-Genon, and Luiz Vale Silva. Short-distance QCD corrections to  $K^0\bar{K}^0$  mixing at next-to-leading order in Left-Right models. *JHEP*, 08:128, 2016. [arXiv:1512.00543](#), [doi:10.1007/JHEP08\(2016\)128](#).
- [34] Andreas Crivellin, Jordi Folch Eguren, and Javier Virto. Next-to-Leading-Order QCD Matching for  $\Delta F = 2$  Processes in Scalar Leptoquark Models. 9 2021. [arXiv:2109.13600](#).
- [35] Frederick J. Gilman and Mark B. Wise.  $K_0$  anti- $K_0$  Mixing in the Six Quark Model. *Phys. Rev. D*, 27:1128, 1983. [doi:10.1103/PhysRevD.27.1128](#).
- [36] Stefan Herrlich and Ulrich Nierste. Enhancement of the K(L) - K(S) mass difference by short distance QCD corrections beyond leading logarithms. *Nucl. Phys. B*, 419:292–322, 1994. [arXiv:hep-ph/9310311](#), [doi:10.1016/0550-3213\(94\)90044-2](#).
- [37] Stefan Herrlich and Ulrich Nierste. The Complete  $-\Delta S = 2$  - Hamiltonian in the next-to-leading order. *Nucl. Phys. B*, 476:27–88, 1996. [arXiv:hep-ph/9604330](#), [doi:10.1016/0550-3213\(96\)00324-0](#).
- [38] Joachim Brod, Martin Gorbahn, and Emmanuel Stamou. Standard-Model Prediction of  $\epsilon_K$  with Manifest Quark-Mixing Unitarity. *Phys. Rev. Lett.*, 125(17):171803, 2020. [arXiv:1911.06822](#), [doi:10.1103/PhysRevLett.125.171803](#).
- [39] Joachim Brod and Martin Gorbahn.  $\epsilon_K$  at Next-to-Next-to-Leading Order: The Charm-Top-Quark Contribution. *Phys. Rev. D*, 82:094026, 2010. [arXiv:1007.0684](#), [doi:10.1103/PhysRevD.82.094026](#).
- [40] Joachim Brod and Martin Gorbahn. Next-to-Next-to-Leading-Order Charm-Quark Contribution to the  $CP$  Violation Parameter  $\epsilon_K$  and  $\Delta M_K$ . *Phys. Rev. Lett.*, 108:121801, 2012. [arXiv:1108.2036](#), [doi:10.1103/PhysRevLett.108.121801](#).
- [41] Joachim Brod, Sandra Kvedaraitė, and Zachary Polonsky. Two-loop Electroweak Corrections to the Top-Quark Contribution to  $\epsilon_K$ . 7 2021. [arXiv:2108.00017](#).
- [42] M. Ciuchini, E. Franco, V. Lubicz, G. Martinelli, L. Silvestrini, and C. Tarantino. Power corrections to the CP-violation parameter  $\epsilon_K$ . 11 2021. [arXiv:2111.05153](#).

- [43] Motoi Endo, Teppei Kitahara, and Daiki Ueda. SMEFT top-quark effects on  $\Delta F = 2$  observables. *JHEP*, 07:182, 2019. [arXiv:1811.04961](#), [doi:10.1007/JHEP07\(2019\)182](#).
- [44] Giampiero Passarino. Field reparametrization in effective field theories. *Eur. Phys. J. Plus*, 132(1):16, 2017. [arXiv:1610.09618](#), [doi:10.1140/epjp/i2017-11291-5](#).
- [45] Giampiero Passarino. XEFT, the challenging path up the hill: dim = 6 and dim = 8. 1 2019. [arXiv:1901.04177](#).
- [46] J. C. Criado and M. Pérez-Victoria. Field redefinitions in effective theories at higher orders. *JHEP*, 03:038, 2019. [arXiv:1811.09413](#), [doi:10.1007/JHEP03\(2019\)038](#).
- [47] Abdurrahman Barzinji, Michael Trott, and Anagha Vasudevan. Equations of Motion for the Standard Model Effective Field Theory: Theory and Applications. *Phys. Rev.*, D98(11):116005, 2018. [arXiv:1806.06354](#), [doi:10.1103/PhysRevD.98.116005](#).
- [48] N. Carrasco, P. Dimopoulos, R. Frezzotti, V. Lubicz, G. C Rossi, S. Simula, and C. Tarantino.  $\Delta S=2$  and  $\Delta C=2$  bag parameters in the standard model and beyond from  $N_f=2+1+1$  twisted-mass lattice QCD. *Phys. Rev. D*, 92(3):034516, 2015. [arXiv:1505.06639](#), [doi:10.1103/PhysRevD.92.034516](#).
- [49] R. J. Dowdall, C. T. H. Davies, R. R. Horgan, G. P. Lepage, C. J. Monahan, J. Shigemitsu, and M. Wingate. Neutral B-meson mixing from full lattice QCD at the physical point. *Phys. Rev. D*, 100(9):094508, 2019. [arXiv:1907.01025](#), [doi:10.1103/PhysRevD.100.094508](#).
- [50] Renae Conlin and Alexey A. Petrov. Muonium-antimuonium oscillations in effective field theory. *Phys. Rev. D*, 102(9):095001, 2020. [arXiv:2005.10276](#), [doi:10.1103/PhysRevD.102.095001](#).
- [51] Jérôme Charles, Sebastien Descotes-Genon, Zoltan Ligeti, Stéphane Monteil, Michele Papucci, and Karim Trabelsi. Future sensitivity to new physics in  $B_d, B_s$ , and K mixings. *Phys. Rev. D*, 89(3):033016, 2014. [arXiv:1309.2293](#), [doi:10.1103/PhysRevD.89.033016](#).
- [52] Jérôme Charles, Sébastien Descotes-Genon, Zoltan Ligeti, Stéphane Monteil, Michele Papucci, Karim Trabelsi, and Luiz Vale Silva. New physics in  $B$  meson mixing: future sensitivity and limitations. *Phys. Rev. D*, 102(5):056023, 2020. [arXiv:2006.04824](#), [doi:10.1103/PhysRevD.102.056023](#).
- [53] A. Lenz, U. Nierste, J. Charles, S. Descotes-Genon, A. Jantsch, C. Kaufhold, H. Lacker, S. Monteil, V. Niess, and S. T’Jampens. Anatomy of New Physics in  $B - \bar{B}$  mixing. *Phys. Rev. D*, 83:036004, 2011. [arXiv:1008.1593](#), [doi:10.1103/PhysRevD.83.036004](#).
- [54] A. Lenz, U. Nierste, J. Charles, S. Descotes-Genon, H. Lacker, S. Monteil, V. Niess, and S. T’Jampens. Constraints on new physics in  $B - \bar{B}$  mixing in the light of recent LHCb data. *Phys. Rev. D*, 86:033008, 2012. [arXiv:1203.0238](#), [doi:10.1103/PhysRevD.86.033008](#).
- [55] Sébastien Descotes-Genon, Adam Falkowski, Marco Fedele, Martín González-Alonso, and Javier Virto. The CKM parameters in the SMEFT. *JHEP*, 05:172, 2019. [arXiv:1812.08163](#), [doi:10.1007/JHEP05\(2019\)172](#).
- [56] Luca Silvestrini and Mauro Valli. Model-independent Bounds on the Standard Model Effective Theory from Flavour Physics. *Phys. Lett. B*, 799:135062, 2019. [arXiv:1812.10913](#), [doi:10.1016/j.physletb.2019.135062](#).
- [57] Tobias Hurth, Sophie Renner, and William Shepherd. Matching for FCNC effects in the flavour-symmetric SMEFT. *JHEP*, 06:029, 2019. [arXiv:1903.00500](#), [doi:10.1007/JHEP06\(2019\)029](#).
- [58] Jason Aebischer, Christoph Bobeth, Andrzej J. Buras, and Jacky Kumar. SMEFT ATLAS of  $\Delta F = 2$  transitions. *JHEP*, 12:187, 2020. [arXiv:2009.07276](#), [doi:10.1007/JHEP12\(2020\)187](#).
- [59] Elizabeth E. Jenkins, Aneesh V. Manohar, and Michael Trott. Renormalization Group Evolution of the Standard Model Dimension Six Operators I: Formalism and lambda Dependence. *JHEP*, 10:087, 2013. [arXiv:1308.2627](#), [doi:10.1007/JHEP10\(2013\)087](#).
- [60] Elizabeth E. Jenkins, Aneesh V. Manohar, and Michael Trott. Renormalization Group Evolution of the Standard Model Dimension Six Operators II: Yukawa Dependence. *JHEP*, 01:035, 2014. [arXiv:1310.4838](#), [doi:10.1007/JHEP01\(2014\)035](#).
- [61] Rodrigo Alonso, Elizabeth E. Jenkins, Aneesh V. Manohar, and Michael Trott. Renormalization Group Evolution of the Standard Model Dimension Six Operators III: Gauge Coupling Dependence and Phenomenology. *JHEP*, 04:159, 2014. [arXiv:1312.2014](#), [doi:10.1007/JHEP04\(2014\)159](#).



- [62] P.A. Zyla et al. Review of Particle Physics. *PTEP*, 2020(8):083C01, 2020. [doi:10.1093/ptep/ptaa104](https://doi.org/10.1093/ptep/ptaa104).
- [63] Konstantin G. Chetyrkin, Mikolaj Misiak, and Manfred Munz. Beta functions and anomalous dimensions up to three loops. *Nucl. Phys. B*, 518:473–494, 1998. [arXiv:hep-ph/9711266](https://arxiv.org/abs/hep-ph/9711266), [doi:10.1016/S0550-3213\(98\)00122-9](https://doi.org/10.1016/S0550-3213(98)00122-9).
- [64] H. Simma. Equations of motion for effective Lagrangians and penguins in rare B decays. *Z. Phys. C*, 61:67–82, 1994. [arXiv:hep-ph/9307274](https://arxiv.org/abs/hep-ph/9307274), [doi:10.1007/BF01641888](https://doi.org/10.1007/BF01641888).
- [65] Luiz Vale Silva. Probing squared four-fermion operators of SMEFT with meson-mixing. *PoS, EPS-HEP2021*:519.
- [66] M. Bona et al. Model-independent constraints on  $\Delta F = 2$  operators and the scale of new physics. *JHEP*, 03:049, 2008. [arXiv:0707.0636](https://arxiv.org/abs/0707.0636), [doi:10.1088/1126-6708/2008/03/049](https://doi.org/10.1088/1126-6708/2008/03/049).
- [67] Andrzej J. Buras, Sebastian Jager, and Jorg Urban. Master formulae for Delta F=2 NLO QCD factors in the standard model and beyond. *Nucl. Phys. B*, 605:600–624, 2001. [arXiv:hep-ph/0102316](https://arxiv.org/abs/hep-ph/0102316), [doi:10.1016/S0550-3213\(01\)00207-3](https://doi.org/10.1016/S0550-3213(01)00207-3).
- [68] S. Aoki et al. FLAG Review 2019: Flavour Lattice Averaging Group (FLAG). *Eur. Phys. J. C*, 80(2):113, 2020. [arXiv:1902.08191](https://arxiv.org/abs/1902.08191), [doi:10.1140/epjc/s10052-019-7354-7](https://doi.org/10.1140/epjc/s10052-019-7354-7).