

Dual Seminorms, Ergodic Coefficients and Semicontraction Theory

Giulia De Pasquale *Student Member, IEEE*, Kevin D. Smith *Student Member, IEEE*, Francesco Bullo *Fellow, IEEE*, M. Elena Valcher *Fellow, IEEE*

Abstract—Dynamical systems that are contracting on a subspace are said to be semicontracting. Semicontraction theory is a useful tool in the study of consensus algorithms and dynamical flow systems such as Markov chains.

To develop a comprehensive theory of semicontracting systems, we investigate seminorms on vector spaces and define two canonical notions: projection and distance seminorms. We show that the well-known ℓ_p ergodic coefficients are induced matrix seminorms and play a central role in stability problems. In particular, we formulate a duality theorem that explains why the Markov-Dobrushin coefficient is the rate of contraction for both averaging and conservation flows in discrete time. Moreover, we obtain parallel results for induced matrix log seminorms. Finally, we propose comprehensive theorems for strong semicontractivity of linear and non-linear time-varying dynamical systems with invariance and conservation properties both in discrete and continuous time.

Index Terms—Semicontraction theory, ergodic coefficients, induced matrix seminorm, logarithmic norm, duality.

I. INTRODUCTION

Problem description and motivation

Before Stefan Banach proved his famous contraction principle in 1922 [2], Andrey Markov started in 1906 [25] the study of stochastic processes. As documented by Eugene Seneta [31], Markov established a key contraction inequality and a corresponding contraction factor now known with the name of *ergodic coefficient* of a Markov chain. This paper aims to provide a modern semicontraction theory approach to explain and generalize ergodic coefficients.

To be concrete, let the matrix A be row-stochastic and consider the discrete-time dynamical systems

$$x(k+1) = Ax(k), \quad (1a)$$

$$\pi(k+1) = A^\top \pi(k). \quad (1b)$$

Similarly, let L be a Laplacian matrix and consider the continuous-time counterparts:

$$\dot{x}(t) = -Lx(t), \quad \dot{\pi}(t) = -L^\top \pi(t). \quad (2)$$

Submitted on 2022/12/22. This work was supported in part by AFOSR grant FA9550-22-1-0059 and by Fondazione Ing. Aldo Gini.

Giulia De Pasquale and Maria Elena Valcher are with the Dipartimento di Ingegneria dell'Informazione Università di Padova, Padova, 35131, Italy, {giulia.depasquale, meme}@dei.unipd.it.

Kevin D. Smith and Francesco Bullo are with the Center for Control, Dynamical Systems, and Computation, UC Santa Barbara, Santa Barbara, CA 93101 USA. {kevinsmith, bullo}@ucsb.edu.

These systems are perhaps the simplest examples of general averaging-based dynamics (e.g., robotic coordination and distributed optimization) and dynamical flow systems (e.g., compartmental and traffic systems). Important generalizations include systems of the form $\dot{x} = f(t, x)$, where f satisfies invariance properties (generalizing $A\mathbb{1}_n = \mathbb{1}_n$) or conservation properties (generalizing $\mathbb{1}_n^\top A^\top = \mathbb{1}_n^\top$); in all these (linear and nonlinear) cases, the system is at most marginally stable.

Markov and later scientists essentially showed that, under a certain connectivity assumption, maps of the form $\pi \mapsto A^\top \pi$ are contraction maps with respect to the total variation distance on the simplex. To be specific, define the simplex $\Delta_n = \{x \in \mathbb{R}^n \mid x \geq 0, \mathbb{1}_n^\top x = 1\}$ and the total variation distance on Δ_n by $d_{TV}(\pi, \sigma) = \frac{1}{2} \sum_i |\pi_i - \sigma_i|$. Then any two solutions $\pi(k), \sigma(k)$ to (1b) satisfy

$$d_{TV}(\pi(k) - \sigma(k)) \leq \tau_1(A)^k d_{TV}(\pi(0) - \sigma(0)), \quad (3)$$

where $\tau_1(A)$ is the so-called *Markov-Dobrushin ergodic coefficient* defined by

$$\tau_1(A) := \max_{\|z\|_1=1, z^\top \mathbb{1}_n=0} \|A^\top z\|_1. \quad (4)$$

In short, when $\tau_1(A) < 1$, existence, uniqueness and global exponential stability of an equilibrium $\pi^* \in \Delta_n$ for system (1b) is ensured.

Now comes a remarkable similarity. If one defines the seminorm $\|x\|_{\text{dist}, \infty} = \frac{1}{2} (\max_i \{x_i\} - \min_j \{x_j\})$, the following fact is also known [13, Theorem 1.1] about averaging systems of the form (1a):

$$\|x(k)\|_{\text{dist}, \infty} \leq \tau_1(A)^k \|x(0)\|_{\text{dist}, \infty}. \quad (5)$$

Despite the extensive research in this field, numerous known related facts remain somehow mysterious and numerous related mathematical questions remain open. For example, why is the same ergodic coefficient τ_1 relevant for the contraction properties of both dynamical flow systems and averaging systems? And is it the tightest such bound? How does one generalize the bounds (3) and (5) to ergodic coefficients τ_p defined with respect to arbitrary ℓ_p norms (instead of the ℓ_1 norm in (4))? How does one provide a unified robust stability analysis for both systems? What are the canonical Lyapunov functions for both systems (1a)-(1b), whose discrete-time variation along the flow is described by $\tau_p(A)$? How does one define ergodic coefficients for continuous-time systems? Is there a contraction theoretic framework that applies to time-varying and nonlinear systems with generalized invariance or conservation properties?

Contributions

This paper provides a comprehensive answer to all the open research questions outlined above.

In order to define Lyapunov functions for averaging, flow systems and their generalizations to nonlinear dynamical systems with invariant subspaces, we study seminorms, induced matrix seminorms for discrete time systems and logarithmic seminorms for continuous-time systems. A key contribution of this paper is to explain precisely in what sense ergodic coefficients are *induced matrix seminorms* and, when less than unity, contraction factors for discrete-time systems. This equality is the fundamental reason why ergodic coefficients play a critical role in robust stability theory for discrete-time dynamical systems with invariance properties. It is surprising that induced norms are widely studied in the matrix theory literature, but induced seminorms much less (e.g., see [15]).

After characterizing various seminorms' properties, we define two canonical sets of seminorms, namely, *distance* and *projection seminorms*, and establish remarkable duality properties between the two. Our first result generalizes and strengthens the so called *Markov contraction inequality* as a duality result between the aforementioned seminorms. Our duality result precisely explains why the induced matrix seminorms for both A and A^T are identical, when computed with respect to dual seminorms. Particular emphasis is given to the case of *consensus seminorms*, that is, seminorms whose kernel is the consensus space (i.e., seminorms that are positive definite about the consensus space). Consensus seminorms appear naturally in averaging algorithms and surprisingly in systems with conservation property (such as Markov chains and dynamical flow systems).

It is an elementary algebraic observation that the total variation distance on the simplex arises from the restriction of the ℓ_1 projection consensus seminorm.

We then leverage all these notions to provide a general nonlinear semicontraction theory, grounded in two key theorems both for continuous and discrete time varying dynamical systems. The semicontraction theory we develop is tailored to systems with invariance or conservation properties. More in detail, when either the system's Jacobian leaves invariant the seminorm kernel (invariance property) or its orthogonal complement (conservation property), there is a well defined notion of *perpendicular dynamics* which is strictly contracting. For both systems, in the linear time varying case, we show how canonical Lyapunov functions (some of which partly known in the literature) naturally arise from seminorms. For the non-linear case, our first key theorem establishes conditions and features of strong semicontracting continuous time, time varying systems that enjoy the invariance property. The theorem extends Theorem 13 in [17] through the formulation of a cascade decomposition and by establishing a strong contractivity property on the orthogonal complement to the seminorm kernel. The second key theorem is entirely novel and pertains semicontraction conditions for continuous time, time varying, dynamical systems that enjoy the conservation property. A discrete time version of these two theorems is also provided.

Literature review

Interest in contractivity of dynamical systems via matrix measures can be traced back to Demidovič [7] and Krasovskii [20]. Logarithmic norms have been exploited in control theory later on by Desoer and Vidyasagar in [8] and applied in the study of contraction theory for dynamical systems for the first time by Lohmiller and Slotine [24]. In the context of control theory, this literature inspired many generalizations of contraction theory such as partial contraction [36], weak- and semi-contraction [17], horizontal contraction on Riemannian and Finsler manifolds [12], [32], etc.

In particular, partial contraction refers to convergence of systems trajectories to a specific behavior, or a manifold [33], see also [9] for a survey on this theory. While partial contraction establishes convergence to a manifold, semicontraction ensures contractivity on the subspace perpendicular to the kernel of the seminorm. For a characterization of partial contraction in the ℓ_2 -norm for the study of synchronization in networked systems, see [36]. The notion of partial contraction is closely related to the one of semicontraction and weak contraction proposed and investigated in [17]. Semicontraction theory relies on a relaxed concept of matrix measure, known as matrix semimeasure. For this reason, contractivity of a dynamical system is only ensured on a certain subspace and the distance between trajectories is allowed to increase along certain directions.

A relevant behavior, to which semicontraction theory applies, is the one of consensus for dynamical systems. Strictly related to consensus, when it comes to stochastic systems, is the concept of (weak) ergodicity [34]. The concept of weak ergodicity was first formalized in 1931 by Kolmogorov [18], who stated that a sequence of stochastic matrices is *weakly ergodic* if the rows of the matrix product tend to become identical as the number of factors increases. The study of ergodicity coefficients is traced back to the pioneering work of Markov [25], in 1906, in which a first expression of ergodicity coefficients was provided in the context of the Weak Law of Large Numbers. Subsequent works from Doeblin [11] and Dobrushin [10] provided conditions for weak ergodicity. The key results in this research area were extended and then reviewed by Seneta in the 80's, see, e.g., [29]. A survey of ergodicity coefficients is given by Ipsen and Selee [16], a historical discussion is given by Hartfiel [13, Chapter 1], and a recent treatment on their connection with spectral graph theory is given by Marsli and Hall [26]. A characterization of "convergability" [23], namely the convergence of a product of an infinite number of stochastic matrices, is studied by Liu et. al in [23], where a different approach, based on optimally deflated matrices, is proposed. Despite the evident relation between ergodicity coefficients, contraction factors and induced matrix seminorms, especially in the context of stochastic and averaging systems [1], to the best of our knowledge none in the past has shed full light on their connections (see [6] for some preliminary work in this direction). This manuscript aims to bridge the existing gap in the scientific literature between semicontraction and ergodicity of dynamical systems.

Paper organization

Section II presents notation and preliminary results. Section III introduces the projection and distance seminorms and establishes their duality relationship. Section IV pertains with induced matrix seminorms and induced matrix log-seminorms. In Section V semicontraction theory is applied to dynamical systems. Finally, Section VII concludes the manuscript.

All theorems in this manuscript are new. Lemmas and Corollaries are either new or simple derivations from known results. This manuscript extends the submitted version in the IEEE Transaction on Automatic Control and includes the proofs of Lemma 13 and Theorem 22, explicit expressions for projection seminorms of columns stochastic matrices (17)-(19), Corollary 24, the property (ii) from Lemma 17, and Remark 3.

II. NOTATION AND PRELIMINARIES

A. Notation

The set $\mathbb{R}_{\geq 0}$ is the set of nonnegative real numbers. Let $I_n \in \mathbb{R}^{n \times n}$ denote the identity matrix of size n . Let $\mathbf{1}_n$ and $\mathbf{0}_n$ denote the n dimensional column vectors all whose entries equal 1 and 0, respectively. Let \mathbf{e}_i denote the i -th vector of the canonical basis in \mathbb{R}^n . For a matrix $A \in \mathbb{R}^{n \times n}$, let A^T denote its transpose, $[A]_{i,j}$ its (i,j) th entry. The matrix A is *nonnegative* if all its entries are nonnegative, it is *row stochastic* if it is nonnegative and $A\mathbf{1}_n = \mathbf{1}_n$, it is *column stochastic* if A^T is row stochastic.

Given $A \in \mathbb{R}^{n \times n}$, a vector subspace $\mathcal{K} \subseteq \mathbb{R}^n$ is *A-invariant* if $AK \subseteq \mathcal{K}$. The symbol $\langle \cdot, \cdot \rangle : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}$ denotes the standard inner product on \mathbb{R}^n . We let Π_{\perp} denote the orthogonal projection matrix onto \mathcal{K}^{\perp} , where the symbol \mathcal{K}^{\perp} denotes the orthogonal complement of \mathcal{K} . Note that $\Pi_{\perp} = \Pi_{\perp}^T$, and if $\mathcal{K} = \text{span}\{\mathbf{1}_n\}$, then $\Pi_{\perp} = I_n - \mathbf{1}_n \mathbf{1}_n^T / n =: \Pi_n$. Given $x \in \mathbb{R}^n$, the perpendicular and parallel components of x to \mathcal{K} are denoted by $x_{\perp} = \Pi_{\perp} x$ and $x_{\parallel} = (I_n - \Pi_{\perp})x$, respectively. Define the n -simplex as $\Delta_n = \{v \in \mathbb{R}_{\geq 0}^n \mid \mathbf{1}_n^T v = 1\}$ and the *sign function*, $\text{sign} : \mathbb{R} \rightarrow \{-1, 0, 1\}$, as $\text{sign}(x) = \frac{x}{|x|}$ if $x \neq 0$, and $\text{sign}(0) = 0$. Given two matrices $A, B \in \mathbb{R}^{n \times n}$ we use the notation $A \preceq B$ to indicate that $A - B$ is a negative semidefinite matrix.

A directed, weighted graph is a triple [5], $\mathcal{G} = (\mathcal{V}, \mathcal{E}, \mathcal{A})$, where $\mathcal{V} = \{1, \dots, n\}$ is the set of vertices, $\mathcal{E} \subseteq \mathcal{V} \times \mathcal{V}$ is the set of arcs and \mathcal{A} is the adjacency matrix. An arc (i, j) belongs to \mathcal{G} if and only if $[A]_{ij} \neq 0$. Two nodes $i, j \in \mathcal{V}$ are *weakly adjacent* if either $(i, j) \in \mathcal{E}$ or $(j, i) \in \mathcal{E}$.

Given a real vector space V , the *dual space* V^* is the vector space of linear maps from V into \mathbb{R} . If $V = \mathbb{R}^n$, then V^* is the vector space of row vectors in \mathbb{R}^n . In this case, it is typical to make a slight abuse of notation and assume $V^* = \mathbb{R}^n$.

B. Basic concepts

We start with some basic useful concepts. For $x \in \mathbb{R}^n$ and $p \in \mathbb{N}$, the ℓ_p -norm of x is

$$\|x\|_p \triangleq \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}},$$

while the ℓ_{∞} -norm is

$$\|x\|_{\infty} = \lim_{p \rightarrow \infty} \left(\sum_{i=1}^n |x_i|^p \right)^{\frac{1}{p}} = \max_i |x_i|.$$

For $A \in \mathbb{R}^{n \times m}$ and $p \in \mathbb{N}$, the ℓ_p -induced norm of A is

$$\|A\|_p = \max_{\substack{x \in \mathbb{R}^m \\ \|x\|_p \leq 1}} \|Ax\|_p.$$

Definition 1 (Seminorms). A function $\|\cdot\| : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ is a seminorm on \mathbb{R}^n if it satisfies the following properties for all $x, y \in \mathbb{R}^n$ and $a \in \mathbb{R}$:

$$\begin{aligned} (\text{homogeneity}): \quad & \|ax\| = |a| \|x\|, \text{ and} \\ (\text{subadditivity}): \quad & \|x + y\| \leq \|x\| + \|y\|. \end{aligned}$$

Consequently, $\|x\| = 0$ does not imply $x = 0$. The kernel of a seminorm is the vector space

$$\mathcal{K} \triangleq \ker(\|\cdot\|) = \{x \in \mathbb{R}^n : \|x\| = 0\}.$$

From now onward, for a seminorm $\|\cdot\|$ on \mathbb{R}^n with kernel \mathcal{K} we will use the symbol $\|\cdot\|^{\mathcal{K}}$.

Lemma 2 (Seminorms of orthogonal projections). Let $\|\cdot\|^{\mathcal{K}}$ be a seminorm on \mathbb{R}^n with kernel \mathcal{K} , and let Π_{\perp} be the orthogonal projection matrix onto \mathcal{K}^{\perp} . For all $x \in \mathbb{R}^n$, $\|x\|^{\mathcal{K}} = \|\Pi_{\perp} x\|^{\mathcal{K}}$.

Proof. The result is a direct consequence of the reverse triangle inequality and the sub additivity property of seminorms applied to the orthogonal decomposition $x = x_{\perp} + x_{\parallel}$, with $x_{\parallel} \in \mathcal{K}$. \square

Remark 3 (Relationship between norm and seminorm). A seminorm on \mathbb{R}^n with kernel \mathcal{K} induces a norm on \mathcal{K}^{\perp} by restriction. Vice-versa, given a subspace \mathcal{K} of \mathbb{R}^n , a norm $\|\cdot\|$ on \mathcal{K}^{\perp} , denoted by $\|\cdot\|_{\perp}$, induces a seminorm on \mathbb{R}^n with kernel \mathcal{K} by projection: $\|x\|^{\mathcal{K}} = \|\Pi_{\perp} x\|_{\perp}$.

Definition 4. (Induced seminorm [17]) Given a seminorm $\|\cdot\|^{\mathcal{K}} : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ with kernel \mathcal{K} , the induced seminorm on $\mathbb{R}^{n \times n}$ is

$$\|A\|^{\mathcal{K}} \triangleq \max_{\substack{\|x\|^{\mathcal{K}} \leq 1 \\ x \perp \mathcal{K}}} \|Ax\|^{\mathcal{K}}.$$

Definition 5 (Matrix logarithmic seminorms [4]). Given a seminorm $\|\cdot\|^{\mathcal{K}} : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ with kernel \mathcal{K} , the induced matrix logarithmic seminorm on $\mathbb{R}^{n \times n}$ is

$$\mu_{\|\cdot\|^{\mathcal{K}}}(A) \triangleq \lim_{h \rightarrow 0^+} \frac{\|I_n + hA\|^{\mathcal{K}} - 1}{h}.$$

Definition 6 (Generalized ℓ_p ergodicity coefficient [28]). Given $p \in [1, \infty]$ and a vector subspace $\mathcal{K} \subset \mathbb{R}^m$, the generalized ℓ_p ergodicity coefficient $\tau_p : \mathcal{K} \times \mathbb{R}^{m \times n} \rightarrow \mathbb{R}_{\geq 0}$ is defined by

$$\tau_p(\mathcal{K}, A) := \max_{\substack{\|z\|_p = 1 \\ z \perp \mathcal{K}}} \|A^T z\|_p. \quad (6)$$

The ergodicity coefficient (6) is the norm of the operator defined on the real (normed) linear space \mathcal{K}^{\perp} by $x \rightarrow A^T x$ [27].

Lemma 7 (ℓ_2 -Norm LMI characterization [4]). *Given any $A \in \mathbb{R}^{n \times n}$,*

$$\|A\|_2 = \min\{b \in \mathbb{R}_{\geq 0} \mid A^\top A \preceq b^2 I_n\}.$$

III. SEMINORMS AND DUALITY

A. Projection and Distance Seminorms

In the following we provide the definition of projection and distance seminorms. These two seminorms will play a fundamental role in the duality result.

Definition 8 (Projection and distance seminorms). *Let $\mathcal{K} \subset \mathbb{R}^n$ be a vector space and $\Pi_\perp \in \mathbb{R}^{n \times n}$ be the orthogonal projection matrix onto \mathcal{K}^\perp . For each $p \in [1, \infty]$, define the ℓ_p -projection seminorm with respect to \mathcal{K} by*

$$\|x\|_{\text{proj},p}^\mathcal{K} \triangleq \|\Pi_\perp x\|_p \quad (7)$$

and the ℓ_p -distance seminorm with respect to \mathcal{K} by

$$\|x\|_{\text{dist},p}^\mathcal{K} \triangleq \text{dist}_p(x, \mathcal{K}) = \min_{u \in \mathcal{K}} \|x - u\|_p. \quad (8)$$

Note that the optimization problem (8) is well posed since the norm function is convex.

Lemma 9 (Basic properties). *For each $p \in [1, \infty]$,*

- (i) $\ker(\|\cdot\|_{\text{proj},p}^\mathcal{K}) = \ker(\|\cdot\|_{\text{dist},p}^\mathcal{K}) = \mathcal{K}$,
- (ii) $\|x\|_{\text{dist},p}^\mathcal{K} \leq \min\{\|x\|_p, \|x\|_{\text{proj},p}^\mathcal{K}\}$ for all $x \in \mathbb{R}^n$.

Proof. Statement (i) is obvious from (7) and (8). Next, we compute

$$\begin{aligned} \min_{u \in \mathcal{K}} \|x - u\|_p &\leq \|x - 0_n\|_p = \|x\|_p, \\ \min_{u \in \mathcal{K}} \|x - u\|_p &\leq \|x - (I_n - \Pi_\perp)x\|_p = \|x\|_{\text{proj},p}^\mathcal{K}. \end{aligned}$$

This completes the proof of statement (ii). \square

It is not true in general that $\|x\|_{\text{proj},p}^\mathcal{K} \leq \|x\|_p$.

Example 10 (Seminorms for consensus and stationary distribution). *When $\mathcal{K} = \text{span}\{\mathbf{1}_n\}$ and $p \in \{1, 2, \infty\}$, explicit formulas for the ℓ_p -projection and distance seminorms are either easily derivable or available in the literature [4], [14], [23]. For each $x \in \mathbb{R}^n$, with the shorthand $x_{\text{avg}} = \frac{1}{n} \mathbf{1}_n^\top x$,*

$$\begin{aligned} \|x\|_{\text{proj},1}^\mathcal{K} &= \sum_{i=1}^n |x_i - x_{\text{avg}}|, \\ \|x\|_{\text{proj},2}^\mathcal{K} &= \left(\frac{1}{n} \sum_{i,j} (x_i - x_j)^2 \right)^{1/2}, \\ \|x\|_{\text{proj},\infty}^\mathcal{K} &= \max_i |x_i - x_{\text{avg}}|. \end{aligned}$$

Next, sort the entries of x according to $x_{(1)} \geq x_{(2)} \geq \dots \geq x_{(n)}$. With this shorthand,

$$\begin{aligned} \|x\|_{\text{dist},1}^\mathcal{K} &= \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} x_{(i)} - \sum_{i=\lceil \frac{n}{2} \rceil+1}^n x_{(i)}, \\ \|x\|_{\text{dist},2}^\mathcal{K} &= \left(\frac{1}{n} \sum_{i,j} (x_i - x_j)^2 \right)^{1/2}, \\ \|x\|_{\text{dist},\infty}^\mathcal{K} &= \frac{1}{2} (x_{(1)} - x_{(n)}) = \frac{1}{2} (\max_i \{x_i\} - \min_i \{x_j\}). \end{aligned}$$

Figure 1 illustrates the unit disks for these seminorms on \mathbb{R}^3 .

Example 11 (Total variation distance). *The total variation [21, Section 4.1] is a metric on the simplex Δ_n defined by*

$$d_{\text{TV}}(x, y) \triangleq \frac{1}{2} \sum_{i=1}^n |x_i - y_i|.$$

Given any two vectors $x, y \in \Delta_n$, a simple derivation shows

$$d_{\text{TV}}(x, y) = \frac{1}{2} \|x - y\|_{\text{proj},1}^\mathcal{K},$$

where $\|\cdot\|_{\text{proj},1}^\mathcal{K}$ is the ℓ_1 -projection seminorm with respect to the kernel $\mathcal{K} = \text{span}\{\mathbf{1}_n\}$.

B. Duality

In this section we establish a useful duality relationship between projection and distance seminorms. We start with the notion of dual seminorm.

Definition 12 (Dual seminorm). *Let $\|\cdot\|^\mathcal{K}$ be a seminorm on a real vector space $V \subseteq \mathbb{R}^n$ with kernel $\mathcal{K} \subset V$. The dual seminorm is the function $\|\cdot\|_\star^\mathcal{K} : V^\star \rightarrow \mathbb{R}$ defined by*

$$\|x\|_\star^\mathcal{K} \triangleq \max_{\|y\|^\mathcal{K} \leq 1, y \perp \mathcal{K}} \langle x, y \rangle.$$

Lemma 13 (Well-posedness of dual seminorms). *Let $\|\cdot\|^\mathcal{K}$ be a seminorm on a real vector space V with kernel \mathcal{K} . Then the dual seminorm $\|\cdot\|_\star^\mathcal{K}$ is a seminorm on V^\star .*

Proof. Let $y, z \in V^\star$, and let $a \in \mathbb{R}$. Since $x = 0_n$ satisfies $\|x\|^\mathcal{K} \leq 1$ and $x \in \mathcal{K}^\perp$, $\|y\|_\star^\mathcal{K} \geq \langle y, 0_n \rangle = 0$, establishing the non-negativity of $\|\cdot\|_\star^\mathcal{K}$. To prove homogeneity,

$$\begin{aligned} \|ay\|_\star^\mathcal{K} &= \max_{\|x\|^\mathcal{K} \leq 1, x \in \mathcal{K}^\perp} \langle ay, x \rangle = \max_{\|x\|^\mathcal{K} \leq 1, x \in \mathcal{K}^\perp} |a| \text{sgn}(a) \langle y, x \rangle \\ &= |a| \max_{\|x\|^\mathcal{K} \leq 1, x \in \mathcal{K}^\perp} \langle y, \text{sgn}(a)x \rangle = |a| \|y\|_\star^\mathcal{K}. \end{aligned}$$

Finally, to prove sub-additivity,

$$\begin{aligned} \|y + z\|_\star^\mathcal{K} &= \max_{\|x\|^\mathcal{K} \leq 1, x \in \mathcal{K}^\perp} \langle y + z, x \rangle = \max_{\|x\|^\mathcal{K} \leq 1, x \in \mathcal{K}^\perp} \langle y, x \rangle + \langle z, x \rangle \\ &\leq \max_{\|x\|^\mathcal{K} \leq 1, x \in \mathcal{K}^\perp} \langle y, x \rangle + \max_{\|x\|^\mathcal{K} \leq 1, x \in \mathcal{K}^\perp} \langle z, x \rangle = \|y\|_\star^\mathcal{K} + \|z\|_\star^\mathcal{K}. \end{aligned}$$

\square

When $V = \mathbb{R}^n$, we make the usual identification $(\mathbb{R}^n)^\star = \mathbb{R}^n$. In this case, the kernel of the dual seminorm is identical to the kernel of the primal seminorm.

Next, we present an important generalization to arbitrary ℓ_p/ℓ_q norms of the Markov contraction inequality from [14, Lemma 2.3].

Lemma 14 (Markov contraction inequality). *Let $p, q \in [1, \infty]$ satisfy $p^{-1} + q^{-1} = 1$ (with the convention $1/\infty = 0$) and consider a vector space $\mathcal{K} \subset \mathbb{R}^n$. For all $x, y \in \mathbb{R}^n$,*

$$x^\top \Pi_\perp y \leq \|x\|_{\text{proj},p}^\mathcal{K} \|y\|_{\text{dist},q}^\mathcal{K}.$$




Fig. 1: Two-dimensional sections of three-dimensional unit disks of projection (solid contours) and distance (dashed contours) consensus seminorms. We plot the section corresponding to $(x_1, x_2, x_3 = 0)$ for $p = 1$ (left), $p = 2$ (center), and $p = \infty$ (right).

Proof. For each $u \in \mathcal{K}$ satisfying $\|y\|_{\text{dist},q}^{\mathcal{K}} = \|y - u\|_q$,

$$x^T \Pi_{\perp} y = x^T \Pi_{\perp} (y - u) \stackrel{\text{(Hölder's ineq)}}{\leq} \|\Pi_{\perp} x\|_p \|y - u\|_q.$$

The result follows from minimizing with respect to u . \square

Remark 15 (Markov contraction and Hölder's inequalities). *For the inner product of vectors perpendicular to a subspace, the Markov contraction inequality provides a tighter bound than the Hölder's inequality $x^T y \leq \|x\|_p \|y\|_q$. In fact, as a consequence of Lemma 9(ii),*

$$x^T \Pi_{\perp} y \leq \|x\|_{\text{proj},p}^{\mathcal{K}} \|y\|_{\text{dist},q}^{\mathcal{K}} \leq \|x\|_{\text{proj},p}^{\mathcal{K}} \|y\|_{\text{proj},q}^{\mathcal{K}}.$$

Next, we recall that, for unconstrained vectors, the Hölder's inequality provides a tight bound in the sense that, for all $x \in \mathbb{R}^n$, there exists $y \in \mathbb{R}^n$ such that $x^T y = \|x\|_p \|y\|_q$. We now show this tightness result also for the Markov contraction inequality, thereby establishing the duality relationship between projection and the distance seminorms.

Theorem 16 (Duality of distance and projection seminorms). *Let $p, q \in [1, \infty]$ satisfy that $p^{-1} + q^{-1} = 1$ (with the convention $1/\infty = 0$) and let $\mathcal{K} \subset \mathbb{R}^n$ be a vector subspace. Then $\|\cdot\|_{\text{dist},p}^{\mathcal{K}}$ and $\|\cdot\|_{\text{proj},q}^{\mathcal{K}}$, with kernel \mathcal{K} , are dual seminorms:*

$$\|\cdot\|_{\text{dist},p}^{\mathcal{K}} = \left(\|\cdot\|_{\text{proj},q}^{\mathcal{K}} \right)_{\star} \quad (9)$$

$$\|\cdot\|_{\text{proj},q}^{\mathcal{K}} = \left(\|\cdot\|_{\text{dist},p}^{\mathcal{K}} \right)_{\star}. \quad (10)$$

Proof. To prove (9), consider two cases. First, if $x \in \mathcal{K}$, then $\|x\|_{\text{dist},p}^{\mathcal{K}} = 0$. On the other hand, $y \in \mathcal{K}^{\perp}$ implies $y^T x = 0$. So, both sides of (9) are zero. Second, if $x \notin \mathcal{K}$, by Lemma 38 in Appendix I, there exists $\psi_p(x) \in \mathcal{K}^{\perp}$ with $\|\psi_p(x)\|_{\text{proj},q}^{\mathcal{K}} = 1$ such that

$$\|x\|_{\text{dist},p}^{\mathcal{K}} = \psi_p(x)^T x \leq \max_{\substack{y \in \mathcal{K}^{\perp} \\ \|y\|_{\text{proj},q}^{\mathcal{K}} \leq 1}} y^T x = \left(\|\cdot\|_{\text{proj},q}^{\mathcal{K}} \right)_{\star}.$$

To prove the opposite inequality, choose any $y \in \mathbb{R}^n$ such that $\|y\|_{\text{proj},q}^{\mathcal{K}} \leq 1$ and $y \in \mathcal{K}^{\perp}$. Then $\|y\|_q = \|y\|_{\text{proj},q}^{\mathcal{K}} \leq 1$, so by Lemma 14,

$$y^T x \leq \|y\|_q \|x\|_{\text{dist},p}^{\mathcal{K}} \leq \|x\|_{\text{dist},p}^{\mathcal{K}}.$$

To prove equality (10) we notice, as in the previous first case, that if $x \in \mathcal{K}$, then $\|x\|_{\text{proj},q}^{\mathcal{K}} = 0$, while $y \in \mathcal{K}^{\perp}$ implies

that $y^T x = 0$, so both sides of (10) are zero. Otherwise, in the second case, if $x \notin \mathcal{K}$, by Lemma 39, there exists $\zeta_q(x) \in \mathcal{K}^{\perp}$ with $\|\zeta_q(x)\|_{\text{dist},p}^{\mathcal{K}} \leq 1$ such that

$$\|x\|_{\text{proj},q}^{\mathcal{K}} = \zeta_q(x)^T x \leq \max_{\substack{y \in \mathcal{K}^{\perp} \\ \|y\|_{\text{dist},p}^{\mathcal{K}} \leq 1}} y^T x = \left(\|\cdot\|_{\text{dist},p}^{\mathcal{K}} \right)_{\star}.$$

To prove the opposite inequality, choose any $y \in \mathbb{R}^n$ such that $\|y\|_{\text{dist},p}^{\mathcal{K}} \leq 1$ and $y \in \mathcal{K}^{\perp}$. Lemma 14 implies

$$x^T y = x^T \Pi_{\perp} y \leq \|x\|_{\text{proj},q}^{\mathcal{K}} \|y\|_{\text{dist},p}^{\mathcal{K}} \leq \|x\|_{\text{proj},q}^{\mathcal{K}}.$$

This concludes the proof. \square

IV. INDUCED MATRIX SEMINORMS AND LOG SEMINORMS

A. Induced Matrix Seminorms

In the following we list some basic properties related to induced matrix seminorms.

Lemma 17 (Properties of induced matrix seminorms). *Let $\|\cdot\|_{\mathcal{K}}$ be a seminorm on \mathbb{R}^n with kernel \mathcal{K} . For any $A, B \in \mathbb{R}^{n \times n}$,*

$$(i) \quad \|Ax\|_{\mathcal{K}} \leq \|A\|_{\mathcal{K}} \|x\|_{\mathcal{K}} \text{ for all } x \in \mathcal{K}^{\perp}.$$

If $A^T \mathcal{K} \subseteq \mathcal{K}$, then

$$(ii) \quad \|BA\|_{\mathcal{K}} \leq \|B\|_{\mathcal{K}} \|A\|_{\mathcal{K}}.$$

Moreover, If $AK \subseteq \mathcal{K}$, then

$$(iii) \quad \|A\|_{\mathcal{K}} = \max_{\|x\|_{\mathcal{K}} \leq 1} \|Ax\|_{\mathcal{K}},$$

$$(iv) \quad \|Ax\|_{\mathcal{K}} \leq \|A\|_{\mathcal{K}} \|x\|_{\mathcal{K}}, \text{ and}$$

$$(v) \quad \|AB\|_{\mathcal{K}} \leq \|A\|_{\mathcal{K}} \|B\|_{\mathcal{K}}.$$

Proof. Property (i) was proven in [19]. To prove property (ii), assume $x^* \in \mathcal{K}^{\perp}$ to be a vector with $\|x^*\|_{\mathcal{K}} = 1$ such that $\|BAx^*\|_{\mathcal{K}} = \|BA\|_{\mathcal{K}}$. Note that, under the assumption $A^T \mathcal{K} \subseteq \mathcal{K}$, $Ax^* \in \mathcal{K}^{\perp}$. Hence

$$\begin{aligned} \|BA\|_{\mathcal{K}} &= \|BAx^*\|_{\mathcal{K}} = \\ &= \left\| B \frac{Ax^*}{\|Ax^*\|_{\mathcal{K}}} \right\|_{\mathcal{K}} \|Ax^*\|_{\mathcal{K}} \leq \|B\|_{\mathcal{K}} \|A\|_{\mathcal{K}}. \end{aligned}$$

To prove property (iii), decompose any vector $x \in \mathbb{R}^n$ as $x = x_\perp + x_\parallel$, with $x_\perp \in \mathcal{K}^\perp$ and $x_\parallel \in \mathcal{K}$, and notice that

$$\begin{aligned} \max_{\|x\|_{\mathcal{K}^\perp} \leq 1} \|Ax\|_{\mathcal{K}} &= \max_{\|x\|_{\mathcal{K}^\perp} \leq 1} \|A(x_\perp + x_\parallel)\|_{\mathcal{K}} \\ &= \max_{\|x_\perp\|_{\mathcal{K}^\perp} \leq 1} \|Ax_\perp\|_{\mathcal{K}} = \max_{\substack{\|y\|_{\mathcal{K}^\perp} \leq 1 \\ y \perp \mathcal{K}}} \|Ay\|_{\mathcal{K}} = \|A\|_{\mathcal{K}}, \end{aligned}$$

where the second equality is based on Lemma 2 and exploits the fact that $A\mathcal{K} \subseteq \mathcal{K}$. To prove property (iv) we notice that, by adopting the same decomposition as before

$$\begin{aligned} \|Ax\|_{\mathcal{K}} &= \|Ax_\perp\|_{\mathcal{K}} = \left\| A(x_\perp / \|x_\perp\|_{\mathcal{K}^\perp}) \right\|_{\mathcal{K}} \|x_\perp\|_{\mathcal{K}^\perp} \\ &\leq \|A\|_{\mathcal{K}} \|x_\perp\|_{\mathcal{K}^\perp} = \|A\|_{\mathcal{K}} \|x\|_{\mathcal{K}} \end{aligned}$$

where the first equality is based on Lemma 2 and exploits the fact that $A\mathcal{K} \subseteq \mathcal{K}$, while the inequality derives from Definition 4. Property (v) can be found in [19] and can be proved by following arguments similar to property (ii). \square

Based on Theorem 16 we are now in the position to provide one of the main results of this manuscript.

For a matrix $A \in \mathbb{R}^{k \times n}$, and for $p, q \in [1, \infty]$, with $p^{-1} + q^{-1} = 1$ it holds that

$$\|A\|_p = \|A^\top\|_q, \quad (11)$$

where $\|\cdot\|_p : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ and $\|\cdot\|_q : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ are dual norms.

The following theorem represents a generalization of the duality relationship between induced matrix norms (11) to seminorms.

Theorem 18 (Duality of induced matrix seminorms). *Let $p, q \in [1, \infty]$ such that $p^{-1} + q^{-1} = 1$. For any matrix $A \in \mathbb{R}^{n \times n}$, and any vector space $\mathcal{K} \subseteq \mathbb{R}^n$,*

$$\|A^\top\|_{\text{proj},q}^{\mathcal{K}} = \|A\|_{\text{dist},p}^{\mathcal{K}}. \quad (12)$$

Additionally, if $A\mathcal{K} \subseteq \mathcal{K}$, then

$$\tau_q(\mathcal{K}, A) = \|A^\top\|_{\text{proj},q}^{\mathcal{K}} = \|A\|_{\text{dist},p}^{\mathcal{K}}. \quad (13)$$

Proof. Eqn. (12) is a direct consequence of Theorem 16:

$$\begin{aligned} \|A^\top\|_{\text{proj},q}^{\mathcal{K}} &= \max_{\substack{\|x\|_{\text{proj},q}^{\mathcal{K}} \leq 1 \\ x \perp \mathcal{K}}} \|A^\top x\|_{\text{proj},q}^{\mathcal{K}} \\ &\stackrel{(10)}{=} \max_{\substack{\|x\|_{\text{proj},q}^{\mathcal{K}} \leq 1 \\ x \perp \mathcal{K}}} \max_{\substack{\|y\|_{\text{dist},p}^{\mathcal{K}} \leq 1 \\ y \perp \mathcal{K}}} y^\top A^\top x \\ &= \max_{\substack{\|y\|_{\text{dist},p}^{\mathcal{K}} \leq 1 \\ y \perp \mathcal{K}}} \max_{\substack{\|x\|_{\text{proj},q}^{\mathcal{K}} \leq 1 \\ x \perp \mathcal{K}}} x^\top A y \\ &\stackrel{(9)}{=} \max_{\substack{\|y\|_{\text{dist},p}^{\mathcal{K}} \leq 1 \\ y \perp \mathcal{K}}} \|Ay\|_{\text{dist},p}^{\mathcal{K}} = \|A\|_{\text{dist},p}^{\mathcal{K}}. \end{aligned}$$

To prove (13) note that

$$\begin{aligned} \|A^\top\|_{\text{proj},q}^{\mathcal{K}} &= \max_{\substack{\|x\|_{\text{proj},q}^{\mathcal{K}} \leq 1 \\ x \perp \mathcal{K}}} \|A^\top x\|_{\text{proj},q}^{\mathcal{K}} \\ &= \max_{\|x\|_{\mathcal{K}^\perp} \leq 1} \|\Pi_\perp A^\top x\|_q = \max_{\substack{\|x\|_q \leq 1 \\ x \perp \mathcal{K}}} \|A^\top x\|_q = \tau_q(\mathcal{K}, A) \end{aligned}$$

where the second-last equality follows from the fact that $A^\top \mathcal{K}^\perp \subseteq \mathcal{K}^\perp$ and, since $x \in \mathcal{K}^\perp$, $x = \Pi_\perp x$. \square

In the following we provide some explicit expressions for the distance seminorm of row-stochastic matrices and the projection seminorm of column stochastic matrices for the case in which the kernel of the seminorms is the consensus subspace. The explicit expressions can be derived by the ones available in the literature for ergodicity coefficients [16], [30] and by the duality result from Theorem 18.

Corollary 19 (Formulas for induced matrix seminorms). *Consider the consensus distance and projection seminorm. Let $A \in \mathbb{R}^{n \times n}$. Let $a_{i,(j)}$ represent the entries of each row $i \in \{1, 2, \dots, n\}$ sorted according to $a_{i,(1)} \geq a_{i,(2)} \geq \dots \geq a_{i,(n)}$ and similarly, let $a_{(i),j}$ represent the entries of each column $j \in \{1, 2, \dots, n\}$ sorted by $a_{(1),j} \geq a_{(2),j} \geq \dots \geq a_{(n),j}$.*

If A is row-stochastic, then

$$\|A\|_{\text{dist},1}^{\mathcal{K}} = \max_j \left\{ \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} a_{i,j} - \sum_{i=\lceil \frac{n}{2} \rceil+1}^n a_{i,j} \right\}, \quad (14)$$

$$\|A\|_{\text{dist},2}^{\mathcal{K}} = \|\Pi_n A\|_2 = \min \{b \geq 0 : A^\top \Pi_n A \preceq b^2 \Pi_n\}, \quad (15)$$

$$\begin{aligned} \|A\|_{\text{dist},\infty}^{\mathcal{K}} &= \frac{1}{2} \max_{i \neq j} \sum_{k=1}^n |a_{ik} - a_{jk}| \\ &= 1 - \min_{i \neq j} \sum_{k=1}^n \min\{a_{ik}, a_{jk}\}. \end{aligned} \quad (16)$$

If A is column-stochastic, then

$$\begin{aligned} \|A\|_{\text{proj},1}^{\mathcal{K}} &= \frac{1}{2} \max_{i \neq j} \sum_{k=1}^n |a_{ki} - a_{kj}| \\ &= 1 - \min_{i \neq j} \sum_{k=1}^n \min\{a_{ki}, a_{kj}\}, \end{aligned} \quad (17)$$

$$\begin{aligned} \|A\|_{\text{proj},2}^{\mathcal{K}} &= \|\Pi_n A\|_2 \\ &= \min \{b \geq 0 : A^\top \Pi_n A \preceq b^2 \Pi_n\}, \end{aligned} \quad (18)$$

$$\|A\|_{\text{proj},\infty}^{\mathcal{K}} = \max_i \left\{ \sum_{j=1}^{\lfloor \frac{n}{2} \rfloor} a_{i,(j)} - \sum_{j=\lceil \frac{n}{2} \rceil+1}^n a_{i,(j)} \right\}. \quad (19)$$

Proof. The formulas (14), (16) and the first equality in (15) follow from the equivalence $\|A\|_{\text{dist},p}^{\mathcal{K}} = \tau_q(\mathbb{1}_n, A)$ in Theorem 18 and by applying the explicit expressions for $\tau_q(\mathbb{1}_n, A)$ provided in Theorem 3.7, Corollary 3.9, Theorem 4.2, and Theorem 6.19 from [16].

The second equality in (15) follows from Lemma 7, since

$$\begin{aligned} \|\Pi_n A\|_2 &= \min_{b \in \mathbb{R}} \{(\Pi_n A)^\top (\Pi_n A) \preceq b^2 I_n\} \\ &= \min_{b \in \mathbb{R}} \{A^\top \Pi_n A \preceq b^2 I_n\}. \end{aligned}$$

Since $b^2 \Pi_n \preceq b^2 I_n$, it is clear that $A^\top \Pi_n A \preceq b^2 \Pi_n$ implies $A^\top \Pi_n A \preceq b^2 I_n$. Conversely, assume $A^\top \Pi_n A \preceq b^2 I_n$, so that $v^\top A^\top \Pi_n A v \leq b^2 v^\top v$ for all $v \in \mathbb{R}^n$. Then for any $u \in \mathbb{R}^n$, we can decompose $u = u_\perp + u_\parallel$, with $u_\perp \in \text{span}\{\mathbb{1}_n\}^\perp$ and $u_\parallel \in \text{span}\{\mathbb{1}_n\}$. Since A is row stochastic,

$$u^\top A^\top \Pi_n A u = u_\perp^\top A^\top \Pi_n A u_\perp \leq b^2 u_\perp^\top u_\perp = b^2 u^\top \Pi_n u$$

and thus $A^\top \Pi_n A \preceq b^2 \Pi_n$. This way we have proved that $A^\top \Pi_n A \preceq b^2 \Pi_n$ if and only if $A^\top \Pi_n A \preceq b^2 \Pi_n$. In turn, this implies

$$\|\Pi_n A\|_2 = \min_{b \in \mathbb{R}} \{A^\top \Pi_n A \preceq b^2 \Pi_n\}.$$

Formulas (17) – (19) are derived by duality. \square

Finally, we include a comparative analysis for induced seminorms and the notion of optimal deflation given by [23].

Definition 20 (*p*-optimal deflation [23]). *For each $p \in [1, \infty]$, the p -optimal deflation of a matrix $A \in \mathbb{R}^{k \times n}$ is*

$$|A|_p \triangleq \min_{v \in \mathbb{R}^n} \|A - \mathbb{1}_n v^\top\|_p. \quad (20)$$

Lemma 21 (Bounds on matrix seminorms). *Given a row-stochastic matrix $A \in \mathbb{R}^{n \times n}$, for each $p \in [1, \infty]$*

$$\|A\|_{\text{dist},p}^{\mathcal{K}} \leq |A|_p \leq \|A\|_p.$$

Proof. We first establish that $|A|_p \geq \|A\|_{\text{dist},p}^{\mathcal{K}}$. By the max-min inequality [3, Section 5.4.1],

$$\begin{aligned} |A|_p &= \min_{v \in \mathbb{R}^n} \max_{\|w\|_p \leq 1} \|(A - \mathbb{1}_n v^\top) w\|_p \\ &\geq \max_{\|w\|_p \leq 1} \min_{v \in \mathbb{R}^n} \|(Aw) - (v^\top w) \mathbb{1}_n\|_p \\ &\geq \max_{\|w\|_p \leq 1} \|Aw\|_{\text{dist},p}^{\mathcal{K}}. \end{aligned}$$

Let $w \in \mathbb{R}^n$ be such that $\|w\|_{\text{dist},p}^{\mathcal{K}} \leq 1$, which implies that $\|w - \alpha \mathbb{1}_n\|_p \leq 1$ for some $\alpha \in \mathbb{R}$. Let $u = w - \alpha \mathbb{1}_n$, and observe that $\|u\|_p \leq 1$, and that $\|Au\|_{\text{dist},p}^{\mathcal{K}} = \|Aw\|_{\text{dist},p}^{\mathcal{K}}$, since A is row-stochastic and $\|\cdot\|_{\text{dist},p}^{\mathcal{K}}$ is invariant with respect to perturbations in $\text{span}\{\mathbb{1}_n\}$. Therefore

$$\begin{aligned} \max_{\|w\|_p \leq 1} \|Aw\|_{\text{dist},p}^{\mathcal{K}} &\geq \max_{\|w\|_{\text{dist},p}^{\mathcal{K}} \leq 1} \|Aw\|_{\text{dist},p}^{\mathcal{K}} \\ &\geq \max_{\substack{\|w\|_{\text{dist},p}^{\mathcal{K}} \leq 1 \\ \mathbb{1}_n^\top w = 0}} \|Aw\|_{\text{dist},p}^{\mathcal{K}} = \|A\|_{\text{dist},p}^{\mathcal{K}}. \end{aligned}$$

The inequality $|A|_p \leq \|A\|_p$ is obtained at $v = \mathbb{0}_n$ in (20). \square

B. Induced Matrix Log Seminorms

We now present a duality result for induced matrix log seminorms which is parallel to the one in Theorem 18.

Theorem 22 (Dual logarithmic seminorms). *Let $p, q \in [1, \infty]$ be such that $p^{-1} + q^{-1} = 1$. For any matrix $M \in \mathbb{R}^{n \times n}$, and any kernel \mathcal{K} ,*

$$\mu_{\text{dist},p}(M) = \mu_{\text{proj},q}(M^\top).$$

Proof. From (5) and Theorem 18,

$$\begin{aligned} \mu_{\text{dist},p}(M) &= \lim_{h \rightarrow 0^+} \frac{\|I_n + hM\|_{\text{dist},p}^{\mathcal{K}} - 1}{h} = \\ &= \lim_{h \rightarrow 0^+} \frac{\|I_n + hM^\top\|_{\text{proj},q}^{\mathcal{K}} - 1}{h} = \mu_{\text{proj},q}(M^\top). \end{aligned}$$

\square

We derive now explicit formulas for ℓ_p -distance logarithmic seminorm of (minus) Laplacian matrices, for $p \in \{1, 2, \infty\}$.

Theorem 23 (Explicit formulas for distance logarithmic seminorms). *Consider the consensus distance and projection seminorms. Let $L \in \mathbb{R}^{n \times n}$ be the Laplacian matrix corresponding to an adjacency matrix $A \in \mathbb{R}^{n \times n}$ without self-loops, and let $d_{\text{out}} = A \mathbb{1}_n$. For each $i \in \{1, 2, \dots, n\}$, sort the off-diagonal entries of $A e_j$ according to*

$$a_{(1),j} \geq a_{(2),j} \geq \dots \geq a_{(n-1),j}.$$

Then

$$\mu_{\text{dist},1}(-L) = -\min_j \left\{ [d_{\text{out}}]_j - \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor - 1} a_{(i),j} + \sum_{i=\lceil \frac{n}{2} \rceil}^{n-1} a_{(i),j} \right\},$$

$$\mu_{\text{dist},2}(-L) = \min_{b \in \mathbb{R}} \{b : \Pi_n L + L^\top \Pi_n \succeq -2b \Pi_n\},$$

$$\mu_{\text{dist},\infty}(-L) = -\min_{i \neq j} \left\{ a_{ij} + a_{ji} + \sum_{k \neq i,j} \min\{a_{ik}, a_{jk}\} \right\}.$$

Proof. Set $S_h = I_n - hL$. Observe that S_h is row-stochastic for every $h > 0$, and its entries are

$$[S_h]_{ij} = \begin{cases} 1 - h[d_{\text{out}}]_i, & i = j, \\ h a_{ij}, & i \neq j. \end{cases}$$

Also, $\mu_{\|\cdot\|_{\mathcal{K}}}(-L) = \lim_{h \rightarrow 0^+} h^{-1} (\|S_h\|_{\mathcal{K}} - 1)$ for any seminorm $\|\cdot\|_{\mathcal{K}}$.

Case $\|\cdot\|_{\text{dist},1}^{\mathcal{K}}$. For each $j \in \{1, 2, \dots, n\}$, sort the entries of $S_h e_j$ as

$$(S_h)_{(1),j} \geq (S_h)_{(2),j} \geq \dots \geq (S_h)_{(n),j}$$

Assume h is so small that $(S_h)_{(1),j} = (S_h)_{j,j}$. Then by (14),

$$\begin{aligned} \|S_h\|_{\text{dist},1}^{\mathcal{K}} &= \max_j \left\{ \sum_{i=1}^{\lfloor \frac{n}{2} \rfloor} s_{(i),j} - \sum_{i=\lceil \frac{n}{2} \rceil+1}^n s_{(i),j} \right\} \\ &= 1 + h \max_j \left\{ -[d_{\text{out}}]_j + \sum_{i=2}^{\lfloor \frac{n}{2} \rfloor} a_{(i),j} - \sum_{i=\lceil \frac{n}{2} \rceil+1}^n a_{(i),j} \right\}. \end{aligned}$$

Substituting into (5) yields the formula for $\mu_{\text{dist},1}(-L)$, since the order of the off-diagonal elements of $A e_j$ is identical to the order of the off-diagonal elements of $S_h e_j$ for all $h > 0$.

Case $\|\cdot\|_{\text{dist},2}^{\mathcal{K}}$. By (15),

$$\begin{aligned} \|S_h\|_{\text{dist},2}^{\mathcal{K}} &= \min_{b \geq 0} \{b : S_h^\top \Pi_n S_h \preceq b^2 \Pi_n\} \\ &= \min \{b \geq 0 : (I_n - hL)^\top \Pi_n (I_n - hL) \preceq b^2 \Pi_n\} \\ &= \min \{b \geq 0 : h^2 L^\top \Pi_n L - h \Pi_n L - h L^\top \Pi_n \preceq (b^2 - 1) \Pi_n\} \end{aligned}$$

Therefore $\frac{\|S_h\|_{\text{dist},2}^{\mathcal{K}} - 1}{h}$ is equal to

$$\min \left\{ \frac{b-1}{h} : b \geq 0, h^2 L^\top \Pi_n L - h \Pi_n L - h L^\top \Pi_n \preceq (b^2 - 1) \Pi_n \right\}.$$

Let $\bar{b} = h^{-1}(b-1)$, so that $b \geq 0$ if and only if $\bar{b} \geq -h^{-1}$, and $(b^2 - 1) = h\bar{b}(2+h\bar{b})$. Performing this change of variables,

$$\begin{aligned} \mu_{\text{dist},2}(L) &= \lim_{h \rightarrow 0^+} \frac{\|S_h\|_{\text{dist},2}^{\mathcal{K}} - 1}{h} \\ &= \lim_{h \rightarrow 0^+} \min \{\bar{b} \geq -h^{-1} : h L^\top \Pi_n L - \Pi_n L - L^\top \Pi_n \preceq \bar{b}(2+h\bar{b}) \Pi_n\} \\ &= \min \{\bar{b} \geq 0 : -\Pi_n L - L^\top \Pi_n \preceq 2\bar{b} \Pi_n\}, \end{aligned}$$

which is equivalent to the formula for $\mu_{\text{dist},2}(-L)$.

Case $\|\cdot\|_{\text{dist},\infty}^{\mathcal{K}}$: Assume h is sufficiently small that $1 - h[d_{\text{out}}]_i > ha_{ji}$ for all i, j . Applying (16),

$$\begin{aligned} \|S_h\|_{\text{dist},\infty}^{\mathcal{K}} &= 1 - \min_{i \neq j} \{ \min\{1 - h[d_{\text{out}}]_i, ha_{ji}\} \\ &\quad + \min\{1 - h[d_{\text{out}}]_i, ha_{ij}\} + h \sum_{k \neq i,j} \min\{a_{ik}, a_{jk}\} \} \\ &= 1 - h \min_{i \neq j} \left\{ a_{ij} + a_{ji} + \sum_{k \neq i,j} \min\{a_{ik}, a_{jk}\} \right\}. \end{aligned}$$

Substituting into (5) yields the formula for $\mu_{\text{dist},\infty}(-L)$. \square

We also notice that, when $L = L^T$, one can also show that $\mu_{\text{dist},2}(-L) = -\lambda_2(L)$ (e.g., see [5, Exercise 6.3]). Explicit expressions for the ℓ_p -projection logarithmic seminorm of Laplacian matrices, for $p \in \{1, 2, \infty\}$, are derived by duality.

Corollary 24 (Explicit formulas for projection logarithmic seminorms). *Consider the distance and the projection seminorms with $\ker(\|\cdot\|_{\text{dist}}^{\mathcal{K}}) = \ker(\|\cdot\|_{\text{proj}}^{\mathcal{K}}) = \text{span}\{\mathbf{1}_n\}$. Let $L \in \mathbb{R}^{n \times n}$ be the Laplacian matrix corresponding to an adjacency matrix $A \in \mathbb{R}^{n \times n}$ without self-loops, and let $d_{\text{out}} = A\mathbf{1}_n$. For each $j \in \{1, 2, \dots, n\}$, sort the off-diagonal entries of $e_i^T A$ according to $a_{i,(1)} \geq a_{i,(2)} \geq \dots \geq a_{i,(n-1)}$. Then*

$$\begin{aligned} \mu_{\text{proj},1}(-L) &= -\min_{i \neq j} \left\{ a_{ji} + a_{ij} + \sum_{k \neq i,j} \min\{a_{ki}, a_{kj}\} \right\}, \\ \mu_{\text{proj},2}(-L) &= \min_{b \in \mathbb{R}} \{ b : \Pi_n L + L^T \Pi_n \succeq -2b \Pi_n \}, \\ \mu_{\text{proj},\infty}(-L) &= -\min_i \left\{ [d_{\text{out}}]_i - \sum_{j=1}^{\lfloor \frac{n}{2} \rfloor - 1} a_{i,(j)} + \sum_{j=\lceil \frac{n}{2} \rceil}^{n-1} a_{i,(j)} \right\}. \end{aligned}$$

V. SEMICONTRACTING DYNAMICAL SYSTEMS

We exploit now the duality result of induced matrix seminorms and induced matrix logarithmic seminorms for the study of strong semicontractivity of dynamical systems. We also provide some theoretical results that formalize semicontractivity conditions for linear and nonlinear dynamical systems both in discrete and continuous time.

Given a vector subspace $\mathcal{K} \subset \mathbb{R}^n$ and a vector field $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$, the *perpendicular vector field* $f_{\perp} : \mathbb{R}^n \rightarrow \mathcal{K}^{\perp}$ and the *parallel vector field* $f_{\parallel} : \mathbb{R}^n \rightarrow \mathcal{K}$ are denoted for all $x \in \mathbb{R}^n$ by $f_{\perp}(x) = \Pi_{\perp} f(x)$ and $f_{\parallel}(x) = (I_n - \Pi_{\perp})f(x)$, respectively. Given a seminorm $\|\cdot\|_{\mathcal{K}} : \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$, with kernel \mathcal{K} , the domain restriction of $\|\cdot\|_{\mathcal{K}}$ to \mathcal{K}^{\perp} , will be denoted by $\|\cdot\|_{\perp} : \mathcal{K}^{\perp} \rightarrow \mathbb{R}_{\geq 0}$.

Definition 25 (Invariant sets). *Let $f : \mathbb{R}^n \rightarrow \mathbb{R}^n$. A subspace $V \subset \mathbb{R}^n$ is f -invariant on a domain $C \subseteq \mathbb{R}^n$ if $f(x+v) - f(x) \in V$ for all $x \in C$ and $v \in V$.*

Lemma 26 (Differential characterization of invariance). *Given a continuously differentiable map $f : C \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^n$, a subspace $V \subset \mathbb{R}^n$ is f -invariant if and only if $Df(x)V \subseteq V$ for all $x \in C$.*

Proof. If V is f -invariant, then $f(x+hv) - f(x) \in V$ for all $x \in C$, $v \in V$, and $h \in \mathbb{R}$, which implies that

$$Df(x)v = \lim_{h \rightarrow 0} \frac{f(x+hv) - f(x)}{h} \in V$$

thus $Df(x)V \subseteq V$ for all $x \in C$. To prove the converse, assume $Df(x)V \subseteq V$; then for all $v \in V$,

$$f(x+v) - f(x) = \int_0^1 Df(x+\alpha v)v \, d\alpha \in V.$$

\square

A. Discrete Time Semicontraction

Let us consider the discrete time, time varying, nonlinear dynamics

$$x(k+1) = f(k, x(k)) \quad (21)$$

with $k \in \mathbb{Z}_{\geq 0}$, $x \in \mathbb{R}^n$. We assume f to be continuously differentiable in the second argument. In the following we give a generalized definition of strongly semicontracting discrete time system with respect to the one in [17]. The generalization applies to systems with arbitrary contraction step.

Definition 27 (Semicontracting discrete time systems). *Let $\|\cdot\|_{\mathcal{K}}$ be a seminorm on \mathbb{R}^n with kernel \mathcal{K} . If there exists $m \in \mathbb{N}$, $\rho < 1$ and a domain $C \subseteq \mathbb{R}^n$ for which the time-varying vector field $f : \mathbb{Z}_{\geq 0} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is such that ¹*

$$\|D(f^m(k, x))\|_{\mathcal{K}} \leq \rho \quad (22)$$

for all $k \in \mathbb{Z}_{\geq 0}$ and $x \in C$, then the vector field is strongly semicontracting on C with rate $\sqrt[m]{\rho}$.

Lemma 28 provides sufficient conditions for two fundamental discrete-time systems to be strongly semicontracting.

Lemma 28 (Strong semicontractivity of discrete-time affine systems). *Given a subspace $\mathcal{K} \subset \mathbb{R}^n$ and $p, q \in [1, \infty]$ with $p^{-1} + q^{-1} = 1$, consider a sequence of matrices $\{A(k)\}_{k \in \mathbb{Z}_{\geq 0}} \subset \mathbb{R}^{n \times n}$ satisfying:*

$$A(k)\mathcal{K} \subseteq \mathcal{K} \quad \text{for all } k \in \mathbb{Z}_{\geq 0}, \quad (\text{invariance})$$

$$\rho \triangleq \sup_{k \in \mathbb{Z}_{\geq 0}} \tau_p(\mathcal{K}, A(k)) < 1. \quad (\text{semicontractivity})$$

(i) Then the system

$$x(k+1) = A(k)x(k) + b(k), \quad b(k) \in \mathbb{R}^n, \quad (23)$$

is strongly semicontracting with rate ρ in the distance ℓ_q seminorm with kernel \mathcal{K} . Moreover

$$\|x(k) - y(k)\|_{\text{dist},q}^{\mathcal{K}} \leq \rho^k \|x(0) - y(0)\|_{\text{dist},q}^{\mathcal{K}}.$$

(ii) The system

$$x(k+1) = A^T(k)x(k) + b(k), \quad b(k) \in \mathbb{R}^n, \quad (24)$$

is strongly semicontracting with rate ρ in the projection ℓ_p seminorm with kernel \mathcal{K} . Moreover, for any $x(0), y(0)$ satisfying $x(0) - y(0) \in \mathcal{K}^{\perp}$,

$$\|x(k) - y(k)\|_{\text{proj},p}^{\mathcal{K}} \leq \rho^k \|x(0) - y(0)\|_{\text{proj},p}^{\mathcal{K}}.$$

¹ f^m is the m -th iterate of f defined recursively by $f^m(k, x) = f(f^{m-1}(k, x))$.

Proof. The proof of part (i) follows from equation (13) in Theorem 18, and from the conditional submultiplicative property (iii) Lemma 17:

$$\begin{aligned} & \|x(k+1) - y(k+1)\|_{\text{dist},q}^{\mathcal{K}} \\ & \leq \|A(k)\|_{\text{dist},q}^{\mathcal{K}} \|x(k) - y(k)\|_{\text{dist},q}^{\mathcal{K}} \\ & = \tau_p(\mathcal{K}, A(k)) \|x(k) - y(k)\|_{\text{dist},q}^{\mathcal{K}}. \end{aligned} \quad (25)$$

The proof of part (ii) follows from Lemma 17 part (i) since $x(k) - y(k) \in \mathcal{K}^\perp$, $\forall k \in \mathbb{Z}_{\geq 0}$, as a consequence of the invariance assumption (invariance) and therefore

$$\begin{aligned} & \|x(k+1) - y(k+1)\|_{\text{proj},p}^{\mathcal{K}} \\ & \leq \|A^T(k)\|_{\text{proj},p}^{\mathcal{K}} \|x(k) - y(k)\|_{\text{proj},p}^{\mathcal{K}} \\ & = \tau_p(\mathcal{K}, A(k)) \|x(k) - y(k)\|_{\text{proj},p}^{\mathcal{K}}. \end{aligned} \quad (26)$$

□

For example, when the subspace \mathcal{K} is the consensus subspace, the matrices $\{A(k)\}_{k=0}^\infty$ are row-stochastic and the term $b(k) \equiv 0_n \forall k \in \mathbb{Z}_{\geq 0}$, the systems (23) and (24) are the standard averaging (1a) and flow systems (1b) in the Introduction and the bounds (25) and (26) are precisely the bounds (3) and (5) stated in the Introduction.

The following theorem focuses on strong semicontractivity of discrete-time dynamical systems that enjoy the invariance property of the kernel of the seminorm.

Theorem 29 (Discrete time semicontracting dynamics with invariance property). *Consider a system as in (21). Let $\mathcal{K} \subset \mathbb{R}^n$ be an f -invariant subspace, and suppose that f is strongly semicontracting with rate $\rho < 1$, with respect to a seminorm $\|\cdot\|_{\mathcal{K}}$ on \mathbb{R}^n with kernel \mathcal{K} . Then,*

(i) *the system admits the cascade decomposition*

$$x_{\parallel}(k+1) = f_{\parallel}(k, x_{\parallel}(k) + x_{\perp}(k)), \quad (27)$$

$$x_{\perp}(k+1) = f_{\perp}(k, x_{\perp}(k)); \quad (28)$$

(ii) *the perpendicular dynamics (28) are strongly contracting on \mathcal{K}^\perp with rate ρ , with respect to $\|\cdot\|_{\perp} : \mathcal{K}^\perp \rightarrow \mathbb{R}_{\geq 0}$; and*

(iii) *for any two trajectories $x(k), y(k)$ of (21),*

$$\|x(k) - y(k)\|_{\mathcal{K}}^{\mathcal{K}} \leq \rho^k \|x(0) - y(0)\|_{\mathcal{K}}^{\mathcal{K}}$$

for all $k \in \mathbb{Z}_{\geq 0}$.

Proof. Regarding part (i), the cascade decomposition (27)-(28) follows from the observation that

$$\begin{aligned} x_{\perp}(k+1) &= \Pi_{\perp} f(k, x_{\parallel}(k) + x_{\perp}(k)) \\ &= \Pi_{\perp} f(k, x_{\perp}(k)) = f_{\perp}(k, x_{\perp}(k)) \end{aligned}$$

where the second equality is due to the f -invariance of \mathcal{K} . Part (ii) follows from

$$\begin{aligned} \max_{\substack{y \perp \mathcal{K} \\ k \geq 0}} \|Df_{\perp}(k, y)\|_{\mathcal{K}}^{\mathcal{K}} &= \max_{\substack{y \perp \mathcal{K} \\ k \geq 0}} \|\Pi_{\perp} Df(k, y)\|_{\mathcal{K}}^{\mathcal{K}} \\ &= \max_{\substack{y \perp \mathcal{K} \\ k \geq 0}} \|Df(k, y)\|_{\mathcal{K}}^{\mathcal{K}} \leq \max_{\substack{x \in \mathbb{R}^n \\ k \geq 0}} \|Df(k, x)\|_{\mathcal{K}}^{\mathcal{K}} \leq \rho \end{aligned}$$

where the second equality follows from the fact that for a generic matrix A , $\|A\|_{\mathcal{K}}^{\mathcal{K}} = \|\Pi_{\perp} A\|_{\mathcal{K}}^{\mathcal{K}}$. Part (iii) is a direct consequence of (ii). □

The following theorem focuses on strong semicontractivity of discrete-time dynamical systems that enjoy the invariance property of the orthogonal complement of the kernel of the seminorm.

Theorem 30 (Discrete time semicontracting dynamics with conservation property). *Consider a system as in (21). Let $\mathcal{K} \subset \mathbb{R}^n$ such that \mathcal{K}^\perp is an f -invariant subspace. Let $f : \mathbb{Z}_{\geq 0} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ be strongly semicontracting with rate $\rho < 1$ with respect to a seminorm $\|\cdot\|_{\mathcal{K}}$ on \mathbb{R}^n with kernel \mathcal{K} . Then,*

(i) *the system admits the cascade decomposition*

$$x_{\parallel}(k+1) = f_{\parallel}(k, x_{\parallel}(k)), \quad (29)$$

$$x_{\perp}(k+1) = f_{\perp}(k, x_{\parallel}(k) + x_{\perp}(k)); \quad (30)$$

(ii) *for each $x_{\parallel} \in \mathcal{K}$, the vector field $x_{\perp} \mapsto f_{\perp}(k, x_{\parallel} + x_{\perp})$ is strongly contracting with rate ρ , with respect to $\|\cdot\|_{\perp} : \mathcal{K}^\perp \rightarrow \mathbb{R}_{\geq 0}$;*

(iii) *if the map $x_{\parallel} \mapsto f_{\perp}(k, x_{\parallel} + x_{\perp})$ is Lipschitz² with constant $\ell \in \mathbb{R}$ with respect to some metric $d_{\mathcal{K}}$ on \mathcal{K} , then for any two trajectories $x(k), y(k)$ of (21), satisfying $x(0) - y(0) \in \mathcal{K}^\perp$*

$$\begin{aligned} & \|x(k+1) - y(k+1)\|_{\mathcal{K}}^{\mathcal{K}} \\ & \leq \rho \|x(k) - y(k)\|_{\mathcal{K}}^{\mathcal{K}} + \ell d_{\mathcal{K}}(x_{\parallel}(k), y_{\parallel}(k)) \end{aligned}$$

for all $k \in \mathbb{Z}_{\geq 0}$.

Proof. Regarding part (i), the cascade decomposition (29)-(30) follows from the observation that

$$\begin{aligned} x_{\parallel}(k+1) &= (I_n - \Pi_{\perp})f(k, x_{\parallel}(k) + x_{\perp}(k)) \\ &= (I_n - \Pi_{\perp})f(k, x_{\parallel}(k)) = f_{\parallel}(k, x_{\parallel}(k)) \end{aligned}$$

where the second equality is due to the f -invariance of \mathcal{K}^\perp . To prove (ii), fix $x_{\parallel} \in \mathcal{K}$, and pick any $x_{\perp}, y_{\perp} \in \mathcal{K}^\perp$. Then

$$\begin{aligned} & \|f_{\perp}(k, x_{\parallel} + x_{\perp}(k)) - f_{\perp}(k, x_{\parallel} + y_{\perp}(k))\|_{\perp} \\ &= \|f_{\perp}(k, x_{\parallel} + x_{\perp}(k)) - f_{\perp}(k, x_{\parallel} + y_{\perp}(k))\|_{\mathcal{K}}^{\mathcal{K}} \\ &\leq \rho \|x_{\perp}(k) - y_{\perp}(k)\|_{\mathcal{K}}^{\mathcal{K}} = \rho \|x_{\perp}(k) - y_{\perp}(k)\|_{\perp} \end{aligned}$$

To prove (iii), let $x_{\parallel}, y_{\parallel} \in \mathcal{K}$ and $x_{\perp}, y_{\perp} \in \mathcal{K}^\perp$. Then

$$\begin{aligned} & \|f(k, x_{\parallel}(k) + x_{\perp}(k)) - f(k, y_{\parallel}(k) + y_{\perp}(k))\|_{\mathcal{K}}^{\mathcal{K}} \\ & \leq \|f(k, x_{\parallel}(k) + x_{\perp}(k)) - f(k, y_{\parallel}(k) + x_{\perp}(k))\|_{\mathcal{K}}^{\mathcal{K}} \\ & \quad + \|f(k, y_{\parallel}(k) + x_{\perp}(k)) - f(k, y_{\parallel}(k) + y_{\perp}(k))\|_{\mathcal{K}}^{\mathcal{K}} \\ & \leq \ell d_{\mathcal{K}}(x_{\parallel}(k), y_{\parallel}(k)) + \rho \|x(k) - y(k)\|_{\mathcal{K}}^{\mathcal{K}} \end{aligned}$$

where the first inequality is due to the subadditivity property and the second one follows from point (ii) and the invariance of \mathcal{K}^\perp . □

²That is, for all $x_{\parallel}, y_{\parallel} \in \mathcal{K}$, $z_{\perp} \in \mathcal{K}^\perp$, and $k \in \mathbb{Z}_{\geq 0}$, we have $\|f_{\perp}(k, x_{\parallel} + z_{\perp}) - f_{\perp}(k, y_{\parallel} + z_{\perp})\|_{\perp} \leq \ell d_{\mathcal{K}}(x_{\parallel}, y_{\parallel})$.

B. Continuous Time Semicontraction

Let us consider the continuous time, time varying, nonlinear dynamics

$$\dot{x}(t) = f(t, x(t)) \quad (31)$$

with $t \in \mathbb{R}_{\geq 0}, x \in \mathbb{R}^n$. We assume f to be continuously differentiable in the second argument.

Definition 31 (Semicontracting continuous time systems). *Let $\|\cdot\|^\mathcal{K}$ be a seminorm on \mathbb{R}^n with kernel \mathcal{K} . The time-varying vector field $f : \mathbb{R}_{\geq 0} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ is strongly infinitesimally semicontracting with rate $c > 0$ on a domain $C \subseteq \mathbb{R}^n$ if $\forall t \in \mathbb{R}_{\geq 0}$ and $x \in C$,*

$$\mu_{\|\cdot\|^\mathcal{K}}(Df(t, x)) \leq -c.$$

Lemma 32 provides sufficient conditions for two fundamental continuous time dynamical systems to be strongly infinitesimally semicontracting.

Lemma 32 (Strong semicontractivity of continuous-time affine systems). *Given a subspace $\mathcal{K} \subset \mathbb{R}^n$ and $p, q \in [1, \infty]$ with $p^{-1} + q^{-1} = 1$, consider a sequence of matrices $\{A(t)\}_{t \in \mathbb{R}_{\geq 0}} \subset \mathbb{R}^{n \times n}$ satisfying:*

$$A(t)\mathcal{K} \subseteq \mathcal{K} \quad \text{for all } t \in \mathbb{R}_{\geq 0}, \quad (\text{invariance})$$

$$c \triangleq - \sup_{t \in \mathbb{R}_{\geq 0}} \mu_{\text{dist}, p}(A(t)) > 0. \quad (\text{semicontractivity})$$

(i) *The system*

$$\dot{x}(t) = A(t)x(t) + b(t), \quad b(t) \in \mathbb{R}^n,$$

is strongly infinitesimally semicontracting with rate c in the distance ℓ_p seminorm with kernel \mathcal{K} , moreover

$$\|x(t) - y(t)\|_{\text{dist}, p}^\mathcal{K} \leq e^{-ct} \|x(0) - y(0)\|_{\text{dist}, p}^\mathcal{K}, \quad \forall t$$

(ii) *the system*

$$\dot{x}(t) = A^\top(t)x(t) + b(t), \quad b(t) \in \mathbb{R}^n, \quad (32)$$

is strongly infinitesimally semicontracting with rate c in the projection ℓ_q seminorm with kernel \mathcal{K} , moreover, for any $x(0), y(0)$ satisfying $x(0) - y(0) \in \mathcal{K}^\perp$,

$$\|x(t) - y(t)\|_{\text{proj}, q}^\mathcal{K} \leq e^{-ct} \|x(0) - y(0)\|_{\text{proj}, q}^\mathcal{K}, \quad \forall t. \quad (33)$$

Proof. The proof of part (i) follows from Theorem 13, part i) in [17]. To prove part (ii) we follow a similar reasoning as in Theorem 11 from [17]. In fact, for all $x(0), y(0)$ such that $x(0) - y(0) \in \mathcal{K}^\perp$, since the solutions $t \mapsto x(t)$ of (32) are differentiable, by defining $z(t) \triangleq x(t) - y(t)$, for small h , one can write

$$z(t+h) = z(t) + h(A^\top(t)(z(t))) + o(h) = \Pi_\perp(z(t+h))$$

since $z(t) \in \mathcal{K}^\perp$ and $A^\top \mathcal{K}^\perp \subseteq \mathcal{K}^\perp$ by hypothesis. Therefore, by Lemma 2 and Lemma 17 part (i)

$$\frac{\|z(t+h)\|^\mathcal{K} - \|z(t)\|^\mathcal{K}}{h} \leq \frac{\|I_n + hA^\top(t)\|^\mathcal{K} - 1}{h} \|z(t)\|^\mathcal{K} + \frac{o(h)}{h}.$$

Taking the limit as $h \rightarrow 0^+$, one gets $\frac{d}{dt} \|z(t)\|^\mathcal{K} \leq \mu_{\|\cdot\|^\mathcal{K}}(A^\top(t)) \|z(t)\|^\mathcal{K}$. Finally, from the Grönwall comparison inequality (e.g., see [4, Exercise 2.1])

$$\|x(t) - y(t)\|^\mathcal{K} \leq \exp\left(\int_0^t \mu_{\|\cdot\|^\mathcal{K}}(A^\top(\tau)) d\tau\right) \|x(0) - y(0)\|^\mathcal{K}.$$

Eq. (33) follows from the fact that $\mu_{\text{dist}, p}(A(t)) = \mu_{\text{proj}, q}(A^\top(t)) \leq -c$ for all t . \square

The following theorem focuses on strong infinitesimal semicontractivity of continuous-time dynamical systems that enjoy the invariance property of the kernel of the seminorm. This theorem extends Theorem 13 from [17] through the formulation of a cascade decomposition and by establishing a strong contractivity property on the orthogonal complement to the seminorm kernel.

Theorem 33 (Continuous time semicontracting dynamics with invariance property, partially from [17]). *Consider a system as in (31). Let $\mathcal{K} \subset \mathbb{R}^n$ be an f -invariant subspace and suppose that f is strongly infinitesimally semicontracting with rate $c > 0$, with respect to a seminorm $\|\cdot\|^\mathcal{K}$ in \mathbb{R}^n with kernel \mathcal{K} . Then,*

(i) *the system admits the cascade decomposition*

$$\dot{x}_\parallel(t) = f_\parallel(t, x_\parallel(t) + x_\perp(t)), \quad (34)$$

$$\dot{x}_\perp(t) = f_\perp(t, x_\perp(t)); \quad (35)$$

(ii) *the perpendicular dynamics (35) are strongly infinitesimally contracting on \mathcal{K}^\perp with rate c , with respect to $\|\cdot\|_\perp : \mathcal{K}^\perp \rightarrow \mathbb{R}_{\geq 0}$;*

(iii) *for any two trajectories $x(t), y(t)$ of (31),*

$$\|x(t) - y(t)\|^\mathcal{K} \leq e^{-ct} \|x(0) - y(0)\|^\mathcal{K}$$

for all $t \in \mathbb{R}_{\geq 0}$.

Proof. Regarding part (i), the cascade decomposition is obtained by following the same reasoning as in Theorem 29. Part (ii) follows from

$$\begin{aligned} \mu_{\|\cdot\|^\mathcal{K}}(Df_\perp(t, y)) &= \mu_{\|\cdot\|^\mathcal{K}}(\Pi_\perp Df(t, y)) \\ &\leq \mu_{\|\cdot\|^\mathcal{K}}(Df(t, x)) \leq -c \end{aligned}$$

where the first equality follows from the fact that for a generic matrix A , $\mu_{\|\cdot\|^\mathcal{K}}(A) = \mu_{\|\cdot\|^\mathcal{K}}(\Pi_\perp A)$. Part (iii) is a direct consequence of part (ii). \square

The following theorem focuses on strong semicontractivity of continuous-time dynamical systems that enjoy the invariance property of the orthogonal complement of the kernel of the seminorm.

Theorem 34 (Continuous time semicontracting dynamics with conservation property). *Consider a system as in (31). Let $\mathcal{K} \subset \mathbb{R}^n$ be such that \mathcal{K}^\perp is an f -invariant subspace. Let $f : \mathbb{R}_{\geq 0} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$ be strongly infinitesimally semicontracting with rate $c > 0$ with respect to a seminorm $\|\cdot\|^\mathcal{K}$ on \mathbb{R}^n with kernel \mathcal{K} . Then,*

(i) the system admits the cascade decomposition

$$\dot{x}_{\parallel}(t) = f_{\parallel}(t, x_{\parallel}(t)), \quad (36)$$

$$\dot{x}_{\perp}(t) = f_{\perp}(t, x_{\parallel}(t) + x_{\perp}(t)); \quad (37)$$

- (ii) for each $x_{\parallel} \in \mathcal{K}$, the vector field $x_{\perp} \mapsto f_{\perp}(t, x_{\parallel} + x_{\perp})$ is strongly infinitesimally contracting with rate c , with respect to $\|\cdot\|_{\perp} : \mathcal{K}^{\perp} \rightarrow \mathbb{R}_{\geq 0}$;
- (iii) if the map $x_{\parallel} \mapsto f_{\perp}(t, x_{\parallel} + x_{\perp})$ is Lipschitz³ continuous with constant $\ell \in \mathbb{R}$ with respect to some metric $d_{\mathcal{K}}$ on \mathcal{K} , then for any two trajectories $x(t), y(t)$ of (31), satisfying $x(0) - y(0) \in \mathcal{K}^{\perp}$

$$D^+ \|x(t) - y(t)\|^{\mathcal{K}} \leq -c \|x(t) - y(t)\|^{\mathcal{K}} + \ell d_{\mathcal{K}}(x_{\parallel}(t), y_{\parallel}(t))$$

for all $t \in \mathbb{R}_{\geq 0}$, where $D^+(\cdot)$ indicates the upper right Dini derivative [4, Section 2.1].

Proof. The proof follows the same arguments as Theorem 30 for discrete time systems. \square

VI. GRAPH THEORETICAL CONDITIONS FOR SEMICONTRACTIVITY

We now provide graph theoretical conditions for the systems (1) and (2) to be semicontracting with respect to ℓ_p distance and projection seminorms, for $p \in \{1, 2, \infty\}$. For the discrete time case, the following conditions are topological abstractions of algebraic conditions in [16], [23]. Lemma 36 is novel.

Lemma 35 (Topological conditions for discrete-time averaging systems). *The averaging system (1a) $x(k+1) = Ax(k)$ with A row stochastic is strongly semicontracting in the*

- (i) ℓ_1 distance consensus seminorm if A is doubly stochastic and $\mathcal{G}(A)$ is strongly connected and aperiodic;
- (ii) ℓ_2 distance consensus seminorm if A is doubly stochastic and $\mathcal{G}(A)$ is weakly connected with self loops at each node;
- (iii) ℓ_{∞} distance consensus seminorm if $\mathcal{G}(A)$ has self loops at each node and a globally reachable node.

Proof. Condition (i) ensures, in particular, that there exists $m \in \mathbb{N}$ such that A^m has at least $\lfloor \frac{n}{2} \rfloor + 1$ nonzero entries in each column so the expression in (14) takes value less than one. Consequently, the system is strongly semicontracting according to condition (22) in Definition 27.

Condition (ii) directly follows from Lemma 21 and Theorem 8 in [23]. Finally, according to Corollary 4.5 in [5], condition (iii) ensures that there exists $m \in \mathbb{N}$ such that A^m (has a column with all nonzero entries and hence) is scrambling. Consequently, according to Corollary 3.9 in [16] and condition (22) in Definition 27 the system is strongly semicontracting. \square

Strong semicontractivity of Markov chains in the ℓ_p projection seminorms, $p \in \{1, 2, \infty\}$, can be derived by duality.

³That is, for all $x_{\parallel}, y_{\parallel} \in \mathcal{K}$, $z_{\perp} \in \mathcal{K}^{\perp}$, and $t \in \mathbb{R}_{\geq 0}$, we have $\|f_{\perp}(t, x_{\parallel} + z_{\perp}) - f_{\perp}(t, y_{\parallel} + z_{\perp})\|_{\perp} \leq \ell d_{\mathcal{K}}(x_{\parallel}, y_{\parallel})$.

Lemma 36 (Topological conditions for continuous-time averaging). *The averaging system (2) $\dot{x} = -Lx$ with L the Laplacian of a graph with adjacency matrix A and without self-loops, is strongly infinitesimally semicontracting in the*

- (i) ℓ_1 distance consensus seminorm if A is doubly stochastic and every node has at least $\lfloor \frac{n}{2} \rfloor$ in-neighbors,
- (ii) ℓ_2 distance consensus seminorm if A is doubly stochastic and $\mathcal{G}(A)$ is weakly connected,
- (iii) ℓ_{∞} distance consensus seminorm if every two nodes are either (weakly) adjacent or have a common out-neighbor.

Proof. To prove (i), note that $\mu_{\|\cdot\|_{\perp}, \mathcal{K}_{\text{dist},1}}(-L) < 0$ if and only if

$$\sum_{i=1}^{\lfloor \frac{n}{2} \rfloor - 1} a_{(i),j} - \sum_{j=\lceil \frac{n}{2} \rceil}^{n-1} a_{(i),j} < 1, \quad \forall i$$

that for A doubly stochastic is fulfilled if and only if each node has at least $\lfloor \frac{n}{2} \rfloor$ in-neighbors.

To prove (ii) note that for A doubly stochastic $L\Pi_n = \Pi_n L$ and hence the formula for $\mu_{\text{dist},2}(-L)$ in Theorem 24 reads as

$$\mu_{\text{dist},2}(-L) = \min_b \left\{ b : \frac{L + L^T}{2} + bI_n \succeq 0 \text{ on } \mathcal{K}^{\perp} \right\}.$$

The minimum is obtained for $b = -\lambda_2\left(\frac{L+L^T}{2}\right)$ so that, for $\mathcal{G}(A)$ weakly connected, $\mu_{\text{dist},2}(-L) < 0$.

To prove (iii) note that $\mu_{\|\cdot\|_{\perp}, \mathcal{K}_{\text{dist},\infty}}(-L) < 0$ if and only if

$$a_{ij} + a_{ji} + \sum_{k \neq i,j} \min\{a_{ik}, a_{jk}\} > 0 \quad \forall i \neq j$$

that for nonnegative adjacency matrices is true if and only if: (i, j) is an edge or, (j, i) is an edge or (i, k) and (j, k) are an edge for some third node k . \square

VII. CONCLUSIONS

We have studied seminorms on vector spaces and induced matrix seminorms for discrete- and continuous-time dynamical systems. We have shown how the natural distance and projection seminorms are dual and how the long-studied ℓ_p ergodic coefficients of a row-stochastic matrix are precisely induced matrix seminorms. We have provided a comprehensive treatment of semicontraction for discrete- and continuous-time systems with invariance or conservation properties. Future research directions include the application of semicontraction theory to systems with symmetries, such as robotic vehicles ($SE(3)$ symmetry) and coupled oscillators (torus symmetry), as well as systems with invariance properties, such as population games and evolutionary dynamics (whose state space is the simplex). A long-term elusive task is the definition of an ergodic coefficient that is strictly less than unity for row-stochastic matrices satisfying weak connectivity properties.

APPENDIX I SEMINORM COEFFICIENTS

Here we recall some useful properties of standard p -norms. In the following, for a differentiable function $f : \mathbb{R}^n \rightarrow \mathbb{R}$, we denote by $\nabla(f)$ its gradient.

Lemma 37 (Properties of differentiable p -norms [37]). *Let $p \in (1, \infty)$, then $\|\cdot\|_p$ has the following properties:*

- (i) $\|x\|_p$ is differentiable on \mathbb{R}^n ;
- (ii) $\|x\|_p = x^\top \nabla(\|x\|_p)$ for all $x \in \mathbb{R}^n$;
- (iii) $\|\nabla(\|x\|_p)\|_q = 1$ for all $x \neq 0_n$.

Proof. See the final remark and Equation (18) from [37]. \square

Based on Lemma 37, we establish a novel and useful characterization of the distance and projection seminorms.

Lemma 38 (Coefficients for distance seminorms). *Let $p, q \in [1, \infty]$ be such that $p^{-1} + q^{-1} = 1$ and let $\mathcal{K} \subset \mathbb{R}^n$ be a vector subspace. There exists a distance coefficient map $\psi_p : \mathbb{R}^n \rightarrow \mathcal{K}^\perp$ such that, for all $x \in \mathbb{R}^n$,*

- (i) $\psi_p(x) = 0_n$ if $x \in \mathcal{K}$ and $\|\psi_p(x)\|_{\text{proj},q}^\mathcal{K} = 1$ otherwise, and
- (ii) $\|x\|_{\text{dist},p}^\mathcal{K} = \psi_p(x)^\top x$.

Proof. Let $V \in \mathbb{R}^{n \times k}$ be a matrix whose columns are a basis for \mathcal{K} , so that we can write

$$\|x\|_{\text{dist},p}^\mathcal{K} = \min_{\alpha \in \mathbb{R}^k} \|x - V\alpha\|_p$$

At the optimum α^* , 0_k is a subgradient of $\|x - V\alpha^*\|_p$:

$$0_n \in \partial\|x - V\alpha^*\|_p = -V^\top G_p(x - V\alpha^*)$$

where G_p is the subdifferential $G_p = \partial\|\cdot\|_p \subset \mathbb{R}^n$. Consequently, there exists a vector $\psi_p(x) \in G_p(x - V\alpha^*)$ such that $\psi_p(x) \in \ker(V^\top) = \mathcal{K}^\perp$. Note that $\|\psi_p(x)\|_{\text{proj},q}^\mathcal{K} = \|\psi_p(x)\|_q$, that $\psi_p(x)^\top x = \psi_p(x)^\top (x - V\alpha^*)$, and that $\|x\|_{\text{dist},p}^\mathcal{K} = \|x - V\alpha^*\|_p$, so we need only to show for each $p \in [1, \infty]$ that $\|\psi_p(x)\|_q = 1$ and that $\psi_p(x)^\top (x - V\alpha^*) = \|x - V\alpha^*\|_p$.

Case $p = 1$: Using the standard formula for the subgradient of the absolute value function [3], $\psi_1(x) \in G_1(x - V\alpha^*)$ implies that

$$(\psi_1(x))_i = \begin{cases} \text{sgn}((x - V\alpha^*)_i) & (x - V\alpha^*)_i \neq 0 \\ -1 \text{ or } +1, & (x - V\alpha^*)_i = 0 \end{cases}, \quad \forall i$$

If $x \notin \mathcal{K}$, then $x - V\alpha^* \neq 0_n$, so $\|\psi_1(x)\|_\infty = 1$. Furthermore,

$$\begin{aligned} (x - V\alpha^*)^\top \psi_1(x) &= \sum_{i:(x - V\alpha^*)_i \neq 0} (x - V\alpha^*)_i (\psi_1(x))_i \\ &= \sum_{i:(x - V\alpha^*)_i \neq 0} (x - V\alpha^*)_i \text{sgn}((x - V\alpha^*)_i) \\ &= \|x - V\alpha^*\|. \end{aligned}$$

Case $p \in (1, \infty)$: If $p \in (1, \infty)$, then $\|\cdot\|_p$ is differentiable, so $G_p(z) = \nabla\|z\|_p$ for all $z \in \mathbb{R}^n$, and thus $\psi_p(x) = \nabla\|x - V\alpha^*\|_p$ (where the gradient is taken with respect to $x - V\alpha^*$). If $x \notin \mathcal{K}^\perp$, then $x - V\alpha^* \neq 0_n$, so $\|\psi_p(x)\|_q = 1$ due to Lemma 37. A further consequence of this lemma is that

$$(x - V\alpha^*)^\top \psi_p(x) = \|x - V\alpha^*\|_p = \|x\|_{\text{dist},p}^\mathcal{K}$$

Case $p = \infty$: Let $\mathcal{I} \subseteq \{1, 2, \dots, n\}$ be the set of indices such that $\|x - V\alpha^*\|_\infty = |x - V\alpha^*|_i$. Using a standard formula for the subdifferential of a pointwise maximum [3], $\psi_\infty(x) \in G_\infty(x - V\alpha^*)$ implies that

$$\psi_\infty(x) \in \text{conv} \bigcup_{i \in \mathcal{I}} \partial|x - V\alpha^*|_i$$

where conv denotes the convex hull, and the subdifferential of each absolute value is with respect to its argument. Therefore, there exist $g_i \in \partial|x - V\alpha^*|_i$ for each $i \in \mathcal{I}$, as well as convex weights λ_i , such that

$$\psi_\infty(x) = \sum_{i \in \mathcal{I}} \lambda_i g_i$$

For each g_i , we have $[g_i]_j = 0$ for $j \neq i$, since $|z|_i$ only depends on z_i for any $z \in \mathbb{R}^n$. Furthermore, if $x \notin \mathcal{K}$, then $x - V\alpha^* \neq 0_n$, so $|x - V\alpha^*|_i > 0$ for all $i \in \mathcal{I}$, which implies that $[g_i]_i = \text{sgn}(x - V\alpha^*)_i$. Together, these two observations imply that

$$\|\psi_\infty(x)\|_1 = \sum_{i \in \mathcal{I}} \lambda_i \|g_i\|_1 = \sum_{i \in \mathcal{I}} \lambda_i = 1$$

Finally,

$$\begin{aligned} (x - V\alpha^*)^\top \psi_\infty(x) &= \sum_{i \in \mathcal{I}} (x - V\alpha^*)_i \sum_{j \in \mathcal{I}} \lambda_j [g_j]_i \\ &= \sum_{i \in \mathcal{I}} \lambda_i (x - V\alpha^*)_i \text{sgn}(x - V\alpha^*)_i = \|x - V\alpha^*\|_\infty \end{aligned}$$

\square

Lemma 39 (Coefficients for projection seminorms). *Let $p, q \in [1, \infty]$ be such that $p^{-1} + q^{-1} = 1$ and $\mathcal{K} \subset \mathbb{R}^n$ be a vector subspace. There exists a projection coefficient map $\zeta_p : \mathbb{R}^n \rightarrow \mathcal{K}^\perp$ such that, for all $x \in \mathbb{R}^n$,*

- (i) $\zeta_p(x) = 0_n$ if $x \in \mathcal{K}$ and $\|\zeta_p(x)\|_{\text{dist},q}^\mathcal{K} \leq 1$ otherwise, and
- (ii) $\|x\|_{\text{proj},p}^\mathcal{K} = \zeta_p(x)^\top x$.

Proof. Let $x \in \mathbb{R}^n$ and define $x_\perp = \Pi_\perp x$.

Case $p = 1$: Let $\zeta_1(x) = \Pi_\perp \text{sgn}(x_\perp)$. By Lemmas 2 and 9 (ii),

$$\|\zeta_1(x)\|_{\text{dist},\infty}^\mathcal{K} = \|\text{sgn}(x_\perp)\|_{\text{dist},\infty}^\mathcal{K} \leq \|\text{sgn}(x_\perp)\|_\infty \leq 1,$$

where $x \in \mathcal{K}$ implies that $\text{sgn}(x_\perp) = 0_n$. Furthermore,

$$\zeta_1(x)^\top x = \text{sgn}(x_\perp)^\top \Pi_\perp x = \|x_\perp\|_1 = \|x\|_{\text{proj},1}^\mathcal{K}.$$

Case $p \in (1, \infty)$: Let $\zeta_p(x) = \Pi_\perp \nabla(\|x_\perp\|_p)$. By Lemmas 2, 9 (ii), and 37 (iii), if $x \notin \mathcal{K}$, then

$$\|\zeta_p(x)\|_{\text{dist},q}^\mathcal{K} = \|\nabla(\|x_\perp\|_p)\|_{\text{dist},q}^\mathcal{K} \leq \|\nabla(\|x_\perp\|_p)\|_q = 1.$$

But if $x \in \mathcal{K}$, then $x_\perp = 0_n$, so $\zeta_p(x) = 0_n$. Furthermore, as a consequence of Lemma 37 (ii),

$$\zeta_p(x)^\top x = (\nabla(\|x_\perp\|_p))^\top \Pi_\perp x = \|x_\perp\|_p = \|x\|_{\text{proj},p}^\mathcal{K}.$$

Case $p = \infty$: Let $i \in \{1, 2, \dots, n\}$ be such that $\|x_\perp\|_\infty = |x_\perp|_i$, and let $\zeta_\infty(x) = \text{sgn}(x_\perp)_i \Pi_\perp e_i$. By Lemmas 2 and 9 (ii),

$$\|\zeta_\infty(x)\|_{\text{dist},1}^{\mathcal{K}} = \|\text{sgn}(x_\perp)_i e_i\|_{\text{dist},1}^{\mathcal{K}} \leq \|\text{sgn}(x_\perp)_i e_i\|_1 \leq 1$$

where $x \in \mathcal{K}$ implies that $\text{sgn}(x_\perp) = 0_n$. Furthermore,

$$\zeta_\infty(x)^\top x = \text{sgn}(x_\perp)_i e_i^\top x_\perp = \|x_\perp\|_\infty = \|x\|_{\text{proj},\infty}^{\mathcal{K}}.$$

□

APPENDIX II CONJECTURES

Conjecture 40 (Optimal deflation and distance seminorm). *Here a conjecture on the equivalence between ℓ_p distance seminorm and p -optimal deflation as in Definition 20. For each $p \in [1, \infty]$ and row-stochastic matrix $A \in \mathbb{R}^{n \times n}$,*

$$|A|_p = \|A\|_{\text{dist},p}^{\mathcal{K}}.$$

Here some reasons in support of this conjecture.


- (i) Expressions given in [22] for $p \in \{1, 2, \infty\}$ of $x \in \mathbb{R}^n$ and of $A \in \mathbb{R}^{n \times n}$ row stochastic, of $|x|_p$ and $|A|_p$ coincide with the ones of $\|x\|_{\text{dist},p}^{\mathcal{K}}$ and $\|A\|_{\text{dist},p}^{\mathcal{K}}$, respectively.
- (ii) If the envelope theorem [35, Theorem 1.F.1] could be applied⁴ to the projection seminorm, with kernel $\mathcal{K} = \text{span}\{\mathbb{1}_n\}$, it would lead to the orthogonality constraint with respect to \mathcal{K} and consequently to the equivalence between $|A|_p$ and $\|A\|_{\text{proj},q}^{\mathcal{K}}$.

REFERENCES

- [1] Z. Askarzadeh, R. Fu, A. Halder, Y. Chen, and T. T. Georgiou. Stability theory of stochastic models in opinion dynamics. *IEEE Transactions on Automatic Control*, 65(2):522–533, 2020. doi:10.1109/TAC.2019.2912490.
- [2] Stefan Banach. Sur les opérations dans les ensembles abstraits et leur application aux équations intégrales. *Fundamenta Mathematicae*, 3(1):133–181, 1922. doi:10.4064/fm-3-1-133-181.
- [3] S. Boyd and L. Vandenberghe. *Convex Optimization*. Cambridge University Press, 2004, ISBN 0521833787.
- [4] F. Bullo. *Contraction Theory for Dynamical Systems*. Kindle Direct Publishing, 1.0 edition, 2022, ISBN 979-8836646806. URL: <http://motion.me.ucsb.edu/book-ctds>.
- [5] F. Bullo. *Lectures on Network Systems*. Kindle Direct Publishing, 1.6 edition, January 2022, ISBN 978-1986425643. URL: <http://motion.me.ucsb.edu/book-lns>.
- [6] B. Charron-Bost. Orientation and connectivity based criteria for asymptotic consensus, March 2013. Available at <https://arxiv.org/abs/1303.2043>.
- [7] B. P. Demidovič. Dissipativity of a nonlinear system of differential equations. *Uspekhi Matematicheskikh Nauk*, 16(3(99)):216, 1961.
- [8] C. A. Desoer and M. Vidyasagar. *Feedback Systems: Input-Output Properties*. Academic Press, 1975, ISBN 978-0-12-212050-3. doi:10.1137/1.9780898719055.
- [9] M. Di Bernardo, D. Fiore, G. Russo, and F. Scafoli. Convergence, consensus and synchronization of complex networks via contraction theory. In *Complex Systems and Networks*, pages 313–339. Springer, 2016. doi:10.1007/978-3-662-47824-0_12.
- [10] R. L. Dobrushin. Central limit theorem for nonstationary Markov chains. I. *Theory of Probability & Its Applications*, 1(1):65–80, 1956. doi:10.1137/1101006.
- [11] W. Doeblin. Le cas discontinu des probabilités en chaîne. *Publ. Faculty of Science University Masaryk (Brno)*, pages 3–13, 1937.
- [12] F. Forni and R. Sepulchre. A differential Lyapunov framework for contraction analysis. *IEEE Transactions on Automatic Control*, 59(3):614–628, 2014. doi:10.1109/TAC.2013.2285771.
- [13] D. J. Hartfiel. *Markov Set-Chains*. Springer, 1998, ISBN 978-3-540-68711-5. doi:10.1007/BFb0094586.
- [14] M. Haviv and L. van der Heyden. Perturbation bounds for the stationary probabilities of a finite Markov chain. *Advances in Applied Probability*, 16(4):804–818, 1984. doi:10.2307/1427341.
- [15] R. A. Horn and C. R. Johnson. *Matrix Analysis*. Cambridge University Press, 1985, ISBN 0521386322.
- [16] I. C. F. Ipsen and T. M. Selee. Ergodicity coefficients defined by vector norms. *SIAM Journal on Matrix Analysis and Applications*, 32(1):153–200, 2011. doi:10.1137/090752948.
- [17] S. Jafarpour, P. Cisneros-Velarde, and F. Bullo. Weak and semi-contraction for network systems and diffusively-coupled oscillators. *IEEE Transactions on Automatic Control*, 67(3):1285–1300, 2022. doi:10.1109/TAC.2021.3073096.
- [18] A. N. Kolmogorov. Über die analytischen Methoden in der Wahrscheinlichkeitsrechnung. *Mathematische Annalen*, 104:415–158, 1931. doi:10.1007/BF01457949.
- [19] V. V. Korpakov. Matrix seminorms and related inequalities. *Journal of Soviet Mathematics*, 23:2094–2106, 1983. doi:10.1007/BF01093289.
- [20] N. N. Krasovskii. *Stability of Motion. Applications of Lyapunov's Second Method to Differential Systems and Equations with Delay*. Stanford University Press, 1963. Translation of the 1959 edition in Russian by J. L. Brenner.
- [21] D. A. Levin and Y. Peres. *Markov Chains and Mixing Times*. American Mathematical Society, 2 edition, 2017, ISBN 978147042962. With contributions by E. L. Wilmer.
- [22] J. Liu, A. S. Morse, B. D. O. Anderson, and C. Yu. Contractions for consensus processes. In *IEEE Conf. on Decision and Control and European Control Conference*, pages 1974–1979, 2011. doi:10.1109/CDC.2011.6160989.
- [23] J. Liu, S. Mou, A. S. Morse, B. D. O. Anderson, and C. Yu. Deterministic gossiping. *Proceedings of the IEEE*, 99(9):1505–1524, 2011. doi:10.1109/JPROC.2011.2159689.
- [24] W. Lohmiller and J.-J. E. Slotine. On contraction analysis for nonlinear systems. *Automatica*, 34(6):683–696, 1998. doi:10.1016/S0005-1098(98)00019-3.
- [25] Andrey A. Markov. Extensions of the law of large numbers to dependent quantities. *Izvestiya Fiziko-matematicheskogo obschestva pri Kazanskom universitete*, 15, 1906. (in Russian).
- [26] R. Marsli and F. J. Hall. Some properties of ergodicity coefficients with applications in spectral graph theory. *Linear and Multilinear Algebra*, 0(0):1–21, 2020. doi:10.1080/03081087.2020.1777251.
- [27] U. G. Rothblum. Algebraic eigenspaces of nonnegative matrices. *Linear Algebra and its Applications*, 12(3):282–292, 1975. doi:10.1016/0024-3795(75)90050-6.
- [28] U. G. Rothblum and C. P. Tan. Upper bounds and the maximum modulus of subdominant eigenvalues of nonnegative matrices. In *Linear Algebra and its Applications*, volume 66, pages 45–86, 1985. doi:10.1016/0024-3795(85)90125-9.
- [29] E. Seneta. *Non-negative Matrices and Markov Chains*. Springer, 2 edition, 1981, ISBN 0387297650.
- [30] E. Seneta. Explicit forms for ergodicity coefficients and spectrum localization. *Linear Algebra and its Applications*, 60:187–197, 1984. doi:10.1016/0024-3795(84)90079-X.
- [31] E. Seneta. Markov and the creation of Markov chains. In A. N. Langville and W. J. Stewart, editors, *Markov Anniversary Meeting*, pages 1–20. C & M Online Media, 2006. URL: <https://www.csc2.ncsu.edu/conferences/nsmc>.
- [32] J. W. Simpson-Porco and F. Bullo. Contraction theory on Riemannian manifolds. *Systems & Control Letters*, 65:74–80, 2014. doi:10.1016/j.sysconle.2013.12.016.
- [33] J.-J. E. Slotine. Modular stability tools for distributed computation and control. *International Journal of Adaptive Control and Signal Processing*, 17(6):397–416, 2003. doi:10.1002/acs.754.
- [34] A. Tahbaz-Salehi and A. Jadbabaie. A necessary and sufficient condition for consensus over random networks. *IEEE Transactions on Automatic Control*, 53(3):791–795, 2008. doi:10.1109/TAC.2008.917743.
- [35] A. Takayama. *Mathematical Economics*. Cambridge University Press, 1985, ISBN 0521314984.
- [36] W. Wang and J. J. Slotine. On partial contraction analysis for coupled nonlinear oscillators. *Biological Cybernetics*, 92(1):38–53, 2005. doi:10.1007/s00422-004-0527-x.

⁴It requires to prove the continuous differentiability of v in (20) as function of the inner optimization vector related to the induced matrix norm.

- [37] W. H. Yang. On generalized Hölder inequality. *Nonlinear Analysis, Theory, Methods & Applications*, 16(5), 1990. doi:10.1016/0362-546X(91)90072-9.



Giulia De Pasquale (S-'19) received the B.Sc. and the M.Sc degree in Information Engineering and Systems and Control Engineering from the University of Padova, Italy, in 2017 and 2019, respectively, where she is currently pursuing the Ph.D. degree with the Department of Information Engineering. In 2022 she was visiting research scholar at the University of California, Santa Barbara. In 2018 and 2019 she was a visiting student at LTU, Sweden and ETH Zürich, respectively. Her current research interests are in

the analysis and control of networked systems, with a special focus on opinion dynamics.



Kevin D. Smith (S-'16) received the B.S. degree in physics from Harvey Mudd College, Claremont, CA, USA, in 2017 and the M.S. degree in 2019 from the University of California, Santa Barbara, where he is currently working toward the Ph.D. degree with the Center for Control, Dynamical Systems, and Computation. His research interests include dynamics, control, and identification of network systems, particularly infrastructure networks.



Francesco Bullo (S-'95 M-'99 SM-'03 F-'10) is a Distinguished Professor of Mechanical Engineering with the University of California, Santa Barbara, CA, USA. He was previously with the University of Padova (Laurea degree, 1994), Italy, the California Institute of Technology (Ph.D. degree, 1998), Pasadena, CA, and the University of Illinois at Urbana-Champaign, IL, USA. His research interests include contraction theory, network systems, and distributed control. He is the author or coauthor of *Geometric Control*

of Mechanical Systems (Springer, 2004), *Distributed Control of Robotic Networks* (Princeton, 2009), *Lectures on Network Systems* (KDP, 2022), and *Contraction Theory for Dynamical Systems* (KDP, 2022, v1.0). He served as IEEE CSS President and SIAG CST Chair. He is a Fellow of ASME, IEEE, IFAC, and SIAM.



M. Elena Valcher (M-'99 SM-'03 F-'12) received the Master degree (1991) and the PhD degree (1995) from the University of Padova. Since 2005 she is full professor at the University of Padova. She is author/co-author of 87 journal papers, 105 conference papers, 2 text-books and several book chapters. Her research interests include social networks, cooperative control and consensus, positive switched systems and Boolean control networks. She is the Founding Editor in Chief of the *IEEE Control Systems*

Letters (2017-). She was IEEE CSS President (2015). She received the 2011 IEEE CSS Distinguished Member Award and she is an IEEE Fellow since 2012. She is a member of the IFAC Technical Board (2017-2020) and of the EUCA Board (2017-).