

An elementary alternative to ECH capacities

Michael Hutchings*

Abstract

The ECH capacities are a sequence of numerical invariants of symplectic four-manifolds which give (sometimes sharp) obstructions to symplectic embeddings. These capacities are defined using embedded contact homology, and establishing their basic properties currently requires Seiberg-Witten theory. In this note we define a new sequence of symplectic capacities in four dimensions using only basic notions of holomorphic curves. The new capacities satisfy the same basic properties as ECH capacities and agree with the ECH capacities for the main examples for which the latter have been computed, namely convex and concave toric domains. The new capacities are also useful for obstructing symplectic embeddings into closed symplectic four-manifolds. This work is inspired by a recent preprint of McDuff-Siegel giving a similar elementary alternative to symplectic capacities from rational SFT.

Contents

1	Introduction	2
2	Definition of the capacities c_k	3
3	Proof of the monotonicity lemma	6
4	Properties of the capacities c_k	8
5	Computation for convex toric domains	11
6	Comparison with ECH capacities	13
7	Additional computations using Seiberg-Witten theory	15
8	An even simpler definition of capacities	18

*Partially supported by NSF grant DMS-2005437.

1 Introduction

We define a *symplectic capacity* to be a function c which maps some set of symplectic manifolds (possibly noncompact, disconnected, and/or with boundary or corners) to $[0, \infty]$. We assume the following two properties:

(Monotonicity) If (X, ω) and (X', ω') are symplectic manifolds of the same dimension for which c is defined, and if there exists a symplectic embedding $\varphi : (X, \omega) \rightarrow (X', \omega')$, then $c(X, \omega) \leq c(X', \omega')$.

(Conformality) If $r > 0$ then $c(X, r\omega) = rc(X, \omega)$.

Various symplectic capacities are used to study symplectic embedding problems, in particular to give obstructions via the Monotonicity property. See e.g. the survey [5].

Perhaps the most basic example of a symplectic capacity is the *Gromov width* c_{Gr} . For $a > 0$, define the ball

$$B^{2n}(a) = \{z \in \mathbb{C}^n \mid \pi|z|^2 \leq a\}$$

with the restriction of the standard symplectic form on $\mathbb{C}^n = \mathbb{R}^{2n}$. If $\dim(X) = 2n$, then $c_{\text{Gr}}(X, \omega)$ is defined to be the supremum over a such that there exists a symplectic embedding $B^{2n}(a) \rightarrow (X, \omega)$. The celebrated Gromov nonsqueezing theorem [14] is equivalent to the statement that the cylinder

$$Z^{2n}(a) = \{z \in \mathbb{C}^n \mid \pi|z_1|^2 \leq a\}$$

has Gromov width equal to a .

While the Gromov width has a very simple definition, it is difficult to use by itself for studying symplectic embedding problems, since it is defined in terms of symplectic embeddings. In general, there is a gap we would like to bridge between (1) symplectic capacities with simple geometric definitions that can be hard to compute, such as the Gromov width; and (2) symplectic capacities defined using Floer-theoretic or related machinery which are more computable, but whose definition requires substantial technical work.

One example of the latter type of capacity is the sequence of Ekeland-Hofer capacities defined using variational methods in [11], or the conjecturally equivalent capacities defined in [16] using positive S^1 -equivariant symplectic homology.

Another example, which is the focus of the present paper, is the sequence of ECH capacities introduced in [20]; see the expositions in [21, 22] and §6 below. Let (X, ω) be a symplectic four-manifold, not necessarily closed or connected. The ECH capacities of (X, ω) are a sequence of real numbers

$$0 = c_0^{\text{ECH}}(X, \omega) < c_1^{\text{ECH}}(X, \omega) \leq c_2^{\text{ECH}}(X, \omega) \leq \cdots \leq +\infty.$$

Monotonicity of ECH capacities means that if (X', ω') is another symplectic four-manifold, and if there exists a symplectic embedding $(X, \omega) \rightarrow (X', \omega')$, then

$$c_k^{\text{ECH}}(X, \omega) \leq c_k^{\text{ECH}}(X', \omega') \quad (1.1)$$

for all k . This obstruction is known to be sharp in some cases, for example when X and X' are open ellipsoids in \mathbb{R}^4 with the restriction of the standard symplectic form, as shown by McDuff [31]. More generally, Cristofaro-Gardiner [7] showed that the obstruction is sharp when X is an open “concave toric domain” and X' is a “convex toric domain”; see the definitions in §5 below. The ECH capacities are defined using embedded contact homology, and the proof of the symplectic embedding obstruction (1.1) uses cobordism maps on embedded contact homology, which currently need to be defined using Seiberg-Witten theory¹.

More recently, Siegel [35] used rational symplectic field theory [12], whose technical foundations in general are still a work in progress, to define a set of symplectic capacities which are well suited to studying stabilized symplectic embedding problems. Subsequently, McDuff-Siegel [32] showed that for key applications, some of Siegel’s capacities can be replaced by an alternate set of capacities with a more elementary definition directly in terms of holomorphic curves with local tangency constraints. More generally, one can hope that capacities extracted from Floer theories can be understood geometrically without passing through Floer theory, or at least can be replaced by more elementary capacities with the same applications.

The purpose of this note is to pursue this direction for the ECH capacities. Namely we give an elementary definition of a sequence of symplectic capacities for symplectic four-manifolds, which we denote by c_k , which are defined directly in terms of holomorphic curves constrained to pass through k points. We show that the capacities c_k have the same basic properties as ECH capacities and agree with them in important examples. In particular, this allows some of the applications of ECH capacities to be re-proved without using Seiberg-Witten theory. The capacities c_k also give good obstructions to symplectic embeddings into some closed symplectic four-manifold such as $\mathbb{C}P^2$ or $S^2 \times S^2$, whose ECH capacities are not known. At the end, we define an even simpler sequence of capacities \widehat{c}_k in any dimension, which conjecturally agree with the capacities c_k in the main four-dimensional cases.

Acknowledgment. Thanks to Grisha Mikhalkin and Kyler Siegel for helpful discussion and comments on an initial draft of this note.

2 Definition of the capacities c_k

We begin by recalling some basic definitions.

¹Heuristically one might expect to define such a cobordism map just by counting holomorphic curves. Although this is possible in some special cases [13, 34], in general there are severe transversality difficulties with multiply covered curves; see [22, §5.5] for explanation.

Let Y be a three-manifold and let λ be a contact form on Y . Let $\xi = \text{Ker}(\lambda)$ denote the associated contact structure, and let R denote the associated Reeb vector field. Define an *orbit set* to be a finite set of pairs $\alpha = \{(\alpha_i, m_i)\}$ where the α_i are distinct simple Reeb orbits, and the m_i are positive integers. Define the *symplectic action* of the orbit set α by

$$\mathcal{A}(\alpha) = \sum_i m_i \int_{\alpha_i} \lambda.$$

The contact form λ is *nondegenerate* if every Reeb orbit (simple or multiply covered) is nondegenerate, i.e. the linearized return map does not have 1 as an eigenvalue.

We say that an almost complex structure J on $\mathbb{R} \times Y$ is λ -*compatible* if $J\partial_s = R$, where s denotes the \mathbb{R} coordinate; J sends the contact structure ξ to itself, rotating positively with respect to $d\lambda$; and J is \mathbb{R} -invariant.

We define a four-dimensional *Liouville domain* to be a compact symplectic four-manifold (X, ω) with boundary Y such that there exists a primitive of ω which restricts to a contact form λ on Y . A basic example is a star-shaped domain in \mathbb{R}^4 . Here a “star-shaped domain” is a compact domain in \mathbb{R}^{2n} with smooth boundary which is transverse to the radial vector field, with the restriction of the standard symplectic form. We say that the Liouville domain (X, ω) is *nondegenerate* if the contact form λ on Y is nondegenerate; this notion does not depend on the choice of primitive of ω .

Given a Liouville domain as above, and given $\varepsilon > 0$, a choice of primitive of ω determines a neighborhood N_ε of Y in X , and an identification

$$N_\varepsilon \simeq (-\varepsilon, 0] \times Y, \tag{2.1}$$

under which $\omega|_{N_\varepsilon}$ is identified with $d(e^s\lambda)$, where s denotes the $(-\varepsilon, 0]$ coordinate. Using this identification, we can glue to obtain a smooth manifold

$$\overline{X} = X \cup_Y ([0, \infty) \times Y), \tag{2.2}$$

which we call the “symplectization completion” of X . This has a symplectic form $\overline{\omega}$ which agrees with ω on X and with $d(e^s\lambda)$ on $[0, \infty) \times Y$. Strictly speaking, this completion depends on the choice of primitive of ω , which we suppress from the notation.

We say that an almost complex structure J on \overline{X} is *cobordism-compatible* if $J|_X$ is ω -compatible, and if $J|_{[0, \infty) \times Y}$ is the restriction of a λ -compatible almost complex structure on $\mathbb{R} \times Y$.

Define an *admissible symplectic four-manifold* to be a (possibly disconnected) compact symplectic four-manifold (X, ω) such that each component is either closed or a nondegenerate Liouville domain. Define \overline{X} to be the union of the closed components and the symplectization completions of the Liouville domain components.

Define $\mathcal{J}(\overline{X}, \omega)$ to be the set of almost complex structures on \overline{X} which are ω -compatible on the closed components and cobordism-compatible on the completed Liouville domain components.

Let $J \in \mathcal{J}(\overline{X}, \omega)$. We consider holomorphic maps

$$u : (\Sigma, j) \longrightarrow (\overline{X}, J)$$

where Σ is a punctured compact Riemann surface (possibly disconnected), such that for each puncture in Σ , there is a Reeb orbit γ on ∂X and a neighborhood of the puncture mapping asymptotically to $[0, \infty) \times \gamma$ as $s \rightarrow \infty$. To avoid trivialities we assume that the restriction of u to each component of the domain Σ is nonconstant. Let $\mathcal{M}^J(\overline{X})$ denote the set of J -holomorphic maps as above, modulo reparametrization by biholomorphic maps $(\Sigma', j') \xrightarrow{\cong} (\Sigma, j)$. If $x_1, \dots, x_k \in X$ are distinct points, let $\mathcal{M}^J(\overline{X}; x_1, \dots, x_k)$ denote the set of $u \in \mathcal{M}^J(\overline{X})$ such that $x_1, \dots, x_k \in u(\Sigma)$.

Define the *energy* $\mathcal{E}(u)$ as follows. If Σ is connected and u maps to a closed component of X , then $\mathcal{E}(u) = \int_{\Sigma} u^* \omega$. If Σ is connected and u maps to a completed Liouville domain component, then $\mathcal{E}(u)$ is the sum over the punctures of Σ of the symplectic actions of the corresponding Reeb orbits. If Σ is disconnected, then $\mathcal{E}(u)$ is the sum of the energies of the connected components.

Definition 2.1. Let (X, ω) be an admissible symplectic four-manifold and let k be a nonnegative integer. Define

$$c_k(X, \omega) = \sup_{\substack{J \in \mathcal{J}(\overline{X}) \\ x_1, \dots, x_k \in X \text{ distinct}}} \inf_{u \in \mathcal{M}^J(\overline{X}; x_1, \dots, x_k)} \mathcal{E}(u) \in [0, \infty]. \quad (2.3)$$

Remark 2.2. A key observation, which avoids various technical difficulties, is that in (2.3), we can restrict attention to holomorphic curves u that *do not have any multiply covered components*². This is because we can always replace a multiply covered component by the underlying somewhere injective curve to reduce energy without invalidating the point constraints.

Lemma 2.3. (proved in §3) *Let (X, ω) and (X', ω') be admissible symplectic four-manifolds and let k be a nonnegative integer. If there exists a symplectic embedding $\varphi : (X, \omega) \rightarrow (X', \omega')$, then*

$$c_k(X, \omega) \leq c_k(X', \omega').$$

To extend the definition of c_k to more general symplectic four-manifolds, we use the following basic procedure; compare [20, §4.2].

²We say that $u : \Sigma \rightarrow \overline{X}$ “has no multiply covered components” if the restriction of the map u to each component of the domain Σ is not multiply covered (which means that it must be somewhere injective), and no two components of Σ have the same image under u .

Definition 2.4. Let (X', ω') be any symplectic four-manifold (possibly noncompact, disconnected, and/or with boundary or corners) and let k be a nonnegative integer. Define

$$c_k(X', \omega') = \sup\{c_k(X, \omega)\}$$

where the supremum is over admissible symplectic four-manifolds (X, ω) for which there exists a symplectic embedding $\varphi : (X, \omega) \rightarrow (X', \omega')$.

It follows from Lemma 2.3 that Definition 2.4 agrees with Definition 2.1 when (X', ω') is already an admissible symplectic four-manifold.

Remark 2.5. The definition of c_k is inspired by the paper of McDuff-Siegel [32], which gives a similar elementary definition of a sequence of symplectic capacities $\tilde{\mathfrak{g}}_k$, as an alternative to symplectic capacities that were defined in [35] using rational symplectic field theory [12]. The capacities $\tilde{\mathfrak{g}}_k$ are defined for symplectic manifolds of any dimension using genus zero holomorphic curves that are constrained to have contact of order k with a local divisor.

Some variants of Definition 2.1 are possible. For example one could require each component of the domain of u to have genus zero; the resulting capacities may be related to the capacities $\tilde{\mathfrak{g}}_k$.

3 Proof of the monotonicity lemma

Before discussing the basic properties of the capacities c_k in §4 below, we now prove Lemma 2.3.

The following notation will be useful. Let (X, ω) be an admissible symplectic four-manifold and let $\alpha = \{(\alpha_i, m_i)\}$ be an orbit set for $Y = \partial X$. Let $H_2(X, \alpha)$ denote the set of relative homology classes of 2-chains Z in X with $\partial Z = \alpha$. This set is an affine space over $H_2(X)$.

Given $J \in \mathcal{J}(\overline{X}, \omega)$, let $\mathcal{M}^J(\overline{X}, \alpha; x_1, \dots, x_k)$ denote the set of holomorphic curves in $\mathcal{M}^J(\overline{X}; x_1, \dots, x_k)$ such that for each i , there are punctures asymptotic to covers of α_i with total multiplicity m_i , and there are no other punctures. Note that each $u \in \mathcal{M}^J(\overline{X}, \alpha; x_1, \dots, x_k)$ has a well-defined relative homology class $[u] \in H_2(X, \alpha)$.

Proof of Lemma 2.3. For $\varepsilon > 0$, let N_ε denote the neighborhood of ∂X in (2.1). The time ε flow of the Liouville vector field (coming from the primitive of ω) defines a symplectomorphism

$$(X \setminus N_\varepsilon, \omega|_{X \setminus N_\varepsilon}) \simeq (X, e^{-\varepsilon} \omega). \quad (3.1)$$

It follows from Definition 2.1 that c_k satisfies the Conformality property, so we deduce from (3.1) that

$$c_k(X \setminus N_\varepsilon, \omega|_{X \setminus N_\varepsilon}) = e^{-\varepsilon} c_k(X, \omega) \quad (3.2)$$

Consequently, by replacing X with $X \setminus N_\varepsilon$ for $\varepsilon > 0$ small if necessary, we can assume without loss of generality that $\varphi(X) \subset \text{int}(X')$.

Now fix $x_1, \dots, x_k \in X$ distinct, $J \in \mathcal{J}(\overline{X}, \omega)$, and $\varepsilon > 0$. To prove the lemma, we need to show that there exists $u \in \mathcal{M}^J(\overline{X}; x_1, \dots, x_k)$ with

$$\mathcal{E}(u) < c_k(X', \omega') + \varepsilon.$$

We will use a “neck stretching” argument.

Write $Y = \partial X$ and let λ denote the contact form on Y . Since $\varphi(X) \subset \text{int}(X')$, there exists a neighborhood \mathcal{U} of $\varphi(Y)$ in $X' \setminus \varphi(\text{int}(X))$ and an identification

$$(\mathcal{U}, \omega'|_{\mathcal{U}}) \simeq ([0, \delta) \times Y, d(e^s \lambda))$$

for some $\delta > 0$, where s denotes the $[0, \delta)$ coordinate. For each $R > 0$, we can choose an almost complex structure $J_R \in \mathcal{J}(\overline{X'}, \omega')$ such that φ extends to a biholomorphism

$$\varphi_R : (X \cup_Y ([0, R) \times Y), J) \xrightarrow{\simeq} (\varphi(X) \cup \mathcal{U}, J_R). \quad (3.3)$$

We can further assume that J_R is independent of R outside of $\varphi(X) \cup \mathcal{U}$.

By the definition of c_k , for each R we can choose

$$u_R \in \mathcal{M}^{J_R}(\overline{X'}; \varphi(x_1), \dots, \varphi(x_k))$$

with

$$\mathcal{E}(u_R) < c_k(X', \omega') + \varepsilon \quad (3.4)$$

Let u_R^φ denote the intersection of the curve u_R with $\varphi(X) \cup \mathcal{U}$, composed with φ_R^{-1} . We now want to argue that there is a sequence $R_i \rightarrow \infty$ such that the intersections of the curves $u_{R_i}^\varphi$ converge in some sense to the desired curve u . This task is complicated by the fact that we do not have an a priori bound on the genus of the components of the domains of the curves u_R , so we cannot directly use SFT compactness as in [1, 6].

Fortunately, there is a local version of Gromov compactness using currents which does not require any genus bound. This was proved in the four-dimensional case by Taubes [36, Prop. 3.3], and an updated version which works in arbitrary dimension was proved by Doan-Wapulski [10, Prop. 1.9]. By this local Gromov compactness and the energy bound (3.4), as applied in [18, §9.4], we can find a sequence $R_i \rightarrow \infty$ such that the curves $u_{R_i}^\varphi$ converge as currents to a proper holomorphic map u to \overline{X} which passes through the points x_1, \dots, x_k , is asymptotic as a current as $s \rightarrow \infty$ to an orbit set for Y , and has energy less than $c_k(X', \omega') + \varepsilon$. A priori, components of the domain of u may have infinite genus, and to complete the proof of the lemma we need to arrange that they have finite genus.

We can choose the sequence R_i so that each u_{R_i} is in $\mathcal{M}^{J_{R_i}}(\overline{X'}, \alpha'; \varphi(x_1), \dots, \varphi(x_k))$ for a fixed orbit set α' for $\partial X'$, because there are only finitely many orbit sets for

$\partial X'$ with action less than $c_k(X', \omega') + \varepsilon$. When applying Gromov compactness above, we can further use the arguments in [18, §9.4] to chase down the rest of the energy of the holomorphic curves u_{R_i} and pass to a subsequence such that the relative homology class $[u_{R_i}] \in H_2(X', \alpha')$ does not depend on i .

By Remark 2.2, we can assume that each u_{R_i} has no multiply covered components. Since we are in four dimensions, the relative adjunction formula of [19, Prop. 4.9] and the asymptotic writhe bound of [19, Lem. 4.20] imply that there is a lower bound on the Euler characteristic of the domain of u_{R_i} depending only on the orbit set α' and the relative homology class $[u_{R_i}]$. We can also assume that each u_{R_i} has at most k components, since otherwise some components can be discarded without violating the requirement to pass through the points x_1, \dots, x_k . Consequently we obtain an i -independent upper bound on the genus of each component of u_{R_i} , and hence of the limiting curve u . \square

4 Properties of the capacities c_k

Theorem 4.1. *The capacities c_k of four-dimensional symplectic manifolds have the following properties:*

(Conformality) If $r > 0$ then

$$c_k(X, r\omega) = r c_k(X, \omega). \quad (4.1)$$

(Increasing)

$$0 = c_0(X, \omega) < c_1(X, \omega) \leq c_2(X, \omega) \leq \dots \leq +\infty.$$

(Disjoint Union)

$$c_k \left(\prod_{i=1}^m (X_i, \omega_i) \right) = \max_{k_1 + \dots + k_m = k} \sum_{i=1}^m c_{k_i}(X_i, \omega_i).$$

(Sublinearity)

$$c_{k+l}(X, \omega) \leq c_k(X, \omega) + c_l(X, \omega).$$

(Monotonicity) If there exists a symplectic embedding $\varphi : (X, \omega) \rightarrow (X', \omega')$, then

$$c_k(X, \omega) \leq c_k(X', \omega').$$

(C^0 -Continuity) For each k , the capacity c_k defines a continuous function on the set of star-shaped domains in \mathbb{R}^4 with respect to the Hausdorff metric on compact sets.

(Spectrality) If (X, ω) is a four-dimensional Liouville domain with boundary Y , then for each k with $c_k(X, \omega) < \infty$, there exists an orbit set α in Y with $c_k(X, \omega) = \mathcal{A}(\alpha)$.

(ECH Index) If X is a nondegenerate star-shaped domain in \mathbb{R}^4 , then $c_k(X) < \infty$, and in the Spectrality property, we can choose α so that its ECH index³ satisfies $I(\alpha) \geq 2k$.

(Ball)

$$c_k(B^4(a)) = da$$

where d is the unique nonnegative integer with

$$d^2 + d \leq 2k \leq d^2 + 3d.$$

(Asymptotics) If $X \subset \mathbb{R}^4$ is a compact domain with smooth boundary, then

$$c_k(X) = 2 \operatorname{vol}(X)^{1/2} k^{1/2} + O(k^{1/4}).$$

Proof. For admissible symplectic four-manifolds, the Conformality, Increasing, Disjoint Union, and Sublinearity properties follow immediately from Definition 2.1. It then follows from Lemma 2.3 and Definition 2.4 that these properties, as well as the Monotonicity property, also hold for general symplectic four-manifolds.

The C^0 -Continuity property follows from Conformality and Monotonicity, since if two star-shaped domains are close in the Hausdorff metric, then each is contained in the scaling of the other by a number slightly larger than 1. Note here that if X is a star-shaped domain and $r > 0$, then Conformality implies that $c_k(rX) = r^2 c_k(X)$.

To prove the Spectrality property, suppose first that (X, ω) is a nondegenerate Liouville domain with $c_k(X, \omega) < \infty$. It follows from the definition of c_k that there is an orbit set α with $c_k(X, \omega) = \mathcal{A}(\alpha)$, because in (2.3), for every curve u , the energy $\mathcal{E}(u)$ is the action of some orbit set α , and the set of all such actions is discrete.

If (X, ω) is a degenerate Liouville domain, then the Spectrality property follows by approximating with nondegenerate Liouville domains and using (3.2) and Monotonicity as in the proof of C^0 continuity.

To prove the ECH Index property, first note that $c_k(X) < \infty$ by Monotonicity and the upper bound on c_k of a ball proved in (4.3) below. Recall from Remark 2.2 that in (2.3), we can restrict attention to holomorphic curves that do not have any multiply covered components. Let $\mathcal{M}_*^J(\overline{X}, \alpha; x_1, \dots, x_k)$ denote the set of curves in $\mathcal{M}^J(\overline{X}, \alpha; x_1, \dots, x_k)$ without multiply covered components. The hypothesis that

³See e.g. [24, Def. 5.2] for the definition of the ECH index of α . The definition there is stated for ECH generators (a special kind of orbit set, see Remark 4.2), but is valid for arbitrary orbit sets.

X is nondegenerate implies that the set of symplectic actions of orbit sets in ∂X is discrete, so we can rewrite (2.3) as

$$c_k(X) = \max_{\substack{J \in \mathcal{J}(\overline{X}) \\ x_1, \dots, x_k \in X \text{ distinct}}} \min\{\mathcal{A}(\alpha) \mid \mathcal{M}_*^J(\overline{X}, \alpha; x_1, \dots, x_k) \neq \emptyset\}. \quad (4.2)$$

If $u \in \mathcal{M}_*^J(\overline{X}, \alpha; x_1, \dots, x_k)$, then it follows from the ECH index inequality, see e.g. [22, §3.4], that

$$\text{ind}(u) \leq I(\alpha).$$

Here $\text{ind}(u)$ denotes the Fredholm index of u , which for generic $J \in \mathcal{J}(\overline{X})$ is the dimension of the component of the moduli space $\mathcal{M}_*^J(\overline{X}, \alpha)$ containing u . It follows that for generic $J \in \mathcal{J}(\overline{X})$ and for generic $x_1, \dots, x_k \in X$, the minimum in (4.2) has the form $\mathcal{A}(\alpha)$ where $I(\alpha) \geq 2k$. By Gromov compactness as in the proof of Lemma 2.3, the maximum in (4.2) must be realized by generic $J \in \mathcal{J}(\overline{X})$ and generic $x_1, \dots, x_k \in X$.

To prepare for the proof of the Ball property, if $a, b > 0$, define the ellipsoid

$$E(a, b) = \left\{ z \in \mathbb{C}^2 \mid \frac{\pi|z_1|^2}{a} + \frac{\pi|z_2|^2}{b} \leq 1 \right\}.$$

Calculations e.g. in [22, §3.7] show that for any ellipsoid $E(a, b)$ with a/b irrational, there are just two simple Reeb orbits, which have symplectic action a and b , and the ECH index defines a bijection from the set of orbit sets to the set of nonnegative even integers. Furthermore the symplectic action is an increasing function of the ECH index.

To prove the Ball property, by the Conformality property we can assume that $a = 1$. Let $\varepsilon > 0$ be irrational and consider the ellipsoid

$$E(1 - \varepsilon, 1) \subset E(1, 1) = B^4(1).$$

For a given nonnegative integer d , if ε is sufficiently small, then by the previous paragraph, the orbit set of ECH index $d^2 + d$ has symplectic action $d(1 - \varepsilon)$. Taking $\varepsilon \rightarrow 0$, it follows from the ECH index and Monotonicity properties that

$$c_{(d^2+d)/2}(B^4(1)) \geq d.$$

To complete the proof of the Ball property, by the Increasing property, we need to show that

$$c_{(d^2+3d)/2}(B^4(1)) \leq d. \quad (4.3)$$

By Monotonicity, it is enough to show that

$$c_{(d^2+3d)/2}(\mathbb{C}P^2, \omega_{FS}) \leq d. \quad (4.4)$$

Here ω_{FS} denotes the Fubini-Study form on $\mathbb{C}P^2$, normalized so that a line has symplectic area 1. As reviewed in §7, the upper bound (4.4) follows from Taubes’s “Seiberg-Witten = Gromov” theorem⁴.

Finally, the Asymptotics property was shown for ECH capacities in [24, Thm. 1.1]. The proof there just uses the Monotonicity and Disjoint Union properties and the formula for the capacities of a cube. Theorem 5.1 below implies that for a cube, the ECH capacities and the capacities c_k agree. Hence the Asymptotics property also holds for the capacities c_k . \square

Remark 4.2. The properties of the capacities c_k in Theorem 4.1, aside from the Sublinearity property, are also known to hold for ECH capacities. These properties of ECH capacities were proved in [20], except for the Asymptotics property, which is a later refinement proved in [24].

For the ECH capacities, a slightly stronger version of the ECH Index property follows from the definition of ECH capacities reviewed in §6 below: namely one can arrange that $I(\alpha) = 2k$, and furthermore that the orbit set α is an ECH generator. Here we say that an orbit set $\alpha = \{(\alpha_i, m_i)\}$ is an *ECH generator* if $m_i = 1$ whenever α_i is hyperbolic (meaning that the linearized return map has real eigenvalues).

Remark 4.3. Some applications of ECH capacities only need the properties in Theorem 4.1, and thus can be re-proved using the capacities c_k . For example, Irie [29] proved a C^∞ closing lemma for Reeb vector fields on closed three-manifolds, using the asymptotics of the ECH spectrum [8]. In the case of S^3 with the standard contact structure, which corresponds to star-shaped hypersurfaces in \mathbb{R}^4 , the ECH spectrum agrees with the ECH capacities of the corresponding star-shaped domain, and Irie’s proof of the closing lemma works using only the Spectrality and Asymptotics properties in Theorem 4.1.

5 Computation for convex toric domains

We now show that for “convex toric domains”, the capacities c_k agree with a known combinatorial formula for their ECH capacities⁵. In fact, the capacities c_k for these examples are uniquely determined by the properties in Theorem 4.1.

Let Ω be a compact domain in $\mathbb{R}_{\geq 0}^2$. Define the *toric domain*

$$X_\Omega = \{z \in \mathbb{C}^n \mid \pi(|z_1|^2, |z_2|^2) \in \Omega\}.$$

Define a (four-dimensional) *convex toric domain* to be a toric domain X_Ω as above such that the set

$$\widehat{\Omega} = \{\mu \in \mathbb{R}^2 \mid (|\mu_1|, |\mu_2|) \in \Omega\}$$

⁴If one is trying to avoid using Seiberg-Witten theory, one could prove the same result using Gromov-Witten theory instead.

⁵This formula appears in [23, Prop. 5.6]. It is a specialization of a result in [7, Cor. A.12] computing the ECH capacities of a more general notion of “convex toric domain”.

is convex⁶. Define a (four-dimensional) *concave toric domain* to be a toric domain X_Ω such that the set $\mathbb{R}_{\geq 0}^2 \setminus \Omega$ is convex.

If X_Ω is a four-dimensional convex toric domain, let $\|\cdot\|_\Omega^*$ denote the norm on \mathbb{R}^2 defined by

$$\|v\|_\Omega^* = \max \left\{ \langle v, w \rangle \mid w \in \widehat{\Omega} \right\}.$$

Define a *convex integral path* to be a polygonal path in the nonnegative quadrant from the point $(0, b)$ to the point $(a, 0)$, for some nonnegative integers a and b , with vertices at lattice points, such that the region bounded by Λ and the line segments from $(0, 0)$ to $(a, 0)$ and from $(0, 0)$ to $(0, b)$ is convex. Define $\widehat{\mathcal{L}}(\Lambda)$ to be the number of lattice points in this region, including lattice points on the boundary. Let $\ell_\Omega(\Lambda)$ denote the length of Λ in the norm $\|\cdot\|_\Omega^*$, that is the sum of the norms of the vectors coming from the line segments in the path.

Theorem 5.1. *If X_Ω is a four-dimensional convex toric domain, then*

$$c_k(X_\Omega) = \min \{ \ell_\Omega(\Lambda) \mid \widehat{\mathcal{L}}(\Lambda) = k + 1 \} \quad (5.1)$$

where the minimum is over convex integral paths Λ .

Proof. It is shown in [23, Lem. 5.4] that given $L, \varepsilon > 0$, there is a nondegenerate star-shaped domain X' with $\text{dist}_{C^0}(X', X_\Omega) < \varepsilon$ with the following property: Every orbit set α for X' with action $\mathcal{A}(\alpha) < L$ determines a convex generator Λ such that $|\mathcal{A}(\alpha) - \ell_\Omega(\Lambda)| < \varepsilon$ and the ECH index $I(\alpha) \leq 2(\widehat{\mathcal{L}}(\Lambda) - 1)$. It then follows from the C^0 -Continuity and ECH Index properties in Theorem 4.1 that

$$c_k(X_\Omega) \geq \min \{ \ell_\Omega(\Lambda) \mid \widehat{\mathcal{L}}(\Lambda) \geq k + 1 \}. \quad (5.2)$$

We now prove the reverse inequality. Let $a > 0$ be the smallest real number such that $X_\Omega \subset B^4(a)$. In [7, §2.2], a “negative weight sequence” is defined; this is a nondecreasing (possibly finite) sequence of positive real numbers (a_1, a_2, \dots) . It has the property that there is a symplectic embedding

$$X_\Omega \sqcup \coprod_i \text{int}(B^4(a_i)) \longrightarrow B^4(a)$$

which fills the volume of $B^4(a)$. It follows from the Disjoint Union property that

$$c_k(X_\Omega) \leq \inf_{l \geq 0} \left(c_{k+l}(B^4(a)) - c_l \left(\coprod_{i \leq l} B^4(a_i) \right) \right).$$

⁶This is slightly misleading terminology, as a “convex toric domain” is not the same thing as a toric domain that is convex; see [17, §2] for clarification.

Furthermore, c_k agrees with c_k^{ECH} for a disjoint union of balls by the Disjoint Union and Ball properties, so we can rewrite the above inequality as

$$c_k(X_\Omega) \leq \inf_{l \geq 0} \left(c_{k+l}^{\text{ECH}}(B^4(a)) - c_l^{\text{ECH}} \left(\prod_{i \leq l} B^4(a_i) \right) \right). \quad (5.3)$$

Finally, a combinatorial calculation in [7, §A.3] shows that the right hand side of (5.3) is less than or equal to the right hand side of (5.2).

To complete the proof, we observe that

$$\min\{\ell_\Omega(\Lambda) \mid \widehat{\mathcal{L}}(\Lambda) \geq k + 1\} = \min\{\ell_\Omega(\Lambda) \mid \widehat{\mathcal{L}}(\Lambda) = k + 1\},$$

as explained in [7, §A.3]. □

Remark 5.2. Theorem 5.1 shows that the capacities c_k agree with the ECH capacities for convex toric domains. It follows from the Monotonicity property that all obstructions to symplectic embeddings between convex toric domains coming from ECH capacities can be recovered using the capacities c_k .

Remark 5.3. Going beyond ECH capacities, it is shown in [23, Thm. 1.19] that if X_Ω and $X_{\Omega'}$ are four-dimensional convex toric domains, and if there exists a symplectic embedding $X_\Omega \rightarrow X_{\Omega'}$, then a certain combinatorial criterion holds. This leads to stronger symplectic obstructions in some cases where ECH capacities do not give sharp obstructions, for example to symplectically embedding a polydisk into a ball or ellipsoid; see [23, 4, 9].

The proof of [23, Thm. 1.19] rests on the existence of a certain ECH index 0 holomorphic curve in (the completion of) a symplectic cobordism between the (perturbed) boundaries of X_Ω and $X_{\Omega'}$ which is produced using Seiberg-Witten theory. One can re-prove the existence of this curve using methods of this paper, namely by using the existence of curves in $\overline{X_{\Omega'}}$ with point constraints in the image of X_Ω , as guaranteed by the capacities c_k , and then neck stretching as in the proof of Lemma 2.3.

6 Comparison with ECH capacities

Aside from the examples of toric domains, we do not know to what extent c_k agrees with c_k^{ECH} , but we do have the following general fact, whose proof (and statement) use Seiberg-Witten theory:

Theorem 6.1. *Let X be a star-shaped domain in \mathbb{R}^4 and let k be a nonnegative integer. Then*

$$c_k(X) \leq c_k^{\text{ECH}}(X).$$

To prepare for the proof of Theorem 6.1, we now recall the definition of the ECH capacities c_k^{ECH} , for the simplest case of four-dimensional nondegenerate Liouville domains with connected boundary.

Let Y be a closed oriented three-manifold and let λ be a nondegenerate contact form on Y . The following is an outline of the definition of the *embedded contact homology* $ECH(Y, \lambda)$. We define $ECC(Y, \lambda)$ to be the free $\mathbb{Z}/2$ -module⁷ generated by the ECH generators; see Remark 4.2. For a generic λ -compatible almost complex structure J on $\mathbb{R} \times Y$, the ECH differential

$$\partial_J : ECC(Y, \lambda) \longrightarrow ECC(Y, \lambda)$$

is defined as follows. If α and β are ECH generators, then the coefficient of β in $\partial_J \alpha$, which we denote by $\langle \partial_J \alpha, \beta \rangle$, is a mod 2 count of “ J -holomorphic currents” \mathcal{C} in $\mathbb{R} \times Y$, modulo \mathbb{R} translation, that are asymptotic to α as $s \rightarrow +\infty$ and to β as $s \rightarrow -\infty$, and that have ECH index $I(\mathcal{C}) = 1$. See [22, §3] for detailed definitions. It is shown in [25] that $\partial_J^2 = 0$. We define $ECH(Y, \lambda)$ to be the homology of the chain complex $(ECC(Y, \lambda), \partial_J)$.

It follows from the definition of λ -compatible almost complex structure that the ECH differential decreases symplectic action:

$$\langle \partial_J \alpha, \beta \rangle \neq 0 \implies \mathcal{A}(\alpha) > \mathcal{A}(\beta). \quad (6.1)$$

As a result, for each $L \in \mathbb{R}$, the ECH generators with action less than L span a subcomplex of $(ECC(Y, \lambda), \partial_J)$. We define the *filtered ECH*, which we denote by $ECH^L(Y, \lambda)$, to be the homology of this subcomplex.

It was shown by Taubes [38] that $ECH(Y, \lambda)$ is isomorphic to a version of Seiberg-Witten Floer cohomology defined by Kronheimer-Mrowka [30]. Taubes’s isomorphism was used in [28, Thm. 1.3] to show that $ECH(Y, \lambda)$ and $ECH^L(Y, \lambda)$ do not depend on J ; that is, the homologies for different choices of J are canonically isomorphic to each other.

There is also a map

$$U : ECH^L(Y, \lambda) \longrightarrow ECH^L(Y, \lambda)$$

induced by a chain map which counts J -holomorphic currents with ECH index 2 passing through a base point in $\mathbb{R} \times Y$. This map does not depend on the choice of base point when Y is connected; otherwise it depends on a choice of connected component of Y . See [27, §2.5] for more details.

Now let (X, ω) be a four-dimensional nondegenerate Liouville domain with connected boundary Y and associated contact form λ . In this case the k^{th} ECH capacity is defined by

$$c_k^{\text{ECH}}(X, \omega) = \inf\{L \in \mathbb{R} \mid \exists \eta \in ECH^L(Y, \lambda) : U^k \eta = [\emptyset]\}. \quad (6.2)$$

⁷It is also possible to define ECH with integer coefficients [26, §9].

Here $[\emptyset]$ is the homology class in $ECH^L(Y, \lambda)$ of the empty set of Reeb orbits, which is a cycle by (6.1).

Proof of Theorem 6.1. Since c_k and c_k^{ECH} are both C^0 -continuous, we can assume without loss of generality that X is nondegenerate.

Let Y denote the boundary of X and let λ denote the restriction of the standard Liouville form to Y . As explained for example in [20, Thm. 2.3], for each L , the filling X of Y induces a cobordism map

$$\Phi : ECH^L(Y, \lambda) \longrightarrow \mathbb{Z}/2,$$

defined using Seiberg-Witten theory, which sends $[\emptyset]$ to 1.

Now suppose that $J \in \mathcal{J}(\overline{X})$ and $x_1, \dots, x_k \in X$. Heuristically one might expect that if J and x_1, \dots, x_k are generic, then the composition

$$\Phi \circ U^k : ECH^L(Y, \lambda) \longrightarrow \mathbb{Z}/2 \tag{6.3}$$

is induced by a cocycle

$$\phi : ECC^L(Y, \lambda) \longrightarrow \mathbb{Z}/2$$

that counts J -holomorphic curves in \overline{X} with ECH index $2k$ passing through x_1, \dots, x_k . What one can actually prove, as in the “holomorphic curves axiom” for ECH cobordism maps in [28, Thm. 1.9] and the comparison of U maps in [39, Thm. 1.1], is that for any $J \in \mathcal{J}(\overline{X})$ and any $x_1, \dots, x_k \in X$ (not necessarily generic), the composition (6.3) is induced by a (noncanonical) cocycle ϕ with the following property: If α is an ECH generator and $\phi(\alpha) \neq 0$, then there exists a “broken J -holomorphic current” in \overline{X} passing through x_1, \dots, x_k . This last statement implies that there is an orbit set α' with $\mathcal{A}(\alpha') \leq \mathcal{A}(\alpha)$ and a holomorphic curve in $\mathcal{M}^J(\overline{X}, \alpha'; x_1, \dots, x_k)$.

Now suppose that $L > c_k^{\text{ECH}}(X)$. Then by the definition of ECH capacities (6.2), there exists $\eta \in ECH^L(Y, \lambda)$ with $U^k \eta = [\emptyset]$. It follows that $(\Phi \circ U^k)(\eta) = 1$. By the previous paragraph, for any $J \in \mathcal{J}(X)$ and any $x_1, \dots, x_k \in X$, there exists an ECH generator α' with $\mathcal{A}(\alpha') < L$ such that $\mathcal{M}^J(X, \alpha'; x_1, \dots, x_k) \neq \emptyset$. It then follows from equation (4.2) that $c_k(X) \leq L$. Since $L > c_k^{\text{ECH}}(X)$ was arbitrary, the theorem follows. \square

7 Additional computations using Seiberg-Witten theory

We now compute some additional examples of the capacities c_k using Seiberg-Witten theory (which could perhaps be avoided with more work).

If X_Ω is a four-dimensional concave toric domain as defined in §5, define an “anti-norm” on \mathbb{R}^2 by

$$[v]_\Omega = \min\{\langle (|v_1|, |v_2|), w \rangle \mid w \in \partial_+ \Omega\}$$

where $\partial_+\Omega$ denotes the closure of the portion of $\partial\Omega$ not on the axes. Define a *concave integral path* to be a polygonal path in the nonnegative quadrant from the point $(0, b)$ to the point $(a, 0)$, for some nonnegative integers a and b , with vertices at lattice points, which is the graph of a convex function. Define $\check{\mathcal{L}}(\Lambda)$ to be the number of lattice points in the region bounded by Λ and the axes, this time (in contrast to the case of convex toric domains) not including lattice points on Λ . Let $\ell_\Omega(\Lambda)$ denote the length of Λ defined using $[\cdot]_\Omega$.

Theorem 7.1. *If X_Ω is a four-dimensional concave toric domain, then*

$$c_k(X_\Omega) = \max\{\ell_\Omega(\Lambda) \mid \check{\mathcal{L}}(\Lambda) = k\} \quad (7.1)$$

where the maximum is over concave integral paths Λ .

Remark 7.2. It is shown in [3, Thm. 1.12] that the same formula holds for the ECH capacities $c_k^{\text{ECH}}(X_\Omega)$.

Proof of Theorem 7.1. In [3, §1.3], see also [24, §1.3], a “weight expansion” of X_Ω is defined; this is a nondecreasing (possibly finite) sequence of positive real numbers (a_1, a_2, \dots) . There is a symplectic embedding

$$\coprod_i \text{int } B^4(a_i) \longrightarrow X_\Omega$$

which fills the volume of X_Ω . It follows from the Monotonicity property that

$$c_k(X_\Omega) \geq c_k \left(\coprod_{i \leq k} B^4(a_i) \right).$$

By the Ball and Disjoint Union properties, we have

$$c_k \left(\coprod_{i \leq k} B^4(a_i) \right) = c_k^{\text{ECH}} \left(\coprod_{i \leq k} B^4(a_i) \right).$$

It is shown in [3, §2] by a combinatorial calculation that

$$c_k^{\text{ECH}} \left(\coprod_{i \leq k} B^4(a_i) \right) \geq \max\{\ell_\Omega(\Lambda) \mid \check{\mathcal{L}}(\Lambda) = k\}.$$

By Remark 7.2 and Theorem 6.1, the above inequalities are equalities. \square

We now consider some closed symplectic manifolds. Given $a > 0$, let $\mathbb{C}P^2(a)$ denote $\mathbb{C}P^2$ with the Fubini-Study form, scaled so that a line has symplectic area a . Let $S^2(a)$ denote S^2 with a symplectic form of area a .

Theorem 7.3. *Let $a, b > 0$ and let k be a nonnegative integer.*

$$(a) \ c_k(\mathbb{C}P^2(a)) = da \text{ where } d \text{ is the unique nonnegative integer with } d^2 + d \leq 2k \leq d^2 + 3d.$$

$$(b) \ c_k(S^2(a) \times S^2(b)) = \min\{am + bn \mid m, n \in \mathbb{Z}_{\geq 0}, (m+1)(n+1) \geq k+1\}.$$

To prepare for the proof of this theorem, if (X, ω) is a closed symplectic four-manifold with $b_2^+(X) = 1$, and if $A \in H_2(X)$, let $SW(X, \omega, A) \in \mathbb{Z}/2$ denote the mod 2 Seiberg-Witten invariant of X , for the spin-c structure determined by ω and A , in the symplectic chamber; see the review in [2, §2]. Define the ECH index

$$I(A) = A \cdot A + \langle c_1(TX), A \rangle \in \mathbb{Z}.$$

Lemma 7.4. *Let (X, ω) be a closed symplectic four-manifold with $b_2^+(X) = 1$ and let $A \in H_2(X)$. If $SW(X, \omega, A) \neq 0$ and $I(A) = 2k$, then $c_k(X, \omega) \leq \langle [\omega], A \rangle$.*

Proof. If $J \in \mathcal{J}(X, \omega)$ and $x_1, \dots, x_k \in X$ are generic, then it follows from Taubes's "Seiberg-Witten = Gromov" theorem [37] that there exists a (possibly disconnected) J -holomorphic curve in the homology class A passing through the points x_1, \dots, x_k . Thus

$$\inf_{u \in \mathcal{M}^J(X; x_1, \dots, x_k)} \mathcal{E}(u) \leq \langle [\omega], A \rangle$$

when J, x_1, \dots, x_k are generic. A Gromov compactness argument shows that the supremum in the definition (2.3) of $c_k(X, \omega)$ is realized for generic J, x_1, \dots, x_k . \square

Proof of Theorem 7.3. (a) Let d be the unique nonnegative integer with $d^2 + d \leq 2k \leq d^2 + 3d$. If $A \in H_2(\mathbb{C}P^2)$ is d times the homology class of a line, then $I(A) = d^2 + 3d$. As reviewed in [2, §2.4], we have $SW(A) \neq 0$. It follows from Lemma 7.4 that $c_k(\mathbb{C}P^2(a)) \leq da$. The reverse inequality follows from Monotonicity, since it was shown in the proof of the Ball property in Theorem 4.1 that $c_k(B^4(a)) \geq da$.

(b) Let L denote the right hand side of the equation in (b). If m and n are nonnegative integers, and if $A = (m, n) \in H_2(S^2 \times S^2)$, then $I(A) = 2(mn + m + n)$. As reviewed in [2, §2.4], we have $SW(A) \neq 0$. It follows from Lemma 7.4 that

$$c_k(S^2(a) \times S^2(b)) \leq L.$$

To prove the reverse inequality, consider the polydisk

$$P(a, b) = \{z \in \mathbb{C}^2 \mid \pi|z_1|^2 \leq a, \pi|z_2|^2 \leq b\}.$$

A calculation using Theorem 5.1 shows that

$$c_k(P(a, b)) = L.$$

Since the interior of $P(a, b)$ symplectically embeds into $S^2(a) \times S^2(b)$, we are done by Monotonicity. \square

Remark 7.5. Theorem 7.3 shows that the capacities c_k are the same for $\mathbb{C}P^2(a)$ and the ball $B^4(a)$. Likewise they are the same for $S^2(a) \times S^2(b)$ and the polydisk $P(a, b)$. This means that if the capacities c_k obstruct a symplectic embedding of a symplectic four-manifold (X, ω) into $B^4(a)$ or $P(a, b)$ respectively, then a symplectic embedding of (X, ω) into $\mathbb{C}P^2(a)$ or $S^2(a) \times S^2(b)$ respectively is not possible either. The same statement is true for the ECH capacities c_k^{ECH} when X is a star-shaped domain by [2, Thm. 1.4].

8 An even simpler definition of capacities

To conclude, we now define an even simpler series of symplectic capacities, for symplectic manifolds of any dimension.

If (X, ω) is symplectic manifold, let $\mathcal{J}(X, \omega)$ denote the set of ω -compatible almost complex structures on X . Given $J \in \mathcal{J}(X, \omega)$, let $\mathcal{P}^J(X)$ denote the set of proper holomorphic maps

$$u : (S, j) \longrightarrow (X, J)$$

where (S, j) is a one-dimensional complex manifold (not necessarily compact or connected), and we assume that the restriction of u to each component of S is nonconstant. Given u as above, define the *energy*

$$\mathcal{E}(u) = \int_S u^* \omega \in [0, \infty].$$

Note that the energy is well-defined because u is proper and $u^* \omega$ is pointwise non-negative. If $x_1, \dots, x_k \in X$ are distinct, let $\mathcal{P}^J(X; x_1, \dots, x_k)$ denote the set of proper maps u as above such that $x_1, \dots, x_k \in u(S)$.

Definition 8.1. Let (X, ω) be a compact symplectic manifold (possibly disconnected and/or with boundary), and let k be a nonnegative integer. Define

$$\widehat{c}_k(X, \omega) = \sup_{\substack{J \in \mathcal{J}(X, \omega) \\ x_1, \dots, x_k \in \text{int}(X) \text{ distinct}}} \inf_{u \in \mathcal{P}^J(\text{int}(X); x_1, \dots, x_k)} \mathcal{E}(u) \in [0, \infty]. \quad (8.1)$$

Remark 8.2. It follows immediately from the definition that the capacities \widehat{c}_k satisfy the Conformality, Increasing, Disjoint Union, and Sublinearity properties in Theorem 4.1.

We can also quickly show that they satisfy Monotonicity under symplectic embeddings $\varphi : (X, \omega) \rightarrow (X', \omega')$ between symplectic manifolds of the same dimension, without using Gromov compactness. This is because since X is compact, any $J \in \mathcal{J}(X, \omega)$ can be extended to $J' \in \mathcal{J}(X', \omega')$ with $J'|_{\varphi(X)} = \varphi_* J$.

One can now further deduce that each \widehat{c}_k is a C^0 -continuous function on the set of star-shaped domains in \mathbb{R}^{2n} .

Remark 8.3. When $k = 1$, the capacity \widehat{c}_1 agrees with the “symplectic width” defined by Gromov in [15, §4.2]. In particular, $\widehat{c}_1(B^{2n}(a)) = a$. The symplectic width should not be confused with the Gromov width c_{Gr} from §1. The monotonicity lemma for minimal surfaces implies that $c_{\text{Gr}} \leq \widehat{c}_1$.

In a sense the capacities \widehat{c}_k are more natural than the c_k , because for domains that are not Liouville domains, they are defined directly, without taking a supremum over symplectic embeddings as in Definition 2.4. However the price for this is that we have to consider holomorphic curves without nice boundary conditions, which makes computations more difficult.

Remark 8.4. Suppose that $\dim(X) = 4$. If X is closed, then $\widehat{c}_k(X, \omega) = c_k(X, \omega)$ by definition. If (X, ω) is a Liouville domain, then we have

$$\widehat{c}_k(X, \omega) \leq c_k(X, \omega). \quad (8.2)$$

This is because if $\varepsilon > 0$, then any almost complex structure $J \in \mathcal{J}(X, \omega)$ can be extended to an $\overline{\omega}$ -compatible almost complex structure on \overline{X} whose restriction to $[\varepsilon, \infty) \times Y$ agrees with an $e^\varepsilon \lambda$ -compatible almost complex structure on $\mathbb{R} \times Y$. It follows from this as in (3.2) that

$$\widehat{c}_k(X, \omega) \leq e^\varepsilon c_k(X, \omega).$$

We can choose $\varepsilon > 0$ arbitrarily small, and this proves (8.2).

We conjecture that in fact $\widehat{c}_k(X, \omega) = c_k(X, \omega)$ when (X, ω) is a four-dimensional Liouville domain.

Example 8.5. The simplest example of \widehat{c}_k that we do not know how to compute is \widehat{c}_3 of a four-dimensional ball. We currently just know that

$$\frac{3}{2} \leq \widehat{c}_3(B^4(1)) \leq 2.$$

Here the first inequality holds because three copies of $\text{int}(B^4(1/2))$ can be symplectically embedded into $B^4(1)$, and the second inequality holds because $\text{int}(B^4(1))$ can be symplectically embedded into $\mathbb{C}P^2(1)$.

A number of additional capacities along the lines of the \widehat{c}_k , defined using proper holomorphic maps satisfying various constraints, are studied in [33].

References

- [1] F. Bourgeois, Y. Eliashberg, H. Hofer, K. Wysocki, and E. Zehnder, *Compactness results in symplectic field theory*, *Geom. Topol.* **7** (2003), 799–888.
- [2] J. Chaidez and B. Wormleighton, *ECH embedding obstructions for rational surfaces*, arXiv:2008.10125.

- [3] K. Choi, D. Cristofaro-Gardiner, D. Frenkel, M. Hutchings, and V. Ramos, *Symplectic embeddings into four-dimensional concave toric domains*, J. Topol. **7** (2014), 1054–1076.
- [4] K. Christianson and J. Nelson, *Symplectic embeddings of four-dimensional polydisks into balls*, Alg. Geom. Topol. **18** (2018), 2151–2178.
- [5] K. Cieliebak, H. Hofer, J. Latschev, and F. Schlenk, *Quantitative symplectic geometry*, in Dynamics, Ergodic Theory, and Geometry, Math. Sci. Res. Inst. Publ. **54** (2007), Cambridge University Press, 1–44.
- [6] K. Cieliebak and K. Mohnke, *Compactness for punctured holomorphic curves*, J. Symplectic Geom. **3** (2005), 589–654.
- [7] D. Cristofaro-Gardiner (appendix with K. Choi), *Symplectic embeddings of concave toric domains into convex ones*, J. Diff. Geom. **112** (2019), 199–232.
- [8] D. Cristofaro-Gardiner, M. Hutchings, and V. Ramos, *The asymptotics of ECH capacities*, Invent. Math. **199** (2015), 187–214.
- [9] L. Digiosia, J. Nelson, H. Ning, M. Weiler, and Y. Yang, *Symplectic embeddings of four-dimensional polydisks into half integer ellipsoids*, arXiv:2010.06687.
- [10] A. Doan and T. Walpuski, *Castelnuovo’s bound and rigidity in almost complex geometry*, arXiv:1809.04731, to appear in Adv. Math.
- [11] I. Ekeland and H. Hofer, *Symplectic topology and Hamiltonian dynamics, II*, Math Z. **203** (1990), 553–567.
- [12] Y. Eliashberg, A. Givental, and H. Hofer, *Introduction to symplectic field theory*, Geom. Funct. Anal. (2000), Special Volume, Part II, 560–673.
- [13] C. Gerig, *Taming the pseudoholomorphic beasts in $\mathbb{R} \times (S^1 \times S^2)$* , Geom. Topol. **24** (2020), 1791–1839.
- [14] M. Gromov, *Pseudoholomorphic curves in symplectic manifolds*, Invent. Math. **82** (1985), 307–347.
- [15] M. Gromov, *Soft and hard symplectic geometry*, Proceedings of the ICM, Berkeley, 1986.
- [16] J. Gutt and M. Hutchings, *Symplectic capacities from positive S^1 -equivariant symplectic homology*, Alg. Geom. Topol. **18** (2018), 3537–3600.
- [17] J. Gutt, M. Hutchings, and V. Ramos, *Examples around the strong Viterbo conjecture*, arXiv:2003.10854, to appear in Journal of Fixed Point Theory and Applications.

- [18] M. Hutchings, *An index inequality for embedded pseudoholomorphic curves in symplectizations*, J. Eur. Math. Soc. **4** (2002), 313–361.
- [19] M. Hutchings, *The embedded contact homology index revisited*, in New Perspectives and Challenges in Symplectic Field Theory, CRM Proc. Lecture Notes, vol. 49 (AMS, 2009), pp. 263–297.
- [20] M. Hutchings, *Quantitative embedded contact homology*, J. Diff. Geom. **88** (2011), 231–266.
- [21] M. Hutchings, *Recent progress on symplectic embedding problems in four dimensions*, Proc. Natl. Acad. Sci. USA **108** (2011), 8093–8099.
- [22] M. Hutchings, *Lecture notes on embedded contact homology*, Contact and symplectic topology, 389–484, Bolyai Soc. Math. Stud. bf 26, Springer, 2014.
- [23] M. Hutchings, *Beyond ECH capacities*, Geom. Topol. **20** (2016), 1085–1126.
- [24] M. Hutchings, *ECH capacities and the Ruelle invariant*, arXiv:1910.08260, to appear in Journal of Fixed Point Theory and Applications.
- [25] M. Hutchings and C. H. Taubes, *Gluing pseudoholomorphic curves along branched covered cylinders I*, J. Sympl. Geom. **5** (2007), 43–137.
- [26] M. Hutchings and C. H. Taubes, *Gluing pseudoholomorphic curves along branched covered cylinders II*, J. Sympl. Geom. **7** (2009), 29–133.
- [27] M. Hutchings and C. H. Taubes, *The Weinstein conjecture for stable Hamiltonian structures*, Geom. Topol. **13** (2009), 901–941.
- [28] M. Hutchings and C. H. Taubes, *Proof of the Arnold chord conjecture in three dimensions, II*, Geom. Topol. **17** (2013), 2601–2688.
- [29] K. Irie *Dense existence of periodic orbits and ECH spectral invariants*, J. Mod. Dyn. **9** (2015), 357–363.
- [30] P. B. Kronheimer and T. S. Mrowka, *Monopoles and three-manifolds*, Cambridge University Press, 2008.
- [31] D. McDuff, *The Hofer conjecture on embedding symplectic ellipsoids*, J. Diff. Geom. **88** (2011), 519–532.
- [32] D. McDuff and K. Siegel, *Symplectic capacities, unperturbed curves, and convex toric domains*, arXiv:2111.00515.
- [33] G. Mikhalkin, *Exotic enumerative capacities and symplectic camel paths*, in preparation.

- [34] J. Rooney, *Cobordism maps in embedded contact homology*, arXiv:1912.01048.
- [35] K. Siegel, *Higher symplectic capacities*, arXiv:1902.01490.
- [36] C. H. Taubes, *The structure of pseudoholomorphic subvarieties for a degenerate almost complex structure and symplectic form on $S^1 \times B^3$* , *Geom. Topol.* **2** (1998), 221-332.
- [37] C. H. Taubes, *Seiberg-Witten and Gromov invariants for symplectic 4-manifolds*, First International Press Lecture Series 2, International Press, 2000.
- [38] C. H. Taubes, *Embedded contact homology and Seiberg-Witten Floer cohomology I*, *Geom. Topol.* **14** (2010), 2497–2581.
- [39] C. H. Taubes, *Embedded contact homology and Seiberg-Witten Floer cohomology V*, *Geom. Topol.* **14** (2010), 2961–3000.