

BKT transitions in classical and quantum long-range systems

Guido Giachetti*

SISSA and INFN Sezione di Trieste, Via Bonomea 265, I-34136 Trieste, Italy

Andrea Trombettoni

*Department of Physics, University of Trieste, Strada Costiera 11, I-34151 Trieste, Italy and
SISSA and INFN Sezione di Trieste, Via Bonomea 265, I-34136 Trieste, Italy*

Stefano Ruffo

*SISSA and INFN Sezione di Trieste, Via Bonomea 265, I-34136 Trieste, Italy and
Istituto dei Sistemi Complessi, Consiglio Nazionale delle Ricerche,
Via Madonna del Piano 10, I-50019 Sesto Fiorentino, Italy*

Nicolò Defenu

Institut für Theoretische Physik, ETH Zürich, Wolfgang-Pauli-Str. 27 Zürich, Switzerland

Abstract

In the past decades considerable efforts have been made in order to understand the critical features of both classical and quantum long-range interacting models. The case of the Berezinskii-Kosterlitz-Thouless (BKT) universality class, as in the $2d$ classical XY model, is considerably complicated by the presence, for short-range interactions, of a line of renormalization group fixed points. In this paper we discuss a field theoretical treatment of the $2d$ XY model with long-range couplings and we compare it with results from the self-consistent harmonic approximation. These methods lead to a rich phase diagram, where both power-law BKT scaling and spontaneous symmetry breaking appear for the same (intermediate) decay rates of long-range interactions. We also discuss the Villain approximation for the $2d$ XY model with power-law couplings, providing hints that, in the long-range regime, it fails to reproduce the correct critical behavior. The obtained results are then applied to the long-range quantum XXZ spin chain at zero temperature. We discuss the relation between the phase diagrams of the two models and we give predictions about the scaling of the order parameter of the quantum chain close to the transition.

I. INTRODUCTION

In the context of statistical mechanics and condensed matter, it is well known that the presence of long-range (LR) interactions, such as slow-decaying couplings with power-law behavior at large distances, gives rise to plenty of new physical phenomena in the equilibrium and non-equilibrium properties of both classical [1] and quantum [2] systems. These properties have recently sparked a new wave of interest, due to the possibility of experimental realizations in atomic, molecular and optical (AMO) systems [2–9].

If the decay of the interaction is slow enough, LR effects influence the universal critical properties and even induce a spontaneous symmetry breaking (SSB) in a low-dimensional system, since the celebrated Hohenberg-Mermin-Wagner theorem does not hold in presence of LR couplings, as discussed since the classic papers [10–12].

A qualitative and quantitative understanding of the critical properties of LR interacting systems follows from the Sak criterion [13]. Let us consider power-law couplings of the form $\sim 1/r^{d+\sigma}$, where d is the system dimension. For the sake of clarity, in the following considerations we explicitly refer to classical $O(n)$ models [14]. In the gaussian non-interacting case ($n \rightarrow \infty$),

for large distances and low energies, one has to compare the short-range (SR) critical propagator, $1/k^2$, with the LR one, $1/k^\sigma$. However, interacting systems possess a finite anomalous dimension η at the critical point, so that the propagator scales as $1/k^{2-\eta}$ [15]. Denoting by η_{sr} the anomalous dimension in the SR limit of the theory, i.e. $\sigma \rightarrow \infty$, one concludes that *at criticality* for $\sigma > 2 - \eta_{\text{sr}}$ the system behaves as its SR counterpart, while for $\sigma < 2 - \eta_{\text{sr}}$ the LR nature of the couplings modifies the critical behaviour. This argument is at the heart of Sak's seminal paper [13] and it implies that one can define a value σ_* such that for $\sigma > \sigma_*$ the system is in the universality class of its SR counterpart, while for $\sigma < \sigma_*$ the system changes universality class. The value of σ_* is given by

$$\sigma_* = 2 - \eta_{\text{sr}}. \quad (1)$$

Sak's criterion was formulated for classical $O(n)$ models, such as the Ising model, a playground where it has been the subject of thorough investigation. Regarding its validity it is fair to conclude that, to this date, it has to be considered valid and well tested, see reviews [16, 17] and Refs therein. Despite the fact that there is no rigorous proof for it, and numerical results can only put an upper bound on its violation, its usefulness is unquestioned. For the quantum $O(n)$ models, one finds $\sigma_* = 2$ at mean-field level [18] and one can obtain beyond-mean-field results by applying renormalization group techniques [19].

Sak's criterion applies to all d and n , except for $d = 2$

* ggiachet@sisssa.it

and $n = 2$, i.e. the $2d$ XY universality class. There, it is well known that SSB occurs at low temperature for $\sigma < 2$ [20] and, therefore, several questions arise concerning the applicability of Sak's criterion in the SR limit, where the model does not present SSB. Moreover, in the $2d$ XY model, η_{sr} – unlike the other cases covered by Sak's criterion – actually depends on the temperature, raising further concerns about the applicability of the criterion to the $2d$ XY model with LR couplings.

In the recent paper [21], the fate of the BKT line of critical points has been discussed. For $\sigma > 2$ simple arguments (not directly applicable to $\sigma < 2$) show the existence of the BKT transition [22], while what happens for $\sigma < 2$ is considerably more subtle. In Ref. [21] SSB was found to persist in the range $7/4 < \sigma < 2$ up to a finite critical temperature T_c , beyond which a quasi-ordered BKT-like phase appears below a larger temperature T_{BKT} . These results are compatible with the application of Sak's criterion with the temperature dependent η_{sr} of the XY model in the SR limit.

Since the classical $2d$ XY model at finite temperature lies in the same universality class of the quantum XXZ chain at $T = 0$ for nearest-neighbors-interactions, a natural question is how to apply these results for the $2d$ LR XY model to the $1d$ LR XXZ chain. We remind that, in the SR limit, a mapping can be obtained via the transfer matrix technique [23]. However, this method does not straightforwardly apply to the LR case, nor to any non-local couplings, in general. While it is natural to think the two models to be in the same universality class for fast-decaying couplings as well, it is not clear what happens in the LR regime. We observe that the $2D$ boson gas at finite temperature with $1/r^3$ density-density interactions is equivalent to a quantum XXZ chain with $z - z$ LR couplings and it has been found to follow the usual BKT scenario [24] with no signatures of SSB.

The LR $1d$ XXZ chain has been studied in Ref. [25], where evidence were provided showing the existence of three phases in the $T = 0$ phase diagram as a function of σ and Δ (with Δ the anisotropy parameter). One phase features SSB, another is disordered and the third is gapless. The comparison between the phase diagrams of Ref. [21] for the $2d$ classical finite temperature XY model and of Ref. [25] for the $1d$ quantum zero temperature XXZ model is one of the motivations of the present paper.

The study of the $2d$ XY model with LR interactions is of great interest *per se*: the model is indeed paradigmatic for the study of $O(2)$ -symmetric systems in $d = 2$ (including thin ^4He films [26], quasi-2D layered superconductors [27–31], exciton-polariton systems [32], cold atoms in $2D$ traps [33, 34] and $2D$ electron gases at the interface between insulating oxides in artificial heterostructures [35–37]).

In the present paper, the scenario presented in Ref. [21] for the $2d$ XY model is analyzed in detail and investigated using different methods: the low-temperature expansion, an extension of the traditional self-consistent

harmonic approximation (SCHA) [22, 38, 39] to $\sigma < 2$ and renormalization group (RG) field theoretical arguments. The difficulties inherent in the use of the Villain approximation for $\sigma < 2$ are pointed out and contrasted with the SR case, where, at variance, the $2d$ Villain and XY model are related through controllable approximations and lie in the same universality class. Using these results, we closely examine the link between the classical XY model in $2d$ and the quantum XXZ chain, leading to a detailed analysis of the quantum model. The comparison with the results of Ref. [25] is presented. Comments on the BKT transition in LR spin- S quantum chains are also presented.

The paper is structured as follows: after presenting the model in Sec. II, in Sec. III we describe its low-temperature limit. In Sec. IV and V, we examine the peculiar nature of the crossover between LR and SR regimes. In particular, Sec. IV is devoted to extending the self-consistent approximation to the proper $\sigma < 2$ LR regime; while in Sec V we examine the field-theory picture. In Sec. VI we discuss the relation between our analysis and the Villain method used to understand the BKT phase transition in the SR regime, showing the difficulties encountered to describe this crossover. Finally, in Sec. VII we examine the relationship between the $d = 2$ classical XY model and the $1d$ quantum XXZ chain, showing how our results can be used to make predictions about the $T = 0$ phase diagram of the latter.

II. THE XY MODEL

The model consists of a set of N planar rotators, arranged in a 2-dimensional square lattice, interacting through the Hamiltonian

$$\beta H = \frac{1}{2} \sum_{\mathbf{i}, \mathbf{j}} J(r) [1 - \cos(\theta_{\mathbf{i}} - \theta_{\mathbf{j}})] \quad (2)$$

with $\mathbf{i}, \mathbf{j} \in \mathbb{Z}^2$, $\mathbf{r} = \mathbf{i} - \mathbf{j}$, $r = |\mathbf{r}|$, $J(r) \sim Jr^{-2-\sigma}$ for $r \gg 1$ where $J = \beta J_0$. Here we used a particular instance of the general convention $J(r) \sim r^{-d-\sigma}$ used in the d -dimensional case (see e.g. [19]), which assures that the thermodynamical quantities remain additive for any $\sigma > 0$ [1]. The interaction between the spins $\mathbf{S}_{\mathbf{x}} \equiv (\cos \theta(\mathbf{x}), \sin \theta(\mathbf{x}))$ is invariant under global $O(2)$ rotations. In what follows we take our energy units so that $J_0 = 1$ and we will limit ourself to the $\sigma > 0$ case.

In the SR regime, although the possibility of a SSB is ruled out by the Hohenberg-Mermin-Wagner theorem, the low-temperature phase is characterized by quasi-long-range-order, i.e. power-law behavior in the connected correlations functions with a temperature-dependent exponent. The transition between this phase and the disordered, high-temperature one, i.e. the well-known BKT mechanism, may be understood in terms of the unbinding of topological defects, leading to the celebrated mapping to a Coulomb gas of charged particles [40–44].

When the decay of the couplings is slow enough, we expect a SSB phase to appear at low temperature. The accepted criterion for understanding whether the LR interactions do actually modify the critical behavior is the one provided by Sak [13]. However, as discussed in the introduction, this criterion cannot be straightforwardly applied to the BKT mechanism, due to the temperature-dependent anomalous dimension η_{sr} . Moreover, also the duality construction [45], which allows the mapping between the $2d$ SR XY model, the Coulomb gas [46, 47] and the sine-Gordon model [47, 48], breaks down, already in the case of next-to-nearest-neighbors couplings. The other way of obtaining the same mappings is the so-called Villain approximation [49–51] (i.e. the substitution of Hamiltonian (3) with a quadratic one which takes into account the phase periodicity). However, as shown in Sec. VI of this paper, there is reason to believe that this latter Hamiltonian is no longer in the same universality class as soon as the interaction becomes LR, so that the Coulomb gas picture cannot be straightforwardly used to describe the physics of the $2d$ LR XY model.

III. LOW TEMPERATURE EXPANSION

In analogy with the SR regime, we expect the low-temperature behavior of the model to be well described by an approximation *à la* Berezinskii [41]. Since we expect our thermal fluctuations to be small in this regime, we expand the cosine in the Hamiltonian (3) to second order, obtaining, up to an immaterial constant term,

$$\beta H \sim \frac{1}{4} \sum_{\mathbf{i}, \mathbf{j}} J(r) (\theta_{\mathbf{i}} - \theta_{\mathbf{j}})^2. \quad (3)$$

Being this theory quadratic and translationally invariant, we can diagonalize the above Hamiltonian by means of a Fourier transform

$$\theta_{\mathbf{q}} = \frac{1}{\sqrt{N}} \sum_{\mathbf{j}} e^{-i\mathbf{q} \cdot \mathbf{j}} \theta_{\mathbf{j}} \quad \theta_{\mathbf{j}} = \frac{1}{\sqrt{N}} \sum_{\mathbf{q} \in IBZ} e^{i\mathbf{q} \cdot \mathbf{j}} \theta_{\mathbf{q}}. \quad (4)$$

We obtain

$$\beta H \sim \frac{1}{2} \sum_{\mathbf{q} \in IBZ} K(\mathbf{q}) |\theta_{\mathbf{q}}|^2, \quad (5)$$

where we introduced

$$K(\mathbf{q}) = \sum_{\mathbf{r}} J(\mathbf{r}) (1 - \cos(\mathbf{q} \cdot \mathbf{r})). \quad (6)$$

If we are interested in the long-wavelength modes, we can approximate the above sum with a continuous integral, obtaining:

$$K(q) = \int_{r>a} d^2 \mathbf{r} J(r) (1 - \cos(\mathbf{q} \cdot \mathbf{r})). \quad (7)$$

As shown in Appendix A, this quantity grows as $K(q) \sim q^2$ if $\sigma > 2$, while it shows a non-analytic behavior $K(q) \sim q^\sigma$ for $\sigma < 2$. The difference between the two regimes can be fully appreciated if we consider the two-point correlation function:

$$\langle \mathbf{s}_{\mathbf{r}} \cdot \mathbf{s}_0 \rangle = \langle \cos(\theta_{\mathbf{r}} - \theta_0) \rangle. \quad (8)$$

Being the theory quadratic in this approximation, the above quantity can be evaluated quite easily by exploiting the identity $\langle \cos A \rangle_0 = e^{-\frac{1}{2} \langle A^2 \rangle_0}$. From Eq. (5), it follows immediately that $\langle \theta_{\mathbf{q}} \theta_{\mathbf{q}'} \rangle_0 = \delta_{\mathbf{q}+\mathbf{q}'} K(\mathbf{q})^{-1}$ so that we find $\langle \cos(\theta_{\mathbf{r}} - \theta_0) \rangle = e^{-G(\mathbf{r})}$ with

$$G(\mathbf{r}) = \frac{1}{2} \langle (\theta_0 - \theta_{\mathbf{r}})^2 \rangle_0 = \frac{1}{N} \sum_{\mathbf{q} \in IBZ} \frac{1 - \cos(\mathbf{q} \cdot \mathbf{r})}{K(\mathbf{q})}. \quad (9)$$

The long-distance behavior can be once again captured by replacing the sum with an integral

$$G(r) = a^2 \int_{q<\Lambda} \frac{d^2 \mathbf{q}}{(2\pi)^2} \frac{1 - \cos(\mathbf{q} \cdot \mathbf{r})}{K(q)}, \quad (10)$$

where we approximated the first Brillouin zone with a sphere of radius $\Lambda = \sqrt{8\pi}a^{-1}$, so that its volume is preserved, i.e. $\int_{q<\Lambda} d^2 \mathbf{q} = (2\pi)^2 a^{-2}$. As shown in Appendix B, the asymptotic behavior of $G(r)$ depends on σ . In particular, if $\sigma > 2$ we have

$$G(r) \sim \eta(J) \ln \frac{r}{a} + AJ^{-1} \quad (11)$$

with $\eta(J) = p/J$, and A, p are non-universal constants (see a study of these constants in [22]). If $\sigma < 2$ instead, we have that

$$G(r) \sim AJ^{-1}. \quad (12)$$

Then, depending on whether $\sigma < 2$ or $\sigma > 2$ we have that the correlation $\langle \mathbf{s}_{\mathbf{r}} \cdot \mathbf{s}_0 \rangle \sim \text{const}$ or $\langle \mathbf{s}_{\mathbf{r}} \cdot \mathbf{s}_0 \rangle \sim r^{-\eta(J)}$ respectively. In the latter case then, we recover the SR low-temperature BKT behavior [41], in which the correlations decay as a power law with a temperature-dependent exponent. The former case, instead, gives rise to a finite-magnetization ordered phase, with

$$m^2 = \lim_{r \rightarrow \infty} \langle \mathbf{s}_{\mathbf{r}} \cdot \mathbf{s}_0 \rangle = \lim_{r \rightarrow \infty} e^{-G(r)}. \quad (13)$$

This argument then suggests that $\sigma_* = 2$. If compared with Sak's criterion, this implies an effective anomalous exponent $\eta = 0$. Let us notice, however, how the low-temperature approximation *per se* is unable to describe the topological configurations, since $\theta_{\mathbf{i}}$ is no longer a phase, and thus is no longer defined up to multiples of 2π . As a consequence, even in the SR case, the approximation is unable to correctly reproduce all the phenomenology of the BKT transition.

IV. SELF-CONSISTENT APPROACH

For the nearest-neighbor case, the results of the low-temperature expansion can be improved by the self-consistent-harmonic approximation (SCHA), which is able to correctly foresee the existence of the BKT phase transition. The idea behind the SCHA approximation is to replace the original Hamiltonian with a quadratic one, whose couplings are optimized according to a variational principle.

In the SR case the natural variational parameter is an effective coupling \tilde{J} , which replaces the bare one J in the effective theory. As shown in Refs. [38, 39], in this case \tilde{J} jumps discontinuously to zero as the temperature increases, implying that the exponent of the correlations becomes infinite in the high-temperature region. This path has been pursued for the case of a power-law interaction as well: in this case both the coupling J and the exponent of the interaction σ are replaced by variational parameters \tilde{J} and s . The results are unambiguous as long as $\sigma > 2$, predicting a phenomenology analogous to the SR case [22]. Our aim is to generalize such an analysis in order to correctly deal with the $\sigma \in (0, 2]$ regime.

To achieve this scope, we replace the cosine in the original Hamiltonian (3) with a quadratic term

$$\beta H_0 = \frac{1}{4} \sum_{\mathbf{i}, \mathbf{j}} \tilde{J}(\mathbf{r}) (\theta_{\mathbf{i}} - \theta_{\mathbf{j}})^2, \quad (14)$$

$\tilde{J}(r)$ being a completely arbitrary function of $\mathbf{r} = \mathbf{i} - \mathbf{j}$, to be determined from free energy minimization in a self-consistent way.

The quadratic Hamiltonian H_0 induces the Boltzmann measure

$$\langle \cdot \rangle_0 = \frac{1}{Z_0} \int \prod_{\mathbf{j}} d\theta_{\mathbf{j}} e^{-\beta H_0}, \quad (15)$$

where

$$Z_0 = e^{-\beta F_0} = \int \prod_{\mathbf{j}} d\theta_{\mathbf{j}} e^{-\beta H_0} \quad (16)$$

is the partition function of the model. The variational principle establishes that our best ansatz minimizes the variational free energy

$$\mathcal{F} = \beta F_0 + \beta \langle H \rangle_0 - \beta \langle H_0 \rangle_0. \quad (17)$$

On the other hand, from the equipartition theorem it follows that $\langle H_0 \rangle_0 = \frac{N}{2\beta}$ is independent on the choice of $\tilde{J}(\mathbf{r})$, so that we can safely ignore it. Since H_0 has the same quadratic structure of the low-temperature Hamiltonian, we can diagonalize H_0 as well by means of the Fourier transform

$$\beta H_0 = \frac{1}{2} \sum_{\mathbf{q} \in IBZ} \tilde{K}(\mathbf{q}) |\theta_{\mathbf{q}}|^2, \quad (18)$$

where $\tilde{K}(\mathbf{q})$ is given by Eq. (6) with $J(r)$ replaced by $\tilde{J}(r)$. As shown in Appendix B, the variational free energy takes the form

$$\mathcal{F} = \frac{1}{2} \sum_{\mathbf{q} \in IBZ} \ln \tilde{K}(\mathbf{q}) - \frac{N}{2} \sum_{\mathbf{r}} J(r) e^{-\tilde{G}(\mathbf{r})}, \quad (19)$$

where, once again, $\tilde{G}(\mathbf{r})$ is given by Eq. (9) with $J(r)$ replaced by $\tilde{J}(r)$. In order to simplify the notation, in the following we will drop the \sim symbol for $K(\mathbf{q})$.

Since in Eq. (19) $\tilde{J}(\mathbf{r})$ appears only through the $K(\mathbf{q})$, in order to find the minimum is sufficient to derive \mathcal{F} with respect to the latter. By exploiting the fact that

$$\frac{\delta \tilde{G}(\mathbf{r})}{\delta K(\mathbf{q})} = -\frac{1}{N} \frac{1 - \cos(\mathbf{q} \cdot \mathbf{r})}{K(\mathbf{q})^2}, \quad (20)$$

we find

$$\frac{\delta \mathcal{F}}{\delta K(\mathbf{q})} = \frac{K(\mathbf{q}) - \sum_{\mathbf{r}} J(r) (1 - \cos(\mathbf{q} \cdot \mathbf{r})) e^{-G(\mathbf{r})}}{2K(\mathbf{q})^2}. \quad (21)$$

By using the definition (6) of $K(q)$ and noticing that the above expression has to be valid for each value of $\mathbf{q} \in IBZ$ (first Brillouin zone), we find the condition

$$J(r) = \tilde{J}(\mathbf{r}) e^{\tilde{G}(\mathbf{r})}. \quad (22)$$

Since we are interested in the large length-scales regime (i.e in the continuous limit), we can assume \tilde{J} to be a function of r only: in this case, indeed, $\tilde{G}(\mathbf{r})$ only depends on the modulus r as well, so that the above condition can be written in terms of single variable functions

$$J(r) = \tilde{J}(r) e^{\tilde{G}(r)}. \quad (23)$$

Let us notice how, in this limit, one has to redefine $\mathbf{r} \rightarrow a\mathbf{r}$, $\mathbf{q} \rightarrow a^{-1}\mathbf{q}$, $J, \tilde{J} \rightarrow a^{-2}J, a^{-2}\tilde{J}$.

Let us now discuss the possible solutions of Eq. (23). The possible asymptotic behaviors of $K(q)$ and $\tilde{G}(q)$ in terms of σ are examined in Appendix A. We find: we find that:

- If $\tilde{J}(r)$ decays at infinity faster than r^{-4} (e.g. an exponential or a fast-decaying power law), then we have that $\tilde{G}(r) \sim \ln(r)$ as $r \rightarrow \infty$. Since in this case $e^{\tilde{G}(r)}$ is a power law, it follows from Eq. (23) that $\tilde{J}(r)$ must behave asymptotically as a power-law as well. We can then assume that for large r

$$\tilde{J}(r) \sim \tilde{J} r^{-2-s} \quad (24)$$

for some $s > 2$ to be determined, so that

$$\tilde{G}(r) \sim \eta(\tilde{J}) \ln \frac{r}{a} + A\tilde{J}^{-1}, \quad (25)$$

where $\eta(\tilde{J}) = p\tilde{J}^{-1}$ and A, p are non-universal constants. Finally, from Eq. (23), we have the conditions

$$\sigma = s - \eta(\tilde{J}); \quad J = \tilde{J} e^{A/\tilde{J}}. \quad (26)$$

In this case the correlation functions decay as $e^{-\tilde{G}(r)} \sim r^{-\eta(\tilde{J})}$. Then, as long as \tilde{J} is non-vanishing, we find a quasi-long-range order, characteristic of the BKT phenomenology.

- If the variational coupling behaves as $\tilde{J}(r) \sim \tilde{J}r^{-2-s}$ with $s \in (0, 2)$, then $G(r) = A\tilde{J}^{-1} + O(r^{s-2})$. In this case we then have:

$$\sigma = s; \quad J = \tilde{J}e^{A/\tilde{J}}, \quad (27)$$

leading to correlations behaving as $e^{-\tilde{G}(r)} \sim e^{-A\tilde{J}^{-1}}$. Then as long as \tilde{J} is non-vanishing, we find a finite magnetization $\sim e^{-A\tilde{J}^{-1}/2}$.

In both cases, the equation for \tilde{J} has the same form of the nearest-neighbour case. By introducing $\tilde{J} = Ax$, $J = Ay$ it can be written as:

$$y = xe^{1/x}. \quad (28)$$

The minimum of the r.h.s. term is in correspondence of $x = 1$, $y = e$ so that we have two solutions for $J > J_c \equiv eA$, and only the trivial solution for $J < J_c$, signaling a jump in \tilde{J} from $\tilde{J}_c = A$ to zero. However, only the larger of the two solutions present for $J > J_c$ is physically acceptable. Indeed it is the only one to have the correct asymptotic behavior $\tilde{J}(r) \sim J(r)$ in the large J regime, in which our SCHA becomes the low-temperature approximation studied in Sec. III. The meaning of this low-temperature, finite \tilde{J} phase, however, depends on the whether $s < 2$ (ordered phase) or $s > 2$ (quasi-long-range-ordered BKT phase). Let us notice that, in the latter case, we have that $\eta = p\tilde{J}^{-1}$ of Eq. (25) cannot be larger than a given value $\eta_c = p\tilde{J}_c^{-1}$. As a consequence we have that:

- For $\sigma > 2$, the only possibility is that $s = \sigma + \eta_{SR} > 2$. We are then in the first case, so that we get the usual BKT phenomenology. This in agreement with the findings of Ref. [22].
- For $\sigma < 2 - \eta_c$, the only possibility is that $s = \sigma$ with $s < 2$. We are then in the second case, i.e. we find a finite magnetization for low temperature.
- For $2 - \eta_c < \sigma < 2$ both solutions are actually viable so that it is unclear whether the system is in the ordered or in the quasi-long-range ordered phase. To establish which solution should we choose we should compute \mathcal{F} on both solutions. The dependence of \mathcal{F} on the non-universal details of the model, and in particular on the short-range behavior of $\tilde{J}(r)$, hinders the possibility to reach a definite conclusion for this regime within the SCHA.

Although non-conclusive, our self-consistent analysis accounts for the possibility of the existence of an intermediate region, in which the ordered phase or the BKT behavior can prevail, depending on the temperature. Keeping in mind the results of the Berezinskii approximation,

it appears sensible to think that the magnetized phase will prevail at lower temperatures, while the quasi-long-range ordered phase will, possibly, correspond to an intermediate range of temperature. Let us notice, however, that the predictions of our analysis are non-universal and, thus, their quantitative outcome depends on the peculiar model under study. Moreover, the first-order phase transition foreseen for $\sigma < 2 - \eta_c$ is could be an artifact of the approximation, since the critical behavior of the model is known to be captured by the mean-field approximation for $\sigma < 1$, which foresees a second-order phase transition, see e.g. Ref. [19].

V. FIELD-THEORETICAL APPROACH

In order to go beyond the limits of the SCHA, and capture the universal quantities we are interested in, we resort to a field-theoretical approach, introducing a continuous action, which encodes the same physics of our Hamiltonian (3). First, we write the coupling as $J(r) = J_S(r) + J_{LR}r^{-(2+\sigma)}$, J_S being a SR term which accounts for the small-distances behavior. This decomposition allows us to refine our low-temperature approximation. This, in fact, is fully justified for the SR coupling, even at intermediate temperatures, since it couples neighboring sites. At the same time, it can become too crude for the long-range part of the Hamiltonian. Since this couples far-away pairs, there can be smooth configurations for which the phase difference $\theta_i - \theta_j$ is not necessarily small, and these configurations may give a significant contribution to the Hamiltonian in an intermediate range of temperatures. However, even for the SR term, the approximation *à la* Berezinskii is unable to capture the presence of vortices.

To further proceed, we expand the cosine in the SR part

$$1 - \cos(\theta_i - \theta_j) \approx \frac{1}{2}(\theta_i - \theta_j)^2 \approx \frac{1}{2}|\nabla\theta|^2, \quad (29)$$

where the last substitution is justified by the fact that in the SR part of the Hamiltonian only neighbouring lattice sites are important. We then find

$$S[\theta] = \frac{J}{2} \int d^2x |\nabla\theta|^2 + S_{LR}[\theta], \quad (30)$$

where we introduced the LR perturbation

$$S_{LR} = J_{LR} \int d^2x \int_{r>a} \frac{d^2r}{r^{2+\sigma}} [1 - \cos(\theta(\mathbf{x}) - \theta(\mathbf{x} + \mathbf{r}))]. \quad (31)$$

In turn, this term can be rewritten in terms of the fractional Laplacian, whose definition is given, along with the details of the calculation, in Appendix C. We obtain

$$S_{LR} = \frac{g}{2} \int d^2x e^{-i\theta} \nabla^\sigma e^{i\theta}, \quad (32)$$




FIG. 1. Behavior of the RG flow for $y = 0$ in the $\frac{7}{4} < \sigma < 2$ regime. The dotted black line separates the region in which vortices are relevant (right, $\dot{y}_\ell > 0$), or not (left, $\dot{y}_\ell < 0$). We find three regions, corresponding to the three phases: for high temperatures (grey), the vortices push the system towards the disordered phase, for intermediate temperatures (gold) the system flows towards a SR $g = 0$ fixed point, while for low-temperatures (blue) the LR perturbation becomes relevant, pushing the system towards an ordered region. In bold black we plot the separatrices between the different phases.

with $g = J_{\text{LR}}/\gamma_{2,\sigma}$ and $\gamma_{2,\sigma} = 2^\sigma \Gamma(\frac{1+\sigma}{2}) \pi^{-1} |\Gamma(-\frac{\sigma}{2})|^{-1}$. This term is intrinsically interacting. Moreover it is invariant under global translations $\theta(\mathbf{x}) \rightarrow \theta(\mathbf{x}) + \alpha$, and it correctly catches the fact that the field θ is periodic. However, because of the kinetic term in $S[\theta]$, the whole action still does not correctly describes topological phenomena.

Let us remind briefly of the $g = 0$ case. Here one can introduce vortices in Eq. (30) i.e. evaluating the energy cost of a vortex configuration, and the core energy cost of a single vortex ε_c , which can be absorbed into the definition of the vortex fugacity $y = \exp(-\varepsilon_c)$. Since topological and non-topological configurations decouple in the quadratic part of the action, this leads to the Coulomb gas picture and to the well-known Kosterlitz-Thouless RG equations [42]. In turn, this implies that all the fixed points for $y = 0$ and $J > \frac{2}{\pi}$, which correspond to a Gaussian massless action, are stable. There, the low-temperature approximation becomes correct (with the original $J(T)$ replaced by the one corresponding to the fixed point), so that the power-law scaling observed for the two-point functions is recovered. For $J < \frac{2}{\pi}$, however, vortices become relevant and the theory flows towards the disordered regime.

In order to study the transition between the SR and the LR regime, we can consider the effect of a small g deformation of the Gaussian fixed points. In this case, we expect to be able to parametrize our theory in terms of three parameters: J , g and y . In Appendix D we derive the RG flow perturbatively in y , g . However, we

provide here an argument to understand what one can expect on general grounds. First, we notice that the fixed points with $J < \frac{2}{\pi}$ are already unstable under topological perturbations, so that we will stick to the stable fixed line with $J > \frac{2}{\pi}$. Here, the LR perturbation can be relevant or not depending on the scaling dimension Δ_g of the coupling g , $g_\ell \approx \exp(\Delta_g \ell)$ for $\ell \gg 1$, where $\ell = \ln(r/a)$ is the RG time.

Due to the quadratic nature of the measure, one has

$$\left\langle e^{i(\theta(\mathbf{x}) - \theta(\mathbf{x}'))} \right\rangle = e^{-\frac{1}{2} \langle (\theta(\mathbf{x}) - \theta(\mathbf{x}'))^2 \rangle} = |\mathbf{x} - \mathbf{x}'|^{-\eta_{\text{sr}}(J)}, \quad (33)$$

where $\eta_{\text{sr}}(J) = \frac{1}{2\pi J}$ corresponds to the exponent of the two-point function for $g = 0$ [42, 43, 51]. Then

$$\Delta_g = 2 - \sigma - \eta_{\text{sr}}(J). \quad (34)$$

The LR perturbation is relevant only in the regime $\sigma < 2 - \eta_{\text{sr}}(J)$. This bears some similarity with the SSB case [14], but with the main difference that the anomalous dimension is temperature dependent.

For $\sigma > 2$, the LR term is always irrelevant. For $\sigma < 2$ Eq. (34) predicts that the LR perturbation is always relevant at low enough temperatures. Since the points which are stable under the proliferation of vortices are those with $J > \frac{2}{\pi}$, we have that $0 < \eta_{\text{sr}} < \frac{1}{4}$, as in the usual BKT theory (which indeed corresponds to the $g = 0$ theory). As a consequence, for $\frac{7}{4} < \sigma < 2$, we have that it exists an intermediate range of value of J for which the Gaussian theory is stable with respect to both the topological and the LR perturbations, leading to conventional quasi-long-range order in a given temperature window. Instead, for $\sigma < 7/4$ the BKT stable fixed line is completely swallowed by the action of the LR perturbation, so that our perturbative picture breaks down. However, it is sensible to assume that in this regime the system simply undergoes an order-disorder phase transition.

This picture is essentially in agreement with the results of the SCHA. However, our results no longer depends on the regularization procedure or the exact form of $J_S(r)$ and are genuinely universal. Our observation can be strengthened by the renormalization equations (see Appendix D for details) which, at the leading order in y and g are given by

$$\begin{aligned} \frac{dy_\ell}{d\ell} &= (2 - \pi J)y, \\ \frac{dg_\ell}{d\ell} &= \left(2 - \sigma - \frac{1}{2\pi J_\ell}\right)g_\ell, \\ \frac{dJ_\ell}{d\ell} &= c_\sigma \eta_{\text{sr}}(J_\ell)g_\ell, \end{aligned} \quad (35)$$

with $c_\sigma = \frac{\pi}{2} a^{2-\sigma} \int_1^\infty du u^{1-\sigma} \mathcal{J}_0(2\pi u)$ and $\mathcal{J}_0(x)$ is the zeroth-order Bessel function of the first kind. Our argument correctly predicts the first two equations, but not the third one, i.e. the renormalization of the spin-waves stiffness. In particular, for $7/4 < \sigma < 2$, we find a line of stable quadratic fixed points for $g = y = 0$ and

$J_{\text{BKT}} \equiv \frac{2}{\pi} < J < J_\sigma \equiv \frac{1}{2\pi(2-\sigma)}$, as expected. The behavior of y_ℓ , at this order, is completely determined by J and, as long as $J > J_{\text{BKT}}$, $y_\ell \rightarrow 0$.

Then, for $7/4 < \sigma < 2$ we can characterize the transition between the ordered phase and the quasi-long-range ordered one by looking at the $y = 0$ plane. As long as g is small, we can explicitly identify the form of the flow trajectories of Eqs. (35):

$$g_\ell(J) = \frac{\pi(2-\sigma)}{c_\sigma} \left[(J_\ell - J_\sigma)^2 + k \right]. \quad (36)$$

If $k < 0$ the trajectory arrives at the fixed point line $g = 0$ for some $J < J_\sigma$, while for $k > 0$ the trajectory starts from the $g = 0$ line for $J > J_\sigma$ and it goes to infinity, signaling the existence of a new, low-temperature, phase. The separatrix is given by the semi-parabola with $k = 0$, $J \leq J_\sigma$. The graphical depiction of the RG flow of Eqs. (35), in the $y = 0$ plane, is shown in Fig. 1.

Being Eqs. (35) a perturbative result we derived for small g_ℓ and y_ℓ , its use in the low temperature region ($T < T_c$) is not justified, as g_ℓ grows indefinitely. However, for $T \rightarrow T_c^-$, the amount of time spent by the flow in the region close to $J = J_\sigma$, $g = 0$ becomes larger and larger, so that the scaling behavior of g_ℓ with T in this regime can be reliably obtained from Eqs. (35). In particular, for fixed ℓ such that $J_\ell > J_\sigma$ we have

$$g_\ell \sim e^{-B(T-T_c)^{-1/2}} \quad (37)$$

where B is a non universal constant (see Appendix E for details).

Although the infrared regime is beyond the reach of our perturbative analysis, it is possible to guess the corresponding form taken by the action on physical grounds. Indeed, the coupling J diverges there, suppressing spatial fluctuations of the phase $\theta(\mathbf{x})$, as confirmed by the rigorous results of Ref. [20], which predict a finite magnetization for low enough temperatures. This suggests that, in the infrared region corresponding to the low-temperature phase, we are allowed to Taylor expand the exponential in Eq. (30), so that the action becomes

$$S_{\text{LR}} = -\frac{g}{2} \int d^2\mathbf{x} \, \theta \nabla^\sigma \theta, \quad (38)$$

where we absorbed in g some immaterial constants. A further evidence in favour of this action comes from the fact that, in the limit $J \rightarrow \infty$, we have $\Delta_g = 2 - \sigma - \eta_{sr}(J) \rightarrow 2 - \sigma$, which is exactly the scaling dimension of g in Eq. (38). This also suggests that in this regime topological excitations are suppressed.

Physically speaking, this is due to the fact that, as $J \rightarrow \infty$, the energy cost of highly non-local excitations like the topological one becomes higher and higher, and the presence of relevant LR perturbation further contributes to this. Let us notice that the action of Eq. (38) is nothing but the continuous version of the approximation *à la* Berezinskii. This tells us that, for small enough




FIG. 2. Qualitative phase diagram for the long-range XY model in $d = 2$, in which the three phases of the model (ordered, quasi-long-range-ordered, disordered) are shown. For $\sigma > 2$ the system undergoes the usual BKT transition (gray line). The value σ^* at which the LR term becomes relevant is here a function of the temperature and varies from $\sigma = 2$ (at $T = 0$) to $7/4$ (when it met the BKT transition temperature). In this range of σ then, as the temperature varies, beyond the BKT transition, we find an infinite order symmetry-breaking transition (blue line). For $\sigma < 7/4$ the quasi-long-range ordered phase disappears so that we only have an order-disorder transition.

temperature, the approximation is indeed reliable. In particular, we can use Eq. (13) to derive the magnetization

$$\ln m \sim g^{-1} \int q^{-\sigma} d^2\mathbf{q} \quad (39)$$

Finally, from Eq. (37) we find the scaling of the magnetization

$$\ln m \sim -e^{B(T_c-T)^{-1/2}} \quad (40)$$

Here, the phase transition is of infinite order as all the derivatives of the order parameter with respect to the temperature vanish at $T = T_c$. The same can be said for the free energy which would as well exhibit an essential singularity. A similar behavior in T is found approaching T_{BKT} from above, so that this seems to be a general property of BKT phase. The presence of a SSB phase is, however, proper of the LR regime. A similar, albeit not identical, coexistence of finite order parameter and infinite order scaling signatures has been also observed in other long-range statistical mechanics models [52–56], see also Ref. [57] for more examples in this direction.

As $\sigma \rightarrow 7/4^+$, T_c reaches T_{BKT} from below, leaving only a SSB phase transition. Unfortunately however, our set of equations (35) is not reliable in this regime: the RG flow spends a considerable close to the $g = 0$, $J = J_\sigma$ fixed point, which, in this regime, corresponds to $\dot{y}_\ell > 0$. As a consequence, y_ℓ will not remain small.

Summarizing, when $7/4 < \sigma < 2$, we find *i*) for $T < T_c$, finite magnetization (ordered phase); *ii*) for $T_c < T < T_{\text{BKT}}$ quasi-long-range-order with zero magnetization and temperature-dependent power-law decay in the two-point correlation function (BKT phase); *iii*) for $T > T_{\text{BKT}}$ zero magnetization (disordered phase). The system, due to the power-law character of the interactions, exhibits power-law decaying two-point functions, also in the high-temperature phase, although with the same exponent of the coupling $\langle \mathbf{S}(\mathbf{r}) \cdot \mathbf{S}(0) \rangle \sim r^{-2-\sigma}$ [58–60]. The qualitative form of the phase diagram of the model is shown in Fig. 2.

VI. VILLAIN APPROXIMATION

The study of the SR XY model in $d = 2$ is greatly simplified by the possibility of performing the so-called Villain approximation, in which we replace in the Hamiltonian in Eq. (2)

$$1 - \cos(\theta_i - \theta_j) \rightarrow \frac{1}{2} (\theta_i - \theta_j - 2\pi n_{i,j})^2 \quad (41)$$

where $n_{i,j}$ is an auxiliary link variable which assumes integer values and which has to be traced out. Although the variable appears only quadratically, the model exhibits a BKT phenomenology, since it correctly reproduces the $\theta_i \rightarrow \theta_i + 2\pi m$ symmetry with $m \in \mathbb{Z}$. In particular, by integrating out the θ_i , the Villain Hamiltonian can be exactly mapped into the Coulomb gas Hamiltonian (in any dimension).

While the same approximation could be, in principle, performed even for the case of long-range-decaying couplings, it is not guaranteed for the resulting phenomenology to be in the same universality class of the LR XY model. There are, in fact, reasons to believe this is not true, as we are going to argue. First, let us notice that, if on the one hand the Villain approximation improves Berezinskii's result by taking into account the vortices, on the other hand, it can only account for interaction between topological excitations which is quadratic in their charges. As already seen in the context of the field theory, however, such a quadratic approximation is not always justified.

To be more concrete, let us express the action $S[\theta]$ of the XY model in terms of the topological charges. We start by splitting the field in a topological and a non-topological spin-waves part:

$$\theta(\mathbf{x}) = \theta_0(\mathbf{x}) + \theta_{\text{top}}(\mathbf{x}) \quad (42)$$

where, for every closed circuit,

$$\oint \nabla \theta_0 \cdot d\mathbf{x} = 0 \quad \oint \nabla \theta_{\text{top}} \cdot d\mathbf{x} = 2\pi m_{\text{enc}} \quad (43)$$

m_{enc} being the total topological charge enclosed in the circuit. The ambiguity in the decomposition can be lifted if we impose the further constraint $\nabla \cdot \nabla \theta_{\text{top}} = 0$. This,

in turn, allows us to write θ_{top} in terms of the vortex configuration so that we can finally write Eq. (30) as

$$S[\theta] = \frac{J}{2} \int d^2 \mathbf{x} |\nabla \theta_0|^2 - \pi J \sum_{j \neq k} m_j m_k \ln |\mathbf{x}_k - \mathbf{x}_j| + \varepsilon_c \sum_k m_k^2 + \frac{g}{2} \int d^2 \mathbf{x} e^{-i(\theta_0 + \theta_{\text{top}})} \nabla^\sigma e^{i(\theta_0 + \theta_{\text{top}})} \quad (44)$$

where m_k and \mathbf{x}_k are respectively the charges and the positions of each vortex (see Appendix F for details). For small values of g we can integrate out the non-topological component of the field, θ_0 , finding

$$S_{\text{eff}} \sim -\pi J \sum_{j \neq k} m_j m_k \ln |\mathbf{x}_k - \mathbf{x}_j| + \varepsilon_c \sum_k m_k^2 + \frac{g}{2} \int d^2 \mathbf{x} e^{-i\theta_{\text{top}}} \nabla^{\sigma + \eta_{\text{sr}}(J)} e^{i\theta_{\text{top}}} \quad (45)$$

where it is implied that, if $\sigma + \eta_{\text{sr}}(J) > 2$, the above operator has to be interpreted as the usual Laplacian (see Appendix F). In the latter case, in particular,

$$-e^{-i\theta_{\text{top}}} \nabla^2 e^{i\theta_{\text{top}}} = |\nabla e^{i\theta_{\text{top}}}|^2 = |\nabla \theta_{\text{top}}|^2 \quad (46)$$

so that, by replacing θ_{top} with its expression in terms of the m_i , we recover the usual Coulomb gas interaction

$$S_{\text{eff}} \sim -\pi(J + g) \sum_{j \neq k} m_j m_k \ln |\mathbf{x}_k - \mathbf{x}_j| + \varepsilon_c \sum_k m_k^2 \quad (47)$$

We thus recovered the main result of our analysis, namely, the fact that the system behaves as its SR counterpart even for $\sigma < 2$, provided that $\sigma + \eta_{\text{sr}}(J) > 2$.

If $\sigma + \eta_{\text{sr}}(J) < 2$, it is not possible to derive a simple charge-charge interaction, unless we expand the exponential. This expansion is actually not easily justified, since the topological configurations θ_{top} are spatially extended. This suggests that the higher-order terms in θ_{top} in the expansion are relevant, so that we have to keep in the Hamiltonian higher-order interaction terms in the m_k (e.g. proportional to $m_i m_j m_k m_p$), which cannot be obtained within the Villain approximation.

Moreover, let us notice that $\sigma + \eta_{\text{sr}}(J) < 2$ is precisely the condition under which the LR term becomes relevant. In this regime, our field-theoretical analysis foresees a boundless growth for the coupling g , so that the integration on θ_0 cannot be performed perturbatively as we did in Eq. (45). The fact that spin-waves and topological contributions do not decouple is a further proof that the long-range phenomenology cannot be captured by a quadratic model in θ , as the Villain model.

VII. QUANTUM LONG-RANGE XXZ CHAIN

The mapping between the nearest-neighbors $2d$ classical XY model and the nearest-neighbors, spin-1/2, $1d$




FIG. 3. Qualitative phase diagram of the LR XXZ quantum chain studied in Ref. [25]. The CSB (continuous symmetry breaking) phase corresponds to the SSB phase of the LR XY model, the AF (anti-ferromagnetic) phase to the disordered one, the gapless XY phase to the BKT phase. The topology of the phase diagram is the same of the LR XY case (Fig. 2), but here the border between the new broken phase and the gapless XY phase of the chain goes from $\sigma = 2$ to $\sigma = 1$, as discussed in the text. In the region $\sigma < 1$ the intermediate phase disappears.

quantum XXZ model relies on the local nature of the couplings [23]. However, the extension of the equivalence in presence of LR couplings is not obvious. Indeed, the original derivation in Ref. [23], neglects the z - z interaction terms with a range ≥ 4 lattice sites in the resulting Hamiltonian and introduces a suitable, averaged, interaction term for distances lower than three lattice sites. This allows introducing a bosonic hard-core condition and the mapping onto the $1d$ quantum XXZ Hamiltonian. This approach cannot be straightforwardly applied to the case of the $2d$ XY model with LR interactions, as one should show the RG irrelevance of terms violating the hard-core condition.

Therefore, an interesting question is to ascertain whether and how the $T = 0$ phase diagram of the $1d$ XXZ Hamiltonian with LR interactions

$$H = - \sum_{i,r} \frac{1}{r^{1+\sigma}} (S_i^x S_{i+r}^x + S_i^y S_{i+r}^y - \Delta S_i^z S_{i+r}^z) \quad (48)$$

(where $\sigma > 0$ and $S^{x,y,z}$ are the components of the spin $1/2$ operators) is related to that of the $2d$ classical XY model. It is worth noting that Hamiltonian (48) has been studied in Ref. [25] through the bosonization technique and numerical simulations. The effective action derived there describing the XXZ model bears a strong resemblance to the one of Eq. (30). The resulting RG flow is therefore similar to the one of Eq. (35) indicating that the LR-SR crossover of the two models is somehow analogous. In particular, identifying in Eqs. (4) and (8) of Ref. [25] K, g_{LR}, g with $\pi J/4, g, y$ respectively, these flow

equations becomes

$$\begin{aligned} \frac{dy_\ell}{d\ell} &= (2 - \pi J)y \\ \frac{dg_\ell}{d\ell} &= \left(2 - \sigma - \frac{2}{\pi J_\ell}\right)g_\ell \\ \frac{dJ_\ell}{d\ell} &= A(J)g_\ell \end{aligned} \quad (49)$$

$A(J)$ being a function of J , depending on the RG-scheme. The similarity between Eq. (35) and Eq. (49) can be traced back both to the fact that the z - z LR interaction is actually irrelevant (for $\sigma > 0$) and to the fact that the coefficient of the time derivative in the bosonized action does not renormalize (see the corresponding discussion for $\sigma < 0$ in Ref. [61]). This also explains why both phase diagrams feature three phases: the SSB, the disorder and the BKT one, which are called in Ref. [25] continuous symmetry breaking (CSB) phase, anti-ferromagnetic (AF) phase and gapless XY phase, respectively.

Let us notice that in the LR regime, the usual mapping based on the coherent spin states, would result in a highly anisotropic action, due to the fact that one still has SR interactions along the Euclidean time direction [62]. In principle, a similar anisotropy would be present in the bosonized action of Ref. [25] due to the presence of the z - z LR contribution. As already noticed, however, this term turns out to be irrelevant so that the effective infrared action is actually isotropic.

Notice that Eqs. (49) predict that the BKT phase extends here up to $\sigma = 1$, rather than $\sigma = 7/4$. This difference boils down to the fact that, in the SR regime, in correspondence to the border between the XY and the anti-ferromagnetic phase ($\Delta = 1$), the correlation functions of the x, y components decay as r^{-1} , rather than $r^{-1/4}$, so that the corresponding anomalous dimension is $\eta_{sr} = 1$ and therefore, using the formalism introduced with Sak's criterion, the smallest value of σ_* is 1 and σ_* ranges between 1 and 2. This has to be compared with the range for σ_* between $7/4$ and 1 for the classical LR XY at finite temperature in $d = 2$.

Even if the ranges of σ_* do not coincide, the qualitative form of the phase diagrams of the two models is, therefore, the same, as it is possible to see from Fig. 3. This is similar to the results in Fig. 1 of Ref. [25] (let us notice that $1/\alpha \equiv 1/(1 + \sigma)$ on the vertical axis is replaced by $1/\sigma$ in our Fig. 3). In that figure two lines, one separating the BKT and the SSB phases, and the other separating the BKT and the disordered phases, are drawn as extracted from numerical DMRG simulations. It is not clear by the figure whether and where the two lines are going to merge in correspondence of $\sigma = 1$, as foreseen by Eqs. (49) and in agreement with the picture presented in this paper. Therefore, more extensive numerical simulations are needed to confirm the predictions from Eqs. (49) in the vicinity of $\sigma = 1$.

Moreover, the substantial similarity between the field theory descriptions of the two models implies that the

magnetization m_{\parallel} in the x - y plane of the XXZ chain, scales as

$$\ln m_{\parallel} \sim -e^{B(\Delta_c - \Delta)^{-1/2}} \quad (50)$$

i.e. it follows the same scaling of Eq (40). This is, to the best of our knowledge, a new prediction for the XXZ LR model.

The present result fits within the effective-dimensionality picture described in Ref. [19], for LR quantum systems. In particular, the crossover between various LR regimes of the quantum $O(n)$ model in dimension d can be described by introducing the effective dimension

$$D_{\text{eff}} = \frac{2 - \eta_{\text{sr}}(D_{\text{eff}})}{\sigma} d + 1 \quad (51)$$

valid for any $\sigma < 2 - \eta_{\text{sr}}(d)$, $\eta_{\text{sr}}(D)$ being the anomalous dimension of the D -dimensional action which describes the SR version of our model. For the case the LR $d = 1$ XXZ phase-diagram, when $D_{\text{eff}} = 2$, we know that the system undergoes a BKT transition, for which η_{sr} is not defined. If, however, we take $\eta_{\text{sr}} \in [0, 1]$, rather than a single value, we find, from Eq. (51) we have

$$2 > \sigma > 1 \quad (52)$$

that is, exactly the range of coexistence of the SSB and the BKT phases.

For $\sigma < 2/3$, we find $D_{\text{eff}} > 4$ (with $\eta_{\text{sr}}(D_{\text{eff}}) = 0$), so that the order-disorder transition is expected to be captured by the mean-field picture. The same thing it is supposed to happen for the $d = 2$ classical XY model in the $\sigma < 1$ region. Therefore one could expect that the critical behavior of the quantum chain in the $2/3 < \sigma < 1$ interval can be related to the one of the classical model for $1 < \sigma < 7/4$. However, given that Eqs. (49) and Eq. (35) do not, respectively, apply in these regimes, further studies would be necessary to clarify this point. A table summarizing the analogous phases of the quantum XXZ and the classical XY model is presented, see Tab. I.

The present analysis and, especially, the extension of the SCHA approach to LR couplings, discussed in Sec. IV, could be extended to study generic quantum spin- S systems with LR interactions. According to the Haldane conjecture [63–65] antiferromagnetic half-integer spin chains have a gapless excitation spectrum and they are known to display BKT scaling (e.g., for spin-1/2 XXZ model in the SR regime at $\Delta = 1$), so that we expect the generalization of our picture to higher, half-integer, values of the spin to be straightforward.

Regarding the integer-spin case, instead, one can have BKT transitions on the ferromagnetic side, as shown for the spin-1 chain in [66]: therefore, one could also expect a phase diagram like the one in Fig. 3, also for integer spin chains; more investigation is, anyway, needed. The application of our studies to ferromagnetic couplings requires the generalization of the transfer-matrix approach

$2d \ XY$	$1d \ XXZ$
$\sigma < 1$	$\sigma < 2/3$
$1 < \sigma < 7/4$	$2/3 < \sigma < 1$
$7/4 < \sigma < 2$	$1 < \sigma < 2$
$\sigma > 2$	$\sigma > 2$

TABLE I. Correspondence between the phases of the $2d \ XY$ LR model at finite temperature (left) and in the $1d \ XXZ$ LR model at zero temperature (right). For the values of σ corresponding to the first line we have a mean-field SSB transition; in correspondence of the second line an interacting SSB transition; in correspondence of the third we have both the order-disorder phase transition and the BKT one; in correspondence of the fourth, finally, only the BKT transition is present.

beyond the antiferromagnetic case [23]. In this perspective, one can also resort to SCHA in order to reduce the quantum system to an effective classical one, which can then be simulated via Monte Carlo [67, 68].

VIII. CONCLUSIONS

We have shown how the rich and unusual phase diagram of the $2d$ classical XY model in presence of long-range (LR) interactions can be obtained both using a self-consistent approach and in a field-theoretical way. We also showed how this phase diagram can be related to the one of the LR XXZ quantum chain.

While the mapping between the two models is well established in the nearest-neighbors case [23], the similarity between the phase diagrams of the classical and quantum models in the LR case is remarkable. We provided reasons to believe that our results cannot be obtained via the Coulomb gas mapping or via the Villain approximation, as in the case of SR interactions, so that the latter models appear to be in a different universality class with respect to the $2d \ XY$ model when LR interactions are present. Along the same lines, numerical results, obtained in Refs. [69, 70], seem to indicate that the diluted version of the LR $d = 2 \ XY$ model has a different phase diagram as well, although further investigation would be needed in order to confirm this expectation.

We were also able to use the results obtained for the classical $2d$ system to foresee a new, non-analytic, scaling for the order parameter of the LR XXZ chain, which eludes the current classification [71]. Comments on higher-spin quantum chains with LR couplings were also provided.

The appearance of exotic scaling phenomena by the addition of complex interaction patterns in two-dimensional systems is not unique. Indeed, several interesting interplays between topological defect unbinding and other critical phenomena have been described, e.g. in the case of coupled XY planes [72], $2d$ systems with anisotropic dipolar interactions [73, 74], high-dimensional systems

with Lifshitz criticality [75, 76], and anisotropic 3d XY model [77].

Let us notice that we cannot derive any prediction from our analysis of the 2d XY model in the region $\sigma < 7/4$ ($\sigma < 1$ for the case of the XXZ quantum chain). It would be important to extend our analysis to the $\sigma < 7/4$ region, in order to determine whether the transition is a second-order one, as one would expect, and also whether the quantum to classical mapping we established is still valid.

Finally, it would be interesting to study the LR Villain model, in order to establish the failure of the Villain approximation to reproduce the phase diagram of the 2d LR XY universality.

Appendix A: $K(q)$ and $G(r)$

We derive the asymptotic expression for $K(q)$ and $G(r)$, presented in the text. In the following we will consider a $J(r)$ with a general σ , so that our results can be applied to the self-consistent-harmonic calculation as well.

First, we notice that the continuous definition of $G(r)$ and $K(q)$, given in Eq. (10) and Eq. (7) respectively, can be expressed, once we integrate on the solid angle, as

$$K(q) = 2\pi \int_a^\infty dr r J(r) (1 - \mathcal{J}_0(qr)) \quad (\text{A1})$$

and

$$G(r) = a^2 \int_0^\Lambda \frac{dq}{2\pi} \frac{q(1 - \mathcal{J}_0(qr))}{K(q)}, \quad (\text{A2})$$

where $\mathcal{J}_0(x)$ is the zeroth-order Bessel function of the first kind. As long as $\tilde{J}(r)$ decays slower than r^{-4} , then $K''(0)$ is finite, so that $K(q) \sim q^2$, the proportionality constant being non-universal. As a consequence, in this case

$$G(r) \sim \int_0^\Lambda dq \frac{1 - \mathcal{J}_0(qr)}{q}, \quad (\text{A3})$$

which can be rewritten, through the substitution $x = qr$, as

$$G(r) \sim \int_0^{\Lambda r} dx \frac{1 - \mathcal{J}_0(x)}{x}. \quad (\text{A4})$$

For large r , the dominant term in the integral above is $\ln(\Lambda r)$. We then conclude that:

$$G(r) \sim \eta \ln \frac{r}{a} + B, \quad (\text{A5})$$

η and B being two non-universal, cutoff dependent, constants. In particular, if we suppose $J(r)$ behaves at infinity as $Jr^{-2-\sigma}$, with $\sigma > 2$, we have $\eta, B \propto J^{-1}$, so that we can write

$$G(r) \sim \eta(J) \ln \frac{r}{a} + AJ^{-1} \quad (\text{A6})$$

with $\eta(J) = p/J$ for some p .

Let us now consider instead the case $J(r) \sim \frac{J}{r^{2+\sigma}}$, with $\sigma \in (0, 2)$. By operating the substitution $x = qr$ we obtain

$$\begin{aligned} K(q) &= 2\pi J q^\sigma \int_{aq}^\infty dx x^{-2-\sigma} (1 - \mathcal{J}_0(x)) \\ &= 2\pi J q^\sigma \int_0^\infty dx x^{-2-\sigma} (1 - \mathcal{J}_0(x)) + O(q^2), \end{aligned} \quad (\text{A7})$$

so that we have $K(q) = c_\sigma J q^\sigma + O(q^2)$ with $c_\sigma = 2^{-\sigma} \pi |\Gamma(-\sigma/2)|/\Gamma(1+\sigma/2)$. The expression for $G(r)$ becomes then

$$G(r) \sim \frac{1}{J} \int_0^\Lambda dq q^{1-\sigma} (1 - \mathcal{J}_0(qr)) \quad (\text{A8})$$

which asymptotically goes as

$$G(r) = AJ^{-1} + O(r^{\sigma-2}). \quad (\text{A9})$$

Appendix B: Variational free energy

We derive in this Appendix Eq. (19), starting from the definition of \mathcal{F} in Eq. (17). Since $\langle \cos A \rangle_0 = e^{-\frac{1}{2}\langle A^2 \rangle_0}$, as valid for every Gaussian measure, we find

$$\begin{aligned} \beta \langle H \rangle_0 &= -\frac{1}{2} \sum_{\mathbf{i}, \mathbf{j}} J(r) \langle \cos(\theta_{\mathbf{i}} - \theta_{\mathbf{j}}) \rangle_0 \\ &= -\frac{1}{2} \sum_{\mathbf{i}, \mathbf{j}} J(r) e^{-\frac{1}{2}\langle (\theta_{\mathbf{i}} - \theta_{\mathbf{j}})^2 \rangle_0} \\ &= -\frac{N}{2} \sum_{\mathbf{r}} J(r) e^{-G(|\mathbf{j}-\mathbf{i}|)} \\ &= -\frac{N}{2} \sum_{\mathbf{r}} J(r) e^{-G(r)}, \end{aligned} \quad (\text{B1})$$

where we used the definition (9) of $G(\mathbf{r})$. By the diagonalized form (18) of H_0 , it follows immediately

$$\beta F_0 = \frac{1}{2} \sum_{\mathbf{q} \in \text{IBZ}} \ln K(\mathbf{q}). \quad (\text{B2})$$

Plugging this result into the expression for \mathcal{F} , we find the desired result.

Appendix C: Fractional Laplacian and S_{LR}

We want now to provide the definition of the fractional Laplacian, that we used, and derive the form (32) of S_{LR} .

For any $\sigma \in (0, 2)$, and a function $f(\mathbf{x}) : \mathbb{R}^d \rightarrow \mathbb{R}$ one can define $\nabla^\sigma f(\mathbf{x})$ as:

$$\nabla^\sigma f(\mathbf{x}) \equiv \gamma_{d,\sigma} \int d^d r \frac{f(\mathbf{x} + \mathbf{r}) - f(\mathbf{x})}{r^{d+\sigma}}, \quad (\text{C1})$$

with $\gamma_{d,\sigma} = \frac{2^\sigma \Gamma(\frac{d+\sigma}{2})}{\pi^{d/2} |\Gamma(-\frac{\sigma}{2})|}$. One can derive an alternative expression for this quantity in the momentum space. In terms of Fourier transform of $f(\mathbf{x})$, $f(\mathbf{q})$, one finds

$$\nabla^\sigma f(\mathbf{x}) = -\gamma_{d,\sigma} \int d^d q f(\mathbf{q}) e^{i\mathbf{q}\cdot\mathbf{x}} \int d^d r \frac{1 - e^{i\mathbf{q}\cdot\mathbf{r}}}{r^{d+\sigma}}. \quad (\text{C2})$$

and, exploiting the fact that,

$$\int d^d r \frac{1 - e^{i\mathbf{q}\cdot\mathbf{r}}}{r^{d+\sigma}} = \gamma_{d,\sigma}^{-1} q^\sigma, \quad (\text{C3})$$

one have

$$\nabla^\sigma f(\mathbf{x}) = - \int d^d q q^\sigma f(\mathbf{q}) e^{i\mathbf{q}\cdot\mathbf{x}}. \quad (\text{C4})$$

(from which, in the limit $\sigma \rightarrow 2$ one recovers the usual behavior of the standard Laplacian).

In our case, we can notice that the quantity present in Eq (31), namely

$$\int d^2 x \int_{r>a} \frac{d^2 r}{r^{2+\sigma}} [1 - \cos(\theta(\mathbf{x}) - \theta(\mathbf{x} + \mathbf{r}))]. \quad (\text{C5})$$

naturally fits into the definition of a two-dimensional fractional Laplacian. Indeed, let us notice that, provided that $\sigma < 2$, one can actually extend the integral on r on the whole space, and absorb the contribution coming from the $r < a$ region into the definition of the SR term. Then, we can write the additional LR term as the real part of

$$\begin{aligned} & \int d^2 x \int \frac{d^2 r}{r^{2+\sigma}} [1 - e^{i\theta(\mathbf{x}+\mathbf{r}) - i\theta(\mathbf{x})}] = \\ & e^{-i\theta(\mathbf{x})} \int d^2 x \int \frac{d^2 r}{r^{2+\sigma}} [e^{i\theta(\mathbf{x})} - e^{i\theta(\mathbf{x}+\mathbf{r})}]. \end{aligned} \quad (\text{C6})$$

In turn, this can be rewritten by exploiting the definition (C1) of the fractional Laplacian,

$$-\gamma_{2,\sigma}^{-1} \int d^2 x e^{-i\theta} \nabla^\sigma e^{i\theta}. \quad (\text{C7})$$

The expression above is already real, so that we recover the form of the long-range term given in the main text.

Appendix D: Renormalization procedure

We provide here a derivation of Eqs. (35) through the RG procedure. Let us consider the action $S[\theta]$ written as

$$S[\theta] = \int d^2 x \left(\frac{J_\ell}{2} |\nabla\theta|^2 + \frac{g_\ell}{2} \int_{r>a} \frac{d^2 r}{r^{2+\sigma}} [1 - \cos(\Delta_{\mathbf{r}}\theta(\mathbf{x}))] \right) \quad (\text{D1})$$

with $\Delta_{\mathbf{r}}\theta(\mathbf{x}) = \theta(\mathbf{x} + \mathbf{r}) - \theta(\mathbf{x})$. We are going to compute the RG flow equations perturbatively in g and in the vortex fugacity y . Let us notice, however, how the effect of vortices is not encoded in Eq. (D1). However, at the

first perturbative order in y and g the two perturbations act independently, so that we can consider the effect of the renormalization of the vortices on the SR kinetic term only. As usual in this case, one can map this theory into the Sine-Gordon action:

$$S_{\text{SG}} = \int d^2 \mathbf{x} \left(\frac{1}{2J_\ell} |\nabla\varphi|^2 - y_\ell \cos 2\pi\varphi \right). \quad (\text{D2})$$

At the first order of the renormalization group, we get that y varies accordingly the scaling dimension of the vertex operator $\cos(2\pi\varphi)$

$$\frac{dy_\ell}{d\ell} = (2 - \pi J) y, \quad (\text{D3})$$

while J_ℓ is left unchanged. This is in agreement with the Kosterlitz-Thouless RG for the SR, in which $\dot{J} = O(y^2)$.

Let us now consider the first-order effect of the long-range perturbation: in this case we can set $y = 0$. We can then write the field $\theta(\mathbf{x})$ as the sum of a fast-varying and a slow-varying component. In particular, introducing the momentum cutoff $\Lambda = \frac{2\pi}{a}$, we have $\theta = \theta^> + \theta^<$ with

$$\begin{aligned} \theta^<(\mathbf{x}) &= \int_{q < \Lambda e^{-d\ell}} \frac{d^2 q}{(2\pi)^2} \theta(\mathbf{q}) e^{i\mathbf{q}\cdot\mathbf{x}} \\ \theta^>(\mathbf{x}) &= \int_{\Lambda > q > \Lambda e^{-d\ell}} \frac{d^2 q}{(2\pi)^2} \theta(\mathbf{q}) e^{i\mathbf{q}\cdot\mathbf{x}}, \end{aligned} \quad (\text{D4})$$

and integrate out $\theta^>$. The non-Gaussian part of the action can be expanded in cumulant: at the first order we have

$$S_{\text{eff}}[\theta^<] = S_0[\theta^<] + \langle S_{\text{LR}} \rangle_> + O(g^2). \quad (\text{D5})$$

Because of the \mathbb{Z}_2 simetry in θ symmetry, we can replace $\cos(\Delta_{\mathbf{r}}\theta)$ with $\cos(\Delta_{\mathbf{r}}\theta^>) \cos(\Delta_{\mathbf{r}}\theta^<)$, finding, up to immaterial constants

$$\langle S_{\text{LR}} \rangle_> = \frac{g_\ell}{2} \int d^2 x \int \frac{d^2 r}{r^{2+\sigma}} \langle \cos(\Delta_{\mathbf{r}}\theta^>) \rangle_> [1 - \cos(\Delta_{\mathbf{r}}\theta^<)] \quad (\text{D6})$$

(from now on we let the integration run from $r = 0$). Exploiting the identity $\langle \cos(\Delta_{\mathbf{r}}\theta^>) \rangle_> = e^{-\frac{1}{2} \langle (\theta(\mathbf{r}) - \theta(0))^2 \rangle_>}$, valid on the Gaussian measure, one has

$$\begin{aligned} & \frac{1}{2} \langle (\theta(\mathbf{r}) - \theta(0))^2 \rangle_> \\ &= \int_{\Lambda > q > \Lambda e^{-d\ell}} \frac{d^2 q}{(2\pi)^2} \frac{1 - \cos(\mathbf{q} \cdot \mathbf{r})}{J_\ell q^2} \\ &= \frac{d\ell}{2\pi J_\ell} \left(1 - \mathcal{J}_0(\Lambda r) \right), \end{aligned} \quad (\text{D7})$$

where, as before, we denoted with $\mathcal{J}_0(x)$ the zeroth-order Bessel function of the first kind. Then, remembering that $\eta_{\text{sr}}(J) = \frac{1}{2\pi J_\ell}$, we have

$$\begin{aligned} \langle \cos(\Delta_{\mathbf{r}}\theta^>) \rangle_> &= e^{-\eta_{\text{sr}}(J_\ell) d\ell (1 - \mathcal{J}_0(\Lambda r))} \\ &= 1 - \eta_{\text{sr}}(J_\ell) d\ell + \eta_{\text{sr}}(J_\ell) d\ell \mathcal{J}_0(\Lambda r) \end{aligned} \quad (\text{D8})$$

The first two terms provide, as expected, an anomalous dimension of the coupling g , namely $g_{\ell+d\ell} = g_{\ell}e^{-\eta_{\text{sr}}(J_{\ell})d\ell}$. The last term, however, results in a new term in the action of the form

$$\langle S_g \rangle = \frac{1}{2} \int d^2x \left\{ \int \frac{d^2r}{r^{2+\sigma}} g e^{-\eta_{\text{sr}}(J_{\ell})d\ell} [1 - \cos(\Delta_{\mathbf{r}}\theta^<)] + g \eta_{\text{sr}}(J_{\ell})d\ell \int \frac{d^2r}{r^{2+\sigma}} \mathcal{J}_0(\Lambda r) [1 - \cos(\Delta_{\mathbf{r}}\theta^<)] \right\}. \quad (\text{D9})$$

The second term of l.h.s. has the same form of the original XY form. However, for $x \gg 1$ $\mathcal{J}_0(x) \sim x^{-1/2} \cos(x - \pi/4)$ so that this effective interaction has a fast-decaying oscillating behavior. Since this act as a natural cutoff for $r \sim \Lambda^{-1} \sim a$ this can be interpreted as an additional local interaction, and reabsorbed into the SR part of $S[\theta]$. One natural way to procede is to expand $1 - \cos(\Delta_{\mathbf{r}}\theta) \approx \frac{1}{2}(\mathbf{r} \cdot \nabla_{\mathbf{x}}\theta)^2$ so that

$$\int \frac{d^2r}{r^{2+\sigma}} \mathcal{J}_0(\Lambda r) (\mathbf{r} \cdot \nabla_{\mathbf{x}}\theta^<)^2 = \pi |\nabla_{\mathbf{x}}\theta^<|^2 \int_a^{\Lambda^{-1}} dr r^{1-\sigma} \mathcal{J}_0(\Lambda r). \quad (\text{D10})$$

For $\sigma > \frac{1}{2}$ the integral is infrared convergent so that one can neglect the cutoff (this is safe, since we are interested in the $\sigma > 7/4$ regime). Finally, putting $r = au$, we find the correction of the action as

$$\frac{c_{\sigma}}{2} (g_{\ell} a^{2-\sigma}) \eta_{\text{sr}}(J_{\ell}) d\ell \int d^2x |\nabla_{\mathbf{x}}\theta^<|^2, \quad (\text{D11})$$

where $c_{\sigma} = \frac{\pi}{2} \int_1^{\infty} du u^{1-\sigma} \mathcal{J}_0(2\pi u) > 0$. However, the precise expression of the coefficient is influential important for what follows.

Summarizing, we found that the effect of the integration over the fast modes can be reabsorbed, at the first order in g, y , into a redefinition of the couplings $g \rightarrow g + dg$, $J \rightarrow J + dJ$ with

$$\begin{aligned} dg &= -\eta_{\text{sr}}(J_{\ell}) g_{\ell} d\ell \\ dJ &= c_{\sigma} \eta_{\text{sr}}(J_{\ell}) (g_{\ell} a^{2-\sigma}) d\ell. \end{aligned} \quad (\text{D12})$$

Finally, we ought to perform the rescaling $\mathbf{x} \rightarrow \mathbf{x} e^{-d\ell}$, in order to obtain a theory with the same cutoff as the original one. Then g, J are further modified by their own bare length-dimension (namely $2 - \sigma$ and 0)

$$\begin{aligned} dg &= (2 - \sigma - \eta_{\text{sr}}(J_{\ell})) g_{\ell} d\ell \\ dJ &= c_{\sigma} \eta_{\text{sr}}(J_{\ell}) (g_{\ell} a^{2-\sigma}) d\ell. \end{aligned} \quad (\text{D13})$$

The RG equations then take the form

$$\begin{aligned} \frac{dg}{d\ell} &= (2 - \sigma - \eta_{\text{sr}}(J_{\ell})) g_{\ell} \\ \frac{dJ}{d\ell} &= c_{\sigma} \eta_{\text{sr}}(J_{\ell}) g_{\ell}, \end{aligned} \quad (\text{D14})$$

(we absorbed the dimensional part $a^{2-\sigma}$ into the definition of the constant c_{σ}). Together with Eq. (D3), those constitute the desired set of equations.

Appendix E: Derivation of Eq. (37)

Let us consider the flow equations (35) in the $y = 0$ plane. It follows that

$$g_{\ell} = g e^{(2-\sigma)\ell} e^{-\int \eta_{\text{sr}}(J_{\ell})d\ell}, \quad (\text{E1})$$

which is a reliable result as long as g_{ℓ} is small. Now let us consider a RG flow which starts very close to the critical temperature T_c : the corresponding trajectory will follow the separatrix up to the vicinity of the fixed point $g = 0, J = J_{\sigma}$. According to Eq. (36), this is given by the equation

$$g = \frac{\pi(2-\sigma)}{c_{\sigma}} [(J - J_{\sigma})^2 + k] \quad (\text{E2})$$

with $k \rightarrow 0^+$. We now consider a point in the flow ℓ^* such that $g(\ell^*)$ is small and $J(\ell^*) > J_{\sigma}$. Then

$$\begin{aligned} \int_0^{\ell^*} \eta_{\text{sr}}(J_{\ell}) d\ell &= \int_{J_0}^{\ell^*} \eta_{\text{sr}}(J) \frac{dJ}{J} \\ &= c_{\sigma}^{-1} \int_{J_0}^{\ell^*} \frac{dJ}{g(J)} \\ &= \pi(2-\sigma) \int_{J_0}^{J(\ell^*)} \frac{dJ}{(J - J_{\sigma})^2 + k} \end{aligned} \quad (\text{E3})$$

Let us consider now the physical bare parameter J_0 as a function of the temperature. This will cross the separatrix ($k \rightarrow 0^+$) for some $J_c < J_{\sigma}$ (which corresponds to T_c) so that $k \sim T_c - T$. In this case the second order singularity J_{σ} lies within the integration interval of Eq. (E3), so that the integral diverges as $k^{-1/2}$ as $k \rightarrow 0^+$. Then we have, as expected

$$g_{\ell^*} \sim e^{-B(T-T_c)^{-1/2}} \quad (\text{E4})$$

where B is a non universal constant.

Appendix F: Vortex gas representation of the action

We discuss in this Appendix the vortex gas representation of the action (30), starting from the decomposition of Eq. (42). At first let us notice how, from the definition of $\theta_0, \theta_{\text{top}}$ we get a local condition of the form

$$\nabla \times \nabla \theta_0 = 0 \quad \nabla \times \nabla \theta_{\text{top}} = 2\pi n(\mathbf{x}) \quad (\text{F1})$$

where $n(\mathbf{x}) = \sum_k m_k \delta(\mathbf{x} - \mathbf{x}_k)$ is the vortex density. Let us notice that $\nabla \theta_{\text{top}}$ is not irrotational because θ_{top} is not a single variable function. The curl of a vector field, in $d = 2$, is a scalar defined as $\nabla \times \mathbf{a} = \epsilon_{ij} \partial_i a_j$, where the Einstein summation convention has been used and ϵ_{ij} is the rank-two completely antisymmetric tensor. Exploiting the condition $\nabla \cdot \nabla \theta_{\text{top}} = 0$ we can write $\partial_i \theta_{\text{top}} = \epsilon_{ij} \partial_j \bar{\theta}$ where $\bar{\theta}$ is singled valued. It follows that

$$\nabla^2 \bar{\theta}(\mathbf{x}) = 2\pi n(\mathbf{x}) \quad (\text{F2})$$

that can be solved introducing the $d = 2$ Green function of the Laplacian $G_c(r) = -\frac{1}{2\pi} \ln r$, finding

$$\bar{\theta}(\mathbf{x}) = 2\pi \sum_k m_k G_c(|\mathbf{x} - \mathbf{x}_k|) \quad (\text{F3})$$

In terms of the decomposition (42) then, the kinetic term in action (30) decouples into the two terms

$$\begin{aligned} \frac{J}{2} |\nabla \theta|^2 &= \frac{J}{2} |\nabla \theta_0|^2 + \frac{J}{2} |\nabla \theta_{\text{top}}|^2 \\ &= \frac{J}{2} |\nabla \theta_0|^2 + \frac{J}{2} |\nabla \bar{\theta}|^2 \end{aligned} \quad (\text{F4})$$

Replacing the solution (F3) in the second term we get the usual Coulomb gas interaction, so that $S[\theta]$ can be written as Eq. (44).

Let us now integrate out the non-topological field perturbatively in g . At the first order in g the cumulant expansion gives the correction:

$$\langle S_{\text{LR}}[\theta_0 + \theta_{\text{top}}] \rangle_0 \quad (\text{F5})$$

where the average is computed on the quadratic non-topological term. By exploiting the form (31) of S_{LR} , and considering that only the even terms in θ_0 survives we have:

$$\begin{aligned} &\int d^2 \mathbf{x} \int_{r>a} d^2 \mathbf{r} \frac{\langle \cos(\Delta_{\mathbf{r}} \theta_0) \rangle_0}{r^{2+\sigma}} (1 - \cos \Delta_{\mathbf{r}} \theta_{\text{top}}) \\ &\sim \int d^2 \mathbf{x} \int_{r>a} d^2 \mathbf{r} \frac{1 - \cos \Delta_{\mathbf{r}} \theta_{\text{top}}(\mathbf{x})}{r^{2+\sigma+\eta_{\text{sr}}(J)}} \end{aligned} \quad (\text{F6})$$

up to immaterial additive constants. This has the same form of Eq. (31) so that, provided that $\sigma + \eta_{\text{sr}}(J) < 2$, it can be written as well as

$$\frac{g}{2} \int d^2 \mathbf{x} e^{-i\theta_{\text{top}} \nabla^{\sigma+\eta_{\text{sr}}(J)}} e^{i\theta_{\text{top}}} \quad (\text{F7})$$

where we absorbed some multiplicative proportionality factor into the coupling g . If $\sigma + \eta_{\text{sr}}(J) > 2$, then the dominant term is the Laplacian.

REFERENCES

- [1] A. Campa, T. Dauxois, D. Fanelli, and S. Ruffo, *Physics of Long-Range Interacting Systems* (Oxford Univ. Press, 2014).
- [2] N. Defenu, T. Donner, T. Macrì, G. Pagano, S. Ruffo, and A. Trombettoni, “Long-range interacting quantum systems,” arXiv preprint arXiv:2109.01063 (2021).
- [3] H. Häffner, C. Ross, and R. Blatt, “Quantum computing with trapped ions,” *Phys. Rep.* **469**, 155–203 (2008).
- [4] T. Lahaye, C. Menotti, L. Santos, M. Lewenstein, and T. Pfau, “The physics of dipolar bosonic quantum gases,” *Rep. Prog. Phys.* **72**, 126401 (2009).
- [5] M. Saffman, T. G. Walker, and K. Mølmer, “Quantum information with rydberg atoms,” *Rev. Mod. Phys.* **82**, 2313–2363 (2010).
- [6] H. Ritsch, P. Domokos, F. Brennecke, and T. Esslinger, “Cold atoms in cavity-generated dynamical optical potentials,” *Rev. Mod. Phys.* **85**, 553–601 (2013).
- [7] H. Bernien, S. Schwartz, A. Keesling, H. Levine, A. Omran, H. Pichler, S. Choi, A. S. Zibrov, M. Endres, M. Greiner, V. Vuletić, and M. D. Lukin, “Probing many-body dynamics on a 51-atom quantum simulator,” *Nature* **551**, 579–584 (2017).
- [8] C. Monroe, W. C. Campbell, L.-M. Duan, Z.-X. Gong, A. V. Gorshkov, P. W. Hess, R. Islam, K. Kim, N. M. Linke, G. Pagano, P. Richerme, C. Senko, and N. Y. Yao, “Programmable quantum simulations of spin systems with trapped ions,” *Rev. Mod. Phys.* **93**, 025001 (2021).
- [9] F. Mivehvar, F. Piazza, T. Donner, and H. Ritsch, “Cavity qed with quantum gases: new paradigms in many-body physics,” *Adv. Phys.* **70**, 1–153 (2021).
- [10] F. J. Dyson, “Existence of a phase-transition in a one-dimensional ising ferromagnet,” *Comm. Math. Phys.* **12**, 91–107 (1969).
- [11] D. J. Thouless, “Long-range order in one-dimensional ising systems,” *Phys. Rev.* **187**, 732–733 (1969).
- [12] J. M. Kosterlitz, “Phase transitions in long-range ferromagnetic chains,” *Phys. Rev. Lett.* **37**, 1577–1580 (1976).
- [13] J. Sak, “Recursion relations and fixed points for ferromagnets with long-range interactions,” *Phys. Rev. B* **8**, 281–285 (1973).
- [14] N. Defenu, A. Trombettoni, and A. Codello, “Fixed-point structure and effective fractional dimensionality for $o(n)$ models with long-range interactions,” *Phys. Rev. E* **92**, 052113 (2015).
- [15] J. Cardy, *Scaling and Renormalization in Statistical Physics* (Cambridge University Press, 1996).
- [16] E. Luijten, *Monte Carlo Simulation of Spin Models with Long-Range Interactions*, Ph.D. thesis, Technische Universiteit Delft (1997).
- [17] N. Defenu, A. Codello, S. Ruffo, and A. Trombettoni, “Criticality of spin systems with weak long-range interactions,” *Journal of Physics A Mathematical General* **53**, 143001 (2020).
- [18] A. Dutta and J. Bhattacharjee, “Phase transitions in the quantum ising and rotor models with a long-range interaction,” *Physical Review B* **64**, 184106 (2001).
- [19] N. Defenu, A. Trombettoni, and S. Ruffo, “Criticality and phase diagram of quantum long-range $o(n)$ models,” *Phys. Rev. B* **96**, 104432 (2017).
- [20] H. Kunz and C. E. Pfister, “First order phase transition in the plane rotator ferromagnetic model in two dimensions,” *Comm. Math. Phys.* **46**, 245–251 (1976).
- [21] G. Giachetti, N. Defenu, S. Ruffo, and A. Trombettoni, “Berezinskii-kosterlitz-thouless phase transitions with long-range couplings,” *Phys. Rev. Lett.* **127**, 156801 (2021).
- [22] G. Giachetti, N. Defenu, S. Ruffo, and A. Trombettoni, “Self-consistent harmonic approximation in presence of non-local couplings(a),” *EPL* **133**, 57004 (2021).
- [23] D. C. Mattis, “Transfer matrix in plane-rotator model,” *Phys. Lett. A* **104**, 357–360 (1984).
- [24] A. Filinov, N. V. Prokof’ev, and M. Bonitz, “Berezinskii-kosterlitz-thouless transition in two-dimensional dipole systems,” *Phys. Rev. Lett.* **105**, 070401 (2010).
- [25] M. F. Maghrebi, Z.-X. Gong, and A. V. Gorshkov, “Continuous symmetry breaking in 1d long-range interacting quantum systems,” *Phys. Rev. Lett.* **119**, 023001 (2017).

- [26] D. J. Bishop and J. D. Reppy, “Study of the superfluid transition in two-dimensional 4-He films,” *Phys. Rev. Lett.* **40**, 1727–1730 (1978).
- [27] J. Corson, R. Mallozzi, J. Orenstein, J. N. Eckstein, and I. Bozovic, “Vanishing of phase coherence in underdoped Bi_2Sr ,” *Nature (London)* **398**, 221–223 (1999).
- [28] I. Hetel, T. R. Lemberger, and M. Randeira, “Quantum critical behaviour in the superfluid density of strongly underdoped ultrathin copper oxide films,” *Nat. Phys.* **3**, 700 (2007).
- [29] L. S. Bilbro, R. V. Aguilar, G. Logve11, O. Pelleg, I. Božović, and N. P. Armitage, “Temporal correlations of superconductivity above the transition temperature in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ probed by terahertz spectroscopy,” *Nat. Phys.* **7**, 298 (2011).
- [30] J. Yong, M. J. Hinton, A. McCray, M. Randeria, M. Naamneh, A. Kanigel, and T. R. Lemberger, “Evidence of two-dimensional quantum critical behavior in the superfluid density of deeply underdoped $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+x}$,” *Phys. Rev. B* **85**, 180507 (2012).
- [31] P. G. Baity, X. Shi, Z. Shi, L. Benfatto, and D. Popović, “Effective two-dimensional thickness for the berezinskii-kosterlitz-thouless-like transition in a highly underdoped $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$,” *Phys. Rev. B* **93**, 024519 (2016).
- [32] W. H. Nitsche, N. Y. Kim, G. Roumpos, C. Schneider, M. Kamp, S. Höfling, A. Forchel, and Y. Yamamoto, “Algebraic order and the berezinskii-kosterlitz-thouless transition in an exciton-polariton gas,” *Phys. Rev. B* **90**, 205430 (2014).
- [33] Z. Hadzibabic, P. Krüger, M. Cheneau, B. Battelier, and J. Dalibard, “Berezinskii-kosterlitz-thouless crossover in a trapped atomic gas,” *Nature (London)* **441**, 1118 (2006).
- [34] P. A. Murthy, I. Boettcher, L. Bayha, M. Holzmann, D. Kedar, M. Neidig, M. G. Ries, A. N. Wenz, G. Zürn, and S. Jochim, “Observation of the berezinskii-kosterlitz-thouless phase transition in an ultracold fermi gas,” *Phys. Rev. Lett.* **115**, 010401 (2015).
- [35] N. Reyren, S. Thiel, A. D. Caviglia, L. F. Kourkoutis, G. Hammerl, C. Richter, C. W. Schneider, T. Kopp, A.-S. Rüetschi, D. Jaccard, M. Gabay, D. A. Muller, J.-M. Triscone, and J. Mannhart, “Electron scattering at dislocations in $\text{LaAlO}_3/\text{SrTiO}_3$ interfaces,” *Science* **317**, 1196–1199 (2007).
- [36] G. N. Daptary, S. Kumar, P. Kumar, A. Dogra, N. Mohanta, A. Taraphder, and A. Bid, “Correlated non-gaussian phase fluctuations in $\text{LaAlO}_3/\text{SrTiO}_3$ heterointerfaces,” *Phys. Rev. B* **94**, 085104 (2016).
- [37] A. M. R. V. L. Monteiro, D. J. Groenendijk, I. Groen, J. de Bruijckere, R. Gaudenzi, H. S. J. van der Zant, and A. D. Caviglia, “Two-dimensional superconductivity at the $(111)\text{LaAlO}_3/\text{SrTiO}_3$ interface,” *Phys. Rev. B* **96**, 020504(R) (2017).
- [38] V. Pokrovsky and G. Uimin, “Magnetic properties of two-dimensional and layered systems,” *Phys. Lett. A* **45**, 467–468 (1973).
- [39] A. S. T. Pires, “A quantum self-consistent harmonic approximation,” *Solid State Commun.* **104**, 771–773 (1997).
- [40] J. M. Kosterlitz, “Topological defects and phase transitions,” *Rev. Mod. Phys.* **89**, 040501 (2017).
- [41] V. L. Berezinskii, “Destruction of long-range order in one-dimensional and two-dimensional systems possessing a continuous symmetry group. ii. quantum systems,” *Sov. Phys. JETP* **34**, 610 (1972).
- [42] J. M. Kosterlitz and D. J. Thouless, “Ordering, metastability and phase transitions in two-dimensional systems,” *J. Phys. C* **6**, 1181–1203 (1973).
- [43] J. M. Kosterlitz, “The critical properties of the two-dimensional xy model,” *J. Phys. C* **7**, 1046 (1974).
- [44] J. V. José, ed., *40 Years Of Berezinskii-kosterlitz-thouless Theory* (World Scientific, Singapore, 2013).
- [45] R. Savit, “Duality in field theory and statistical systems,” *Rev. Mod. Phys.* **52**, 453–487 (1980).
- [46] P. Minnhagen, “The two-dimensional coulomb gas, vortex unbinding, and superfluid-superconducting films,” *Rev. Mod. Phys.* **59**, 1001–1066 (1987).
- [47] Z. Gulacsi and M. Gulacsi, “Theory of phase transitions in two-dimensional systems,” *Adv. Phys.* **47**, 1–89 (1998).
- [48] Amit, Y. Y. Goldschmidt, and S. Grinstein, “Renormalisation group analysis of the phase transition in the 2d coulomb gas, sine-gordon theory and xy-model,” *J. Phys. A: Math. Gen.* **13**, 585–620 (1980).
- [49] J. Villain, “Theory of one-dimensional and two-dimensional magnets with an easy magnetization plane. 2. The Planar, classical, two-dimensional magnet,” *J. Phys. France* **36**, 581–590 (1975).
- [50] J. Villain, “Theory of one- and two-dimensional magnets with an easy magnetization plane. the planar, two-dimensional magnet,” *J. Phys. France* **36**, 581–590 (1975).
- [51] J. V. José, L. P. Kadanoff, S. Kirkpatrick, and D. R. Nelson, *Phys. Rev. B* **16**, 1217–1241 (1977).
- [52] P. W. Anderson and G. Yuval, “Exact results in the kondo problem: Equivalence to a classical one-dimensional coulomb gas,” *Phys. Rev. Lett.* **23**, 89–92 (1969).
- [53] D. J. Thouless, “Long-range order in one-dimensional ising systems,” *Phys. Rev.* **187**, 732–733 (1969).
- [54] F. J. Dyson, “An Ising ferromagnet with discontinuous long-range order,” *Communications in Mathematical Physics* **21**, 269–283 (1971).
- [55] J. L. Cardy, “One-dimensional models with $1/r^2$ interactions,” *Journal of Physics A: Mathematical and General* **14**, 1407–1415 (1981).
- [56] M. Aizenman, J. T. Chayes, L. Chayes, and C. M. Newman, “Discontinuity of the magnetization in one-dimensional $1/|x-y|^2$ Ising and Potts models,” *Journal of Statistical Physics* **50**, 1–40 (1988).
- [57] M. Barma, S. N. Majumdar, and D. Mukamel, “Fluctuation-dominated phase ordering at a mixed order transition,” *Journal of Physics A: Mathematical and Theoretical* **52**, 254001 (2019).
- [58] H. Spohn and W. Zwerger, “Decay of the two-point function in one-dimensional $o(n)$ spin models with long-range interactions,” *J. Stat. Phys.* **94**, 1037–1043 (1999).
- [59] A. Kargol, “Decay of correlations in multicomponent ferromagnets with long-range interaction,” *Rep. Math. Phys.* **56**, 379–386 (2005).
- [60] A. Kargol, “Decay of correlations in n -vector ferromagnetic quantum models with long-range interactions,” *J. Stat. Mech.: Theory Exp.* **2014**, 10006 (2014).
- [61] T. Botzung, D. Hagenmüller, G. Masella, J. Dubail, N. Defenu, A. Trombettoni, and G. Pupillo, “Effects of energy extensivity on the quantum phases of long-range interacting systems,” *Phys. Rev. B* **103**, 155139 (2021).

- [62] N. Defenu, A. Trombettoni, and S. Ruffo, “Anisotropic long-range spin systems,” *Phys. Rev. B* **94**, 224411 (2016).
- [63] F. Haldane, “Continuum dynamics of the 1-d heisenberg antiferromagnet: Identification with the $o(3)$ nonlinear sigma model,” *Physics Letters A* **93**, 464–468 (1983).
- [64] F. D. M. Haldane, “Nonlinear field theory of large-spin heisenberg antiferromagnets: Semiclassically quantized solitons of the one-dimensional easy-axis néel state,” *Phys. Rev. Lett.* **50**, 1153–1156 (1983).
- [65] I. Affleck and E. H. Lieb, “A proof of part of Haldane’s conjecture on spin chains,” *Letters in Mathematical Physics* **12**, 57–69 (1986).
- [66] W. Chen, K. Hida, and B. C. Sanctuary, “Ground-state phase diagram of $s = 1$ XXZ chains with uniaxial single-ion-type anisotropy,” *Phys. Rev. B* **67**, 104401 (2003).
- [67] A. Cuccoli, R. Giachetti, V. Tognetti, R. Vaia, and P. Verrucchi, “The effective potential and effective hamiltonian in quantum statistical mechanics,” *Journal of Physics: Condensed Matter* **7**, 7891–7938 (1995).
- [68] A. Cuccoli, G. Gori, R. Vaia, and P. Verrucchi, “Phase diagram of the two-dimensional quantum antiferromagnet in a magnetic field,” *Journal of Applied Physics* **99**, 08H503 (2006).
- [69] M. I. Berganza and L. Leuzzi, “Critical behavior of the x y model in complex topologies,” *Physical Review B* **88**, 144104 (2013).
- [70] F. Cescatti, M. Ibáñez-Berganza, A. Vezzani, and R. Burioni, “Analysis of the low-temperature phase in the two-dimensional long-range diluted xy model,” *Phys. Rev. B* **100**, 054203 (2019).
- [71] A. Raju, C. B. Clement, L. X. Hayden, J. P. Kent-Dobias, D. B. Liarte, D. Z. Rocklin, and J. P. Sethna, “Normal form for renormalization groups,” *Phys. Rev. X* **9**, 021014 (2019).
- [72] G. Bighin, N. Defenu, I. Nándori, L. Salasnich, and A. Trombettoni, “Berezinskii-kosterlitz-thouless paired phase in coupled xy models,” *Phys. Rev. Lett.* **123**, 100601 (2019).
- [73] P. G. Maier and F. Schwabl, “Ferromagnetic ordering in the two-dimensional dipolar xy model,” *Phys. Rev. B* **70**, 134430 (2004).
- [74] A. Vasilyev, A. Tarkhov, L. Men’shikov, P. Fedichev, and U. Fischer, “Universality of the berezinskii-kosterlitz-thouless type of phase transition in the dipolar xy -model,” *New J. Phys.* **16**, 053011 (2014).
- [75] L. Jacobs and R. Savit, “Absence of long-range order above two dimensions,” *Ann. N. Y. Acad. Sci.* **410**, 281–290 (1983).
- [76] N. Defenu, A. Trombettoni, and D. Zappalà, “Topological phase transitions in four dimensions,” *Nucl. Phys. B* **964**, 115295 (2021).
- [77] S. R. Shenoy and B. Chattopadhyay, “Anisotropic three-dimensional xy model and vortex-loop scaling,” *Phys. Rev. B* **51**, 9129–9147 (1995).