Heat Kernels and Hardy Spaces on Non-Tangentially Accessible Domains with Applications to Global Regularity of Inhomogeneous Dirichlet Problems

Sibei Yang and Dachun Yang*

Abstract. Let $n \ge 2$ and Ω be a bounded non-tangentially accessible domain (for short, NTA domain) of \mathbb{R}^n . Assume that L_D is a second-order divergence form elliptic operator having realvalued, bounded, measurable coefficients on $L^2(\Omega)$ with the Dirichlet boundary condition. The main aim of this article is threefold. First, the authors prove that the heat kernels $\{K_t^{L_D}\}_{t>0}$ generated by L_D are Hölder continuous. Second, for any $p \in (0, 1]$, the authors introduce the 'geometrical' Hardy space $H^p_r(\Omega)$ by restricting any element of the Hardy space $H^p(\mathbb{R}^n)$ to Ω , and show that, when $p \in (\frac{n}{n+\delta_0}, 1]$, $H_r^p(\Omega) = H^p(\Omega) = H_{L_0}^p(\Omega)$ with equivalent quasi-norms, where $H^p(\Omega)$ and $H_{L_D}^p(\Omega)$ respectively denote the Hardy space on Ω and the Hardy space associated with L_D , and $\delta_0 \in (0, 1]$ is the critical index of the Hölder continuity for the kernels $\{K_t^{L_D}\}_{t>0}$. Third, as applications, the authors obtain the global gradient estimates in both $L^p(\Omega)$, with $p \in (1, p_0)$, and $H_z^p(\Omega)$, with $p \in (\frac{n}{n+1}, 1]$, for the inhomogeneous Dirichlet problem of second-order divergence form elliptic equations on bounded NTA domains, where $p_0 \in (2, \infty)$ is a constant depending only on n, Ω , and the coefficient matrix of L_D . Here, the 'geometrical' Hardy space $H_z^p(\Omega)$ is defined by restricting any element of the Hardy space $H^p(\mathbb{R}^n)$ supported in Ω to Ω , where Ω denotes the closure of Ω in \mathbb{R}^n . It is worth pointing out that the range $p \in (1, p_0)$ for the global gradient estimate in the scale of Lebesgue spaces $L^{p}(\Omega)$ is sharp and the above results are established without any additional assumptions on both the coefficient matrix of L_D , and the domain Ω.

1 Introduction

The study of elliptic value problems on non-smooth domains of \mathbb{R}^n has a long history (see, for instance, [21, 39, 43] and the references therein). In recent years, the research of the global regularity for elliptic equations with rough coefficients on non-smooth domains of \mathbb{R}^n has aroused great interest (see, for instance, [11, 21, 22, 23, 24, 25, 30, 57, 59]). The global regularity estimates of elliptic equations with rough coefficients on the non-smooth domain Ω of \mathbb{R}^n in the scale of Lebesgue spaces $L^p(\Omega)$, with $p \in (1, \infty)$, have been extensively studied in the existing literatures (see, for instance, the recent survey article [21], the monograph [57], and the references therein).

²⁰²⁰ Mathematics Subject Classification. Primary 42B35; Secondary 35J05, 35J25, 47B06.

Key words and phrases. NTA domain, heat kernel, Hardy space, divergence form elliptic operator, global gradient estimate, Dirichlet problem.

This project is partially supported by the National Natural Science Foundation of China (Grant Nos. 11871254, 12071431, 11971058 and 12071197), the National Key Research and Development Program of China (Grant No. 2020YFA0712900) and the Fundamental Research Funds for the Central Universities (Grant No. lzujbky-2021-ey18).

^{*}Corresponding author, E-mail: dcyang@bnu.edu.cn/January 10, 2022/Final version.

However, there exist very few literatures on global regularity estimates of elliptic equations with rough coefficients on the non-smooth domain Ω of \mathbb{R}^n in the scale of Hardy spaces $H^p(\Omega)$ with $p \in (0, 1]$.

Let $n \ge 2$ and Ω be a bounded non-tangentially accessible domain (for short, NTA domain) of \mathbb{R}^n . Assume that L_D is a second-order divergence form elliptic operator having real-valued, bounded, measurable coefficients on $L^2(\Omega)$ with the Dirichlet boundary condition. The main aim of this article is threefold. First, we prove that the heat kernels $\{K_t^{L_D}\}_{t>0}$ generated by L_D are Hölder continuous. Second, for any $p \in (0, 1]$, we introduce the 'geometrical' Hardy space $H_r^p(\Omega)$ by restricting any element of the Hardy space $H^p(\mathbb{R}^n)$ to Ω , and show that, when $p \in (\frac{n}{n+\delta_0}, 1]$, $H_r^p(\Omega) = H^p(\Omega) = H_{L_D}^p(\Omega)$ with equivalent quasi-norms, where $H^p(\Omega)$ and $H_{L_D}^p(\Omega)$ respectively denote the Hardy space on Ω and the Hardy space associated with L_D , and $\delta_0 \in (0, 1]$ is the critical index of the Hölder continuity for the kernels $\{K_t^{L_D}\}_{t>0}$. Third, as applications, for the inhomogeneous Dirichlet boundary value problem

(1.1)
$$\begin{cases} -\operatorname{div} (A\nabla u) = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases}$$

where the matrix A is real-valued, bounded, and measurable, and satisfies the uniform ellipticity condition [see (1.3) below for the details], and $\partial\Omega$ denotes the boundary of Ω , we obtain the global gradient estimates of the weak solution u in both Lebesgue spaces $L^p(\Omega)$, with $p \in (1, p_0)$, and Hardy spaces $H_z^p(\Omega)$, with $p \in (\frac{n}{n+1}, 1]$, where $p_0 \in (2, \infty)$ is a constant depending only on n, Ω , and the coefficient matrix A. Here, the 'geometrical' Hardy space $H_z^p(\Omega)$ is defined by restricting any element of the Hardy space $H^p(\mathbb{R}^n)$ supported in $\overline{\Omega}$ to Ω , where $\overline{\Omega}$ denotes the closure of Ω in \mathbb{R}^n . Meanwhile, it is worth pointing out that the range $p \in (1, p_0)$ of p for the global gradient estimate in the scale of the Lebesgue space $L^p(\Omega)$ is sharp [see Remark 1.10(i) below for the details].

Compared with the global regularity estimate of elliptic equations on the non-smooth domain Ω of \mathbb{R}^n in Lebesgue spaces $L^p(\Omega)$ established in [1, 11, 20, 22, 59], we obtain the global regularity estimate for the Dirichlet problem (1.1) without any additional assumptions on both the coefficient matrix A and the domain Ω . Recall that the global gradient estimate in $L^p(\Omega)$ with any given $p \in (1, \infty)$ for the Dirichlet problem (1.1), with f replaced by div (f), was established by Di Fazio [20], under the assumptions that $A \in \text{VMO}(\mathbb{R}^n; \mathbb{R}^{n^2})$ (see, for instance, [56]) and $\partial \Omega \in C^{1,1}$, which was weakened to $\partial \Omega \in C^1$ by Auscher and Qafsaoui [1]. Moreover, the global gradient estimate in $L^p(\Omega)$ with any given $p \in (1, \infty)$ for the problem (1.1), with f replaced by div (f), was obtained by Byun and Wang [11], under the assumptions that A satisfies the (δ , R)-BMO condition (see, for instance, [11] or Definition 2.6 below for its definition) for sufficiently small $\delta \in (0, \infty)$, and that Ω is a bounded Reifenberg flat domain of \mathbb{R}^n (see, for instance, [53, 64] or Remark 2.5(i) below for its definition). Furthermore, for the Dirichlet problem (1.1) with f replaced by div (f), the global gradient estimate in $L^p(\Omega)$ with any given $p \in (1, \infty)$ was established by Dong and Kim [22, 23], under the assumptions that A has partial sufficiently small BMO coefficients and that $\Omega \subset \mathbb{R}^n$ is a bounded Lipschitz domain with small Lipschitz constant, or a bounded Reifenberg flat domain. Meanwhile, for the problem (1.1) with f replaced by div (f), the global gradient estimate in $L^{p}(\Omega)$, with any given $p \in (\frac{3}{2} - \varepsilon, 3 + \varepsilon)$ when $n \ge 3$, or $p \in (\frac{4}{3} - \varepsilon, 4 + \varepsilon)$ when n = 2, was obtained by Shen [59], under the assumptions that $A \in \text{VMO}(\mathbb{R}^n; \mathbb{R}^{n^2})$ and that $\Omega \subset \mathbb{R}^n$ is a bounded Lipschitz domain, where $\varepsilon \in (0, \infty)$ is a constant depending only on *n* and Ω .

Moreover, NTA domains considered in this article were originally introduced by Jerison and Kenig [38] when studying the boundary behavior of harmonic functions. We point out that NTA domains have a wide generality and contain Lipschitz domains, BMO₁ domains, Zygmund domains, quasi-spheres, and some Reifenberg flat domains as special examples (see, for instance, [38, 44, 64]). Furthermore, NTA domains are closely related to the theory of quasi-conformal mappings (see, for instance, [38, 41] and the references therein).

To describe the main results of this article, we first recall some necessary notions. Let Ω be a bounded NTA domain of \mathbb{R}^n as in Definition 2.1 below (see also [38]) and $p \in (0, \infty)$. Recall that the *Lebesgue space* $L^p(\Omega)$ is defined by setting

$$L^{p}(\Omega) := \left\{ f \text{ is measurable on } \Omega : ||f||_{L^{p}(\Omega)} := \left[\int_{\Omega} |f(x)|^{p} dx \right]^{\frac{1}{p}} < \infty \right\}.$$

Moreover, for any given $m \in \mathbb{N}$, let

(1.2)
$$L^{p}(\Omega; \mathbb{R}^{m}) := \{ \mathbf{f} := (f_{1}, \dots, f_{m}) : \text{ for any } i \in \{1, \dots, m\}, f_{i} \in L^{p}(\Omega) \}$$

and

$$\|\mathbf{f}\|_{L^p(\Omega;\mathbb{R}^m)} := \sum_{i=1}^m \|f_i\|_{L^p(\Omega)}.$$

Denote by $W^{1,p}(\Omega)$ the Sobolev space on Ω equipped with the norm

$$||f||_{W^{1,p}(\Omega)} := ||f||_{L^{p}(\Omega)} + ||\nabla f||_{L^{p}(\Omega;\mathbb{R}^{n})},$$

where ∇f is the *distributional gradient* of f. Furthermore, $W_0^{1,p}(\Omega)$ is defined to be the *closure* of $C_c^{\infty}(\Omega)$ in $W^{1,p}(\Omega)$, where $C_c^{\infty}(\Omega)$ denotes the set of all *infinitely differentiable functions on* Ω *with compact support contained in* Ω .

For any given $x \in \Omega$, let $A(x) := \{a_{ij}(x)\}_{i,j=1}^n$ denote an $n \times n$ matrix with real-valued, bounded, and measurable entries. Then *A* is said to satisfy the *uniform ellipticity condition* if there exists a positive constant $\mu_0 \in (0, 1]$ such that, for any $x \in \Omega$ and $\xi := (\xi_1, \ldots, \xi_n) \in \mathbb{R}^n$,

(1.3)
$$\mu_0 |\xi|^2 \le \sum_{i,j=1}^n a_{ij}(x) \xi_i \xi_j \le \mu_0^{-1} |\xi|^2.$$

Denote by L_D the maximal-accretive operator (see, for instance, [51, p. 23, Definition 1.46] for the definition) on $L^2(\Omega)$ with the largest domain $\mathcal{D}(L_D) \subset W_0^{1,2}(\Omega)$ such that, for any $f \in \mathcal{D}(L_D)$ and $g \in W_0^{1,2}(\Omega)$,

$$(L_D f, g) = \int_{\Omega} A(x) \nabla f(x) \cdot \nabla g(x) \, dx,$$

where (\cdot, \cdot) denotes the *interior product* in $L^2(\Omega)$. In this sense, for any $f \in \mathcal{D}(L_D)$, we write

(1.4)
$$L_D f := -\operatorname{div}(A\nabla f).$$

Let $\{K_t^{L_D}\}_{t>0}$ be the kernels of the semigroup $\{e^{-tL_D}\}_{t>0}$. By [19, Corollary 3.2.8] (see also [3]), we find that there exist positive constants *C* and *c* such that, for any $t \in (0, \infty)$ and $x, y \in \Omega$,

(1.5)
$$\left| K_t^{L_D}(x, y) \right| \le \frac{C}{t^{n/2}} \exp\left\{ -\frac{|x - y|^2}{ct} \right\}.$$

Furthermore, it is worth pointing out that the upper and the lower bound estimates, and the Hölder continuity of the heat kernels play a key roles in the study of the well-posedness of some parabolic partial differential equations, real-variable characterizations of some function spaces, and some Sobolev-type inequalities (see, for instance, [19, 33, 55]).

Now, we state the main results of this article as follows; see Definitions 2.6 and 2.4 below, respectively, for the definitions of both the (γ, R) -BMO condition and the (γ, σ, R) quasi-convex domain.

Theorem 1.1. Let $n \ge 2$, Ω be a bounded NTA domain of \mathbb{R}^n , the real-valued, bounded, and measurable matrix A satisfy (1.3), and L_D be as in (1.4). Denote by $\{K_t^{L_D}\}_{t>0}$ the heat kernels generated by L_D .

(i) Then there exists a constant δ₀ ∈ (0, 1], depending only on n, A, and Ω, such that, for any given δ ∈ (0, δ₀), there exist constants C, c ∈ (0,∞) such that, for any t ∈ (0,∞) and x, y₁, y₂ ∈ Ω with |y₁ - y₂| ≤ √t/2,

(1.6)
$$\left|K_t^{L_D}(x, y_1) - K_t^{L_D}(x, y_2)\right| \le \frac{C}{t^{n/2}} \left[\frac{|y_1 - y_2|}{\sqrt{t}}\right]^{\delta} \exp\left\{-\frac{|x - y_1|^2}{ct}\right\}.$$

(ii) For any given δ₀ ∈ (0, 1], there exists a constant γ₀ ∈ (0, ∞), depending only on δ₀, n, and Ω, such that, if A satisfies the (γ, R)-BMO condition and Ω is a (γ, σ, R) quasi-convex domain for some γ ∈ (0, γ₀), σ ∈ (0, 1), and R ∈ (0, ∞), then, for any given δ ∈ (0, δ₀), there exist constants C, c ∈ (0, ∞) such that, for any t ∈ (0, ∞) and x, y₁, y₂ ∈ Ω with |y₁ − y₂| ≤ √t/2, (1.6) holds true.

Remark 1.2. We point out that, when Ω is a bounded Lipschitz domain of \mathbb{R}^n , the conclusion of Theorem 1.1(i) is well known (see, for instance, [3]). Moreover, when Ω is a bounded semi-convex domain of \mathbb{R}^n (see, for instance, [48, 49] or Remark 2.5(ii) below for the details), and $L_D := -\Delta_D$ with Δ_D being the Laplace operator with the Dirichlet boundary condition on Ω , Theorem 1.1(ii) was obtained in [26, Lemma 2.7]. Recall that the bounded semi-convex domain Ω is a (γ, σ, R) quasi-convex domain for any $\gamma \in (0, 1)$, some $\sigma \in (0, 1)$, and some $R \in (0, \infty)$ (see, for instance, [69]). Thus, Theorem 1.1(ii) essentially improves [26, Lemma 2.7] by weakening the assumptions on both the matrix A and the domain Ω .

When $n \ge 3$, we prove Theorem 1.1(i) by using an upper estimate for the Green function associated with L_D in terms of distance functions (see, for instance, [35, Remark 4.9]), the Harnack inequality (see, for instance, [31, Theorem 8.22]), and the functional calculus associated with L_D . Precisely, using the upper estimate for the Green function of L_D in terms of distance functions, and the Harnack inequality, and borrowing some ideas from the proof of Grüter and Widman [32,

Theorem (1.9)], we prove the Hölder continuity of the Green function associated with L_D . Moreover, applying the Hölder continuity of the Green function, and the functional calculus associated with L_D , and borrowing some ideas from the proofs of Duong et al. [26, Lemmas 2.6 and 2.7], we prove Theorem 1.1(i) in the case of $n \ge 3$. When n = 2, we show Theorem 1.1(i) via establishing the global gradient estimate for the Dirichlet problem (1.1) in $L^p(\Omega)$ with some $p \in (2, \infty)$ (see Lemma 3.6 below), and using the Sobolev embedding theorem. Furthermore, we prove Theorem 1.1(ii) by establishing the global gradient estimate for the Dirichlet problem (1.1) in $L^p(\Omega)$ for sufficiently large $p \in (n, \infty)$ (see Lemma 3.9 below), and applying the Sobolev embedding theorem.

Next, we recall the definitions of the Hardy space $H^p(\mathbb{R}^n)$, the 'geometrical' Hardy spaces $H^p_z(\Omega)$ and $H^p(\Omega)$, the Hardy space $H^p(\Omega)$, and the Hardy space $H^p_{L_D}(\Omega)$ associated with L_D .

Denote by $S(\mathbb{R}^n)$ the space of all Schwartz functions equipped with the well-known topology determined by a countable family of norms, and by $S'(\mathbb{R}^n)$ its dual space (namely, the space of all tempered distributions) equipped with the weak-* topology. Let $\mathcal{D}(\Omega)$ denote the space of all infinitely differentiable functions with compact support in Ω equipped with the inductive topology, and $\mathcal{D}'(\Omega)$ its topological dual equipped with the weak-* topology, which is called the space of distributions on Ω .

In what follows, for any $x \in \mathbb{R}^n$ and $r \in (0, \infty)$, we always let $B(x, r) := \{y \in \mathbb{R}^n : |y - x| < r\}$.

Definition 1.3. Let $p \in (0, 1]$ and Ω be a domain of \mathbb{R}^n .

(i) The *Hardy space* $H^p(\mathbb{R}^n)$ is defined to be the set of all $f \in \mathcal{S}'(\mathbb{R}^n)$ such that $f^+ \in L^p(\mathbb{R}^n)$ equipped with the *quasi-norm* $||f||_{H^p(\mathbb{R}^n)} := ||f^+||_{L^p(\mathbb{R}^n)}$, where the *radial maximal function* f^+ of f is defined by setting, for any $x \in \mathbb{R}^n$,

$$f^+(x) := \sup_{t \in (0,\infty)} \left| e^{-t\Delta}(f)(x) \right|.$$

Here, $\{e^{-t\Delta}\}_{t>0}$ denotes the heat semigroup generated by the *Laplace operator* Δ on \mathbb{R}^n . (ii) The *Hardy space* $H_{\tau}^p(\Omega)$ is defined by setting

$$H_z^p(\Omega) := \left\{ f \in H^p(\mathbb{R}^n) : f = 0 \text{ on } \overline{\Omega}^{\mathbb{C}} \right\} / \{ f \in H^p(\mathbb{R}^n) : f = 0 \text{ on } \Omega \}.$$

Here and thereafter, $\overline{\Omega}$ and $\overline{\Omega}^{\mathbb{C}}$ denote, respectively, the *closure* of Ω in \mathbb{R}^n , and the *complementary set* of $\overline{\Omega}$ in \mathbb{R}^n . Moreover, the *quasi-norm* of the element in $H_z^p(\Omega)$ is defined to be the quotient norm, namely, for any $f \in H_z^p(\Omega)$,

$$||f||_{H^p_z(\Omega)} := \inf \left\{ ||F||_{H^p(\mathbb{R}^n)} : F \in H^p(\mathbb{R}^n), F = 0 \text{ on } \overline{\Omega}^{\mathbb{C}}, \text{ and } F|_{\Omega} = f \right\},\$$

where the infimum is taken over all $F \in H^p(\mathbb{R}^n)$ satisfying F = 0 on $\overline{\Omega}^{\mathbb{C}}$, and F = f on Ω .

(iii) A distribution $f \in \mathcal{D}'(\Omega)$ is said to belong to the *Hardy space* $H_r^p(\Omega)$ if f is the restriction to Ω of a distribution F in $H^p(\mathbb{R}^n)$, namely,

$$H^p_r(\Omega) := \{ f \in \mathcal{D}'(\Omega) : \text{ there exists an } F \in H^p(\mathbb{R}^n) \text{ such that } F|_{\Omega} = f \}$$

$$= H^p(\mathbb{R}^n) / \{ F \in H^p(\mathbb{R}^n) : F = 0 \text{ on } \Omega \}.$$

Moreover, for any $f \in H^p_r(\Omega)$, the quasi-norm $||f||_{H^p_r(\Omega)}$ of f in $H^p_r(\Omega)$ is defined by setting

$$||f||_{H^{p}(\Omega)} := \inf \{ ||F||_{H^{p}(\mathbb{R}^{n})} : F \in H^{p}(\mathbb{R}^{n}) \text{ and } F|_{\Omega} = f \},\$$

where the infimum is taken over all $F \in H^p(\mathbb{R}^n)$ satisfying F = f on Ω .

(iv) Let $\phi \in C_{c}^{\infty}(\mathbb{R}^{n})$ be a nonnegative function such that

$$\operatorname{supp}(\phi) := \{ x \in \mathbb{R}^n : \phi(x) \neq 0 \} \subset B(\mathbf{0}, 1)$$

and $\int_{\mathbb{R}^n} \phi(x) dx = 1$, here and thereafter, **0** denotes the *origin* of \mathbb{R}^n . For any $f \in \mathcal{D}'(\Omega)$, the *radial maximal function* f_{Ω}^+ is defined by setting, for any $x \in \Omega$,

$$f_{\Omega}^+(x) := \sup_{t \in (0,\delta(x)/2)} |\phi_t * f(x)|,$$

where, for any $x \in \Omega$, $\delta(x) := \text{dist}(x, \Omega^{\complement})$ and, for any $t \in (0, \infty)$, $\phi_t(\cdot) := \frac{1}{t^n} \phi(\frac{\cdot}{t})$. Then the *Hardy space* $H^p(\Omega)$ is defined by setting

$$H^{p}(\Omega) := \left\{ f \in \mathcal{D}'(\Omega) : \|f\|_{H^{p}(\Omega)} := \left\|f_{\Omega}^{+}\right\|_{L^{p}(\Omega)} < \infty \right\}.$$

Let $p \in (0, 1]$. From the definitions of both $H_r^p(\Omega)$ and $H_z^p(\Omega)$, it follows that $H_z^p(\Omega) \subset H_r^p(\Omega)$. Moreover, by [15], we find that $H_z^p(\Omega) \subsetneq H_r^p(\Omega)$.

Definition 1.4. Let $n \ge 2$, Ω be a bounded NTA domain of \mathbb{R}^n , $p \in (0, 1]$, and L_D be as in (1.4). For any $f \in L^2(\Omega)$, the *Lusin area function*, $S_{L_D}(f)$, associated with L_D , is defined by setting, for any $x \in \Omega$,

$$S_{L_D}(f)(x) := \left[\int_{\Gamma(x)} \left| t^2 L_D e^{-t^2 L_D}(f)(y) \right|^2 \frac{dy \, dt}{|B_\Omega(x,t)|t} \right]^{1/2}$$

where

$$\Gamma(x) := \{ (y,t) \in \Omega \times (0,\infty) : |x-y| < t \}$$

and $B_{\Omega}(x,t) := B(x,t) \cap \Omega$.

A function $f \in L^2(\Omega)$ is said to be in the set $\mathbb{H}^p_{L_D}(\Omega)$ if $S_{L_D}(f) \in L^p(\Omega)$; moreover, for any $f \in \mathbb{H}^p_{L_D}(\Omega)$, define $||f||_{H^p_{L_D}(\Omega)} := ||S_{L_D}(f)||_{L^p(\Omega)}$. Then the Hardy space $H^p_{L_D}(\Omega)$ is defined to be the completion of $\mathbb{H}^p_{L_D}(\Omega)$ with respect to the quasi-norm $|| \cdot ||_{H^p_{L_D}(\Omega)}$.

It is well known that the real-variable theory of Hardy spaces on \mathbb{R}^n , initiated by Stein and Weiss [62] and then systematically developed by Fefferman and Stein [29], plays important roles in various fields of analysis and partial differential equations. In recent years, the study on the real-variable theory of Hardy spaces on \mathbb{R}^n or its domains, associated with different differential operators, has aroused great interests (see, for instance, [10, 28, 36, 37, 60] for Hardy spaces on \mathbb{R}^n , and [2, 7, 9, 12, 26, 46, 65, 66, 68] for Hardy spaces on domains). Moreover, the Hardy space $H^p(\Omega)$ on the domain Ω of \mathbb{R}^n was introduced and studied by Miyachi [50]. Furthermore, when Ω is a Lipschitz domain of \mathbb{R}^n , the 'geometrical' Hardy spaces $H^p_r(\Omega)$ and $H^p_z(\Omega)$ on domains were

introduced by Chang et al. [15, 16] which naturally appear in the study of the regularity of the Dirichlet and the Neumann boundary value problems of second-order elliptic equations (see, for instance, [2, 13, 15, 26]).

Then we have the following equivalence relation between $H_r^p(\Omega)$, $H^p(\Omega)$, and $H_{L_p}^p(\Omega)$.

Theorem 1.5. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, and $\delta_0 \in (0, 1]$ as in Theorem 1.1. Then, for any given $p \in (\frac{n}{n+1}, 1]$, $H_r^p(\Omega) = H^p(\Omega)$ with equivalent quasi-norms. Moreover, for any given $p \in (\frac{n}{n+\delta_0}, 1]$, $H_{L_D}^p(\Omega) = H_r^p(\Omega) = H^p(\Omega)$ with equivalent quasi-norms.

Remark 1.6. Let $\delta_0 \in (0, 1]$ be as in Theorem 1.1. When Ω is a bounded Lipschitz domain of \mathbb{R}^n , the equivalence of the Hardy spaces $H_{L_D}^1(\Omega)$ and $H_r^1(\Omega)$ in Theorem 1.5 was obtained by Auscher and Russ [2, Theorem 1 and Proposition 5]. Moreover, when Ω is a bounded Lipschitz domain of \mathbb{R}^n , the operator L_D is non-negative and adjoint, and $p \in (\frac{n}{n+\delta_0}, 1]$, the equivalence of the spaces $H_{L_D}^p(\Omega)$, $H_r^p(\Omega)$, and $H^p(\Omega)$ in Theorem 1.5 was established by Bui et al. [9, Theorem 4.4 and Remark 4.5(c)] (see also [68, Corollary 4.5]). Thus, Theorem 1.5 improves the results of Auscher and Russ [2, Theorem 1 and Proposition 5] and Bui et al. [9, Theorem 4.4 and Remark 4.5(c)] by weakening the assumptions on both the domain Ω and the operator L_D .

Furthermore, when Ω is a bounded Lipschitz domain, the equivalence of $H_r^p(\Omega)$ and $H^p(\Omega)$ for any given $p \in (\frac{n}{n+1}, 1]$ was essentially proved by Chang et al. [15, Theorem 2.7]. Recall that NTA domains contain Lipschitz domains as special examples (see, for instance, [38, 64] or Remark 2.5(iii) below). Therefore, the equivalence of $H_r^p(\Omega)$ and $H^p(\Omega)$ for any given $p \in (\frac{n}{n+1}, 1]$ obtained in Theorem 1.5 improves [15, Theorem 2.7] via weakening the assumption on the domain Ω under consideration.

By subtly using some geometrical properties of NTA domains, obtained in Lemma 2.2 below, the reflection technology related to NTA domains, the atomic characterizations of both $H^p(\mathbb{R}^n)$ and $H^p(\Omega)$, and the Hölder continuity of the heat kernels $\{K_t^{L_D}\}_{t>0}$ given in Theorem 1.1, we show Theorem 1.5.

Let $n \ge 2$, $p \in (1, \infty)$, and $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain. Assume that

(1.7)
$$p_* := \begin{cases} \frac{np}{n+p} & \text{when } p \in \left(\frac{n}{n-1}, \infty\right), \\ 1+\epsilon & \text{when } p \in \left(1, \frac{n}{n-1}\right], \end{cases}$$

where $\epsilon \in (0, \infty)$ is an arbitrary given constant. Let $p \in (1, \infty)$, $f \in L^{p_*}(\Omega)$, and the real-valued, bounded, and measurable matrix *A* satisfy (1.3). Then a function *u* is called a *weak solution* of the *Dirichlet boundary value problem* (1.1) if $u \in W_0^{1,p}(\Omega)$ and, for any $\varphi \in C_c^{\infty}(\Omega)$,

(1.8)
$$\int_{\Omega} A(x)\nabla u(x) \cdot \nabla \varphi(x) \, dx = \int_{\Omega} f(x)\varphi(x) \, dx.$$

Moreover, the Dirichlet problem (1.1) is said to be *uniquely solvable* if, for any $f \in L^{p_*}(\Omega)$, there exists a unique $u \in W_0^{1,p}(\Omega)$ such that (1.8) holds true for any $\varphi \in C_c^{\infty}(\Omega)$.

Remark 1.7. Assume that $f \in L^{2_*}(\Omega)$, where 2_* is as in (1.7) with *p* replaced by 2. Then, by the Sobolev inequality (see, for instance, [5, Theorem 1.1]) and the Lax–Milgram theorem (see, for

instance, [31, Theorem 5.8]), we find that the Dirichlet problem (1.1) is uniquely solvable and the weak solution u satisfies

(1.9)
$$\|\nabla u\|_{L^{2}(\Omega;\mathbb{R}^{n})} \leq C \|f\|_{L^{2*}(\Omega)},$$

where C is a positive constant independent of u and f.

For the Dirichlet problem (1.1) on bounded NTA domains, we have the following global regularity estimates in both Lebesgue spaces $L^q(\Omega)$ with $q \in (1, p_0)$, and Hardy spaces $H_z^q(\Omega)$ with $q \in (\frac{n}{n+1}, 1]$, where $p_0 \in (2, \infty)$ is a constant depending only on *n*, the domain Ω , and the matrix *A*.

Theorem 1.8. Let $n \ge 2$, Ω be a bounded NTA domain of \mathbb{R}^n , and the real-valued, bounded, and measurable matrix A satisfy (1.3).

(i) Then there exists a $p_0 \in (2, \infty)$, depending only on n, A, and Ω , such that, for any given $q \in (\frac{n}{n-1}, p_0)$ and $p \in (1, n)$ satisfying $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$, and any $f \in L^p(\Omega) \cap L^2(\Omega)$, the weak solution u of the problem (1.1) uniquely exists and satisfies $\nabla u \in L^q(\Omega; \mathbb{R}^n)$ and

 $\|\nabla u\|_{L^q(\Omega;\mathbb{R}^n)} \le C \|f\|_{L^p(\Omega)},$

where *C* is a positive constant independent of *u* and *f*.

(ii) Let $q \in (1, \frac{n}{n-1}]$ and $p \in (\frac{n}{n+1}, 1]$ satisfy $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$, and $f \in H^p_{L_D}(\Omega) \cap L^2(\Omega)$. Then the weak solution u of the problem (1.1) uniquely exists and satisfies $\nabla u \in L^q(\Omega; \mathbb{R}^n)$ and

$$\|\nabla u\|_{L^q(\Omega;\mathbb{R}^n)} \le C \|f\|_{H^p_{I_-}(\Omega)},$$

where C is a positive constant independent of u and f.

(iii) Let $q \in (\frac{n}{n+1}, 1]$ and $p \in (\frac{n}{n+2}, \frac{n}{n+1}]$ satisfy $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$, and $f \in H^p_{L_D}(\Omega) \cap L^2(\Omega)$. Then the weak solution u of the problem (1.1) uniquely exists and satisfies $\nabla u \in H^q_z(\Omega; \mathbb{R}^n)$ and

$$\|\nabla u\|_{H^q_{\mathcal{I}}(\Omega;\mathbb{R}^n)} \le C \|f\|_{H^p_{\mathcal{I}_n}(\Omega)}$$

where *C* is a positive constant independent of *u* and *f*. Here and thereafter, the space $H^q_{\tau}(\Omega; \mathbb{R}^n)$ is defined as $L^p(\Omega; \mathbb{R}^n)$ via $L^p(\Omega)$ replaced by $H^q_{\tau}(\Omega)$ [see (1.2)].

We prove Theorem 1.8 by the following strategy. We first obtain the global gradient estimate for the problem (1.1) in $L^q(\Omega)$ with $q \in (2, p_0)$, by using (1.9), a reverse Hölder inequality for the gradient of the weak solution of some local Dirichlet boundary value problems (see Lemma 3.8 below for the details), and a real-variable argument for $L^q(\Omega)$ estimates, essentially established by Shen [58, Theorem 3.4] (see also [57, Theorem 4.2.6] and [59, Theorem 3.3]). Then we show (i) by the global gradient estimate in $L^q(\Omega)$ with $q \in (2, p_0)$, the conclusion of (ii), and the complex interpolation theory of Hardy spaces (see, for instance, [42, Theorem 8.1 and (9.3)]). Moreover, we prove (ii) via establishing some fine estimate for the kernels $\{K_t^{L_D}\}_{t>0}$ (see Lemma 5.1 below), and using the global gradient estimate in $L^q(\Omega)$ with $q \in (2, p_0)$, and the molecular characterization of $H_{L_D}^p(\Omega)$. Finally, we show (iii) via using the global gradient estimate in $L^q(\Omega)$ with $q \in (2, p_0)$, and $H^p(\mathbb{R}^n)$.

As a corollary of Theorems 1.5 and 1.8, we have the following conclusion.

Corollary 1.9. Let $n \ge 2$, Ω be a bounded NTA domain of \mathbb{R}^n , and $\delta_0 \in (0, 1]$ as in Theorem 1.1. Assume that $q \in (1, \frac{n}{n-1}]$ and $p \in (\frac{n}{n+\delta_0}, 1]$ satisfy $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$. Let $f \in H^p_r(\Omega) \cap L^2(\Omega)$. Then the weak solution u of the problem (1.1) uniquely exists and satisfies $\nabla u \in L^q(\Omega; \mathbb{R}^n)$ and

$$\|\nabla u\|_{L^q(\Omega;\mathbb{R}^n)} \le C \|f\|_{H^p_r(\Omega)},$$

where C is a positive constant independent of u and f.

- **Remark 1.10.** (i) By an example given in [4, p. 120] (which is essentially due to C. E. Kenig, as was pointed out in [4, p. 119, Theorem 7]), we find that, even when $\Omega \subset \mathbb{R}^2$ is a bounded smooth domain, there exists a real-valued, bounded, and measurable matrix A satisfying (1.3) such that, for any given $p \in (2, \infty)$, the weak solution u of the problem (1.1) with some $f \in L^2(\Omega)$ satisfies $|\nabla u| \notin L^p(\Omega)$. Therefore, the range $q \in (\frac{n}{n-1}, p_0)$ of q obtained in Theorem 1.8(i) is sharp.
 - (ii) When A := I (the identity matrix) and Ω is a bounded Lipschitz domain of ℝⁿ, Dahlberg [18, Theorem 1] obtained Theorem 1.8(i) with p₀ := 3 + ε₀ when n ≥ 3, and with p₀ := 4 + ε₀ when n = 2, where ε₀ ∈ (0,∞) is a constant depending only on n and Ω. Meanwhile, Dahlberg [18] also showed that the range q ∈ (ⁿ/_{n-1}, p₀) of q is sharp in this case. Thus, Theorem 1.8(i) extends the results of Dahlberg [18, Theorem 1] via weakening the assumptions on both the coefficient matrix A and the domain Ω.

Moreover, to the best of our knowledge, the global gradient estimates obtained in (ii) and (iii) of Theorem 1.8 and Corollary 1.9 are new even when $\Omega \subset \mathbb{R}^n$ is a bounded Lipschitz domain.

The organization of the remainder of this article is as follows.

In Section 2, we present the definitions of both NTA domains and quasi-convex domains, some geometrical properties of NTA domains, the definition of the (γ, R) -BMO condition, and the atomic and the molecular characterizations of the Hardy spaces $H^p(\mathbb{R}^n)$, $H^p(\Omega)$, and $H^p_{L_D}(\Omega)$ associated with L_D .

Section 3 is devoted to the proof of Theorem 1.1. The proofs of Theorems 1.5 and 1.8 are presented, respectively, in Sections 4 and 5.

Finally, we make some conventions on notation. Throughout the whole article, we always denote by *C* or *c* a *positive constant* which is independent of the main parameters, but it may vary from line to line. We also use $C_{(\gamma,\beta,...)}$ to denote a *positive constant* depending on the indicated parameters γ, β, \ldots . The symbol $f \leq g$ means that $f \leq Cg$. If $f \leq g$ and $g \leq f$, then we write $f \sim g$. If $f \leq Cg$ and g = h or $g \leq h$, we then write $f \leq g \sim h$ or $f \leq g \leq h$, rather than $f \leq g = h$ or $f \leq g \leq h$. For each ball $B := B(x_B, r_B)$ in \mathbb{R}^n , with $x_B \in \mathbb{R}^n$ and $r_B \in (0, \infty)$, and $\alpha \in (0, \infty)$, let $\alpha B := B(x_B, \alpha r_B)$; furthermore, denote by B_Ω the ball $B \cap \Omega$ in Ω with *B* being a ball of \mathbb{R}^n . For any subset *E* of \mathbb{R}^n , we denote by $E^{\mathbb{C}}$ the set $\mathbb{R}^n \setminus E$, and by $\mathbf{1}_E$ its *characteristic function*. We also let $\mathbb{N} := \{1, 2, \ldots\}$ and $\mathbb{Z}_+ := \mathbb{N} \cup \{0\}$. The symbol $\lfloor s \rfloor$ for any $s \in \mathbb{R}$ denotes the largest integer not greater than *s*. For any multi-index $\alpha := (\alpha_1, \ldots, \alpha_n) \in \mathbb{Z}_+^n := (\mathbb{Z}_+)^n$, let $|\alpha| := \alpha_1 + \cdots + \alpha_n$. For any ball *B* of \mathbb{R}^n or of Ω , let $S_j(B) := (2^{j+1}B) \setminus (2^jB)$ for any $j \in \mathbb{N}$, and $S_0(B) := 2B$. Moreover, for any $q \in [1, \infty]$, we denote by q' its *conjugate exponent*, namely, 1/q + 1/q' = 1. Finally, for any measurable set $E \subset \mathbb{R}^n$ with $|E| < \infty$, and any $f \in L^1(E)$, we let

$$\int_{E} f(x) \, dx := \frac{1}{|E|} \int_{E} f(x) \, dx$$

2 Preliminaries

In this section, we first recall the definitions of NTA domains, quasi-convex domains, and the (γ, R) -BMO condition, and then give some geometrical properties of NTA domains. Moreover, we present the atomic and the molecular characterizations of the Hardy spaces $H^p(\mathbb{R}^n)$, $H^p(\Omega)$, and $H^p_{L_D}(\Omega)$ associated with L_D .

2.1 NTA Domains

In this subsection, we first recall the definitions of NTA domains introduced by Jerison and Kenig [38] (see also [44, 64]) and quasi-convex domains introduced by Jia et al. [40], and then state some geometrical properties of NTA domains. We begin with recalling several notions. For any given $x \in \mathbb{R}^n$ and measurable subset $E \subset \mathbb{R}^n$, let dist $(x, E) := \inf\{|x - y| : y \in E\}$. Meanwhile, for any measurable subsets $E, F \subset \mathbb{R}^n$, let dist $(E, F) := \inf\{|x - y| : x \in E, y \in F\}$ and diam $(E) := \sup\{|x - y| : x, y \in E\}$.

Definition 2.1. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a *domain* which means that Ω is a connected open set, and $\Omega^{\complement} := \mathbb{R}^n \setminus \Omega$.

- (i) The domain Ω is said to satisfy the *interior* [resp., *exterior*] *corkscrew condition* if there exist constants $R \in (0, \infty)$ and $\sigma \in (0, 1)$ such that, for any $x \in \partial \Omega$ and $r \in (0, R)$, there exists a point $x_0 \in \Omega$ [resp., $x_0 \in \Omega^{\complement}$], depending on x, such that $B(x_0, \sigma r) \subset \Omega \cap B(x, r)$ [resp., $B(x_0, \sigma r) \subset \Omega^{\complement} \cap B(x, r)$].
- (ii) The domain Ω is said to satisfy the *Harnack chain condition* if there exist a constant $m_1 \in (1, \infty)$ and a constant $m_2 \in (0, \infty)$ such that, for any $x_1, x_2 \in \Omega$ satisfying

$$M := \frac{|x_1 - x_2|}{\min\{\operatorname{dist}(x_1, \partial \Omega), \operatorname{dist}(x_2, \partial \Omega)\}} > 1,$$

there exists a chain $\{B_i\}_{i=1}^N$ of open Harnack balls, with $B_i \subset \Omega$ for any $i \in \{1, ..., N\}$, that connects x_1 to x_2 ; namely, $x_1 \in B_1$, $x_2 \in B_N$, $B_i \cap B_{i+1} \neq \emptyset$ for any $i \in \{1, ..., N-1\}$, and, for any $i \in \{1, ..., N\}$,

$$m_1^{-1}$$
diam $(B_i) \leq$ dist $(B_i, \partial \Omega) \leq m_1$ diam $(B_i),$

where the integer N satisfies $N \le m_2 \log_2 M$.

(iii) The domain Ω is called a *non-tangentially accessible domain* (for short, NTA *domain*) if Ω satisfies the interior and the exterior corkscrew conditions, and the Harnack chain condition.

We point out that NTA domains include Lipschitz domains, BMO₁ domains, Zygmund domains, quasi-spheres, and some Reifenberg flat domains as special examples (see, for instance, [38, 44, 64]). Moreover, it is well known that NTA domains are $W^{1,p}$ -extension domains with $p \in [1, \infty)$ (see, for instance, [34, 41]). Meanwhile, NTA domains have the following geometrical properties.

Lemma 2.2. Let $n \ge 2$ and Ω be a bounded NTA domain of \mathbb{R}^n .

(i) Then there exists a constant $C \in (0, 1]$, depending only on n and Ω , such that, for any ball $B \subset \mathbb{R}^n$ with the center $x_B \in \overline{\Omega}$ and the radius $r_B \in (0, \text{diam}(\Omega))$,

$$|B \cap \Omega| \ge C|B|.$$

- (ii) Let $B := B(x_B, r_B) \subset \mathbb{R}^n$ be a ball satisfying that $2B \subset \Omega$ and $4B \cap \Omega^{\complement} \neq \emptyset$. Then there exists a ball $\widetilde{B} \subset \Omega^{\complement}$ such that $r_{\widetilde{B}} \sim r_B$ and dist $(B, \widetilde{B}) \sim r_B$, where the positive equivalence constants are independent of both B and \widetilde{B} .
- (iii) For any ball $B \subset \mathbb{R}^n$ with the center $x_B \in \partial \Omega$ and the radius $r_B \in (0, \operatorname{diam}(\Omega))$, there exists a ball $\widetilde{B} \subset \Omega^{\mathbb{C}} \cap B$ such that $r_{\widetilde{B}} \sim r_B$, where the positive equivalence constants are independent of both B and \widetilde{B} . In particular, there exists a constant $C \in (0, 1]$, depending only on n and Ω , such that, for any ball $B \subset \mathbb{R}^n$ with the center $x_B \in \partial \Omega$ and the radius $r_B \in (0, \operatorname{diam}(\Omega))$,

$$|B \cap \Omega^{\cup}| \ge C|B|.$$

Proof. We first show (i). By the fact that Ω is a $W^{1,p}$ -extension domain with $p \in [1, \infty)$, and [34, Theorem 2], we conclude that there exist constants $R_0 \in (0, \infty)$ and $C_1 \in (0, 1]$ such that, for any given ball $B := B(x_B, r_B)$ of \mathbb{R}^n with $x_B \in \overline{\Omega}$ and $r_B \in (0, R_0]$,

$$(2.1) |B \cap \Omega| \ge C_1|B|.$$

If diam $(\Omega) \leq R_0$, then, from (2.1), it follows that (i) holds true in this case. If diam $(\Omega) > R_0$, then, for any ball $B := B(x_B, r_B)$ of \mathbb{R}^n with $x_B \in \overline{\Omega}$ and $r_B \in (R_0, \text{diam}(\Omega))$,

$$|B \cap \Omega| \ge |B(x_B, R_0) \cap \Omega| \ge C_1 |B(x_B, R_0)|$$

= $C_1 \left[\frac{R_0}{\operatorname{diam}(\Omega)} \right]^n |B(x_B, \operatorname{diam}(\Omega))| \ge C_1 \left[\frac{R_0}{\operatorname{diam}(\Omega)} \right]^n |B|,$

which, together with (2.1), further implies that (i) holds true in this case. This finishes the proof of (i).

Now, we prove (ii). Let $y_B \in \partial \Omega$ be such that $|x_B - y_B| = \text{dist}(x_B, \partial \Omega)$. Define $\ell := |x_B - y_B|$. Then $\ell \in (2r_B, 4r_B)$. By the exterior corkscrew condition of Ω , we find that there exists a ball $\widetilde{B} := B(x_0, \sigma r) \subset [\Omega^{\complement} \cap B(y_B, r)]$, where $r \in (\frac{1}{2}\min\{R, \ell\}, \min\{R, \ell\})$, and σ and R are as in Definition 2.1. From this, we deduce that

$$\frac{r_B}{r_{\widetilde{B}}} \le \max\left\{\frac{\operatorname{diam}\left(\Omega\right)}{2\sigma R}, \frac{1}{\sigma}\right\}$$

and

$$\frac{r_B}{r_{\widetilde{B}}} \ge \frac{1}{4\sigma},$$

which implies that $r_B \sim r_{\widetilde{B}}$. Moreover, it is easy to see that dist $(B, \widetilde{B}) \geq r_B$ and

dist
$$(B, \widetilde{B}) \leq |x_B - y_B| + r = \ell + \frac{1}{\sigma} r_{\widetilde{B}} \leq r_B.$$

Therefore, dist $(B, \tilde{B}) \sim r_B$. This finishes the proof of (ii).

Finally, we show (iii). Let $B := B(x_B, r_B)$ be a ball of \mathbb{R}^n with the center $x_B \in \partial \Omega$ and the radius $r_B \in (0, \operatorname{diam}(\Omega))$. Then, when $r_B \in (0, \min\{R, \operatorname{diam}(\Omega)\})$, where $R \in (0, \infty)$ is as in Definition 2.1, from the exterior corkscrew condition of Ω , we deduce that there exists a ball $\widetilde{B} := B(x_0, \sigma r_B) \subset [\Omega^{\mathbb{C}} \cap B(x_B, r_B)]$, which further implies that $r_{\widetilde{R}} \sim r_B$ and

$$\left| B(x_B, r_B) \cap \Omega^{\mathbb{C}} \right| \ge \left| B(x_0, \sigma r_B) \right| \ge \left| B(x_B, r_B) \right|$$

Thus, (iii) holds true in the case that $r_B \in (0, \min\{R, \operatorname{diam}(\Omega)\})$.

Moreover, when $r_B \in (\min\{R, \operatorname{diam}(\Omega)\}, \operatorname{diam}(\Omega))$, let $\ell := \min\{R, \operatorname{diam}(\Omega)\}/2$. By the exterior corkscrew condition of Ω again, we conclude that there exists a ball $\widetilde{B} := B(x_0, \sigma \ell) \subset [\Omega^{\complement} \cap B(x_B, \ell)]$, which implies that $r_{\widetilde{B}} \sim r_B$ and

$$\left| B(x_B, r_B) \cap \Omega^{\mathbb{C}} \right| \ge \left| B(x_B, \ell) \cap \Omega^{\mathbb{C}} \right| \ge \left| B(x_B, \ell) \right|$$
$$\sim \left| B(x_B, \operatorname{diam}(\Omega)) \right| \ge \left| B(x_B, r_B) \right|.$$

From this, it follows that (iii) holds true in the case that $r_B \in (\min\{R, \operatorname{diam}(\Omega)\}, \operatorname{diam}(\Omega))$. This finishes the proof of (iii) and hence of Lemma 2.2.

Remark 2.3. Let $n \ge 2$ and Ω be a bounded NTA domain of \mathbb{R}^n . By Lemma 2.2(i), we find that $(\Omega, |\cdot|, dx)$ is a space of homogeneous type in the sense of Coifman and Weiss [17], where $|\cdot|$ denotes the usual norm on \mathbb{R}^n , and dx the Lebesgue measure on \mathbb{R}^n . Moreover, as a space of homogeneous type, the collection of all balls of Ω is given by the set

$$\{B_{\Omega} := B \cap \Omega : \text{ any ball } B \subset \mathbb{R}^n \text{ satisfying } x_B \in \overline{\Omega} \text{ and } r_B \in (0, \operatorname{diam}(\Omega))\},\$$

where x_B denotes the center of B and r_B the radius of B.

Now, we recall the definition of the (γ, σ, R) quasi-convex domain as follows.

Definition 2.4. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a domain, $\gamma, \sigma \in (0, 1)$, and $R \in (0, \infty)$. Then Ω is called a (γ, σ, R) quasi-convex domain if, for any $x \in \partial \Omega$ and $r \in (0, R]$,

- (a) there exists an $x_0 \in \Omega$, depending on *x*, such that $B(x_0, \sigma r) \subset [\Omega \cap B(x, r)]$;
- (b) there exists a convex domain V := V(x, r), depending on x and r, such that $[B(x, r) \cap \Omega] \subset V$ and

$$d_H(\partial(B(x,r)\cap\Omega),\partial V) \leq \gamma r,$$

where $d_H(\cdot, \cdot)$ denotes the *Hausdorff distance* which is defined by setting, for any non-empty measurable subsets E_1 and E_2 of \mathbb{R}^n ,

$$d_H(E_1, E_2) := \max \left\{ \sup_{x \in E_1} \inf_{y \in E_2} |x - y|, \sup_{x \in E_2} \inf_{y \in E_1} |x - y| \right\}.$$

The concept of quasi-convex domains was introduced by Jia et al. [40] to study the global regularity of second-order elliptic equations. Roughly speaking, a quasi-convex domain is a domain satisfying that the local boundary is close to be convex at small scales. It is easy to find that, if Ω is a convex domain, then Ω is a (γ , σ , R) quasi-convex domain for any $\gamma \in (0, 1)$, some $\sigma \in (0, 1)$, and some $R \in (0, \infty)$.

Remark 2.5. (i) Let $n \ge 2$, $\gamma \in (0, 1)$, and $R \in (0, \infty)$. A domain $\Omega \subset \mathbb{R}^n$ is called a (γ, R) -*Reifenberg flat domain* if, for any $x_0 \in \partial \Omega$ and $r \in (0, R_0]$, there exists a system $\{y_1, \ldots, y_n\}$ of coordinates, which may depend on x_0 and r, such that, in this coordinate system, $x_0 = \mathbf{0}$ and

 $\left[B(\mathbf{0},r) \cap \{y \in \mathbb{R}^n : y_n > \gamma r\}\right] \subset \left[B(\mathbf{0},r) \cap \Omega\right] \subset \left[B(\mathbf{0},r) \cap \{y \in \mathbb{R}^n : y_n > -\gamma r\}\right].$

The Reifenberg flat domain was introduced by Reifenberg [53], which naturally appears in the theory of both minimal surfaces and free boundary problems. A typical example of Reifenberg flat domains is the well-known Van Koch snowflake (see, for instance, [64]). Moreover, it was shown by Kenig and Toro [44, Theorem 3.1] that there exists a $\gamma_0(n) \in$ (0, 1), depending only on *n*, such that, if Ω is a (γ, R) -Reifenberg flat domain for some $\gamma \in (0, \gamma_0(n)]$ and $R \in (0, \infty)$, then Ω is an NTA domain. Meanwhile, it is easy to see that a (γ, R) -Reifenberg flat domain is also a (γ, σ, R) quasi-convex domain for some $\sigma \in (0, 1)$. In recent years, boundary value problems of elliptic or parabolic equations on Reifenberg flat domains have been widely concerned and studied (see, for instance, [8, 11, 23, 24, 25]).

- (ii) It is known that, for any open set $\Omega \subset \mathbb{R}^n$ with compact boundary, Ω is a semi-convex domain of \mathbb{R}^n if and only if Ω is a Lipschitz domain satisfying a uniform exterior ball condition (see, for instance, [48, 49, 67] for the definitions of both the semi-convex domain and the uniform exterior ball condition). It is worth pointing out that convex domains of \mathbb{R}^n are semi-convex domains and (semi-)convex domains are special cases of Lipschitz domains (see, for instance, [48, 49, 67]). Furthermore, a (semi-)convex domain is also a (γ, σ, R) quasi-convex domain for any $\gamma \in (0, 1)$, some $\sigma \in (0, 1)$, and some $R \in (0, \infty)$ (see, for instance, [69]).
- (iii) On NTA domains, Lipschitz domains, quasi-convex domains, Reifenberg flat domains, C^1 domains, and (semi-)convex domains, we have the following relations (see, for instance, [38, 40, 44, 64, 67, 69]).
 - (a) class of C^1 domains
 - \subsetneq class of Lipschitz domains with small Lipschitz constants
 - \subsetneq class of Lipschitz domains
 - \subsetneq class of NTA domains;
 - (b) class of C^1 domains
 - ⊊ class of Lipschitz domains with small Lipschitz constants
 - \subsetneq class of Reifenberg flat domains
 - \subsetneq class of quasi-convex domains;
 - (c) class of (semi-)convex domains \subsetneq class of quasi-convex domains.

Next, we recall the definition of the (γ, R) -BMO condition (see, for instance, [11]). For any given domain $\Omega \subset \mathbb{R}^n$, denote by $L^1_{loc}(\Omega)$ the set of all locally integrable functions on Ω .

Definition 2.6. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a domain, and γ , $R \in (0, \infty)$. A function $f \in L^1_{loc}(\Omega)$ is said to satisfy the (γ, R) -BMO *condition* if

$$||f||_{*,R} := \sup_{B(x,r) \subset \Omega, r \in (0,R)} \int_{B(x,r)} |f(y) - f_{B(x,r)}| \, dy \le \gamma,$$

where the supremum is taken over all balls $B(x, r) \subset \Omega$ with $r \in (0, R)$, and $f_{B(x,r)} := \oint_{B(x,r)} f(z) dz$. Moreover, a matrix $A := \{a_{ij}\}_{i,j=1}^{n}$ is said to satisfy the (γ, R) -BMO *condition* if, for any $i, j \in \mathbb{R}$ {1, ..., *n*}, a_{ij} satisfies the (γ , *R*)-BMO condition.

Hardy Spaces 2.2

In this subsection, we recall the atomic and the molecular characterizations of the Hardy spaces $H^p(\mathbb{R}^n)$, $H^p(\Omega)$, and $H^p_{L_D}(\Omega)$ associated with L_D .

Definition 2.7. Let $p \in (0, 1]$, $q \in (1, \infty]$, and $s \in \mathbb{Z}_+$ with $s \ge \lfloor n(\frac{1}{p} - 1) \rfloor$.

- (i) A function $a \in L^q(\mathbb{R}^n)$ is called a (p, q, s)-atom if there exists a ball B of \mathbb{R}^n such that
 - (i)₁ supp $(a) \subset B$;
 - (i)₂ $||a||_{L^q(\mathbb{R}^n)} \leq |B|^{1/q-1/p};$
 - (i)₃ $\int_{\mathbb{R}^n} a(x) x^{\alpha} dx = 0$ for any $\alpha \in \mathbb{Z}^n_+$ with $|\alpha| \le s$.
- (ii) The *atomic Hardy space* $H_{at}^{p, q, s}(\mathbb{R}^n)$ is defined to be the set of all $f \in \mathcal{S}'(\mathbb{R}^n)$ satisfying that $f = \sum_{j=1}^{\infty} \lambda_j a_j$ in $\mathcal{S}'(\mathbb{R}^n)$, where $\{a_j\}_{j=1}^{\infty}$ is a sequence of (p, q, s)-atoms, and $\{\lambda_j\}_{j=1}^{\infty} \subset \mathbb{C}$ satisfies $\sum_{j=1}^{\infty} |\lambda_j|^p < \infty$. Moreover, for any $f \in H^{p,q,s}_{at}(\mathbb{R}^n)$, the quasi-norm $||f||_{H^{p,q,s}_{at}(\mathbb{R}^n)}$ of f is defined by setting

$$||f||_{H^{p,q,s}_{\mathrm{at}}(\mathbb{R}^n)} := \inf\left\{ \left(\sum_{j=1}^{\infty} |\lambda_j|^p \right)^{1/p} \right\},\$$

where the infimum is taken over all decompositions of f as above.

Definition 2.8. Let $p \in (0, 1]$, $q \in (1, \infty]$, $s \in \mathbb{Z}_+$ with $s \ge \lfloor n(\frac{1}{p} - 1) \rfloor$, and $\varepsilon \in (0, \infty)$.

- (i) A function $\alpha \in L^q(\mathbb{R}^n)$ is called a (p, q, s, ε) -molecule associated with the ball B of \mathbb{R}^n if (i)₁ for any $j \in \mathbb{Z}_+$, $\|\alpha\|_{L^q(S_j(B))} \le 2^{-j\varepsilon} |2^j B|^{1/q} |B|^{-1/p}$.
 - (i)₂ $\int_{\mathbb{R}^n} \alpha(x) x^{\beta} dx = 0$ for any $\beta \in \mathbb{Z}^n_+$ with $|\beta| \le s$.
- (ii) The molecular Hardy space $H^{p,q,s,\varepsilon}_{mol}(\mathbb{R}^n)$ is defined to be the set of all $f \in \mathcal{S}'(\mathbb{R}^n)$ satisfying that $f = \sum_{j=1}^{\infty} \lambda_j \alpha_j$ in $\mathcal{S}'(\mathbb{R}^n)$, where $\{\alpha_j\}_{j=1}^{\infty}$ is a sequence of (p, q, s, ε) -molecules, and $\{\lambda_j\}_{j=1}^{\infty} \subset \mathbb{C}$ satisfies $\sum_{j=1}^{\infty} |\lambda_j|^p < \infty$. Moreover, define

$$\|f\|_{H^{p,q,s,\varepsilon}_{\mathrm{mol}}(\mathbb{R}^n)} := \inf\left\{ \left(\sum_{j=1}^{\infty} |\lambda_j|^p \right)^{1/p} \right\},\$$

where the infimum is taken over all decompositions of f as above.

Then we have the following atomic and molecular characterizations of the Hardy space $H^p(\mathbb{R}^n)$, respectively (see, for instance, [45, 61, 63]).

Lemma 2.9. Let $p \in (0, 1]$, $q \in (1, \infty]$, and $s \in \mathbb{Z}_+$ with $s \ge \lfloor n(\frac{1}{p} - 1) \rfloor$, and $\varepsilon \in (\max\{n + 1\}, n$ s, n/p, ∞). Then the spaces $H^p(\mathbb{R}^n) = H^{p, q, s}_{at}(\mathbb{R}^n) = H^{p, q, s, \varepsilon}_{mol}(\mathbb{R}^n)$ with equivalent quasi-norms.

Definition 2.10. Let $n \ge 2$, Ω be a domain of \mathbb{R}^n , $p \in (0, 1]$, $q \in (1, \infty]$, and $s \in \mathbb{Z}_+$ with $s \ge \lfloor n(\frac{1}{p} - 1) \rfloor$.

- (i) A ball $B \subset \Omega$ is called a *type* (a) *ball* of Ω if $4B \subset \Omega$, and a ball $B \subset \Omega$ is called a *type* (b) *ball* of Ω if $2B \cap \Omega^{\complement} = \emptyset$ but $4B \cap \Omega^{\complement} \neq \emptyset$.
- (ii) A function a ∈ L^q(Ω) is called a *type* (a) (p, q, s)_Ω-atom if there exists a type (a) ball B ⊂ Ω such that supp (a) ⊂ B and a is a (p, q, s)-atom.
 Moreover, a function b ∈ L^q(Ω) is called a *type* (b) (p, q)_Ω-atom if there exists a type (b)

ball $B \subset \Omega$ such that supp $(b) \subset B$ and $||b||_{L^q(\Omega)} \leq |B|^{1/q-1/p}$.

(iii) The *atomic Hardy space* $H_{at}^{p,q,s}(\Omega)$ is defined to be the set of all distributions $f \in \mathcal{D}'(\Omega)$ satisfying that there exist two sequences $\{\lambda_j\}_{j=1}^{\infty} \subset \mathbb{C}$ and $\{\kappa_j\}_{j=1}^{\infty} \subset \mathbb{C}$, a sequence $\{a_j\}_{j=1}^{\infty}$ of type (a) $(p, q, s)_{\Omega}$ -atoms, and a sequence $\{b_j\}_{j=1}^{\infty}$ of type (b) $(p, q)_{\Omega}$ -atoms such that

(2.2)
$$f = \sum_{j=1}^{\infty} \lambda_j a_j + \sum_{j=1}^{\infty} \kappa_j b_j$$

in $\mathcal{D}'(\Omega)$, and

$$\sum_{j=1}^\infty |\lambda_j|^p + \sum_{j=1}^\infty |\kappa_j|^p < \infty.$$

Furthermore, for any given $f \in H^{p,q,s}_{at}(\Omega)$, let

$$||f||_{H^{p,q,s}_{\mathrm{at}}(\Omega)} := \inf\left\{ \left(\sum_{j=1}^{\infty} |\lambda_j|^p \right)^{1/p} + \left(\sum_{j=1}^{\infty} |\kappa_j|^p \right)^{1/p} \right\},\$$

where the infimum is taken over all decompositions of f as in (2.2).

Then we have the following atomic characterization of the Hardy space $H^p(\Omega)$ on the domain Ω (see, for instance, [50, Theorem 1]).

Lemma 2.11. Let $n \ge 2$ and $\Omega \subset \mathbb{R}^n$ be a proper domain, $p \in (0, 1]$, $q \in (1, \infty]$, and $s \in \mathbb{Z}_+$ with $s \ge \lfloor n(\frac{1}{n} - 1) \rfloor$. Then $H^p(\Omega) = H^{p,q,s}_{at}(\Omega)$ with equivalent quasi-norms.

Definition 2.12. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, L_D as in (1.4), and $p \in (0, 1]$. Assume that $q \in (1, \infty]$, $M \in \mathbb{N}$, $\epsilon \in (0, \infty)$, $B := B(x_B, r_B)$ with $x_B \in \Omega$ and $r_B \in (0, \text{diam } (\Omega))$, and $B_{\Omega} := B \cap \Omega$.

(i) A function α ∈ L^q(Ω) is called a (p, q, M, ε)_{L_D}-molecule associated with the ball B_Ω if, for any k ∈ {0, ..., M} and j ∈ Z₊, it holds true that

$$\left\| \left(r_B^{-2} L_D^{-1} \right)^k \alpha \right\|_{L^q(S_j(B_\Omega))} \le 2^{-j\epsilon} |2^j B_\Omega|^{1/q} |B_\Omega|^{-1/p}.$$

(ii) For any $f \in L^2(\Omega)$,

(2.3)
$$f = \sum_{j=1}^{\infty} \lambda_j \alpha_j$$

is called a *molecular* $(p, q, M, \epsilon)_{L_D}$ -representation of the function f if, for any $j \in \mathbb{N}, \alpha_j$ is a $(p, q, M, \epsilon)_{L_D}$ -molecule associated with the ball $B_{\Omega, j} \subset \Omega$, the summation (2.3) converges in $L^2(\Omega)$, and $\{\lambda_j\}_{j=1}^{\infty} \subset \mathbb{C}$ satisfies $\sum_{j=1}^{\infty} |\lambda_j|^p < \infty$.

Let

$$\mathbb{H}_{L_{D}, \operatorname{mol}}^{p, q, M, \epsilon}(\Omega) := \left\{ f \in L^{2}(\Omega) : f \text{ has a molecular } (p, q, M, \epsilon)_{L} \text{-representation} \right\}$$

equipped with the quasi-norm

$$\|f\|_{H^{p,q,M,\epsilon}_{L,\mathrm{mol}}(\Omega)} := \inf\left\{ \left(\sum_{j=1}^{\infty} |\lambda_j|^p \right)^{1/p} : \sum_{j=1}^{\infty} \lambda_j \alpha_j \text{ is a } (p, q, M, \epsilon)_{L_D} \text{-representation of } f \right\},$$

where the infimum is taken over all molecular $(p, q, M, \epsilon)_{L_D}$ -representations of f as above. Then the *molecular Hardy space* $H_{L_D, \text{mol}}^{p, q, M, \epsilon}(\Omega)$ is then defined as the completion of the set $\mathbb{H}_{L_D, \text{mol}}^{p, q, M, \epsilon}(\Omega)$ with respect to the *quasi-norm* $\|\cdot\|_{H_{L_D, \text{mol}}^{p, q, M, \epsilon}(\Omega)}$.

By Remark 2.3, we find that bounded NTA domains of \mathbb{R}^n are spaces of homogeneous type. Thus, for the Hardy space $H^p_{L_D}(\Omega)$, we have the following molecular characterization (see, for instance, [6]).

Lemma 2.13. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, L_D as in (1.4), and $p \in (0, 1]$. Then, for any $q \in (1, \infty)$, $M \in \mathbb{N} \cap (\frac{n}{2p}, \infty)$, and $\epsilon \in (\frac{n}{p}, \infty)$, $H_{L_D}^p(\Omega) = H_{L_D, \text{mol}}^{p, q, M, \epsilon}(\Omega)$ with equivalent quasi-norms.

3 Proof of Theorem 1.1

In this section, we show Theorem 1.1. Let L_D be as in (1.4). It was shown by Grüter and Widman [32] that, when $n \ge 3$, the Green function associated with the operator L_D exists; in this case, denote by $G(\cdot, \cdot)$ the Green function associated with L_D . Then we have the following estimates for $G(\cdot, \cdot)$ which are known (see, for instance, [35, Remark 4.9]).

Lemma 3.1. Let $n \ge 3$ and Ω be a bounded NTA domain of \mathbb{R}^n . Then there exist constants $C \in (0, \infty)$ and $\beta \in (0, 1]$, depending only on n, the matrix A appeared in the definition of L_D , and Ω , such that, for any $x, y \in \Omega$,

(3.1)
$$G(x,y) \le C[\delta(y)]^{\beta} |x-y|^{2-n-\beta},$$

here and thereafter, $\delta(y) := \text{dist}(y, \partial \Omega)$ with $\partial \Omega$ being the boundary of Ω .

Now, we prove the following Hölder continuity of the Green function G via using Lemma 3.1 and the Harnack inequality (see, for instance, [31, Theorem 8.22]).

Lemma 3.2. Let $n \ge 3$ and Ω be a bounded NTA domain of \mathbb{R}^n . Then there exist constants $C \in (0, \infty)$ and $\beta \in (0, 1]$, depending only on n, the matrix A appeared in the definition of L_D , and Ω , such that, for any $x, y_1, y_2 \in \Omega$,

$$|G(x, y_1) - G(x, y_2)| \le C |y_1 - y_2|^{\beta} \left[|x - y_1|^{2 - n - \beta} + |x - y_2|^{2 - n - \beta} \right].$$

Proof. We show the present lemma via borrowing some ideas from the proof of [32, Theorem (1.9)]. Fix $x, y_1, y_2 \in \Omega$. Without loss of generality, we may assume that $G(x, y_1) \ge G(x, y_2)$. Now, we prove the present lemma by considering the following three cases on $|y_1 - y_2|$.

Case 1) $|y_1 - y_2| \ge |x - y_1|/2$. In this case, by the upper estimate that $G(x, z) \le |x - z|^{2-n}$ for any $x, z \in \Omega$ (see, for instance, [32, (1.8)]) and $1 \le |y_1 - y_2||x - y_1|^{-1}$, we conclude that, for any given $\beta \in (0, 1]$,

$$(3.2) |G(x, y_1) - G(x, y_2)| = G(x, y_1) - G(x, y_2) \le G(x, y_1) \le |x - y_1|^{2-n} \le |y_1 - y_2|^{\beta} |x - y_1|^{2-n-\beta} \le |y_1 - y_2|^{\beta} \left[|x - y_1|^{2-n-\beta} + |x - y_2|^{2-n-\beta} \right].$$

Case 2) $|y_1 - y_2| < |x - y_1|/2$ and $\delta(y_1) \le |y_1 - y_2|$. In this case, from Lemma 3.1 and $\delta(y_1) \le |y_1 - y_2|$, it follows that there exists a $\beta \in (0, 1]$ such that

(3.3)
$$|G(x, y_1) - G(x, y_2)| \le G(x, y_1) \le [\delta(y_1)]^{\beta} |x - y_1|^{2 - n - \beta}$$
$$\le |y_1 - y_2|^{\beta} \left[|x - y_1|^{2 - n - \beta} + |x - y_2|^{2 - n - \beta} \right]$$

Case 3) $|y_1 - y_2| < |x - y_1|/2$ and $\delta(y_1) > |y_1 - y_2|$. In this case, let $R := \min\{\delta(y_1), |x - y_1|/2\}$. It is easy to see that $L_D^*G(x, \cdot) = 0$ in $B(y_1, R)$. Applying the Harnack inequality (see, for instance, [31, Theorem 8.22]), we find that there exists a $\beta \in (0, 1]$ such that, for any $0 < r_1 \le r_2 \le R$,

$$\max_{B(y_1,r_1)} G(x,\cdot) - \min_{B(y_1,r_1)} G(x,\cdot) \lesssim \left(\frac{r_1}{r_2}\right)^{\beta} \left[\max_{B(y_1,r_2)} G(x,\cdot)\right],$$

which further implies that

(3.4)
$$|G(x, y_1) - G(x, y_2)| \leq |y_1 - y_2|^{\beta} R^{-\beta} \max_{B(y_1, R)} G(x, \cdot).$$

If $R = |x_1 - y|/2$, then, by (3.4) and the upper estimate $G(x, y_1) \leq |x - y_1|^{2-n}$, we conclude that

(3.5)
$$|G(x, y_1) - G(x, y_2)| \leq |y_1 - y_2|^{\beta} |x - y_1|^{2-n-\beta} \\ \leq |y_1 - y_2|^{\beta} \left[|x - y_1|^{2-n-\beta} + |x - y_2|^{2-n-\beta} \right].$$

If $R = \delta(y_1)$, then $\delta(y_1) \le |x-y_1|/2$. From this and Lemma 3.1, we deduce that, for any $z \in B(y_1, R)$,

$$\begin{split} G(x,z) &\lesssim [\delta(z)]^{\beta} |x-z|^{2-n-\beta} \lesssim [|z-y_1| + \delta(y_1)]^{\beta} [|x-y_1| - |y_1-z|]^{2-n-\beta} \\ &\lesssim [\delta(y_1)]^{\beta} |x-y_1|^{2-n-\beta}, \end{split}$$

which, combined with (3.4), further implies that

$$|G(x, y_1) - G(x, y_2)| \leq |y_1 - y_2|^{\beta} \left[|x - y_1|^{2 - n - \beta} + |x - y_2|^{2 - n - \beta} \right].$$

This, together with (3.2), (3.3), and (3.5), then finishes the proof of Lemma 3.2.

In what follows, for any given $z \in \mathbb{C}$, denote by $\Re(z)$ and $\arg z$, respectively, the real part of z and the argument in $(-\pi, \pi]$ of z. Recall that L_D is a ω_0 -accretive operator on $L^2(\Omega)$ with some $\omega_0 \in [0, \pi/2)$ (see, for instance, [4] for the definition of the ω_0 -accretive operator). For any given $\theta \in [0, \pi)$, let $\Gamma_{\theta} := \{z \in \mathbb{C} \setminus \{0\}, |\arg z| < \theta\}$. It is well known that $-L_D$ generates a holomorphic semigroup $\{e^{-zL_D}\}_{z \in \mathbb{C}, |\arg z| < \frac{\pi}{2} - \omega_0}$ in $\Gamma_{\frac{\pi}{2} - \omega_0}$ (see, for instance, [4, 51]).

Moreover, it is known that, for any $\lambda \in \Gamma_{\pi-\omega_0}$, the operator $L_D + \lambda I$ is bounded on $L^2(\Omega)$, and has the integral kernel G_{λ} , where I denotes the identity operator. Namely, for any given $f \in L^2(\Omega)$ and $x \in \Omega$,

$$(L_D + \lambda I)^{-1} f(x) = \int_{\Omega} G_{\lambda}(x, y) f(y) \, dy.$$

Lemma 3.3. Assume that $n \ge 3$ and Ω is a bounded NTA domain of \mathbb{R}^n . Let $\mu \in (\omega_0, \pi/2)$. Then there exists a $\delta \in (0, 1]$ such that, for any $\lambda \in \Gamma_{\pi-\mu}$ and $x, y_1, y_2 \in \Omega$ with $x \ne y_i$ for $i \in \{1, 2\}$,

$$|G_{\lambda}(x, y_1) - G_{\lambda}(x, y_2)| \le C \left[\max_{i \in \{1, 2\}} e^{-\gamma \sqrt{|\lambda|} ||x - y_i|} \right] |y_1 - y_2|^{\delta} \left[\frac{1}{|x - y_1|^{n-2+\delta}} + \frac{|\lambda|^{\delta/2}}{|x - y_2|^{n-2}} \right],$$

where *C* is a positive constant depending only on μ , *n*, and Ω , and γ is a positive constant depending only on μ and δ .

Proof. Let $\lambda \in \Gamma_{\pi-\mu}$. By the resolvent identity (see, for instance, [52, p. 36, (9.1)]), we conclude that, for any $y, z \in \Omega$ with $y \neq z$,

(3.6)
$$G_{\lambda}(y,z) = G(y,z) + \lambda \int_{\Omega} G(t,z) G_{\lambda}(y,t) dt.$$

Moreover, since L_D generates a holomorphic semigroup $\{e^{-zL_D}\}_{z\in\Gamma_{\frac{\pi}{2}-\omega_0}}$ (see, for instance, [51, Theorem 1.52]), and the kernels $\{K_t^{L_D}\}_{t>0}$ of the semigroup $\{e^{-tL_D}\}_{t>0}$ satisfy the Gaussian upper bound estimate (1.5), similarly to the proof of [26, Lemma 2.3], it follows that there exists a positive constant *c* such that, for any $y, z \in \Omega$ with $y \neq z$,

(3.7)
$$|G_{\lambda}(y,z)| \leq e^{-c \sqrt{|\lambda|}|y-z|} \frac{1}{|y-z|^{n-2}}$$

Thus, by (3.6), (3.7), and Lemma 3.2, we find that there exists a $\beta \in (0, 1]$ such that, for any given $x, y_1, y_2 \in \Omega$ with $x \neq y_i$ for any $i \in \{1, 2\}$,

$$(3.8) |G_{\lambda}(x, y_1) - G_{\lambda}(x, y_2)| \\
\leq |G(x, y_1) - G(x, y_2)| + |\lambda| \int_{\Omega} |G(z, y_1) - G(z, y_2)| |G_{\lambda}(x, z)| dz \\
\lesssim |y_1 - y_2|^{\beta} \left[|x - y_1|^{2-n-\beta} + |x - y_2|^{2-n-\beta} \right] \\
+ |\lambda||y_1 - y_2|^{\beta} \int_{\Omega} \frac{e^{-c\sqrt{|\lambda|}|x-z|}}{|x - z|^{n-2}} \left[|z - y_1|^{2-n-\beta} + |z - y_2|^{2-n-\beta} \right] dz.$$

For any $i \in \{1, 2\}$, let $\Omega_1 := \{z \in \Omega : |x - z| \le |z - y_i|\}$ and $\Omega_2 := \{z \in \Omega : |x - z| > |z - y_i|\}$. Then, for any $i \in \{1, 2\}$, we have

(3.9)
$$\int_{\Omega} \frac{e^{-c\sqrt{|\lambda|}|x-z|}}{|x-z|^{n-2}} \frac{1}{|z-y_i|^{n-2+\beta}} dz$$

$$= \int_{\Omega_1} \frac{e^{-c \sqrt{|\lambda|}|x-z|}}{|x-z|^{n-2}} \frac{1}{|z-y_i|^{n-2+\beta}} dz + \int_{\Omega_2} \dots =: I_1 + I_2.$$

For I₁, from the fact that, for any $z \in \Omega_1$, $2|z - y_i| \ge |x - z| + |z - y_j| \ge |x - y_i|$, we deduce that

(3.10)
$$I_{1} \leq \frac{2}{|x - y_{i}|^{n-2+\beta}} \int_{\Omega_{1}} \frac{e^{-c\sqrt{|\lambda|}|x - z|}}{|x - z|^{n-2}} dz \leq \frac{1}{|x - y_{i}|^{n-2+\beta}} \frac{1}{|\lambda|}$$

Moreover, it is easy to see that, for any $z \in \Omega_2$, $2|x - z| \ge |x - z| + |z - y_i| \ge |x - y_i|$. By this, we find that

(3.11)
$$I_{2} \leq \frac{2}{|x - y_{i}|^{n-2}} \int_{\Omega_{2}} \frac{e^{-c\sqrt{|\lambda|}|z - y_{i}|}}{|z - y_{i}|^{n-2+\beta}} dz \lesssim \frac{1}{|x - y_{i}|^{n-2}} \frac{1}{|\lambda|^{1-\beta/2}}.$$

To finish the proof of the present lemma, we fix x, y_1 , $y_2 \in \Omega$ with $x \neq y_i$ for any $i \in \{1, 2\}$, and, without loss of generally, we may assume that $|x - y_1| \leq |x - y_2|$. Thus, from (3.8), (3.9), (3.10), (3.11), and $|x - y_1| \leq |x - y_2|$, it follows that

$$(3.12) \qquad |G_{\lambda}(x, y_{1}) - G_{\lambda}(x, y_{2})| \\ \lesssim |y_{1} - y_{2}|^{\beta} \left[\frac{1}{|x - y_{1}|^{n-2+\beta}} + \frac{1}{|x - y_{2}|^{n-2+\beta}} \right] \\ + |y_{1} - y_{2}|^{\beta} \left[\frac{|\lambda|^{\beta/2}}{|x - y_{1}|^{n-2}} + \frac{|\lambda|^{\beta/2}}{|x - y_{2}|^{n-2}} \right] \\ \lesssim |y_{1} - y_{2}|^{\beta} \left[\frac{1}{|x - y_{1}|^{n-2+\beta}} + \frac{|\lambda|^{\beta/2}}{|x - y_{1}|^{n-2}} \right].$$

Furthermore, by (3.7) and $|x - y_1| \le |x - y_2|$, we conclude that

$$|G_{\lambda}(x,y_1) - G_{\lambda}(x,y_2)| \lesssim \sum_{i=1}^{2} e^{-c\sqrt{|\lambda|}|x-y_i|} \frac{1}{|x-y_i|^{n-2}} \lesssim e^{-c\sqrt{|\lambda|}|x-y_1|} \frac{1}{|x-y_1|^{n-2}},$$

which, together with (3.12), further implies that, for any $\delta \in (0, \beta)$,

$$\begin{split} &|G_{\lambda}(x,y_{1}) - G_{\lambda}(x,y_{2})| \\ &\lesssim \left\{ |y_{1} - y_{2}|^{\beta} \left[\frac{1}{|x - y_{1}|^{n-2+\beta}} + \frac{|\lambda|^{\beta/2}}{|x - y_{1}|^{n-2}} \right] \right\}^{\frac{\delta}{\beta}} \left[e^{-c \sqrt{|\lambda|}|x - y_{1}|} \frac{1}{|x - y_{1}|^{n-2}} \right]^{1-\frac{\delta}{\beta}} \\ &\lesssim e^{-\gamma \sqrt{|\lambda|}|x - y_{1}|} |y_{1} - y_{2}|^{\delta} \left[\frac{1}{|x - y_{1}|^{n-2+\beta}} + \frac{|\lambda|^{\beta/2}}{|x - y_{1}|^{n-2}} \right]^{\frac{\delta}{\beta}} \left[\frac{1}{|x - y_{1}|^{n-2}} \right]^{1-\frac{\delta}{\beta}} \\ &\sim e^{-\gamma \sqrt{|\lambda|}|x - y_{1}|} |y_{1} - y_{2}|^{\delta} \left[\frac{|x - y_{1}|^{-\beta}}{|x - y_{1}|^{n-2}} + \frac{|\lambda|^{\beta/2}}{|x - y_{1}|^{n-2}} \right]^{\frac{\delta}{\beta}} \left[\frac{1}{|x - y_{1}|^{n-2}} \right]^{1-\frac{\delta}{\beta}} \\ &\sim e^{-\gamma \sqrt{|\lambda|}|x - y_{1}|} |y_{1} - y_{2}|^{\delta} \frac{1}{|x - y_{1}|^{n-2}} \left[\frac{1 + |x - y_{1}|^{\beta} |\lambda|^{\beta/2}}{|x - y_{1}|^{\beta}} \right]^{\frac{\delta}{\beta}} \\ &\lesssim e^{-\gamma \sqrt{|\lambda|}|x - y_{1}|} |y_{1} - y_{2}|^{\delta} \frac{1}{|x - y_{1}|^{n-2+\delta}} \left[1 + |x - y_{1}|^{\delta} |\lambda|^{\delta/2} \right] \end{split}$$

SIBEI YANG AND DACHUN YANG

$$\sim e^{-\gamma \sqrt{|\lambda|}|x-y_1|} |y_1 - y_2|^{\delta} \left[\frac{1}{|x-y_1|^{n-2+\delta}} + \frac{|\lambda|^{\delta/2}}{|x-y_1|^{n-2}} \right]$$

This finishes the proof of Lemma 3.3.

Lemma 3.4. Let $n \ge 3$, Ω be a bounded NTA domain of \mathbb{R}^n , $\mu \in (\omega_0, \pi/2)$, and $\lambda \in \Gamma_{\pi-\mu}$. Then, for any $m \in \mathbb{N}$ with m > (n+2)/2, the operator $(L_D + \lambda I)^{-m}$ has a kernel $R_{\lambda,m}$ and there exists a constant $\delta \in (0, 1)$ such that, for any $x, y_1, y_2 \in \Omega$ with $x \ne y_i$ for $i \in \{1, 2\}$,

$$\left| R_{\lambda,m}(x,y_1) - R_{\lambda,m}(x,y_2) \right| \le C |\lambda|^{-m+\frac{n}{2}+\frac{\delta}{2}} |y_1 - y_2|^{\delta} \left| \max_{i \in \{1,2\}} e^{-c\sqrt{|\lambda|}|x-y_i|} \right|,$$

where *C* is a positive constant depending only on μ , δ , *n*, and Ω , and *c* is a positive constant depending only on μ and δ .

Proof. We prove the present lemma via borrowing some ideas from the proof of [26, Lemma 2.6]. Similarly to the proof of [27, Theorem 1], using (1.5), we find that, for any $m \in \mathbb{N}$ with m > n/2, $(L_D + \lambda I)^{-m}$ has a kernel $R_{\lambda,m}$ and there exists a positive constant κ such that, for any $x, y \in \Omega$ with $x \neq y$,

$$(3.13) \left| R_{\lambda,m}(x,y) \right| \lesssim |\lambda|^{-m+\frac{n}{2}} e^{-\kappa \sqrt{|\lambda|}|x-y|}$$

Let $m \in \mathbb{N}$ and m > (n+2)/2. Then, from $(L_D + \lambda I)^{-(m+1)} = (L_D + \lambda I)^{-m} (L_D + \lambda I)^{-1}$, we deduce that, for any $x, y \in \Omega$ with $x \neq y$,

(3.14)
$$R_{\lambda,m+1}(x,y) = \int_{\Omega} R_{\lambda,m}(x,z) G_{\lambda}(z,y) \, dz.$$

Therefore, by (3.13), (3.14), and Lemma 3.3, we find that there exists a $\delta \in (0, 1)$ such that, for any $x, y_1, y_2 \in \Omega$ with $x \neq y_i$ for any $i \in \{1, 2\}$,

$$(3.15) \qquad \begin{aligned} |R_{\lambda,m+1}(x,y_1) - R_{\lambda,m+1}(x,y_2)| \\ &\leq \int_{\Omega} \left| R_{\lambda,m}(x,z) \right| |G_{\lambda}(z,y_1) - G_{\lambda}(z,y_2)| dz \\ &\leq |\lambda|^{-m+\frac{n}{2}} |y_1 - y_2|^{\delta} \int_{\Omega} e^{-\kappa \sqrt{|\lambda|}|x-z|} e^{-\gamma \sqrt{|\lambda|}|z-y_1|} \frac{1}{|z-y_1|^{n-2+\delta}} dz \\ &+ |\lambda|^{-m+\frac{n}{2}} |y_1 - y_2|^{\delta} \int_{\Omega} e^{-\kappa \sqrt{|\lambda|}|x-z|} e^{-\gamma \sqrt{|\lambda|}|z-y_1|} \frac{|\lambda|^{\delta/2}}{|x-y_1|^{n-2}} dz \\ &+ |\lambda|^{-m+\frac{n}{2}} |y_1 - y_2|^{\delta} \int_{\Omega} e^{-\kappa \sqrt{|\lambda|}|x-z|} e^{-\gamma \sqrt{|\lambda|}|z-y_2|} \frac{1}{|z-y_2|^{n-2+\delta}} dz \\ &+ |\lambda|^{-m+\frac{n}{2}} |y_1 - y_2|^{\delta} \int_{\Omega} e^{-\kappa \sqrt{|\lambda|}|x-z|} e^{-\gamma \sqrt{|\lambda|}|z-y_2|} \frac{1}{|z-y_2|^{n-2+\delta}} dz \\ &+ |\lambda|^{-m+\frac{n}{2}} |y_1 - y_2|^{\delta} \int_{\Omega} e^{-\kappa \sqrt{|\lambda|}|x-z|} e^{-\gamma \sqrt{|\lambda|}|z-y_2|} \frac{|\lambda|^{\delta/2}}{|x-y_2|^{n-2}} dz \\ &=: II_1 + II_2 + II_3 + II_4. \end{aligned}$$

Let $c := \min\{\kappa, \gamma/2\}$, where κ is as in (3.13). From the fact that, for any $z \in \Omega$, $|x - z| + |z - y_1| \ge |x - y_1|$, it follows that

$$\int_{\Omega} e^{-\kappa \sqrt{|\lambda|}|x-z|} e^{-\gamma \sqrt{|\lambda|}|z-y_1|} \frac{1}{|z-y_1|^{n-2+\delta}} dz$$

$$\begin{split} &\leq \int_{\Omega} e^{-c \sqrt{|\lambda|} (|x-z|+|z-y_1|)} e^{-\frac{\gamma}{2} \sqrt{|\lambda|} |z-y_1|} \frac{1}{|z-y_1|^{n-2+\delta}} \, dz \\ &\leq e^{-c \sqrt{|\lambda|} |x-y_1|} \int_{\Omega} e^{-\frac{\gamma}{2} \sqrt{|\lambda|} |z-y_1|} \frac{1}{|z-y_1|^{n-2+\delta}} \, dz \\ &\lesssim e^{-c \sqrt{|\lambda|} |x-y_1|} |\lambda|^{-1+\frac{\delta}{2}} \int_{\Omega} e^{-\frac{\gamma}{2} |z-y_1|} \frac{1}{|z-y_1|^{n-2+\delta}} \, dz \\ &\lesssim e^{-c \sqrt{|\lambda|} |x-y_1|} |\lambda|^{-1+\frac{\delta}{2}}, \end{split}$$

which further implies that

(3.16)
$$II_1 \leq |\lambda|^{-(m+1)+\frac{n}{2}+\frac{\delta}{2}} |y_1 - y_2|^{\delta} e^{-c\sqrt{|\lambda|}|x-y_1|}.$$

Similarly to the estimation of (3.16), for any $j \in \{2, 3, 4\}$, we also have

$$II_j \leq |\lambda|^{-(m+1)+\frac{n}{2}+\frac{\delta}{2}}|y_1 - y_2|^{\delta}e^{-c\sqrt{|\lambda|}|x-y_1|}$$

which, combined with (3.16) and (3.15), implies that

$$\left| R_{\lambda, m+1}(x, y_1) - R_{\lambda, m+1}(x, y_2) \right| \lesssim |\lambda|^{-(m+1) + \frac{n}{2} + \frac{\delta}{2}} |y_1 - y_2|^{\delta} e^{-c \sqrt{|\lambda|} |x - y_1|}.$$

This finishes the proof of Lemma 3.4.

Lemma 3.5. Let $n \ge 3$, Ω be a bounded NTA domain of \mathbb{R}^n , the real-valued, bounded, and measurable matrix A satisfy (1.3), and L_D be as in (1.4). Then there exists a constant $\delta_0 \in (0, 1]$, depending only on n, A, and Ω , such that, for any given $\delta \in (0, \delta_0)$, there exist constants $C, c \in (0, \infty)$ such that, for any $t \in (0, \infty)$ and $x, y_1, y_2 \in \Omega$ with $|y_1 - y_2| \le \sqrt{t}/2$,

$$\left| K_t^{L_D}(x, y_1) - K_t^{L_D}(x, y_2) \right| \le \frac{C}{t^{n/2}} \left[\frac{|y_1 - y_2|}{\sqrt{t}} \right]^{\delta} \exp\left\{ -\frac{|x - y_1|^2}{ct} \right\}.$$

Proof. For any given $\mu \in (\pi/2, \pi - \omega_0)$ and $R \in (0, \infty)$, define

$$\Gamma_1 := \{ re^{-i\mu} : r \ge R \}, \quad \Gamma_2 := \{ Re^{-i\phi} : |\phi| \le \mu \},$$

$$\Gamma_3 := \{ re^{i\mu} : r \ge R \}, \quad \text{and} \quad \Gamma_R := \Gamma_1 \cup \Gamma_2 \cup \Gamma_3.$$

Let $m \in \mathbb{N}$ and $m \ge \frac{n+3}{2}$. Using the inverse Laplace transform (see, for instance, [52, p. 30, Theorem 7.7]), we find that, for any given $t \in (0, \infty)$ and any $x, y \in \Omega$ with $x \ne y$,

(3.17)
$$K_t^{L_D}(x,y) = (-1)^m \frac{(m-1)!}{2\pi i t^{m-1}} \int_{\Gamma_R} e^{\lambda t} R_{\lambda,m}(x,y) \, d\lambda,$$

where $R \in [R(x, y, t), \infty)$ and

(3.18)
$$R(x, y, t) := \max\left\{\frac{1}{t}, \frac{|x - y|^2}{t^2}\right\}.$$

Fix $x, y_1, y_2 \in \Omega$ with $x \neq y_i$ for any $i \in \{1, 2\}$, a $t \in (0, \infty)$, and an

$$R \in [\max \{R(x, y_1, t), R(x, y_2, t)\}, \infty),\$$

where $R(x, y_1, t)$ and $R(x, y_2, t)$ are as in (3.18). Then, by (3.17) and Lemma 3.4, we conclude that

(3.19)
$$\begin{aligned} \left| K_{t}^{L_{D}}(x, y_{1}) - K_{t}^{L_{D}}(x, y_{2}) \right| \\ &\lesssim \sum_{i=1}^{2} \frac{1}{t^{m-1}} |y_{1} - y_{2}|^{\delta} \int_{\Gamma_{R}} e^{\Re(\lambda t)} |\lambda|^{-m + \frac{n}{2} + \frac{\delta}{2}} e^{-c \sqrt{|\lambda|} |x - y_{i}|} d|\lambda| \\ &=: E + F, \end{aligned}$$

where $\delta \in (0, 1)$ is as in Lemma 3.4. From the assumption that $R \ge \max\{\frac{1}{t}, \frac{|x-y_1|^2}{t^2}\}$, it follows that there exists a positive constant \tilde{c} such that

$$(3.20) \qquad \frac{1}{t^{m-1}} \int_{\Gamma_{1} \cup \Gamma_{2}} e^{\Re(\lambda t)} |\lambda|^{-m+\frac{n}{2}+\frac{\delta}{2}} e^{-c\sqrt{|\lambda|}|x-y_{1}|} d|\lambda| \lesssim t^{-\frac{n+\delta}{2}} \int_{R}^{\infty} e^{-\overline{c}|\lambda|t} (|\lambda|t)^{-m+\frac{n}{2}+\frac{\delta}{2}+1} e^{-c\sqrt{|\lambda|}|x-y_{1}|} \frac{d|\lambda|}{|\lambda|} \lesssim t^{-\frac{n+\delta}{2}} e^{-c\sqrt{R}|x-y_{1}|} e^{-\frac{\widetilde{c}}{2}Rt} \int_{1}^{\infty} e^{-\frac{\widetilde{c}}{2}s} s^{-m+\frac{n}{2}+\frac{\delta}{2}} ds \lesssim t^{-\frac{n+\delta}{2}} e^{-c\sqrt{R}|x-y_{1}|} e^{-\frac{\widetilde{c}}{2}Rt} \lesssim t^{-\frac{n+\delta}{2}} e^{-c_{1}\frac{|x-y_{1}|^{2}}{t}},$$

where *c* is as in Lemma 3.4, and $c_1 := c + \frac{\tilde{c}}{2}$. Moreover, by the assumptions that $-m + \frac{n}{2} + \frac{\delta}{2} + 1 < 0$ and $R \ge \max\{\frac{1}{t}, \frac{|x-y_1|^2}{t^2}\}$ again, we find that

$$\begin{aligned} \frac{1}{t^{m-1}} & \int_{\Gamma_3} e^{\Re(\lambda t)} |\lambda|^{-m+\frac{n}{2}+\frac{\delta}{2}} e^{-c\sqrt{|\lambda|}|x-y_1|} d|\lambda| \\ & \lesssim t^{-\frac{n+\delta}{2}} \int_{\{\lambda \in \mathbb{C}: \ |\lambda|=R\}} e^{-\widetilde{c}Rt} (Rt)^{-m+\frac{n}{2}+\frac{\delta}{2}+1} e^{-c\sqrt{R}|x-y_1|} \frac{d|\lambda|}{|\lambda|} \\ & \lesssim t^{-\frac{n+\delta}{2}} (Rt)^{-m+\frac{n}{2}+\frac{\delta}{2}+1} e^{-c_1\frac{|x-y_1|^2}{t}} \lesssim t^{-\frac{n+\delta}{2}} e^{-c_1\frac{|x-y_1|^2}{t}}, \end{aligned}$$

which, together with (3.20), further implies that

(3.21)
$$\mathbf{E} \lesssim t^{-\frac{n}{2}} e^{-c_1 \frac{|x-y_1|^2}{t}} \left[\frac{|y_1 - y_2|}{t^{1/2}} \right]^{\delta}.$$

Similarly to the estimation of (3.21), we also have

$$\mathbf{F} \lesssim t^{-\frac{n}{2}} e^{-c_1 \frac{|x-y_2|^2}{t}} \left[\frac{|y_1 - y_2|}{t^{1/2}} \right]^{\delta}.$$

From this, (3.21), and (3.19), we deduce that, for any $t \in (0, \infty)$ and $x, y_1, y_2 \in \Omega$ satisfying $|y_1 - y_2| \leq \sqrt{t/2}$,

$$\left|K_{t}^{L_{D}}(x,y_{1})-K_{t}^{L_{D}}(x,y_{2})\right| \lesssim t^{-\frac{n}{2}} \left(\max_{i\in\{1,2\}} e^{-c_{1}\frac{|x-y_{i}|^{2}}{t}}\right) \left[\frac{|y_{1}-y_{2}|}{t^{1/2}}\right]^{\delta} \lesssim t^{-\frac{n}{2}} e^{-c_{1}\frac{|x-y_{1}|^{2}}{t}} \left[\frac{|y_{1}-y_{2}|}{t^{1/2}}\right]^{\delta}.$$

This finishes the proof of Lemma 3.5.

To show Theorem 1.1, we also need the following global regularity estimate for the Dirichlet problem (1.1).

Lemma 3.6. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, and the real-valued, bounded, and measurable matrix A satisfy (1.3). Then there exists a positive constant $p_0 \in (2, \infty)$, depending only on n, A, and Ω , such that, for any given $p \in [2, p_0)$, the Dirichlet problem (1.1), with $f \in L^{p_*}(\Omega)$, is uniquely solvable and, moreover, for any weak solution u of the problem (1.1), $u \in W_0^{1,p}(\Omega)$ and there exists a positive constant C, depending only on n, p, and Ω , such that

$$(3.22) ||\nabla u||_{L^p(\Omega;\mathbb{R}^n)} \le C||f||_{L^{p_*}(\Omega)}.$$

To prove Lemma 3.6, we need the global regularity estimate for the following Dirichlet problem (3.23).

Let $p \in (1, \infty)$ and $\mathbf{f} \in L^p(\Omega; \mathbb{R}^n)$. Then a function *u* is called a *weak solution* of the following *Dirichlet boundary value problem*

(3.23)
$$\begin{cases} -\operatorname{div} (A\nabla u) = \operatorname{div} (\mathbf{f}) & \text{in } \Omega, \\ u = 0 & \text{on } \partial \Omega \end{cases}$$

if $u \in W_0^{1,p}(\Omega)$ and, for any $\varphi \in C_c^{\infty}(\Omega)$,

(3.24)
$$\int_{\Omega} A(x)\nabla u(x) \cdot \nabla \varphi(x) \, dx = -\int_{\Omega} \mathbf{f}(x) \cdot \nabla \varphi(x) \, dx.$$

Moreover, the Dirichlet problem (3.23) is said to be *uniquely solvable* if, for any $\mathbf{f} \in L^p(\Omega; \mathbb{R}^n)$, there exists a unique $u \in W_0^{1,p}(\Omega)$ such that (3.24) holds true for any $\varphi \in C_c^{\infty}(\Omega)$.

By the Lax–Milgram theorem, we conclude that, when p = 2, the Dirichlet problem (3.23), with $\mathbf{f} \in L^2(\Omega; \mathbb{R}^n)$, is uniquely solvable and the weak solution u satisfies

$$\|\nabla u\|_{L^2(\Omega;\mathbb{R}^n)} \le \mu_0^{-1} \|\mathbf{f}\|_{L^2(\Omega;\mathbb{R}^n)},$$

where μ_0 is as in (1.3).

Lemma 3.7. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, and the real-valued, bounded, and measurable matrix A satisfy (1.3). Then there exists a positive constant $p_0 \in (2, \infty)$, depending only on n, A, and Ω , such that, for any given $p \in (p'_0, p_0)$ with $1/p'_0 + 1/p_0 = 1$, the Dirichlet problem (3.23), with $\mathbf{f} \in L^p(\Omega; \mathbb{R}^n)$, is uniquely solvable and, moreover, for any weak solution u of the problem (3.23), $u \in W_0^{1,p}(\Omega)$ and there exists a positive constant C, depending only on n, p, and Ω , such that

$$\|\nabla u\|_{L^p(\Omega;\mathbb{R}^n)} \le C \|\mathbf{f}\|_{L^p(\Omega;\mathbb{R}^n)}$$

Via replacing [69, Lemma 4.3] by the following Lemma 3.8, using a real-variable argument for $L^{p}(\Omega)$ estimates, which was essentially established by Shen [58, Theorem 3.4] (see also [57, Theorem 4.2.6] and [59, Theorem 3.3]), and then repeating the proof of [69, Theorem 1.10], we can prove Lemma 3.7; we omit the details here.

Lemma 3.8. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, and $B(x_0, r)$ a ball such that $r \in (0, r_0/4)$ and either $x_0 \in \partial \Omega$ or $B(x_0, 2r) \subset \Omega$, where $r_0 \in (0, \operatorname{diam}(\Omega))$ is a constant. Assume that the real-valued, bounded, and measurable matrix A satisfies (1.3), and $u \in W^{1,2}(B_{\Omega}(x_0, 2r))$ is a weak solution of the following Dirichlet problem

$$\begin{cases} \operatorname{div} (A\nabla u) = 0 & \text{ in } B_{\Omega}(x_0, 2r), \\ u = 0 & \text{ on } B(x_0, 2r) \cap \partial \Omega \end{cases}$$

Then there exists a constant $p_0 \in (2, \infty)$, depending on Ω , n, and μ_0 in (1.3), such that

$$\left[\int_{B_{\Omega}(x_0,r)} |\nabla u(x)|^{p_0} dx\right]^{1/p_0} \le C \left[\int_{B_{\Omega}(x_0,2r)} |\nabla u(x)|^2 dx\right]^{1/2},$$

where *C* is a positive constant depending only on *n*, Ω , and p_0 .

Lemma 3.8 was established in [47, Lemma 3.2 and Corollary 4.1]. Now, we show Lemma 3.6 by using Lemma 3.7.

Proof of Lemma 3.6. Let $p_0 \in (2, \infty)$ be as in Lemma 3.8, $p \in [2, p_0)$, and $f \in L^{p_*}(\Omega)$. Consider the Dirichlet problem

(3.25)
$$\begin{cases} -\operatorname{div} (A^* \nabla v) = \operatorname{div} (\mathbf{g}) & \text{in } \Omega, \\ v = 0 & \text{on } \partial \Omega \end{cases}$$

with $\mathbf{g} \in L^{p'}(\Omega; \mathbb{R}^n)$, where $\frac{1}{p} + \frac{1}{p'} = 1$. Then, by Lemma 3.7 and $\mathbf{g} \in L^{p'}(\Omega; \mathbb{R}^n)$ with $p' \in (p'_0, p_0)$, we conclude that the Dirichlet problem (3.25) is uniquely solvable and the weak solution v satisfies

$$(3.26) \|\nabla v\|_{L^{p'}(\Omega;\mathbb{R}^n)} \lesssim \|\mathbf{g}\|_{L^{p'}(\Omega;\mathbb{R}^n)}.$$

Moreover, from the assumptions that $p \ge 2$ and $f \in L^{p_*}(\Omega)$, and Remark 1.7, we deduce that there exists a weak solution $u \in W_0^{1,2}(\Omega)$ for the Dirichlet problem (1.1) with $f \in L^{p_*}(\Omega)$. Moreover, we have

$$\int_{\Omega} \nabla u(x) \cdot \mathbf{g}(x) \, dx = -\int_{\Omega} A^*(x) \nabla v(x) \cdot \nabla u(x) \, dx$$
$$= -\int_{\Omega} A(x) \nabla u(x) \cdot \nabla v(x) \, dx = -\int_{\Omega} f(x) v(x) \, dx,$$

which, together with the Hölder inequality, the Sobolev embedding theorem on NTA domains (see, for instance, [5, Theorem 1.1]), and (3.26), further implies that

$$\begin{split} \|\nabla u\|_{L^{p}(\Omega;\mathbb{R}^{n})} &= \sup_{\|\mathbf{g}\|_{L^{p'}(\Omega;\mathbb{R}^{n})} \leq 1} \left| \int_{\Omega} \nabla u(x) \cdot \mathbf{g}(x) \, dx \right| = \sup_{\|\mathbf{g}\|_{L^{p'}(\Omega;\mathbb{R}^{n})} \leq 1} \left| \int_{\Omega} f(x) v(x) \, dx \right| \\ &\leq \sup_{\|\mathbf{g}\|_{L^{p'}(\Omega;\mathbb{R}^{n})} \leq 1} \|f\|_{L^{p_{\ast}}(\Omega)} \|v\|_{L^{(p_{\ast})'}(\Omega)} \\ &\lesssim \sup_{\|\mathbf{g}\|_{L^{p'}(\Omega;\mathbb{R}^{n})} \leq 1} \|f\|_{L^{p_{\ast}}(\Omega)} \|\nabla v\|_{L^{p'}(\Omega;\mathbb{R}^{n})} \end{split}$$

$$\lesssim \sup_{\|\mathbf{g}\|_{L^{p'}(\Omega;\mathbb{R}^n)} \leq 1} \|f\|_{L^{p_*}(\Omega)} \|\mathbf{g}\|_{L^{p'}(\Omega;\mathbb{R}^n)} \lesssim \|f\|_{L^{p_*}(\Omega)}.$$

By this and Remark 1.7, we find that the Dirichlet problem (1.1) with $f \in L^{p_*}(\Omega)$ is uniquely solvable and (3.22) holds true. This finishes the proof of Lemma 3.6.

Furthermore, to prove Theorem 1.1, we also need the following global gradient estimate for the problem (1.1) on (γ , σ , R) quasi-convex domains.

Lemma 3.9. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, and the real-valued, bounded, and measurable matrix A satisfy (1.3). Assume that $p \in (2, \infty)$. Then there exists a positive constant $\gamma_0 \in (0, 1)$, depending only on n, p, and Ω , such that, if A satisfies the (γ, R) -BMO condition and Ω is a (γ, σ, R) quasi-convex domain for some $\gamma \in (0, \gamma_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, then the Dirichlet problem (1.1), with $f \in L^{p_*}(\Omega)$, is uniquely solvable and, moreover, for any weak solution u of the problem (1.1), $u \in W_0^{1,p}(\Omega)$ and there exists a positive constant C, depending only on n, p, and Ω , such that

$$\|\nabla u\|_{L^p(\Omega;\mathbb{R}^n)} \le C \|f\|_{L^{p_*}(\Omega)}$$

Lemma 3.10. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, and $r_0 \in (0, \operatorname{diam}(\Omega))$. Assume that the matrix A is as in Lemma 3.9. Let $v \in W^{1,2}(B_{\Omega}(x_0, 4r))$ be a weak solution of the equation $\operatorname{div}(A\nabla v) = 0$ in $B_{\Omega}(x_0, 4r)$ satisfying v = 0 on $B(x_0, 4r) \cap \partial\Omega$, where $r \in (0, r_0/4)$ and either $x_0 \in \partial\Omega$ or $B(x_0, 4r) \subset \Omega$. Then, for any given $p \in (2, \infty)$, there exists a constant $\gamma_0 \in (0, 1)$, depending only on n, p, and Ω , such that, if A satisfies the (γ, R) -BMO condition and Ω is a (γ, σ, R) quasi-convex domain for some $\gamma \in (0, \gamma_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, then the weak reverse Hölder inequality

$$\left[\int_{B_{\Omega}(x_0,2r)} |\nabla v(x)|^p \, dx\right]^{\frac{1}{p}} \le C \left[\int_{B_{\Omega}(x_0,4r)} |\nabla v(x)|^2 \, dx\right]^{\frac{1}{2}}$$

holds true, where C is a positive constant independent of v, x_0 , and r.

Lemma 3.10 was established in [69].

Proof of Lemma 3.9. Via replacing Lemma 3.8 by Lemma 3.10, and repeating the proof of Lemma 3.6, we prove the present lemma, which completes the proof of Lemma 3.9.

Finally, we prove Theorem 1.1 by using Lemmas 3.5, 3.6, and 3.9.

Proof of Theorem 1.1. We first show (i). When $n \ge 3$, by Lemma 3.5, we find that the desired conclusion of (i) holds true in this case.

Now, we assume that n = 2. Denote by L_D^* the adjoint operator of L_D . Then $L_D^* = -\text{div}(A^*\nabla \cdot)$, where A^* is the transpose of the matrix A. Denote by $\{e^{-tL_D^*}\}_{t>0}$ the semigroup generated by L_D^* . Then, for any $t \in (0, \infty)$, $(e^{-tL_D})^* = e^{-tL_D^*}$ (see, for instance, [52, p. 41, Corollary 10.6]), where $(e^{-tL_D})^*$ denotes the adjoint operator e^{-tL_D} , which implies that, for any $t \in (0, \infty)$ and $x, y \in \Omega$, $K_t^{L_D}(x, y) = K_t^{L_D^*}(y, x)$. Here and thereafter, $\{K_t^{L_D^*}\}_{t>0}$ denote the kernels of the semigroup $\{e^{-tL_D^*}\}_{t>0}$. Thus, to prove (i) in the case of n = 2, it suffices to show that there exists a $\delta_0 \in (0, 1]$ such that, for any given $\delta \in (0, \delta_0)$, there exists a constant $c \in (0, \infty)$ such that, for any $t \in (0, \infty)$ and $x, y_1, y_2 \in \Omega$ with $|y_1 - y_2| \le \sqrt{t/2}$,

(3.27)
$$\left| K_t^{L_D^*}(y_1, x) - K_t^{L_D^*}(y_2, x) \right| \lesssim \frac{1}{t^{n/2}} \left[\frac{|y_1 - y_2|}{\sqrt{t}} \right]^{\delta} \exp\left\{ -\frac{|x - y_1|^2}{ct} \right\}.$$

For any given $t \in (0, \infty)$ and $x \in \Omega$, let $u(\cdot) := K_t^{L_D^*}(\cdot, x)$ and $f(\cdot) := -\frac{d}{dt}K_t^{L_D^*}(\cdot, x)$. From the facts that, for any $t \in (0, \infty)$ and $x \in \Omega$, $K_t^{L_D^*}(\cdot, x)$, $\frac{d}{dt}K_t^{L_D^*}(\cdot, x) \in \mathcal{D}(L_D^*) \subset W_0^{1,2}(\Omega)$, and $L_D^*e^{-tL_D^*}g = -\frac{d}{dt}e^{-tL_D^*}g$ for any $g \in L^2(\Omega)$, we deduce that

(3.28)
$$\begin{cases} \operatorname{div} (A^* \nabla u) = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial \Omega. \end{cases}$$

Let $p_0 \in (2, \infty)$ be as in Lemma 3.6, and $\delta_0 := 1 - 2/p_0$. For any given $\delta \in (0, \delta_0)$, take $\delta_1 \in (\delta, \delta_0)$ sufficiently large such that there exists a $p_1 \in (2, p_0)$ satisfying $\delta_1 := 1 - 2/p_1$. By (3.28) and Lemma 3.6, we conclude that

(3.29)
$$\left\|\nabla K_t^{L_D^*}(\cdot, x)\right\|_{L^{p_1}(\Omega; \mathbb{R}^n)} \lesssim \left\|\frac{d}{dt} K_t^{L_D^*}(\cdot, x)\right\|_{L^{(p_1)_*}(\Omega)}$$

Then, from n = 2, the Sobolev embedding theorem (see, for instance, [31, Theorem 7.26]), the Poincaré inequality (see, for instance, [5, Theorem 1.1]), and (3.29), it follows that $K_t^{L_D^*}(\cdot, x) \in C^{0,\delta_1}(\overline{\Omega})$ and, for any $t \in (0, \infty)$ and $x \in \Omega$,

$$(3.30) \left\| K_t^{L_D^*}(\cdot, x) \right\|_{C^{0,\delta_1}(\overline{\Omega})} \lesssim \left\| K_t^{L_D^*}(\cdot, x) \right\|_{W^{1,p_1}(\Omega)} \lesssim \left\| \nabla K_t^{L_D^*}(\cdot, x) \right\|_{L^{p_1}(\Omega;\mathbb{R}^n)} \\ \lesssim \left\| \frac{d}{dt} K_t^{L_D^*}(\cdot, x) \right\|_{L^{(p_1)*}(\Omega)}.$$

Here, $C^{0,\delta_1}(\overline{\Omega})$ denotes the *Hölder space* on Ω , which is defined by setting

$$C^{0,\delta_1}(\overline{\Omega}) := \left\{ g \text{ is bounded and continuous on } \Omega : \|g\|_{C^{0,\delta_1}(\overline{\Omega})} < \infty \right\}$$

with

$$||g||_{C^{0,\delta_1}(\overline{\Omega})} := \sup_{x, y \in \Omega, x \neq y} \frac{|g(x) - g(y)|}{|x - y|^{\delta_1}}.$$

Furthermore, by [51, Theorem 6.17] and (1.5), we find that, for any $t \in (0, \infty)$ and $x, y \in \Omega$,

$$\left|\frac{d}{dt}K_t^{L_D^*}(x,y)\right| \lesssim \frac{1}{t^{1+n/2}} \exp\left\{-\frac{|x-y|^2}{ct}\right\},\,$$

which further implies that

(3.31)
$$\left\|\frac{d}{dt}K_{t}^{L_{D}^{*}}(\cdot,x)\right\|_{L^{(p_{1})_{*}}(\Omega)} \lesssim \left[\int_{\Omega} \frac{1}{t^{1+n/2}} e^{-\frac{(p_{1})_{*}|x-y|^{2}}{ct}} dy\right]^{1/(p_{1})_{*}} \lesssim \frac{1}{t^{1+\frac{n}{2}[1-\frac{1}{(p_{1})_{*}}]}}.$$

From (3.30), (3.31), and $\frac{1}{(p_1)_*} - \frac{1}{p_1} = \frac{1}{n}$, we deduce that, for any $t \in (0, \infty)$ and $x \in \Omega$,

(3.32)
$$\left\|K_{t}^{L_{D}^{*}}(\cdot, x)\right\|_{C^{0,\delta_{1}}(\overline{\Omega})} \lesssim \frac{1}{t^{n/2}} \frac{1}{t^{\frac{1}{2}(1-\frac{n}{p_{1}})}} \sim \frac{1}{t^{(n+\delta_{1})/2}}$$

Thus, by (3.32), we conclude that, for any $t \in (0, \infty)$ and $x, y_1, y_2 \in \Omega$,

(3.33)
$$\left| K_t^{L_D^*}(y_1, x) - K_t^{L_D^*}(y_2, x) \right| \lesssim \frac{1}{t^{n/2}} \left[\frac{|y_1 - y_2|}{t^{1/2}} \right]^{\delta_1}$$

On the other hand, from (1.5), it follows that, for any $t \in (0, \infty)$ and $x, y_1, y_2 \in \Omega$ with $|y_1 - y_2| \le \sqrt{t/2}$,

$$\left|K_{t}^{L_{D}^{*}}(y_{1},x) - K_{t}^{L_{D}^{*}}(y_{2},x)\right| \leq \left|K_{t}^{L_{D}^{*}}(y_{1},x)\right| + \left|K_{t}^{L_{D}^{*}}(y_{2},x)\right| \lesssim \frac{1}{t^{n/2}} \exp\left\{-\frac{|x-y_{1}|^{2}}{ct}\right\},$$

which, together with (3.33), further implies that

$$\left|K_t^{L_D^*}(y_1, x) - K_t^{L_D^*}(y_2, x)\right| \lesssim \frac{1}{t^{n/2}} \left[\frac{|y_1 - y_2|}{t^{1/2}}\right]^{\delta} \exp\left\{-\frac{|x - y_1|^2}{ct}\right\}.$$

This finishes the proof of (3.27), and hence of (i).

Next, we show (ii). To prove this, similarly to the proof of (i) in the case of n = 2, it suffices to show that, for any given $\delta_0 \in (0, 1)$, there exists a constant $\gamma_0 \in (0, \infty)$, depending on δ_0 , n, and Ω , such that, if A satisfies the (γ, R) -BMO condition and Ω is a (γ, σ, R) quasi-convex domain for some $\gamma \in (0, \gamma_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, then, for any given $\delta \in (0, \delta_0)$, there exists a positive constant $c \in (0, \infty)$ such that, for any $t \in (0, \infty)$ and $x, y_1, y_2 \in \Omega$ with $|y_1 - y_2| \le \sqrt{t}/2$,

(3.34)
$$\left| K_t^{L_D^*}(y_1, x) - K_t^{L_D^*}(y_2, x) \right| \lesssim \frac{1}{t^{n/2}} \left[\frac{|y_1 - y_2|}{\sqrt{t}} \right]^{\delta} \exp\left\{ -\frac{|x - y_1|^2}{ct} \right\}.$$

For any given $\delta_0 \in (0, 1)$, let $p \in (n, \infty)$ be such that $\delta_0 = 1 - n/p$. By Lemma 3.9, we find that there exists a constant $\gamma_0 \in (0, 1)$, depending on p, n, and Ω , such that, if A satisfies the (γ, R) -BMO condition and Ω is a (γ, σ, R) quasi-convex domain for some $\gamma \in (0, \gamma_0)$, $\sigma \in (0, 1)$, and $R \in (0, \infty)$, then

(3.35)
$$\left\|\nabla K_t^{L_D^*}(\cdot, x)\right\|_{L^p(\Omega; \mathbb{R}^n)} \lesssim \left\|\frac{d}{dt} K_t^{L_D^*}(\cdot, x)\right\|_{L^{p_*}(\Omega)}$$

Replacing (3.29) by (3.35), and repeating the estimation of (3.27), we conclude that (3.34) holds true. This finishes the proof of (ii), and hence of Theorem 1.1. \Box

4 **Proof of Theorem 1.5**

In this section, we show Theorem 1.5. We begin with recalling some concepts on the tent space $T^p(\Omega \times (0, \infty))$ (see, for instance, [54]).

Definition 4.1. Let $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain and $p \in (0, \infty)$. Then the *tent space* $T^p(\Omega \times (0, \infty))$ is defined to be the set of all measurable functions f on $\Omega \times (0, \infty)$ such that $\|f\|_{T^p(\Omega \times (0,\infty))} := \|\mathcal{A}(f)\|_{L^p(\Omega)} < \infty$, where, for any $x \in \Omega$,

$$\mathcal{A}(f)(x) := \left[\int_0^\infty \int_{\Gamma(x)} |f(y,t)|^2 \frac{dy \, dt}{|B_\Omega(x,t)|t|}\right]^{1/2}$$

where $\Gamma(x) := \{y \in \Omega : |x - y| < t\}.$

Moreover, assume that O is an open subset of Ω . Then the *tent* over O, denoted by T(O), is defined by setting

$$T(O) := \left\{ (x, t) \in \Omega \times (0, \infty) : \operatorname{dist} \left(x, O^{\mathbb{U}} \cap \Omega \right) \ge t \right\}.$$

Let $p \in (0, 1]$ and $q \in (1, \infty)$. A measurable function a on $\Omega \times (0, \infty)$ is called a (p, q)-atom if there exists a ball B_{Ω} of Ω , which means $B_{\Omega} := B \cap \Omega$ and $B := B(x_B, r_B)$ is a ball of \mathbb{R}^n with $x_B \in \Omega$ and $r_B \in (0, \text{diam}(\Omega))$, such that

- (i) supp (a) := { $(x, t) \in \Omega \times (0, \infty)$: $a(x, t) \neq 0$ } $\subset T(B_{\Omega})$;
- (ii) $||a||_{T^q(\Omega \times (0,\infty))} \le |B_{\Omega}|^{1/q-1/p}$.

Then we have the following atomic decomposition of $T^p(\Omega \times (0, \infty))$, which is a special case of [54, Theorem 1.1].

Lemma 4.2. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, $p \in (0, 1]$, and $q \in (1, \infty)$. Then, for any $f \in T^p(\Omega \times (0, \infty))$, there exist $\{\lambda_j\}_{j \in \mathbb{N}} \subset \mathbb{C}$ and a sequence $\{a_j\}_{j \in \mathbb{N}}$ of (p, q)-atoms such that, for almost every $(x, t) \in \Omega \times (0, \infty)$, $f(x, t) = \sum_{i \in \mathbb{N}} \lambda_i a_i(x, t)$ and

$$C^{-1}||f||_{T^p(\Omega\times(0,\infty))} \leq \left(\sum_{j=1}^\infty |\lambda_j|^p\right)^{1/p} \leq C||f||_{T^p(\Omega\times(0,\infty))},$$

where *C* is a positive constant independent of *f*. Moreover, if $f \in T^p(\Omega \times (0, \infty)) \cap T^2(\Omega \times (0, \infty))$, then $f(x, t) = \sum_{j \in \mathbb{N}} \lambda_j a_j(x, t)$ holds true in both $T^p(\Omega \times (0, \infty))$ and $T^2(\Omega \times (0, \infty))$.

Next, we prove Theorem 1.5 by using Lemmas 2.2, 2.9, 2.11, 2.13, and 4.2.

Proof of Theorem 1.5. Let $p \in (\frac{n}{n+1}, 1]$. We first prove that

(4.1)
$$H_r^p(\Omega) = H^p(\Omega)$$

with equivalent quasi-norms. Let $f \in H_r^p(\Omega)$. Then there exists an $F \in H^p(\mathbb{R}^n)$ such that $F|_{\Omega} = f$ and

(4.2)
$$||f||_{H^p_r(\Omega)} \sim ||F||_{H^p(\mathbb{R}^n)}.$$

Moreover, by Lemma 2.9, we find that there exist a sequence $\{a_j\}_{j\in\mathbb{N}}$ of $(p, \infty, 0)$ -atoms, and $\{\lambda_j\}_{j\in\mathbb{N}} \subset \mathbb{C}$ satisfying $\sum_{j=1}^{\infty} |\lambda_j|^p < \infty$ such that $F = \sum_{j=1}^{\infty} \lambda_j a_j$ in $\mathcal{S}'(\mathbb{R}^n)$, and

(4.3)
$$||F||_{H^p(\mathbb{R}^n)} \sim \left(\sum_{j=1}^{\infty} |\lambda_j|^p\right)^{1/p}.$$

To show $f \in H^p(\Omega)$ and $||f||_{H^p(\Omega)} \leq ||f||_{H^p_r(\Omega)}$, it suffices to prove that, for any $(p, \infty, 0)$ -atom *a* supported in the ball $B := B(x_B, r_B) \subset \mathbb{R}^n$ with $x_B \in \mathbb{R}^n$ and $r_B \in (0, \infty)$, $a \in H^p(\Omega)$ and

$$(4.4) ||a||_{H^p(\Omega)} \leq 1$$

Indeed, if (4.4) holds true, from (4.2), (4.3), and (4.4), it follows that $f \in H^p(\Omega)$ and

$$||f||_{H^{p}(\Omega)}^{p} = ||F||_{H^{p}(\Omega)}^{p} \lesssim \sum_{j=1}^{\infty} |\lambda_{j}|^{p} ||a_{j}||_{H^{p}(\Omega)}^{p} \lesssim \sum_{j=1}^{\infty} |\lambda_{j}|^{p} \sim ||f||_{H^{p}_{r}(\Omega)}^{p},$$

which further implies that $||f||_{H^p(\Omega)} \leq ||f||_{H^p_r(\Omega)}$.

Now, we prove (4.4) by considering the following two cases on the ball *B*.

Case 1) $4B \subset \Omega$. In this case, by the definitions of $(p, \infty, 0)$ -atoms, we conclude that *a* is a type (a) $(p, \infty, 0)_{\Omega}$ -atom, which, combined with Lemma 2.11, implies that $a \in H^p(\Omega)$ and (4.4) holds true.

Case 2) $4B \cap \partial\Omega \neq \emptyset$. In this case, let ϕ be as in Definition 1.3(iv). We first claim that supp $(a_{\Omega}^{+}) \subset 8B_{\Omega}$. We prove this claim via borrowing some ideas from the proof of [7, Theorem 3.1]. Indeed, for any $x \in (8B_{\Omega})^{\mathbb{C}}$,

(4.5)
$$a_{\Omega}^{+}(x) = \sup_{t \in (0,\delta(x)/2)} \left| \int_{B_{\Omega}} \frac{1}{t^{n}} \phi\left(\frac{x-y}{t}\right) a(y) \, dy \right|$$
$$\leq \left[\sup_{t \in (0,7r_{B})} + \sup_{t \in (7r_{B},\delta(x)/2)} \right] \left| \int_{B_{\Omega}} \frac{1}{t^{n}} \phi\left(\frac{x-y}{t}\right) a(y) \, dy \right|$$
$$=: I_{1} + I_{2},$$

where $\delta(x) := \text{dist}(x, \partial\Omega)$ with $\partial\Omega$ being the boundary of Ω . From the fact that $|x - y| > 7r_B$ for any $y \in B_{\Omega}$ and $x \in (8B_{\Omega})^{\mathbb{C}}$, we deduce that, for any $t \in (0, 7r_B)$, $y \in B_{\Omega}$, and $x \in (8B_{\Omega})^{\mathbb{C}}$, $\phi(\frac{x-y}{t}) = 0$, which implies that $I_1 = 0$. Moreover, it is easy to see that, when $\delta(x) \le 14r_B$, $I_2 = 0$. We assume that $\delta(x) > 14r_B$. In this case, we find that, for any $y \in B_{\Omega}$ and $x \in (8B_{\Omega})^{\mathbb{C}}$ with $\delta(x) > 14r_B$,

$$(4.6) |x-y| \ge \delta(x) - \delta(y).$$

Since $4B \cap \partial \Omega \neq \emptyset$, it follows that, for any $y \in B_{\Omega}$, $\delta(y) \leq 4r_B$, which, together with (4.6) implies that, for any $y \in B_{\Omega}$ and $x \in (8B_{\Omega})^{\complement}$ with $\delta(x) > 14r_B$,

$$|x - y| \ge \delta(x) - \delta(y) \ge \delta(x) - 4r_B > \delta(x) - \delta(x)/2 = \delta(x)/2 > t.$$

By this, we conclude that, for any $t \in [7r_B, \delta(x)/2)$, $y \in B_\Omega$, and $x \in (8B_\Omega)^{\mathbb{C}}$, $\phi(\frac{x-y}{t}) = 0$, which further implies that $I_2 = 0$. This, combined with (4.5) and $I_1 = 0$, concludes that, for any $x \in (8B_\Omega)^{\mathbb{C}}$, $a_\Omega^+(x) = 0$. Thus, supp $(a_\Omega^+) \subset 8B_\Omega$.

Recall that the *Hardy–Littlewood maximal operator* \mathcal{M} on \mathbb{R}^n is defined by setting, for any $f \in L^1_{loc}(\mathbb{R}^n)$ and $x \in \mathbb{R}^n$,

$$\mathcal{M}(f)(x) := \sup_{B \ni x} \int_{B} |f(y)| \, dy,$$

where the supremum is taken over all balls $B \subset \mathbb{R}^n$ containing *x*. Then, from supp $(a_{\Omega}^+) \subset 8B_{\Omega}$, the fact that $a_{\Omega}^+ \leq \mathcal{M}(a)$, the Hölder inequality, and the boundedness of \mathcal{M} on $L^q(\mathbb{R}^n)$ with $q \in (1, \infty]$, we deduce that

$$\begin{aligned} \|a\|_{H^{p}(\Omega)} &= \left\|a_{\Omega}^{+}\right\|_{L^{p}(\Omega)} = \left\|a_{\Omega}^{+}\right\|_{L^{p}(8B_{\Omega})} \lesssim \|\mathcal{M}(a)\|_{L^{q}(8B_{\Omega})} |8B_{\Omega}|^{1/p-1/q} \\ &\lesssim \|a\|_{L^{q}(B_{\Omega})} |B_{\Omega}|^{1/p-1/q} \lesssim |B_{\Omega}|^{1/q-1/p} |B_{\Omega}|^{1/p-1/q} \sim 1. \end{aligned}$$

Therefore, (4.4) holds true.

Next, we assume that $f \in H^p(\Omega)$. By Lemma 2.11, we find that there exist two sequences $\{\lambda_j\}_{j=1}^{\infty} \subset \mathbb{C}$ and $\{\kappa_j\}_{j=1}^{\infty} \subset \mathbb{C}$, a sequence $\{a_j\}_{j=1}^{\infty}$ of type (a) $(p, \infty, 0)_{\Omega}$ -atoms, and a sequence $\{b_j\}_{j=1}^{\infty}$ of type (b) $(p, \infty)_{\Omega}$ -atoms such that

$$f = \sum_{j=1}^{\infty} \lambda_j a_j + \sum_{j=1}^{\infty} \kappa_j b_j$$

in $\mathcal{D}'(\Omega)$, and

(4.7)
$$||f||_{H^{p}(\Omega)} \sim \left(\sum_{j=1}^{\infty} |\lambda_{j}|^{p}\right)^{1/p} + \left(\sum_{j=1}^{\infty} |\kappa_{j}|^{p}\right)^{1/p}.$$

Let *b* be a type (b) $(p, \infty)_{\Omega}$ -atom supported in the ball $B := B(x_B, r_B) \subset \Omega$ with $x_B \in \Omega$ and $r_B \in (0, \infty)$. From Lemma 2.2(ii), it follows that there exists a ball $\widetilde{B}(x_{\widetilde{B}}, r_{\widetilde{B}}) \subset \Omega^{\mathbb{C}}$ with $x_{\widetilde{B}} \in \Omega^{\mathbb{C}}$ and $r_{\widetilde{B}} \in (0, \infty)$ such that $r_{\widetilde{B}} \sim r_B$ and dist $(B, \widetilde{B}) \sim r_B$. Assume that $B_0(x_{B_0}, r_{B_0}) \subset \mathbb{R}^n$ with $x_{B_0} \in \mathbb{R}^n$ and $r_{B_0} \in (0, \infty)$ is a ball such that $B \cup \widetilde{B} \subset B_0$ and $r_{B_0} \sim r_B$. Let

(4.8)
$$\widetilde{b} := b - \left[\frac{1}{|\widetilde{B}|} \int_{B} b(x) \, dx\right] \mathbf{1}_{\widetilde{B}}.$$

Then $\widetilde{b}|_{\Omega} = b$, supp $(\widetilde{b}) \subset B_0$, and $\int_{\mathbb{R}^n} \widetilde{b}(x) dx = 0$. Moreover, it is easy to see that

$$\left\|\widetilde{b}\right\|_{L^{\infty}(\mathbb{R}^n)} \leq \|b\|_{L^{\infty}(B)} \left[1 + \frac{|B|}{|\widetilde{B}|}\right] \leq \|b\|_{L^{\infty}(B)} \leq |B|^{-1/p} \sim |B_0|^{-1/p}.$$

Therefore, \tilde{b} is a harmless constant multiple of a $(p, \infty, 0)$ -atom supported in the ball B_0 .

For any $j \in \mathbb{N}$, let \tilde{b}_j be as in (4.8) with *b* replaced by b_j . Define

$$\widetilde{f} := \sum_{j=1}^{\infty} \lambda_j a_j + \sum_{j=1}^{\infty} \kappa_j \widetilde{b}_j.$$

Then $\widetilde{f}|_{\Omega} = f, \ \widetilde{f} \in H^p(\mathbb{R}^n)$, and

$$\left\|\widetilde{f}\right\|_{H^p(\mathbb{R}^n)} \lesssim \left(\sum_{j=1}^{\infty} |\lambda_j|^p\right)^{1/p} + \left(\sum_{j=1}^{\infty} |\kappa_j|^p\right)^{1/p},$$

which, together with (4.7), further implies that $f \in H_r^p(\Omega)$ and $||f||_{H_r^p(\Omega)} \leq ||f||_{H^p(\Omega)}$. This finishes the proof of (4.1).

Let $p \in (\frac{n}{n+\delta_0}, 1]$, where $\delta_0 \in (0, 1]$ is as in Theorem 1.1. Now, we prove that $H_r^p(\Omega) = H_{L_D}^p(\Omega)$ with equivalent quasi-norms. To this end, we first show that

(4.9)
$$\left[H_r^p(\Omega) \cap L^2(\Omega)\right] \subset \left[H_{L_D}^p(\Omega) \cap L^2(\Omega)\right].$$

Let $f \in [H_r^p(\Omega) \cap L^2(\Omega)]$. By the definition of $H_r^p(\Omega)$, we conclude that there exists an $\tilde{f} \in H^p(\mathbb{R}^n)$ such that $\tilde{f}|_{\Omega} = f$ and

(4.10)
$$\left\| \widetilde{f} \right\|_{H^p(\mathbb{R}^n)} \lesssim \|f\|_{H^p_r(\Omega)}$$

Then there exist a sequence $\{a_j\}_{j\in\mathbb{N}}$ of $(p, \infty, 0)$ -atoms, and $\{\lambda_j\}_{j\in\mathbb{N}} \subset \mathbb{C}$ such that

(4.11)
$$\widetilde{f} = \sum_{j \in \mathbb{N}} \lambda_j a_j$$

in both $\mathcal{S}'(\mathbb{R}^n)$ and $(H^p(\mathbb{R}^n))^*$, and

(4.12)
$$\left\| \widetilde{f} \right\|_{H^p(\mathbb{R}^n)} \sim \left(\sum_{j \in \mathbb{N}} |\lambda_j|^p \right)^{1/p},$$

where $(H^p(\mathbb{R}^n))^*$ denotes the dual space of $H^p(\mathbb{R}^n)$. Denote by $\{H_t^{L_D}\}_{t>0}$ the kernels of the family $\{tL_De^{-tL_D}\}_{t>0}$ of operators. From the fact that, for any $t \in (0, \infty)$,

$$tL_D e^{-tL_D} = 2\left(\frac{t}{2}L_D e^{-t/2L_D}\right)e^{-t/2L_D},$$

we deduce that, for any $t \in (0, \infty)$ and $x, y \in \Omega$,

(4.13)
$$H_t^{L_D}(x,y) = 2 \int_{\Omega} H_{t/2}^{L_D}(x,z) K_{t/2}^{L_D}(z,y) dz$$

Moreover, by [51, Theorem 6.17] and (1.5), we find that, for any $t \in (0, \infty)$ and $x, y \in \Omega$,

(4.14)
$$\left|H_t^{L_D}(x,y)\right| \lesssim \frac{1}{t^{n/2}} \exp\left\{-\frac{|x-y|^2}{ct}\right\},$$

which, combined with (4.13) and Theorem 1.1, implies that, for any given $\delta \in (0, \delta_0)$, and any $t \in (0, \infty)$ and $x, y, z \in \Omega$ with $|y - z| \le \sqrt{t/2}$,

(4.15)
$$\left| H_t^{L_D}(x,y) - H_t^{L_D}(x,z) \right| \lesssim \frac{1}{t^{n/2}} \left[\frac{|y-z|}{\sqrt{t}} \right]^{\delta} \exp\left\{ -\frac{|x-y|^2}{ct} \right\}.$$

For any $t \in (0, \infty)$ and $x \in \Omega$, denote by $\widetilde{H}_t^{L_D}(x, \cdot)$ the zero extension of $H_t^{L_D}(x, \cdot)$ from Ω to \mathbb{R}^n . Let $\gamma \in (0, \infty)$. Then the *Campanato space* $\mathcal{L}^{\gamma, 1, 0}(\mathbb{R}^n)$ is defined by setting

$$\mathcal{L}^{\gamma,\,1,\,0}(\mathbb{R}^n) := \left\{ g \in L^1_{\text{loc}}(\mathbb{R}^n) : \ \|g\|_{\mathcal{L}^{\gamma,\,1,\,0}(\mathbb{R}^n)} < \infty \right\},\,$$

where

$$||g||_{\mathcal{L}^{\gamma,1,0}(\mathbb{R}^n)} := \sup_{B \subset \mathbb{R}^n} \left[|B|^{-\gamma} \int_B |g(x) - g_B| \, dx \right]$$

with the supremum taken over all balls $B \subset \mathbb{R}^n$. Then, similarly to the proof of [14, Lemma 3.9], via applying Theorem 1.1 and Lemma 2.2, we find that, for any given $t \in (0, \infty), x \in \Omega$, and $\delta \in (0, \delta_0), \widetilde{H}_t^{L_D}(x, \cdot) \in \mathcal{L}^{\delta/n, 1, 0}(\mathbb{R}^n)$. From this, the fact that $\mathcal{L}_t^{\frac{1}{p}-1, 1, 0}(\mathbb{R}^n)$ is the dual space of $H^p(\mathbb{R}^n)$ (see, for instance, [61]), $p \in (\frac{n}{n+\delta_0}, 1]$, and (4.11), it follows that, for any $t \in (0, \infty)$ and $x \in \Omega$,

$$\begin{split} \int_{\Omega} H_{t^2}^{L_D}(x, y) f(y) \, dy &= \int_{\mathbb{R}^n} \widetilde{H}_{t^2}^{L_D}(x, y) \widetilde{f}(y) \, dy = \sum_{j \in \mathbb{N}} \lambda_j \int_{\mathbb{R}^n} \widetilde{H}_{t^2}^{L_D}(x, y) a_j(y) \, dy \\ &= \sum_{j \in \mathbb{N}} \lambda_j \int_{\Omega} H_{t^2}^{L_D}(x, y) a_j(y) \, dy, \end{split}$$

which, together with the boundedness of S_{L_D} on $L^2(\Omega)$ (see, for instance, [6, Theorem 2.13]), further implies that, for almost every $x \in \Omega$,

(4.16)
$$S_{L_D}(f)(x) \le \sum_{j \in \mathbb{N}} |\lambda_j| S_{L_D}(a_j)(x).$$

To prove (4.9), we only need to show that, for any $(p, \infty, 0)$ -atom a supported in the ball B := $B(x_B, r_B)$ with $x_B \in \mathbb{R}^n$ and $r_B \in (0, \infty)$,

(4.17)
$$\int_{\Omega} \left[S_{L_D}(a)(x) \right]^p \, dx \lesssim 1.$$

Indeed, if (4.17) holds true, then, by (4.16), (4.17), (4.10), (4.11), and (4.12), we conclude that $f \in H^p_{L_D}(\Omega)$ and

$$\begin{split} \|f\|_{H^p_{L_D}(\Omega)} &\lesssim \left\{ \sum_{j=1} |\lambda_j|^p \int_{\Omega} \left[S_{L_D}(a_j)(x) \right]^p \, dx \right\}^{1/p} \lesssim \left(\sum_{j \in \mathbb{N}} |\lambda_j|^p \right)^{1/p} \\ &\sim \left\| \widetilde{f} \right\|_{H^p(\mathbb{R}^n)} \lesssim \|f\|_{H^p(\Omega)}. \end{split}$$

Thus, (4.9) holds true.

Next, we show (4.17) by considering the following three cases on the ball *B*. *Case 1*) $B \cap \Omega = \emptyset$. In this case, we find that, for any $x \in \Omega$,

ise 1)
$$B \cap \Omega = \emptyset$$
. In this case, we find that, for any $x \in \Omega$,

$$t^{2}L_{D}e^{-t^{2}L_{D}}(a)(x) = \int_{B\cap\Omega} H_{t^{2}}^{L_{D}}(x, y)a(y)dy = 0,$$

which implies that $S_{L_D}(a) = 0$. From this, we deduce that (4.17) holds true in this case.

Case 2) $B \subset \Omega$. In this case, by the fact that $S_{L_{\Omega}}$ is bounded on $L^{q}(\Omega)$ with $q \in (1, \infty)$ (see, for instance, [6, Theorem 2.13]), we conclude that

(4.18)
$$\int_{4B_{\Omega}} \left[S_{L_{D}}(a)(x) \right]^{p} dx \leq \left\| S_{L_{D}}(a) \right\|_{L^{q}(4B_{\Omega})}^{p} \left| 4B_{\Omega} \right|^{1-\frac{p}{q}} \leq \left\| a \right\|_{L^{q}(4B_{\Omega})}^{p} \left| B \right|^{1-\frac{p}{q}} \leq 1.$$

Furthermore, for any $x \in (4B)^{\complement} \cap \Omega$, we have

(4.19)
$$[S_{L_D}(a)(x)]^2 = \int_0^{r_B} \int_{|y-x| < t} \left| t^2 L_D e^{-t^2 L_D}(a)(y) \right|^2 \frac{dy \, dt}{|B_\Omega(x,t)|t} + \int_{r_B}^\infty \int_{|y-x| < t} \cdots$$
$$=: E + F.$$

It is easy to see that, for any $x \in (4B)^{\mathbb{C}} \cap \Omega$, and any $t \in (0, \infty)$ and $y, z \in \Omega$ satisfying |x - y| < tand $|z - x_B| < r_B$,

(4.20)
$$t + |y - z| \ge t + |x - x_B| - |x - y| - |x_B - z| > |x - x_B| - r_B \ge \frac{3|x - x_B|}{4}.$$

Take $\delta_1 \in (0, \delta)$ such that $p > \frac{n}{n+\delta_1}$. From (4.14), (4.20), and the Hölder inequality, it follows that

$$(4.21) E = \int_{0}^{r_{B}} \int_{|y-x| < t} \left| \int_{B \cap \Omega} H_{t^{2}}^{L_{D}}(y, z) a(z) dz \right|^{2} \frac{dy dt}{|B_{\Omega}(x, t)|t} \\ \lesssim \int_{0}^{r_{B}} \int_{|y-x| < t} \left| \int_{B} \frac{1}{t^{n}} e^{-\frac{|y-z|^{2}}{ct^{2}}} a(z) dz \right|^{2} \frac{dy dt}{t^{n+1}} \\ \lesssim \int_{0}^{r_{B}} \int_{|y-x| < t} \left[\int_{B} \frac{t^{\delta_{1}}}{|y-z|^{n+\delta_{1}}} |a(z)| dz \right]^{2} \frac{dy dt}{t^{n+1}} \\ \lesssim \frac{1}{|x-x_{B}|^{2(n+\delta_{1})}} \left[\int_{B} |a(z)| dz \right]^{2} \int_{0}^{r_{B}} t^{2\delta_{1}-1} dt \\ \lesssim \left[\frac{r_{B}}{|x-x_{B}|} \right]^{2(n+\delta_{1})} |B|^{-2/p}.$$

Next, we deal with the term *F*. By $\int_B a(x) dx = 0$, (4.15), (4.20), $t > r_B$, and the Hölder inequality, we find that, for any $y \in \Omega$,

$$\begin{split} \left| t^{2} L_{D} e^{-t^{2} L_{D}}(a)(y) \right| \\ &\leq \int_{B} \left| H_{t^{2}}^{L_{D}}(y,z) - K_{t^{2}}^{L_{D}}(y,x_{B}) \right| |a(z)| \ dz \\ &\lesssim \int_{B} \frac{1}{t^{n}} \left[\frac{|z - x_{B}|}{t} \right]^{\delta} e^{-\frac{|y-z|^{2}}{ct^{2}}} |a(z)| \ dz \\ &\lesssim \int_{B} \left[\frac{r_{B}}{t} \right]^{\delta} \frac{t^{\delta_{1}}}{|y - z|^{n + \delta_{1}}} |a(z)| \ dz \\ &\lesssim \int_{B} \frac{r_{B}^{\delta}}{t^{\delta - \delta_{1}} |x - x_{B}|^{n + \delta_{1}}} |a(z)| \ dz \lesssim \frac{r_{B}^{n + \delta}}{t^{\delta - \delta_{1}} |x - x_{B}|^{n + \delta_{1}}} |B|^{-1/p}. \end{split}$$

From this, we deduce that

$$F \lesssim \frac{r_B^{2(n+\delta)} |B|^{-2/p}}{|x - x_B|^{2(n+\delta_1)}} \int_{r_B}^{\infty} \int_{|y - x| < t} t^{-2(\delta - \delta_1)} \frac{dy \, dt}{|B_{\Omega}(x, t)|t}$$

SIBEI YANG AND DACHUN YANG

$$\lesssim \frac{r_B^{2(n+\delta)}|B|^{-2/p}}{|x-x_B|^{2(n+\delta_1)}} \int_{r_B}^{\infty} t^{-2(\delta-\delta_1)-1} dt \lesssim \left[\frac{r_B}{|x-x_B|}\right]^{2(n+\delta_1)} |B|^{-2/p},$$

which, combined with (4.19) and (4.21), further implies that, for any $x \in (4B)^{\complement} \cap \Omega$,

$$S_{L_D}(a)(x) \lesssim \left[\frac{r_B}{|x-x_B|}\right]^{n+\delta_1} |B|^{-1/p}.$$

By this and $p > \frac{n}{n+\delta_1}$, we obtain

$$\begin{split} \int_{\Omega \setminus (4B_{\Omega})} \left[S_{L_D}(a)(x) \right]^p \, dx &= \sum_{j=2}^{\infty} \int_{S_j(B_{\Omega})} \left[S_{L_D}(a)(x) \right]^p \, dx \\ &\lesssim \sum_{j=2}^{\infty} \int_{S_j(B_{\Omega})} 2^{-(n+\delta_1)jp} |B|^{-1} \, dx \lesssim \sum_{j=2}^{\infty} 2^{-[(n+\delta_1)p-n]j} \lesssim 1, \end{split}$$

which, together with (4.18), further implies that (4.17) holds true in this case.

Case 3) $B \cap \Omega \neq \emptyset$. In this case, take $y_B \in B \cap \partial \Omega$. From the fact that, for any given $t \in (0, \infty)$ and $x \in \Omega$, $H_t^{L_D}(x, \cdot) \in W_0^{1,2}(\Omega)$, it follows that $H_t^{L_\Omega}(x, y_B) = 0$, which implies that, for any given $t \in (0, \infty)$ and any $x \in \Omega$,

$$t^{2}L_{D}e^{-t^{2}L_{D}}(a)(x) = \int_{B\cap\Omega} \left[H_{t^{2}}^{L_{D}}(x,y) - H_{t^{2}}^{L_{D}}(x,y_{B})\right]a(y)dy.$$

The remaining estimations are similar to those of Case 2), we omit the details here. This finishes the proof of (4.17), and hence of (4.9).

Now, we prove that

(4.22)
$$\left[H^p_{L_D}(\Omega) \cap L^2(\Omega)\right] \subset \left[H^p_r(\Omega) \cap L^2(\Omega)\right].$$

Let $f \in [H_{L_D}^p(\Omega) \cap L^2(\Omega)]$. Then, by the H^{∞} -functional calculus associated with L_{Ω} (see, for instance, [4, p. 8]), we find that

(4.23)
$$f = 8 \int_0^\infty \left(t^2 L_D e^{-t^2 L_D} \right) \left(t^2 L_D e^{-t^2 L_D} \right) (f) \frac{dt}{t}$$

in $L^2(\Omega)$. From the assumption $f \in [H^p_{L_D}(\Omega) \cap L^2(\Omega)]$, we deduce that $S_{L_D}(f) \in L^p(\Omega) \cap L^2(\Omega)$, which implies that

(4.24)
$$t^{2}L_{D}e^{-t^{2}L_{D}}(f) \in T^{p}(\Omega \times (0,\infty)) \cap T^{2}(\Omega \times (0,\infty))$$

and

(4.25)
$$\|f\|_{H^p_{L_D}(\Omega)} = \left\|t^2 L_D e^{-t^2 L_D}(f)\right\|_{T^p(\Omega \times (0,\infty))}$$

By (4.24) and Lemma 4.2, we conclude that there exist $\{\lambda_j\}_{j\in\mathbb{N}} \subset \mathbb{C}$ and a sequence $\{a_j\}_{j\in\mathbb{N}}$ of (p, 2)-atoms associated, respectively, with the balls $\{B_j \cap \Omega\}_{j\in\mathbb{N}}$ of Ω such that, for almost every $(x, t) \in \Omega \times (0, \infty)$,

(4.26)
$$t^2 L_D e^{-t^2 L_D}(f) = \sum_{j \in \mathbb{N}} \lambda_j a_j$$

and

(4.27)
$$\left\| t^2 L_D e^{-t^2 L_D}(f) \right\|_{T^p(\Omega \times (0,\infty))} \sim \left(\sum_{j=1}^\infty |\lambda_j|^p \right)^{1/p}.$$

For any $j \in \mathbb{N}$, let $\alpha_j := 8 \int_0^\infty t^2 L_\Omega e^{-t^2 L_\Omega} (a_j(\cdot, t)) \frac{dt}{t}$. Then, from (4.23) and (4.26), it follows that

(4.28)
$$f = \sum_{j \in \mathbb{N}} \lambda_j \alpha_j$$

in $L^2(\Omega)$.

For any (p, 2)-atom *a* associated with the ball B_{Ω} , let

$$\alpha := 8 \int_0^\infty t^2 L_D e^{-t^2 L_D}(a) \frac{dt}{t}.$$

To show (4.22), it suffices to prove that there exist a function $\tilde{\alpha}$ on \mathbb{R}^n such that

(4.29)
$$\widetilde{\alpha}|_{\Omega} = \alpha,$$

and a sequence $\{\kappa_i\}_i \subset \mathbb{C}$ and a sequence $\{b_i\}_i$ of (p, 2, 0)-atoms such that $\widetilde{\alpha} = \sum_i \kappa_i b_i$ in $L^2(\mathbb{R}^n)$, and

(4.30)
$$\sum_{i} |\kappa_i|^p \lesssim 1.$$

Indeed, if (4.29) and (4.30) hold true, then, by (4.28), we find that, for any $j \in \mathbb{N}$, there exists a function $\widetilde{\alpha}_j$ on \mathbb{R}^n such that $\widetilde{\alpha}_j|_{\Omega} = \alpha_j$. Let

$$\widetilde{f} := \sum_{j \in \mathbb{N}} \lambda_j \widetilde{\alpha}_j.$$

Then $\widetilde{f}|_{\Omega} = f$. Furthermore, from (4.30), we deduce that there exist a sequence $\{\kappa_{j,i}\}_{j \in \mathbb{N}, i} \subset \mathbb{C}$ and a sequence $\{b_{j,i}\}_{j \in \mathbb{N}, i}$ of (p, 2, 0)-atoms such that

$$\widetilde{f} = \sum_{j \in \mathbb{N}} \sum_{i} \lambda_{j} \kappa_{j,i} b_{j,i}$$

and

$$\sum_{j\in\mathbb{N},\,i}\left|\lambda_{j}\kappa_{j,\,i}\right|^{p}\lesssim\sum_{j\in\mathbb{N}}\left|\lambda_{j}\right|^{p}.$$

By this, Lemma 2.9, (4.25), and (4.27), we conclude that $\tilde{f} \in H^p(\mathbb{R}^n)$ and

$$\left\| \widetilde{f} \right\|_{H^{p}(\mathbb{R}^{n})} \sim \left\| \widetilde{f} \right\|_{H^{p,2,0}_{\mathrm{at}}(\mathbb{R}^{n})} \lesssim \| f \|_{H^{p}_{L_{D}}(\Omega)}.$$

Thus, $f \in H^p_r(\Omega)$ and

$$||f||_{H^p_r(\Omega)} \lesssim ||f||_{H^p_{L_D}(\Omega)},$$

which, combined with the arbitrariness of $f \in H^p_{L_D}(\Omega) \cap L^2(\Omega)$, implies that (4.22) holds true.

Next, we prove (4.29) and (4.30) by considering the following two cases on the ball $B := B(x_B, r_B)$, with $x_B \in \mathbb{R}^n$ and $r_B \in (0, \infty)$, which appears in the support of a.

Case 1) $8B \cap \Omega^{\complement} \neq \emptyset$. In this case, let $\widetilde{S}_0(B_\Omega) := \bigcup_{i=0}^2 S_i(B_\Omega)$ and

$$J_{\Omega} := \{k \in \mathbb{N} : k \ge 3, |S_k(B_{\Omega})| > 0\}.$$

Moreover, assume that $\mathbf{1}_0 := \mathbf{1}_{\widetilde{S}_0(B_\Omega)}, m_0 := \int_{\widetilde{S}_0(B_\Omega)} \alpha(x) dx$, and, for any $k \in J_\Omega, \mathbf{1}_k := \mathbf{1}_{S_k(B_\Omega)}$ and $m_k := \int_{S_k(B_\Omega)} \alpha(x) dx$. Then

$$\alpha = \alpha \mathbf{1}_0 + \sum_{k \in J_\Omega} \alpha \mathbf{1}_k$$

holds true almost everywhere and also in $L^2(\Omega)$. From the assumption $8B \cap \Omega^{\mathbb{C}} \neq \emptyset$, it follows that there exists a $y_B \in \partial \Omega$ such that $B(y_B, 16r_B) \supset 8B$. By this and Lemma 2.2(iii), we find that there exists a ball $\widetilde{B} \subset \Omega^{\mathbb{C}}$ such that $r_{\widetilde{B}} \sim r_B$ and dist $(B_\Omega, \widetilde{B}) \sim r_B$. Therefore, there exists a ball $B_0^* \subset \mathbb{R}^n$ such that $(B \cup \widetilde{B}) \subset B_0^*$ and

(4.31)
$$r_{B_0^*} \sim r_B$$

Let

$$b_0 := \alpha \mathbf{1}_0 - \left[\frac{1}{|\widetilde{B}|} \int_{\widetilde{S}_{(B_\Omega)}} \alpha(x) \, dx\right] \mathbf{1}_{\widetilde{B}}.$$

Then $\int_{\mathbb{R}^n} b_0(x) dx = 0$ and $\operatorname{supp}(b_0) \subset B_0^*$. Moreover, it is known that $||\alpha||_{L^2(\Omega)} \leq ||a||_{T^2(\Omega \times (0,\infty))}$ (see, for instance, [6, Proposition 4.5]), which further implies that

(4.32)
$$\|\alpha\|_{L^{2}(\Omega)} \leq \|a\|_{T^{2}(\Omega \times (0,\infty))} \leq |B_{\Omega}|^{1/2 - 1/p}$$

From this, $r_{\tilde{B}} \sim r_B$, Lemma 2.2(i), and (4.31), we deduce that

$$\begin{aligned} \|b_0\|_{L^2(\mathbb{R}^n)} &\leq \|\alpha\|_{L^2(\Omega)} + \|\alpha\|_{L^2(\Omega)} \left|\widetilde{B}\right|^{-1/2} |B|^{1/2} \\ &\lesssim \|\alpha\|_{L^2(\Omega)} \leq |B_\Omega|^{1/2 - 1/p} \sim \left|B_0^*\right|^{1/2 - 1/p} \end{aligned}$$

Therefore, b_0 is a harmless constant multiple of a (p, 2, 0)-atom.

For any $k \in J_{\Omega}$, by the definition of J_{Ω} , we find that $2^{k}r_{B} \leq \text{diam}(\Omega)$ and $2^{k}B \cap \partial\Omega \neq \emptyset$. By this, we conclude that there exists a $y_{B} \in \partial\Omega$ such that $B(y_{B}, 2^{k+1}r_{B}) \supset 2^{k}B$, which, together with Lemma 2.2(iii), further implies that there exists a ball $\widetilde{B}_{k} \subset \Omega^{\mathbb{C}}$ such that $r_{\widetilde{B}_{k}} \sim 2^{k}r_{B}$ and dist $(S_{k}(B_{\Omega}), \widetilde{B}_{k}) \leq 2^{k}r_{B}$. Let the ball $B_{k}^{*} \subset \mathbb{R}^{n}$ satisfy that $\widetilde{B}_{k} \cup S_{k}(B_{\Omega}) \subset B_{k}^{*}$ and $r_{B_{k}^{*}} \sim r_{2^{k}B}$. Assume that, for any $k \in J_{\Omega}$,

$$b_k := \alpha \mathbf{1}_k - \left[\frac{1}{|\widetilde{B}_k|} \int_{S_k(B_\Omega)} \alpha(x) \, dx\right] \mathbf{1}_{\widetilde{B}_k}.$$

Then, for any $k \in J_{\Omega}$, supp $(b_k) \subset B_k^*$ and $\int_{\mathbb{R}^n} b_k(x) dx = 0$. Furthermore, from the Hölder inequality and Lemma 2.2(i), it follows that

(4.33)
$$\|b_k\|_{L^2(\mathbb{R}^n)} \le \|\alpha\|_{L^2(S_k(B_\Omega))} + \|\alpha\|_{L^2(S_k(B_\Omega))} \left|\widetilde{B}_k\right|^{-1/2} |S_k(B_\Omega)|^{1/2}$$

 $\lesssim \|\alpha\|_{L^2(S_k(B_\Omega))}.$

Moreover, we have, for any $k \in J_{\Omega}$ with $k \ge 3$, and any $x \in S_k(B_{\Omega})$,

$$\begin{aligned} (4.34) \qquad |\alpha(x)| &\lesssim \int_{0}^{r_{B}} \int_{B_{\Omega}} \left| H_{t^{2}}^{L_{D}}(x,y) \right| |a(y,t)| \frac{dy \, dt}{t} \\ &\lesssim \int_{0}^{r_{B}} \int_{B_{\Omega}} \frac{1}{t^{n}} e^{-\frac{|x-y|^{2}}{ct^{2}}} |a(y,t)| \frac{dy \, dt}{t} \\ &\lesssim ||a||_{T^{2}(\Omega \times (0,\infty))} \left\{ \int_{0}^{r_{B}} \int_{B_{\Omega}} \frac{t^{2}}{|x-y|^{2(n+1)}} \frac{dy \, dt}{t} \right\}^{1/2} \\ &\lesssim |x-x_{B}|^{-(n+1)} r_{B} |B_{\Omega}|^{1/2} ||a||_{T^{2}(\Omega \times (0,\infty))} \lesssim 2^{-k(n+1)} |B_{\Omega}|^{-1/p}, \end{aligned}$$

which, together with (4.33), further implies that

(4.35)
$$||b_k||_{L^2(\mathbb{R}^n)} \lesssim 2^{-(n/2+1)k} |B_{\Omega}|^{1/2-1/p} \lesssim 2^{-s_0 k} |B_k^*|^{1/2-1/p}$$

where $s_0 := n(1 + \frac{1}{n} - \frac{1}{p})$. By this, $\operatorname{supp}(b_k) \subset B_k^*$, and $\int_{\mathbb{R}^n} b_k(x) dx = 0$, we conclude that $2^{s_0k}b_k$ is a harmless constant multiple of a (p, 2, 0)-atom. Let

$$\widetilde{\alpha} := b_0 + \sum_{k \in J_\Omega} 2^{-s_0 k} (2^{s_0 k} b_k).$$

Then $\widetilde{\alpha}|_{\Omega} = \alpha$, $\widetilde{\alpha} \in H^p(\mathbb{R}^n)$, and $\|\widetilde{\alpha}\|_{H^p(\mathbb{R}^n)} \leq 1$. *Case 2*) $8B \subset \Omega$. In this case, let $k_0 \in \mathbb{N}$ be such that $2^{k_0}B \subset \Omega$ but $(2^{k_0+1}B) \cap \partial\Omega \neq \emptyset$. Then $k_0 \geq 3$. Let

$$J_{\Omega,k_0} := \{k \in \mathbb{N} : k \ge k_0 + 1, |S_k(B_\Omega)| > 0\}.$$

For any $k \in \mathbb{Z}_+$, let $\mathbf{1}_k := \mathbf{1}_{S_k(B_{\Omega})}$,

$$m_k := \int_{S_k(B_\Omega)} \alpha(x) \, dx,$$

 $M_k := \alpha \mathbf{1}_k - m_k |S_k(B_\Omega)|^{-1} \mathbf{1}_k$, and $\widetilde{M}_k := \alpha \mathbf{1}_k$. Then

$$\alpha = \sum_{k=0}^{k_0} M_k + \sum_{k \in J_{\Omega,k_0}} \widetilde{M}_k + \sum_{k=0}^{k_0} m_k |S_k(B_\Omega)|^{-1} \mathbf{1}_k.$$

For any $k \in \{0, ..., k_0\}$, from the definition of M_k , we deduce that

$$\int_{\mathbb{R}^n} M_k(x) \, dx = 0 \quad \text{and} \quad \text{supp} \left(M_k \right) \subset 2^{k+1} B.$$

Moreover, if k = 0, by the Hölder inequality and (4.32), we find that

(4.36)
$$||M_0||_{L^2(\mathbb{R}^n)} \lesssim ||\alpha||_{L^2(\Omega)} \lesssim ||\alpha||_{L^2(\Omega)} \lesssim |B|^{1/2 - 1/p}$$

and, if $k \in \{1, \ldots, k_0\}$, from (4.34), it follows that

(4.37)
$$||M_k||_{L^2(\mathbb{R}^n)} \lesssim ||\alpha||_{L^2(S_k(B_\Omega))} \lesssim 2^{-s_0 k} \left|2^{k+1}B\right|^{1/2-1/p},$$

where s_0 is as in (4.35). Thus, for any $k \in \{0, ..., k_0\}$, $2^{s_0k}M_k$ is a harmless constant multiple of a (p, 2, 0)-atom.

For any $k \in J_{\Omega,k_0}$, by the definitions of both k_0 and J_{Ω,k_0} , we have $2^k r_B \leq \text{diam}(\Omega)$ and $2^k B \cap \partial \Omega \neq \emptyset$. By this, we conclude that there exists a $y_B \in \partial \Omega$ such that $B(y_B, 2^{k+1}r_B) \supset 2^k B$, which, combined with Lemma 2.2(iii), implies that there exists a ball $\widetilde{B}_k \subset \Omega^{\mathbb{C}}$ such that $r_{\widetilde{B}_k} \sim 2^k r_B$ and dist $(S_k(B_\Omega), \widetilde{B}_k) \leq 2^k r_B$. Then there exists a ball B_k^* such that $\widetilde{B}_k \cup S_k(B_\Omega) \subset B_k^*$ and $r_{B_k^*} \sim 2^k r_B$. Let

$$a_k := \alpha \mathbf{1}_{S_k(B_{\Omega})} - \left[\frac{1}{|\widetilde{B}_k|} \int_{S_k(B_{\Omega})} \alpha(x) \, dx\right] \mathbf{1}_{\widetilde{B}_k}.$$

Then supp $(a_k) \subset B_k^*$ and $\int_{\mathbb{R}^n} a_k(x) dx = 0$. Moreover, from (4.34) and Lemma 2.2(i), we deduce that

(4.38)
$$\|a_k\|_{L^2(\mathbb{R}^n)} \lesssim \|\alpha\|_{L^2(S_k(B_\Omega))} \lesssim 2^{-s_0 k} |B_k^*|^{1/2 - 1/p} .$$

Thus, for any $k \in J_{\Omega, k_0}$, $2^{s_0 k} a_k$ is a harmless constant multiple of a (p, 2, 0)-atom. For any $j \in \{0, \ldots, k_0\}$, let $N_j := \sum_{k=j}^{k_0} m_k$. It is easy to see that

$$\sum_{k=0}^{k_0} m_k |S_k(B_\Omega)|^{-1} \mathbf{1}_k = \sum_{k=1}^{k_0} \left[|S_k(B_\Omega)|^{-1} \mathbf{1}_k - |S_{k-1}(B_\Omega)|^{-1} \mathbf{1}_{k-1} \right] N_k + N_0 |2B|^{-1} \mathbf{1}_0.$$

For any $k \in \{1, ..., k_0\}$, by

$$||S_k(B_{\Omega})|^{-1} \mathbf{1}_k - |S_{k-1}(B_{\Omega})|^{-1} \mathbf{1}_{k-1}| \leq |2^k B|^{-1},$$

the Hölder inequality, and (4.34), we find that

(4.39)
$$\left\| \left[|S_{k}(B_{\Omega})|^{-1} \mathbf{1}_{k} - |S_{k-1}(B_{\Omega})|^{-1} \mathbf{1}_{k-1} \right] N_{k} \right\|_{L^{2}(\mathbb{R}^{n})} \\ \lesssim |2^{k}B|^{-1/2} |N_{k}| \lesssim |2^{k}B|^{-1/2} \left[\sum_{j=k}^{k_{0}} ||\alpha||_{L^{2}(S_{j}(B_{\Omega}))} |S_{j}(B_{\Omega})|^{1/2} \right] \\ \lesssim |2^{k}B|^{-1/2} \left[\sum_{j=k}^{k_{0}} 2^{-(n+1)j} |B|^{1/2-1/p} \left| 2^{j}B \right|^{1/2} \right] \lesssim 2^{-s_{0}k} \left| 2^{k}B \right|^{1/2-1/p}$$

where s_0 is as in (4.35), which, together with

$$\int_{\mathbb{R}^n} \left[|S_k(B_\Omega)|^{-1} \mathbf{1}_k(x) - |S_{k-1}(B_\Omega)|^{-1} \mathbf{1}_{k-1}(x) \right] \, dx = 0$$

and supp $(|S_k(B_{\Omega})|^{-1}\mathbf{1}_k - |S_{k-1}(B_{\Omega})|^{-1}\mathbf{1}_{k-1}) \subset 2^k B$, implies that, for any $k \in \{1, ..., k_0\}$, the function $2^{s_0k}[|S_k(B_{\Omega})|^{-1}\mathbf{1}_k - |S_{k-1}(B_{\Omega})|^{-1}\mathbf{1}_{k-1}]N_k$ is a harmless constant multiple of a (p, 2, 0)-atom.

Finally, we deal with $N_0|2B|^{-1}\mathbf{1}_0$. From $2^{k_0-1}r_0 < \text{dist}(x_0, \partial\Omega) \le 2^{k_0}r_0$, it follows that there exist a positive integer K and a sequence $\{B_{0,i}\}_{i=1}^K$ of balls such that

(i)
$$K \sim 2^{k_0}$$
;

- (ii) for any $i \in \{1, ..., K\}$, $r_{B_{0,i}} = 2r_0$ and $B_{0,i} \subset \Omega$;
- (iii) for any $i \in \{1, \ldots, K-1\}$, $B_{0,i} \cap B_{0,i+1} \neq \emptyset$ and dist $(B_{0,i}, \partial\Omega) \ge \text{dist}(B_{0,i+1}, \partial\Omega)$;
- (iv) $2B_{0,K} \cap \partial \Omega \neq \emptyset$.

By Lemma 2.2(ii), we conclude that there exists a ball $B_{0,K+1} \subset \Omega^{\complement}$ such that $r_{B_{0,K+1}} \sim r_0$ and dist $(B_{0,K}, B_{0,K+1}) \sim r_0$. Let

$$a_{0,1} := N_0 |2B|^{-1} \mathbf{1}_0 - N_0 |B_{0,1}|^{-1} \mathbf{1}_{B_{0,1}}$$

and

$$a_{0,i} := N_0 |B_{0,i-1}|^{-1} \mathbf{1}_{B_{0,i-1}} - N_0 |B_{0,i}|^{-1} \mathbf{1}_{B_{0,i}}$$

with $i \in \{2, ..., K + 1\}$. Obviously, for any $i \in \{1, ..., K + 1\}$, from the definition of $a_{0,i}$, we deduce that $\int_{\mathbb{R}^n} a_{0,i}(x) dx = 0$ and there exists a ball $B_{0,i}^* \subset \mathbb{R}^n$ such that $\sup(a_{0,i}) \subset B_{0,i}^*$ and

(4.40)
$$r_{B_{0,i}^*} \sim r_B.$$

Moreover, similarly to the estimation of [65, (3.66)], we have

$$(4.41) |N_0| \lesssim 2^{-\frac{n+1}{n}k_0} |B|^{1-1/p}.$$

For any $i \in \{1, \ldots, K+1\}$, by the definition of $a_{0,i}$, (4.40), and (4.41), we conclude that

(4.42)
$$\left\|a_{0,i}\right\|_{L^{2}(\mathbb{R}^{n})} \lesssim |N_{0}||B|^{-1/2} \lesssim 2^{-\frac{n+1}{n}k_{0}}|B|^{1/2-1/p} \sim 2^{-\frac{n+1}{n}k_{0}} \left|B_{0,i}^{*}\right|^{1/2-1/p}$$

which, combined with the facts that $\int_{\mathbb{R}^n} a_{0,i}(x) dx = 0$ and $\sup(a_{0,i}) \subset B^*_{0,i}$, further implies that $2^{\frac{n+1}{n}k_0}a_{0,i}$ is a harmless constant multiple of a (p, 2, 0)-atom. Let

$$\widetilde{\alpha} := \sum_{k=1}^{k_0} M_k + \sum_{k \in J_{\Omega, k_0}} a_k + \sum_{k=1}^{k_0} \left[|S_k(B_{\Omega})|^{-1} \mathbf{1}_k - |S_{k-1}(B_{\Omega})|^{-1} \mathbf{1}_{k-1} \right] N_k + \sum_{i=1}^{K+1} a_{0, i}.$$

It is easy to find that $\tilde{\alpha}|_{\Omega} = \alpha$. Moreover, from (4.36), (4.37), (4.38), (4.39), and (4.42), it follows that $\tilde{\alpha}$ has the following atomic decomposition

$$\widetilde{\alpha} = \sum_{k=1}^{k_0} 2^{-s_0 k} \left(2^{s_0 k} M_k \right) + \sum_{k \in J_{\Omega, k_0}} 2^{-s_0 k} \left(2^{s_0 k} a_k \right) + \sum_{k=1}^{k_0} 2^{-s_0 k} \left\{ 2^{s_0 k} \left[|S_k(B_{\Omega})|^{-1} \mathbf{1}_k - |S_{k-1}(B_{\Omega})|^{-1} \mathbf{1}_{k-1} \right] N_k \right\} + \sum_{i=1}^{K+1} 2^{-\frac{n+1}{n} k_0} \left[2^{\frac{n+1}{n} k_0} a_{0, i} \right]$$

and

$$\sum_{k=1}^{k_0} 2^{-s_0 k p} + \sum_{k \in J_{\Omega, k_0}} 2^{-s_0 k p} + \sum_{k=1}^{k_0} 2^{-s_0 k p} + \sum_{i=1}^{K+1} 2^{-\frac{n+1}{n} k_0 p}$$

SIBEI YANG AND DACHUN YANG

$$\lesssim \sum_{k=1}^{\infty} 2^{-s_0 k p} + 2^{k_0} 2^{-\frac{n+1}{n} k_0 p} \lesssim 1,$$

which further implies that $\tilde{\alpha} \in H^p(\mathbb{R}^n)$ and $\|\tilde{\alpha}\|_{H^p(\mathbb{R}^n)} \leq 1$. This finishes the proofs of both (4.29) and (4.30), and hence of (4.22).

By (4.9) and (4.22), we obtain

$$\left[H^p_{L_D}(\Omega) \cap L^2(\Omega)\right] = \left[H^p_r(\Omega) \cap L^2(\Omega)\right],$$

which, together with the fact that $H^p_{L_D}(\Omega) \cap L^2(\Omega)$ and $H^p_r(\Omega) \cap L^2(\Omega)$ are dense, respectively, in the spaces $H^p_{L_D}(\Omega)$ and $H^p_r(\Omega)$, and a density argument, implies that $H^p_{L_D}(\Omega)$ and $H^p_r(\Omega)$ coincide with equivalent quasi-norms. This finishes the proof of Theorem 1.5.

5 **Proof of Theorem 1.8**

In this section, we prove Theorem 1.8. We begin with establishing the following estimates for the kernels of the family $\{(tL_D)^k e^{-tL_D}\}_{t>0}$ of operators.

Lemma 5.1. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, the real-valued, bounded, and measurable matrix A satisfy (1.3), and L_D be as in (1.4). Assume that $p_0 \in (2, \infty)$ is as in Lemma 3.6 and $q \in (2, p_0)$. For any given $k \in \mathbb{N}$, denote by $\{K_{t,k}^{L_D}\}_{t>0}$ the kernels of the family $\{(tL_D)^k e^{-tL_D}\}_{t>0}$ of operators. Then there exist positive constants C and c, depending on n, p, k, and Ω , such that, for any $r \in (0, \operatorname{diam}(\Omega)), y \in \Omega$, and $t \in (0, \infty)$,

$$\left[\int_{\{x\in\Omega: \ r\le|x-y|\le 2r\}} \left|\nabla_x K_{t,k}^{L_D}(x,y)\right|^q \, dx\right]^{1/q} \le Cr^{\frac{n}{q}-1}t^{-\frac{n}{2}}e^{-\frac{r^2}{ct}}.$$

Proof. Assume that $f \in L^2(\Omega)$ and $u \in W_0^{1,2}(\Omega)$ is a weak solution of the Dirichlet boundary value problem (1.1). Let $\eta \in C_c^{\infty}(\mathbb{R}^n)$. Then

(5.1)
$$-\operatorname{div}\left(A\nabla(u\eta)\right) = -\operatorname{div}\left(uA\nabla\eta\right) - A\nabla u \cdot \nabla\eta + f\eta$$

in the sense of (1.8). Indeed, for any $\varphi \in C_c^{\infty}(\Omega)$,

$$\begin{split} &\int_{\Omega} A(x)\nabla(u\eta)(x)\cdot\nabla\varphi(x)\,dx\\ &=\int_{\Omega} A(x)\eta(x)\nabla u(x)\cdot\nabla\varphi(x)\,dx + \int_{\Omega} A(x)u(x)\nabla\eta(x)\cdot\nabla\varphi(x)\,dx\\ &=\int_{\Omega} A(x)\nabla u(x)\cdot\nabla(\eta\varphi)(x)\,dx - \int_{\Omega} A(x)\nabla u(x)\cdot\nabla\eta(x)\varphi(x)\,dx\\ &+\int_{\Omega} A(x)u(x)\nabla\eta(x)\cdot\nabla\varphi(x)\,dx\\ &=\int_{\Omega} f(x)\eta(x)\varphi(x)\,dx - \int_{\Omega} A(x)\nabla u(x)\cdot\nabla\eta(x)\varphi(x)\,dx \end{split}$$

40

+
$$\int_{\Omega} A(x)u(x)\nabla\eta(x)\cdot\nabla\varphi(x)\,dx$$

which implies that (5.1) holds true.

Let $t \in (0, \infty)$, $y \in \Omega$, $r \in (0, \text{diam}(\Omega))$, $u_t := K_{t,k}^{L_D}(\cdot, y)$, and $f := -\frac{d}{dt}K_{t,k}^{L_D}(\cdot, y)$. Then $L_D u_t = f$ in the sense of (1.8). Take $\eta \in C_c^{\infty}(\mathbb{R}^n)$ satisfying $\eta \equiv 1$ on $\{x \in \mathbb{R}^n : r \le |x - y| \le 2r\}$,

$$\operatorname{supp}(\eta) \subset \left\{ x \in \mathbb{R}^n : \frac{5r}{6} \le |x - y| \le \frac{13r}{6} \right\},\$$

and $|\nabla \eta| \leq r^{-1}$. Assume that $q \in (2, p_0)$ and $p \in (1, n)$ satisfy $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$. Then, by Lemmas 3.6 and 3.7, and (5.1), we conclude that

(5.2)
$$\|\nabla(u_t\eta)\|_{L^q(\Omega;\mathbb{R}^n)} \lesssim \|u_tA\nabla\eta\|_{L^q(\Omega;\mathbb{R}^n)} + \|A\nabla u_t\cdot\nabla\eta\|_{L^p(\Omega)} + \|f\eta\|_{L^p(\Omega)}.$$

Let $S_1 := \{x \in \Omega : r \le |x - y| \le 2r\}$ and $S_2 := \{x \in \Omega : \frac{5r}{6} \le |x - y| \le \frac{13r}{6}\}$. Then, from (5.2), we deduce that

(5.3)
$$\|\nabla u_t\|_{L^q(S_1;\mathbb{R}^n)} \lesssim r^{-1} \|u_t\|_{L^q(S_2)} + r^{-1} \|\nabla u_t\|_{L^p(S_2;\mathbb{R}^n)} + \left\|\frac{du_t}{dt}\right\|_{L^p(S_2)}$$

We first assume that $p \le 2$. Similarly to the proof of [3, Proposition 16], we have

$$\|\nabla u_t\|_{L^2(S_2;\mathbb{R}^n)} \lesssim t^{-\frac{1}{2}-\frac{n}{4}} \left(\frac{r}{t^{1/2}}\right)^{\frac{n-2}{2}} e^{-\frac{r^2}{ct}},$$

which, together with the Hölder inequality, implies that

(5.4)
$$r^{-1} \|\nabla u_t\|_{L^p(S_2;\mathbb{R}^n)} \lesssim r^{-1} \|\nabla u_t\|_{L^2(S_2;\mathbb{R}^n)} \|S_2\|^{\frac{1}{p}-\frac{1}{2}} \lesssim r^{\frac{n}{q}-1} t^{-\frac{n}{2}} e^{-\frac{r^2}{ct}}.$$

Furthermore, by [51, Theorem 6.17] and (1.5), we find that, for any $x \in \Omega$,

(5.5)
$$|u_t(x)| + t \left| \frac{du_t(x)}{dt} \right| \lesssim t^{-\frac{n}{2}} e^{-\frac{|x-y|^2}{ct}},$$

which further implies that

$$r^{-1} ||u_t||_{L^q(S_2)} + \left\| \frac{du_t}{dt} \right\|_{L^p(S_2)} \lesssim r^{-1} r^{\frac{n}{q}} t^{-\frac{n}{2}} e^{-\frac{r^2}{ct}} + t^{-1} r^{\frac{n}{p}} t^{-\frac{n}{2}} e^{-\frac{r^2}{ct}}$$
$$\lesssim r^{\frac{n}{q}-1} t^{-\frac{n}{2}} e^{-\frac{r^2}{ct}}.$$

From this, (5.3), and (5.4), it follows that

(5.6)
$$\|\nabla u_t\|_{L^q(S_1;\mathbb{R}^n)} \lesssim r^{\frac{n}{q}-1} t^{-\frac{n}{2}} e^{-\frac{r^2}{ct}}.$$

This finishes the proof of the present lemma in the case that $p \leq 2$.

When p > 2, take $i_0 \in \mathbb{N}$ and $1 < p_{i_0} \le 2 < p_{i_0-1} < \cdots < p_1 := p < q$ such that $\frac{1}{p_{i+1}} - \frac{1}{p_i} = \frac{1}{n}$ for any $i \in \{1, \ldots, i_0 - 1\}$. Then, using a simple iteration argument, (5.3), (5.5), and (5.6), we conclude that (5.6) also holds true in this case that p > 2. This finishes the proof of Lemma 5.1.

Moreover, to show Theorem 1.8, we need the following uniform boundedness of the family $\{\sqrt{t}\nabla e^{-tL_D}\}_{t>0}$ of operators on $L^p(\Omega)$, whose proof is similar to that of [4, Proposition 22]; we omit the details here.

Lemma 5.2. Let $n \ge 2$, $\Omega \subset \mathbb{R}^n$ be a bounded NTA domain, the real-valued, bounded, and measurable matrix A satisfy (1.3), and L_D be as in (1.4). Then there exists an $\epsilon_0 \in (0, \infty)$, depending only on n and μ_0 , such that, for any given $p \in (2 - \epsilon_0, 2 + \epsilon_0)$, the family $\{\sqrt{t} \nabla e^{-tL_D}\}_{t>0}$ of operators is uniformly bounded on $L^p(\Omega)$.

Now, we prove Theorem 1.8 by using Lemmas 3.6, 5.1, and 5.2.

Proof of Theorem 1.8. We first show (i). By Lemma 3.6, we find that the conclusion of (i) holds true when $q \in (2, p_0)$. We assume that (ii) holds true, and use the conclusion of (ii) to finish the proof of (i). Define a linear operator ∇L_D^{-1} on \mathbb{R}^n as follows. Let $p \in [1, n)$ and $q \in [\frac{n}{n-1}, \infty)$ satisfy $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$. For any $f \in L^p(\mathbb{R}^n)$ when $p \in (1, n)$, or $f \in H^1(\mathbb{R}^n)$, and any $x \in \mathbb{R}^n$, let

$$\widetilde{\nabla L_D^{-1}}(f)(x) := \begin{cases} \nabla L_D^{-1}(f|_{\Omega})(x) & \text{when } x \in \Omega, \\ 0 & \text{when } x \in \Omega^{\complement}. \end{cases}$$

Then, from Lemma 3.6, it follows that, when $q \in (2, p_0)$,

(5.7)
$$\left\|\widetilde{\nabla L_D^{-1}}(f)\right\|_{L^q(\mathbb{R}^n;\mathbb{R}^n)} = \left\|\nabla L_D^{-1}(f|_{\Omega})\right\|_{L^q(\Omega;\mathbb{R}^n)} \lesssim \|f|_{\Omega}\|_{L^p(\Omega)} \lesssim \|f\|_{L^p(\mathbb{R}^n)}.$$

Thus, the operator $\widetilde{\nabla L_D^{-1}}$ is bounded from $L^p(\mathbb{R}^n)$ to $L^q(\mathbb{R}^n;\mathbb{R}^n)$. Moreover, by (ii), we conclude that

(5.8)
$$\left\|\widetilde{\nabla L_D^{-1}}(f)\right\|_{L^{\frac{n}{n-1}}(\mathbb{R}^n;\mathbb{R}^n)} = \left\|\nabla L_D^{-1}(f|_{\Omega})\right\|_{L^{\frac{n}{n-1}}(\Omega;\mathbb{R}^n)} \lesssim \|f|_{\Omega}\|_{H^1(\Omega)} \lesssim \|f\|_{H^1(\mathbb{R}^n)}$$

which further implies that $\widetilde{\nabla L_D^{-1}}$ is bounded from $H^1(\mathbb{R}^n)$ to $L^{\frac{n}{n-1}}(\mathbb{R}^n; \mathbb{R}^n)$. Recall that, for any $q \in (1, \infty)$, the Hardy space $H^q(\mathbb{R}^n)$ is just the Lebesgue space $L^q(\mathbb{R}^n)$. From this, (5.7), (5.8), and the complex interpolation theory of Hardy spaces on \mathbb{R}^n (see, for instance, [42, Theorem 8.1 and (9.3)]), we deduce that $\widetilde{\nabla L_D^{-1}}$ is bounded from $L^p(\mathbb{R}^n)$ to $L^q(\mathbb{R}^n; \mathbb{R}^n)$ for any given $p \in (1, \frac{np_0}{n+p_0})$ and $q \in (\frac{n}{n-1}, p_0)$ satisfying $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$, which, combined with the definition of $\widetilde{\nabla L_D^{-1}}$, implies that the operator ∇L_D^{-1} is bounded from $L^p(\Omega)$ to $L^q(\Omega; \mathbb{R}^n)$ for any given $p \in (1, \frac{np_0}{n+p_0})$ and $q \in (\frac{n}{n-1}, p_0)$ satisfying $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$. This finishes the proof of (i).

Now, we prove (ii). Let $p \in (\frac{n}{n+1}, 1]$ and $q \in (1, \frac{n}{n-1}]$ satisfy $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$, and $f \in H^p_{L_D}(\Omega) \cap L^2(\Omega)$. Take $1 < p_1 < 2 < q_1 < \min\{p_0, 2+\epsilon_0\}$ such that $\frac{1}{q_1} - \frac{1}{p_1} = \frac{1}{q} - \frac{1}{p}$, where p_0 and ϵ_0 are, respectively, as in Lemmas 3.6 and 5.2. Let $\epsilon \in (\frac{n}{p}, \infty)$ and $M \in \mathbb{N} \cap (\max\{\frac{n}{2p}, \frac{1+\epsilon}{2}\}, \infty)$. Then there exist $\{\lambda_j\}_{j=1}^{\infty} \subset \mathbb{C}$ and a sequence $\{\alpha_j\}_{j=1}^{\infty}$ of $(p, q_1, M, \epsilon)_{L_D}$ -molecules associated, respectively, with the balls $\{B_{\Omega,j}\}_{j=1}^{\infty}$ such that

(5.9)
$$f = \sum_{j=1}^{\infty} \lambda_j \alpha_j$$

in $L^2(\Omega)$, and

(5.10)
$$||f||_{H^p_{L_D}(\Omega)} \sim \left(\sum_{j=0}^{\infty} |\lambda_j|^p\right)^{1/p},$$

where, for any $j \in \mathbb{N}$, $B_{\Omega,j} := B_j \cap \Omega$, and $B_j := B(x_j, r_j)$ with $x_j \in \Omega$ and $r_j \in (0, \text{diam}(\Omega))$ is a ball of \mathbb{R}^n . To finish the proof of (ii), it suffices to prove that, for any $(p, q_1, M, \epsilon)_{L_D}$ -molecule α associated with the ball $B_\Omega := B \cap \Omega$,

(5.11)
$$\left\|\nabla L_D^{-1}(\alpha)\right\|_{L^q(\Omega;\mathbb{R}^n)} \lesssim 1.$$

Indeed, if (5.11) holds true, then, by (5.9), (5.10), (5.11), and $q > 1 \ge p$, we find that

$$\begin{split} \|\nabla u\|_{L^{q}(\Omega;\mathbb{R}^{n})} &= \left\|\nabla L_{D}^{-1}(f)\right\|_{L^{q}(\Omega;\mathbb{R}^{n})} \leq \sum_{j=1}^{\infty} |\lambda_{j}| \left\|\nabla L_{D}^{-1}(\alpha_{j})\right\|_{L^{q}(\Omega;\mathbb{R}^{n})} \\ &\lesssim \sum_{j=0}^{\infty} |\lambda_{j}| \leq \left(\sum_{j=0}^{\infty} |\lambda_{j}|^{p}\right)^{1/p} \sim \|f\|_{H^{p}_{L_{D}}(\Omega)}, \end{split}$$

which implies that (ii) holds true.

Next, we prove (5.11). From Lemma 3.6, it follows that ∇L_D^{-1} is bounded from $L^{p_1}(\Omega)$ to $L^{q_1}(\Omega; \mathbb{R}^n)$, which, together with the Hölder inequality and $\frac{1}{p_1} - \frac{1}{q_1} = \frac{1}{p} - \frac{1}{q}$, implies that

(5.12)
$$\left\| \nabla L_D^{-1}(\alpha) \right\|_{L^q(8B_{\Omega};\mathbb{R}^n)} \le \left\| \nabla L_D^{-1}(\alpha) \right\|_{L^{q_1}(8B_{\Omega};\mathbb{R}^n)} \left| 8B_{\Omega} \right|^{\frac{1}{q} - \frac{1}{q_1}} \\ \lesssim \|\alpha\|_{L^{p_1}(8B_{\Omega})} |8B_{\Omega}|^{\frac{1}{p} - \frac{1}{p_1}} \lesssim 1.$$

For any $j \in \mathbb{N}$ with $j \ge 3$, let $\widetilde{S}_j(B_{\Omega}) := (2^{j+3}B_{\Omega}) \setminus (2^{j-3}B_{\Omega})$ and $E_j(B_{\Omega}) := \Omega \setminus \widetilde{S}_j(B_{\Omega})$. Then, by the equation

$$\nabla L_D^{-1}(\alpha) = \int_0^\infty \nabla e^{-tL_D}(\alpha) \, dt$$

and the Minkowski inequality, we conclude that

(5.13)
$$\left\|\nabla L_D^{-1}(\alpha)\right\|_{L^q(S_j(B_\Omega);\mathbb{R}^n)} \le \int_0^{r_B^2} \left\|\nabla e^{-tL_D}(\alpha)\right\|_{L^q(S_j(B_\Omega);\mathbb{R}^n)} dt + \int_{r_B^2}^{\infty} \cdots$$
$$=: \mathbf{I} + \mathbf{II}.$$

For the term I, we have

(5.14)
$$\mathbf{I} = \int_{0}^{r_{B}^{2}} \left\| \nabla e^{-tL_{D}} \left(\alpha \mathbf{1}_{\widetilde{S}_{j}(B_{\Omega})} \right) \right\|_{L^{q}(S_{j}(B_{\Omega});\mathbb{R}^{n})} dt$$
$$+ \int_{0}^{r_{B}^{2}} \left\| \nabla e^{-tL_{D}} \left(\alpha \mathbf{1}_{E_{j}(B_{\Omega})} \right) \right\|_{L^{q}(S_{j}(B_{\Omega});\mathbb{R}^{n})} dt$$
$$=: \mathbf{I}_{1} + \mathbf{I}_{2}.$$

From Lemma 5.2, the Hölder inequality, and $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$, we deduce that

(5.15)
$$I_{1} \lesssim \int_{0}^{r_{B}^{2}} \left\| \nabla e^{-tL_{D}} \left(\alpha \mathbf{1}_{\widetilde{S}_{j}(B_{\Omega})} \right) \right\|_{L^{q_{1}}(S_{j}(B_{\Omega});\mathbb{R}^{n})} \left| 2^{j}B_{\Omega} \right|^{\frac{1}{q}-\frac{1}{q_{1}}} dt$$
$$\lesssim 2^{-j\epsilon} \left| 2^{j}B_{\Omega} \right|^{\frac{1}{q_{1}}} \left| B_{\Omega} \right|^{-\frac{1}{p}} \left| 2^{j}B_{\Omega} \right|^{\frac{1}{q}-\frac{1}{q_{1}}} \int_{0}^{r_{B}^{2}} t^{-1/2} dt$$
$$\lesssim 2^{-j(\epsilon-\frac{n}{q})} \left| B_{\Omega} \right|^{-\frac{1}{n}} r_{B} \lesssim 2^{-j(\epsilon-\frac{n}{q})}.$$

Moreover, by Lemma 5.1, the Minkowski inequality, and $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$, we conclude that

(5.16)
$$I_{2} \lesssim \int_{0}^{r_{B}^{2}} \left\| \nabla e^{-tL_{D}} \left(\alpha \mathbf{1}_{E_{j}(B_{\Omega})} \right) \right\|_{L^{q_{1}}(S_{j}(B_{\Omega});\mathbb{R}^{n})} \left| 2^{j}B_{\Omega} \right|^{\frac{1}{q} - \frac{1}{q_{1}}} dt$$
$$\lesssim \left| 2^{j}B_{\Omega} \right|^{\frac{1}{q} - \frac{1}{q_{1}}} \int_{0}^{r_{B}^{2}} \int_{E_{j}(B_{\Omega})} \left\| \nabla K_{t}^{L_{D}}(\cdot, y) \right\|_{L^{q_{1}}(S_{j}(B_{\Omega});\mathbb{R}^{n})} \left| \alpha(y) \right| dy dt$$
$$\lesssim \left| 2^{j}B_{\Omega} \right|^{\frac{1}{q} - \frac{1}{q_{1}}} \int_{0}^{r_{B}^{2}} (2^{j}r_{B})^{\frac{n}{q_{1}} - 1} t^{-\frac{n}{2}} e^{-\frac{(2^{j}r_{B})^{2}}{ct}} \left| B_{\Omega} \right|^{1 - \frac{1}{p}} dt \lesssim 2^{-j(\epsilon - \frac{n}{q})}.$$

For the term II, we have

(5.17)
$$\begin{aligned} \Pi &= \int_{r_B^2}^{\infty} \frac{1}{t^M} \left\| \nabla (tL_D)^M e^{-tL_D} \left(\left(L^{-M} \alpha \right) \mathbf{1}_{\widetilde{S}_j(B_\Omega)} \right) \right\|_{L^q(S_j(B_\Omega);\mathbb{R}^n)} dt \\ &+ \int_{r_B^2}^{\infty} \frac{1}{t^M} \left\| \nabla (tL_D)^M e^{-tL_D} \left(\left(L_D^{-M} \alpha \right) \mathbf{1}_{E_j(B_\Omega)} \right) \right\|_{L^q(S_j(B_\Omega);\mathbb{R}^n)} dt \\ &=: \Pi_1 + \Pi_2. \end{aligned}$$

From the facts that, for any $t \in (0, \infty)$,

$$\sqrt{t}\nabla(tL_D)^M e^{-tL_D} = 2^{M+\frac{1}{2}} \left(\left(\frac{t}{2}\right)^{1/2} \nabla e^{-\frac{t}{2}L_D} \right) \left(\frac{t}{2}L_D\right)^M e^{-\frac{t}{2}L_D},$$

 $\{(\frac{t}{2}L_D)^M e^{-\frac{t}{2}L_D}\}_{t>0}$ is uniformly bounded on $L^s(\Omega)$ for any given $s \in [1, \infty)$ (see, for instance, [51]), and Lemma 5.2, it follows that, for any given $s \in (2 - \epsilon_0, 2 + \epsilon_0)$, $\{\sqrt{t}\nabla(tL_D)^M e^{-tL_D}\}_{t>0}$ is uniformly bounded on $L^s(\Omega)$. By this and the Hölder inequality, we find that

(5.18)
$$\begin{aligned} \Pi_{1} \lesssim \int_{r_{B}^{2}}^{\infty} \frac{1}{t^{M}} \left\| \nabla (tL_{D})^{M} e^{-tL_{D}} \left(\left(L^{-M} \alpha \right) \mathbf{1}_{\widetilde{S}_{j}(B_{\Omega})} \right) \right\|_{L^{q_{1}}(S_{j}(B_{\Omega});\mathbb{R}^{n})} \left| 2^{j} B_{\Omega} \right|^{\frac{1}{q} - \frac{1}{q_{1}}} dt \\ \lesssim 2^{-j\epsilon} r_{B}^{2M} \left| 2^{j} B_{\Omega} \right|^{\frac{1}{q_{1}}} \left| B_{\Omega} \right|^{-\frac{1}{p}} \left| 2^{j} B_{\Omega} \right|^{\frac{1}{q} - \frac{1}{q_{1}}} \int_{r_{B}^{2}}^{\infty} t^{-(M+1/2)} dt \\ \lesssim 2^{-j(\epsilon - \frac{n}{q})} r_{B}^{2M} \left| B_{\Omega} \right|^{-\frac{1}{n}} r_{B}^{-2M+1} \lesssim 2^{-j(\epsilon - \frac{n}{q})}. \end{aligned}$$

Moreover, from the Hölder inequality, the Minkowski inequality, and Lemma 5.1, we deduce that

(5.19)
$$II_{2} \lesssim \int_{r_{B}^{2}}^{\infty} \frac{1}{t^{M}} \left\| \nabla (tL_{D})^{M} e^{-tL_{D}} \left(\left(L_{D}^{-M} \alpha \right) \mathbf{1}_{E_{j}(B_{\Omega})} \right) \right\|_{L^{q_{1}}(S_{j}(B_{\Omega});\mathbb{R}^{n})} \left| 2^{j} B_{\Omega} \right|^{\frac{1}{q} - \frac{1}{q_{1}}} dt$$

$$\begin{split} &\lesssim \left|2^{j}B_{\Omega}\right|^{\frac{1}{q}-\frac{1}{q_{1}}} \int_{r_{B}^{2}}^{\infty} \frac{1}{t^{M}} \int_{E_{j}(B_{\Omega})} \left\|\nabla K_{t,M}^{L_{D}}(\cdot,y)\right\|_{L^{q_{1}}(S_{j}(B_{\Omega});\mathbb{R}^{n})} \left|L_{D}^{-M}\alpha(y)\right| \, dy dt \\ &\lesssim \left|2^{j}B_{\Omega}\right|^{\frac{1}{q}-\frac{1}{q_{1}}} \int_{r_{B}^{2}}^{\infty} (2^{j}r_{B})^{\frac{n}{q_{1}}-1} t^{-(M+\frac{n}{2})} e^{-\frac{(2^{j}r_{B})^{2}}{c^{l}}} r_{B}^{2M} |B_{\Omega}|^{1-\frac{1}{p}} \, dt \\ &\lesssim \left|2^{j}B_{\Omega}\right|^{\frac{1}{q}-\frac{1}{n}} r_{B}^{2M} |B_{\Omega}|^{1-\frac{1}{p}} \int_{r_{B}^{2}}^{\infty} t^{-(M+\frac{n}{2})} \left[\frac{t}{(2^{j}r_{B})^{2}}\right]^{M-1} \, dt \\ &\lesssim 2^{-j(2M-1-\frac{n}{q})} \lesssim 2^{-j(\epsilon-\frac{n}{q})}. \end{split}$$

Thus, by (5.13) through (5.19), we conclude that, for any $j \in \mathbb{N}$ with $j \ge 3$,

(5.20)
$$\left\|\nabla L_D^{-1}(\alpha)\right\|_{L^q(S_j(B_\Omega);\mathbb{R}^n)} \lesssim 2^{-j(\epsilon-\frac{n}{q})},$$

which, combined with (5.12) and $\epsilon > \frac{n}{p} > \frac{n}{q}$, further implies that

$$\left\|\nabla L_D^{-1}(\alpha)\right\|_{L^q(\Omega;\mathbb{R}^n)} \le \left\|\nabla L_D^{-1}(\alpha)\right\|_{L^q(\mathbb{B}_\Omega;\mathbb{R}^n)} + \sum_{j=3}^{\infty} \left\|\nabla L_D^{-1}(\alpha)\right\|_{L^q(S_j(B_\Omega);\mathbb{R}^n)} \le 1.$$

Thus, (5.11) holds true. This finishes the proof of (ii).

Finally, we show (iii). Let $p \in (\frac{n}{n+2}, \frac{n}{n+1}]$ and $q \in (\frac{n}{n+1}, 1]$ satisfy $\frac{1}{p} - \frac{1}{q} = \frac{1}{n}$, and $f \in H_{L_D}^p(\Omega) \cap L^2(\Omega)$. Take $1 < p_2 < 2 < q_2 < \min\{p_0, 2 + \epsilon_0\}$ such that $\frac{1}{q_2} - \frac{1}{p_2} = \frac{1}{q} - \frac{1}{p}$, where p_0 and ϵ_0 are, respectively, as in Lemmas 3.6 and 5.2. Let $\epsilon \in (\frac{n}{p}, \infty)$ and $M \in \mathbb{N} \cap (\max\{\frac{n}{2p}, \frac{1+\epsilon}{2}\}, \infty)$. Then there exist $\{\lambda_j\}_{j=1}^{\infty} \subset \mathbb{C}$ and a sequence $\{\alpha_j\}_{j=1}^{\infty}$ of $(p, q_2, M, \epsilon)_{L_D}$ -molecules associated, respectively, with the balls $\{B_{\Omega, j}\}_{j=1}^{\infty}$ such that (5.9) and (5.10) hold true. To finish the proof of (iii), it suffices to show that, for any $(p, q_2, M, \epsilon)_{L_D}$ -molecule α , the zero extension of $\nabla L_D^{-1}(\alpha)$ from Ω to \mathbb{R}^n , denoted by $\nabla \widetilde{L_D^{-1}(\alpha)}$, is a harmless constant multiple of a $(q, q_2, 0, \epsilon)$ -molecule associated with the ball *B*. Indeed, if this claim holds true, then, by this, p < q, (5.9), and (5.10), we find that the zero extension of $\nabla L_D^{-1}(f)$ from Ω to \mathbb{R}^n , denoted by $\nabla \widetilde{L_D^{-1}(f)}$ from Ω to \mathbb{R}^n , denoted by $\nabla \widetilde{L_D^{-1}(f)}$ form Ω to \mathbb{R}^n , denoted by $\nabla \widetilde{L_D^{-1}(f)}$ form Ω to \mathbb{R}^n , denoted by $\nabla \widetilde{L_D^{-1}(f)}$, belongs to $H^q(\mathbb{R}^n; \mathbb{R}^n)$, and

$$\left\|\nabla \widetilde{L_D^{-1}(f)}\right\|_{H^q(\mathbb{R}^n;\mathbb{R}^n)} \lesssim \left(\sum_{j=0}^\infty |\lambda_j|^q\right)^{1/q} \lesssim \left(\sum_{j=0}^\infty |\lambda_j|^p\right)^{1/p} \sim \|f\|_{H^p_{L_D}(\Omega)},$$

which further implies that $\nabla u = \nabla L_D^{-1}(f) \in H_z^q(\mathbb{R}^n; \mathbb{R}^n)$ and

$$\|\nabla u\|_{H^q_z(\Omega;\mathbb{R}^n)} = \left\|\nabla L_D^{-1}(f)\right\|_{H^q_z(\Omega;\mathbb{R}^n)} \lesssim \|f\|_{H^p_{L_D}(\Omega)}.$$

Thus, (iii) holds true.

Let α be a $(p, q_2, M, \epsilon)_{L_D}$ -molecule. Now, we prove that $\nabla L_D^{-1}(\alpha)$ is a harmless constant multiple of a $(q, q_2, 0, \epsilon)$ -molecule associated with the ball *B*. Take a ball $B_0 \subset \mathbb{R}^n$ such that $\Omega \subset B_0$, and $\varphi \in C_c^{\infty}(\mathbb{R}^n)$ satisfies that $\varphi \equiv 1$ on B_0 . From $\alpha \in L^{p_2}(\Omega)$ and (i), we deduce that $L_D^{-1}(\alpha) \in W_0^{1,q_2}(\Omega)$, which, together with that $\varphi \equiv 1$ on B_0 , further implies that

(5.21)
$$\int_{\mathbb{R}^n} \nabla \widetilde{L_D^{-1}(\alpha)}(x) \, dx = \int_{\mathbb{R}^n} \nabla \widetilde{L_D^{-1}(\alpha)}(x) \varphi(x) \, dx = -\int_{\mathbb{R}^n} L_D^{-1}(\alpha)(x) \nabla \varphi(x) \, dx = 0.$$

Furthermore, by (i), we find that ∇L_D^{-1} is bounded from $L^{p_2}(\Omega)$ to $L^{q_2}(\Omega; \mathbb{R}^n)$, which, combined with $\frac{1}{q_2} - \frac{1}{p_2} = \frac{1}{q} - \frac{1}{p}$, implies that

(5.22)
$$\left\| \nabla \widetilde{L^{-1}}(\alpha) \right\|_{L^{q_2}(4B;\mathbb{R}^n)} = \left\| \nabla L^{-1}(\alpha) \right\|_{L^{q_2}(4B_\Omega;\mathbb{R}^n)} \lesssim \|\alpha\|_{L^{p_2}(\Omega)}$$
$$\lesssim |B_\Omega|^{\frac{1}{p_2} - \frac{1}{p}} \sim |B_\Omega|^{\frac{1}{q_2} - \frac{1}{q}} \lesssim |4B|^{\frac{1}{q_2} - \frac{1}{q}}.$$

Moreover, similarly to the proof of (5.20), for any $j \in \mathbb{N}$ with $j \ge 3$, we have

(5.23)
$$\left\|\nabla \widetilde{L^{-1}(\alpha)}\right\|_{L^{q_2}(S_j(B);\mathbb{R}^n)} = \left\|\nabla L^{-1}(\alpha)\right\|_{L^{q_2}(S_j(B_\Omega);\mathbb{R}^n)} \lesssim 2^{-j\epsilon} \left|2^j B\right|^{\frac{1}{q_2}} |B|^{-\frac{1}{q}}.$$

Thus, from (5.21), (5.22), and (5.23), it follows that $\nabla L^{-1}(\alpha)$ is a harmless constant multiple of a $(q, q_2, 0, \epsilon)$ -molecule. This finishes the proof of (iii), and hence of Theorem 1.8.

Acknowledgements. Sibei Yang would like to thank Professors Jun Cao and Jun Geng for some helpful discussions on the topic of this article.

References

- [1] P. Auscher and M. Qafsaoui, Observations on $W^{1,p}$ estimates for divergence elliptic equations with VMO coefficients, Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8) 5 (2002), 487-509.
- [2] P. Auscher and E. Russ, Hardy spaces and divergence operators on strongly Lipschitz domains of Rⁿ, J. Funct. Anal. 201 (2003), 148-184.
- [3] P. Auscher and P. Tchamitchian, Gaussian estimates for second order elliptic divergence operators on Lipschitz and C¹ domains, in: Evolution Equations and Their Applications in Physical and Life Sciences (Bad Herrenalb, 1998), 15-32, Lecture Notes in Pure and Applied Mathematics 215, Dekker, New York, 2001.
- [4] P. Auscher and P. Tchamitchian, Square Root Problem for Divergence Operators and Related Topics, Astérisque 249 (1998), viii+172 pp.
- [5] S. Buckley and P. Koskela, Sobolev–Poincaré implies John, Math. Res. Lett. 2 (1995), 577-593.
- [6] T. A. Bui, J. Cao, L. D. Ky, D. Yang and S. Yang, Musielak–Orlicz–Hardy spaces associated with operators satisfying reinforced off-diagonal estimates, Anal. Geom. Metr. Spaces 1 (2013), 69-129.
- [7] T. A. Bui and X. T. Duong, Regularity estimates for Green operators of Dirichlet and Neumann problems on weighted Hardy spaces, J. Math. Soc. Japan 73 (2021), 597-631.
- [8] T. A. Bui and X. T. Duong, Global Lorentz estimates for nonlinear parabolic equations on nonsmooth domains, Calc. Var. Partial Differential Equations 56 (2017), no. 2, Art. 47, 24 pp.
- [9] T. A. Bui, X. T. Duong and F. K. Ly, Maximal function characterizations for Hardy spaces on spaces of homogeneous type with finite measure and applications, J. Funct. Anal. 278 (2020), no. 8, 108423, 55 pp.
- [10] T. A. Bui, X. T. Duong and F. K. Ly, Maximal function characterizations for new local Hardy-type spaces on spaces of homogeneous type, Trans. Amer. Math. Soc. 370 (2018), 7229-7292.

- [11] S.-S. Byun and L. Wang, Elliptic equations with BMO coefficients in Reifenberg domains, Comm. Pure Appl. Math. 57 (2004), 1283-1310.
- [12] J. Cao, D.-C. Chang, D. Yang and S. Yang, Weighted local Orlicz–Hardy spaces on domains and their applications in inhomogeneous Dirichlet and Neumann problems, Trans. Amer. Math. Soc. 365 (2013), 4729-4809.
- [13] D.-C. Chang, G. Dafni and E. M. Stein, Hardy spaces, BMO, and boundary value problems for the Laplacian on a smooth domain in \mathbb{R}^n , Trans. Amer. Math. Soc. 351 (1999), 1605-1661.
- [14] D.-C. Chang, Z. Fu, D. Yang and S. Yang, Real-variable characterizations of Musielak– Orlicz–Hardy spaces associated with Schrödinger operators on domains, Math. Methods Appl. Sci. 39 (2016), 533-569.
- [15] D.-C. Chang, S. G. Krantz and E. M. Stein, H^p theory on a smooth domain in R^N and elliptic boundary value problems, J. Funct. Anal. 114 (1993), 286-347.
- [16] D.-C. Chang, S. G. Krantz and E. M. Stein, Hardy spaces and elliptic boundary value problems, in: The Madison Symposium on Complex Analysis (Madison, WI, 1991), 119-131, Contemp. Math. 137, Amer. Math. Soc., Providence, RI, 1992.
- [17] R. R. Coifman and G. Weiss, Analyse Harmonique Non-Commutative sur Certains Espaces Homogénes, (French) Étude de certaines intégrales singuliéres, Lecture Notes in Mathematics 242, Springer-Verlag, Berlin–New York, 1971.
- [18] B. E. J. Dahlberg, L^q-estimates for Green potentials in Lipschitz domains, Math. Scand. 44 (1979), 149-170.
- [19] E. B. Davies, Heat Kernels and Spectral Theory, Cambridge Tracts in Mathematics 92, Cambridge University Press, Cambridge, 1989.
- [20] G. Di Fazio, L^p estimates for divergence form elliptic equations with discontinuous coefficients, Boll. Un. Mat. Ital. A (7) 10 (1996), 409-420.
- [21] H. Dong, Recent progress in the L^p theory for elliptic and parabolic equations with discontinuous coefficients, Anal. Theory Appl. 36 (2020), 161-199.
- [22] H. Dong and D. Kim, Elliptic equations in divergence form with partially BMO coefficients, Arch. Ration. Mech. Anal. 196 (2010), 25-70.
- [23] H. Dong and D. Kim, The conormal derivative problem for higher order elliptic systems with irregular coefficients, in: Recent Advances in Harmonic Analysis and Partial Differential Equations, 69-97, Contemp. Math. 581, Amer. Math. Soc., Providence, RI, 2012.
- [24] H. Dong and D. Kim, On L^p -estimates for elliptic and parabolic equations with A_p weights, Trans. Amer. Math. Soc. 370 (2018), 5081-5130.
- [25] H. Dong and Z. Li, The conormal and Robin boundary value problems in nonsmooth domains satisfying a measure condition, J. Funct. Anal. 281 (2021), no. 9, Paper No. 109167, 32 pp.
- [26] X. T. Duong, S. Hofmann, D. Mitrea, M. Mitrea and L. Yan, Hardy spaces and regularity for the inhomogeneous Dirichlet and Neumann problems, Rev. Mat. Iberoam. 29 (2013), 183-236.
- [27] X. T. Duong and E. M. Ouhabaz, Gaussian upper bounds for heat kernels of a class of nondivergence operators, in: International Conference on Harmonic Analysis and Related Topics (Sydney, 2002), 35-45, Proc. Centre Math. Appl. Austral. Nat. Univ., 41, Austral. Nat. Univ., Canberra, 2003.

- [28] X. T. Duong and L. Yan, Duality of Hardy and BMO spaces associated with operators with heat kernel bounds, J. Amer. Math. Soc. 18 (2005), 943-973.
- [29] C. Fefferman and E. M. Stein, H^p spaces of several variables, Acta Math. 129 (1972), 137-193.
- [30] J. Geng, *W*^{1,*p*} estimates for elliptic problems with Neumann boundary conditions in Lipschitz domains, Adv. Math. 229 (2012), 2427-2448.
- [31] D. Gilbarg and N. S. Trudinger, Elliptic Partial Differential Equations of Second Order, Reprint of the 1998 edition, Springer-Verlag, Berlin, 2001.
- [32] M. Grüter and K.-O. Widman, The Green function for uniformly elliptic equations, Manuscripta Math. 37 (1982), 303-342.
- [33] P. Gyrya and L. Saloff-Coste, Neumann and Dirichlet Heat Kernels in Inner Uniform Domains, Astérisque 336 (2011), viii+144 pp.
- [34] P. Hajłasz, P. Koskela and H. Tuominen, Sobolev embeddings, extensions and measure density condition, J. Funct. Anal. 254 (2008), 1217-1234.
- [35] S. Hofmann and S. Kim, The Green function estimates for strongly elliptic systems of second order, Manuscripta Math. 124 (2007), 139-172.
- [36] S. Hofmann, G. Lu, D. Mitrea, M. Mitrea and L. Yan, Hardy spaces associated to nonnegative self-adjoint operators satisfying Davies–Gaffney estimates, Mem. Amer. Math. Soc. 214 (2011), no. 1007, vi+78 pp.
- [37] S. Hofmann, S. Mayboroda and A. McIntosh, Second order elliptic operators with complex bounded measurable coefficients in L^p, Sobolev and Hardy spaces, Ann. Sci. Éc. Norm. Super. (4) 44 (2011), 723-800.
- [38] D. Jerison and C. E. Kenig, Boundary behavior of harmonic functions in nontangentially accessible domains, Adv. Math. 46 (1982), 80-147.
- [39] D. Jerison and C. E. Kenig, The inhomogeneous Dirichlet problem in Lipschitz domains, J. Funct. Anal. 130 (1995), 161-219.
- [40] H. Jia, D. Li and L. Wang, Global regularity for divergence form elliptic equations on quasiconvex domains, J. Differential Equations 249 (2010), 3132-3147.
- [41] P. W. Jones, Quasiconformal mappings and extendability of functions in Sobolev spaces, Acta Math. 147 (1981), 71-88.
- [42] N. Kalton, S. Mayboroda and M. Mitrea, Interpolation of Hardy–Sobolev–Besov–Triebel– Lizorkin spaces and applications to problems in partial differential equations, in: Interpolation Theory and Applications, 121-177, Contemp. Math. 445, Amer. Math. Soc., Providence, RI, 2007.
- [43] C. E. Kenig, Harmonic Analysis Techniques for Second Order Elliptic Boundary Value Problems, CBMS Regional Conference Series in Mathematics 83, American Mathematical Society, Providence, RI, 1994.
- [44] C. E. Kenig and T. Toro, Harmonic measure on locally flat domains, Duke Math. J. 87 (1997), 509-551.
- [45] R. H. Latter, A characterization of $H^p(\mathbb{R}^n)$ in terms of atoms, Studia Math. 62 (1978), 93-101.
- [46] J. Li and B. D. Wick, Characterizations of $H^1_{\Delta_N}(\mathbb{R}^n)$ and $BMO_{\Delta_N}(\mathbb{R}^n)$ via weak factorizations and commutators, J. Funct. Anal. 272 (2017), 5384-5416.
- [47] L. Li and J. Pipher, Boundary behavior of solutions of elliptic operators in divergence form with a BMO anti-symmetric part, Comm. Partial Differential Equations 44 (2019), 156-204.

- [48] D. Mitrea, I. Mitrea, M. Mitrea and L. Yan, Coercive energy estimates for differential forms in semi-convex domains, Commun. Pure Appl. Anal. 9 (2010), 987-1010.
- [49] D. Mitrea, M. Mitrea and L. Yan, Boundary value problems for the Laplacian in convex and semiconvex domains, J. Funct. Anal. 258 (2010), 2507-2585.
- [50] A. Miyachi, H^p spaces over open subsets of \mathbb{R}^n , Studia Math. 95 (1990), 205-228.
- [51] E. M. Ouhabaz, Analysis of Heat Equations on Domains, London Mathematical Society Monographs Series 31, Princeton University Press, Princeton, NJ, 2005.
- [52] A. Pazy, Semigroups of Linear Operators and Applications to Partial Differential Equations, Applied Mathematical Sciences 44, Springer-Verlag, New York, 1983.
- [53] E. R. Reifenberg, Solution of the Plateau Problem for *m*-dimensional surfaces of varying topological type, Acta Math. 104 (1960), 1-92.
- [54] E. Russ, The atomic decomposition for tent spaces on spaces of homogeneous type, in: CMA/AMSI Research Symposium "Asymptotic Geometric Analysis, Harmonic Analysis, and Related Topics", 125-135, Proc. Centre Math. Appl. 42, Austral. Nat. Univ., Canberra, 2007.
- [55] L. Saloff-Coste, The heat kernel and its estimates, in: Probabilistic approach to geometry, 405-436, Adv. Stud. Pure Math. 57, Math. Soc. Japan, Tokyo, 2010.
- [56] D. Sarason, Functions of vanishing mean oscillation, Trans. Amer. Math. Soc. 207 (1975), 391-405.
- [57] Z. Shen, Periodic Homogenization of Elliptic Systems, Operator Theory: Advances and Applications 269, Advances in Partial Differential Equations (Basel), Birkhäuser/Springer, Cham, 2018.
- [58] Z. Shen, The L^p boundary value problems on Lipschitz domains, Adv. Math. 216 (2007), 212-254.
- [59] Z. Shen, Bounds of Riesz transforms on L^p spaces for second order elliptic operators, Ann. Inst. Fourier (Grenoble) 55 (2005), 173-197.
- [60] L. Song and L. Yan, A maximal function characterization for Hardy spaces associated to nonnegative self-adjoint operators satisfying Gaussian estimates, Adv. Math. 287 (2016), 463-484.
- [61] E. M. Stein, Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory integrals, Princeton Univ. Press, Princeton, NJ, 1993.
- [62] E. M. Stein and G. Weiss, On the theory of harmonic functions of several variables. I. The theory of H^p-spaces, Acta Math. 103 (1960), 25-62.
- [63] M. H. Taibleson and G. Weiss, The molecular characterization of certain Hardy spaces, in: Representation Theorems for Hardy Spaces, pp. 67-149, Astérisque 77 (1980).
- [64] T. Toro, Doubling and flatness: geometry of measures, Notices Amer. Math. Soc. 44 (1997), 1087-1094.
- [65] D. Yang and S. Yang, Real-variable characterizations of Orlicz–Hardy spaces on strongly Lipschitz domains of \mathbb{R}^n , Rev. Mat. Iberoam. 29 (2013), 237-292.
- [66] D. Yang and S. Yang, Orlicz–Hardy spaces associated with divergence operators on unbounded strongly Lipschitz domains of \mathbb{R}^n , Indiana Univ. Math. J. 61 (2012), 81-129.
- [67] S. Yang, D.-C. Chang, D. Yang and W. Yuan, Weighted gradient estimates for elliptic problems with Neumann boundary conditions in Lipschitz and (semi-)convex domains, J. Differential Equations 268 (2020), 2510-2550.

- [68] S. Yang and D. Yang, Atomic and maximal function characterizations of Musielak–Orlicz– Hardy spaces associated to non-negative self-adjoint operators on spaces of homogeneous type, Collect. Math. 70 (2019), 197-246.
- [69] S. Yang, D. Yang and W. Yuan, Global gradient estimates for Dirichlet problems of elliptic operators with a BMO anti-symmetric part, Submitted or arXiv: 2201.00909v1.

Sibei Yang

School of Mathematics and Statistics, Gansu Key Laboratory of Applied Mathematics and Complex Systems, Lanzhou University, Lanzhou 730000, People's Republic of China *E-mail:* yangsb@lzu.edu.cn (S. Yang)

Dachun Yang (Corresponding author)

Laboratory of Mathematics and Complex Systems (Ministry of Education of China), School of Mathematical Sciences, Beijing Normal University, Beijing 100875, People's Republic of China *E-mail*: dcyang@bnu.edu.cn (D. Yang)