

ESTIMATES FOR FUNDAMENTAL SOLUTIONS OF PARABOLIC EQUATIONS IN NON-DIVERGENCE FORM

HONGJIE DONG, SEICK KIM, AND SUNGJIN LEE

ABSTRACT. We construct the fundamental solution of second order parabolic equations in non-divergence form under the assumption that the coefficients are of Dini mean oscillation in the spatial variables. We also prove that the fundamental solution satisfies a sub-Gaussian estimate. In the case when the coefficients are Dini continuous in the spatial variables and measurable in the time variable, we establish the Gaussian bounds for the fundamental solutions. We present a method that works equally for second order parabolic systems in non-divergence form.

1. INTRODUCTION AND MAIN RESULTS

We consider second order parabolic operator P in non-divergence form

$$Pu = \partial_t u - a^{ij}(t, x) D_{ij} u$$

in \mathbb{R}^{d+1} . Here and below, we use the summation convention over repeated indices. We assume that the coefficients $\mathbf{A} = (a^{ij})$ are symmetric and satisfy the uniform parabolicity condition

$$\lambda |\xi|^2 \leq a^{ij}(t, x) \xi_i \xi_j \leq \Lambda |\xi|^2, \quad \forall \xi \in \mathbb{R}^d, \quad \forall (t, x) \in \mathbb{R}^{d+1}. \quad (1.1)$$

In this article, we are concerned with the fundamental solution of the operator P . By the fundamental solution, we mean a function $\Gamma(t, x, s, y)$ formally satisfying

$$P\Gamma(\cdot, \cdot, s, y) = \delta_{s,y}(\cdot, \cdot) \text{ in } \mathbb{R}^{d+1},$$

or equivalently

$$P\Gamma(\cdot, \cdot, s, y) = 0 \text{ in } (s, \infty) \times \mathbb{R}^d, \quad \lim_{t \rightarrow s+} \Gamma(t, \cdot, s, y) = \delta_y(\cdot) \text{ on } \mathbb{R}^d.$$

We show that if the coefficients $\mathbf{A} = (a^{ij})$ are of Dini mean oscillation in x , then the fundamental solution $\Gamma(t, x, s, y)$ exists and satisfies certain estimates, in particular a sub-Gaussian estimate. Moreover, if the coefficients are Dini continuous in x , then the fundamental solution enjoys the usual Gaussian bounds. We emphasize that our methods are also applicable to parabolic systems of second order and this is one of the novelties of the paper.

Key words and phrases. Fundamental solution; Parabolic equation in non-divergence form; Dini mean oscillation.

H. Dong was partially supported by the Simons Foundation, grant no. 709545, a Simons fellowship, grant no. 007638, and the NSF under agreement DMS-2055244.

S. Kim was partially supported by the National Research Foundation of Korea under agreements NRF-2019R1A2C2002724 and NRF-2022R1A2C1003322.

Before we state our main theorems, let us introduce some basic definitions. We define the parabolic distance between $X = (t, x)$ and $Y = (s, y)$ in \mathbb{R}^{d+1} by

$$|X - Y| = \max(|x - y|, \sqrt{|t - s|}).$$

We define the $(d + 1)$ -dimensional cylinders $Q_r(X)$, $Q_r^+(X)$, and $Q_r^-(X)$, by

$$\begin{aligned} Q_r(X) &= \{Y \in \mathbb{R}^{d+1} : |Y - X| < r\} = (s - r^2, s + r^2) \times B_r(x), \\ Q_r^+(X) &= (s, s + r^2) \times B_r(x), \quad \text{and} \quad Q_r^-(X) = (s - r^2, s) \times B_r(x). \end{aligned}$$

For $X = (t, x) \in \mathbb{R}^{d+1}$ and $r > 0$, we define

$$\omega_{\mathbf{A}}^x(r, X) := \oint_{Q_r^-(X)} |\mathbf{A}(s, y) - \bar{\mathbf{A}}_{x,r}^x(s)| dy ds, \quad \text{where} \quad \bar{\mathbf{A}}_{x,r}^x(s) := \oint_{B_r(x)} \mathbf{A}(s, y) dy.$$

Then for a subset Q of \mathbb{R}^{d+1} , we define

$$\omega_{\mathbf{A}}^x(r, Q) := \sup \{ \omega_{\mathbf{A}}^x(r, X) : X \in Q \} \quad \text{and} \quad \omega_{\mathbf{A}}^x(r) := \omega_{\mathbf{A}}^x(r, \mathbb{R}^{d+1}).$$

We say that \mathbf{A} is of *Dini mean oscillation in x* over Q and write $\mathbf{A} \in \text{DMO}_x(Q)$ if $\omega_{\mathbf{A}}^x(r, Q)$ satisfies the Dini condition

$$\int_0^1 \frac{\omega_{\mathbf{A}}^x(r, Q)}{r} dr < +\infty.$$

The adjoint operator P^* is given by

$$P^*u = -\partial_t u - D_{ij}(a^{ij}(t, x)u).$$

We are now ready to state the main results.

Theorem 1.1. *Assume that $\mathbf{A} = (a^{ij})$ satisfies (1.1) and belongs to $\text{DMO}_x(\mathbb{R}^{d+1})$. Then, there exist unique fundamental solutions $\Gamma(X, Y) = \Gamma(t, x, s, y)$ and $\Gamma^*(X, Y) = \Gamma^*(t, x, s, y)$ for the operators P and P^* , respectively, and they satisfy the symmetry relation*

$$\Gamma(t, x, s, y) = \Gamma^*(s, y, t, x). \quad (1.2)$$

The fundamental solution Γ is continuous in $\mathbb{R}^{d+1} \times \mathbb{R}^{d+1} \setminus \{(X, X) : X \in \mathbb{R}^{d+1}\}$ and

$$\Gamma(t, x, s, y) = 0 \quad \text{if } t < s.$$

Also, for each $Y \in \mathbb{R}^{d+1}$, $D_x \Gamma(\cdot, Y)$ and $D_x^2 \Gamma(\cdot, Y)$ are continuous in $\mathbb{R}^{d+1} \setminus \{Y\}$; if \mathbf{A} is continuous, then $\partial_t \Gamma(\cdot, Y)$ is continuous in $\mathbb{R}^{d+1} \setminus \{Y\}$ as well. Moreover, for any $R_0 > 0$, there exist constants $C = C(d, \lambda, \Lambda, \omega_{\mathbf{A}}^x, R_0)$ such that we have

$$|\Gamma(X, Y)| \leq C|X - Y|^{-d} \quad (1.3)$$

for any $X, Y \in \mathbb{R}^{d+1}$ satisfying $0 < |X - Y| < R_0$.

Remark 1.2. In Theorem 1.1, in addition to (1.3), we also have pointwise bounds for the derivatives of the fundamental solutions, that is,

$$|D_x \Gamma(X, Y)| \leq C|X - Y|^{-d-1}, \quad |\partial_t \Gamma(X, Y)| + |D_x^2 \Gamma(X, Y)| \leq C|X - Y|^{-d-2}$$

for any $X, Y \in \mathbb{R}^{d+1}$ satisfying $0 < |X - Y| < R_0$. These estimates follow directly from (1.3) and [3, Theorem 3.2] applied to $\Gamma(\cdot, Y)$ in $Q_R^-(X)$ with $R = \frac{1}{2}|X - Y|$.

We recall that \mathbf{A} belongs to $\text{VMO}_x(\mathbb{R}^{d+1})$ if and only if $\lim_{r \rightarrow 0} \omega_{\mathbf{A}}^x(r) = 0$ (see, e.g., [13]) and thus $\text{VMO}_x(\mathbb{R}^{d+1})$ contains $\text{DMO}_x(\mathbb{R}^{d+1})$.

Theorem 1.3 (Sub-Gaussian estimate). *Assume that $\mathbf{A} = (a^{ij})$ satisfies (1.1) and belongs to $\text{VMO}_x(\mathbb{R}^{d+1})$. Suppose there exists a fundamental solution $\Gamma(t, x, s, y)$ for the operator P , which satisfies (1.3). Then, for any $T > 0$ and $\delta \in (0, 1)$, there exist a constant $C = C(d, \lambda, \Lambda, \omega_{\mathbf{A}}^x, \delta, T)$ and a universal constant $\beta > 0$ such that for any $x, y \in \mathbb{R}^d$ and $t, s \in \mathbb{R}$ satisfying $0 < t - s < T$ we have*

$$|\Gamma(t, x, s, y)| \leq \frac{C}{(t-s)^{d/2}} \exp \left\{ -\beta \left(\frac{|x-y|}{\sqrt{t-s}} \right)^{2-\delta} \right\}. \quad (1.4)$$

We shall say that \mathbf{A} is uniformly Dini continuous in x over \mathbb{R}^{d+1} if its modulus of continuity in x defined by

$$\varrho_{\mathbf{A}}^x(r) := \sup \{ |\mathbf{A}(t, x) - \mathbf{A}(t, y)| : x, y \in \mathbb{R}^d, t \in \mathbb{R}, |x - y| \leq r \}$$

satisfies the Dini condition

$$\int_0^1 \frac{\varrho_{\mathbf{A}}^x(r)}{r} dr < +\infty.$$

It is clear that if \mathbf{A} is uniformly Dini continuous in x over \mathbb{R}^{d+1} , then it is of Dini mean oscillation in x over \mathbb{R}^{d+1} .

Theorem 1.4 (Gaussian estimate). *Assume that $\mathbf{A} = (a^{ij})$ satisfies (1.1) and $\mathbf{A} = (a^{ij})$ is uniformly Dini continuous in x over \mathbb{R}^{d+1} . Then the fundamental solution satisfies the Gaussian bounds, that is, for any $T > 0$, there exists $C = C(d, \lambda, \Lambda, T, \varrho_{\mathbf{A}}^x)$ and $\kappa = \kappa(\lambda, \Lambda)$ such that for any $x, y \in \mathbb{R}^d$ and $t, s \in \mathbb{R}$ satisfying $0 < t - s < T$ we have*

$$|\Gamma(t, x, s, y)| \leq \frac{C}{(t-s)^{d/2}} \exp \left\{ -\kappa \frac{|x-y|^2}{t-s} \right\}. \quad (1.5)$$

A few remarks are in order. The fundamental solutions are topics in many classical books. See, e.g., [4, 8, 11, 16] and references therein. It is well known that the fundamental solutions of second order parabolic equations in divergence form have two-sided Gaussian bounds even in the case when the coefficients are just bounded and measurable; see [1]. In contrast to parabolic equations in divergence form, the fundamental solutions of parabolic equations in non-divergence form do not necessarily have the Gaussian bounds if the coefficients do not possess some kind of regularity, although certain pointwise bounds are available in terms of so-called normalized adjoint solutions; see [5]. As a matter of fact, even if the coefficients are continuous in t and x , the following weaker estimate may not hold:

$$|\Gamma(t, x, s, y)| \leq C(t-s)^{-d/2} \quad \text{for } 0 < t - s < T.$$

A counterexample is given in [10] for the equation (in one space variable) $\partial u / \partial t = a(t, x) \partial^2 u / \partial x^2$ with a coefficient $a(t, x)$, continuous in t and x , and satisfying $1/2 \leq a(t, x) \leq 3/2$, whose fundamental solution is unbounded at any given point x_0 for any $t > 0$. See also [7] and [19] for examples of equations with continuous coefficients, whose fundamental solutions (as measure) are singular with respect to Lebesgue measure for any $t > 0$. On the other hand, if the coefficients are of Dini mean oscillation in x , then the fundamental solutions have the usual Gaussian bounds; see [3]. However, the proof there relies heavily on the Harnack type properties of nonnegative (adjoint) solutions and is not applicable to the systems setting. To the best of our knowledge, the Gaussian bounds for the fundamental solutions were available to the non-scalar setting if the coefficients are continuous

in t and x , and additionally, if they are doubly Dini continuous in x . See [18] and [4].

We give a brief description of the methods we use in the proofs. To show Theorem 1.1, we adapt an argument in [12] for non-divergence form elliptic equations, by using the pointwise estimates of solutions established in [3]. In the proof of Theorem 1.3, we first establish an exponential decay estimate by using the $W_p^{1,2}$ estimate and an iteration argument. We then improve the exponential decay estimate to the sub-Gaussian estimate (1.4) by exploiting the semi-group property of the fundamental solution together with a delicate re-scaling argument. Finally, we modify the parametrix method of Levi [17] to prove the Gaussian estimate (1.5) in Theorem 1.4. The main difference between Levi's original method and ours is that Levi's procedure was intended to construct the fundamental solution and thus required more restriction on the coefficients while in our approach, we construct the fundamental solution by different means and prove that it is identical with the resulting kernel produced by our modified parametrix method, which inherits the Gaussian bounds from the fundamental solutions of parabolic operators with coefficients depending only on t . It is also worth mentioning that in contrast to the scalar case, we are only able to get a one-sided Gaussian estimate.

Finally, the organization of the paper is as follows. In Section 2, we state some preliminary definition and lemmas. The proofs of Theorems 1.1, 1.3, and 1.4 are given in Sections 3, 4, and 5, respectively.

2. PRELIMINARIES

For any domain $Q \subset \mathbb{R}^{d+1}$ and $p \in [1, \infty]$, we shall denote by $L_p(Q)$ the standard Lebesgue class. We define the function space

$$W_p^{1,2}(Q) = \{u : u, \partial_t u, Du, D^2u \in L_p(Q)\},$$

which are equipped with norm

$$\|u\|_{W_p^{1,2}(Q)} = \|u\|_{L_p(Q)} + \|Du\|_{L_p(Q)} + \|D^2u\|_{L_p(Q)} + \|\partial_t u\|_{L_p(Q)}.$$

We deal with the adjoint problem

$$P^*u = \operatorname{div}^2 \mathbf{g} + f \text{ in } (t_0, t_1) \times \mathbb{R}^d, \quad u(t_1, \cdot) = 0 \text{ on } \mathbb{R}^d, \quad (2.1)$$

where $\mathbf{g} = (g^{kl})$ is a symmetric $d \times d$ matrix-valued function and $\operatorname{div}^2 \mathbf{g} = D_{kl}g^{kl}$.

Definition 2.1. Assume that $\mathbf{g} \in L_p((t_0, t_1) \times \mathbb{R}^d)$ and $f \in L_p((t_0, t_1) \times \mathbb{R}^d)$, where $1 < p < \infty$. We say that $u \in L_p((t_0, t_1) \times \mathbb{R}^d)$ is a solution to (2.1) if u satisfies

$$\int_{t_0}^{t_1} \int_{\mathbb{R}^d} u P v = \int_{t_0}^{t_1} \int_{\mathbb{R}^d} f v + \operatorname{tr}(\mathbf{g} D^2 v) \quad (2.2)$$

for any $v \in W_{p'}^{1,2}((t_0, t_1) \times \mathbb{R}^d)$ satisfying $u(t_0, \cdot) = 0$, where $\frac{1}{p} + \frac{1}{p'} = 1$.

Lemma 2.2. Assume that $\mathbf{g} \in L_p((t_0, t_1) \times \mathbb{R}^d)$ and $f \in L_p((t_0, t_1) \times \mathbb{R}^d)$, where $1 < p < \infty$. Then there exists a unique solution v of the adjoint problem (2.1) in $L_p((t_0, t_1) \times \mathbb{R}^d)$ and it satisfies

$$\|v\|_{L_p((t_0, t_1) \times \mathbb{R}^d)} \leq C \left(\|\mathbf{g}\|_{L_p((t_0, t_1) \times \mathbb{R}^d)} + \|f\|_{L_p((t_0, t_1) \times \mathbb{R}^d)} \right),$$

where C is a constant depending only on $d, \lambda, \Lambda, p, t_0, t_1$, and $\omega_{\mathbf{A}}^x$.

Proof. Recall that $\text{DMO}_x(\mathbb{R}^{d+1}) \subset \text{VMO}_x(\mathbb{R}^{d+1})$. The existence and uniqueness of the solution to (2.1) is simple to derive by transposition from the unique existence of a solution $v \in W_{p'}^{1,2}((t_0, t_1) \times \mathbb{R}^d)$ to the direct problem

$$Pv = g \text{ in } (t_0, t_1) \times \mathbb{R}^d, \quad v(t_0, \cdot) = 0 \text{ in } \mathbb{R}^d,$$

and the corresponding $L_{p'}$ estimates:

$$\|D^2 v\|_{L_{p'}((t_0, t_1) \times \mathbb{R}^d)} + \|\partial_t v\|_{L_{p'}((t_0, t_1) \times \mathbb{R}^d)} + \|v\|_{L_{p'}((t_0, t_1) \times \mathbb{R}^d)} \leq C \|g\|_{L_{p'}((t_0, t_1) \times \mathbb{R}^d)}, \quad (2.3)$$

where $C = C(d, \lambda, \Lambda, t_0, t_1, \omega_A^x)$. See [13] and [6]. \blacksquare

Lemma 2.3. *Let $R_0 > 0$ and $\mathbf{g} = (g^{ij}) \in \text{DMO}_x(Q_{R_0}^+(X_0))$. Suppose v is an L_2 solution of*

$$P^* u = \text{div}^2 \mathbf{g} \text{ in } Q_{2r}^+(X_0),$$

where $0 < r \leq \frac{1}{2}R_0$. Then we have

$$\|u\|_{L_\infty(Q_r^+(X_0))} \leq C \left(\int_{Q_{2r}^+(X_0)} |u| + \int_0^r \frac{\omega_{\mathbf{g}}^x(\tau, Q_{2r}^+(X_0))}{\tau} d\tau \right),$$

where $C = C(d, \lambda, \Lambda, \omega_A^x, R_0)$.

Proof. The proof is essentially given in [3, Theorem 3.3]. cf. [12, Appendix]. \blacksquare

3. PROOF OF THEOREM 1.1

By adapting the argument in [12], we shall first construct the fundamental solution $\Gamma^*(Y, X) = \Gamma^*(s, y, t, x)$ for the adjoint operator P^* in Section 3.1. We then establish in Section 3.2 that

$$|\Gamma^*(Y, X)| \leq C|X - Y|^{-d} \text{ for all } X, Y \text{ satisfying } 0 < |X - Y| < R_0.$$

In Section 3.3, we construct the fundamental solution $\Gamma(t, x, s, y)$ of the operator P and show the symmetry relation (1.2), which in particular implies (1.3).

3.1. Construction of the adjoint fundamental solution. Fix a point $X_0 = (t_0, x_0)$ in \mathbb{R}^{d+1} . We construct fundamental solution $\Gamma^*(\cdot, X_0) = \Gamma^*(\cdot, \cdot, t_0, x_0)$ for the adjoint operator P^* with a pole at $X_0 = (t_0, x_0)$.

Lemma 3.1. *For any $r > 0$, $\{\bar{\mathbf{A}}_{x_0, 2^{-k}r}^x(\cdot)\}_{k=0}^\infty$ converges in $L_1((t_0 - r^2, t_0))$ to a function $\mathbf{A}_{x_0}(\cdot)$, which is symmetric and satisfies (1.1). Moreover,*

$$\int_{Q_r^-(X_0)} |\mathbf{A} - \mathbf{A}_{x_0}| \leq c(d) \int_0^r \frac{\omega_{\mathbf{A}}^x(s)}{s} ds. \quad (3.1)$$

Proof. By the triangle inequality,

$$\sum_{k=0}^\infty |\bar{\mathbf{A}}_{x_0, 2^{-k}r}^x(t) - \bar{\mathbf{A}}_{x_0, 2^{-k-1}r}^x(t)| \leq \sum_{k=0}^\infty 2^d \int_{B_{2^{-k}r}(x_0)} |\mathbf{A}(t, x) - \bar{\mathbf{A}}_{x_0, 2^{-k}r}^x(t)| dx.$$

Therefore, by the Fubini theorem, we have

$$\begin{aligned}
\sum_{k=0}^{\infty} \int_{t_0-r^2}^{t_0} |\bar{\mathbf{A}}_{x_0,2^{-k}r}^x(t) - \bar{\mathbf{A}}_{x_0,2^{-k-1}r}^x(t)| dt &\leq 2^d \sum_{k=0}^{\infty} \int_{t_0-r^2}^{t_0} \int_{B_{2^{-k}r}(x_0)} |\mathbf{A} - \bar{\mathbf{A}}_{x_0,2^{-k}r}^x(t)| dx dt \\
&\leq 2^d \sum_{k=0}^{\infty} \sum_{j=0}^{2^{2k}-1} \int_{t_0-(j+1)(2^{-k}r)^2}^{t_0-j(2^{-k}r)^2} \int_{B_{2^{-k}r}(x_0)} |\mathbf{A} - \bar{\mathbf{A}}_{x_0,2^{-k}r}^x(t)| dx dt \\
&\leq 2^d \sum_{k=0}^{\infty} \sum_{j=0}^{2^{2k}-1} (2^{-k}r)^2 \omega_{\mathbf{A}}^x(2^{-k}r) = 2^d r^2 \sum_{k=0}^{\infty} \omega_{\mathbf{A}}^x(2^{-k}r). \tag{3.2}
\end{aligned}$$

In view of the proof on [15, p. 495], we have $\omega_{\mathbf{A}}^x(t) \simeq \omega_{\mathbf{A}}^x(s)$ when $t \simeq s$. Thus, (3.2) implies

$$\sum_{k=0}^{\infty} \int_{t_0-r^2}^{t_0} |\bar{\mathbf{A}}_{x_0,2^{-k}r}^x(t) - \bar{\mathbf{A}}_{x_0,2^{-k-1}r}^x(t)| dt \leq C(d)r^2 \int_0^r \frac{\omega_{\mathbf{A}}^x(s)}{s} ds < \infty. \tag{3.3}$$

Therefore, $\{\bar{\mathbf{A}}_{x_0,2^{-k}r}^x(\cdot)\}$ is a Cauchy sequence in $L_1((t_0-r^2, t_0))$. Let $\mathbf{A}_{x_0}(\cdot)$ be the limit. Thus, from (3.3), we have

$$\int_{t_0-r^2}^{t_0} |\bar{\mathbf{A}}_{x_0,r}^x(t) - \mathbf{A}_{x_0}(t)| dt \leq C(d)r^2 \int_0^r \frac{\omega_{\mathbf{A}}^x(s)}{s} ds. \tag{3.4}$$

Finally, by using the triangle inequality and (3.4),

$$\begin{aligned}
\int_{Q_r^-(X_0)} |\mathbf{A} - \mathbf{A}_{x_0}| &\leq \int_{Q_r^-(X_0)} |\mathbf{A} - \bar{\mathbf{A}}_{x_0,r}^x(t)| + \int_{Q_r^-(X_0)} |\bar{\mathbf{A}}_{x_0,r}^x(t) - \mathbf{A}_{x_0}(t)| \\
&\leq \omega_{\mathbf{A}}^x(r) + \int_{t_0-r^2}^{t_0} |\bar{\mathbf{A}}_{x_0,r}^x(t) - \mathbf{A}(t, x_0)| dt \leq c(d) \int_0^r \frac{\omega_{\mathbf{A}}^x(s)}{s} ds. \quad \blacksquare
\end{aligned}$$

Remark 3.2. By a slight modification of the proof above, it is easily seen that \mathbf{A}_{x_0} is independent of t_0 and r . Moreover, if \mathbf{A} is continuous in x , then clearly $\mathbf{A}_{x_0}(t) = \mathbf{A}(t, x_0)$ for a.e. t .

We now consider the parabolic operator P_0 defined by

$$P_0 u := \partial_t u - a_0^{ij}(t) D_{ij} u = \partial_t u - \text{tr}(\mathbf{A}_{x_0}(t) D^2 u).$$

Let $\Phi(t, x, s, y)$ be the fundamental solution for P_0 . It is well known that there are positive constants $C_0 = C_0(d, \lambda, \Lambda)$ and $\kappa_0 = \kappa_0(\lambda, \Lambda)$ such that

$$|\Phi(t, x, s, y)| \leq C_0(t-s)^{-d/2} e^{-\kappa_0 \frac{|x-y|^2}{t-s}} \quad \text{for } t > s \tag{3.5}$$

and $\Phi(t, x, s, y) \equiv 0$ if $t < s$. See, for instance, [8, Chapter 9] and [14, Chapter 2]. Since \mathbf{A}_{x_0} does not depend on x , we also have

$$\Phi(t, x, s, y) = \Phi^*(s, y, t, x), \tag{3.6}$$

where Φ^* is fundamental solution for the adjoint operator P_0^* given by

$$P_0^* u := -\partial_t u - D_{ij}(a_0^{ij}(t)u) = -\partial_t u - \text{div}^2(\mathbf{A}_{x_0}(t)u).$$

Note that, if we set $v = \Gamma^*(\cdot, X_0) - \Phi^*(\cdot, X_0)$, then it would satisfy

$$\begin{aligned}
P^* v &= P^* \Gamma^*(\cdot, X_0) - P^* \Phi^*(\cdot, X_0) + P_0^* \Phi^*(\cdot, X_0) - P_0^* \Phi^*(\cdot, X_0) \\
&= \text{div}^2((\mathbf{A} - \mathbf{A}_{x_0})\Phi^*(\cdot, X_0)).
\end{aligned}$$

We are thus lead to consider the problem

$$P^*v = \operatorname{div}^2 \mathbf{g} \text{ in } (t_0 - T, t_0) \times \mathbb{R}^d, \quad v(t_0, \cdot) = 0 \text{ on } \mathbb{R}^d, \quad (3.7)$$

where $T > 0$ and

$$\mathbf{g} := (\mathbf{A} - \mathbf{A}_{x_0})\Phi^*(\cdot, X_0).$$

By a straightforward computation using (3.5) and (3.6), for $p \in (0, \frac{d+2}{d})$, we have

$$\begin{aligned} \int_{t_0-T}^{t_0} \int_{\mathbb{R}^d} |\mathbf{g}|^p dx dt &\leq \|\mathbf{A} - \mathbf{A}_{x_0}\|_\infty^p \int_0^T \int_{\mathbb{R}^d} (C_0 t^{-d/2} e^{-\kappa_0 |x|^2/t})^p dx dt \\ &= \|\mathbf{A} - \mathbf{A}_{x_0}\|_\infty^p C_0^p \int_0^T t^{d/2-dp/2} \int_{\mathbb{R}^d} t^{-d/2} e^{-\kappa_0 p |x|^2/t} dx dt \\ &= \|\mathbf{A} - \mathbf{A}_{x_0}\|_\infty^p C_0^p \int_{\mathbb{R}^d} e^{-\kappa_0 p |x|^2} dx \int_0^T t^{d/2-dp/2} dt = CT^{\frac{2+d-dp}{2}}, \end{aligned}$$

where $C = C(d, \lambda, \Lambda, p)$. We just proved that

$$\mathbf{g} \in L_p((t_0 - T, t_0) \times \mathbb{R}^d), \quad \forall T > 0, \quad \forall p \in (0, \frac{d+2}{d}). \quad (3.8)$$

Therefore, for $1 < p < \frac{d+2}{d}$, by Lemma 2.2 there is a unique L_p solution v of the problem (3.7). By extending $v = 0$ on $(t_0, \infty) \times \mathbb{R}^d$ and letting $T \rightarrow \infty$, we may assume that v is defined on the entire \mathbb{R}^{d+1} .

Lemma 3.3. *Let v be as above. The function $\Gamma^*(\cdot, X_0)$ defined by*

$$\Gamma^*(\cdot, X_0) = v + \Phi^*(\cdot, X_0)$$

is the fundamental solution of P^ with a pole at $X_0 = (t_0, x_0)$.*

Proof. For any $f \in C_c^\infty(\mathbb{R}^{d+1})$, fix a $T > |t_0|$ such that $(-T, T) \times \mathbb{R}^d$ contains the support of f . For $p' > \frac{d+2}{2}$, let $u \in W_p^{1,2}((-T, T) \times \mathbb{R}^d)$ be the solution of the problem $Pu = f$ with $u(-T, \cdot) = 0$. Then, by (3.7) we have

$$\begin{aligned} \int_{-T}^T \int_{\mathbb{R}^d} v f &= \int_{-T}^T \int_{\mathbb{R}^d} v Pu = \int_{-T}^T \int_{\mathbb{R}^d} (P^*v)u \\ &= \int_{-T}^T \int_{\mathbb{R}^d} \operatorname{tr}((\mathbf{A} - \mathbf{A}_{x_0})\Phi^*(\cdot, X_0)D^2u) = \int_{-T}^T \int_{\mathbb{R}^d} \Phi^*(\cdot, X_0)(P_0u - Pu) \\ &= \int_{-T}^T \int_{\mathbb{R}^d} \Phi^*(\cdot, X_0)P_0u - \int_{-T}^T \int_{\mathbb{R}^d} \Phi^*(\cdot, X_0)f = u(X_0) - \int_{-T}^T \int_{\mathbb{R}^d} \Phi^*(\cdot, X_0)f, \end{aligned}$$

where in the last equality we use the fact that $\Phi^*(\cdot, X_0)$ is the fundamental solution for P_0^* . Therefore, we have

$$u(X_0) = \int_{-T}^T \int_{\mathbb{R}^d} \Gamma^*(\cdot, X_0)f = \int_{\mathbb{R}^{d+1}} \Gamma^*(\cdot, X_0)f, \quad (3.9)$$

which means that $\Gamma^*(\cdot, X_0)$ is the fundamental solution for P^* with a pole at X_0 . ■

Noting that $\Gamma^*(s, y, t, s) = 0$ for $t < s$, we actually proved the following.

Proposition 3.4. *For $p > \frac{d+2}{2}$ and $f \in L_p((t_0, t_1) \times \mathbb{R}^d)$, if $u \in W_p^{1,2}((t_0, t_1) \times \mathbb{R}^d)$ is the solution of $Pu = f$ in $(t_0, t_1) \times \mathbb{R}^d$ satisfying $u(t_0, \cdot) = 0$, then we have the representation formula*

$$u(t, x) = \int_{t_0}^t \int_{\mathbb{R}^d} \Gamma^*(s, y, t, x)f(s, y) dy ds.$$

3.2. Pointwise bound for the adjoint fundamental solution. Let $R_0 > 0$ be fixed but arbitrary. We shall show that there exists a constant C depending R_0 as well as on d, λ, Λ , and ω_A^\times such that we have

$$|\Gamma^*(X, X_0)| \leq C|X - X_0|^{-d} \quad \text{for all } X \text{ satisfying } 0 < |X - X_0| < R_0. \quad (3.10)$$

Define \mathbf{g}_1 and \mathbf{g}_2 by setting

$$\mathbf{g}_1 = \zeta(\mathbf{A} - \mathbf{A}_{x_0})\Phi^*(\cdot, X_0) \quad \text{and} \quad \mathbf{g}_2 = (1 - \zeta)(\mathbf{A} - \mathbf{A}_{x_0})\Phi^*(\cdot, X_0), \quad (3.11)$$

where ζ is a smooth function on \mathbb{R}^{d+1} such that

$$0 \leq \zeta \leq 1, \quad \zeta = 0 \text{ in } Q_R(X_0), \quad \zeta = 1 \text{ in } \mathbb{R}^{d+1} \setminus Q_{2R}(X_0), \quad |D_x \zeta| \leq 4/R,$$

and $R > 0$ is a constant to be fixed later. Since $\Phi^*(\cdot, X_0)$ vanished on $(t_0, \infty) \times \mathbb{R}^d$, we see that

$$\mathbf{g}_1 = \mathbf{g}_2 \equiv 0 \quad \text{on } (t_0, \infty) \times \mathbb{R}^d. \quad (3.12)$$

Also, by (3.5) and (3.6), there is a positive constant $C'_0 = C'_0(d, \lambda, \Lambda)$ such that

$$|\Phi^*(X, X_0)| \leq C'_0|X - X_0|^{-d}, \quad \forall X \neq X_0. \quad (3.13)$$

In the following lemmas, we show that $\mathbf{g}_1 \in L_{p_1}$ for $p_1 > \frac{d+2}{d}$ and $\mathbf{g}_2 \in L_{p_2}$ for $1 \leq p_2 < \frac{d+2}{d}$.

Lemma 3.5. *For $p > \frac{d+2}{d}$, there is a constant $C = C(d, \lambda, \Lambda, p)$ such that*

$$\|\mathbf{g}_1\|_{L_p(\mathbb{R}^{d+1})} \leq CR^{\frac{d+2}{p}-d}. \quad (3.14)$$

For $1 \leq p < \frac{d+2}{d}$, there is a constant $C = C(d, \lambda, \Lambda, p)$ such that

$$\|\mathbf{g}_2\|_{L_p(\mathbb{R}^{d+1})} \leq C \left(\int_0^{2R} \frac{\omega_A^\times(s)}{s} ds \right)^{1/p} R^{\frac{d+2}{p}-d}. \quad (3.15)$$

Proof. Note that $\|\mathbf{A} - \mathbf{A}_{x_0}\|_\infty \leq C(d, \Lambda)$. Therefore, if $p > \frac{d+2}{d}$, we get from (3.13) that

$$\begin{aligned} \int_{\mathbb{R}^{d+1}} |\mathbf{g}_1|^p &\leq C \sum_{k=0}^{\infty} \int_{Q_{2^{k+1}R}(X_0) \setminus Q_{2^kR}(X_0)} |X - X_0|^{-dp} dX \\ &\leq C \sum_{k=0}^{\infty} (2^k R)^{-dp} (2^{k+1} R)^{d+2} \leq CR^{d+2-dp}. \end{aligned}$$

When $1 \leq p < \frac{d+2}{d}$, by (3.12), (3.13), the properties of ζ , and (3.1), we have

$$\begin{aligned} \int_{\mathbb{R}^{d+1}} |\mathbf{g}_2|^p &\leq C \sum_{k=0}^{\infty} \int_{Q_{2^{1-k}R}^-(X_0) \setminus Q_{2^{-k}R}^-(X_0)} |\mathbf{A} - \mathbf{A}_{x_0}|^p |X - X_0|^{-dp} dX \\ &\leq C \sum_{k=0}^{\infty} (2^{-k} R)^{-dp} \int_{Q_{2^{1-k}R}^-(X_0)} |\mathbf{A} - \mathbf{A}_{x_0}| \\ &\leq C \sum_{k=0}^{\infty} (2^{-k} R)^{-dp} (2^{1-k} R)^{d+2} \int_0^{2^{1-k}R} \frac{\omega_A^\times(s)}{s} ds \\ &\leq C \sum_{k=0}^{\infty} (2^{-k} R)^{d+2-dp} \int_0^{2R} \frac{\omega_A^\times(s)}{s} ds \leq CR^{d+2-dp} \int_0^{2R} \frac{\omega_A^\times(s)}{s} ds \end{aligned}$$

and the lemma follows. \blacksquare

Let v be the solution of the problem (3.7). Fix $p_1 \in (\frac{d+2}{d}, \infty)$ and $p_2 \in (1, \frac{d+2}{d})$ and let $v_i \in L_{p_i}((t_0 - R_0^2, t_0) \times \mathbb{R}^d)$ be the solution of the problems

$$P^*v_i = \operatorname{div}^2 \mathbf{g}_i \text{ in } (t_0 - R_0^2, t_0) \times \mathbb{R}^d, \quad v_i(t_0, \cdot) = 0 \text{ on } \mathbb{R}^d. \quad (i = 1, 2).$$

Then by Lemma 2.2 together with (3.14) and (3.15), respectively, we have

$$\|v_1\|_{L_{p_1}((t_0 - R_0^2, t_0) \times \mathbb{R}^d)} \leq CR^{(d+2)/p_1 - d} \quad (3.16)$$

and

$$\|v_2\|_{L_{p_2}((t_0 - R_0^2, t_0) \times \mathbb{R}^d)} \leq C \left(\int_0^{2R} \frac{\omega_{\mathbf{A}}^x(r)}{r} dr \right)^{1/p_2} R^{(d+2)/p_2 - d}. \quad (3.17)$$

We note that the constant C in the above depends on R_0 as well as on $d, \lambda, \Lambda, p_1, p_2$, and $\omega_{\mathbf{A}}^x$. By the same computation as in (3.8), we find $\mathbf{g}_1 \in L_{p_2}((t_0 - R_0^2, t_0) \times \mathbb{R}^d)$ as well, and thus $v_1 \in L_{p_2}((t_0 - R_0^2, t_0) \times \mathbb{R}^d)$. Therefore, by the uniqueness, we see that

$$v = v_1 + v_2.$$

We extend v_1 and v_2 by zero on $(t_0, \infty) \times \mathbb{R}^d$.

Now, for any fixed $Y_0 = (s_0, y_0)$ with $0 < |Y_0 - X_0| < R_0$, we take

$$R = \frac{1}{5}|Y_0 - X_0|$$

and estimate $v_1(Y_0)$ and $v_2(Y_0)$ by using Lemma 2.3 as follows:

$$|v_i(Y_0)| \leq C \int_{Q_{2R}^+(Y_0)} |v_i| + C \int_0^R \frac{\omega_{\mathbf{g}_i}^x(r, Q_{2R}^+(Y_0))}{r} dr. \quad (i = 1, 2). \quad (3.18)$$

By Hölder's inequality, (3.16), and (3.17), we have

$$\begin{aligned} \int_{Q_{2R}^+(Y_0)} |v_1| &\leq CR^{-(d+2)/p_1} \|v_1\|_{L_{p_1}} \leq CR^{-d}, \\ \int_{Q_{2R}^+(Y_0)} |v_2| &\leq CR^{-(d+2)/p_2} \|v_2\|_{L_{p_2}} \leq CR^{-d} \left(\int_0^{2R} \frac{\omega_{\mathbf{A}}^x(r)}{r} dr \right)^{1/p_2}. \end{aligned} \quad (3.19)$$

Lemma 3.6. Suppose $R := \frac{1}{5}|Y_0 - X_0| > 0$ and let η be a Lipschitz function on \mathbb{R}^{d+1} such that $0 \leq \eta \leq 1$ and $|D_x \eta| \leq 4/R$. Set

$$\mathbf{g} = \eta(\mathbf{A} - \mathbf{A}_{x_0})\Phi^*(\cdot, X_0).$$

Then, for any $r \in (0, R]$ we have

$$\omega_{\mathbf{g}}^x(r, Q_{2R}^+(Y_0)) \leq CR^{-d} \left(\omega_{\mathbf{A}}^x(r) + \frac{r}{R} \int_0^r \frac{\omega_{\mathbf{A}}^x(s)}{s} ds \right),$$

where $C = C(d, \lambda, \Lambda)$.

Proof. Let us denote

$$\Phi_0(X) = \Phi_0(t, x) = \Phi^*(t, x, t_0, x_0) = \Phi^*(X, X_0).$$

For $Z = (\tau, \xi) \in Q_{2R}^+(Y_0)$ and $0 < r \leq R$, we have

$$\begin{aligned} \int_{Q_r^-(Z)} |\mathbf{g} - \bar{\mathbf{g}}_{\xi,r}^x| &= \int_{Q_r^-(Z)} \left| \eta(\mathbf{A} - \mathbf{A}_{x_0})\Phi_0 - \overline{(\eta(\mathbf{A} - \mathbf{A}_{x_0})\Phi_0)}_{\xi,r}^x \right| \\ &\leq \int_{Q_r^-(Z)} \left| (\mathbf{A} - \mathbf{A}_{x_0})\eta\Phi_0 - \overline{(\mathbf{A} - \mathbf{A}_{x_0})}_{\xi,r}^x \eta\Phi_0 \right| \\ &\quad + \int_{Q_r^-(Z)} \left| \overline{(\mathbf{A} - \mathbf{A}_{x_0})}_{\xi,r}^x \eta\Phi_0 - \overline{((\mathbf{A} - \mathbf{A}_{x_0})\eta\Phi_0)}_{\xi,r}^x \right| \\ &=: I + II. \end{aligned}$$

Note that by the triangle inequality, $|X - X_0| \geq 2R$ for any $X \in Q_r^-(Z)$ and thus, we have

$$|\Phi_0(X)| + R|D_x\Phi_0(X)| \leq CR^{-d}, \quad \forall X \in Q_r^-(Z), \quad (3.20)$$

where $C = C(d, \lambda, \Lambda)$. Here the bound of $D_x\Phi_0(X)$ is due to the fact that \mathbf{A}_{x_0} only depends on t . See (3.13) and (5.1). Therefore, we have

$$\begin{aligned} I &\leq \int_{Q_r^-(Z)} \left| (\mathbf{A} - \mathbf{A}_{x_0}) - \overline{(\mathbf{A} - \mathbf{A}_{x_0})}_{\xi,r}^x \right| |\Phi_0| \\ &\leq \int_{Q_r^-(Z)} CR^{-d} |\mathbf{A} - \bar{\mathbf{A}}_{\xi,r}^x| \leq CR^{-d} \omega_{\mathbf{A}}^x(r). \end{aligned} \quad (3.21)$$

Also, we have

$$\begin{aligned} II &= \int_{Q_r^-(Z)} \left| \int_{B_r(\xi)} (\mathbf{A}(t, y) - \mathbf{A}_{x_0}(t)) (\eta(t, x)\Phi_0(t, x) - \eta(t, y)\Phi_0(t, y)) dy \right| dx dt \\ &\leq \int_{Q_r^-(Z)} \int_{B_r(\xi)} |\mathbf{A}(t, y) - \mathbf{A}_{x_0}(t)| |\eta(t, x)\Phi_0(t, x) - \eta(t, y)\Phi_0(t, y)| dy dx dt. \end{aligned} \quad (3.22)$$

By using (3.20), and the properties of η , for $(t, x) \in Q_r^-(Z)$ and $y \in B_r(\xi)$, we have

$$\begin{aligned} &|\eta(t, x)\Phi_0(t, x) - \eta(t, y)\Phi_0(t, y)| \\ &\leq |\eta(t, x)| |\Phi_0(t, x) - \Phi_0(t, y)| + |\eta(t, x) - \eta(t, y)| |\Phi_0(t, y)| \\ &\leq CrR^{-d-1} + C(r/R)R^{-d} \leq CrR^{-d-1}. \end{aligned} \quad (3.23)$$

Plugging (3.23) into (3.22), we obtain

$$II \leq CrR^{-d-1} \int_{Q_r^-(Z)} |\mathbf{A}(t, y) - \mathbf{A}_{x_0}(t)| dy dt.$$

We claim that

$$\int_{Q_r^-(Z)} |\mathbf{A}(t, x) - \mathbf{A}_{x_0}(t)| dx dt \leq C \left(\frac{R\omega_{\mathbf{A}}^x(r)}{r} + \int_0^r \frac{\omega_{\mathbf{A}}^x(s)}{s} ds \right), \quad (3.24)$$

where $C = C(d, \lambda, \Lambda)$. Assume the claim for now. Then, we have

$$II \leq CrR^{-d-1} \left(\frac{R\omega_{\mathbf{A}}^x(r)}{r} + \int_0^r \frac{\omega_{\mathbf{A}}^x(s)}{s} ds \right). \quad (3.25)$$

Combining (3.21) and (3.25), we have (recall $r \leq R$)

$$\omega_{\mathbf{g}}^x(r, Z) \leq I + II \leq CR^{-d} \left(\omega_{\mathbf{A}}^x(r) + \frac{r}{R} \int_0^r \frac{\omega_{\mathbf{A}}^x(s)}{s} ds \right).$$

The lemma is proved by taking supremum over $Z \in Q_{2R}^+(Y_0)$.

It remains to prove the claim (3.24). Note that we can choose a sequence of points x_1, x_2, \dots, x_N in \mathbb{R}^d with $x_N = \xi$ so that $|x_{i-1} - x_i| \leq r$ for $i = 1, \dots, N$ and

$$N = \lceil 7R/r \rceil \leq 8R/r. \quad (3.26)$$

Then by using the triangle inequality, we have

$$|\mathbf{A}(t, x) - \mathbf{A}_{x_0}(t)| \leq |\mathbf{A}(t, x) - \bar{\mathbf{A}}_{\xi, r}^x(t)| + \sum_{i=1}^N |\bar{\mathbf{A}}_{x_i, r}^x(t) - \bar{\mathbf{A}}_{x_{i-1}, r}^x(t)| + |\bar{\mathbf{A}}_{x_0, r}^x(t) - \mathbf{A}_{x_0}(t)|. \quad (3.27)$$

Note that by (3.4), we have

$$\int_{\tau-r^2}^{\tau} |\bar{\mathbf{A}}_{x_0, r}^x(t) - \mathbf{A}_{x_0}(t)| dt \leq c(d) \int_0^r \frac{\omega_{\mathbf{A}}^x(s)}{s} ds. \quad (3.28)$$

Also, by averaging the following triangle inequality

$$|\bar{\mathbf{A}}_{x_i, r}^x(t) - \bar{\mathbf{A}}_{x_{i-1}, r}^x(t)| \leq |\mathbf{A}(t, x) - \bar{\mathbf{A}}_{x_i, r}^x(t)| + |\mathbf{A}(t, x) - \bar{\mathbf{A}}_{x_{i-1}, r}^x(t)|$$

over $x \in B_r(x_{i-1}) \cap B_r(x_i)$ and using $|x_{i-1} - x_i| \leq r$, we find that

$$|\bar{\mathbf{A}}_{x_i, r}^x(t) - \bar{\mathbf{A}}_{x_{i-1}, r}^x(t)| \leq c(d) \left(\int_{B_r(x_i)} |\mathbf{A}(t, x) - \bar{\mathbf{A}}_{x_i, r}^x(t)| dx + \int_{B_r(x_{i-1})} |\mathbf{A}(t, x) - \bar{\mathbf{A}}_{x_{i-1}, r}^x(t)| dx \right).$$

Then, by averaging the last inequality over $t \in (\tau - r^2, \tau)$, we get

$$\int_{\tau-r^2}^{\tau} |\bar{\mathbf{A}}_{x_i, r}^x(t) - \bar{\mathbf{A}}_{x_{i-1}, r}^x(t)| dt \leq c(d) \omega_{\mathbf{A}}^x(r), \quad i = 1, \dots, N. \quad (3.29)$$

Finally, averaging the inequality (3.27) over $X = (t, x) \in Q_r^-(Z)$ and using (3.28), (3.29), and (3.26), we obtain

$$\int_{Q_r^-(Z)} |\mathbf{A}(t, x) - \mathbf{A}_{x_0}(t)| dx dt \leq \omega_{\mathbf{A}}^x(r) + c(d) \frac{8R}{r} \omega_{\mathbf{A}}^x(r) + c(d) \int_0^r \frac{\omega_{\mathbf{A}}^x(s)}{s} ds,$$

from which (3.24) follows. \blacksquare

Applying Lemma 3.6 with $\eta = \zeta$ and $\eta = 1 - \zeta$, respectively, we get

$$\begin{aligned} \int_0^R \frac{\omega_{\mathbf{g}}^x(r, Q_{2R}^+(Y_0))}{r} dr &\leq CR^{-d} \left(\int_0^R \frac{\omega_{\mathbf{A}}^x(r)}{r} dr + \frac{1}{R} \int_0^R \int_0^r \frac{\omega_{\mathbf{A}}^x(s)}{s} ds dr \right) \\ &\leq CR^{-d} \int_0^R \frac{\omega_{\mathbf{A}}^x(s)}{s} ds. \end{aligned} \quad (3.30)$$

Putting (3.30) back to (3.18) together with (3.19), we get

$$\begin{aligned} |v_1(Y_0)| + |v_2(Y_0)| &\leq CR^{-d} \left(1 + \left(\int_0^{2R} \frac{\omega_{\mathbf{A}}^x(s)}{s} ds \right)^{1/p_2} + \int_0^R \frac{\omega_{\mathbf{A}}^x(s)}{s} ds \right) \\ &\leq C \left(1 + \int_0^{R_0} \frac{\omega_{\mathbf{A}}^x(s)}{s} ds \right) R^{-d} \leq CR^{-d}. \end{aligned} \quad (3.31)$$

Therefore, by using (3.31) and recalling that $v = v_1 + v_2$ and $R = \frac{1}{5}|X_0 - Y_0|$, we have

$$|v(Y_0)| \leq C|X_0 - Y_0|^{-d}, \quad (3.32)$$

where $C = C(d, \lambda, \Lambda, \omega_{\mathbf{A}}^x, R_0)$. Since

$$\Gamma^*(Y_0, X_0) = \Phi^*(Y_0, X_0) + v(Y_0)$$

and Y_0 satisfies $0 < |Y_0 - X_0| < R_0$, the estimate (3.10) follows from (3.32) and (3.13).

3.3. Construction of fundamental solution and the symmetry relation. We shall prove that the function $\Gamma(t, x, s, y)$ given by the formula (1.2) is the fundamental solution for the operator P .

For $Y = (s, y) \in \mathbb{R}^{d+1}$ and $\varepsilon > 0$, we first construct the approximate fundamental solution $\Gamma_\varepsilon(\cdot, Y)$ by following the strategy in [2]. Let $u = \Gamma_\varepsilon(\cdot, Y)$ be the solution of the problem

$$Pu = \frac{1}{|Q_\varepsilon^-(Y)|} \chi_{Q_\varepsilon^-(Y)} \text{ in } (s - \varepsilon^2, s + T) \times \mathbb{R}^d, \quad u(s - \varepsilon^2, \cdot) = 0 \text{ on } \mathbb{R}^d, \quad (3.33)$$

where $T \geq 1$ is fixed but arbitrary. By setting $\Gamma_\varepsilon(\cdot, Y) = 0$ on $(-\infty, s - \varepsilon^2) \times \mathbb{R}^d$ and letting $T \rightarrow \infty$, we extend the domain of $\Gamma_\varepsilon(\cdot, Y)$ to the entire \mathbb{R}^{d+1} .

Then by Proposition 3.4, we have

$$\Gamma_\varepsilon(X, Y) = \oint_{Q_\varepsilon^-(Y)} \Gamma^*(Z, X) dZ. \quad (3.34)$$

We conclude from (3.34) and (3.10) that for any $X, Y \in \mathbb{R}^{d+1}$ with $0 < |X - Y| < R_0$, we have

$$|\Gamma_\varepsilon(X, Y)| \leq C|X - Y|^{-d}, \quad \forall \varepsilon \in \left(0, \frac{1}{5}|X - Y|\right),$$

where C is a constant depending only on $d, \lambda, \Lambda, \omega_A^x$, and R_0 .

We construct fundamental solution for the operator P by modifying the method in [2]. Let $Y = (s, y) \in \mathbb{R}^{d+1}$ be fixed. For any $T \geq 1$, let us denote

$$R_T^{d+1} = (s - T, s + T) \times \mathbb{R}^d.$$

The following two lemmas are the adaptation of Lemmas 2.13 and 2.19 in [9] to the parabolic setting.

Lemma 3.7. *Let $p \in (1, \infty)$. For any $\varepsilon \in (0, 1)$, we have*

$$\int_{\mathbb{R}_T^{d+1} \setminus \overline{Q}_r(Y)} |\Gamma_\varepsilon(t, x, s, y)|^p dx dt \leq C r^{-pd+d+2}, \quad \forall r > 0 \text{ when } p > (d+2)/d, \quad (3.35)$$

$$\int_{\mathbb{R}_T^{d+1} \setminus \overline{Q}_r(Y)} |\partial_t \Gamma_\varepsilon(t, x, s, y)|^p + |D_x^2 \Gamma_\varepsilon(t, x, s, y)|^p dx dt \leq C r^{-(d+2)(p-1)}, \quad \forall r > 0, \quad (3.36)$$

where $C = C(d, \lambda, \Lambda, p, T, \omega_A^x)$.

Proof. We first establish (3.36). It is enough to consider the case when $r > 4\varepsilon$. Indeed, if $r \leq 4\varepsilon$, then by (2.3), we have

$$\int_{\mathbb{R}_T^{d+1}} |\partial_t \Gamma_\varepsilon(X, Y)|^p + |D_x^2 \Gamma_\varepsilon(X, Y)|^p dX \leq C \varepsilon^{-(d+2)(p-1)} \leq C r^{-(d+2)(p-1)}.$$

For $\mathbf{g} \in C_c^\infty(\mathbb{R}_T^{d+1} \setminus \overline{Q}_r(Y))$, let $u \in L_q(\mathbb{R}_T^{d+1})$ be the solution of the problem

$$P^*u = \operatorname{div}^2 \mathbf{g} \text{ in } \mathbb{R}_T^{d+1}, \quad u(s + T, \cdot) = 0 \text{ on } \mathbb{R}^d,$$

where $q = p/(p-1)$. Then by (2.2) we have

$$\oint_{Q_\varepsilon^-(Y)} u = \int_{\mathbb{R}_T^{d+1}} \operatorname{tr}(\mathbf{g} D^2 \Gamma_\varepsilon(\cdot, Y)). \quad (3.37)$$

Since $\mathbf{g} = 0$ in $Q_r(Y)$, we see that u is continuous on $\overline{Q_{r/2}}(Y)$ by [3, Theorem 3.3]. Note that if $Z \in Q_\varepsilon^-(Y)$, then $Q_{r/2}^+(Z) \subset Q_r(Y)$. It follows from Lemma 2.3 that

$$\|u\|_{L_\infty(Q_{r/4}^+(Z))} \leq Cr^{-d-2}\|u\|_{L_1(Q_{r/2}^+(Z))} \leq Cr^{-d-2}\|u\|_{L_1(Q_r(Y))}. \quad (3.38)$$

Therefore, by Hölder's inequality and Lemma 2.2, we have

$$\|u\|_{L_\infty(Q_\varepsilon^-(Y))} \leq Cr^{-\frac{d+2}{q}}\|u\|_{L_q(Q_r(Y))} \leq Cr^{-\frac{d+2}{q}}\|u\|_{L_q(\mathbb{R}_T^{d+1})} \leq Cr^{-\frac{d+2}{q}}\|\mathbf{g}\|_{L_q(\mathbb{R}_T^{d+1})}.$$

Since \mathbf{g} is supported in $\mathbb{R}_T^{d+1} \setminus \overline{Q_r}(Y)$, by (3.37) and the above estimate, we have

$$\left| \int_{\mathbb{R}_T^{d+1} \setminus \overline{Q_r}(Y)} \text{tr}(\mathbf{g} D^2 \Gamma_\varepsilon(\cdot, Y)) \right| \leq Cr^{-\frac{d+2}{q}}\|\mathbf{g}\|_{L_q(\mathbb{R}_T^{d+1} \setminus \overline{Q_r}(Y))}.$$

Therefore, by duality, we have

$$\int_{\mathbb{R}_T^{d+1} \setminus \overline{Q_r}(Y)} |D_x^2 \Gamma_\varepsilon(t, x, s, y)|^p dx dt \leq Cr^{-(d+2)(p-1)}.$$

Then the estimate (3.36) follows from the last inequality and the fact that $P \Gamma_\varepsilon(\cdot, Y) = 0$ in $\mathbb{R}^{d+1} \setminus \overline{Q_r}(Y)$.

Next, we turn to the proof of (3.35). Again, it is enough to consider the case when $r > 4\varepsilon$ because by (2.3) and the parabolic Sobolev embedding, we have

$$\|\Gamma_\varepsilon(\cdot, Y)\|_{L_p(\mathbb{R}_T^{d+1})} \leq C\|\Gamma_\varepsilon(\cdot, Y)\|_{W_{p(d+2)/(d+2+2p)}^{1,2}(\mathbb{R}_T^{d+1})} \leq C\varepsilon^{-d+(d+2)/p} \leq Cr^{-d+(d+2)/p},$$

where in the last inequality we used the fact that $-d + (d+2)/p < 0$. For $f \in C_c^\infty(\mathbb{R}_T^{d+1} \setminus \overline{Q_r}(Y))$, let $u \in L_q(\mathbb{R}_T^{d+1})$ be the solution of the problem

$$P^*u = f \text{ in } \mathbb{R}_T^{d+1}, \quad u(s+T, \cdot) = 0 \text{ on } \mathbb{R}^d.$$

Then by (2.2) we have

$$\int_{Q_\varepsilon^-(Y)} u = \int_{\mathbb{R}_T^{d+1}} f \Gamma_\varepsilon(\cdot, Y). \quad (3.39)$$

Similar to (3.38), for $Z \in Q_\varepsilon^-(Y)$, we have

$$\|u\|_{L_\infty(Q_{r/4}^+(Z))} \leq Cr^{-d-2}\|u\|_{L_1(Q_r(Y))}. \quad (3.40)$$

Let v be the solution of

$$-\partial_t v - \Delta v = f \text{ in } \mathbb{R}_T^{d+1}, \quad v(s+T, \cdot) = 0 \text{ on } \mathbb{R}^d.$$

By the L_p estimates (cf. (2.3)) and the parabolic Sobolev embedding, we have

$$\|v\|_{L_{q(d+2)/(d+2-q)}(\mathbb{R}_T^{d+1})} \leq C\|v\|_{W_q^{1,2}(\mathbb{R}_T^{d+1})} \leq C\|f\|_{L_q(\mathbb{R}_T^{d+1})}.$$

Note that $w = u - v$ satisfies

$$P^*w = -\text{div}^2((\mathbf{A} - \mathbf{I})v) \text{ in } \mathbb{R}_T^{d+1}, \quad w(s+T, \cdot) = 0 \text{ on } \mathbb{R}^d.$$

Therefore, by Lemma 2.2 and the last inequality, we have

$$\|w\|_{L_{q(d+2)/(d+2-q)}(\mathbb{R}_T^{d+1})} \leq C\|\mathbf{A} - \mathbf{I}\|\|v\|_{L_{q(d+2)/(d+2-q)}(\mathbb{R}_T^{d+1})} \leq C\|f\|_{L_q(\mathbb{R}_T^{d+1})},$$

which in turn implies that

$$\|u\|_{L_{q(d+2)/(d+2-q)}(\mathbb{R}_T^{d+1})} \leq \|v\|_{L_{q(d+2)/(d+2-q)}(\mathbb{R}_T^{d+1})} + \|w\|_{L_{q(d+2)/(d+2-q)}(\mathbb{R}_T^{d+1})} \leq C\|f\|_{L_q(\mathbb{R}_T^{d+1})}.$$

Then by (3.40) and Hölder's inequality, we have

$$\|u\|_{L_\infty(Q_\varepsilon^-(Y))} \leq Cr^{-d+(d+2)/p}\|u\|_{L_{q(d+2)/(d+2-2q)}(Q_r(Y))} \leq Cr^{-d+(d+2)/p}\|f\|_{L_q(\mathbb{R}_T^{d+1})}.$$

Therefore, it follows from (3.39) and the assumption that $f = 0$ in $Q_r(Y)$, that

$$\left| \int_{\mathbb{R}_T^{d+1} \setminus \overline{Q}_r(Y)} f \Gamma_\varepsilon(\cdot, Y) \right| \leq C r^{-d+(d+2)/p} \|f\|_{L_q(\mathbb{R}_T^{d+1} \setminus \overline{Q}_r(Y))}.$$

Again, we obtain (3.35) from the last inequality by duality. \blacksquare

Lemma 3.8. *For any $\varepsilon \in (0, 1)$, we have*

$$\begin{aligned} \left| \left\{ (t, x) \in \mathbb{R}_T^{d+1} : |\Gamma_\varepsilon(t, x, s, y)| > \alpha \right\} \right| &\leq C \alpha^{-\frac{d+2}{d}}, \quad \forall \alpha > 0, \\ \left| \left\{ (t, x) \in \mathbb{R}_T^{d+1} : |\partial_t \Gamma_\varepsilon(t, x, s, y)| + |D_x^2 \Gamma_\varepsilon(t, x, s, y)| > \alpha \right\} \right| &\leq C \alpha^{-1}, \quad \forall \alpha > 0, \end{aligned}$$

where $C = C(d, \lambda, \Lambda, T, \omega_A^\times)$.

Proof. These follow from (3.35) and (3.36), respectively. See the proof of [2, Lemma 3.4]. \blacksquare

With Lemmas 3.7 and 3.8 available, one can modify the argument of [2] to construct the fundamental solution $\Gamma(X, Y)$ for the operator P out of the family $\{\Gamma_\varepsilon(X, Y)\}$. We claim that for any $p \in (1, \infty)$, $r > 0$, and $T \geq 1$, we have

$$\sup_{0 < \varepsilon < 1} \|\Gamma_\varepsilon(\cdot, Y)\|_{W_p^{1,2}(\mathbb{R}_T^{d+1} \setminus \overline{Q}_r(Y))} < +\infty. \quad (3.41)$$

Indeed, by using the fact that $\Gamma_\varepsilon(s - T, \cdot) \equiv 0$, it follows from the Poincaré inequality and Lemma 3.7 that

$$\int_{s-T}^{s+T} \int_{\mathbb{R}^d \setminus B_r(y)} |\Gamma_\varepsilon(t, x, s, y)|^p dx dt \leq C \int_{s-T}^{s+T} \int_{\mathbb{R}^d \setminus B_r(y)} |\partial_t \Gamma_\varepsilon(t, x, s, y)|^p dx dt \leq C,$$

where C is a constant that depends on the parameters including p , r , and T but is independent of ε . Then, by the interpolation inequality, we have

$$\begin{aligned} \int_{s-T}^{s+T} \int_{\mathbb{R}^d \setminus B_r(y)} |D_x \Gamma_\varepsilon(t, x, s, y)|^p dx dt \\ \leq C \int_{s-T}^{s+T} \int_{\mathbb{R}^d \setminus B_r(y)} |\Gamma_\varepsilon(t, x, s, y)|^p + |D_x^2 \Gamma_\varepsilon(t, x, s, y)|^p dx dt \leq C. \end{aligned}$$

Let $\eta = \eta(x)$ be a smooth function such that

$$0 \leq \eta \leq 1, \quad \eta = 1 \text{ in } B_r(y), \quad \eta = 0 \text{ in } \mathbb{R}^d \setminus B_{2r}(y), \quad |D\eta| \leq 2/r.$$

We apply the Poincaré inequality in the space variable to $\eta D_x \Gamma(\cdot, \cdot, s, y)$ on $I \times B_{2r}(y)$ for $I = (s - T, s - r^2)$ and $I = (s + r^2, s + T)$, separately, to get

$$\begin{aligned} \int_I \int_{B_r(y)} |D_x \Gamma_\varepsilon(t, x, s, y)|^p dx dt &\leq C \int_I \int_{B_{2r}(y)} |D_x^2 \Gamma_\varepsilon(t, x, s, y)|^p dx dt \\ &\quad + C r^{-p} \int_I \int_{B_{2r}(y) \setminus B_r(y)} |D_x \Gamma_\varepsilon(t, x, s, y)|^p dx dt \leq C, \end{aligned}$$

and similarly with $\eta \Gamma(\cdot, \cdot, s, y)$ in place of $\eta D_x \Gamma(\cdot, \cdot, s, y)$, we get

$$\begin{aligned} \int_I \int_{B_r(y)} |\Gamma_\varepsilon(t, x, s, y)|^p dx dt &\leq C \int_I \int_{B_{2r}(y)} |D_x \Gamma_\varepsilon(t, x, s, y)|^p dx dt \\ &\quad + C r^{-p} \int_I \int_{B_{2r}(y) \setminus B_r(y)} |\Gamma_\varepsilon(t, x, s, y)|^p dx dt \leq C. \end{aligned}$$

Combining these together, we obtain (3.41). Therefore, by applying a diagonalization process, we see that there exists a sequence of positive numbers $\{\varepsilon_i\}_{i=1}^\infty$ with $\lim_{i \rightarrow \infty} \varepsilon_i = 0$ and a function $\Gamma(\cdot, Y)$ on $\mathbb{R}^{d+1} \setminus \{Y\}$, which belongs to $W_2^{1,2}(\mathbb{R}_T^{d+1} \setminus \overline{Q_r}(Y))$ for any $T \geq 1$ and $r > 0$, such that

$$\Gamma_{\varepsilon_i}(\cdot, Y) \rightharpoonup \Gamma(\cdot, Y) \text{ weakly in } W_2^{1,2}(\mathbb{R}_T^{d+1} \setminus \overline{Q_r}(Y)). \quad (3.42)$$

On the other hand, Lemma 3.8 implies that for $1 < p < \frac{d+2}{d}$, we have

$$\sup_{0 < \varepsilon < 1} \|\Gamma_\varepsilon(\cdot, Y)\|_{L_p(\overline{Q_r}(Y))} < +\infty,$$

which together with (3.41) implies that

$$\sup_{0 < \varepsilon < 1} \|\Gamma_\varepsilon(\cdot, Y)\|_{L_p(\mathbb{R}_T^{d+1})} < +\infty,$$

Therefore, by passing to a subsequence if necessary, we see that

$$\Gamma_{\varepsilon_i}(\cdot, Y) \rightharpoonup \Gamma(\cdot, Y) \text{ weakly in } L_p(\mathbb{R}_T^{d+1}), \quad \forall p \in (1, \frac{d+2}{d}).$$

Finally, from (3.42) and (3.33), we find that $\Gamma(\cdot, Y)$ belongs to $W_2^{1,2}(\mathbb{R}_T^{d+1} \setminus \overline{Q_r}(Y))$ and satisfies $P\Gamma(\cdot, Y) = 0$ in $\mathbb{R}_T^{d+1} \setminus \overline{Q_r}(Y)$. Since we assume that \mathbf{A} belongs to $\text{DMO}_X \subset \text{VMO}_X$, we see that for any $r > 0$, $\Gamma_\varepsilon(\cdot, Y)$ is locally uniformly continuous in $\mathbb{R}^{d+1} \setminus Q_r(Y)$ for sufficiently small ε 's, with a uniform modulus of continuity. Thus, by the Arzela-Ascoli theorem and passing to another subsequence if necessary, we see that

$$\Gamma_{\varepsilon_i}(\cdot, Y) \rightarrow \Gamma(\cdot, Y) \text{ locally uniformly on } \mathbb{R}^{d+1} \setminus Q_r(Y), \quad \forall r > 0.$$

Recall that $\Gamma^*(\cdot, X)$ satisfies

$$P^*\Gamma^*(\cdot, X) = 0 \text{ in } \mathbb{R}^{d+1} \setminus Q_r(X) \text{ for any } r > 0,$$

and thus by [3, Theorem 3.3], we see that $\Gamma^*(\cdot, X)$ is continuous in $\mathbb{R}^{d+1} \setminus \{X\}$. Therefore, we obtain the identity (1.2) by taking limit $\varepsilon \rightarrow 0$ in (3.34).

Note that we have just shown that $\Gamma(X, Y)$ is continuous in $\mathbb{R}^{d+1} \times \mathbb{R}^{d+1}$ away from the diagonal $\{(X, X) : X \in \mathbb{R}^{d+1}\}$. The property that $\Gamma(t, x, s, y) = 0$ for $t < s$ follows from the fact that $\Gamma_\varepsilon(t, x, s, y) = 0$ if $t \leq s - \varepsilon^2$. Also, it follows from [3, Theorem 3.2] that $D_x^2 \Gamma(\cdot, Y)$ is continuous in $\mathbb{R}^{d+1} \setminus \{Y\}$ and that $\partial_t \Gamma(\cdot, Y)$ is continuous in $\mathbb{R}^{d+1} \setminus \{Y\}$ if \mathbf{A} is continuous. We obtain (1.3) immediately from (3.10). \blacksquare

4. PROOF OF THEOREM 1.3

For the sake of simplicity, let us assume that $Y = 0$ and $T = 1$. Also, let us denote

$$u(t, x) = \Gamma(t, x, 0, 0).$$

In Section 4.1, we first show that $u(t, x)$ has the exponential decay

$$|u(t, x)| \leq C_0 t^{-d/2} \exp(-\kappa_0 |x| / \sqrt{t})$$

for some $\kappa_0 > 0$ and $C_0 > 1$. Then in Section 4.2, by using the semigroup property

$$\Gamma(t, x, s, y) = \int_{\mathbb{R}^d} \Gamma(t, x, \tau, \xi) \Gamma(\tau, \xi, s, y) d\xi, \quad \text{for } s < \tau < t, \quad (4.1)$$

iteratively with appropriately chosen time steps, we establish the almost Gaussian estimate (1.4).

4.1. Exponential decay of the fundamental solution. For $k = 1, 2, \dots$, let $\eta_k = \eta_k(x)$ be a smooth function in \mathbb{R}^d such that

$$\eta_k = 0 \text{ in } B_k(0), \quad \eta_k = 1 \text{ in } \mathbb{R}^d \setminus B_{k+1}(0), \quad \|D\eta\|_\infty \leq 2, \quad \|D^2\eta\|_\infty \leq 4.$$

Let $v = ue^{-\mu t}$, where $\mu \geq 1$ is a constant to be specified. Note that

$$v_k = v_k(t, x) := \eta_k(x)v(t, x)$$

satisfies

$$Pv_k + \mu v_k = f_k := -2a^{ij}D_i\eta_k D_j v - va^{ij}D_{ij}\eta_k \text{ in } (0, 1) \times \mathbb{R}^d, \quad v_k(0, \cdot) = 0 \text{ on } \mathbb{R}^d.$$

Let us denote

$$B_k = B_k(0), \quad B_k^c = \mathbb{R}^d \setminus B_k(0).$$

By the $W_p^{1,2}$ -estimates (see, for instance, [13]), we have

$$\begin{aligned} \mu \|v_k\|_{L_p((0,1) \times \mathbb{R}^d)} + \sqrt{\mu} \|Dv_k\|_{L_p((0,1) \times \mathbb{R}^d)} + \|D^2v_k\|_{L_p((0,1) \times \mathbb{R}^d)} \\ \leq N_0 \|f_k\|_{L_p((0,1) \times \mathbb{R}^d)} \leq N_0 \left(\|Dv\|_{L_p((0,1) \times (B_{k+1} \setminus B_k))} + \|v\|_{L_p((0,1) \times (B_{k+1} \setminus B_k))} \right), \end{aligned}$$

where $N_0 = N_0(d, \lambda, \Lambda, p, \omega_A^x)$ is independent of μ . On the other hand, note that

$$\|v\|_{L_p((0,1) \times B_{k+1}^c)} + \|Dv\|_{L_p((0,1) \times B_{k+1}^c)} \leq \|v_k\|_{L_p((0,1) \times \mathbb{R}^d)} + \|Dv_k\|_{L_p((0,1) \times \mathbb{R}^d)}.$$

Combining the last two inequalities, we have

$$\begin{aligned} \|v\|_{L_p((0,1) \times B_{k+1}^c)} + \|Dv\|_{L_p((0,1) \times B_{k+1}^c)} &\leq N_0 \mu^{-\frac{1}{2}} \left(\|v\|_{L_p((0,1) \times (B_{k+1} \setminus B_k))} + \|Dv\|_{L_p((0,1) \times (B_{k+1} \setminus B_k))} \right) \\ &\leq N_0 \mu^{-\frac{1}{2}} \left(\|v\|_{L_p((0,1) \times B_k^c)} + \|Dv\|_{L_p((0,1) \times B_k^c)} \right). \end{aligned} \quad (4.2)$$

Taking μ so large that $N_0 \mu^{-1/2} \leq 1/2$ and iterating on $k = 1, 2, 3, \dots$ in (4.2), we get

$$\begin{aligned} \|v\|_{L_p((0,1) \times B_{k+1}^c)} + \|Dv\|_{L_p((0,1) \times B_{k+1}^c)} \\ \leq 2^{-k} \left(\|v\|_{L_p((0,1) \times (B_2 \setminus B_1))} + \|Dv\|_{L_p((0,1) \times (B_2 \setminus B_1))} \right) \leq C 2^{-k} \end{aligned} \quad (4.3)$$

for $k = 1, 2, 3, \dots$, where we used the local $W_p^{1,2}$ estimate and the pointwise estimate (1.3) in the last inequality.

Then, by using (4.3), the fact that $Pu = 0$ in $(0, 1) \times \mathbb{R}^d$, and (1.3) we find that there are constants $C_0 > 1$ and $\kappa_0 > 0$ such that

$$|u(1, x)| \leq C_0 e^{-\kappa_0 |x|}, \quad \forall x \in \mathbb{R}^d. \quad (4.4)$$

We remark that in the proof of (4.4) above, we only used the bound (1.3) with $Y = 0$.

Notice that for $\varepsilon \in (0, 1]$, if we set \tilde{u} and \tilde{a}^{ij} by

$$\tilde{u}(t, x) = \varepsilon^d u(\varepsilon^2 t, \varepsilon x), \quad \tilde{a}^{ij}(t, x) = a^{ij}(\varepsilon^2 t, \varepsilon x),$$

and define the operator \tilde{P} by

$$\tilde{P}\tilde{u} := \partial_t \tilde{u} - \tilde{a}^{ij} D_{ij} \tilde{u},$$

then it is easily seen that $\tilde{u}(t, x)$ satisfies $\tilde{P}\tilde{u} = 0$ in $(0, 1) \times \mathbb{R}^d$ and that \tilde{u} satisfies the bound (1.3) with $Y = 0$, i.e.,

$$|\tilde{u}(t, x)| \leq C \max(\sqrt{t}, |x|)^{-d}.$$

Since $0 < \varepsilon \leq 1$, we can keep the same the constants C_0 and κ_0 in (4.4) for \tilde{u} and obtain

$$|\Gamma(\varepsilon^2, x, 0, 0)| = \varepsilon^{-d} |\varepsilon^d u(\varepsilon^2, \varepsilon x / \varepsilon)| = \varepsilon^{-d} |\tilde{u}(1, x / \varepsilon)| \leq C_0 \varepsilon^{-d} e^{-\kappa_0 |x| / \varepsilon}. \quad (4.5)$$

Also, since translation does not alter the constants κ_0 and C_0 in the estimate (4.5), for any $x, y \in \mathbb{R}^d$ and $s \in \mathbb{R}$, we have

$$|\Gamma(s + \varepsilon^2, x, s, y)| \leq C_0 \varepsilon^{-d} e^{-\kappa_0 \frac{|x-y|}{\varepsilon}}, \quad \forall \varepsilon \in (0, 1]. \quad (4.6)$$

4.2. Almost Gaussian estimate. For $(t, x) \in (0, 1] \times \mathbb{R}^d$, let $N = N(t, x) > 1$ be an integer to be chosen later. We partition the interval $(0, 1)$ into N^2 subintervals of equal length t/N^2 . Let us denote

$$t_j = j(t/N^2), \quad j = 1, 2, \dots, N^2.$$

By using (4.6) and (4.1), we have

$$\Gamma(t_{j+1}, x_{j+1}, 0, 0) = \int_{\mathbb{R}^d} \Gamma(t_{j+1}, x_{j+1}, t_j, x_j) \Gamma(t_j, x_j, 0, 0) dx_j.$$

Inductively, we have

$$\Gamma(t_{N^2}, x_{N^2}, 0, 0) = \int_{\mathbb{R}^d} \cdots \int_{\mathbb{R}^d} \prod_{j=1}^{N^2-1} \Gamma(t_{j+1}, x_{j+1}, t_j, x_j) \Gamma(t_1, x_1, 0, 0) dx_1 \cdots dx_{N^2-1}.$$

Therefore, by using (4.6) with $\varepsilon = \sqrt{t}/N$, we have

$$\begin{aligned} |\Gamma(t_{N^2}, x_{N^2}, 0, 0)| &\leq \left(\frac{C_0 N^d}{t^{d/2}} \right)^{N^2} \int_{(\mathbb{R}^d)^{N^2-1}} \left(\prod_{j=1}^{N^2-1} e^{-\kappa_0 \frac{N|x_{j+1}-x_j|}{\sqrt{t}}} \right) e^{-\kappa_0 \frac{N|x_1|}{\sqrt{t}}} dx_1 \cdots dx_{N^2-1} \\ &\leq C_0^{N^2} \left(\frac{N^d}{t^{d/2}} \right) \int_{(\mathbb{R}^d)^{N^2-1}} e^{-\kappa_0 \sum_{j=1}^{N^2-1} |y_j| - \kappa_0 \left| \frac{N}{\sqrt{t}} x_{N^2} - \sum_{j=1}^{N^2-1} y_j \right|} dy_1 \cdots dy_{N^2-1}, \end{aligned} \quad (4.7)$$

where we used the change of variables

$$y_1 = \frac{N}{\sqrt{t}} x_1; \quad y_j = \frac{N}{\sqrt{t}} (x_j - x_{j-1}), \quad j = 2, \dots, N^2 - 1.$$

By the triangle inequality, for any $(y_1, \dots, y_{N^2-1}) \in (\mathbb{R}^d)^{N^2-1}$, we have

$$\sum_{j=1}^{N^2-1} |y_j| + \left| \frac{N}{\sqrt{t}} x - \sum_{j=1}^{N^2-1} y_j \right| \geq \sum_{j=1}^{N^2-1} |y_j| + \left| \frac{N}{\sqrt{t}} x \right| - \left| \sum_{j=1}^{N^2-1} y_j \right| \geq \frac{N}{\sqrt{t}} |x|. \quad (4.8)$$

For $n = 0, 1, 2, \dots$, let us denote

$$\Omega_n = \left\{ (y_1, \dots, y_{N^2-1}) \in (\mathbb{R}^d)^{N^2-1} : n \frac{N}{\sqrt{t}} |x| \leq \sum_{j=1}^{N^2-1} |y_j| < (n+1) \frac{N}{\sqrt{t}} |x|, \quad \sum_{j=1}^{N^2-1} y_j = \frac{N}{\sqrt{t}} x \right\}.$$

If $(y_1, \dots, y_{N^2-1}) \in \Omega_n$, then we have

$$\sum_{j=1}^{N^2-1} |y_j| + \left| \frac{N}{\sqrt{t}} x - \sum_{j=1}^{N^2-1} y_j \right| = \sum_{j=1}^{N^2-1} |y_j| \geq n \frac{N}{\sqrt{t}} |x|. \quad (4.9)$$

Notice that $d(N^2 - 1)$ -dimensional Lebesgue measure $|\Omega_n|$ is bounded by

$$|\Omega_n| \leq \left(2(n+1) \frac{N}{\sqrt{t}} |x| \right)^{d(N^2-1)} \quad (4.10)$$

and $\Omega_0 = \emptyset$.

By taking $x_{N^2} = x$ and decomposing the last integral in (4.7) into the sums of integrals over Ω_n , we obtain from (4.9), (4.8), and (4.10) that

$$\begin{aligned} |\Gamma(t, x, 0, 0)| &\leq C_0^{N^2} \left(\frac{N^d}{t^{d/2}} \right) \left(e^{-\kappa_0 \frac{N|x|}{\sqrt{t}}} |\Omega_1| + \sum_{n=2}^{\infty} e^{-\kappa_0 n \frac{N|x|}{\sqrt{t}}} |\Omega_n| \right) \\ &\leq C_0^{N^2} \left(\frac{N^d}{t^{d/2}} \right) \left(\frac{2N|x|}{\sqrt{t}} \right)^{d(N^2-1)} e^{-\kappa_0 \frac{N|x|}{\sqrt{t}}} \left(2^{d(N^2-1)} + \sum_{n=2}^{\infty} (n+1)^{d(N^2-1)} e^{-\kappa_0(n-1) \frac{N|x|}{\sqrt{t}}} \right). \end{aligned} \quad (4.11)$$

By the integral comparison, the binomial formula, and Stirling's formula, we have

$$\begin{aligned} \sum_{n=2}^{\infty} (n+1)^k e^{-\alpha(n-1)} &\leq \sum_{n=2}^{\infty} \int_{n-2}^{n-1} (s+3)^k e^{-\alpha s} ds = \int_0^{\infty} (s+3)^k e^{-\alpha s} ds \\ &= \int_0^{\infty} \sum_{m=0}^k \binom{k}{m} s^m 3^{k-m} e^{-\alpha s} ds = \sum_{m=0}^k \binom{k}{m} 3^{k-m} \alpha^{-m-1} \int_0^{\infty} s^m e^{-s} ds \\ &= \sum_{m=0}^k \binom{k}{m} 3^{k-m} \alpha^{-m-1} m! \leq \frac{3^k}{\alpha} k! \sum_{m=0}^k \binom{k}{m} \left(\frac{1}{3\alpha} \right)^m \\ &= \frac{k!}{\alpha} \left(3 + \frac{1}{\alpha} \right)^k \leq c_0 \frac{\sqrt{k}}{\alpha} \left(\frac{k}{e} \right)^k \left(3 + \frac{1}{\alpha} \right)^k, \end{aligned} \quad (4.12)$$

where c_0 is an absolute constant. By combining (4.11) and (4.12), we have

$$\begin{aligned} |\Gamma(t, x, 0, 0)| &\leq \frac{1}{t^{d/2}} e^{-\kappa_0 \frac{N|x|}{\sqrt{t}}} C_0^{N^2} N^d \left(\frac{4N|x|}{\sqrt{t}} \right)^{d(N^2-1)} \\ &\quad + c_0 \frac{1}{t^{d/2}} e^{-\kappa_0 \frac{N|x|}{\sqrt{t}}} \sqrt{d(N^2-1)} \left(\frac{\sqrt{t}}{\kappa_0 N|x|} \right) C_0^{N^2} N^d \left\{ \frac{2d(N^2-1)}{e} \left(\frac{3N|x|}{\sqrt{t}} + \frac{1}{\kappa_0} \right) \right\}^{d(N^2-1)} \\ &\leq \frac{1}{t^{d/2}} e^{-\kappa_0 \frac{N|x|}{\sqrt{t}}} C_0^{N^2} N^d \left(\frac{4N|x|}{\sqrt{t}} \right)^{d(N^2-1)} \\ &\quad + \frac{c_0 \sqrt{d}}{t^{d/2}} e^{-\kappa_0 \frac{N|x|}{\sqrt{t}}} \left(\frac{\sqrt{t}}{\kappa_0 |x|} \right) C_0^{N^2} N^d \left\{ \frac{2d(N^2-1)}{e} \left(\frac{3N|x|}{\sqrt{t}} + \frac{1}{\kappa_0} \right) \right\}^{d(N^2-1)}. \end{aligned} \quad (4.13)$$

Let us write $\xi = x/\sqrt{t}$ and take $N = \lceil |\xi|^{1-\delta} \rceil$, where $\delta \in (0, 1)$ is fixed but arbitrary. Note that

$$|\xi|^{1-\delta} \leq N < |\xi|^{1-\delta} + 1.$$

Let us consider

$$\begin{aligned} A &= -\kappa_0 |\xi|^{2-\delta} + (\log C_0) (|\xi|^{1-\delta} + 1)^2 + d \log (|\xi|^{1-\delta} + 1) \\ &\quad + d \{ (|\xi|^{1-\delta} + 1)^2 - 1 \} \log (4 (|\xi|^{1-\delta} + 1) |\xi|), \\ B &= -\kappa_0 |\xi|^{2-\delta} - \log (\kappa_0 |\xi|) + (\log C_0) (|\xi|^{1-\delta} + 1)^2 + d \log (|\xi|^{1-\delta} + 1) \\ &\quad + d \{ (|\xi|^{1-\delta} + 1)^2 - 1 \} \log (2d (|\xi|^{1-\delta} + 1)^2 - 1) (3|\xi| (|\xi|^{1-\delta} + 1) + \kappa_0^{-1}) e^{-1}. \end{aligned}$$

Note that there exist $R_0 = R_0(\delta, C_0, d, \kappa_0) \geq 1$ such that if $|\xi| > R_0$, then

$$A \leq -\beta |\xi|^{2-\delta}, \quad B \leq -\beta |\xi|^{2-\delta},$$

where $\beta = \kappa_0/2$. Then, it follows from (4.13) that for any $(t, x) \in (0, 1] \times \mathbb{R}^d$ with $|x|/\sqrt{t} > R_0$, we have

$$|\Gamma(t, x, 0, 0)| \leq C_1 t^{-d/2} \exp\left(-\beta(|x|/\sqrt{t})^{2-\delta}\right), \quad \text{where } C_1 = C_1(d).$$

On the other hand, in the case when $(t, x) \in (0, 1) \times \mathbb{R}^d$ satisfies $|x|/\sqrt{t} \leq R_0$, then we can use (4.5) to bound $\Gamma(t, x, 0, 0)$.

In conclusion, we have the following: For any $\delta \in (0, 1)$, there exists $C = C(d, \lambda, \Lambda, \omega_A^x, \delta)$ such that

$$|\Gamma(t, x, 0, 0)| \leq C t^{-d/2} \exp\left(-\beta(|x|/\sqrt{t})^{2-\delta}\right) \quad \text{on } (0, 1] \times \mathbb{R}^d.$$

Finally, by translation and the semigroup property (4.1), we get (1.4). \blacksquare

5. PROOF OF THEOREM 1.4

Let $\Gamma(t, x, \tau, \xi)$ be the fundamental solution of the operator P constructed in Section 3. Let $y \in \mathbb{R}^n$ be fixed and let \bar{P}^y be given by

$$\bar{P}^y u = \partial_t u - a^{ij}(t, y) D_{ij} u.$$

Let $\Phi^y(t, x, \tau, \xi)$ be the fundamental solution of the operator \bar{P}^y . Notice that the coefficients of \bar{P}^y depend only on t and thus one can compute $\Phi^y(t, x, \tau, \xi)$ by using the Fourier transform. However, we do not need its explicit form and will just make use of the following fact. For $t > \tau$ we have

$$\begin{aligned} |\Phi^y(t, x, \tau, \xi)| &\leq \frac{C_0}{(t - \tau)^{d/2}} e^{-\kappa_0 \frac{|x - \xi|^2}{t - \tau}}, \\ |D_x^2 \Phi^y(t, x, \tau, \xi)| &\leq \frac{C'_0}{(t - \tau)^{d/2}} \left(\frac{1}{t - \tau} + \frac{|x - \xi|^2}{(t - \tau)^2} \right) e^{-\kappa_0 \frac{|x - \xi|^2}{t - \tau}}, \end{aligned} \quad (5.1)$$

where $C_0 = C_0(d, \lambda, \Lambda)$, $C'_0 = C'_0(d, \lambda, \Lambda)$, and $\kappa_0 = \kappa_0(\lambda, \Lambda)$ are positive constants.

5.1. Modified parametrix method. Notice that we have

$$\begin{aligned} P\Gamma(t, x, \tau, \xi) - P\Phi^y(t, x, \tau, \xi) &= P\Gamma(t, x, \tau, \xi) - P\Phi^y(t, x, \tau, \xi) + \bar{P}^y\Phi^y(t, x, \tau, \xi) - \bar{P}^y\Phi^y(t, x, \tau, \xi) \\ &= -(P - \bar{P}^y)\Phi^y(t, x, \tau, \xi) = (a_{ij}(t, x) - a_{ij}(t, y))D_{ij}\Phi^y(t, x, \tau, \xi). \end{aligned}$$

In particular, by taking $y = \xi$ and setting

$$v(t, x, \tau, \xi) := \Gamma(t, x, \tau, \xi) - \Phi^\xi(t, x, \tau, \xi), \quad (5.2)$$

we have

$$Pv(t, x, \tau, \xi) = (a_{ij}(t, x) - a_{ij}(t, \xi))D_{ij}\Phi^\xi(t, x, \tau, \xi).$$

We shall shortly show that the following representation formula is available:

$$v(t, x, \tau, \xi) = \int_\tau^t \int_{\mathbb{R}^d} \Gamma(t, x, s, y) (a_{ij}(s, y) - a_{ij}(s, \xi)) D_{ij}\Phi^\xi(s, y, \tau, \xi) dy ds. \quad (5.3)$$

It then follows from (5.3) that v satisfies the relation

$$\begin{aligned} v(t, x, \tau, \xi) &= \int_\tau^t \int_{\mathbb{R}^d} \Phi^y(t, x, s, y) (a_{ij}(s, y) - a_{ij}(s, \xi)) D_{ij}\Phi^\xi(s, y, \tau, \xi) dy ds \\ &\quad + \int_\tau^t \int_{\mathbb{R}^d} v(t, x, s, y) (a_{ij}(s, y) - a_{ij}(s, \xi)) D_{ij}\Phi^\xi(s, y, \tau, \xi) dy ds. \end{aligned} \quad (5.4)$$

We note that both integrals in (5.4) are absolutely convergent. See (5.17). The last formula is reminiscent of the classical parametrix method for constructing the fundamental solutions. First, we set

$$w_0(t, x, \tau, \xi) = \int_{\tau}^t \int_{\mathbb{R}^d} \Phi^y(t, x, s, y) (a_{ij}(s, y) - a_{ij}(s, \xi)) D_{ij} \Phi^{\xi}(s, y, \tau, \xi) dy ds \quad (5.5)$$

and inductively define for $k = 0, 1, 2, \dots$,

$$w_{k+1}(t, x, \tau, \xi) = \int_{\tau}^t \int_{\mathbb{R}^d} w_k(t, x, s, y) (a_{ij}(s, y) - a_{ij}(s, \xi)) D_{ij} \Phi^{\xi}(s, y, \tau, \xi) dy ds. \quad (5.6)$$

Suppose that

$$w(t, x, \tau, \xi) := \sum_{k=0}^{\infty} w_k(t, x, \tau, \xi) \quad (5.7)$$

converges uniformly. Then by summing over $k = 0, 1, 2, \dots$ in (5.6), we find

$$w(t, x, \tau, \xi) = w_0(t, x, \tau, \xi) + \int_{\tau}^t \int_{\mathbb{R}^d} w(t, x, s, y) (a_{ij}(s, y) - a_{ij}(s, \xi)) D_{ij} \Phi^{\xi}(s, y, \tau, \xi) dy ds.$$

Since we also have (5.4), it is plausible that

$$v(t, x, \tau, \xi) = w(t, x, \tau, \xi). \quad (5.8)$$

We shall verify (5.8) after we establish the Gaussian estimate for w .

5.2. Gaussian estimate for w . Recall that we assume \mathbf{A} is uniformly Dini continuous in x , that is,

$$\varrho_{\mathbf{A}}^x(r) := \sup \left\{ |\mathbf{A}(t, x) - \mathbf{A}(t, y)| : x, y \in \mathbb{R}^d, t \in \mathbb{R}, |x - y| \leq r \right\}$$

satisfies the Dini condition

$$\int_0^1 \frac{\varrho_{\mathbf{A}}^x(r)}{r} dr < +\infty.$$

It follows from (5.5) and (5.1) that

$$\begin{aligned} |w_0(t, x, \tau, \xi)| &\leq \int_{\tau}^t \int_{\mathbb{R}^d} |\Phi^y(t, x, s, y)| |\mathbf{A}(s, y) - \mathbf{A}(s, \xi)| |D^2 \Phi^{\xi}(s, y, \tau, \xi)| dy ds \\ &\leq \int_{\tau}^t \int_{\mathbb{R}^d} \frac{C_0 C'_0 e^{-\kappa_0 \frac{|x-y|^2}{t-s}}}{(t-s)^{d/2} (s-\tau)^{d/2}} \varrho_{\mathbf{A}}^x(|y - \xi|) \left(\frac{1}{s-\tau} + \frac{|y - \xi|^2}{(s-\tau)^2} \right) e^{-\kappa_0 \frac{|y-\xi|^2}{s-\tau}} dy ds. \end{aligned} \quad (5.9)$$

Since $\varrho_{\mathbf{A}}^x$ is increasing and by the triangle inequality, we have

$$\varrho_{\mathbf{A}}^x(r_1 + r_2) \leq \varrho_{\mathbf{A}}^x(r_1) + \varrho_{\mathbf{A}}^x(r_2), \quad \forall r_1, r_2 \geq 0,$$

it follows that

$$\frac{\varrho_{\mathbf{A}}^x(|y - \xi|)}{|y - \xi|} \leq 2 \frac{\varrho_{\mathbf{A}}^x(\sqrt{s-\tau})}{\sqrt{s-\tau}} \quad \text{for } |y - \xi| \geq \sqrt{s-\tau}.$$

Therefore, in the case when $|y - \xi| \geq \sqrt{s-\tau}$, we have

$$\varrho_{\mathbf{A}}^x(|y - \xi|) \left(\frac{1}{s-\tau} + \frac{|y - \xi|^2}{(s-\tau)^2} \right) \leq 2 \frac{\varrho_{\mathbf{A}}^x(\sqrt{s-\tau})}{s-\tau} \left(\frac{|y - \xi|^2}{s-\tau} \right)^{\frac{1}{2}} \left(1 + \frac{|y - \xi|^2}{s-\tau} \right).$$

On the other hand, if $|y - \xi| < \sqrt{s - \tau}$, then we have

$$\varrho_{\mathbf{A}}^x(|y - \xi|) \left(\frac{1}{s - \tau} + \frac{|y - \xi|^2}{(s - \tau)^2} \right) \leq \frac{\varrho_{\mathbf{A}}^x(\sqrt{s - \tau})}{s - \tau} \left(1 + \frac{|y - \xi|^2}{s - \tau} \right).$$

In both cases, notice that for any $\kappa'_0 \in (0, \kappa_0)$, there is a constant $C_1 = C_1(\kappa_0, \kappa'_0) > 0$ such that we have

$$\varrho_{\mathbf{A}}^x(|y - \xi|) \left(\frac{1}{s - \tau} + \frac{|y - \xi|^2}{(s - \tau)^2} \right) e^{-\kappa'_0 \frac{|y - \xi|^2}{s - \tau}} \leq C_1 \frac{\varrho_{\mathbf{A}}^x(\sqrt{s - \tau})}{s - \tau} e^{-\kappa'_0 \frac{|y - \xi|^2}{s - \tau}}. \quad (5.10)$$

We recall the following identity, which is a simple consequence of the Fourier transform: For $\tau < s < t$, we have

$$\int_{\mathbb{R}^d} \frac{1}{(t - s)^{d/2}} e^{-\kappa'_0 \frac{|x - y|^2}{t - s}} \frac{1}{(s - \tau)^{d/2}} e^{-\kappa'_0 \frac{|y - \xi|^2}{s - \tau}} dy = C_2 \frac{1}{(t - \tau)^{d/2}} e^{-\kappa'_0 \frac{|x - \xi|^2}{t - \tau}}, \quad (5.11)$$

where

$$C_2 = \int_{\mathbb{R}^d} e^{-\kappa'_0 |y|^2} dy = (\pi/\kappa'_0)^{d/2}.$$

Therefore, by plugging in (5.10) into (5.9) and using the identity (5.11), we get

$$\begin{aligned} |w_0(t, x, \tau, \xi)| &\leq C_0 C'_0 C_1 \int_{\tau}^t \frac{\varrho_{\mathbf{A}}^x(\sqrt{s - \tau})}{s - \tau} \left(\int_{\mathbb{R}^d} \frac{1}{(t - s)^{d/2}} e^{-\kappa'_0 \frac{|x - y|^2}{t - s}} \frac{1}{(s - \tau)^{d/2}} e^{-\kappa'_0 \frac{|y - \xi|^2}{s - \tau}} dy \right) ds \\ &\leq \left(C_0 C'_0 C_1 C_2 \int_{\tau}^t \frac{\varrho_{\mathbf{A}}^x(\sqrt{s - \tau})}{s - \tau} ds \right) \frac{1}{(t - \tau)^{d/2}} e^{-\kappa'_0 \frac{|x - \xi|^2}{t - \tau}}. \end{aligned} \quad (5.12)$$

Note that

$$\int_{\tau}^t \frac{\varrho_{\mathbf{A}}^x(\sqrt{s - \tau})}{s - \tau} ds = 2 \int_0^{\sqrt{t - \tau}} \frac{\varrho_{\mathbf{A}}^x(s)}{s} ds.$$

Let $\varepsilon_0 \in (0, 1)$ be to fixed later. Take $\delta_0 > 0$ such that

$$2C'_0 C_1 C_2 \int_0^{\delta_0} \frac{\varrho_{\mathbf{A}}^x(s)}{s} ds \leq \varepsilon_0. \quad (5.13)$$

Then we find from (5.12) and (5.13) that

$$|w_0(t, x, \tau, \xi)| \leq \varepsilon_0 C_0 \frac{1}{(t - \tau)^{d/2}} e^{-\kappa'_0 \frac{|x - \xi|^2}{t - \tau}} \quad \text{provided } 0 < t - \tau \leq \delta_0^2. \quad (5.14)$$

Now using (5.6), (5.14), and (5.1), we get

$$\begin{aligned} |w_1(t, x, \tau, \xi)| &\leq \int_{\tau}^t \int_{\mathbb{R}^d} |w_0(t, x, s, y)| |\mathbf{A}(s, y) - \mathbf{A}(s, \xi)| |D^2 \Phi^{\xi}(s, y, \tau, \xi)| dy ds \\ &\leq \varepsilon_0 \int_{\tau}^t \int_{\mathbb{R}^d} \frac{C_0 C'_0 e^{-\kappa'_0 \frac{|x - y|^2}{t - s}}}{(t - s)^{d/2} (s - \tau)^{d/2}} \varrho_{\mathbf{A}}^x(|y - \xi|) \left(\frac{1}{s - \tau} + \frac{|y - \xi|^2}{(s - \tau)^2} \right) e^{-\kappa'_0 \frac{|y - \xi|^2}{s - \tau}} dy ds. \end{aligned}$$

By using (5.10) and repeating the same computation as in (5.12), we get

$$|w_1(t, x, \tau, \xi)| \leq \varepsilon_0^2 C_0 \frac{1}{(t - \tau)^{d/2}} e^{-\kappa'_0 \frac{|x - \xi|^2}{t - \tau}} \quad \text{provided } 0 < t - \tau \leq \delta_0^2.$$

Inductively, we have

$$|w_k(t, x, \tau, \xi)| \leq \varepsilon_0^{k+1} C_0 \frac{1}{(t - \tau)^{d/2}} e^{-\kappa'_0 \frac{|x - \xi|^2}{t - \tau}} \quad \text{provided } 0 < t - \tau \leq \delta_0^2.$$

Then by (5.7), we have for $0 < t - \tau \leq \delta_0^2$ that

$$|w(t, x, \tau, \xi)| \leq \sum_{k=0}^{\infty} \varepsilon_0^{k+1} C_0 \frac{1}{(t - \tau)^{d/2}} e^{-\kappa'_0 \frac{|x - \xi|^2}{t - \tau}} \leq \frac{\varepsilon_0 C_0}{1 - \varepsilon_0} \frac{1}{(t - \tau)^{d/2}} e^{-\kappa'_0 \frac{|x - \xi|^2}{t - \tau}}. \quad (5.15)$$

5.3. Verification of (5.3) and (5.8). We shall prove (5.3) first. Let us denote

$$f(s, y) := (a_{ij}(s, y) - a_{ij}(s, \xi)) D_{ij} \Phi^\xi(s, y, \tau, \xi).$$

Notice that in deriving (5.12), we have seen that

$$|f(s, y)| \leq C \frac{\varrho_{\mathbf{A}}^x(\sqrt{s - \tau})}{s - \tau} \frac{1}{(s - \tau)^{d/2}} e^{-\kappa'_0 \frac{|y - \xi|^2}{s - \tau}}. \quad (5.16)$$

Write $Z = (\tau, \xi)$ and let ζ be a smooth function on \mathbb{R}^{d+1} such that

$$0 \leq \zeta \leq 1, \quad \zeta = 0 \text{ in } Q_{r/2}(Z), \quad \zeta = 1 \text{ in } \mathbb{R}^{d+1} \setminus Q_r(Z), \quad |\partial_t \zeta| + |D\zeta|^2 + |D^2 \zeta| \leq Cr^{-2},$$

where, $0 < r < \frac{1}{4}(t - \tau)$. Then, $\tilde{v} = \zeta v(\cdot, \cdot, \tau, \xi)$ satisfies

$$P\tilde{v} = \zeta f + vP\zeta - 2a^{ij}D_i v D_j \zeta \text{ in } (\tau, t) \times \mathbb{R}^d, \quad \tilde{v}(\tau, \cdot) = 0 \text{ on } \mathbb{R}^d.$$

Notice that $\zeta f + vP\zeta - 2a^{ij}D_i v D_j \zeta \in L_p((\tau, t) \times \mathbb{R}^d)$ with $p > (d + 2)/2$. Therefore, by Proposition 3.4 and the symmetry relation (1.2), we have

$$\begin{aligned} v(t, x, \tau, \xi) &= \tilde{v}(t, x) = I + II := \int_{\tau}^t \int_{\mathbb{R}^d} \Gamma(t, x, s, y) \zeta(s, y) f(s, y) dy ds \\ &\quad + \int_{Q_r(Z) \setminus Q_{r/2}(Z)} \Gamma(t, x, s, y) \{P\zeta(s, y) v(s, y, \tau, \xi) - 2a^{ij}D_i \zeta(s, y) D_j v(s, y, \tau, \xi)\} dy ds. \end{aligned}$$

We claim that $II \rightarrow 0$ as $r \rightarrow 0$. Assume the claim for now. By (5.16) and (1.4), we see that I is absolutely convergent, that is,

$$\begin{aligned} \int_{\tau}^t \int_{\mathbb{R}^d} |\Gamma(t, x, s, y) f(s, y)| dy ds \\ = \int_{\tau}^{\frac{t+\tau}{2}} \int_{\mathbb{R}^d} |\Gamma(t, x, s, y) f(s, y)| dy ds + \int_{\frac{t+\tau}{2}}^t \int_{\mathbb{R}^d} |\Gamma(t, x, s, y) f(s, y)| dy ds < +\infty, \end{aligned} \quad (5.17)$$

and thus we obtain (5.3) by the dominated convergence theorem applied to I .

Now, we prove the claim that $II \rightarrow 0$. For $Y = (s, y) \in (\tau, t) \times \mathbb{R}^d$, let

$$\tilde{r} = \frac{1}{5}|Y - Z|.$$

We set $\delta = \alpha \tilde{r}$, where $\alpha > 1$ is to be specified. Recall that

$$v(s, y, \tau, \xi) = \Gamma(s, y, \tau, \xi) - \Phi^\xi(s, y, \tau, \xi) = \Gamma^*(\tau, \xi, s, y) - (\Phi^\xi)^*(\tau, \xi, s, y),$$

and note that

$$v^* = v^*(\cdot, \cdot) = \Gamma(s, y, \cdot, \cdot) - \Phi^\xi(s, y, \cdot, \cdot) \quad (5.18)$$

satisfies

$$P^* v^* = \operatorname{div}^2((\mathbf{A} - \tilde{\mathbf{A}}_0) \tilde{\Phi}_0) \text{ in } (\tau - 1, s) \times \mathbb{R}^d, \quad v^*(s, \cdot) = 0 \text{ on } \mathbb{R}^d,$$

where we set

$$\tilde{\mathbf{A}}_0 = \tilde{\mathbf{A}}_0(\cdot) = \mathbf{A}(\cdot, \xi) \quad \text{and} \quad \tilde{\Phi}_0 = \tilde{\Phi}_0(\cdot, \cdot) = (\Phi^\xi)^*(\cdot, \cdot, s, y).$$

Let $\tilde{\zeta}$ be a smooth function on \mathbb{R}^{d+1} such that

$$0 \leq \tilde{\zeta} \leq 1, \quad \tilde{\zeta} = 0 \text{ in } Q_{\delta/2}(Y), \quad \tilde{\zeta} = 1 \text{ in } \mathbb{R}^{d+1} \setminus Q_\delta(Y), \quad |D\tilde{\zeta}| \leq 4/\delta,$$

and define \mathbf{g}_1 and \mathbf{g}_2 by (cf. (3.11))

$$\mathbf{g}_1 = \tilde{\zeta}(\mathbf{A} - \tilde{\mathbf{A}}_0)\tilde{\Phi}_0 \quad \text{and} \quad \mathbf{g}_2 = (1 - \tilde{\zeta})(\mathbf{A} - \tilde{\mathbf{A}}_0)\tilde{\Phi}_0.$$

Noting that $\|\mathbf{A} - \tilde{\mathbf{A}}_0\|_\infty \leq C(d, \Lambda)$ and using (3.13) and properties of $\tilde{\zeta}$, we have

$$\int_{\mathbb{R}^{d+1}} |\mathbf{g}_1|^{\frac{2(d+2)}{d}} \leq \sum_{k=0}^{\infty} \int_{Q_{2^k\delta}(Y) \setminus Q_{2^{k-1}\delta}(Y)} |\mathbf{g}_1|^{\frac{2(d+2)}{d}} \leq C \sum_{k=0}^{\infty} (2^k\delta)^{-d-2} \leq C\delta^{-d-2}. \quad (5.19)$$

Note that we have

$$|\mathbf{A} - \tilde{\mathbf{A}}_0| \leq \varrho_{\mathbf{A}}^{\mathbf{x}}(\delta) + \varrho_{\mathbf{A}}^{\mathbf{x}}(5\tilde{r}) \leq (\alpha + 6)\varrho_{\mathbf{A}}^{\mathbf{x}}(\tilde{r}) \quad \text{in } Q_\delta(Y),$$

and thus we have

$$\int_{\mathbb{R}^{d+1}} |\mathbf{g}_2|^{\frac{d+1}{d}} \leq ((\alpha + 6)\varrho_{\mathbf{A}}^{\mathbf{x}}(\tilde{r}))^{\frac{d+1}{d}} \int_{Q_\delta(Y)} |\mathbf{g}_2|^{\frac{d+1}{d}} \leq C((\alpha + 6)\varrho_{\mathbf{A}}^{\mathbf{x}}(\tilde{r}))^{\frac{d+1}{d}} \delta. \quad (5.20)$$

Let v_i ($i = 1, 2$) be the solutions of the problems

$$P^*v_i = \operatorname{div}^2 \mathbf{g}_i \quad \text{in } (\tau - 1, s) \times \mathbb{R}^d, \quad v_i(s, \cdot) = 0 \quad \text{on } \mathbb{R}^d \quad (i = 1, 2).$$

We extend v_1 and v_2 to be zero on $(s, \infty) \times \mathbb{R}^d$. By Lemma 2.2 together with (5.19) and (5.20), we have

$$\|v_1\|_{L_{2(d+2)/d}((\tau-1,s) \times \mathbb{R}^d)} \leq C\delta^{-\frac{d}{2}} \quad \text{and} \quad \|v_2\|_{L_{(d+1)/d}((\tau-1,s) \times \mathbb{R}^d)} \leq C(\alpha + 6)\varrho_{\mathbf{A}}^{\mathbf{x}}(\tilde{r})\delta^{\frac{d}{d+1}}. \quad (5.21)$$

By (3.8), we see that both v_1 and v_2 also belong to $L_p((\tau - 1, s) \times \mathbb{R}^d)$ for any $p \in (1, (d + 2)/d)$. Therefore, by the uniqueness, we have

$$v^* = v_1 + v_2. \quad (5.22)$$

We now estimate $v_1(Z)$ and $v_2(Z)$. By using Lemma 2.3, we have

$$|v_i(Z)| \leq C \int_{Q_{2\tilde{r}}^+(Z)} |v_i| + C \int_0^{\tilde{r}} \frac{\omega_{\mathbf{g}_i}^{\mathbf{x}}(\tilde{t}, Q_{2\tilde{r}}^+(Z))}{\tilde{t}} d\tilde{t} \quad (i = 1, 2). \quad (5.23)$$

Using (5.21) together with Hölder's inequalities, we have

$$\begin{aligned} \int_{Q_{2\tilde{r}}^+(Z)} |v_1| &\leq C\tilde{r}^{-\frac{d}{2}} \|v_1\|_{L_{2(d+2)/d}(Q_{2\tilde{r}}^+(Z))} \leq C\tilde{r}^{-\frac{d}{2}} \delta^{-\frac{d}{2}}, \\ \int_{Q_{2\tilde{r}}^+(Z)} |v_2| &\leq C\tilde{r}^{-\frac{(d+2)d}{d+1}} \|v_2\|_{L_{(d+1)/d}(Q_{2\tilde{r}}^+(Z))} \leq C\tilde{r}^{-d} (\alpha + 6) \alpha^{\frac{d}{d+1}} \varrho_{\mathbf{A}}^{\mathbf{x}}(\tilde{r}). \end{aligned} \quad (5.24)$$

By using the bound of $\tilde{\Phi}_0$, we have

$$\omega_{\mathbf{g}_i}^{\mathbf{x}}(\tilde{t}, Q_{2\tilde{r}}^+(Z)) \leq C(d, \lambda, \Lambda) \tilde{r}^{-d} \varrho_{\mathbf{A}}^{\mathbf{x}}(\tilde{t}), \quad \forall \tilde{t} \in (0, \tilde{r}] \quad (i = 1, 2). \quad (5.25)$$

By combining (5.22), (5.23), (5.24), and (5.25), we obtain

$$|v^*(Z)| \leq C\tilde{r}^{-d} \left(\alpha^{-\frac{d}{2}} + (\alpha + 6) \alpha^{\frac{d}{d+1}} \varrho_{\mathbf{A}}^{\mathbf{x}}(\tilde{r}) + \int_0^{\tilde{r}} \frac{\varrho_{\mathbf{A}}^{\mathbf{x}}(\tilde{t})}{\tilde{t}} d\tilde{t} \right),$$

where C is a constant independent of \tilde{r} . Recall that $|Y - Z| = 5\tilde{r}$. Now for any $\varepsilon \in (0, 1)$, we can take $\alpha > 1$ sufficiently large and then \tilde{r} sufficiently small such that

$$|v^*(Z)| \leq \varepsilon |Y - Z|^{-d}. \quad (5.26)$$

Therefore, we conclude from (5.26) and (5.18) that for all small $r > 0$,

$$r^d |v(s, y, \tau, \xi)| = o(r), \quad \forall (s, y) \in Q_{2r}(Z) \setminus Q_{r/4}(Z),$$

where we use $o(r)$ to denote some bounded quantity that tends to 0 as $r \rightarrow 0$.

To estimate Dv , we use the equation $Pv = f$. Notice that

$$|f(s, y)| \leq C\varrho_A^X(|y - \xi|)|Z - Y|^{-d-2} \leq C\varrho_A^X(r)r^{-d-2} \quad \text{in } Q_{2r}(Z) \setminus Q_{r/4}(Z).$$

Therefore, by using (5.26), the local $W_p^{1,2}$ estimate

$$\begin{aligned} & \left\| r^2 |\partial_s v(\cdot, \cdot, \tau, \xi)| + r^2 |D_y^2 v(\cdot, \cdot, \tau, \xi)| + r |D_y v(\cdot, \cdot, \tau, \xi)| \right\|_{L_p(Q_r(Z) \setminus Q_{r/2}(Z))} \\ & \leq C \|v(\cdot, \cdot, \tau, \xi)\|_{L_p(Q_{2r}(Z) \setminus Q_{r/4}(Z))} + Cr^2 \|f\|_{L_p(Q_{2r}(Z) \setminus Q_{r/4}(Z))}, \quad p > d + 2, \end{aligned}$$

and the Sobolev embedding, we have

$$r^{d+1} |Dv(s, y, \tau, \xi)| = o(r), \quad \forall (s, y) \in Q_r(Z) \setminus Q_{r/2}(Z). \quad (5.27)$$

Therefore, by using (5.26), (5.27), and the properties of ζ , we get $II \rightarrow 0$ as $r \rightarrow 0$, which completes the proof of (5.3).

To show (5.8), we invoke the contraction mapping theorem. For $(t, x) \in \mathbb{R}^{d+1}$, let $\mathcal{B} = L_1((t - \delta_0^2, t) \times \mathbb{R}^d)$, where δ_0 is as in (5.13). We shall show that the mapping $T : \mathcal{B} \rightarrow \mathcal{B}$ defined by

$$Tu(\tau, \xi) = w_0(t, x, \tau, \xi) + \int_{\tau}^t \int_{\mathbb{R}^d} u(s, y) (a_{ij}(s, y) - a_{ij}(s, \xi)) D_{ij} \Phi^{\xi}(s, y, \tau, \xi) dy ds$$

is a contraction. Indeed, by (5.14), (5.10), Fubini's theorem, and (5.13), we find that

$$\begin{aligned} & \int_{t-\delta_0^2}^t \int_{\mathbb{R}^d} |Tu(\tau, \xi)| d\xi d\tau \leq \int_{t-\delta_0^2}^t \int_{\mathbb{R}^d} |w_0(t, x, \tau, \xi)| d\xi d\tau \\ & + \int_{t-\delta_0^2}^t \int_{\mathbb{R}^d} \int_{\tau}^t \int_{\mathbb{R}^d} |u(s, y)| C_0' \frac{\varrho_A^X(|y - \xi|)}{(s - \tau)^{d/2}} \left(\frac{1}{s - \tau} + \frac{|y - \xi|^2}{(s - \tau)^2} \right) e^{-\kappa_0 \frac{|y - \xi|^2}{s - \tau}} dy ds d\xi d\tau \\ & \leq \int_{t-\delta_0^2}^t \int_{\mathbb{R}^d} \varepsilon_0 C_0 \frac{1}{(t - \tau)^{d/2}} e^{-\kappa_0' \frac{|x - \xi|^2}{t - \tau}} d\xi d\tau \\ & + \int_{t-\delta_0^2}^t \int_{\mathbb{R}^d} |u(s, y)| \int_{t-\delta_0^2}^s \frac{\varrho_A^X(\sqrt{s - \tau})}{s - \tau} \int_{\mathbb{R}^d} \frac{C_0' C_1}{(s - \tau)^{d/2}} e^{-\kappa_0' \frac{|y - \xi|^2}{s - \tau}} d\xi d\tau dy ds \\ & \leq \varepsilon_0 C_0 C_2 \delta_0^2 + \varepsilon_0 \int_{t-\delta_0}^t \int_{\mathbb{R}^d} |u(s, y)| dy ds. \end{aligned}$$

Therefore, we have $Tu \in \mathcal{B}$ for all $u \in \mathcal{B}$. By a similar calculation, we also find that

$$\|Tu_1 - Tu_2\|_{\mathcal{B}} \leq \varepsilon_0 \|u_1 - u_2\|_{\mathcal{B}},$$

which implies T is a contraction mapping on \mathcal{B} since we assume $\varepsilon_0 \in (0, 1)$. We now fix $\varepsilon_0 = 1/2$. Note that it follows from (1.4) and (5.15), respectively, that $v \in \mathcal{B}$ and $w \in \mathcal{B}$, which establishes the equality (5.8).

5.4. Conclusion. Therefore, by (5.2) (5.8), (5.15), and (5.1), we find that

$$|\Gamma(t, x, s, y)| \leq \frac{C}{(t - s)^{d/2}} e^{-\kappa_0' \frac{|x - y|^2}{t - s}} \quad \text{provided } 0 < t - s \leq \delta_0^2. \quad (5.28)$$

We can take $\kappa_0' = \kappa_0/2$ in the above and call it κ . It is clear that κ then depends only on λ and Λ . By using (4.1) and (5.28), we establish the Gaussian bound (1.5). See e.g., [2, Section 5.5] for the details. \blacksquare

REFERENCES

- [1] Aronson, D. G. *Bounds for the fundamental solution of a parabolic equation*. Bull. Amer. Math. Soc. **73** (1967), 890–896.
- [2] Cho, Sungwon; Dong, Hongjie; Kim, Seick. *On the Green's matrices of strongly parabolic systems of second order*. Indiana Univ. Math. J. **57** (2008), no. 4, 1633–1677.
- [3] Dong, Hongjie; Escauriaza, Luis; Kim, Seick. *On $C^{1/2,1}$, $C^{1,2}$, and $C^{0,0}$ estimates for linear parabolic operators*. J. Evol. Equ. **21** (2021), no. 4, 4641–4702.
- [4] Eidel'man, S. D. *Parabolic systems*. Translated from the Russian by Scripta Technica, London North-Holland Publishing Co., Amsterdam-London; Wolters-Noordhoff Publishing, Groningen 1969.
- [5] Escauriaza, Luis. *Bounds for the fundamental solution of elliptic and parabolic equations in nondivergence form*. Comm. Partial Differential Equations **25** (2000), no. 5-6, 821–845.
- [6] Escauriaza, Luis; Montaner, Santiago. *Some remarks on the L^p regularity of second derivatives of solutions to non-divergence elliptic equations and the Dini condition*. Rend. Lincei Mat. Appl. **28** (2017) 49–63.
- [7] Fabes, Eugene B.; Kenig, Carlos E. *Examples of singular parabolic measures and singular transition probability densities*. Duke Math. J. **48** (1981), no. 4, 845–856.
- [8] Friedman, Avner. *Partial differential equations of parabolic type*. Prentice-Hall, Inc., Englewood Cliffs, N.J. 1964.
- [9] Hwang, Sukjung; Kim, Seick. *Green's function for second order elliptic equations in non-divergence form*. Potential Anal. **52** (2020), no. 1, 27–39.
- [10] Il'in, A. M. *On the fundamental solution of a parabolic equation*. (Russian) Dokl. Akad. Nauk SSSR **147** 1962 768–771.
- [11] Il'in, A. M.; Kalašnikov, A. S.; Oleinik, O. A. *Second-order linear equations of parabolic type*. (Russian) Uspehi Mat. Nauk **17** 1962 no. 3 (105), 3–146.
- [12] Kim, Seick; Lee, Sungjin. *Estimates for Green's functions of elliptic equations in non-divergence form with continuous coefficients*. Ann. Appl. Math. **37** (2021), no. 2, 111–130.
- [13] Krylov, Nicolai V. *Parabolic and Elliptic Equations with VMO Coefficients*. Comm. Partial Differential Equations. **32** (2007) 453–475.
- [14] Krylov, N. V. *Lectures on elliptic and parabolic equations in Sobolev spaces*. Graduate Studies in Mathematics, **96**. American Mathematical Society, Providence, RI, 2008.
- [15] Li, Yanyan. *On the C^1 regularity of solutions to divergence form elliptic systems with Dini-continuous coefficients*. Chin. Ann. Math. Ser. B **38** (2017), no. 2, 489–496.
- [16] Ladyženskaja, O. A.; Solonnikov, V. A.; Ural'ceva, N. N. *Linear and quasilinear equations of parabolic type*. (Russian) Translated from the Russian by S. Smith. Translations of Mathematical Monographs, Vol. 23 American Mathematical Society, Providence, R.I. 1968.
- [17] Levi, E. E. *Sulle equazioni lineari totalmente ellittiche alle derivate parziali*. Rend. Circ. Mat. Palermo, **24** (1907), 275–317.
- [18] Porper, F. O.; Eidel'man, S. D. *Two-sided estimates of the fundamental solutions of second-order parabolic equations and some applications of them*. (Russian) Uspekhi Mat. Nauk **39** (1984), no. 3(237), 107–156. English translation: Russian Math. Surveys **39** (1984), no. 3, 119–179.
- [19] Safonov, M. V. *An example of diffusion process with singular distribution at some given time*. In: Abstr. Comm. Third Vilnius Conference on Probability Theory and Math. Statistics, June 22–27, Vilnius: Institut matematiki i kibernetiki AN Litovskoi SSR, (1981), 133–134 (in Russian).

(H. Dong) DIVISION OF APPLIED MATHEMATICS, BROWN UNIVERSITY, 182 GEORGE STREET, PROVIDENCE, RI 02912, UNITED STATES OF AMERICA
 Email address: Hongjie.Dong@brown.edu

(S. Kim) DEPARTMENT OF MATHEMATICS, YONSEI UNIVERSITY, 50 YONSEI-RO, SEODAEMUN-GU, SEOUL 03722, REPUBLIC OF KOREA.
 Email address: kimseick@yonsei.ac.kr

(S. Lee) DEPARTMENT OF MATHEMATICS, YONSEI UNIVERSITY, 50 YONSEI-RO, SEODAEMUN-GU, SEOUL 03722, REPUBLIC OF KOREA
 Email address: sungjinlee@yonsei.ac.kr