

# NATURAL ALMOST HERMITIAN STRUCTURES ON CONFORMALLY FOLIATED 4-DIMENSIONAL LIE GROUPS WITH MINIMAL LEAVES

EMMA ANDERSDOTTER SVENSSON AND SIGMUNDUR GUDMUNDSSON

**ABSTRACT.** Let  $(G, g)$  be a 4-dimensional Riemannian Lie group with a 2-dimensional left-invariant, conformal foliation  $\mathcal{F}$  with minimal leaves. Let  $J$  be an almost Hermitian structure on  $G$  adapted to the foliation  $\mathcal{F}$ . We classify such structures  $J$  which are almost Kähler ( $\mathcal{AK}$ ), integrable ( $\mathcal{I}$ ) or Kähler ( $\mathcal{K}$ ). Hereby we construct several new multi-dimensional examples in each class.

## 1. INTRODUCTION

The theory of *harmonic morphisms* was initiated by Jacobi, in the 3-dimensional Euclidean geometry, with his famous paper [8] from 1848. In the late 1970s this was then generalised to Riemannian geometry in [3] and [7] by Fuglede and Ishihara, independently. This has lead to a vibrant development which can be traced back in the standard reference [2], by Baird and Wood, and the regularly updated online bibliography [6], maintained by the second author. The following result of Baird and Eells gives the theory of harmonic morphisms, with values in a surface, a strong geometric flavour.

**Theorem 1.1.** [1] *Let  $\phi : (M^m, g) \rightarrow (N^2, h)$  be a horizontally conformal submersion from a Riemannian manifold to a surface. Then  $\phi$  is harmonic if and only if  $\phi$  has minimal fibres.*

In their work [5], the authors classify the 4-dimensional Riemannian Lie groups  $(G, g)$  equipped with a 2-dimensional, conformal and left-invariant foliation  $\mathcal{F}$  with minimal leaves. Such a foliation is locally given by a submersive harmonic morphism into a surface. For such a Riemannian Lie group there exist natural almost Hermitian structures  $J$  adopted to the foliation structure.

The purpose of this work is to classify these structures which are *almost Kähler* ( $\mathcal{AK}$ ), *integrable* ( $\mathcal{I}$ ) or *Kähler* ( $\mathcal{K}$ ), see [4]. Our classification result is the following.

**Theorem 1.2.** *Let  $(G, g, J)$  be a 4-dimensional almost Hermitian Lie group equipped with a 2-dimensional, conformal and left-invariant foliation  $\mathcal{F}$  with*

---

2010 *Mathematics Subject Classification.* 53C43, 58E20.

*Key words and phrases.* harmonic morphisms, foliations, almost complex structures.

*minimal leaves. If a natural Hermitian structure  $J$ , adapted to the foliation  $\mathcal{F}$ , is almost Kähler ( $\mathcal{AK}$ ), integrable ( $\mathcal{I}$ ) or Kähler ( $\mathcal{K}$ ), then the corresponding Lie algebra  $\mathfrak{g}_k^{\mathcal{AK}}$ ,  $\mathfrak{g}_k^{\mathcal{I}}$  or  $\mathfrak{g}_k^{\mathcal{K}}$  of  $G$  is one of those given below ( $k = 1, 2, \dots, 20$ ).*

## 2. TWO DIMENSIONAL CONFORMAL FOLIATIONS $\mathcal{F}$ ON $(G^4, g)$

Let  $(M, g)$  be a Riemannian manifold,  $\mathcal{V}$  be an integrable distribution on  $M$  and denote by  $\mathcal{H}$  its orthogonal complementary distribution. As customary, we shall also by  $\mathcal{V}$  and  $\mathcal{H}$  denote the orthogonal projections onto the corresponding subbundles of the tangent bundle  $TM$  of  $M$  and denote by  $\mathcal{F}$  the foliation tangent to  $\mathcal{V}$ . Then the second fundamental form for  $\mathcal{V}$  is given by

$$B^{\mathcal{V}}(Z, W) = \frac{1}{2} \mathcal{H}(\nabla_Z W + \nabla_W Z) \quad (Z, W \in \mathcal{V}),$$

while the second fundamental form for  $\mathcal{H}$  satisfies

$$B^{\mathcal{H}}(X, Y) = \frac{1}{2} \mathcal{V}(\nabla_X Y + \nabla_Y X) \quad (X, Y \in \mathcal{H}).$$

The foliation  $\mathcal{F}$  tangent to  $\mathcal{V}$  is said to be *conformal* if there exists a vector field  $V \in \mathcal{V}$  such that the second fundamental form  $B^{\mathcal{H}}$  satisfies

$$B^{\mathcal{H}} = g \otimes V$$

and  $\mathcal{F}$  is said to be *Riemannian* if  $V = 0$ . Furthermore,  $\mathcal{F}$  is said to be *minimal* if  $\text{trace } B^{\mathcal{V}} = 0$  and *totally geodesic* if  $B^{\mathcal{V}} = 0$ . This is equivalent to the leaves of  $\mathcal{F}$  being minimal or totally geodesic submanifolds of  $M$ , respectively.

It is well-known that the fibres of a horizontally conformal map (resp. Riemannian submersion) give rise to a conformal foliation (resp. Riemannian foliation). Conversely, the leaves of any conformal foliation (resp. Riemannian foliation) are locally the fibres of a horizontally conformal (resp. Riemannian) submersion, see [2].

Let  $(G, g)$  be a 4-dimensional Lie group equipped with a left-invariant Riemannian metric  $g$  and  $K$  be a 2-dimensional subgroup of  $G$ . Let  $\mathfrak{k}$  and  $\mathfrak{g}$  be the Lie algebras of  $K$  and  $G$ , respectively. Let  $\mathfrak{m}$  be the two dimensional orthogonal complement of  $\mathfrak{k}$  in  $\mathfrak{g}$  with respect to the Riemannian metric  $g$  on  $G$ . By  $\mathcal{V}$  we denote the integrable distribution generated by  $\mathfrak{k}$  and by  $\mathcal{H}$  its orthogonal distribution given by  $\mathfrak{m}$ . Further let  $\mathcal{F}$  be the foliation of  $G$  tangent to  $\mathcal{V}$ . Let the set

$$\mathcal{B} = \{X, Y, Z, W\}$$

be an orthonormal basis for the Lie algebra  $\mathfrak{g}$  of  $G$ , such that  $Z, W$  generate the subalgebra  $\mathfrak{k}$  and  $[W, Z] = \lambda W$  for some  $\lambda \in \mathbb{R}$ . The elements  $X, Y \in \mathfrak{m}$  can clearly be chosen such that  $\mathcal{H}[X, Y] = \rho X$  for some  $\rho \in \mathbb{R}$ .

If the 2-dimensional foliation  $\mathcal{F}$  is conformal with minimal leaves then, following [5], the Lie bracket relations of  $\mathfrak{g}$  take the following form

$$\begin{aligned}
[W, Z] &= \lambda W, \\
[Z, X] &= \alpha X + \beta Y + z_1 Z + w_1 W, \\
[Z, Y] &= -\beta X + \alpha Y + z_2 Z + w_2 W, \\
[W, X] &= aX + bY + z_3 Z - z_1 W, \\
[W, Y] &= -bX + aY + z_4 Z - z_2 W, \\
[Y, X] &= rX + \theta_1 Z + \theta_2 W,
\end{aligned} \tag{2.1}$$

with real structure coefficients. To ensure that these equations actually define a Lie algebra they must satisfy the Jacobi equations. These are equivalent to a system of 14 second order homogeneous polynomial equations in 14 variables, see Appendix A. This has been solved in [5] and the solutions give a complete classifications of such Lie algebras. They form 20 multi-dimensional families which can be found therein.

For the foliation  $\mathcal{F}$  we state the following easy result describing the geometry of the situation.

**Proposition 2.1.** *Let  $(G, g)$  be a 4-dimensional Lie group with Lie algebra  $\mathfrak{g}$  as above. Then*

- (i)  $\mathcal{F}$  is totally geodesic if and only if  $z_1 = z_2 = z_3 + w_1 = z_4 + w_2 = 0$ ,
- (ii)  $\mathcal{F}$  is Riemannian if and only if  $\alpha = a = 0$ , and
- (iii)  $\mathcal{H}$  is integrable if and only if  $\theta_1 = \theta_2 = 0$ .

### 3. THE ADAPTED ALMOST HERMITIAN STRUCTURES

Let  $(G, g)$  be a 4-dimensional Riemannian Lie group as above. Further, let  $J$  be a left-invariant almost Hermitian structure on the tangent bundle  $TG$  adapted to the decomposition  $TG = \mathcal{V} \oplus \mathcal{H}$  i.e. a bundle isomorphism  $J : TG \rightarrow TG$  such that

$$J^2 = -\text{id}, \quad J(\mathcal{V}) = \mathcal{V} \quad \text{and} \quad J(\mathcal{H}) = \mathcal{H}.$$

The almost Hermitian structure  $J$  is left-invariant and can therefore be fully described by its action on the orthonormal basis

$$\mathcal{B} = \{X, Y, Z, W\}$$

of the Lie algebra  $\mathfrak{g}$  of  $G$ . Here the vector fields  $X, Y \in \mathcal{H}$  and  $Z, W \in \mathcal{V}$  can be chosen such that

$$J(X) = Y \quad \text{and} \quad J(Z) = W. \tag{3.1}$$

**Lemma 3.1.** *Let  $(G, g, J)$  be a 4-dimensional almost Hermitian Lie group as above. Then the structure  $J$  is almost Kähler (AK) if and only if*

$$\theta_1 = 2a \quad \text{and} \quad \theta_2 = -2\alpha.$$

*Proof.* The Kähler form  $\omega$  satisfies  $\omega(E, F) = g(JE, F)$  for all  $E, F \in \mathfrak{g}$ . It is well known that  $J$  is almost Kähler if and only if the corresponding Kähler form  $\omega$  is closed i.e.  $d\omega = 0$ .

$$\begin{aligned}
d\omega(X, Y, Z) &= -g(J[X, Y], Z) - g(J[Y, Z], X) - g(J[Z, X], Y) \\
&= -g(J(-rX - \theta_1 Z - \theta_2 W), Z) \\
&\quad -g(J(\beta X - \alpha Y - z_2 Z - w_2 W), X) \\
&\quad -g(J(\alpha X + \beta Y + z_1 Z + w_1 W), Y) \\
&= -g(-rY - \theta_1 W + \theta_2 Z, Z) \\
&\quad -g(\beta Y + \alpha X - z_2 W + w_2 Z, X) \\
&\quad -g(\alpha Y - \beta X + z_1 W - w_1 Z, Y) \\
&= -\theta_2 - 2\alpha.
\end{aligned}$$

The statement is then easily obtained by performing the corresponding computations in the other cases  $d\omega(W, X, Y)$ ,  $d\omega(Z, W, X)$  and  $d\omega(Y, Z, W)$ .  $\square$

**Lemma 3.2.** *Let  $(G, g, J)$  be a 4-dimensional almost Hermitian Lie group as above. Then the almost complex structure  $J$  is integrable ( $\mathcal{I}$ ) if and only if*

$$2z_1 - z_4 - w_2 = 0 \quad \text{and} \quad 2z_2 + z_3 + w_1 = 0.$$

*Proof.* It is well known that  $J$  is integrable if and only if the corresponding skew-symmetric Nijenhuis tensor  $N$  vanishes i.e. if

$$N(E, F) = [E, F] + J[JE, F] + J[E, JF] - [JE, JF] = 0,$$

for all  $E, F \in \mathfrak{g}$ . It is obvious that  $N(X, Y) = N(Z, W) = 0$ . Now

$$\begin{aligned}
N(X, Z) &= [X, Z] + J[JX, Z] + J[X, JZ] - [JX, JZ] \\
&= [X, Z] + J[Y, Z] + J[X, W] - [Y, W] \\
&= -\alpha X - \beta Y - z_1 Z - w_1 W + J(\beta X - \alpha Y - z_2 Z - w_2 W) \\
&\quad - J(\alpha X + \beta Y + z_3 Z - z_1 W) - bX + aY + z_4 Z - z_2 W \\
&= -\alpha X - \beta Y - z_1 Z - w_1 W + \beta Y + \alpha X - z_2 W + w_2 Z \\
&\quad - aY + bX - z_3 W - z_1 Z - bX + aY + z_4 Z - z_2 W \\
&= -(2z_1 - z_4 - w_2)Z - (2z_2 + z_3 + w_1)W.
\end{aligned}$$

Similar calculations for  $N(X, W)$ ,  $N(Y, Z)$  and  $N(Y, W)$  easily provide the result.  $\square$

**Corollary 3.3.** *Let  $(G, g, J)$  be a 4-dimensional almost Hermitian Lie group as above. Then the structure  $J$  is Kähler ( $\mathcal{K}$ ) if and only if*

$$2z_1 - z_4 - w_2 = 0, \quad 2z_2 + z_3 + w_1 = 0, \quad \theta_1 = 2a \quad \text{and} \quad \theta_2 = -2\alpha.$$

*Proof.* The almost Hermitian structure  $J$  is Kähler if and only if it is almost Kähler and integrable. Hence the statement is an immediate consequence of Lemmas 3.1 and 3.2.  $\square$

**Remark 3.4.** The 2-dimensional Lie algebra  $\mathfrak{k}$  of the subgroup  $K$  of  $G$  has orthonormal basis  $\{Z, W\}$ , satisfying  $[W, Z] = \lambda W$ . This means that its simply connected universal covering group  $\tilde{K}$  has constant Gaussian curvature  $\kappa = -\lambda^2$ . Thus it is either a hyperbolic disk  $H_\lambda^2$  or the flat Euclidean plane  $\mathbb{R}^2$ .

The 4-dimensional Riemannian Lie groups  $(G, g)$ , with a left-invariant conformal foliation  $\mathcal{F}$  with minimal leaves, were classified in [5]. The purpose of this work is to investigate their adapted almost Hermitian structure  $J$ , given by (3.1), in each case. Here we adopt the notation introduced in [5].

#### 4. CASE (A) - $(\lambda \neq 0 \text{ AND } (\lambda - \alpha)^2 + \beta^2 \neq 0)$

**Example 4.1** ( $\mathfrak{g}_1(\lambda, r, w_1, w_2)$ ). This is a 4-dimensional family of *solvable* Lie algebras obtained by assuming that  $r \neq 0$ , see [5]. The Lie bracket relations are given by

$$\begin{aligned} [W, Z] &= \lambda W, \\ [Z, X] &= w_1 W, \\ [Z, Y] &= w_2 W, \\ [Y, X] &= rX + \frac{rw_1}{\lambda} W. \end{aligned}$$

( $\mathcal{AK}$ ) : According to Lemma 3.1, the adapted almost Hermitian structure  $J$  is *almost Kähler* if and only if  $\theta_1 = 2a$  and  $\theta_2 = -2\alpha$ . We see from the bracket relations that  $a = \theta_1 = \alpha = 0$ , hence  $\theta_2 = 0$ . But  $\lambda\theta_2 = rw_1$  which implies that  $rw_1 = 0$ . Since  $r \neq 0$  we conclude that the family is in the almost Kähler class if and only if  $w_1 = 0$ . This provides the 3-dimensional family  $\mathfrak{g}_1^{\mathcal{AK}}(\lambda, r, w_2)$ , given by

$$\begin{aligned} [W, Z] &= \lambda W, \\ [Z, Y] &= w_2 W, \\ [Y, X] &= rX. \end{aligned}$$

Here the corresponding simply connected Lie groups are semidirect products  $H_r^2 \ltimes H_\lambda^2$  of hyperbolic disks, see Remark 3.4.

( $\mathcal{I}$ ) : According to Lemma 3.2 the structure  $J$  is *integrable* if and only if

$$2z_2 + z_3 + w_1 = 0 \quad \text{and} \quad 2z_1 - z_4 - w_2 = 0.$$

Because  $z_1 = z_2 = z_3 = z_4 = 0$  we conclude that  $w_1 = w_2 = 0$ . From this we yield the 2-dimensional family  $\mathfrak{g}_1^{\mathcal{I}}(\lambda, r)$  satisfying

$$\begin{aligned} [W, Z] &= \lambda W, \\ [Y, X] &= rX. \end{aligned}$$

( $\mathcal{K}$ ) : The family  $\mathfrak{g}_1^{\mathcal{K}}(\lambda, r) = \mathfrak{g}_1^{\mathcal{I}}(\lambda, r)$  is contained in  $\mathfrak{g}_1^{\mathcal{AK}}(\lambda, r, w_2)$  and thus *Kähler*, according to Corollary 3.3. Here the corresponding simply connected Lie groups are direct products  $H_r^2 \ltimes H_\lambda^2$  of hyperbolic disks.

**Example 4.2** ( $\mathfrak{g}_2(\lambda, \alpha, \beta, w_1, w_2)$ ). Here we have the 5-dimensional family of *solvable* Lie algebras with the bracket relations

$$\begin{aligned} [W, Z] &= \lambda W, \\ [Z, X] &= \alpha X + \beta Y + w_1 W, \\ [Z, Y] &= -\beta X + \alpha Y + w_2 W. \end{aligned}$$

( $\mathcal{AK}$ ) : Since  $\theta_1 = \theta_2 = a = 0$ , the family  $\mathfrak{g}_2$  is *almost Kähler* if and only if  $\alpha = 0$ . This yields a 4-dimensional family  $\mathfrak{g}_2^{\mathcal{AK}}(\lambda, \beta, w_1, w_2)$  with the Lie bracket relations

$$\begin{aligned} [W, Z] &= \lambda W, \\ [Z, X] &= \beta Y + w_1 W, \\ [Z, Y] &= -\beta X + w_2 W. \end{aligned}$$

( $\mathcal{I}$ ) : Because  $z_1 = z_2 = z_3 = z_4 = 0$ , the structure  $J$  is *integrable* if and only if  $w_1 = w_2 = 0$ . This provides us with a 3-dimensional family  $\mathfrak{g}_2^{\mathcal{I}}(\lambda, \alpha, \beta)$  given by

$$\begin{aligned} [W, Z] &= \lambda W, \\ [Z, X] &= \alpha X + \beta Y, \\ [Z, Y] &= -\beta X + \alpha Y. \end{aligned}$$

The corresponding simply connected Lie group is clearly a semidirect product  $H_\lambda^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{K}$ ) : The family  $\mathfrak{g}_2$  is *Kähler* if and only if it is both almost Kähler and integrable i.e. if  $\alpha = w_1 = w_2 = 0$ . Here we obtain the 2-dimensional family  $\mathfrak{g}_2^{\mathcal{K}}(\lambda, \beta)$  given by

$$\begin{aligned} [W, Z] &= \lambda W, \\ [Z, X] &= \beta Y, \\ [Z, Y] &= -\beta X. \end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $H_\lambda^2 \ltimes \mathbb{R}^2$ .

**Example 4.3** ( $\mathfrak{g}_3(\alpha, \beta, w_1, w_2, \theta_2)$ ). This is a 5-dimensional family of *solvable* Lie algebras with  $r = \theta_1 = 0$ ,  $\theta_2 \neq 0$  and  $\lambda = -2\alpha$ . The bracket relations are given by

$$\begin{aligned} [W, Z] &= -2\alpha W, \\ [Z, X] &= \alpha X + \beta Y + w_1 W, \\ [Z, Y] &= -\beta X + \alpha Y + w_2 W, \\ [Y, X] &= \theta_2 W. \end{aligned}$$

( $\mathcal{AK}$ ) : Here  $a = \theta_1 = 0$ ,  $\lambda = -2\alpha \neq 0$  and  $\theta_2 \neq 0$ . Thus, the family  $\mathfrak{g}_3$  is *almost Kähler* if and only if  $\theta_2 = -2\alpha = \lambda \neq 0$ , which gives the 4-dimensional family  $\mathfrak{g}_3^{\mathcal{AK}}(\alpha, \beta, w_1, w_2)$  with

$$[W, Z] = -2\alpha W,$$

$$\begin{aligned}
[Z, X] &= \alpha X + \beta Y + w_1 W, \\
[Z, Y] &= -\beta X + \alpha Y + w_2 W, \\
[Y, X] &= -2\alpha W.
\end{aligned}$$

( $\mathcal{I}$ ) : Because  $z_1 = z_2 = z_3 = z_4 = 0$ , we see that the structure  $J$  is *integrable* if and only if  $w_1 = w_2 = 0$ , and obtain the family  $\mathfrak{g}_3^{\mathcal{I}}(\alpha, \beta, \theta_2)$  with

$$\begin{aligned}
[W, Z] &= -2\alpha W, \\
[Z, X] &= \alpha X + \beta Y, \\
[Z, Y] &= -\beta X + \alpha Y, \\
[Y, X] &= \theta_2 W.
\end{aligned}$$

( $\mathcal{K}$ ) : The family  $\mathfrak{g}_3$  is *Kähler* if and only if  $\theta_2 = -2\alpha \neq 0$  and  $w_1 = w_2 = 0$ . This provides  $\mathfrak{g}_3^{\mathcal{K}}(\alpha, \beta)$  with

$$\begin{aligned}
[W, Z] &= -2\alpha W, \\
[Z, X] &= \alpha X + \beta Y, \\
[Z, Y] &= -\beta X + \alpha Y, \\
[Y, X] &= -2\alpha W.
\end{aligned}$$

#### 5. CASE (B) - ( $\lambda \neq 0$ AND $(\lambda - \alpha)^2 + \beta^2 = 0$ )

**Example 5.1** ( $\mathfrak{g}_4(\lambda, z_2, w_1, w_2)$ ). Here we have the 4-dimensional family of *solvable* Lie algebras with the bracket relations

$$\begin{aligned}
[W, Z] &= \lambda W, \\
[Z, X] &= \lambda X + w_1 W, \\
[Z, Y] &= \lambda Y + z_2 Z + w_2 W, \\
[W, Y] &= -z_2 W, \\
[Y, X] &= -z_2 X - \frac{z_2 w_1}{\lambda} W.
\end{aligned}$$

( $\mathcal{AK}$ ) : From  $\theta_1 = 2a = 0$ ,  $\alpha = \lambda \neq 0$  and  $\lambda\theta_2 = -z_2 w_1$ , we see that  $\mathfrak{g}_4$  is *almost Kähler* if and only if  $\theta_2 = -2\alpha = -2\lambda$ . This implies that  $2\lambda^2 = w_1 z_2 \neq 0$  and we yield the 3-dimensional  $\mathfrak{g}_4^{\mathcal{AK}}(\lambda, z_2, w_2)$  with the following Lie bracket relations

$$\begin{aligned}
[W, Z] &= \lambda W, \\
[Z, X] &= \lambda X + \frac{2\lambda^2}{z_2} W, \\
[Z, Y] &= \lambda Y + z_2 Z + w_2 W, \\
[W, Y] &= -z_2 W, \\
[Y, X] &= -z_2 X - 2\lambda W.
\end{aligned}$$

( $\mathcal{I}$ ) : Because  $z_1 = z_3 = z_4 = 0$ , the structure  $J$  is *integrable* if and only if  $2z_2 + w_1 = w_2 = 0$ . This gives a 2-dimensional family  $\mathfrak{g}_4^{\mathcal{I}}(\lambda, z_2)$  with bracket

relations

$$\begin{aligned}
[W, Z] &= \lambda W, \\
[Z, X] &= \lambda X - 2z_2 W, \\
[Z, Y] &= \lambda Y + z_2 Z, \\
[W, Y] &= -z_2 W, \\
[Y, X] &= -z_2 X + \frac{2z_2^2}{\lambda} W.
\end{aligned}$$

( $\mathcal{K}$ ) : Here the *Kähler* condition is never satisfied. This requires  $\lambda^2 + z_2^2 = 0$ , which clearly has no real solutions for  $\lambda \neq 0$ . Hence  $\mathfrak{g}_4^{\mathcal{K}} = \{0\}$ .

#### 6. CASE (C) - ( $\lambda = 0$ , $r \neq 0$ AND $(a\beta - \alpha b) \neq 0$ )

**Example 6.1** ( $\mathfrak{g}_5(\alpha, a, \beta, b, r)$ ). This is the 5-dimensional family of *solvable* Lie algebras satisfying the bracket relations

$$\begin{aligned}
[Z, X] &= \alpha X + \beta Y - \frac{r(\beta b - \alpha a)}{2(\alpha b - a\beta)} Z - \frac{r(\alpha^2 - \beta^2)}{2(\alpha b - a\beta)} W, \\
[Z, Y] &= -\beta X + \alpha Y - \frac{r(\alpha b + \beta a)}{2(\alpha b - a\beta)} Z + \frac{r\alpha\beta}{(\alpha b - a\beta)} W, \\
[W, X] &= aX + bY - \frac{r(b^2 - a^2)}{2(\alpha b - a\beta)} Z - \frac{r(\alpha a - \beta b)}{2(\alpha b - a\beta)} W, \\
[W, Y] &= -bX + aY - \frac{rab}{(\alpha b - a\beta)} Z + \frac{r(\alpha b + \beta a)}{2(\alpha b - a\beta)} W, \\
[Y, X] &= rX + \frac{ar^2}{2(\alpha b - a\beta)} Z - \frac{\alpha r^2}{2(\alpha b - a\beta)} W.
\end{aligned}$$

( $\mathcal{AK}$ ) : In this case we have the two relations

$$\theta_1 = \frac{ar^2}{2(\alpha b - a\beta)} \quad \text{and} \quad \theta_2 = -\frac{\alpha r^2}{2(\alpha b - a\beta)}.$$

They imply that the structure  $J$  is *almost Kähler* if and only if

$$2a = \frac{ar^2}{2(\alpha b - a\beta)} \quad \text{and} \quad 2\alpha = \frac{\alpha r^2}{2(\alpha b - a\beta)}.$$

The condition  $a\beta - \alpha b \neq 0$  shows that not both  $a$  and  $\alpha$  can be zero, which implies that  $r^2 = 4(\alpha b - a\beta) > 0$ . If we substitute this into the above system, we then yield the following two solutions

$$\begin{aligned}
[Z, X] &= \alpha X + \beta Y \mp \frac{(\beta b - \alpha a)}{\sqrt{\alpha b - a\beta}} Z \mp \frac{(\alpha^2 - \beta^2)}{\sqrt{\alpha b - a\beta}} W, \\
[Z, Y] &= -\beta X + \alpha Y \mp \frac{(\alpha b + \beta a)}{\sqrt{\alpha b - a\beta}} Z \pm \frac{2\alpha\beta}{\sqrt{\alpha b - a\beta}} W, \\
[W, X] &= aX + bY \mp \frac{(b^2 - a^2)}{\sqrt{\alpha b - a\beta}} Z \mp \frac{(\alpha a - \beta b)}{\sqrt{\alpha b - a\beta}} W,
\end{aligned}$$



$$\begin{aligned}
[W, Y] &= -bX + aY \mp \frac{2ab}{\sqrt{\alpha b - a\beta}} Z \pm \frac{(\alpha b + \beta a)}{\sqrt{\alpha b - a\beta}} W, \\
[Y, X] &= \pm 2\sqrt{\alpha b - a\beta} X + 2a Z - 2\alpha W.
\end{aligned}$$

Here we have obtained two 4-dimensional families  $\mathfrak{g}_5^{AK}(\alpha, a, \beta, b)^\pm$  satisfying the almost Kähler condition.

( $\mathcal{I}$ ) : Here we are assuming that  $r \neq 0$ ,  $a\beta - \alpha b \neq 0$  and have

$$\begin{aligned}
z_1 &= -\frac{r(\beta b - \alpha a)}{2(\alpha b - a\beta)}, \quad z_2 = -\frac{r(\alpha b + \beta a)}{2(\alpha b - a\beta)}, \quad z_3 = -\frac{r(b^2 - a^2)}{2(\alpha b - a\beta)}, \\
z_4 &= -\frac{2rab}{2(\alpha b - a\beta)}, \quad w_1 = -\frac{r(\alpha^2 - \beta^2)}{2(\alpha b - a\beta)}, \quad w_2 = \frac{2r\alpha\beta}{2(\alpha b - a\beta)}.
\end{aligned}$$

From this we observe that  $J$  is *integrable* if and only if

$$2(\alpha b + \beta a) + b^2 - a^2 + \alpha^2 - \beta^2 = 0$$

and

$$\beta b - \alpha a - ab + \alpha\beta = 0.$$

A simple calculation shows that  $J$  is integrable if and only if  $a = \beta$  and  $b = -\alpha$ . This provides us with the 3-dimensional family  $\mathfrak{g}_5^{\mathcal{I}}(\alpha, \beta, r)$  fulfilling the bracket relations

$$\begin{aligned}
[Z, X] &= \alpha X + \beta Y - \frac{r\alpha\beta}{(\beta^2 + \alpha^2)} Z + \frac{r(\alpha^2 - \beta^2)}{2(\beta^2 + \alpha^2)} W, \\
[Z, Y] &= -\beta X + \alpha Y - \frac{r(\alpha^2 - \beta^2)}{2(\beta^2 + \alpha^2)} Z - \frac{r\alpha\beta}{(\beta^2 + \alpha^2)} W, \\
[W, X] &= \beta X - \alpha Y + \frac{r(\alpha^2 - \beta^2)}{2(\beta^2 + \alpha^2)} Z + \frac{r\alpha\beta}{(\beta^2 + \alpha^2)} W, \\
[W, Y] &= \alpha X + \beta Y - \frac{r\alpha\beta}{(\beta^2 + \alpha^2)} Z + \frac{r(\alpha^2 - \beta^2)}{2(\beta^2 + \alpha^2)} W, \\
[Y, X] &= rX - \frac{\beta r^2}{2(\beta^2 + \alpha^2)} Z + \frac{\alpha r^2}{2(\beta^2 + \alpha^2)} W.
\end{aligned}$$

( $\mathcal{K}$ ) : Combining the observations  $r^2 = 4(\alpha b - a\beta) > 0$  from ( $\mathcal{AK}$ ) and  $a = \beta$ ,  $b = -\alpha$  from ( $\mathcal{I}$ ) we see that the *Kähler* condition is never satisfied for the Lie algebra  $\mathfrak{g}_5$ , so  $\mathfrak{g}_5^{\mathcal{K}} = \{0\}$ .

#### 7. CASE (D) - ( $\lambda = 0$ , $r \neq 0$ AND $(a\beta - \alpha b) = 0$ )

**Example 7.1** ( $\mathfrak{g}_6(z_1, z_2, z_3, r, \theta_1, \theta_2)$ ). Here we have a 6-dimensional family of *solvable* Lie algebras with  $z_1^2 = -w_1 z_3 \neq 0$ ,

$$z_4 = \frac{z_3(r + 2z_2)}{2z_1}, \quad w_1 = -\frac{z_1^2}{z_3} \quad \text{and} \quad w_2 = \frac{z_1(r - 2z_2)}{2z_3}.$$

In this case the Lie bracket relations are given by

$$[Z, X] = z_1 Z - \frac{z_1^2}{z_3} W,$$

$$\begin{aligned}
[Z, Y] &= z_2 Z + \frac{z_1(r - 2z_2)}{2z_3} W, \\
[W, X] &= z_3 Z - z_1 W, \\
[W, Y] &= \frac{z_3(r + 2z_2)}{2z_1} Z - z_2 W, \\
[Y, X] &= rX + \theta_1 Z + \theta_2 W.
\end{aligned}$$

( $\mathcal{AK}$ ) : Because  $a = \alpha = 0$ , the structure  $J$  is *almost Kähler* if and only if  $\theta_1 = \theta_2 = 0$ . This provides the 4-dimensional family  $\mathfrak{g}_6^{\mathcal{AK}}(z_1, z_2, z_3, r)$  with the bracket relations

$$\begin{aligned}
[Z, X] &= z_1 Z - \frac{z_1^2}{z_3} W, \\
[Z, Y] &= z_2 Z + \frac{z_1(r - 2z_2)}{2z_3} W, \\
[W, X] &= z_3 Z - z_1 W, \\
[W, Y] &= \frac{z_3(r + 2z_2)}{2z_1} Z - z_2 W, \\
[Y, X] &= rX.
\end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $H_r^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{I}$ ) : We note that

$$z_1 \neq 0, \quad z_4 = \frac{z_3(r + 2z_2)}{2z_1}, \quad w_1 = -\frac{z_1^2}{z_3} \quad \text{and} \quad w_2 = \frac{z_1(r - 2z_2)}{2z_3}.$$

Here the structure  $J$  is *integrable* if and only if

$$z_1^2 = (2z_2 + z_3)z_3 \quad \text{and} \quad z_1^2(4z_3 - r + 2z_2) = z_3^2(r + 2z_2).$$

By multiplying the first equation by  $(4z_3 - r + 2z_2)$ , we see that these two conditions imply

$$(2z_2 + z_3)(4z_3 - r + 2z_2) = z_3(r + 2z_2),$$

which can be rewritten as

$$(z_2 + z_3)(2(z_2 + z_3) - r) = 0.$$

If  $z_2 + z_3 = 0$ , our earlier conditions give  $z_1^2 + z_3^2 = 0$ , which is not possible for nonzero real constants. Thus we must have  $z_2 + z_3 \neq 0$ . This leaves us with  $r = 2(z_2 + z_3)$ . We conclude that  $J$  is *integrable* if and only if

$$r = \frac{z_1^2 + z_3^2}{z_3} \quad \text{and} \quad z_2 = \frac{z_1^2 - z_3^2}{2z_3}.$$

This leads to the following bracket relations for the 4-dimensional family  $\mathfrak{g}_6^{\mathcal{I}}(z_1, z_3, \theta_1, \theta_2)$

$$[Z, X] = z_1 Z - \frac{z_1^2}{z_3} W,$$

$$\begin{aligned}
[Z, Y] &= \frac{z_1^2 - z_3^2}{2z_3} Z + z_1 W, \\
[W, X] &= z_3 Z - z_1 W, \\
[W, Y] &= z_1 Z - \frac{z_1^2 - z_3^2}{2z_3} W, \\
[Y, X] &= \frac{z_1^2 + z_3^2}{z_3} X + \theta_1 Z + \theta_2 W.
\end{aligned}$$

( $\mathcal{K}$ ) : In this case the *Kähler* condition is fulfilled if and only if

$$\theta_1 = \theta_2 = 0, \quad r = \frac{z_1^2 + z_3^2}{z_3} \quad \text{and} \quad z_2 = \frac{z_1^2 - z_3^2}{2z_3}.$$

Hence we have the 2-dimensional family  $\mathfrak{g}_6^{\mathcal{K}}(z_1, z_3)$  with bracket relations

$$\begin{aligned}
[Z, X] &= z_1 Z - \frac{z_1^2}{z_3} W, \\
[Z, Y] &= \frac{z_1^2 - z_3^2}{2z_3} Z + z_1 W, \\
[W, X] &= z_3 Z - z_1 W, \\
[W, Y] &= z_1 Z - \frac{z_1^2 - z_3^2}{2z_3} W, \\
[Y, X] &= \frac{z_1^2 + z_3^2}{z_3} X.
\end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $H_r^2 \ltimes \mathbb{R}^2$ .

**Example 7.2** ( $\mathfrak{g}_7(z_2, w_1, w_2, \theta_1, \theta_2)$ ). This is a 5-dimensional family of *solvable* Lie algebras with  $z_1 = z_3 = z_4 = 0$ ,  $r = 2z_2$  and  $w_1 \neq 0$ . The bracket relations are

$$\begin{aligned}
[Z, X] &= w_1 W, \\
[Z, Y] &= z_2 Z + w_2 W, \\
[W, Y] &= -z_2 W, \\
[Y, X] &= 2z_2 X + \theta_1 Z + \theta_2 W.
\end{aligned}$$

( $\mathcal{AK}$ ) : Because  $\alpha = a = 0$ , the family belongs to the *almost Kähler* class if and only if  $\theta_1 = \theta_2 = 0$ . This gives the 3-dimensional family  $\mathfrak{g}_7^{\mathcal{AK}}(z_2, w_1, w_2)$  with

$$\begin{aligned}
[Z, X] &= w_1 W, \\
[Z, Y] &= z_2 Z + w_2 W, \\
[W, Y] &= -z_2 W, \\
[Y, X] &= 2z_2 X.
\end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $H_{2z_2}^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{I}$ ) : We have  $z_1 = z_3 = z_4 = 0$ . Thus the structure  $J$  is *integrable* if and only if  $2z_2 + w_1 = w_2 = 0$  and we yield a 3-dimensional family  $\mathfrak{g}_7^{\mathcal{I}}(z_2, \theta_1, \theta_2)$  given by

$$\begin{aligned} [Z, X] &= -2z_2W, \\ [Z, Y] &= z_2Z, \\ [W, Y] &= -z_2W, \\ [Y, X] &= 2z_2X + \theta_1Z + \theta_2W. \end{aligned}$$

Note that  $z_2 \neq 0$ .

( $\mathcal{K}$ ) : The family  $\mathfrak{g}_7$  is of the *Kähler* class if and only if  $\theta_1 = \theta_2 = 0$ ,  $w_2 = 0$  and  $w_1 = -2z_2 \neq 0$  providing the 1-dimensional family  $\mathfrak{g}_7^{\mathcal{K}}(z_2)$  with

$$\begin{aligned} [Z, X] &= -2z_2W, \\ [Z, Y] &= z_2Z, \\ [W, Y] &= -z_2W, \\ [Y, X] &= 2z_2X. \end{aligned}$$

The corresponding simply connected Lie group is clearly a semidirect product  $H_{2z_2}^2 \ltimes \mathbb{R}^2$ .

**Example 7.3** ( $\mathfrak{g}_8(z_2, z_4, w_2, r, \theta_1, \theta_2)$ ). Here we have the 6-dimensional family of *solvable* Lie algebras with bracket relations

$$\begin{aligned} [Z, Y] &= z_2Z + w_2W, \\ [W, Y] &= z_4Z - z_2W, \\ [Y, X] &= rX + \theta_1Z + \theta_2W. \end{aligned}$$

( $\mathcal{AK}$ ) : Since  $\alpha = a = 0$ , the family is almost Kähler if and only if  $\theta_1 = \theta_2 = 0$ , giving the 4-dimensional family  $\mathfrak{g}_8^{\mathcal{AK}}(z_2, z_4, w_2, r)$  with

$$\begin{aligned} [Z, Y] &= z_2Z + w_2W, \\ [W, Y] &= z_4Z - z_2W, \\ [Y, X] &= rX. \end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $H_r^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{I}$ ) : We have  $z_1 = z_3 = w_1 = 0$ , so  $J$  is integrable if and only if  $z_2 = z_4 + w_2 = 0$ , so we yield the family  $\mathfrak{g}_8^{\mathcal{I}}(w_2, r, \theta_1, \theta_2)$  with

$$\begin{aligned} [Z, Y] &= w_2W, \\ [W, Y] &= -w_2Z, \\ [Y, X] &= rX + \theta_1Z + \theta_2W. \end{aligned}$$

( $\mathcal{K}$ ) : The family is Kähler if and only if  $\theta_1 = \theta_2 = z_2 = z_4 + w_2 = 0$ . We get the 2-dimensional family  $\mathfrak{g}_8^{\mathcal{K}}(w_2, r)$  with

$$\begin{aligned} [Z, Y] &= w_2W, \\ [W, Y] &= -w_2Z, \end{aligned}$$

$$[Y, X] = rX.$$

The corresponding simply connected Lie group is clearly a semidirect product  $H_r^2 \ltimes \mathbb{R}^2$

**Example 7.4** ( $\mathfrak{g}_9(z_2, z_3, z_4, \theta_1, \theta_2)$ ). This is the 5-dimensional family of *solvable* Lie algebras with  $z_3 \neq 0$  and bracket relations

$$\begin{aligned} [Z, Y] &= z_2 Z, \\ [W, X] &= z_3 Z, \\ [W, Y] &= z_4 Z - z_2 W, \\ [Y, X] &= -2z_2 X + \theta_1 Z + \theta_2 W. \end{aligned}$$

( $\mathcal{AK}$ ) : Because  $\alpha = a = 0$ , we see that the family is in the *almost Kähler* class if and only if  $\theta_1 = \theta_2 = 0$ , providing  $\mathfrak{g}_9^{\mathcal{AK}}(z_2, z_3, z_4)$  with

$$\begin{aligned} [Z, Y] &= z_2 Z, \\ [W, X] &= z_3 Z, \\ [W, Y] &= z_4 Z - z_2 W, \\ [Y, X] &= -2z_2 X. \end{aligned}$$

( $\mathcal{I}$ ) : From  $z_1 = w_1 = w_2 = 0$ , we find that  $J$  is *integrable* if and only if  $2z_2 + z_3 = z_4 = 0$ , from which we yield  $\mathfrak{g}_9^{\mathcal{I}}(z_2, \theta_1, \theta_2)$  with the relations

$$\begin{aligned} [Z, Y] &= z_2 Z, \\ [W, X] &= -2z_2 Z, \\ [W, Y] &= -z_2 W, \\ [Y, X] &= -2z_2 X + \theta_1 Z + \theta_2 W. \end{aligned}$$

( $\mathcal{K}$ ) : The family is *Kähler* if and only if  $\theta_1 = \theta_2 = 0$ ,  $z_3 = -2z_2$  and  $z_4 = 0$ . Here we obtain the 1-dimensional family  $\mathfrak{g}_9^{\mathcal{K}}(z_2)$  with

$$\begin{aligned} [Z, Y] &= z_2 Z, \\ [W, X] &= -2z_2 Z, \\ [W, Y] &= -z_2 W, \\ [Y, X] &= -2z_2 X. \end{aligned}$$

The corresponding simply connected Lie group is clearly a semidirect product  $H_{2z_2}^2 \ltimes \mathbb{R}^2$

#### 8. CASE (E) - ( $\lambda = 0$ , $r = 0$ AND $\alpha b - a\beta \neq 0$ )

**Example 8.1** ( $\mathfrak{g}_{10}(\alpha, a, \beta, b)$ ). Here we have the 4-dimensional family of *solvable* Lie algebras with the bracket relations

$$\begin{aligned} [Z, X] &= \alpha X + \beta Y, \\ [Z, Y] &= -\beta X + \alpha Y, \\ [W, X] &= aX + bY, \\ [W, Y] &= -bX + aY. \end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{AK}$ ) : Since  $\theta_1 = \theta_2 = 0$ , we see that the family is *almost Kähler* if and only if  $a = \alpha = 0$ . But then the requirement  $\alpha b - a\beta \neq 0$  is not true, so this family can not be almost Kähler, so  $\mathfrak{g}_{10}^{\mathcal{AK}} = \{0\}$ .

( $\mathcal{I}$ ) : Because  $z_1 = z_2 = z_3 = z_4 = w_1 = w_2 = 0$ , we can see that the structure  $J$  is always *integrable* in this family, therefore  $\mathfrak{g}_{10}^{\mathcal{I}} = \mathfrak{g}_{10}(\alpha, a, \beta, b)$ .

( $\mathcal{K}$ ) : The Lie algebra  $\mathfrak{g}_{10}$  is never in the *Kähler* class, hence  $\mathfrak{g}_{10}^{\mathcal{K}} = \{0\}$ .

### 9. CASE (F) - ( $\lambda = 0$ , $r = 0$ AND $\alpha b - a\beta = 0$ )

In this case our analysis divides into disjoint cases parametrized by  $\Lambda = (\alpha, a, \beta, b)$ . The variables are assumed to be zero if and only if they are marked by 0. For example, if  $\Lambda = (0, a, \beta, 0)$  then the two variable  $\alpha$  and  $b$  are assumed to be zero and  $a$  and  $\beta$  to be non-zero.

**Example 9.1** ( $\mathfrak{g}_{11}(z_1, z_2, z_3, w_1, \theta_1, \theta_2)$ ). This is a 6-dimensional family of *solvable* Lie algebras. Here  $\Lambda = (0, 0, 0, 0)$  and  $z_1 \neq 0$ , giving the Lie bracket relations

$$\begin{aligned} [Z, X] &= z_1 Z + w_1 W, \\ [Z, Y] &= z_2 Z + \frac{z_2 w_1}{z_1} W, \\ [W, X] &= z_3 Z - z_1 W, \\ [W, Y] &= \frac{z_2 z_3}{z_1} Z - z_2 W, \\ [Y, X] &= \theta_1 Z + \theta_2 W. \end{aligned}$$

( $\mathcal{AK}$ ) : Because  $\alpha = a = 0$ , we see that  $\mathfrak{g}_{11}$  is in the *almost Kähler* class if and only if  $\theta_1 = \theta_2 = 0$ . This gives a 4-dimensional family  $\mathfrak{g}_{11}^{\mathcal{AK}}(z_1, z_2, z_3, w_1)$  with

$$\begin{aligned} [Z, X] &= z_1 Z + w_1 W, \\ [Z, Y] &= z_2 Z + \frac{z_2 w_1}{z_1} W, \\ [W, X] &= z_3 Z - z_1 W, \\ [W, Y] &= \frac{z_2 z_3}{z_1} Z - z_2 W. \end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{I}$ ) : From the relations

$$z_1 \neq 0, \quad z_4 = \frac{z_2 z_3}{z_1} \quad \text{and} \quad w_2 = \frac{z_2 w_1}{z_1},$$

we see that the structure  $J$  is *integrable* if and only if

$$z_3 + w_1 = -2z_2 \quad \text{and} \quad 2z_1^2 - z_2(z_3 + w_1) = 0.$$

This implies the impossible  $z_1^2 + z_2^2 = 0$ , so the structure  $J$  is never integrable, therefore  $\mathfrak{g}_{11}^{\mathcal{K}} = \{0\}$ .

( $\mathcal{K}$ ) : This family  $\mathfrak{g}_{11}$  is never in the *Kähler* class, so  $\mathfrak{g}_{11}^{\mathcal{K}} = \{0\}$ .

**Example 9.2** ( $\mathfrak{g}_{12}(z_3, w_1, w_2, \theta_1, \theta_2)$ ). Here we have the 5-dimensional family of *solvable* Lie algebras. Here  $\Lambda = (0, 0, 0, 0)$ ,  $z_1 = 0$  and  $w_1 \neq 0$ , which gives

$$z_2 = 0 \quad \text{and} \quad z_4 = \frac{z_3 w_2}{w_1}$$

and the Lie bracket relations

$$\begin{aligned} [Z, X] &= w_1 W, \\ [Z, Y] &= w_2 W, \\ [W, X] &= z_3 Z, \\ [W, Y] &= \frac{z_3 w_2}{w_1} Z, \\ [Y, X] &= \theta_1 Z + \theta_2 W. \end{aligned}$$

( $\mathcal{AK}$ ) : Since  $\alpha = a = 0$ , this family is *almost Kähler* if and only if  $\theta_1 = \theta_2 = 0$ . This yields the 3-dimensional family  $\mathfrak{g}_{12}^{\mathcal{AK}}(z_3, w_1, w_2)$  with the relations

$$\begin{aligned} [Z, X] &= w_1 W, \\ [Z, Y] &= w_2 W, \\ [W, X] &= z_3 Z, \\ [W, Y] &= \frac{z_3 w_2}{w_1} Z. \end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{I}$ ) : For  $\mathfrak{g}_{12}$  we have

$$z_1 = z_2 = 0, \quad z_4 = \frac{z_3 w_2}{w_1} \quad \text{and} \quad w_1 \neq 0.$$

We see that the structure  $J$  is *integrable* if and only if

$$z_3 + w_1 = \frac{z_3 w_2}{w_1} + w_2 = 0,$$

or equivalently,  $z_3 = -w_1 \neq 0$ . Here we yield the 4-dimensional family  $\mathfrak{g}_{12}^{\mathcal{I}}(w_1, w_2, \theta_1, \theta_2)$  which has the Lie bracket relations

$$\begin{aligned} [Z, X] &= w_1 W, \\ [Z, Y] &= w_2 W, \\ [W, X] &= -w_1 Z, \\ [W, Y] &= -w_2 Z, \\ [Y, X] &= \theta_1 Z + \theta_2 W. \end{aligned}$$

( $\mathcal{K}$ ) : The family is *Kähler* if and only if  $\theta_1 = \theta_2 = 0$  and  $z_3 = -w_1 \neq 0$ , which gives the 2-dimensional family  $\mathfrak{g}_{12}^{\mathcal{K}}(w_1, w_2)$  with relations

$$\begin{aligned} [Z, X] &= w_1 W, \\ [Z, Y] &= w_2 W, \\ [W, X] &= -w_1 Z, \\ [W, Y] &= -w_2 Z. \end{aligned}$$

The corresponding simply connected Lie group is clearly a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

**Example 9.3** ( $\mathfrak{g}_{13}(z_3, z_4, \theta_1, \theta_2)$ ). This is a 4-dimensional family of *nilpotent* Lie algebras. Here  $\Lambda = (0, 0, 0, 0)$ ,  $z_1 = w_1 = 0$  and  $z_3 \neq 0$ , which gives  $z_2 = w_2 = 0$  and the solutions

$$\begin{aligned} [W, X] &= z_3 Z, \\ [W, Y] &= z_4 Z, \\ [Y, X] &= \theta_1 Z + \theta_2 W. \end{aligned}$$

( $\mathcal{AK}$ ) : Because  $\alpha = a = 0$ , we see that  $\mathfrak{g}_{13}$  family is in the *almost Kähler* class if and only if  $\theta_1 = \theta_2 = 0$ . We get the 2-dimensional family  $\mathfrak{g}_{13}^{\mathcal{AK}}(z_3, z_4)$  with

$$\begin{aligned} [W, X] &= z_3 Z, \\ [W, Y] &= z_4 Z. \end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{I}$ ) : The condition  $z_1 = z_2 = w_1 = w_2 = 0$  gives that  $J$  is *integrable* if and only if  $z_3 = z_4 = 0$ , contradicting  $z_3 \neq 0$ . Hence  $\mathfrak{g}_{13}^{\mathcal{I}} = \{0\}$

( $\mathcal{K}$ ) : This family  $\mathfrak{g}_{13}$  is never in the *Kähler* class, so  $\mathfrak{g}_{13}^{\mathcal{K}} = \{0\}$ .

**Example 9.4** ( $\mathfrak{g}_{14}(z_2, z_4, w_2, \theta_1, \theta_2)$ ). Here we have the 5-dimensional family of *solvable* Lie algebras with  $\Lambda = (0, 0, 0, 0)$  and  $z_1 = z_3 = w_1 = 0$  and the bracket relations

$$\begin{aligned} [Z, Y] &= z_2 Z + w_2 W, \\ [W, Y] &= z_4 Z - z_2 W, \\ [Y, X] &= \theta_1 Z + \theta_2 W. \end{aligned}$$

( $\mathcal{AK}$ ) : The condition  $\alpha = a = 0$  implies that this family is *almost Kähler* if and only if  $\theta_1 = \theta_2 = 0$ . We get the 3-dimensional family  $\mathfrak{g}_{14}^{\mathcal{AK}}(z_2, z_4, w_2)$  given by

$$\begin{aligned} [Z, Y] &= z_2 Z + w_2 W, \\ [W, Y] &= z_4 Z - z_2 W. \end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .



( $\mathcal{I}$ ) : From  $z_1 = z_3 = w_1 = 0$  we see that  $J$  is *integrable* if and only if  $z_2 = 0$  and  $z_4 + w_2 = 0$ . This gives a 3-dimensional family  $\mathfrak{g}_{14}^{\mathcal{I}}(w_2, \theta_1, \theta_2)$  with Lie bracket relations

$$\begin{aligned} [Z, Y] &= w_2 W, \\ [W, Y] &= -w_2 Z, \\ [Y, X] &= \theta_1 Z + \theta_2 W. \end{aligned}$$

( $\mathcal{K}$ ) : The family is *Kähler* if and only if  $\theta_1 = \theta_2 = 0$ ,  $z_2 = 0$  and  $z_4 + w_2 = 0$ . We get a 1-dimensional family  $\mathfrak{g}_{14}^{\mathcal{K}}(w_2)$  with

$$\begin{aligned} [Z, Y] &= w_2 W, \\ [W, Y] &= -w_2 Z. \end{aligned}$$

The corresponding simply connected Lie group is clearly a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

**Example 9.5** ( $\mathfrak{g}_{15}(\alpha, w_1, w_2)$ ). This is a 3-dimensional family of *solvable* Lie algebras with  $\Lambda = (\alpha, 0, 0, 0)$ , which gives

$$z_1 = z_2 = z_3 = z_4 = \theta_1 = \theta_2 = 0$$

and

$$\begin{aligned} [Z, X] &= \alpha X + w_1 W, \\ [Z, Y] &= \alpha Y + w_2 W. \end{aligned}$$

( $\mathcal{AK}$ ) : Here,  $a = \theta_1 = \theta_2 = 0$  and  $\alpha \neq 0$ , so we see that the *almost Kähler* is never fulfilled, hence  $\mathfrak{g}_{15}^{\mathcal{AK}} = \{0\}$ .

( $\mathcal{I}$ ) : The conditions  $z_1 = z_2 = z_3 = z_4 = 0$  imply that  $J$  is *integrable* if and only if  $w_1 = w_2 = 0$ . We yield a 1-dimensional family  $\mathfrak{g}_{15}^{\mathcal{I}}(\alpha)$  with

$$\begin{aligned} [Z, X] &= \alpha X, \\ [Z, Y] &= \alpha Y. \end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{K}$ ) : Here the *Kähler* condition is never satisfied, so  $\mathfrak{g}_{15}^{\mathcal{K}} = \{0\}$ .

**Example 9.6** ( $\mathfrak{g}_{16}(\beta, w_1, w_2, \theta_1, \theta_2)$ ). Here we have a 5-dimensional family of Lie algebras. They are *not solvable* in general. Here  $\Lambda = (0, 0, \beta, 0)$ , which implies  $z_1 = z_2 = z_3 = z_4 = 0$ . This gives

$$\begin{aligned} [Z, X] &= \beta Y + w_1 W, \\ [Z, Y] &= -\beta X + w_2 W, \\ [Y, X] &= \theta_1 Z + \theta_2 W. \end{aligned}$$

( $\mathcal{AK}$ ) : We have  $\alpha = a = 0$ , so this family is *almost Kähler* if and only if  $\theta_1 = \theta_2 = 0$ . We get the 3-dimensional family  $\mathfrak{g}_{16}^{\mathcal{AK}}(\beta, w_1, w_2)$  with

$$\begin{aligned} [Z, X] &= \beta Y + w_1 W, \\ [Z, Y] &= -\beta X + w_2 W. \end{aligned}$$

( $\mathcal{I}$ ) : We have  $z_1 = z_2 = z_3 = z_4 = 0$ , so  $J$  is *integrable* if and only if  $w_1 = w_2 = 0$ , giving the 3-dimensional family  $\mathfrak{g}_{16}^{\mathcal{I}}(\beta, \theta_1, \theta_2)$  with

$$\begin{aligned} [Z, X] &= \beta Y, \\ [Z, Y] &= -\beta X, \\ [Y, X] &= \theta_1 Z + \theta_2 W. \end{aligned}$$

( $\mathcal{K}$ ) : The family  $\mathfrak{g}_{16}$  is *Kähler* if and only if  $\theta_1 = \theta_2 = 0$  and  $w_1 = w_2 = 0$ . We obtain the 1-dimensional family  $\mathfrak{g}_{16}^{\mathcal{K}}(\beta)$  with Lie bracket relations

$$\begin{aligned} [Z, X] &= \beta Y, \\ [Z, Y] &= -\beta X. \end{aligned}$$

The corresponding simply connected Lie group is clearly a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

**Example 9.7** ( $\mathfrak{g}_{17}(\alpha, a, w_1, w_2)$ ). This is a 4-dimensional family of *solvable* Lie algebras  $\Lambda = (\alpha, a, 0, 0)$ , which gives

$$z_1 = -\frac{aw_1}{\alpha}, \quad z_2 = -\frac{aw_2}{\alpha}, \quad z_3 = -\frac{a^2w_1}{\alpha^2}, \quad z_4 = -\frac{a^2w_2}{\alpha^2}, \quad \theta_1 = 0, \quad \theta_2 = 0.$$

Thus the bracker relations are

$$\begin{aligned} [Z, X] &= \alpha X - \frac{aw_1}{\alpha} Z + w_1 W, \\ [Z, Y] &= \alpha Y - \frac{aw_2}{\alpha} Z + w_2 W, \\ [W, X] &= aX - \frac{a^2w_1}{\alpha^2} Z + \frac{aw_1}{\alpha} W, \\ [W, Y] &= aY - \frac{a^2w_2}{\alpha^2} Z + \frac{aw_2}{\alpha} W. \end{aligned}$$

( $\mathcal{AK}$ ) : Because  $\theta_1 = \theta_2 = 0$ ,  $\alpha \neq 0$  and  $a \neq 0$ , the *almost Kähler* condition is never satisfied, hence  $\mathfrak{g}_{17}^{\mathcal{AK}} = \{0\}$ .

( $\mathcal{I}$ ) : From

$$z_1 = -\frac{aw_1}{\alpha}, \quad z_2 = -\frac{aw_2}{\alpha}, \quad z_3 = -\frac{a^2w_1}{\alpha^2} \quad \text{and} \quad z_4 = -\frac{a^2w_2}{\alpha^2}$$

we see that the structure  $J$  is *integrable* if and only if

$$(\alpha^2 - a^2) w_1 = 2a\alpha w_2 \quad \text{and} \quad (\alpha^2 - a^2) w_2 = -2a\alpha w_1. \quad (9.1)$$

Since  $2a\alpha \neq 0$ , we can divide and get

$$w_2 = \frac{\alpha^2 - a^2}{2a\alpha} w_1 = -\frac{(\alpha^2 - a^2)^2}{4a^2\alpha^2} w_2.$$

If  $w_2 \neq 0$ , then we yield the impossible  $(\alpha^2 + a^2)^2 = 0$ . Thus  $J$  is integrable if and only if  $w_1 = w_2 = 0$ . This gives the 2-dimensional family  $\mathfrak{g}_{17}^{\mathcal{I}}(\alpha, a)$  with

$$\begin{aligned} [Z, X] &= \alpha X, \\ [Z, Y] &= \alpha Y, \end{aligned}$$

$$\begin{aligned}[W, X] &= aX, \\ [W, Y] &= aY.\end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{K}$ ) : The family  $\mathfrak{g}_{17}$  never satisfies the *Kähler* condition, so  $\mathfrak{g}_{17}^{\mathcal{K}} = \{0\}$ .

**Example 9.8** ( $\mathfrak{g}_{18}(\beta, b, z_3, z_4, \theta_1, \theta_2)$ ). Here we have a 6-dimensional family of Lie algebras. They are *not solvable* in general. Here  $\Lambda = (0, 0, \beta, b)$ , which gives

$$z_1 = \frac{\beta z_3}{b}, \quad z_2 = \frac{\beta z_4}{b}, \quad w_1 = -\frac{\beta^2 z_3}{b^2} \quad \text{and} \quad w_2 = -\frac{\beta^2 z_4}{b^2}.$$

The bracket relations satisfy

$$\begin{aligned}[Z, X] &= \beta Y + \frac{\beta z_3}{b} Z - \frac{\beta^2 z_3}{b^2} W, \\ [Z, Y] &= -\beta X + \frac{\beta z_4}{b} Z - \frac{\beta^2 z_4}{b^2} W, \\ [W, X] &= bY + z_3 Z - \frac{\beta z_3}{b} W, \\ [W, Y] &= -bX + z_4 Z - \frac{\beta z_4}{b} W, \\ [Y, X] &= \theta_1 Z + \theta_2 W.\end{aligned}$$

( $\mathcal{AK}$ ) : Since  $a = \alpha = 0$ , this family is *almost Kähler* if and only if  $\theta_1 = \theta_2 = 0$ . We get the 4-dimensional family  $\mathfrak{g}_{18}^{\mathcal{K}}(\beta, b, z_3, z_4)$  with

$$\begin{aligned}[Z, X] &= \beta Y + \frac{\beta z_3}{b} Z - \frac{\beta^2 z_3}{b^2} W, \\ [Z, Y] &= -\beta X + \frac{\beta z_4}{b} Z - \frac{\beta^2 z_4}{b^2} W, \\ [W, X] &= bY + z_3 Z - \frac{\beta z_3}{b} W, \\ [W, Y] &= -bX + z_4 Z - \frac{\beta z_4}{b} W.\end{aligned}$$

( $\mathcal{I}$ ) : For the family  $\mathfrak{g}_{18}$  we have

$$z_1 = \frac{\beta z_3}{b}, \quad z_2 = \frac{\beta z_4}{b}, \quad w_1 = -\frac{\beta^2 z_3}{b^2} \quad \text{and} \quad w_2 = -\frac{\beta^2 z_4}{b^2}$$

and we can easily see that  $J$  is *integrable* if and only if

$$(\beta^2 - b^2)z_3 = 2\beta b z_4 \quad \text{and} \quad (\beta^2 - b^2)z_4 = -2\beta b z_3.$$

We notice that this system has the same form as that of (9.1). Thus the structure  $J$  is integrable if and only if  $z_3 = z_4 = 0$ . Here we yield the 4-dimensional family  $\mathfrak{g}_{18}^{\mathcal{I}}(\beta, b, \theta_1, \theta_2)$ , where

$$\begin{aligned}[Z, X] &= \beta Y, \\ [Z, Y] &= -\beta X,\end{aligned}$$

$$\begin{aligned}
[W, X] &= bY, \\
[W, Y] &= -bX, \\
[Y, X] &= \theta_1 Z + \theta_2 W.
\end{aligned}$$

( $\mathcal{K}$ ) : The family is *Kähler* if and only if  $\theta_1 = \theta_2 = 0$  and  $z_3 = z_4 = 0$ . Hence we obtain the 2-dimensional family  $\mathfrak{g}_{18}^{\mathcal{K}}(\beta, b)$  with

$$\begin{aligned}
[Z, X] &= \beta Y, \\
[Z, Y] &= -\beta X, \\
[W, X] &= bY, \\
[W, Y] &= -bX.
\end{aligned}$$

The corresponding simply connected Lie group is clearly a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

**Example 9.9** ( $\mathfrak{g}_{19}(\alpha, \beta, w_1, w_2)$ ). This is a 4-dimensional family of *solvable* Lie algebras. Here  $\Lambda = (\alpha, 0, \beta, 0)$ , which implies that

$$z_1 = z_2 = z_3 = z_4 = \theta_1 = \theta_2 = 0$$

and the bracket relations are

$$\begin{aligned}
[Z, X] &= \alpha X + \beta Y + w_1 W, \\
[Z, Y] &= -\beta X + \alpha Y + w_2 W.
\end{aligned}$$

( $\mathcal{AK}$ ) : Here the *almost Kähler* condition is never satisfied since  $\alpha \neq 0$  and  $\theta_2 = 0$ , hence  $\mathfrak{g}_{19}^{\mathcal{AK}} = \{0\}$ .

( $\mathcal{I}$ ) : Because  $z_1 = z_2 = z_3 = z_4 = 0$ , the structure  $J$  is *integrable* if and only if  $w_1 = w_2 = 0$ . We yield the 2-dimensional family  $\mathfrak{g}_{19}^{\mathcal{H}}(\alpha, \beta)$  given by

$$\begin{aligned}
[Z, X] &= \alpha X + \beta Y, \\
[Z, Y] &= -\beta X + \alpha Y.
\end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{K}$ ) : The *Kähler* condition is never satisfied, so  $\mathfrak{g}_{19}^{\mathcal{K}} = \{0\}$ .

**Example 9.10** ( $\mathfrak{g}_{20}(\alpha, a, \beta, w_1, w_2)$ ). Here we have a 5-dimensional family of *solvable* Lie algebras. Now  $\Lambda = (\alpha, a, \beta, b)$ , which gives

$$\begin{aligned}
z_1 &= -\frac{aw_1}{\alpha}, \quad z_2 = -\frac{aw_2}{\alpha}, \quad z_3 = -\frac{a^2 w_1}{\alpha^2}, \quad z_4 = -\frac{a^2 w_2}{\alpha^2}, \\
b &= \frac{\beta a}{\alpha}, \quad \theta_1 = 0, \quad \theta_2 = 0.
\end{aligned}$$

The bracket relations fulfill

$$\begin{aligned}
[Z, X] &= \alpha X + \beta Y - \frac{aw_1}{\alpha} Z + w_1 W, \\
[Z, Y] &= -\beta X + \alpha Y - \frac{aw_2}{\alpha} Z + w_2 W,
\end{aligned}$$

$$\begin{aligned}
[W, X] &= aX + \frac{\beta a}{\alpha}Y - \frac{a^2 w_1}{\alpha^2}Z + \frac{a}{\alpha}w_1 W, \\
[W, Y] &= -\frac{\beta a}{\alpha}X + aY - \frac{a^2 w_2}{\alpha^2}Z + \frac{a}{\alpha}w_2 W.
\end{aligned}$$

( $\mathcal{AK}$ ) : Since  $\theta_1 = \theta_2 = 0$ ,  $\alpha \neq 0$  and  $a \neq 0$ , the *almost Kähler* condition is never satisfied, hence  $\mathfrak{g}_{20}^{\mathcal{AK}} = \{0\}$ .

( $\mathcal{I}$ ) : Here the coefficients  $z_1, \dots, z_4$ ,  $w_1$  and  $w_2$  are the same as for the family  $\mathfrak{g}_{17}$ . Thus the structure  $J$  is *integrable* if and only if  $w_1 = w_2 = 0$ . This gives the 3-dimensional family  $\mathfrak{g}_{20}^{\mathcal{I}}(\alpha, a, \beta)$  with

$$\begin{aligned}
[Z, X] &= \alpha X + \beta Y, \\
[Z, Y] &= -\beta X + \alpha Y, \\
[W, X] &= aX + \frac{\beta a}{\alpha}Y, \\
[W, Y] &= -\frac{\beta a}{\alpha}X + aY.
\end{aligned}$$

Here the corresponding simply connected Lie group is a semidirect product  $\mathbb{R}^2 \ltimes \mathbb{R}^2$ .

( $\mathcal{K}$ ) : The *Kähler* condition is never satisfied, therefore  $\mathfrak{g}_{20}^{\mathcal{K}} = \{0\}$ .

#### APPENDIX A. THE HOMOGENEOUS SECOND ORDER SYSTEM

In Section 2 we presented the general bracket relations (2.1). To ensure that these actually define a Lie algebra they must satisfy the Jacobi equation. A standard computation shows that this is equivalent to the following system of 14 second order homogeneous polynomial equations in the 14 variables involved

$$\begin{aligned}
0 &= \lambda a, \\
0 &= \lambda b, \\
0 &= -w_2 z_3 + w_1 z_4 - 2\alpha \theta_1 + r z_1, \\
0 &= -2z_4 z_1 + 2z_3 z_2 - 2a \theta_1 + r z_3, \\
0 &= -\lambda z_3 - z_2 b + z_4 \beta - z_1 a + z_3 \alpha, \\
0 &= -\lambda z_4 - z_2 a + z_4 \alpha + z_1 b - z_3 \beta, \\
0 &= \lambda \theta_1 - w_1 z_4 + w_2 z_3 - 2a \theta_2 - r z_1, \\
0 &= -\lambda \theta_2 + 2z_1 w_2 - 2z_2 w_1 - 2\alpha \theta_2 + r w_1, \\
0 &= -w_2 a - w_1 b - z_2 \alpha - z_1 \beta - \alpha r, \\
0 &= -w_2 b + w_1 a - z_2 \beta + z_1 \alpha + r \beta, \\
0 &= \lambda z_1 - w_2 b - z_2 \beta - w_1 a - z_1 \alpha, \\
0 &= z_2 a + z_1 b - z_4 \alpha - z_3 \beta - a r, \\
0 &= z_2 b - z_1 a - z_4 \beta + z_3 \alpha + r b, \\
0 &= \lambda z_2 - w_2 a - z_2 \alpha + w_1 b + z_1 \beta.
\end{aligned}$$

## REFERENCES

- [1] P. Baird and J. Eells, *A conservation law for harmonic maps*, Geometry Symposium Utrecht 1980, Lecture Notes in Mathematics **894**, 1-25, Springer (1981).
- [2] P. Baird, J.C. Wood, *Harmonic morphisms between Riemannian manifolds*, The London Mathematical Society Monographs **29**, Oxford University Press (2003).
- [3] B. Fuglede, *Harmonic morphisms between Riemannian manifolds*, Ann. Inst. Fourier **28** (1978), 107-144.
- [4] A. Gray, L. M. Hervella, *The sixteen classes of almost Hermitian manifolds and their linear invariants*. Ann. Mat. Pura Appl. **123** (1980), 35-58.
- [5] S. Gudmundsson, M. Svensson, *Harmonic morphisms from four-dimensional Lie groups*, J. Geom. Phys. **83** (2014), 1-11.
- [6] S. Gudmundsson, *The Bibliography of Harmonic Morphisms*, [www.matematik.lu.se/matematiklu/personal/sigma/harmonic/bibliography.html](http://www.matematik.lu.se/matematiklu/personal/sigma/harmonic/bibliography.html)
- [7] T. Ishihara, *A mapping of Riemannian manifolds which preserves harmonic functions*, J. Math. Kyoto Univ. **19** (1979), 215-229.
- [8] C. G. J. Jacobi, *Über eine Lösung der partiellen Differentialgleichung  $\Delta(V) = 0$* , J. Reine Angew. Math. **36** (1848), 113-134.

MATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF LUND, BOX 118, LUND 221 00, SWEDEN  
*Email address:* `emmaasv@hotmail.com`

MATHEMATICS, FACULTY OF SCIENCE, UNIVERSITY OF LUND, BOX 118, LUND 221 00, SWEDEN  
*Email address:* `Sigmundur.Gudmundsson@math.lu.se`