

Probing spin-dependent dark matter interactions with ${}^6\text{Li}$

G. Angloher¹, G. Benato², A. Bento^{1,9}, E. Bertoldo^{*a,1}, A. Bertolini¹, R. Breier³, C. Bucci², L. Canonica¹, A. D’Addabbo², S. Di Lorenzo², L. Einfalt^{5,6}, A. Erb^{4,10}, F. v. Feilitzsch⁴, N. Ferreiro Iachellini^{1,11}, S. Fichtinger⁵, D. Fuchs¹, A. Fuss^{5,6}, A. Garai¹, V.M. Ghete⁵, P. Gorla², S. Gupta⁵, D. Hauff¹, M. Jeřkovský³, J. Jochum⁷, M. Kaznacheeva⁴, A. Kinast⁴, H. Kluck⁵, H. Kraus⁸, A. Langenkämper⁴, M. Mancuso^{*b,1}, L. Marini^{2,12}, V. Mokina⁵, A. Nilima¹, M. Olmi², T. Ortmann⁴, C. Pagliarone^{2,13}, V. Palušová³, L. Pattavina^{2,4}, F. Petricca¹, W. Potzel⁴, P. Povinec³, F. Pröbst¹, F. Pucci¹, F. Reindl^{5,6}, J. Rothe⁴, K. Schäffner¹, J. Schieck^{5,6}, D. Schmiedmayer^{5,6}, S. Schönert⁴, C. Schwertner^{5,6}, M. Stahlberg¹, L. Stodolsky¹, C. Strandhagen⁷, R. Strauss⁴, I. Usherov⁷, F. Wagner⁵, M. Willers⁴, V. Zema¹ (CRESST Collaboration)

¹Max-Planck-Institut für Physik, D-80805 München, Germany

²INFN, Laboratori Nazionali del Gran Sasso, I-67100 Assergi, Italy

³Comenius University, Faculty of Mathematics, Physics and Informatics, 84248 Bratislava, Slovakia

⁴Physik-Department and ORIGINS Excellence Cluster, Technische Universität München, D-85747 Garching, Germany

⁵Institut für Hochenergiephysik der Österreichischen Akademie der Wissenschaften, A-1050 Wien, Austria

⁶Atominstytut, Technische Universität Wien, A-1020 Wien, Austria

⁷Eberhard-Karls-Universität Tübingen, D-72076 Tübingen, Germany

⁸Department of Physics, University of Oxford, Oxford OX1 3RH, United Kingdom

⁹also at: LIBPhys-UC, Departamento de Física, Universidade de Coimbra, P3004 516 Coimbra, Portugal

¹⁰also at: Walther-Meißner-Institut für Tieftemperaturforschung, D-85748 Garching, Germany

¹¹also at: Excellence Cluster Origins, D-85748 Garching, Germany

¹²also at: GSSI-Gran Sasso Science Institute, I-67100 L’Aquila, Italy

¹³also at: Dipartimento di Ingegneria Civile e Meccanica, Università degli Studi di Cassino e del Lazio Meridionale, I-03043 Cassino, Italy

Received: date / Accepted: date

Abstract CRESST is one of the most prominent direct detection experiments for dark matter particles with sub-GeV/c² mass. One of the advantages of the CRESST experiment is the possibility to include a large variety of nuclides in the target material used to probe dark matter interactions. In this work, we discuss in particular the interactions of dark matter particles with protons and neutrons of ${}^6\text{Li}$. This is now possible thanks to new calculations on nuclear matrix elements of this specific isotope of Li. To show the potential of using this particular nuclide for probing dark matter interactions, we used the data collected previously by a CRESST prototype based on LiAlO_2 and operated in an above ground test-facility at Max-Planck-Institut für Physik in Munich, Germany. In particular, the inclusion of ${}^6\text{Li}$ in the limit calculation drastically improves the result obtained for spin-dependent interactions with neutrons in the whole mass range. The improvement is significant, greater than two order of magnitude for dark matter masses below 1 GeV/c²,

compared to the limit previously published with the same data.

Keywords Dark Matter · Li-6 · Spin-Dependent · Lithium · Crystal

1 Introduction

CRESST is a direct detection dark matter experiment located in the underground laboratory of Gran Sasso (LNGS) [1]. CRESST operates cryogenic calorimeters consisting of a scintillating target crystal instrumented with a Transition Edge Sensor (TES) coupled to an auxiliary cryogenic calorimeter to measure the scintillation light, called Light Detector (LD). These detectors guarantee a low energy threshold along with a high energy resolution for particle interactions with the target crystal. Furthermore, the employment of scintillating crystals allows to achieve an effective particle discrimination through the simultaneous detection of the light and phonon signals. As a result of these remarkable features, CRESST is one of the leading experiments for light dark matter search.

*Corresponding authors

^abertoldo@mpp.mpg.de

^bmancuso@mpp.mpg.de

Light dark matter refers to dark matter particles with masses $\lesssim 1 \text{ GeV}/c^2$. There are various theoretical scenarios which could motivate particle candidates in this mass range, such as asymmetric dark matter [2–4], scalar dark matter [5], and hidden sector dark matter [6]. These scenarios together with the absence of positive signals from experiments specialized in the detection of dark matter particles with weak-scale coupling motivate the exploration of the low-mass parameter space.

The CRESST-III experiment reported its first results for spin-independent dark matter interactions with nuclei of detectors employing CaWO_4 target crystals [1]. In parallel to this effort, the CRESST collaboration has also investigated new suitable crystals for its detectors, with particular focus on lithium-containing crystals [7–9]. The motivation for the development of these novel cryogenic detectors stems from the ideal properties of lithium for the study of the spin-dependent interactions of sub- GeV/c^2 dark matter with its nuclei. As an evidence of this, those prototypes achieved competitive results studying spin-dependent interactions of dark matter particles with nuclei of ^7Li , the most abundant isotope of lithium.

In this work, we calculate a new limit on spin-dependent interactions of dark matter particles with ^6Li using the data collected with these prototypes [9]. This analysis is now possible thanks to a recent study [10] on the ^6Li nuclear ground state: there, the authors report the results obtained for the computation of proton and neutron spin matrix elements which we use in this work to calculate the expected rate of dark matter interactions with ^6Li .

2 Dark matter interactions with lithium

Lithium is currently the lightest element which can be employed in a CRESST-like detector. When considering elastic interactions of dark matter particles with nuclei, its light mass offers a clear advantage for the exploration of the sub- GeV/c^2 dark matter parameter space over heavier elements, due to the kinematics of such interactions. However, in the case of the classic spin-independent cross-section, the expected rate of interactions for dark matter particles with nuclei of lithium is significantly lower compared to heavier targets. This is due to the fact that the expected rate is proportional to A^2 [11], which significantly favors heavier nuclei. Overall, the employment of lithium could still offer a slight improvement over oxygen for CRESST in the study of spin-independent interactions of very low mass dark matter particles ($\lesssim 0.1 \text{ GeV}/c^2$), but the main advantage is the additional sensitivity to spin-dependent dark matter interactions.

In order to successfully probe spin-dependent interactions with a direct dark matter search experiment, the nuclides of the experimental target must hold a non-zero nuclear spin

($J_N \neq 0$) [12–14]. Furthermore, as opposed to the spin-independent standard scenario, which does not distinguish between the type of nucleons, spin-dependent interaction are usually studied for the two different cases of *proton-only* and *neutron-only* interactions. These *proton-only* and *neutron-only* interactions are proportional respectively to the square of the mean value of proton spin $\langle S_p \rangle^2$ and neutron spin $\langle S_n \rangle^2$, where $\langle S_p \rangle$ and $\langle S_n \rangle$ are the spin matrix elements for protons and neutrons of a given nucleus. In practical terms, it means that a suitable nuclide is usually particularly sensitive to only one of the two types of interactions, depending on its nuclear composition.

There is a small amount of stable nuclides, beside ^6Li , composed by an odd number of protons and an odd number of neutrons which also have the property of holding a non-zero nuclear spin. These nuclides can be used to study simultaneously spin-dependent dark matter interactions with the protons and the neutrons composing their nuclei with high sensitivity. A list of such nuclides with their natural abundance is given in Table 1: of these, only ^6Li and ^{10}B can straightforwardly be employed in a CRESST-like detector.

In the case of standard CRESST-III detectors [15] made by CaWO_4 , the isotopes which can be employed to study spin-dependent interactions are either with significantly low natural abundance (^{17}O and ^{43}Ca) or large mass (^{183}W). Despite these limitations, in a previous work we derived a leading limit on spin-dependent interactions of dark matter particles with neutrons of ^{17}O nuclei below $1.5 \text{ GeV}/c^2$ [1]. Thus, we found a strong motivation to find more suitable target crystals for the study of spin-dependent interactions with CRESST-like detectors.

There has been a general lack of interest for the calculation of spin-dependent matrix elements of these isotopes listed in Table 1, since, for historical reasons, the community has mostly focused on nuclides composed by an odd number of nucleons [16]. However, it is important to stress that ^6Li is arguably the best isotope which can be employed to study spin-dependent interactions of sub- GeV/c^2 dark matter particles with ordinary matter. In fact, ^6Li is light in mass, has a not negligible natural abundance, and, mean values for both proton and neutron spin operators extremely close to the maximum value as we can see in [10].

This recent study is providing the first precise calculation of $\langle S_p \rangle$ and $\langle S_n \rangle$, the spin matrix elements arising from the *proton-only* and *neutron-only* interactions, for an isotope of lithium. Unfortunately, up-to-date calculations for ^7Li are still missing and we will refer in this work to the ones presented in [16, 17].

A good target material compatible with cryogenic detectors is LiAlO_2 , which was already employed in CRESST-like detectors [9]. LiAlO_2 contains several isotopes with $J_N \neq 0$: both isotopes of lithium (^6Li , ^7Li), ^{27}Al , and, ^{17}O . The case study for ^7Li was already presented in detail in [7] and now

Nuclide	Natural Abundance (%)
^2H	0.01
^6Li	7.59
^{10}B	19.97
^{14}N	99.63

Table 1 Stable nuclides with odd-numbered neutrons and protons and non-zero nuclear spin.

we are going to present the new results obtained with the addition of ^6Li to the calculation.

3 Data

For this analysis we use the data collected in a previous work [9] with the prototype detector labelled *module B*. The final spectrum obtained in this work has an energy threshold for particle interactions of $E_T = (213.02 \pm 1.48)$ eV. The energy threshold has been calculated using the method presented in [18]. The energy calibration is implemented using the 5.895 keV peak from the ^{55}Fe source and heater pulses of four different known amplitudes to interpolate the energy calibration in the whole energy region of interest, which was chosen from E_T to 4 keV. A total of 22.2 hours of data without any radioactive source ("background data") were collected.

The exposure time after analysis cuts is 17.2 hours, which corresponds to a total exposure of $2.01 \cdot 10^{-3}$ kg·d, with an exposure for ^7Li of $1.95 \cdot 10^{-4}$ kg·d and an exposure for ^{27}Al of $8.22 \cdot 10^{-4}$ kg·d.

The data are publicly available at [9] (see ancillary files on arXiv).

4 Dark Matter results

To be fully consistent, the new results are calculated in the same fashion as the one presented in the previous work, with the only difference being the addition of the cross section for dark matter interactions with ^6Li alongside to the ones for ^7Li and ^{27}Al . As such, the exposure of ^6Li is equal to $1.60 \cdot 10^{-5}$ kg·d.

The limits are calculated using Yellin's optimal interval method [19, 20] and are shown in Figures 1 and 2. The calculation adopts the standard dark matter halo model, with a Maxwellian velocity distribution and a local dark matter density of $\rho_{\text{DM}} = 0.3 (\text{GeV}/c^2)/\text{cm}^3$ [21], a galactic escape velocity of $v_{\text{esc}} = 544 \text{ km/s}$ [22] and $v_{\odot} = 220 \text{ km/s}$ for the solar orbit velocity [23]. To calculate the *neutron-only* and *proton-only* exclusion limits for ^6Li , we adopted $\langle S_n \rangle = \langle S_p \rangle = 0.472$ [10].

As is possible to see, the addition of ^6Li slightly improves




Fig. 1 Exclusion limits set by various direct detection experiments for spin-dependent interactions of dark matter particles with protons. The result obtained from *module B* data with $^7\text{Li}+^{27}\text{Al}$ is shown with a dashed black line [9]. The new result obtained with $^6\text{Li}+^7\text{Li}+^{27}\text{Al}$ is shown with a solid red line. Additionally, limits from other experiments are also shown: CDMSlite with ^{73}Ge [24]; PICO with ^{19}F [25]; XENON1T (Migdal effect) with $^{129}\text{Xe}+^{131}\text{Xe}$ [26]; J. I. Collar with ^1H [27]. Finally, a constraint from Borexino data derived in [28] is shown in black.

the result obtained for *proton-only* interactions, due to a modest increase in the overall exposure. Instead, for *neutron-only* interactions we can observe a clear improvement (Figure 2), by at least one order of magnitude over the whole mass range with respect to the previous result [9]. In particular, there is a drastic improvement in the sub- GeV/c^2 dark matter mass region, by more than 2 orders of magnitude at $1.0 \text{ GeV}/c^2$ and over 3 orders at $0.4 \text{ GeV}/c^2$. This is exclusively due to the added sensitivity provided by the inclusion of ^6Li in the limit calculation.

For a dark matter particle mass of $1.0 \text{ GeV}/c^2$ the limit on the cross section reaches $7.59 \cdot 10^{-32} \text{ cm}^2$ for *proton-only* interactions and $9.23 \cdot 10^{-31} \text{ cm}^2$ for *neutron-only* interactions.

5 Conclusions

In this work we present a detailed analysis of the potential to probe spin-dependent dark matter interactions with ^6Li for both *proton-only* and *neutron-only* interactions. Since ^6Li is always present in nature alongside ^7Li , an ideal candidate to probe *proton-only* spin-dependent interactions, the major improvement offered by the presence of ^6Li is a clear enhancement in sensitivity for *neutron-only* interactions.

As such, we have shown a real case scenario with a CRESST-like detector employing a LiAlO_2 target crystal: the total improvement with the simple addition of ^6Li in the limit calculation is equal to more than two orders of magnitude for a dark matter particle mass of $1.0 \text{ GeV}/c^2$ compared to the




Fig. 2 Exclusion limits set by various direct detection experiments for spin-dependent interactions of dark matter particles with neutrons. The result obtained from *module B* data with ${}^7\text{Li}+{}^{27}\text{Al}$ is shown with a dashed black line [9]. The new result obtained with ${}^6\text{Li}+{}^7\text{Li}+{}^{27}\text{Al}$ is shown with a solid red line. The result obtained with ${}^{17}\text{O}$ in CRESST-III is shown with a dashed red line [1]. For comparison, limits from other experiments are also shown: EDELWEISS [29] and CDM-Slite [24] using ${}^{73}\text{Ge}$, LUX [30] and XENONIT (Migdal effect) [26] using ${}^{129}\text{Xe}+{}^{131}\text{Xe}$.

result previously obtained.

This result strengthens the case for lithium-containing crystals as the ideal target material for cryogenic detectors designed to probe sub-GeV/ c^2 spin-dependent dark matter interactions with ordinary matter.

Acknowledgements We thank Alex Gnech for the useful discussion which gave a fundamental contribution to this work.

This work is supported through the DFG by SFB1258 and the Origins Cluster, and by the BMBF05A17WO4.

References

1. CRESST, A. H. Abdelhameed *et al.*, Phys. Rev. **D100**, 102002 (2019), 1904.00498.
2. D. E. Kaplan, M. A. Luty, and K. M. Zurek, Phys. Rev. **D79**, 115016 (2009), 0901.4117.
3. K. Petraki and R. R. Volkas, Int. J. Mod. Phys. **A28**, 1330028 (2013), 1305.4939.
4. K. M. Zurek, Phys. Rept. **537**, 91 (2014), 1308.0338.
5. C. Boehm and P. Fayet, Nucl. Phys. **B683**, 219 (2004), hep-ph/0305261.
6. J. L. Feng and J. Kumar, Phys. Rev. Lett. **101**, 231301 (2008), 0803.4196.
7. CRESST, A. H. Abdelhameed *et al.*, Eur. Phys. J. **C79**, 630 (2019), 1902.07587.
8. CRESST, E. Bertoldo *et al.*, J. Low Temp. Phys. **199**, 510 (2019).
9. CRESST, A. H. Abdelhameed *et al.*, Eur. Phys. J. **C80**, 834 (2020), 2005.02692.
10. A. Gnech, M. Viviani, and L. E. Marcucci, Phys. Rev. **C102**, 014001 (2020), 2004.05814.
11. A. Kurylov and M. Kamionkowski, Phys. Rev. **D69**, 063503 (2004), hep-ph/0307185.

12. M. W. Goodman and E. Witten, Phys. Rev. **D31**, 3059 (1985).
13. J. R. Ellis and R. A. Flores, Phys. Lett. **B263**, 259 (1991).
14. J. Engel, S. Pittel, and P. Vogel, Int. J. Mod. Phys. **E1**, 1 (1992).
15. R. S. *et al.*, Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment **845**, 414 (2017), Proceedings of the Vienna Conference on Instrumentation 2016.
16. V. A. Bednyakov and F. Simkovic, Phys. Part. Nucl. **36**, 131 (2005), hep-ph/0406218, [Fiz. Elem. Chast. Atom. Yadra36,257(2005)].
17. A. F. Pacheco and D. Strottman, Phys. Rev. **D40**, 2131 (1989).
18. M. Mancuso *et al.*, Nuclear Instruments and Methods in Physics Research Section A **940**, 492 (2019), 1711.11459.
19. S. Yellin, Physical Review D **66**, 032005 (2002), 0203002.
20. S. Yellin, (2007), 0709.2701.
21. P. Salucci, F. Nesti, G. Gentile, and C. F. Martins, Astronomy and Astrophysics **523**, A83 (2010), 1003.3101.
22. M. C. Smith *et al.*, Monthly Notices of the Royal Astronomical Society **379**, 755 (2007), 0611671.
23. D. Kerr, F. J.; Lynden-Bell, Monthly Notices of the Royal Astronomical Society **221**, 1023 (1986).
24. SuperCDMS, R. Agnese *et al.*, Physical Review D **97**, 022002 (2018), 1707.01632.
25. PICO, C. Amole *et al.*, Physical Review D **100**, 022001 (2019), 1902.04031.
26. XENON, E. Aprile *et al.*, Physical Review Letters **123**, 241803 (2019), 1907.12771.
27. J. I. Collar, Physical Review D **98**, 023005 (2018), 1805.02646.
28. T. Bringmann and M. Pospelov, Physical Review Letters **122**, 171801 (2019), 1810.10543.
29. EDELWEISS, E. Armengaud *et al.*, Physical Review D **99**, 082003 (2019), 1901.03588.
30. LUX, D. S. Akerib *et al.*, Physical Review Letters **118**, 251302 (2017), 1705.03380.