OPERATOR-VALUED p-APPROXIMATE SCHAUDER FRAMES K. MAHESH KRISHNA

Statistics and Mathematics Unit Indian Statistical Institute, Bangalore Centre Bengaluru, Karnataka 560 059 India Email: kmaheshak@gmail.com

and

P. SAM JOHNSON

Department of Mathematical and Computational Sciences National Institute of Technology Karnataka, Surathkal Mangaluru 575 025, India Email: sam@nitk.edu.in Date: January 12, 2022

Abstract: We give an operator-algebraic treatment of theory of p-approximate Schuader frames which includes the theory of operator-valued frames by Kaftal, Larson, and Zhang [Trans. AMS., 2009], G-frames by Sun [JMAA, 2006], factorable weak operator-valued frames by Krishna and Johnson [Annals of FA, 2022] and p-approximate Schuader frames by Krishna and Johnson [J. Pseudo-Differ. Oper. Appl, 2021] as particular cases. We show that a sufficiently rich theory can be developed even for Banach spaces. We achieve this by defining various concepts and characterizations in Banach spaces. These include duality, approximate duality, equivalence, orthogonality and stability.

Keywords: Frame, operator-valued frame, p-approximate Schauder frame.

Mathematics Subject Classification (2020): 42C15, 47A05, 46B25.

Contents

1.	Introduction	1
2.	Operator-valued p-approximate Schauder frames and their characterizations	5
3.	Duality and orthogonality	11
4.	Approximate duality	15
5.	Equivalence	19
6.	Perturbations	21
7.	Operator-valued Feichtinger conjectures	23
8.	Acknowledgements	24
Re	eferences	24

1. INTRODUCTION

Both theoretically and practically successful theory of frames for Hilbert spaces which originated from the seminal work of Duffin and Schaeffer [27,34,37,54,85] demanded the development of frame theory for Banach spaces. It was in 1991, when Grochenig [48] introduced the notion of frames for Banach spaces, called as Banach frames, in connection with the theory of atomic decompositions [38–40]. In the sequel \mathcal{X}, \mathcal{Y} always denote Banach spaces. Dual of \mathcal{X} is denoted by \mathcal{X}^* . The identity operator on \mathcal{X} is denoted by $I_{\mathcal{X}}$. Field real or complex numbers is denoted by \mathbb{K} . The space of all bounded linear operators from \mathcal{X} to \mathcal{Y} is denoted by $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. We set $\mathcal{B}(\mathcal{X}) \coloneqq \mathcal{B}(\mathcal{X}, \mathcal{X})$.

Definition 1.1. [48] Let \mathcal{X} be a Banach space, \mathcal{X}_d be an associated BK-space, $S : \mathcal{X}_d \to \mathcal{X}$ be a bounded linear operator and $\{f_n\}_n$ be a collection in \mathcal{X}^* . The pair $(\{f_n\}_n, S)$ is said to be a **Banach frame** for \mathcal{X} if the following conditions hold.

- (i) $\{f_n(x)\}_n \in \mathcal{X}_d$, for each $x \in \mathcal{X}$.
- (ii) There are a, b > 0 such that

$$a\|x\| \le \|\{f_n(x)\}_n\| \le b\|x\|, \quad \forall x \in \mathcal{X}.$$

Constants a and b are called as Banach frame lower and upper bounds, respectively.

(iii) $S{f_n(x)}_n = x$, for each $x \in \mathcal{X}$.

Definition 1.2. [48] Let \mathcal{X} be a Banach space, \mathcal{X}_d be an associated BK-space, $\{\tau_n\}_n$ be a collection in \mathcal{X} and $\{f_n\}_n$ be a collection in \mathcal{X}^* . The pair $(\{f_n\}_n, \{\tau_n\}_n)$ is said to be an **atomic decomposition** for \mathcal{X} if the following conditions hold.

- (i) $\{f_n(x)\}_n \in \mathcal{X}_d$, for each $x \in \mathcal{X}$.
- (ii) There are a, b > 0 such that

$$a||x|| \le ||\{f_n(x)\}_n|| \le b||x||, \quad \forall x \in \mathcal{X}.$$

Constants a and b are called as lower and upper atomic bounds, respectively.

(iii) $\sum_{n=1}^{\infty} f_n(x)\tau_n = x$, for each $x \in \mathcal{X}$.

Hilbert space frame theory mainly concentrates around representing every element of the space using frames as a series but this is missing in the definition of Banach frames. After a decade from the work of Grochenig, it is Aldroubi, Sun, Tang, Han, Larson, Casazza, Christensen, and Stoeva [2, 10, 15, 30] analyzed when it is possible to write every element as a series by using a sequence in the space or in the dual space. In [10], it was particularly analyzed the series representation only by using first two conditions in Definition 1.1. Moreover, using the approximation properties of Banach space [11], it was proved by Casazza and Christensen [12] that there are Banach spaces \mathcal{X} such that there is no collection $\{f_n\}_n$ in \mathcal{X}^* and no collection in $\{\tau_n\}_n$ in \mathcal{X} satisfying

$$x = \sum_{n=1}^{\infty} f_n(x)\tau_n, \quad \forall x \in \mathcal{X}.$$

Motivated from the dilation theory of frames [32,50,58], in 1999, Casazza, Han, and Larson [15] introduced three other notions of frames for Banach spaces called as framing or unconditional framing, projective frame and modeled frame. It is then proved in [15] that all these three notions are equivalent. Around the same time, Terekhin also introduced another notion of frames for Banach spaces, again using dilation viewpoint [79–81].

Important paper of Daubechies and DeVore [33] led Casazza, Dilworth, Odell, Schlumprecht, and Zsak [9] to weaken the unconditional convergence condition in the definition of framing and they defined Schauder frames for Banach spaces as follows.

Definition 1.3. [9] Let $\{\tau_n\}_n$ be a collection in \mathcal{X} and $\{f_n\}_n$ be a collection in \mathcal{X}^* . The pair $(\{f_n\}_n, \{\tau_n\}_n)$ is said to be a Schauder frame for \mathcal{X} if

$$x = \sum_{n=1}^{\infty} f_n(x)\tau_n, \quad \forall x \in \mathcal{X}.$$

Definition 1.3 has been extended by Thomas, Freeman, Odell, Schlumprecht, and Zsak in 2014 [44,82].

Definition 1.4. [44, 82] Let $\{\tau_n\}_n$ be a collection in \mathcal{X} and $\{f_n\}_n$ be a collection in \mathcal{X}^* . The pair $(\{f_n\}_n, \{\tau_n\}_n)$ is said to be an approximate Schauder frame (ASF) for \mathcal{X} if

$$S_{f,\tau}: \mathcal{X} \ni x \mapsto S_{f,\tau}x \coloneqq \sum_{n=1}^{\infty} f_n(x)\tau_n \in \mathcal{X}$$

is a well-defined bounded linear, invertible operator.

Again looking back to the theory of frames for Hilbert spaces, following generalizations have been proposed.

- (i) frames for subspaces/fusion frames [18,19].
- (ii) outer frames [1].
- (iii) oblique frames [24].
- (iv) pseudo frames [69].
- (v) quasi-projectors [43].

In 2009, Kaftal, Larson, Zhang [57] introduced the notion of operator-valued frames which unified all the notions mentioned earlier.

Definition 1.5. [57] Let $\mathcal{H}, \mathcal{H}_0$ be Hilbert spaces. A collection $\{A_n\}_n$ in $\mathcal{B}(\mathcal{H}, \mathcal{H}_0)$ is said to be an operator-valued frame in $\mathcal{B}(\mathcal{H}, \mathcal{H}_0)$ if the series

$$S_A \coloneqq \sum_{n=1}^{\infty} A_n^* A_n$$

converges in the strong-operator topology on $\mathcal{B}(\mathcal{H})$ to a bounded invertible operator.

In 2006, Sun [78] introduced G-frames which is equivalent to the notion of operator-valued frames.

Definition 1.6. [78] Let $\mathcal{H}, \mathcal{H}_n$ be Hilbert spaces and let $A_n \in \mathcal{B}(\mathcal{H}, \mathcal{H}_n)$, for each $n \in \mathbb{N}$. The collection $\{A_n\}_n$ is said to be a **G-frame** if there are a, b > 0 such that

$$a\|h\|^{2} \leq \sum_{n=1}^{\infty} \|A_{n}h\|^{2} \leq b\|h\|^{2}, \quad \forall h \in \mathcal{H}.$$

It is noticed by the authors of this paper that there are some other notions of frames even for Hilbert spaces, which will not include in the theory of operator-valued frames. These notions are as follow.

- (i) Framings for Hilbert spaces [15].
- (ii) Schauder frames and approximate Schauder frames for Hilbert spaces [9,44,82].
- (iii) Atomic decompositions for Hilbert spaces [15].
- (iv) cb-frames for Hilbert spaces [73].
- (v) Signed frames for Hilbert spaces [75].
- (vi) Pair frames for Hilbert space [42].
- (vii) Controlled frames for Hilbert spaces [8].
- (viii) K-frames for Hilbert spaces [45]

4

(ix) Semi-frames for Hilbert spaces [4].

To include most of the notions of frames for Hilbert spaces, the notion of weak operator-valued frames for Hilbert spaces has been introduced in [66]. Unfortunately, the frame operator in this case may not split and to get a reasonable theory like that of Hilbert space frames, a proper subclass of weak operator-valued frames called as factorable weak operator-valued frames has been introduced and is defined as follows.

Definition 1.7. [66] Given $n \in \mathbb{N}$, define

$$L_n: \mathcal{H}_0 \ni h \mapsto L_n h \coloneqq e_n \otimes h \in \ell^2(\mathbb{N}) \otimes \mathcal{H}_0$$

where $\{e_n\}_n$ is the standard orthonormal basis for $\ell^2(\mathbb{N})$. Let $\{A_n\}_n$ and $\{\Psi_n\}_n$ be collections in $\mathcal{B}(\mathcal{H},\mathcal{H}_0)$. The pair $(\{A_n\}_n, \{\Psi_n\}_n)$ is said to be a **factorable weak operator-valued frame** in $\mathcal{B}(\mathcal{H},\mathcal{H}_0)$ if the following conditions hold.

(i) The series

$$S_{A,\Psi} \coloneqq \sum_{n=1}^{\infty} \Psi_n^* A_n$$

converges in the strong-operator topology on $\mathcal{B}(\mathcal{H})$ to a bounded invertible operator.

(ii) The map

$$\theta_A : \mathcal{H} \ni h \mapsto \theta_A h \coloneqq \sum_{n=1}^{\infty} L_n A_n h \in \ell^2(\mathbb{N}) \otimes \mathcal{H}_0$$

is a well-defined bounded linear operator.

(iii) The map

$$\theta_A: \mathcal{H} \ni h \mapsto \theta_A h \coloneqq \sum_{n=1}^{\infty} L_n \Psi_n h \in \ell^2(\mathbb{N}) \otimes \mathcal{H}_0$$

is a well-defined bounded linear operator.

Earlier, we listed various notions of frames for Hilbert spaces. There are corresponding notions for Banach spaces.

- (i) Framings for Banach spaces [15].
- (ii) Schauder frames and approximate Schauder frames for Banach spaces [9,44,82].
- (iii) Atomic decompositions for Banach spaces [15].
- (iv) cb-frames for operator spaces [73].
- (v) p-frames for Banach spaces [2, 30]
- (vi) p-approximate Schauder frames for Banach spaces [64].
- (vii) pseudo-Schauder frames for Banach spaces [71].
- (viii) K-frames for Banach spaces [46].
- (ix) Hilbert-Schauder frames for Banach spaces [72].

Most of the results in frame theory for Hilbert spaces arise by doing analysis in the standard separable Hilbert space $\ell^2(\mathbb{N})$ and switching between a given Hilbert space and $\ell^2(\mathbb{N})$. Therefore a natural space to work for Banach spaces is the classical sequence space $\ell^p(\mathbb{N})$ for $p \in [1, \infty)$. This is the main motivation for the introduction of p-approximate Schauder frames for Banach spaces. **Definition 1.8.** [64] An ASF $(\{f_n\}_n, \{\tau_n\}_n)$ for \mathcal{X} is said to be a *p*-approximate Schauder frame (written as *p*-ASF), $p \in [1, \infty)$ if both the maps

$$\theta_f : \mathcal{X} \ni x \mapsto \theta_f x \coloneqq \{f_n(x)\}_n \in \ell^p(\mathbb{N}) \text{ and}$$
$$\theta_\tau : \ell^p(\mathbb{N}) \ni \{a_n\}_n \mapsto \theta_\tau \{a_n\}_n \coloneqq \sum_{n=1}^\infty a_n \tau_n \in \mathcal{X}$$

are well-defined bounded linear operators.

In this paper, we give operator-algebraic version of both factorable weak operator-valued frames and p-approximate Schauder frames in the setting of Banach spaces. In Section 2 we define the notion of operator-valued p-approximate Schauder frames and then show that our notion includes factorable weak operator-valued frames, operator-valued frames/G-frames and p-approximate Schauder frames. We also introduce the notion of operator-valued p-Riesz bases. Through an explicit construction we show that influential Naimark-Han-Larson dilation result can be proved even for the operator-version. We then show that a result based on the result of Holub for Hilbert spaces can be derived for Banach spaces which characterizes operator-valued p-approximate Schauder frames. In Section 3 we study two important notions associated with a given frame, namely, duality and orthogonality. We characterize dual frames and use orthogonality to generate new frames. In Section 4, we study the notion of approximate duality and characterize them. Section 5 describes equivalence of frames with an explicit description of operator giving similarity. Section 6 derives perturbation results. Section 7 states some conjectures.

2. Operator-valued p-approximate Schauder frames and their characterizations

Let \mathcal{X} , \mathcal{Y} be Banach spaces and $p \in [1, \infty)$. By $\ell^p(\mathbb{N}) \otimes \mathcal{Y}$, we mean the Banach space obtained by tensoring $\ell^p(\mathbb{N})$ and \mathcal{Y} w.r.t. any cross norm (see [35, 36, 76] for the tensor product of Banach spaces). In the development of paper [57], certain collection of operators played important role. Here we define corresponding operators for Banach spaces. Given $n \in \mathbb{N}$, we define

$$L_n: \mathcal{Y} \ni x \mapsto L_n x \coloneqq e_n \otimes x \in \ell^p(\mathbb{N}) \otimes \mathcal{Y},$$

where $\{e_n\}_n$ is the standard Schauder basis for $\ell^p(\mathbb{N})$. Since the norm is a cross norm, it then follows that L_n 's are isometries from \mathcal{Y} to $\ell^p(\mathbb{N}) \otimes \mathcal{Y}$. We also define Γ_n as the (unique) tensor product of bounded linear operators

$$\ell^p(\mathbb{N}) \ni \{a_m\}_m \mapsto a_n \in \mathbb{K}, \quad \mathcal{Y} \ni y \mapsto y \in \mathcal{Y}.$$

Since $\mathbb{K} \otimes \mathcal{Y}$ is isometrically isomorphic to \mathcal{Y} , we write codomain of Γ_n as \mathcal{Y} . We also note that $a_n \otimes y = a_n y$. Hence

$$\Gamma_n(\{a_m\}_m \otimes y) = a_n y, \quad \forall \{a_m\}_m \in \ell^p(\mathbb{N}), \forall y \in \mathcal{Y}, \forall n \in \mathbb{N}.$$

Therefore, for $n, m \in \mathbb{N}$ we have

(1)
$$\Gamma_n L_m = \begin{cases} I_{\mathcal{Y}} & \text{if } n = m \\ 0 & \text{if } n \neq m \end{cases} \text{ and } \sum_{n=1}^{\infty} L_n \Gamma_n z = z, \quad \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}.$$

We now introduce the notion of operator-valued p-approximate Schauder frames and demonstrate that it includes all known notions of frames for which frame operator factors. **Definition 2.1.** Let $\{A_n\}_n$ be collection in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ and $\{\Psi_n\}_n$ be a collection in $\mathcal{B}(\mathcal{Y}, \mathcal{X})$. The pair $(\{A_n\}_n, \{\Psi_n\}_n)$ is said to be an **operator-valued p-approximate Schauder frame** (we write **operator-valued p-ASF**) in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ if the following conditions hold.

(i) The frame operator

$$S_{A,\Psi}: \mathcal{X} \ni x \mapsto \sum_{n=1}^{\infty} \Psi_n A_n x \in \mathcal{X}$$

is a well-defined bounded linear invertible operator. Constants a, b > 0 satisfying

$$a||x|| \le \left\|\sum_{n=1}^{\infty} \Psi_n A_n x\right\| \le b||x||, \quad \forall x \in \mathcal{X}$$

are called as lower and upper frame bounds, respectively.

(ii) The analysis operator

$$\theta_A: \mathcal{X} \ni x \mapsto \theta_A x \coloneqq \sum_{n=1}^{\infty} L_n A_n x \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}$$

is a well-defined bounded linear operator. Constant c > 0 satisfying

$$\left\|\sum_{n=1}^{\infty} L_n A_n x\right\| \le c \|x\|, \quad \forall x \in \mathcal{X}$$

is called as analysis bound.

(iii) The synthesis operator

$$\theta_{\Psi}: \ell^p(\mathbb{N}) \otimes \mathcal{Y} \ni z \mapsto \theta_{\Psi} z \coloneqq \sum_{n=1}^{\infty} \Psi_n \Gamma_n z \in \mathcal{X}$$

is a well-defined bounded linear operator. Constant d > 0 satisfying

$$\left\|\sum_{n=1}^{\infty} \Psi_n \Gamma_n z\right\| \le c \|z\|, \quad \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}$$

is called as synthesis bound.

If $S_{A,\Psi} = I_{\mathcal{X}}$, then the frame is called as a **Parseval operator-valued p-ASF**. If we do not demand the invertibility of $S_{A,\Psi}$, then we say that $(\{A_n\}_n, \{\Psi_n\}_n)$ is an **operator-valued p-approximate Bessel sequence** (we write **operator-valued p-ABS**).

Example 2.2. (i) Let $\mathcal{H}, \mathcal{H}_0$ be Hilbert spaces and $(\{A_n\}_n, \{\Psi_n\}_n)$ be a factorable weak-OVF in $\mathcal{B}(\mathcal{H}, \mathcal{H}_0)$. Define

$$B_n \coloneqq A_n, \ \Phi_n \coloneqq \Psi_n^*, \quad \forall n \in \mathbb{N}.$$

Then $(\{B_n\}_n, \{\Phi_n\}_n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{H}, \mathcal{H}_0)$. Note that $\Gamma_n = L_n^*, \forall n \in \mathbb{N}$. (ii) Let \mathcal{X} be a Banach space and $(\{f_n\}_n, \{\tau_n\}_n)$ be a p-ASF for \mathcal{X} . Define

$$A_n \coloneqq f_n, \ \Psi_n : \mathbb{K} \ni \alpha \mapsto \Psi_n \alpha \coloneqq \alpha \tau_n \in \mathcal{X}, \quad \forall n \in \mathbb{N}.$$

Then

$$\sum_{n=1}^{\infty} \Psi_n A_n x = \sum_{n=1}^{\infty} \Psi_n f_n(x) = \sum_{n=1}^{\infty} f_n(x) \tau_n, \quad \forall x \in \mathcal{X}.$$

Hence $(\{A_n\}_n, \{\Psi_n\}_n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathbb{K})$.

(iii) Let \mathcal{X} , \mathcal{Y} be Banach spaces and $U : \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ and $V : \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ be bounded linear operators such that VU is bounded invertible. Define

$$A_n = \Gamma_n U, \quad \Psi_n = V L_n, \quad \forall n \in \mathbb{N}.$$

Then

$$\sum_{n=1}^{\infty} \Psi_n A_n x = \sum_{n=1}^{\infty} V L_n \Gamma_n U x = V \left(\sum_{n=1}^{\infty} L_n \Gamma_n \right) U x = V U x, \quad \forall x \in \mathcal{X}.$$

Hence $(\{A_n\}_n, \{\Psi_n\}_n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$.

(iv) This example is motivated from the Cuntz algebra [31]. Let \mathcal{X} be a Banach space, $n \in \mathbb{N}$ and $A_1, \ldots, A_n, \Psi_1, \ldots, \Psi_n : \mathcal{X} \to \mathcal{X}$ be bounded linear operators such that $\sum_{j=1}^n \Psi_n A_n = I_{\mathcal{X}}$. Then $(\{A_j\}_{j=1}^n, \{\Psi_j\}_{j=1}^n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X})$.

It is clear that an operator-valued p-ABS need not be an operator-valued p-ASF. Therefore the next attempt is to see that whether we can convert an operator-valued p-ABS to operator-valued p-ASF. In the case of Hilbert spaces, it was Li and Sun [67] who showed it is possible to expand every G-Bessel sequence to a G-frame just by adding one element (for a proof in finite dimensional Hilbert spaces, see [20]). Surprisingly, using a result of Casazza and Christensen [12], it is proved in [65] that for Banach spaces one may not able to convert Bessel sequences to frames. Here is a partial answer for operator-valued p-ASFs.

Theorem 2.3. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ABS in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. If there are $B \in \mathcal{B}(\mathcal{X}, \mathcal{Y})$ and $\Phi \in \mathcal{B}(\mathcal{Y}, \mathcal{X})$ such that $I_{\mathcal{X}} - S_{A,\Psi} = \Phi B$, then

$$(\{A_n\}_n \cup \{B\}, \{\Psi_n\}_n \cup \{\Phi\})$$

is a Parseval operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. In particular, if $\mathcal{Y} = \mathcal{X}$ and $I_{\mathcal{X}} - S_{A,\Psi}$ is a square, then every operator-valued p-ABS can be turned into an operator-valued p-ASF.

Proof. We only need to observe that

$$\sum_{n=1}^{\infty} \Psi_n A_n x + \Phi B x = S_{A,\Psi} x + (I_{\mathcal{X}} - S_{A,\Psi}) x = x, \quad \forall x \in \mathcal{X}.$$

In general, it is well known in Hilbert space frame theory that it is difficult to invert frame operator and get an expansion of every element of the Hilbert space using frame. Thus one needs an algorithm to approximate every element using frames but without using invertibility of frame operator. First such an algorithm was given by Duffin and Schaeffer [37] and later two more algorithms were given by Grochenig [47]. We now derive an algorithm for operator-valued p-ASFs under a condition.

Proposition 2.4. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ with bounds a and b. For $x \in \mathcal{X}$, define

$$x_0 \coloneqq 0, \ x_n \coloneqq x_{n-1} + \frac{2}{a+b} S_{A,\Psi}(x - x_{n-1}), \quad \forall n \ge 1.$$

If $||I_{\mathcal{X}} - \frac{2}{b+a}S_{A,\Psi}|| \le \frac{b-a}{b+a}$, then

$$||x_n - x|| \le \left(\frac{b-a}{b+a}\right)^n ||x||, \quad \forall n \ge 1.$$

In particular, $x_n \to x$ as $n \to \infty$.

Proof. We observe

$$\begin{aligned} x - x_n &= x - x_{n-1} - \frac{2}{a+b} S_{A,\Psi}(x - x_{n-1}) \\ &= \left(I_{\mathcal{X}} - \frac{2}{b+a} S_{A,\Psi} \right) (x - x_{n-1}) \\ &= \dots = \left(I_{\mathcal{X}} - \frac{2}{b+a} S_{A,\Psi} \right)^n x, \quad \forall x \in \mathcal{X}, \forall n \ge 1. \end{aligned}$$

Hence

$$||x_n - x|| \le \left||I_{\mathcal{X}} - \frac{2}{b+a}S_{A,\Psi}||^n ||x|| \le \left(\frac{b-a}{b+a}\right)^n ||x||, \quad \forall n \ge 1.$$

Most important result in the development of theory is the following result.

Theorem 2.5. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$.

- (i) The analysis operator $\theta_A : \mathcal{X} \ni x \mapsto \theta_A x = \sum_{n=1}^{\infty} L_n A_n x \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ is injective. (ii) The synthesis operator $\theta_{\Psi} : \ell^p(\mathbb{N}) \otimes \mathcal{Y} \ni z \mapsto \sum_{n=1}^{\infty} \Psi_n \Gamma_n z \in \mathcal{X}$ surjective. (iii) $(\{A_n S_{A,\Psi}^{-1}\}_n, \{S_{A,\Psi}^{-1}\Psi_n\}_n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$.

- (iv) Frame operator factors as $S_{A,\Psi} = \theta_{\Psi} \theta_A$.
- (v) $P_{A,\Psi} \coloneqq \theta_A S_{A,\Psi}^{-1} \theta_{\Psi} : \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y} \text{ is a projection onto } \theta_A(\mathcal{X}).$

Proof. Since $S_{A,\Psi}$ is invertible, we get (i), (ii) and (iii). For (iv), we use Equation (1) and get

$$\theta_{\Psi}\theta_{A}x = \left(\sum_{n=1}^{\infty} \Psi_{n}\Gamma_{n}\right) \left(\sum_{k=1}^{\infty} L_{k}A_{k}x\right) = \sum_{n=1}^{\infty} \Psi_{n} \left(\sum_{k=1}^{\infty} \Gamma_{n}L_{k}A_{k}x\right)$$
$$= \sum_{n=1}^{\infty} \Psi_{n}A_{n}x = S_{A,\Psi}x, \quad \forall x \in \mathcal{X}.$$

For (v),

$$P_{A,\Psi}^2 = \theta_A S_{A,\Psi}^{-1} \theta_\Psi \theta_A S_{A,\Psi}^{-1} \theta_\Psi = \theta_A S_{A,\Psi}^{-1} I_\mathcal{X} \theta_\Psi = P_{A,\Psi}.$$

In the next result we are proving that projections give frames to subspaces.

Theorem 2.6. Let \mathcal{Z} be a closed complementable subspace of \mathcal{X} and $P: \mathcal{X} \to \mathcal{Z}$ be an onto projection.

- (i) If $(\{A_n\}_n, \{\Psi_n\}_n)$ is a Parseval operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$, then $(\{A_nP\}_n, \{P\Psi_n\}_n)$ is a Parseval operator-valued p-ASF in $\mathcal{B}(\mathcal{Z}, \mathcal{Y})$.
- (ii) If $(\{A_n\}_n, \{\Psi_n\}_n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$, then $(\{A_nP\}_n, \{P\Psi_n\}_n)$ is an operatorvalued p-ASF in $\mathcal{B}(\mathcal{Z}, \mathcal{Y})$.
- *Proof.* (i) We have to show that the frame operator for $(\{A_nP\}_n, \{P\Psi_n\}_n)$ is the identity on \mathcal{Z} . Let $z \in \mathcal{Z}$. Then

$$\sum_{n=1}^{\infty} P\Psi_n A_n P z = P\left(\sum_{n=1}^{\infty} \Psi_n A_n z\right) = P z = z.$$

(ii) We observe that the frame operator $S_{AP,P\Psi}$ is $PS_{A,\Psi}P$. Since $S_{A,\Psi}$ is invertible on \mathcal{X} , $PS_{A,\Psi}P$ is invertible on \mathcal{Z} with inverse $PS_{A,\Psi}^{-1}P$.

Definition 2.7. (i) An operator-valued p-ASF $(\{A_n\}_n, \{\Psi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is said to be an operatorvalued *p*-approximate Riesz basis (we write operator-valued *p*-ARB) if $P_{A,\Psi} = I_{\ell^p(\mathbb{N})} \otimes I_{\mathcal{Y}}$.

(ii) A collection $(\{A_n\}_n, \{\Psi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is said to be an operator-valued *p*-approximate Riesz sequence if it is an operator-valued *p*-ARB in $\mathcal{B}(\mathcal{Z}, \mathcal{Y})$, where

$$\mathcal{Z} \coloneqq \overline{\operatorname{span}} \cup_{n=1}^{\infty} \Psi_n(\mathcal{Y}).$$

Riesz bases play important role in the dilation theory of frames. First dilation result of frames was obtained independently by Han and Larson [50] and Kashin and Kulikova [58] (see [32] for the history of this theorem). The result was also known to Daubechies [17]. Dilation of framings for Banach spaces were obtained in [15]. This work later led to the notion of general dilation theory of operator-valued measures in [51, 52]. For p-ASFs, dilation theorem is derived in [62]. For OVFs, general Naimark-Han-Larson dilation theorem was obtained in [53]. Here we obtain dilation result for operator-valued p-ASFs. This is basically a kind of converse to Theorem 2.6.

Theorem 2.8. (Dilation theorem) Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. Then there exist a Banach space \mathcal{X}_1 which contains \mathcal{X} isometrically and bounded linear operators $B_n : \mathcal{X}_1 \to \mathcal{Y}$, $\Phi_n : \mathcal{Y} \to \mathcal{X}_1$, $\forall n$ such that $(\{B_n\}_n, \{\Phi_n\}_n)$ is an operator-valued p-ARB in $\mathcal{B}(\mathcal{X}_1, \mathcal{Y})$ and $B_n|_{\mathcal{X}} = A_n, \forall n \in \mathbb{N}$.

Proof. Since $I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi}$ is idempotent, $(I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})(\mathcal{X})$ is a Banach space. Define

$$\mathcal{X}_1 \coloneqq \mathcal{X} \oplus (I_{\ell^p(\mathbb{N}) \otimes \mathcal{Y}} - P_{A,\Psi})(\mathcal{X}).$$

Then the map $\mathcal{X} \ni x \mapsto x \oplus 0 \in \mathcal{X}_1$ is an isometry. Now define

$$B_n: \mathcal{X}_1 \ni x \oplus y \mapsto A_n x + \Gamma_n (I_{\ell^p(\mathbb{N}) \otimes \mathcal{Y}} - P_{A, \Psi}) y \in \mathcal{Y},$$

$$\Phi_n: \mathcal{Y} \ni y \mapsto \Psi_n y \oplus (I_{\ell^p(\mathbb{N}) \otimes \mathcal{Y}} - P_{A, \Psi}) L_n y \in \mathcal{X}_1, \quad \forall n \in \mathbb{N}.$$

Then we have $B_n|_{\mathcal{X}} = A_n, \forall n \in \mathbb{N}$. We now find

$$\theta_B(x \oplus y) = \sum_{n=1}^{\infty} L_n A_n x + \sum_{n=1}^{\infty} L_n \Gamma_n (I_{\ell^p(\mathbb{N}) \otimes \mathcal{Y}} - P_{A, \Psi}) y$$
$$= \theta_A x + (I_{\ell^p(\mathbb{N}) \otimes \mathcal{Y}} - P_{A, \Psi}) y, \quad \forall x \oplus y \in \mathcal{X}_1.$$

and

$$\theta_{\Phi} z = \sum_{n=1}^{\infty} \Psi_n \Gamma_n z \oplus \sum_{n=1}^{\infty} (I_{\ell^p(\mathbb{N}) \otimes \mathcal{Y}} - P_{A,\Psi}) L_n \Gamma_n z$$
$$= \theta_{\Psi} z \oplus (I_{\ell^p(\mathbb{N}) \otimes \mathcal{Y}} - P_{A,\Psi}) z, \quad \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}.$$

We also notice

$$(I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})\theta_A = \theta_A - \theta_A S_{A,\Psi}^{-1}\theta_\Psi \theta_A = \theta_A - \theta_A = 0$$

10

and

$$\theta_{\Psi}(I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}}-P_{A,\Psi})=\theta_{\Psi}-\theta_{\Psi}\theta_{A}S_{A,\Psi}^{-1}\theta_{\Psi}=\theta_{\Psi}-\theta_{\Psi}=0$$

Using these expressions we get the frame operator of $(\{B_n\}_n, \{\Phi_n\}_n)$ as

$$\begin{split} S_{B,\Phi}(x\oplus y) &= \theta_{\Phi}\theta_{B}(x\oplus y) = \theta_{\Phi}(\theta_{A}x + (I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})y) \\ &= \theta_{\Psi}(\theta_{A}x + (I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})y) \oplus (I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})(\theta_{A}x + (I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})y) \\ &= (\theta_{\Psi}\theta_{A}x + \theta_{\Psi}(I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})y) \oplus ((I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})\theta_{A}x + (I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})^{2}y) \\ &= (S_{A,\Psi}x + 0) \oplus (0 + (I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})y) \\ &= (S_{A,\Psi} \oplus (I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi}))(x\oplus y), \quad \forall x\oplus y \in \mathcal{X}_{1}. \end{split}$$

Therefore $S_{B,\Phi}$ invertible with inverse $S_{A,\Psi}^{-1} \oplus (I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})$. Hence $(\{B_n\}_n, \{\Phi_n\}_n)$ is an operatorvalued p-ASF. Next we need to show that it is Riesz. Let $z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}$. Then

$$P_{B,\Phi}z = \theta_B S_{B,\Phi}^{-1} \Phi_{\Phi}z = \theta_B S_{B,\Phi}^{-1} (\theta_{\Psi}z \oplus (I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})z)$$

$$= \theta_B (S_{A,\Psi}^{-1} \oplus (I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi}))(\theta_{\Psi}z \oplus (I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})z)$$

$$= \theta_B (S_{A,\Psi}^{-1} \theta_{\Psi}z \oplus (I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})z)$$

$$= \theta_A S_{A,\Psi}^{-1} \theta_{\Psi}z + (I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})^2 z$$

$$= P_{A,\Psi}z + (I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - P_{A,\Psi})z = z.$$

It is possible to characterize frames for Hilbert spaces using the standard orthonormal basis $\{e_n\}_n$ for $\ell^2(\mathbb{N})$ [56]. This result has been extended in [64] which shows that p-ASFs can be characterized using the standard Schauder basis $\{e_n\}_n$ for $\ell^p(\mathbb{N})$. We further generalize this result for operator-valued p-ASFs.

Theorem 2.9. A pair $(\{A_n\}_n, \{\Psi_n\}_n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ if and only if

$$A_n = \Gamma_n U, \quad \Psi_n = V L_n, \quad \forall n \in \mathbb{N},$$

where $U: \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ and $V: \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ are bounded linear operators such that VU is bounded invertible.

Proof. (\Leftarrow) Given $x \in \mathcal{X}$, Equation (1) gives

$$S_{A,\Psi}x = \sum_{n=1}^{\infty} \Psi_n A_n x = \sum_{n=1}^{\infty} V L_n \Gamma_n U x = V \left(\sum_{n=1}^{\infty} L_n \Gamma_n\right) U x = V U x.$$

Therefore $S_{A,\Psi}$ is bounded invertible which says $(\{A_n\}_n, \{\Psi_n\}_n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. (\Rightarrow) Define $Ux \coloneqq \sum_{n=1}^{\infty} L_n A_n x, \forall x \in \mathcal{X}, Vz \coloneqq \sum_{n=1}^{\infty} \Psi_n \Gamma_n z, \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}.$ Then

$$\Gamma_n Ux = \Gamma_n \left(\sum_{k=1}^{\infty} L_k A_k x \right) = \sum_{k=1}^{\infty} \Gamma_n L_k A_k x = A_n x, \quad \forall x \in \mathcal{X}, \forall n \in \mathbb{N}$$

and

$$VL_n z = \sum_{k=1}^{\infty} \Psi_k \Gamma_k L_n z = \Psi_n z, \quad \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}, \forall n \in \mathbb{N}.$$

We are left with showing that VU is invertible. For this we find

$$VUx = \left(\sum_{n=1}^{\infty} \Psi_n \Gamma_n\right) \left(\sum_{k=1}^{\infty} L_k A_k x\right) = \sum_{n=1}^{\infty} \Psi_n A_n x = S_{A,\Psi} x, \quad \forall x \in \mathcal{X}$$

which says VU is invertible.

Corollary 2.10. A pair $(\{A_n\}_n, \{\Psi_n\}_n)$ is an operator-valued p-ABS in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ if and only if

$$A_n = \Gamma_n U, \quad \Psi_n = V L_n, \quad \forall n \in \mathbb{N},$$

where $U: \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ and $V: \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ are bounded linear operators.

Corollary 2.11. A pair $(\{A_n\}_n, \{\Psi_n\}_n)$ is an operator-valued p-ARB in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ if and only if

$$A_n = \Gamma_n U, \quad \Psi_n = V L_n, \quad \forall n \in \mathbb{N}$$

where $U: \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ and $V: \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ are bounded linear operators such that VU is bounded invertible and $U(VU)^{-1}V = I_{\ell^p(\mathbb{N}) \otimes \mathcal{Y}}$.

3. DUALITY AND ORTHOGONALITY

Given an operator-valued p-ASF $(\{A_n\}_n, \{\Psi_n\}_n)$, consider the frame $(\{B_n \coloneqq A_n S_{A,\Psi}^{-1}\}_n, \{\Phi_n \coloneqq S_{A,\Psi}^{-1}\Psi_n\}_n)$ in (iii) of Theorem 2.5. This frame has the property that $\theta_{\Psi}\theta_B = \theta_{\Phi}\theta_A = I_{\mathcal{X}}$. In general, there are other frames satisfying this property. This brings us to the notion of dual frames.

Definition 3.1. An operator-valued p-ASF $(\{B_n\}_n, \{\Phi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is said to be a **dual** for an operator-valued p-ASF $(\{A_n\}_n, \{\Psi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ if

$$\theta_{\Psi}\theta_B = \theta_{\Phi}\theta_A = I_{\mathcal{X}}.$$

If both $(\{B_n\}_n, \{\Phi_n\}_n)$ and $(\{A_n\}_n, \{\Psi_n\}_n)$ are operator-valued p-ABS, then we say that they are dual operator-valued p-ABSs.

By using the definition of analysis and synthesis operators we will have the following result.

Proposition 3.2. An operator-valued p-ASF $(\{B_n\}_n, \{\Phi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is a dual for an operator-valued p-ASF $(\{A_n\}_n, \{\Psi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ if and only if

$$\sum_{n=1}^{\infty} \Psi_n B_n x = \sum_{n=1}^{\infty} \Phi_n A_n x = x, \quad \forall x \in \mathcal{X}$$

The frame $(\{A_n S_{A,\Psi}^{-1}\}_n, \{S_{A,\Psi}^{-1}\Psi_n\}_n)$ is called as the canonical dual for $(\{A_n\}_n, \{\Psi_n\}_n)$. They have the following simple property whose proof follows from the calculation of its frame operator and we leave the proof.

Theorem 3.3. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ with frame bounds a and b. Then

- (i) The canonical dual for the canonical dual for $(\{A_n\}_n, \{\Psi_n\}_n)$ is itself.
- (ii) $\frac{1}{b}, \frac{1}{a}$ are frame bounds for the canonical dual of $(\{A_n\}_n, \{\Psi_n\}_n)$.
- (iii) If a, b are optimal frame bounds for $(\{A_n\}_n, \{\Psi_n\}_n)$, then $\frac{1}{b}, \frac{1}{a}$ are optimal frame bounds for its canonical dual.

Complete description of dual frames for Hilbert spaces was described Li [68]. This description has been generalized in the context of factorable weak OVFs in [66] and p-ASFs in [64]. Here we describe duals which cover all of them.

Lemma 3.4. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. Then an operator-valued p-ASF $(\{B_n\}_n, \{\Phi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is a dual for $(\{A_n\}_n, \{\Psi_n\}_n)$ if and only if

$$B_n = \Gamma_n U, \quad \Phi_n = V L_n, \quad \forall n \in \mathbb{N}$$

where $U : \mathcal{H} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ is a bounded right-inverse of θ_{Ψ} , $V : \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ is a bounded left-inverse of θ_A such that VU is bounded invertible.

Proof. (\Rightarrow) Let $(\{B_n\}_n, \{\Phi_n\}_n)$ be a dual operator-valued p-ASF for $(\{A_n\}_n, \{\Psi_n\}_n)$. Then $\theta_{\Psi}\theta_B = I_{\mathcal{X}} = \theta_{\Phi}\theta_A$. Define $U \coloneqq \theta_B, V \coloneqq \theta_{\Phi}$. Then $U : \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ is a right-inverse of $\theta_{\Psi}, V : \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ is a left-inverse of θ_A such that $VU = \theta_{\Phi}\theta_B = S_{B,\Phi}$ is invertible. We find

$$\Gamma_n U x = \Gamma_n \left(\sum_{k=1}^{\infty} L_k B_k x \right) = B_n x, \quad \forall x \in \mathcal{X},$$
$$V L_n z = \sum_{k=1}^{\infty} \Phi_k \Gamma_k L_n z = \Phi_n z, \quad \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}, \forall n \in \mathbb{N}.$$

(\Leftarrow) It is clear that $(\{B_n\}_n, \{\Phi_n\}_n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ and a calculation gives $\theta_B = U, \ \theta_\Phi = V.$ For duality, $\theta_\Phi \theta_A = V \theta_A = I_{\mathcal{X}}, \ \theta_\Psi \theta_B = \theta_\Psi U = I_{\mathcal{X}}.$

Lemma 3.5. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. Then

(i) $R: \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ is a bounded right-inverse of θ_{Ψ} if and only if

$$R = \theta_A S_{A,\Psi}^{-1} + (I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - \theta_A S_{A,\Psi}^{-1}\theta_\Psi)U,$$

where $U: \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ is a bounded linear operator.

(ii) $L: \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ is a bounded left-inverse of θ_A if and only if

$$L = S_{A,\Psi}^{-1} \theta_{\Psi} + V(I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - \theta_{A}S_{A,\Psi}^{-1}\theta_{\Psi}),$$

where $V : \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ is a bounded linear operator.

Proof. (i) (\Leftarrow) Let $U : \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ be a bounded linear operator. Then $\theta_{\Psi}(\theta_A S_{A,\Psi}^{-1} + (I_{\ell^p(\mathbb{N}) \otimes \mathcal{Y}} - \theta_A S_{A,\Psi}^{-1} \theta_{\Psi})U) = I_{\mathcal{X}} + \theta_{\Psi}U - \theta_{\Psi}U = I_{\mathcal{X}}$. Therefore $R \coloneqq \theta_A S_{A,\Psi}^{-1} + (I_{\ell^p(\mathbb{N}) \otimes \mathcal{Y}} - \theta_A S_{A,\Psi}^{-1} \theta_{\Psi})U$ is a bounded right-inverse of θ_{Ψ} .

 $(\Rightarrow) \text{ Let } R: \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y} \text{ be a bounded right-inverse of } \theta_{\Psi}. \text{ Define } U \coloneqq R. \text{ Then } \theta_A S_{A,\Psi}^{-1} + (I_{\ell^p(\mathbb{N}) \otimes \mathcal{Y}} - \theta_A S_{A,\Psi}^{-1} \theta_{\Psi})R = \theta_A S_{A,\Psi}^{-1} + R - \theta_A S_{A,\Psi}^{-1} = R.$

(ii) (\Leftarrow) Let $V : \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ be a bounded linear operator. Then $(S_{A,\Psi}^{-1}\theta_{\Psi} + V(I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - \theta_A S_{A,\Psi}^{-1}\theta_{\Psi}))\theta_A = I_{\mathcal{X}} + V\theta_A - V\theta_A I_{\mathcal{X}} = I_{\mathcal{X}}$. Therefore $V \coloneqq S_{A,\Psi}^{-1}\theta_{\Psi} + V(I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - \theta_A S_{A,\Psi}^{-1}\theta_{\Psi})$ is a bounded left-inverse of θ_A .

$$(\Rightarrow) \text{ Let } L: \ell^{p}(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X} \text{ be a bounded left-inverse of } \theta_{A}. \text{ Define } V \coloneqq L. \text{ Then } S_{A,\Psi}^{-1}\theta_{\Psi} + V(I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - \theta_{A}S_{A,\Psi}^{-1}\theta_{\Psi}) = S_{A,\Psi}^{-1}\theta_{\Psi} + L - I_{\mathcal{X}}S_{A,\Psi}^{-1}\theta_{\Psi} = L.$$

Theorem 3.6. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. Then an operator-valued p-ASF $(\{B_n\}_n, \{\Phi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is a dual for $(\{A_n\}_n, \{\Psi_n\}_n)$ if and only if

$$B_n = A_n S_{A,\Psi}^{-1} + \Gamma_n U - A_n S_{A,\Psi}^{-1} \theta_{\Psi} U,$$

$$\Phi_n = S_{A,\Psi}^{-1} \Psi_n + V L_n - V \theta_A S_{A,\Psi}^{-1} \Psi_n, \quad \forall n \in \mathbb{N}$$

such that the operator

$$S_{A,\Psi}^{-1} + VU - V\theta_A S_{A,\Psi}^{-1} \theta_\Psi U$$

is bounded invertible, where $U : \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ and $V : \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ are bounded linear operators. *Proof.* Lemmas 3.4 and 3.5 give the characterization of dual operator-valued p-ASF for $(\{A_n\}_n, \{\Psi_n\}_n)$ as

$$B_n = \Gamma_n(\theta_A S_{A,\Psi}^{-1} + (I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - \theta_A S_{A,\Psi}^{-1}\theta_\Psi)U) = A_n S_{A,\Psi}^{-1} + \Gamma_n U - A_n S_{A,\Psi}^{-1}\theta_\Psi U,$$

$$\Phi_n = (S_{A,\Psi}^{-1}\theta_\Psi + V(I_{\ell^p(\mathbb{N})\otimes\mathcal{Y}} - \theta_A S_{A,\Psi}^{-1}\theta_\Psi))L_n = S_{A,\Psi}^{-1}\Psi_n + VL_n - V\theta_A S_{A,\Psi}^{-1}\Psi_n, \quad \forall n \in \mathbb{N}$$

such that the operator

$$(S_{A,\Psi}^{-1}\theta_{\Psi} + V(I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - \theta_{A}S_{A,\Psi}^{-1}\theta_{\Psi}))(\theta_{A}S_{A,\Psi}^{-1} + (I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - \theta_{A}S_{A,\Psi}^{-1}\theta_{\Psi})U)$$

is bounded invertible, where $U : \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ and $V : \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ are bounded linear operators. We make an expansion and get

$$(S_{A,\Psi}^{-1}\theta_{\Psi} + V(I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - \theta_{A}S_{A,\Psi}^{-1}\theta_{\Psi}))(\theta_{A}S_{A,\Psi}^{-1} + (I_{\ell^{p}(\mathbb{N})\otimes\mathcal{Y}} - \theta_{A}S_{A,\Psi}^{-1}\theta_{\Psi})U)$$

= $S_{A,\Psi}^{-1} + VU - V\theta_{A}S_{A,\Psi}^{-1}\theta_{\Psi}U.$

Notion which is opposite to the notion of duality is the notion of orthogonality of frames which is very useful in the construction of new frames from two frames. Balan [7] introduced this notion and later studied by Han and Larson [50]. A detailed study in the finite dimensional case can be found in [49]. Orthogonality will be defined as follows based in Definition 3.1.

Definition 3.7. An operator-valued p-ASF $(\{B_n\}_n, \{\Phi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is said to be **orthogonal** to an operator-valued p-ASF $(\{A_n\}_n, \{\Psi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ if

$$\theta_{\Psi}\theta_B = \theta_{\Phi}\theta_A = 0$$

Similar to Proposition 3.2, we have the following result for orthogonality.

Proposition 3.8. An operator-valued p-ASF $(\{B_n\}_n, \{\Phi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is orthogonal to an operatorvalued p-ASF $(\{A_n\}_n, \{\Psi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ if and only if

$$\sum_{n=1}^{\infty} \Psi_n B_n x = \sum_{n=1}^{\infty} \Phi_n A_n x = 0, \quad \forall x \in \mathcal{X}.$$

As mentioned earlier, we can get new frames using orthogonal frames. Following two results will describe them.

Proposition 3.9. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ be two Parseval operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ which are orthogonal. If $C, D, E, F \in \mathcal{B}(\mathcal{X})$ are such that $EC + FD = I_{\mathcal{X}}$, then

$$(\{A_nC+B_nD\}_n, \{E\Psi_n+F\Phi_n\}_n)$$

is a Parseval operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. In particular, if scalars c, d, e, f satisfy ce + df = 1, then $(\{cA_n + dB_n\}_n, \{e\Psi_n + f\Phi_n\}_n)$ is a Parseval operator-valued p-ASF.

Proof. For each $x \in \mathcal{X}$,

$$S_{AC+BD,E\Psi+F\Phi}x = \sum_{n=1}^{\infty} (E\Psi_n + F\Phi_n)(A_nC + B_nD)x$$

= $ES_{A,\Psi}Cx + E\left(\sum_{n=1}^{\infty}\Psi_nB_n\right)Dx + F\left(\sum_{n=1}^{\infty}\Phi_nA_n\right)Cx + FS_{B,\Phi}Dx$
= $ECx + 0 + 0 + FDx = x.$

Proposition 3.10. If $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ are orthogonal operator-valued *p*-ASFs in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$, then $(\{A_n \oplus B_n\}_n, \{\Psi_n \oplus \Phi_n\}_n)$ is an operator-valued *p*-ASF in $\mathcal{B}(\mathcal{X} \oplus \mathcal{X}, \mathcal{Y})$. Further, if both $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ are Parseval, then $(\{A_n \oplus B_n\}_n, \{\Psi_n \oplus \Phi_n\}_n)$ is Parseval.

Proof. Let $x \oplus x_1 \in \mathcal{X} \oplus \mathcal{X}$. Then

$$S_{A\oplus B,\Psi\oplus\Phi}(x\oplus x_1) = \sum_{n=1}^{\infty} (\Psi_n \oplus \Phi_n)(A_n \oplus B_n)(x\oplus x_1) = \sum_{n=1}^{\infty} (\Psi_n \oplus \Phi_n)(A_nx + B_nx_1)$$
$$= \sum_{n=1}^{\infty} (\Psi_n(A_nx + B_nx_1) \oplus \Phi_n(A_nx + B_nx_1))$$
$$= \left(\sum_{n=1}^{\infty} \Psi_nA_nx + \sum_{n=1}^{\infty} \Psi_nB_nx_1\right) \oplus \left(\sum_{n=1}^{\infty} \Phi_nA_nx + \sum_{n=1}^{\infty} \Phi_nB_nx_1\right)$$
$$= (S_{A,\Psi}x + 0) \oplus (0 + S_{B,\Phi}x_1) = (S_{A,\Psi} \oplus S_{B,\Phi})(x \oplus x_1).$$

In the case of Hilbert spaces, it is known that orthogonal operator-valued frames have a common dual operator-valued frame. However, for Banach spaces, we have the following weaker result.

Proposition 3.11. Two orthogonal operator-valued p-ASFs have a common dual operator-valued p-ABS.

Proof. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ be orthogonal operator-valued p-ASFs in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. Define

$$C_n \coloneqq A_n S_{A,\Psi}^{-1} + B_n S_{B,\Phi}^{-1}, \quad \Xi_n \coloneqq S_{A,\Psi}^{-1} \Psi_n + S_{B,\Phi}^{-1} \Phi_n \quad \forall n \in \mathbb{N}.$$

Clearly $(\{C_n\}_n, \{\Xi_n\}_n)$ is an operator-valued p-ABS in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. We need to show that this is a common dual for $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$. Let $x \in \mathcal{X}$. Then

$$\sum_{n=1}^{\infty} \Xi_n A_n x = \sum_{n=1}^{\infty} (S_{A,\Psi}^{-1} \Psi_n + S_{B,\Phi}^{-1} \Phi_n) A_n x = x + 0 = x,$$

$$\sum_{n=1}^{\infty} \Psi_n C_n x = \sum_{n=1}^{\infty} \Psi_n (A_n S_{A,\Psi}^{-1} + B_n S_{B,\Phi}^{-1}) x = x + 0 = x,$$

$$\sum_{n=1}^{\infty} \Xi_n B_n x = \sum_{n=1}^{\infty} (S_{A,\Psi}^{-1} \Psi_n + S_{B,\Phi}^{-1} \Phi_n) B_n x = 0 + x = x,$$

$$\sum_{n=1}^{\infty} \Phi_n C_n x = \sum_{n=1}^{\infty} \Phi_n (A_n S_{A,\Psi}^{-1} + B_n S_{B,\Phi}^{-1}) x = 0 + x = x.$$

We end this section by showing that there is no need of any conditions on two operator-valued p-ASFs to make the tensor product as an operator-valued p-ASFs. For frames for Hilbert spaces, this result was proved by Feichtinger and Grochenig [41].

Proposition 3.12. If $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ are operator-valued p-ASFs in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$, then $(\{A_n \otimes B_m\}_{n,m}, \{\Psi_n \otimes \Phi_m\}_{n,m})$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X} \otimes \mathcal{X}, \mathcal{Y})$. Further, if both $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ are Parseval, then $(\{A_n \otimes B_m\}_{n,m}, \{\Psi_n \otimes \Phi_m\}_{n,m})$ is Parseval.

Proof. We need to show that the frame operator for the tensor product is invertible. We first find it at elementary tensors as follows.

$$S_{A\otimes B,\Psi\otimes\Phi}(x\otimes x_1) = \sum_{n,m=1}^{\infty} (\Psi_n \otimes \Phi_m)(A_n \otimes B_m)(x\otimes x_1)$$

=
$$\sum_{n,m=1}^{\infty} (\Psi_n A_n x \otimes \Phi_m B_m x_1) = \left(\sum_{n=1}^{\infty} \Psi_n A_n x\right) \otimes \left(\sum_{m=1}^{\infty} \Phi_m B_m x_1\right)$$

=
$$S_{A,\Psi} x \otimes S_{B,\Phi} x_1 = (S_{A,\Psi} \otimes S_{B,\Phi})(x \otimes x_1), \quad \forall x, x_1 \in \mathcal{X}.$$

We therefore have $S_{A \otimes B, \Psi \otimes \Phi} = S_{A, \Psi} \otimes S_{B, \Phi}$ which completes the proof.

4. Approximate duality

Even though all duals for an operator-valued p-ASF are characterized completely in the previous section, even in the case of Hilbert spaces, it is known that given a collection, it is difficult to verify whether it is a dual of a frame. This led Christensen and Laugesen [29] to introduce the notion of approximate duals (also see [70]). This notion has then been introduced and characterized for Banach spaces in [61]. Here we give operator-valued version of those results for Banach spaces.

Definition 4.1. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ be operator-valued *p*-ABSs in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. We say that they are **approximately dual p-ABSs** if

$$||I_{\mathcal{X}} - \theta_{\Phi}\theta_A|| < 1$$
 and $||I_{\mathcal{X}} - \theta_{\Psi}\theta_B|| < 1$.

In addition, if $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ are p-ASFs in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$, then we say that they are approximately dual p-ASFs.

Definition 4.1 gives the approximate duality using analysis and synthesis operators but the following proposition says that expansion of elements using approximate duals is close to the expansion using duals as described in Proposition 3.2.

Proposition 4.2. If $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ are approximately dual operator-valued *p*-ABSs in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$, then

$$\left\|x - \sum_{n=1}^{\infty} \Phi_n A_n x\right\| < \|x\| \quad and \quad \left\|x - \sum_{n=1}^{\infty} \Psi_n B_n x\right\| < \|x\|, \quad \forall x \in \mathcal{X} \setminus \{0\}.$$

Proof. Given $x \in \mathcal{X} \setminus \{0\}$,

$$\left\| x - \sum_{n=1}^{\infty} \Phi_n A_n x \right\| = \| I_{\mathcal{X}} x - \theta_{\Phi} \theta_A x \| \le \| I_{\mathcal{X}} - \theta_{\Phi} \theta_A \| \| x \| < \| x \|,$$
$$\left\| x - \sum_{n=1}^{\infty} \Psi_n B_n x \right\| = \| I_{\mathcal{X}} x - \theta_{\Psi} \theta_B x \| \le \| I_{\mathcal{X}} - \theta_{\Psi} \theta_B \| \| x \| < \| x \|.$$

One of the most important properties associated with approximate duals is that they generate duals.

Proposition 4.3. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ be approximately dual operator-valued *p*-ABSs in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. Then $(\{B_nS_{B,\Psi}^{-1}\}_n, \{S_{A,\Phi}^{-1}\Phi_n\}_n)$ is a dual operator-valued *p*-ABS for $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{A_nS_{A,\Phi}^{-1}\}_n, \{S_{B,\Psi}^{-1}\Psi_n\}_n)$ is a dual operator-valued *p*-ABS for $(\{B_n\}_n, \{\Phi_n\}_n)$.

Proof. Let $x \in \mathcal{X}$. Then

$$\sum_{n=1}^{\infty} \Psi_n B_n S_{B,\Psi}^{-1} x = S_{B,\Psi} S_{B,\Psi}^{-1} x = x, \quad \sum_{n=1}^{\infty} S_{A,\Phi}^{-1} \Phi_n A_n x = S_{A,\Phi}^{-1} S_{A,\Phi} x = x,$$
$$\sum_{n=1}^{\infty} \Phi_n A_n S_{A,\Phi}^{-1} x = S_{A,\Phi} S_{A,\Phi}^{-1} x = x, \quad \sum_{n=1}^{\infty} S_{B,\Psi}^{-1} \Psi_n B_n x = S_{B,\Psi}^{-1} S_{B,\Psi} x = x.$$

Motivated from a characterization of duals, we try to get an operator-theoretic characterization of approximate duality.

Theorem 4.4. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ABS in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. An operator-valued p-ABS $(\{B_n\}_n, \{\Phi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is an approximately dual operator-valued p-ABS for $(\{A_n\}_n, \{\Psi_n\}_n)$ if and only if there exist bounded linear operators $U, V : \mathcal{X} \to \mathcal{X}$ satisfying $||I_{\mathcal{X}} - U|| < 1$ and $||I_{\mathcal{X}} - V|| < 1$ such that $(\{C_n \coloneqq B_n U^{-1}\}_n, \{\Xi_n \coloneqq V^{-1}\Phi_n\}_n)$ is a dual for $(\{A_n\}_n, \{\Psi_n\}_n)$. Statement holds even if p-ABS is replaced by p-ASF.

Proof. (\Rightarrow) Define $U \coloneqq \theta_{\Psi} \theta_B$ and $V \coloneqq \theta_{\Phi} \theta_A$. We then have $||I_{\chi} - U|| < 1$ and $||I_{\chi} - V|| < 1$ which say that U and V are invertible, and hence

$$\sum_{n=1}^{\infty} \Psi_n C_n x = \sum_{n=1}^{\infty} \Psi_n B_n U^{-1} x = \theta_{\Psi} \theta_B U^{-1} x = U U^{-1} x = x,$$

$$\sum_{n=1}^{\infty} \Xi_n A_n x = \sum_{n=1}^{\infty} V^{-1} \Phi_n A_n x = V^{-1} \theta_{\Phi} \theta_A x = V^{-1} V x = x, \quad \forall x \in \mathcal{X}.$$

Hence $(\{C_n\}_n, \{\Xi_n\}_n)$ is a dual for $(\{A_n\}_n, \{\Psi_n\}_n)$.

(\Leftarrow) We see that $\theta_C = \theta_B U^{-1}$ and $\theta_{\Xi} = V^{-1} \theta_{\Phi}$. Since $(\{C_n\}_n, \{\Xi_n\}_n)$ is a dual for $(\{A_n\}_n, \{\Psi_n\}_n)$, we get that

$$\|I_{\mathcal{X}} - \theta_{\Phi}\theta_A\| = \|I_{\mathcal{X}} - V\theta_{\Xi}\theta_A\| = \|I_{\mathcal{X}} - V\| < 1,$$

$$\|I_{\mathcal{X}} - \theta_{\Psi}\theta_B\| = \|I_{\mathcal{X}} - \theta_{\Psi}\theta_C U\| = \|I_{\mathcal{X}} - U\| < 1.$$

Hence $(\{B_n\}_n, \{\Phi_n\}_n)$ is an approximately dual operator-valued p-ABS for $(\{A_n\}_n, \{\Psi_n\}_n)$.

Theorem 4.4 combined with Theorem 3.6 gives the following result.

Theorem 4.5. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. An operator-valued p-ASF $(\{B_n\}_n, \{\Phi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is an approximately dual operator-valued p-ASF for $(\{A_n\}_n, \{\Psi_n\}_n)$ if and only if there exist bounded linear operators $U, V : \mathcal{X} \to \mathcal{X}, A : \mathcal{X} \to \ell^p(\mathbb{N}), B : \ell^p(\mathbb{N}) \to \mathcal{X}$ satisfying $\|I_{\mathcal{X}} - U\| < 1, \|I_{\mathcal{X}} - V\| < 1,$

$$\begin{split} B_n &= A_n S_{A,\Psi}^{-1} U + \Gamma_n A U - A_n S_{A,\Psi}^{-1} \theta_{\Psi} A U, \\ \Phi_n &= V S_{A,\Psi}^{-1} \Psi_n + V B L_n - V B \theta_A S_{A,\Psi}^{-1} \Psi_n, \quad \forall n \in \mathbb{N} \end{split}$$

such that the operator

$$S_{A,\Psi}^{-1} + BA - B\theta_A S_{A,\Psi}^{-1} \theta_\Psi A$$

is bounded invertible, where $A: \mathcal{X} \to \ell^p(\mathbb{N}) \otimes \mathcal{Y}$ and $B: \ell^p(\mathbb{N}) \otimes \mathcal{Y} \to \mathcal{X}$ are bounded linear operators.

In [29], Christensen and Laugesen gave a method to construct approximately duals iteratively. In [61], this result was derived for Banach spaces. Here we have a similar result for operator-valued p-ABSs.

Theorem 4.6. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ABS in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ be an operator-valued p-ABS in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ which is an approximately dual for $(\{A_n\}_n, \{\Psi_n\}_n)$.

(i) The dual operator-valued p-ABS $(\{B_n S_{B,\Psi}^{-1}\}_n, \{S_{A,\Phi}^{-1}\Phi_n\}_n)$ for $(\{A_n\}_n, \{\Psi_n\}_n)$ can be written as

$$B_n S_{B,\Psi}^{-1} = B_n + \sum_{m=1}^{\infty} B_n (I_{\mathcal{X}} - S_{B,\Psi})^m,$$

$$S_{A,\Phi}^{-1} \Phi_n = \Phi_n + \sum_{m=1}^{\infty} (I_{\mathcal{X}} - S_{A,\Phi})^m \Phi_n, \quad \forall n \in \mathbb{N}.$$

(ii) For $N \in \mathbb{N}$, define

$$C_n^{(N)} \coloneqq B_n + \sum_{m=1}^N B_n (I_{\mathcal{X}} - S_{B,\Psi})^m,$$

$$\Xi_n^{(N)} \coloneqq \Phi_n + \sum_{m=1}^N (I_{\mathcal{X}} - S_{A,\Phi})^m \Phi_n, \quad \forall n \in \mathbb{N}.$$

Then $(\{C_n^{(N)}\}_n, \{\Xi_n^{(N)}\}_n)$ is an operator-valued p-ABS and is an approximately dual for $(\{A_n\}_n, \{\Psi_n\}_n)$, for each $N \in \mathbb{N}$. Moreover,

$$\begin{aligned} \|I_{\mathcal{X}} - \theta_{\Xi}^{(N)} \theta_A\| &\leq \|I_{\mathcal{X}} - S_{A,\Phi}\|^{N+1} \to 0 \quad \text{as } N \to \infty, \\ \|I_{\mathcal{X}} - \theta_{\Psi} \theta_C^{(N)}\| &\leq \|I_{\mathcal{X}} - S_{B,\Psi}\|^{N+1} \to 0 \quad \text{as } N \to \infty. \end{aligned}$$

Proof. (i) We get these from the Neumann series expansion

$$S_{B,\Psi}^{-1} = \sum_{m=0}^{\infty} (I_{\mathcal{X}} - S_{B,\Psi})^m, \quad S_{A,\Phi}^{-1} = \sum_{m=0}^{\infty} (I_{\mathcal{X}} - S_{A,\Phi})^m.$$

(ii) Clearly $(\{C_n^{(N)}\}_n, \{\Xi_n^{(N)}\}_n)$ is an operator-valued p-ABS. Now consider

$$\begin{aligned} \theta_{\Xi}^{(N)} \theta_{A} x &= \sum_{n=1}^{\infty} \Xi_{n}^{(N)} A_{n} x = \sum_{n=1}^{\infty} \sum_{m=0}^{N} (I_{\mathcal{X}} - S_{A,\Phi})^{m} \Phi_{n} A_{n} x \\ &= \sum_{m=0}^{N} \sum_{n=1}^{\infty} (I_{\mathcal{X}} - S_{A,\Phi})^{m} \Phi_{n} A_{n} x = \sum_{m=0}^{N} (I_{\mathcal{X}} - S_{A,\Phi})^{m} \sum_{n=1}^{\infty} \Phi_{n} A_{n} x \\ &= \sum_{m=0}^{N} (I_{\mathcal{X}} - S_{A,\Phi})^{m} S_{A,\Phi} x = \sum_{m=0}^{N} (I_{\mathcal{X}} - S_{A,\Phi})^{m} (I_{\mathcal{X}} - (I_{\mathcal{X}} - S_{A,\Phi})) x \\ &= x - (I_{\mathcal{X}} - S_{A,\Phi})^{N+1} x, \quad \forall x \in \mathcal{X} \\ &\implies \|I_{\mathcal{X}} - \theta_{\Xi}^{(N)} \theta_{A}\| \leq \|I_{\mathcal{X}} - S_{A,\Phi}\|^{N+1} \end{aligned}$$

and

$$\begin{aligned} \theta_{\Psi} \theta_{C}^{(N)} x &= \sum_{n=1}^{\infty} \Psi_{n} C_{n}^{(N)} x = \sum_{n=1}^{\infty} \sum_{m=0}^{N} \Psi_{n} B_{n} (I_{\mathcal{X}} - S_{B,\Psi})^{m} x \\ &= \sum_{m=0}^{N} \sum_{n=1}^{\infty} \Psi_{n} B_{n} (I_{\mathcal{X}} - S_{B,\Psi})^{m} x = \sum_{m=0}^{N} S_{B,\Psi} (I_{\mathcal{X}} - S_{B,\Psi})^{m} x \\ &= \sum_{m=0}^{N} (I_{\mathcal{X}} - (I_{\mathcal{X}} - S_{B,\Psi})) (I_{\mathcal{X}} - S_{B,\Psi})^{m} x = x - (I_{\mathcal{X}} - S_{B,\Psi})^{N+1} x, \quad \forall x \in \mathcal{X} \\ &\implies \|I_{\mathcal{X}} - \theta_{\Psi} \theta_{C}^{(N)}\| \leq \|I_{\mathcal{X}} - S_{B,\Psi}\|^{N+1}. \end{aligned}$$

Since $||I_{\mathcal{X}} - S_{A,\Phi}|| < 1$ and $||I_{\mathcal{X}} - S_{B,\Psi}|| < 1$, $(\{C_n^{(N)}\}_n, \{\Xi_n^{(N)}\}_n)$ is an approximately dual for $(\{A_n\}_n, \{\Psi_n\}_n)$.

In [29], it is also showed by Christensen and Laugesen that by perturbing frame we can get approximate duals. In [61], it is showed that this result is valid for Banach spaces. Here is the operator-valued version of that result.

Theorem 4.7. Let $\{A_n\}_n$ be a collection in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ and $\{\Psi_n\}_n$ be a collection in $\mathcal{B}(\mathcal{Y}, \mathcal{X})$. Let $(\{C_n\}_n, \{\Xi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ such that for some R > 0, Q > 0

(2)
$$\left\|\sum_{n=1}^{\infty} L_n (A_n - C_n) x\right\| \le R \|x\|, \quad \forall x \in \mathcal{X}$$

and

(3)
$$\left\|\sum_{n=1}^{\infty} (\Xi_n - \Psi_n) \Gamma_n z\right\| \le Q \|z\|, \quad \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}$$

Let $(\{B_n\}_n, \{\Phi_n\}_n)$ be a dual frame for $(\{C_n\}_n, \{\Xi_n\}_n)$ with analysis bound c and synthesis bound d. If dR < 1, cQ < 1, then $(\{B_n\}_n, \{\Phi_n\}_n)$ is an approximately dual for $(\{A_n\}_n, \{\Psi_n\}_n)$.

Proof. Inequalities (2) and (3) say that $(\{A_n\}_n, \{\Psi_n\}_n)$ is an operator-valued p-ABS in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. Now Inequality (2) can be written as $\|\theta_A x - \theta_C x\| \leq R \|x\|$, $\forall x \in \mathcal{X}$. Similarly, Inequality (3) can be written as $\|\theta_{\Xi} z - \theta_{\Psi} z\| \leq Q \|z\|, \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}$. Therefore

$$\|I_{\mathcal{X}} - \theta_{\Phi}\theta_A\| = \|\theta_{\Phi}\theta_C - \theta_{\Phi}\theta_A\| \le \|\theta_{\Phi}\| \|\theta_C - \theta_A\| \le dR < 1,$$

and

$$\|I_{\mathcal{X}} - \theta_{\Psi}\theta_B\| = \|\theta_{\Xi}\theta_B - \theta_{\Psi}\theta_B\| \le \|\theta_{\Xi} - \theta_{\Psi}\|\|\theta_B\| \le cQ < 1.$$

5. Equivalence

Given two frames for Hilbert space, one tries to get one from another. It is also clear that action of every element of frame by a fixed invertible bounded linear operator again gives a frame. This led Balan to introduce the notion of similarity or equivalence for frames for Hilbert spaces [6]. This notion has been generalized in [57] for operator-valued frames, in [66] for factorable weak-OVFs and in [64] for p-ASFs. Here we define the same notion for operator-valued p-ASFs.

Definition 5.1. An operator-valued p-ASF $(\{B_n\}_n, \{\Phi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is said to be similar or equivalent to an operator-valued p-ASF $(\{A_n\}_n, \{\Psi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ if there exist invertible operators $R_{A,B}, L_{\Psi,\Phi} \in \mathcal{B}(\mathcal{X})$ such that

$$B_n = A_n R_{A,B}, \quad \Phi_n = L_{\Psi,\Phi} \Psi_n, \quad \forall n \in \mathbb{N}.$$

We first observe that for every operator-valued p-ASF $(\{A_n\}_n, \{\Psi_n\}_n)$, each of operator-valued p-ASF $(\{A_nS_{A,\Psi}^{-1}\}_n, \{\Psi_n\}_n)$ and $(\{A_n\}_n, \{S_{A,\Psi}^{-1}\Psi_n\}_n)$ is a Parseval operator-valued p-ASF which is similar to $(\{A_n\}_n, \{\Psi_n\}_n)$. Like in the case of Hilbert spaces, since $R_{A,B}, R_{\Psi,\Phi}$ are bounded invertible, the relation "similarity" is an equivalence relation on the set

 $\{(\{A_n\}_n, \{\Psi_n\}_n) : (\{A_n\}_n, \{\Psi_n\}_n) \text{ is an operator-valued p-ASF in } \mathcal{B}(\mathcal{X}, \mathcal{Y})\}.$

Next lemma shows that we can easily get analysis, synthesis and frame operators of similar frames from one another.

Lemma 5.2. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ be similar operator-valued p-ASFs in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ and $B_n = A_n R_{A,B}$, $\Phi_n = L_{\Psi,\Phi} \Psi_n$, $\forall n \in \mathbb{N}$, for some invertible $R_{A,B}$, $L_{\Psi,\Phi} \in \mathcal{B}(\mathcal{X})$. Then

- (i) $\theta_B = \theta_A R_{A,B}, \theta_\Phi = L_{\Psi,\Phi} \theta_\Psi.$
- (ii) $S_{B,\Phi} = L_{\Psi,\Phi} S_{A,\Psi} R_{A,B}$.
- (iii) $P_{B,\Phi} = P_{A,\Psi}$.

Proof. Let $x \in \mathcal{X}$ and $z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}$. Then $\theta_B x = \sum_{n=1}^{\infty} L_n B_n x = \sum_{n=1}^{\infty} L_n A_n R_{A,B} x = \theta_A R_{A,B} x$ and $\theta_{\Phi} z = \sum_{n=1}^{\infty} \Phi_n \Gamma_n z = \sum_{n=1}^{\infty} L_{\Psi,\Phi} \Psi_n \Gamma_n z = L_{\Psi,\Phi} \theta_{\Psi} z$. Therefore

$$S_{B,\Phi} = \theta_{\Phi}\theta_{B} = L_{\Psi,\Phi}\theta_{\Psi}\theta_{A}R_{A,B} = L_{\Psi,\Phi}S_{A,\Psi}R_{A,B}$$

and

$$P_{B,\Phi} = \theta_B S_{B,\Phi}^{-1} \theta_\Phi = (\theta_A R_{A,B}) (L_{\Psi,\Phi} S_{A,\Psi} R_{A,B})^{-1} (L_{\Psi,\Phi} \theta_\Psi) = P_{A,\Psi}.$$

In the next result we characterize similarity using operators which do not depend on natural numbers. Further, we give a formula for operators which give similarity.

Theorem 5.3. For two operator-valued p-ASFs $(\{A_n\}_n, \{\Psi_n\}_n)$ and $(\{B_n\}_n, \{\Phi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$, the following are equivalent.

- (i) $B_n = A_n R_{A,B}, \Phi_n = L_{\Psi,\Phi} \Psi_n, \forall n \in \mathbb{N}, \text{ for some invertible } R_{A,B}, L_{\Psi,\Phi} \in \mathcal{B}(\mathcal{X}).$
- (ii) $\theta_B = \theta_A R_{A,B}, \ \theta_\Phi = L_{\Psi,\Phi} \theta_\Psi$ for some invertible $R_{A,B}, L_{\Psi,\Phi} \in \mathcal{B}(\mathcal{X})$.
- (iii) $P_{B,\Phi} = P_{A,\Psi}$.

If one of the above conditions is satisfied, then invertible operators in (i) and (ii) are unique and are given by $R_{A,B} = S_{A,\Psi}^{-1} \theta_{\Psi} \theta_{B}$, $L_{\Psi,\Phi} = \theta_{\Phi} \theta_{A} S_{A,\Psi}^{-1}$. In the case that $(\{A_n\}_n, \{\Psi_n\}_n)$ is Parseval, then $(\{B_n\}_n, \{\Phi_n\}_n)$ is Parseval if and only if $R_{\Psi,\Phi} L_{A,B} = I_{\mathcal{X}}$ if and only if $L_{A,B} R_{\Psi,\Phi} = I_{\mathcal{X}}$.

Proof. Lemma 5.2 gives (i) \Rightarrow (ii) \Rightarrow (iii). We show (ii) \Rightarrow (i). Assume (ii). We try to recover B_n and Φ_n . Using Equation (1),

$$\begin{split} B_n x &= \Gamma_n \theta_B x = \Gamma_n \theta_A R_{A,B} x = A_n R_{A,B} x, \quad \forall x \in \mathcal{X}, \forall n \in \mathbb{N}, \\ \Phi_n z &= \theta_\Phi L_n z = L_{\Psi, \Phi} \theta_\Psi L_n z = L_{\Psi, \Phi} \Psi_n, \quad \forall n \in \mathbb{N}, \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}. \end{split}$$

We show (iii) \Rightarrow (ii). Assume (iii). A calculation says that $\theta_B = P_{B,\Phi}\theta_B$ and $\theta_{\Phi} = \theta_{\Phi}P_{B,\Phi}$. This observation says that

$$\theta_B = P_{B,\Phi}\theta_B = P_{A,\Psi}\theta_B = \theta_A(S_{A,\Psi}^{-1}\theta_\Psi\theta_B)$$
$$\theta_\Phi = \theta_\Phi P_{B,\Phi} = \theta_\Phi P_{A,\Psi} = (\theta_\Phi\theta_A S_{A,\Psi}^{-1})\theta_\Psi.$$

To obtain (ii) it suffices to show that both $S_{A,\Psi}^{-1}\theta_{\Psi}\theta_{B}$ and $\theta_{\Phi}\theta_{A}S_{A,\Psi}^{-1}$ are invertible. For,

$$(S_{A,\Psi}^{-1}\theta_{\Psi}\theta_{B})(S_{B,\Phi}^{-1}\theta_{\Phi}\theta_{A}) = S_{A,\Psi}^{-1}\theta_{\Psi}P_{B,\Phi}\theta_{A} = S_{A,\Psi}^{-1}\theta_{\Psi}P_{A,\Psi}\theta_{A} = S_{A,\Psi}^{-1}\theta_{\Psi}\theta_{A} = I_{\mathcal{X}},$$
$$(S_{B,\Phi}^{-1}\theta_{\Phi}\theta_{A})(S_{A,\Psi}^{-1}\theta_{\Psi}\theta_{B}) = S_{B,\Phi}^{-1}\theta_{\Phi}P_{A,\Psi}\theta_{B} = S_{B,\Phi}^{-1}\theta_{\Phi}P_{B,\Phi}\theta_{B} = S_{B,\Phi}^{-1}\theta_{\Phi}\theta_{B} = I_{\mathcal{X}}$$

and

$$\begin{aligned} (\theta_{\Phi}\theta_{A}S_{A,\Psi}^{-1})(\theta_{\Psi}\theta_{B}S_{B,\Phi}^{-1}) &= \theta_{\Phi}P_{A,\Psi}\theta_{B}S_{B,\Phi}^{-1} = \theta_{\Phi}P_{B,\Phi}\theta_{B}S_{B,\Phi}^{-1} = \theta_{\Phi}\theta_{B}S_{B,\Phi}^{-1} = I_{\mathcal{X}}, \\ (\theta_{\Psi}\theta_{B}S_{B,\Phi}^{-1})(\theta_{\Phi}\theta_{A}S_{A,\Psi}^{-1}) &= \theta_{\Psi}P_{B,\Phi}\theta_{A}S_{A,\Psi}^{-1} = \theta_{\Psi}P_{A,\Psi}\theta_{A}S_{A,\Psi}^{-1} = \theta_{\Psi}\theta_{A}S_{A,\Psi}^{-1} = I_{\mathcal{X}}. \end{aligned}$$

Let $R_{A,B}$, $L_{\Psi,\Phi} \in \mathcal{B}(\mathcal{X})$ be invertible. We saw that $R_{A,B}$ and $L_{\Psi,\Phi}$ satisfy (i) if and only if they satisfy (ii). Let $B_n = A_n R_{A,B}$, $\Phi_n = L_{\Psi,\Phi} \Psi_n$, $\forall n \in \mathbb{N}$. Using (ii) $\theta_B = \theta_A R_{A,B}$, $\theta_{\Phi} = L_{\Psi,\Phi} \theta_{\Psi}$. Therefore $\theta_{\Psi} \theta_B = \theta_{\Psi} \theta_A R_{A,B} = S_{A,\Psi} R_{A,B}$ and $\theta_{\Phi} \theta_A = L_{\Psi,\Phi} \theta_{\Psi} \theta_A = L_{\Psi,\Phi} S_{A,\Psi}$ which give the required formulas.

Corollary 5.4. For any given operator-valued p-ASF $(\{A_n\}_n, \{\Psi_n\}_n)$, the canonical dual of $(\{A_n\}_n, \{\Psi_n\}_n)$ is the only dual operator-valued p-ASF that is similar to $(\{A_n\}_n, \{\Psi_n\}_n)$.

Proof. Let $(\{B_n\}_n, \{\Phi_n\}_n)$ be an operator-valued p-ASF which is both dual and similar to $(\{A_n\}_n, \{\Psi_n\}_n)$. Then we have $\theta_{\Psi}\theta_B = I_{\mathcal{X}} = \theta_{\Phi}\theta_A$ and there exist invertible $R_{A,B}, L_{\Psi,\Phi} \in \mathcal{B}(\mathcal{X})$ such that $B_n = A_nR_{A,B}, \Phi_n = L_{\Psi,\Phi}\Psi_n, \forall n \in \mathbb{N}$. Theorem 5.3 gives $R_{A,B} = S_{A,\Psi}^{-1}\theta_{\Psi}\theta_B, L_{\Psi,\Phi} = \theta_{\Phi}\theta_A S_{A,\Psi}^{-1}$. But then $R_{A,B} = S_{A,\Psi}^{-1}I_{\mathcal{X}} = S_{A,\Psi}^{-1}, L_{\Psi,\Phi} = I_{\mathcal{X}}S_{A,\Psi}^{-1} = S_{A,\Psi}^{-1}$. Therefore $(\{B_n\}_n, \{\Phi_n\}_n)$ is the canonical dual for $(\{A_n\}_n, \{\Psi_n\}_n)$.

Corollary 5.5. Two similar operator-valued p-ASFs cannot be orthogonal.

Proof. Let an operator-valued p-ASF $(\{B_n\}_n, \{\Phi_n\}_n)$ be similar to $(\{A_n\}_n, \{\Psi_n\}_n)$. Then there are invertible $R_{A,B}, L_{\Psi,\Phi} \in \mathcal{B}(\mathcal{X})$ such that $B_n = A_n R_{A,B}, \Phi_n = L_{\Psi,\Phi} \Psi_n, \forall n \in \mathbb{N}$. Using Theorem 5.3 and

the invertibility of $R_{A,B}$ and $S_{A,\Psi}$, we have

$$\theta_{\Psi}\theta_B = \theta_{\Psi}\theta_A R_{A,B} = S_{A,\Psi}R_{A,B} \neq 0.$$

6. Perturbations

Most useful Paley-Wiener theorem [5] in Hilbert spaces says that sequences which are close to orthonormal bases are Riesz bases and in Banach spaces it states that sequences which are close to Schauder bases are again Schauder bases. This motivated the perturbation of frames for Hilbert spaces. First result about stability of frames for Hilbert spaces was derived by Christensen in [25] and improved in papers [22, 26]. For Banach frames, framings and Schauder frames, stability results are derived in [23, 28, 77, 86]. We obtained stability of p-ASFs in [63]. Here we derive stability results of operator-valued p-ASFs. First we recall a result which is an improvement of result of Hilding [55].

Theorem 6.1. [16, 22, 83] Let \mathcal{X}, \mathcal{Y} be Banach spaces, $U : \mathcal{X} \to \mathcal{Y}$ be a bounded invertible operator. If a bounded linear operator $V : \mathcal{X} \to \mathcal{Y}$ is such that there exist $\alpha, \beta \in [0, 1)$ with

$$\|Ux - Vx\| \le \alpha \|Ux\| + \beta \|Vx\|, \quad \forall x \in \mathcal{X},$$

then V is bounded invertible and

$$\frac{1-\alpha}{1+\beta} \|Ux\| \le \|Vx\| \le \frac{1+\alpha}{1-\beta} \|Ux\|, \quad \forall x \in \mathcal{X};$$
$$\frac{1-\beta}{1+\alpha} \frac{1}{\|U\|} \|y\| \le \|V^{-1}y\| \le \frac{1+\beta}{1-\alpha} \|U^{-1}\| \|y\|, \quad \forall y \in \mathcal{Y}.$$

Theorem 6.2. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. Assume that a collection $\{\Psi_n\}_n$ in $\mathcal{B}(\mathcal{Y}, \mathcal{X})$ is such that there exist $\alpha, \beta, \gamma \geq 0$ with $\max\{\alpha + \frac{\gamma}{\sqrt{a}}, \beta\} < 1$ and

(4)
$$\left\|\sum_{n=1}^{m} (\Psi_n - \Phi_n) \Gamma_n z\right\| \le \alpha \left\|\sum_{n=1}^{m} \Psi_n \Gamma_n z\right\| + \gamma \|z\| + \beta \left\|\sum_{n=1}^{m} \Phi_n \Gamma_n z\right\|, \quad \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}, m = 1, \dots$$

Then $(\{A_n\}_n, \{\Phi_n\}_n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ with frame bounds

$$\frac{\|S_{A,\Psi}^{-1}\|(1+\beta)}{1-(\alpha+\gamma\|\theta_A S_{A,\Psi}^{-1}\|)} \quad and \quad \frac{\|\theta_A\|((1+\alpha)\|\theta_\Psi\|+\gamma)}{1-\beta}.$$

Proof. For $m = 1, \ldots,$

$$\left\|\sum_{n=1}^{m} \Phi_{n} \Gamma_{n} z\right\| \leq \left\|\sum_{n=1}^{m} (\Phi_{n} - \Psi_{n}) \Gamma_{n} z\right\| + \left\|\sum_{n=1}^{m} \Psi_{n} \Gamma_{n} z\right\|$$
$$\leq (1+\alpha) \left\|\sum_{n=1}^{m} \Psi_{n} \Gamma_{n} z\right\| + \beta \left\|\sum_{n=1}^{m} \Phi_{n} \Gamma_{n} z\right\| + \gamma \|z\|, \quad \forall z \in \ell^{p}(\mathbb{N}) \otimes \mathcal{Y}.$$

Hence

$$\left\|\sum_{n=1}^{m} \Phi_n \Gamma_n z\right\| \leq \frac{1+\alpha}{1-\beta} \left\|\sum_{n=1}^{m} \Psi_n \Gamma_n z\right\| + \frac{\gamma}{1-\beta} \|z\|, \quad \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}, m = 1, \dots$$

Therefore θ_{Φ} is a well-defined bounded linear operator with norm estimate

$$\|\theta_{\Phi}\| \leq \frac{1+\alpha}{1-\beta} \|\theta_{\Psi}\| + \frac{\gamma}{1-\beta}.$$

Now Equation (4) gives

$$\left\|\sum_{n=1}^{\infty} (\Psi_n - \Phi_n) \Gamma_n z\right\| \le \alpha \left\|\sum_{n=1}^{\infty} \Psi_n \Gamma_n z\right\| + \gamma \|z\| + \beta \left\|\sum_{n=1}^{\infty} \Phi_n \Gamma_n z\right\|, \quad \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}.$$

This is same as

(5)
$$\|\theta_{\Psi}z - \theta_{\Phi}z\| \le \alpha \|\theta_{\Psi}z\| + \gamma \|z\| + \beta \|\theta_{\Phi}z\|, \quad \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}.$$

For every $x \in \mathcal{X}$, by taking $z = \theta_A S_{A,\Psi}^{-1} x$ in Equation (5), we get

$$\|\theta_{\Psi}\theta_{A}S_{A,\Psi}^{-1}x - \theta_{\Phi}\theta_{A}S_{A,\Psi}^{-1}x\| \le \alpha \|\theta_{\Psi}\theta_{A}S_{A,\Psi}^{-1}x\| + \gamma \|\theta_{A}S_{A,\Psi}^{-1}x\| + \beta \|\theta_{\Phi}\theta_{A}S_{A,\Psi}^{-1}x\|, \quad \forall x \in \mathcal{X}.$$

That is,

$$\begin{aligned} \|x - \theta_{\Phi} \theta_A S_{A,\Psi}^{-1} x\| &\leq \alpha \|x\| + \gamma \|\theta_A S_{A,\Psi}^{-1} x\| + \beta \|\theta_{\Phi} \theta_A S_{A,\Psi}^{-1} x\| \\ &\leq (\alpha + \gamma \|\theta_A S_{A,\Psi}^{-1}\|) \|x\| + \beta \|\theta_{\Phi} \theta_A S_{A,\Psi}^{-1} x\|, \quad \forall x \in \mathcal{X}. \end{aligned}$$

Since $\max\{\alpha + \gamma \| \theta_A S_{A,\Psi}^{-1} \|, \beta\} < 1$, Theorem 6.1 says that operator $S_{A,\Phi} S_{A,\Psi}^{-1}$ is invertible and

$$\|(S_{A,\Phi}S_{A,\Psi}^{-1})^{-1}\| \le \frac{1+\beta}{1-(\alpha+\gamma\|\theta_A S_{A,\Psi}^{-1}\|)}$$

Hence the operator $S_{A,\Phi} = (S_{A,\Phi}S_{A,\Psi}^{-1})S_{A,\Psi}$ is invertible. Therefore $(\{A_n\}_n, \{\Phi_n\}_n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. To get frame bounds we calculate:

$$\|S_{A,\Phi}^{-1}\| \le \|S_{A,\Psi}^{-1}\| \|S_{A,\Psi}S_{A,\Phi}^{-1}\| \le \frac{\|S_{A,\Psi}^{-1}\|(1+\beta)}{1-(\alpha+\gamma\|\theta_A S_{A,\Psi}^{-1}\|)} \quad \text{and} \\ \|S_{A,\Phi}\| \le \|\theta_{\Phi}\| \|\theta_A\| \le \frac{\|\theta_A\|((1+\alpha)\|\theta_{\Psi}\|+\gamma)}{1-\beta}.$$

Theorem 6.3. Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. Assume that a collection $\{B_n\}_n$ in $\mathcal{B}(\mathcal{X},\mathcal{Y})$ and a collection $\{\Phi_n\}_n$ in $\mathcal{B}(\mathcal{Y},\mathcal{X})$ are such that there exist $r, s, t, \alpha, \beta, \gamma \geq 0$ with $\max\{\beta, s\} < 1$ and

$$\left\|\sum_{n=1}^{m} L_n (A_n - B_n) x\right\| \le r \left\|\sum_{n=1}^{m} L_n A_n x\right\| + t \|x\| + s \left\|\sum_{n=1}^{m} L_n B_n x\right\|, \quad \forall x \in \mathcal{X}, m = 1, \dots, \\\left\|\sum_{n=1}^{m} (\Psi_n - \Phi_n) \Gamma_n z\right\| \le \alpha \left\|\sum_{n=1}^{m} \Psi_n \Gamma_n z\right\| + \gamma \|z\| + \beta \left\|\sum_{n=1}^{m} \Phi_n \Gamma_n z\right\|, \quad \forall z \in \ell^p(\mathbb{N}) \otimes \mathcal{Y}, m = 1, \dots.$$

Assume that one of the following holds.

(1)
$$\sum_{n=1}^{\infty} \| (\Psi_n A_n - \Phi_n B_n) S_{A,\Psi}^{-1} \| < 1.$$

(2)
$$\sum_{n=1}^{\infty} \|S_{A,\Psi}^{-1}(\Psi_n A_n - \Phi_n B_n)\| < 1$$

- $\begin{array}{l} (2) \quad \sum_{n=1}^{\infty} \|S_{A,\Psi}^{-1}\Psi_{n}A_{n} \Phi_{n}B_{n}S_{A,\Psi}^{-1}\| < 1. \\ (4) \quad \sum_{n=1}^{\infty} \|\Psi_{n}A_{n}S_{A,\Psi}^{-1} S_{A,\Psi}^{-1}\Phi_{n}B_{n}\| < 1. \end{array}$

Then $(\{B_n\}_n, \{\Phi_n\}_n)$ is an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$. Moreover, an upper frame bound is

$$\left(\frac{1+\alpha}{1-\beta}\|\theta_{\Psi}\| + \frac{\gamma}{1-\beta}\right)\left(\frac{1+r}{1-s}\|\theta_{A}\| + \frac{t}{1-s}\right)$$

Proof. Similar to the arguments given in the beginning of proof of Theorem 6.2, we see that θ_B and θ_{Φ} are well-defined bounded linear operators. We deal with four cases.

Assume (1). Then

$$\begin{aligned} \left\| x - \sum_{n=1}^{\infty} \Phi_n B_n S_{A,\Psi}^{-1} x \right\| &= \left\| \sum_{n=1}^{\infty} \Psi_n A_n S_{A,\Psi}^{-1} x - \sum_{n=1}^{\infty} \Phi_n B_n S_{A,\Psi}^{-1} x \right\| \\ &= \left\| \sum_{n=1}^{\infty} (\Psi_n A_n S_{A,\Psi}^{-1} - \Phi_n B_n S_{A,\Psi}^{-1}) x \right\| \\ &\leq \sum_{n=1}^{\infty} \| (\Psi_n A_n - \Phi_n B_n) S_{A,\Psi}^{-1} \| \| x \|, \quad \forall x \in \mathcal{X} \end{aligned}$$

Therefore the operator $S_{B,\Phi}S_{A,\Psi}^{-1}$ is invertible. Assume (2). Then

$$\begin{aligned} \left\| x - \sum_{n=1}^{\infty} S_{A,\Psi}^{-1} \Phi_n B_n x \right\| &= \left\| \sum_{n=1}^{\infty} S_{A,\Psi}^{-1} \Psi_n A_n x - \sum_{n=1}^{\infty} S_{A,\Psi}^{-1} \Phi_n B_n x \right\| \\ &= \left\| \sum_{n=1}^{\infty} (S_{A,\Psi}^{-1} \Psi_n A_n - S_{A,\Psi}^{-1} \Phi_n B_n) x \right\| \\ &\leq \sum_{n=1}^{\infty} \| S_{A,\Psi}^{-1} (\Psi_n A_n - \Phi_n B_n) \| \| x \|, \quad \forall x \in \mathcal{X} \end{aligned}$$

Therefore the operator $S_{A,\Psi}^{-1}S_{B,\Phi}$ is invertible. Assume (3). Then

$$\begin{aligned} \left\| x - \sum_{n=1}^{\infty} \Phi_n B_n S_{A,\Psi}^{-1} x \right\| &= \left\| \sum_{n=1}^{\infty} S_{A,\Psi}^{-1} \Psi_n A_n x - \sum_{n=1}^{\infty} \Phi_n B_n S_{A,\Psi}^{-1} x \right\| \\ &= \left\| \sum_{n=1}^{\infty} (S_{A,\Psi}^{-1} \Psi_n A_n - \Phi_n B_n S_{A,\Psi}^{-1}) x \right\| \\ &\leq \sum_{n=1}^{\infty} \| S_{A,\Psi}^{-1} \Psi_n A_n - \Phi_n B_n S_{A,\Psi}^{-1} \| \| x \|, \quad \forall x \in \mathcal{X} \end{aligned}$$

Therefore the operator $S_{B,\Phi}S_{A,\Psi}^{-1}$ is invertible. Assume (4). Then

$$\begin{aligned} \left\| x - \sum_{n=1}^{\infty} S_{A,\Psi}^{-1} \Phi_n B_n x \right\| &= \left\| \sum_{n=1}^{\infty} \Psi_n A_n S_{A,\Psi}^{-1} x - \sum_{n=1}^{\infty} S_{A,\Psi}^{-1} \Phi_n B_n x \right\| \\ &= \left\| \sum_{n=1}^{\infty} (\Psi_n A_n S_{A,\Psi}^{-1} - S_{A,\Psi}^{-1} \Phi_n B_n) x \right\| \\ &\leq \sum_{n=1}^{\infty} \|\Psi_n A_n S_{A,\Psi}^{-1} - S_{A,\Psi}^{-1} \Phi_n B_n\| \|x\|, \quad \forall x \in \mathcal{X} \end{aligned}$$

Therefore the operator $S_{A,\Psi}^{-1}S_{B,\Phi}$ is invertible.

7. Operator-valued Feichtinger conjectures

Recently first author formulated a variety of conjectures and problems for p-approximate Schauder frames for Banach spaces [59, 60]. Here we present operator-valued versions of some of them.

Conjecture 7.1. (Feichtinger conjecture for operator-valued p-ASFs) Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ such that

$$0 < \inf_{n \in \mathbb{N}} \|\Psi_n\| \le \sup_{n \in \mathbb{N}} \|\Psi_n\| < \infty \quad \textit{ and } \quad 0 < \inf_{n \in \mathbb{N}} \|A_n\| \le \sup_{n \in \mathbb{N}} \|A_n\| < \infty$$

Then $(\{A_n\}_n, \{\Psi_n\}_n)$ can be partitioned into a finite union of operator-valued p-approximate Riesz sequences. Moreover, what is the number of partitions required?

Conjecture 7.2. (Feichtinger conjecture for operator-valued p-ABSs) Let $(\{A_n\}_n, \{\Psi_n\}_n)$ be an operator-valued p-ABS in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ such that

$$0 < \inf_{n \in \mathbb{N}} \|\Psi_n\| \le \sup_{n \in \mathbb{N}} \|\Psi_n\| < \infty \quad \textit{ and } \quad 0 < \inf_{n \in \mathbb{N}} \|A_n\| \le \sup_{n \in \mathbb{N}} \|A_n\| < \infty$$

Then $(\{A_n\}_n, \{\Psi_n\}_n)$ can be partitioned into a finite union of operator-valued p-approximate Riesz sequences. Moreover, what is the number of partitions required?

Definition 7.3. A operator-valued p-ASF $(\{A_n\}_n, \{\Psi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ is said to be **p-scalable** if there exist sequences of operators $\{T_n\}_n$ and $\{S_n\}_n$ in $\mathcal{B}(\mathcal{X})$ such that

 $(\{T_nC_n\}_n, \{S_n\Psi_n\}_n)$ is a Parseval operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$.

Problem 7.4. (Scaling problem for Banach spaces) For given Banach spaces \mathcal{X} , \mathcal{Y} , classify operator-valued p-ASFs $(\{A_n\}_n, \{\Psi_n\}_n)$ in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$ so that there are sequences of operators $\{T_n\}_n$, $\{S_n\}_n$ in $\mathcal{B}(\mathcal{X})$ such that $(\{A_nT_n\}_n, \{S_n\Psi_n\}_n)$ is a Parseval operator-valued p-ASF in $\mathcal{B}(\mathcal{X}, \mathcal{Y})$, i.e., $(\{A_n\}_n, \{\Psi_n\}_n)$ is p-scalable.

Note that, due to Theorem 3.1 in [78] (and due to the fundamental works of Marcus, Spielman, Srivastava, Casazza, Edidin, Weaver, Fickus, Tremain, Vershynin, Anderson, and Weber [3,13,14,21,74,84]), operator-valued Feichtinger conjectures for operator-valued frames/G-frames are true for Hilbert spaces.

8. Acknowledgements

First author thanks Prof. B. V. Rajarama Bhat for the Post Doc position through his J. C. Bose Fellowship (SERB).

References

- Akram Aldroubi, Carlos Cabrelli, and Ursula M. Molter. Wavelets on irregular grids with arbitrary dilation matrices and frame atoms for L²(R^d). Appl. Comput. Harmon. Anal., 17(2):119–140, 2004.
- [2] Akram Aldroubi, Qiyu Sun, and Wai-Shing Tang. p-frames and shift invariant subspaces of L^p. J. Fourier Anal. Appl., 7(1):1–21, 2001.
- [3] Joel Anderson. A conjecture concerning the pure states of B(H) and a related theorem. In Topics in modern operator theory (Timişoara/Herculane, 1980), volume 2 of Operator Theory: Advances and Applications, pages 27–43. Birkhäuser, Basel-Boston, Mass., 1981.
- [4] Jean-Pierre Antoine and Peter Balazs. Frames and semi-frames. J. Phys. A, 44(20):205201, 25, 2011.
- [5] Maynard G. Arsove. The Paley-Wiener theorem in metric linear spaces. Pacific J. Math., 10:365–379, 1960.
- [6] Radu Balan. Equivalence relations and distances between Hilbert frames. Proc. Amer. Math. Soc., 127(8):2353–2366, 1999.
- [7] Radu Victor Balan. A study of Weyl-Heisenberg and wavelet frames. ProQuest LLC, Ann Arbor, MI, 1998. Thesis (Ph.D.)-Princeton University.
- [8] Peter Balazs, Jean-Pierre Antoine, and Anna Gryboś. Weighted and controlled frames: mutual relationship and first numerical properties. Int. J. Wavelets Multiresolut. Inf. Process., 8(1):109–132, 2010.

- [9] P. G. Casazza, S. J. Dilworth, E. Odell, Th. Schlumprecht, and A. Zsák. Coefficient quantization for frames in Banach spaces. J. Math. Anal. Appl., 348(1):66–86, 2008.
- [10] Pete Casazza, Ole Christensen, and Diana T. Stoeva. Frame expansions in separable Banach spaces. J. Math. Anal. Appl., 307(2):710–723, 2005.
- [11] Peter G. Casazza. Approximation properties. In Handbook of the geometry of Banach spaces, Vol. I, pages 271–316. North-Holland, Amsterdam, 2001.
- [12] Peter G. Casazza and Ole Christensen. The reconstruction property in Banach spaces and a perturbation theorem. Canad. Math. Bull., 51(3):348–358, 2008.
- [13] Peter G. Casazza and Dan Edidin. Equivalents of the Kadison-Singer problem. In *Function spaces*, volume 435 of *Contemp. Math.*, pages 123–142. Amer. Math. Soc., Providence, RI, 2007.
- [14] Peter G. Casazza, Matthew Fickus, Janet C. Tremain, and Eric Weber. The Kadison-Singer problem in mathematics and engineering: a detailed account. In Operator theory, operator algebras, and applications, volume 414 of Contemp. Math., pages 299–355. Amer. Math. Soc., Providence, RI, 2006.
- [15] Peter G. Casazza, Deguang Han, and David R. Larson. Frames for Banach spaces. In *The functional and harmonic analysis of wavelets and frames (San Antonio, TX, 1999)*, volume 247 of *Contemp. Math.*, pages 149–182. Amer. Math. Soc., Providence, RI, 1999.
- [16] Peter G. Casazza and Nigel J. Kalton. Generalizing the Paley-Wiener perturbation theory for Banach spaces. Proc. Amer. Math. Soc., 127(2):519–527, 1999.
- [17] Peter G. Casazza and Jelena Kovačević. Equal-norm tight frames with erasures. Adv. Comput. Math., 18(2-4):387–430, 2003.
- [18] Peter G. Casazza and Gitta Kutyniok. Frames of subspaces. In Wavelets, frames and operator theory, volume 345 of Contemp. Math., pages 87–113. Amer. Math. Soc., Providence, RI, 2004.
- [19] Peter G. Casazza, Gitta Kutyniok, and Shidong Li. Fusion frames and distributed processing. Appl. Comput. Harmon. Anal., 25(1):114–132, 2008.
- [20] Peter G. Casazza and Nicole Leonhard. Classes of finite equal norm Parseval frames. In Frames and operator theory in analysis and signal processing, volume 451 of Contemp. Math., pages 11–31. Amer. Math. Soc., Providence, RI, 2008.
- [21] Peter G. Casazza and Janet Crandell Tremain. The Kadison-Singer problem in mathematics and engineering. Proc. Natl. Acad. Sci. USA, 103(7):2032–2039, 2006.
- [22] Peter G. Cazassa and Ole Christensen. Perturbation of operators and applications to frame theory. J. Fourier Anal. Appl., 3(5):543–557, 1997.
- [23] Dong Yang Chen, Lei Li, and Ben Tuo Zheng. Perturbations of frames. Acta Math. Sin. (Engl. Ser.), 30(7):1089–1108, 2014.
- [24] O. Christensen and Y. C. Eldar. Oblique dual frames and shift-invariant spaces. Appl. Comput. Harmon. Anal., 17(1):48–68, 2004.
- [25] Ole Christensen. Frame perturbations. Proc. Amer. Math. Soc., 123(4):1217–1220, 1995.
- [26] Ole Christensen. A Paley-Wiener theorem for frames. Proc. Amer. Math. Soc., 123(7):2199–2201, 1995.
- [27] Ole Christensen. An introduction to frames and Riesz bases. Applied and Numerical Harmonic Analysis. Birkhäuser/Springer, second edition, 2016.
- [28] Ole Christensen and Christopher Heil. Perturbations of Banach frames and atomic decompositions. Math. Nachr., 185:33–47, 1997.
- [29] Ole Christensen and Richard S. Laugesen. Approximately dual frames in Hilbert spaces and applications to Gabor frames. Sampl. Theory Signal Image Process., 9(1-3):77–89, 2010.
- [30] Ole Christensen and Diana T. Stoeva. p-frames in separable Banach spaces. Adv. Comput. Math., 18(2-4):117–126, 2003.
- [31] Joachim Cuntz. Simple C*-algebras generated by isometries. Comm. Math. Phys., 57(2):173–185, 1977.
- [32] Wojciech Czaja. Remarks on Naimark's duality. Proc. Amer. Math. Soc., 136(3):867–871, 2008.
- [33] Ingrid Daubechies and Ron DeVore. Approximating a bandlimited function using very coarsely quantized data: a family of stable sigma-delta modulators of arbitrary order. Ann. of Math. (2), 158(2):679–710, 2003.
- [34] Ingrid Daubechies, A. Grossmann, and Y. Meyer. Painless nonorthogonal expansions. J. Math. Phys., 27(5):1271–1283, 1986.
- [35] Andreas Defant and Klaus Floret. Tensor norms and operator ideals, volume 176 of North-Holland Mathematics Studies. North-Holland Publishing Co., Amsterdam, 1993.

- [36] Joe Diestel, Jan H. Fourie, and Johan Swart. The metric theory of tensor products: Grothendieck's resume revisited. American Mathematical Society, Providence, RI, 2008.
- [37] R. J. Duffin and A. C. Schaeffer. A class of nonharmonic Fourier series. Trans. Amer. Math. Soc., 72:341–366, 1952.
- [38] Hans G. Feichtinger and K. H. Gröchenig. Banach spaces related to integrable group representations and their atomic decompositions. I. J. Funct. Anal., 86(2):307–340, 1989.
- [39] Hans G. Feichtinger and K. H. Gröchenig. Banach spaces related to integrable group representations and their atomic decompositions. II. Monatsh. Math., 108(2-3):129–148, 1989.
- [40] Hans G. Feichtinger and Karlheinz Gröchenig. A unified approach to atomic decompositions via integrable group representations. In *Function spaces and applications (Lund, 1986)*, volume 1302 of *Lecture Notes in Math.*, pages 52–73. Springer, Berlin, 1988.
- [41] Hans G. Feichtinger and Karlheinz Gröchenig. Theory and practice of irregular sampling. In Wavelets: mathematics and applications, Stud. Adv. Math., pages 305–363. CRC, Boca Raton, FL, 1994.
- [42] Abolhassan Fereydooni and Ahmad Safapour. Pair frames. Results Math., 66(1-2):247–263, 2014.
- [43] Massimo Fornasier. Quasi-orthogonal decompositions of structured frames. J. Math. Anal. Appl., 289(1):180–199, 2004.
- [44] D. Freeman, E. Odell, Th. Schlumprecht, and A. Zsák. Unconditional structures of translates for $L_p(\mathbb{R}^d)$. Israel J. Math., 203(1):189–209, 2014.
- [45] Laura Gavruta. Frames for operators. Appl. Comput. Harmon. Anal., 32(1):139-144, 2012.
- [46] Ramu Geddavalasa and P. Sam Johnson. Frames for operators in Banach spaces. Acta Math. Vietnam., 42(4):665–673, 2017.
- [47] K. Grochenig. Acceleration of the frame algorithm. IEEE Transactions on Signal Processing, 41(12):1331–1340, 1993.
- [48] Karlheinz Gröchenig. Describing functions: atomic decompositions versus frames. Monatsh. Math., 112(1):1-42, 1991.
- [49] Deguang Han, Keri Kornelson, David Larson, and Eric Weber. Frames for undergraduates, volume 40 of Student Mathematical Library. American Mathematical Society, Providence, RI, 2007.
- [50] Deguang Han and David R. Larson. Frames, bases and group representations. Mem. Amer. Math. Soc., 147(697):x+94, 2000.
- [51] Deguang Han, David R. Larson, Bei Liu, and Rui Liu. Dilations of frames, operator-valued measures and bounded linear maps. In Operator methods in wavelets, tilings, and frames, volume 626 of Contemp. Math., pages 33–53. Amer. Math. Soc., Providence, RI, 2014.
- [52] Deguang Han, David R. Larson, Bei Liu, and Rui Liu. Operator-valued measures, dilations, and the theory of frames. Mem. Amer. Math. Soc., 229(1075):viii+84, 2014.
- [53] DeGuang Han, PengTong Li, Bin Meng, and WaiShing Tang. Operator valued frames and structured quantum channels. Sci. China Math., 54(11):2361–2372, 2011.
- [54] Christopher Heil. A basis theory primer. Applied and Numerical Harmonic Analysis. Birkhäuser/Springer, New York, expanded edition, 2011.
- [55] Sven H. Hilding. Note on completeness theorems of Paley-Wiener type. Ann. of Math. (2), 49:953–955, 1948.
- [56] James R. Holub. Pre-frame operators, Besselian frames, and near-Riesz bases in Hilbert spaces. Proc. Amer. Math. Soc., 122(3):779–785, 1994.
- [57] Victor Kaftal, David R. Larson, and Shuang Zhang. Operator-valued frames. Trans. Amer. Math. Soc., 361(12):6349– 6385, 2009.
- [58] B. S. Kashin and T. Yu. Kulikova. A remark on the description of frames of general form. Mat. Zametki, 72(6):941–945, 2002.
- [59] K. Mahesh Krishna. Feichtinger conjectures, R_{ε} -conjecture and Weaver's conjecture for Banach spaces. arXiv: 2201.00125 v1 [math.FA], 1 January, 2022.
- [60] K. Mahesh Krishna. Paulsen and projection problems for Banach spaces. arXiv: 2201.00991v1 [math.FA], 4 January, 2022.
- [61] K. Mahesh Krishna and P. Sam Johnson. Approximately dual p-approximate Schauder frames. arXiv:2110.10121 [math.FA] 19 October 2021.
- [62] K. Mahesh Krishna and P. Sam Johnson. Dilation theorem for p-approximate Schauder frames for separable Banach spaces. Palestine Journal of Mathematics (accepted).
- [63] K. Mahesh Krishna and P. Sam Johnson. Perturbation of p-approximate Schauder frames for separable Banach spaces. Poincare Journal of Analysis and Applications (accepted).

- [64] K. Mahesh Krishna and P. Sam Johnson. Towards characterizations of approximate Schauder frame and its duals for Banach spaces. J. Pseudo-Differ. Oper. Appl., 12(1):Paper No. 9, 13, 2021.
- [65] K. Mahesh Krishna and P. Sam Johnson. Expansion of weak reconstruction sequences to approximate Schauder frames for Banach spaces. Asian-European Journal of Mathematics, (2250060), 2022.
- [66] K. Mahesh Krishna and P. Sam Johnson. Factorable weak operator-valued frames. Annals of Functional Analysis, 13(11), 2022.
- [67] Deng Feng Li and Wen Chang Sun. Expansion of frames to tight frames. Acta Math. Sin. (Engl. Ser.), 25(2):287–292, 2009.
- [68] Shidong Li. On general frame decompositions. Numer. Funct. Anal. Optim., 16(9-10):1181–1191, 1995.
- [69] Shidong Li and Hidemitsu Ogawa. Pseudoframes for subspaces with applications. J. Fourier Anal. Appl., 10(4):409–431, 2004.
- [70] Shidong Li and Dunyan Yan. Frame fundamental sensor modeling and stability of one-sided frame perturbation. Acta Appl. Math., 107(1-3):91–103, 2009.
- [71] Bei Liu, Rui Liu, and Bentuo Zheng. Parseval p-frames and the Feichtinger conjecture. J. Math. Anal. Appl., 424(1):248– 259, 2015.
- [72] Rui Liu. Hilbert-Schauder frame operators. Oper. Matrices, 7(1):91–99, 2013.
- [73] Rui Liu and Zhong-Jin Ruan. Cb-frames for operator spaces. J. Funct. Anal., 270(11):4280–4296, 2016.
- [74] Adam W. Marcus, Daniel A. Spielman, and Nikhil Srivastava. Interlacing families II: Mixed characteristic polynomials and the Kadison-Singer problem. Ann. of Math. (2), 182(1):327–350, 2015.
- [75] Irine Peng and Shayne Waldron. Signed frames and Hadamard products of Gram matrices. Linear Algebra Appl., 347:131–157, 2002.
- [76] Raymond A. Ryan. Introduction to tensor products of Banach spaces. Springer Monographs in Mathematics. Springer-Verlag London, Ltd., London, 2002.
- [77] Diana T. Stoeva. Perturbation of frames in Banach spaces. Asian-Eur. J. Math., 5(1):1250011, 15, 2012.
- [78] Wenchang Sun. G-frames and g-Riesz bases. J. Math. Anal. Appl., 322(1):437-452, 2006.
- [79] P. A. Terëkhin. Representation systems and projections of bases. Mat. Zametki, 75(6):944–947, 2004.
- [80] P. A. Terëkhin. Banach frames in the affine synthesis problem. Mat. Sb., 200(9):127–146, 2009.
- [81] P. A. Terekhin. Frames in a Banach space. Funktsional. Anal. i Prilozhen., 44(3):50–62, 2010.
- [82] S.M. Thomas. Approximate Schauder Frames for \mathbb{R}^n . Masters Thesis, St. Louis University, St. Louis, MO, 2012.
- [83] S.J.L. van Eijndhoven. Hilding's theorem for Banach spaces. RANA: reports on applied and numerical analysis, volume 9612: Technische Universiteit Eindhoven, pages 1–6, 1996.
- [84] Nik Weaver. The Kadison-Singer problem in discrepancy theory. Discrete Math., 278(1-3):227–239, 2004.
- [85] Robert M. Young. An introduction to nonharmonic Fourier series, volume 93 of Pure and Applied Mathematics. Academic Press, Inc., New York-London, 1980.
- [86] Yu Can Zhu and Si Yuan Wang. The stability of Banach frames in Banach spaces. Acta Math. Sin. (Engl. Ser.), 26(12):2369–2376, 2010.