## AN APOCALYPSE-FREE FIRST-ORDER LOW-RANK OPTIMIZATION ALGORITHM\*

GUILLAUME OLIKIER<sup>†</sup>, KYLE A. GALLIVAN<sup>‡</sup>, AND P.-A. ABSIL<sup>†</sup>

**Abstract.** We consider the problem of minimizing a differentiable function with locally Lipschitz continuous gradient on the real determinantal variety, and present a first-order algorithm designed to find stationary points of that problem. This algorithm applies steps of steepest descent with back-tracking line search on the variety, as proposed by Schneider and Uschmajew (2015), but by taking the numerical rank into account to perform suitable rank reductions. We prove that this algorithm produces sequences of iterates the accumulation points of which are stationary, and therefore does not follow the so-called apocalypses described by Levin, Kileel, and Boumal (2021).

Key words. Convergence analysis  $\cdot$  Stationarity  $\cdot$  Low-rank optimization  $\cdot$  Determinantal variety  $\cdot$  Steepest descent  $\cdot$  Tangent cones.

AMS subject classifications. 14M12, 65K10, 90C26, 90C30, 40A05.

1. Introduction. We consider the problem

(1) 
$$\min_{X \in \mathbb{R}_{< r}^{m \times n}} f(X)$$

of minimizing a differentiable function  $f : \mathbb{R}^{m \times n} \to \mathbb{R}$  with locally Lipschitz continuous gradient on the *determinantal variety* [6, Lecture 9]

(2) 
$$\mathbb{R}^{m \times n}_{< r} := \{ X \in \mathbb{R}^{m \times n} \mid \operatorname{rank} X \le r \},$$

m, n, and r being positive integers such that  $r < \min\{m, n\}$ . This problem has been shown to appear in several applications such as matrix equations, model reduction, matrix sensing, and matrix completion; see, e.g., [15, 5] and the references therein. As problem (1) is in general intractable [3], our goal is just to find a *stationary point* of this problem, i.e., a zero of the *stationarity measure* 

(3) 
$$\mathbf{s}_f: \mathbb{R}^{m \times n}_{\leq r} \to \mathbb{R}: X \mapsto \|P_{T_{\mathbb{R}^{m \times n}_{\leq r}}(X)}(-\nabla f(X))\|$$

that returns the norm of any projection of  $-\nabla f(X)$  onto the tangent cone to  $\mathbb{R}_{\leq r}^{m \times n}$  at X; the notation is introduced in Section 2.

To the best of our knowledge, no first-order algorithm in the literature (see Proposition A.2) has been proved to produce, when given an arbitrary initial iterate, a feasible sequence  $(X_i)_{i\in\mathbb{N}}$  such that  $s_f$  goes to zero along every convergent subsequence. Furthermore, this property would not be sufficient to guarantee that every accumulation point of  $(X_i)_{i\in\mathbb{N}}$  is a stationary point. Indeed, even if  $(X_i)_{i\in\mathbb{N}}$  is convergent and  $\lim_{i\to\infty} s_f(X_i) = 0$ , because of the discontinuity of  $T_{\mathbb{R}^{m\times n}_{\leq r}}(\cdot)$  [12, Theorem 4.1],  $s_f$  may fail to be lower semicontinuous at the limit X of  $(X_i)_{i\in\mathbb{N}}$ , i.e., it may happen that  $s_f(X) > 0$ , the triplet  $(X, (X_i)_{i\in\mathbb{N}}, f)$  being then called an *apocalypse* according

<sup>\*</sup>This work was supported by the Fonds de la Recherche Scientifique – FNRS and the Fonds Wetenschappelijk Onderzoek – Vlaanderen under EOS Project no 30468160.

<sup>&</sup>lt;sup>†</sup>ICTEAM Institute, UCLouvain, Avenue Georges Lemaître 4, 1348 Louvain-la-Neuve, Belgium (guillaume.olikier@uclouvain.be, pa.absil@uclouvain.be).

<sup>&</sup>lt;sup>‡</sup>Department of Mathematics, Florida State University, 1017 Academic Way, Tallahassee, FL 32306-4510, USA (kgallivan@fsu.edu).

to [10, Definition 2.7]. As a matter of fact, in [10, §2.2] is presented an example of a polynomial function  $f : \mathbb{R}_{\leq 2}^{3 \times 3} \to \mathbb{R}$  and of an initial iterate  $X_0 \in \mathbb{R}_{\leq 2}^{3 \times 3}$  such that the sequence  $(X_i)_{i \in \mathbb{N}}$  produced by [15, Algorithm 3]—dubbed P<sup>2</sup>GD—converges to X, and  $(X, (X_i)_{i \in \mathbb{N}}, f)$  is an apocalypse. In [10, §3], a second-order algorithm is proposed that, under some assumptions on f, produces sequences the accumulation points of which are stationary. In [10, §4], the following question is raised: "Is there an algorithm running directly on  $\mathbb{R}_{\leq r}^{m \times n}$  that only uses first-order information about the cost function and which is guaranteed to converge to a stationary point?"

In this paper, we introduce a first-order algorithm on  $\mathbb{R}_{\leq r}^{m \times n}$  (Algorithm 2) that is apocalypse-free, in the sense that every accumulation point of the generated sequence is a stationary point (Theorem 5.2), which also implies that  $s_f$  goes to zero along every convergent subsequence (Corollary 5.3). This algorithm applies the main step of P<sup>2</sup>GD but by taking the numerical rank into account to perform suitable rank reductions. As mentioned in [10, Remark 2.11], low-rank optimization algorithms using rank reductions can already be found in the literature; see, e.g., [17, Algorithm 3] and [9, §3.3]. However, the proposed algorithm uses rank reductions to exploit the continuity of both the singular values and the restriction of  $s_f$  to the smooth manifold [7, Proposition 4.1]

(4) 
$$\mathbb{R}_r^{m \times n} := \{ X \in \mathbb{R}^{m \times n} \mid \operatorname{rank} X = \underline{r} \}$$

for every  $\underline{r} \in \{1, \ldots, r\}$ ; see Corollary 2.11 for the latter. This allows us to prove Theorem 5.2. An overview of Algorithm 2's design and analysis strategy can be found in Section 4.

This paper is organized as follows. After preliminaries in Section 2, we analyze  $P^2GD$  in Section 3 under the assumption of local Lipschitz continuity of  $\nabla f$ . Then, we introduce the proposed algorithm in Section 4, and conduct a convergence analysis in Section 5 based on the results collected in the preceding sections. Finally, Section 6 gathers concluding remarks, and Appendix A complementary results.

2. Preliminaries. This section gathers notation and preliminary results that will be used throughout the paper. Section 2.1 recalls basic facts about the projection onto closed cones in  $\mathbb{R}^{m \times n}$ , and Section 2.2 focuses in particular on the projection onto  $\mathbb{R}_{\leq \underline{r}}^{m \times n}$  for every nonnegative integer  $\underline{r} < \min\{m, n\}$ , with the convention  $\mathbb{R}_{\leq 0}^{m \times n} := \mathbb{R}_{0}^{m \times n} := \{0_{m \times n}\}$ . Based on Section 2.1, we review in Section 2.3 the notion of tangent cone and the stationarity measure  $s_f$  defined in (3). Then, in Section 2.4, we derive an upper bound on the distance to  $\mathbb{R}_{\leq \underline{r}}^{m \times n}$  from any of its tangent lines. Next, in Section 2.5, we prove based on [12, Theorem 4.1] that, for every  $\underline{r} \in \{1, \ldots, r\}$ , the restriction of  $s_f$  to the smooth manifold  $\mathbb{R}_{\underline{r}}^{m \times n}$  defined in (4) is continuous. Finally, in Section 2.6, we prove that  $\mathbb{R}_{\leq r}^{m \times n}$  has no serendipitous point in the sense of [10, Definition 2.8].

**2.1. Projection onto closed cones.** In this paper,  $\mathbb{R}^{m \times n}$  is endowed with the Frobenius inner product  $\langle \cdot, \cdot \rangle$ ,  $\| \cdot \|$  denotes the Frobenius norm, and, for every  $X \in \mathbb{R}^{m \times n}$  and every  $\rho \in (0, \infty)$ ,  $B[X, \rho] := \{Y \in \mathbb{R}^{m \times n} \mid \|X - Y\| \le \rho\}$  is the closed ball of center X and radius  $\rho$  in  $\mathbb{R}^{m \times n}$ . For every nonempty subset S of  $\mathbb{R}^{m \times n}$  and every  $X \in \mathbb{R}^{m \times n}$ ,  $d(X, S) := \inf_{Y \in S} \|X - Y\|$  is the distance from X to S, and  $P_{\mathcal{S}}(X) := \operatorname{argmin}_{Y \in S} \|X - Y\|$  is the projection of X onto S;  $P_{\mathcal{S}}(X)$  can be empty in general but not if S is closed, as formulated in the following proposition.

PROPOSITION 2.1 ([14, Example 1.20]). For every nonempty closed subset S of  $\mathbb{R}^{m \times n}$  and every  $X \in \mathbb{R}^{m \times n}$ ,  $P_{S}(X)$  is nonempty and compact.

A nonempty subset  $\mathcal{C}$  of  $\mathbb{R}^{m \times n}$  is said to be a *cone* if, for every  $X \in \mathcal{C}$  and every  $\lambda \in [0, \infty), \lambda X \in \mathcal{C}$ . In this paper, we will mostly project onto closed cones and, in that case, the preceding proposition can be completed as follows.

PROPOSITION 2.2 ([10, Proposition A.6]). Let  $\mathcal{C} \subseteq \mathbb{R}^{m \times n}$  be a closed cone. For every  $X \in \mathbb{R}^{m \times n}$  and every  $Y \in P_{\mathcal{C}}(X)$ ,

$$||Y||^{2} = ||X||^{2} - d(X, \mathcal{C})^{2}$$

**2.2.** Numerical rank and rank reduction. The goal of this section is to prove Proposition 2.5 on which the proposed algorithm crucially relies. To this end, we first need to review basic facts about singular values, numerical rank, and rank reduction.

In what follows, the singular values of  $X \in \mathbb{R}^{m \times n}$  are denoted by  $\sigma_1(X) \geq \cdots \geq \sigma_{\min\{m,n\}}(X) \geq 0$ , as in [4, §2.4.1]. Moreover, if  $X \neq 0_{m \times n}$ , then  $\sigma_1(X)$  and  $\sigma_{\operatorname{rank} X}(X)$  are respectively denoted by  $\sigma_{\max}(X)$  and  $\sigma_{\min}(X)$ . The following proposition shows that the singular values are Lipschitz continuous with Lipschitz constant 1, a property that will be used in the proof of Proposition 2.5 and in the convergence analysis in Section 5.

PROPOSITION 2.3 ([4, Corollary 8.6.2]). For every  $X, Y \in \mathbb{R}^{m \times n}$  and every  $j \in \{1, \ldots, \min\{m, n\}\},\$ 

$$|\sigma_j(X) - \sigma_j(Y)| \le \sigma_1(X - Y) \le ||X - Y||.$$

The numerical rank can be defined as follows.

DEFINITION 2.4 ([4, p. 276]). Given  $\Delta \in [0, \infty)$  and  $X \in \mathbb{R}^{m \times n}$ , the  $\Delta$ -rank of X is

$$\operatorname{rank}_{\Delta} X := \begin{cases} 0 & \text{if } X = 0_{m \times n} \\ \max\{j \in \{1, \dots, \operatorname{rank} X\} \mid \sigma_j(X) > \Delta\} & \text{otherwise.} \end{cases}$$

By reducing the rank of  $X \in \mathbb{R}^{m \times n}$ , we mean computing an element of  $P_{\mathbb{R}_{\leq \underline{r}}^{m \times n}}(X)$  for some nonnegative integer  $\underline{r} < \operatorname{rank} X$ . According to the Eckart–Young theorem [1], this can be achieved by truncating the SVD of X. In particular, for every nonnegative integer  $\underline{r} < \min\{m, n\}$  and every  $X \in \mathbb{R}^{m \times n}$ :

- 1. if rank  $X \leq \underline{r}$ , then  $P_{\mathbb{R}_{\leq \underline{r}}^{m \times n}}(X) = \{X\};$
- 2. if rank  $X > \underline{r}$ , then  $d(X, \mathbb{R}_{\leq \underline{r}}^{m \times n}) = d(X, \mathbb{R}_{\underline{r}}^{m \times n}) = \sqrt{\sum_{j=\underline{r}+1}^{\operatorname{rank} X} \sigma_j^2(X)}$  and  $P_{\mathbb{R}_{\leq \underline{r}}^{m \times n}}(X) = P_{\mathbb{R}_{\underline{r}}^{m \times n}}(X).$

We can now introduce Proposition 2.5; it will be invoked in the convergence analysis conducted in Section 5. For convenience, we introduce the notation  $\mathbb{R}_{\leq \underline{r}}^{m \times n} := \mathbb{R}_{\leq \underline{r}}^{m \times n} \setminus \mathbb{R}_{\underline{r}}^{m \times n}$  and  $\mathbb{R}_{\geq \underline{r}}^{m \times n} := \mathbb{R}^{m \times n} \setminus \mathbb{R}_{\leq \underline{r}}^{m \times n}$  for every positive integer  $\underline{r} \leq \min\{m, n\}$ .

PROPOSITION 2.5. Let  $\underline{r} \in \{1, \ldots, \min\{m, n\}\}$  and  $X \in \mathbb{R}_{\underline{r}}^{m \times n}$ . For every  $\varepsilon \in (0, \infty)$  and every  $Y \in B[X, \varepsilon]$ ,  $P_{\mathbb{R}_{\leq \underline{r}}^{m \times n}}(Y) \subset B[X, 2\varepsilon]$ . Moreover, for every  $\Delta \in (0, \infty)$ , every  $\varepsilon \in (0, \sigma_{\underline{r}}(X)) \cap (0, \Delta]$ , and every  $Y \in B[X, \varepsilon]$ , it holds that  $\operatorname{rank}_{\Delta} Y \leq \underline{r} \leq \operatorname{rank} Y$ .

*Proof.* The inclusion holds because, for every  $\hat{Y} \in P_{\mathbb{R}^{m \times n}}(Y)$ ,

$$||X - \hat{Y}|| \le ||X - Y|| + ||Y - \hat{Y}|| = ||X - Y|| + d(Y, \mathbb{R}_{\underline{r}}^{m \times n}) \le ||X - Y|| + ||X - Y|| \le 2\varepsilon.$$

The second inequality holds because  $B[X, \varepsilon] \subset \mathbb{R}_{\geq \underline{r}}^{m \times n}$  since  $\varepsilon < \sigma_{\underline{r}}(X) = d(X, \mathbb{R}_{\leq \underline{r}}^{m \times n})$ . Besides, the first inequality is trivial if the second one is an equality. Let us therefore consider  $Y \in B[X, \varepsilon]$  such that  $\underline{r} < \operatorname{rank} Y$ . Then, the first inequality holds because

$$\sigma_{\underline{r}+1}(Y) = |\sigma_{\underline{r}+1}(Y) - \sigma_{\underline{r}+1}(X)| \le ||X - Y|| \le \varepsilon \le \Delta,$$

where the first inequality follows from Proposition 2.3.

**2.3.** Tangent cone and stationarity measure on the determinantal variety. The concept of tangent cone plays a fundamental role in constrained optimization.

PROPOSITION 2.6. For every nonempty subset S of  $\mathbb{R}^{m \times n}$  and every  $X \in S$ , the set

$$T_{\mathcal{S}}(X) := \left\{ V \in \mathbb{R}^{m \times n} \mid \exists \begin{array}{l} (t_i)_{i \in \mathbb{N}} \text{ in } (0, \infty) \text{ converging to } 0\\ (V_i)_{i \in \mathbb{N}} \text{ in } \mathbb{R}^{m \times n} \text{ converging to } V \end{array} \right. : X + t_i V_i \in \mathcal{S} \; \forall i \in \mathbb{N} \right\}$$

is a closed cone, not necessarily convex however, called the tangent cone to S at X.

*Proof.* We refer to  $[12, \S2.3]$  and the references therein for a more complete review of the concept of tangent cone.

We review in the forthcoming Proposition 2.7 formulas describing  $T_{\mathbb{R}^{m \times n}_{\leq r}}(X)$  and  $P_{T_{\mathbb{R}^{m \times n}_{\leq r}}(X)}(G)$  for every  $X \in \mathbb{R}^{m \times n}_{\leq r}$  and every  $G \in \mathbb{R}^{m \times n}$  based on the SVD of X. Before that, we can already observe that Propositions 2.2 and 2.6 together imply that the stationarity measure  $s_f$  introduced in (3) is well defined with, for every  $X \in \mathbb{R}^{m \times n}_{< r}$ ,

(5) 
$$s_f(X) = \sqrt{\|\nabla f(X)\|^2 - d(-\nabla f(X), T_{\mathbb{R}^{m \times n}_{\leq r}}(X))^2}$$

The formulas given in the following proposition can be obtained from the definition in Proposition 2.6.

PROPOSITION 2.7 ([15, Theorem 3.2 and Corollary 3.3]). Let  $\underline{r} \in \{1, \ldots, r\}$ ,  $X \in \mathbb{R}_{\underline{r}}^{m \times n}$ , and

$$X = \begin{bmatrix} U \ U_{\perp} \end{bmatrix} \begin{bmatrix} \Sigma & \\ & 0_{m-\underline{r} \times n-\underline{r}} \end{bmatrix} \begin{bmatrix} V \ V_{\perp} \end{bmatrix}^{\top}$$

be an SVD. Then,

$$T_{\mathbb{R}^{m \times n}_{\leq r}}(X) = [U \ U_{\perp}] \begin{bmatrix} \mathbb{R}^{\underline{r} \times \underline{r}} & \mathbb{R}^{\underline{r} \times n - \underline{r}} \\ \mathbb{R}^{m - \underline{r} \times \underline{r}} & \mathbb{R}^{\underline{m} - \underline{r} \times n - \underline{r}} \\ \leq r - \underline{r} \end{bmatrix} [V \ V_{\perp}]^{\top}.$$

Moreover, for every  $G \in \mathbb{R}^{m \times n}$ ,

$$G = \begin{bmatrix} U \ U_{\perp} \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} V \ V_{\perp} \end{bmatrix}^{\top}$$

with  $A = U^{\top}GV$ ,  $B = U^{\top}GV_{\perp}$ ,  $C = U_{\perp}^{\top}GV$ , and  $D = U_{\perp}^{\top}GV_{\perp}$ , and

$$P_{T_{\mathbb{R}^{m \times n}_{\leq r}}(X)}(G) = \begin{bmatrix} U \ U_{\perp} \end{bmatrix} \begin{bmatrix} A & B \\ C & P_{\mathbb{R}^{m-\underline{r} \times n-\underline{r}}_{\leq r-\underline{r}}}(D) \end{bmatrix} \begin{bmatrix} V \ V_{\perp} \end{bmatrix}^{\top}.$$

Finally,

(6) 
$$\|\nabla f(X)\| \ge \mathbf{s}_f(X) \ge \sqrt{\frac{r-\underline{r}}{\min\{m,n\}-\underline{r}}} \|\nabla f(X)\|.$$

We are now going to review a fundamental result concerning  $s_f$ . To this end, we need to recall a basic definition from set-valued analysis. A correspondence, or a set-valued mapping, is a triple F := (A, B, G) where A and B are sets respectively called the set of departure and the set of destination of F, and G is a subset of  $A \times B$ called the graph of F. If F := (A, B, G) is a correspondence, written  $F : A \multimap B$ , then the image of  $x \in A$  by F is  $F(x) := \{y \in B \mid (x, y) \in G\}$ , and the domain of F is dom  $F := \{x \in A \mid F(x) \neq \emptyset\}$ .

PROPOSITION 2.8. For every differentiable function  $\phi : \mathbb{R}^{m \times n} \to \mathbb{R}$ : 1. the correspondence

$$\mathbb{R}^{m\times n}_{\leq r} \multimap \mathbb{R}^{m\times n}: X \mapsto P_{T_{\mathbb{R}^{m\times n}_{\leq r}}(X)}(-\nabla \phi(X))$$

depends on  $\phi$  only through its restriction  $\phi|_{\mathbb{R}^{m \times n}_{< r}}$ ;

2. if  $X \in \mathbb{R}_{\leq r}^{m \times n}$  is a local minimizer of  $\phi|_{\mathbb{R}_{< r}^{m \times n}}$ , then  $s_{\phi}(X) = 0$ .

*Proof.* The first part corresponds to [10, Theorem A.9(a)], and the second follows from [14, Theorem 6.12] and [10, Proposition A.6].

2.4. An upper bound on the distance to the determinantal variety from a tangent line. In this section, we look for an upper bound on  $d(X + G, \mathbb{R}_{\leq r}^{m \times n})$ holding for every  $X \in \mathbb{R}_{\leq r}^{m \times n} \setminus \{0_{m \times n}\}$  and every  $G \in T_{\mathbb{R}_{\leq r}^{m \times n}}(X)$ . A trivial bound is  $\|G\|$ , and [15, Proposition 3.6] tightens it to  $\frac{1}{\sqrt{2}}\|G\|$ . However, we will need an upper bound proportional to  $\|G\|^2$  in the proof of Proposition 3.1; such a bound is given in Proposition 2.9 and shown to be tight in Proposition A.1.

PROPOSITION 2.9. For every  $X \in \mathbb{R}_{\leq r}^{m \times n} \setminus \{0_{m \times n}\}$  and every  $G \in T_{\mathbb{R}_{< r}^{m \times n}}(X)$ ,

$$d(X+G, \mathbb{R}_{\leq r}^{m \times n}) \leq \frac{\sqrt{\operatorname{rank} X}}{2\sigma_{\min}(X)} \|G\|^2.$$

*Proof.* Let  $\underline{r} := \operatorname{rank} X$ . Let

$$X = \begin{bmatrix} U \ U_{\perp} \end{bmatrix} \begin{bmatrix} \Sigma & \\ & 0_{m-\underline{r} \times n-\underline{r}} \end{bmatrix} \begin{bmatrix} V \ V_{\perp} \end{bmatrix}^{\top}$$

be an SVD. By Proposition 2.7, there are  $A \in \mathbb{R}^{\underline{r} \times \underline{r}}$ ,  $B \in \mathbb{R}^{\underline{r} \times n - \underline{r}}$ ,  $C \in \mathbb{R}^{m - \underline{r} \times \underline{r}}$ , and  $D \in \mathbb{R}^{\underline{m} - \underline{r} \times n - \underline{r}}_{\leq r - \underline{r}}$  such that

$$G = \begin{bmatrix} U \ U_{\perp} \end{bmatrix} \begin{bmatrix} A & B \\ C & D \end{bmatrix} \begin{bmatrix} V \ V_{\perp} \end{bmatrix}^{\top}.$$

Let us define  $\gamma : [0, \infty) \to \mathbb{R}_{\leq r}^{m \times n}$  by

$$\gamma(t) := \left(U + t(U_{\perp}C + \frac{1}{2}UA)\Sigma^{-1}\right)\Sigma\left(V + t(V_{\perp}B^{\top} + \frac{1}{2}VA^{\top})\Sigma^{-1}\right)^{\top} + tU_{\perp}DV_{\perp}^{\top}$$

where the first term is inspired from [17, (13)];  $\gamma$  is well defined since the ranks of the

two terms are respectively upper bounded by  $\underline{r}$  and  $r - \underline{r}$ . For every  $t \in [0, \infty)$ ,

$$\gamma(t) = X + tG + \frac{t^2}{4} \begin{bmatrix} U & U_{\perp} \end{bmatrix} \begin{bmatrix} A \\ 2C \end{bmatrix} \Sigma^{-1} \begin{bmatrix} A & 2B \end{bmatrix} \begin{bmatrix} V & V_{\perp} \end{bmatrix}^{\top}$$

thus

$$d(X + tG, \mathbb{R}_{\leq r}^{m \times n}) \le \|(X + tG) - \gamma(t)\| = \frac{t^2}{4} \left\| \begin{bmatrix} A\Sigma^{-1}A & 2A\Sigma^{-1}B\\ 2C\Sigma^{-1}A & 4C\Sigma^{-1}B \end{bmatrix} \right\|.$$

Observe that

$$\begin{split} \left\| \begin{bmatrix} A\Sigma^{-1}A & 2A\Sigma^{-1}B \\ 2C\Sigma^{-1}A & 4C\Sigma^{-1}B \end{bmatrix} \right\|^2 \\ &= \|A\Sigma^{-1}A\|^2 + 4\|A\Sigma^{-1}B\|^2 + 4\|C\Sigma^{-1}A\|^2 + 16\|C\Sigma^{-1}B\|^2 \\ &\leq \|\Sigma^{-1}\|^2 \left(\|A\|^4 + 4\|A\|^2\|B\|^2 + 4\|A\|^2\|C\|^2 + 16\|B\|^2\|C\|^2\right) \\ &\leq \|\Sigma^{-1}\|^2\|G\|^4 \max_{\substack{x,y,z\in\mathbb{R}\\ x^2+y^2+z^2=1}} x^4 + 4x^2y^2 + 4x^2z^2 + 16y^2z^2 \\ &= 4\|\Sigma^{-1}\|^2\|G\|^4. \end{split}$$

Furthermore,

$$\|\Sigma^{-1}\| \le \frac{\sqrt{\underline{r}}}{\sigma_{\underline{r}}(X)}$$

Therefore, for every  $t \in [0, \infty)$ ,

$$d(X + tG, \mathbb{R}_{\leq r}^{m \times n}) \leq t^2 \frac{\sqrt{\underline{r}}}{2\sigma_{\underline{r}}(X)} \|G\|^2.$$

Choosing t = 1 yields the result.

**2.5.** Continuity of the restriction of the stationarity measure to any fixed-rank manifold. In this section, we prove that, for every  $\underline{r} \in \{1, \ldots, r\}$ , the restriction of  $s_f$  to  $\mathbb{R}_{\underline{r}}^{m \times n}$  is continuous; this will play a fundamental role in the convergence analysis conducted in Section 5. We refer to [12, §2.2] and the references therein for the definition of continuity of correspondences between metric spaces that appears in this paper only in the two following results.

PROPOSITION 2.10. Let  $G : \mathbb{R}^{m \times n} \to \mathbb{R}^{m \times n}$  be continuous,  $\mathcal{T} : \mathbb{R}^{m \times n} \multimap \mathbb{R}^{m \times n}$ be closed-valued, and  $\mathcal{S}$  be a nonempty subset of dom  $\mathcal{T}$ . If  $\mathcal{T}$  is continuous at  $X \in \mathcal{S}$ relative to  $\mathcal{S}$ , then the function

dom 
$$\mathcal{T} \to \mathbb{R} : Y \mapsto d(G(Y), \mathcal{T}(Y))$$

is continuous at X relative to  $\mathcal{S}$ .

*Proof.* Let  $X \in \mathcal{S}$ . For every  $Y \in \mathcal{S}$ ,

$$|d(G(X), \mathcal{T}(X)) - d(G(Y), \mathcal{T}(Y))| \le |d(G(X), \mathcal{T}(X)) - d(G(X), \mathcal{T}(Y))| + |d(G(X), \mathcal{T}(Y)) - d(G(Y), \mathcal{T}(Y))|$$

and, by [16, Proposition 1.3.17],

$$|d(G(X), \mathcal{T}(Y)) - d(G(Y), \mathcal{T}(Y))| \le ||G(X) - G(Y)||_{\mathcal{H}}$$

Let  $\varepsilon \in (0, \infty)$ . First, by [14, Proposition 5.11(c)], the function

$$\operatorname{dom} \mathcal{T} \to \mathbb{R} : Y \mapsto d(G(X), \mathcal{T}(Y))$$

is continuous at X relative to S. Thus, there exists  $\delta_1 \in (0, \infty)$  such that, for every  $Y \in B[X, \delta_1] \cap S$ ,

$$|d(G(X),\mathcal{T}(X)) - d(G(X),\mathcal{T}(Y))| \le \frac{\varepsilon}{2}.$$

Second, since G is continuous at X, there exists  $\delta_2 \in (0, \infty)$  such that, for every  $Y \in B[X, \delta_2]$ ,

$$\|G(X) - G(Y)\| \le \frac{\varepsilon}{2}.$$

Therefore, if  $\delta := \min\{\delta_1, \delta_2\}$ , then, for every  $Y \in B[X, \delta] \cap S$ ,

$$|d(G(X), \mathcal{T}(X)) - d(G(Y), \mathcal{T}(Y))| \le \varepsilon,$$

which completes the proof.

COROLLARY 2.11. For every  $\underline{r} \in \{1, \ldots, r\}$  and every continuously differentiable function  $\phi : \mathbb{R}^{m \times n} \to \mathbb{R}$ , the restriction  $s_{\phi}|_{\mathbb{R}^{m \times n}_{r}}$  is continuous.

*Proof.* Let  $X \in \mathbb{R}_{\underline{r}}^{m \times n}$ . We have to prove that  $s_{\phi}$  is continuous at X relative to  $\mathbb{R}_{\underline{r}}^{m \times n}$ . By Proposition 2.6 and [12, Theorem 4.1],  $T_{\mathbb{R}_{\leq r}^{m \times n}}(\cdot)$  is closed-valued and continuous at X relative to  $\mathbb{R}_{\underline{r}}^{m \times n}$ . Thus, by the preceding proposition, the function

$$\mathbb{R}_{\leq r}^{m \times n} \to \mathbb{R} : Y \mapsto d(-\nabla \phi(Y), T_{\mathbb{R}_{\leq r}^{m \times n}}(Y))$$

is continuous at X relative to  $\mathbb{R}_{\underline{r}}^{m \times n}$ . The result then follows from (5).

**2.6.** The determinantal variety has no serendipitous point. According to [10, Definition 2.8],  $X \in \mathbb{R}_{\leq r}^{m \times n}$  is said to be *serendipitous* if there exist a sequence  $(X_i)_{i \in \mathbb{N}}$  in  $\mathbb{R}_{\leq r}^{m \times n}$  converging to X, a continuously differentiable function  $\phi : \mathbb{R}^{m \times n} \to \mathbb{R}$ , and  $\varepsilon \in (0, \infty)$  such that  $s_{\phi}(X_i) > \varepsilon$  for every  $i \in \mathbb{N}$ , yet  $s_{\phi}(X) = 0$ . The next result will be invoked in the proof of Corollary 5.3.

PROPOSITION 2.12. No point of  $\mathbb{R}_{\leq r}^{m \times n}$  is serendipitous.

*Proof.* Let  $X \in \mathbb{R}_{\leq r}^{m \times n}$ . Let us prove that X is not serendipitous using [10, Theorem 2.17]. To this end, we need to recall two definitions for which we refer to [12, §2] and the references therein:

• for every nonempty subset S of  $\mathbb{R}^{m \times n}$ ,

$$\mathcal{S}^{-} := \{ Y \in \mathbb{R}^{m \times n} \mid \langle Y, Z \rangle \le 0 \; \forall Z \in \mathcal{S} \}$$

is a closed convex cone called the *(negative)* polar of S;

• for every sequence  $(\mathcal{S}_i)_{i\in\mathbb{N}}$  of nonempty subsets of  $\mathbb{R}^{m\times n}$ ,

$$\overline{\lim_{i \to \infty}} \, \mathcal{S}_i := \left\{ Y \in \mathbb{R}^{m \times n} \mid \liminf_{i \to \infty} d(Y, \mathcal{S}_i) = 0 \right\}$$

is a closed set called the *outer limit* of  $(\mathcal{S}_i)_{i \in \mathbb{N}}$ .

Let  $(X_i)_{i \in \mathbb{N}}$  be a sequence in  $\mathbb{R}_{\leq r}^{m \times n}$  converging to X. If rank X = r, then [12, Proposition 4.3] yields  $\overline{\lim}_{i \to \infty} T_{\mathbb{R}_{\leq r}^{m \times n}}(X_i) = T_{\mathbb{R}_{\leq r}^{m \times n}}(X)$ , and thus  $(\overline{\lim}_{i \to \infty} T_{\mathbb{R}_{\leq r}^{m \times n}}(X_i))^- = (T_{\mathbb{R}_{\leq r}^{m \times n}}(X))^-$ . If rank X < r, then

$$\left(T_{\mathbb{R}^{m\times n}_{\leq r}}(X)\right)^{-} = \{0_{m\times n}\} \subseteq \left(\overline{\lim_{i\to\infty}} T_{\mathbb{R}^{m\times n}_{\leq r}}(X_i)\right)^{-},$$

where the equality follows from [12, (25)].

3. Projected steepest descent with backtracking line search on  $\mathbb{R}_{\leq r}^{m \times n}$ . In this section, we analyze Algorithm 1—that corresponds to the main step of  $\mathbb{P}^2 \mathrm{GD}$  [15, Algorithm 3] except that the initial step size for the backtracking procedure is chosen in a given bounded interval—under the assumption that f is differentiable with  $\nabla f$  locally Lipschitz continuous. This will serve as a basis for the convergence analysis conducted in Section 5 since Algorithm 1 is used as a subroutine in Algorithm 2.

Algorithm 1 One step of P<sup>2</sup>GD (projected steepest descent with backtracking line search on  $\mathbb{R}^{m \times n}_{\leq r}$ )

**Require:**  $(f, X, \underline{\alpha}, \overline{\alpha}, \beta, c)$  where  $f : \mathbb{R}^{m \times n} \to \mathbb{R}$  is differentiable with locally Lipschitz continuous gradient,  $X \in \mathbb{R}_{\leq r}^{m \times n}$  is such that  $s_f(X) > 0, 0 < \underline{\alpha} < \overline{\alpha} < \infty$ , and  $\beta, c \in (0, 1)$ . 1: Choose  $G \in P_{T_{\mathbb{R}_{\leq r}^{m \times n}}(X)}(-\nabla f(X)), \alpha \in [\underline{\alpha}, \overline{\alpha}]$ , and  $Y \in P_{\mathbb{R}_{\leq r}^{m \times n}}(X + \alpha G)$ ; 2: while  $f(Y) > f(X) - c\alpha s_f(X)^2$  do 3:  $\alpha \leftarrow \alpha \beta$ ; 4: Choose  $Y \in P_{\mathbb{R}_{\leq r}^{m \times n}}(X + \alpha G)$ ; 5: end while 6: Return Y.

Following [10], the set of all possible outputs of Algorithm 1 is denoted by  $P^2GD(f, X, \underline{\alpha}, \overline{\alpha}, \beta, c)$  in the rest of the paper.

Let us recall that, since  $\nabla f$  is locally Lipschitz continuous, for every closed ball  $\mathcal{B} \subset \mathbb{R}^{m \times n}$ ,

$$\operatorname{Lip}_{\mathcal{B}}(\nabla f) := \sup_{\substack{X,Y \in \mathcal{B} \\ X \neq Y}} \frac{\|\nabla f(X) - \nabla f(Y)\|}{\|X - Y\|} < \infty$$

which implies, by [11, Lemma 1.2.3], that, for every  $X, Y \in \mathcal{B}$ ,

(7) 
$$|f(Y) - f(X) - \langle \nabla f(X), Y - X \rangle| \leq \frac{\operatorname{Lip}_{\mathcal{B}}(\nabla f)}{2} ||Y - X||^2.$$

PROPOSITION 3.1. Let  $X \in \mathbb{R}_{\leq r}^{m \times n}$  and  $\bar{\alpha} \in (0, \infty)$ . Let  $\mathcal{B} \subset \mathbb{R}^{m \times n}$  be a closed ball such that, for every  $G \in P_{T_{\mathbb{R}_{\leq r}^{m \times n}}(X)}(-\nabla f(X))$  and every  $\alpha \in [0, \bar{\alpha}]$ ,  $P_{\mathbb{R}_{\leq r}^{m \times n}}(X + \alpha G) \subset \mathbb{R}_{\leq r}^{m \times n}$ 

 $\mathcal{B}$ ; an example of such a ball is  $B[X, \rho(X)]$  with

$$\underline{\rho}(X) := \begin{cases} \bar{\alpha} \mathbf{s}_f(X) & \text{if } X = \mathbf{0}_{m \times n} \\ (1 + \frac{1}{\sqrt{2}}) \bar{\alpha} \mathbf{s}_f(X) & \text{otherwise.} \end{cases}$$

Then, for every  $G \in P_{\mathbb{T}_{\mathbb{R}^{m \times n}_{< r}}(X)}(-\nabla f(X))$  and every  $\alpha \in [0, \bar{\alpha}]$ ,

(8) 
$$\sup f(P_{\mathbb{R}^{m \times n}_{\leq r}}(X + \alpha G)) \leq f(X) + \mathrm{s}_f(X)^2 \alpha \left(-1 + \kappa_{\mathcal{B}}(f, X, \bar{\alpha})\alpha\right),$$

where

$$\kappa_{\mathcal{B}}(f, X, \bar{\alpha}) := \begin{cases} \frac{1}{2} \operatorname{Lip}(\nabla f) & \text{if } X = 0_{m \times n}, \\ \frac{\mathcal{B}}{2\sigma_{\min}(X)} \|\nabla f(X)\| + \frac{1}{2} \operatorname{Lip}(\nabla f) \left(\frac{\sqrt{\operatorname{rank} X}}{2\sigma_{\min}(X)} \bar{\alpha} \operatorname{s}_{f}(X) + 1\right)^{2} \text{otherwise.} \end{cases}$$

Proof. The example  $B[X, \underline{\rho}(X)]$  is correct by [15, Proposition 3.6]. The proof of (8) is based on (7) and the equality  $\langle \nabla f(X), G \rangle = -\mathbf{s}_f(X)^2$  given in the proof of [15, Proposition 2.6]. The result follows readily from (7) if  $X = 0_{m \times n}$  since  $T_{\mathbb{R}_{\leq r}^{m \times n}}(0_{m \times n}) = \mathbb{R}_{\leq r}^{m \times n}$ . Let us therefore consider the case where  $X \neq 0_{m \times n}$ . Let  $L := \operatorname{Lip}_{\mathcal{B}}(\nabla f)$ . For every  $Y \in P_{\mathbb{R}_{\leq r}^{m \times n}}(X + \alpha G)$ ,

$$\begin{split} &f(Y) - f(X) \\ &\leq \langle \nabla f(X), Y - X \rangle + \frac{L}{2} \| Y - X \|^2 \\ &= \langle \nabla f(X), Y - (X + \alpha G) + \alpha G \rangle + \frac{L}{2} \| Y - (X + \alpha G) + \alpha G \|^2 \\ &= -\alpha s_f(X)^2 + \langle \nabla f(X), Y - (X + \alpha G) \rangle + \frac{L}{2} \| Y - (X + \alpha G) + \alpha G \|^2 \\ &\leq -\alpha s_f(X)^2 + \| \nabla f(X) \| d(X + \alpha G, \mathbb{R}_{\leq r}^{m \times n}) + \frac{L}{2} \left( d(X + \alpha G, \mathbb{R}_{\leq r}^{m \times n}) + \alpha s_f(X) \right)^2 \\ &\leq -\alpha s_f(X)^2 + \| \nabla f(X) \| \frac{\sqrt{\operatorname{rank} X}}{2\sigma_{\min}(X)} \alpha^2 s_f(X)^2 + \frac{L}{2} \left( \frac{\sqrt{\operatorname{rank} X}}{2\sigma_{\min}(X)} \alpha^2 s_f(X)^2 + \alpha s_f(X) \right)^2 \\ &= \alpha s_f(X)^2 \left( -1 + \alpha \left( \| \nabla f(X) \| \frac{\sqrt{\operatorname{rank} X}}{2\sigma_{\min}(X)} + \frac{L}{2} \left( \frac{\sqrt{\operatorname{rank} X}}{2\sigma_{\min}(X)} \alpha s_f(X) + 1 \right)^2 \right) \right) \\ &\leq \alpha s_f(X)^2 \left( -1 + \alpha \kappa_{\mathcal{B}}(f, X, \bar{\alpha}) \right), \end{split}$$

where the third inequality follows from Proposition 2.9.

COROLLARY 3.2. Each point Y produced by Algorithm 1 satisfies the Armijo condition

$$f(Y) \le f(X) - c\alpha s_f(X)^2$$

for some  $\alpha \in \left[\min\{\underline{\alpha}, \beta \frac{1-c}{\kappa_{\mathcal{B}}(f, X, \bar{\alpha})}\}, \bar{\alpha}\right]$ , where  $\mathcal{B}$  is any closed ball as in Proposition 3.1. Proof. For all  $\alpha \in (0, \infty)$ ,

$$f(X) + s_f(X)^2 \alpha \left( -1 + \kappa_{\mathcal{B}}(f, X, \bar{\alpha}) \alpha \right) \le f(X) - c s_f(X)^2 \alpha \iff \alpha \le \frac{1 - c}{\kappa_{\mathcal{B}}(f, X, \bar{\alpha})}.$$

Since the left-hand side is an upper bound on  $f(P_{\mathbb{R}_{\leq r}^{m \times n}}(X + \alpha G))$  for every  $\alpha \in (0, \bar{\alpha}]$ , the Armijo condition is necessarily satisfied if  $\alpha \in (0, \min\{\bar{\alpha}, \frac{1-c}{\kappa_{\mathcal{B}}(f, X, \bar{\alpha})}\}]$ . Therefore, either the initial step size chosen in  $[\underline{\alpha}, \bar{\alpha}]$  satisfies the Armijo condition or the while loop ends with  $\alpha$  such that  $\frac{\alpha}{\beta} > \frac{1-c}{\kappa_{\mathcal{B}}(f, X, \bar{\alpha})}$ .

4. The proposed algorithm. We now introduce a first-order algorithm for low-rank optimization as Algorithm 2. At each iteration, this algorithm applies Algorithm 1 (i.e., one step of P<sup>2</sup>GD) to the current iterate Y but also, if rank<sub> $\Delta$ </sub> Y < rank Y, to  $\hat{Y}^{j} \in P_{\mathbb{R}_{\text{rank }Y-j}^{m \times n}}(Y)$  for every  $j \in \{1, \ldots, \text{rank }Y - \text{rank}_{\Delta}Y\}$ . Then, it forms the next iterate by choosing the result that decreases f the most; in particular, it produces sequences along which f is strictly decreasing. This will allow us to prove in the next section that it is apocalypse-free based on the following idea: whenever the current iterate Y is sufficiently close to a non-stationary point  $X \in \mathbb{R}_{\text{rank }X}^{m \times n}$ , Algorithm 2 will apply Algorithm 1 in particular to a projection of Y onto  $\mathbb{R}_{\text{rank }X}^{m \times n}$  which, by continuity of  $s_f|_{\mathbb{R}_{\text{rank }X}^{m \times n}}$  and  $\sigma_{\text{rank }X}$ , will produce a sufficient decrease in the cost function f in view of Corollary 3.2.

## Algorithm 2 $P^2GD$ with rank reduction

<b>Require:</b> $(f, X_0, \underline{\alpha}, \overline{\alpha}, \beta, c, \Delta)$ where $f : \mathbb{R}^{m \times n} \to \mathbb{R}$ is different values.	ferentiable with locally
Lipschitz continuous gradient, $X_0 \in \mathbb{R}^{m \times n}_{\leq r}, 0 < \underline{\alpha} < \overline{\alpha}$	$<\infty, \beta, c \in (0,1), \text{ and }$
$\Delta \in (0,\infty).$	
1: $i \leftarrow 0;$	
2: while $s_f(X_i) > 0$ do	
3: <b>for</b> $j \in \{0, \dots, \operatorname{rank} X_i - \operatorname{rank}_\Delta X_i\}$ <b>do</b>	$\triangleright$ See Definition 2.4.
4: Choose $\hat{X}_i^j \in P_{\mathbb{R}^{m \times n}_{\operatorname{rank} X_i - j}}(X_i);$	
5: Choose $\tilde{X}_i^j \in \mathrm{P}^2\mathrm{GD}(f, \hat{X}_i^j, \underline{\alpha}, \bar{\alpha}, \beta, c);$	$\triangleright$ See Algorithm 1.
6: end for	
7: Choose $X_{i+1} \in \operatorname{argmin}_{\{\tilde{X}_i^j   j \in \{0, \dots, \operatorname{rank} X_i - \operatorname{rank}_\Delta X_i\}\}} f;$	
8: $i \leftarrow i+1;$	
9: end while	

5. Convergence analysis. The purpose of this section is to prove Theorem 5.2. To this end, we will use the abstract framework proposed in [13, §1.3]. Indeed, the problem considered in this paper can be formulated as follows: find  $X \in \mathbb{R}_{\leq r}^{m \times n}$  such that  $s_f(X) = 0$ . It is thus a particular instance of [13, Abstract Problem 1] where the Banach space is  $\mathbb{R}^{m \times n}$ , its closed subset is  $\mathbb{R}_{\leq r}^{m \times n}$  and "desirable" means "stationary". Moreover, Algorithm 2 is a particular instance of [13, Algorithm Model 9] where the stop rule is f and the search function is Algorithm 3.

Algorithm 3 Search function of Algorithm 2

**Require:**  $(f, X, \underline{\alpha}, \overline{\alpha}, \beta, c, \Delta)$  where  $f : \mathbb{R}^{m \times n} \to \mathbb{R}$  is differentiable with locally Lipschitz continuous gradient,  $X \in \mathbb{R}_{\leq r}^{m \times n}$  is such that  $s_f(X) > 0, 0 < \underline{\alpha} < \overline{\alpha} < \infty$ ,  $\beta, c \in (0, 1)$ , and  $\Delta \in (0, \infty)$ . 1: for  $j \in \{0, \dots, \operatorname{rank} X - \operatorname{rank}_{\Delta} X\}$  do 2: Choose  $\hat{X}^j \in P_{\mathbb{R}_{\operatorname{rank} X-j}^{m \times n}}(X)$ ; 3: Choose  $\tilde{X}^j \in P^2 \operatorname{GD}(f, \hat{X}^j, \underline{\alpha}, \overline{\alpha}, \beta, c)$ ; 4: end for 5: Return  $Y \in \operatorname{argmin}_{\{\tilde{X}^j \mid j \in \{0, \dots, \operatorname{rank} X - \operatorname{rank}_{\Delta} X\}} f$ .

Thus, to prove Theorem 5.2, it suffices to verify that Algorithm 2 satisfies the two assumptions of [13, Theorem 10], which we do below.

The first assumption is that the objective function f is continuous at each nondesirable point or bounded from below on  $\mathbb{R}^{m \times n}_{\leq r}$ . It is thus satisfied since f is continuous. The following proposition exactly states that the second assumption is satisfied.

PROPOSITION 5.1. For every  $X \in \mathbb{R}_{\leq r}^{m \times n}$  such that  $s_f(X) > 0$ , there exist real numbers  $\varepsilon(X), \delta(X) > 0$  such that, for every  $Y \in B[X, \varepsilon(X)] \cap \mathbb{R}_{\leq r}^{m \times n}$  and every Z produced by Algorithm 3 applied to  $(f, Y, \underline{\alpha}, \overline{\alpha}, \beta, c, \Delta)$ ,

$$f(Z) - f(Y) \le -\delta(X)$$

Proof. Let  $X \in \mathbb{R}_{\leq r}^{m \times n}$  be such that  $s_f(X) > 0$ . Let us first consider the case where  $X = 0_{m \times n}$ . Let  $\varepsilon(0_{m \times n}) := \Delta$ . Then, for every  $Y \in B[0_{m \times n}, \varepsilon(0_{m \times n})] \cap \mathbb{R}_{\leq r}^{m \times n}$ , rank  $\Delta Y = 0$  since  $\sigma_1(Y) \leq ||Y|| \leq \varepsilon(0_{m \times n}) = \Delta$ . Therefore, Algorithm 3 will consider  $\hat{Y}^{\mathrm{rank}\,Y} = 0_{m \times n}$  and  $\hat{Y}^{\mathrm{rank}\,Y} \in \mathrm{P}^2\mathrm{GD}(f, 0_{m \times n}, \underline{\alpha}, \overline{\alpha}, \beta, c)$ . Thus, by Corollary 3.2,  $\delta(0_{m \times n}) := cs_f(0_{m \times n})^2 \min\{\underline{\alpha}, \frac{\beta(1-c)}{\kappa_{B[0_{m \times n}, \alpha]}(f, 0_{m \times n}, \overline{\alpha})}\}$  is a valid choice.

Let us now consider the case where  $X \neq \bar{0}_{m \times n}$ . Let  $\underline{r} := \operatorname{rank} X$ . By Corollary 2.11,  $\operatorname{sf}|_{\mathbb{R}_{\underline{r}}^{m \times n}}$  is continuous at X and thus there exists  $\rho(X) \in (0, \infty)$  such that  $\operatorname{sf}(B[X, \rho(X)] \cap \mathbb{R}_{\underline{r}}^{m \times n}) \subseteq [\frac{1}{2}\operatorname{sf}(X), \frac{3}{2}\operatorname{sf}(X)]$ . Let  $\varepsilon(X) := \min\{\Delta, \frac{1}{3}\sigma_{\underline{r}}(X), \frac{1}{2}\rho(X)\}$ . By Proposition 2.5, for every  $Y \in B[X, \varepsilon(X)] \cap \mathbb{R}_{\leq r}^{m \times n}$ ,  $\operatorname{rank}_{\Delta} Y \leq \underline{r} \leq \operatorname{rank} Y$  and  $P_{\mathbb{R}_{\underline{r}}^{m \times n}}(Y) \subset B[X, 2\varepsilon(X)]$ . Thus,  $0 \leq \operatorname{rank} Y - \underline{r} \leq \operatorname{rank} Y - \operatorname{rank}_{\Delta} Y$  and Algorithm 3 will therefore consider  $\hat{Y}^{\operatorname{rank} Y - \underline{r}} \in P_{\mathbb{R}_{\underline{r}}^{m \times n}}(Y) \subset B[X, 2\varepsilon(X)] \cap \mathbb{R}_{\underline{r}}^{m \times n}$  and  $\tilde{Y}^{\operatorname{rank} Y - \underline{r}} \in P^2 \operatorname{GD}(f, \hat{Y}^{\operatorname{rank} Y - \underline{r}}, \underline{\alpha}, \bar{\alpha}, \beta, c)$ . Since  $2\varepsilon(X) \leq \rho(X), \frac{1}{2}\operatorname{sf}(X) \leq \operatorname{sf}(\hat{Y}^{\operatorname{rank} Y - \underline{r}}) \leq \frac{3}{2}\operatorname{sf}(X)$ . Let  $\bar{\rho}(X) := \frac{3}{2}\underline{\rho}(X) + 2\varepsilon(X)$ . For every  $\alpha \in [0, \bar{\alpha}]$  and every  $G \in P_{T_{\mathbb{R}_{\underline{s}r}^{m \times n}}(\hat{Y}^{\operatorname{rank} Y - \underline{r}} + \alpha G), \|Z - X\| \leq \|Z - \hat{Y}^{\operatorname{rank} Y - \underline{r}}\| + \|\hat{Y}^{\operatorname{rank} Y - \underline{r}} - X\| \leq \underline{\rho}(\hat{Y}^{\operatorname{rank} Y - \underline{r}}) + 2\varepsilon(X)$  and  $\underline{\rho}(\hat{Y}^{\operatorname{rank} Y - \underline{r}}) \leq \frac{3}{2}\underline{\rho}(X)$ . Thus, by Corollary 3.2,  $\tilde{Y}^{\operatorname{rank} Y - \underline{r}}$  satisfies the Armijo condition with a step size at least  $\min\{\underline{\alpha}, \frac{\beta(1-c)}{\kappa_{B[X,\bar{\rho}(X)]}(f, \hat{Y}^{\operatorname{rank} Y - \underline{r}, \overline{\alpha})}\}$ .

$$\begin{aligned} |\nabla f(\hat{Y}^{\operatorname{rank} Y - \underline{r}})|| &\leq ||\nabla f(X)|| + L||X - \hat{Y}^{\operatorname{rank} Y - \underline{r}}|| \\ &\leq ||\nabla f(X)|| + 2L\varepsilon(X) \\ &\leq ||\nabla f(X)|| + \frac{2}{3}L\sigma_{\underline{r}}(X), \end{aligned}$$

and that, by Proposition 2.3,

$$\sigma_{\underline{r}}(\hat{Y}^{\operatorname{rank}Y-\underline{r}}) \ge \sigma_{\underline{r}}(X) - \|X - \hat{Y}^{\operatorname{rank}Y-\underline{r}}\| \ge \sigma_{\underline{r}}(X) - 2\varepsilon(X) \ge \frac{1}{3}\sigma_{\underline{r}}(X).$$

Thus,

$$\begin{aligned} \kappa_{B[X,\bar{\rho}(X)]}(f,\hat{Y}^{\operatorname{rank}Y-\underline{r}},\bar{\alpha}) &\leq \bar{\kappa}(f,X,\bar{\alpha}) \\ &:=\sqrt{\underline{r}}\left(\frac{3\|\nabla f(X)\|}{2\sigma_{\underline{r}}(X)} + L\right) + \frac{L}{2}\left(\frac{9\bar{\alpha}\sqrt{\underline{r}}\mathbf{s}_{f}(X)}{4\sigma_{\underline{r}}(X)} + 1\right)^{2}.\end{aligned}$$

It follows that

$$\delta(X) := c \min\left\{\underline{\alpha}, \frac{\beta(1-c)}{\bar{\kappa}(f, X, \bar{\alpha})}\right\} \frac{\mathbf{s}_f(X)^2}{4}$$

is a valid choice.

We have thus proved the following.

THEOREM 5.2. Consider the sequence constructed by Algorithm 2. If this sequence is finite, then its last element is stationary. If it is infinite, then each of its accumulation points is stationary, i.e., is a zero of the stationarity measure  $s_f$  defined in (3).

COROLLARY 5.3. Assume that Algorithm 2 produces a sequence  $(X_i)_{i\in\mathbb{N}}$ . The sequence has at least one accumulation point if and only if  $\liminf_{i\to\infty} ||X_i|| < \infty$ . For every convergent subsequence  $(X_{i_k})_{k\in\mathbb{N}}$ ,  $\lim_{k\to\infty} s_f(X_{i_k}) = 0$ . If  $(X_i)_{i\in\mathbb{N}}$  is bounded, which is the case notably if the sublevel set  $\{X \in \mathbb{R}^{m\times n}_{\leq r} \mid f(X) \leq f(X_0)\}$  is bounded, then  $\lim_{i\to\infty} s_f(X_i) = 0$ , and all accumulation points have the same image by f.

*Proof.* The "if and only if" statement is a classical result. The two limits follow from Proposition 2.12. The final claim follows from the argument given in the proof of [13, Theorem 65].  $\Box$ 

6. Conclusion. We close this work with four concluding remarks.

- 1. As in [10], everything said in this paper remains true if f is only defined on an open subset of  $\mathbb{R}^{m \times n}$  containing  $\mathbb{R}_{\leq r}^{m \times n}$ .
- 2. It is possible to prove Proposition 5.1 without relying on Corollary 2.11. Indeed, using the same notation, if rank X = r, then  $s_f$  is continuous at X since  $\mathbb{R}_{\leq r}^{m \times n}$  is identical to the smooth manifold  $\mathbb{R}_r^{m \times n}$  around X, and  $s_f$  therefore coincides with the norm of the Riemannian gradient of  $f|_{\mathbb{R}_r^{m \times n}}$ , which is continuous. If rank X < r, then, in view of (6) and the continuity of  $\nabla f$ ,  $s_f$  is bounded away from zero on the intersection of  $\mathbb{R}_{< r}^{m \times n}$  and a sufficiently small ball centered at X. However, we chose to keep the proof of Proposition 5.1 as it stands because it allows us to treat the cases where rank X = r and rank X < r together.
- 3. In many practical situations, when  $\Delta$  is chosen reasonably small, all the iterates of Algorithm 2 satisfy  $\operatorname{rank}_{\Delta} X_i = \operatorname{rank} X_i$ . In this case, the range of values of j in the for-loop of Algorithm 2 always reduces to  $\{0\}$ , and Algorithm 2 generates the same sequence  $(X_i)_{i \in \mathbb{N}}$  as P<sup>2</sup>GD. In this scenario, the only computational overhead in Algorithm 2 is the computation of rank  $X_i$  and  $\operatorname{rank}_{\Delta} X_i$ . However, for all  $i \geq 1$ , in view of line 4 of Algorithm 1, it is reasonable to assume that  $X_i$  has been obtained by a truncated SVD, in which case rank  $X_i$  and  $\operatorname{rank}_{\Delta} X_i$  are immediately available, making the overhead

12

insignificant. In summary, Algorithm 2 offers stronger convergence properties than  $P^2GD$ , and while incurring an insignificant overhead in many practical situations.

4. However, the range of values of j in the for-loop of Algorithm 2 can also be as large as  $\{0, \ldots, r\}$ , and there are situations where this occurs each time the for-loop is reached (e.g., in the case of a bounded sublevel set, when  $\Delta$  is chosen so large that rank $\Delta X = 0$  for all X in the sublevel set). One can thus wonder if is possible to restrict (conditionally or not) the range of values of jwhile preserving the apocalypse-free property of Algorithm 2. This is an open question. We can just recall that the bound given in Proposition 2.9 is tight (Proposition A.1), and observe that, in the neighborhood of any  $X \in \mathbb{R}_{< r}^{m \times n}$ ,  $\sigma_j$  can be arbitrarily small for every  $j \in \{\operatorname{rank} X + 1, \ldots, r\}$ . Hence it seems unlikely that Algorithm 2 with a restricted for-loop can be analyzed along the lines of Section 5. On the other hand, should the answer of the open question be negative, another counterexample as the one of [10, §2.2] would be required in view of [10, Remark 2.11].

Appendix A. Complementary results. The following proposition shows that the  $\frac{1}{\sigma_{\min}}$  factor cannot be removed in the upper bound of Proposition 2.9.

PROPOSITION A.1. For all  $\varepsilon \in (0, \infty)$ , there exist  $X \in \mathbb{R}^{m \times n}_r$  and  $G \in T_{\mathbb{R}^{m \times n}_{\leq r}}(X)$  such that

$$d(X+G,\mathbb{R}^{m\times n}_{\leq r}) \ge \left(\frac{1}{2\sigma_{\min}(X)} - \varepsilon\right) \|G\|^2.$$

*Proof.* Let  $\varepsilon \in (0,\infty)$ ,  $\sigma := \frac{1}{4\varepsilon}$ ,

$$\begin{aligned} X &:= \sigma \operatorname{diag}(2I_{r-1}, \begin{bmatrix} 1 & \\ 0 \end{bmatrix}, 0_{m-r-1 \times n-r-1}), \\ G &:= \sigma \operatorname{diag}(0_{r-1 \times r-1}, \begin{bmatrix} 1 & \\ 1 \end{bmatrix}, 0_{m-r-1 \times n-r-1}) \end{aligned}$$

Then,  $\sigma_{\min}(X) = \sigma$  and

$$X + G = \sigma \operatorname{diag}(2I_{r-1}, \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, 0_{m-r-1 \times n-r-1})$$

The nonzero singular values of X + G are the absolute values of the eigenvalues of  $\sigma \operatorname{diag}(2I_{r-1}, \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix})$ , i.e.,  $2\sigma$ , with multiplicity r-1,  $\frac{\sqrt{5}+1}{2}\sigma$ , and  $\frac{\sqrt{5}-1}{2}\sigma$ . Thus,  $d(X+G, \mathbb{R}_{\leq r}^{m \times n}) = \frac{\sqrt{5}-1}{2}\sigma$ ,  $\|G\| = \sqrt{2}\sigma$ , and

$$\frac{d(X+G,\mathbb{R}_{\leq r}^{m\times n})}{\|G\|^2} = \frac{\sqrt{5}-1}{4\sigma} = \frac{\sqrt{5}}{4\sigma} - \varepsilon > \frac{1}{2\sigma} - \varepsilon,$$

which completes the proof.

As an aside, we point out that Proposition A.1 can be deduced from geometric principles. Let  $\gamma$  be a curve on the submanifold  $\mathbb{R}_r^{m \times n}$  of  $\mathbb{R}^{m \times n}$ . In view of the Gauss formula along a curve [8, Corollary 8.3], the normal part of the acceleration of  $\gamma$  is given by  $\mathbb{I}(\gamma', \gamma')$ , where  $\mathbb{I}$  denotes the second fundamental form. In view of [2, §4], the largest principal curvature of  $\mathbb{R}_r^{m \times n}$  at X is  $1/\sigma_r(X)$ ; hence  $\|\mathbb{I}(\gamma'(t), \gamma'(t))\| \leq \|\gamma'(t)\|^2/\sigma_r(\gamma(t))$ , and the bound is attained when  $\gamma'(t)$  is along the corresponding principal direction.

PROPOSITION A.2. There exists no rank-related retraction [17, Definition 2] such that [17, Assumption 6] holds for every analytic cost function  $f : \mathbb{R}^{m \times n} \to \mathbb{R}$ .

*Proof.* Let  $\tilde{R} : \mathbb{R}^{2 \times 2} \times \mathbb{R}^{2 \times 2} \to \mathbb{R}^{2 \times 2}$  be a rank-related retraction [17, Definition 2], where we have identified the tangent bundle of  $\mathbb{R}^{2\times 2}$  with  $\mathbb{R}^{2\times 2} \times \mathbb{R}^{2\times 2}$ . Let us prove that, for the determinantal variety  $\mathbb{R}^{2\times 2}_{\leq 1}$  and the cost function  $f: \mathbb{R}^{2\times 2} \to \mathbb{R}: X \to \tilde{X}$  $X_{2,2}$ , the lifted function  $f \circ \hat{R}$  does not satisfy [17, Assumption 6]. For  $X_* := 0_{2 \times 2}$ , let  $\delta_{X_*}$  and  $\mathcal{U}_*$  be respectively the positive real number and the neighborhood of  $X_*$ in  $\mathbb{R}_{\leq 1}^{2\times 2}$  that correspond to  $X_*$  in [17, Definition 2]. Let  $\beta_{\mathrm{RL}} \in (0, \infty)$ ,  $\delta_{\mathrm{RL}} \in (0, \delta_{X_*})$ , and  $\mathcal{U} \subseteq \mathcal{U}_*$  be a neighborhood of  $X_*$  in  $\mathbb{R}^{2\times 2}_{\leq 1}$ . Let  $\xi_1, \xi_2 \in (0,1)$  be such that  $\xi_1^2 + 2\xi_2^2 = 1$ . Let  $\sigma \in (0, \frac{\xi_2^2}{3\beta_{\mathrm{RL}}})$  be such that  $X := \operatorname{diag}(\sigma, 0) \in \mathcal{U}$ . Let  $\tilde{R}_X : \mathbb{R}^{2 \times 2} \to \mathbb{R}^2$  $\mathbb{R}^{2\times 2}: \xi \mapsto \tilde{R}(X,\xi). \text{ Then, } \xi := \begin{bmatrix} -\xi_1 & \xi_2 \\ \xi_2 & 0 \end{bmatrix} \in T_{\mathbb{R}^{2\times 2}_{<1}}(X), \ \|\xi\| = 1, \text{ the update-rank } [17, \mathbb{R}^{2\times 2}_{<1}]$ Definition 1] of  $\xi$  is 1, and the following properties hold:  $\tilde{R}_X(0_{2\times 2}) = X$ , the function  $[0, \delta_{X_*}) \to \mathbb{R}^{2\times 2} : t \mapsto \tilde{R}_X(t\xi)$  is continuously differentiable, its image is contained in  $\mathbb{R}^{2\times 2}_{\leq 1}$ , and  $\frac{\mathrm{d}}{\mathrm{d}t}\tilde{R}_X(t\xi)|_{t=0} = \xi$ . Since the function  $[0, \delta_{X_*}) \to \mathbb{R}^{2\times 2} : t \mapsto \tilde{R}_X(t\xi)$ is continuous and  $(\tilde{R}_X(t\xi))_{1,1} = \sigma$ , there exists  $\tilde{\delta} \in (0, \delta_{\mathrm{RL}}]$  such that, for every  $t \in [0, \tilde{\delta}], (\tilde{R}_X(t\xi))_{1,1} \in [\frac{1}{2}\sigma, \frac{3}{2}\sigma].$  Therefore, for every  $t \in [0, \tilde{\delta}]$ , since det  $\tilde{R}_X(t\xi) = 0$ ,  $\tilde{R}_X(t\xi) = \begin{bmatrix} x(t) & y(t) \\ z(t) & \frac{y(t)z(t)}{x(t)} \end{bmatrix}, \text{ where } x : [0, \tilde{\delta}] \to \mathbb{R}, \ y : [0, \tilde{\delta}] \to \mathbb{R}, \text{ and } z : [0, \tilde{\delta}] \to \mathbb{R} \text{ are } t$ continuously differentiable, and such that  $x(0) = \sigma$ , y(0) = z(0) = 0,  $\dot{x}(0) = -\xi_1$ ,  $\dot{y}(0) = \dot{z}(0) = \xi_2$ , and  $x([0,\tilde{\delta}]) \subseteq [\frac{1}{2}\sigma, \frac{3}{2}\sigma]$ . Let  $\hat{f} : [0,\tilde{\delta}] \to \mathbb{R} : t \mapsto f(\tilde{R}_X(t\xi))$ . Then,  $\hat{f} = \frac{yz}{x}$  and  $\dot{\hat{f}} = \frac{y\dot{z}}{x} + \frac{\dot{y}z}{x} - \frac{\dot{x}yz}{x^2}$ . By continuity, there exists  $\delta \in (0, \tilde{\delta}]$  such that  $\dot{y}([0,\delta]), \dot{z}([0,\delta]) \subseteq [\frac{1}{2}\xi_2, \frac{3}{2}\xi_2], \text{ and } -\dot{x}([0,\delta]) \subseteq [\frac{1}{2}\xi_1, \frac{3}{2}\xi_1].$  Thus, since  $\dot{f}(0) = 0$ , for every  $t \in (0, \delta]$ ,

$$\frac{\left|\hat{f}(t) - \hat{f}(0)\right|}{t} = \frac{\hat{f}(t)}{t} \ge \frac{\xi_2^2}{3\sigma} + \frac{\xi_1\xi_2^2}{18\sigma^2}t > \frac{\xi_2^2}{3\sigma} > \beta_{\rm RL},$$

which completes the proof.

## REFERENCES

- C. ECKART AND G. YOUNG, The approximation of one matrix by another of lower rank, Psychometrika, 1 (1936), pp. 211–218, https://doi.org/https://doi.org/10.1007/BF02288367.
- [2] F. FEPPON AND P. F. J. LERMUSIAUX, A Geometric Approach to Dynamical Model Order Reduction, SIAM Journal on Matrix Analysis and Applications, 39 (2018), pp. 510–538, https://doi.org/https://doi.org/10.1137/16M1095202.
- N. GILLIS AND F. GLINEUR, Low-Rank Matrix Approximation with Weights or Missing Data Is NP-Hard, SIAM Journal on Matrix Analysis and Applications, 32 (2011), pp. 1149–1165, https://doi.org/https://doi.org/10.1137/110820361.
- [4] G. H. GOLUB AND C. F. VAN LOAN, *Matrix Computations*, Johns Hopkins Studies in the Mathematical Sciences, Johns Hopkins University Press, Baltimore, MD, USA, 4th ed., 2013.
- W. HA, H. LIU, AND R. FOYGEL BARBER, An Equivalence between Critical Points for Rank Constraints Versus Low-Rank Factorizations, SIAM Journal on Optimization, 30 (2020), pp. 2927–2955, https://doi.org/https://doi.org/10.1137/18M1231675.
- [6] J. HARRIS, Algebraic Geometry, vol. 133 of Graduate Texts in Mathematics, Springer-Verlag New York, 1992.
- U. HELMKE AND M. A. SHAYMAN, Critical Points of Matrix Least Squares Distance Functions, Liner Algebra and its Applications, 215 (1995), pp. 1–19, https://doi.org/https://doi.org/ 10.1016/0024-3795(93)00070-G.

- [8] J. M. LEE, Introduction to Riemannian Manifolds, vol. 176 of Graduate Texts in Mathematics, Springer, Cham, 2nd ed., 2018.
- [9] E. LEVIN, Towards optimization on varieties, master's thesis, Princeton University, September 2020.
- [10] E. LEVIN, J. KILEEL, AND N. BOUMAL, Finding stationary points on bounded-rank matrices: A geometric hurdle and a smooth remedy, tech. report, 2021, https://arxiv.org/abs/2107. 03877v1.
- [11] Y. NESTEROV, Lectures on Convex Optimization, vol. 137 of Springer Optimization and Its Applications, Springer, Cham, 2nd ed., 2018.
- [12] G. OLIKIER AND P.-A. ABSIL, On the continuity of the tangent cone to the determinantal variety, tech. report, UCLouvain, 2021, https://sites.uclouvain.be/absil/2021.06. Accepted for publication in Set-Valued and Variational Analysis: Theory and Applications.
- [13] E. POLAK, Computational Methods in Optimization, vol. 77 of Mathematics in Science and Engineering, Academic Press, 1971.
- [14] R. T. ROCKAFELLAR AND R. J.-B. WETS, Variational Analysis, vol. 317 of Grundlehren der mathematischen Wissenschaften, Springer-Verlag Berlin Heidelberg, 1998. Corrected 3rd printing 2009.
- [15] R. SCHNEIDER AND A. USCHMAJEW, Convergence Results for Projected Line-Search Methods on Varieties of Low-Rank Matrices Via Lojasiewicz Inequality, SIAM Journal on Optimization, 25 (2015), pp. 622–646, https://doi.org/https://doi.org/10.1137/140957822.
- [16] M. WILLEM, Functional Analysis: Fundamentals and Applications, Cornerstones, Birkhäuser Basel, Basel, Switzerland, 2013.
- [17] G. ZHOU, W. HUANG, K. A. GALLIVAN, P. VAN DOOREN, AND P.-A. ABSIL, A Riemannian rank-adaptive method for low-rank optimization, Neurocomputing, 192 (2016), pp. 72–80, https://doi.org/https://doi.org/10.1016/j.neucom.2016.02.030.