### A hybrid estimation of distribution algorithm for joint stratification and sample allocation

Mervyn O'Luing,<sup>1</sup> Steven Prestwich,<sup>1</sup> S. Armagan Tarim<sup>2</sup>

#### Abstract

In this study we propose a hybrid estimation of distribution algorithm (HEDA) to solve the joint stratification and sample allocation problem. This is a complex problem in which each the quality of each stratification from the set of all possible stratifications is measured its optimal sample allocation. EDAs are stochastic black-box optimization algorithms which can be used to estimate, build and sample probability models in the search for an optimal stratification. In this paper we enhance the exploitation properties of the EDA by adding a simulated annealing algorithm to make it a hybrid EDA. Results of empirical comparisons for atomic and continuous strata show that the HEDA attains the bests results found so far when compared to benchmark tests on the same data using a grouping genetic algorithm, simulated annealing algorithm or hill-climbing algorithm. However, the execution times and total execution are, in general, higher for the HEDA.

**Keywords:** Hybrid estimation of distribution algorithm; Optimal stratification; Sample allocation; R software.

<sup>&</sup>lt;sup>1</sup> Insight Centre for Data Analytics, Department of Computer Science, University College Cork, Ireland. Email: mervyn.oluing@insight-centre.org, steven.prestwich@insight-centre.org

<sup>&</sup>lt;sup>2</sup> Cork University Business School, University College Cork, Ireland. Email: armagan.tarim@ucc.ie

### 1 Introduction

The joint determination of stratification and sample allocation designs is a complex problem in which each solution is a stratification of basic strata (either atomic or continuous). The quality of each solution is measured by the optimal sample size that can be allocated to this stratification and still meet the precision constraints set by the survey designer. This quality is evaluated by the Bethel-Chromy algorithm (Bethel, 1985; Chromy, 1987; Bethel, 1989) which is expensive (in computational terms). Previous contributions in this area includes work carried out by (Kozak et al., 2007; Keskintürk and Er, 2007; Benedetti et al., 2008; Baillargeon and Rivest, 2009, 2011; Ballin and Barcaroli, 2013; O'Luing et al., 2019; Ballin and Barcaroli, 2020; O'Luing et al., 2020, 2021).

In this paper, we propose a hybrid estimation of distribution algorithm (HEDA) to solve this problem. EDAs are stochastic black-box optimization algorithms which can be used to estimate, build and sample probability models in the search for an optimal solution. We, therefore, describe the search for the lowest cost stratification as that of black-box optimization, in order to tie the problem in with existing EDA literature.

Rather than other methodologies where the strata have already been determined (e.g. administrative strata or the cumulative root frequency method) before evaluating the optimal sample allocation - the basic premise of this problem is that the optimal solution is unknown. It is a non linear problem with a rugged search space for which there are many near-optimal sample allocations (or local minima) and also perhaps more than one (i.e. attainable through multiple stratifications) optimal sample allocation (global minimum).

However, we cannot see the sample allocation for each stratification in advance, and thus cannot say *a priori* which stratification provides the optimal allocation. We could say that we are dealing with a black box. The only sure way of determining the optimal stratification is to evaluate each solution. This is known as grid-search and is intractable for large problems. Indeed, for representative surveys, especially in official statistics, for any practical sized sampling frame the number of basic strata to be stratified (given that they may be derived from multiple auxiliary variables) can be quite large.

A black-box optimization procedure explores the search space by generating solutions, evaluating them, and processing the results of this evaluation in order to generate new promising solutions (Gonzalez-Fernandez and Soto, 2012). Such procedures tend to find local minima of varying proximity to a global minimum. However they tend to be faster alternatives to grid-search. Some are deterministic, e.g. the direct search approach of (Hooke and Jeeves, 1961) or the simplex method of (Nelder and Mead, 1965) and some stochastic, e.g. k-means (Hartigan and Wong, 1979), grouping genetic algorithm (John, 1975), simulated annealing (Kirkpatrick et al., 1983; Černý, 1985) and hill-climbing (Lin, 1965; Lin and Kernighan, 1973). The latter set of procedures provide a means of attaining a solution that is "good enough" in a computing time that is "small enough" (Sörensen and Glover, 2013).

That has been our focus in earlier papers in which we presented evolutionary (a grouping genetic algorithm -(O'Luing et al., 2019)) and local search (simulated annealing algorithm - (O'Luing et al., 2020) algorithms and explored multi-stage combinations of clustering algorithms with a hill-climbing algorithm ((O'Luing et al., 2021)). We have evaluated the performance of these algorithms by the solution quality and computation time taken to find a local minimum. In (O'Luing et al., 2020) and (O'Luing et al., 2021) we also considered training times. The performance has varied according to each data stratification problem.

To expand on the details mentioned above, we explore a new paradigm of evolutionary algorithms named estimation of distribution algorithms (EDAs) (Baluja, 1994; Mühlenbein and Paaß, 1996; Larrañaga and Lozano, 2001; Pelikan et al., 2003) in the context of this problem. In particular, we consider the application of a hybridised version of an EDA and compare it against the best results achieved so far with the algorithms discussed in (O'Luing et al., 2019), (O'Luing et al., 2020) and (O'Luing et al., 2021). Section 2 provides some background details on EDAs. Section 3 motivates the use of a HEDA for this problem. Section 4 summarises the objective function.

paper. Section 6 discusses the evaluation approach. Section 7 demonstrates how the EDA component of the HEDA works. Sections 8 and 9 give results of empirical comparisons of the HEDA for atomic and continuous strata. Sections 10 and 11 describe our conclusions and suggestions for further work.

#### 2 Estimation of distribution algorithms (EDAs)

EDAs, which are also known as probabilistic model-building genetic algorithms (PMB-GAs), belong to a family of evolutionary algorithms. They are, in effect, stochastic blackbox optimization algorithms which are characterized by iteratively estimating, building and sampling probability models in the search for optimal solutions.

To put it another way, EDAs work by selecting promising solutions from a population (similar to elitism in evolutionary algorithms) and building probabilistic models of those solutions. They then sample from the corresponding probability distributions to obtain new solutions (Lima et al., 2011). It is these characteristics of probabilistic model building and sampling from that model that most differentiates the EDA from other evolutionary algorithms such as the genetic algorithm.

If the model built in each generation captures the important features of selected solutions and generates new solutions with these features, then the EDA should be able to quickly converge to the optimum (Mühlenbein and Mahnig, 1999). Indeed, assuming that the population size is large enough to ensure reliable convergence (Harik et al., 1999; Goldberg, 2002), the EDA based on the probability model provides an efficient and reliable approach to solving many optimization problems (Hauschild and Pelikan, 2011).

Furthermore, as EDAs iteratively refine the probabilistic model using elite solutions, thereby increasing the probability of generating the global optimum, after a reasonable number of iterations, the procedure should locate the global optimum or its accurate approximation (Hauschild and Pelikan, 2011). Empirically, EDAs have been shown to outperform other optimisation techniques in problems such as the multi-objective knapsack problem (Shah and Reed, 2011), multi-objective monitoring network design (Kollat et al., 2008) or optimising cancer chemotherapy (Petrovski et al., 2006).

Basic EDAs use a simple probability model that has a fixed structure and learns the parameters for that model, e.g. the univariate marginal distribution algorithm (UMDA) (Mühlenbein and Paaß, 1996), population-based incremental learning (PBIL) (Baluja, 1994) and compact genetic algorithm (cGA) (Harik et al., 1999). Such EDAS are suitable for univariate problems.

On the other hand, for more complex problems there are EDAs with adaptive multivariate models such as Bayesian networks (BNs) (Pearl, 1988), which can model complex multivariate interactions. Examples of Bayesian EDAs include the Bayesian optimization algorithm (BOA) (Pelikan et al., 1999, 2003), the estimation of Bayesian networks algorithm (EBNA) Etxeberria (1999), and the learning factorized distribution algorithm (LFDA)(Mühlenbein and Mahnig, 1999).

For this problem, as we are dealing with mutually exclusive basic strata, we assume that the probability distribution of any basic stratum belonging to a particular stratum is independent of that for another basic stratum, and for this reason we will implement a univariate EDA based closely on the UMDA.

### **3** A hybrid estimation of distribution algorithm (HEDA)

Even though we pointed out in section 2 that EDAs should locate the global optimum or its near approximation in a reasonable amount of iterations, they have two main computational bottlenecks: (1) fitness evaluation and (2) model building. These must be addressed by efficiency enhancement techniques suitable for EDAs (Hauschild and Pelikan, 2011).

The latter bottleneck we have addressed by using the simple model building approach of the UMDA - thus avoiding the computational overhead of more complex model building. The former bottleneck, clearly, is not only an issue for the EDA as it relates not only to the size of the search space but also the complexity of the problem. However, we can address this using hybridisation (Hauschild and Pelikan, 2011). This is an efficiency technique the purpose of which is to speed up the performance of the EDA.

In many real-world applications, evolutionary algorithms are combined with other search optimization algorithms such as local search, tabu search, simulated annealing or hill-climbing algorithms. Such metaheuristic hybrids combine the advantages of population based exploration methods (which search regions of the search space) with trajectory methods (which optimise locally and are often very successful). Simulated annealing is one example of a trajectory method which we focus on in this paper.

Simulated annealing resembles hill-climbing in that it makes small adjustments to a candidate solution - keeping those which lead to an improvement in solution quality and gradually morphing the solution towards the closest local optimum. Hill-climbing algorithms generally attain good results especially when combined with population based search procedures (Lima et al., 2009).

As an example, to solve the minimal switching graph problem (Tang and Lau, 2005) combined the exploration properties of the UMDA and the exploitation properties of the hill-climbing algorithm into a hybrid EDA to find an optimal or near-optimal solution efficiently and effectively. They compared the hybrid algorithm with the UMDA and the hill-climbing algorithm separately, and found that the performance of the hybrid EDA is significantly better than both the UMDA and the hill-climbing algorithm.

Although, the literature on hybridisation of EDAs does not appear to have considered simulated annealing, we expect a hybrid EDA using simulated annealing to perform well. This is because hill-climbing solutions gravitate towards the nearest local minima, where they become trapped until the algorithm reaches a pre-defined end-point. On the other hand, simulated annealing allows for a probabilistic acceptance of inferior quality solutions, enabling escape from local minima and thus attaining better quality solutions than hill-climbing. As it relates to our problem, for the reasons outlined in this section and in section 2, combining an EDA with simulated annealing should generate less solutions and thus translate into calling the Bethel-Chromy evaluation algorithm less often to attain optimal or near-optimal solutions.

### 4 Objective function

A detailed consideration of the objective function is provided in (O'Luing et al., 2019). We provide an outline below:

$$\min n = \sum_{h=1}^{H} n_h$$
  
s.t.  $CV\left(\hat{T}_g\right) \le \varepsilon_g \quad (g = 1, \dots, G)$   
 $2 \le n_h \le N_h$ 

where n is the sample size,  $n_h$  is the sample size for stratum h,  $N_h$  is the number of units in stratum h and H is the number of strata.  $\hat{T}_g$  is the estimator of the population total for each one of G target variable columns. The upper limit of precision  $\varepsilon_g$  is expressed as a coefficient of variation CV for each  $\hat{T}_g$ . We use the Bethel-Chromy algorithm (Bethel, 1985, 1989; Chromy, 1987) to solve the allocation problem for a particular stratification.

### 5 Outline of the HEDA

We base our description of the hyrbid estimation of distribution aglorithm on that found in (Gonzalez-Fernandez and Soto, 2012). We start with a population of  $N_p$  solutions either generated at random, or generated with random permutations of a starting solution (accompanying that solution). The population is evaluated and the best solutions are selected. Any selection method can be used (e.g. roulette wheel, tournament selection, Boltzmann selection, etc.) however for our algorithm we use elitism. We construct a probability model of the best solutions. We then use that model to generate new solutions to replace those not selected at the elitism stage. After a tunable number of iterations a simulated annealing algorithm is applied to a tunable number of solutions from the population (for the experiments in sections 8 and 9 we apply simulated annealing to the top ranked solution). This process continues until a stopping rule has been reached.

Algorithm 1	Hybrid EDA
-------------	------------

- 1: Set  $i \leftarrow 1$  and generate initial population  $P_1$  of  $N_p$  solutions
- 2: Apply simulated annealing algorithm

 $\triangleright$  tunable conditions

- 3: Select  $N_{Elite}$  promising solutions  $S_i$  from  $P_i$
- 4: Construct a probability model  $M_i$  of the best solution from  $S_i$ .
- 5: Create a new population  $P_{i+1}$  by sampling a set of  $N_p N_{Elite}$  new solutions according to the distribution encoded by  $M_i$
- 6: set  $i \leftarrow i + 1$
- 7: If stopping criteria (number of iterations, I) not met repeat steps 2 to 6

#### 6 Evaluation approach

Each of the  $N_p$  solutions in the population are evaluated in a two step-approach using the aggrStrata function (Ballin and Barcaroli, 2020) and Bethel algorithm (Ballin and Barcaroli, 2020; De Meo and De Meo, 2009). The aggrStrata function obtains summary statistics (including point estimates) for the basic strata. The Bethel function uses those statistics to compute the optimal sample allocation for that stratification. We have coded these functions in C++ and integrated them, along with the remainder of the HEDA, into the R software language (R Core Team, 2021) using the Rcpp package (Eddelbuettel, 2013) in order to speed up computation times. Accordingly, the traditional set.seed() function in R which was useful for reproducbility of experiments is not easily applied in Rcpp meaning that some further work is required before we can reproduce results exactly.

Nonetheless, we found this two-step approach to be a more useful evaluation metric for the HEDA than the Bayesian Information Criterion (BIC) or total within sums of squares (TWSS) which are more suited to clustering problems and faster than the evaluation algorithm. We initially considered these as "surrogate" functions. However, solutions which attain better BIC or TWSS scores might not attain better sample allocations. This is because minimising TWSS eventually leads towards one cluster per atomic stratum whereas maximising BIC scores is useful for selecting the optimal number of clusters, but not necessarily the optimal arrangement of items within clusters.

# 7 Example of EDA component of the HEDA on the iris data set

To demonstrate how an estimation of distribution algorithm works we use the iris data set (Anderson, 1935; Fisher, 1936; R Core Team, 2021) and atomic strata. We select Petal Length and Petal Width as the target variables. We use Sepal Length and Species as auxiliary variables. We convert Sepal Length to a categorical variable with 3 bins using the k-means algorithm (Hartigan and Wong, 1979) and a seed of 1234. We use an upper coefficient of variation level of 0.05 for the target variables. The cross product of the categorical version of Sepal Length with Species results in 8 atomic strata. We group these 8 atomic strata into H strata, in this example H = 3 or H = 4, i.e. some of the solutions have three strata and some have four.

Atomic Stratum, $l$	Ν	M1	M2	<b>S</b> 1	S2
а	40	1.46	0.24	0.17	0.10
b	5	3.40	1.10	0.30	0.15
с	1	4.50	1.70	0.00	0.00
d	10	1.48	0.29	0.17	0.09
е	31	4.23	1.31	0.36	0.19
f	12	5.07	1.88	0.22	0.27
g	14	4.64	1.45	0.21	0.11
h	37	5.74	2.08	0.50	0.25

Table 7.1: Summary statistics for the L atomic strata

We initialise a population size of  $N_p$  solutions (in this case  $N_p = 5$ ) each of size L, i.e. the number of atomic strata, where l = 1, 2, ..., L. As can be seen in table 7.1 we initially use letters of the alphabet to differentiate between atomic strata, however in subsequent tables an integer denotes to which one of the H strata each atomic stratum belongs. Each solution is evaluated and ranked.

		Initi	al Po	opula		Quality	Rank		
3	2	2	3	2	1	2	2	19.19	3
3	2	4	3	2	1	2	3	46.77	5
3	2	4	1	2	1	2	4	14.48	2
3	2	4	3	2	1	1	4	10.65	1
3	3	2	3	2	1	2	2	30.30	4

**Table 7.2:** Population of size  $N_p$  with solution quality and rank

We create a new selected population of the best solutions (in this example we select 2 solutions) in a process that is analysus with elitism in evolutionary algorithms.

Table 7.3: Selected population of best elite solutions

Selected Population							
3	2	4	3	2	1	1	4
3	2	4	1	2	1	2	4

We then construct a probabilistic model of the solutions in the selected population with the aim of estimating the probability distribution for each of the H strata for each atomic stratum. The probability of a candidate solution (or vector, V) where ( $V = V_1, V_2, ..., V_L$ ) is the product of probabilities of individual strata for each atomic stratum:

$$p(V_1, V_2, ..., V_L) = p(V_1) \times p(V_2), ..., \times p(V_L)$$
(7.1)

where  $p(V_l)$  is the probability of variable  $(V_l = v_l)$ ,  $v_l$  is an integer between 1 and H and  $p(V_1, V_2, ..., V_L)$  is the probability of the candidate solution  $(V_1, V_2, ..., V_L)$  and H represents the number of strata which are labelled individually by an integer between 1 and H. The univariate model for the L variables consists of L vectors containing probabilities of an atomic stratum belonging to the different strata. The probabilities different strata sum to 1.

The joint probability distribution is factorized as a product of independent univariate marginal distributions, which are estimated from marginal frequencies (Paul and Iba, 2002):

$$p(V_l) = \frac{\sum_{j=1}^{N_{Elite}} \delta_j (V_l = v_l | N_{Elite}^{iter-1})}{N_{Elite}^{iter-1}}$$
(7.2)

where  $N_{Elite}^{iter-1}$  is the number of elite solutions from the previous iteration,  $\delta_j(V_l = v_l | N_{Elite}^{iter-1}) = 1$  if  $V_l = v_l$  in the  $j^{th}$  case of  $N_{Elite}^{iter-1}$ , and 0 otherwise.

**Table 7.4:** Model estimating the probability of each atomic stratum *l* belonging to each stratum *h* 

Stratum, $h$		Probability Model						
1	0	0	0	0.5	0	1	0.5	0
2	0	1	0	0	1	0	0.5	0
3	1	0	0	0.5	0	0	0	0
4	0	0	1	0	0	0	0	1
Total	1	1	1	1	1	1	1	1

We generate new solutions from this model to replace the non-elite solutions by sampling from that distribution. We evaluate the new solutions and rank all solutions by their quality. This offspring population returns a solution quality of 9.34 which is the global minimum (to find the global minimum we evaluated each of the 4,140 possible partitions of the 8 atomic strata).

	0	ffspr	ing	Quality	Rank				
3	2	4	3	2	1	1	4	10.65	2
3	2	4	1	2	1	2	4	14.48	3
3	2	4	3	2	1	1	4	10.65	2
3	2	4	3	2	1	2	4	9.34	1
3	2	4	1	2	1	2	4	14.48	3

Table 7.5: Offspring population with solution quality and rank

Normally the algorithm repeats the above process until a stopping criterion has been met.

The extreme probabilities — 0 and 1 — mean that unless some mitigating measure is introduced the atomic stratum l would remain fixed forever to stratum h at a probability of either 0 or 1, obstructing some regions of the search space (Dang et al., 2019). However, at advanced stages of the search, e.g. when using a starting solution, the stratifications are of good quality and atomic stratum may already be in the correct stratum (i.e. the stratum it would be allocated to in the optimal solution).

In the above experiment it is clear that for all solutions in the population there are a number of cases where there is only one stratum choice available for an atomic stratum (i.e. it is assigned to that stratum with a probability of 1). Here we are very close to a global minimum and we get there on the second step - without adjusting the probability of strata assignments. On a more general level however, the hybrid EDA uses mutation and add strata components (see (O'Luing et al., 2019)), as well as simulated annealing (see (O'Luing et al., 2020) which enable escape from this occurrence. The UMDA also has alternative approaches of avoiding 1,0 probabilities (such as capping the probability interval to between  $\frac{1}{L}$  and  $(1 - \frac{1}{L})$ ) e.g. (Doerr and Krejca, 2020), however, for our purposes they are not required.

#### 8 Empirical comparisons for atomic strata

#### 8.1 Background details

We compare the performance of the HEDA with the best solution qualities from the experiments we carried out on the same data sets, with the grouping genetic algorithm, simulated annealing algorithm and combinations of clustering algorithms with hill-climbing. We use the results from these experiments as a benchmark for comparing the HEDA. More details on these experiments are provided in O'Luing et al. (2019, 2020, 2021).

The data sets including target and auxiliary variables are summarised in table 8.1. We explain the meaning of each variable in table 1.1 in section 1.

Data set	Target Variables	Auxiliary Variables	Number of Records	Number of Atomic Strata
Swiss Municipalities	Surfacesbois, Airbat	POPTOT, HApoly	2,896	579
American Community Survey, 2015	HINCP, VALP, SMOCP, INSP	BLD, TEN, WKEXREL, WORKSTAT, HFL, YBL	619,747	123,007
Kiva Loans	terminmonths, lendercount, loanamount	sector, currency, activity, region, partnerid	614,361	84,897

Table 8.1: Summary by data set of the variables, number of records and atomic strata

Table 8.2 provides details of the best sample sizes and the corresponding algorithms from the aforementioned which they were attained with.

**Table 8.2:** Summary by data set of the sample size and evaluation time for the final stage in multi stage combinations of clustering algorithms with the simulated annealing algorithm and hill-climbing algorithm

Dataset	Initial Stage(s)	Final Stage	Sample Size	Execution Time	Total Execution Time
Swiss Municipalities	Kmeans	SAA	125.17	248.91	8,808.63
American Community Survey, 2015	SOM, FC	HC	9,065.45	6,298.81	6,298.81
Kiva Loans	FC	HC	6,326.35	6,018.61	6,018.61

However, as we have sought to develop algorithms that provide solutions that are good enough in a computing time that is small enough we also report evaluation (execution) and training (total execution) times to provide context to the results. In table 8.2, SAArefers to simulated annealing algorithm, SOM applies to self-organising map, FC relates to Fuzzy Clustering and HC indicates hill-climbing.

For comparison purposes execution time and total execution time relates to the final stage algorithms only. Note: there was no fine-tuning required for the hill-climbing algorithm. Details of the fine-tuning time for the initial stage solutions are available in (O'Luing et al., 2020) and (O'Luing et al., 2021).

#### 8.2 Hyperparameters

The hyperparameters used in the experiments for the HEDA are provided in Table 8.3.

Dataset	Iterations, I	SAA Frequency	Mutation Chance	Elitism Rate	Add Strata Factor	Temperature, T
Swiss Municipalities	3,000	100	0.000185221	0.20	0.00000146	0.000000027
American Community Survey, 2015	200	10	0.000000016	0.30	0.00000054	0.000655857
Kiva Loans	600	100	0.000038838	0.30	0.000000739	0.001400962
Dataset	Decrement Constant,DC	Max number of sequences, maxit	Length of sequence, J	% of L for maximum q value, Lmax%	Probability of New Stratum, P(H + 1)	Population Size, $N_p$
Swiss Municipalities	0.941411680	2	20,000	0.10	0.009183601	20
American Community Survey, 2015	0.815042754	20	5,000	0.01	0.000000609	20
Kiva Loans	0.954218273	2	10,000	0.02	0.000052373	20

**Table 8.3:** Hyperparameters for the hybrid estimation of distribution algorithm (HEDA)

The hyperparameters operate in this way, *Iterations*, I indicates the number of iterations the algorithm runs for. The algorithm stops at the last iteration. The HEDA builds and samples from a *probability model*,  $M_i$  once every iteration. The simulated annealing algorithm also runs after a tunable number of iterations, i.e. for the Swiss Municipalities and Kiva Loans experiments it runs once every 100 iterations whereas for the American Community Survey experiment it runs once every 10 iterations. The simulated annealing algorithm runs each time for a maximum number of sequences with the length of each sequence varying for each experiment. See (O'Luing et al., 2020) for more details on the hyperparameters for the simulated annealing algorithm.

The following hyperparameters are described in more detail in (O'Luing et al., 2019). However in brief, *Elitism rate* indicates the proportion of the ranked population that is carried over to the next iteration, *Mutation chance* indicates the probability of each atomic stratum being moved to another stratum every iteration. Similarly, *Add Strata Factor* is the probability of an atomic stratum moving to a new stratum. *Population size*,  $N_p$  is the number of solutions used for exploration and evolution each iteration.

The hyperparameters often need to be fine-tuned to suit the characteristics of a particular data set. We fine-tuned the hyperparameters for these experiments using the methodology on as described by (Bischla et al., 2017) in the R software language (R Core Team, 2021). Finally, although we have stated the hyperparameters, the solution quality attained with them in the HEDA may not be reproduced exactly on their first application. See table 2.1 for details of the hyperparameters, execution and total execution times in the training process.

#### 8.3 Results

The results of these experiments are provided in Table 8.4.

the multi stage combinations of clustering algorithms with the HEDA Data set Initial Stage(s) Final Stage Sample Size Execution Time Total Execution Time 122.39 47,045.87 Swiss Municipalities Kmeans HEDA 1,316.11 American Community Survey, 2015 SOM and FC HEDA 28,857.52 550,676.58 8,800.53 HEDA 6,246.74 53,006.37 3,292.97

**Table 8.4:** Summary by data set of the benchmark sample size and evaluation time for

We compare the sample sizes attained in table 8.2 and table 8.4 and present the results

Kmeans

below in table 8.5.

Kiva Loans

Table 8.5: Ratio comparison of the sample sizes for the benchmark results and the HEDA results

Data set	Benchmark results	HEDA Results	Ratio comparisons
Swiss Municipalities	125.17	122.39	0.98
American Community Survey, 2015	9,065.45	8,800.53	0.97
Kiva Loans	6,326.35	6,246.74	0.99

The ratio of the HEDA results to the benchmark results indicate that, although they are similar, the HEDA results are of better quality in all cases. However, as our focus has been on obtaining results that are good enough in a time that is small enough these results should be considered in the context of execution and total execution times.

Table 8.6 compares the execution times for the HEDA and those for the final stage algorithm in the benchmark results.

Table 8.6: Ratio comparison of the execution times and total execution times for the benchmark results and the HEDA results

	Benchmark Results	HEDA Results		Benchmark Results	HEDA Results	
Data set	Execution Time	Execution Time	Ratio Comparisons	Total Execution Time	Total Execution Time	Ratio Comparisons
Swiss Municipalities	248.91	1,316.11	5.29	8,808.63	47,045.87	5.34
American Community Survey, 2015	6,298.81	28,857.52	4.58	6,298.81	550,676.58	87.43
Kiva Loans	6,018.61	3,292.97	0.55	6,018.61	53,006.37	8.81

In brief the results indicate that overall, for these experiments, the HEDA takes longer to execute and fine-tune than the SAA for the Swiss Municipalities, or the combination of clustering algorithms with hill-climbing for the American Community Survey, 2015 and Kiva Loans experiments.

#### 9 Empirical comparisons for continuous strata

Using the same methodology applied in section 8 we ran experiments using the same data sets comparing the performance of the HEDA on the stratification of continuous strata with the benchmark results from experiments we described in (O'Luing et al., 2019), (O'Luing et al., 2020) and (O'Luing et al., 2021). As before we first provide a summary of the target and auxiliary variables in table 9.1.

Data set	Target variables Auxiliary variabl	Auxiliary variables							
Swiss Municipalities	Airbat Surfacesbois								
American Community Survey, 2015	HINCP	VALP	SMOCP	INSP					
US Census, 2000	HHINCOME								
Kiva Loans	term_in_months	lender_count	loan_amount						
UN Commodity Trade Statistics data	trade_usd								

Table 9.1: Target and auxiliary variables for the Continuous method

Likewise table 9.2 provides details of the best sample sizes and the corresponding algorithms from the aforementioned which they were attained with.

**Table 9.2:** Summary by data set of the sample size and evaluation time for the final stageof multi stage combinations of clustering algorithms with the hill-climbingalgorithm

Dataset	Initial Stage(s)	Final Stage	Sample Size	Execution Time	Total Execution Time
Swiss Municipalities	NG and EM	HC	110	38.61	38.61
American Community Survey, 2015	NG and FC	HC	2,753.55	28,632.09	28,632.09
	EN (	на	1 550.00		F 050 F0
Kiva Loans	EM	HC	1,752.82	7,370.53	7,370.53

#### 9.1 Hyperparameters

The hyperparameters used in the HEDA are provided below in Table 9.3.

Dataset	Iterations, I	SAA Frequency	Mutation Chance	Elitism Rate	Add Strata Factor	Temperature, T
Swiss Municipalities	1,000	100	0.0000008	0.2	0.0000000	0.0000058
American Community Survey, 2015	30	10	0.0000003	0.3	0.0000010	0.0000010
Kiva Loans	90	10	0.0000004	0.3	0.0000001	0.0000001
Dataset	Decrement Constant, DC	Max number of sequences, maxit	Length of sequence, J	% of L for maximum q value, Lmax%	$\begin{array}{l} {\rm Probability}\\ {\rm of}\\ {\rm New}\\ {\rm Stratum},\\ {\rm P}({\rm H}+1) \end{array}$	Population Size, N_p
Swiss Municipalities	0.6266293	10	3,000	0.0195154	0.0000377	20
American Community Survey, 2015	0.9849021	2	50,000	0.0113557	0.0000001	20
Kiva Loans	0.9844605	2	20,000	0.0156086	0.0000008	20

**Table 9.3:** Hyperparameters for the hybrid estimation of distribution algorithm (HEDA)

Table 3.1 outlines the hyperaparameters used, along with the execution and total execution times resulting from the training process.

#### 9.2 Results

The results of the HEDA experiments on continuous strata are provided in Table 9.4.

**Table 9.4:** Summary by data set of the sample size and evaluation time for the final stage in multi stage combinations of clustering algorithms with the HEDA

Data set	Initial $Stage(s)$	Final Stage	Sample Size	Execution Time	Total Execution Time
Swiss Municipalities	Kmeans	HEDA	104.93	207.11	6,792.40
American Community Survey, 2015	NG and FC	HEDA	2,619.06	22,663.52	247,314.63
Kiva Loans	NG and Kmeans	HEDA	1,576.80	8,791.12	88,951.97

As before, we compare the sample sizes attained in table 9.2 and table 9.4 and present the results below in table 9.5.

 Table 9.5: Ratio comparison of the sample sizes for the benchmark results and the HEDA results

Data set	Benchmark Results	HEDA Results	Ratio Comparisons
Swiss Municipalities	110	104.930419822273	0.95
American Community Survey, 2015	2,753.55	2619.06326917666	0.95
Kiva Loans	1,752.82	1,576.80	0.90

The ratio of the HEDA results to the benchmark results indicate that, the HEDA results are of better quality in all three experiments.

Table 9.6 indicates that the execution times and total execution times are, overall, significantly greater for the HEDA.

	Benchmark Results	HEDA Results		Benchmark Results	HEDA Results	
Data set	Execution Time	Execution Time	Ratio Comparisons	Total Execution Time	Total Execution Time	Ratio Comparisons
Swiss Municipalities	38.61	207.11	5.36	95.33	6,792.40	71.25
American Community Survey, 2015	28,632.09	22,663.52	0.79	41,740.96	247,314.63	5.92
Kiva Loans	7,370.53	8,791.12	1.19	7,428.64	88,951.97	11.97

**Table 9.6:** Ratio comparison of the execution times and total execution times for the benchmark results and the HEDA results

### **10** Conclusions

Although these experiments are intended to be demonstrative rather than exhaustive, the multi-stage combination of an initial solution with the HEDA generally attains better sample sizes than the combinations of initial solutions with either the SAA or HC algorithm which had attained the best results in earlier experiments (that also included the GGA).

The execution times and total execution are, generally, significantly higher for the HEDA than some of the best alternative performing algorithms (especially the clustering algorithms considered in (O'Luing et al., 2021)). However, it is worth recalling that the HC algorithm in the above experiments had converged on a solution and would not have attained better results without adjusting the stopping conditions for algorithm or adjusting the solution. We should also bear in mind that the HEDA has found the best results so far for these experiments, and the execution and total times for the benchmark algorithms to find solutions of that quality (or better), using either the same or alternative starting solution, have not been evaluated.

That said, the HEDA is useful for a survey designer with expert knowledge of the sampling frame and the stratification of basic strata (atomic and continuous)- for whom the requirement to fine-tune hyperparameters is mitigated by prior knowledge. It is also useful for survey designers for whom the fine-tuning times can be included in their computation budget.

Nonetheless, because of the combination of probability models, explorative and exploitative search we feel the HEDA is well placed to generally outperform when it comes to attaining results which are good enough in a time that is small enough.

#### 11 Further Work

in the case of the HEDA adjustments can be made in order to use a seed function for reproducibility of results. Optimised versions of the GGA, SAA and basic EDA can also be implemented in *Rcpp*. It might be useful to extend the functionality of the HEDA to the stratification approach used for spatial sampling by Ballin and Barcaroli (2020). Furthermore, as we kept the population size, and the number of solutions to which we applied simulated annealing to every iteration, the same for all experiments, there is scope to fine those hyperparameters. Lastly, we could investigate more complex techniques such as the Bayesian optimisation algorithm, the estimation of Bayesian networks algorithm, the learning factorized distribution algorithm or mutual information maximising input clustering (MIMIC) (De Bonet et al., 1997) on this problem.

#### 12 Acknowledgements

We acknowledge the useful discussions with Dr. Thierry Vallée of the Central Statistics Office in general with respect to the progress of the PhD, but also with respect to tweaking the *Rcpp* code outlined above.

This material is based upon work supported by the Insight Centre for Data Analytics and Science Foundation Ireland under Grant No. 12/RC/2289-P2 which is co-funded under the European Regional Development Fund. Also, this publication has emanated from research supported in part by a grant from Science Foundation Ireland under Grant number  $16/\mathrm{RC}/3918$  which is co-funded under the European Regional Development Fund.

## Bibliography

- Anderson, E. (1935). The irises of the gaspe peninsula. Bulletin of the American Iris society 59, 2–5.
- Baillargeon, S. and L.-P. Rivest (2009). A general algorithm for univariate stratification. International Statistical Review 77(3), 331–344.
- Baillargeon, S. and L.-P. Rivest (2011). The construction of stratified designs in r with the package stratification. Survey Methodology 37(1), 53–65.
- Ballin, M. and G. Barcaroli (2013). Joint determination of optimal stratification and sample allocation using genetic algorithm. *Survey Methodology* 39(2), 369–393.
- Ballin. М. G. Barcaroli (2020).and Optimization of sampling with the SamplingStrata package. Accessed May 2021.strata 3 https://barcaroli.github.io/SamplingStrata/articles/SamplingStrata.html
- Baluja, S. (1994). Population-based incremental learning. a method for integrating genetic search based function optimization and competitive learning. Technical report, Carnegie-Mellon Univ Pittsburgh Pa Dept Of Computer Science.
- Benedetti, R., G. Espa, and G. Lafratta (2008). A tree-based approach to forming strata in multipurpose business surveys. *Survey Methodology* 34(2), 195–203.
- Bethel, J. W. (1985). An optimum allocation algorithm for multivariate surveys. American Statistical Proceedings of the Survey Research Methods, Section, 209-212.
- Bethel, J. (1989). Sample allocation in multivariate surveys. Survey methodology 15(1), 47-57.
- Bischla, B., J. Richterb, J. Bossekc, D. Hornb, J. Thomasa, and M. Langb (2017). mlrmbo: A modular framework for model-based optimization of expensive black-box functions. *Gradient Boosting in Automatic Machine Learning: Feature Selection and Hyperparameter Optimization*, 36.

- Černý, V. (1985). Thermodynamical approach to the traveling salesman problem: An efficient simulation algorithm. Journal of optimization theory and applications 45(1), 41-51.
- Chromy, J. R. (1987). Design optimization with multiple objectives. *Proceedings of the* Survey Research Methods Section.
- De Bonet, J. S., C. L. Isbell, P. Viola, et al. (1997). Mimic: Finding optima by estimating probability densities. Advances in neural information processing systems, 424–430.
- De Meo, M. and M. M. De Meo (2009). Package 'bethel'.
- Dang, D.-C., P. K. Lehre, and P. T. H. Nguyen (2019). Level-based analysis of the univariate marginal distribution algorithm. *Algorithmica* 81(2), 668–702.
- Doerr, B. and M. S. Krejca (2020). The univariate marginal distribution algorithm copes well with deception and epistasis. In *Proceedings of the 2020 Genetic and Evolutionary Computation Conference Companion*, pp. 17–18.
- Eddelbuettel, E. (2013). Seamless R and C++ Integration with Rcpp ISBN 978-1-4614-6867-7 10.1007/978-1-4614-6868-4
- Etxeberria, R. (1999). Global optimization using bayesian networks. In Proc. 2nd Symposium on Artificial Intelligence (CIMAF-99).
- Fisher, R. A. (1936). The use of multiple measurements in taxonomic problems. *Annals of Eugenics* 7(2), 179–188.
- Goldberg, D. E. (2002). The design of innovation: Lessons from and for competent genetic algorithms, Volume 1. Springer.
- Gonzalez-Fernandez, Y. and M. Soto (2012). copulaedas: An r package for estimation of distribution algorithms based on copulas. arXiv preprint arXiv:1209.5429.
- Harik, G. R., F. G. Lobo, and D. E. Goldberg (1999). The compact genetic algorithm. *IEEE transactions on evolutionary computation* 3(4), 287–297.
- Harik, G., E. Cantú-Paz, D. E. Goldberg, and B. L. Miller (1999). The gambler's ruin problem, genetic algorithms, and the sizing of populations. *Evolutionary computa*tion 7(3), 231–253.
- Hartigan, J. A. and M. A. Wong (1979). Hybrid Evolutionary Algorithms for Graph Coloring.Algorithm as 136: A k-means clustering algorithm *Journal of the Royal Statistical Society.* Series C (Applied Statistics) 28(1), 100–108.
- Hauschild, M. and M. Pelikan (2011). An introduction and survey of estimation of distribution algorithms. *Swarm and evolutionary computation* 1(3), 111–128.
- Hooke, R. and T. A. Jeeves (1961). "direct search" solution of numerical and statistical problems. *Journal of the ACM (JACM)* 8(2), 212–229.

- Holland, J. (1975). Adaptation in natural and artificial systems, 1992. Ann Arbor, MI: University of Michigan Press.
- Keskintürk, T. and Ş. Er (2007). A genetic algorithm approach to determine stratum boundaries and sample sizes of each stratum in stratified sampling. Computational Statistics & Data Analysis 52(1), 53–67.
- Kirkpatrick, S., C. D. Gelatt, and M. P. Vecchi (1983). Optimization by simulated annealing. science 220(4598), 671–680.
- Kollat, J. B., P. M. Reed, and J. R. Kasprzyk (2008). A new epsilon-dominance hierarchical bayesian optimization algorithm for large multiobjective monitoring network design problems. Advances in Water Resources 31(5), 828–845.
- Kozak, M., M. R. Verma, and A. Zielinski (2007). Modern approach to optimum stratification: Review and perspectives. *Statistics in Transition-new series* 8(2), 223–250.
- Larrañaga, P. and J. A. Lozano (2001). *Estimation of distribution algorithms: A new tool for evolutionary computation*, Volume 2. Springer Science & Business Media.
- Lima, C. F., M. Pelikan, F. G. Lobo, and D. E. Goldberg (2009). Loopy substructural local search for the bayesian optimization algorithm. In *International Workshop on Engineering Stochastic Local Search Algorithms*, pp. 61–75. Springer.
- Lima, C. F., F. G. Lobo, M. Pelikan, and D. E. Goldberg (2011). Model accuracy in the bayesian optimization algorithm. *Soft Computing* 15(7), 1351–1371.
- Lin, S. (1965). Computer solutions of the traveling salesman problem. *Bell System Technical Journal* 44(10), 2245–2269.
- Lin, S. and B. W. Kernighan (1973). An effective heuristic algorithm for the travelingsalesman problem. *Operations research* 21(2), 498–516.
- Mühlenbein, H. and G. Paaß (1996). From recombination of genes to the estimation of distributions i. binary parameters. In *International conference on parallel problem* solving from nature, pp. 178–187. Springer.
- Mühlenbein, H. and T. Mahnig (1999). Fda-a scalable evolutionary algorithm for the optimization of additively decomposed functions. *Evolutionary computation* 7(4), 353–376.
- Mühlenbein, H. and T. Mahnig (2001). Evolutionary algorithms: From recombination to search distributions. In *Theoretical Aspects of Evolutionary Computing*, pp. 135–173. Springer.
- Nelder, J. A. and R. Mead (1965). A simplex method for function minimization. *The computer journal* 7(4), 308–313.

- O'Luing, M., S. Prestwich, and S. Armagan Tarim (2019). A grouping genetic algorithm for joint stratification and sample allocation designs. *Survey Methodology* 45(3), 513–531.
- O'Luing, M., S. Prestwich, and S. A. Tarim (2020). A simulated annealing algorithm for joint stratification and sample allocation designs. arXiv preprint arXiv:2011.13006.
- O'Luing, M., S. Prestwich, and S. A. Tarim (2021). Combining k-means type algorithms with hill climbing for joint stratification and sample allocation designs. *arXiv preprint* arXiv:2108.08038.
- Paul, T. K. and H. Iba (2002). Linear and combinatorial optimizations by estimation of distribution algorithms. In 9th MPS symposium on Evolutionary Computation, IPSJ, Japan, Volume 12.
- Pearl, J. (1988). Probabilistic reasoning in intelligent systems: networks of plausible inference. San Mateo, CA: Morgan Kaufmann.
- Pelikan, M., D. E. Goldberg, E. Cantú-Paz, et al. (1999). Boa: The bayesian optimization algorithm. In Proceedings of the genetic and evolutionary computation conference GECCO-99, Volume 1, pp. 525–532. Citeseer.
- Pelikan, M., D. E. Goldberg, and S. Tsutsui (2003). Hierarchical bayesian optimization algorithm: toward a new generation of evolutionary algorithms. In SICE 2003 Annual Conference (IEEE Cat. No. 03TH8734), Volume 3, pp. 2738–2743. IEEE.
- Petrovski, A., S. Shakya, and J. McCall (2006). Optimising cancer chemotherapy using an estimation of distribution algorithm and genetic algorithms. In *Proceedings of the 8th annual conference on Genetic and evolutionary computation*, pp. 413–418.
- R Core Team (2021). R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing.
- Robles, V., J. M. Peña, P. Larrañaga, M. S. Pérez, and V. Herves (2006). Ga-eda: A new hybrid cooperative search evolutionary algorithm. In *Towards a New Evolutionary Computation*, pp. 187–219. Springer.
- Shah, R. and P. Reed (2011). Comparative analysis of multiobjective evolutionary algorithms for random and correlated instances of multiobjective d-dimensional knapsack problems. *European Journal of Operational Research* 211(3), 466–479.
- Sörensen, K. and F. Glover (2013). Metaheuristics. *Encyclopedia of operations research* and management science 62, 960–970.
- Tang, M. and R. Lau (2005). A hybrid estimation of distribution algorithm for the minimal switching graph problem. In International Conference on Computational Intelligence for Modelling, Control and Automation and International Conference on Intelligent Agents, Web Technologies and Internet Commerce (CIMCA-IAWTIC'06), Volume 1, pp. 708–713.

# 1 Appendix

# 1 Descriptions of target and auxiliary variables for the HEDA experiment data sets

Data set	Variable	Description
SwissMunicipalities	Surfacebois	wood area.
${f SwissMunicipalities}$	Airbat	area with buildings.
${f SwissMunicipalities}$	POPTOT	total population
${f SwissMunicipalities}$	Hapoly	municipality area.
American Community Survey, 2015	HINCP	Household income (past 12 months)
American Community Survey, 2015	VALP	Property value
American Community Survey, 2015	SMOCP	Selected monthly owner costs
American Community Survey, 2015	INSP	Fire/hazard/flood insurance (yearly amount)
American Community Survey, 2015	BLD	Units in structure
American Community Survey, 2015	TEN	TEN
American Community Survey, 2015	WKEXREL	Work experience of householder and spouse
American Community Survey, 2015	WORKSTAT	Work status of householder or spouse in family households
American Community Survey, 2015	HFL	House heating fuel
American Community Survey, 2015	YBL	When structure first built
Kiva Loans	term_in_months	The duration for which the loan was disbursed in months
Kiva Loans	lender_count	The total number of lenders that contributed to this loan
Kiva Loans	loan_amount	The amount disbursed by the field agent to the borrower (USD)
Kiva Loans	sector	High level category, e.g. Food
Kiva Loans	currency	The currency in which the loan was disbursed
Kiva Loans	activity	More granular category, e.g. Fruits & Vegetables
Kiva Loans	region	Full region name within the country
Kiva Loans	partner_id	ID of partner organization

Table 1.1: Breakdown by data set of descriptions for target and auxiliary variables

# 2 Fine-tuning the hyperparameters for the hybrid Estimation of Distribution Algorithm - Atomic Strata

			Swi	ss Municipaliti	es		
Fine-tuning		SAA	Mutation	Elitism	Add	Temperature,	Decrement Constant,
iteration	Iterations, I	Frequency	Chance	Rate	Strata	T	DC
No.		Frequency	Chance	nate	Factor	T	ЪС
1	5,000	100	0.0004011	0.15	0.0000008	0.00000002	0.6933468
2	3,000	100	0.0009943	0.15	0.0000002	0.00000005	0.5736595
3	5,000	100	0.0002985	0.2	0.0000005	0.00000009	0.9229554
4	5,000	50	0.0004681	0.1	0.0000008	0.00000003	0.7242638
5	3,000	100	0.0001686	0.1	0.0000005	0.00000007	0.8150428
6	3,000	50	0.0000172	0.1	0.0000000	0.00000006	0.6266293
7	4,000	50	0.0006155	0.1	0.0000004	0.00000004	0.8724350
8	4,000	50	0.0005785	0.15	0.0000002	0.00000003	0.9842787
9	5,000	100	0.0008674	0.2	0.0000006	0.00000008	0.7707891
10	4,000	50	0.0007308	0.2	0.0000010	0.00000010	0.5456765
11	3,000	100	0.0007510	0.1	0.0000006	0.00000007	0.8459806
12	4,000	50	0.0007705	0.1	0.0000004	0.00000006	0.8436530
13	3,000	100	0.0001289	0.15	0.0000006	0.00000007	0.8117457
14	3,000	100	0.0001827	0.2	0.0000003	0.00000007	0.8184057
15	3,000	100	0.0001847	0.2	0.0000003	0.00000003	0.8292686
16	3,000	50	0.0001903	0.2	0.0000003	0.00000004	0.8295554
17	3,000	100	0.0001852	0.2	0.0000001	0.00000003	0.9414117
18	4,000	100	0.0001856	0.2	0.0000001	0.00000001	0.8659521
19	3,000	100	0.0001844	0.2	0.0000008	0.00000002	0.8288662
20	3,000	100	0.0001343	0.2	0.0000002	0.00000003	0.9526036
			% of				
	Max	Length	L	Probability			
Fine-tuning	number	of	for	of	Population	Sample	Execution
iteration	of sequences,	sequence,	maximum	New	Size,	Size	Time
No.	maxit	J	q value,	Stratum, P(H + 1)	N_p		
			Lmax%	F (II + I)			
1	1	10,000	0.0801744	0.0007444	20	128.98	840.66
2	2	30,000	0.0546600	0.0096341	20	131.21	3,752.61
3	1	20,000	0.0458527	0.0073244	20	128.19	1,314.53
4	1	30,000	0.0287394	0.0048503	20	129.06	5,527.23
5	2	20,000	0.0120236	0.0060850	20	133.58	1,353.60
6	2	10,000	0.0670922	0.0037592	20	126.99	886.40
7	2	20,000	0.0821999	0.0028418	20	125.84	3,697.88
8	2	30,000	0.0625123	0.0057328	20	127.06	7,791.48
9	1	20,000	0.0946376	0.0082044	20	126.42	1,367.88
10	1	10,000	0.0233130	0.0018504	20	130.34	659.67
11	2	20,000	0.0954986	0.0064802	20	125.10	1,304.16
12	2	20,000	0.0135087	0.0033416	20	130.00	3,382.07
13	2	20,000	0.0880779	0.0050088	20	126.38	1,379.07
14	2	20,000	0.0958774	0.0076491	20	122.92	1,316.11
15	2	20,000	0.0101602	0.0088849	20	138.17	1,240.32
16	2	30,000	0.0988187	0.0085451	20	125.56	5,776.53
17	2	20,000	0.0990880	0.0091836	20	122.39	1,337.31
18	2	20,000	0.0105801	0.0090396	20	133.49	1,587.33
19	2	20,000	0.0992476	0.0093288	20	128.43	1,319.65
20	2	20,000	0.0102580	0.0082125	20	133.65	1,211.38
					Minimum	122.39	
						Total	47,045.87
			American C	Community Sur	vey, 2015		

Table 2.1: Hyperparameters for the HEDA in the case of Atomic Strata

Fine-tuning		SAA	Mutation	Elitism	Add	Temperature,	Decrement Constant,
iteration	Iterations, I	Frequency	Chance	Rate	Strata	T	DC
No.		Frequency	Chance	nate	Factor	1	DC
1	300	10	0.0000000	0.4	0.0000001	0.00009222	0.6933468
2	200	10	0.0000001	0.4	0.0000000	0.00042573	0.5736595
3	300	10	0.0000000	0.5	0.0000000	0.00085991	0.9229554
4	300	5	0.0000000	0.3	0.0000001	0.00019888	0.7242638
5	200	10	0.0000000	0.3	0.0000001	0.00065586	0.8150428
6	200	5	0.0000000	0.3	0.0000000	0.00057615	0.6266293
7	200	5	0.0000001	0.3	0.0000000	0.00033515	0.8724350
8	300	5	0.0000001	0.4	0.0000000	0.00021361	0.9842787
9	300	10	0.0000001	0.5	0.0000001	0.00077701	0.7707891
10	200	5	0.0000001	0.5	0.0000001	0.00095672	0.5456765
11	200	10	0.0000000	0.3	0.0000001	0.00013141	0.8217492
12	300	10	0.0000000	0.5	0.0000001	0.00004408	0.7949279
13	300	10	0.0000001	0.3	0.0000001	0.00011081	0.7006825
14	300	10	0.0000000	0.4	0.0000001	0.00001201	0.8424332
15	300	10	0.0000000	0.5	0.0000001	0.00015274	0.8056020
16	300	10	0.0000001	0.3	0.0000001	0.00009295	0.6690758
17	200	10	0.0000000	0.4	0.0000001	0.00007020	0.7462928
18	300	10	0.0000000	0.3	0.0000001	0.00076447	0.8238593
19	300	10	0.0000001	0.5	0.0000001	0.00009108	0.6111786
20	200	10	0.0000001	0.4	0.0000001	0.00043680	0.7087167
			% of	<b>D</b>			
<b>D</b> ( )	Max	Length	L	Probability	<b>D</b>		
Fine-tuning	number	of	for	of	Population	Sample	Execution
iteration	of sequences,	sequence,	maximum	New	Size,	Size	Time
No.	maxit	J	q value,	Stratum,	N_p		
			Lmax%	P(H + 1)			
1	10	2,500	0.0216957	0.0000001	20	8,849.13	8,542.77
2	20	5,000	0.0174433	0.0000010	20	8,829.25	28,978.64
3	10	2,500	0.0159755	0.0000007	20	8,834.01	6,984.81
4	10	5,000	0.0131232	0.0000005	20	8,805.62	36,489.00
5	20	5,000	0.0103373	0.0000006	20	8,800.53	28,857.52
6	20	2,500	0.0195154	0.0000004	20	8,824.74	20,805.10
7	20	2,500	0.0220333	0.0000003	20	8,816.72	20,756.65
8	20	5,000	0.0187521	0.0000006	20	8,811.21	89,838.25
9	10	5,000	0.0241063	0.0000008	20	8,838.34	17,576.73
10	10	2,500	0.0122188	0.0000002	20	8,809.30	9,145.87
11	20	5,000	0.0110847	0.0000001	20	8,801.18	28,832.86
12	20	5,000	0.0100309	0.0000001	20	8,810.14	44,465.19
13	20	5,000	0.0143632	0.0000001	20	8,807.34	44,395.63
14	10	2,500	0.0109556	0.0000001	20	8,822.30	7,088.03
15	20	5,000	0.0217767	0.0000000	20	8,819.96	44,605.16
16	10	5,000	0.0154209	0.0000006	20	8,824.89	17,641.07
17	20	5,000	0.0153991	0.0000007	20	8,815.61	28,870.49
18	10	5,000	0.0111377	0.0000007	20	8,811.88	17,556.67
19	20	5,000	0.0105166	0.0000000	20	8,804.58	44,660.20
13	10	2,500	0.0103235	0.0000004	20	8,850.46	4,585.94
20			1		Minimum	8,800.53	
						Total	550,676.58
				Kiva Loans		Total	550,676.58
20				Kiva Loans	Add	Total	550,676.58
	Iterations, I	SAA	Mutation	Kiva Loans Elitism	Add Strata	Total Temperature, T	550,676.58 Decrement Constant, DC

	600	200	0.0000000	0.0	0.0000000	0.00100202	0.0022469
1	600	300	0.0000396	0.2	0.0000008	0.00100396	0.6933468
2	400	300	0.0000994	0.3	0.0000002	0.00430903	0.5736595
3	600	300	0.0000292	0.3	0.0000005	0.00861177	0.9229554
4	600	100	0.0000463	0.2	0.0000008	0.00206098	0.7242638
5	400	300	0.0000161	0.2	0.0000005	0.00658958	0.8150428
6	400	200	0.0000008	0.2	0.0000000	0.00579972	0.6266293
7	500	100	0.0000612	0.2	0.0000003	0.00341138	0.8724350
8	500	100	0.0000575	0.3	0.0000002	0.00220699	0.9842787
9	600	200	0.0000866	0.3	0.0000006	0.00779018	0.7707891
10	500	200	0.0000728	0.3	0.0000010	0.00957111	0.5456765
11	600	100	0.0000388	0.3	0.0000007	0.00140096	0.9542183
12	600	100	0.0000058	0.3	0.0000007	0.00025835	0.9846227
13	600	100	0.0000402	0.3	0.0000008	0.00950009	0.9444742
14	600	300	0.0000919	0.3	0.0000005	0.00164402	0.9690933
15	600	100	0.0000229	0.2	0.0000010	0.00040293	0.6803285
16	600	300	0.0000342	0.3	0.0000009	0.00065333	0.9685014
17	600	100	0.0000345	0.3	0.0000008	0.00030204	0.9093287
18	500	100	0.0000228	0.3	0.0000004	0.00848591	0.9809824
19	600	100	0.0000375	0.3	0.0000009	0.00109957	0.6936422
20	600	100	0.0000281	0.3	0.0000009	0.00166877	0.6873736
			% of	Probability			
Fine-tuning	Max	Length	L	of	Population		
iteration	number	of	for	New	Size,	Sample	Execution
No.	of sequences,	sequence,	maximum	Stratum,	N_p	Size	Time
	maxit	J	q value,	P(H + 1)	<u>-</u> F		
			Lmax%	,			
1	1	5,000	0.0216957	0.0000075	20	6,536.39	7,325.52
2	3	10,000	0.0174433	0.0000963	20	6,338.61	6,190.59
3	2	5,000	0.0159755	0.0000733	20	6,559.13	3,720.53
4	1	10,000	0.0131232	0.0000486	20	6,285.78	2,837.05
5	3	10,000	0.0103373	0.0000609	20	6,374.74	1,422.56
6	3	5,000	0.0195154	0.0000377	20	6,441.66	1,111.02
7	2	5,000	0.0220333	0.0000285	20	6,321.22	1,768.59
8	3	10,000	0.0187521	0.0000574	20	6,265.44	3,174.16
9	2	10,000	0.0241063	0.0000821	20	6,349.25	2,045.19
10							
11	1	5,000	0.0122188	0.0000186	20	6,447.99	1,280.92
ļ	2	5,000 10,000	0.0122188 0.0151998	0.0000186 0.0000524	20 20	6,447.99 6,246.74	1,280.92 3,292.97
12							
	2	10,000	0.0151998	0.0000524	20	6,246.74	3,292.97
12	2 2	10,000 5,000	0.0151998 0.0162442	0.0000524	20 20	6,246.74 6,287.11	3,292.97 2,154.07
12 13	2 2 2	10,000 5,000 5,000	0.0151998 0.0162442 0.0152621	0.0000524 0.0000388 0.0000566	20 20 20	6,246.74 6,287.11 6,439.26	3,292.97 2,154.07 2,109.17
12 13 14	2 2 2 2	10,000 5,000 5,000 5,000	0.0151998 0.0162442 0.0152621 0.0150536	0.0000524 0.0000388 0.0000566 0.0000973	20 20 20 20	6,246.74           6,287.11           6,439.26           6,466.34	3,292.97 2,154.07 2,109.17 1,400.07
12 13 14 15	2 2 2 2 1	10,000 5,000 5,000 5,000 10,000	0.0151998           0.0162442           0.0152621           0.0150536           0.0187736	0.0000524 0.0000388 0.0000566 0.0000973 0.0000071	20 20 20 20 20 20	6,246.74           6,287.11           6,439.26           6,466.34           6,250.33	3,292.97 2,154.07 2,109.17 1,400.07 2,703.06
12 13 14 15 16	2 2 2 2 1 2	10,000 5,000 5,000 5,000 10,000 10,000	0.0151998 0.0162442 0.0152621 0.0150536 0.0187736 0.0195352	0.0000524 0.0000388 0.0000566 0.0000973 0.0000071 0.0000050	20 20 20 20 20 20 20	6,246.74           6,287.11           6,439.26           6,466.34           6,250.33           6,365.74	3,292.97 2,154.07 2,109.17 1,400.07 2,703.06 1,649.00
12 13 14 15 16 17	2 2 2 1 2 1	10,000 5,000 5,000 5,000 10,000 10,000 5,000	0.0151998           0.0162442           0.0152621           0.0150536           0.0187736           0.0195352           0.0162050	0.0000524 0.0000388 0.0000566 0.0000973 0.0000071 0.0000050 0.0000050	20 20 20 20 20 20 20 20	6,246.74           6,287.11           6,439.26           6,466.34           6,250.33           6,365.74           6,313.38	3,292.97 2,154.07 2,109.17 1,400.07 2,703.06 1,649.00 1,888.96
12       13       14       15       16       17       18	2 2 2 2 1 2 1 1 1	10,000         5,000         5,000         5,000         10,000         10,000         5,000         10,000         10,000	0.0151998           0.0162442           0.0152621           0.0150536           0.0187736           0.0195352           0.0162050           0.0141372	0.0000524           0.0000388           0.0000566           0.0000973           0.0000071           0.0000050           0.0000050           0.0000098	20 20 20 20 20 20 20 20 20 20	6,246.74         6,287.11         6,439.26         6,466.34         6,250.33         6,365.74         6,313.38         6,428.67	3,292.97 2,154.07 2,109.17 1,400.07 2,703.06 1,649.00 1,888.96 2,138.39
12       13       14       15       16       17       18       19	2 2 2 1 2 1 1 1 1	10,000         5,000         5,000         5,000         10,000         10,000         5,000         10,000         10,000         10,000	0.0151998           0.0162442           0.0152621           0.0150536           0.0187736           0.0195352           0.0162050           0.0141372           0.0159892	0.0000524           0.0000388           0.0000566           0.0000973           0.0000071           0.0000050           0.0000050           0.00000678           0.0000338	20 20 20 20 20 20 20 20 20 20 20	6,246.74         6,287.11         6,439.26         6,466.34         6,250.33         6,365.74         6,313.38         6,428.67         6,250.00	3,292.97 2,154.07 2,109.17 1,400.07 2,703.06 1,649.00 1,888.96 2,138.39 2,677.30

# 3 Fine-tuning the hyperparameters for the hybrid Estimation of Distribution Algorithm - Continuous Strata

Functional branc         Functional Procession         Functional Processin <thf< th=""><th></th><th>-</th><th>1 1</th><th>Swi</th><th>ss Municipaliti</th><th>es</th><th>-</th><th></th></thf<>		-	1 1	Swi	ss Municipaliti	es	-	
21,0002000.0000940.30.00000230.00000430.573650533,0002000.00000230.20.00000640.0000210.72403851,0001000.00000080.20.00000030.00006840.851042861,0001000.00000080.20.00000030.00006840.85243072,0001000.00005750.30.00000200.0000220.85427893,0002000.00006860.30.00000080.0000780.770781102,0001000.0000180.30.00000080.30000780.5356765112,0001000.0000180.30.00000020.00000820.533399123,0001000.0000010.20.00000030.00000820.5326765133,0001000.0000010.20.00000030.00000820.532676141,0001000.0000010.20.00000030.00000820.532623152,0002000.0000080.20.00000030.6000680.545023161,0001000.0000080.20.00000080.545023171,0001000.0000080.20.00000880.654023181,0001000.0000080.20.00000880.654023191,0001000.0000080.20.00000880.654023191,0001000.0000080	iteration	Iterations, I				Strata		,
33,0002000,00002920.30,00000490,0000860,22953443,0001000,00001610,20,00000670,0000660,72433851,0001000,00000610,20,00000500,00003640,81504272,0001000,00001610,20,000000500,0000220,81504272,0001000,0000720,30,00000000,0000720,00000000,0000720,00000700,942787102,0001000,0000720,30,00000000,00000000,5057550,00000000,00000700,537057112,0001000,0000010,30,00000000,00000000,5303670,53386133,0001000,0000010,00000000,00000000,00000700,533869141,0001000,0000000,00000000,00000000,834874152,0002000,00000000,00000000,00000000,540575161,0001000,0000000,00000000,00000000,834874161,0001000,00000000,00000000,00000000,540575171,0001000,00000000,00000000,00000000,00000000,540575181,0001000,00000000,00000000,00000000,540575191,0001000,00000000,00000000,00000000,540575191,0001000,0000000<	1	3,000	200	0.0000396	0.2	0.00000082	0.0000010	0.6933468
4         3,000         100         0.000043         0.2         0.000076         0.000061         0.724268           5         1,000         200         0.0000151         0.2         0.0000053         0.000065         0.55623           6         1,000         100         0.0000052         0.2         0.0000053         0.0000054         0.856233           7         2,000         100         0.0000671         0.3         0.0000022         0.9842787           9         3,000         100         0.000075         0.3         0.0000083         0.0000082         0.377781           10         2,000         100         0.000071         0.3         0.0000083         0.3708787           11         2,000         100         0.000071         0.3         0.0000083         0.3708787           12         3,000         100         0.0000011         0.3         0.0000003         0.0000082         0.323389           14         1,000         100         0.000007         0.3         0.000003         0.000008         0.5545723           15         2,000         100         0.000007         0.2         0.000003         0.000008         0.454552           16	2	1,000	200	0.0000994	0.3	0.00000023	0.0000043	0.5736595
5         1,000         200         0.000011         0.2         0.0000084         0.20000085         0.266293           7         2,000         100         0.0000085         0.2         0.0000035         0.0000084         0.826293           7         2,000         100         0.0000675         0.3         0.0000035         0.0000024         0.826293           9         3,000         200         0.000066         0.3         0.0000035         0.0000078         0.3707891           10         2,000         100         0.000072         0.3         0.0000066         0.446765           11         2,000         100         0.0000011         0.3         0.0000020         0.000089         0.323369           12         3,000         100         0.0000011         0.2         0.0000020         0.0000020         0.828658           14         1,000         100         0.0000002         0.3         0.0000020         0.0000056         0.54923           15         2,000         200         0.0000002         0.2         0.0000030         0.0000056         0.545023           16         1,000         100         0.0000067         0.2         0.0000003         0.0000057	3	3,000	200	0.0000292	0.3	0.00000049	0.0000086	0.9229554
61,0001000.0000080.20.00000330.0000880.626629372,0001000.0000510.20.00000350.00000340.87245082,0001000.00005730.30.00000350.00000220.984278793,0002000.00005660.30.00000360.00000560.5456765102,0001000.0000720.30.00000060.56567112,0001000.0000010.30.00000020.0000080.537087123,0002000.0000010.20.00000020.0000050.5233369133,0001000.0000010.20.0000030.0000760.8248724141,0001000.0000070.20.0000030.0000760.8248724152,0002000.0000070.20.0000030.0000760.5248724161,0001000.0000070.20.0000030.0000780.5545023171,0002000.00000770.20.0000030.0000580.5545023181,0001000.00000770.20.00000080.5459252191,0001000.00000770.20.00000080.545925191,0001000.0016770.20.0000080.64415131153,0000.2118570.000007820107.94218.911155,0000.017433200	4	3,000	100	0.0000463	0.2	0.00000076	0.0000021	0.7242638
7         2,000         100         0.000612         0.2         0.0000035         0.0000032         0.0000032         0.0342787           9         3,000         200         0.000858         0.3         0.0000063         0.000078         0.3000078         0.0000078         0.0000078         0.0000078         0.0000078         0.0000078         0.0000078         0.0000078         0.0000078         0.0000078         0.0000078         0.0000078         0.0000078         0.056767           11         2,000         100         0.000010         0.3         0.0000008         0.0000080         0.5323369           13         3,000         200         0.000001         0.2         0.0000003         0.0000078         0.5245724           14         1,000         100         0.0000002         0.3         0.0000003         0.0000078         0.545523           16         1,000         100         0.000007         0.2         0.0000003         0.0000058         0.545523           17         1,000         100         0.0000077         0.2         0.0000005         0.0000075         0.5000075         0.0000075         0.0000075         0.5000075         0.0000075         0.0000075         0.5000075         0.0000075         0.	5	1,000	200	0.0000161	0.2	0.00000054	0.0000066	0.8150428
82,0001000.0005750.30.00000200.0000220.94278793,0002000.0000660.30.00000680.00000880.777891102,0001000.0000140.30.00000080.00000880.545765123,0002000.0000110.30.00000000.00000830.5336987133,0001000.0000020.30.00000020.00000700.828658141,0001000.0000010.00000020.00000020.00000700.52927152,0002000.0000070.20.00000020.0000080.545023161,0001000.0000070.20.00000000.0000680.545023171,0002000.0000070.20.00000000.0000680.545023181,0001000.0000770.20.00000000.0000680.545023191,0001000.0000770.20.00000000.0000870.54503100.0000870.20.00000000.0000870.545030.0000871010,0001000.0000770.20.00000000.0000870.54503111,0001000.0000770.20.00000000.0000870.54503111,0001000.0000780.0000870.0000870.0000870.54503111,0001,0000.0000780.0000870.0000870.0000870.0000870	6	1,000	100	0.0000008	0.2	0.00000003	0.0000058	0.6266293
93,0002000.0000860.30.00000630.0000780.777891102,0001000.00007880.30.00000060.00000800.545675112,0001000.00000180.30.00000020.00000200.533369133,0001000.00000180.30.00000020.00000760.5233769141,0001000.0000010.20.00000020.00000760.523474152,0002000.0000070.20.00000030.0000760.5545023161,0001000.0000070.20.0000030.0000080.545523171,0001000.0000070.20.00000030.0000080.545023181,0001000.0000070.20.00000030.0000080.545523191,0001000.0000070.20.00000020.0000080.641813191,0001000.0000070.20.00000030.0000080.641813101,0001000.0140030.0000750.00000740.00000740.441811153,0000.0157550.00007520107.80458.671155,0000.0157550.00007320107.91218.9112104,0000.0137370.00007420107.94218.911355,0000.0137310.00007420107.80364.22145	7	2,000	100	0.0000612	0.2	0.00000035	0.0000034	0.8724350
102,0001000.00007280.30.00000980.0000080.5456765112,0001000.00000110.30.00000020.00000820.523369123,0002000.00000110.30.00000020.00000820.523369133,0001000.00000110.20.00000030.00000760.8246724152,0002000.0000020.30.00000020.0000030.602297161,0001000.0000070.20.00000030.0000080.6756674181,0001000.0000020.30.00000030.0000080.545023191,0001000.0000070.20.00000030.0000080.545023201,0001000.0000070.20.00000030.0000080.549252191,0001000.0000070.20.00000030.0000080.549252201,0001000.0000080.20.00000030.0000080.6441813maximus $y^*$ of sequence, navit $y^*$ of sequence, navi	8	2,000	100	0.0000575	0.3	0.00000020	0.0000022	0.9842787
112,0001000.0000410.30.0000060.0000980.6376087123,0002000.0000180.30.00000020.00000280.5233369133,0001000.0000020.30.00000030.00000760.8248724141,0001000.0000010.20.00000020.00000760.622977152,0002000.0000020.30.0000030.0000080.6750674161,0001000.0000070.20.00000030.0000080.545523171,0001000.0000070.20.00000030.0000080.545023181,0001000.00000070.20.00000030.0000080.545023191,0001000.00000080.20.00000030.00000500.5490809201,0001000.00000770.20.00000030.00000500.6441513101,0001000.00000770.20.00000080.00000570.64415131153,0000.02169570.00007520107.94218.911254,0000.0137330.00005820107.94218.913104,0000.02169570.00073320107.94218.91350.0000.0131320.00005820107.94218.91350.0000.0131330.00005820107.80364.225104	9	3,000	200	0.0000866	0.3	0.00000063	0.0000078	0.7707891
123,0002000.0000180.30.0000020.0000020.0000020.000002133,0001000.0000010.20.00000030.0000060.892658141,0001000.0000010.20.00000030.00000130.602927152,0002000.00000070.20.00000030.0000560.5545023161,0001000.0000070.20.00000030.0000080.545523171,0002000.0000080.20.00000030.0000080.545523181,0001000.0000070.20.00000030.0000080.545023191,0002000.0000080.20.00000030.0000080.545023191,0002000.0000080.20.00000030.0000080.545023101,0002000.0000080.20.00000030.0000080.545023101,0002000.0000080.20.00000030.0000080.641513101,0002000.000080.20.0000080.6415131153,0000.0194730.0007520107.80\$45.6712105,0000.0174330.00007520107.94218.9113104,0000.0137370.00007520107.80364.221455,0000.0131330.00008720107.80364.3215104,000 </td <td>10</td> <td>2,000</td> <td>100</td> <td>0.0000728</td> <td>0.3</td> <td>0.00000098</td> <td>0.0000096</td> <td>0.5456765</td>	10	2,000	100	0.0000728	0.3	0.00000098	0.0000096	0.5456765
133,0001000.0000020.30.0000040.0000690.8926658141,0001000.0000110.20.00000030.0000070.824874152,0002000.0000080.30.00000020.0000130.6029297161,0001000.00000070.20.00000030.0000560.5545023171,0002000.00000020.30.00000030.0000050.545023181,0001000.0000080.20.00000020.00000570.5490297191,0001000.0000080.20.00000020.00000570.5490297201,0001000.0000080.20.00000020.00000570.5490297201,0002000.0000080.20.00000020.00000570.549089201,0002000.0000070.20.00000080.0000570.549089201,0002000.0000070.20.00000850.64415133000550.0000750.00000750.00000850.64415131153,0000.0154570.00007320107.80458.6721105,0000.0154530.00007320107.91629.753104,0000.0131220.00048620107.82161.03455,0000.0131340.00003720107.83364.225104,0000.013515	11	2,000	100	0.0000041	0.3	0.00000006	0.0000098	0.6376087
141,0001000.0000110.20.0000030.0000760.8248724152,0002000.0000080.30.00000020.0000130.602927161,0001000.00000070.20.00000030.0000080.5545023171,0002000.00000080.20.00000040.0000080.675674181,0001000.0000080.20.00000040.0000080.5450252191,0001000.0000070.20.00000020.0000570.549089201,0002000.00000670.20.00000080.0000550.6441513191,0002000.0000080.20.00000080.0000550.6441513201,0002000.0000080.20.00000550.6441513201,00020010.0000080.0000550.6441513201,0002000.0000550.0000550.0000550.6441513201,00020010.0000710.50.0000560.0000550.000055115.0000.0144330.00005620107.80458.6721105.0000.0151540.00007320107.91629.7551003.0000.0151540.00007520107.91629.7551003.0000.0151540.00025420107.4874.806103.0000.0151540.00005720104.93<	12	3,000	200	0.0000018	0.3	0.00000002	0.0000082	0.5233369
152,0002000.0000000.30.00000020.0000030.0000130.602927161,0001000.00000020.30.0000030.0000080.554503171,0002000.00000020.30.00000030.0000080.549522181,0001000.0000070.20.00000020.0000080.5495252191,0002000.0000070.20.00000020.0000080.641513201,0002000.0000080.20.0000080.0000080.641513501,0002000.0000051.20.0000080.0000850.64151350500.0000071.20.0000080.0000850.6415135051505,0000.0174330.0007320107.94218.91515,0000.0137330.00008520107.94216.0361103,0000.0137330.00037420107.94216.0351104,0000.0137330.00037420107.4351.9554105,0000.0187510.00037420107.4351.9564100.0137330.00037420107.4351.9555104,000 <td>13</td> <td>3,000</td> <td>100</td> <td>0.0000002</td> <td>0.3</td> <td>0.00000004</td> <td>0.0000069</td> <td>0.8926658</td>	13	3,000	100	0.0000002	0.3	0.00000004	0.0000069	0.8926658
161,0001000.0000070.20.0000030.0000050.545023171,0002000.0000080.20.00000030.0000080.60000750.545023181,0001000.0000080.20.0000020.00000570.549029191,0001000.0000080.20.00000020.00000570.5490809201,0002000.0000080.20.00000080.00000570.6441513Fine-tuning iteration No.Max number of of sequences, maxitLength of for maxitProbability of for maxitProbability of probability of (P(H + 1))Population Size, N_PSample Size, N_PExecution Time153.0000.0159750.00007520107.94458.672105.0000.0159750.00007520107.94218.91355.0000.0131230.00008520107.94364.22455.0000.013730.00007720104.93207.115104.0000.0193730.00027720104.93207.117104.0000.0210330.00008520107.7451.958105.0000.0154510.00037720104.93207.117104.0000.0210330.000085120107.7451.958105.0000.0154530.00037720	14	1,000	100	0.0000011	0.2	0.00000003	0.0000076	0.8248724
171,0002000.0000020.30.0000030.0000080.6750674181,0001000.0000800.20.0000020.00000570.5495252191,0002000.00000670.20.00000020.0000570.5490809201,0002000.0000080.20.0000080.0000850.6441513Fine-tuning iterationMax number of sequencey maxitKength 	15	2,000	200	0.0000008	0.3	0.00000002	0.0000013	0.6029297
181,0001000.0000080.20.00000400.0000080.5495252191,0001000.0000670.20.00000020.00000570.5490809201,0002000.0000080.20.00000080.00000850.6441513201,0002000.0000080.20.00000080.00000850.644151320MaxLengthforProbabilityofSize,SampleSampleExecutionnumberofsequence,y avince,YCH + 1)N_MPopulationSize,SampleSampleExecution153,0000.02169570.00007520107.80458.67Size,2105,0000.0174330.00006320107.91218.91355,0000.0137250.00007320107.82161.03455,0000.013730.00005720107.82161.035104,0000.0220330.00025720107.82161.036103,0000.0151540.00037420107.48728.80953,0000.0216080.00005120108.03359.16105,0000.0121880.000082120108.32296.151153,0000.0125680.00035020107.76296.911253,0000.0156580.00016020107.89371.97 <td>16</td> <td>1,000</td> <td>100</td> <td>0.0000007</td> <td>0.2</td> <td>0.00000003</td> <td>0.0000056</td> <td>0.5545023</td>	16	1,000	100	0.0000007	0.2	0.00000003	0.0000056	0.5545023
191,0001000.0006770.20.00000020.00000570.5490809201,0002000.0000080.20.0000080.0000850.6441513201,00020010000080.20.0000080.0000850.64151320Max of sequence, of sequence, maxin% of rq value, p(u) Lmax%Probability of stratum, P(H + 1)Population Size, N_PSample Size, Size, N_PSample Size, Size, N_PSample Size, Size, Size, N_PSample Size, Size, Size, N_PSample Size, Size, Size, N_PSample Size, Size, Size, Size, Size, Size,Sample Size,<	17	1,000	200	0.0000002	0.3	0.00000003	0.0000088	0.6750674
201,0002000.0000080.20.0000080.0000850.6441513Fine-tuning iteration No.Max number of sequences, maxitLength of sequence, JProbability of sequence, JProbability of New P(H + 1)Population Size, P(H + 1)Sample Sample Size, N_PExecution Time153,0000.0216970.00007520107.80458.672105,0000.0174330.00096320107.94218.91354,0000.0159750.00047320107.9162.975455,0000.0131230.00046620107.9162.9755103,0000.0191540.00037720104.93207.116103,0000.01951540.0007720104.93207.117104,0000.01951540.0007720107.4251.958105,0000.01875210.0007720107.43359.16953,0000.0121880.00016420108.32296.1510153,0000.0152540.00037520107.76296.9111153,0000.0152540.00037520107.76296.9112254,0000.0152540.00037520107.76296.91133100.0121880.00037520107.76296.911415	18	1,000	100	0.0000008	0.2	0.00000040	0.0000008	0.5495252
Fine-tuning iteration No.Max number of sequences, maxitLength of sequence, J% of L for maximum q value, Lmax%Probability of New P(H + 1)Population Size, N_PSample Size, N_PExecution Time153,0000.02169570.00007520107.80458.672105,0000.0174330.00096320107.94218.91354,0000.01597550.00007320108.06364.22455,0000.01312320.000086320107.91629.755104,0000.0133730.000037320107.82161.036103,0000.0193130.000037120107.82161.036103,0000.0187510.000037520107.43571.958105,0000.0187520.000057420107.43571.95954,0000.0210330.00028520107.74571.95104,0000.0187520.000057420107.48728.80953,0000.0121880.00018620108.03359.161153,0000.01564530.00039520107.76296.911254,0000.0179240.00039520107.76296.9113103,0000.01564530.00039520107.76296.91145 <td>19</td> <td>1,000</td> <td>100</td> <td>0.0000677</td> <td>0.2</td> <td>0.00000002</td> <td>0.0000057</td> <td>0.5490809</td>	19	1,000	100	0.0000677	0.2	0.00000002	0.0000057	0.5490809
Max inmber No.         Max inmber isequence maxit         Length if if isequence indication         Length if if if indication         Probability of if if if         Populati of isequence indication         Populati ise, ise, N_P         Sample ise, ise, N_P         Execution if ise, N_P           1         5         3,000         4,000         5,00000         0.000070         20100         107.90         45.07           1         5         3,000         0.010430         0.000070         20100         107.90         21.01           1         5         3,000         0.010430         0.000730         20.00         107.91         21.02           1         5         3,000         0.010430         20.000730         20.00         107.91         21.02           1         5         4,000         0.013132         0.000730         20.00         107.91         20.91           1         1         4,000         0.013132         0.000370         20.00         107.91         20.11           1         1         3,000         0.010373         20.000         107.91         20.11           1         1         3,000         0.010370         20.000         107.40         20.11           1         1 <t< td=""><td>20</td><td>1,000</td><td>200</td><td>0.0000008</td><td>0.2</td><td>0.00000008</td><td>0.0000085</td><td>0.6441513</td></t<>	20	1,000	200	0.0000008	0.2	0.00000008	0.0000085	0.6441513
2         10         5,000         0.0174433         0.000963         20         107.94         218.91           3         5         4,000         0.0159755         0.000733         20         108.06         364.22           4         5         5,000         0.0131232         0.000486         20         107.91         629.75           5         10         4,000         0.013373         0.000609         20         107.82         161.03           6         10         3,000         0.0195154         0.000377         20         104.93         207.11           7         10         4,000         0.0220333         0.000285         20         107.74         571.95           8         10         5,000         0.0187521         0.000574         20         107.48         728.80           9         5         4,000         0.0221063         0.000821         20         108.03         359.16           10         5,000         0.015218         0.000186         20         108.32         296.15           11         5         3,000         0.0156453         0.0000375         20         107.76         296.91           12         5	iteration	number of sequences,	of sequence,	L for maximum q value,	of New Stratum,	Size,	-	
3         5         4,000         0.0159755         0.0000733         20         108.06         364.22           4         5         5,000         0.0131232         0.0000486         20         107.91         629.75           5         10         4,000         0.013373         0.000609         20         107.82         161.03           6         10         3,000         0.0195154         0.000377         20         104.93         207.11           7         10         4,000         0.0220333         0.000285         20         107.74         571.95           8         10         5,000         0.0187521         0.000574         20         107.48         728.80           9         5         4,000         0.0221063         0.000821         20         108.03         359.16           10         5,000         0.0122188         0.000186         20         108.32         296.15           11         5         3,000         0.0156453         0.0000395         20         107.76         296.91           12         5         4,000         0.0179254         0.0000770         20         107.89         371.97           13         10	1	5	3,000	0.0216957	0.0000075	20	107.80	458.67
4         5         5,000         0.0131232         0.0000486         20         107.91         629.75           5         10         4,000         0.0103373         0.000609         20         107.82         161.03           6         10         3,000         0.0195154         0.000377         20         104.93         207.11           7         10         4,000         0.0220333         0.000285         20         107.74         571.95           8         10         5,000         0.0187521         0.000574         20         107.48         728.80           9         5         4,000         0.0221063         0.000821         20         108.03         359.16           10         5,000         0.012188         0.000186         20         108.32         296.15           11         5         3,000         0.0156453         0.000395         20         107.76         296.91           12         5         4,000         0.0179254         0.0000770         20         107.89         371.97           13         10         3,000         0.0156468         0.0000166         20         106.97         674.13	2	10	5,000	0.0174433	0.0000963	20	107.94	218.91
5         10         4,000         0.0103373         0.000609         20         107.82         161.03           6         10         3,000         0.0195154         0.000377         20         104.93         207.11           7         10         4,000         0.0220333         0.000285         20         107.74         571.95           8         10         5,000         0.0187521         0.000574         20         107.48         728.80           9         5         4,000         0.0221063         0.000821         20         108.03         359.16           10         5,000         0.012188         0.000186         20         108.32         296.15           10         5         3,000         0.0156453         0.000395         20         107.76         296.91           11         5         3,000         0.0179254         0.0000770         20         107.76         296.91           12         5         4,000         0.0179254         0.0000770         20         107.89         371.97           13         10         3,000         0.015686         0.000116         20         106.97         674.13	3	5	4,000	0.0159755	0.0000733	20	108.06	364.22
6         10         3,000         0.0195154         0.0000377         20         104.93         207.11           7         10         4,000         0.0220333         0.000285         20         107.74         571.95           8         10         5,000         0.0187521         0.0000574         20         107.48         728.80           9         5         4,000         0.0221063         0.0000821         20         108.03         359.16           10         5         3,000         0.012188         0.0000821         20         108.03         359.16           10         5         3,000         0.0122188         0.0000375         20         108.32         296.15           11         5         3,000         0.0156453         0.0000375         20         107.76         296.91           12         5         4,000         0.0179254         0.0000770         20         107.89         371.97           13         10         3,000         0.0155686         0.0000116         20         106.97         674.13	4	5	5,000	0.0131232	0.0000486	20	107.91	629.75
7         10         4,000         0.0220333         0.0000285         20         107.74         571.95           8         10         5,000         0.0187521         0.0000574         20         107.48         728.80           9         5         4,000         0.0241063         0.0000821         20         108.03         359.16           10         5         3,000         0.012188         0.0000861         20         108.32         296.15           11         5         3,000         0.0156453         0.000395         20         107.76         296.91           12         5         4,000         0.0179254         0.0000770         20         107.89         371.97           13         10         3,000         0.0155686         0.0000116         20         106.97         674.13	5	10	4,000	0.0103373	0.0000609	20	107.82	161.03
8         10         5,000         0.0187521         0.0000574         20         107.48         728.80           9         5         4,000         0.0241063         0.000821         20         108.03         359.16           10         5         3,000         0.0122188         0.0000362         20         108.32         296.15           11         5         3,000         0.0156453         0.000395         20         107.76         296.91           12         5         4,000         0.0179254         0.0000770         20         107.89         371.97           13         10         3,000         0.0155686         0.0000116         20         106.97         674.13	6	10	3,000	0.0195154	0.0000377	20	104.93	207.11
9         5         4,000         0.0241063         0.0000821         20         108.03         359.16           10         5         3,000         0.0122188         0.000186         20         108.32         296.15           11         5         3,000         0.0156453         0.000395         20         107.76         296.91           12         5         4,000         0.0179254         0.0000770         20         107.89         371.97           13         10         3,000         0.0155686         0.0000116         20         106.97         674.13	7	10	4,000	0.0220333	0.0000285	20	107.74	571.95
10         5         3,000         0.0122188         0.0000186         20         108.32         296.15           11         5         3,000         0.0156453         0.0000395         20         107.76         296.91           12         5         4,000         0.0179254         0.0000770         20         107.89         371.97           13         10         3,000         0.0155686         0.0000116         20         106.97         674.13	8	10	5,000	0.0187521	0.0000574	20	107.48	728.80
11         5         3,000         0.0156453         0.0000395         20         107.76         296.91           12         5         4,000         0.0179254         0.0000770         20         107.89         371.97           13         10         3,000         0.0155686         0.0000116         20         106.97         674.13	9	5	4,000	0.0241063	0.0000821	20	108.03	359.16
12         5         4,000         0.0179254         0.0000770         20         107.89         371.97           13         10         3,000         0.0155686         0.0000116         20         106.97         674.13	10	5	3,000	0.0122188	0.0000186	20	108.32	296.15
13         10         3,000         0.0155686         0.0000116         20         106.97         674.13	11	5	3,000	0.0156453	0.0000395	20	107.76	296.91
	12	5	4,000	0.0179254	0.0000770	20	107.89	371.97
14         5         3,000         0.0213493         0.0000442         20         106.65         139.98	13	10	3,000	0.0155686	0.0000116	20	106.97	674.13
	14	5	3,000	0.0213493	0.0000442	20	106.65	139.98

<i>Table 3.1:</i>	Hyperparameters	for the	HEDA ·	in the	case a	of Continue	ous Strata
-------------------	-----------------	---------	--------	--------	--------	-------------	------------

		1			1		
15	10	3,000	0.0191401	0.0000816	20	107.40	280.55
16	10	4,000	0.0119913	0.0000363	20	107.00	277.91
17	10	5,000	0.0195217	0.0000438	20	106.90	201.23
18	10	3,000	0.0188604	0.0000191	20	107.65	213.88
19	5	4,000	0.0235152	0.0000413	20	107.03	174.75
20	10	4,000	0.0218042	0.0000257	20	105.95	165.34
					Minimum	104.93	
						Total	6,792.40
	ſ	Γ	American C	Community Sur	vey, 2015		I
Fine-tuning		SAA	Mutation	Elitism	Add	Temperature,	Decrement Constant,
iteration	Iterations, I	Frequency	Chance	Rate	Strata	т	DC
No.					Factor		
1	30	15	0.0000004	0.2	0.0000008	0.0000001	0.6933468
2	20	15	0.0000010	0.3	0.0000002	0.0000004	0.5736595
3	30	15	0.0000003	0.3	0.0000005	0.0000009	0.9229554
4	30	10	0.0000005	0.2	0.0000008	0.0000002	0.7242638
5	20	15	0.0000002	0.2	0.0000005	0.0000007	0.8150428
6	20	10	0.0000000	0.2	0.0000000	0.0000006	0.6266293
7	20	10	0.0000006	0.2	0.0000003	0.0000003	0.8724350
8	30	10	0.0000006	0.3	0.0000002	0.0000002	0.9842787
9	30	15	0.0000009	0.3	0.0000006	0.0000008	0.7707891
10	20	10	0.0000007	0.3	0.0000010	0.0000010	0.5456765
11	30	10	0.0000003	0.3	0.0000002	0.0000009	0.9659689
12	30	10	0.0000005	0.3	0.0000002	0.0000009	0.9542911
13	30	15	0.0000002	0.3	0.0000002	0.0000009	0.9813440
14	30	10	0.0000002	0.3	0.0000001	0.0000001	0.9533995
15	30	10	0.0000003	0.3	0.0000002	0.0000009	0.9863629
16	30	10	0.0000003	0.3	0.0000002	0.0000010	0.9702858
17	30	10	0.0000003	0.3	0.0000010	0.0000010	0.9849021
18	20	15	0.0000004	0.3	0.0000010	0.0000008	0.9729872
19	30	10	0.0000000	0.3	0.0000002	0.0000010	0.5353839
		10	0.0000003	0.3	0.0000008	0.0000010	0.5662247
20	30	10					
20	30	10	% of	Probability			
	30 Max	Length	% of L	Probability			
Fine-tuning				of	Population	Sample	Execution
Fine-tuning iteration	Max	Length	L	of New	Population Size,	Sample Size	Execution Time
Fine-tuning	Max number	Length of	L for	of New Stratum,	Population	-	
Fine-tuning iteration	Max number of sequences,	Length of sequence,	L for maximum	of New	Population Size,	-	
Fine-tuning iteration	Max number of sequences,	Length of sequence,	L for maximum q value,	of New Stratum,	Population Size,	-	
Fine-tuning iteration No.	Max number of sequences, maxit 1 2	Length of sequence, J 25,000 50,000	L for maximum q value, Lmax% 0.0216957 0.0174433	of New Stratum, P(H + 1) 0.00000008 0.00000096	Population Size, N_p 20 20	Size 3,026.35 2,762.46	Time 4,003.11 11,470.11
Fine-tuning iteration No. 1	Max number of sequences, maxit 1 2 1	Length of sequence, J 25,000 50,000 25,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755	of New Stratum, P(H + 1) 0.00000008 0.00000096 0.00000073	Population Size, N_p 20 20 20	Size 3,026.35	Time 4,003.11
Fine-tuning iteration No. 1 2 3 4	Max number of sequences, maxit 1 2 1 1 1	Length of sequence, J 25,000 50,000 25,000 50,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232	of New Stratum, P(H + 1) 0.00000008 0.00000096 0.00000073 0.00000049	Population Size, N_P 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32	Time 4,003.11 11,470.11
Fine-tuning iteration No. 1 2 3 4 5	Max number of sequences, maxit 1 2 1 1 2 2	Length of sequence, J 25,000 50,000 25,000 50,000 50,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755	of New Stratum, P(H + 1) 0.00000008 0.00000096 0.00000073 0.00000049 0.00000061	Population Size, N_P 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18	Time 4,003.11 11,470.11 3,987.64
Fine-tuning iteration No. 1 2 3 4	Max number of sequences, maxit 1 2 1 1 2 2 2	Length of sequence, J 25,000 50,000 25,000 50,000 25,000 25,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154	of New Stratum, P(H + 1) 0.0000008 0.00000096 0.00000049 0.00000061 0.00000038	Population Size, N_p 20 20 20 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32	Time 4,003.11 11,470.11 3,987.64 15,910.83
Fine-tuning iteration No. 1 2 3 4 5	Max number of sequences, maxit 1 2 1 1 2 2	Length of sequence, J 25,000 50,000 25,000 50,000 50,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373	of New Stratum, P(H + 1) 0.00000008 0.00000096 0.00000073 0.00000049 0.00000061	Population Size, N_P 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32 2,766.39	Time 4,003.11 11,470.11 3,987.64 15,910.83 11,459.90
Fine-tuning iteration No. 1 2 3 4 5 6	Max number of sequences, maxit 1 2 1 1 2 2 2	Length of sequence, J 25,000 50,000 25,000 50,000 25,000 25,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154	of New Stratum, P(H + 1) 0.0000008 0.00000096 0.00000049 0.00000061 0.00000038	Population Size, N_p 20 20 20 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32 2,766.39 2,932.12	Time 4,003.11 11,470.11 3,987.64 15,910.83 11,459.90 5,156.96
Fine-tuning iteration No. 1 2 3 4 5 6 7	Max number of sequences, maxit 1 2 1 1 2 2 2 2 2	Length of sequence, J 25,000 50,000 25,000 50,000 25,000 25,000 25,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333	of New Stratum, P(H + 1) 0.00000096 0.00000096 0.00000049 0.00000049 0.00000061 0.00000038	Population Size, N_p 20 20 20 20 20 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32 2,766.39 2,932.12 2,928.78	Time           4,003.11           11,470.11           3,987.64           15,910.83           11,459.90           5,156.96           5,177.00
Fine-tuning iteration No. 1 2 3 4 5 6 7 8	Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 1 1 1 1 1	Length of sequence, J 25,000 50,000 25,000 25,000 25,000 25,000 50,000 25,000 25,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0187521 0.0241063 0.0122188	of New Stratum, P(H + 1) 0.00000008 0.00000096 0.00000073 0.00000049 0.00000049 0.00000049 0.00000057 0.00000057 0.00000082 0.00000019	Population Size, N_p 20 20 20 20 20 20 20 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32 2,766.39 2,932.12 2,932.12 2,928.78 2,620.40	Time         4,003.11         11,470.11         3,987.64         15,910.83         11,459.90         5,156.96         5,177.00         22,859.85
Fine-tuning iteration No. 1 2 3 4 5 6 6 7 8 9	Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 1	Length of sequence, J 25,000 50,000 25,000 25,000 25,000 25,000 50,000 50,000 50,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0187521 0.0241063	of New Stratum, P(H + 1) 0.00000008 0.00000096 0.00000073 0.00000049 0.00000049 0.00000049 0.00000057 0.00000057	Population Size, N_p 20 20 20 20 20 20 20 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32 2,766.39 2,932.12 2,928.78 2,620.40 2,861.24	Time         4,003.11         11,470.11         3,987.64         15,910.83         11,459.90         5,156.96         5,177.00         22,859.85         8,281.68
Fine-tuning iteration No. 1 2 3 4 5 6 7 8 9 9 10	Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 1 1 1 1 1	Length of sequence, J 25,000 50,000 25,000 25,000 25,000 25,000 50,000 50,000 50,000 50,000 50,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0187521 0.0241063 0.0122188	of New Stratum, P(H + 1) 0.00000008 0.00000096 0.00000073 0.00000049 0.00000049 0.00000049 0.00000057 0.00000057 0.00000082 0.00000019	Population Size, N_p 20 20 20 20 20 20 20 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32 2,766.39 2,932.12 2,928.78 2,620.40 2,861.24 3,027.27	Time         4,003.11         11,470.11         3,987.64         15,910.83         11,459.90         5,156.96         5,177.00         22,859.85         8,281.68         3,872.97
Fine-tuning iteration No. 1 2 3 4 5 6 7 8 9 10 11	Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 2 1 1 1 2 2 2 2	Length of sequence, J 25,000 50,000 25,000 25,000 25,000 25,000 50,000 50,000 50,000 50,000 50,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0195154 0.0220333 0.0187521 0.0241063 0.0122188 0.0157505	of New Stratum, P(H + 1) 0.0000008 0.00000096 0.00000049 0.00000049 0.00000049 0.00000057 0.00000057 0.00000057 0.00000052	Population Size, N_P 20 20 20 20 20 20 20 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32 2,766.39 2,932.12 2,928.78 2,620.40 2,861.24 3,027.27 2,621.62	Time 4,003.11 11,470.11 3,987.64 15,910.83 11,459.90 5,156.96 5,177.00 22,859.85 8,281.68 3,872.97 22,758.15
Fine-tuning iteration No. 1 2 3 4 5 6 7 8 9 10 11 12	Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 2 1 1 1 2 2 1 1 1 2 2 1 1 1 2 1 1 2 1 1 2 2 1 1 1 2 2 1 1 1 2 2 1 1 1 2 1 2 1 1 2 1 1 2 1 1 2 1 1 2 1	Length of sequence, J 25,000 50,000 25,000 25,000 25,000 25,000 50,000 50,000 50,000 50,000 50,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0187521 0.0241063 0.0122188 0.0157505 0.0165162	of New Stratum, P(H + 1) 0.0000008 0.00000096 0.00000049 0.00000049 0.00000049 0.00000049 0.00000057 0.00000057 0.00000057 0.00000043 0.00000043 0.0000003	Population Size, N_P 20 20 20 20 20 20 20 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32 2,766.39 2,932.12 2,928.78 2,620.40 2,861.24 3,027.27 2,621.62 2,704.42	Time         4,003.11         11,470.11         3,987.64         15,910.83         11,459.90         5,156.96         5,177.00         22,859.85         8,281.68         3,872.97         22,758.15         15,941.84
Fine-tuning iteration No. 1 2 3 4 5 6 7 8 9 10 11 12 13	Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 1 1 1 2 2 1 1 1 2 1 1 2 1 1 1 2 1 1 1 2 1 1 1 2 1 1 1 2 2 1 1 1 1 2 2 1 1 1 1 2 1 1 1 1 2 1	Length of sequence, J 25,000 50,000 25,000 25,000 25,000 25,000 50,000 25,000 50,000 25,000 50,000 25,000 25,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0195154 0.0220333 0.0187521 0.0241063 0.0122188 0.0157505 0.0165162 0.0155316	of New Stratum, P(H + 1) 0.0000008 0.00000096 0.00000049 0.00000049 0.00000043 0.00000057 0.00000057 0.00000057 0.00000043 0.00000043 0.00000043	Population Size, N_P 20 20 20 20 20 20 20 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32 2,766.39 2,932.12 2,928.78 2,620.40 2,861.24 3,027.27 2,621.62 2,704.42 3,015.87	Time         4,003.11         11,470.11         3,987.64         15,910.83         11,459.90         5,156.96         5,177.00         22,859.85         8,281.68         3,872.97         22,758.15         15,941.84         4,023.10
Fine-tuning iteration No. 1 2 3 4 5 6 7 8 9 10 11 12 13 14	Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 1 1 1 2 2 1 1 2 2 1 1 2 2 1 1 2 2 1 2 2 2 2 1 1 1 2 2 2 2 2 2 1 2 2 2 2 1 2	Length of sequence, J 25,000 50,000 25,000 25,000 25,000 25,000 50,000 50,000 50,000 50,000 25,000 25,000 25,000 25,000 25,000	L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0187521 0.0241063 0.0122188 0.0157505 0.0165162 0.0155316 0.0145631	of New Stratum, P(H + 1) 0.0000008 0.00000096 0.00000049 0.00000049 0.00000043 0.00000028 0.00000028 0.00000028 0.00000028 0.00000043 0.00000043 0.00000043 0.00000041 0.00000047	Population Size, N_p 20 20 20 20 20 20 20 20 20 20 20 20 20	Size 3,026.35 2,762.46 3,029.18 2,703.32 2,766.39 2,932.12 2,928.78 2,620.40 2,861.24 3,027.27 2,621.62 2,704.42 3,015.87 2,759.56	Time         4,003.11         11,470.11         3,987.64         15,910.83         11,459.90         5,156.96         5,177.00         22,859.85         8,281.68         3,872.97         22,758.15         15,941.84         4,023.10         10,261.55

18	2	50,000	0.0166339	0.0000014	20	2 763 12	11 136 54
18	2	50,000 50,000	0.0166339	0.00000014	20 20	2,763.13 2,619.99	11,136.54 22,400.28
							15,818.99
20	1	50,000	0.0121512	0.00000007	20 Minimum	2,701.10	15,818.99
					Willinum	2,619.06 Total	247,314.63
				Kiva Loans		Iotai	247,314.03
Fine-tuning			1	Itiva Loans	Add		
iteration	Iterations, I	SAA	Mutation	Elitism	Strata	Temperature,	Decrement Constant,
No.	, -	Frequency	Chance	Rate	Factor	т	DC
1	90	20	0.0000004	0.2	0.0000008	0.0000001	0.6933468
2	50	20	0.0000010	0.3	0.0000002	0.0000004	0.5736595
3	90	20	0.0000003	0.3	0.0000005	0.0000009	0.9229554
4	100	10	0.0000005	0.2	0.0000008	0.0000002	0.7242638
5	60	20	0.0000002	0.2	0.0000005	0.0000007	0.8150428
6	50	10	0.0000000	0.2	0.0000000	0.0000006	0.6266293
7	70	10	0.0000006	0.2	0.0000003	0.0000003	0.8724350
8	80	10	0.0000006	0.3	0.0000002	0.0000002	0.9842787
9	100	20	0.0000009	0.3	0.0000006	0.0000008	0.7707891
10	70	10	0.0000007	0.3	0.0000010	0.0000010	0.5456765
11	80	10	0.0000004	0.3	0.0000008	0.0000001	0.9727142
12	80	20	0.0000003	0.3	0.0000008	0.0000000	0.9796571
13	90	10	0.0000004	0.3	0.0000008	0.0000002	0.9820326
14	90	20	0.0000004	0.3	0.0000004	0.0000002	0.9887106
15	90	10	0.0000004	0.3	0.0000001	0.0000001	0.9844605
16	90	20	0.0000003	0.3	0.0000000	0.0000001	0.9838737
17	90	10	0.0000004	0.3	0.0000007	0.0000001	0.9778088
18							
10	90	10	0.0000001	0.3	0.0000005	0.0000001	0.9318817
18 19	90 90	10 10	0.0000001 0.0000004	0.3 0.3	0.0000005 0.0000004	0.0000001 0.0000001	0.9318817 0.9233493
			0.0000004 0.0000004				
19	90 90	10 10	0.0000004 0.0000004 % of	0.3	0.0000004	0.0000001	0.9233493
19	90 90 Max	10 10 Length	0.0000004 0.0000004 % of L	0.3	0.0000004	0.0000001 0.0000002	0.9233493 0.7065010
19 20	90 90 Max number	10 10 Length of	0.0000004 0.0000004 % of L for	0.3 0.3 Probability	0.0000004	0.0000001 0.0000002 Sample	0.9233493 0.7065010 Execution
19 20 Fine-tuning	90 90 Max number of sequences,	10 10 Length of sequence,	0.0000004 0.0000004 % of L for maximum	0.3 0.3 Probability of	0.0000004 0.0000008 Population	0.0000001 0.0000002	0.9233493 0.7065010
19 20 Fine-tuning iteration	90 90 Max number	10 10 Length of	0.0000004 0.0000004 % of L for maximum q value,	0.3 0.3 Probability of New	0.0000004 0.0000008 Population Size,	0.0000001 0.0000002 Sample	0.9233493 0.7065010 Execution
19 20 Fine-tuning iteration No.	90 90 Max number of sequences, maxit	10 10 Length of sequence, J	0.0000004 0.0000004 % of L for maximum q value, Lmax%	0.3 0.3 Probability of New Stratum, P(H + 1)	0.0000004 0.0000008 Population Size, N_p	0.0000001 0.0000002 Sample Size	0.9233493 0.7065010 Execution Time
19 20 Fine-tuning iteration No.	90 90 Max number of sequences, maxit 1	10 10 Length of sequence, J 10,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001	0.0000004 0.0000008 Population Size, N_p 20	0.0000001 0.0000002 Sample Size 1,806.82	0.9233493 0.7065010 Execution Time 1,150.00
19 20 Fine-tuning iteration No. 1 2	90 90 Max number of sequences, maxit 1 2	10 10 Length of sequence, J 10,000 20,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433	0.3 0.3 Probability of New Stratum, P(H + 1)	0.0000004 0.0000008 Population Size, N_p 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59
19 20 Fine-tuning iteration No. 1 2 3	90 90 Max number of sequences, maxit 1 2 1	10 10 Length of sequence, J 10,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000001 0.0000007	0.0000004 0.0000008 Population Size, N_P 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60
19 20 Fine-tuning iteration No. 1 2 3 4	90 90 Max number of sequences, maxit 1 2	10 10 Length of sequence, J 10,000 20,000 10,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000001 0.0000007 0.0000005	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39
19 20 Fine-tuning iteration No. 1 2 3 3 4 5	90 90 Max number of sequences, maxit 1 2 1 1 2 2	10 10 Length of sequence, J 10,000 20,000 10,000 20,000 20,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000001 0.0000007 0.0000005 0.0000006	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81
19 20 Fine-tuning iteration No. 1 2 3 4 5 5 6	90 90 Max number of sequences, maxit 1 2 1 1 2 2 2	10 10 Length of sequence, J 10,000 20,000 10,000 20,000 10,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0174433 0.0159755	0.3 0.3 Probability of New Stratum, P(H + 1) 0.000001 0.0000001 0.0000005 0.0000005 0.0000006	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39
19 20 Fine-tuning iteration No. 1 2 3 3 4 5	90 90 Max number of sequences, maxit 1 2 1 1 2 2	10 10 Length of sequence, J 10,000 20,000 10,000 20,000 20,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000001 0.0000007 0.0000005 0.0000006	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30
19 20 Fine-tuning iteration No. 1 2 3 4 5 5 6 7	90 90 Max number of sequences, maxit 1 2 1 1 2 2 2 2 2	10 10 Length of sequence, J 10,000 20,000 10,000 20,000 10,000 10,000 10,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000001 0.0000005 0.0000005 0.0000006 0.0000004 0.0000003	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86 1,681.44	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30 2,239.25
19 20 Fine-tuning iteration No. 1 2 3 4 5 5 6 7 8	90 90 Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2	10 10 Length of sequence, J 10,000 20,000 10,000 20,000 10,000 10,000 20,000 10,000 20,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0187521	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000001 0.0000007 0.0000005 0.0000006 0.0000003 0.0000003	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86 1,681.44 1,597.02	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30 2,239.25 7,702.03
19 20 Fine-tuning iteration No. 1 2 3 4 5 5 6 6 7 7 8 9	90 90 Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 1	10 10 Length of sequence, J 10,000 20,000 10,000 20,000 10,000 10,000 20,000 20,000 20,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0187521 0.0241063	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000001 0.0000007 0.0000005 0.0000006 0.0000004 0.0000003 0.0000006 0.0000008	0.0000004 0.0000008 Population Size, N_p 20 20 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86 1,681.44 1,597.02 1,647.24	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30 2,239.25 7,702.03 2,869.39
19 20 Fine-tuning iteration No. 1 2 3 4 5 6 7 8 9 9 10	90 90 Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 1 1 1 1 2 2 2 2	10 10 Length of sequence, J 10,000 20,000 10,000 20,000 10,000 20,000 10,000 20,000 10,0	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0195154 0.0220333 0.0195154	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000001 0.0000007 0.0000005 0.0000006 0.0000006 0.0000008 0.0000008 0.0000002	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86 1,681.44 1,597.02 1,647.24 1,683.68	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30 2,239.25 7,702.03 2,869.39 1,517.16
19 20 20 Fine-tuning iteration No. 1 2 3 4 5 5 6 7 5 6 7 8 9 9 10 11	90 90 Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 1 1 1 1 1 1 1 1	10 10 Length of sequence, J 10,000 20,000 10,000 20,000 10,000 20,000 10,000 20,000 10,000 20,000 20,000 10,000 20,000 20,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0195154 0.0220333 0.0187521 0.0241063 0.0122188 0.0183654	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000001 0.0000005 0.0000005 0.0000006 0.0000003 0.0000003 0.0000008 0.0000002 0.0000007	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86 1,681.44 1,597.02 1,647.24 1,683.68 1,606.42	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30 2,239.25 7,702.03 2,869.39 1,517.16 4,810.26
19 20 20 Fine-tuning iteration No. 1 2 3 4 5 5 6 7 8 9 9 10 11 11 12	90 90 Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 1 1 1 1 1 1 1 1	10 10 Length of sequence, J 10,000 20,000 20,000 20,000 10,000 20,000 10,000 20,000 10,000 20,000 20,000 20,000 20,000 20,000 20,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0195154 0.0220333 0.0187521 0.0241063 0.0187521 0.0241063	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000007 0.0000005 0.0000005 0.0000006 0.0000006 0.0000008 0.0000002 0.0000007 0.0000007 0.0000007	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86 1,681.44 1,597.02 1,647.24 1,683.68 1,606.42 1,739.74	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30 2,239.25 7,702.03 2,869.39 1,517.16 4,810.26 2,112.80
19 20 20 Fine-tuning iteration No. 1 2 3 3 4 5 5 6 7 7 8 9 9 10 11 11 12 13	90 90 Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 2 1 1 1 1 1 1 1	10 10 Length of sequence, J 10,000 20,000 20,000 20,000 10,000 20,000 10,000 20,000 10,000 20,000 20,000 20,000 20,000 20,000 20,000 20,000 20,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0195154 0.0220333 0.0187521 0.0241063 0.0187521 0.0241063 0.018763	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000007 0.0000005 0.0000006 0.0000006 0.0000008 0.0000008 0.0000007 0.0000007 0.0000007 0.0000007 0.0000006 0.0000006	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86 1,681.44 1,597.02 1,647.24 1,683.68 1,606.42 1,739.74 1,592.34	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30 2,239.25 7,702.03 2,869.39 1,517.16 4,810.26 2,112.80 8,885.01
19 20 20 5 1 1 2 3 3 4 5 5 6 7 3 4 5 5 6 7 8 9 9 10 11 12 12 13 13 14	90 90 Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 1 1 1 1 1 1 1 1	10 10 Length of sequence, J 10,000 20,000 10,000 20,000 10,000 20,000 10,000 20,0	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0195154 0.0220333 0.0187521 0.0241063 0.018763 0.0183654 0.0187089 0.0137673 0.0138855	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000007 0.0000005 0.0000006 0.0000006 0.0000006 0.0000008 0.0000008 0.0000002 0.0000007 0.0000006 0.0000001 0.0000001	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86 1,681.44 1,597.02 1,647.24 1,683.68 1,606.42 1,739.74 1,592.34 1,629.67	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30 2,239.25 7,702.03 2,869.39 1,517.16 4,810.26 2,112.80 8,885.01 4,429.31
19 20 20 Fine-tuning iteration No. 1 2 3 3 4 5 6 7 6 7 8 9 9 10 10 11 10 12 13 13 14 15	90 90 90 Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 2 2 1 1 1 1 1 1	10 10 Length of sequence, J 10,000 20,000 10,000 20,000 10,000 20,000 10,000 20,000 20,000 20,000 20,000 20,000 20,000 20,000 20,000 20,000 20,000 20,000 20,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0174433 0.0159755 0.0174433 0.0159755 0.0174433 0.0159755 0.0174433 0.0159755 0.0174433 0.0159755 0.0174433 0.0122188 0.0187654 0.0183654 0.0183654 0.0187673 0.0138555 0.0156086	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000005 0.0000005 0.0000006 0.0000006 0.0000008 0.0000008 0.0000002 0.0000007 0.0000007 0.0000007 0.0000001 0.0000001 0.0000001 0.0000001	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86 1,681.44 1,597.02 1,647.24 1,683.68 1,606.42 1,739.74 1,592.34 1,629.67 1,576.80	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30 2,239.25 7,702.03 2,869.39 1,517.16 4,810.26 2,112.80 8,885.01 4,429.31 8,791.12
19 20 20 Fine-tuning iteration No. 1 2 3 4 5 6 7 8 9 10 10 11 10 11 12 13 13 14 15 16	90 90 90 Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 2 2 1 1 1 1 1 2 2 2 2 2 2 1 1 1 1 2	10 10 Length of sequence, J 10,000 20,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0195154 0.0220333 0.0187521 0.0241063 0.0122188 0.0187639 0.0137673 0.0138855 0.0156086 0.0152018	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000007 0.0000005 0.0000006 0.0000006 0.0000008 0.0000008 0.0000007 0.0000007 0.0000007 0.0000006 0.0000007 0.0000000 0.0000001 0.0000001 0.0000008 0.0000008	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86 1,681.44 1,597.02 1,647.24 1,683.68 1,606.42 1,739.74 1,592.34 1,592.34 1,592.34 1,629.67 1,576.80 1,622.36	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30 2,239.25 7,702.03 2,869.39 1,517.16 4,810.26 2,112.80 8,885.01 4,429.31 8,791.12 4,377.68
19 20 20 Fine-tuning iteration No. 1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17	90 90 90 Max number of sequences, maxit 1 2 1 1 2 2 2 2 2 2 2 2 2 2 1 1 1 1 1	10 10 Length of sequence, J 10,000 20,000	0.0000004 0.0000004 % of L for maximum q value, Lmax% 0.0216957 0.0174433 0.0159755 0.0174433 0.0159755 0.0131232 0.0103373 0.0195154 0.0220333 0.0195154 0.0220333 0.0195154 0.022188 0.0122188 0.0122188 0.0138654 0.0137673 0.0138855 0.0156086 0.0152018 0.0197092	0.3 0.3 Probability of New Stratum, P(H + 1) 0.0000001 0.0000007 0.0000005 0.0000006 0.0000006 0.0000003 0.0000008 0.0000002 0.0000007 0.0000007 0.0000006 0.0000007 0.0000008 0.0000001 0.0000008 0.0000008	0.0000004 0.0000008 Population Size, N_P 20 20 20 20 20 20 20 20 20 20	0.0000001 0.0000002 Sample Size 1,806.82 1,747.71 1,797.86 1,613.25 1,719.17 1,753.86 1,681.44 1,597.02 1,647.24 1,683.68 1,606.42 1,739.74 1,592.34 1,629.67 1,576.80 1,622.36 1,600.29	0.9233493 0.7065010 Execution Time 1,150.00 2,157.59 1,120.60 6,316.39 2,172.81 1,463.30 2,239.25 7,702.03 2,869.39 1,517.16 4,810.26 2,112.80 8,885.01 4,429.31 8,791.12 4,377.68 5,549.41

		Minimum	1,576.80	
			Total	88,951.97