

SIMULTANEOUS RECOVERY OF PIECEWISE ANALYTIC COEFFICIENTS IN A SEMILINEAR ELLIPTIC EQUATION

BASTIAN HARRACH AND YI-HSUAN LIN

ABSTRACT. In this short note, we investigate simultaneous recovery inverse problems for semilinear elliptic equations with partial data. The main technique is based on suitable linearization and monotonicity approaches. With these method at hand, we can determine the diffusion, cavity and coefficients simultaneously by knowing the corresponding Dirichlet-Neumann operators.

Keywords. Inverse boundary value problems, inverse obstacle problem, semilinear elliptic equations, simultaneous recovery, partial data, higher order linearization, monotonicity method, localized potentials

CONTENTS

1. Introduction	1
2. The main results	3
3. Proofs of main results	5
3.1. Local well-posedness result for the forward problem.	7
3.2. Unique identifiability result for the inverse problem.	7
3.3. Unique identifiability result for the inverse obstacle problem.	12
Appendix A. Piecewise analyticity	12
References	13

1. INTRODUCTION

In this note, we investigate some inverse problems for semilinear elliptic equations. Inverse problems for nonlinear partial differential equations (PDEs) have been paid a lot of attentions in past few decades. The main method to study inverse problems for nonlinear PDEs is relied on suitable linearization techniques, and the linearization approaches can be traced back to the pioneer work by Isakov [Isa93]. In [Isa93], he demonstrated that the first linearization of the corresponding DN map of the semilinear parabolic equation agrees to the DN map of the associated linearized equation. Hence, related known results of inverse boundary value problems for linear equations can be expected to apply, such that one is able to solve inverse problems for the nonlinear equations.

For the semilinear elliptic equation $\Delta u + a(x, u) = 0$ in a domain, the inverse problem of determining the coefficient $a(x, u)$ was proved in [IS94, Sun10], for $n \geq 3$, and in [IN95, Sun10, IY13] for $n = 2$. In addition, for quasilinear elliptic equations, related inverse boundary value problems have also been studied by [Sun96, SU97, KN02, LW07, MU20, KKU20, CFK⁺21]. Meanwhile, researchers also solved some inverse problems for the degenerate p -Laplace equation [SZ12, BHKS18], and for the fractional semilinear Schrödinger equation [LL19, LL21, Lin21]. The interior unique determination problem for quasilinear equations on Riemannian manifolds was recently studied in [LLS20]. We refer the readers to [Sun05, Uhl09] for more

introduction and discussions on inverse problems for nonlinear elliptic equations related to the more developments mentioned above.

In order to show the results in this work, we utilize the method has been developed in [FO20, LLLS21, LLLS20, LLST21], which benefits from the *nonlinearity* so that one can solve inverse problems for a class of semilinear elliptic equation. The method is called the *higher order linearization*, which introduces particular parameters to reduce a semilinear elliptic equation into different linearized elliptic equations. In [FO20, LLLS21], the authors studied inverse boundary value problem with full boundary measurements. In addition, simultaneous recovering inverse problems for semilinear PDEs were also considered in [LLS20, LLST21, LLL21, LLLZ21], and [KU20b, LLLS20, KU20a] studied the Calderón type problem with partial data independently. A similar approach was utilized to study some related inverse problems for fractional semilinear elliptic equations [LL21, Lin21].

In this note, we will not utilize some special solutions (such as the complex geometrical optics solutions) to prove the global uniqueness result. Instead, we are going to introduce the *monotonicity method* combining with the *localized potentials* to demonstrate the unique determination consequence. Roughly speaking, by using the first linearization, we can obtain a second order conductivity equation. Via the known boundary determination from the localized DN map, one is able to recover σ on the specific region of the periphery. Since σ is piecewise real analytic, one can recover the conductivity piece by piece by utilizing the analyticity of coefficients. This implies that the global uniqueness result for σ holds. The arguments of combining monotonicity method with localized potentials was initiated by [Geb08] as applied to various inverse problems, such as [Har09, HS10, Har12, AH13, HU13, BHHM17, HU17, BHKS18, GH18, HPSar, HLL18, SKJ⁺19, HL19, HPS19]. Meanwhile, there are some related works build practical reconstruction methods on monotonicity properties [TR02, HLU15, HU15, HM16, MVVT16, TSV⁺16, Gar17, GS17, SUG⁺17, VMC⁺17, HM18, ZHS18, GS19].

We next introduce the mathematical model in this work. Let $\Omega \subset \mathbb{R}^n$ be a bounded connected domain with a C^∞ -smooth boundary $\partial\Omega$, for $n \geq 2$, and $D \Subset \Omega$ is an open subset such that $\Omega \setminus \overline{D}$ is connected. Let us consider the following semilinear elliptic equation with diffusion

$$(1.1) \quad \begin{cases} -\nabla \cdot (\sigma(x)\nabla u) + a(x, u) = 0 & \text{in } \Omega, \\ u = f & \text{on } \partial\Omega, \end{cases}$$

and

$$(1.2) \quad \begin{cases} -\nabla \cdot (\sigma(x)\nabla u) + a(x, u) = 0 & \text{in } \Omega \setminus \overline{D}, \\ u = 0 & \text{on } \partial D, \\ u = f & \text{on } \partial\Omega. \end{cases}$$

In this work, we first demonstrate the local well-posedness result of (1.1) and (1.2), whenever the diffusivity $\sigma = \sigma(x)$, and the lower order coefficients $a = a(x, y)$, which are *piecewise real-analytic* with respect to the x -variable, for any $y \in \mathbb{R}$. Throughout this paper, we shall use the term "analytic" as a synonym for "real-analytic", and the rigorous definition of the piecewise analytic is given in Appendix A.

Note that the local well-posedness of (1.1) and (1.2) was shown in [FO20, KU20a, LLLS21] for semilinear elliptic equations under different regularity assumptions. As a matter of fact, due to the discontinuities of the coefficients, one needs to treat the well-posedness of the Dirichlet boundary value problem (1.1) carefully, and we offer the proof in Section 2. Assuming the well-posedness, let $\Gamma \subset \partial\Omega$ be a relatively

open subset, $0 < \alpha < 1$, then there must exist constants $\varepsilon > 0$ and $C > 0$ such that

$$f \in N_\varepsilon := \{f \in C_0^\alpha(\Gamma) : \|f\|_{C^\alpha(\Gamma)} \leq \varepsilon\}.$$

In fact, we will show that given any boundary data $f \in N_\varepsilon$, there is a (unique) solution $u = u_f \in V$ solving (1.1), which satisfies

$$\|u\|_V \leq C\delta,$$

where

$$(1.3) \quad V := \{v \in H^1(\Omega) : -\nabla \cdot (\sigma \nabla v) \in L^\infty(\Omega), v|_{\partial\Omega} \in C_0^\alpha(\Gamma)\},$$

and

$$(1.4) \quad \|v\|_V := \|v\|_{H^1(\Omega)} + \|\nabla \cdot (\sigma \nabla v)\|_{L^\infty(\Omega)} + \|v|_{\partial\Omega}\|_{C^\alpha(\Gamma)}.$$

Moreover, the local well-posedness also holds for (1.2), when the domain Ω in (1.3) and (1.4) is replaced by $\Omega \setminus \overline{D}$.

With these notations at hand, by using the local well-posedness of (1.1) and (1.2), given $0 < \alpha < 1$, then one can define the corresponding (partial) *Dirichlet-Neumann*(DN) operator

$$\Lambda_{\sigma,a}^\Gamma : N_\varepsilon \rightarrow H^{-1/2}(\Gamma), \quad \Lambda_{\sigma,a}^\Gamma(f) := \sigma \partial_\nu u_f|_\Gamma,$$

for some sufficiently small number $\varepsilon > 0$, where u_f is the unique solution of (1.1) and ν is the unit outer normal on $\partial\Omega$. Meanwhile, one can also define the DN operator

$$\Lambda_{\sigma,a,D}^\Gamma : N_\varepsilon \rightarrow H^{-1/2}(\Gamma), \quad \Lambda_{\sigma,a,D}^\Gamma(f) := \sigma \partial_\nu u_f|_\Gamma,$$

for some sufficiently small number $\varepsilon > 0$, where u_f is the unique solution of (1.2) and ν is the unit outer normal on $\partial\Omega$. Then our inverse problems become that:

- (1) Can one identify both σ and a by knowing the partial measurements $\Lambda_{\sigma,a}^\Gamma$?
- (2) Can one identify both σ , a and D by knowing the partial measurements $\Lambda_{\sigma,a,D}^\Gamma$?

We will give affirmative answers to the preceding two questions in the rest of this paper.

The paper is structured as follows. In Section 2, we state our main results of this note, and the proofs are given in Section 3. The main methods depend on suitable linearization and monotonicity methods combining with localized potentials.

2. THE MAIN RESULTS

In this section, we will formulate our two main results: The semilinear elliptic equation (1.1) is uniquely solvable (for sufficiently small Dirichlet data), and the associated Dirichlet-Neumann operator uniquely determines the coefficients in equation (1.1).

Our results will be valid under the following assumptions on the domain and the coefficients.

Assumption 2.1. *We assume that $\Omega \subset \mathbb{R}^n$, $n \geq 2$ is a bounded domain with C^∞ -smooth boundary $\partial\Omega$, and that $\sigma \in L_+^\infty(\Omega)$ is a piecewise analytic function in the sense of [KV85, Section 3]. The function $a : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ is assumed to fulfill*

$$a(x, y) = \sum_{k=0}^{\infty} a_k(x) \frac{y^k}{k!},$$

with $a_k \in L^\infty(\Omega)$, $a_0 = a_1 = 0$, and $\sup_{k \geq 2} \|a_k\|_{L^\infty(\Omega)} < \infty$. Moreover each function a_k is assumed to be piecewise analytic in the sense of [KV85, Sect. 3].

Note that [KV85, Sect. 3] implies that two piecewise analytic functions are piecewise analytic with respect to the same partition, and this naturally extends to every finite number of piecewise analytic functions. However, assumption 2.1 contains infinitely many such functions σ, a_k ($k \in \mathbb{N}$), and we do not assume that they are piecewise analytic with respect to the same partition.

For our solvability result for the forward problem, we use the following solution space

$$V := \{v \in H^1(\Omega) : -\nabla \cdot (\sigma \nabla v) \in L^\infty(\Omega), v|_{\partial\Omega} \in C^\alpha(\partial\Omega)\}.$$

equipped with the norm

$$\|v\|_V := \|v\|_{H^1(\Omega)} + \|\nabla \cdot (\sigma \nabla v)\|_{L^\infty(\Omega)} + \|v|_{\partial\Omega}\|_{C^\alpha(\partial\Omega)}.$$

Clearly, V is a Banach space. Moreover, by a result of Li and Vogelius [LV00, Corollary 7.3],

$$(2.1) \quad V \subseteq H^1(\Omega) \cap L^\infty(\Omega) \quad \text{is continuously embedded.}$$

Theorem 2.1 (Local well-posedness of the forward problem). *Let $\Omega \subset \mathbb{R}^n$, $\sigma : \Omega \rightarrow \mathbb{R}$, and $a : \Omega \times \mathbb{R} \rightarrow \mathbb{R}$ fulfill assumption 2.1. Then there exists $\varepsilon > 0$ so that for all*

$$(2.2) \quad f \in N_\varepsilon := \{\phi \in C^\alpha(\partial\Omega) : \|\phi\|_{C^\alpha(\partial\Omega)} < \varepsilon\},$$

there exists a solution $u \in V \subseteq H^1(\Omega) \cap L^\infty(\Omega)$ to the Dirichlet problem

$$(2.3) \quad \begin{cases} -\nabla \cdot (\sigma \nabla u) + a(x, u) = 0 & \text{in } \Omega, \\ u = f & \text{on } \partial\Omega. \end{cases}$$

Moreover, there exists $\delta > 0$, so that, for all $f \in N_\varepsilon$, the solution is unique in the set of all

$$H_\delta := \{v \in H^1(\Omega) \cap L^\infty(\Omega) : \|v\|_{H^1(\Omega)} + \|v\|_{L^\infty(\Omega)} \leq \delta\},$$

and that the solution operator

$$\mathcal{S} : N_\varepsilon \rightarrow V \subseteq H^1(\Omega) \cap L^\infty(\Omega), \quad f \mapsto u, \quad \text{where } u \in H_\delta \text{ solves (2.3)}$$

is infinitely differentiable.

Clearly, $V \subseteq L^\infty(\Omega)$ implies that $x \mapsto a(x, u)$ is a $L^\infty(\Omega)$ -function for all $u \in V$. Hence, a solution $u \in V$ of (2.3) has well-defined Neumann boundary values, so that we can define the (non-linear) Dirichlet-Neumann-Operator

$$\Lambda_{\sigma,a} : N_\varepsilon \rightarrow H^{-1/2}(\partial\Omega), \quad \Lambda_{\sigma,a}(f) := \sigma \partial_\nu u|_{\partial\Omega},$$

where u solves (2.3). For an open boundary part $\Gamma \subseteq \partial\Omega$, the restriction of $\Lambda_{\sigma,a}$ to Γ is denoted by $\Lambda_{\sigma,a}^\Gamma$.

Theorem 2.2 (Unique identifiability from local boundary data). *Let $\Omega \subset \mathbb{R}^n$, and two set of coefficients (σ, a) , and $(\tilde{\sigma}, \tilde{a})$ each fulfill Assumption 2.1. Let $\varepsilon > 0$ be sufficiently small, so that both, $\Lambda_{\sigma,a}$ and $\Lambda_{\tilde{\sigma},\tilde{a}}$, are defined on N_ε .*

If there exists an open boundary part $\Gamma \subseteq \partial\Omega$, so that

$$\Lambda_{\sigma,a}^\Gamma(f) = \Lambda_{\tilde{\sigma},\tilde{a}}^\Gamma(f)$$

for all $f \in N_\varepsilon$ with $\text{supp}(f) \subseteq \Gamma$, then

$$\sigma = \tilde{\sigma} \quad \text{and} \quad a = \tilde{a}.$$

Remark 2.2. *Note that even for the full data case, that is, when $\Gamma = \partial\Omega$, Theorem 2.2 is also a new result to our best knowledge. Furthermore, for the case $\Gamma = \partial\Omega$, we can reduce the regularity of σ , by using the first linearization and results in [HT⁺13] when σ is Lipschitz continuous for $n \geq 3$, and [AP06] when $\sigma \in L^\infty(\Omega)$ for $n = 2$.*

Furthermore, we are also interested in a simultaneous recovering inverse obstacle problem. Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with a connected C^∞ boundary $\partial\Omega$ and $D \Subset \Omega$ be an open set with C^∞ boundary ∂D such that $\Omega \setminus \overline{D}$ is connected. Consider the following Dirichlet problem

$$(2.4) \quad \begin{cases} -\nabla \cdot (\sigma \nabla u) + a(x, u) = 0 & \text{in } \Omega \setminus \overline{D}, \\ u = 0 & \text{on } \partial D, \\ u = f & \text{on } \partial\Omega, \end{cases}$$

where $\sigma \in L_+^\infty(\Omega)$, and $a(x, z)$ are piecewise real analytic functions defined for any x in the domain $\overline{\Omega} \setminus D$ are piecewise analytic with respect to the x -variable. The well-posedness of (2.4) can be found in [KU20a, Appendix] and [LLLS21, Proposition 2.1] in a slightly different settings. In [KU20a, LLLS21], the proofs of the well-posedness under sufficiently smooth conditions of coefficients. However, in this paper, we need to deal with the regularity issues for our semilinear elliptic equations with piecewise analytic coefficients (see Section 2). More precisely, (2.4) admits a unique (small) solution $u \in C^{1,\alpha}(\overline{\Omega_m})$ provided that $f \in C^\alpha(\partial\Omega)$. Here we abuse the notations to denote that Ω_m 's are mutually disjoint subdomains of $\Omega \setminus \overline{D}$ such that $\overline{\Omega} \setminus D = \bigcup_{m=1}^M \overline{\Omega_m}$. Without loss of generality, one may consider

$\Gamma \subseteq \partial\Omega$ to be an arbitrary nonempty open subset. Thus, we are able to define the localized DN operator

$$\Lambda_{\sigma,a,D}^\Gamma : C_0^{1,\alpha}(\Gamma) \rightarrow H^{-1/2}(\Gamma), \quad \Lambda_{\sigma,a,D}^\Gamma(f) := \sigma \partial_\nu u_f|_\Gamma,$$

for $f \in N_\varepsilon$ and $\text{supp}(f) \subset \Gamma$, where u_f is the unique solution of (2.4) and ν is the unit outer normal on $\partial\Omega$. We want to determine the unknown diffusion σ , nonlinear term a and obstacle D simultaneously from the localized DN map $\Lambda_{\sigma,a,D}^\Gamma$.

The following theorem is analogous to [LLLS20, Theorem 1.2] and [KU20a, Theorem 1.6]. Here we are able to allow the diffusion coefficient σ to be piecewise analytic in $\overline{\Omega}$, but unknown apriori and that we can use partial DN operators on $\Gamma \subseteq \partial\Omega$.

Theorem 2.3 (Simultaneous recovering of coefficients and obstacle). *Let $\Omega \subset \mathbb{R}^n$, and two set of coefficients (σ, a) , and $(\tilde{\sigma}, \tilde{a})$ each fulfill Assumption 2.1 in $\Omega \setminus \overline{D}$ and $\Omega \setminus \overline{\tilde{D}}$, respectively. Let $\varepsilon > 0$ be sufficiently small, so that both, $\Lambda_{\sigma,a,D}^\Gamma$ and $\Lambda_{\tilde{\sigma},\tilde{a},\tilde{D}}^\Gamma$, are defined on N_ε .*

If there exists an open boundary part $\Gamma \subseteq \partial\Omega$, so that

$$\Lambda_{\sigma,a,D}^\Gamma(f) = \Lambda_{\tilde{\sigma},\tilde{a},\tilde{D}}^\Gamma(f)$$

for all $f \in N_\varepsilon$ with $\text{supp}(f) \subseteq \Gamma$, then

$$\sigma = \tilde{\sigma}, \quad a = \tilde{a} \quad \text{and} \quad D = \tilde{D}.$$

To our best knowledge, the preceding theorems are new results. The proof will be given prove in the next section. Since the DN map in Theorem 2.3 is incomplete, so there is a connection between the simultaneous recovery and the partial data inverse problems.

3. PROOFS OF MAIN RESULTS

To prove our two main results, we start with the following lemma, which will be used for our results.

Lemma 3.1. *The mapping*

$$G : V \rightarrow L^\infty(\Omega), \quad v(x) \mapsto a(x, v(x))$$

is infinitely differentiable and its l -th Fréchet derivative fulfills

$$G^{(l)}(v)(w_1, \dots, w_l) = \sum_{k=0}^{\infty} a_{k+l}(x) \frac{v(x)^k}{k!} w_1 \dots w_l \quad \text{for all } v, w \in V.$$

Proof. We also define for $l \in \mathbb{N}_0$

$$G_l : V \rightarrow L^\infty(\Omega), \quad v(x) \mapsto \sum_{k=0}^{\infty} a_{k+l}(x) \frac{v(x)^k}{k!}$$

Then $G_0 = G$, and for all $v \in V$, $l \in \mathbb{N}_0$, $G_l(v) \in L^\infty(\Omega)$ follows from $V \subseteq L^\infty(\Omega)$.

We will prove that G_l is one-time Fréchet differentiable and that its derivative $G'_l : V \rightarrow \mathcal{L}(V, L^\infty(\Omega))$ is given by

$$(3.1) \quad G'_l(v)w = M_w G_{l+1}(v) \quad \text{for all } v, w \in V,$$

where, for $w \in V \subseteq L^\infty(\Omega)$

$$M_w : L^\infty(\Omega) \rightarrow L^\infty(\Omega), \quad u \mapsto wu$$

denotes the continuous linear multiplication operator. Then the assertion follows by trivial induction.

Clearly, (3.1) defines a continuous linear operator $G'_l(v) \in \mathcal{L}(V, L^\infty(\Omega))$. To prove that this is indeed the Fréchet derivative of G_l , let $v, w \in V$, $x \in \Omega$, and define

$$\begin{aligned} \psi_x : \mathbb{R} &\rightarrow \mathbb{R}, \\ \psi_x(t) &:= G_l(v + t(w - v))(x) = \sum_{k=0}^{\infty} a_{k+l}(x) \frac{(v(x) + t(w(x) - v(x)))^k}{k!}. \end{aligned}$$

Then ψ is infinitely differentiable with

$$\begin{aligned} \psi'_x(t) &= \sum_{k=1}^{\infty} a_{k+l}(x) \frac{(v(x) + t(w(x) - v(x)))^{k-1}}{(k-1)!} (w(x) - v(x)) \\ \psi''_x(t) &= \sum_{k=2}^{\infty} a_{k+l}(x) \frac{(v(x) + t(w(x) - v(x)))^{k-2}}{(k-2)!} (w(x) - v(x))^2. \end{aligned}$$

Using

$$|v(x) + t(w(x) - v(x))| \leq |v(x)| + |w(x)| \quad \text{for all } t \in [0, 1],$$

and that a_k are uniformly bounded, and that $V \subseteq L^\infty(\Omega)$ is continuously embedded we have that

$$|\psi''_x(t)| \leq C \|w - v\|_V^2 \exp(\|v\|_V + \|w\|_V) \quad \text{for all } x \in \Omega.$$

Using Taylor's formula

$$|\psi_x(1) - \psi_x(0) - \psi'_x(0)| \leq \frac{1}{2} \max_{\tau \in [0, 1]} |\psi''_x(\tau)|,$$

we thus obtain

$$\|G_l(v) - G_l(w) - G'_l(v)(w - v)\|_{L^\infty(\Omega)} \leq \frac{C}{2} \|w - v\|_V^2 \exp(\|v\|_V + \|w\|_V),$$

so that the assertion is proven. \square

3.1. Local well-posedness result for the forward problem.

Proof of Theorem 2.1. We will apply the implicit function theorem to the map

$$\begin{aligned} F : C^\alpha(\partial\Omega) \times V &\rightarrow W := L^\infty(\Omega) \times C^\alpha(\partial\Omega), \\ F : (f, v) &\mapsto \nabla \cdot (-\sigma \nabla v) + a(x, v), v|_{\partial\Omega} - f. \end{aligned}$$

By Lemma 3.1 this mapping is well-defined and infinitely differentiable, and its derivative with respect to $v \in V$ is the continuous linear operator

$$D_v F(0, 0) : V \rightarrow W$$

with

$$D_v F(0, 0)u = (-\sigma \nabla u, u|_{\partial\Omega}).$$

Given $w = (w_1, w_2) \in W$ there exists a unique solution $u \in H^1$ of

$$-\sigma \nabla u = w_1, \quad \text{and} \quad u|_{\partial\Omega} = w_2.$$

Then $u \in V$ holds by definition of V , which shows that $D_v F(0, 0)$ is surjective. Since u is the unique solution, $D_v F(0, 0)$ is also injective. Since also $F(0, 0) = 0$ is fulfilled, we can apply the implicit function theorem (cf., e.g., [RR06, Sect. 10.1.1]), which yields an infinitely differentiable function

$$S : N_\varepsilon \rightarrow V$$

defined on a neighborhood of the origin $C^\alpha(\partial\Omega)$,

$$N_\varepsilon := \{\phi \in C^\alpha(\partial\Omega) : \|\phi\|_{C^\alpha(\partial\Omega)} < \varepsilon\},$$

so that

$$F(f, S(f)) = 0 \quad \text{for all } f \in N_\varepsilon,$$

and $S(f)$ is the only such element in a neighborhood of the origin in V . Since $F(f, S(f)) = 0$ implies that $S(f) \in V$ solves (2.3), the existence of a solution in $V \subseteq H^1(\Omega) \cap L^\infty(\Omega)$ is proven. Moreover, since every solution $u \in H^1(\Omega) \cap L^\infty(\Omega)$ of (2.3) fulfills

$$u \in V, \quad \text{and} \quad \|u\|_V = \|u\|_{H^1(\Omega)} + \sup_{k \geq 2} \|a_k\|_{L^\infty(\Omega)} e^{\|u\|_{L^\infty(\Omega)}} + \|f\|_{C^\alpha(\partial\Omega)},$$

the solution is unique in H_δ with sufficiently small δ . Finally, the solution operator S is also infinitely differentiable as a function from $N_\varepsilon \rightarrow H^1(\Omega) \cap L^\infty(\Omega)$. \square

3.2. Unique identifiability result for the inverse problem. To prove Theorem 2.2 we first derive some auxilliary results on the higher-order derivatives of the solution of (2.3). In the rest of this note, let us fix $\varepsilon > 0$ to be a sufficiently small number, such that the well-posedness for (1.1) and (1.2) hold, for any $f \in N_\varepsilon$.

Lemma 3.2. *Let $f_1, f_2 \in N_\varepsilon$, and define*

$$F : (-1/2, 1/2) \times (-1/2, 1/2) \rightarrow V, \quad F(t_1, t_2) := \mathcal{S}(t_1 f_1 + t_2 f_2).$$

Then

$$u_1^{(\ell)} := \partial_{t_\ell} F(0, 0) = \partial_{t_\ell} \mathcal{S}(t_1 f_1 + t_2 f_2)|_{t_1=t_2=0} \in V$$

solves

$$\begin{cases} \nabla \cdot (\sigma \nabla u_1^{(\ell)}) = 0 & \text{in } \Omega \\ u_1^{(\ell)} = f_\ell & \text{on } \partial\Omega, \end{cases}$$

for $\ell = 1, 2$. Moreover, for all $m > 1$, $m \in \mathbb{N}$,

$$u_m := \partial_{t_1}^2 \partial_{t_2}^{m-2} F(0, 0) = \partial_{t_1}^2 \partial_{t_2}^{m-2} \mathcal{S}(t_1 f_1 + t_2 f_2)|_{t_1=t_2=0} \in V$$

solves

$$\begin{cases} \nabla \cdot (\sigma \nabla u_m) = \partial_{t_1}^2 \partial_{t_2}^{m-2} G(F(t))|_{t_1=t_2=0} & \text{in } \Omega \\ u_m = 0 & \text{on } \partial\Omega. \end{cases}$$

Moreover, for all $f \in N_\varepsilon$, the mapping

$$(t_1, t_2) \mapsto \Lambda(t_1 f_1 + t_2 f_2), \quad (-1/2, 1/2) \times (-1/2, 1/2) \rightarrow H^{-1/2}(\partial\Omega)$$

is infinitely differentiable, and

$$\partial_{t_1}^2 \partial_{t_2}^{m-2} \Lambda(t_1 f_1 + t_2 f_2)|_{t_1=t_2=0} = \sigma \partial_\nu u_m|_{\partial\Omega}.$$

Proof. Note that F , G , and \mathcal{S} are infinitely differentiable functions by Lemma 3.1, and Theorem 2.1. Moreover, since the trace operator $u \mapsto u|_{\partial\Omega}$ is a continuous linear function from V to $H^{1/2}(\Omega)$, it follows that

$$u_1^{(\ell)}|_{\partial\Omega} = f_\ell, \text{ for } \ell = 1, 2, \quad \text{and} \quad u_m|_{\partial\Omega} = 0, \text{ for } m \geq 2.$$

Let $v \in H_0^1(\Omega)$. Since $u := \mathcal{S}(t_1 f_1 + t_2 f_2)$ solves (2.3) we have that

$$\begin{aligned} 0 &= \int_{\Omega} \sigma \nabla u \cdot \nabla v \, dx + \int_{\Omega} a(x, u) v \, dx \\ &= \int_{\Omega} \sigma \nabla \mathcal{S}(t_1 f_1 + t_2 f_2) \cdot \nabla v \, dx + \int_{\Omega} G(\mathcal{S}(t_1 f_1 + t_2 f_2)) v \, dx. \end{aligned}$$

Noting that the first summand is a linear continuous functional with respect to $\mathcal{S}(t_1 f_1 + t_2 f_2) \in V$, and the second summand is linear and continuous with respect to $G(\mathcal{S}(t_1 f_1 + t_2 f_2)) \in L^\infty(\Omega)$, we obtain by differentiation

$$\begin{aligned} 0 &= \int_{\Omega} \sigma \nabla \partial_{t_1}^2 \partial_{t_2}^{m-2} \mathcal{S}(t_1 f_1 + t_2 f_2)|_{t_1=t_2=0} \cdot \nabla v \, dx \\ &\quad + \int_{\Omega} \partial_{t_1}^2 \partial_{t_2}^{m-2} G(\mathcal{S}(t_1 f_1 + t_2 f_2))|_{t_1=t_2=0} v \, dx \\ &= \int_{\Omega} \sigma \nabla u_m \cdot \nabla v \, dx + \int_{\Omega} \partial_{t_1}^2 \partial_{t_2}^{m-2} G(\mathcal{S}(t_1 f_1 + t_2 f_2))|_{t_1=t_2=0} v \, dx. \end{aligned}$$

This proves that, for all $m > 1$,

$$\nabla \cdot (\sigma \nabla u_m) = \partial_{t_1}^2 \partial_{t_2}^{m-2} G(\mathcal{S}(t_1 f_1 + t_2 f_2))|_{t_1=t_2=0}.$$

On the other hand, for $\ell = 1, 2$, it follows from $S(0) = 0$, and $G'(0) = 0$, that

$$\partial_{t_\ell} G(\mathcal{S}(t_1 f_1 + t_2 f_2))|_{t_1=t_2=0} = G'(\mathcal{S}(0)) (\partial_{t_\ell} \mathcal{S}(0)) f_\ell = 0.$$

Hence, $u_1^{(\ell)} \in V$ solves $\nabla \cdot (\sigma \nabla u_1^{(\ell)}) = 0$, for $\ell = 1, 2$.

Moreover, for all $f_1, f_2 \in N_\varepsilon$, and $(t_1, t_2) \in (-1/2, 1/2) \times (-1/2, 1/2)$, the Neumann data $\Lambda_{\sigma,a}(t_1 f_1 + t_2 f_2) \in H^{-1/2}(\partial\Omega)$ fulfills

$$\langle \Lambda_{\sigma,a}(t_1 f_1 + t_2 f_2), g \rangle = \int_{\Omega} \sigma \nabla F(t_1, t_2) \cdot \nabla v \, dx + \int_{\Omega} G(F(t_1, t_2)) v(x) \, dx$$

for all $v \in H^1(\Omega)$. As above, the first summand is a linear continuous functional with respect to $F(t_1, t_2) \in V$, and the second summand is linear and continuous with respect to $G(F(t_1, t_2)) \in L^\infty(\Omega)$. Hence, $(t_1, t_2) \mapsto \Lambda(t_1 f_1 + t_2 f_2)$ is infinitely

differentiable with respect to t_1 and t_2 , and, for all $g \in H^{1/2}(\partial\Omega)$,

$$\begin{aligned} & \left\langle \partial_{t_1}^2 \partial_{t_2}^{m-2} \Lambda_{\sigma,a}(t_1 f_1 + t_2 f_2) \Big|_{t_1=t_2=0} : g \right\rangle \\ &= \partial_{t_1}^2 \partial_{t_2}^{m-2} \langle \Lambda_{\sigma,a}(t_1 f_1 + t_2 f_2), g \rangle \Big|_{t_1=t_2=0} \\ &= \int_{\Omega} \sigma \nabla u_m \cdot \nabla v \, dx + \int_{\Omega} \partial_{t_1}^2 \partial_{t_2}^{m-2} G(F(t_1, t_2)) \Big|_{t_1=t_2=0} v(x) \, dx \\ &= \langle \sigma \partial_{\nu} u_m \Big|_{\partial\Omega}, g \rangle, \end{aligned}$$

which proves that $\partial_{t_1}^2 \partial_{t_2}^{m-2} \Lambda_{\sigma,a}(t_1 f_1 + t_2 f_2) \Big|_{t_1=t_2=0} = \partial_{\nu} u_m \Big|_{\partial\Omega}$ as desired. \square

Lemma 3.3. *Let $m \in \mathbb{N}$. There exist numbers $a_{m,j}^{p_1, p'_1, \dots, p_j, p'_j} \in \mathbb{N}_0$ (depending on $j = 1, \dots, m$, and $p_1, p'_1, \dots, p_j, p'_j \in \mathbb{N} \cup \{0\}$ with $p_1 + p'_1 + \dots + p_j + p'_j = m$) so that*

$$\begin{aligned} & \partial_{t_1}^2 \partial_{t_2}^{m-2} G(F(t_1, t_2)) \\ &= G^{(m)}(F(t_1, t_2)) \underbrace{(\partial_{t_1} F(t_1, t_2), \partial_{t_1} F(t_1, t_2), \partial_{t_2} F(t_1, t_2), \dots, \partial_{t_2} F(t_1, t_2))}_{m\text{-tuples}} \\ &+ \sum_{j=2}^{m-1} \sum_{\substack{p_1, p'_1, \dots, p_j, p'_j \in \mathbb{N} \cup \{0\}, \\ p_1 + \dots + p_j = 2, \quad p'_1 + p'_2 + \dots + p'_j = m-2}} a_{m,j}^{p_1, p'_1, \dots, p_j, p'_j} G^{(j)}(F(t_1, t_2)) \\ &\quad \times \underbrace{\left(\partial_{t_1}^{p_1} \partial_{t_2}^{p'_1} F(t_1, t_2), \dots, \partial_{t_1}^{p_j} \partial_{t_2}^{p'_j} F(t_1, t_2) \right)}_{j\text{-tuples}} \\ &+ G'(F(t_1, t_2)) (\partial_{t_1}^2 \partial_{t_2}^{m-2} F(t_1, t_2)). \end{aligned}$$

Proof. This follows by induction using the chain rule for the Fréchet derivative. \square

We also need the following variant of the localized potentials result in [Geb08]:

Lemma 3.4. *Let D_1, D_2 be two disjoint non-empty sets, where $D_1 \subseteq \Omega$ is open, $D_2 \subseteq \bar{\Omega}$ is closed, $\Omega \setminus D_2$ is connected, and $\Gamma \cap \bar{\Omega} \setminus D_2 \neq \emptyset$. Then there exists a sequence $(\phi_k)_{k \in \mathbb{N}} \subset C^\alpha(\partial\Omega)$ with $\text{supp}(\phi_k) \subseteq \Gamma$, and*

$$\int_{D_2} |v_k|^2 \, dx \rightarrow 0, \quad \text{and} \quad \int_{D_1} |v_k|^2 \, dx \rightarrow \infty,$$

Proof. Let $C_0^\alpha(\Gamma)$ denote the closure of the space of all $\phi \in C^\alpha(\partial\Omega)$ with $\phi \subseteq \Gamma$ with respect to the $C^\alpha(\Gamma)$ -norm. For $j = 1, 2$, we define $A_j \in \mathcal{L}(C_0^\alpha(\Gamma), L^2(D_j))$ by $A_j : \phi \mapsto v|_{D_j}$, where $v \in V$ solves

$$\begin{cases} \nabla \cdot (\sigma \nabla v) = 0 & \text{in } \Omega, \\ v = \phi & \text{on } \partial\Omega. \end{cases}$$

Note that V is continuously embedded in $L^2(D_j)$, so that A_j are well-defined.

The assertion is proven if we can show that

$$\exists C > 0 : \|A_1 \phi\|_{L^2(D_1)} \leq C \|A_2 \phi\|_{L^2(D_2)},$$

and, by [Geb08, Lemma 2.5], this is equivalent to proving

$$\mathcal{R}(A'_1) \not\subseteq \mathcal{R}(A'_2).$$

The operators A'_j are easily checked to map a source term $\psi \in L^2(D_j)$ to the Neumann boundary values of the solution of $\nabla \cdot (\sigma \nabla w) = \psi$ with zero Dirichlet data. By a standard unique continuation argument, it then follows that $0 = \mathcal{R}(A'_1) \cap \mathcal{R}(A'_2)$, which proves the assertion. \square

Now we can prove our simultaneously unique identifiability result for the inverse coefficient problem.

Proof of Theorem 2.2. Let $\Omega \subset \mathbb{R}^n$, and two set of coefficients (σ, a) , and $(\tilde{\sigma}, \tilde{a})$ each fulfill assumption 2.1. Let $\epsilon > 0$ be sufficiently small, so that both, $\Lambda_{\sigma, a}$ and $\Lambda_{\tilde{\sigma}, \tilde{a}}$, are defined on N_ϵ .

Given $f \in N_\epsilon$, we define the operators F, G, \mathcal{S} , and the functions $u_m \in V$ ($m \in \mathbb{N}$) as in Lemma 3.2, Lemma 3.1, and Theorem 2.1 using the coefficient pair (σ, a) . The corresponding entities with (σ, a) replaced by $(\tilde{\sigma}, \tilde{a})$ will be denoted by $\tilde{F}, \tilde{G}, \tilde{\mathcal{S}}$, and \tilde{u}_m ($m \in \mathbb{N}$).

We will show that

(a) If

$$(3.2) \quad \partial_{t_\ell} \Lambda_{\sigma, a}^\Gamma(t_1 f_1 + t_2 f_2) \Big|_{\Gamma} \Big|_{t_1=t_2=0} = \partial_{t_\ell} \Lambda_{\tilde{\sigma}, \tilde{a}}^\Gamma(t_1 f_1 + t_2 f_2) \Big|_{\Gamma} \Big|_{t_1=t_2=0},$$

for all $f_1, f_2 \in N_\epsilon$, then $\sigma = \tilde{\sigma}$, and $u_1^{(\ell)} = \tilde{u}_1^{(\ell)}$, for $\ell = 1, 2$.

(b) If, for some $m \geq 2$, $\sigma = \tilde{\sigma}$, $a_j = \tilde{a}_j$, $u_j = \tilde{u}_j$ for all $j = 1, \dots, m-1$, and

$$\partial_{t_1}^2 \partial_{t_2}^{m-2} \Lambda_{\sigma, a}^\Gamma(t_1 f_1 + t_2 f_2) \Big|_{\Gamma} \Big|_{t_1=t_2=0} = \partial_{t_1}^2 \partial_{t_2}^{m-2} \Lambda_{\tilde{\sigma}, \tilde{a}}^\Gamma(t_1 f_1 + t_2 f_2) \Big|_{\Gamma} \Big|_{t_1=t_2=0},$$

for all $f_1, f_2 \in N_\epsilon$, then $a_m = \tilde{a}_m$, and $u_m = \tilde{u}_m$.

Clearly, this proves Theorem 2.2 by induction, since we assumed that $a_0 = 0 = \tilde{a}_0$ and $a_1 = 0 = \tilde{a}_1$.

To show (a), note that (3.2) implies that the local DN operator for the linear elliptic equation $\nabla \cdot (\sigma \nabla v) = 0$ is the same for the two coefficients σ and $\tilde{\sigma}$. This implies $\sigma = \tilde{\sigma}$ by the classical Kohn-Vogelius result [KV85], and the uniqueness of solutions yield that $u_1^{(\ell)} = \tilde{u}_1^{(\ell)}$ in Ω , for $\ell = 1, 2$.

To prove (b), note that

$$G'(F(0, 0))(\partial_{t_1}^2 \partial_{t_2}^{m-2} F(0, 0)) = 0 = \tilde{G}'(\tilde{F}(0, 0))(\partial_{t_1}^2 \partial_{t_2}^{m-2} \tilde{F}(0, 0))$$

since $F(0, 0) = 0 = \tilde{F}(0, 0)$ and $G'(0) = 0 = \tilde{G}'(0)$. Moreover,

$$\partial_{t_1}^{p_1} \partial_{t_2}^{p_2} F(0, 0) = u_p = \tilde{u}_p = \partial_{t_1}^{p_1} \partial_{t_2}^{p_2} \tilde{F}(0, 0), \quad \text{for all } p_1 + p_2 = 1, \dots, m-1,$$

and, for all w_1, \dots, w_j ,

$$\begin{aligned} G^{(j)}(F(0, 0))(w_1, \dots, w_l) &= a_j(x) w_1 \dots w_l \\ &= \tilde{a}_j(x) w_1 \dots w_l = \tilde{G}^{(j)}(\tilde{F}(0))(w_1, \dots, w_l). \end{aligned}$$

Using Lemma 3.3 we thus obtain

$$\begin{aligned} & \partial_{t_1}^2 \partial_{t_2}^{m-2} G(F(t_1, t_2)) \Big|_{\Gamma} \Big|_{t_1=t_2=0} - \partial_{t_1}^2 \partial_{t_2}^{m-2} \tilde{G}(\tilde{F}(t_1, t_2)) \Big|_{\Gamma} \Big|_{t_1=t_2=0} \\ &= G^{(m)}(0) (\partial_{t_1} F(t_1, t_2), \partial_{t_1} F(t_1, t_2), \partial_{t_2} F(t_1, t_2), \dots, \partial_{t_2} F(t_1, t_2)) \Big|_{\Gamma} \Big|_{t_1=t_2=0} \\ & \quad - \tilde{G}^{(m)}(0) (\partial_{t_1} \tilde{F}(t_1, t_2), \partial_{t_1} \tilde{F}(t_1, t_2), \partial_{t_2} \tilde{F}(t_1, t_2), \dots, \partial_{t_2} \tilde{F}(t_1, t_2)) \Big|_{\Gamma} \Big|_{t_1=t_2=0} \\ &= a_m u_1^m - \tilde{a}_m \tilde{u}_1^m = (a_m - \tilde{a}_m) u_1^m. \end{aligned}$$

Hence, $u_m - \tilde{u}_m \in V$ solves

$$(3.3) \quad \begin{aligned} 0 &= -\nabla \cdot (\sigma \nabla u_m) + \partial_{t_1}^2 \partial_{t_2}^{m-2} G(F(t)) \Big|_{\Gamma} \Big|_{t_1=t_2=0} \\ & \quad + \nabla \cdot (\sigma \nabla \tilde{u}_m) - \partial_{t_1}^2 \partial_{t_2}^{m-2} \tilde{G}(\tilde{F}(t)) \Big|_{\Gamma} \Big|_{t_1=t_2=0} \\ &= -\nabla \cdot (\sigma \nabla (u_m - \tilde{u}_m)) + (a_m - \tilde{a}_m) \left(u_1^{(1)} \right)^2 \left(u_1^{(2)} \right)^{m-2}. \end{aligned}$$

For all $g \in C^\alpha(\Gamma)$ with $\text{supp}(g) \subset \Gamma$, we thus obtain

$$\begin{aligned}
(3.4) \quad 0 &= \left\langle \partial_{t_1}^2 \partial_{t_2}^{m-2} \Lambda_{\sigma,a}(t_1 f_1 + t_2 f_2)|_{\Gamma}|_{t_1=t_2=0}, g \right\rangle \\
&\quad - \left\langle \partial_{t_1}^2 \partial_{t_2}^{m-2} \Lambda_{\tilde{\sigma},\tilde{a}}(t_1 f_1 + t_2 f_2)|_{\Gamma}|_{t_1=t_2=0}, g \right\rangle \\
&= \langle \sigma \partial_\nu(u_m - \tilde{u}_m)|_{\Gamma}, g \rangle - \langle \sigma \partial_\nu \tilde{u}_m|_{\Gamma}, g \rangle \\
&= \int_{\Omega} \sigma \nabla(u_m - \tilde{u}_m) \cdot \nabla v_1 \, dx + \int_{\Omega} (a_m - \tilde{a}_m) \left(u_1^{(1)}\right)^2 \left(u_1^{(2)}\right)^{m-2} v_1 \, dx \\
&= \int_{\Omega} (a_m - \tilde{a}_m) \left(u_1^{(1)}\right)^2 \left(u_1^{(2)}\right)^{m-2} v_1 \, dx,
\end{aligned}$$

where $v_1 \in V$ solves

$$\begin{cases} \nabla \cdot (\sigma \nabla v_1) = 0 & \text{in } \Omega, \\ v_1 = g & \text{on } \partial\Omega. \end{cases}$$

We will now show that this implies $a_m = \tilde{a}_m$. Clearly this also implies $u_m = \tilde{u}_m$ by using (3.3) and $u_m|_{\partial\Omega} = f = \tilde{u}_m|_{\partial\Omega}$.

Assume that this is not the case. Since a_m and \tilde{a}_m are piecewise analytic, we can choose two disjoint non-empty sets D_1, D_2 , where $D_1 \subseteq \Omega$ is open, $D_2 \subseteq \overline{\Omega}$ is closed, $\Omega \setminus D_2$ is connected, and $\Gamma \cap \overline{\Omega} \setminus D_2 \neq \emptyset$ such that either

- (i) $a_m|_{\Omega \setminus D_2} \geq \tilde{a}_m|_{\Omega \setminus D_2}$ and $(a_m - \tilde{a}_m)|_{D_1} \in L_+^\infty(D_1)$, or
- (ii) $a_m|_{\Omega \setminus D_2} \leq \tilde{a}_m|_{\Omega \setminus D_2}$ and $(\tilde{a}_m - a_m)|_{D_1} \in L_+^\infty(D_1)$,

cf. [HU13, Appendix A] for a proof for $\Gamma = \partial\Omega$ that also holds for arbitrarily small open boundary pieces $\Gamma \subset \partial\Omega$. Without loss of generality we will assume that (i) holds true in the following.

For $m \in \mathbb{N}$, let us choose a non-negative, but not identically zero, function $\psi \in N_\varepsilon \subset C^\alpha(\partial\Omega)$ with $\text{supp}(\psi) \subseteq \Gamma$. By the strong maximum principle, the corresponding solution $w \in V$ of $\nabla \cdot (\sigma \nabla w) = 0$ in Ω , with $w|_{\partial\Omega} = \psi$, will be positive inside Ω . Then we can use the localized potentials result in Lemma 3.4 to obtain a sequence $(\phi_k)_{k \in \mathbb{N}} \subset C^\alpha(\partial\Omega)$ with $\text{supp}(\phi_k) \subseteq \Gamma$, and

$$\int_{D_2} w_{1,k}^2 \, dx \rightarrow 0, \quad \text{and} \quad \int_{D_1} w_{1,k}^2 \, dx \rightarrow \infty,$$

where $w_{1,k} \in V$ solves $\nabla \cdot (\sigma \nabla w_{1,k}) = 0$ in Ω with $w_{1,k}|_{\partial\Omega} = \phi_k$.

Using the Dirichlet data

$$f_{1,k} := \frac{\varepsilon}{2\|\phi_k\|_{C^\alpha(\partial\Omega)}} \phi_k \in N_\varepsilon, \quad f_2 = \psi \quad \text{and} \quad g_k := \left(\frac{2\|\phi_k\|_{C^\alpha(\partial\Omega)}}{\varepsilon} \right)^2 \psi,$$

such that solutions of the first linearized equation satisfy $u_{1,k}|_{\partial\Omega} = f_{1,k}$, $u_2|_{\partial\Omega} = f_2$ and $v_{1,k}|_{\partial\Omega} = g_k$, then we can obtain from (3.4) that

$$\begin{aligned}
0 &= \int_{\Omega} (a_m - \tilde{a}_m) u_{1,k}^2 u_2^{m-2} v_{1,k} \, dx \\
&\geq \int_{D_2} (a_m - \tilde{a}_m) u_{1,k}^2 u_2^{m-2} v_{1,k} \, dx + \int_{D_1} (a_m - \tilde{a}_m) u_{1,k}^2 u_2^{m-2} v_{1,k} \, dx \rightarrow \infty,
\end{aligned}$$

as $k \rightarrow \infty$. Here we have used the nonnegativity of ψ such that u_2 and $v_{1,k}$ are positive for any $k \in \mathbb{N}$. This contradiction shows that (b) holds. Hence, the proof is complete. \square

3.3. Unique identifiability result for the inverse obstacle problem. Finally, let us prove Theorem 2.3.

Proof of Theorem 2.3. Similar to the proof of Theorem 2.2, one can determine that $\sigma = \tilde{\sigma}$ in $\Omega \setminus (D \cup \tilde{D})$ by applying the boundary determination to piecewise analytic functions. Let us denote that u_1 and \tilde{u}_1 to be the solution of $\nabla \cdot (\sigma \nabla u_1) = 0$ and $\nabla \cdot (\tilde{\sigma} \nabla \tilde{u}_1) = 0$ in Ω , respectively. By using the strategy introduced in [LLLS20], let G be the connected component of $\Omega \setminus (\overline{D_1} \cup \overline{D_2})$ whose boundary contains $\partial\Omega$. Let $U := u_1 - \tilde{u}_1$, then U is a solution of

$$\begin{cases} \nabla \cdot (\sigma \nabla U) = 0 & \text{in } G, \\ U = \sigma \partial_\nu U = 0 & \text{on } \Gamma, \end{cases}$$

where we have utilized $\Lambda_{\sigma, a, D}^\Gamma(f) = \Lambda_{\tilde{\sigma}, \tilde{a}, \tilde{D}}^\Gamma(f)$, for any $f \in N_\varepsilon$. By the unique continuation for second order elliptic equations, one has that $U \equiv 0$ in G . Therefore,

$$(3.5) \quad u_1 = \tilde{u}_1 \text{ in } G.$$

We next prove $D = \tilde{D}$ via a contradiction argument. Suppose not, i.e., $D \neq \tilde{D} \Subset \Omega$. By using [LLLS20, Lemma A.1] or [KU20a, Section 4], without loss of generality, we may assume that there exists a point x_1 such that

$$x_1 \in \partial G \cap (\Omega \setminus \overline{D}) \cap \partial \tilde{D}.$$

$\tilde{u}_1(x_1) = 0$ since $x_1 \in \partial D_2$. By (3.5), we have $u_1(x_1) = 0$. Note that x_1 is an interior point of the open set $\Omega \setminus \overline{D_1}$. Consider the boundary values $u_1|_\Gamma \geq 0$ such that $u_1|_\Gamma \not\equiv 0$. Now, since $u_1(x_1) = 0$, by the maximum principle, we have that $u_1 \equiv 0$ in the connected open set $\Omega \setminus \overline{D_1}$, which contradicts with the nonzero boundary condition on Γ . Therefore, the conclusion $D = \tilde{D}$ must hold. Furthermore, we have by (3.5) that

$$(3.6) \quad u_1 = \tilde{u}_1 \text{ in } \Omega \setminus \overline{D},$$

as desired. Finally, by repeating the same arguments as in the proof of Theorem 2.2, we can show that $a = \tilde{a}$ in $\Omega \setminus \overline{D} \times \mathbb{R}$ as we wish. This proves the assertion. \square

APPENDIX A. PIECEWISE ANALYTICITY

For the sake of completeness, let us recall the piecewise analyticity introduced in [KV85, Section 3]. The piecewise analytic conditions of both $\sigma(x)$ and $a(x, y)$, for $x \in \overline{\Omega}$, there must exist a piecewise analytic cover $\{\Omega_m\}_{m=1}^M$ such that

$$\overline{\Omega} = \bigcup_{m=1}^M \overline{\Omega}_m,$$

such that σ , and $a(x, z)$ are analytic for $x \in \Omega_m$, for $m = 1, 2, \dots, M$, for some $M \in \mathbb{N}$. Throughout this work, we further assume that each boundaries of $\partial\Omega_m$ is C^2 . Notice that σ and $a(x, z)$ might have jump discontinuities across each boundaries of subdomains $\partial\Omega_m \subset \overline{\Omega}$, for $m = 1, 2, \dots, M$.

Acknowledgement. Y.-H. Lin is partially supported by the Ministry of Science and Technology Taiwan, under the Columbus Program: MOST-110-2636-M-009-007.

REFERENCES

- [AH13] Lilian Arnold and Bastian Harrach. Unique shape detection in transient eddy current problems. *Inverse Problems*, 29(9):095004, 2013.
- [AP06] Kari Astala and Lassi Päivärinta. Calderón’s inverse conductivity problem in the plane. *Annals of Mathematics*, pages 265–299, 2006.
- [BHHM17] Andrea Barth, Bastian Harrach, Nuutti Hyvönen, and Lauri Mustonen. Detecting stochastic inclusions in electrical impedance tomography. *Inverse Problems*, 33(11):115012, 2017.
- [BHKS18] Tommi Brander, Bastian Harrach, Manas Kar, and Mikko Salo. Monotonicity and enclosure methods for the p -Laplace equation. *SIAM J. Appl. Math.*, 78(2):742–758, 2018.
- [CFK⁺21] Cătălin I Cârstea, Ali Feizmohammadi, Yavar Kian, Katya Krupchyk, and Gunther Uhlmann. The Calderón inverse problem for isotropic quasilinear conductivities. *Advanced in Mathematics*, 391:Paper No. 107956., 2021.
- [FO20] Ali Feizmohammadi and Lauri Oksanen. An inverse problem for a semi-linear elliptic equation in Riemannian geometries. *Journal of Differential Equations*, 296(6):4683–4719, 2020.
- [Gar17] Henrik Garde. Comparison of linear and non-linear monotonicity-based shape reconstruction using exact matrix characterizations. *Inverse Problems in Science and Engineering*, pages 1–18, 2017.
- [Geb08] Bastian Gebauer. Localized potentials in electrical impedance tomography. *Inverse Probl. Imaging*, 2(2):251–269, 2008.
- [GH18] Roland Griesmaier and Bastian Harrach. Monotonicity in inverse medium scattering on unbounded domains. *SIAM J. Appl. Math.*, 78(5):2533–2557, 2018.
- [GS17] Henrik Garde and Stratos Staboulis. Convergence and regularization for monotonicity-based shape reconstruction in electrical impedance tomography. *Numerische Mathematik*, 135(4):1221–1251, 2017.
- [GS19] Henrik Garde and Stratos Staboulis. The regularized monotonicity method: Detecting irregular indefinite inclusions. *Inverse Probl. Imaging*, 13(1):93–116, 2019.
- [Har09] Bastian Harrach. On uniqueness in diffuse optical tomography. *Inverse Problems*, 25:055010 (14pp), 2009.
- [Har12] Bastian Harrach. Simultaneous determination of the diffusion and absorption coefficient from boundary data. *Inverse Probl. Imaging*, 6(4):663–679, 2012.
- [HL19] Bastian Harrach and Yi-Hsuan Lin. Monotonicity-based inversion of the fractional Schrödinger equation I. Positive potentials. *SIAM Journal on Mathematical Analysis*, 51(4):3092–3111, 2019.
- [HLL18] Bastian Harrach, Yi-Hsuan Lin, and Hongyu Liu. On localizing and concentrating electromagnetic fields. *SIAM J. Appl. Math.*, 78(5):2558–2574, 2018.
- [HLU15] Bastian Harrach, Eunjung Lee, and Marcel Ullrich. Combining frequency-difference and ultrasound modulated electrical impedance tomography. *Inverse Problems*, 31(9):095003, 2015.
- [HM16] Bastian Harrach and Mach Nguyet Minh. Enhancing residual-based techniques with shape reconstruction features in electrical impedance tomography. *Inverse Problems*, 32(12):125002, 2016.
- [HM18] Bastian Harrach and Mach Nguyet Minh. Monotonicity-based regularization for phantom experiment data in electrical impedance tomography. In *New Trends in Parameter Identification for Mathematical Models*, pages 107–120. Springer, 2018.
- [HPS19] Bastian Harrach, Valter Pohjola, and Mikko Salo. Dimension bounds in monotonicity methods for the Helmholtz equation. *SIAM Journal on Mathematical Analysis*, 51(4):2995–3019, 2019.
- [HPSar] Bastian Harrach, Valter Pohjola, and Mikko Salo. Monotonicity and local uniqueness for the Helmholtz equation. *Anal. PDE*, to appear.
- [HS10] Bastian Harrach and Jin Keun Seo. Exact shape-reconstruction by one-step linearization in electrical impedance tomography. *SIAM Journal on Mathematical Analysis*, 42(4):1505–1518, 2010.
- [HT⁺13] Boaz Haberman, Daniel Tataru, et al. Uniqueness in Calderón’s problem with Lipschitz conductivities. *Duke Mathematical Journal*, 162(3):497–516, 2013.
- [HU13] Bastian Harrach and Marcel Ullrich. Monotonicity-based shape reconstruction in electrical impedance tomography. *SIAM Journal on Mathematical Analysis*, 45(6):3382–3403, 2013.
- [HU15] Bastian Harrach and Marcel Ullrich. Resolution guarantees in electrical impedance tomography. *IEEE Trans. Med. Imaging*, 34:1513–1521, 2015.

- [HU17] Bastian Harrach and Marcel Ullrich. Local uniqueness for an inverse boundary value problem with partial data. *Proceedings of the American Mathematical Society*, 145(3):1087–1095, 2017.
- [IN95] Victor Isakov and A Nachman. Global uniqueness for a two-dimensional elliptic inverse problem. *Trans. of AMS*, 347:3375–3391, 1995.
- [IS94] Victor Isakov and John Sylvester. Global uniqueness for a semilinear elliptic inverse problem. *Communications on Pure and Applied Mathematics*, 47(10):1403–1410, 1994.
- [Isa93] Victor Isakov. On uniqueness in inverse problems for semilinear parabolic equations. *Archive for Rational Mechanics and Analysis*, 124(1):1–12, 1993.
- [IY13] Oleg Imanuvilov and Masahiro Yamamoto. Unique determination of potentials and semilinear terms of semilinear elliptic equations from partial Cauchy data. *J. Inverse Ill-Posed Probl.*, 21(1):85–108, 2013.
- [KKU20] Yavar Kian, Katya Krupchyk, and Gunther Uhlmann. Partial data inverse problems for quasilinear conductivity equations. *arXiv preprint arXiv:2010.11409*, 2020.
- [KN02] Hyeonbae Kang and Gen Nakamura. Identification of nonlinearity in a conductivity equation via the Dirichlet-to-Neumann map. *Inverse Problems*, 18(4):1079, 2002.
- [KU20a] Katya Krupchyk and Gunther Uhlmann. Partial data inverse problems for semilinear elliptic equations with gradient nonlinearities. *Mathematical Research Letters*, 27(6), 2020.
- [KU20b] Katya Krupchyk and Gunther Uhlmann. A remark on partial data inverse problems for semilinear elliptic equations. *Proceedings of the American Mathematical Society*, 148(2):681–685, 2020.
- [KV85] Robert V Kohn and Michael Vogelius. Determining conductivity by boundary measurements II. Interior results. *Communications on Pure and Applied Mathematics*, 38(5):643–667, 1985.
- [Lin21] Yi-Hsuan Lin. Monotonicity-based inversion of fractional semilinear elliptic equations with power type nonlinearities. *Calc. Var. Partial Differential Equations*, *accepted for publication*, 2021.
- [LL19] Ru-Yu Lai and Yi-Hsuan Lin. Global uniqueness for the fractional semilinear Schrödinger equation. *Proceedings of the American Mathematical Society*, 147(3):1189–1199, 2019.
- [LL21] Ru-Yu Lai and Yi-Hsuan Lin. Inverse problems for fractional semilinear elliptic equations. *Nonlinear Analysis*, *accepted for publication*, 2021.
- [LLL21] Yi-Hsuan Lin, Hongyu Liu, and Xu Liu. Determining a nonlinear hyperbolic system with unknown sources and nonlinearity. *arXiv preprint arXiv:2107.10219*, 2021.
- [LLLS20] Matti Lassas, Tony Liimatainen, Yi-Hsuan Lin, and Mikko Salo. Partial data inverse problems and simultaneous recovery of boundary and coefficients for semilinear elliptic equations. *Revista Matemática Iberoamericana*, 37(4):1553–1580, 2020.
- [LLLS21] Matti Lassas, Tony Liimatainen, Yi-Hsuan Lin, and Mikko Salo. Inverse problems for elliptic equations with power type nonlinearities. *Journal de mathématiques pures et appliquées*, 145:44–82, 2021.
- [LLLZ21] Yi-Hsuan Lin, Hongyu Liu, Xu Liu, and Shen Zhang. Simultaneous recoveries for semilinear parabolic systems. *arXiv preprint arXiv:2111.05242*, 2021.
- [LLS20] Matti Lassas, Tony Liimatainen, and Mikko Salo. The Poisson embedding approach to the Calderón problem. *Mathematische Annalen*, 377(1):19–67, 2020.
- [LLST21] Tony Liimatainen, Yi-Hsuan Lin, Mikko Salo, and Teemu Tyni. Inverse problems for elliptic equations with fractional power type nonlinearities. *J. Differential Equations*, *accepted for publication*, 2021.
- [LV00] Yan Yan Li and Michael Vogelius. Gradient estimates for solutions to divergence form elliptic equations with discontinuous coefficients. *Archive for rational mechanics and analysis*, 153(2):91–151, 2000.
- [LW07] Xiaosheng Li and Jenn-Nan Wang. Determination of viscosity in the stationary Navier-Stokes equations. *J. Differential Equations*, 242(1):24–39, 2007.
- [MU20] Claudio Munoz and Gunther Uhlmann. The Calderón problem for quasilinear elliptic equations. In *Annales de l’Institut Henri Poincaré C, Analyse non linéaire*, volume 37, pages 1143–1166. Elsevier, 2020.
- [MVVT16] Antonio Maffucci, Antonio Vento, Salvatore Ventre, and Antonello Tamburrino. A novel technique for evaluating the effective permittivity of inhomogeneous interconnects based on the monotonicity property. *IEEE Transactions on Components, Packaging and Manufacturing Technology*, 6(9):1417–1427, 2016.
- [RR06] Michael Renardy and Robert C Rogers. *An introduction to partial differential equations*, volume 13. Springer Science & Business Media, 2006.

- [SKJ⁺19] Jin Keun Seo, Kang Cheol Kim, Ariungerel Jargal, Kyoungun Lee, and Bastian Harrach. A learning-based method for solving ill-posed nonlinear inverse problems: a simulation study of lung eit. *SIAM journal on Imaging Sciences*, 12(3):1275–1295, 2019.
- [SU97] Ziqi Sun and Gunther Uhlmann. Inverse problems in quasilinear anisotropic media. *American Journal of Mathematics*, 119(4):771–797, 1997.
- [SUG⁺17] Zhiyi Su, Lalita Udpa, Gaspare Giovinco, Salvatore Ventre, and Antonello Tamburrino. Monotonicity principle in pulsed eddy current testing and its application to defect sizing. In *Applied Computational Electromagnetics Society Symposium-Italy (ACES), 2017 International*, pages 1–2. IEEE, 2017.
- [Sun96] Ziqi Sun. On a quasilinear inverse boundary value problem. *Math. Z.*, 221(2):293–305, 1996.
- [Sun05] Ziqi Sun. Conjectures in inverse boundary value problems for quasilinear elliptic equations. *Cubo*, 7(3):65–73, 2005.
- [Sun10] Ziqi Sun. An inverse boundary-value problem for semilinear elliptic equations. *Electronic Journal of Differential Equations (EJDE)[electronic only]*, 37:1–5, 2010.
- [SZ12] Mikko Salo and Xiao Zhong. An inverse problem for the p-Laplacian: boundary determination. *SIAM journal on mathematical analysis*, 44(4):2474–2495, 2012.
- [TR02] Antonello Tamburrino and Guglielmo Rubinacci. A new non-iterative inversion method for electrical resistance tomography. *Inverse Problems*, 18(6):1809, 2002.
- [TSV⁺16] Antonello Tamburrino, Zhiyi Sua, Salvatore Ventre, Lalita Udpa, and Satish S Udpa. Monotonicity based imang method in time domain eddy current testing. *Electromagnetic Nondestructive Evaluation (XIX)*, 41:1, 2016.
- [Uhl09] Gunther Uhlmann. Electrical impedance tomography and Calderón’s problem. *Inverse Problems*, 25:123011, 2009.
- [VMC⁺17] Salvatore Ventre, Antonio Maffucci, François Caire, Nechtan Le Lostec, Antea Perrotta, Guglielmo Rubinacci, Bernard Sartre, Antonio Vento, and Antonello Tamburrino. Design of a real-time eddy current tomography system. *IEEE Transactions on Magnetism*, 53(3):1–8, 2017.
- [ZHS18] Liangdong Zhou, Bastian Harrach, and Jin Keun Seo. Monotonicity-based electrical impedance tomography for lung imaging. *Inverse Problems*, 34(4):045005, 2018.

INSTITUTE FOR MATHEMATICS, GOETHE UNIVERSITY FRANKFURT, GERMANY
Email address: harrach@math.uni-frankfurt.de

DEPARTMENT OF APPLIED MATHEMATICS, NATIONAL YANG MING CHIAO TUNG UNIVERSITY,
TAIWAN
Email address: yihuanlin3@gmail.com