

On partial uniqueness of complete non-compact Ricci flat metrics

Yuanqi Wang*

Abstract

Using techniques for Caccioppoli inequality, on a fairly general class of complete non-compact Kähler manifolds with sub-quadratic volume growth, we show uniqueness of bounded $C^{1,1}$ solution to Monge-Ampere equation. This does not a priori require any decay of the solution.

1 Introduction

In an arbitrary Kähler class on a closed Kähler manifold with vanishing c_1 , uniqueness of Ricci flat metric can be proved via integration by parts [2, 18]. On a non-compact complete Kähler manifold, we show that under sub-quadratic volume growth and other mild conditions, even without decay, there is still an integration by parts argument for uniqueness of bounded solution to Monge-Ampere equation. The technique follows Caccioppoli inequality on regularity for elliptic equations. For example, see [1, 7, 8, 14].

A sub-harmonic function on a compact manifold must be a constant. In contrast and in general, the same fails on Euclidean domains even if we prescribe constant boundary value. For example, the function $|X|^2 - 1$ is sub-harmonic and vanishes on the boundary of the unit ball centered at the origin. Via this viewpoint, a complete non-compact Riemannian manifold is in between: it is not compact, but it is complete as a metric space and has no boundary. Our result shows, under the volume growth condition, a sub-solution to Monge-Ampere equation with constant density 1 is still a constant.

Theorem 1.1. *Let (M, ω) be a complete non-compact Kähler manifold of complex dimension $n \geq 2$ and sub-quadratic volume growth. Let f be a continuous function bounded from above on M .*

1. *Suppose ω has strictly sub-quadratic volume growth. Then any bounded $C^{1,1}$ sub-solution, super-solution, or solution ϕ to*

$$(\omega + i\partial\bar{\partial}\phi)^n = \omega^n \tag{1}$$

is a constant.

*University of Kansas, Lawrence, KS, USA. yqwang@ku.edu.

2. Suppose $e^f - 1$ is integrable i.e. $\int_M |e^f - 1| \omega^n < +\infty$. Then any bounded $C^{1,1}$ solution ϕ to

$$(\omega + i\partial\bar{\partial}\phi)^n = e^f \omega^n \quad (2)$$

has bounded Dirichlet energy i.e. $\int_M |\nabla_\omega \phi|^2 \omega^n \leq c_{DE} < +\infty$, where

$$c_{DE} = 1000n^{4n+4} |\phi|_{C^0(M)} [(1+K)^{2n} c_{vol} + |e^f - 1|_{L^1(M, \omega^n)}]. \quad (3)$$

3. Suppose additionally that ω satisfies weak Neumann Poincare inequality on annulus. Then any bounded $C^{1,1}$ solution ϕ to (1) is a constant.

The constant (3) might not be optimal, but is effective. The terms involved are defined below.

Definition 1.2. A bounded $C^{1,1}$ solution, sub-solution, or super-solution to (2) is a real-valued function ϕ with the following properties.

- ϕ is twice continuously differentiable under the holomorphic (smooth) manifold structure (not necessarily with norm bound).
- ϕ satisfies (2), $(\omega + i\partial\bar{\partial}\phi)^n \geq e^f \omega^n$, or $(\omega + i\partial\bar{\partial}\phi)^n \leq e^f \omega^n$ respectively.
- There is a positive number $K (< \infty)$ such that

$$|\phi|_{C^0(M)} = \sup_M |\phi| < \infty \text{ and } 0 < \omega_\phi \leq K\omega \text{ on the whole } M. \quad (4)$$

Fix a point $o \in M$ as distance origin and center of balls. We say a complete non-compact Kähler metric ω has *sub-quadratic volume growth*, if there is a positive sequence $\rho_i \rightarrow +\infty$ such that

$$\limsup_{i \rightarrow \infty} \frac{\text{Vol}[B(2\rho_i) \setminus B(\rho_i)]}{\rho_i^2} < +\infty. \quad (5)$$

The value of the existing limit superior is denoted by c_{vol} . We say such an ω has *strictly sub-quadratic volume growth* if $c_{vol} = 0$ i.e.

$$\lim_{i \rightarrow \infty} \frac{\text{Vol}[B(2\rho_i) \setminus B(\rho_i)]}{\rho_i^2} = 0. \quad (6)$$

On the other hand, we say it satisfies *weak Neumann Poincare inequality on annulus* if there is $\underline{\rho}_i \rightarrow \infty$ and μ_i , such that $\underline{\rho}_i \leq \rho_i$ when i is large, and

$$\int_{B(2\rho_i) \setminus B(\rho_i)} |\phi - \mu_i|^2 \leq c_P \rho_i^2 \int_{\{r \geq \underline{\rho}_i\}} |\nabla \phi|^2, \quad (7)$$

where c_P is independent of i or ϕ , as long as ϕ is twice continuously differentiable. Our argument (19) below is independent of μ_i .

Suppose we have two solutions ϕ_1 and ϕ_2 to the general volume form equation (2). For uniqueness, as long as the conditions hold, we can apply Theorem 1.1.1 or 1.1.3 with reference metric being $\omega + i\partial\bar{\partial}\phi_1$ or $\omega + i\partial\bar{\partial}\phi_2$, and ϕ being $\pm(\phi_2 - \phi_1)$ respectively.

Theorem 1.1 partially addresses the uniqueness of Tian-Yau solutions [17, Theorem 1.1]. This particular result is under sub-quadratic volume growth, and their solution is bounded $C^{1,1}$. Let “unique...(up to constant)” abbreviates “unique...up to addition by a real constant”.

Corollary 1.3. *Let $2 > \alpha \geq 1$. If the $(K, 2, \beta)$ -polynomial growth condition is strengthened to (K, α, β) , the Tian-Yau solution φ in [17, Theorem 1.1] is the unique bounded $C^{1,1}$ solution (up to constant) to the Monge-Ampere equation [17, (1.1)]. The solution u in Hein’s version [10, Prop 4.1], under $SOB(\beta)$ -condition, $\beta \leq 2$, is the unique bounded $C^{1,1}$ solution (up to constant) to the Monge-Ampere equation therein. Consequently, the following holds.*

- *The Tian-Yau Ricci-flat space [17, Theorem 4.1] of volume growth $O(r^{\frac{2n}{n+1}})$ is the unique solution (up to constant) to the Monge-Ampere equation [17, (1.1)] with reference metric ω_N [17, (4.4)].*
- *All the gravitational instantons in Hein’s construction [11, Theorem 1.5] are unique solutions (up to constant) to corresponding Monge-Ampere equations in [10, Prop 4.1].*

Our sub-quadratic volume growth is a sequential condition, and is apparently implied by that ball of radius R (centered at the base point) has volume $\leq CR^2$, for large R cf. [17, Definition 1.1]. Suppose the weak Poincare inequality (7) on annulus is implied by $(K, 2, \beta)$ -polynomial growth and other conditions in [17, Theorem 1.1]. Then uniqueness of Tian-Yau solution [17, Theorem 1.1] holds in full generality. Nevertheless, for Hein’s version [10, Prop 4.1], we do have Poincare inequality [10, Prop 3.4]. Therefore uniqueness also holds for rigorous quadratic volume growth i.e. $SOB(2)$ -case.

Under faster than quadratic volume growth, uniqueness of Monge-Ampere solutions is implied by certain decay on the Kähler potential ϕ . See [13, 8.5 Theorem A4] for example. Geometric uniqueness of Ricci flat metrics, as in [5, 6, 9, 10, 11, 13, 16], usually involves both Monge-Ampere uniqueness and $i\partial\bar{\partial}$ -lemma under decay conditions. For recent work on Liouville theorem of Monge-Ampere equations on product manifolds, see Hein [12] for example. For recent work on Liouville theorem of metric Laplacians on certain non-compact complete manifolds, see Sun-Zhang [15] and Carron [3] for examples. For earlier work on rigidity of harmonic functions on non-compact complete Riemannian manifolds, see Cheng-Yau [4] for example.

2 Proof

Convention: Unless otherwise specified, the metric for gradient is ω . The integrals and volumes are with respect to the top degree form ω^n .

For Theorem 1.1.1

Write ω_ϕ for $\omega + i\partial\bar{\partial}\phi$. We have the difference

$$\omega_\phi^n - \omega^n = i\partial\bar{\partial}\phi \wedge Q. \quad (8)$$

where $Q = \omega_\phi^{n-1} + \omega_\phi^{n-2} \wedge \omega + \dots + \omega^{n-1}$. By positivity and $C^{1,1}$ -condition (4), we verify

$$\omega^{n-1} \leq Q \leq n^{2n}(1+K)^n \omega^{n-1}.$$

Actually, any constant depending only on the data in Theorem 1.1 and Definition 1.2 suffices, but we want explicit constant, though not necessarily optimal. Moreover,

$$\begin{cases} i\partial\bar{\partial}\phi \wedge Q \geq 0 \text{ for subsolution, and} \\ i\partial\bar{\partial}\phi \wedge Q \leq 0 \text{ for super-solution.} \end{cases} \quad (9)$$

Because ϕ is bounded,

- if it is a sub-solution, by adding a constant if necessary, we assume $\min \phi \geq 1$;
- if it is a super-solution, by adding a constant if necessary, we assume $\max \phi \leq -1$.

Let χ be compactly supported Lipschitz function. We multiply both hand sides in (9) by $\chi^2\phi$. In either case, because of the definite sign of $\chi^2\phi$, we find

$$\chi^2\phi \cdot i\partial\bar{\partial}\phi \wedge Q \geq 0. \quad (10)$$

We integrate (10) by parts:

$$\int_M \chi^2 i\partial\phi \wedge \bar{\partial}\phi \wedge Q \leq -2 \int_M \phi \chi \cdot i\partial\chi \wedge \bar{\partial}\phi \wedge Q. \quad (11)$$

The left side of (11) is bounded from below by

$$\int_M \chi^2 i\partial\phi \wedge \bar{\partial}\phi \wedge Q \geq \int_M \chi^2 i\partial\phi \wedge \bar{\partial}\phi \wedge \omega^{n-1} \geq \frac{1}{2n} \int_M \chi^2 |\nabla\phi|^2.$$

On the other hand, Cauchy-Schwartz yields an upper bound on the right side of (11):

$$\begin{aligned} | -2 \int_M \phi \chi i\partial\chi \wedge \bar{\partial}\phi \wedge Q | &\leq 2n^{2n}(1+K)^n \int_M |\phi| |\chi| |\nabla\chi| |\nabla\phi| \\ &\leq 2n^{2n}(1+K)^n \left(\int_M \chi^2 |\nabla\phi|^2 \right)^{\frac{1}{2}} \left(\int_M \phi^2 |\nabla\chi|^2 \right)^{\frac{1}{2}}. \end{aligned}$$

The above two inequalities and (11) imply

$$\int_M \chi^2 |\nabla \phi|^2 \leq 4n^{2n+1} (1+K)^n \left(\int_M \chi^2 |\nabla \phi|^2 \right)^{\frac{1}{2}} \left(\int_M \phi^2 |\nabla \chi|^2 \right)^{\frac{1}{2}}. \quad (12)$$

Either $\int_M \chi^2 |\nabla \phi|^2$ is 0 or not, we find

$$\int_M \chi^2 |\nabla \phi|^2 \leq 16n^{4n+2} (1+K)^{2n} \int_M \phi^2 |\nabla \chi|^2. \quad (13)$$

Now let χ be the following piece-wise linear function in the distance r from the base point o , and $\rho > 0$.

$$\chi(r) = \begin{cases} 1 & \text{when } r \leq \rho, \\ 2 - \frac{r}{\rho} & \text{when } \rho \leq r \leq 2\rho, \\ 0 & \text{when } r \geq 2\rho. \end{cases} \quad (14)$$

Apparently, χ is Lipschitz. We find

$$\begin{aligned} \int_{B(\rho)} |\nabla \phi|^2 &\leq \frac{16n^{4n+2} (1+K)^{2n}}{\rho^2} \int_{B(2\rho) \setminus B(\rho)} |\phi|^2 \\ &\leq 16n^{4n+2} (1+K)^{2n} |\phi|_{C^0(M)} \cdot \frac{\text{Vol}[B(2\rho) \setminus B(\rho)]}{\rho^2}. \end{aligned} \quad (15)$$

Let $\rho = \rho_i$ and $i \rightarrow \infty$. Monotone convergence theorem implies $|\nabla \phi|^2$ is integrable on M and

$$\int_M |\nabla \phi|^2 \leq 16n^{4n+2} (1+K)^{2n} |\phi|_{C^0(M)} \cdot \limsup_{i \rightarrow \infty} \frac{\text{Vol}[B(2\rho_i) \setminus B(\rho_i)]}{\rho_i^2} = 0. \quad (16)$$

This means ϕ is a constant.

For Theorem 1.1.2

In this case we do not add any constant to ϕ . It is a solution by assumption. The equality still holds if we multiply $i\partial\bar{\partial}\phi \wedge Q = (e^f - 1)\omega^n$ by $\chi^2\phi$. The same argument (8)—(12) with $e^f\omega^n$ instead of ω^n yields

$$\begin{aligned} &\int_M \chi^2 |\nabla \phi|^2 \\ &\leq 4n^{2n+1} (1+K)^n \left(\int_M \chi^2 |\nabla \phi|^2 \right)^{\frac{1}{2}} \left(\int_M \phi^2 |\nabla \chi|^2 \right)^{\frac{1}{2}} + \int_M \chi^2 |\phi(e^f - 1)| \\ &\leq 4n^{2n+1} (1+K)^n \left(\int_M \chi^2 |\nabla \phi|^2 \right)^{\frac{1}{2}} \left(\int_M \phi^2 |\nabla \chi|^2 \right)^{\frac{1}{2}} + |\phi|_{C^0(M)} \cdot |e^f - 1|_{L^1(M)}. \end{aligned}$$

The following elementary claim is proved by completing square.

Claim 2.1. *Let a, b, c, d be non-negative numbers such that*

$$a \leq 2d\sqrt{a} \cdot \sqrt{b} + c.$$

Then $a \leq 2d^2b + c + 2\sqrt{d^2b(d^2b + c)} \leq 4d^2b + 3c$.

We then find

$$\int_M \chi^2 |\nabla \phi|^2 \leq 64n^{4n+2}(1+K)^{2n} \int_M \phi^2 |\nabla \chi|^2 + 3|\phi|_{C^0(M)} \cdot |e^f - 1|_{L^1(M)} \quad (17)$$

Still let $\rho = \rho_i$ and $i \rightarrow \infty$. Same argument as (15) and (16) yields

$$\int_M |\nabla \phi|^2 \leq 64n^{4n+2}(1+K)^{2n} c_{vol} |\phi|_{C^0(M)} + 3|\phi|_{C^0(M)} \cdot |e^f - 1|_{L^1(M)}. \quad (18)$$

We enlarge the constant to (3).

For Theorem 1.1.3

By monotone convergence theorem, the established Dirichlet energy bound implies

$$\lim_{\rho \rightarrow \infty} \int_{\{r \geq \rho\}} |\nabla \phi|^2 = 0.$$

We apply the argument last section to $\phi - \mu_i$, which is also a solution. Multiplying $i\partial\bar{\partial}\phi \wedge Q = 0$ by $\chi^2(\phi - \mu_i)$, we still have equality. The assumed Poincaré inequality (7) and derivation of (15) shows

$$\begin{aligned} \int_{B(\rho_i)} |\nabla \phi|^2 &= \int_{B(\rho_i)} |\nabla(\phi - \mu_i)|^2 \leq \frac{16n^{4n+2}(1+K)^{2n}}{\rho_i^2} \int_{B(2\rho_i) \setminus B(\rho_i)} |\phi - \mu_i|^2 \\ &\leq 16n^{4n+2}(1+K)^{2n} c_P \int_{\{r \geq \rho_i\}} |\nabla \phi|^2 \text{ which } \rightarrow 0 \text{ as } i \rightarrow \infty. \end{aligned} \quad (19)$$

This again says $\int_M |\nabla \phi|^2 = 0$ and ϕ is a constant.

For Corollary 1.3

It is obvious from the general uniqueness 1.1 and the volume growth condition. Because the density data f in [17, Theorem 1.1] and [10, Proposition 3.4] are bounded on the whole M i.e. there exists a positive (finite) number c_0 such that $|f| \leq c_0$, any bounded $C^{1,1}$ -solution v yields $\omega_0 + i\partial\bar{\partial}v$ quasi-isometric to the reference metric ω_0 i.e. there is a $c_1 > 0$ possibly depending on v and f such that

$$\frac{\omega_0}{c_1} \leq \omega_0 + i\partial\bar{\partial}v \leq c_1\omega_0.$$

In conjunction with the remark above Corollary 1.3, if there are two solutions ϕ_1 and ϕ_2 , use ω_{ϕ_1} as the new reference metric and denote it by ω , and denote $\phi_2 - \phi_1$ by ϕ . The volume form of ω and ω_ϕ coincide. We apply Theorem 1.1.

When $\alpha < 2$ in the (K, α, β) -condition [17, Definition 1.1], or $\beta < 2$ in Hein's $SOB(\beta)$ -condition, by quasi-isometry, ω has strict sub-quadratic volume growth. Then Theorem 1.1.1 yields the result.

We elaborate more for $SOB(2)$. We do not know whether the solution ω is $SOB(2)$, though ω_0 is by assumption. Nevertheless, *the interior Neumann Poincare inequality* [10, Proposition 3.4] for $SOB(2)$ -reference metric ω_0 and the quasi-isometry

$$\frac{\omega}{c_2} \leq \omega_\phi \leq c_2 \omega$$

still implies the weak Neumann Poincare inequality (7) for ω . Namely, fix a single base point o for both ω_0 and ω . For large enough c_3 independent of i such that the ball $B(4\rho_i)$ with respect to ω is contained in the ball $B(c_3\rho_i)$ with respect to ω_0 , and the ball $B(\frac{\rho_i}{2})$ with respect to ω contains the ball $B(\frac{\rho_i}{c_3})$ with respect to ω_0 , we assign the data

$$r_1 = \frac{\rho_i}{c_3}, \quad s = \frac{\rho_i}{1000c_3}, \quad r_2 = c_3\rho_i, \quad \kappa = 0$$

on radius and other to the ball and annuli in [10, Proposition 3.4] for the $SOB(2)$ -reference metric ω_0 . When i is large, [10, (3.4)] implies

$$\begin{aligned} & \int_{B(2\rho_i) \setminus B(\rho_i)} |h - h_{A(\frac{\rho_i}{c_3}, c_3\rho_i)}|^2 \omega^n \\ & \leq c'_3 \int_{B(c_3\rho_i) \setminus B(\frac{\rho_i}{c_3})} |h - h_{A(\frac{\rho_i}{c_3}, c_3\rho_i)}|^2 \omega_0^n \leq c_4 \rho_i^2 \int_{\{r \geq \frac{\rho_i}{10c_3}\}} |\nabla_{\omega_0} f|^2 \omega_0^n \\ & \leq c_5 \rho_i^2 \int_{\{r \geq \frac{\rho_i}{c_5}\}} |\nabla f|^2 \omega^n, \end{aligned} \tag{20}$$

where the large enough positive constants c'_3 , c_4 , c_5 are independent of ρ_i or the arbitrary twice differentiable function h , and $h_{A(\frac{\rho_i}{c_3}, c_3\rho_i)}$ is the ω_0^n -average of h on the closed annulus therein. Note $h_{A(\frac{\rho_i}{c_3}, c_3\rho_i)}$ is also applied in the first line of (20) in the integration against ω^n . Related to c_4 , the volume ratio N in [10, (3.4)] is bounded (from above) by the volume constants in $SOB(2)$ -condition [10, Definition 3.1] (including the Ricci lower bound). This fulfills requirement (7) because $\frac{\rho_i}{c_5}$ still approaches ∞ as $i \rightarrow \infty$. Then apply Theorem 1.1.3.

References

- [1] A. Björn, N. Marola. *Moser iteration for (quasi) minimizers on metric spaces*. Manuscripta Math. 121 (2006), no. 3, 339–366.
- [2] E. Calabi. *On Kähler manifolds with vanishing canonical class*, *Algebraic Geometry and Topology*. Algebraic geometry and topology. A symposium in honor of S. Lefschetz, pp. 78–89. Princeton University Press, Princeton, N.J., 1957.

- [3] G. Carron. *Harmonic functions on manifolds whose large spheres are small*. Ann. Math. Blaise Pascal 23 (2016), no. 2, 249-261.
- [4] S-Y. Cheng, S-T. Yau, *Differential equations on Riemannian manifolds and their geometric applications*. Communications on Pure and Applied Mathematics 28 (1975), no. 3, 333-354.
- [5] R. Conlon, H.J. Hein. *Asymptotically conical Calabi-Yau manifolds I*. Duke Math. J. 162 (2013), no. 15, 2855-2902.
- [6] R. Conlon, H.J. Hein. *Asymptotically conical Calabi-Yau metrics on quasi-projective varieties*. Geom. Funct. Anal. 25 (2015), no. 2, 517-552.
- [7] E. DiBenedetto, N.S. Trudinger. *Harnack inequalities for quasiminima of variational integrals*. Ann. Inst. H. Poincaré Anal. Non Linéaire 1 (1984), no. 4, 295-308.
- [8] M. Giaquinta, L. Martinazzi . (2012) *L^2 -regularity: The Caccioppoli inequality*. In: *An Introduction to the Regularity Theory for Elliptic Systems, Harmonic Maps and Minimal Graphs*. Publications of the Scuola Normale Superiore. Edizioni della Normale, Pisa.
- [9] M. Haskins, H.-J. Hein, J. Nordström. *Asymptotically cylindrical Calabi-Yau manifolds*. J. Differential Geom. 101 (2015), no. 2, 213-265.
- [10] H.-J. Hein. *On gravitational instantons*. Thesis (Ph.D.)-Princeton University. 2010. 129 pp. ISBN: 978-1124-34891-9, ProQuest LLC
- [11] H.-J. Hein. *Gravitational instantons from rational elliptic surfaces*. J. Amer. Math. Soc. 25 (2012), no. 2, 355-393.
- [12] H.-J. Hein. *A Liouville theorem for the complex Monge-Ampère equation on product manifolds*. Comm. Pure Appl. Math. 72 (2019), 122-135.
- [13] D. Joyce. *Compact manifolds with special holonomy*. Oxford Mathematical Monographs.
- [14] J. Kinnunen, N. Shanmugalingam. *Regularity of quasi-minimizers on metric spaces*. Manuscripta Math. 105 (2001), no. 3, 401-423.
- [15] S. Sun, R.B. Zhang. *A Liouville theorem on asymptotically Calabi spaces*. Calculus of Variations and Partial Differential Equations 60 (3), 1-43.
- [16] Gábor Székelyhidi. *Uniqueness of some Calabi-Yau metrics on $\mathbb{C}^n$* . Geom. Funct. Anal. 30 (2020), no. 4, 1152-1182.
- [17] G. Tian, S.T. Yau. *Complete Kähler manifolds with zero Ricci curvature*. I. J. Amer. Math. Soc. 3 (1990), no. 3, 579-609.
- [18] S.T. Yau. *On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation*. I. Comm. Pure Appl. Math. 31 (1978), no. 3, 339-411.