

GEOMETRIC DESCRIPTION OF $\langle 2 \rangle$ -POLARISED HILBERT SQUARES OF GENERIC K3 SURFACES

SIMONE NOVARIO

ABSTRACT. A generic K3 surface of degree $2t$ is a general complex projective K3 surface S_{2t} whose Picard group is generated by the class of an ample divisor $H \in \text{Div}(S_{2t})$ such that $H^2 = 2t$ with respect to the intersection form. We show that if X is the Hilbert square of a generic K3 surface of degree $2t$ with $t \neq 2$ which admits an ample divisor $D \in \text{Div}(X)$ with $q_X(D) = 2$, where q_X is the Beauville–Bogomolov–Fujiki form, then X is a double EPW sextic.

1. INTRODUCTION

A classical result in the theory of complex projective K3 surfaces says that if a K3 surface admits an ample divisor D with $D^2 = 2$ with respect to the intersection form, then the complete linear system $|D|$ is basepoint free and the morphism that it induces is a double cover of the plane \mathbb{P}^2 ramified on a sextic curve. This was proved by Saint–Donat in [SD74].

It is natural to study a similar problem for a projective IHS (Irreducible Holomorphic Symplectic) manifold of dimension $2n$ with $n \geq 2$, a sort of higher dimensional generalization of K3 surfaces. The importance of these varieties is linked to the Beauville–Bogomolov decomposition theorem, see [Bea83b]: up to a finite étale cover, any compact Kähler manifold with trivial first Chern class is the product of a complex torus, irreducible Calabi–Yau manifolds and IHS manifolds.

Let S be a K3 surface. We denote by $T(S)$ its transcendental lattice, which coincides with $\text{NS}(S)^\perp$, where $\text{NS}(S)$ is the Néron–Severi group of S and the orthogonal is taken with respect to the intersection form. Then $T(S)_\mathbb{Q} := T(S) \otimes \mathbb{Q}$ is a rational Hodge structure of weight 2 and we denote by

$$E_S := \text{Hom}_0(T(S)_\mathbb{Q}, T(S)_\mathbb{Q})$$

the algebra of endomorphisms of weight 0 on $T(S)_\mathbb{Q}$. See [Huy16, §3] for details on Hodge structures. By a result of Zahrin, see [Zar83], E_S is either \mathbb{Q} , or a totally real field or a CM field. We say that S is a K3 surface *general in its rank*, or simply *general*, if $E_S \cong \mathbb{Q}$. If E_S is a totally real field then S has *real multiplication* (RM), while if E_S is a CM field then S has *complex multiplication* (CM). Then K3 surfaces with RM or CM describe a locus of positive codimension in the analytic moduli space of K3 surfaces of fixed Picard rank, except for K3 surfaces of Picard rank 20, which have all CM. See [vG08] and [EJ16] for details.

A *generic K3 surface* is a general projective K3 surface whose Picard group has rank 1. We say that a generic K3 surface S_{2t} has *degree* $2t$ if $\text{Pic}(S_{2t}) = \mathbb{Z}H$ with $H^2 = 2t$. In this paper we focus on Hilbert schemes of 2 points on a generic K3 surface, also known as *Hilbert squares* of generic K3 surfaces: the Hilbert square of a K3 surface S will be denoted by $S^{[2]}$. The main problem of this paper is to determine the geometric description of a $\langle 2 \rangle$ -polarised Hilbert square X of a generic K3 surface, i.e., X admits an ample divisor $D \in \text{Div}(X)$ with $q_X(D) = 2$, where q_X is the *Beauville–Bogomolov–Fujiki* quadratic form on $H^2(X, \mathbb{Z})$. This can be seen as a generalization of the problem studied by Saint–Donat presented in the first lines. We denote by $|D|$ the complete linear system associated to $D \in \text{Div}(X)$, i.e., the set of effective divisors linearly equivalent to D , and by $\text{Bs}|D|$ the base locus of $|D|$, which is the set of points on X where all the global sections in $H^0(X, \mathcal{O}_X(D))$ vanish. Our aim is to study the rational map $\varphi_{|D|}$ induced by the complete linear system $|D|$, i.e.,

$$\varphi_{|D|} : X \dashrightarrow \mathbb{P}(H^0(X, \mathcal{O}_X(D))^\vee),$$

where by definition $x \in X \setminus \text{Bs}|D|$ is mapped to the hyperplane in $H^0(X, \mathcal{O}_X(D))$ consisting of sections vanishing at x . If $\text{Bs}|D|$ is empty, we say that $|D|$ is *basepoint free*: in this case $\varphi_{|D|}$ is a morphism.

Let X and D be as above. Then there exists an anti-symplectic involution ι which generates the group $\text{Aut}(X)$ of biregular automorphisms on X by a result obtained by Boissière, Cattaneo, Nieper-Wißkirchen and Sarti in [BCNWS16], and ι is such that $\iota^*[D] = [D]$ in the Néron–Severi group $\text{NS}(X)$. Here anti-symplectic means that $\iota^*\sigma_X = -\sigma_X$, where $\sigma_X \in H^0(X, \Omega_X^2)$ is a symplectic form of X . The main theorem of this paper, which we now

present, gives a geometrical description of the map induced by the complete linear system $|D|$, cf. Theorem 6.4 and Theorem 8.5.

Theorem. *Let X be the Hilbert square of a generic K3 surface S_{2t} of degree $2t$ such that X admits an ample divisor $D \in \text{Div}(X)$ with $q_X(D) = 2$. Suppose that $t \neq 2$, and denote by ι the anti-symplectic involution which generates the group $\text{Aut}(X)$. Then the complete linear system $|D|$ is basepoint free, the morphism*

$$\varphi_{|D|} : X \rightarrow Y \subset \mathbb{P}^5$$

is a double cover whose ramification locus is the surface F of points fixed by ι , and $Y \cong X/\langle \iota \rangle$ is an EPW sextic, in particular X is a double EPW sextic. Moreover, if $H^{2,2}(X, \mathbb{Z}) := H^4(X, \mathbb{Z}) \cap H^{2,2}(X)$, then

$$[F] = 5D^2 - \frac{1}{3}c_2(X) \in H^{2,2}(X, \mathbb{Z}),$$

where $c_2(X) \in H^4(X, \mathbb{Z})$ is the second Chern class of X .

Here a double EPW sextic is a double cover of an EPW sextic ramified in its singular locus: see [EPW01] for the definition of EPW sextic and [O'G06] for details on double EPW sextics. The paper [O'G08], where O'Grady gives a classification, up to deformation equivalence, of *numerical* $K3^{[2]}$, will play an important role. A numerical $K3^{[2]}$ is by definition an IHS manifold M which admits an isomorphism of abelian groups $\psi : H^2(M, \mathbb{Z}) \rightarrow H^2(S^{[2]}, \mathbb{Z})$ for some K3 surface S such that $\int_M \alpha^4 = \int_{S^{[2]}} \psi(\alpha)^4$ for every $\alpha \in H^2(M, \mathbb{Z})$. In particular he showed that a numerical $K3^{[2]}$ is deformation equivalent either to a double EPW sextic or to an IHS manifold Z admitting a rational map $f : Z \dashrightarrow \mathbb{P}^5$ which is birational onto its image Y , with $6 \leq \deg(Y) \leq 12$. The link between our problem and the one studied in [O'G08] is given by the fact that O'Grady proved the following: a numerical $K3^{[2]}$ is deformation equivalent to an IHS manifold Z of $K3^{[2]}$ -type such that $\text{Pic}(Z)$ is generated by the class of an ample divisor $H \in \text{Pic}(Z)$ with $q_Z(H) = 2$. Let X be the Hilbert square of a general K3 surface of degree $2t$ and suppose that there is an ample divisor $D \in \text{Div}(X)$ with $q_X(D) = 2$. In this specific case the result obtained by O'Grady implies that X is deformation equivalent to a double EPW sextic, while our main theorem is stronger, since it shows that $X = S_{2t}^{[2]}$, if $t \neq 2$, is exactly a double EPW sextic, giving a more precise geometric description of these varieties. The case $t = 2$ was already studied by Welters and Beauville in [Wel81] and [Bea83a]: the map $\varphi_{|D|}$ is a finite morphism of degree 6 with image $\mathbb{G}(1, \mathbb{P}^3)$, the Grassmannian of lines in \mathbb{P}^3 .

Our strategy is to follow [O'G08], using the anti-symplectic involution which generates $\text{Aut}(X)$ given by [BCNWS16] in order to get as much information as possible on the geometry of the complete linear system.

The paper is organised as follows. In Section 2 we give basics on Pell equations and Pell-type equations. In Section 3 we recall the definition of IHS manifold, together with main properties of this family of varieties. In Section 4 we present some useful results on IHS manifolds of $K3^{[2]}$ -type, in particular we recall the description of the lattice $H^{2,2}(S^{[2]}, \mathbb{Z}) := H^4(S^{[2]}, \mathbb{Z}) \cap H^{2,2}(S^{[2]})$ of integral Hodge classes of type $(2, 2)$ on $S^{[2]}$ obtained in [Nov21] for a general projective K3 surface S : this will be extremely useful in several steps of this paper. In Section 5 we recall the most important results on the Hilbert square X of a generic K3 surface, in particular the description of the group of biregular automorphisms obtained in [BCNWS16]: $\text{Aut}(X)$ is not trivial and generated by a non-natural anti-symplectic involution ι when X is the Hilbert square of a generic K3 surface of degree $2t$ admitting an ample divisor $D \in \text{Div}(X)$ with $q_X(D) = 2$. We also briefly describe the case $t = 2$, already studied by Welters and Beauville in [Wel81] and [Bea83a]. From now on X is the Hilbert square of a generic K3 surface admitting an ample divisor D with $q_X(D) = 2$. In Section 6 we study the surface F of points on X fixed by the anti-symplectic involution ι : we compute the class of F in $H^{2,2}(X, \mathbb{Z})$ and we show that the rational map $\varphi_{|D|}$ factors through the quotient with respect to the involution ι . In Section 7 we prove the *irreducibility property*: if $D_1, D_2 \in |D|$ are distinct and $t \neq 2$, where $2t$ is the degree of the underlying generic K3 surface, then $D_1 \cap D_2$ is a reduced and irreducible surface. This is a fundamental step in the geometrical analysis of the rational map $\varphi_{|D|}$. When $t = 2$, the surface $D_1 \cap D_2$ can be reducible: we give a geometrical interpretation of the two irreducible components. We conclude with Section 8, where we show the main result of this paper: keeping notation as above, if $t \neq 2$ then X is a double EPW sextic and $\varphi_{|D|}$ is the double cover associated.

The topic of this paper is related to [DM19, §7]. The approach and the techniques that we use are different: main tools are results on integral Hodge classes of type $(2, 2)$ on the Hilbert square of a generic projective K3 surface obtained in [Nov21], which are exploited in particular in Theorem 6.4 and in Theorem 7.3, and the papers [O'G08] and [BCNWS16]. Moreover, the irreducibility property of Theorem 7.3 explains geometrically why the case $t \neq 2$ is special, compare with [DM19, Remark 7.7].

This paper is based on Chapter 4 and Chapter 5 of the author's PhD thesis, see [Nov].

Acknowledgements: I thank my advisors Samuel Boissière and Bert van Geemen for the constant support I received. I would like to thank Michela Artebani and Alice Garbagnati for useful discussions on Section 6. I also thank Pietro Beri and Ángel David Ríos Ortiz for many useful discussions.

2. PELL EQUATIONS AND PELL-TYPE EQUATIONS

We give a brief overview of Pell equations and Pell-type equations.

Definition 2.1. A *Pell-type equation* is a diophantine equation of the form

$$x^2 - dy^2 = n,$$

where $d \in \mathbb{Z}_{>0}$ is a positive integer, $n \in \mathbb{Z} \setminus \{0\}$ is a non-zero integer and x, y are variables. We denote an equation of this form by $P_d(n)$. We call $P_d(1)$ a *Pell equation* and $P_d(-1)$ a *negative Pell equation*.

We are interested in integral solutions of Pell equations and Pell-type equations. If $d = c^2$ is a square, the only solutions of the Pell equation $P_d(1)$ are $(x, y) = (\pm 1, 0)$, and the Pell-type equation $P_d(n)$ can be written as $(x + cy)(x - cy) = n$, so it can be easily solved. From now on, we will assume that d is not a square. In this case the Pell-type equation $P_d(n)$ can be written in $\mathbb{Z}[\sqrt{d}] := \mathbb{Z}[x] / (x^2 - d)$ as

$$(x + y\sqrt{d})(x - y\sqrt{d}) = n.$$

Let $z = x + y\sqrt{d} \in \mathbb{Z}[\sqrt{d}]$: we define its *conjugate* as $\bar{z} := x - y\sqrt{d} \in \mathbb{Z}[\sqrt{d}]$, and its *norm* as $N(z) := z\bar{z} = x^2 - dy^2 \in \mathbb{Z}$. With this notation, $P_d(1)$ and $P_d(n)$ can be written respectively as $N(z) = 1$ and $N(z) = n$, where $z = x + y\sqrt{d}$.

Definition 2.2. Given a Pell equation $P_d(1)$ or a Pell-type equation $P_d(n)$, two solutions (X, Y) and (X', Y') are said to be *equivalent* if

$$\frac{XX' - dYY'}{n} \in \mathbb{Z}, \quad \frac{XY' - X'Y}{n} \in \mathbb{Z}.$$

Note that two solutions (X, Y) and (X', Y') of a Pell-type equation $P_d(n)$ are equivalent if $\frac{z_1 \bar{z}_2}{n} \in \mathbb{Z}[\sqrt{d}]$, where $z_1 = X + Y\sqrt{d}$ and $z_2 = X' + Y'\sqrt{d}$.

Definition 2.3. Consider a Pell equation or a Pell-type equation. The *fundamental* solution (X, Y) in an equivalence class of solutions is the one with smallest non-negative Y if such a solution is unique in its class. Otherwise, there are two solutions (X, Y) , $(-X, Y)$, which are said to be *conjugated*, with smallest non-negative Y : the fundamental solution is the one of the form (X, Y) with $X > 0$.

One can show that a Pell equation $P_d(1)$ is always solvable and all its solutions are equivalent. If $z_0 = a + b\sqrt{d}$ is the fundamental solution, then all the other solutions are of the form

$$z = \pm z_0^m, \quad m \in \mathbb{Z}_{>0}.$$

Similarly, if z_0 is a fundamental solution of $P_d(n)$ and \tilde{z}_0 is the fundamental solution of $P_d(1)$, all the other solutions equivalent to z_0 are of the form

$$z = \pm \tilde{z}_0^m \cdot z_0, \quad m \in \mathbb{Z}_{>0}.$$

We now recall the definition of *positive* and *minimal* solution of a Pell-type equation.

Definition 2.4. Let $P_d(n)$ be a Pell-type equation. A solution (X, Y) is *positive* if $X, Y > 0$. The *minimal* solution of $P_d(n)$ is the positive solution with smallest X .

Remark 2.5. The minimal solution of the Pell equation $P_t(1)$ coincides with the square of the minimal solution of the negative Pell equation $P_t(-1)$, if this exists, i.e., if $a + b\sqrt{t} \in \mathbb{Z}[\sqrt{t}]$ is the minimal solution of $N(z) = -1$ and $c + d\sqrt{t} \in \mathbb{Z}[\sqrt{t}]$ is the minimal solution of $N(z) = 1$, then $c + d\sqrt{t} = (a + b\sqrt{t})^2 = a^2 + b^2 + 2ab\sqrt{t}$, hence

$$a^2 + tb^2 = c, \quad d = 2ab. \tag{2.1}$$

We conclude this section with the following elementary result, which will be useful in Section 6 and in Section 7.

Proposition 2.6. Let (a, b) be the minimal solution of the Pell-type equation $P_d(-1)$. Then b is odd.

Proof. Suppose that b is even. Then $a^2 + 1$ is divisible by 4, since $a^2 - db^2 = -1$, i.e., $a^2 + 1 = 4X$ for some $X \in \mathbb{Z}$.

- If a is even, i.e., $a = 2Y$ for some $Y \in \mathbb{Z}$, then $4Y^2 + 1 = 4X$, which is not possible.
- If a is odd, i.e., $a = 2Y + 1$ for some $Y \in \mathbb{Z}$, then $4Y^2 + 1 + 4Y + 1 = 4X$, which gives $4(X - Y^2 - Y) = 2$, which is not possible.

We conclude that b is odd. □

3. GENERALITIES ON IHS MANIFOLDS

In this section we recall basics on irreducible holomorphic symplectic manifolds.

Definition 3.1. An *irreducible holomorphic symplectic* (IHS) *manifold* is a simply connected compact complex Kähler manifold X such that $H^0(X, \Omega_X^2)$ is generated by a non-degenerate holomorphic 2-form, called *symplectic form*.

The definition of IHS manifold generalises to higher dimensions the one of K3 surface, the only example of IHS manifold of dimension 2, as shown by the Enriques–Kodaira classification of compact complex surfaces. The existence of a symplectic form implies that the dimension of an IHS manifold is necessarily even. If X is an IHS manifold, then the \mathbb{C} -vector space $H^0(X, \Omega_X^p)$ is zero if p is odd, and it is generated by $\sigma^{\frac{p}{2}}$ if $0 \leq p \leq \dim(X)$ is even, where σ is a symplectic form, see [Bea83b, Proposition 3]. The Picard group $\text{Pic}(X)$ and the Néron–Severi group, defined as $\text{NS}(X) := H^{1,1}(X)_{\mathbb{R}} \cap H^2(X, \mathbb{Z})$, are isomorphic: $\text{Pic}(X)$ then embeds in the second cohomology group $H^2(X, \mathbb{Z})$. We recall that a *lattice* is a free \mathbb{Z} -module L of finite rank with a symmetric bilinear form $b : L \times L \rightarrow \mathbb{Z}$: we denote by $q : L \rightarrow \mathbb{Z}$ the quadratic form $q(x) := b(x, x)$ for every $x \in L$. If $\mathcal{B} := \{e_1, \dots, e_n\}$ is a \mathbb{Z} -basis of L , the *Gram matrix* of L associated to \mathcal{B} is the $n \times n$ symmetric matrix

$$\begin{pmatrix} b(e_1, e_1) & \cdots & b(e_1, e_n) \\ \vdots & \ddots & \vdots \\ b(e_n, e_1) & \cdots & b(e_n, e_n) \end{pmatrix}.$$

We say that a lattice L of rank n is *non-degenerate* if for any non-zero $l \in L$ there exists $l' \in L$ such that $b(l, l') \neq 0$, equivalently, $\det(G) \neq 0$ if G is a Gram matrix of L . A lattice L is *even* if $b(l, l) \in 2\mathbb{Z}$ for every $l \in L$, and *odd* if it is not even. The *determinant* of a lattice L is the determinant of a Gram matrix G of the lattice, and the *discriminant* of L is $\text{disc}(L) := |\det(G)|$. A lattice L is *unimodular* if $\text{disc}(L) = 1$. If L is a unimodular lattice, for every $x \in L$ there exists $y \in L$ such that $b(x, y) = 1$. A *sublattice* of a lattice L is a free submodule $L' \subseteq L$ with symmetric bilinear form $b' := b|_{L' \times L'}$. A sublattice $L' \subseteq L$ is *primitive* if L/L' is a free module. We define the *direct sum* of two lattices L_1 and L_2 as the lattice $L_1 \oplus L_2$ whose bilinear form is $b(v_1 + v_2, w_1 + w_2) := b_1(v_1, w_1) + b_2(v_2, w_2)$ for every $v_1, w_1 \in L_1$ and $v_2, w_2 \in L_2$, where b_1 and b_2 are the bilinear forms of L_1 and L_2 respectively. If L and L' are two lattices with bilinear forms b and b' respectively, we call *morphism of lattices* $\varphi : L \rightarrow L'$ a morphism of \mathbb{Z} -modules such that for every $l_1, l_2 \in L$ we have $b(l_1, l_2) = b'(\varphi(l_1), \varphi(l_2))$. Note that morphisms between two non-degenerate lattices are injective. We say that a lattice *embeds primitively* in a lattice L' if there is a morphism $\varphi : L \rightarrow L'$ such that $\varphi(L)$ is a primitive sublattice of L' . An *isometry* is a bijective morphism of lattices. The *divisibility* of an element $l \in L$ in a lattice L is the positive generator of the ideal $\{b(l, m) \mid m \in L\} \subseteq \mathbb{Z}$.

For a lattice L of rank n we write $L_{\mathbb{R}} := L \otimes_{\mathbb{Z}} \mathbb{R}$ and we extend \mathbb{R} -bilinearly the bilinear form b to $L_{\mathbb{R}}$, similarly we extend q to $L_{\mathbb{R}}$. If the lattice is non-degenerate, the *signature* of L is the signature $(l_{(+)}, l_{(-)})$ of the quadratic form on $L_{\mathbb{R}}$. A non-degenerate lattice is *positive definite* if $l_{(-)} = 0$, similarly it is *negative definite* if $l_{(+)} = 0$, while it is *indefinite* if $l_{(+)}, l_{(-)} \neq 0$.

Example 3.2. If k is a non-zero integer, let $\langle k \rangle$ be the rank one lattice $L = \mathbb{Z}e$ with bilinear form $b(e, e) = k$.

Example 3.3. Let U be the *hyperbolic lattice*, i.e., the unique unimodular lattice of rank 2 and signature $(1, 1)$. Its Gram matrix is the following:

$$\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}.$$

Example 3.4. Let $E_8(-1)$ be the even unimodular lattice of signature $(0, 8)$ whose Gram matrix is the following:

$$\begin{pmatrix} -2 & 1 & & & & & & \\ 1 & -2 & 1 & & & & & \\ & 1 & -2 & 1 & & & & \\ & & 1 & -2 & 1 & & & \\ & & & 1 & -2 & 1 & & \\ & & & & 1 & -2 & 1 & \\ & & & & & 1 & -2 & 1 \\ & & & & & & 1 & -2 \end{pmatrix}.$$

Let X be an IHS manifold. By the universal coefficient theorem, the second singular cohomology group $H^2(X, \mathbb{Z})$ is torsion free, and this can be equipped with a non-degenerate integral quadratic form by the following result due to Beauville, Bogomolov and Fujiki, see [Bea83b] and [Fuj87].

Theorem 3.5 (Beauville–Bogomolov–Fujiki form). *Let X be an IHS manifold of dimension $2n$. Then there exists an integral quadratic form $q_X : H^2(X, \mathbb{Z}) \rightarrow \mathbb{Z}$ and a constant $c_X \in \mathbb{Q}_{>0}$ such that*

$$\int_X \alpha^{2n} = c_X \frac{(2n)!}{n!2^n} q_X(\alpha)^n \quad \text{for all } \alpha \in H^2(X, \mathbb{Z}).$$

The quadratic form q_X is called *Beauville–Bogomolov–Fujiki (BBF) form*, and c_X is called *Fujiki constant* of X . Moreover, $(H^2(X, \mathbb{Z}), q_X)$ is a lattice of signature $(3, b_2(X) - 3)$, where $b_2(X)$ is the second Betti number of X .

An example of IHS manifold of dimension $2n$, where $n \geq 2$, is given by the *Hilbert scheme of n points on a K3 surface S* , the scheme which parametrises zero-dimensional closed subschemes of length n on a K3 surface: we denote it by $S^{[n]}$. Let $S^{(n)}$ be the quotient of $S^n = S \times \cdots \times S$ by the symmetric group of n elements, so $S^{(n)}$ is the variety of 0-cycles of degree n . Then the *Hilbert–Chow morphism* $\rho : S^{[n]} \rightarrow S^{(n)}$ is defined as follows: a point $[\xi] \in S^{[n]}$ is mapped to the cycle $\sum_x l(\mathcal{O}_{\xi, x}) \cdot x$, see for instance [Ive06]. The singular locus of $S^{(n)}$ is the so-called diagonal, i.e., the set of cycles $p_1 + \cdots + p_n$ such that there exist distinct i and j with $p_i = p_j$. The Hilbert–Chow morphism is a desingularization of $S^{(n)}$, and the pre-image of the diagonal is an irreducible divisor E on $S^{[n]}$. The Hilbert scheme of n points on a K3 surface is an IHS manifold by [Bea83b, Théorème 3]. There exists a primitive class $\delta \in \text{Pic}(S^{[n]})$ such that $2\delta = [E]$. Moreover, there is a primitive embedding of lattices

$$i : H^2(S, \mathbb{Z}) \hookrightarrow H^2(S^{[n]}, \mathbb{Z})$$

such that $H^2(S^{[n]}, \mathbb{Z}) = i(H^2(S, \mathbb{Z})) \oplus \mathbb{Z}\delta$, and $q_{S^{[n]}}(\delta) = -2(n-1)$. Recall that $H^2(S, \mathbb{Z}) \cong U^{\oplus 3} \oplus E_8(-1)^{\oplus 2}$, in particular $H^2(S, \mathbb{Z})$ is an even unimodular lattice of signature $(3, 19)$, see for instance [BHPVdV15, §VII.3]. Then there is an isometry of lattices

$$H^2(S^{[n]}, \mathbb{Z}) \cong U^{\oplus 3} \oplus E_8(-1)^{\oplus 2} \oplus \langle -2(n-1) \rangle,$$

similarly $\text{Pic}(S^{[n]}) = i(\text{Pic}(S)) \oplus \mathbb{Z}\delta$: see [Bea83b, §6] for details. The Fujiki constant of the Hilbert scheme of n points on a K3 surface S is $c_{S^{[n]}} = 1$, see [Bea83b, §9]. In particular for K3 surfaces the BBF form coincides with the intersection form. Moreover, the singular cohomology ring $H^*(S^{[n]}, \mathbb{Z})$ for a K3 surface S and $n \geq 1$ is torsion free by [Mar07, Theorem 1]. When $n = 2$, the variety $S^{[2]}$ is usually called *Hilbert square of a K3 surface*. An IHS manifold which is deformation equivalent to the Hilbert square of a K3 surface is said to be of *K3^[2]-type*.

The other known examples of IHS manifolds up to deformation equivalence are *generalised Kummer varieties*, see [Bea83b, §7], an isolated example of dimension 10 and second Betti number $b_2 = 24$, see [O’G99], and an isolated example of dimension 6 and second Betti number $b_2 = 8$, see [O’G03]. We do not discuss details on these examples since in this paper we deal only with Hilbert squares of K3 surfaces.

We conclude this section with the following useful correspondence between primitive elements in $H^2(X, \mathbb{Z})$ and primitive elements in $H_2(X, \mathbb{Z})_f$, where X is an IHS manifold and $H_2(X, \mathbb{Z})_f$ is the torsion free quotient group of the second singular homology $H_2(X, \mathbb{Z})$. See [HT01] for details.

Proposition 3.6 (Hassett–Tschinkel). *Let X be an IHS manifold and denote by $H_2(X, \mathbb{Z})_f$ the torsion free quotient group of $H_2(X, \mathbb{Z})$. Let (\cdot, \cdot) be the BBF bilinear form. Then there is a correspondence between primitive elements in $H^2(X, \mathbb{Z})$ and primitive elements in $H_2(X, \mathbb{Z})_f$. In particular:*

(i) For every primitive element $R \in H_2(X, \mathbb{Z})_f$ there exists a unique class $\omega \in H^2(X, \mathbb{Q})$ such that

$$\epsilon(R \cap v) = (\omega, v) \quad \text{for every } v \in H^2(X, \mathbb{Z}),$$

where $\epsilon : H_0(X, \mathbb{Z}) \xrightarrow{\sim} \mathbb{Z}$ is the isomorphism in [Bre13, Theorem IV.2.1] and \cap is the cap product. The primitive $\rho \in H^2(X, \mathbb{Z})$ associated to R is the primitive element such that $c\rho = \omega$ for some $c \in \mathbb{Q}_{>0}$.

(ii) For every primitive element $\rho \in H^2(X, \mathbb{Z})$ of divisibility $\text{div}(\rho) = d$ in $(H^2(X, \mathbb{Z}), q_X)$, there exists a unique primitive $R \in H_2(X, \mathbb{Z})_f$ such that

$$d \cdot \epsilon(R \cap v) = (\rho, v) \quad \text{for every } v \in H^2(X, \mathbb{Z}).$$

4. GENERALITIES ON IHS MANIFOLDS OF $K3^{[2]}$ -TYPE

Let X be an IHS manifold of dimension 4 of $K3^{[2]}$ -type. In this section we recall the link between the intersection pairing on $H^4(X, \mathbb{Q})$ and the \mathbb{Q} -bilinear extension on $H^2(X, \mathbb{Q})$ of the BBF form, and we introduce the dual q_X^\vee of the BBF quadratic form. We refer to [O'G08, §2].

First of all, we state the following corollary of Verbitsky's results in [Ver96], obtained by Guan in [Gua01], see also [O'G10, Corollary 2.5]. We denote by $b_i(X)$ the i -th Betti number of X .

Proposition 4.1. *Let X be an IHS manifold of dimension 4. Then $b_2(X) \leq 23$. If equality holds then $b_3(X) = 0$ and the map*

$$\text{Sym}^2 H^2(X, \mathbb{Q}) \rightarrow H^4(X, \mathbb{Q})$$

induced by the cup product is an isomorphism. In particular this happens when X is an IHS fourfold of $K3^{[2]}$ -type.

Since X is a compact complex manifold of dimension $\dim_{\mathbb{C}}(X) = 4$, the singular cohomology group $H^4(X, \mathbb{Z})$ has an intersection pairing induced by the cup product:

$$\langle \cdot, \cdot \rangle : H^4(X, \mathbb{Z}) \times H^4(X, \mathbb{Z}) \rightarrow \mathbb{Z}, \quad \langle \alpha, \beta \rangle := \int_X \alpha \beta.$$

We write $\langle \cdot, \cdot \rangle$ also for the \mathbb{Q} -bilinear extension of the intersection pairing above to $H^4(X, \mathbb{Q}) \times H^4(X, \mathbb{Q})$, obtaining a \mathbb{Q} -valued intersection pairing on $\text{Sym}^2 H^2(X, \mathbb{Q})$. Let X be an IHS manifold of $K3^{[2]}$ -type: the following relation between $\langle \cdot, \cdot \rangle$ and the \mathbb{Q} -extension of the BBF form on $H^2(X, \mathbb{Q})$ holds, see [O'G08, Remark 2.1].

Proposition 4.2 (O'Grady). *Let X be an IHS fourfold of $K3^{[2]}$ -type. The intersection pairing $\langle \cdot, \cdot \rangle$ defined above is the bilinear form on $\text{Sym}^2 H^2(X, \mathbb{Q})$ given by*

$$\langle \alpha_1 \alpha_2, \alpha_3 \alpha_4 \rangle = (\alpha_1, \alpha_2)(\alpha_3, \alpha_4) + (\alpha_1, \alpha_3)(\alpha_2, \alpha_4) + (\alpha_1, \alpha_4)(\alpha_2, \alpha_3)$$

for every $\alpha_1, \alpha_2, \alpha_3, \alpha_4 \in H^2(X, \mathbb{Q})$, where (\cdot, \cdot) denotes the BBF bilinear form on $H^2(X, \mathbb{Q})$.

Let q_X be the BBF quadratic form on X . Let $\{e_1, \dots, e_{23}\}$ be a basis of $H^2(X, \mathbb{Q})$ and $\{e_1^\vee, \dots, e_{23}^\vee\}$ be the dual basis in $H^2(X, \mathbb{Q})^\vee$, i.e., $e_i^\vee(e_j) = \delta_{i,j}$. Then we have

$$q_X = \sum_{i,j} g_{i,j} e_i^\vee \otimes e_j^\vee, \quad q_X^\vee = \sum_{i,j} m_{i,j} e_i e_j,$$

where $g_{i,j} := (e_i, e_j)$, the matrix $(g_{i,j})$ is symmetric and $(m_{i,j}) = (g_{i,j})^{-1}$. The values of the products $\langle q_X^\vee, \alpha \rangle$ for every $\alpha \in H^4(X, \mathbb{Q})$ are given by the following proposition, see [O'G08, Proposition 2.2].

Proposition 4.3 (O'Grady). *Let X be an IHS fourfold of $K3^{[2]}$ -type. Let $\langle \cdot, \cdot \rangle$ be the bilinear form described in Proposition 4.2. Then $\langle \cdot, \cdot \rangle$ is non-degenerate and*

$$\begin{aligned} \langle q_X^\vee, \alpha \beta \rangle &= 25(\alpha, \beta) \quad \text{for all } \alpha, \beta \in H^2(X, \mathbb{Q}), \\ \langle q_X^\vee, q_X^\vee \rangle &= 23 \cdot 25. \end{aligned}$$

Recall that *rational Hodge classes* and *integral Hodge classes* of type (k, k) on a projective manifold Y are elements belonging respectively to

$$H^{k,k}(Y, \mathbb{Q}) := H^{2k}(Y, \mathbb{Q}) \cap H^{k,k}(Y), \quad H^{k,k}(Y, \mathbb{Z}) := H^{2k}(Y, \mathbb{Z}) \cap H^{k,k}(Y).$$

O'Grady has shown in [O'G08, §3] that q_X^\vee is a rational multiple of $c_2(X)$, the second Chern class of the tangent bundle of X , in particular it is an element of $H^{2,2}(X, \mathbb{Q})$, i.e., it is a rational Hodge class of type $(2, 2)$.

Proposition 4.4 (O’Grady). *Let X be an IHS fourfold of $K3^{[2]}$ -type. Then $q_X^\vee \in H^{2,2}(X, \mathbb{Q})$, i.e., q_X^\vee is a rational Hodge class of X of type $(2, 2)$, and*

$$\frac{6}{5}q_X^\vee = c_2(X) \in H^{2,2}(X, \mathbb{Z}).$$

Moreover, $\frac{2}{5}q_X^\vee \in H^{2,2}(X, \mathbb{Z})$ is an integral Hodge class of X of type $(2, 2)$.

If S is a projective K3 surface, then $H^{2,2}(S^{[2]}, \mathbb{Z})$ is a lattice, where the bilinear form considered is the cup product. We recall [Nov21, Theorem 8.3], which gives a basis of the lattice $H^{2,2}(S^{[2]}, \mathbb{Z})$ when S is general and the Picard group of S is known.

Theorem 4.5 (N.). *Let S be a general projective K3 surface, $\{b_1, \dots, b_r\}$ be a basis of $\text{Pic}(S)$ and $X := S^{[2]}$. Then:*

(i) $\text{rk}(H^{2,2}(X, \mathbb{Z})) = \frac{(r+1)r}{2} + r + 2$.

(ii) *The following is a basis of the lattice $H^{2,2}(X, \mathbb{Z})$, which is odd:*

$$\left\{ b_i b_j, \frac{b_i^2 - b_i \delta}{2}, \frac{1}{8} \left(\delta^2 + \frac{2}{5} q_X^\vee \right), \delta^2 \right\}_{1 \leq i \leq j \leq r}.$$

In particular, if $S = S_{2t}$ is a generic K3 surface of degree $2t$, and $h \in \text{Pic}(S_{2t}^{[2]})$ is the class induced by the ample generator of $\text{Pic}(S_{2t})$, then

$$H^{2,2}(S_{2t}^{[2]}, \mathbb{Z}) = \mathbb{Z}h^2 \oplus \mathbb{Z} \frac{h^2 - h\delta}{2} \oplus \mathbb{Z} \frac{1}{8} \left(\delta^2 + \frac{2}{5} q_X^\vee \right) \oplus \mathbb{Z} \frac{2}{5} q_X^\vee.$$

Moreover, $\text{disc}(H^{2,2}(S_{2t}^{[2]}, \mathbb{Z})) = 84t^3$ and the Gram matrix in the basis given above is the following:

$$\begin{pmatrix} 12t^2 & 6t^2 & 2t & 20t \\ 6t^2 & t(3t-1) & t & 10t \\ 2t & t & 1 & 9 \\ 20t & 10t & 9 & 92 \end{pmatrix}.$$

Note that in the case of the Hilbert square of a generic K3 surface S_{2t} we have substituted the element δ^2 in a basis of $H^{2,2}(X, \mathbb{Z})$ with $\frac{2}{5}q_X^\vee$: this choice slightly simplifies computations in Section 7.

5. GENERALITIES ON HILBERT SQUARES OF GENERIC K3 SURFACES

In this section we recall some important results on Hilbert squares of generic K3 surfaces of degree $2t$ and we present the main problem of this paper. Recall that the *nef cone* of a projective variety Y is the convex cone $\text{Nef}(Y) \subseteq \text{NS}(Y)_{\mathbb{R}}$ generated by classes of nef divisors in $\text{NS}(Y)_{\mathbb{R}} = \text{NS}(Y) \otimes \mathbb{R}$, where $\text{NS}(Y)$ is the Néron–Severi group of Y , the *moving cone* is the convex cone $\text{Mov}(Y) \subseteq \text{NS}(Y)_{\mathbb{R}}$ generated by classes of divisors whose complete linear system has base locus of codimension greater than 1, and the *pseudoeffective cone* is the convex cone $\overline{\text{Eff}}(Y) \subseteq \text{NS}(Y)_{\mathbb{R}}$ generated by classes of pseudoeffective divisors. We present the following version of the Bayer–Macrì theorem, see [BM14, Proposition 13.1], describes the closure of the moving cone, the nef cone and the pseudoeffective cone of the Hilbert square of a generic K3 surface.

Theorem 5.1 (Bayer–Macrì). *Let S_{2t} be a projective K3 surface with $\text{Pic}(S_{2t}) = \mathbb{Z}H$ and $H^2 = 2t$. Let $X := S_{2t}^{[2]}$ be its Hilbert square. Denote by $h \in \text{Pic}(X)$ the line bundle induced by the ample generator of $\text{Pic}(S_{2t})$. Then the cones of classes of divisors can be described as follows.*

- (1) *The extremal rays of the closure of the moving cone $\text{Mov}(X)$ are spanned by h and $h - \mu_t \delta$, where:*
 - if t is a perfect square, $\mu_t = \sqrt{t}$;
 - if t is not a perfect square and (c, d) is the minimal solution of the Pell equation $P_t(1)$, then $\mu_t = t \cdot \frac{d}{c}$.
- (2) *The extremal rays of the nef cone $\text{Nef}(X)$ are spanned by h and $h - \nu_t \delta$, where:*
 - if the equation $P_{4t}(5)$ is not solvable, $\nu_t = \mu_t$;
 - if the equation $P_{4t}(5)$ is solvable and (a_5, b_5) is its minimal solution, $\nu_t = 2t \cdot \frac{b_5}{a_5}$.
- (3) *The extremal rays of the pseudoeffective cone $\overline{\text{Eff}}(X)$ are spanned by δ and $h - \omega_t \delta$, where:*
 - if t is a perfect square, $\omega_t = \sqrt{t}$;
 - if t is not a perfect square, $\omega_t = \frac{c}{d}$, where (c, d) is the minimal solution of $P_t(1)$.

Let X be as in Theorem 5.1. We recall the description of the group of biregular automorphisms $\text{Aut}(X)$ given by Boissière, Cattaneo, Nieper-Wißkirchen and Sarti in [BCNWS16].

Theorem 5.2 (Proposition 4.3, Proposition 5.1, Lemma 5.3, Theorem 5.5 in [BCNWS16]). *Let S_{2t} be a projective K3 surface with $\text{Pic}(S_{2t}) = \mathbb{Z}H$ and $H^2 = 2t$. Let $h \in \text{Pic}(S_{2t}^{[2]})$ be the line bundle induced by the ample generator of the Picard group $\text{Pic}(S_{2t})$.*

- (1) *If $t = 1$, then S_2 is the double cover of \mathbb{P}^2 branched along a smooth sextic curve and if ι is the covering involution then*

$$\text{Aut}(S_2^{[2]}) = \{\text{id}_{S_2^{[2]}}, \iota^{[2]}\} \cong \mathbb{Z}/2\mathbb{Z}.$$

- (2) *If $t \geq 2$, then $X := S_{2t}^{[2]}$ admits a non-trivial automorphism if and only if one of the following equivalent conditions is satisfied.*

- *The integer t is not a square, the Pell-type equation $P_{4t}(5)$ has no solution and the negative Pell equation $P_t(-1)$ has a solution.*
- *There exists an ample class $D \in \text{NS}(S_{2t}^{[2]})$ such that $q_X(D) = 2$.*

In this case the class D is unique, the automorphism ι is unique and it is a non-natural anti-symplectic involution. Its action on $\text{NS}(S_{2t}^{[2]})$ is the reflection in the span of the class D of square 2 represented in the basis $\{h, -\delta\}$ by the matrix

$$\begin{pmatrix} c & -d \\ td & -c \end{pmatrix}, \quad (5.1)$$

i.e., $\iota^(xh - y\delta) = (cx - dy)h - (tdx - cy)\delta$, where (c, d) is the minimal solution of the Pell equation $P_t(1)$.*

Let X be the Hilbert square of a generic K3 surface of degree $2t$, and suppose that X admits an ample class $D \in \text{Pic}(X)$ with $q_X(D) = 2$. By Theorem 5.1 the nef cone $\text{Nef}(X)$ and the closure of the moving cone $\text{Mov}(X)$ coincide, and by Theorem 5.2 the action on $\text{NS}(S_{2t}^{[2]}) \otimes \mathbb{R}$ of the anti-symplectic involution ι preserves the nef cone of $S_{2t}^{[2]}$, exchanging the two extremal rays. This is true also for a non-natural non-trivial birational automorphism $\iota \in \text{Bir}(S_{2t}^{[2]})$ on the Hilbert square of a generic K3 surface S_{2t} of degree $2t$, whose action on $\text{NS}(S_{2t}^{[2]}) \otimes \mathbb{R}$ preserves the closure of the moving cone exchanging the two extremal rays, see [Mar11, Lemma 6.22]. For a description of $\text{Bir}(S_{2t}^{[2]})$ see [DM19, Proposition B.3], for a description of $\text{Aut}(S_{2t}^{[n]})$ for $n \geq 3$ see [Cat19] and for a description of $\text{Bir}(S_{2t}^{[n]})$ for $n \geq 3$ see [BC20].

Let $X = S_{2t}^{[2]}$ be as above: the ample divisor D with $q_X(D) = 2$ is unique by Theorem 5.2 and its class in $\text{Pic}(X)$ is $bh - a\delta$, where $h \in \text{Pic}(X)$ is the line bundle induced by the ample generator of $\text{Pic}(S_{2t})$, and (a, b) is the minimal solution of the negative Pell equation $P_t(-1)$. By the Kodaira vanishing theorem and a Riemann–Roch formula for IHS fourfolds of $K3^{[2]}$ -type, see [O’G10, Formula (2.2.7)], we have $\dim(H^0(X, \mathcal{O}_X(D))) = 6$. Hence the rational map $\varphi_{|D|}$ induced by the complete linear system $|D|$ has \mathbb{P}^5 as codomain, i.e.,

$$\varphi_{|D|} : X \dashrightarrow \mathbb{P}^5.$$

We can now state the main problem of this paper.

Problem. *Let X be the Hilbert square of a generic K3 surface of degree $2t$ such that X admits an ample divisor D with $q_X(D) = 2$. Describe the base locus of the complete linear system $|D|$ and describe the rational map*

$$\varphi_{|D|} : X \dashrightarrow \mathbb{P}^5.$$

As observed in [BCNWS16, §1], the first values of t such that $X = S_{2t}^{[2]}$ as in Theorem 5.2 has an ample divisor D with $q_X(D) = 2$ are $t = 2, 10, 13$, etc. The case $t = 2$ was studied by Welters in [Wel81] and Beauville in [Bea83a]: in this case $S_4 \subset \mathbb{P}^3$ is a smooth quartic surface of \mathbb{P}^3 with Picard rank 1, in particular S_4 does not contain any line, the map $\varphi_{|D|}$ is a finite morphism of degree 6 with image $\mathbb{G}(1, \mathbb{P}^3) \subset \mathbb{P}^5$, the Grassmannian of lines in \mathbb{P}^3 , and $\varphi_{|D|}(x) := l_x$ is the unique line in \mathbb{P}^3 which contains the support of the subscheme x of S_4 . Moreover, if $l_x \cap S_4 = x \cup x'$, where $x, x' \in S_4^{[2]}$, then the anti-symplectic involution $\iota \in \text{Aut}(X)$ is the *Beauville involution*, defined as $\iota(x) := x'$. Since S_4 does not contain any line, ι is everywhere well defined. Note that $\varphi_{|D|}$ factors through the quotient $\pi : S_4^{[2]} \rightarrow S_4^{[2]}/\langle \iota \rangle$ with respect to the Beauville involution.

A first step in the analysis of the base locus of $|D|$ in the Problem above is given by the following result, which is a corollary of [Rie18, Theorem 4.7].

Proposition 5.3. *Let X be the Hilbert square of a projective K3 surface of Picard rank 1. Consider a big and nef divisor $D \in \text{Div}(X)$. Then $|D|$ has no fixed part, i.e., the base locus of $|D|$ has codimension greater than 1.*

Proof. By [Rie18, Theorem 4.7] the complete linear system has a fixed part if and only if $D = mL + F$, where $m \geq 2$, the class of L is movable with $q_X(L) = 0$ and F is a reduced and irreducible divisor with $q_X(F) < 0$ and $(L, F) = 1$. Since $\text{Pic}(X) = \mathbb{Z}h \oplus \mathbb{Z}\delta$, where h is the line bundle induced by the ample generator of the Picard group of the underlying K3 surface, and $(h, \delta) = 0$, there are no elements L and F in $\text{Pic}(X)$ such that $(L, F) = 1$. We conclude that $|D|$ has no fixed part. \square

We now describe the set $H^{3,3}(S_{2t}^{[2]}, \mathbb{Z})$ of integral Hodge classes of type $(3, 3)$ on the Hilbert square of a K3 surface S_{2t} with $\text{Pic}(S_{2t}) = \mathbb{Z}H$ and $H^2 = 2t$: this will be useful in the proof of Proposition 8.3.

Theorem 5.4. *Let S be a K3 surface and suppose that $\{b_1, \dots, b_r\}$ is a basis of $\text{Pic}(S^{[2]})$. Then a basis of the \mathbb{Z} -module $H^{3,3}(S^{[2]}, \mathbb{Z})$ is given by $\{cl(b_1^\vee), \dots, cl(b_r^\vee)\}$, where $b_i^\vee \in H_2(S^{[2]}, \mathbb{Z})$ is the primitive element associated to $b_i \in H^2(S^{[2]}, \mathbb{Z})$ by Proposition 3.6, and $cl(b_i^\vee)$ is the inverse of the Poincaré dual of b_i^\vee . In particular, let $X = S_{2t}^{[2]}$ be the Hilbert square of a projective K3 surface with $\text{Pic}(S_{2t}) = \mathbb{Z}H$ and $H^2 = 2t$, and let $h \in \text{Pic}(X)$ be the class induced by the ample generator of $\text{Pic}(S_{2t})$. Consider $h_6^\vee := cl(h^\vee)$ and $\delta_6^\vee := cl(\delta^\vee) \in H^6(X, \mathbb{Z})$. Then*

$$H^{3,3}(X, \mathbb{Z}) \cong \mathbb{Z}h_6^\vee \oplus \mathbb{Z}\delta_6^\vee.$$

Moreover, $h_6^\vee = \frac{1}{6t}h^3$ and $\delta_6^\vee = \frac{1}{4t}h^2\delta$, and the following equalities hold in $H^{3,3}(X, \mathbb{Z})$:

$$\delta^3 = -\frac{3}{t}h^2\delta, \quad h\delta^2 = -\frac{1}{3t}h^3, \quad q_X^\vee h = \frac{25}{6t}h^3, \quad q_X^\vee \delta = \frac{25}{2t}h^2\delta.$$

Proof. The torsion free quotient group $H_2(S^{[2]}, \mathbb{Z})_f$ is a Hodge structure of weight -2 . Moreover the cap product

$$H^2(S^{[2]}, \mathbb{Z}) \otimes H_2(S^{[2]}, \mathbb{Z})_f \rightarrow H_0(S^{[2]}, \mathbb{Z})$$

is a morphism of Hodge structures of weight 0, see for instance [PS08], hence if $R \in H_2(S^{[2]}, \mathbb{Z})_f$ is primitive of type $(-1, -1)$, the corresponding $\rho \in H^2(S^{[2]}, \mathbb{Z})$ is of type $(1, 1)$. Thus $\{b_1^\vee, \dots, b_r^\vee\}$ is a basis of

$$H_{-1,-1}(S^{[2]}, \mathbb{Z}) := H_2(S^{[2]}, \mathbb{Z}) \cap H_{-1,-1}(S^{[2]}),$$

where $H_{-1,-1}(S^{[2]})$ is the component of type $(-1, -1)$ of $H_2(S^{[2]}, \mathbb{C})$. The Poincaré duality

$$PD : H^6(S^{[2]}, \mathbb{Z}) \xrightarrow{\sim} H_2(S^{[2]}, \mathbb{Z})_f$$

is an isomorphism of Hodge structures of weight -4 , hence $H^{3,3}(S^{[2]}, \mathbb{Z}) \cong H_{-1,-1}(S^{[2]}, \mathbb{Z})$, so $\{cl(b_1^\vee), \dots, cl(b_r^\vee)\}$ is a basis of $H^{3,3}(S^{[2]}, \mathbb{Z})$.

Let now $X = S_{2t}^{[2]}$ be the Hilbert square of a projective K3 surface with $\text{Pic}(S_{2t}) = \mathbb{Z}H$ and $H^2 = 2t$, and let $h \in \text{Pic}(X)$ be the line bundle induced by the ample generator of $\text{Pic}(S_{2t})$. The cup product with h^2 gives an isomorphism between $H^{1,1}(X, \mathbb{Q})$ and $H^{3,3}(X, \mathbb{Q})$. Then $H^{3,3}(X, \mathbb{Q})$ is 2-dimensional and it is generated by h^3 and $h^2\delta$ since $H^{1,1}(X, \mathbb{Q}) \cong \text{Pic}(X) \otimes \mathbb{Q}$. We now represent δ^3 , $h\delta^2$, $q_X^\vee h$ and $q_X^\vee \delta$ in terms of this basis. If $\delta^3 = xh^3 + yh^2\delta$ with $x, y \in \mathbb{Q}$, by Proposition 4.2 we have

$$h\delta^3 = 0 = x\langle h^2, h^2 \rangle + y\langle h^2, h\delta \rangle = 12t^2x, \quad \delta^4 = 3 \cdot (-2)^2 = x\langle h^2, h\delta \rangle + y\langle h^2, \delta^2 \rangle = -4ty,$$

which give $x = 0$ and $y = -\frac{3}{t}$, hence

$$\delta^3 = -\frac{3}{t}h^2\delta.$$

Similarly we obtain

$$h\delta^2 = -\frac{1}{3t}h^3, \quad q_X^\vee h = \frac{25}{6t}h^3, \quad q_X^\vee \delta = \frac{25}{2t}h^2\delta.$$

Consider now $H^{3,3}(X, \mathbb{Z})$. The cohomology group $H^6(X, \mathbb{Z})$ is torsion free, and $H^{3,3}(X, \mathbb{Q})$ has dimension 2, hence $\text{rk}(H^{3,3}(X, \mathbb{Z})) = 2$. Let $h^\vee, \delta^\vee \in H_2(X, \mathbb{Z})$ be the primitive classes which correspond, by Proposition 3.6, respectively to the primitive classes $h, \delta \in H^2(X, \mathbb{Z})$. Using Proposition 3.6 one gets $h^\vee = h$ and $\delta^\vee = \frac{\delta}{2}$ seen as elements in $H^2(X, \mathbb{Q})$. Let PD be the Poincaré duality: then $h_6^\vee := PD^{-1}(h^\vee)$ and $\delta_6^\vee := PD^{-1}(\delta^\vee)$ give a basis of $H^{3,3}(X, \mathbb{Z})$, so we have

$$H^{3,3}(X, \mathbb{Z}) \cong \mathbb{Z}h_6^\vee \oplus \mathbb{Z}\delta_6^\vee.$$

Moreover, if \cdot denotes the cup product and \cap the cap product, we get

$$\int_X h_6^\vee \cdot x = \epsilon(h^\vee \cap x), \quad \int_X \delta_6^\vee \cdot x = \epsilon(\delta^\vee \cap x) \quad (5.2)$$

for every $x \in H^2(X, \mathbb{Z})$, see [Hat05, p.249], and $\epsilon : H^0(X, \mathbb{Z}) \xrightarrow{\sim} \mathbb{Z}$. Again by Proposition 3.6 we have

$$\epsilon(h^\vee \cap h) = (h, h) = 2t, \quad \epsilon(h^\vee \cap \delta) = (h, \delta) = 0. \quad (5.3)$$

If we write $h_6^\vee = \alpha h^3 + \beta h^2 \delta$ for some $\alpha, \beta \in \mathbb{Q}$, by Proposition 4.2 we obtain

$$\int_X h_6^\vee \cdot h = \alpha \langle h^2, h^2 \rangle + \beta \langle h^2, h\delta \rangle = 12\alpha t^2, \quad \int_X h_6^\vee \cdot \delta = \alpha \langle h^2, h\delta \rangle + \beta \langle h^2, \delta^2 \rangle = -4\beta t. \quad (5.4)$$

Then (5.2), (5.3) and (5.4) give $\alpha = \frac{1}{6t}$ and $\beta = 0$, hence $h_6^\vee = \frac{1}{6t}h^3$. Similarly we obtain $\delta_6^\vee = \frac{1}{4t}h^2\delta$. \square

6. FIXED LOCUS OF THE ANTI-SYMPLECTIC INVOLUTION

Let X be the Hilbert square of a generic K3 surface S_{2t} of degree $2t$ such that X admits an ample divisor D with $q_X(D) = 2$. We recall some properties of $F := \text{Fix}(\iota)$, the locus of points of X fixed by the anti-symplectic involution $\iota \in \text{Aut}(X)$, then we compute its integral Hodge class in $H^{2,2}(X, \mathbb{Z})$ and we show that the rational map $\varphi_{|D|}$ induced by the complete linear system $|D|$ factors through the quotient $\pi : X \rightarrow X/\langle \iota \rangle$ with respect to ι . As usual we denote by $h \in \text{Pic}(X)$ the line bundle induced by the ample generator of $\text{Pic}(S_{2t})$. Recall that the class of D in $\text{Pic}(X)$ is $bh - a\delta$, where (a, b) is the minimal solution of the negative Pell equation $P_t(-1)$.

First of all, we prove that F is a connected surface in our case.

Proposition 6.1. *Keep notation as above. Then $F = \text{Fix}(\iota)$ is a connected Lagrangian surface in X .*

Proof. By [Bea11, Lemma 1] the fixed locus F is a Lagrangian surface. We now show that F is connected. Let $\mathcal{M}_{\langle 2 \rangle}^\rho$ be the moduli space which parametrizes triples (X, ι_X, i_X) , where X is an IHS manifold of $K3^{[2]}$ -type, $\iota_X \in \text{Aut}(X)$ is an anti-symplectic involution whose action on $H^2(X, \mathbb{Z})$ is the reflection in the class of an ample divisor D with $q_X(D) = 2$ and $i_X : \langle 2 \rangle \hookrightarrow \text{NS}(X)$ is a primitive embedding such that $i(\langle 2 \rangle)$ contains the class of D . Any two points in the moduli space $\mathcal{M}_{\langle 2 \rangle}^\rho$ are deformation equivalent by [BCMS19, Corollary 4.1, Theorem 5.1]. Since deformation equivalence preserves topological properties of $\text{Fix}(\iota)$, it suffices to find a point $(X, \iota_X, i_X) \in \mathcal{M}_{\langle 2 \rangle}^\rho$ such that $\text{Fix}(\iota_X)$ is connected. Let $X = S_4^{[2]}$ be the Hilbert square of a smooth quartic surface of \mathbb{P}^3 with Picard rank 1, and let $\iota_X \in \text{Aut}(X)$ be the Beauville involution: the surface $\text{Fix}(\iota_X)$ is connected by [Wel81, Corollary 3.4.4]. Alternatively, if we show that the class of D , which is $bh - a\delta$, has divisibility 1 in the lattice $(H^2(X, \mathbb{Z}), q_X)$, then the surface $\text{Fix}(\iota)$ is connected by [FMOS20, Main Theorem]. Since (a, b) is the minimal solution of the negative Pell equation $P_t(-1)$, the integers a and b are coprime and b is odd by Proposition 2.6. Since $H^2(S_{2t}, \mathbb{Z})$ is unimodular, there exists $x \in H^2(X, \mathbb{Z})$ such that $(h, x) = 1$, hence there exist $\alpha, \beta \in \mathbb{Z}$ such that $(D, \alpha\delta + \beta x) = 1$. We conclude that $\text{div}(bh - a\delta) = 1$, and F is connected. \square

Keep notation as above. By [CGM19, §3] the quotient variety $X/\langle \iota \rangle$ is singular along $\pi'(F)$, where $\pi' : X \rightarrow X/\langle \iota \rangle$ is the quotient map. The desingularization of $X/\langle \iota \rangle$, which we call W , is the blow-up of $X/\langle \iota \rangle$ along its singular locus, and it is a Calabi–Yau variety. Equivalently, consider the blow-up $\text{Bl}_F(X)$ of X in the locus $F = \text{Fix}(\iota)$ of points fixed by ι . The involution ι gives rise to an involution $\tilde{\iota}$ on $\text{Bl}_F(X)$ which fixes the exceptional divisor $E \subset \text{Bl}_F(X)$. One can show that the quotient $\text{Bl}_F(X)/\langle \tilde{\iota} \rangle$ is isomorphic to W , obtaining the following commutative diagram, see [CGM19, Theorem 3.6] for more details:

$$\begin{array}{ccc} \text{Bl}_F(X) & \xrightarrow{\pi} & W \\ \downarrow \beta & & \downarrow \beta' \\ X & \xrightarrow{\pi'} & X/\langle \iota \rangle. \end{array} \quad (6.1)$$

We now state the following useful technical lemma. See [DHH⁺15, Lemma 4.3] for a more general statement and for the proof.

Lemma 6.2. *Let $\pi : X \rightarrow Y$ be a double cover of a smooth projective variety such that the branch locus is a smooth prime divisor $B \in \text{Div}(Y)$, and denote by $\iota \in \text{Aut}(X)$ the involution associated to the double cover. If $\text{Pic}(X)^\iota$ is the subgroup of ι -invariant line bundles on X , then $\pi^*\text{Pic}(Y) \cong \text{Pic}(X)^\iota$.*

Similarly to the case in [vGS07, §2.6], since $\iota^*D \cong D$ there is an involution induced by ι on $\mathbb{P}(H^0(X, \mathcal{O}_X(D))^\vee)$ which has two fixed spaces \mathbb{P}^a and \mathbb{P}^b , where $a + 1 + b + 1 = 6$ and $a = -1$ if the corresponding eigenspace of ι^* on $H^0(X, \mathcal{O}_X(D))$ is zero, similarly for b . See [MFK94, p.32] for details on the action of ι on $H^0(X, \mathcal{O}_X(D))$. In particular the rational map $\varphi_{|D|}$ induced by the complete linear system $|D|$ factors through the quotient $X \rightarrow X/\langle \iota \rangle$ if and only if the action induced by ι on $\mathbb{P}(H^0(X, \mathcal{O}_X(D))^\vee)$ is trivial.

Let $B \in \text{Div}(W)$ be the branch divisor of $\pi : \text{Bl}_F(X) \rightarrow W$. By [BHPVdV15, Lemma I.17.1] there exists a divisor $N \in \text{Div}(W)$ such that $\mathcal{O}_W(2N) = \mathcal{O}_W(B)$ in $\text{Pic}(W)$. Let D be the ample divisor on X with $q_X(D) = 2$. We show the following result, similar to [vGS07, Proposition 2.7, Item (2)] obtained in the case of K3 surfaces admitting a symplectic involution.

Proposition 6.3. *Keep notation as above. Consider diagram (6.1). Let $\mathcal{D} := \mathcal{O}_X(D)$ and $\mathcal{N} := \mathcal{O}_W(N)$. There exists a line bundle $\mathcal{D}_W \in \text{Pic}(W)$ such that $\pi^*\mathcal{D}_W \cong \beta^*\mathcal{D}$. Moreover, the vector space $H^0(X, \mathcal{D})$ decomposes as*

$$H^0(X, \mathcal{D}) \cong H^0(W, \mathcal{D}_W) \oplus H^0(W, \mathcal{D}_W - \mathcal{N}), \quad (6.2)$$

which is the decomposition of $H^0(X, \mathcal{D})$ into ι^* -eigenspaces.

Proof. Consider $\pi : \text{Bl}_F(X) \rightarrow W$ appearing in diagram (6.1): by Lemma 6.2 we obtain a divisor $D_W \in \text{Div}(W)$ whose class \mathcal{D}_W in $\text{Pic}(W)$ is such that $\pi^*\mathcal{D}_W \cong \beta^*\mathcal{D}$ in $\text{Pic}(\text{Bl}_F(X))$. Note that Lemma 6.2 can be applied: $F \subset X$ is smooth and connected by Proposition 6.1, so the exceptional divisor of β and the branch divisor of π are smooth prime divisors. Since β is a birational morphism, we have $H^0(X, \mathcal{D}) \cong H^0(\text{Bl}_F(X), \beta^*\mathcal{D})$, which is isomorphic to $H^0(\text{Bl}_F(X), \pi^*\mathcal{D}_W)$, being $\beta^*\mathcal{D} = \pi^*\mathcal{D}_W$. Moreover,

$$\begin{aligned} \pi_*(\pi^*\mathcal{D}_W) &\cong \pi_*(\pi^*\mathcal{D}_W \otimes \mathcal{O}_{\text{Bl}_F(X)}) \\ &\cong \mathcal{D}_W \otimes \pi_*\mathcal{O}_{\text{Bl}_F(X)}, \end{aligned}$$

where in the second isomorphism we use the projection formula. By [BHPVdV15, Lemma I.17.2] we have

$$\pi_*\mathcal{O}_{\text{Bl}_F(X)} \cong \mathcal{O}_W \oplus \mathcal{O}_W(-N),$$

where $B \in \text{Div}(W)$ is the branch divisor and $N \in \text{Div}(W)$ is the divisor such that $\mathcal{O}_W(2N) = \mathcal{O}_W(B)$ in $\text{Pic}(W)$. Thus we obtain the isomorphism

$$\pi_*(\pi^*\mathcal{D}_W) \cong \mathcal{D}_W \otimes (\mathcal{O}_W \oplus \mathcal{O}_W(-N)) \cong \mathcal{D}_W \oplus (\mathcal{D}_W - \mathcal{N}).$$

Hence decomposition (6.2) holds. Moreover, this is the decomposition of $H^0(X, \mathcal{D})$ in ι^* -eigenspaces. Indeed, two global sections $s, t \in H^0(X, \mathcal{D})$ are in the same eigenspace if and only if the rational function $f = s/t$ is ι -invariant, and this is true when the sections belong both to $H^0(W, \mathcal{D}_W)$ or $H^0(W, \mathcal{D}_W - \mathcal{N})$ in the decomposition, hence each of these two spaces is contained in an eigenspace of $H^0(X, \mathcal{D})$. We conclude that $H^0(W, \mathcal{D}_W)$ and $H^0(W, \mathcal{D}_W - \mathcal{N})$ are isomorphic to the two eigenspaces by decomposition (6.2). \square

We then see from Proposition 6.3 that the action induced by ι on $\mathbb{P}(H^0(X, \mathcal{O}_X(D)))^\vee$ is trivial, i.e., $\varphi_{|D|}$ factors through the quotient $X \rightarrow X/\langle \iota \rangle$, if and only if either $H^0(X, \mathcal{D}_W)$ or $H^0(W, \mathcal{D}_W - \mathcal{N})$ is zero. We can now prove the main theorem of this section, which is the first step of the geometrical description of the rational map $\varphi_{|D|}$ induced by the complete linear system $|D|$.

Theorem 6.4. *Let $X = S_{2t}^{[2]}$ be the Hilbert square of a generic K3 surface of degree $2t$. Suppose that X admits an ample divisor D with $q_X(D) = 2$. Let $\iota : X \rightarrow X$ be the anti-symplectic involution which generates $\text{Aut}(X)$ and $F := \text{Fix}(\iota)$ be the locus of points of X fixed by ι . Then*

$$[F] = 5D^2 - \frac{2}{5}q_X^\vee \in H^{2,2}(X, \mathbb{Z}), \quad (6.3)$$

where $[F]$ denotes the fundamental cohomological class of F in $H^{2,2}(X, \mathbb{Z})$. Moreover, let $\varphi_{|D|} : X \dashrightarrow \mathbb{P}^5$ be the rational map induced by the complete linear system $|D|$. Then $\varphi_{|D|}$ factors through the quotient $X \rightarrow X/\langle \iota \rangle$, i.e., the following diagram is commutative:

$$\begin{array}{ccc} X & \xrightarrow{\varphi_{|D|}} & \mathbb{P}^5 \\ & \searrow & \uparrow \\ & X/\langle \iota \rangle & \end{array} \quad (6.4)$$

Proof. By Proposition 6.3 there exists $\mathcal{D}_W \in \text{Pic}(W)$ such that $\pi^*\mathcal{D}_W \cong \beta^*\mathcal{D}$, using the notation of diagram (6.1). Applying [CGM19, Proposition 7.3] we have

$$\dim(H^0(W, \mathcal{D}_W)) = \frac{7}{2} + \frac{1}{16}(D|_F)^2,$$

hence by decomposition (6.2) and $\dim(H^0(X, \mathcal{D})) = 6$ we get $\dim(H^0(W, \mathcal{D}_W)) \in \{0, 1, \dots, 6\}$. Thus we obtain the following possible values for $(D|_F)^2$:

$$\begin{aligned} \dim(H^0(W, \mathcal{D}_W)) = 0 &\Leftrightarrow (D|_F)^2 = -56, & \dim(H^0(W, \mathcal{D}_W)) = 4 &\Leftrightarrow (D|_F)^2 = 8, \\ \dim(H^0(W, \mathcal{D}_W)) = 1 &\Leftrightarrow (D|_F)^2 = -40, & \dim(H^0(W, \mathcal{D}_W)) = 5 &\Leftrightarrow (D|_F)^2 = 24, \\ \dim(H^0(W, \mathcal{D}_W)) = 2 &\Leftrightarrow (D|_F)^2 = -24, & \dim(H^0(W, \mathcal{D}_W)) = 6 &\Leftrightarrow (D|_F)^2 = 40. \\ \dim(H^0(W, \mathcal{D}_W)) = 3 &\Leftrightarrow (D|_F)^2 = -8, \end{aligned} \quad (6.5)$$

Since D is ample, $(D|_F)^2 > 0$ by the Nakai–Moishezon criterion. This implies that $\dim(H^0(W, \mathcal{D}_W)) \in \{4, 5, 6\}$. We show that $\dim(H^0(W, \mathcal{D}_W)) = 6$ by computing $[F] \in H^{2,2}(X, \mathbb{Z})$, where $[F]$ is the fundamental cohomological class of the fixed locus $F = \text{Fix}(\iota)$ of the involution ι . Let $h \in \text{Pic}(X)$ be the line bundle induced by the ample generator of $\text{Pic}(S_{2t})$. We can write, see for instance [Nov21, Proposition 7.1],

$$[F] = xh^2 + yh\delta + z\delta^2 + w\frac{2}{5}q_X^\vee \in H^{2,2}(X, \mathbb{Z}), \quad (6.6)$$

with $x, y, z, w \in \mathbb{Q}$ to determine. We write D also for its class in $\text{Pic}(X)$: then $D = bh - a\delta$, with (a, b) minimal solution of the Pell-type equation $P_t(-1)$. We denote by $\langle \cdot, \cdot \rangle$ the bilinear form of $H^4(X, \mathbb{Z})$ of Proposition 4.2. We have the following four conditions.

- (1) $\langle [F], (\sigma + \bar{\sigma})^2 \rangle = 0$, where σ is the symplectic form $\sigma \in H^0(X, \Omega_X^2)$, since by [Bea11, Lemma 1] the surface F is Lagrangian. If $\eta := (\sigma + \bar{\sigma}, \sigma + \bar{\sigma})$, we have:

$$\langle h^2, (\sigma + \bar{\sigma})^2 \rangle = 2t\eta, \quad \langle \delta^2, (\sigma + \bar{\sigma})^2 \rangle = -2\eta, \quad \langle h\delta, (\sigma + \bar{\sigma})^2 \rangle = 0, \quad \langle \frac{2}{5}q_X^\vee, (\sigma + \bar{\sigma})^2 \rangle = 10\eta. \quad (6.7)$$

Note that $\eta \neq 0$, see [GHJ12, Definition 22.10, Theorem 23.14], so we obtain from $\langle [F], (\sigma + \bar{\sigma})^2 \rangle = 0$ and (6.6) the following condition:

$$tx - z + 5w = 0. \quad (6.8)$$

- (2) Applying [Bea11, Theorem 2] in our case we have $c_2(F) = 192$, i.e., $\langle [F], [F] \rangle = 192$. This gives, together with (6.6), the following condition:

$$3t^2x^2 - ty^2 + 3z^2 + 23w^2 - 2txz + 10txw - 10zw = 48. \quad (6.9)$$

- (3) Recall that by Theorem 5.2 the action of ι on $\text{Pic}(X) \otimes \mathbb{Q} \cong H^{1,1}(X, \mathbb{Q})$ is described in the basis $\{h, -\delta\}$ by the matrix (5.1). Consider the action induced by ι on $H^{2,2}(X, \mathbb{Q})$, described in the basis $\{h^2, h\delta, \delta^2, q_X^\vee\}$ by

$$\iota^*(h^2) = \iota^*h \cdot \iota^*h, \quad \iota^*(h\delta) = \iota^*h \cdot \iota^*\delta, \quad \iota^*(\delta^2) = \iota^*\delta \cdot \iota^*\delta, \quad \iota^*(q_X^\vee) = q_X^\vee, \quad (6.10)$$

where \cdot denotes the cup product: the last equation comes from the equality $c_2(X) = \frac{6}{5}q_X^\vee$ of Proposition 4.4 and the fact that ι is an automorphism. Since F is the locus of points fixed by ι , we have $\iota^*([F]) = [F]$, i.e., if

$$\iota^*([F]) = \tilde{x}h^2 + \tilde{y}h\delta + \tilde{z}\delta^2 + \frac{2}{5}\tilde{w}q_X^\vee,$$

with $\tilde{x}, \tilde{y}, \tilde{z}, \tilde{w} \in \mathbb{Q}$, then $x = \tilde{x}$, $y = \tilde{y}$, $z = \tilde{z}$, $w = \tilde{w}$. Imposing $x = \tilde{x}$ we obtain the following condition:

$$x - c^2x - cdy - d^2z = 0, \quad (6.11)$$

where (c, d) is the minimal solution of the Pell equation $P_t(1)$. One remarks a posteriori that $y = \tilde{y}$, $z = \tilde{z}$ and $w = \tilde{w}$ give the same condition (6.11).

- (4) By $(D|_F)^2 \in \{8, 24, 40\}$ we have $\langle [F], D^2 \rangle \in \{8, 24, 40\}$, since the bilinear form $\langle \cdot, \cdot \rangle$ represents the intersection form by Proposition 4.2. This gives, together with (6.6), the following possibilities:

$$\begin{aligned} (t + 2t^2b^2)x + 2abty + (2a^2 - 1)z + 5w = 2 &\iff \langle [F], D^2 \rangle = 8, \\ (t + 2t^2b^2)x + 2abty + (2a^2 - 1)z + 5w = 6 &\iff \langle [F], D^2 \rangle = 24, \\ (t + 2t^2b^2)x + 2abty + (2a^2 - 1)z + 5w = 10 &\iff \langle [F], D^2 \rangle = 40. \end{aligned} \quad (6.12)$$

If $\langle [F], D^2 \rangle = 8$, which is equivalent to $\dim(H^0(W, \mathcal{D}_W)) = 4$ by (6.5), the system given by (6.8), (6.9), (6.11) and the first condition in (6.12) has solutions with $x, y, z, w \notin \mathbb{Q}$, which is impossible. Similarly we cannot have $\langle [F], D^2 \rangle = 24$, i.e., $\dim(H^0(W, \mathcal{D}_W)) = 5$. We conclude that $\langle [F], D^2 \rangle = 40$, which is equivalent to $\dim(H^0(W, \mathcal{D}_W)) = 6$. With the help of a computer, the system given by (6.8), (6.9), (6.11) and the third condition in (6.12) implies $w \in \{-\frac{13}{12}, -1\}$.

By Theorem 4.5 we cannot have $w = -\frac{13}{12}$, hence $w = -1$. Imposing $w = -1$, we necessarily obtain only one admissible solution, which is the following:

$$\begin{cases} x = 5b^2, \\ y = -10ab, \\ z = 5a^2, \\ w = -1. \end{cases}$$

We conclude that

$$[F] = 5D^2 - \frac{2}{5}q_X^\vee \in H^{2,2}(X, \mathbb{Z}).$$

Moreover, we have obtained $H^0(X, \mathcal{D}) \cong H^0(W, \mathcal{D}_W)$, so $H^0(W, \mathcal{D}_W - \mathcal{N}) = \{0\}$, which shows that the action induced by ι on $\mathbb{P}(H^0(X, \mathcal{D})^\vee)$ is trivial, i.e., $\varphi_{|D|}$ factors through the quotient $X \rightarrow X/\langle \iota \rangle$. \square

Remark 6.5. Ferretti in [Fer12, Lemma 4.1] obtained the same relation of (6.3) in the Chow ring of a smooth double EPW sextic X : in that case F is the branch locus of the double cover $f : X \rightarrow Y \subset \mathbb{P}^5$, where Y is an EPW sextic.

7. IRREDUCIBILITY PROPERTY

Let X be the Hilbert square of a generic K3 surface of degree $2t$ such that X admits an ample divisor D with $q_X(D) = 2$. In this section we prove the *irreducibility property* when $t \neq 2$, which will be, together with Theorem 6.4, fundamental to obtain the main result of this paper. This can be seen as an analogue of the irreducibility property in [OG08, Proposition 4.1.(1)]. First of all we show that every divisor in the complete linear system $|D|$ is a prime divisor.

Proposition 7.1. *Keep notation as above. Then every divisor $D' \in |D|$ is reduced and irreducible.*

Proof. By abuse of notation, we write D for an effective divisor $D' \in |D|$. Since $q_X(D) = 2$, the divisor D is reduced, i.e., it is not of the form $D = \alpha E$ with $\alpha \in \mathbb{Z}$, $\alpha \neq \pm 1$, and $E \in \text{Div}(X)$. Suppose that

$$D = D_1 + D_2 \tag{7.1}$$

where $D_1 = \sum_i n_i D_{1,i}$ and $D_2 = \sum_j m_j D_{2,j}$ are effective divisors with $D_{1,i}, D_{2,j}$ prime divisors which are pairwise distinct, and $n_i, m_j \in \mathbb{Z}_{>0}$. Without loss of generality we can assume that D_1 has only one component. By abuse of notation, we write D_i and $D_{i,j}$ also for their classes in $H^2(X, \mathbb{Z})$: we show that (7.1) seen in $H^2(X, \mathbb{Z})$ gives a contradiction. We have

$$q_X(D) = 2 = q_X(D_1) + q_X(D_2) + 2(D_1, D_2). \tag{7.2}$$

Since D_1 and D_2 are effective divisors with no common components, we can apply [Bou04, Proposition 4.2.(i)], obtaining $(D_1, D_2) \geq 0$. Note that we cannot have neither $q_X(D_1) = 0$ nor $q_X(D_2) = 0$. Indeed, the equation $q_X(xh - y\delta) = 0$ has a non zero solution $(x, y) \in \mathbb{Z}^2$ only when t is a perfect square. If t was a perfect square, the Pell-type equation $P_t(-1)$ would be solvable only for $t = 1$, which is a case that we are not considering. Hence t is not a perfect square and $q_X(D_1) \neq 0$, $q_X(D_2) \neq 0$.

Since $H^2(X, \mathbb{Z})$ is an even lattice, we have only two possibilities: either one between D_1 and D_2 is zero, or (at least) one of the two has negative square with respect to the BBF form. We show that $q_X(D_i) > 0$ for $i = 1, 2$. Assume by contradiction that at least one between D_1 and D_2 has negative square. This can happen only if there exists a component $D_{1,i}$ or $D_{2,j}$ whose square with respect to the BBF form is negative. Without loss of generality, we can suppose that $q_X(D_{1,1}) < 0$. Recall that $H^2(X, \mathbb{Z}) \cong H^2(S_{2t}, \mathbb{Z}) \oplus \langle -2 \rangle$, where S_{2t} is a K3 surface, and $H^2(S_{2t}, \mathbb{Z})$ is a unimodular lattice. Then the divisibility in $H^2(X, \mathbb{Z})$ of a primitive class can be only 1 or 2. Hence we have either $q_X(D_{1,1}) = -2$ or $q_X(D_{1,1}) = -4$ by [Mar13, Lemma 3.7], see also [Rie18, Lemma 3.5]. We show that $q_X(D_{1,1}) = -4$ is not possible. Indeed, if $t = 2$, the class of $D_{1,1}$ in $\text{NS}(X)$ is necessarily $h - 2\delta$, which is outside the pseudoeffective cone, whose extremal rays are generated by δ and $2h - 3\delta$ by Theorem 5.1, obtaining a contradiction. If $t \neq 2$, there exists a (-4) -class if and only if the Pell-type equation $P_t(2)$ is solvable: since by assumption $P_t(-1)$ has solutions, by [Per13, p.106-109], see also [Yok94, Proposition 1], we have that $P_t(2)$ has no solution, hence there are no (-4) -classes. We conclude that $D_{1,1}$ must be a (-2) -class, hence it is either δ or $\iota^*\delta$, where ι is the anti-symplectic involution on X which generates $\text{Aut}(X)$, see Theorem 5.2. Since $\iota^*D = D$, it is enough to show that $D_{1,1} = \delta$ is not possible. If $D_{1,1} = \delta$, from (7.1) we get $D = n_1\delta + D_2$, where $n_1 \geq 1$ since D_1 is effective by assumption, hence

$$D_2 = D - n_1\delta = bh - (a + n_1)\delta,$$

where as usual (a, b) is the minimal solution of the negative Pell equation $P_t(-1)$. We show that D_2 is outside the pseudoeffective cone. By Theorem 5.1 the extremal rays of the pseudoeffective cone are generated by the classes δ and $\iota^*\delta = dh - c\delta$, where (c, d) is the minimal solution of the Pell equation $P_t(1)$. Then $\iota^*\delta = 2abh - (a^2 + tb^2)\delta$ by (2.1), hence we need to check that $a + n_1 > \frac{a^2 + tb^2}{2a}$ in order to show that D_2 is outside the pseudoeffective cone. This is true since $a^2 - tb^2 = -1$. We obtain a contradiction, so $q_X(D_i) \geq 0$ for $i = 1, 2$. Moreover, we have already remarked that $q_X(D_i) \neq 0$ for $i = 1, 2$, thus $q_X(D_i) > 0$. We get a contradiction with (7.2), so one between D_1 and D_2 is zero. If $D_2 = 0$, then $n_1 = 1$ since $q_X(D) = n_1^2 q_X(D_{1,1}) = 2$, so D is reduced and irreducible. If $D_1 = 0$, we repeat the argument for $D = D_2$ until we obtain a D which is reduced and irreducible. \square

Suppose now that $D_1, D_2 \in |D|$ are two distinct divisors. We want to study the surface $D_1 \cap D_2$ and see if it is reduced and irreducible. We first show what happens when $t = 2$, i.e., when $X = S_4^{[2]}$ is the Hilbert square of a smooth quartic surface of \mathbb{P}^3 with Picard rank 1, a case already studied in [Wel81] and [Bea83a]. We briefly recall some notation from [GH78, §1.5, §6.2] on Schubert varieties of $\mathbb{G}(1, \mathbb{P}^3)$. Fix a complete flag in \mathbb{P}^3 , i.e., let v_0 be a point in \mathbb{P}^3 , let L_0 be a line in \mathbb{P}^3 and let H_0 be a plane in \mathbb{P}^3 such that $v_0 \in L_0 \subset H_0$. Consider the following two Schubert varieties of dimension 2 of the Grassmannian $\mathbb{G}(1, \mathbb{P}^3)$,

$$\Sigma_{1,1} := \{L \in \mathbb{G}(1, \mathbb{P}^3) \mid L \subset H_0\}, \quad \Sigma_2 := \{L \in \mathbb{G}(1, \mathbb{P}^3) \mid v_0 \in L\}, \quad (7.3)$$

and denote by $\sigma_{1,1}$ and σ_2 the corresponding classes in $H^4(\mathbb{G}(1, \mathbb{P}^3), \mathbb{Z})$. Recall that $\mathbb{G}(1, \mathbb{P}^3)$ can be embedded in \mathbb{P}^5 as a quadric. Then $\sigma_{1,1} + \sigma_2 = \sigma_1^2$, where σ_1 is the hyperplane class of \mathbb{P}^5 restricted to $\mathbb{G}(1, \mathbb{P}^3)$, moreover $\int_{\mathbb{G}(1, \mathbb{P}^3)} \sigma_{1,1} \cdot \sigma_2 = 0$ and $\int_{\mathbb{G}(1, \mathbb{P}^3)} \sigma_{1,1}^2 = \int_{\mathbb{G}(1, \mathbb{P}^3)} \sigma_2^2 = 1$. Recall that S_4 does not contain any line, since it has Picard rank 1. Let $f := \varphi|_{D|}$ be the map seen in Section 5, i.e.,

$$f : S_4^{[2]} \rightarrow \mathbb{G}(1, \mathbb{P}^3) \subset \mathbb{P}^5, \quad x \mapsto l_x, \quad (7.4)$$

where l_x is the unique line in \mathbb{P}^3 which contains the support of the subscheme x . Moreover, the class of D is $h - \delta$, where as usual $h \in \text{Pic}(S_4^{[2]})$ is the class induced by the ample generator of $\text{Pic}(S_4)$. Since f is the map induced by the complete linear system $|D|$, we have $f^*\sigma_1 = c_1(\mathcal{O}_X(D))$, hence $f^*(\sigma_{1,1} + \sigma_2) = c_1(\mathcal{O}_X(D))^2$. This shows that, given two distinct divisors $D_1, D_2 \in |D|$, the surface $D_1 \cap D_2$ can be reducible, and if so, we have $[D_1 \cap D_2] = A + B \in H^{2,2}(X, \mathbb{Z})$, where $A, B \in H^{2,2}(X, \mathbb{Z})$ are the effective classes of the two irreducible components of $D_1 \cap D_2$. Moreover, $A = f^*\sigma_{1,1}$ and $B = f^*\sigma_2$. If S_4 is generic it is possible to compute the classes of $f^*\sigma_{1,1}$ and $f^*\sigma_2$, obtaining the following equalities:

$$f^*\sigma_{1,1} = \frac{1}{2}h^2 - \frac{1}{4}\delta^2 - \frac{1}{2}h\delta - \frac{1}{10}q_X^\vee, \quad f^*\sigma_2 = \frac{1}{2}h^2 + \frac{5}{4}\delta^2 - \frac{3}{2}h\delta + \frac{1}{10}q_X^\vee. \quad (7.5)$$

See [Nov, Proposition 4.4.4] for details. It is also possible to geometrically describe the surface $F \subset S_4^{[2]}$ of points fixed by the Beauville involution. We call *bitangent* of $S_4 \subset \mathbb{P}^3$ a line of \mathbb{P}^3 which intersects S_4 either in two points both with multiplicity 2 or in a point with multiplicity 4. Let $\text{Bit}(S_4) \subset \mathbb{G}(1, \mathbb{P}^3)$ be the surface of bitangents of $S_4 \subset \mathbb{P}^3$. Then we have $f|_F : F \xrightarrow{\sim} \text{Bit}(S_4)$. If S_4 is generic, starting from $[\text{Bit}(S_4)] = 12\sigma_2 + 28\sigma_{1,1} \in H^{2,2}(\mathbb{G}(1, \mathbb{P}^3), \mathbb{Z})$, see for instance [ABT01, Proposition 3.3], it is possible to show that $f^*[\text{Bit}(S_4)] = 20h^2 + 8\delta^2 - 32h\delta - \frac{8}{5}q_X^\vee \in H^{2,2}(S_4^{[2]}, \mathbb{Z})$, see [Nov, Proposition 4.4.4].

We come back to the general problem of the Hilbert square X of a generic K3 surface admitting an ample divisor $D \in \text{Div}(X)$ with $q_X(D) = 2$. Before stating the irreducibility property, we show the following technical lemma.

Lemma 7.2. *Let X and D be as in Theorem 6.4, and let $D_1, D_2 \in |D|$ be two distinct divisors. Denote by ι the anti-symplectic involution which generates the automorphism group $\text{Aut}(X)$. Suppose that there is no decomposition of the form*

$$[D_1 \cap D_2] = A + B \in H^{2,2}(X, \mathbb{Z}),$$

where $[D_1 \cap D_2]$ is the fundamental cohomological class of the surface $D_1 \cap D_2$, and $A, B \in H^{2,2}(X, \mathbb{Z})$ are effective classes such that $\iota^(A) = A$ and $\iota^*(B) = B$ with ι^* described in (6.10). Then the surface $D_1 \cap D_2$ is reduced and irreducible.*

Proof. Suppose that

$$[D_1 \cap D_2] = A_1 + A_2 + \cdots + A_n \in H^{2,2}(X, \mathbb{Z}),$$

where $A_i \in H^{2,2}(X, \mathbb{Z})$ are effective classes, not necessarily pairwise distinct. By Theorem 5.2 we have $\iota^*[D] = [D]$, hence $\iota^*[D_1 \cap D_2] = [D_1 \cap D_2]$. If $n > 1$ is odd, then there exists i such that $\iota^*A_i = A_i$ since ι is an involution, hence we take $A := A_i$ and $B := A_1 + \cdots + A_{i-1} + A_{i+1} + \cdots + A_n$. Thus $\iota^*A = A$ and $\iota^*B = B$. We obtain a contradiction

with the assumption that there is no decomposition of this form. Suppose now that $n = 2$ and $\iota^* A_1 = A_2$. We show that this is not possible. Recall that the class of D in $\text{Pic}(X)$ is $bh - a\delta$, with (a, b) minimal solution of the Pell-type equation $P_t(-1)$. By Theorem 4.5 the classes A_1 and A_2 in $H^{2,2}(X, \mathbb{Z})$ are of the form

$$\begin{aligned} A_1 &= \left(x + \frac{y}{2}\right) h^2 + \frac{z}{8} \delta^2 - \frac{y}{2} h\delta + \left(\frac{1}{8}z + w\right) \frac{2}{5} q_X^\vee, \\ A_2 &= \left(b^2 - x - \frac{y}{2}\right) h^2 + \left(a^2 - \frac{z}{8}\right) \delta^2 + \left(-2ab + \frac{y}{2}\right) h\delta - \left(\frac{1}{8}z + w\right) \frac{2}{5} q_X^\vee, \end{aligned}$$

for some $x, y, z, w \in \mathbb{Z}$. By Theorem 5.2 and (6.10), we obtain

$$\begin{aligned} \iota^* A_1 &= \left(c^2 \left(x + \frac{y}{2}\right) + cd \left(-\frac{y}{2}\right) + \frac{z}{8} d^2\right) h^2 \\ &\quad + \left(t^2 d^2 \left(x + \frac{y}{2}\right) - cdt \frac{y}{2} + c^2 \frac{z}{8}\right) \delta^2 \\ &\quad + \left(-2cdt \left(x + \frac{y}{2}\right) + c^2 \frac{y}{2} + td^2 \frac{y}{2} - 2cd \frac{z}{8}\right) h\delta \\ &\quad + \left(\frac{1}{8}z + w\right) \frac{2}{5} q_X^\vee, \end{aligned}$$

and similarly

$$\begin{aligned} \iota^* A_2 &= \left(c^2 \left(b^2 - x - \frac{y}{2}\right) + cd \left(\frac{y}{2} - 2ab\right) + d^2 \left(a^2 - \frac{z}{8}\right)\right) h^2 \\ &\quad + \left(t^2 d^2 \left(b^2 - x - \frac{y}{2}\right) + \left(\frac{y}{2} - 2ab\right) cdt + c^2 \left(a^2 - \frac{z}{8}\right)\right) \delta^2 \\ &\quad + \left(-2cdt \left(b^2 - x - \frac{y}{2}\right) + (c^2 + td^2) \left(2ab - \frac{y}{2}\right) - 2cd \left(a^2 - \frac{z}{8}\right)\right) h\delta \\ &\quad - \left(\frac{1}{8}z + w\right) \frac{2}{5} q_X^\vee. \end{aligned}$$

Imposing $\iota^* A_1 = A_2$, we obtain a system whose solution is

$$\begin{cases} x = \frac{1}{2}b^2 - ab, \\ y = 2ab, \\ z = 4a^2, \\ w = -\frac{a^2}{2}. \end{cases}$$

Recall that b is odd by Proposition 2.6, so $x \notin \mathbb{Z}$, which is not possible. Since by assumption we cannot have $\iota^* A_1 = A_1$, one between A_1 and A_2 is zero, so $D_1 \cap D_2$ is reduced and irreducible. If $n > 1$ is even, if there exists an i such that $\iota^* A_i = A_i$, we proceed as in the case of n odd, otherwise without loss of generality we can assume that $\iota^* A_1 = A_2$. Then, taking $A := A_1 + A_2$ and $B := A_3 + \dots + A_n$, we have $\iota^* A = A$ and $\iota^* B = B$, which contradicts the assumption. We conclude that $D_1 \cap D_2$ is reduced and irreducible. \square

We can now state the main theorem of this section, the *irreducibility property*.

Theorem 7.3. *Let X be the Hilbert square of a projective K3 surface S_{2t} with $\text{Pic}(S_{2t}) = \mathbb{Z}H$ and $H^2 = 2t$ and such that X admits an ample divisor $D \in \text{Div}(X)$ with $q_X(D) = 2$. Let $D_1, D_2 \in |D|$ be two distinct divisors.*

- (i) *If $t = 2$, then the surface $D_1 \cap D_2$ can be reducible. If $[D_1 \cap D_2] = A + B$, where $[D_1 \cap D_2]$ is the fundamental cohomological class of $D_1 \cap D_2$ in $H^{2,2}(X, \mathbb{Z})$, then $A = f^* \sigma_{1,1}$ and $B = f^* \sigma_2$, where $f : X \rightarrow \mathbb{G}(1, \mathbb{P}^3)$ is the map given in (7.4) and $\sigma_{1,1}, \sigma_2$ are the classes in $H^4(\mathbb{G}(1, \mathbb{P}^3), \mathbb{Z})$ of the Schubert varieties in (7.3). Moreover, if S_4 is generic, then the effective classes A and B in $H^{2,2}(X, \mathbb{Z})$ are*

$$A = \frac{1}{2}h^2 - \frac{1}{4}\delta^2 - \frac{1}{2}h\delta - \frac{1}{10}q_X^\vee, \quad B = \frac{1}{2}h^2 + \frac{5}{4}\delta^2 - \frac{3}{2}h\delta + \frac{1}{10}q_X^\vee.$$

- (ii) *If $t \neq 2$ and S_{2t} is generic, then $D_1 \cap D_2$ is a reduced and irreducible surface.*

Proof. We begin with $t = 2$, so $X = S_4^{[2]}$ is the Hilbert square of a smooth quartic surface of \mathbb{P}^3 with Picard rank 1 and the class of D is $h - \delta \in \text{Pic}(X)$, where h is the class induced by the ample generator of $\text{Pic}(S_4)$. For the reader's convenience, we work under the assumption that S_4 is generic: the procedure that we will develop in this way is exactly the same that we will use for Item (ii), where S_{2t} is generic by assumption, but easier to follow for the case $t = 2$. By Lemma 7.2, the surface $D_1 \cap D_2$ can be reducible only if there exist effective classes $A, B \in H^{2,2}(X, \mathbb{Z})$ such that $[D_1 \cap D_2] = A + B$ and $\iota^*(A) = A, \iota^*(B) = B$, where ι is the Beauville involution. Since $(h - \delta)^2 = h^2 + \delta^2 - 2h\delta$, by Theorem 4.5 we can write, for some $x, y, z, w \in \mathbb{Z}$,

$$\begin{aligned} A &= \left(x + \frac{y}{2}\right) h^2 + \frac{z}{8} \delta^2 - \frac{y}{2} h\delta + \left(\frac{1}{8}z + w\right) \frac{2}{5} q_X^\vee \in H^{2,2}(X, \mathbb{Z}), \\ B &= \left(1 - x - \frac{y}{2}\right) h^2 + \left(1 - \frac{z}{8}\right) \delta^2 + \left(\frac{y}{2} - 2\right) h\delta - \left(\frac{1}{8}z + w\right) \frac{2}{5} q_X^\vee \in H^{2,2}(X, \mathbb{Z}). \end{aligned}$$

- By assumption A and B are effective. Moreover, $h \in \text{Pic}(X)$ is nef by Theorem 5.1. By Kleiman's theorem we have $\langle A, h^2 \rangle \geq 0$ and $\langle B, h^2 \rangle \geq 0$, where $\langle \cdot, \cdot \rangle$ is the bilinear form of $H^4(X, \mathbb{Z})$ given in Proposition 4.2, which coincides with the intersection pairing. We obtain the following condition:

$$0 \leq 12x + 6y + z + 10w \leq 10. \quad (7.6)$$

- Let $\sigma \in H^0(X, \Omega_X^2)$ be the symplectic form. Then, since A and B are effective classes in $H^{2,2}(X, \mathbb{Z})$ by assumption, we have

$$\int_A (\sigma + \bar{\sigma})^2 = 2 \int_A \sigma \wedge \bar{\sigma} \geq 0, \quad \int_B (\sigma + \bar{\sigma})^2 = 2 \int_B \sigma \wedge \bar{\sigma} \geq 0,$$

since $\sigma \wedge \bar{\sigma} \in H^{2,2}(X)$ is a volume form on A and on B . Note that $\sigma \wedge \bar{\sigma}$ can be zero on A or B , for instance when A or B are Lagrangian. Hence $\langle A, (\sigma + \bar{\sigma})^2 \rangle \geq 0$ and $\langle B, (\sigma + \bar{\sigma})^2 \rangle \geq 0$. Using (6.7) we obtain the following condition:

$$0 \leq 4x + 2y + z + 10w \leq 2. \quad (7.7)$$

- By abuse of notation we write D also for its class in $\text{Pic}(X)$. Since D is ample and by assumption A and B are effective, by the Nakai–Moishezon criterion we have $\langle A, D^2 \rangle > 0$ and $\langle B, D^2 \rangle > 0$, and we obtain the following condition:

$$0 < 40x + 12y + 3z + 20w < 12. \quad (7.8)$$

- By Theorem 5.2 and (6.10) we have

$$\iota^* A = \left(9x + \frac{3}{2}y + \frac{1}{2}z\right) h^2 + \left(16x + 2y + \frac{9}{8}\right) \delta^2 - \left(24x + \frac{7}{2}y + \frac{3}{2}z\right) h\delta + \left(\frac{1}{8}z + w\right) \frac{2}{5}q_X^\vee.$$

Since $\iota^* A = A$, we obtain the system

$$\begin{cases} 9x + \frac{3}{2}y + \frac{1}{2}z = x + \frac{y}{2}, \\ 16x + 2y + \frac{9}{8}z = \frac{z}{8}, \\ 24x + \frac{7}{2}y + \frac{3}{2}z = \frac{y}{2}, \end{cases}$$

which gives the following condition:

$$16x + 2y + z = 0. \quad (7.9)$$

We look for $x, y, z, w \in \mathbb{Z}$ which satisfy (7.6), (7.7), (7.8) and (7.9). Since $-2 \leq 8x + 4y \leq 10$ by (7.6) and (7.7), and $x, y \in \mathbb{Z}$, we have

$$2x + y \in \{0, 1, 2\}.$$

- Suppose that $2x + y = 0$. By (7.9) we have $z = -12x$, and (7.8) becomes $0 < w - x < \frac{3}{5}$. Since $w - x \in \mathbb{Z}$, this condition is never satisfied.
- Suppose that $2x + y = 1$. By (7.9) we have $z = -12x - 2$, and (7.8) becomes $-\frac{3}{10} < w - x < \frac{3}{10}$. Since $w - x \in \mathbb{Z}$, we get $x = w$, and (7.7) gives $x \in \{-1, 0\}$. We obtain the following two solutions:

$$\begin{cases} x = 0, \\ y = 1, \\ z = -2, \\ w = 0, \end{cases} \quad \text{and} \quad \begin{cases} x = -1, \\ y = 3, \\ z = 10, \\ w = -1. \end{cases}$$

which coincide with the effective classes $f^*\sigma_{1,1}$ and $f^*\sigma_2$ respectively, see (7.5) and [Nov, Proposition 4.4.4]. Moreover, with the same technique it is possible to show that $f^*\sigma_{1,1}$ and $f^*\sigma_2$ are reduced and irreducible.

- The case $2x + y = 2$ is symmetric to the case $2x + y = 0$, i.e., if A is a class obtained in this case, then A coincides with a class B obtained in the case $2x + y = 0$. Since there are no classes in the case $2x + y = 0$, there are no classes in the cases $2x + y = 2$.

We conclude that if $D_1 \cap D_2$ is a reducible surface, then it is reduced with two irreducible components whose classes in $H^{2,2}(X, \mathbb{Z})$ are $f^*\sigma_{1,1}$ and $f^*\sigma_2$.

Suppose now that $t \neq 2$ and S_{2t} is generic. We want to show that the surface $D_1 \cap D_2$ is reduced and irreducible. By Lemma 7.2 it is enough to show that we cannot have $[D_1 \cap D_2] = A + B$ for effective $A, B \in H^{2,2}(X, \mathbb{Z})$ such that $\iota^* A = A$ and $\iota^* B = B$. The technique is the same seen before. By abuse of notation we write D also for its class in $\text{Pic}(X)$. Recall that $D = bh - a\delta$, where (a, b) is the minimal solution of the Pell-type equation $P_t(-1)$. We have already remarked that the first value of t different from 2 which satisfies our assumptions is $t = 10$, and in this

case $a = 3$, hence for other values of t that we consider we have $a > 3$. Thus we can suppose that $t \geq 10$ and $a \geq 3$. Since S_{2t} is generic by assumption, by Theorem 4.5 we can write

$$\begin{aligned} A &= \left(x + \frac{y}{2}\right) h^2 + \frac{z}{8} \delta^2 - \frac{y}{2} h \delta + \left(\frac{1}{8} z + w\right) \frac{2}{5} q_X^\vee, \\ B &= \left(b^2 - x - \frac{y}{2}\right) h^2 + \left(a^2 - \frac{z}{8}\right) \delta^2 + (-2ab + \frac{y}{2}) h \delta - \left(\frac{1}{8} z + w\right) \frac{2}{5} q_X^\vee, \end{aligned}$$

for some $x, y, z, w \in \mathbb{Z}$. Proceeding as for the case $t = 2$ we obtain the following conditions:

$$0 \leq 6tx + 3ty + z + 10w \leq 6tb^2 - 2a^2, \quad (7.10)$$

$$0 \leq 2tx + ty + z + 10w \leq 2, \quad (7.11)$$

$$0 < (4t + 8t^2b^2)x + (2t + 4t^2b^2 - 4abt)y + (1 + tb^2)z + 20w < 12, \quad (7.12)$$

$$8tdx + 4(td - c)y + dz = 0. \quad (7.13)$$

where condition (7.10) is given by $\langle A, h^2 \rangle \geq 0$ and $\langle B, h^2 \rangle \geq 0$, condition (7.11) is given by $\langle \sigma + \bar{\sigma} \rangle^2 \geq 0$ and $\langle B, (\sigma + \bar{\sigma})^2 \rangle \geq 0$, condition (7.12) is given by $\langle A, D^2 \rangle > 0$ and $\langle B, D^2 \rangle > 0$ and condition (7.13) is given by $\iota^* A = A$. Note that (7.10) and (7.11) implies $-\frac{1}{t} \leq 2x + y \leq 2b^2 + \frac{1}{t}$. Since $2x + y \in \mathbb{Z}$ and $t \geq 10$ we have

$$2x + y \in \{0, 1, \dots, 2b^2\}.$$

Note that, similarly to the case $t = 2$ seen above, a class A obtained by imposing $2x + y \in \{b^2 + 1, \dots, 2b^2\}$ coincide with a class B obtained for $2x + y \in \{0, 1, \dots, b^2\}$. Hence it suffices to study $2x + y \in \{0, 1, \dots, b^2\}$. Suppose that $2x + y = k$, where $k \in \{0, 1, \dots, b^2\}$. By (7.11) we have $-tk \leq z + 10w \leq 2 - tk$, and since $z + 10w \in \mathbb{Z}$ we have

$$z + 10w \in \{-tk, -tk + 1, -tk + 2\}.$$

Suppose that $z + 10w = -tk$. Then (7.12) gives, after some computations, $-4t^2b^2k < -4abty + a^2z < -4t^2b^2k + 12$. Since $-4abty + a^2z \in \mathbb{Z}$, we have $-4abty + a^2z = -4t^2b^2k + h$, where $h \in \{1, 2, \dots, 11\}$. With the help of a computer we obtain

$$w = \frac{-5a^2tk + 2ha^2 - 4tk + h}{10a^2}.$$

Then w is an integer only if $4tk - h \equiv 0 \pmod{a^2}$. If $k = 0$, then $-h \equiv_{a^2} 0$ only if $h = 9$ and $a = 3$, being $a \geq 3$. This happens only when $t = 10$, and we get $w = \frac{19}{10}$, which is not an integer. Suppose now that $k \neq 0$. Since $k \leq b^2$, we have $4tk - h \leq 4a^2 + 4 - h$, hence in order to get $4tk - h \equiv_{a^2} 0$ we must have $4tk - h \in \{a^2, 2a^2, 3a^2, 4a^2\}$. If $4tk - h = a^2$, then $k = \frac{tb^2 + h - 1}{4t}$. If $h \neq 1, 11$, then $h - 1$ is not divisible by $t \geq 10$, and k is not an integer. If $h = 1$, then $k = \frac{b^2}{4}$, which is not an integer by Proposition 2.6. If $h = 11$, then $h - 1 = 10$ is divisible by $t \geq 10$ if and only if $t = 10$, which implies $b = 1$: thus $k = \frac{1}{2}$, which is not an integer.

In a similar way it is possible to show that all the other remaining cases are not possible: we omit the details, which can be found in [Nov, Appendix B]. We conclude that there are no effective classes $A, B \in H^{2,2}(X, \mathbb{Z})$ such that $[D_1 \cap D_2] = A + B$, hence $D_1 \cap D_2$ is a reduced and irreducible surface. \square

8. GEOMETRIC DESCRIPTION

Let X be the Hilbert square of a generic K3 surface of degree $2t$. Suppose that X admits an ample divisor D with $q_X(D) = 2$ and $t \neq 2$. In this section, using Theorem 6.4 and Theorem 7.3, we show that $\varphi_{|D|} : X \rightarrow Y \subset \mathbb{P}^5$ is a double cover of an EPW sextic Y , hence X is a double EPW sextic. Our idea is to follow the strategy developed by O'Grady in [O'G08] with remarkable simplifications obtained thanks to the existence of the anti-symplectic involution given by Theorem 5.2. We will omit proofs which are identical to the ones in [O'G08].

Let X and D be as above. We introduce the following notation from [O'G08, §4]. We choose an isomorphism $|D|^\vee \xrightarrow{\sim} \mathbb{P}^5$ and we denote by $f : X \dashrightarrow \mathbb{P}^5$ the composition $X \dashrightarrow |D|^\vee \xrightarrow{\sim} \mathbb{P}^5$: this is basically the rational map $\varphi_{|D|}$. Let B be the base locus of $|D|$, and $\beta_B : \tilde{X} \rightarrow X$ be the blow-up of X in B . We denote by $\tilde{f} : \tilde{X} \rightarrow \mathbb{P}^5$ the regular map which resolves the indeterminacies of f . Let $Y := \text{Im}(\tilde{f})$, which is a closed subset of \mathbb{P}^5 . We obtain a dominant map, which we call f by abuse of notation:

$$f : X \dashrightarrow Y.$$

Note that $\deg(f)$ is even by Theorem 6.4. Let Y_0 be the interior of $\tilde{f}(X \setminus B)$, thus $Y_0 \subseteq Y$ is open and dense. Let $X_0 := (X \setminus B) \cap \tilde{f}^{-1}(Y_0)$, so $X_0 \subseteq X$ is open and dense. We call f_0 the restriction of \tilde{f} to X_0 , which is a regular surjective map:

$$f_0 : X_0 \rightarrow Y_0.$$

We now summarize the main steps which will lead us to the main theorem of this section:

- (i) $\dim(Y) = 4$.
- (ii) $\deg(f) \cdot \deg(Y) \leq 12$ and the equality holds if and only if $\text{Bs}|D| = \emptyset$.
- (iii) Either $\deg(f) = 2$ and $\deg(Y) = 6$ or $\deg(f) = 4$ and $\deg(Y) = 3$. In particular $\text{Bs}|D| = \emptyset$ and f is a morphism.
- (iv) The case $\deg(f) = 4$ and $\deg(Y) = 3$ never holds.
- (v) The variety X is a double EPW sextic.

The following corollary of Theorem 7.3 will be fundamental to prove the other results of this section.

Corollary 8.1. *Let X and D be as in Theorem 7.3.(ii), and keep notation as above.*

- (i) *If $L \subset \mathbb{P}^5$ is a linear subset of codimension at most 2, then $L \cap Y_0$ is reduced and irreducible and, if non empty, it has pure codimension equal to $\text{cod}(L, \mathbb{P}^5)$.*
- (ii) *The base locus $B = \text{Bs}|D|$ has dimension at most 1. Let B_{red} be the reduced scheme associated to B and B_{red}^1 be the union of the irreducible components of B_{red} of dimension 1. If $D_1, D_2, D_3 \in |D|$ are linearly independent, then $D_1 \cap D_2 \cap D_3$ has pure dimension 1 and there exists a unique decomposition*

$$[D_1 \cap D_2 \cap D_3] = \Gamma + \Sigma,$$

where Γ, Σ are (classes of) effective 1-cycles such that

- $\text{Supp}(\Gamma) \cap B_{\text{red}}$ is either 0-dimensional or empty.
- $\text{Supp}(\Sigma) = B_{\text{red}}^1$.

We only prove Item (i), even if it is almost the same as [O'G08, Corollary 4.2.(i)], since in our setting it is a direct consequence of Proposition 7.1 and of Theorem 7.3.(ii).

Proof. If $L \cong \mathbb{P}^5$, there is nothing to prove. If $\text{cod}(L, \mathbb{P}^5) = 1$, let $D_1 \in |D|$ be the divisor which corresponds to L in the isomorphism $|D|^\vee \cong \mathbb{P}^5$. Then $[D_1 \cap X_0] = f_0^*[L]$, where $[D_1 \cap X_0]$ and $[L]$ are the fundamental cohomological classes of $D_1 \cap X_0$ and L respectively. Since X_0 is open and dense in X and f_0 is surjective, Proposition 7.1 implies that $L \cap Y_0$ is reduced and irreducible of pure codimension 1 if non-empty. If $\text{cod}(L, \mathbb{P}^5) = 2$, then $L = L_1 \cap L_2$ with $L_1, L_2 \subset \mathbb{P}^5$ hyperplanes, hence $[D_1 \cap D_2 \cap X_0] = f_0^*[L]$, where $D_1, D_2 \in |D|$ corresponds to L_1 and L_2 respectively. Hence $L \cap Y_0$ is reduced and irreducible of pure codimension 2 if non-empty by Theorem 7.3.(ii). For Item (ii) see [O'G08, Corollary 4.2.(ii)] (one needs Theorem 7.3.(ii) once again). \square

Note that $[D_1 \cap D_2 \cap D_3]$ in the statement of Corollary 8.1.(ii) denotes the fundamental cycle of $D_1 \cap D_2 \cap D_3$, see [Ful13, §1.5]. The following proposition corresponds to [O'G08, Corollary 4.3].

Proposition 8.2. *Let X and D be as in Theorem 7.3.(ii). Keep notation as above. Then $\dim(Y) \in \{3, 4\}$.*

We want to prove that the case $\dim(Y) = 3$ never holds. First of all, similarly to [O'G08, Proposition 4.5], we give boundaries to $\deg(Y)$. We show the proof, which has little differences with the one by O'Grady: the anti-symplectic involution ι of Theorem 5.2 and Theorem 6.4 are strongly used.

Proposition 8.3. *Keep notation as above and suppose that $\dim(Y) = 3$. Then $3 \leq \deg(Y) \leq 6$. Moreover, if $\deg(Y) = 6$, then the base locus $\text{Bs}|D|$ is 0-dimensional.*

Proof. Let $d_Y := \deg(Y)$. Since $Y \subset \mathbb{P}^5$ is a non-degenerate subvariety, by [EH16, Proposition 0] we have $d_Y \geq 3$. Let now $L_1, L_2, L_3 \subset \mathbb{P}^5$ be three generic hyperplanes. Then the intersection $Y \cap L_1 \cap L_2 \cap L_3$ is transverse and it is given by d_Y points, which we call p_1, \dots, p_{d_Y} . Let $\Gamma_{0,i} := f_0^{-1}(p_i)$ for $i = 1, \dots, d_Y$, and let Γ_i be the closure of $\Gamma_{0,i}$ in X . Let $D_1, D_2, D_3 \in |D|$ be the divisors which correspond to L_1, L_2, L_3 in the isomorphism $|D|^\vee \cong \mathbb{P}^5$. Since L_1, L_2, L_3 are generic, D_1, D_2, D_3 are linearly independent, and by Corollary 8.1.(ii) we have

$$[D_1 \cap D_2 \cap D_3] = \sum_{i=1}^{d_Y} \Gamma_i + \Sigma,$$

where the equality is in $H^{3,3}(X, \mathbb{Z})$ and we write by abuse of notation Γ_i and Σ for their classes in $H^{3,3}(X, \mathbb{Z})$. We write again D also for its class in $\text{Pic}(X)$: recall that $D = bh - a\delta$ with (a, b) minimal solution of the Pell-type equation $P_t(-1)$, and b is odd by Proposition 2.6. Using Proposition 3.6 one obtains $D^\vee = bh^\vee - 2a\delta^\vee$ with $h^\vee = h$ and $\delta^\vee = \frac{\delta}{2}$ in $H^2(X, \mathbb{Q})$. Moreover, as already remarked, the divisibility of D in $H^2(X, \mathbb{Z})$ is $\text{div}(D) = 1$, hence by Proposition 3.6 we have

$$D \cdot D^\vee = \left(bh - a\delta, bh - 2\frac{a}{2}\delta \right) = 2.$$

Since $D_1, D_2, D_3 \in |D|$ and $\iota^*D \cong D$, we have $\iota^*[D_1 \cap D_2 \cap D_3] = [D_1 \cap D_2 \cap D_3]$, hence

$$\sum_{i=1}^{d_Y} \iota^* \Gamma_i + \iota^* \Sigma = \sum_{i=1}^{d_Y} \Gamma_i + \Sigma.$$

By Theorem 5.2 and Theorem 5.4 the only class fixed by the action induced by ι on $H^{3,3}(X, \mathbb{Z})$ is D^\vee . Moreover, from Theorem 6.4 we know that f factors through the quotient by ι , so $\iota^* \Gamma_i \cong \Gamma_i$, since $\Gamma_i = f_0^{-1}(p_i)$. Then the class of Γ_i is either some positive multiple of D^\vee , or it is of the form $\Gamma_i = \Gamma'_i + \iota^* \Gamma'_i$, where Γ'_i is an effective class. Since $q_X(D) = 2$, which gives $\int_X c_1(\mathcal{O}_X(D))^4 = 12$ by Theorem 3.5, we have

$$12 = \langle D, \sum_{i=1}^{d_Y} \Gamma_i + \Sigma \rangle.$$

Note that $\langle D, \Gamma_i \rangle$ is either $\langle D, \alpha D^\vee \rangle = 2\alpha$, where $\alpha \in \mathbb{Z}_{\geq 1}$, or $\langle D, \Gamma'_i + \iota^* \Gamma'_i \rangle \geq 2$, being D ample. Hence we have $12 \geq \langle D, \Sigma \rangle + 2d_Y$, i.e., $d_Y \leq 6$. Moreover, if $d_Y = 6$, then $\Sigma = 0$, otherwise $\langle D, \Sigma \rangle > 0$ by the Nakai–Moishezon criterion, thus the base locus $\text{Bs}|D|$ is 0-dimensional, since the 1-dimensional component of $\text{Bs}|D|$ is contained in Σ by Corollary 8.1.(ii). \square

Using exactly the same techniques in [O'G08], with important simplifications due to Theorem 6.4, we obtain the following proposition.

Proposition 8.4. *Let X and D be as in Theorem 7.3.(ii). Consider the map $f : X \dashrightarrow Y \subset \mathbb{P}^5$ induced by the complete linear system $|D|$. Then $\dim(Y) = 4$. Moreover, $|D|$ is basepoint free, i.e., f is a morphism, and one of the following holds.*

- (i) $\deg(f) = 2$ and $\deg(Y) = 6$.
- (ii) $\deg(f) = 4$ and $\deg(Y) = 3$.

Proof. After having obtained Proposition 8.3, following [O'G08, §5] one shows that $\dim(Y) = 4$. Then, one obtains results which correspond to [O'G08, Proposition 4.6, Corollary 4.7], and proceeding as in [O'G08, §4] we have the following possibilities:

- (1) $\deg(Y) = 2$.
- (2) $\deg(Y) = 3$, $\deg(f) = 3$, $\text{Bs}|D| \neq \emptyset$.
- (3) $\deg(Y) = 3$, $\deg(f) = 4$, $\text{Bs}|D| = \emptyset$.
- (4) $\deg(Y) = 4$, $\deg(f) = 3$, $\text{Bs}|D| = \emptyset$.
- (5) $\deg(Y) = 6$, $\deg(f) = 2$, $\text{Bs}|D| = \emptyset$.
- (6) $\deg(f) = 1$.

As already remarked, $\deg(f)$ is even by Theorem 6.4, hence (2), (4) and (6) are not possible. Case (1) does not hold, otherwise one can find a linear subset $L \subset \mathbb{P}^5$ of dimension 3 such that $L \cap Y_0$ is reducible, which contradicts Corollary 8.1.(i), see [O'G08, §5.2] for details. \square

Note that in [O'G08] much longer discussions are needed to show that (2) and (4) in the proof of Proposition 8.4 never hold: the existence of the anti-symplectic involution and Theorem 6.4 simplifies the situation a lot. Moreover, recall that Corollary 8.1.(i) is a consequence of the irreducibility property of Theorem 7.3.(ii). When $t = 2$, the irreducibility property does not hold, see Theorem 7.3.(i), hence Corollary 8.1.(i) is not true and case (1) in the proof of Proposition 8.4 is possible, actually it holds: as we have already seen in Section 7, in this case $\deg(f) = 6$ and $\deg(Y) = 2$, in particular Y is the Grassmannian $\mathbb{G}(1, \mathbb{P}^3)$ of lines in \mathbb{P}^3 .

The final step is to show that Item (ii) of Proposition 8.4 is impossible and in Item (i) the variety $Y \subset \mathbb{P}^5$ is an EPW sextic, X is a double EPW sextic and f is the double cover associated. See [EPW01] for the definition of EPW sextic and [O'G06] for details on double EPW sextics. We remark that we cannot proceed following [O'G08, §5.5]: in that case, the fact that the variety is a *deformation* of the Hilbert square of a K3 surface plays a central role in the proof. If M is an IHS manifold of $K3^{[2]}$ -type and $h \in H^{1,1}(M, \mathbb{Q})$ is an element such that $q_M(h) \neq 0$, O'Grady shows that $H^4(M, \mathbb{C})$ can be decomposed as

$$H^4(M, \mathbb{C}) = (\mathbb{C}h^2 \oplus \mathbb{C}q_M^\vee) \oplus (\mathbb{C}h \otimes h^\perp) \oplus W(h),$$

where the orthogonality is with respect to the BBF form and $W(h) := (q_M^\vee)^\perp \cap \text{Sym}^2(h^\perp)$, where $(q_M^\vee)^\perp$ is the orthogonal with respect to the intersection product. Then O'Grady takes an IHS manifold X deformation equivalent

to M such that properties (1)-(6) of [O'G08, Proposition 3.2] hold. In particular property (4) says that if $V \subset H^4(X)$ is a rational sub Hodge structure, then $V_{\mathbb{C}} = V_1 \oplus V_2 \oplus V_3$, where $V_1 \subset (\mathbb{C}h^2 \oplus \mathbb{C}q_X^{\vee})$, V_2 is either 0 or equal to $\mathbb{C}h \otimes h^{\perp}$ and V_3 is either 0 or equal to $W(h)$. In our case, there is no reason why a similar result holds, without deforming the variety $X = S_{2t}^{[2]}$ that we are studying: instead we exploit once again the anti-symplectic involution ι and Theorem 6.4. We can finally prove the main theorem of this paper.

Theorem 8.5. *Let X be the Hilbert square of a generic K3 surface S_{2t} of degree $2t$ such that X admits an ample divisor D with $q_X(D) = 2$. Suppose that $t \neq 2$, and denote by ι the anti-symplectic involution which generates $\text{Aut}(X)$. Then the complete linear system $|D|$ is basepoint free, the morphism*

$$\varphi_{|D|} : X \rightarrow Y \subset \mathbb{P}^5$$

is a double cover whose ramification locus is the surface F of points fixed by ι , and $Y \cong X/\langle \iota \rangle$ is an EPW sextic, so X is a double EPW sextic.

Proof. We show that $\deg(f) = 4$ and $\deg(Y) = 3$ in Proposition 8.4 never holds. Suppose by contradiction that $\deg(f) = 4$ and $\deg(Y) = 3$. First we show that $Y \subset \mathbb{P}^5$ is a normal variety. Since $Y \subset \mathbb{P}^5$ is a hypersurface, by [Har13, Proposition II.8.23] we have that Y is normal if and only if $\text{cod}_Y(\text{Sing}(Y)) \geq 2$. Suppose now that $\dim(\text{Sing}(Y)) = 3$. Note that Y does not contain planes, otherwise we get a contradiction with Corollary 8.1.(i). As observed by O'Grady in [O'G08, Claim 5.10], since in \mathbb{P}^5 hyperplanes and quadrics contain planes, a cubic hypersurface of \mathbb{P}^5 which is either non-reduced or reducible contains planes. Hence in our case Y is reduced and irreducible. Then, as in [O'G08, Lemma 5.17], the intersection between the variety Y and a generic plane of \mathbb{P}^5 is a singular cubic curve which is reduced and irreducible. In particular this has only one singular point, so $\text{Sing}(Y)$ has exactly one irreducible component $\Sigma \cong \mathbb{P}^3$ of degree 1. Thus $Y \supset \Sigma$, and it contains planes. This contradicts Corollary 8.1.(i). We conclude that $\text{cod}_Y(\text{Sing}(Y)) \geq 2$, hence Y is normal. The commutative diagram (6.4) gives the following:

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \subset \mathbb{P}^5 \\ & \searrow \pi' & \nearrow \bar{f} \\ & X/\langle \iota \rangle. & \end{array}$$

Since $\deg(f) = 4$ by assumption, by the commutativity of the diagram above we have $\deg(\bar{f}) = 2$. Let $\tilde{Y} := X/\langle \iota \rangle$. Since X is smooth and $\text{Aut}(X) \cong \langle \iota \rangle$ is a finite group, the quotient \tilde{Y} is a normal variety. Both \tilde{Y} and Y are normal varieties, hence by [DK19, Remark 2.4] there is a non trivial involution $\tau : \tilde{Y} \rightarrow \tilde{Y}$ such that $Y \cong \tilde{Y}/\langle \tau \rangle$. We show that τ lifts to an automorphism on X . We use a technique from [O'G13, p.179] and [DK18, Proposition B.8]. Let $F = \text{Fix}(\iota)$ be the fixed locus of the anti-symplectic involution ι . Since $\text{Sing}(\tilde{Y}) = \pi'(F)$ and τ is an automorphism on \tilde{Y} , we have $\tau(\text{Sing}(\tilde{Y})) = \text{Sing}(\tilde{Y})$. Let $Y' := \tilde{Y} \setminus \text{Sing}(\tilde{Y})$. Then the restriction of τ to Y' gives an involution $\tau|_{Y'} : Y' \rightarrow Y'$ of Y' . We set $\tau' := \tau|_{Y'}$. Since $\text{cod}_X(F) = 2$ and X is simply connected, we have that $X' := X \setminus F$ is simply connected. Thus if we restrict π' to X' we obtain the following universal cover:

$$\pi'|_{X'} : X' \rightarrow Y'.$$

By the lifting criterion τ' lifts to an automorphism on X' . Thus we obtain a birational map $\tilde{\tau} : X \dashrightarrow X$ which is not defined a priori on the fixed locus F . By [DM19, Proposition B.3] we have $\text{Bir}(X) \cong \text{Aut}(X) \cong \langle \iota \rangle$, so $\tilde{\tau}$ is either the identity or ι . This implies that the involution $\tau : \tilde{Y} \rightarrow \tilde{Y}$ is the identity, which is a contradiction. We conclude that $\deg(\bar{f})$ cannot be 2, hence we get $\deg(f) = 2$ and $\deg(Y) = 6$. It remains to show that Y is an EPW sextic and f is a double cover ramified over F , so that X is a double EPW sextic. This is true by [O'G06, Theorem 1.1], and we are done. \square

REFERENCES

- [ABT01] E. Arrondo, M. Bertolini, and C. Turrini. A focus on focal surfaces. *Asian Journal of Mathematics*, 5(3):535–560, 2001.
- [BC20] Pietro Beri and Alberto Cattaneo. On birational transformations of Hilbert schemes of points on K3 surfaces. *arXiv preprint arXiv:2005.05187*, 2020.
- [BCMS19] Samuel Boissière, Andrea Cattaneo, Dmitrii Genrikhovich Markushevich, and Alessandra Sarti. On the non-symplectic involutions of the Hilbert square of a K3 surface. *Izvestiya: Mathematics*, 83(4):731, 2019.
- [BCNWS16] Samuel Boissiere, Andrea Cattaneo, Marc Nieper-Wißkirchen, and Alessandra Sarti. The automorphism group of the Hilbert scheme of two points on a generic projective K3 surface. In *K3 surfaces and their moduli*, pages 1–15. Springer, 2016.
- [Bea83a] Arnaud Beauville. Some remarks on Kahler manifolds with $c_1 = 0$. *Classification of algebraic and analytic manifolds, Katata, 1982*, 1983.

- [Bea83b] Arnaud Beauville. Variétés Kähleriennes dont la première classe de Chern est nulle. *Journal of Differential Geometry*, 18(4):755–782, 1983.
- [Bea11] Arnaud Beauville. Antisymplectic involutions of holomorphic symplectic manifolds. *Journal of Topology*, 4(2):300–304, 2011.
- [BHPVdV15] Wolf Barth, Klaus Hulek, Chris Peters, and Antonius Van de Ven. *Compact complex surfaces*, volume 4. Springer, 2015.
- [BM14] Arend Bayer and Emanuele Macrì. MMP for moduli of sheaves on K3s via wall-crossing: nef and movable cones, Lagrangian fibrations. *Inventiones mathematicae*, 198(3):505–590, 2014.
- [Bou04] Sébastien Boucksom. Divisorial Zariski decompositions on compact complex manifolds. In *Annales scientifiques de l’Ecole normale supérieure*, volume 37, pages 45–76, 2004.
- [Bre13] Glen E Bredon. *Topology and geometry*, volume 139. Springer Science & Business Media, 2013.
- [Cat19] Alberto Cattaneo. Automorphisms of Hilbert schemes of points on a generic projective K3 surface. *Mathematische Nachrichten*, 292(10):2137–2152, 2019.
- [CGM19] Chiara Camere, Alice Garbagnati, and Giovanni Mongardi. Calabi–Yau quotients of hyperkähler four-folds. *Canadian Journal of Mathematics*, 71(1):45–92, 2019.
- [DHH⁺15] Ulrich Derenthal, Jürgen Hausen, Armand Heim, Simon Keicher, and Antonio Laface. Cox rings of cubic surfaces and Fano threefolds. *Journal of Algebra*, 436:228–276, 2015.
- [DK18] Olivier Debarre and Alexander Kuznetsov. Gushel-Mukai varieties: Classification and birationalities. *Algebraic Geometry*, pages 15–76, 2018.
- [DK19] Olivier Debarre and Alexander Kuznetsov. Double covers of quadratic degeneracy and Lagrangian intersection loci. *Mathematische Annalen*, pages 1–35, 2019.
- [DM19] Olivier Debarre and Emanuele Macrì. On the period map for polarized hyperkähler fourfolds. *International Mathematics Research Notices*, 2019(22):6887–6923, 2019.
- [EH16] David Eisenbud and Joe Harris. *3264 and all that: A second course in algebraic geometry*. Cambridge University Press, 2016.
- [EJ16] Andreas-Stephan Elsenhans and Jörg Jahnel. Point counting on k3 surfaces and an application concerning real and complex multiplication. *LMS Journal of Computation and Mathematics*, 19(A):12–28, 2016.
- [EPW01] David Eisenbud, Sorin Popescu, and Charles Walter. Lagrangian subbundles and codimension 3 subcanonical subschemes. *Duke Mathematical Journal*, 107(3), 2001.
- [Fer12] Andrea Ferretti. The Chow ring of double EPW sextics. *Algebra & Number Theory*, 6(3):539–560, 2012.
- [FMOS20] Laure Flapan, Emanuele Macrì, Kieran O’Grady, and Giulia Saccà. Antisymplectic involutions on hyperkähler manifolds. In *2020 Fall Eastern Virtual Sectional Meeting*. AMS, 2020.
- [Fuj87] Akira Fujiki. On the de Rham cohomology group of a compact Kähler symplectic manifold. In *Algebraic Geometry, Sendai, 1985*, pages 105–165. Mathematical Society of Japan, 1987.
- [Ful13] William Fulton. *Intersection theory*, volume 2. Springer Science & Business Media, 2013.
- [GH78] Phillip Griffiths and Joseph Harris. *Principles of algebraic geometry*, volume 19. Wiley Online Library, 1978.
- [GHJ12] Mark Gross, Daniel Huybrechts, and Dominic Joyce. *Calabi–Yau manifolds and related geometries: lectures at a summer school in Nordfjordeid, Norway, June 2001*. Springer Science & Business Media, 2012.
- [Gua01] Daniel Guan. On the Betti numbers of irreducible compact hyperkähler manifolds of complex dimension four. *Mathematical Research Letters*, 8(5):663–669, 2001.
- [Har13] Robin Hartshorne. *Algebraic geometry*, volume 52. Springer Science & Business Media, 2013.
- [Hat05] Allen Hatcher. *Algebraic topology*. Cambridge, 2005.
- [HT01] Brendan Hassett and Yuri Tschinkel. Rational curves on holomorphic symplectic fourfolds. *Geometric And Functional Analysis*, 11(6):1201–1228, 2001.
- [Huy16] Daniel Huybrechts. *Lectures on K3 surfaces*, volume 158. Cambridge University Press, 2016.
- [Ive06] Birger Iversen. *Linear determinants with applications to the Picard scheme of a family of algebraic curves*, volume 174. Springer, 2006.
- [Mar07] Eyal Markman. Integral generators for the cohomology ring of moduli spaces of sheaves over Poisson surfaces. *Advances in Mathematics*, 208(2):622–646, 2007.
- [Mar11] Eyal Markman. A survey of Torelli and monodromy results for holomorphic-symplectic varieties. In *Complex and differential geometry*, pages 257–322. Springer, 2011.
- [Mar13] Eyal Markman. Prime exceptional divisors on holomorphic symplectic varieties and monodromy reflections. *Kyoto Journal of Mathematics*, 53(2):345–403, 2013.
- [MFK94] David Mumford, John Fogarty, and Frances Kirwan. *Geometric invariant theory*, volume 34. Springer Science & Business Media, 1994.
- [Nov21] Simone Novario. Hodge classes of type $(2, 2)$ on Hilbert squares of projective K3 surfaces. *arXiv preprint arXiv:2112.11306*, 2021.
- [Nov] Simone Novario. *Linear systems on irreducible holomorphic symplectic manifolds*. PhD thesis, Università degli studi di Milano, Université de Poitiers, available at https://air.unimi.it/bitstream/2434/886303/2/phd_unimi_R12353.pdf, 2021.
- [O’G99] K. G. O’Grady. Desingularized moduli spaces of sheaves on a K3. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 1999(512):49–117, 1999.
- [O’G03] Kieran G. O’Grady. A new six-dimensional irreducible symplectic variety. *Journal of Algebraic Geometry*, 12(3):435–505, 2003.

- [O'G06] Kieran G. O'Grady. Irreducible symplectic 4-folds and Eisenbud–Popescu–Walter sextics. *Duke Mathematical Journal*, 134(1):99–137, 2006.
- [O'G08] Kieran G O'Grady. Irreducible symplectic 4-folds numerically equivalent to $(K3)^{[2]}$. *Communications in Contemporary Mathematics*, 10(04):553–608, 2008.
- [O'G10] Kieran G O'Grady. Higher-dimensional analogues of K3 surfaces. *Current developments in algebraic geometry*, 59:257–293, 2010.
- [O'G13] Kieran O'Grady. Double covers of EPW-sextics. *Michigan Mathematical Journal*, 62(1), 2013.
- [Per13] Oskar Perron. *Die Lehre von den Kettenbrüchen: B and I: Elementare Kettenbrüche*. Springer-Verlag, 2013.
- [PS08] Chris AM Peters and Joseph HM Steenbrink. *Mixed hodge structures*, volume 52. Springer Science & Business Media, 2008.
- [Rie18] Ulrike Rieß. Base divisors of big and nef line bundles on irreducible symplectic varieties. *arXiv preprint arXiv:1807.05192*, 2018.
- [SD74] Bernard Saint-Donat. Projective models of K-3 surfaces. *American Journal of Mathematics*, 96(4):602–639, 1974.
- [Ver96] Mikhail Verbitsky. Cohomology of compact hyperkähler manifolds and its applications. *Geometric & Functional Analysis GAFA*, 6(4):601–611, 1996.
- [vG08] Bert van Geemen. Real multiplication on k3 surfaces and kuga satake varieties. *Michigan Mathematical Journal*, 56(2):375–399, 2008.
- [vGS07] Bert van Geemen and Alessandra Sarti. Nikulin involutions on K3 surfaces. *Mathematische Zeitschrift*, 255(4):731–753, 2007.
- [Wel81] Gerald E Welters. Abel–Jacobi isogenies for certain types of Fano threefolds. *MC Tracts*, 1981.
- [Yok94] Hideo Yokoi. Solvability of the diophantine equation $x^2 - dy^2 = \pm 2$ and new invariants for real quadratic fields. *Nagoya Mathematical Journal*, 134:137–149, 1994.
- [Zar83] Yu.G. Zarhin. Hodge groups of k3 surfaces. *Journal für die reine und angewandte Mathematik*, 341:193–220, 1983.

UNIVERSITÀ DEGLI STUDI DI MILANO, DIPARTIMENTO DI MATEMATICA “F. ENRIQUES”, VIA CESARE SALDINI 50, 20133 MILANO, ITALY.

UNIVERSITÉ DE POITIERS, LABORATOIRE DE MATHÉMATIQUES ET APPLICATIONS, TÉLÉPORT 2, BOULEVARD MARIE ET PIERRE CURIE, 86962 FUTUROSCOPE CHASSENEUIL, FRANCE.

Email address: `simone.novario@unimi.it`, `simone.novario@math.univ-poitiers.fr`