

2-Dimensional Groups with Action : The Category of Crossed Module of Groups with Action

A. Odabaşı^a and E. Soylu Yılmaz^a

^aDepartment of Mathematics and Computer Science, Osmangazi University, Eskişehir, Turkey

Abstract

In this paper, we define the notion of crossed modules of groups with action and investigate related structures. Functions for computing of these structures have been written using the GAP computational discrete algebra programming language.

Key Words: Group with action; category; Gap computational discrete algebra programming language.

Classification: 18-08, 18G50.

1 Introduction

Crossed modules can be considered as two-dimensional of algebraic structures were first mentioned by Whitehead in 1941 [21]. Later Whitehead named crossed modules in [22] as an additional note of his early work. These concepts aroused in the paper entitled 'Combinatorial homotopy II' [23] paper which also introduced the substantial concept of a free crossed module. Many generalizations of crossed module are given in the paper of Janelidze [12].

The term 2-group (two-dimensional group) can be considered as a cluster of well-matched categories of crossed modules and also the cat-1 groups which are the pair of groups. Both crossed modules and cat-1 groups can be viewed as a Moore complex of a simplicial group.

The notion of the group with action first appeared in Datuashvili's paper [9]. Datuashvili demonstrated this structure by one of the three problems of Loday mentioned in [15, 16]. The problem is associated with the Leibniz algebras to define the algebraic correspondence called 'coquecigrue' object as Lie group structure. Leibniz algebras are closely related with the lower central series of a group constructed and examined in [24] by Witt.

Leibniz algebras was firstly identified with the research of Loday in 1989 [14]. A Leibniz algebra is a non-associative equivalent of a Lie algebra. To explain this association, the functor

$$F : \mathbf{Gr}_{\mathbf{Lie}} \longrightarrow \mathbf{Lie}$$

was constructed by Witt [24] in 1937 where $\mathbf{Gr}_{\mathbf{Lie}}$ is the category of Lie groups and \mathbf{Lie} is the category of Lie algebras. A group with action arises from the analogous version of the above functor taking the Leibniz algebras instead of Lie algebras. Since Leibniz algebras are non-associative congruence of Lie algebras, a group with action takes place of Lie groups. Therefore, Datuashvili defined the functor

$$\mathbf{Gr}^\bullet \longrightarrow \mathbf{Leibniz}$$

from the category, \mathbf{Gr}^\bullet , of group with action to the category, $\mathbf{Leibniz}$, of Leibniz algebras (see for details [9, 10]).

Likewise obtaining a group version of Lie algebras, the category of Leibniz algebras are equivalent to the category of group with action.

A shared package XMod [1], for the GAP [11], computational discrete algebra system was described by Wensley et al. which contains functions for computing crossed modules of groups and cat^1 -groups and their morphisms. Thereafter, the algebraic version of a GAP package XModAlg [5] was given by Arvasi and Odabas (see [4]). In this paper, we describe a package XModGwA for GAP which constructs crossed modules of groups with action (see [18]).

In this paper, we investigate the simplicial group with action corresponding to a crossed module of group with action. We also give a natural equivalence for these structures.

Shortly, we can summarize the purpose of this paper as:

- To construct the two - dimensional group with action as crossed module of group with action,
- To determine the action conditions between two group with actions,
- To compose a group with action crossed module by an action,
- To give the categorical equivalences between other structures.

2 The Main Text

The concept of crossed module, generalizing the notion of a G -module, was introduced by Whitehead [23] in the course of his studies on the algebraic structure on the second relative homotopy group. We first recall the crossed modules of groups from [19, 3]. A crossed module of groups is a homomorphism $\partial : S \rightarrow R$ where S and R are groups and there exist an action of R on S denoted by $(r, s) \mapsto r \cdot s$. These data must satisfy the following two conditions,

CM1) ∂ is R -equivariant, so $\partial(r \cdot s) = r\partial(s)r^{-1}$

CM2) Peiffer rule; $\partial(s) \cdot s_1 = ss_1s^{-1}$

where $r \in R$ and $s, s_1 \in S$.

A crossed module is written by $\mathcal{X} = (\partial : S \rightarrow R)$ notation. The groups S, R and the group homomorphism ∂ are called the *source*, *range* and *boundary* of \mathcal{X} respectively. When only the first of these conditions is satisfied, the resulting structure is a *pre-crossed module*. Given a pre-crossed module $\mathcal{X} = (\partial : S \rightarrow R)$, one can form an internal directed graph in the category of groups simply by forming the semidirect product $S \rtimes R$ and taking the source and target to send an element (s, r) to r and $\partial(s)r$ respectively.

Example 1. • Let G be a group and N be a normal subgroup of G . $\mathcal{X} = (\text{inc} : N \hookrightarrow G)$ is a crossed module. Where G acts on N by conjugation.

- Let M be a G -module. \mathcal{X} is a crossed module with trivial morphism $0_M : M \rightarrow G, m \mapsto 1_G$.
- For a group $H, \alpha : H \rightarrow \text{Aut}(H)$ homomorphism represents the action so that $\mathcal{X} = (\alpha : H \rightarrow \text{Aut}(H))$ is a crossed module.
- A central extension crossed module has boundary as surjection $\partial : S \rightarrow R$ with the central kernel, where $r \in R$ acts on S by conjugation with $\partial^{-1}r$.
- The direct product of $\mathcal{X} = (\partial : S \rightarrow R)$ and $\mathcal{X}' = (\partial' : S' \rightarrow R')$ is $\mathcal{X} \times \mathcal{X}' = (\partial \times \partial' : S \times S' \rightarrow R \times R')$ with R, R' acting trivially on S, S' respectively.

Let $\mathcal{X} = (\partial : S \rightarrow R)$ and $\mathcal{X}' = (\partial' : S' \rightarrow R')$ be two crossed modules. The diagram

$$\begin{array}{ccc} S & \xrightarrow{\theta} & S' \\ \partial \downarrow & & \downarrow \partial' \\ R & \xrightarrow{\psi} & R' \end{array}$$

is commutative. That is for $r \in R$ and $s \in S$

$$\partial'(\theta(s)) = \psi(\partial(s))$$

and θ preserves the action of R on S .

$$\theta(r \cdot s) = \psi(r) \cdot \theta(s)$$

then (θ, ψ) is called the morphism of crossed modules. This states a category of crossed modules, **XMod**.

A simplicial group \mathbf{G} consists of a family of groups $\{G_n\}$ together with face and degeneracy maps

$$\begin{aligned} d_i &= d_i^n : G_n \rightarrow G_{n-1}, \quad 0 \leq i \leq n, \quad (n \neq 0) \\ s_i &= s_i^n : G_n \rightarrow G_{n+1}, \quad 0 \leq i \leq n, \quad (n \neq 0) \end{aligned}$$

satisfying the usual simplicial identities given in [17, 8]. The category of simplicial groups is denoted by **SimpGrp**.

The Moore complex \mathbf{NG} of a simplicial group \mathbf{G} is defined to be the normal chain complex (\mathbf{NG}, ∂) with

$$NG_n = \bigcap_{i=0}^{n-1} \ker d_i$$

and with $\partial_n : NG_n \rightarrow NG_{n-1}$ induced from d_n by restriction. The n^{th} homotopy group $\pi_n(\mathbf{G})$ of \mathbf{G} is the n^{th} homology of the Moore complex of \mathbf{G} , i.e.

$$\pi_n(\mathbf{G}) \cong H_n(\mathbf{NG}, \partial) = \bigcap_{i=0}^n \ker d_i^n / d_{n+1}^{n+1} \left(\bigcap_{i=0}^n \ker d_i^{n+1} \right).$$

We say that the Moore complex \mathbf{NG} of a simplicial group \mathbf{G} is of length k if $\mathbf{NG}_n = 1$ for all $n \geq k + 1$. We denote the category of simplicial groups with Moore complex of length k by **SimpGrp** $_{\leq k}$.

Theorem 1. The category of crossed modules is equivalent to the category of simplicial groups with Moore complex of length 1. (see [13])

3 Group with Action

In this section, we recall the definition of group with action given by Datuashvili in [9].

Let G be a group. A map $\varepsilon : G \times G \rightarrow G$ represents the right action on itself. For $g, g', g'' \in G$,

$$\begin{aligned} \varepsilon(g, g' + g'') &= \varepsilon(\varepsilon(g, g'), g'') \\ \varepsilon(g, 0) &= g \\ \varepsilon(g' + g'', g) &= \varepsilon(g', g) + \varepsilon(g'', g) \\ \varepsilon(0, g) &= 0 \end{aligned}$$

with the above conditions, G^\bullet is called the group with action. The action is denoted by $\varepsilon(g, h) = g^h$, for $g, h \in G$. Here the group operation is addition.

Let (G, ε) and (G', ε') be group with actions. A morphism between group with actions is denoted with the following diagram.

$$\begin{array}{ccc} G \times G & \xrightarrow{\varepsilon} & G \\ (\varphi, \varphi) \downarrow & & \downarrow \varphi \\ G' \times G' & \xrightarrow{\varepsilon'} & G' \end{array}$$

with $\phi : G \longrightarrow G'$ map. Furthermore for $g, h \in G$

$$(G, \varepsilon_G) \longrightarrow (G', \varepsilon'_G)$$

Examples

- Let G be a group. Consider G^\bullet as a group with action with the (right) action by conjugation.
- Consider the group klein four $Kl_4 = \{e, a, b, ab\}$. We have ten groups with action obtained from Kl_4 . Three of them are denoted by (Kl_4, ε_i) , $i = 1, 2, 3$. The tables of the actions ε_i are as follows:

ε_1	e	a	b	ab
e	e	a	b	ab
a	e	a	b	ab
b	e	a	ab	b
ab	e	a	ab	b

ε_2	e	a	b	ab
e	e	a	b	ab
a	e	ab	b	a
b	e	a	b	ab
ab	e	ab	b	a

ε_3	e	a	b	ab
e	e	a	b	ab
a	e	a	b	ab
b	e	a	b	ab
ab	e	a	b	ab

where the ij th element shows the right action of the i th element on the j th element for $i, j \in \{1, 2, 3, 4\}$

Definition 1. Let G^\bullet be a group with action and A be a nonempty subset of G . If the conditions

- A is a normal subgroup of G as a group,
- $a^g \in A$, for $a \in A$ and $g \in G$,
- $-g + g^a \in A$, for $a \in A$ and $g \in G$,

satisfied, then A^\bullet is called an ideal of G^\bullet [9].

Condition 1: For each $x, y, z \in G$,

$$x - x^{(z^x)} + x^{y+z^x} - x + x^z - x^{z+y^z} = 0.$$


In [9], category of Abelian groups with action satisfying this condition and category of Lie-Leibniz algebras were defined. Then it was proved that the analogue of Witt's construction defined a functor from the category of groups with action to category of Lie-Leibniz algebras which gave rise to Leibniz algebras (introduced in [15]) over the ring of integers.

Example 2. Each group with the trivial action satisfies Condition 1.

4 Crossed Modules of Groups with Action

In this section, we will define an action between two group with actions and a new category \mathbf{XMod}^\bullet called the category of crossed module of groups with action. Let $S^\bullet = (S, \varepsilon_S)$ and $R^\bullet = (R, \varepsilon_R)$ be groups with action. We can use the exact sequence to define the action. For $s \in S, r \in R$ the action $(r, s) \mapsto r \cdot s$ of R on S can be represented with the following sequence

$$0 \longrightarrow S \xrightarrow{i} K \xrightarrow{j} R \longrightarrow 0$$



Existence of the function t with $jt = 1_R$ is the main property for the described exact sequence and the equation $r \cdot s = -t(r) + i(s) + t(r)$ denotes the R -action on S . For $s, s_1 \in S$ and $r, r_1 \in R$

$$i) \quad (r + r_1) \cdot s = r \cdot (r_1 \cdot s)$$

$$ii) \quad r \cdot (s + s_1) = r \cdot s + r \cdot s_1$$

$$\text{iii) } 0_R \cdot s = s, r \cdot 0_S = 0_s$$

In this category, there must be two derived actions of R on S corresponding to the group operations. The First one is defined above. We define the second action as $r * s = s^{t(r)}$.

The exact sequences with left group actions on itself can be represented by the following diagram.

$$\begin{array}{ccccc}
S & \xrightarrow{i} & K & \xrightarrow{j} & R \\
i_S \downarrow & & \parallel & & \downarrow i_R \\
B & & & & D \\
j_S \downarrow \quad \curvearrowright \quad t_S & & id & & \downarrow j_R \quad \curvearrowright \quad t_R \\
S & \xrightarrow{i} & K & \xrightarrow{j} & R
\end{array}$$

We obtained the following equalities via the diagram.

$$iv) \quad (r + r_1) * s = r * (r_1 * s)$$

$$v) \quad r * (s + s_1) = r * s + r * s_1$$

$$vi) \quad 0_R * s = s, r * 0_S = 0_s$$

$$vii) \quad r * (r_1 \cdot s) = (r_1^r) \cdot (r * s)$$

$$viii) \quad (r * s)^{(r \cdot s_1)} = r * s^{s_1}$$

Definition 2. Let $S^\bullet = (S, \varepsilon_S)$ and $R^\bullet = (R, \varepsilon_R)$ be groups with action. Denote $\varepsilon_S(s, s_1) = s_1^s$ and $\varepsilon_R(r, r_1) = r_1^r$ for $s, s_1 \in S$ and $r, r_1 \in R$. Let

$$\begin{array}{ccc} \cdot : R \times S & \longrightarrow & S \\ (r, s) & \mapsto & r \cdot s \end{array} \quad \text{and} \quad \begin{array}{ccc} \star : R \times S & \longrightarrow & S \\ (r, s) & \mapsto & r \star s \end{array}$$

be actions of R on S . We denote the group operation additively, nevertheless the group is not abelian. Moreover, let φ_S and φ_R be conjugate actions on S and R respectively. Using commutativity of the diagrams,

$$\begin{array}{ccc}
S \times S & \xrightarrow{\varphi_S} & S \\
\partial \times 1_S \downarrow & & \downarrow 1_S \\
R \times S & \xrightarrow{\cdot} & S \\
1_R \times \partial \downarrow & & \downarrow \partial \\
R \times R & \xrightarrow{\varphi_R} & R
\end{array}
\quad
\begin{array}{ccc}
S \times S & \xrightarrow{\varepsilon_S} & S \\
\partial \times 1_S \downarrow & & \downarrow 1_S \\
R \times S & \xrightarrow{\star} & S \\
1_R \times \partial \downarrow & & \downarrow \partial \\
R \times R & \xrightarrow{\varepsilon_R} & R
\end{array}$$

$\partial : (S, \varepsilon_S) \longrightarrow (R, \varepsilon_R)$ is a crossed module in the category of groups with action by following conditions

$$\begin{aligned}
\text{CM1)} \quad \partial(r \cdot s) &= \varphi_R(r, \partial(s)) \\
&= -r + \partial(s) + r \\
\text{CM2)} \quad \partial(s) \cdot s_1 &= 1_S(\varphi_S(s, s_1)) \\
&= \varphi_S(s, s_1) \\
&= -s + s_1 + s \\
\text{CM3)} \quad \partial(r \star s) &= \varepsilon_R(r, \partial(s)) \\
&= \partial(s)^r \\
\text{CM4)} \quad \partial(s) \star s_1 &= 1_S(\varepsilon_S(s, s_1)) \\
&= \varepsilon_S(s, s_1) \\
&= s_1^s
\end{aligned}$$

A crossed module of groups with action is written by $\mathcal{X}^\bullet = (\partial : S^\bullet \rightarrow R^\bullet)$ notation. When only **CM1** and **CM3** conditions are satisfied, the resulting structure is a pre-crossed module.

Examples

- Let R^\bullet be a group with action and S^\bullet be an ideal of R^\bullet . $\mathcal{X}^\bullet = (\text{inc} : S^\bullet \hookrightarrow R^\bullet)$ is a crossed module of groups with action. Where R acts on S by $r \cdot s = -r + s + r$ and $r \star s = \varepsilon_R(r, s) = s^r$.
- Let G^\bullet be a group with action and M be any G -module. Using $\varepsilon_M(m, m_1) = m_1^m = m_1$ trivial action M^\bullet is a group with action. Then $\mathcal{X}^\bullet = (0 : M^\bullet \rightarrow G^\bullet)$ is a crossed module of groups with action. Where the boundary of \mathcal{X}^\bullet is $(m, g) \mapsto e_G$ zero morphism and G acts on M by $(g, m) \rightarrow g \cdot m$ and $(g, m) \rightarrow g \star m$ any two actions.
- The direct product of $\mathcal{X}_1^\bullet = (\partial_1 : S_1^\bullet \rightarrow R_1^\bullet)$ and $\mathcal{X}_2^\bullet = (\partial_2 : S_2^\bullet \rightarrow R_2^\bullet)$ is $\mathcal{X}_1^\bullet \times \mathcal{X}_2^\bullet = (\partial_1 \times \partial_2 : S_1^\bullet \times S_2^\bullet \rightarrow R_1^\bullet \times R_2^\bullet)$ a crossed module of groups with action with R_1, R_2 acting trivially on S_1, S_2 respectively.

Definition 3. Let $\mathcal{X}_1^\bullet = (\partial_1 : S_1^\bullet \rightarrow R_1^\bullet)$ and $\mathcal{X}_2^\bullet = (\partial_2 : S_2^\bullet \rightarrow R_2^\bullet)$ be crossed modules of groups with action. A crossed module of groups with action morphism

$$(\alpha, \beta) : (\partial_1 : S_1^\bullet \rightarrow R_1^\bullet) \longrightarrow (\partial_2 : S_2^\bullet \rightarrow R_2^\bullet)$$

is a pair of homomorphisms $\alpha : S_1^\bullet \rightarrow S_2^\bullet, \beta : R_1^\bullet \rightarrow R_2^\bullet$ such that

$$\begin{aligned}
\text{i)} \quad & \beta \partial_1(s_1) = \partial_2 \alpha(s_1) && \text{for all } s_1 \in S_1 \\
\text{ii)} \quad & \alpha(r_1 \cdot_1 s_1) = \beta(r_1) \cdot_2 \alpha(s_1) && \text{for all } s_1 \in S_1, r_1 \in R_1 \\
\text{iii)} \quad & \alpha(r_1 \star_1 s_1) = \beta(r_1) \star_2 \alpha(s_1)
\end{aligned}$$

So we get the category of crossed module of groups with action. It is denoted by \mathbf{XMod}^\bullet . If $(\alpha, \beta) : (\partial_1 : S_1^\bullet \rightarrow R_1^\bullet) \longrightarrow (\partial_2 : S_2^\bullet \rightarrow R_2^\bullet)$ is a crossed module of groups with action morphism such that α and β both

isomorphisms then (α, β) is called an isomorphism. The kernel of (α, β) , $\ker(\alpha, \beta)$ is the crossed module of groups with action $(\partial : \ker \alpha \rightarrow \ker \beta)$.

5 Simplicial Group with Action

Definition 4. Let $[n]$ be an ordered set. For $[n]$ and $[m]$ ordered sets, $f : [n] \rightarrow [m]$ monotone function is called the operator. $\Delta[n]$ category have $[n]$ objects and f operator morphism. $\Delta^{op}[n]$ is the dual of the $\Delta[n]$ category.

Let \mathbf{Gr}^\bullet be the category of group with action. The functor

$$G^\bullet : \Delta^{op}[n] \rightarrow \mathbf{Gr}^\bullet$$

is the simplicial group with action.

The operator of the simplicial category can be represented with two special operators, δ_i^n and σ_j^n :

$$G^\bullet([n]) = G_n^\bullet, G^\bullet(\delta_i^n) = d_i^n \text{ and } G^\bullet(\sigma_j^n) = s_j^n$$

The category of simplicial group with action can be defined and is denoted by $\mathbf{SimpGrp}^\bullet$.

Definition 5. Let G^\bullet be a simplicial group with action. Let

$$NG_n^\bullet = \bigcap_{i=0}^{n-1} \ker d_i^n$$

and the restriction of function d_n^n

$$\partial_n : NG_n^\bullet \rightarrow NG_{n-1}^\bullet$$

is defined. So the chain complex

$$(NG^\bullet) : \dots NG_2^\bullet \xrightarrow{\partial_2} NG_1^\bullet \xrightarrow{\partial_1} NG_0^\bullet = G_0^\bullet$$

is called the Moore chain complex of simplicial group with action. In the chain complex

$$NG_0^\bullet = G_0^\bullet, NG_1^\bullet = \ker d_0', NG_2^\bullet = \ker d_0^2 \cap \ker d_1^2,$$

are denoted.

The following result of different versions (such as for Lie, group, algebra) can be found in [2, 19]

Theorem 2. The category of crossed module of groups with action \mathbf{XMod}^\bullet is equivalent to the category of simplicial group with action $\mathbf{SimpGrp}^\bullet$ with Moore Complex of length 1.

Proof. Let G^\bullet be a simplicial group with action with Moore complex of length 1. For the group with action crossed modules,

$$NG_2^\bullet \cap D_2 = 0 \text{ and } \partial_2(NG_2^\bullet \cap D_2) = \partial_2(G_2^\bullet \cap D_2) = \{0\}$$

is satisfied. D_2 is a subgroup generated by s_j degenerated operator in G_2^\bullet .

$$(NG^\bullet) : \dots NG_2^\bullet \xrightarrow{\partial_2} NG_1^\bullet \xrightarrow{\partial_1} NG_0^\bullet$$

chain complex has a sub-complex as

$$NG_1^\bullet \rightarrow NG_0^\bullet$$

defined sub-complex is a group with action homomorphism. An action NG_0^\bullet on NG_1^\bullet

$$\begin{array}{ccc} NG_0^\bullet \times NG_1^\bullet & \longrightarrow & NG_1^\bullet \\ (r, s) & \longrightarrow & s^r = -(s_0 r) + s + (s_0 r) \end{array}$$

$$\begin{array}{ccc} NG_0^\bullet \times NG_1^\bullet & \longrightarrow & NG_1^\bullet \\ (r, s) & \longmapsto & r * s = s^{s_0(r)} \end{array}$$

is constructed. For the first condition of group with action of crossed modules,

$$\begin{aligned} \partial_1(s^r) &= \partial_1(-(s_0 r) + s + (s_0 r)) = -r + d_1 s + r \\ &= -r + \partial_1 s + r \end{aligned}$$

is satisfied properly. On the other hand, let $a, b \in NG_1^\bullet$.

$$b^{\partial_1 a} = -s_0 d_1 a + b + s_0 d_1 a$$

is defined by the action of group with action. $NG_2^\bullet \cap D_2^\bullet = N_2^\bullet \cap D_2^\bullet$ equality is satisfied with $a, b \in NG_1^\bullet$ and

$$F_{(0)(1)}(a, b) = [s_0 a, s_1 b][s_1 b, s_1 a] \in NG_2^\bullet \cap D_2^\bullet$$

in $NG_2^\bullet \cap D_2^\bullet$. Since the length of Moore complex is 1, we have

$$\partial_2(N_2^\bullet \cap D_2^\bullet) = \{0\}$$

and the following equality

$$\begin{aligned} \partial_2(F_{(0)(1)}(a, b)) &= d_2([s_0 a, s_1 b][s_1 b, s_1 a]) \\ &= -s_0 d_1 a - b + s_0 d_1 a + b(-b - a + b + a) \in NG_2^\bullet \cap D_2^\bullet \end{aligned}$$

so

$$-s_0 d_1 a + b + s_0 d_1 a = -a + b + a$$

is the second condition of group with action crossed modules. For $r \in NG_0^\bullet, s \in NG_1^\bullet$

$$\begin{aligned} \partial_1(s^{s_0(\partial_1 r)}) &= \partial_1(s)^{\partial_1(r)} \\ &= \partial_1(s^r) \end{aligned}$$

Since ∂_1 is one-to-one morphism, crossed module of group with action CM4 condition holds.

CM3)

$$\begin{aligned} \partial_1(r * s) &= \partial_1(s^{s_0(r)}) \\ &= \partial_1(s)^{\partial_1(s_0 r)} \\ &= \partial_1(s)^r \\ &= \partial_1(s) * r \end{aligned}$$

condition is obtained.

Thus, there exists an obvious functor

$$\begin{array}{ccc} \Delta : \mathbf{XMod}^\bullet & \rightarrow & \mathbf{SimpGrp}^\bullet_{\leq 1} \\ R^\bullet & \mapsto & \Delta(R^\bullet) = (S^\bullet \rightarrow R^\bullet) \end{array}$$

can simply be noticed. Conversely, let S^\bullet and R^\bullet be a group with actions. We will show that

$$\partial : S^\bullet \longrightarrow R^\bullet$$

is a crossed module of group with action. Since R^\bullet acts on S^\bullet we far from the semi-direct product $R^\bullet \rtimes S^\bullet$. We define the semi-direct group with action $R^\bullet \rtimes S^\bullet$ by

$$(r, s)(r_1, s_1) = (r + r_1, s^{r_1} + s_1)$$

and

$$(r, s)^{(r_1, s_1)} = (r * r_1, r * s_1 + s * s_1 + s * r_1)$$

an action on S^\bullet by R^\bullet . Defining $S_0^\bullet = R^\bullet$ and $S_1^\bullet = R^\bullet \rtimes S^\bullet$ we get

$$\begin{aligned} d_1(r, s) &= (\partial_1 s) + r \\ d_0(r, s) &= r \\ s_0(r) &= (0, r) \end{aligned}$$

These operators satisfy the following simplicial identities due to being group with action morphisms.

$$\begin{aligned} d_1 s_0(r) &= d_1(r, 0) = (\partial_1 0) + r = r \\ d_0 s_0(r) &= d_0(r, 0) = r \end{aligned}$$

Thus, a 1-truncated simplicial group with action is $\{S_1^\bullet, S_0^\bullet\} = S'^\bullet$. From truncation we have the following equalities

$$\begin{aligned} NS_0'^\bullet &= NS_0^\bullet = S_0^\bullet = N \\ NS_1'^\bullet &= NS_1^\bullet = (\ker d_0)NS_2'^\bullet = \{0\} \end{aligned}$$

Equivalently, if

$$\partial_2(NS_2^\bullet \cap D_2^\bullet) = \partial_2(NS_2^\bullet) = \{0\}$$

is satisfied, S'^\bullet is denoted a simplicial group with action with Moore complex of length 1. By the definition of $F_{\alpha, \beta}$ functions

$$\partial_2(NS_2^\bullet) = [\ker d_0, \ker d_1]$$

is obtained. Also

$$\begin{aligned} d_0 : R^\bullet \rtimes S^\bullet &\longrightarrow R^\bullet \\ (r, s) &\mapsto r \end{aligned}$$

with $\ker d_0 = \{(0, s) : s \in S^\bullet\} \cong S^\bullet$ and

$$\begin{aligned} d_1 : R^\bullet \rtimes S^\bullet &\longrightarrow R^\bullet \\ (r, s) &\mapsto (\partial_1 s) + r \end{aligned}$$

with $\ker d_1 = \{(-\partial_1 s, s) : s \in S^\bullet\}$ are conveniently determined. Since $(0, s_1) \in \ker d_0$ and $(-\partial_1 s, s) \in \ker d_1$, we have

$$[(0, s_1), (-\partial_1 s, s)] = (0, 0).$$

Using the definition of semi-direct product the following equation holds

$$\begin{aligned} [(0, s_1), (-\partial_1 s, s)] &= (0, s_1)(-\partial_1 s, s)(-(0, s_1))(-(-\partial_1 s, s)) \\ &= (-\partial_1 s, (s_1)^{-\partial_1(s)} + s)(0, -s_1)(\partial_1 s, (-s)^{\partial_1 s}) \\ &= (-\partial_1 s, (s_1)^{-\partial_1 s} + s)(\partial_1 s, (-s_1)^{\partial_1 s} + (-s)^{\partial_1 s}) \\ &= (0, ((s_1)^{-\partial_1 s} + s)^{\partial_1 s} + (-s_1)^{\partial_1 s} + (-s)^{\partial_1 s}) \\ &= (0_R, 0_S) \end{aligned}$$

where $(0_{R^\bullet}, 0_{S^\bullet})$ is the identity element.

Furthermore,

$$\begin{aligned}
(0, s_1)^{(-\partial_1 s, s)} &= (0 * (-\partial_1 s), 0 * s + s_1 * s + s_1 * (-\partial_1 s)) \\
&= (0, \partial_1 s_1 * s + \partial_1 s_1 * (-s)) \\
&= (0, s^{s_1} + (-s)^{s_1}) \\
&= (0, s_1)
\end{aligned}$$

and

$$\begin{aligned}
(-\partial_1 s, s)^{(0, s_1)} &= (-\partial_1 s, -\partial_1 s * s_1 + s * s_1 + s * 0) \\
&= (-\partial_1 s, (-s) * s_1 + s * s_1 + s) \\
&= (-\partial_1 s, 0 * s_1 + s) \\
&= (-\partial_1 s, s)
\end{aligned}$$

equations are hold. So,

$$[\ker d_0, \ker d_1] = (0_{R^\bullet}, 0_{S^\bullet})$$

and

$$\partial_2(NS_2^\bullet) = \{0\}$$

are obtained from $S' = \{S_1^\bullet, S_0^\bullet\}$ with Moore complex of length 1. □

6 Computer Implementation

GAP (Groups, Algorithms, Programming [11]) is the leading symbolic computation system for solving computational discrete algebra problems. Symbolic computation has underpinned several key advances in Mathematics and Computer Science, for example, in number theory and coding theory (see [6]). **GAP** which is free, opensource, and extensible system, can deal with different discrete mathematical problems, but it focuses on computational group theory. It is distributed under the GNU Public License. The system is delivered together with the source codes, which are written in two languages: the kernel of the system is written in C, and the library of functions and additional packages is in a special language, also called **GAP**. The **GAP** system and extension packages now comprise 360K lines of C and 900K lines of **GAP** code. The Small Groups library has been used in such landmark computations as the "Millennium Project" to classify all finite groups of order up to 2000 by Besche, Eick and O'Brien in [7].

The **GwA** package for **GAP** contains functions for groups with action and their morphisms and was first described in [20]. In this paper, we have developed new functions which construct (pre) crossed modules of groups with action and renamed the package to **XModGwA**.

The function **GwA** in the package may be used in two ways. **GwA**(G) returns the group with trivial action, while **GwA**(G, act) returns a group with action for this chosen group and an $\text{act} : G \rightarrow \text{Aut}(G)$ action. Functions for groups with action include **IsGwA**, **IsPerfectGwA**, **IsIdeal**, **AllIdealOnGwA**, **IsGwAC1**, **Commutator**, **LowerCentralSeriesOfGwA**, **IsNilpotent**, and **NilpotencyClassOfGwA**. Attributes of a group with action constructed in this way include **BaseGroup** and **BaseAction**.

The function **GwAMorphismObj** defines morphisms of groups with action, and the function **IsGwAMorphism** which controls whether a map satisfies condition morphisms of groups with action or not. The function **AllGwAMorphisms** is used to find all morphisms between two groups with action.

The following **GAP** session illustrates the use of these functions.

```

gap> k14 := Group((1,2), (3,4));;
gap> allgwa_on_k14 := AllGwAOnGroup(k14);;
gap> Length(allgwa_on_k14);
10
gap> k14_A := allgwa_on_k14[4];
GroupWithAction [ Group( [ (1,2), (3,4) ] ), * ]
gap> A4 := AlternatingGroup(4);;
gap> A4_A := GwA(A4);
GroupWithAction [ AlternatingGroup( [ 1 .. 4 ] ), * ]
gap> IsGwA(k14_A); IsGwA(A4_A);
true
true
gap> IsPerfectGwA(k14_A);
false
gap> List(allgwa_on_k14, i -> NilpotencyClassOfGwA(i));
[ 1, 2, 0, 0, 0, 2, 0, 0, 0, 2 ]
gap> LowerCentralSeriesOfGwA(A4_A);
[ GroupWithAction [ AlternatingGroup( [ 1 .. 4 ] ), * ],
GroupWithAction [ Group( [ () , (1,2)(3,4), (1,3)(2,4), (1,4)(2,3) ] ), * ] ]
gap> gen_k14 := GeneratorsOfGroup(k14);;
gap> gen_A4 := GeneratorsOfGroup(A4);;
gap> el_A4 := Elements(A4);;
gap> f := GroupHomomorphismByImages(k14, A4, gen_k14, [el_A4[4], el_A4[4]]);
[ (1,2), (3,4) ] -> [ (1,2)(3,4), (1,2)(3,4) ]
gap> m := GwAMorphismObj(k14_A, A4_A, f);
GroupWithAction [ Group(
[ (1,2), (3,4) ] ), * ] => GroupWithAction [ AlternatingGroup( [ 1 .. 4 ] ), * ]
gap> IsGwAMorphism(m);
true

```

The function **AllGwAOnGroup(G)** constructs a list of all groups with action over G . The function **AreIsomorphicGwA** is used for checking whether or not two groups with action are isomorphic, and **IsomorphicGwAFamily** returns a list of representatives of the isomorphism classes.

In the following GAP session, we compute all 736 groups with action on $C_2 \times C_2 \times C_2$; representatives of the 14 isomorphism classes; and the list of members of a family.

```

gap> C := Range(IsomorphismPermGroup(SmallGroup(8,5)));;
gap> allgwa_onC := AllGwAOnGroup(C);;
gap> Length(allgwa_onC);
736
gap> AreIsomorphicGwA(allgwa_onC[1], allgwa_onC[2]);
false
gap> IsomorphicGwAFamily(allgwa_onC[101], allgwa_onC);
[ 101, 217, 323, 403, 490, 561, 576 ]

```

Six of the isomorphism families satisfy Condition 1 and there are six families with nilpotent and eight are not nilpotent. Other features obtained with these functions are given in the table below.

Family	Number of Members	Representator	Number of Ideals	Nilpotency Class	Condition 1
1	1	1/736	16	1	✓
2	84	2/736	7	0	✗

3	21	4/736	8	2	✓
4	42	9/736	6	2	✓
5	84	65/736	6	0	✗
6	168	67/736	6	0	✗
7	14	71/736	6	2	✓
8	21	81/736	6	2	✓
9	7	101/736	6	2	✓
10	56	107/736	3	0	✗
11	84	108/736	5	0	✗
12	42	110/736	4	0	✗
13	84	112/736	4	0	✗
14	28	122/736	6	0	✗

Function for the action between two groups with action include **IsGwAAction**. The function **IsGwAAction** is implemented for checking the structure of a group with action.

The group R acts on S by an action $\alpha, \beta : R \rightarrow \text{Aut}(S)$ and let S^\bullet and R^\bullet be groups with action on group S and R , respectively. **IsGwAAction**($S^\bullet, R^\bullet, \alpha, \beta$) is used to verify that the conditions of action between two groups with action in page 5 are satisfied. The function **AllXModGwAActions**(S^\bullet, R^\bullet) constructs a list of all actions of group with action R^\bullet on group with action S^\bullet .

The following GAP session illustrates the use of these functions.

```

gap> G := Range(IsomorphismPermGroup(SmallGroup(8,2)));;
gap> allgwa_onG := AllGwAOnGroup(G);;
gap> Length(allgwa_onG);
32
gap> SwA := allgwa_onG[2];
GroupWithAction [ Group( [ (1,2), (3,4), (5,6) ] ), * ]
gap> RwA := allgwa_onG[4];
GroupWithAction [ Group( [ (1,2), (3,4,5,6) ] ), * ]
gap> all_acts := AllXModGwAActions(SwA,RwA);;
gap> Length(all_acts);
256
gap> act_pair := all_acts[13];;
gap> IsGwAAction(SwA,RwA,act_pair[1],act_pair[2]);
true

```

Functions for crossed modules of groups with action include **PreXModGwAObj**, **IsPreXModGwA**, **IsXModGwA** and **XModGwAByIdeal**. Attributes of a group with action constructed in this way include **XModGwAAction**, **Range**, **Source** and **Boundary**.

A structure which has **IsXModGwAC1** is a pre-crossed module or a crossed module of groups with action whose source and range are both satisfies condition 1.

The following GAP session illustrates the use of these functions.

```

gap> all_bdys := AllGwAMorphisms(SwA,RwA);;
gap> bdy := all_bdys[7];
GroupWithAction [ Group( [ (1,2), (3,4,5,6) ] ), * ] =>
GroupWithAction [ Group( [ (1,2), (3,4,5,6) ] ), * ]
gap> XM1 := PreXModGwAObj(bdy,act_pair[1],act_pair[2]);
GroupWithAction [ Group( [ (1,2), (3,4,5,6) ] ), * ] =>

```

```

GroupWithAction [ Group( [ (1,2), (3,4,5,6) ] ), * ]
gap> IsPreXModGwA(XM1);
true
gap> IsXModGwA(XM1);
Condition 4 is fail
For s = (1,2) and s1 = (1,2) => (1,2) <> (1,2) (3,5) (4,6)
false
gap> all_ideals := AllIdealOnGwA(RwA);;
gap> Length(all_ideals);
6
gap> SwA := all_ideals[3];
GroupWithAction [ Group( [ (1,2) (3,4,5,6), (3,5) (4,6) ] ), * ]
gap> XM2 := XModGwABYIdeal(RwA,SwA);
GroupWithAction [ Group( [ (1,2) (3,4,5,6), (3,5) (4,6) ] ), * ] =>
GroupWithAction [ Group([ (1,2), (3,4,5,6) ] ), * ]
gap> IsXModGwA(XM2);
true
gap> IsXModGwAC1(XM2);
true

```

The global function **AllXModsGwA** may be called in two ways: as **AllXModsGwA(S^\bullet, R^\bullet)** to compute all crossed modules with chosen source and range groups with action; as **AllXModsGwA(x, y, m, n)** to compute all crossed modules of groups with actions with given size and numbers of small groups. The function computes both all pre-crossed modules and crossed modules of groups with action.

In the following GAP session, we get crossed modules using the function.

```

gap> all_XM3 := AllXModsGwA(SwA,RwA);;
gap> Length(all_XM3[1]); Length(all_XM3[2]);
66
10
all_XM4 := AllXModsGwA(4,1,4,2);;
Length(all_XM4[1]); Length(all_XM4[2]);
416
184
gap> list := Filtered(all_XM4[2], XM -> IsXModGwAC1(XM));;
gap> Length(list);
88

```

References

- [1] M., Alp, A., Odabas, E.O, Uslu and C.D., Wensley, Crossed modules and cat^1 -groups, (manual for the XMod package for **GAP**, version 2.77) (2019).
- [2] Z., Arvasi and T., Porter, Higher Dimensional Peiffer Elements in Simplicial Commutative Algebras, *Theory and Applications of Categories*, **vol. 3, no. 1** (1997) 1–23.
- [3] Z., Arvasi, T.S., Kuzpinari and E. Ö., Uslu, Three Crossed Modules, *Homology, Homotopy and Applications* **11, no. 2** (2009) 161-187.
- [4] Z., Arvasi and A., Odabas, Computing 2-dimensional algebras: Crossed modules and Cat^1 -algebras, *J. Algebra Appl.* **15** (2016) 165-185.

- [5] Z., Arvasi and A., Odabas, *Crossed Modules and cat^1 -algebras*. (manual for the `texts-f-XModAlg` share package for GAP, version 1.17) (2018).
- [6] R., Behrends, K., Hammond, V., Janjic, A., Kononov, S., Linton, H-W., Loidl, P., Maier and P., Trinder, *HPC-GAP: Engineering a 21st-century high-performance computer algebra system*, *Concurrency and Computation Practice and Experience* **28** (2016) 3606-3636.
- [7] B., Eick, H. U., Besche and E. A., O'brien, *A Millennium Project : Constructing Small Groups*, *International Journal of Algebra and Computation* **Vol. 12, No. 05** (2002) 623-644.
- [8] E.B., Curtis, *Simplicial Homotopy Theory*, *Adv. in Math.* **6** (1971) 107-209.
- [9] T., Datuashvili, *Central Series For Groups With Action and Leibniz Algebras*, *Georgian Mathematical Journal* **Volume 9, Number 4** 2002 671-682.
- [10] T., Datuashvili, *Witt's theorem for groups with action and free Leibniz algebras*, *Georgian Mathematical Journal* **Volume 11, Number 4** (2004) 691-712.
- [11] The GAP Group, *GAP – Groups, Algorithms, and Programming*, version 4.10.2 (<https://www.gap-system.org>) (2019).
- [12] G., Janelidze, *Internal crossed modules*, *Georgian Math. J.* **10 (1)** (2003) 99-114.
- [13] J.-L., Loday, *Spaces having finitely many non-trivial homotopy groups*, *Jour. Pure Appl. Algebra* **24** (1982) 179-202.
- [14] J.-L., Loday, *Operations sur l'homologie cyclique des algebres commutatives*, *Invent. Math.* **96**(1989) 205-230.
- [15] J.-L., Loday, *Une version non commutative des algebres de Lie: les algebres de Leibniz*, *Enseign. Math.* **(2) 39** (1993).
- [16] J.-L., Loday, *Algebraic K-theory and related operads*, *Lecture Notes in Math.* Springer, Berlin (2001).
- [17] J.P., May, *Simplicial Objects in Algebraic Topology*, Van Nostrand, *Math. Studies* **11** (1965).
- [18] A., Odabas and E., Soylu, *Crossed Modules of Groups with Action*, (The `XModGwA` share package for GAP, version 1.12) (<http://fef.ogu.edu.tr/aodabas/xmodgwa/>) (2020).
- [19] T., Porter, *Homotopy Quantum Field Theories meets the Crossed Menagerie: an introduction to HQFTs and their relationship with things simplicial and with lots of crossed gadgetry*, *Lecture Notes* (2011).
- [20] E.O., Uslu, A.F., Aslan and A., Odabas, *On groups with action on itself*, *Georgian Mathematical Journal* **26(3)** (2019) 459-470.
- [21] J.H.C., Whitehead, *On adding relations to homotopy groups*, *Annals of Mathematics* **42 (2)** (1941) 409-428.
- [22] J.H.C., Whitehead, *Note on a previous paper entitled "On adding relations to homotopy groups"*, *Annals of Mathematics* **47 (2)** (1946) 806- 810.
- [23] J.H.C., Whitehead, *Combinatorial homotopy II*, *Bulletin of the American Mathematical Society* **55 (3)** (1949) 213- 245.
- [24] E., Witt, *Treue Darstellung Liescher Ringe*, *J. Reine Angew. Math.* **177** (1937) 152-160.