

A two-strain reaction-diffusion malaria model with seasonality and vector-bias*

Huijie Chu, Zhenguo Bai

School of Mathematics and Statistics, Xidian University, Xi'an, 710126, China

E-mail: zgbai@xidian.edu.cn

Abstract

To investigate the combined effects of drug resistance, seasonality and vector-bias, we formulate a periodic two-strain reaction-diffusion model. It is a competitive system for resistant and sensitive strains, but the single-strain subsystem is cooperative. We derive the basic reproduction number \mathcal{R}_i and the invasion reproduction number $\hat{\mathcal{R}}_i$ for strain i ($i = 1, 2$), and establish the transmission dynamics in terms of these four quantities. More precisely, (i) if $\mathcal{R}_1 < 1$ and $\mathcal{R}_2 < 1$, then the disease is extinct; (ii) if $\mathcal{R}_1 > 1 > \mathcal{R}_2$ ($\mathcal{R}_2 > 1 > \mathcal{R}_1$), then the sensitive (resistant) strains are persistent, while the resistant (sensitive) strains die out; (iii) if $\mathcal{R}_i > 1$ and $\hat{\mathcal{R}}_i > 1$ ($i = 1, 2$), then two strains are coexistent and periodic oscillation phenomenon is observed. We also study the asymptotic behavior of the basic reproduction number with respect to small and large diffusion coefficients. Numerically, we demonstrate the phenomena of coexistence and competitive exclusion for two strains and explore the influences of seasonality and vector-bias on disease spreading.

Key words: Malaria model; Seasonality; Vector-bias; Two strains; Reproduction numbers.

AMS Subject Classification: 92D30, 37N25, 34K13

1 Introduction

Malaria, one of the most common vector-borne diseases, is endemic in over 100 countries worldwide and causes serious public health problems and a significant economic burden worldwide [1]. Human malaria infection is caused by the genus *Plasmodium* parasite, which can be transmitted to humans by the effective bites of adult female *Anopheles* mosquitoes (after taking a blood meal from humans) [2]. According to the 2020 WHO report [3], the global tally of malaria cases was 229 million in 2019, claiming some 409 000 lives compared

*This research is supported by the NSF of China (No. 11971369) and the Fundamental Research Funds for the Central Universities (No. JB210711).

to 411 000 in 2018. Therefore, a deep understanding of malaria transmission mechanisms will undoubtedly contribute to disease control.

Mathematical models have been proposed to study the dynamics of malaria outbreaks in different parts of the world, the earliest model dates back to the Ross-Macdonald model [4, 5]. Since then, various mathematical models have been designed to describe and predict the spreading of malaria (see, e.g., [6, 7, 8, 9, 10, 11, 12, 13, 14]). However, few studies consider the following three biological factors for malaria transmission simultaneously.

Vector-bias effect. The vector-bias describes that mosquitoes prefer biting infectious humans to susceptible ones. Kingsolver [15] first introduced a vector-bias model for the dynamics of malarial transmission. Following Kingsolver's work, Hosack et al. [16] included the incubation time in mosquitoes to study the dynamics of the disease concerning the reproduction number. Further, Chamchod and Britton [7] extended the model from previous authors by defining the attractiveness in a different way. Motivated by these works, Wang and Zhao incorporated the seasonality into a vector-bias model with incubation period [11]. Bai et al. formulated a time-delayed periodic reaction-diffusion model with vector-bias effect [12] and found that the ignorance of the vector-bias effect will underestimate the infection risk. All these results show that the vector-bias has an important impact on the epidemiology of malaria.

Drug-resistance. Currently, due to the lack of effective and safe vaccine, the main strategy in controlling malaria is drugs. However, the use of anti-malarial drugs such as chloroquine, malarquine, nivaquine, aralen and fansidar results in the appearance and spread of resistance in the parasite population [2, 17, 18]. This poses a significant challenge to the global control of malaria transmission or eradication of the disease. Therefore, it is essential to investigate the resistance in malaria transmission.

Seasonality. It is generally believed that climatic factors such as temperature, rainfall, humidity, wind, and duration of daylight greatly influence the transmission and distribution of vector-borne diseases [19, 20, 21]. For example, rising temperatures will reduce the number of days required for breeding, and thereby increase mosquito development rates [22]. There have been some mathematical models and field observations suggesting that the strength and mechanisms of seasonality can change the pattern of infectious diseases [22, 23]. These results are beneficial for forecasting the mosquito abundance and further effectively controlling the disease.

Except these considerations above, human and vector populations have also contributed to the spread of vector-borne diseases [6, 9]. Therefore, this paper will investigate a periodic two-strain malaria model with diffusion, which is an extension of autonomous limiting system in [24]. In view of the intrinsic mathematical structure of the model, we choose a time-varying phase space to carry out dynamical analysis. This idea has also been used in [25]. In particular, we prove that no subset forms a cycle on the boundary with the aim of using uniform persistence theory. Its proof is nontrivial (see Theorem 4.3).

The rest of this paper is organized as follows. In the next section, we formulate the

model and study its well-posedness. In Section 3, we define the basic reproduction number \mathcal{R}_i and the invasion reproduction number $\hat{\mathcal{R}}_i$ ($i = 1, 2$) for the sensitive and resistant strains, respectively. In Section 4, we investigate the uniform persistence and extinction in terms of the reproduction numbers. In Section 5, we analyze the asymptotic behavior of the basic reproduction number concerning small and large diffusion coefficients. In Section 6, we conduct numerical study for our model. And the paper ends with a brief discussion.

2 Model formulation

Motivated by [12, 24], we consider the model with no immunity; that is, individuals who recovered from malaria cannot resist reinfection of the disease and can become susceptible directly. We assume that no susceptible individual or mosquito can be infected by two virus strains. The total human population $N_h(t, x)$ is divided into three groups: susceptible $S_h(t, x)$, infected individuals with drug sensitive strain $I_1(t, x)$ and infected individuals with drug resistant strain $I_2(t, x)$. For the vector population, only adult female mosquitoes can contract the virus due to adult males and immature mosquitoes do not take blood. Thereby, we consider only adult female mosquitoes in our model. The vector population $M(t, x)$ has the epidemiological classes denoted by $S_v(t, x)$, $I_{v1}(t, x)$ and $I_{v2}(t, x)$ for the susceptible, infected with sensitive and resistant strains, respectively.

Assume that all populations remain confined to a bounded domain $\Omega \subset \mathbb{R}^m$ ($m \geq 1$) with smooth boundary $\partial\Omega$ (when $m \geq 1$). Following the line in [12], we suppose that the density of total human population $N_h(t, x) = S_h(t, x) + I_1(t, x) + I_2(t, x)$ satisfies the following reaction-diffusion equation:

$$\begin{cases} \frac{\partial N_h(t, x)}{\partial t} = D_h \Delta N_h(t, x) + B(x, N_h)N_h(t, x) - dN_h(t, x), & t > 0, x \in \Omega, \\ \frac{\partial N_h(t, x)}{\partial \nu} = 0, & t > 0, x \in \partial\Omega, \end{cases} \quad (2.1)$$

with

$$B(x, u) = \begin{cases} b \left[1 - \frac{u}{K(x)} \right], & 0 \leq u \leq K(x), x \in \bar{\Omega}, \\ 0, & u > K(x), x \in \bar{\Omega}, \end{cases}$$

where Δ is the usual Laplacian operator. $D_h > 0$ is the diffusion coefficient of humans, b and d ($0 < d < b$) are respectively the maximal birth rate and the nature mortality rate of humans, and $K(x)$ denotes the local carrying capacity, which is supposed to be a positive continuous function of location x . By employing [26, Theorems 3.1.5 and 3.1.6], we arrive at that system (2.1) admits a globally attractive positive steady state $N(x)$ in $C(\bar{\Omega}, \mathbb{R}_+) \setminus \{0\}$.

We also assume that the equation of the total mosquito population $M(t, x) = S_v(t, x) + I_{v1}(t, x) + I_{v2}(t, x)$ is of the form:

$$\begin{cases} \frac{\partial M(t, x)}{\partial t} = D_v \Delta M(t, x) + \Lambda(t, x) - \eta(t, x)M(t, x), & t > 0, x \in \Omega, \\ \frac{\partial M(t, x)}{\partial \nu} = 0, & t > 0, x \in \partial\Omega, \end{cases} \quad (2.2)$$

where $D_v > 0$ is the diffusion coefficient of mosquitoes, $\Lambda(t, x)$ is the recruitment rate at which adult female mosquitoes emerge from larval at time t and location x , and $\eta(t, x)$ is the natural death rate of mosquitoes at time t and location x . Functions $\Lambda(t, x)$ and $\eta(t, x)$ are Hölder continuous and nonnegative nontrivial on $\mathbb{R} \times \bar{\Omega}$, and ω -periodic in t for some $\omega > 0$. It easily follows that system (2.2) admits a globally stable positive ω -periodic solution $M^*(t, x)$ in $C(\bar{\Omega}, \mathbb{R}_+)$ (see, e.g., [27, Lemma 2.1]). Biologically, we may suppose that the total human and mosquito density at time t and location x respectively stabilize at $N(x)$ and $M^*(t, x)$, that is, $N_h(t, x) \equiv N(x)$ and $M(t, x) \equiv M^*(t, x)$ for all $t \geq 0$ and $x \in \Omega$.

For model parameters, since the impact of climate change on mosquitoes activities is much more than that on humans, the parameters corresponding to mosquitoes are assumed to be time-dependent. To incorporate a vector-bias term into the model, we use the parameters p and l to describe the probabilities that a mosquito arrives at a human at random and picks the human if he is infectious and susceptible, respectively [7, 11]. Since infectious humans are more attractive to mosquitoes, we assume $p \geq l > 0$. Let $\beta(t, x)$ be the biting rate of mosquitoes at time t and location x ; $c_1(\alpha_1)$ be the transmission probability per bite from infectious mosquitoes (humans) with sensitive strain to susceptible humans (mosquitoes), and $c_2(\alpha_2)$ be the transmission probability per bite from infectious mosquitoes (humans) with resistant strain to susceptible humans (mosquitoes). According to the induction in [24], we obtain

$$\begin{aligned} J_1(I_1(t, x), I_2(t, x)) &:= \frac{c_1 \beta(t, x) l (N(x) - I_1(t, x) - I_2(t, x))}{p(I_1(t, x) + I_2(t, x)) + l(N(x) - I_1(t, x) - I_2(t, x))}, \\ J_2(I_1(t, x), I_2(t, x)) &:= \frac{\alpha_1 \beta(t, x) p I_1(t, x)}{p(I_1(t, x) + I_2(t, x)) + l(N(x) - I_1(t, x) - I_2(t, x))}, \\ J_3(I_1(t, x), I_2(t, x)) &:= \frac{c_2 \beta(t, x) l (N(x) - I_1(t, x) - I_2(t, x))}{p(I_1(t, x) + I_2(t, x)) + l(N(x) - I_1(t, x) - I_2(t, x))}, \\ J_4(I_1(t, x), I_2(t, x)) &:= \frac{\alpha_2 \beta(t, x) p I_2(t, x)}{p(I_1(t, x) + I_2(t, x)) + l(N(x) - I_1(t, x) - I_2(t, x))}, \end{aligned}$$

where $J_1(J_3)$ represents the number of newly infectious humans with sensitive (resistant) strain caused by an infected mosquito with sensitive (resistant) strain per unit time at time t and location x ; and $J_2(J_4)$ means the force of infection on mosquitoes due to the contact with infectious humans with sensitive (resistant) strain.

Taking into account all of these assumptions, we obtain the following periodic reaction-

diffusion model:

$$\left\{ \begin{array}{l} \frac{\partial I_1(t, x)}{\partial t} = D_h \Delta I_1(t, x) - (d + \gamma_1) I_1(t, x) + J_1(I_1(t, x), I_2(t, x)) I_{v1}(t, x), \quad t > 0, x \in \Omega, \\ \frac{\partial I_{v1}(t, x)}{\partial t} = D_v \Delta I_{v1}(t, x) - \eta(t, x) I_{v1}(t, x) \\ \quad + J_2(I_1(t, x), I_2(t, x))(M^*(t, x) - I_{v1}(t, x) - I_{v2}(t, x)), \quad t > 0, x \in \Omega, \\ \frac{\partial I_2(t, x)}{\partial t} = D_h \Delta I_2(t, x) - (d + \gamma_2) I_2(t, x) + J_3(I_1(t, x), I_2(t, x)) I_{v2}(t, x), \quad t > 0, x \in \Omega, \\ \frac{\partial I_{v2}(t, x)}{\partial t} = D_v \Delta I_{v2}(t, x) - \eta(t, x) I_{v2}(t, x) \\ \quad + J_4(I_1(t, x), I_2(t, x))(M^*(t, x) - I_{v1}(t, x) - I_{v2}(t, x)), \quad t > 0, x \in \Omega, \\ \frac{\partial I_1(t, x)}{\partial \nu} = \frac{\partial I_{v1}(t, x)}{\partial \nu} = \frac{\partial I_2(t, x)}{\partial \nu} = \frac{\partial I_{v2}(t, x)}{\partial \nu} = 0, \quad t > 0, x \in \partial \Omega, \\ I_1(0, x) = I_1^0(x) \geq 0, \quad I_{v1}(0, x) = I_{v1}^0(x) \geq 0, \\ I_2(0, x) = I_2^0(x) \geq 0, \quad I_{v2}(0, x) = I_{v2}^0(x) \geq 0, \quad x \in \bar{\Omega}. \end{array} \right. \quad (2.3)$$

Here, the positive constants γ_1 and γ_2 denote the recovery rate of the sensitive and resistant strains for humans, respectively. The function $\beta(t, x)$ is Hölder continuous and nonnegative but not zero identically on $\mathbb{R} \times \bar{\Omega}$, and ω -periodic in t . Other parameters are the same as above.

Let $\mathbb{X} := C(\bar{\Omega}, \mathbb{R}^4)$ be the Banach space with supremum norm $\|\cdot\|$ and $\mathbb{X}^+ := C(\bar{\Omega}, \mathbb{R}_+^4)$. For each $t \geq 0$, we define

$$\begin{aligned} X(t) &:= \{\varphi = (\varphi_1, \varphi_2, \varphi_3, \varphi_4) \in \mathbb{X}^+ : 0 \leq \varphi_1(x) + \varphi_3(x) \leq N(x), \\ &\quad 0 \leq \varphi_2(x) + \varphi_4(x) \leq M^*(t, x), \quad \forall x \in \bar{\Omega}\}. \end{aligned}$$

Let $\mathbb{Y} := C(\bar{\Omega}, \mathbb{R})$ and $\mathbb{Y}^+ := C(\bar{\Omega}, \mathbb{R}_+)$. Let $T_1(t, s)$, $T_2(t, s)$, $T_3(t, s) : \mathbb{Y} \rightarrow \mathbb{Y}, t \geq s$, be the linear evolution operators associated with

$$\begin{aligned} \frac{\partial v_1(t, x)}{\partial t} &= D_h \Delta v_1(t, x) - (d + \gamma_1) v_1(t, x) := A_1 v_1(t, x), \\ \frac{\partial v_2(t, x)}{\partial t} &= D_h \Delta v_2(t, x) - (d + \gamma_2) v_2(t, x) := A_2 v_2(t, x), \\ \frac{\partial v_3(t, x)}{\partial t} &= D_v \Delta v_3(t, x) - \eta(t, x) v_3(t, x) := A_3 v_3(t, x), \end{aligned}$$

subject to the Neumann boundary condition, respectively. Noting that $T_j(t, s) = T_j(t - s), j = 1, 2$, we have $T_j(t + \omega, s + \omega) = T_j(t, s)$ for $(t, s) \in \mathbb{R}^2$ with $t \geq s, j = 1, 2$. Since $\eta(t, x)$ is ω -periodic in t , [28, Lemma 6.1] implies that $T_3(t + \omega, s + \omega) = T_3(t, s)$ for $(t, s) \in \mathbb{R}^2$ with $t \geq s$. Moreover, for $(t, s) \in \mathbb{R}^2$ with $t > s$, $T_j(t, s), j = 1, 2, 3$, are compact and strongly positive. Set $T(t, s) = \text{diag}\{T_1(t, s), T_3(t, s), T_2(t, s), T_3(t, s)\}$ and

$A(t) = \text{diag}\{A_1, A_3(t), A_2, A_3(t)\}$. Define $F = (F_1, F_2, F_3, F_4) : [0, \infty) \times \mathbb{X}^+ \rightarrow \mathbb{X}$ by

$$\begin{aligned} F_1(t, \varphi) &= \frac{c_1 \beta(t, \cdot) l(N(\cdot) - \varphi_1(\cdot) - \varphi_3(\cdot))}{p(\varphi_1(\cdot) + \varphi_3(\cdot)) + l(N(\cdot) - \varphi_1(\cdot) - \varphi_3(\cdot))} \varphi_2(\cdot), \\ F_2(t, \varphi) &= \frac{\alpha_1 \beta(t, \cdot) p \varphi_1(\cdot)}{p(\varphi_1(\cdot) + \varphi_3(\cdot)) + l(N(\cdot) - \varphi_1(\cdot) - \varphi_3(\cdot))} (M^*(t, \cdot) - \varphi_2(\cdot) - \varphi_4(\cdot)), \\ F_3(t, \varphi) &= \frac{c_2 \beta(t, \cdot) l(N(\cdot) - \varphi_1(\cdot) - \varphi_3(\cdot))}{p(\varphi_1(\cdot) + \varphi_3(\cdot)) + l(N(\cdot) - \varphi_1(\cdot) - \varphi_3(\cdot))} \varphi_4(\cdot), \\ F_4(t, \varphi) &= \frac{\alpha_2 \beta(t, \cdot) p \varphi_3(\cdot)}{p(\varphi_1(\cdot) + \varphi_3(\cdot)) + l(N(\cdot) - \varphi_1(\cdot) - \varphi_3(\cdot))} (M^*(t, \cdot) - \varphi_2(\cdot) - \varphi_4(\cdot)) \end{aligned}$$

for all $t \geq 0$ and $\varphi = (\varphi_1, \varphi_2, \varphi_3, \varphi_4) \in \mathbb{X}$. Then system (2.3) becomes

$$\begin{cases} \frac{du}{dt} = A(t)u + F(t, u), & t > 0, \\ u(0) = \varphi \in \mathbb{X}^+, \end{cases}$$

which can be written as an integral equation

$$u(t, \varphi) = T(t, 0)\varphi + \int_0^t T(t, s)F(s, u)ds, \quad \forall t \geq 0, \varphi \in \mathbb{X}^+, \quad (2.4)$$

where

$$u(t, x) = (u_1(t, x), u_2(t, x), u_3(t, x), u_4(t, x)) = (I_1(t, x), I_{v1}(t, x), I_2(t, x), I_{v2}(t, x)).$$

As usual, solutions of (2.4) are called mild solutions to system (2.3).

Lemma 2.1. *For any $\varphi \in X(0)$, system (2.3) has a unique solution $u(t, \cdot, \varphi)$ with $u(0, \cdot, \varphi) = \varphi$ such that $u(t, \cdot, \varphi) \in X(t)$ for all $t \in [0, \infty)$. Moreover, system (2.3) generates an ω -periodic semiflow $Q(t) : X(0) \rightarrow X(t)$ defined by $Q(t)\varphi = u(t, \cdot, \varphi), t \geq 0$. In addition, $Q := Q(\omega)$ admits a strong global attractor in $X(0)$.*

Proof. From the expression of F , we see that F is locally Lipschitz continuous. For any

$(t, \psi) \in \mathbb{R}_+ \times \mathbb{X}^+$ and $h > 0$, in view of $p \geq l > 0$, we have

$$\begin{aligned}
& \psi(x) + hF(t, \psi)(x) \\
&= \begin{pmatrix} \psi_1(x) + h \frac{c_1 \beta(t, x) l (N(x) - \psi_1(x) - \psi_3(x))}{p(\psi_1(x) + \psi_3(x)) + l(N(x) - \psi_1(x) - \psi_3(x))} \psi_2(x) \\ \psi_2(x) + h \frac{\alpha_1 \beta(t, x) p \psi_1(x) (M^*(t, x) - \psi_2(x) - \psi_4(x))}{p(\psi_1(x) + \psi_3(x)) + l(N(x) - \psi_1(x) - \psi_3(x))} \\ \psi_3(x) + h \frac{c_2 \beta(t, x) l (N(x) - \psi_1(x) - \psi_3(x))}{p(\psi_1(x) + \psi_3(x)) + l(N(x) - \psi_1(x) - \psi_3(x))} \psi_4(x) \\ \psi_4(x) + h \frac{\alpha_2 \beta(t, x) p \psi_3(x) (M^*(t, x) - \psi_2(x) - \psi_4(x))}{p(\psi_1(x) + \psi_3(x)) + l(N(x) - \psi_1(x) - \psi_3(x))} \end{pmatrix} \\
&\geq \begin{pmatrix} \psi_1(x) \left(1 - h \frac{c_1 \beta(t, x)}{N(x)} \psi_2(x) \right) - h \frac{c_1 \beta(t, x)}{N(x)} \psi_3(x) \psi_2(x) \\ \psi_2(x) \left(1 - h \frac{\alpha_1 \beta(t, x) p}{l N(x)} \psi_1(x) \right) - h \frac{\alpha_1 \beta(t, x) p}{l N(x)} \psi_1(x) \psi_4(x) \\ \psi_3(x) \left(1 - h \frac{c_2 \beta(t, x)}{N(x)} \psi_4(x) \right) - h \frac{c_2 \beta(t, x)}{N(x)} \psi_1(x) \psi_4(x) \\ \psi_4(x) \left(1 - h \frac{\alpha_2 \beta(t, x) p}{l N(x)} \psi_3(x) \right) - h \frac{\alpha_2 \beta(t, x) p}{l N(x)} \psi_3(x) \psi_2(x) \end{pmatrix}.
\end{aligned}$$

This implies that

$$\lim_{h \rightarrow 0^+} \frac{1}{h} \text{dist}(\psi + hF(t, \psi), \mathbb{X}^+) = 0, \quad \forall (t, \psi) \in \mathbb{R}_+ \times \mathbb{X}^+.$$

In addition, $T(t, s)\mathbb{X}^+ \subseteq \mathbb{X}^+, \forall t \geq s \geq 0$. Therefore, by [29, Corollary 4] with $K = \mathbb{X}^+$ and $S(t, s) = T(t, s)$, system (2.3) admits a unique non-continuable mild solution $u(t, \cdot, \psi)$ on its maximal existence interval $[0, t_\psi)$ with $u(0, \cdot, \psi) = \psi$, and $u(t, \cdot, \psi) \geq 0$ for all $t \in [0, t_\psi)$, where $t_\psi \leq \infty$.

Based on the above analysis, we obtain that for any $\varphi \in X(0) \subset \mathbb{X}^+$, system (2.3) has a unique solution $u(t, \cdot, \varphi) \in \mathbb{X}^+$ on $[0, t_\varphi)$ with $u(0, \cdot, \varphi) = \varphi$, where $t_\varphi \leq \infty$. Next we want to show that $u(t, x, \varphi)$ is bounded for all $t \in [0, t_\varphi)$, which then implies $t_\varphi = \infty$. To this end, we set

$$I_h(t, x) = I_1(t, x) + I_2(t, x), \quad I_v(t, x) = I_{v1}(t, x) + I_{v2}(t, x).$$

It turns out that $N(x)$ and $M^*(t, x)$ are respectively the upper solutions of the following two equations

$$\begin{aligned}
\frac{\partial I_h(t, x)}{\partial t} &= D_h \Delta I_h(t, x) - d I_h(t, x) - \gamma_1 I_1(t, x) - \gamma_2 I_2(t, x) \\
&\quad + J_1(I_1(t, x), I_2(t, x)) I_{v1}(t, x) + J_3(I_1(t, x), I_2(t, x)) I_{v2}(t, x),
\end{aligned}$$

and

$$\begin{aligned}
\frac{\partial I_v(t, x)}{\partial t} &= D_h \Delta I_v(t, x) - \eta(t, x) I_v(t, x) \\
&\quad + J_2(I_1(t, x), I_2(t, x)) (M^*(t, x) - I_v(t, x)) \\
&\quad + J_4(I_1(t, x), I_2(t, x)) (M^*(t, x) - I_v(t, x))
\end{aligned}$$

for $t \in (0, t_\varphi)$ and $x \in \bar{\Omega}$. Thus, the comparison principle implies that solutions of (2.3) are bounded on $[0, t_\varphi)$, and thus, $t_\varphi = \infty$. In addition, we also have that $u(t, \cdot, \varphi) \in X(t)$ for all $t \geq 0$, and it is classic for $t > 0$ in light of the analyticity of $T(t, s)$ when $t > s$.

Define a family of operators $\{Q(t)\}_{t \geq 0}$ from $X(0)$ to $X(t)$ by

$$[Q(t)\varphi](x) = u(t, x, \varphi), \quad \forall \varphi \in X(0), \quad x \in \bar{\Omega}.$$

By the proof of [27, Lemma 2.1], we can show that $Q(t)$ is an ω -periodic semiflow, and thus $Q := Q(\omega) : X(0) \rightarrow X(\omega) = X(0)$ is the Poincaré map associated with system (2.3). The fact that $u(t, \cdot, \varphi) \in X(t)$ for all $t \geq 0$ when $\varphi \in X(0)$ also implies that solutions of (2.3) are ultimately bounded. Hence, by [30, Theorem 2.9], Q has a strong global attractor in $X(0)$. \square

Lemma 2.2. *For any $\varphi \in X(0)$, let $u(t, x, \varphi)$ be the solution of system (2.3). If there exists some $t_0 \geq 0$ such that $u_i(t_0, x, \varphi) \neq 0$, $i = 1, 2, 3, 4$, then*

$$u_i(t, x, \varphi) > 0, \quad i = 1, 2, 3, 4, \quad \forall t \geq t_0, \quad x \in \bar{\Omega}.$$

Proof. For any given $\varphi \in X(0)$, one easily sees

$$\begin{cases} \frac{\partial u_1(t, x)}{\partial t} \geq D_h \Delta u_1(t, x) - (d + \gamma_1)u_1(t, x), & t > 0, \quad x \in \Omega, \\ \frac{\partial u_2(t, x)}{\partial t} \geq D_v \Delta u_2(t, x) - \bar{\eta}u_2(t, x), & t > 0, \quad x \in \Omega, \\ \frac{\partial u_3(t, x)}{\partial t} \geq D_h \Delta u_3(t, x) - (d + \gamma_2)u_3(t, x), & t > 0, \quad x \in \Omega, \\ \frac{\partial u_4(t, x)}{\partial t} \geq D_v \Delta u_4(t, x) - \bar{\eta}u_4(t, x), & t > 0, \quad x \in \Omega, \\ \frac{\partial u_1(t, x)}{\partial \nu} = \frac{\partial u_2(t, x)}{\partial \nu} = \frac{\partial u_3(t, x)}{\partial \nu} = \frac{\partial u_4(t, x)}{\partial \nu} = 0, & t > 0, \quad x \in \partial\Omega, \end{cases}$$

where $\bar{\eta} = \max_{(t, x) \in [0, \omega] \times \bar{\Omega}} \eta(t, x)$. If there exists $t_0 \geq 0$ such that $u_i(t_0, x, \varphi) \neq 0$ for some $i \in \{1, 2, 3, 4\}$, it then follows from the parabolic maximum principle [31, Proposition 13.1] that $u_i(t, x, \varphi) > 0$ for all $t > t_0$ and $x \in \bar{\Omega}$. \square

3 Reproduction numbers

In this section, we first define the basic reproduction number \mathcal{R}_0 of (2.3), and then introduce the invasion reproduction number $\hat{\mathcal{R}}_i$ for strain i ($i = 1, 2$).

3.1 Basic reproduction number

In order to derive the basic reproduction number of (2.3), we first consider subsystems: one involves sensitive strains alone and the other involves resistant strains alone. We fix

$i \in \{1, 2\}$ and let $I_j(t, x) \equiv 0, I_{vj}(t, x) \equiv 0, \forall t \geq 0, x \in \bar{\Omega}, j = 1, 2$ and $j \neq i$. Then system (2.3) reduces to the following single-strain model:

$$\left\{ \begin{array}{l} \frac{\partial I_i(t, x)}{\partial t} = D_h \Delta I_i(t, x) - (d + \gamma_i) I_i(t, x) \\ \quad + \frac{c_i \beta(t, x) l (N(x) - I_i(t, x))}{p I_i(t, x) + l (N(x) - I_i(t, x))} I_{vi}(t, x), \quad t > 0, x \in \Omega, \\ \frac{\partial I_{vi}(t, x)}{\partial t} = D_v \Delta I_{vi}(t, x) - \eta(t, x) I_{vi}(t, x) \\ \quad + \frac{\alpha_i \beta(t, x) p I_i(t, x)}{p I_i(t, x) + l (N(x) - I_i(t, x))} (M^*(t, x) - I_{vi}(t, x)), \quad t > 0, x \in \Omega, \\ \frac{\partial I_i(t, x)}{\partial \nu} = \frac{\partial I_{vi}(t, x)}{\partial \nu} = 0, \quad t > 0, x \in \partial \Omega, \end{array} \right. \quad (3.1)$$

Let $\mathbb{E} := C(\bar{\Omega}, \mathbb{R}^2)$ and $\mathbb{E}^+ := C(\bar{\Omega}, \mathbb{R}_+^2)$. Linearizing (3.1) at $(0, 0)$ yields

$$\left\{ \begin{array}{l} \frac{\partial I_i(t, x)}{\partial t} = D_h \Delta I_i(t, x) - (d + \gamma_i) I_i(t, x) \\ \quad + c_i \beta(t, x) I_{vi}(t, x), \quad t > 0, x \in \Omega, \\ \frac{\partial I_{vi}(t, x)}{\partial t} = D_v \Delta I_{vi}(t, x) - \eta(t, x) I_{vi}(t, x) \\ \quad + \frac{\alpha_i \beta(t, x) p M^*(t, x)}{l N(x)} I_i(t, x), \quad t > 0, x \in \Omega, \\ \frac{\partial I_i(t, x)}{\partial \nu} = \frac{\partial I_{vi}(t, x)}{\partial \nu} = 0, \quad t > 0, x \in \partial \Omega. \end{array} \right. \quad (3.2)$$

Define the operator $\mathcal{F}_i(t) : \mathbb{E} \rightarrow \mathbb{E}$ by

$$\mathcal{F}_i(t) \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = \begin{pmatrix} c_i \beta(t, \cdot) \psi_2(\cdot) \\ \frac{\alpha_i \beta(t, \cdot) p M^*(t, \cdot)}{l N(\cdot)} \psi_1(\cdot) \end{pmatrix}, \quad \forall t \in \mathbb{R}, \psi = (\psi_1, \psi_2) \in \mathbb{E}.$$

Let $-V_i(t)v = D\Delta v - W_i(t)v$, where $D = \text{diag}(D_h, D_v)$ and

$$-[W_i(t)](x) = \begin{pmatrix} -(d + \gamma_i) & 0 \\ 0 & -\eta(t, x) \end{pmatrix}, \quad \forall t \in \mathbb{R}, x \in \bar{\Omega}.$$

Then $\Psi_i(t, s) = \text{diag}(T_i(t, s), T_3(t, s))$, $t \geq s$, is the evolution operator on \mathbb{E} associated with the following system

$$\frac{dv}{dt} = -V_i(t)v$$

subject to the Neumann boundary condition. The exponential growth bound of $\Psi_i(t, s)$ is defined as

$$\bar{\omega}(\Psi_i) = \inf\{\bar{\omega}_i : \exists M \geq 1 \text{ such that } \|\Psi_i(t + s, s)\|_{\mathbb{E}} \leq M e^{\bar{\omega}_i t}, \forall s \in \mathbb{R}, t \geq 0\}.$$

By the Krein-Rutman Theorem and [31, Lemma 14.2], we have

$$0 < r(\Psi_i(\omega, 0)) = \max\{r(T_i(\omega, 0)), r(T_3(\omega, 0))\} < 1,$$

where $r(\Psi_i(\omega, 0))$ is the spectral radius of $\Psi_i(\omega, 0)$. Then, it follows from [32, Proposition 5.5] with $s = 0$ that $\bar{\omega}(\Psi_i) < 0$. Note that $\Psi_i(t, s)$ is a positive operator in the sense that $\Psi_i(t, s)\mathbb{E}^+ \subseteq \mathbb{E}^+$ for all $t \geq s$. Therefore, $\mathcal{F}_i(t)$ and $\Psi_i(t, s)$ satisfy

(H1) For each $t \geq 0$, $\mathcal{F}_i(t)$ is a positive operator on \mathbb{E} .

(H2) For any $t \geq s$, $\Psi_i(t, s)$ is a positive operator on \mathbb{E} , and $\bar{\omega}(\Psi_i) < 0$.

Let $C_\omega(\mathbb{R}, \mathbb{E})$ be the Banach space of all ω -periodic and continuous functions from \mathbb{R} to \mathbb{E} equipped with the maximum norm. Following the theory developed in [33, 34], we define two linear operators on $C_\omega(\mathbb{R}, \mathbb{E})$ by

$$[\mathcal{L}_i v](t) := \int_0^\infty \Psi_i(t, t-s) \mathcal{F}_i(t-s) v(t-s) ds, \quad \forall t \in \mathbb{R}, v \in C_\omega(\mathbb{R}, \mathbb{E}), i = 1, 2.$$

Motivated by the concept of next generation operators [32, 35], we define the basic reproduction number as $\mathcal{R}_i := r(\mathcal{L}_i)$, where $r(\mathcal{L}_i)$ is the spectral radius of \mathcal{L}_i .

The disease-free state of (2.3) is $(0, 0, 0, 0)$ and the corresponding linearized system is

$$\begin{cases} \frac{\partial I_1(t, x)}{\partial t} = D_h \Delta I_1(t, x) - (d + \gamma_1) I_1(t, x) + c_1 \beta(t, x) I_{v1}(t, x), & t > 0, x \in \Omega, \\ \frac{\partial I_{v1}(t, x)}{\partial t} = D_v \Delta I_{v1}(t, x) - \eta(t, x) I_{v1}(t, x) + \frac{\alpha_1 \beta(t, x) p M^*(t, x)}{l N(x)} I_1(t, x), & t > 0, x \in \Omega, \\ \frac{\partial I_2(t, x)}{\partial t} = D_h \Delta I_2(t, x) - (d + \gamma_2) I_2(t, x) + c_2 \beta(t, x) I_{v2}(t, x), & t > 0, x \in \Omega, \\ \frac{\partial I_{v2}(t, x)}{\partial t} = D_v \Delta I_{v2}(t, x) - \eta(t, x) I_{v2}(t, x) + \frac{\alpha_2 \beta(t, x) p M^*(t, x)}{l N(x)} I_2(t, x), & t > 0, x \in \Omega, \\ \frac{\partial I_1(t, x)}{\partial \nu} = \frac{\partial I_{v1}(t, x)}{\partial \nu} = \frac{\partial I_2(t, x)}{\partial \nu} = \frac{\partial I_{v2}(t, x)}{\partial \nu} = 0, & t > 0, x \in \partial \Omega. \end{cases} \quad (3.3)$$

Similarly, we can derive the basic reproduction number of (2.3), which is given by

$$\mathcal{R}_0 = \max\{\mathcal{R}_1, \mathcal{R}_2\}.$$

For any given $t \geq 0$, let $P_i(t)$ be the solution map of (3.2) on \mathbb{E} . Then $P_i := P_i(\omega)$ is the associated Poincaré map. Let $r(P_i)$ be the spectral radius of P_i . By [34, Theorem 3.7] with $\tau = 0$, we have the following nice property.

Lemma 3.1. $\mathcal{R}_i - 1$ has the same sign as $r(P_i) - 1$, $i = 1, 2$, and thus $\mathcal{R}_0 - 1$ has the same sign as $r(P) - 1$, where $r(P) = \max\{r(P_1), r(P_2)\}$ is the spectral radius of the Poincaré map P associated with (3.3).

3.2 Invasion reproduction number

In this subsection, we define the invasion reproduction number for each strain. The invasion reproduction number gives the ability of strain i ($i = 1, 2$) to invade strain j ($j = 1, 2, j \neq i$) measured as the number of secondary infections strain i one-infected individual can produce in a population where strain j is at an endemic state [36]. We express it by $\hat{\mathcal{R}}_i$ ($i = 1, 2$)

and give their definition by analyzing the boundary ω -periodic solution of (2.3), that is, the sensitive strain ω -periodic solution or resistant strain ω -periodic solution.

For each $t \geq 0$, let $E(t)$ be subset in \mathbb{E} defined by

$$E(t) := \{\psi = (\psi_1, \psi_2) \in \mathbb{E}^+ : 0 \leq \psi_1(x) \leq N(x), 0 \leq \psi_2(x) \leq M^*(t, x), \forall x \in \bar{\Omega}\}.$$

After a similar process in [25, Lemma 3], we obtain that for any $\psi \in E(0)$, system (3.1) has a unique solution $v_i(t, \cdot, \psi) = (I_i(t, x), I_{vi}(t, x))$ with $v_i(0, \cdot, \psi) = \psi$ such that $v_i(t, \cdot, \psi) \in E(t)$ for all $t \geq 0$. Moreover, by employing the arguments in [25, Theorem 1], one immediately obtains the following result.

Theorem 3.1. *The following statements are valid:*

- (i) *If $\mathcal{R}_i \leq 1$, then $(0, 0)$ is globally asymptotically stable for (3.1) in $E(0)$;*
- (ii) *If $\mathcal{R}_i > 1$, then system (3.1) admits a unique positive ω -periodic solution $(I_i^*(t, x), I_{vi}^*(t, x))$, and it is globally asymptotically stable for (3.1) in $E(0) \setminus \{(0, 0)\}$.*

For ease of presentation, we introduce the following notations:

- $E_0 = (0, 0, 0, 0)$: The disease-free state of (2.3).
- $E_1(t, x) = (I_1^*(t, x), I_{v1}^*(t, x), 0, 0)$: The sensitive strain ω -periodic solution of (2.3).
- $E_2(t, x) = (0, 0, I_2^*(t, x), I_{v2}^*(t, x))$: The resistant strain ω -periodic solution of (2.3).

By Theorem 3.1, we see that when $\mathcal{R}_i > 1$ ($i = 1, 2$), system (2.3) admits a unique semitrivial boundary ω -periodic solution $E_i(t, x)$. Linearizing (2.3) at the $E_j(t, x)$, $j \neq i$, $i, j = 1, 2$, and considering only the equations for $I_i(t, x)$ and $I_{vi}(t, x)$, we get

$$\left\{ \begin{array}{l} \frac{\partial I_i(t, x)}{\partial t} = D_h \Delta I_i(t, x) - (d + \gamma_i) I_i(t, x) \\ \quad + \frac{c_i \beta(t, x) l(N(x) - I_j^*(t, x))}{p I_j^*(t, x) + l(N(x) - I_j^*(t, x))} I_{vi}(t, x), \quad t > 0, x \in \Omega, \\ \frac{\partial I_{vi}(t, x)}{\partial t} = D_v \Delta I_{vi}(t, x) - \eta(t, x) I_{vi}(t, x) \\ \quad + \frac{\alpha_i \beta(t, x) p(M^*(t, x) - I_{vj}^*(t, x))}{p I_j^*(t, x) + l(N(x) - I_j^*(t, x))} I_i(t, x), \quad t > 0, x \in \Omega, \\ \frac{\partial I_i(t, x)}{\partial \nu} = \frac{\partial I_{vi}(t, x)}{\partial \nu} = 0, \quad t > 0, x \in \partial \Omega. \end{array} \right. \quad (3.4)$$

Similar to Section 3.1, we can define the invasion reproduction numbers $\hat{\mathcal{R}}_i$ ($i = 1, 2$). Further, we have the following characterization of $\hat{\mathcal{R}}_i$.

Lemma 3.2. *$\hat{\mathcal{R}}_i - 1$ has the same sign as $r(\hat{P}_i) - 1$, where \hat{P}_i is the Poincaré map associated with (3.4), and $r(\hat{P}_i)$ is the spectral radius of \hat{P}_i .*

4 Disease extinction and uniform persistence

In this section, we establish the dynamics of (2.3) in terms of \mathcal{R}_i and $\hat{\mathcal{R}}_i$, $i = 1, 2$.

4.1 Global extinction

Theorem 4.1. *If $\mathcal{R}_1 < 1$ and $\mathcal{R}_2 < 1$, then E_0 is globally attractive for (2.3) in $X(0)$.*

Proof. Let $(I_1(t, x), I_{v1}(t, x), I_2(t, x), I_{v2}(t, x))$ be the solution of (2.3) with initial data $\varphi \in X(0)$. It is easily seen that

$$\left\{ \begin{array}{l} \frac{\partial I_1(t, x)}{\partial t} \leq D_h \Delta I_1(t, x) - (d + \gamma_1) I_1(t, x) \\ \quad + \frac{c_1 \beta(t, x) l(N(x) - I_1(t, x))}{p I_1(t, x) + l(N(x) - I_1(t, x))} I_{v1}(t, x), \\ \frac{\partial I_{v1}(t, x)}{\partial t} \leq D_v \Delta I_{v1}(t, x) - \eta(t, x) I_{v1}(t, x) \\ \quad + \frac{\alpha_1 \beta(t, x) p I_1(t, x)}{p I_1(t, x) + l(N(x) - I_1(t, x))} (M^*(t, x) - I_{v1}(t, x)), \\ \frac{\partial I_1(t, x)}{\partial \nu} = \frac{\partial I_{v1}(t, x)}{\partial \nu} = 0, \end{array} \right. \quad \begin{array}{l} t > 0, x \in \Omega, \\ t > 0, x \in \Omega, \\ t > 0, x \in \partial\Omega. \end{array} \quad (4.1)$$

and

$$\left\{ \begin{array}{l} \frac{\partial I_2(t, x)}{\partial t} \leq D_h \Delta I_2(t, x) - (d + \gamma_2) I_2(t, x) \\ \quad + \frac{c_2 \beta(t, x) l(N(x) - I_2(t, x))}{p I_2(t, x) + l(N(x) - I_2(t, x))} I_{v2}(t, x), \\ \frac{\partial I_{v2}(t, x)}{\partial t} \leq D_v \Delta I_{v2}(t, x) - \eta(t, x) I_{v2}(t, x) \\ \quad + \frac{\alpha_2 \beta(t, x) p I_2(t, x)}{p I_2(t, x) + l(N(x) - I_2(t, x))} (M^*(t, x) - I_{v2}(t, x)), \\ \frac{\partial I_2(t, x)}{\partial \nu} = \frac{\partial I_{v2}(t, x)}{\partial \nu} = 0, \end{array} \right. \quad \begin{array}{l} t > 0, x \in \Omega, \\ t > 0, x \in \Omega, \\ t > 0, x \in \partial\Omega. \end{array} \quad (4.2)$$

When $\mathcal{R}_1 < 1$, Theorem 3.1 implies that $(0, 0)$ is globally stable for (3.1). Hence the comparison principle applies to (4.1) and ensures that

$$\lim_{t \rightarrow \infty} (I_1(t, x), I_{v1}(t, x)) = (0, 0) \text{ uniformly for } x \in \bar{\Omega}.$$

In the case where $\mathcal{R}_2 < 1$, by using the similar procedure as above to (4.2), one attains

$$\lim_{t \rightarrow \infty} (I_2(t, x), I_{v2}(t, x)) = (0, 0) \text{ uniformly for } x \in \bar{\Omega}.$$

Therefore, the desired result is established. \square

4.2 Competitive exclusion and coexistence

Theorem 4.2. *Let $(I_1(t, \cdot, \varphi), I_{v1}(t, \cdot, \varphi), I_2(t, \cdot, \varphi), I_{v2}(t, \cdot, \varphi))$ be the solution of (2.3) through $\varphi \in X(0)$. Then the following assertions hold.*

(1) If $\mathcal{R}_1 > 1 > \mathcal{R}_2$ and $\varphi_1, \varphi_2 \not\equiv 0$, then

$$\lim_{t \rightarrow \infty} ((I_1(t, x, \varphi), I_{v1}(t, x, \varphi), I_2(t, x, \varphi), I_{v2}(t, x, \varphi)) - E_1(t, x)) = (0, 0, 0, 0)$$

uniformly for $x \in \bar{\Omega}$.

(2) If $\mathcal{R}_2 > 1 > \mathcal{R}_1$ and $\varphi_3, \varphi_4 \not\equiv 0$, then

$$\lim_{t \rightarrow \infty} ((I_1(t, x, \varphi), I_{v1}(t, x, \varphi), I_2(t, x, \varphi), I_{v2}(t, x, \varphi)) - E_2(t, x)) = (0, 0, 0, 0)$$

uniformly for $x \in \bar{\Omega}$.

Proof. We only prove statement (1), since statement (2) can be treated similarly. In the case where $\mathcal{R}_2 < 1$, one immediately has that $\lim_{t \rightarrow \infty} (I_2(t, \cdot, \varphi), I_{v2}(t, \cdot, \varphi)) = (0, 0)$. Then the limiting system of (2.3) is the system (3.1) with $i = 1$. Moreover, by employing the theory of internally chain transitive sets (see, e.g., [26]), we conclude that

$$\lim_{t \rightarrow \infty} (I_1(t, x, \varphi), I_{12}(t, x, \varphi) - (I_1^*(t, x), I_{v1}^*(t, x))) = (0, 0)$$

uniformly for $x \in \bar{\Omega}$. Hence, statement (1) is established. \square

For each $t \geq 0$, define

$$X_0(t) := \{(\varphi_1, \varphi_2, \varphi_3, \varphi_4) \in X(t) : \varphi_i(\cdot) \not\equiv 0, i = 1, 2, 3, 4\},$$

and

$$\partial X_0(t) := X(t) \setminus X_0(t) = \{(\varphi_1, \varphi_2, \varphi_3, \varphi_4) \in X(t) : \varphi_i(\cdot) \equiv 0 \text{ at least for one } i\}.$$

In order to study the coexistence of strains, we first give the following lemma for our subsequent coexistence result.

Lemma 4.1. *Let $(I_1(t, \cdot, \varphi), I_{v1}(t, \cdot, \varphi), I_2(t, \cdot, \varphi), I_{v2}(t, \cdot, \varphi))$ be the solution of (2.3) with the initial value $\varphi \in X_0(0)$. If $\mathcal{R}_1 > 1$ and $\mathcal{R}_2 > 1$, then there exists $\delta > 0$ such that*

$$\limsup_{t \rightarrow \infty} \|(I_1(t, \cdot, \varphi), I_{v1}(t, \cdot, \varphi), I_2(t, \cdot, \varphi), I_{v2}(t, \cdot, \varphi)) - E_0\| \geq \delta.$$

Proof. Suppose, by contradiction, that there exists some $\psi \in X_0(0)$ such that

$$\limsup_{t \rightarrow \infty} \|(I_1(t, \cdot, \psi), I_{v1}(t, \cdot, \psi), I_2(t, \cdot, \psi), I_{v2}(t, \cdot, \psi)) - E_0\| < \delta.$$

Then there exists a $t_1 > 0$ such that

$$0 < I_i(t, x, \psi) < \delta, \quad 0 < I_{vi}(t, x, \psi) < \delta, \quad i = 1, 2$$

for all $t \geq t_1$ and $x \in \bar{\Omega}$. Then $I_1(t, \cdot, \psi)$ and $I_{v1}(t, \cdot, \psi)$ satisfy

$$\left\{ \begin{array}{l} \frac{\partial I_1(t, x)}{\partial t} \geq D_h \Delta I_1(t, x) - (d + \gamma_1) I_1(t, x) \\ \quad + \frac{c_1 \beta(t, x) l(N(x) - 2\delta)}{2p\delta + lN(x)} I_{v1}(t, x), \quad t \geq t_1, x \in \Omega, \\ \frac{\partial I_{v1}(t, x)}{\partial t} \geq D_v \Delta I_{v1}(t, x) - \eta(t, x) I_{v1}(t, x) \\ \quad + \frac{\alpha_1 \beta(t, x) p(M^*(t, x) - 2\delta)}{2p\delta + lN(x)} I_1(t, x), \quad t \geq t_1, x \in \Omega, \\ \frac{\partial I_1(t, x)}{\partial \nu} = \frac{\partial I_{v1}(t, x)}{\partial \nu} = 0, \quad t \geq t_1, x \in \partial\Omega. \end{array} \right.$$

Let $P_1^\delta : X(0) \rightarrow X(0)$ be the Poincaré map associated with the following system:

$$\left\{ \begin{array}{l} \frac{\partial I_1(t, x)}{\partial t} = D_h \Delta I_1(t, x) - (d + \gamma_1) I_1(t, x) \\ \quad + \frac{c_1 \beta(t, x) l(N(x) - 2\delta)}{2p\delta + lN(x)} I_{v1}(t, x), \quad t \geq 0, x \in \Omega, \\ \frac{\partial I_{v1}(t, x)}{\partial t} = D_v \Delta I_{v1}(t, x) - \eta(t, x) I_{v1}(t, x) \\ \quad + \frac{\alpha_1 \beta(t, x) p(M^*(t, x) - 2\delta)}{2p\delta + lN(x)} I_1(t, x), \quad t \geq 0, x \in \Omega, \\ \frac{\partial I_1(t, x)}{\partial \nu} = \frac{\partial I_{v1}(t, x)}{\partial \nu} = 0, \quad t \geq 0, x \in \partial\Omega. \end{array} \right. \quad (4.3)$$

In view of Lemma 3.1, we have that $\mathcal{R}_1 > 1$ is equivalent to $r(P_1) > 1$. By continuity, we see that $\lim_{\delta \rightarrow 0} r(P_1^\delta) = r(P_1) > 1$. Thus, we can fix a sufficiently small number $\delta > 0$ such that

$$0 < \delta < \min\left\{ \min_{(t,x) \in [0, \omega] \times \bar{\Omega}} \frac{M^*(t, x)}{2}, \min_{x \in \bar{\Omega}} \frac{N(x)}{2} \right\} \text{ and } r(P_1^\delta) > 1.$$

Since P_1^δ is compact and strongly positive on $E(0)$, then Krein-Rutman Theorem implies that $r(P_1^\delta)$ is a simple eigenvalue of P_1^δ having a strongly positive eigenvector. It then follows from [37, Theorem 2.16 and Remark 2.20] that there is a positive ω -periodic function $\vartheta^\delta(t, x)$ such that $e^{\mu^\delta t} \vartheta^\delta(t, x)$ is a positive solution of (4.3), where $\mu^\delta = \frac{\ln r(P_1^\delta)}{\omega} > 0$. From Lemma 2.2, we know that

$$I_1(t, x, \psi) > 0 \text{ and } I_{v1}(t, x, \psi) > 0, \quad \forall t > 0, x \in \bar{\Omega}.$$

Thus, we may choose a $c > 0$ such that

$$(I_1(t_1, x, \psi), I_{v1}(t_1, x, \psi)) \geq ce^{\mu^\delta t_1} \vartheta^\delta(t_1, x), \quad \forall x \in \bar{\Omega}.$$

A simple comparison leads to

$$(I_1(t, \cdot, \psi), I_{v1}(t, \cdot, \psi)) \geq ce^{\mu^\delta t} \vartheta^\delta(t, x), \quad \forall t \geq t_1, x \in \bar{\Omega},$$

Since $\mu^\delta > 0$, it follows that

$$\lim_{t \rightarrow \infty} I_1(t, x, \psi) = \infty, \quad \lim_{t \rightarrow \infty} I_{v1}(t, x, \psi) = \infty \quad \text{uniformly for } x \in \bar{\Omega}.$$

By performing a similar analysis on $(I_2(t, x, \psi), I_{v2}(t, x, \psi))$, when $\mathcal{R}_2 > 1$,

$$\lim_{t \rightarrow \infty} I_2(t, x, \psi) = \infty, \quad \lim_{t \rightarrow \infty} I_{v2}(t, x, \psi) = \infty \quad \text{uniformly for } x \in \bar{\Omega}.$$

This contradicts the boundedness of I_i and I_{vi} , $i = 1, 2$. \square

Theorem 4.3. *Suppose that $\mathcal{R}_1 > 1$, $\mathcal{R}_2 > 1$, $\hat{\mathcal{R}}_1 > 1$, and $\hat{\mathcal{R}}_2 > 1$, then system (2.3) admits at least one positive ω -periodic solution, and there exists a constant $\delta^* > 0$ such that for any $\varphi \in X_0(0)$, we have*

$$\liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} I_i(t, x) \geq \delta^*, \quad \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} I_{vi}(t, x) \geq \delta^*, \quad i = 1, 2.$$

Proof. For any $\varphi \in X_0(0)$, by Lemma 2.2, we have

$$I_i(t, x, \varphi) > 0, \quad I_{vi}(t, x, \varphi) > 0, \quad i = 1, 2, \quad \forall t > 0, \quad x \in \bar{\Omega}.$$

Thus, $Q^n(X_0(0)) \subset X_0(0)$, $\forall n \in \mathbb{N}$. Furthermore, Q admits a global attractor on $X(0)$.

Next we prove that Q is uniformly persistent with respect to $(X_0(0), \partial X_0(0))$. Recalling the definitions of $E_0, E_1(t, x), E_2(t, x)$ in Section 3.2, we let

$$M_0 = E_0, \quad M_1 = E_1(0, \cdot), \quad M_2 = E_2(0, \cdot).$$

Then we have the following claims.

Claim 1. There is a $\delta_1 > 0$ such that

$$\limsup_{n \rightarrow \infty} \|Q^n(\varphi) - M_0\| \geq \delta_1, \quad \forall \varphi \in X_0(0).$$

This claim directly follows from Lemma 4.1.

Consider an auxiliary system with parameter ε :

$$\left\{ \begin{array}{l} \frac{\partial I_2(t, x)}{\partial t} = D_h \Delta I_2(t, x) - (d + \gamma_2) I_2(t, x) \\ \quad + \frac{c_2 \beta(t, x) l(N(x) - I_1^*(t, x) - 2\varepsilon)}{p(I_1^*(t, x) + 2\varepsilon) + l(N(x) - I_1^*(t, x) - 2\varepsilon)} I_{v2}(t, x), \\ \frac{\partial I_{v2}(t, x)}{\partial t} = D_v \Delta I_{v2}(t, x) - \eta(t, x) I_{v2}(t, x) \\ \quad + \frac{\alpha_2 \beta(t, x) p(M^*(t, x) - I_{v1}^*(t, x) - 2\varepsilon)}{p(I_1^*(t, x) + 2\varepsilon) + l(N(x) - I_1^*(t, x) - 2\varepsilon)} I_2(t, x), \\ \frac{\partial I_2(t, x)}{\partial \nu} = \frac{\partial I_{v2}(t, x)}{\partial \nu} = 0, \end{array} \right. \quad (4.4)$$

for all $t > 0$. Let $\hat{P}_2^\varepsilon := \hat{P}_2^\varepsilon(\omega)$ be the Poincaré map of (4.4). Since $\lim_{\varepsilon \rightarrow 0} r(\hat{P}_2^\varepsilon) = r(\hat{P}_2) > 1$, we can fix a small number $\varepsilon > 0$ such that $r(\hat{P}_2^\varepsilon) > 1$. As discussed in Lemma 4.1, there is

a positive ω -periodic function $\vartheta^\varepsilon(t, x)$ such that $e^{\mu^\varepsilon t} \vartheta^\varepsilon(t, x)$ is a positive solution of (4.4), where $\mu^\varepsilon = \frac{\ln r(\tilde{P}_2^\varepsilon)}{\omega} > 0$. For $\varepsilon > 0$ above, by the continuous dependence of solutions on the initial value, there exists $\delta_2 = \delta_2(\varepsilon) > 0$ such that for all $\varphi \in X_0(0)$ with $\|\varphi - M_1\| \leq \delta_2$, we have $\|Q(t)\varphi - Q(t)M_1\| < \varepsilon$, $\forall t \in [0, \omega]$.

Claim 2. $\limsup_{n \rightarrow \infty} \|Q^n(\varphi) - M_1\| \geq \delta_2$, $\forall \varphi \in X_0(0)$.

Suppose the claim is false, then $\limsup_{n \rightarrow \infty} \|Q^n(\psi) - M_1\| < \delta_2$ for some $\psi \in X_0(0)$. Then there exists an integer $N_1 \geq 1$ such that $\|Q^n(\psi) - M_1\| < \delta_2$ for $n \geq N_1$. For any $t \geq N_1\omega$, letting $t = n\omega + \tilde{t}$ with $n = [t/\omega]$ and $\tilde{t} \in [0, \omega)$, we have

$$\|Q(t)\psi - Q(t)M_1\| = \|Q(\tilde{t})(Q^n(\psi)) - Q(\tilde{t})M_1\| < \varepsilon.$$

According to the above inequality and Lemma 2.2, we infer that

$$\begin{aligned} 0 < I_1(t, x, \psi) < I_1^*(t, x, \psi) + \varepsilon, \quad 0 < I_{v1}(t, x, \psi) < I_{v1}^*(t, x, \psi) + \varepsilon, \\ 0 < I_2(t, x, \psi) < \varepsilon, \quad 0 < I_{v2}(t, x, \psi) < \varepsilon, \quad \forall t \geq N_1\omega, \quad x \in \bar{\Omega}. \end{aligned}$$

As a result, $I_2(t, x, \psi)$ and $I_{v2}(t, x, \psi)$ satisfy

$$\left\{ \begin{aligned} \frac{\partial I_2(t, x)}{\partial t} &\geq D_h \Delta I_2(t, x) - (d + \gamma_2) I_2(t, x) \\ &\quad + \frac{c_2 \beta(t, x) l(N(x) - I_1^*(t, x) - 2\varepsilon)}{p(I_1^*(t, x) + 2\varepsilon) + l(N(x) - I_1^*(t, x) - 2\varepsilon)} I_{v2}(t, x), \\ \frac{\partial I_{v2}(t, x)}{\partial t} &\geq D_v \Delta I_{v2}(t, x) - \eta(t, x) I_{v2}(t, x) \\ &\quad + \frac{\alpha_2 \beta(t, x) p(M^*(t, x) - I_{v1}^*(t, x) - 2\varepsilon)}{p(I_1^*(t, x) + 2\varepsilon) + l(N(x) - I_1^*(t, x) - 2\varepsilon)} I_2(t, x), \\ \frac{\partial I_2(t, x)}{\partial \nu} &= \frac{\partial I_{v2}(t, x)}{\partial \nu} = 0, \end{aligned} \right. \quad (4.5)$$

for all $t \geq N_1\omega$. Since $\psi \in X_0(0)$, $I_2(t, x, \psi) > 0$ and $I_{v2}(t, x, \psi) > 0$ for all $t \geq 0$ and $x \in \bar{\Omega}$, there exists a $k > 0$ such that

$$(I_2(N_1\omega, x, \psi), I_{v2}(N_1\omega, x, \psi)) \geq k e^{\mu^\varepsilon N_1\omega} \vartheta^\varepsilon(N_1\omega, x), \quad x \in \bar{\Omega}.$$

An application of the comparison theorem to (4.5) yields

$$(I_2(t, x, \psi), I_{v2}(t, x, \psi)) \geq k e^{\mu^\varepsilon t} \vartheta^\varepsilon(t, x), \quad \forall t \geq N_1\omega, \quad x \in \bar{\Omega}.$$

Since $\mu^\varepsilon > 0$, one sees that $I_2(t, \cdot, \psi) \rightarrow \infty$, $I_{v2}(t, \cdot, \psi) \rightarrow \infty$ as $t \rightarrow \infty$. This gives rise to a contradiction, and thereby, the above claim is true.

In a similar way, we can prove the following claim.

Claim 3. There exists a $\delta_3 > 0$ such that

$$\limsup_{n \rightarrow \infty} \|Q^n(\varphi) - M_2\| \geq \delta_3, \quad \forall \varphi \in X_0(0).$$

With the above three claims, we see that M_0, M_1 and M_2 are isolated invariant sets for Q in $X(0)$ and $W^s(M_i) \cap X_0(0) = \emptyset, i = 0, 1, 2$, where $W^s(M_i)$ is the stable set of M_i for Q . Set

$$M_\partial := \{\varphi \in \partial X_0(0) : Q^n(\varphi) \in \partial X_0(0), \forall n \in \mathbb{N}\}.$$

We now show that $M_\partial = \mathcal{M}_0$, where

$$\mathcal{M}_0 := \{\varphi \in \partial X_0(0) : \varphi_1(\cdot) = \varphi_2(\cdot) \equiv 0 \text{ or } \varphi_3(\cdot) = \varphi_4(\cdot) \equiv 0\}.$$

Obviously, it suffices to prove $M_\partial \subset \mathcal{M}_0$. For any given $\varphi \in M_\partial$, we have $I_1(n\omega, \cdot, \varphi) \equiv 0$ or $I_{v1}(n\omega, \cdot, \varphi) \equiv 0$ or $I_2(n\omega, \cdot, \varphi) \equiv 0$ or $I_{v2}(n\omega, \cdot, \varphi) \equiv 0, \forall n \in \mathbb{N}$. Assume that $\varphi \notin \mathcal{M}_0$, then there are eight possibilities as below:

- (i) $\varphi_1(\cdot) = I_1(0, \cdot, \varphi) \equiv 0, \varphi_2(\cdot) = I_{v1}(0, \cdot, \varphi) > 0, \varphi_3(\cdot) = I_2(0, \cdot, \varphi) \equiv 0, \varphi_4(\cdot) = I_{v2}(0, \cdot, \varphi) > 0.$
- (ii) $\varphi_1(\cdot) = I_1(0, \cdot, \varphi) \equiv 0, \varphi_2(\cdot) = I_{v1}(0, \cdot, \varphi) > 0, \varphi_3(\cdot) = I_2(0, \cdot, \varphi) > 0, \varphi_4(\cdot) = I_{v2}(0, \cdot, \varphi) \equiv 0.$
- (iii) $\varphi_1(\cdot) = I_1(0, \cdot, \varphi) \equiv 0, \varphi_2(\cdot) = I_{v1}(0, \cdot, \varphi) > 0, \varphi_3(\cdot) = I_2(0, \cdot, \varphi) > 0, \varphi_4(\cdot) = I_{v2}(0, \cdot, \varphi) > 0.$
- (iv) $\varphi_1(\cdot) = I_1(0, \cdot, \varphi) > 0, \varphi_2(\cdot) = I_{v1}(0, \cdot, \varphi) \equiv 0, \varphi_3(\cdot) = I_2(0, \cdot, \varphi) \equiv 0, \varphi_4(\cdot) = I_{v2}(0, \cdot, \varphi) > 0.$
- (v) $\varphi_1(\cdot) = I_1(0, \cdot, \varphi) > 0, \varphi_2(\cdot) = I_{v1}(0, \cdot, \varphi) \equiv 0, \varphi_3(\cdot) = I_2(0, \cdot, \varphi) > 0, \varphi_4(\cdot) = I_{v2}(0, \cdot, \varphi) \equiv 0.$
- (vi) $\varphi_1(\cdot) = I_1(0, \cdot, \varphi) > 0, \varphi_2(\cdot) = I_{v1}(0, \cdot, \varphi) \equiv 0, \varphi_3(\cdot) = I_2(0, \cdot, \varphi) > 0, \varphi_4(\cdot) = I_{v2}(0, \cdot, \varphi) > 0.$
- (vii) $\varphi_1(\cdot) = I_1(0, \cdot, \varphi) > 0, \varphi_2(\cdot) = I_{v1}(0, \cdot, \varphi) > 0, \varphi_3(\cdot) = I_2(0, \cdot, \varphi) \equiv 0, \varphi_4(\cdot) = I_{v2}(0, \cdot, \varphi) > 0.$
- (viii) $\varphi_1(\cdot) = I_1(0, \cdot, \varphi) > 0, \varphi_2(\cdot) = I_{v1}(0, \cdot, \varphi) > 0, \varphi_3(\cdot) = I_2(0, \cdot, \varphi) > 0, \varphi_4(\cdot) = I_{v2}(0, \cdot, \varphi) \equiv 0.$

By Lemma 2.2, in case (i), we obtain that $I_{v1}(t, x, \varphi) > 0, I_{v2}(t, x, \varphi) > 0$ for all $t > 0$ and $x \in \bar{\Omega}$. Further, using the first and third equation of (2.3), one obtains that $I_1(t, x, \varphi) > 0, I_2(t, x, \varphi) > 0, \forall t > 0, x \in \bar{\Omega}$, which contradicts with the fact $\varphi \in M_\partial$. By performing a similar analysis, we can show that (ii)-(viii) are impossible. Hence, $\varphi \in \mathcal{M}_0$, and hence, $M_\partial \subset \mathcal{M}_0$. This proves $M_\partial = \mathcal{M}_0$.

Let $\omega(\varphi)$ be the omega limit set of the forward orbit $\gamma^+(\varphi) := \{Q^n(\varphi) : \forall n \in \mathbb{N}\}$. We further have the following claims.

Claim 4. $\cup_{\varphi \in M_\partial} \omega(\varphi) \subset \{M_0, M_1, M_2\}.$

Obviously, there are three possibilities for $\varphi \in M_\partial = \mathcal{M}_0$:

Case 1 : $\varphi_1(\cdot) = \varphi_2(\cdot) \equiv 0$, $\varphi_3(\cdot) > 0$ or $\varphi_4(\cdot) > 0$;

Case 2 : $\varphi_3(\cdot) = \varphi_4(\cdot) \equiv 0$, $\varphi_1(\cdot) > 0$ or $\varphi_2(\cdot) > 0$;

Case 3 : $\varphi_1(\cdot) = \varphi_2(\cdot) = \varphi_3(\cdot) = \varphi_4(\cdot) \equiv 0$.

In what follows, we aim to show that claim 4 holds for each of the above three cases.

If **Case 1** happens, then $I_1(t, x, \varphi) = I_{v1}(t, x, \varphi) \equiv 0$ for all $t \geq 0$ and $x \in \bar{\Omega}$. In view of system (2.3), $I_2(t, x, \varphi), I_{v2}(t, x, \varphi)$ satisfy system (3.1) with $i = 2, j = 1$. Since $\mathcal{R}_2 > 1$, it follows from Theorem 3.1 that

$$\lim_{t \rightarrow \infty} (I_2(t, x, \varphi) - I_2^*(t, x)) = 0, \quad \lim_{t \rightarrow \infty} (I_{v2}(t, x, \varphi) - I_{v2}^*(t, x)) = 0$$

uniformly for $x \in \bar{\Omega}$. Hence, $\omega(\varphi) = M_2$ for any $\varphi \in M_\partial$.

For **Case 2**, by repeating arguments similar to **Case 1**, we can show that $\omega(\varphi) = M_1$ for any $\varphi \in M_\partial$. For **Case 3**, one immediately finds that

$$(I_1(t, x, \varphi), I_{v1}(t, x, \varphi), I_2(t, x, \varphi), I_{v2}(t, x, \varphi)) = (0, 0, 0, 0), \quad \forall t \geq 0, x \in \bar{\Omega}.$$

This implies that $\omega(\varphi) = M_0$ for any $\varphi \in M_\partial$. Thus claim 4 is obtained.

Claim 5. M_1 and M_2 are locally stable, and M_0 is unstable for Q in M_∂ .

Suppose that $\varphi \in M_\partial$, we have that $M_\partial = M_\partial^1 \cup M_\partial^2$, where

$$M_\partial^1 := \{\varphi \in \partial X_0(0) : \varphi_1(\cdot) = \varphi_2(\cdot) \equiv 0\}, \quad M_\partial^2 := \{\varphi \in \partial X_0(0) : \varphi_3(\cdot) = \varphi_4(\cdot) \equiv 0\}.$$

If $\varphi \in M_\partial^1$, then system (2.3) restricted on M_∂^1 is a monotone system. Thus, M_2 is locally Lyapunov stable for Q in M_∂^1 due to [26, Lemma 2.2.1], and M_0 is unstable in M_∂^1 . In a similar manner, if $\varphi \in M_\partial^2$, we can prove that M_1 is locally Lyapunov stable for Q in M_∂^2 , and M_0 is unstable in M_∂^2 .

The claim 5 implies that no subset of $\{M_0, M_1, M_2\}$ forms a cycle in $\partial X_0(0)$. Based on the above analysis, it follows from the acyclicity theorem on uniform persistence for maps [26, Theorem 1.3.1 and Remark 1.3.1] that $Q : X(0) \rightarrow X(0)$ is uniformly persistent with respect to $(X_0(0), \partial X_0(0))$ in the sense that there exists $\tilde{\delta} > 0$ such that

$$\liminf_{n \rightarrow \infty} d(Q^n(\varphi), \partial X_0(0)) \geq \tilde{\delta}, \quad \forall \varphi \in X_0(0).$$

By [30, Theorem 4.5] with $\rho(\phi) = d(\phi, \partial X_0(0))$, Q admits a global attractor A_0 in $X_0(0)$, and Q has a fixed point in $\varphi^* \in A_0$. Clearly, $u(t, \cdot, \varphi^*)$ is an ω -periodic solution of (2.3) and it is strictly positive due to Lemma 2.2.

Finally, we use the arguments in [26, Section 11.2] to obtain the practical uniform persistence. Since $A_0 = Q(\omega)A_0 = Q(A_0)$, we have $\varphi_i > 0, i = 1, 2, 3, 4, \forall \varphi \in A_0$. Let $B_0 := \cup_{t \in [0, \omega]} Q(t)A_0$. Then [26, Theorem 3.1.1] implies that $B_0 \subset X_0(0)$, and $\lim_{t \rightarrow \infty} d(Q(t)\varphi, B_0) = 0$ for all $\varphi \in X_0(0)$. Define a continuous function $p : X(0) \rightarrow \mathbb{R}_+$ by

$$p(\varphi) := \min\{\min_{x \in \bar{\Omega}} \varphi_1(x), \min_{x \in \bar{\Omega}} \varphi_2(x), \min_{x \in \bar{\Omega}} \varphi_3(x), \min_{x \in \bar{\Omega}} \varphi_4(x)\}, \quad \forall \varphi \in X(0).$$

Since B_0 is compact, it follows that $\inf_{\varphi \in B_0} p(\varphi) = \min_{\varphi \in B_0} p(\varphi) > 0$. Therefore, there exists a $\delta^* > 0$ such that for any $\varphi \in X_0(0)$,

$$\liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} I_i(t, x, \varphi) \geq \delta^*, \quad \liminf_{t \rightarrow \infty} \min_{x \in \bar{\Omega}} I_{vi}(t, x, \varphi) \geq \delta^*, \quad i = 1, 2.$$

The proof is complete. \square

5 Asymptotic behavior of \mathcal{R}_0

In this section, we use the recent theory developed in [38] to study the asymptotic behavior of the basic reproduction number as the diffusion coefficients go to zero and infinity. To do this, we write

$$\mathcal{R}_0(D_h, D_v) = \max\{\mathcal{R}_1(D_h, D_v), \mathcal{R}_2(D_h, D_v)\}.$$

Observe that for each $x \in \bar{\Omega}$, the equation

$$\frac{\partial M(t, x)}{\partial t} = \Lambda(t, x) - \eta(t, x)M(t, x)$$

admits a globally stable positive ω -periodic solution $M_0(t, x)$, and it is continuous on $\mathbb{R} \times \bar{\Omega}$. Define $\tilde{g}(t) := |\Omega|^{-1} \int_{\Omega} g(t, x) dx$. One immediately sees that the following scalar periodic equation

$$\frac{\partial M(t, x)}{\partial t} = \tilde{\Lambda}(t) - \tilde{\eta}(t)M(t, x)$$

has a unique positive ω -periodic solution

$$\tilde{M}_{\infty}(t) = \left[\int_0^t \tilde{\Lambda}(s) e^{\int_0^s \tilde{\eta}(\xi) d\xi} ds + \frac{\int_0^{\omega} \tilde{\Lambda}(s) e^{\int_0^s \tilde{\eta}(\xi) d\xi} ds}{e^{\int_0^{\omega} \tilde{\eta}(s) ds} - 1} \right] e^{-\int_0^t \tilde{\eta}(s) ds},$$

which is globally asymptotically stable.

It is easy to verify that assumptions (H1)-(H5) in [38] are valid. An direct application of [38, Theorems 5.2 and 5.5] leads to

$$\lim_{D_v \rightarrow 0} \|M^*(t, \cdot) - M_0(t, \cdot)\|_{\mathbb{Y}} = 0, \quad \lim_{D_v \rightarrow \infty} \|M^*(t, \cdot) - \tilde{M}_{\infty}(t)\|_{\mathbb{Y}} = 0$$

hold uniformly on $t \in \mathbb{R}$. For each $x \in \bar{\Omega}$, let $\{\Gamma_{x,0}^i(t, s) : t \geq s\}$ ($i = 1, 2$) be the evolution family on \mathbb{R}^2 associated with the following system:

$$\begin{cases} \frac{\partial I_i(t, x)}{\partial t} = -(d + \gamma_i)I_i(t, x), & t > s, x \in \bar{\Omega}, \\ \frac{\partial I_{vi}(t, x)}{\partial t} = -\eta(t, x)I_{vi}(t, x), & t > s, x \in \bar{\Omega}, \end{cases}$$

and define

$$F_0^i(t, x) \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = \begin{pmatrix} c_i \beta(t, x) \psi_2 \\ \frac{\alpha_i \beta(t, x) p M_0(t, x)}{l N(x)} \psi_1 \end{pmatrix}, \quad \forall t \in \mathbb{R}, x \in \bar{\Omega}, \psi = (\psi_1, \psi_2) \in \mathbb{R}^2.$$

Let $\{\tilde{\Gamma}_\infty^i(t, s) : t \geq s\}$ ($i = 1, 2$) be the evolution family on \mathbb{R}^2 of the following system:

$$\begin{cases} \frac{\partial I_i(t, x)}{\partial t} = -(d + \gamma_i)I_i(t, x), & t > s, \ x \in \bar{\Omega}, \\ \frac{\partial I_{v2}(t, x)}{\partial t} = -\tilde{\eta}(t)I_{vi}(t, x), & t > s, \ x \in \bar{\Omega}, \end{cases}$$

and define

$$\tilde{F}_\infty^i(t) \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix} = \begin{pmatrix} c_i \tilde{\beta}(t) \psi_2 \\ \tilde{f}_{21}^i(t) \tilde{M}_\infty(t) \psi_1 \end{pmatrix}, \quad \forall t \in \mathbb{R}, \ \psi = (\psi_1, \psi_2) \in \mathbb{R}^2,$$

where

$$\tilde{f}_{21}^i(t) := |\Omega|^{-1} \int_{\Omega} \frac{\alpha_i \beta(t, x) p}{lN(x)} dx, \quad \forall t \in \mathbb{R}.$$

Let $C_\omega(\mathbb{R}, \mathbb{R}^2)$ be the Banach space of all continuous and ω -periodic functions from \mathbb{R} to \mathbb{R}^2 , which is endowed with the maximum norm. For each $x \in \bar{\Omega}$, we respectively define bounded linear positive operators $L_{x,0}^i$ and \tilde{L}_∞^i , $i = 1, 2$, on $C_\omega(\mathbb{R}, \mathbb{R}^2)$ by

$$[L_{x,0}^i v](t) := \int_0^\infty \Gamma_{x,0}^i(t, t-s) F_0^i(t-s, x) v(t-s) ds, \quad \forall t \in \mathbb{R}, \ v \in C_\omega(\mathbb{R}, \mathbb{R}^2),$$

and

$$[\tilde{L}_\infty^i v](t) := \int_0^\infty \tilde{\Gamma}_\infty^i(t, t-s) \tilde{F}_\infty^i(t-s) v(t-s) ds, \quad \forall t \in \mathbb{R}, \ v \in C_\omega(\mathbb{R}, \mathbb{R}^2).$$

Then we define $R_i(x, 0) := r(L_{x,0}^i)$, $\forall x \in \bar{\Omega}$, $i = 1, 2$ and $\tilde{R}_i(\infty) := r(\tilde{L}_\infty^i)$, $i = 1, 2$. By [38, Theorem 4.1] with $\kappa = \text{diag}(D_h, D_v)$, $\chi = D_v$ and $\chi_0 = 0$, and $\kappa = \text{diag}(D_h, D_v)$, $\chi = \frac{1}{D_v}$ and $\chi_0 = 0$, respectively, it follows that

$$\lim_{\max(D_h, D_v) \rightarrow 0} \mathcal{R}_i(D_h, D_v) = \max_{x \in \bar{\Omega}} R_i(x, 0),$$

$$\lim_{\min(D_h, D_v) \rightarrow \infty} \mathcal{R}_i(D_h, D_v) = \tilde{R}_i(\infty), \quad i = 1, 2.$$

Therefore,

$$\lim_{\max(D_h, D_v) \rightarrow 0} \mathcal{R}_0(D_h, D_v) = \max_{x \in \bar{\Omega}} \{R_1(x, 0), R_2(x, 0)\},$$

$$\lim_{\min(D_h, D_v) \rightarrow \infty} \mathcal{R}_0(D_h, D_v) = \max\{\tilde{R}_1(\infty), \tilde{R}_2(\infty)\}.$$

6 Numerical simulations

To verify these analytic results and examine the effects of seasonality and vector-bias on the malaria transmission, we perform illustrative numerical investigations.

6.1 Competitive exclusion and coexistence

We choose the period of our model to be $T = 12$ months and concentrate on one dimensional domain $\Omega = [0, \pi]$. For illustrative purpose, we only let $\beta(t, x)$ be the time-dependent parameters, given by

$$\begin{aligned}\beta(t, x) = & 4 \times (5.1492 - 1.83692 \cos(0.523599t) - 0.175817 \cos(1.0472t) \\ & - 0.166233 \cos(1.5708t) - 0.16485 \cos(2.0944t) - 0.17681 \cos(2.61799t) \\ & - 1.37079 \sin(0.523599t) + 0.296267 \sin(1.0472t) + 0.2134 \sin(1.5708t) \\ & - 0.295228 \sin(2.0944t) - 0.201712 \sin(2.61799t)) \text{ month}^{-1},\end{aligned}$$

which is adapted from [8]. Unless stated otherwise, the baseline parameters are seen in Table 1. We use the numerical scheme proposed in [34, Lemma 2.5 and Remark 3.2] to compute the reproduction number of each strain. In order to demonstrate the outcomes of competitive exclusion and coexistence, we consider the following three cases.

Table 1: Parameter values.

Parameter	Value (range)	Dimension	Reference
$N(x)$	110	dimensionless	[12]
$M^*(t, x)$	220	dimensionless	[24]
d	$1/(72 \times 12)$	month^{-1}	[12]
$\eta(t, x)$	0.8	month^{-1}	[24]
D_h	0.4	$\text{km}^2 \cdot \text{month}^{-1}$	[12]
D_v	0.02	$\text{km}^2 \cdot \text{month}^{-1}$	[12]
p	0.8 (0,1)	dimensionless	[12]
l	0.2 (0,1)	dimensionless	[12]

Case 1. $\mathcal{R}_1 > 1, \mathcal{R}_2 > 1, \hat{\mathcal{R}}_1 > 1$ and $\hat{\mathcal{R}}_2 > 1$. We choose $\gamma_1 = 0.096 \text{ month}^{-1}, \gamma_2 = 0.082 \text{ month}^{-1}, \alpha_1 = 0.56, \alpha_2 = 0.6, c_1 = 0.25, c_2 = 0.2$. Then we obtain $\mathcal{R}_1 = 11.1267, \mathcal{R}_2 = 10.3022, \hat{\mathcal{R}}_1 = 2.2919$, and $\hat{\mathcal{R}}_2 = 2.2605$. Fig. 1 shows that the disease is uniformly persistent, and periodic oscillation phenomenon occurs, which is consistent with Theorem 4.3.

Case 2. $\mathcal{R}_1 > 1, \mathcal{R}_2 > 1, \hat{\mathcal{R}}_1 > 1$ and $\hat{\mathcal{R}}_2 < 1$. We choose $\gamma_1 = 0.083 \text{ month}^{-1}, \gamma_2 = 0.082 \text{ month}^{-1}, \alpha_1 = 0.35, \alpha_2 = 0.55, c_1 = 0.2, c_2 = 0.1$. Then we have $\mathcal{R}_1 = 7.8683, \mathcal{R}_2 = 6.9746, \hat{\mathcal{R}}_1 = 1.0934, \hat{\mathcal{R}}_2 = 0.4729$. Fig. 2 shows that the sensitive strains are persistent, but the resistant strains die out.

Case 3. $\mathcal{R}_1 > 1, \mathcal{R}_2 > 1, \hat{\mathcal{R}}_1 < 1$ and $\hat{\mathcal{R}}_2 > 1$. We choose $\gamma_1 = 0.096 \text{ month}^{-1}, \gamma_2 = 0.082 \text{ month}^{-1}, \alpha_1 = 0.55, \alpha_2 = 0.45, c_1 = 0.15, c_2 = 0.2$. Then we get $\mathcal{R}_1 = 8.5413, \mathcal{R}_2 = 8.9219, \hat{\mathcal{R}}_1 = 0.7052, \hat{\mathcal{R}}_2 = 1.9601$. Fig. 3 depicts that the resistant strains persist, but the sensitive strains go extinct.




Figure 1: Two strains coexist: (a) the sensitive strains I_1, I_{v1} , (b) The resistant strains I_2, I_{v2} . The initial data are chosen as $I_1(0, x) = 6(1 + \cos(2x))$, $I_{v1}(0, x) = 10(1 + \cos(2x))$, $I_2(0, x) = 5(1 + \cos(2x))$, $I_{v2}(0, x) = 8(1 + \cos(2x))$, $\forall x \in [0, \pi]$.




Figure 2: (a) I_1 and I_{v1} persist; (b) I_2 and I_{v2} die out. The initial data are the same as Fig. 1.

It should be pointed out that in Figs. 1-3, we only plot the graph of x -intersection with $x = 0.7448$. In addition, for the second and third case, the competitive exclusion phenomena are also observed even though $\mathcal{R}_0 > 1$. It is a pity that we now can not prove it, which is left for future consideration.

6.2 Effects of parameters on \mathcal{R}_0

In order to explore the effect of seasonality, we set the biting rate $\beta(t) \approx a_0(1 - b_0 \cos(0.523599t))$, where a_0 is the average biting rate, and $b_0 \in [0, 1]$ is the strength of seasonal forcing. We




Figure 3: (a) I_1 and I_{v1} die out; (b) I_2 and I_{v2} persist. The initial data are the same as Fig. 1.

use the same parameter values as in Case 1 in Section 6.1. Fig. 4 describes the dependence of \mathcal{R}_0 on a_0 and b_0 . More precisely, Fig. 4(a) shows that \mathcal{R}_0 is an increasing function of a_0 for fixed b_0 . Fig. 4(b) compares the influences of the time-dependent biting rate and the time-averaged biting rate on \mathcal{R}_0 . As can be seen in Fig. 4(b), \mathcal{R}_0 increases as b_0 increases. This implies that the use of the time-averaged biting rate may underestimate the risk of disease transmission. It should be emphasized that this phenomenon is not observed in all malaria models, which is dependent on model parameters.




Figure 4: The effect of seasonality on \mathcal{R}_0 . (a) \mathcal{R}_0 as a function of a_0 when $b_0 = 0.35674$; (b) \mathcal{R}_0 as a function of b_0 when $a_0 = 5.1492$.

Next, we investigate the vector-bias effect. We use $q := l/p$ to measure the relative attractivity of susceptible host versus infection one. Our numerical result in Fig. 5 shows

that \mathcal{R}_0 decreases as q increases, which indicates that the ignorance of the vector-bias effect will underestimate the value of \mathcal{R}_0 . In fact, we can analytically prove the monotonicity of




Figure 5: The effect of vector-bias on \mathcal{R}_0 .

\mathcal{R}_0 with respect to q . Let A^i and B^i ($i = 1, 2$) be two bounded linear operators on $C_\omega(\mathbb{R}, \mathbb{E})$ given by

$$[A^i v](t) := \int_0^\infty \Psi_i(t, t-s)v(t-s)ds, \quad [B^i v](t) := \mathcal{F}_i(t)v, \quad \forall t \in \mathbb{R}, v \in C_\omega(\mathbb{R}, \mathbb{E}),$$

where Ψ_i and $\mathcal{F}_i(t)$ are defined as in Section 3. Inspired by Section 4.2 in [37], we write

$$A^i v = (A_1^i v_1, A_2^i v_2), \text{ and } B^i v = (B_1^i v_2, B_2^i v_1), \quad \forall v = (v_1, v_2) \in C_\omega(\mathbb{R}, \mathbb{E}),$$

where

$$\begin{aligned} [A_1^i v_1](t) &= \int_0^\infty T_i(t, t-s)v_1(t-s)ds, \quad [A_2^i v_2](t) = \int_0^\infty T_3(t, t-s)v_2(t-s)ds, \\ [B_1^i v_2](t) &= c_i \beta(t, \cdot)v_2(\cdot), \quad [B_2^i v_1](t) = \frac{\alpha_i \beta(t, \cdot) p M^*(\cdot)}{l N(\cdot)} v_1(\cdot), \quad i = 1, 2. \end{aligned}$$

According to Section 3.1, $\mathcal{L}_i(q) = A^i B^i v = (A_1^i B_1^i v_2, A_2^i B_2^i v_1)$, it then follows that

$$\mathcal{L}_i^2(q)v = (A_1^i B_1^i A_2^i B_2^i v_1, A_2^i B_2^i A_1^i B_1^i v_2) = \frac{1}{q} \mathcal{L}_i^2(1)v, \quad i = 1, 2,$$

and hence, $\mathcal{L}_i^2(q) = \frac{1}{q} \mathcal{L}_i^2(1)$. In view of $r^2(\mathcal{L}_i(q)) = r(\mathcal{L}_i^2(q))$, we obtain

$$\mathcal{R}_i(q) := r(\mathcal{L}_i(q)) = \frac{1}{\sqrt{q}} r(\mathcal{L}_i(1)) = \frac{1}{\sqrt{q}} \mathcal{R}_i(1), \quad i = 1, 2.$$

Therefore, $\mathcal{R}_0(q) = \max\{\mathcal{R}_1(q), \mathcal{R}_2(q)\} = \frac{1}{\sqrt{q}} \max\{\mathcal{R}_1(1), \mathcal{R}_2(1)\}$. This supports our numerical finding.

7 Discussion

In this paper, we have proposed a two-strain malaria model with seasonality and vector-bias. It is of interest to note that our model is a competitive system for sensitive and resistant strains, but the corresponding subsystem of each strain is cooperative. To characterize this mathematical structure, we define a time-dependent region $X(t)$. Although the introduction of time-varying region brings out some mathematical difficulties, the solution map $Q(t) : X(0) \rightarrow X(t)$ is an ω -periodic semiflow. This nice property makes us use uniform persistence theory for model dynamics. Our results show that the zero solution is global attractiveness if $\mathcal{R}_0 = \max\{\mathcal{R}_1, \mathcal{R}_2\} < 1$ (see Theorem 4.1); sensitive (resistant) strains are uniformly persistent if $\mathcal{R}_1 > 1 > \mathcal{R}_2$ ($\mathcal{R}_2 > 1 > \mathcal{R}_1$) (see Theorem 4.2); and the model is uniformly persistent and admits a positive periodic solution if $\mathcal{R}_1 > 1, \mathcal{R}_2 > 1, \hat{\mathcal{R}}_1 > 1$ and $\hat{\mathcal{R}}_2 > 1$ (see Theorem 4.3). We also have analyzed the asymptotic behavior of the basic reproduction number with small and large diffusion coefficients. Numerically, we have demonstrated the long-time behaviors of solutions: competitive exclusion and coexistence, and revealed the influences of some key parameters on the basic reproduction number. It is found that \mathcal{R}_0 increases as the strength of seasonal forcing increases, but it is a decreasing function of the relative attractivity of susceptible host versus infection one.

Finally, we mention that under certain condition, system (2.3) is a monotone system with respect to the partial order \leq_K , which is induced by the cone $K = \mathbb{E}^+ \times (-\mathbb{E}^+)$. Hence, if we can prove the uniqueness of positive periodic solution in Theorem 4.3, then the positive periodic solution is globally attractive in $X(0) \setminus \{0\}$ by the virtue of the theory of monotone systems. This is a challenging problem and left for future study.

References

- [1] J. B. Gutierrez, M. R. Galinski, S. Cantrell, et al., From within host dynamics to the epidemiology of infectious disease: Scientific overview and challenges, *Math. Biosci.* 270 (2015) 143–155.
- [2] F. Forouzannia, A. B. Gumel, Mathematical analysis of an age-structured model for malaria transmission dynamics, *Math. Biosci.* 247 (2014) 80–94.
- [3] The World Health Report 2020. Website: <https://www.who.int/news/item/30-11-2020-who-calls-for-reinvigorated-action-to-fight-malaria>.
- [4] R. Ross, The prevention of malaria, 2nd edition, Murray, London, 1911.
- [5] G. Macdonald, The epidemiology and control of malaria, Oxford University Press, London, 1957.
- [6] C. Cosner, J. C. Beier, R. S. Cantrell, et al., The effects of human movement on the persistence of vector-borne diseases, *J. Theor. Biol.* 258 (2009) 550–560.

- [7] F. Chamchod, N. F. Britton, Analysis of a vector-bias model on malaria transmission, *Bull. Math. Biol.* 73 (2011) 639–657.
- [8] Y. Lou, X.-Q. Zhao, A climate-based malaria transmission model with structured vector population, *SIAM J. Appl. Math.* 70 (2010) 2023–2044.
- [9] Y. Lou, X.-Q. Zhao, A reaction-diffusion malaria model with incubation period in the vector population, *J. Math. Biol.* 62 (2011) 543–568.
- [10] Y. Xiao, X. Zou, Transmission dynamics for vector-borne diseases in a patchy environment. *J. Math. Biol.* 69 (2014) 113–146.
- [11] X. Wang, X.-Q. Zhao, A periodic vector-bias malaria model with incubation period, *SIAM J. Appl. Math.* 77 (2017) 181–201.
- [12] Z. Bai, R. Peng, X.-Q. Zhao, A reaction-diffusion malaria model with seasonality and incubation period, *J. Math. Biol.* 77 (2018) 201–228.
- [13] R. Wu, X.-Q. Zhao, A reaction-diffusion model of vector-borne disease with periodic delays, *J. Nonlinear Sci.* 29 (2019) 29–64.
- [14] B.-G. Wang, L. Qiang, Z.-C. Wang, An almost periodic Ross-Macdonald model with structured vector population in a patchy environment, *J. Math. Biol.* 80 (2020) 835–863.
- [15] J. G. Kingsolver, Mosquito host choice and the epidemiology of malaria, *Am. Nat.* 130 (1987) 811–827.
- [16] G. R. Hosack, P. A. Rossignol, P. van den Driessche, The control of vector-borne disease epidemics, *J. Theor. Biol.* 255 (2008) 16–25.
- [17] S. J. Aneke, Mathematical modelling of drug resistant malaria parasites and vector populations, *Math. Methods Appl. Sci.* 25 (2002) 335–346.
- [18] E. Y. Klein, Antimalarial drug resistance: a review of the biology and strategies to delay emergence and spread, *Int. J. Antimicrob. Ag.* 41 (2013) 311–317.
- [19] F. B. Augusto, A. B. Gumel, P. E. Parham, Qualitative assessment of the role of temperature variations on malaria transmission dynamics, *J. Biol. Syst.* 23 (2015) 1–34.
- [20] P. Cailly, A. Tran, T. Balenghiene, et al., A climate-driven abundance model to assess mosquito control strategies, *Ecol. Model.* 227 (2012) 7–17.
- [21] M. B. Hoshen, A. P. Morse, A weather-driven model of malaria transmission, *Malaria J.* 3 (2004), article 32.

- [22] D. A. Ewing, C. A. Cobbold, B. V. Purse, et al., Modelling the effect of temperature on the seasonal population dynamics of temperate mosquitoes, *J. Theor. Biol.* 400 (2016) 65–79.
- [23] S. Altizer, A. Dobson, P. Hosseini, et al., Seasonality and the dynamics of infectious diseases, *Ecol. Lett.* 9 (2006) 467–484.
- [24] Y. Shi, H. Zhao, Analysis of a two-strain malaria transmission model with spatial heterogeneity and vector-bias, *J. Math. Biol.* 82 (2021) 24.
- [25] F. Li, X.-Q. Zhao, Global dynamics of a reaction-diffusion model of Zika virus transmission with seasonality, *Bull. Math. Biol.* 83 (2021) 43.
- [26] X.-Q. Zhao, *Dynamical systems in population biology*, 2nd edition, Springer, New York, 2017.
- [27] L. Zhang, Z.-C. Wang, X.-Q. Zhao, Threshold dynamics of a time periodic reaction-diffusion epidemic model with latent period, *J. Differential Equations* 258 (2015) 3011–3036.
- [28] D. Daners, P. K. Medina, *Abstract Evolution Equations, Periodic Problems and Applications*, Pitman Research Notes in Mathematics Series, vol. 279, Longman Scientific and Technical, Harlow, UK, 1992.
- [29] R. H. Martin, H. L. Smith, Abstract functional differential equations and reaction-diffusion systems, *Trans. Amer. Math. Soc.* 321 (1990) 1–44.
- [30] P. Magal, X.-Q. Zhao, Global attractors and steady states for uniformly persistent dynamical systems, *SIAM J. Math. Anal.* 37 (2005) 251–275.
- [31] P. Hess, *Periodic-Parabolic Boundary Value Problems and Positivity*, Pitman Research Notes in Mathematics Series, vol. 247, Longman Scientific and Technical, Harlow, UK, 1991.
- [32] H. R. Thieme, Spectral bound and reproduction number for infinite-dimensional population structure and time heterogeneity, *SIAM J. Appl. Math.* 70 (2009) 188–211.
- [33] X.-Q. Zhao, Basic reproduction ratios for periodic compartmental models with time delay, *J. Dynam. Differential Equations* 29 (2017) 67–82.
- [34] X. Liang, L. Zhang, X.-Q. Zhao, Basic reproduction ratios for periodic abstract functional differential equations (with application to a spatial model for Lyme disease), *J. Dynam. Differential Equations* 31 (2019) 1247–1278.
- [35] N. Bacaër, S. Guernaoui, The epidemic threshold of vector-borne diseases with seasonality, *J. Math. Biol.* 53 (2006) 421–436.

- [36] N. Tuncer, M. Martcheva, Analytical and numerical approaches to coexistence of strains in a two-strain SIS model with diffusion, *J. Biol. Dyn.* 6 (2012) 406–439.
- [37] X. Liang, L. Zhang, X.-Q. Zhao, The principal eigenvalue for degenerate periodic reaction-diffusion systems, *SIAM J. Math. Anal.* 49 (2017) 3603–3636.
- [38] L. Zhang, X.-Q. Zhao, Asymptotic behavior of the basic reproduction ratio for periodic reaction-diffusion systems, *SIAM J. Math. Anal.* 53 (2021) 6873–6909.