

N-qubit states with maximum entanglement across all bipartitions: A graph state approach

Sowrabh Sudevan* and Sourin Das†

Indian Institute of Science Education and Research Kolkata
Mohanpur, Nadia-741 246, West Bengal, India

(Dated: January 19, 2022)

We discuss the construction of N -qubit pure states with maximum bipartite entanglement across all possible choices of k vs $N-k$ bi-partitioning, which implies that the Von Neumann entropy of any k -qubit reduced density matrix corresponding to this state should be $k \ln 2$. We name them k -entangled states. We show that a subset of the 'graph states' satisfy this condition hence providing a recipe for constructing k -entangled states. Finding recipes for construction of k -entangled states using graph states is useful as it paves the way to identification of unitary transformations which when apply to a product state will generate these k -entangled states since such unitary transformations are known for all *graph states*. Though, a priori it is not clear how to construct a graph which corresponds to an arbitrary k -entangled state, but in particular, we show that graphs organized as a circular linear chain corresponds to the case of 2-entangled state, where we show that the minimum number of qubits required to host such a state is $N = 5$. 3-entangled states can be constructed by forming bilayer graphs with $\frac{N}{2}$ qubits ($N = 2Z$) in each layer, such that each layer forms a fully connected graph while inter-layer connections are such that the vertices in one layer has a one to one connectivity to the other layer. 4-entangled states can be formed by taking 2D lattice graphs(popularly referred to as a 2D cluster state) with periodic boundary conditions along both dimensions and both dimensions having at least 5 vertices.

I. INTRODUCTION:

The physical phenomena of entanglement was discussed by Einstein, Podolsky, and Rosen in the famous paper Ref. 1 in 1935. Since then, there has been extensive discussions of various measures of entanglement [2–4]. The entanglement measure this paper uses is bipartite entanglement. We are interested in constructing qubit states that maximise the bipartite entanglement across all bipartitions of given number of qubits. Page [5] showed that the most typical large qubit state has very high entanglement across the largest bipartition, That is, it is trivial to see high bipartite entanglement for large N qubit systems. Our work focus on the opposite limit of finite number of qubits. We demonstrates four specific pure state constructions that give us different amounts of entanglement across many different system sizes(number of qubits). Being able to engineer specific amounts of entanglement into a smaller number of qubits could be useful in quantum computation[6]. The particular kind of entangled pure qubit state that we have worked on in this paper has also been studied in the following papers [7, 8]. The authors in [7, 8] also studied pure qubit states where all reduced density matrices of fixed number of qubits have maximum possible entropy. Arnaud et al.[8] expanded their pure state density matrices in the generalized Bloch basis and equated the entanglement criterion explored here to the presence and absence of certain types of Bloch expansions terms and then connected this to some bounds related to quantum error cor-

recting codes to claim a lower bound on the amount of entanglement that can be seen in a typical large qubit state. We utilize this same connection between Bloch expansions of pure states and their entanglement and for the sake of completeness have proven this result following an independent route. Facchi et al.[7] worked in the qubit basis to prove the existence of an extremal version of our entanglement criterion(Absolutely maximally entangled states[9]) for small number of qubits and also disprove the existence of such absolutely maximally entangled states in n qubits systems with $n \geq 8$. Most of the states we have constructed here do not approach this extremal case. One of the motivation for our work is to identify recipes for constructing unitary transformations which when apply to a product state leads to a k -entangled states, that is states where every k vs $N-k$ bipartition leads to k qubit reduced density matrices that are maximally mixed for different k values. Now we argue that this is achieved once we construct the k -entangled states in terms of graph state[10].

There exists a one to one map from any simple graph[11](undirected, unweighted and without multiple edges or loops) to a pure n -qubit state. The graph edges also tell us what unitary operations need to be applied to realise these states. Hence finding a recipes for construction of k -entangled states using graph state automatically paves the way to identification of unitary transformations which when apply to a product state will generate these k -entangled states. Graph states have been experimentally realised [12] and it has been shown that these states exhibit rich entanglement structures [13–16], have applications in quantum computation for a different model of computation using measurements on graphs states[17] and their local unitary equivalents are also useful in quan-

* ss18ip003@iiserkol.ac.in

† sourin@iiserkol.ac.in

tum error correction[18].

II. MAXIMALLY ENTANGLED STATES:

An n qubit state is k-entangled, if any k qubit reduced density matrix of this state is maximally mixed. It is straightforward to show that a k-entangled state is also a 1,2,3,...,k-1-entangled state[3]. An n qubit state can at most be $\lfloor \frac{n}{2} \rfloor$ -entangled, which follows from constraints on the dimensions of the reduced density matrix and the fact that both reduced density matrices in the bipartition should have the same entropy. An n qubit state is called absolutely maximally entangled [19] if it is $\lfloor \frac{n}{2} \rfloor$ -entangled.

For example, a class of 1-entangled states is given by[8],

$$\frac{1}{\sqrt{2}} (|0\rangle^{\otimes n} + |1\rangle^{\otimes n}) \quad (1)$$

. For $n = 2$, this corresponds to the well known Bell state[20] and for $n \geq 3$ this corresponds to generalized GHZ states[3, 21].

The 3 qubit GHZ state in particular turns out to be absolutely maximally entangled, since $1 = \lfloor \frac{3}{2} \rfloor$. Generalized GHZ states with $n > 3$ are not absolutely maximally entangled.

A. Generalized Bloch basis

In this section we introduce the generalized Bloch basis [3, 8, 22] for describing an N qubit system. In this paper we have restricted our study to systems comprising of n qubits, hence our density matrices(ρ) are of dimensions $2^n \times 2^n$. Since ρ is also Hermitian, it can be expanded in terms of tensor products of Pauli matrices($\sigma_0 = I = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$, $\sigma_1 = X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}$, $\sigma_2 = Y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix}$, $\sigma_3 = Z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$),

$$\rho_{2^n \times 2^n} = \sum_{i_1, i_2, \dots, i_n} c_{i_1, i_2, \dots, i_n} \sigma_{i_1} \otimes \sigma_{i_2} \otimes \dots \otimes \sigma_{i_n} \quad (2)$$

where,

$$c_{i_1, i_2, i_3, \dots, i_n} = \frac{1}{2^n} \text{Trace}(\sigma_{i_1} \otimes \sigma_{i_2} \otimes \sigma_{i_3} \otimes \dots \otimes \sigma_{i_n} \times \rho) \quad (3)$$

For example, in this basis the Bell state, $|\psi\rangle = \frac{1}{\sqrt{2}}(|01\rangle + |10\rangle)$ can be expanded as,

$$(|\psi\rangle)(\langle\psi|) = \frac{1}{4}(I \otimes I + X \otimes X + Y \otimes Y - Z \otimes Z) \quad (4)$$

For the rest of the paper we will omit the explicitly written tensor product symbols and then, the typical terms appearing in an expansion are 'Pauli words' like,

$XX, YY, ZZ, XIII, X$ and so on. These Pauli words can be classified as 1-particle, 2-particle ... n-particle terms, based on how many non identity matrices they contain. Example: $XXY = 3$ -particle term, $XIZZIY = 4$ -particle term and so on. Now we define the weight of a Pauli word, which is equal to the number of non identity Pauli matrices it contains. At this point one should remember $c_{i_1, i_2, i_3, \dots, i_n}$ also corresponds to the expectation value of an operator $\sigma_{i_1} \sigma_{i_2} \sigma_{i_3} \dots \sigma_{i_n}$ in state ρ is also given by the same expression. Therefore expanding density matrices in this basis, directly gives us access to the average values of various physical observables associated with the individual spins for systems in that state.

The expectation value of all one particle operators is zero for 1- entangled states

Proof: From (3), we compute $c_{0,0,\dots,0} = \frac{1}{2^n} \text{Trace}(I_{2^n \times 2^n} \times \rho) = \frac{1}{2^n}$. Note that the rest of the terms is a subset of the set of 1-particle terms, 2-particle terms... n-particle terms. Therefore the first term of the Bloch expansion of any state is $\frac{I_{2^n \times 2^n}}{2^n}$. Partial tracing is distributive across matrix addition, so we can sum over the partial traces of the different matrices in the Bloch expansion. Partial tracing out $(n-1)$ qubits from $\frac{I_{2^n \times 2^n}}{2^n}$ can give us our required fully mixed $\frac{I_{2 \times 2}}{2}$ matrix. Therefore, we need the rest of the matrices in the Bloch expansion of ρ to partial trace out to zero.

Partial tracing distributes across matrix tensor products as matrix multiplications. example: $\text{Trace}_{23}(A_1 \otimes B_2 \otimes C_3) = A_1 \times \text{Trace}(B_2) \times \text{Trace}(C_3)$ An m-particle term in the Bloch expansion of an n-qubit state has m zero-trace matrices(X, Y, Z) and n-m Identities(I). For $m \geq 2$, when we partial trace out $n-1$ qubits, we will be forced to include at least one $\text{trace} = 0$ matrix in the set of matrices being partial traced by pigeon hole principle. This will make that term in the Bloch expansion disappear and so we are left with only the $m = 1$ and $m = 0$ operators. These correspond to the 1-particle and $\frac{I_{2^n \times 2^n}}{2^n}$. So we can see that if a Bloch expansion does not contain any 1-particle terms then, all terms other than $\frac{I_{2^n \times 2^n}}{2^n}$ will partial trace out to zero and hence we will be left with the fully mixed $\frac{I_{2 \times 2}}{2}$ matrix \Rightarrow 1-entangled states do not contain one particle terms in their Bloch expansions \square

This idea of the lack of one particle operators can be extended in a straightforward way to make the following more useful result.

Theorem 1: The expectation values/coefficients in the Bloch expansions of 1,2,3,... k-particle terms will be zero for k-entangled states.

For an n qubit pure state density matrix to become a k qubit maximally mixed density matrix upon




FIG. 1: Example of 4 qubit graph

tracing out $n-k$ qubits, we need the absence of any $1,2,3,..k$ -particle operators in its Bloch expansion. $1,2,3,..k$ -particle operators are the only operators that may not trace out to zero, when tracing out $n-k$ qubits from an n qubit state. All higher particle operators will necessarily partially trace out to zero.

B. Graph states

We start this section by reviewing the definition of graph states for the sake of completeness. Graphs [11] are a collection of vertices and edges (the vertices are connected by the edges), which can be represented as $G = (V, E)$ where, V is the set of vertices and E is the set of edges respectively. In the graphs that we consider, there will be no self loops (an edge from a vertex to itself) and between any two vertices there can be at most one edge. Also we only consider undirected graphs (graphs where the edges are not assigned a sense of direction).

Corresponding to any undirected n vertex graph as in figure 1, we can write down an n qubit pure quantum state called a graph state [10]. The vertices are labelled with certain operators (called correlation operators) and it should be noted that these operators commute. The way these correlation operators are constructed is by first numbering each vertex as representing a qubit. There is one correlation operator per graph vertex. The qubit states we are looking at are not identical in the quantum mechanical sense and so in an n qubit system, each qubit can be numbered. Once we number the vertices/qubits of the graph the correlation operator corresponding to each vertex can be constructed by including a Pauli $X(\sigma_x)$ acting on the Hilbert space of the i 'th qubit and Pauli $Z(\sigma_z)$ matrices at the Hilbert spaces of the qubits that are connected to the i 'th qubit, and Pauli I (Identity) matrices elsewhere. For example, in the graph in figure 1, the correlation operators are given by $XIZZ$, $IXZI$, $ZZXZ$, $ZIZX$. These four operators commute and square to identity and these two properties allow the correlation operators to be the generators of a 16 element Z_2^4 abelian group of operators under matrix multiplication. In general the n correlation operators of an n vertex graph generates 2^n operators called the stabilizer of the state. The sum of all these stabilizer operators multiplied by $\frac{1}{2^n}$ gives the density matrix for the n qubit graph state. The stabilizers of an n qubit state form a Z_2^n group.

As an example we can write the full expansion of the

graph in fig 1, $|G_4\rangle$,

$$|G_4\rangle\langle G_4| = \frac{1}{16}(IIII + XIZZ + IXZI + ZZXZ + ZIZX + XXIZ + YZYI + YIYI + ZYYZ + ZXIX + IZYY + IXIZ - YYXI - IYXY - XZXX + YXZY - XYYX) \quad (5)$$

III. ENTANGLEMENT IN GRAPH STATES:

There have been previous works that have explored the entanglement content of graph states through the use of different entanglement monotones [14, 23]. There has also been previous work exploring entanglement in the same spirit as this paper, where they show the lower limit of k -entanglement in large graph states using connections to quantum error correction as well as establishing that permutation symmetric stabilizer states [8] can be at most 1-entangled. In contrast we are considering graph states that are not permutation symmetric as a possible construction of k -entangled states.

To make progress we start with the following definition. **Definition-1** k -products is the set of elements generated by products between k correlation operators of a graph state. The different k -product sets are disjoint subsets of the full Bloch expansion because the correlation operators are elements of an abelian group. Now note that the smallest particle sector in the Bloch expansion of a k -entangled n qubit state is $k+1$. The weight of a correlation operator corresponding to a vertex with degree d is $d+1$. The correlation operators are themselves part of the Bloch expansion and hence a k -entangled graph state will only contain vertices with *degree*, $d \geq k$.

While *degree*, $d \geq k$ for every vertex of a graph is necessary for the state to be k -entangled, this is not sufficient because it is possible for products of correlation operators to have smaller weight. Since we need at least degree k for each vertex a useful starting point in a search for k -entangled graphs could be to look at k -regular graphs (graphs with *degree* = k for each vertex).

(Since we are interested in studying the weights of different Bloch expansion terms ..) Note that the i 'th correlation operator can be written as,

$$K_i = X_i \otimes \prod_{l \in N_i} Z_l \quad (6)$$

where $i \in 1, 2, 3, \dots, n$ for n qubit system. N_i , the neighbourhood of vertex i , is the set of vertices that are connected to the vertex i with exactly one edge. The set of N_i 's comes from the underlying graph. For example, in our previous 4 qubit graph (figure 1), $N_4 = [1, 3]$. and hence $K_4 = X_4 Z_1 Z_3$.

To understand what a product of k correlation operators looks like, let us first construct a general two-product between two correlation operators K_i and K_j with $N_i =$ neighbourhood of i and $N_j =$ neighbourhood of j ,

$$K_i \times K_j = \left(X_i \prod_{\otimes l}^{l \in N_i} Z_l \right) \times \left(X_j \prod_{\otimes m}^{m \in N_j} Z_m \right) \quad (7)$$

where,

$$i \neq j \in N = \{1, 2, 3, \dots, n\}; l \subset N - \{i\}; m \subset N - \{j\} \quad (8)$$

which can be reduced to,

$$K_i \times K_j = X_i X_j \prod_{\otimes l}^{l \in S_2} Z_l \quad (9)$$

where, $S_2 = N_i \oplus N_j$. " \oplus " is the symmetric difference binary operation between sets, which is defined as, $A \oplus B = (A \cup B) - (A \cap B)$. Note that all Z 's will have distinct indices owing to the fact that Z 's belonging to the same index multiply with each other in pairs to give identity as $Z^2 = I$. As defined earlier, the weight of the above product is equal to the number of distinct indices in the string of Pauli operators. We have unique indices in eqn 9 for the Z operators. If we include X_i and X_j we do not have all unique indices if $S_2 \cap \{i, j\} \neq \phi$, where ϕ denotes the empty set. Note that while N_i cannot contain i and N_j cannot contain j , $N_i \oplus N_j$ can contain both. To ensure that every index in eqn 9 is unique, we take out Z_i and Z_j if present, from the product over Z 's and subtract $\{i, j\}$ from the set S_2 . These Z 's that we have taken out will multiply with X_i and X_j depending on whether they exist in the symmetric sum S_2 or not. Finally the two-product looks like,

$$K_i \times K_j = X_i (Z_i)^{C_i(S_2)} X_j (Z_j)^{C_j(S_2)} \prod_{\otimes l}^{l \in S_2 - \{i, j\}} Z_l \quad (10)$$

Where $C_i(S_2)$ is the cardinality of i in S_2 , which means the number of times element i appears in set S_2 . So here we have a neat way of writing a general 2-product where each Pauli operator has a unique index. We see that this is easily generalizable to 3-products and greater. A general 3-product would look like,

$$K_i K_j K_r = X_i (Z_i)^{C_i(S_3)} X_j (Z_j)^{C_j(S_3)} X_r (Z_r)^{C_r(S_3)} \prod_{\otimes l}^{l \in S_3 - \{i, j, r\}} Z_l \quad (11)$$

where $S_3 = N_i \oplus N_j \oplus N_r$

The general n -product would look like,

$$K_{i_1} K_{i_2} \dots K_{i_n} = X_{i_1} (Z_{i_1})^{C_{i_1}(S_n)} X_{i_2} (Z_{i_2})^{C_{i_2}(S_n)} \dots X_{i_n} (Z_{i_n})^{C_{i_n}(S_n)} \prod_{\otimes l}^{l \in S_n - \{i_1, i_2, \dots, i_n\}} Z_l \quad (12)$$

where $S_n = N_{i_1} \oplus N_{i_2} \oplus N_{i_3} \oplus \dots \oplus N_{i_n}$,

Looking at (10), (11) and (12), we see that the weight of these products regardless of the weight of what appears inside the product (\prod) is at least equal to the number of X 's in front of the \prod . The number of X 's is clearly equal to the number of different correlation operators being multiplied, therefore in general,

Theorem 2: $w(k - \text{product}) \geq k$. k -product has minimum weight of k .

The theorem above is very important to the rest of this paper. What we have shown is that a 2-product has minimum weight of two and a 3-product has a minimum weight of 3 and so on, regardless of what graph you take. This result allows us to avoid computing the full Bloch expansion that corresponds to a given graph. We only need to compute a certain number of terms depending on what entanglement we expect in the state. With the n correlation operators we can read off of any given graph, one has to compute 2^n products to get the full Bloch expansion. But if one only needs to check if the state is k -entangled, since $k \leq \lfloor \frac{n}{2} \rfloor$ we only need to evaluate $\binom{n}{0} + \binom{n}{1} + \binom{n}{2} + \dots + \binom{n}{k} \leq 2^{n-1}$ terms. So, for 1-entanglement we only need weights of 1-products to be more than one, for 2-entanglement we only need weights of 1-products and 2-products to be more than two, for 3-entanglement we only need weights of 1-products, 2-products and 3-products to be more than three. We do not need to evaluate any of the other terms of the Bloch expansion when checking for 1,2 or 3...-entanglement in graph states.

A. 1 - entangled states:

As per theorem 1 we need the correlation operators to have a weight of at least 2 ($w \geq 2$) for 1-entanglement. Because of theorem 2 we only need to ensure the 1-products(correlation operators) of 1-entangled states have weight more than one. This can be easily ensured by connecting every vertex to at least one other vertex. So the recipe for a 1-entangled state is : Graphs where the degree of every vertex is greater than zero are 1-entangled. If one needs to construct a 1-entangled state with a minimum number of controlled-Z unitaries applied to a product state, then one needs to pair off all the vertices(qubits). For an odd number of qubits, we need one extra edge to connect the left out odd qubit with one of the pairs(figure 2a).

n-qubit GHZ state is 1-entangled: N qubit GHZ state is equivalent to an n qubit fully connected graph state up to local unitary transformations [10]. A fully connected graph is one, where every vertex is connected with an edge each, to every other vertex as in figure 2b

Now comparing this graph with the result in the pre-




FIG. 2: 1-entangled states

vious section, we know the GHZ state is at least 1-entangled. Now we show how it is exactly 1-entangled.

Each of the correlation operators have weight n for n qubit complete graph. Now look at the two products,

$$K_i K_j = X_i (Z_i)^{C_i(S_2)} X_j (Z_j)^{C_j(S_2)} \prod_{l \in S_2 - \{i,j\}} Z_l \quad (13)$$

All vertices are connected to all other vertices,

$$N_i \cup \{i\} = N_j \cup \{j\} \quad (14)$$

$$\implies S_2 - \{i,j\} = N_i \oplus N_j - \{i,j\} = \phi \quad (15)$$

ϕ is the null set. So the weight of what's inside the product in eqn(13) is zero. Therefore all two products of GHZ states are two particle terms. By theorem 2 Three and higher products have higher weights, therefore the smallest term in the Bloch expansion of the state has weight two and hence an n qubit GHZ is exactly 1-entangled.

B. 2 - entangled states:

1. *Qubit state with more than four qubits arranged in a circular linear chain is 2-entangled*

For 2- entanglement, we can start with 4-qubit systems, since smaller systems cannot have 2-entanglement (due to the symmetry of the entropy in the two parts after bi-partitioning as well as the fact that the maximum possible entropy of a $2^n \times 2^n$ matrix is $n \log 2$). We take the corresponding correlation operators for a "closed




FIG. 3: 5 qubit 2-entangled state

chain graph" as our ansatz and show that it is indeed 2-entangled. The general form of the correlation operators for a closed chain graph is given by

$$K_i = Z_{i-1} X_i Z_{i+1}. \quad (16)$$

For five qubits this ansatz gives us figure 3.

Now these are three particle terms and the $i+1$ and $i-1$ are modulo n sums so that they go in a cycle. Example for 4 qubits, $(4+1) \bmod(4) = 1$ and $(1-1) \bmod 4 = 4$.

Two-product looks like ,

$$K_i K_j = Z_{i-1} X_i Z_{i+1} Z_{j-1} X_j Z_{j+1} \quad (17)$$

If $i+1 = j$, that is they are neighbours,

$$K_i K_{i+1} = Z_{i-1} Y_i Y_{i+1} Z_{i+2} \quad (18)$$

This is a four body term

if $i+2 = j$, that is they are next nearest neighbours,

$$K_i K_{i+2} = Z_{i-1} X_i X_{i+2} Z_{i+3} \quad (19)$$

This is also a four body term. For 4 qubits, $(i-1) \bmod(4) = (i+3) \bmod(4)$ and hence Z_{i-1} will square to identity with Z_{i+3} and therefore, for 4 qubits, 2 product between next nearest neighbours becomes two particle term and hence 4 qubit state following this recipe cannot be 2-entangled. But for any qubit state above four qubits, this cancellation due overlapping of indices from modular addition does not take place and we again get four particle terms. All two products other than those between nearest neighbours and next nearest neighbours are between correlation operators with no overlap, and hence they give 6 particle terms,

$$K_i K_j = Z_{i-1} X_i Z_{i+1} Z_{j-1} X_j Z_{j+1} \quad (20)$$

To summarise, two products are all giving at least 4 particle terms for systems with more than 4 qubit in our ansatz, three products and higher give at least 3 particle terms and higher due to theorem 2, and individual correlation operators are 3 particle terms, therefore this recipe gives us 2-entangled states.

C. 3 - entangled states:

For 3-entanglement the recipes we have constructed involves arranging a $2n$ qubit state as a bilayer graph,




FIG. 4: 3-entangled states with 6,8 and 10 qubits and their corresponding adjacency matrices. Note the two diagonal blocks in all the matrices correspond to the fully connected sub graphs/layers and the off diagonal blocks correspond to the one to one connections between the two layers.

with n qubits in each layer. Within a layer, all the qubits are fully connected to one another. Between the layers, there are edges in a one to one mapping between the two sets of n qubits. The examples in figures 4a ,4b ,4c for 6,8 and 10 qubits might help clarify our construction.

This construction can be extended for any even number of qubits. To understand why this construction works we need to look at the adjacency matrices(fig:4) for these three graphs. Adjacency matrices are ubiquitous in graph theory, for our purposes here, all we have done is arrange the correlation operators for our 3-entangled graphs in rows, such that the Pauli X's are all along the main diagonal. Note that this necessarily forms a symmetric matrix as we are considering undirected graphs only. A "Z" in row i and column j of the matrix denotes that there is an edge between qubit i and qubit j . An "I" in row i and column j of the matrix denotes the absence of an edge between qubit i and qubit j .

Observing the matrices in fig:4 we can see that even as we scale up this system, there is a persistent structure to these adjacency matrices. There are two diagonal blocks which corresponds to each fully connected layer of the bi-layer and the two off diagonal blocks represent the one to one connections between the two layers. For 3-entanglement we need to analyze 1-products, 2-products and 3-products and ensure they have weight greater than three, all higher products are necessarily four body and higher as per theorem 2.

Now we prove how our ansatz is 3-entangled. For any $2n$ qubit system constructed according to our recipe, 1-products all have weight $n+1$, where n weight comes from connections in the plane and one from the connection be-

tween planes. Due to the symmetry of this system, there are exactly three qualitatively "different" 2-products and 3- products. The three different typical 2- products are,

1) 2-products where both the correlation operators belong to same layer

$$\begin{aligned} & X Z Z \dots Z Z I I \dots I \\ & \times Z X Z \dots Z I Z I \dots I \\ & = Y Y I \dots I Z Z I \dots I \end{aligned} \quad (21)$$

\implies weight, $w = 4$

2) 2-products where the correlation operators are connected but belong to different layers

$$\begin{aligned} & X Z Z \dots Z Z I I \dots I \\ & \times Z I I \dots I X Z Z \dots Z \\ & = Y Z Z \dots Z Y Z Z \dots Z \end{aligned} \quad (22)$$

$\implies 2n$ weight.

3) 2-products where the correlation operators belong to different layers of the bilayer graph and belong to vertices that are not connected

$$\begin{aligned} & X Z Z Z \dots Z Z I I I \dots I \\ & \times I Z I I \dots I Z X Z Z \dots Z \\ & = X I Z Z \dots Z I X Z Z \dots Z \end{aligned} \quad (23)$$

$\implies 2n - 2$ weight.

The three different typical 3- products are,

1) All three correlation operators belong to same layer

$$\begin{aligned} & X Z Z Z Z \dots Z Z I I I I \dots I \\ & \times Z X Z Z Z \dots Z I Z I I I \dots I \\ & \times Z Z X Z Z \dots Z I I Z I I \dots I \\ & = X X X Z Z \dots Z Z Z Z I I \dots I \end{aligned} \quad (24)$$




FIG. 5: (a) 25 qubit periodic lattice (torus) for 4-entangled state. Note that in this periodic lattice it is the edges outgoing from the diagrammatically opposite ends along the two independent directions that are identified to form the torus. This is represented by the double headed arrows that represents the bonds on the edge. (b) Correlation operators separated by distance ≥ 3 cannot overlap. (c) Two correlation operators overlapping at the lattice sites marked yellow. The weight of the product is the number of red sites in this diagram. The reds represent the non identity operators and the yellow dots show where the correlation operators overlap such that Z 's of same index square to identity. (d) In this 5×5 lattice, note the minimum weight product of five correlation operators lies on one of the diagonals.

$\implies n + 3$ weight.

2) Two correlation operators in one layer, third correlation operator in different layer and the corresponding vertex is connected to one of the previous two vertices

$$\begin{aligned}
 & X \ Z \ Z \ Z \ \dots \ Z \ Z \ I \ I \ I \ \dots \ I \\
 \times & Z \ X \ Z \ Z \ \dots \ Z \ I \ Z \ I \ I \ \dots \ I \\
 \times & Z \ I \ I \ I \ \dots \ I \ X \ Z \ Z \ Z \ \dots \ Z \\
 = & X \ Y \ I \ I \ \dots \ I \ Y \ I \ Z \ Z \ \dots \ Z
 \end{aligned} \quad (25)$$

$\implies n + 1$ weight.

3) Two correlation operators in one layer, third correlation operator in different layer and the corresponding vertex is NOT connected to one of the previous two correlation operators

$$\begin{aligned}
 & X \ Z \ Z \ Z \ \dots \ Z \ Z \ I \ I \ I \ \dots \ I \\
 \times & Z \ X \ Z \ Z \ \dots \ Z \ I \ Z \ I \ I \ \dots \ I \\
 \times & I \ I \ Z \ I \ \dots \ I \ Z \ Z \ X \ Z \ \dots \ Z \\
 = & Y \ Y \ Z \ I \ \dots \ I \ I \ I \ X \ Z \ \dots \ Z
 \end{aligned} \quad (26)$$

$\implies n + 1$ weight.

The minimum weight term from the set of 1-products, 2-products and 3-products is equal to the minimum of $[(n + 1), 4, 2n, 2n - 2, (n + 3), (n + 1), (n + 1)]$ which is equal to minimum of $[4, (2n - 2), (n + 1)]$. For $n \geq 3$ this minimum is 4. That is any $2n$ qubit system with $n \geq 3$ that follows this bilayer construction is 3-entangled.

D. 4 - entangled states:

Consider a 2D lattice graph state as shown in fig 5 with periodic boundary conditions imposed along both the independent directions hence forming a torus like geometry. We claim that all such 2D lattice graphs with lattice dimensions more than 5×5 are 4-entangled. All correlation operators (which are nothing but 1-products) has weight

$= 5$ for such a graph, as each lattice site in the graph has four neighbours. Therefore the maximum entanglement that we can expect from this arrangement is $4 \ln 2$. By theorem 1 we need to find out the minimum weights of 2-products, 3-products and 4-products to confirm this $4 \ln 2$ entanglement. If we consider k correlation operators that are far apart (they have no overlapping vertices in the graph), the weight of such a k -product will be $5k$. Like in figure 5 (b) with two correlation operators that are three distance [24] apart.

What can reduce the weight of a k -product from $5k$ is vertices common between multiple correlation operators.

These common vertices correspond to Z operators with the same index squaring to Identity (I). For correlation operators to have common vertices, the distance between them along this lattice graph needs to be at most 2. So when attempting to evaluate the minimum k -product we need to consider all configurations of k operators such that they are all within two distance of each other. Weight of 2-product $= 2 \times 5 - \theta \times 2$, θ = number of overlapping vertices. In the 2D lattice it is possible to exhaustively show that the best way to arrange a 2-product to get minimum weight is to place them diagonal to each other (figure 5(c)).

This produces two overlaps and so the weight is $10 - 4 = 6$. This extends for 3-products, 4-products and so on. That is, the minimum 3-product consists of the product of three correlation operators along the same diagonal, the minimum 4-product consists of the product of four correlation operators along the same diagonal and so on. The minimum k -product will involve k correlation operators along a diagonal. If k is less than both lattice dimensions then there will be $2(k - 1)$ overlaps or a reduction of $4(k - 1)$ weight. Therefore the minimum k -product has weight $= 5k - 4(k - 1) = k + 4$. Since $k \geq 1$, the minimum weight of 1-product $= 5 \leq 2$

product ≤ 3 -product and so on and so forth. Now we need to evaluate what minimum size of the lattice will allow 4-entangled states. Consider a k -product organized along a diagonal where $k = \min(l, m)$ where l, m are the number of vertices along the two independent directions, this gives us two more overlaps (-4 weight) and therefore the weight is $k + 4 - 4 = k$ like in figure 5(d).

We need this k product to have weight ≥ 5 . This argument can be applied to both dimensions (l, m) of this lattice and since this lattice has $l \times m$ qubits with $l, m \geq 5$, the smallest lattice is a square with $l = m = 5$. By this construction we need at least 25 qubits to construct a 4-entangled state.

IV. CONCLUSION:

We constructed 1, 2, 3 and 4-entangled states from graph states. Since the structure of a graph state can tell us what quantum circuit can be used to construct the state starting from a product state [25], hence our findings provides a series of explicit recipes to construct these states starting from product states.

ACKNOWLEDGMENTS

S.S. acknowledges IISER Kolkata, India, for support in the form of a fellowship. S.D. would like to acknowledge the MATRICS grant (Grant No. MTR/ 2019/001 043) from the Science and Engineering Research Board (SERB) for funding.

-
- [1] A. Einstein, B. Podolsky, and N. Rosen, Can quantum-mechanical description of physical reality be considered complete?, *Phys. Rev.* **47**, 777 (1935).
 - [2] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Quantum entanglement, *Reviews of Modern Physics* **81**, 865–942 (2009).
 - [3] M. A. Nielsen and I. Chuang, Quantum computation and quantum information (2002).
 - [4] M. B. Plenio and S. S. Virmani, An introduction to entanglement theory, *Quantum information and coherence*, 173 (2014).
 - [5] D. N. Page, Average entropy of a subsystem, *Phys. Rev. Lett.* **71**, 1291 (1993).
 - [6] A. Ekert and C. Macchiavello, Quantum error correction for communication, *Phys. Rev. Lett.* **77**, 2585 (1996).
 - [7] P. Facchi, G. Florio, G. Parisi, and S. Pascazio, Maximally multipartite entangled states, *Physical Review A* **77**, 10.1103/physreva.77.060304 (2008).
 - [8] L. Arnaud and N. J. Cerf, Exploring pure quantum states with maximally mixed reductions, *Physical Review A* **87**, 10.1103/physreva.87.012319 (2013).
 - [9] W. Helwig, W. Cui, J. I. Latorre, A. Riera, and H.-K. Lo, Absolute maximal entanglement and quantum secret sharing, *Phys. Rev. A* **86**, 052335 (2012).
 - [10] M. Hein, W. Dür, J. Eisert, R. Raussendorf, M. Nest, and H.-J. Briegel, Entanglement in graph states and its applications, arXiv preprint quant-ph/0602096 (2006).
 - [11] D. B. West *et al.*, *Introduction to graph theory*, Vol. 2 (Prentice hall Upper Saddle River, 2001).
 - [12] S. M. Lee, H. S. Park, J. Cho, Y. Kang, J. Y. Lee, H. Kim, D.-H. Lee, and S.-K. Choi, Experimental realization of a four-photon seven-qubit graph state for one-way quantum computation, *Opt. Express* **20**, 6915 (2012).
 - [13] A. H. Shenoy and R. Srikanth, Maximally nonlocal subspaces, *Journal of Physics A: Mathematical and Theoretical* **52**, 095302 (2019).
 - [14] M. Hein, J. Eisert, and H. J. Briegel, Multiparty entanglement in graph states, *Physical Review A* **69**, 10.1103/physreva.69.062311 (2004).
 - [15] N. Gisin and H. Bechmann-Pasquinucci, Bell inequality, bell states and maximally entangled states for n qubits, *Physics Letters A* **246**, 1–6 (1998).
 - [16] J. C. Adcock, S. Morley-Short, A. Dahlberg, and J. W. Silverstone, Mapping graph state orbits under local complementation, *Quantum* **4**, 305 (2020).
 - [17] R. Raussendorf and H. J. Briegel, A one-way quantum computer, *Phys. Rev. Lett.* **86**, 5188 (2001).
 - [18] D. Gottesman, *Stabilizer codes and quantum error correction*. *Caltech Ph. D.*, Ph.D. thesis, Thesis, eprint: quant-ph/9705052 (1997).
 - [19] W. Helwig and W. Cui, Absolutely maximally entangled states: Existence and applications (2013), arXiv:1306.2536 [quant-ph].
 - [20] J. J. Sakurai and E. D. Commins, *Modern quantum mechanics*, revised edition (1995).
 - [21] D. M. Greenberger, M. A. Horne, and A. Zeilinger, Going beyond bell’s theorem (2007), arXiv:0712.0921 [quant-ph].
 - [22] F. Bloch, Nuclear induction, *Phys. Rev.* **70**, 460 (1946).
 - [23] M. Hajdušek and M. Muraio, Direct evaluation of pure graph state entanglement, *New Journal of Physics* **15**, 013039 (2013).
 - [24] The distance between two vertices on a graph is equal to the number of edges on any of the shortest paths between the two vertices.
 - [25] The qubit at each vertex is initially prepared in state $\frac{1}{\sqrt{2}}(|0\rangle + |1\rangle)$. An edge between vertex i and j corresponds to a controlled-Z gate acting between qubits i and j [10].