

The Parameterized Complexity of s -Club with Triangle and Seed Constraints

Jaroslav Garvardt, Christian Komusiewicz^[0000-0003-0829-7032], and Frank Sommer^{*[0000-0003-4034-525X]}

Fachbereich Mathematik und Informatik, Philipps-Universität Marburg,
Marburg, Germany

{garvardt,komusiewicz,fsommer}@informatik.uni-marburg.de

Abstract. The s -CLUB problem asks, for a given undirected graph G , whether G contains a vertex set S of size at least k such that $G[S]$, the subgraph of G induced by S , has diameter at most s . We consider variants of s -CLUB where one additionally demands that each vertex of $G[S]$ is contained in at least ℓ triangles in $G[S]$, that $G[S]$ contains a spanning subgraph G' such that each edge of $E(G')$ is contained in at least ℓ triangles in G' , or that S contains a given set W of seed vertices. We show that in general these variants are $W[1]$ -hard when parameterized by the solution size k , making them significantly harder than the unconstrained s -CLUB problem. On the positive side, we obtain some FPT algorithms for the case when $\ell = 1$ and for the case when $G[W]$, the graph induced by the set of seed vertices, is a clique.

1 Introduction

Finding cohesive subgroups in social or biological networks is a fundamental task in network analysis. A classic formulation of cohesiveness is based on the observation that cohesive groups have small diameter. This observation led to the s -club model originally proposed by Mokken [15]. An s -club in a graph $G = (V, E)$ is a set of vertices S such that $G[S]$, the subgraph of G induced by S , has diameter at most s . The 1-clubs are thus precisely the cliques and the larger the value of s , the more the clique-defining constraint of having diameter one is relaxed. In the s -CLUB problem we aim to decide whether G contains an s -club of size at least k .

A big drawback of s -clubs is that the largest s -clubs are often not very cohesive with respect to other cohesiveness measures such as density or minimum degree. This behavior is particularly pronounced for $s = 2$: the largest 2-club in a graph is often the vertex v of maximum degree together with its neighbors [10]. To avoid these so-called hub-and-spoke structures, it has been proposed to augment the s -club definition with additional constraints [5,14,16,18].

One of these augmented models, proposed by Carvalho and Almeida [5], asks that every vertex is part of a triangle [5]. This property was later generalized to

* Supported by the Deutsche Forschungsgemeinschaft (DFG), project EAGR (KO 3669/6-1).

the *vertex- ℓ -triangle* property, which asks that every vertex of S is in at least ℓ triangles in $G[S]$ [1].

VERTEX TRIANGLE s -CLUB

Input: An undirected graph $G = (V, E)$, and two integers $k, \ell \geq 1$.

Question: Does G contain an s -club S of size at least k that fulfills the vertex- ℓ -triangle property?

The vertex- ℓ -triangle constraint entails some desirable properties for cohesive subgraphs. For example, in a vertex- ℓ -triangle s -club, the minimum degree is larger than $\sqrt{2\ell}$. However, some undesirable behavior of hub-and-spoke structures remains. For example, the graph consisting of two cliques of size $d + 1$ that are connected via one edge is a vertex- $\binom{d}{2}$ -triangle 3-club but it can be made disconnected via one edge deletion. Thus, vertex- ℓ -triangle s -clubs are not robust with respect to edge deletions.

To overcome this problem, we introduce a new model where we put triangle constraints on the edges of the s -club instead of the vertices. More precisely, we say that a vertex set S of a graph G fulfills the *edge- ℓ -triangle* property if $G[S]$ contains a spanning subgraph $G' := (S, E')$ such that every edge in $E(G')$ is in at least ℓ triangles in G' and the diameter of G' is at most s . Next, we introduce the related problem.

EDGE TRIANGLE s -CLUB

Input: An undirected graph $G = (V, E)$, and two integers $k, \ell \geq 1$.

Question: Does G contain a vertex set S of size at least k that fulfills the edge- ℓ -triangle property?

Note that in this definition, the triangle and diameter constraints are imposed on a spanning subgraph of $G[S]$. In contrast, for VERTEX TRIANGLE s -CLUB, they are imposed directly on $G[S]$. The reason for this distinction is that we would like to have properties that are closed under edge insertions. Properties which are closed under edge insertions are also well-motivated from an application point of view since adding a new connection within a group should not destroy this group. If we would impose the triangle constraint on the induced subgraph $G[S]$ instead, then an edge- ℓ -triangle s -club S would not be robust to edge additions. For example, consider a graph G consisting of clique C to which two vertices u and v are attached in such a way that both u and v have exactly 2 neighbors in C which are distinct. The $V(G)$ is an edge-1-triangle 3-club, but after adding the edge uv , the edge uv is contained in no triangle and thus any edge-1-triangle 3-club cannot contain both u and v .

Observe that every set that fulfills the edge- ℓ -triangle property also fulfills the vertex- ℓ -triangle property. Also note that the converse is not true: A vertex- ℓ -triangle s -club is not necessarily also an edge- ℓ -triangle s -club. For this, consider the above mentioned graph consisting of two cliques of size $d + 1$ that are connected via one edge. Then, this graph is an vertex- $\binom{d}{2}$ -triangle-3-club. But after deleting e the graph is disconnected. Moreover, each vertex $v \in S$ has at least $\ell + 1$ neighbors in S : Consider an arbitrary edge uv . Since uv is in at least ℓ

triangles $\{u, v, w_1\}, \dots, \{u, v, w_\ell\}$ we thus conclude that u and v have degree at least ℓ . We can show an even stronger statement: an edge- ℓ -triangle s -club S is robust against up to ℓ edge deletions, as desired.

Proposition 1. *Let $G = (V, E)$ be a graph and let S be an edge- ℓ -triangle s -club in G . More precisely, let G' be a spanning subgraph of $G[S]$ such that every edge in $E(G')$ is in at least ℓ triangles in G' and the diameter of G' is at most s . If ℓ edges are removed from G' , then S is still an $(s + \ell)$ -club and a $(2s)$ -club.*

Proof. We show that if ℓ edges are removed from G' , the diameter of the resulting graph \tilde{G} increases by at most ℓ . Let $P = (v_1, \dots, v_{s+1})$ be a path of length s in G' . Since G' is an edge- ℓ -triangle s -club, every edge $v_i v_{i+1}$ of P is part of at least ℓ triangles in G' . Thus, for two vertices v_i and v_{i+1} in P there is a path of length at most two from v_i to v_{i+1} in G' , either directly through the edge $v_i v_{i+1}$ or via a vertex u that forms one of the ℓ triangles with v_i and v_{i+1} in G' . Thus, $\text{dist}(v_i, v_{i+1})$ increases by at most 1 after one edge deletion and only if $v_i v_{i+1}$ is removed. Since at most ℓ of the edges in P are removed, we have $\text{dist}(v_1, v_{s+1}) \leq \text{dist}(v_1, v_2) + \dots + \text{dist}(v_s, v_{s+1}) \leq s + \ell$ in \tilde{G} . By the same arguments, we also have $\text{dist}(v_1, v_{s+1}) \leq 2s$.

Thus, after deleting ℓ edges in G' , S is an $(s + \ell)$ -club and a $(2s)$ -club. \square

The following further variant of s -CLUB is also practically motivated but not necessarily by concerns about the robustness of the s -club. Here the difference to the standard problem is simply that we are given a set of seed vertices W and aim to find a large s -club that contains all seed vertices.

SEEDED s -CLUB

Input: An undirected graph $G = (V, E)$, a subset $W \subseteq V$, and an integer $k \geq 1$.

Question: Does G contain an s -club S of size at least k such that $W \subseteq S$?

This variant has applications in community detection, where we are often interested in finding communities containing some set of fixed vertices [12,19].

In this work, we study the parameterized complexity of the three above-mentioned problems with respect to the standard parameter solution size k . Our goal is to determine whether FPT results for s -CLUB [6,17] transfer to these practically motivated problem variants.

Known Results. The s -CLUB problem is NP-hard for all $s \geq 1$ [4], even when the input graph has diameter $s + 1$ [2]. For $s = 1$, s -CLUB is equivalent to CLIQUE and thus W[1]-hard with respect to k . In contrast, for every $s > 1$, s -CLUB is fixed-parameter tractable (FPT) with respect to the solution size k [6,17]. This fixed-parameter tractability can be shown via a Turing kernel with $\mathcal{O}(k^2)$ vertices for even s and $\mathcal{O}(k^3)$ vertices for odd s [17,6]. The complexity of s -CLUB has been also studied with respect to different classes of input graphs [9] and with respect to structural parameters such as degeneracy of the input graph [11]. The s -CLUB problem can be solved efficiently in practice, in particular for $s = 2$ [4,6,10]. In

particular, the 2-CLUB problem has efficient branch-and-bound algorithms [6,10] which can compute the optimal solutions on very large sparse graphs.

VERTEX TRIANGLE s -CLUB is NP-hard for all $s \geq 1$ and for all $\ell \geq 1$ [5,1]. We are not aware of any algorithmic studies of EDGE TRIANGLE s -CLUB or SEEDED s -CLUB. NP-hardness of EDGE TRIANGLE s -CLUB for $\ell = 1$ can be shown via the reduction for VERTEX TRIANGLE s -CLUB for $\ell = 1$ [1]. Also, the NP-hardness of SEEDED s -CLUB for $W \neq \emptyset$ follows directly from the fact that an algorithm for the case where $|W| = 1$ can be used as a black box to solve s -CLUB. Further robust models of s -clubs, which are not considered in this work, include t -hereditary s -clubs [16], t -robust s -clubs [18], and t -connected s -clubs [20,14]. For an overview on clique relaxation models and complexity issues for the corresponding subgraph problems we refer to the relevant surveys [13,16].

Our Results. An overview of our results is given in Table 1. For VERTEX TRIANGLE s -CLUB and EDGE TRIANGLE s -CLUB, we provide a complexity dichotomy for all interesting combinations of s and ℓ , that is, for all $s \geq 2$ and $\ell \geq 1$, into cases that are FPT or W[1]-hard with respect to k , respectively. Our W[1]-hardness reduction for EDGE TRIANGLE s -CLUB for $\ell \geq 2$ also shows the NP-hardness of this case. The FPT-algorithms are obtained via adaptations of the Turing kernelization for s -CLUB. Interestingly, VERTEX TRIANGLE s -CLUB with $\ell = 1$ is FPT only for larger s , whereas EDGE TRIANGLE s -CLUB with $\ell = 1$ is FPT for all s . In our opinion, this means that the edge- ℓ -triangle property is preferable not only from a modelling standpoint but also from an algorithmic standpoint as it allows to employ Turing kernelization as a part of the solving procedure, at least for $\ell = 1$. It is easy to see that standard problem kernels of polynomial size are unlikely to exist for VERTEX TRIANGLE s -CLUB and EDGE TRIANGLE s -CLUB: s -clubs are necessarily connected and thus taking the disjoint union of graphs gives a trivial or-composition and, therefore, a polynomial problem kernel implies $\text{coNP} \subseteq \text{NP/poly}$ [3].

All of our hardness results for VERTEX TRIANGLE s -CLUB and EDGE TRIANGLE s -CLUB are shown by a reduction from CLIQUE which is W[1]-hard with respect to k [7,8]. The idea is to replace each vertex of the CLIQUE instance by a vertex gadget. These gadgets are constructed in such a way that if one of these vertices is part of a vertex/edge- ℓ -triangle 2-club S , then the entire vertex gadget is part of S . We can then use the distance constraint to make sure that full vertex gadgets are chosen only if the corresponding vertices are adjacent. While this idea is very natural, using the triangle constraint without creating many vertices that are too close to each other turned out to be technically challenging.

For SEEDED s -CLUB, we provide a kernel with respect to k for clique seeds W and W[1]-hardness with respect to k for some other cases. For $s = 2$, our results provide a dichotomy into FPT and W[1]-hardness with respect to k in terms of the structure of the seed.

The W[1]-hardness of SEEDED s -CLUB is provided by two reductions from CLIQUE. One reduction is for the case $s = 2$ and any seed that contains at least two non-adjacent vertices u and z . The other reduction is for the case $s \geq 3$ and any seed that contains at least two connected components U and Z . In both

Table 1. Overview of our results of the parameterized complexity of the three problems with respect to the parameter solution size k .

	VERTEX TRIANGLE s -CLUB	EDGE TRIANGLE s -CLUB	SEEDED s -CLUB
FPT	$\ell = 1$ and $s \geq 4$	$\ell = 1$ for each s	W is a clique
W[1]-h	$\ell = 1$ and $s \leq 3$	$\ell \geq 2$ for each s	$s = 2$ and $G[W]$ contains at least two non-adjacent vertices
	$\ell \geq 2$ for each s		$s \geq 3$ and $G[W]$ contains at least 2 connected components

cases we add two copies X_1 and X_2 of the graph of the CLIQUE instance to the new instance of SEEDED s -CLUB such that each vertex of X_1 has distance at most s to u or U if its copy in X_2 is also part of the solution. We show a similar property for X_2 and z or Z . This ensures that if $p_1 \in X_1$ is in the solution if and only if $p_2 \in X_2$ is in the solution. Furthermore, the reductions have the property that vertex p_1 in X_1 has distance at most s to vertex q_2 in X_2 if and only if pq is an edge of the CLIQUE instance. This feature will then ensure that the same clique has to be chosen from both copies.

Our W[1]-hardness results, in particular those for SEEDED s -CLUB, show that the FPT results for s -CLUB are quite brittle since the standard argument that we may assume $k \geq \Delta$ fails and that adding even simple further constraints makes finding small-diameter subgraphs much harder.

Preliminaries. For integers p, q , we denote $[p, q] := \{p, p + 1, \dots, q\}$ and $[q] := [1, q]$. For a graph G , we let $V(G)$ denote its vertex set and $E(G)$ its edge set. We let n and m denote the order of G and the number of edges in G , respectively. A *path of length p* is a sequence of pairwise distinct vertices v_1, \dots, v_{p+1} such that $v_i v_{i+1} \in E(G)$ for each $i \in [p]$. The *distance* $\text{dist}_G(u, v)$ is the length of a shortest path between vertices u and v . Furthermore, we define $\text{dist}_G(u, W) := \min_{w \in W} \text{dist}(u, w)$. We denote by $\text{diam}_G(G) := \max_{u, v \in V(G)} \text{dist}_G(u, v)$ the *diameter* of G . Let $S \subseteq V(G)$ be a vertex set. We denote by $N_i(S) := \{u \in V \mid \text{dist}(u, S) = i\}$ the *open i -neighborhood* of S and by $N_i[S] := \bigcup_{j \leq i} N_j(S) \cup S$ the *closed i -neighborhood* of S . For a vertex $v \in V(G)$, we write $N_i(v) := N_i(\{v\})$ and $N_i[v] := N_i[\{v\}]$. A graph $G' := (V', E')$ with $V' \subseteq V$, and $E' \subseteq E(G[V'])$ is a *subgraph* of G . By $G[S] := (S, \{uv \in E(G) \mid u, v \in S\})$ we denote the *subgraph induced by S* . Furthermore, by $G - S := G[V \setminus S]$ we denote the induced subgraph obtained after the deletion of the vertices in S . A vertex set such that each pair of vertices is adjacent is called a *clique* and a clique consisting of three vertices is a *triangle*.

For the definitions of parameterized complexity theory, we refer to the standard monographs [7,8]. All of our hardness results are shown by a reduction from CLIQUE.

CLIQUE

Input: An undirected graph $G = (V, E)$ and an integer k .

Question: Does G contain a clique of size at least k ?

CLIQUE is W[1]-hard with respect to k [7,8].

2 Vertex Triangle s -Club

In this section, we settle the parameterized complexity of VERTEX TRIANGLE s -CLUB with respect to the solution size k . First, we show that this problem is fixed-parameter tractable when $\ell = 1$ and $s \geq 4$. Afterwards, we show W[1]-hardness for all remaining cases, that is, for $\ell \geq 2$ and $s \geq 2$, and also for $\ell = 1$ and $s \in \{2, 3\}$.

2.1 FPT-Algorithms

The overall idea is based on the idea of the Turing kernel for 2-CLUB, that is, bounding the size of $N_s[v]$ for each vertex $v \in V(G)$. The first step is to remove all vertices which are not in a triangle.

Reduction Rule 1. *Let (G, k) be an instance of VERTEX TRIANGLE s -CLUB. Delete all vertices from G which are not part of any triangle.*

Clearly, Reduction Rule 1 is correct and can be exhaustively applied in polynomial time. The application of Reduction Rule 1 has the following effect: if some vertex v is close to many vertices, then (G, k) is a trivial yes-instance.

Lemma 1. *Let (G, k) be an instance of VERTEX TRIANGLE s -CLUB with $\ell = 1$ and $s \geq 4$ to which Reduction Rule 1 is applied. Then, (G, k) is a yes-instance if $|N_{\lfloor s/2 \rfloor - 1}[v]| \geq k$ for some vertex $v \in V(G)$.*

Proof. Let $v \in V(G)$ be a vertex such that $|N_{\lfloor s/2 \rfloor - 1}[v]| \geq k$. We construct a vertex-1-triangle s -club T of size at least $|N_{\lfloor s/2 \rfloor - 1}[v]| \geq k$. Initially, we set $T := N_{\lfloor s/2 \rfloor - 1}[v]$. Now, for each vertex $w \in N_{\lfloor s/2 \rfloor - 1}(v)$ we do the following: Since Reduction Rule 1 is applied, we conclude that there exist two vertices x and y such that $G[\{w, x, y\}]$ is a triangle. We add x and y to the set T . We call the set of vertices added in this step the T -expansion.

Next, we show that T is indeed a vertex-1-triangle s -club for $s \geq 4$. Observe that each vertex in T is either in $N_{\lfloor s/2 \rfloor - 1}[v]$ or a neighbor of a vertex in $N_{\lfloor s/2 \rfloor - 1}(v)$. Hence, each vertex in T has distance at most $\lfloor s/2 \rfloor$ to vertex v . Thus, T is an s -club. It remains to show that each vertex of T is in a triangle. Observe that for each vertex $w \in N_{\lfloor s/2 \rfloor - 2}[v]$ we have $N(w) \subseteq N_{\lfloor s/2 \rfloor - 1}[v]$. Recall that since Reduction Rule 1 is applied, each vertex in G is contained in a triangle. Thus, each vertex of $N_{\lfloor s/2 \rfloor - 2}[v]$ is contained in a triangle in T . Furthermore, all vertices in $N_{\lfloor s/2 \rfloor - 1}(v) \cup (T \setminus N_{\lfloor s/2 \rfloor - 1}[v])$ are in a triangle because of the T -expansion. Since $|T| \geq |N_{\lfloor s/2 \rfloor - 1}[v]| \geq k$, the statement follows. \square

Next, we show that Lemma 1 implies the existence of a Turing kernel for $s \geq 4$. We do this by showing that $N_s[v]$ is bounded for every vertex in the graph. This in turn implies that the problem is fixed-parameter tractable. It is sufficient to bound the size of $N_s[v]$ for each $v \in V(G)$ since we then can query the oracle for an s -club of size k .

Theorem 2. *VERTEX TRIANGLE s -CLUB for $\ell = 1$ admits a k^4 -vertex Turing kernel if $s = 4$ or $s = 7$, a k^5 -vertex Turing kernel if $s = 5$, and a k^3 -vertex Turing kernel if $s = 6$ or $s \geq 8$.*

Proof. First, we apply Reduction Rule 1. Because of Lemma 1 we conclude that (G, k) is a trivial yes-instance if $|N_{\lfloor s/2 \rfloor - 1}[v]| \geq k$ for any vertex $v \in V(G)$. Thus, in the following we can assume that $|N_{\lfloor s/2 \rfloor - 1}[v]| < k$ for each vertex $v \in V(G)$. We use this fact to bound the size of $N_s[v]$ in non-trivial instances.

In this case we have $\lfloor s/2 \rfloor - 1 = 1$. Hence, for each $s \geq 4$ from $|N_{\lfloor s/2 \rfloor - 1}[v]| < k$ we obtain that the size of the neighborhood of each vertex is bounded. Thus, we obtain a k^4 -vertex Turing kernel for $s = 4$ and a k^5 -vertex Turing kernel for $s = 5$. Furthermore, if $s = 7$ we have $\lfloor s/2 \rfloor - 1 = 2$. Thus, we obtain a k^4 -vertex Turing kernel for $s = 7$ since $N_7[v] \subseteq N_8[v] = N_2[N_2[N_2[N_2[v]]]]$.

If $s = 6$ or $s \geq 8$, then $\lfloor s/2 \rfloor - 1 \geq \lceil s/3 \rceil$. Observe that $N_s[v]$ is contained in $N_{\lceil s/3 \rceil}[N_{\lceil s/3 \rceil}[N_{\lceil s/3 \rceil}[v]]$. Thus, we obtain a k^3 -vertex Turing kernel for $s = 6$ or $s \geq 8$. \square

Note that $s \geq 4$ is necessary to ensure $\lfloor s/2 \rfloor - 1 \geq 1$. In our arguments to obtain a Turing kernel $\ell = 1$ is necessary for the following reason: if $\ell \geq 2$, then the remaining vertices of the other triangles of a vertex in the T -expansion may be contained in $N_{\lfloor s/2 \rfloor + 1}$ and, thus, adding them will not necessarily give an s -club. Also note that using $N_t[v]$ for some $t < \lfloor s/2 \rfloor - 1$ does not help: The remaining vertices of the other triangles of a vertex in the T -expansion may be contained in N_{t+1} . But now, another T -expansion for the vertices in N_{t+1} is necessary. This may lead to a cascade of T -expansions where eventually, we add vertices with distance at least $s + 1$ to v . Thus, the constructed set is no s -club anymore.

2.2 Parameterized Hardness

In the following, we prove W[1]-hardness for VERTEX TRIANGLE s -CLUB parameterized by the solution size k for all cases not covered by Theorem 2, that is, $\ell \geq 2$ and $s \geq 2$, and also for $\ell = 1$ and $s \in \{2, 3\}$.

Theorem 3. *VERTEX TRIANGLE s -CLUB is W[1]-hard for parameter k if $\ell \geq 2$, and if $\ell = 1$ and $s \in \{2, 3\}$.*

For some combinations of s and ℓ we provide hardness for restricted input graphs. More precisely, we prove that VERTEX TRIANGLE s -CLUB is W[1]-hard even if each vertex $v \in V(G)$ is contained in *exactly* ℓ triangles in the input graph. In other words, the hardness does not depend on the fact that we could choose

different triangles. We provide this hardness for the case $s \geq 3$ and arbitrary ℓ and also for $s = 2$ when $\ell = \binom{c-1}{2}$ for some integer c .

We prove the theorem by considering four subcases. The proofs for the four cases all use a reduction from the W[1]-hard CLIQUE problem. In these constructions, each vertex v of the CLIQUE instance is replaced by a vertex gadget T^v such that every vertex- ℓ -triangle s -club S either contains T^v completely or contains no vertex of T^v . This property is obtained since each vertex in T^v will be in exactly ℓ triangles and each of these triangles is within T^v . The idea is that if $uv \notin E(G)$ then there exists a vertex $x \in T^u$ and a vertex $y \in T^v$ such that $\text{dist}(x, y) \geq s + 1$.

Vertex Triangle 2-Club. First, we handle the case $s = 2$.

Construction 4. Let (G, k) be an instance of CLIQUE and let c be the smallest number such that $\binom{c-1}{2} \geq \ell$. We construct an instance $(G', c(k+1))$ of VERTEX TRIANGLE 2-CLUB as follows.

- For each vertex $v \in V(G)$, we add a clique $T^v := \{x_1^v, \dots, x_c^v\}$ of size c to G' .
- For each edge $vw \in E(G)$, we connect the cliques T^v and T^w by adding the edge $x_{2i-1}^v x_{2i}^w$ and $x_{2i-1}^w x_{2i}^v$ to G' for each $i \in \lceil [c/2] \rceil$.
- Furthermore, we add a clique $Y := \{y_1, \dots, y_c\}$ of size c to G' .
- We also add, for each $i \in [c]$ and each $v \in V(G)$, the edge $x_i^v y_i$ to G' .

Note that the clique size c ensures that each vertex $x \in V(G')$ is contained in at least $\binom{c-1}{2} \geq \ell$ triangles in G' . Furthermore, note that the clique Y is only necessary when c is odd to ensure that the vertices x_c^v and x_c^w have a common neighbor. We add the clique Y in both cases to unify the construction and the correctness proof. Next, we show that for each vertex gadget T^v the intersection with each vertex- ℓ -triangle 2-club is either empty or T^v .

Lemma 2. *Let S be a vertex- ℓ -triangle 2-club in G' . Then,*

- a) $S \cap T^v \neq \emptyset \Leftrightarrow T^v \subseteq S$, and
- b) $S' := S \cup Y$ is also a vertex- ℓ -triangle 2-club in G' .

Proof. First, we show statement a). Assume that for a vertex $z \in T^v$ for some $v \in V(G)$ we have $z \in S$ for some vertex- ℓ -triangle 2-club S . Note that T^v contains all vertices which form a triangle with vertex z . Since c is minimal such that $\binom{c-1}{2} \geq \ell$ and since T^v is a clique, we conclude that $T^v \subseteq S$ to fulfill the property that vertex z is contained in at least ℓ triangles in $G[S]$. Thus, $T^v \subseteq S$.

Second, we show statement b). Since each vertex $y \in Y$ forms only triangles with vertices in Y and Y has size $\binom{c-1}{2}$, we conclude that $Y \subseteq S^* \Leftrightarrow S^* \cap Y \neq \emptyset$ for each vertex- ℓ -triangle 2-club S^* . In the following, let S be a vertex- ℓ -triangle 2-club such that $Y \cap S = \emptyset$. From statement a) we conclude that $S := \bigcup_{v \in P} T^v$ for some set $P \subseteq V(G)$. Next, we show that $S' := S \cup Y$ is also a vertex- ℓ -triangle 2-club. Since each vertex is contained in a clique of size c in S , it is in sufficiently many triangles. Thus, it remains to prove that S' is a 2-club. Hence, consider some vertex x_i^v and some vertex y_j for some $i, j \in [c]$ and $v \in P$. Then, y_i is a common neighbor of x_i^v and y_j . Hence, S' is a vertex- ℓ -triangle 2-club and thus b) holds. \square

Now, we prove the correctness of Construction 4.

Lemma 3. *For each $\ell \in \mathbb{N}$, the VERTEX TRIANGLE 2-CLUB problem parameterized by k is $W[1]$ -hard.*

Proof. We prove that G contains a clique of size k if and only if G' contains a vertex- ℓ -triangle 2-club of size at least $c(k+1)$.

Let C be a clique of size at least k in G . We argue that $S := Y \cup \bigcup_{v \in C} T^v$ is a vertex- ℓ -triangle 2-club of size $c(k+1)$ in G' . Note that for each vertex $v \in T^v$ we have $|T^v| = c$. Since T^v is a clique, we conclude that each vertex in T^v is contained in exactly $\binom{c-1}{2} \geq \ell$ triangles. The same is true for each vertex in Y . Hence, each vertex in S is contained in at least ℓ triangles. Thus, it remains to show that S is a 2-club. Consider the vertices x_i^v and x_j^w for $v, w \in C$, $i \in [c-1]$, and $j \in [c]$. If i is odd, then $x_{i+1}^w \in N(x_i^v) \cap N(x_j^w)$. Otherwise, if i is even, $x_{i-1}^w \in N(x_i^v) \cap N(x_j^w)$. In both cases, we obtain $\text{dist}(x_i^v, x_j^w) \leq 2$. Next, consider two vertices x_c^v and x_c^w in S . Observe that $y_c \in N(x_c^v) \cap N(x_c^w)$. Since Y is a clique, it remains to consider vertices x_i^v and y_j in S for $i \in [c]$ and $j \in [c]$. Observe that $x_j^v \in N(y_j) \cap N[x_i^v]$. Thus, G' contains a vertex- ℓ -triangle 2-club of size at least $c(k+1)$.

Conversely, suppose that G' contains a vertex- ℓ -triangle 2-club S of size at least $c(k+1)$. By Lemma 2, we can assume that $Y \subseteq S$ and for each vertex gadget $T^v \in G'$ we either have $T^v \subseteq S$ or $T^v \cap S = \emptyset$. Hence, S contains at least k cliques of the form T^v . Assume towards a contradiction that S contains two cliques T^v and T^w such that $vw \notin E(G)$ and consider the two vertices $x_1^v \in T^v$ and $x_2^w \in T^w$. Note that these vertices always exist since $c \geq 3$. Observe that $N[x_1^v] = T^v \cup \{x_2^u \mid uv \in E(G)\} \cup \{y_1\}$ and $N[x_2^w] = T^w \cup \{x_1^u \mid uw \in E(G)\} \cup \{y_2\}$. Thus, $\text{dist}(x_1^v, x_2^w) \geq 3$, a contradiction. Hence, for each two distinct vertex gadgets T^v and T^w that are contained in S , we observe that $vw \in E(G)$. Consequently, the set $\{v \mid T^v \subseteq S\}$ is a clique of size at least k in G . \square

If $\ell = \binom{c-1}{2}$ for some integer c , then Lemma 3 also holds for the restriction that each vertex is contained in exactly ℓ triangles in the input graph G' .

Vertex Triangle s -Club for $s = 3$ and for $\ell \geq 2$ and $s \geq 4$. Now, we provide hardness for the remaining cases. We consider three subcases. Case 1 deals with odd s . Case 2 covers the case that s is even and $\ell \geq 3$. Case 3 deals with the case that s is even and $\ell = 2$. All three cases use the same vertex gadget. Only the edges between these gadgets, called *connector edges*, differ. The idea is to construct the vertex gadgets T^v in such a way that there are pairs of vertices in T^v of distance $2s^*$ which is almost s . Thus, in a vertex- ℓ -triangle s -club, the distance between two different vertex gadgets must be small. The first part of the following construction describes the vertex gadget which is used in all three cases. For an illustration of this construction see Fig. 1.

Construction 5. We set $s^* := \lfloor (s-1)/2 \rfloor$. Let (G, k) be an instance of CLIQUE. We construct an equivalent instance $(G', 3\ell k s^*)$ of VERTEX TRIANGLE s -CLUB. Recall that $\ell \geq 1$ and $s = 3$, or $\ell \geq 2$ and $s \geq 4$. For each vertex $v \in V(G)$ we




Fig. 1. a) The vertex gadgets T^v and T^w for $s \in \{7, 8\}$ and $\ell = 2$. The blue lines are only added if s is odd and the red lines are only added if s is even. b) The vertex gadget T^v for $s \in \{5, 6\}$ and $\ell = 3$.

construct a vertex gadget T^v . This vertex gadget is used for each reduction of the three cases.

- We add the vertices p_i^v , and q_i^v , and an edge $p_i^v q_i^v$ for each $i \in [\ell]$ to G' .
- We add a vertex $x_{j,i}^v$ for each $i \in [\ell]$ and each $j \in [s^*]$ to G' .
- We add an edge $y_{t,i}^v z_{t,i}^v$ for each $i \in [\ell]$ and each $t \in [s^* - 1]$ to G' . Note that these vertices only exist, if $s \geq 5$.

The vertices $x_{j,i}^v$, $y_{t,i}^v$, and $z_{t,i}^v$ are referred to as the *cascading vertices*. They are used to ensure that all vertices in T^v are in exactly ℓ triangles and that there are vertex pairs of distance $2s^*$ within T^v . Note that since $s \geq 3$ we create at least ℓ many x -vertices. We connect these vertices as follows:

- We add the edges $p_i^v x_{1,j}^v$ and $q_i^v x_{1,j}^v$ for each $i \in [\ell - 1]$ and each $j \in [\ell]$ to G' .
- We add the edges $p_\ell^v x_{s^*,j}^v$ and $q_\ell^v x_{s^*,j}^v$ for each $j \in [\ell]$ to G' .
- We add the edges $y_{t,i}^v x_{t,i}^v$ and $z_{t,i}^v x_{t,i}^v$ for each $i \in [\ell]$, and each $t \in [s^* - 1]$ to G' .
- We add the edges $y_{t,i}^v x_{t+1,j}^v$ and $z_{t,i}^v x_{t+1,j}^v$ for each $i \in [\ell]$, each $j \in [\ell] \setminus \{i\}$, and each $t \in [s^* - 1]$ to G' .

Note that if $s = 3$ or $s = 4$, the graph G' is a non-induced biclique where one partite set consists of the vertices $\{x_{1,j}^v \mid j \in [\ell]\}$ and the other partite set consists of the vertices $\{p_i^v, q_i^v \mid i \in [\ell]\}$. Furthermore, the additional edges are $p_i^v q_i^v$ for each $i \in [\ell]$. Also, observe that each vertex gadget T^v consists of exactly $3\ell s^*$ vertices.

From now on, the construction for the three cases differs. We now connect these vertex gadgets by introducing the *connector edges*: For each edge $vw \in$

$E(G)$ we add edges between the vertex gadgets T^v and T^w of the corresponding vertices. We distinguish the three cases.

Case I: s is odd. We add the edges $p_i^v q_i^w$ and $q_i^v p_i^w$ to G' for each $i \in [\ell]$ to G' .

Case II: s is even and $\ell \geq 3$. We add the edges $p_1^v q_1^w$, $q_1^v p_1^w$, $p_\ell^v q_\ell^w$, and $q_\ell^v p_\ell^w$ to G' .

Case III: s is even and $\ell = 2$. We add the edges $p_1^v x_{s^*,1}^w$, and $p_1^w x_{s^*,1}^v$ to G' .

We make the following observation about the connector edges between different vertex gadgets: If s is odd (Case I), or if s is even and $\ell \geq 3$ (Case II), we have $N(p_i^v) \setminus T^v \subseteq \{q_i^w \mid vw \in E(G)\}$ and also $N(q_i^v) \setminus T^v \subseteq \{p_i^w \mid vw \in E(G)\}$ for each $v \in V(G)$ and each $i \in [\ell]$. Otherwise, if s is even and $\ell = 2$ (Case III), observe that we have $N(p_1^v) \setminus T^v = \{x_{s^*,1}^w \mid vw \in E(G)\}$ and also $N(x_{s^*,1}^v) \setminus T^v = \{p_1^w \mid vw \in E(G)\}$ for each $v \in V(G)$. Since only the connector edges have endpoints in different vertex gadgets, we thus observe the following.

Observation 6. *All 3 endpoints of each triangle in G' are contained in exactly one vertex gadget.*

Next, we show that each vertex in a vertex gadget T^v is contained in exactly ℓ triangles. Together with Observation 6 this implies that each vertex in G' is contained in exactly ℓ triangles.

Lemma 4. *Let T^v be a vertex gadget. Each vertex in T^v is contained in exactly ℓ triangles.*

Proof. We make a case distinction, that is, for each vertex $a \in T^v$ we present exactly ℓ triangles containing vertex a . We distinguish the different vertices of a vertex gadget.

Case 1: Consider vertex p_d^v for some $d \in [\ell - 1]$. Because of Observation 6 we only have to consider the neighbors of p_d^v in T^v . By construction we have $N(p_d^v) \cap T^v = \{q_d^v\} \cup \{x_{1,j}^v \mid j \in [\ell]\}$. Observe that the only edges within $N(p_d^v) \cap T^v$ are the edges $q_d^v x_{1,j}^v$ for each $j \in [\ell]$. Thus, p_d^v is contained in the ℓ triangles $\{\{p_d^v, q_d^v, x_{1,i}^v\} \mid i \in [\ell]\}$. By similar arguments the same is true for vertex q_d^v .

Case 2: Consider vertex p_ℓ^v . By construction we have $N(p_\ell^v) = \{q_\ell^v\} \cup \{x_{s^*,i}^v \mid i \in [\ell]\} \cup \{q_\ell^w \mid vw \in E(G)\}$ if s is odd or s is even and $\ell \geq 3$. If s is even and $\ell = 2$ we have $N(p_\ell^v) = \{q_\ell^v\} \cup \{x_{s^*,i}^v \mid i \in [\ell]\}$. Observe that the only edges within $N(p_\ell^v)$ are the edges $q_\ell^v x_{s^*,i}^v$ for each $i \in [\ell]$. Thus, p_ℓ^v is contained in the ℓ triangles $\{\{p_\ell^v, q_\ell^v, x_{s^*,i}^v\} \mid i \in [\ell]\}$. By similar arguments the same is true for vertex q_ℓ^v .

Case 3: Now, consider vertices $x_{1,i}^v$ and $x_{s^*,i}^v$ for some $i \in [\ell]$. Here we have to distinguish if $s \in \{3, 4\}$ or $s \geq 5$ since $y_{t,i}^v$ exists only in the second case.

First, consider the case $s = 3$ or $s = 4$. Note that we now have $x_{1,i}^v = x_{s^*,i}^v$. By construction we have $N(x_{1,i}^v) = \{p_j^v \mid j \in [\ell]\} \cup \{q_j^v \mid j \in [\ell]\}$. Note that the only edges within $N(x_{1,i}^v)$ have the form $p_j^v q_j^v$ for each $j \in [\ell]$. Thus, $x_{1,i}^v$ is contained in the ℓ triangles $\{\{x_{1,i}^v, p_j^v, q_j^v\} \mid j \in [\ell]\}$.

Second, consider the case $s \geq 5$. First, we investigate vertex $x_{1,i}^v$. By construction we have $N(x_{1,i}^v) = \{p_j^v \mid j \in [\ell - 1]\} \cup \{q_j^v \mid j \in [\ell - 1]\} \cup \{y_{1,i}^v, z_{1,i}^v\}$. The only edges within $N(x_{1,i}^v)$ are the edge $p_j^v q_j^v$ for each $j \in [\ell - 1]$ and the edge $y_{1,i}^v z_{1,i}^v$. Thus, $x_{1,i}^v$ is contained in the $\ell - 1$ triangles $\{\{x_{1,i}^v, p_j^v, q_j^v\} \mid j \in [\ell - 1]\}$, and in the triangle $\{x_{1,i}^v, y_{1,i}^v, z_{1,i}^v\}$.

Now, consider vertex $x_{s^*,i}^v$. Because of Observation 6 we only have to consider the neighbors of $x_{s^*,i}^v$ in T^v . By construction we observe that $N(x_{s^*,i}^v) \cap T^v \subseteq \{p_\ell^v, q_\ell^v\} \cup \{y_{s^*-1,j}^v, z_{s^*-1,j}^v \mid j \in [\ell] \setminus \{i\}\}$. Note that the only edges within $N(x_{s^*,i}^v)$ are the edge $p_\ell^v q_\ell^v$ and the edge $y_{s^*-1,j}^v z_{s^*-1,j}^v$ for each $j \in [\ell] \setminus \{i\}$. Thus, $x_{s^*,i}^v$ is contained in the triangle $\{x_{s^*,i}^v, p_\ell^v, q_\ell^v\}$ and in the $\ell - 1$ triangles $\{\{x_{s^*,i}^v, y_{s^*-1,j}^v, z_{s^*-1,j}^v\} \mid j \in [\ell] \setminus \{i\}\}$.

Case 4: Consider vertex $x_{r,i}^v$ for some $i \in [\ell]$ and some $r \in [2, s^* - 1]$. Recall that these vertices only exist if $s \geq 7$. By construction $N(x_{r,i}^v) = \{y_{r,i}^v, z_{r,i}^v\} \cup \{y_{r-1,j}^v, z_{r-1,j}^v \mid j \in [\ell] - 1\}$. The only edges within $N(x_{r,i}^v)$ have the form $y_{r,i}^v z_{r,i}^v$ and $y_{r-1,j}^v z_{r-1,j}^v$ for each $j \in [\ell] \setminus \{i\}$. Thus, $x_{r,i}^v$ is contained in the triangle $\{x_{r,i}^v, y_{r,i}^v, z_{r,i}^v\}$ and in the $\ell - 1$ triangles $\{\{x_{r,i}^v, y_{r-1,j}^v, z_{r-1,j}^v\} \mid j \in [\ell] \setminus \{i\}\}$.

Case 5: Finally, consider vertex $y_{t,i}^v$ for some $i \in [\ell]$ and some $t \in [s^* - 1]$. Recall that these vertices only exist if $s \geq 5$. By construction we have $N(y_{t,i}^v) = \{x_{t,i}^v, z_{t,i}^v\} \cup \{x_{t+1,j}^v \mid j \in [\ell] \setminus \{i\}\}$. The only edges within $N(y_{t,i}^v)$ is the edge $x_{t,i}^v z_{t,i}^v$ and the edge $x_{t+1,j}^v z_{t,i}^v$ for each $j \in [\ell] \setminus \{i\}$. Thus, $y_{t,i}^v$ is contained in the triangle $\{y_{t,i}^v, z_{t,i}^v, x_{t,i}^v\}$ and in the $\ell - 1$ triangles $\{\{y_{t,i}^v, z_{t,i}^v, x_{t+1,j}^v\} \mid j \in [\ell] \setminus \{i\}\}$. By similar arguments the same holds for vertex $z_{t,i}^v$.

Thus, each vertex in T^v is contained in exactly ℓ triangles. \square

From Lemma 4 and Observation 6, we conclude the following.

Observation 7. *Let S be a vertex- ℓ -triangle s -club for $s = 3$ and $\ell \geq 1$ and or for $s \geq 4$ and $\ell \geq 2$ in G' . Then, $S \cap T^v \neq \emptyset \Leftrightarrow T^v \subseteq S$.*

Now, we prove the correctness of Construction 5.

Lemma 5. *For each $s = 3$ and $\ell \geq 1$, and also for each $s \geq 4$ and $\ell \geq 2$ the VERTEX TRIANGLE s -CLUB problem parameterized by k is $W[1]$ -hard, even if each vertex in the input graph is contained in exactly ℓ triangles.*

Proof. We show that G contains a clique of size at least k if and only if G' contains a vertex- ℓ -triangle s -club of size at least $3\ell ks^*$.

Let K be a clique of size at least k in G . We argue that $S := \bigcup_{v \in K} T^v$ is a vertex- ℓ -triangle s -club of size at least $3\ell ks^*$ in G' . The size bound follows from the fact that each T^v consists of exactly $3\ell s^*$ vertices. Furthermore, by Lemma 4 each vertex in T^v for some $v \in V(G)$ is contained in exactly ℓ triangles in $G'[T^v]$. Hence, it remains to show that S is an s -club.

To do so, we first prove the following two claims. To formulate the claims, we need some further notation. We define $T_0^v := \{p_1^v, \dots, p_{\ell-1}^v\} \cup \{q_1^v, \dots, q_{\ell-1}^v\}$ and $T_\ell^v := \{p_\ell^v, q_\ell^v\}$. Recall that if $\ell = 1$, then $T_0^v = \emptyset$. Otherwise, both sets are nonempty. This claim

Claim 1. For $\ell \geq 2$, we have $\text{dist}_{G'}(u, a) + \text{dist}_{G'}(u, b) = 2s^*$ for each vertex $a \in T_0^v$, each vertex $b \in T_\ell^v$, and for each vertex $u \in T^v \setminus (T_0^v \cup T_\ell^v)$.

Proof. By $X_1^v := \{x_{1,j}^v \mid j \in [\ell]\}$ we denote the neighbors of T_0^v . We define the sets $X_2^v, \dots, X_{s^*}^v$ and $Y_1^v, Z_1^v, \dots, Y_{s^*-1}^v, Z_{s^*-1}^v$ analogue. We first make some observations about the neighborhoods of these vertex sets which then allows us to obtain upper and lower bounds on these paths.

- $T_0^v \cup \{y_{1,i}^v, z_{1,i}^v\} \subseteq N(x_{1,i}^v)$ for each $i \in [\ell]$. Hence $N(X_1^v) = T_0^v \cup Y_1^v \cup Z_1^v$.
- $T_\ell^v \cup \{y_{s^*-1,j}^v, z_{s^*-1,j}^v\} \subseteq N(x_{s^*,i}^v)$ for each $i \in [\ell]$ and each $j \in [\ell] \setminus \{i\}$. Hence, $N(X_{s^*}^v) = T_\ell^v \cup Y_{s^*-1}^v \cup Z_{s^*-1}^v$.
- $\{y_{t,i}^v, y_{t-1,j}^v, z_{t,i}^v, z_{t-1,j}^v\} \subseteq N(x_{t,i}^v)$ for each $t \in [2, s^* - 1]$, each $i \in [\ell]$ and each $j \in [\ell] \setminus \{i\}$. Hence, $N(X_t^v) = Y_t^v \cup Z_t^v \cup Y_{t-1}^v \cup Z_{t-1}^v$ for $t \in [2, s^* - 1]$.
- $x_{t,i}^v, x_{t+1,j}^v, z_{t,i}^v \in N(y_{t,i}^v)$ and also $x_{t,i}^v, x_{t+1,j}^v, y_{t,i}^v \in N(z_{t,i}^v)$ for each $t \in [s^* - 1]$, each $i \in [\ell]$ and each $j \in [\ell] \setminus \{i\}$. Hence, $N(Y_t^v) = X_t^v \cup Z_t^v \cup Y_{t-1}^v$ and $N(Z_t^v) = X_t^v \cup Y_t^v \cup Y_{t-1}^v$ for each $t \in [s^* - 1]$.

From the above and a vertex $u \in X_z^v$ we obtain a path to a of length $2z - 1$ by using subsequent vertices from $Y_{z-1}, X_{z-1}, \dots, Y_1, X_1$. A similar observation can be made for a path from u to b . The cases $u \in Y_z \cup Z_z$ is treated similar. This implies that $\text{dist}_{G'}(u, a) + \text{dist}_{G'}(u, b) \leq 2s^*$ for each vertex $a \in T_0^v$, each vertex $b \in T_\ell^v$, and for each vertex $u \in T^v \setminus (T_0^v \cup T_\ell^v)$. Furthermore, note that in each two steps on any path we can increase/decrease the first sub-index by one. Hence $\text{dist}_{G'}(u, a) + \text{dist}_{G'}(u, b) \geq 2s^*$. Thus, $\text{dist}_{G'}(u, a) + \text{dist}_{G'}(u, b) = 2s^*$. ■

Now, we use Claim 1 to show that T^v is a $(2s^*)$ -club.

Claim 2. If s is odd, then T^v is an $(s - 1)$ -club and if s is even, then T^v is an $(s - 2)$ -club for each vertex $v \in V(G)$.

Proof. Recall that for $s \in \{3, 4\}$ the gadget T^v is a non-induced biclique. Hence, the statement is true in this case. In the following, we assume that $s \geq 5$. Note that this implies that $\ell \geq 2$. We only consider the case that s is odd. The case that s is even follows analogously.

Consider a pair of vertices u, \tilde{u} of $T^v \setminus (T_0^v \cup T_\ell^v)$. To bound the distance of u and \tilde{u} , we consider one path via a vertex in T_0^v and one via a vertex in T_ℓ^v . We have $\text{dist}_{G'}(u, \tilde{u}) \leq \min(\text{dist}_{G'}(u, a) + \text{dist}_{G'}(a, \tilde{u}), \text{dist}_{G'}(u, b) + \text{dist}_{G'}(b, \tilde{u}))$ for each vertex $a \in T_0^v$, and each vertex $b \in T_\ell^v$. From Claim 1 we know that $\text{dist}_{G'}(u, a) + \text{dist}_{G'}(u, b) = 2s^*$ and that $\text{dist}_{G'}(\tilde{u}, a) + \text{dist}_{G'}(\tilde{u}, b) = 2s^*$. Hence, $\text{dist}_{G'}(u, \tilde{u}) \leq 2s^* \leq s - 1$. Thus, T^v is indeed an $(s - 1)$ -club. ■

Now, we show that for two vertices v and w in K , a vertex $u \in T^v$, and a vertex $\tilde{u} \in T^w$ we have $\text{dist}_{G'}(u, \tilde{u}) \leq s$. We consider the three cases:

Case I: s is odd. Observe that since $vw \in E(G)$, each vertex $u \in (T_0^v \cup T_\ell^v)$ has one neighbor in T^w . Since T^w is an $(s - 1)$ -club by Claim 2, we obtain that for each vertex $\tilde{u} \in T^w$ we have $\text{dist}_{G'}(u, \tilde{u}) \leq s$ if $u \in T_0^v \cup T_\ell^v$. Hence, it remains to consider the case that $u \in T^v \setminus (T_0^v \cup T_\ell^v)$ and that $\tilde{u} \in T^w \setminus (T_0^w \cup T_\ell^w)$. For this, let $u_1^v := \text{dist}_{G'}(u, p_1^v)$, $\tilde{u}_1^w := \text{dist}_{G'}(q_1^w, \tilde{u})$, $u_\ell^v := \text{dist}_{G'}(u, p_\ell^v)$, and $\tilde{u}_\ell^w :=$

$\text{dist}_{G'}(q_\ell^w, \tilde{u})$. Note that

$$\begin{aligned} \text{dist}_{G'}(u, \tilde{u}) &\leq \min(u_1^v + 1 + \tilde{u}_1^w, u_\ell^v + 1 + \tilde{u}_\ell^w) \\ &= 1 + \min(u_1^v + \tilde{u}_1^w, u_\ell^v + \tilde{u}_\ell^w). \end{aligned}$$

In this inequality, the '+1' is the result of the fact that we have to use an edge to get from gadget T^v to gadget T^w . By Claim 1 we know that $u_1^v + u_\ell^v = 2s^*$ and that $\tilde{u}_1^w + \tilde{u}_\ell^w = 2s^*$. Since s is odd, we obtain that $\text{dist}_{G'}(u, \tilde{u}) \leq 1 + 2s^* = s$. Thus, S is a vertex- ℓ -triangle s -club of size $3\ell ks^*$.

Case II: s is even and $\ell \geq 3$. For each vertex pair $u \in T^v \setminus T_0^v$ and $\tilde{u} \in T^w \setminus T_0^w$ the proof of $\text{dist}_{G'}(u, \tilde{u}) \leq s$ is analogous to the proof in Case 1 handling odd values of s . Hence, it remains to show that each vertex $u \in T_0^v$ has distance at most s to each vertex $\tilde{u} \in T^w$. Observe that $\text{dist}_{G'}(u, p_1^v) \leq 2$ since $T_0^v \subseteq N(x_{1,1}^v)$ and $x_{1,1}^v$ is a neighbor of p_1^v . Thus, $\text{dist}_{G'}(u, q_1^w) = 3$ since $vw \in E(G)$. Furthermore, for each vertex $\tilde{u} \in T^w \setminus T_\ell^w$ we have $\text{dist}_{G'}(\tilde{u}, q_1^w) \leq s - 3$ by the proof of Claim 2. Hence, $\text{dist}(u, \tilde{u}) \leq s$. Thus, it remains to consider the case that $u \in T_0^v$ and that $\tilde{u} \in T_\ell^w = \{p_\ell^w, q_\ell^w\}$. Since $vw \in E(G)$ we conclude that \tilde{u} has a neighbor in T^v . Since T^v is an $(s - 2)$ -club by Claim 2 we conclude that $\text{dist}_{G'}(u, \tilde{u}) = s - 1$. Hence, S is a vertex- ℓ -triangle s -club of size $3\ell ks^*$.

Case III: s is even and $\ell = 2$. Let $x^w := x_{s^*,1}^w$ and $x^v := x_{s^*,1}^v$. Furthermore, let $u_1^v := \text{dist}_{G'}(u, p_1^v)$, $\tilde{u}_x^w := \text{dist}_{G'}(x^w, \tilde{u})$, $u_x^v := \text{dist}_{G'}(u, x^v)$, and $\tilde{u}_1^w := \text{dist}_{G'}(p_1^w, \tilde{u})$. We have

$$\begin{aligned} \text{dist}_{G'}(u, \tilde{u}) &\leq \min(u_1^v + 1 + \tilde{u}_x^w, u_x^v + 1 + \tilde{u}_1^w) \\ &= 1 + \min(u_1^v + \tilde{u}_x^w, u_x^v + \tilde{u}_1^w). \end{aligned}$$

Again, the '+1' is the result that we have to use an edge to go from T^v to gadget T^w . Consider the following claim.

Claim 3. For each vertex $u \in T^v$ we have $u_1^v + u_x^v \leq s - 1$.

Claim 3 directly implies that

$$\min(u_1^v + \tilde{u}_x^w, u_x^v + \tilde{u}_1^w) \leq s - 1$$

and thus $\text{dist}_{G'}(u, \tilde{u}) \leq s$. This in turn implies that S is a vertex- ℓ -triangle s -club of size $3\ell ks^*$. Thus, it remains to prove Claim 3.

Proof. Since T^v is an $(s - 2)$ -club by Claim 2, we conclude that each vertex $u \in (N[p_1^v] \cup N[x_{s^*,1}^v]) \cap T^v$ has distance at most s to each vertex in T^v . Hence, it remains to show that each vertex $u \in T^v \setminus (N[p_1^v] \cup N[x_{s^*,1}^v])$ has distance at most s to each vertex $u' \in T^v$. If $s = 4$, then $T^v \subseteq (N[p_1^v] \cup N[x_{s^*,1}^v])$ and hence the claim is proven. Hence, in the following, we consider the case $s \geq 6$.

Next, we consider the case that $u \in T_0^v$. Since $T_0^v \subseteq N(x_{1,1}^v)$ we conclude that $u_1^v \leq 2$. By the proof of Claim 2 we conclude that $u_x^v = s - 3$. Hence, Claim 3 is true in this case.

For all remaining cases, it is sufficient to prove the claim for each vertex $u \in T^v \setminus (T_0^v \cup T_\ell^v)$. First, we consider the case that $u := x_{t,i}^v$ for some $t \in [s^*]$ and

some $i \in [\ell]$. Then we have $\text{dist}_{G'}(x_{t,i}^v, p_1^v) = 2t - 1$ and $\text{dist}_{G'}(x_{t,i}^v, x_{s^*,1}^v) \leq 2(s^* - t) + 2$ and hence $\text{dist}_{G'}(x_{t,i}^v, p_1^v) + \text{dist}_{G'}(x_{t,i}^v, x_{s^*,1}^v) \leq 2t - 1 + s^* - 2t + 2 = s^* + 1 \leq s - 1$.

Second, we consider the case that $u := y_{t,i}^v$ for some $t \in [s^* - 1]$ and some $i \in [\ell]$. Then we have $\text{dist}_{G'}(y_{t,i}^v, p_1^v) = 2t$ and $\text{dist}_{G'}(y_{t,i}^v, x_{s^*,1}^v) \leq 2(s^* - t) + 1$ and hence we obtain that $\text{dist}_{G'}(y_{t,i}^v, p_1^v) + \text{dist}_{G'}(y_{t,i}^v, x_{s^*,1}^v) \leq 2t + s^* - 2t + 1 = s^* + 1 \leq s - 1$. The case that $u := z_{t,i}^v$ follows by the same argumentation. \blacksquare

Conversely, suppose that G' contains a vertex- ℓ -triangle s -club of size at least $3\ell k s^*$. Because of Observation 7 for each vertex-gadget T^v we either have $T^v \subseteq S$ or $T^v \cap S = \emptyset$. Hence, S contains at least k vertex gadgets. We assume towards a contradiction that S contains two vertex gadgets T^v and T^w such that $vw \notin E(G)$. In each case, we will determine a vertex $u_v \in T^v$ and a vertex $u_w \in T^w$ such that $\text{dist}_{G'}(u_v, u_w) \geq s + 1$. This contradiction to the s -club property allows us to conclude that the set $\{v \mid T^v \subseteq S\}$ is a clique of size at least k in G .

Case I: s is odd. We define vertex u_v as follows. The vertex u_w is defined analogously. Recall that in this case we have $s^* = (s - 1)/2$.

- If $s \equiv 3 \pmod{4}$, we set $u_v := x_{(s+1)/4,1}^v$.
- Otherwise, if $s \equiv 1 \pmod{4}$, we set $u_v := y_{(s-1)/4,1}^v$.

Observe that for each vertex $u \in T_0^v \cup T_\ell^v$ we have $\text{dist}_{G'}(u_v, u) = (s - 1)/2$. Furthermore, recall that the vertices in $T_0^v \cup T_\ell^v$ are the only vertices in T^v with neighbors in other vertex gadgets.

Similarly, for each vertex $u' \in T_0^w \cup T_\ell^w$ we have $\text{dist}_{G'}(u_w, u') = (s - 1)/2$. But since $vw \notin E(G)$ there are no edges between T^v and T^w , we obtain for each choice of u and u' that $\text{dist}_{G'}(u_v, u_w) \geq \text{dist}_{G'}(u_v, u) + 2 + \text{dist}_{G'}(u', u_w) = (s - 1)/2 + 2 + (s - 1)/2 = s + 1$, a contradiction.

Case II: s is even and $\ell \geq 3$. We set $u_v := p_2^v$ and $u_w := x_{s^*,1}^w$. Note that $u_v \in T_0^v$ since $\ell \geq 3$. By the construction we obtain that for each vertex $u_0 \in \{p_1^v, q_1^v\}$ we have $\text{dist}_{G'}(u_v, u_0) = 2$ since $T_0^v \subseteq N(x_{1,1}^v)$. Furthermore, for each vertex $u_\ell \in \{p_\ell^v, q_\ell^v\}$ we have $\text{dist}_{G'}(u_v, u_\ell) = s - 2$. Next, observe that for each vertex $u'_0 \in \{p_1^w, q_1^w\}$ we have $\text{dist}_{G'}(u_w, u'_0) = s - 3$ and for each vertex $u'_\ell \in \{p_\ell^w, q_\ell^w\}$ we have $\text{dist}(u_w, u'_\ell) = 1$.

Since $vw \notin E(G)$, there are no edges between the gadgets T^v and T^w . Hence, $\text{dist}(u_v, u_w) \geq \min(\text{dist}(u_v, u_0) + 2 + \text{dist}_{G'}(u'_0, u_w), \text{dist}_{G'}(u_v, u_\ell) + 2 + \text{dist}_{G'}(u'_\ell, u_w))$ for each vertex $u_0 \in \{p_1^v, q_1^v\}$, each $u'_0 \in \{p_1^w, q_1^w\}$, each $u_\ell \in T_\ell^v$, and each $u'_\ell \in T_\ell^w$. By the above argumentation we obtain $\text{dist}_{G'}(u_v, u_w) \geq \min(2 + 2 + s - 3, s - 2 + 2 + 1) = s + 1$, a contradiction.

Case III: s is even and $\ell = 2$. We define the vertices u_v and u_w as follows:

- If $s = 4$, we set $u_v := x_{1,2}^v$ and $u_w := p_2^w$.
- If $s \equiv 0 \pmod{4}$ and $s \geq 8$, we set $u_v := x_{s/4,1}^v$ and $u_w := y_{s/4,1}^w$.
- If $s \equiv 2 \pmod{8}$, we set $u_v := y_{(s-2)/4,2}^v$ and $u_w := x_{(s+2)/4,1}^w$.
- If $s \equiv 6 \pmod{8}$, we set $u_v := y_{(s-2)/4,1}^v$ and $u_w := x_{(s+2)/4,2}^w$.

From the definition of these vertices we obtain that $\text{dist}_{G'}(u_v, p_1^v) = s/2 - 1$, $\text{dist}_{G'}(u_v, x_{s^*,1}^v) = s/2$, $\text{dist}_{G'}(u_w, p_1^w) = s/2$, and that $\text{dist}_{G'}(u_w, x_{s^*,1}^w) = s/2 - 1$. Recall that the vertices p_1^u and $x_{s^*,1}^u$ are the only vertices in T^u which have neighbors outside T^u for each $u \in V(G)$. Furthermore, observe that all neighbors of p_1^u which are not contained in T^u are the vertices $x_{s^*,1}^b$ where $ub \in E(G)$. Similar, all neighbors of $x_{s^*,1}^u$ which are not in T^u are of the form p_1^b where $ub \in E(G)$. We conclude that $\text{dist}_{G'}(u_v, u_w) \geq \text{dist}(u_v, p_1^v) + 3 + \text{dist}(x_{s^*,1}^w, u_w)$. Here, the '+3' results from the fact that at least 3 edges to switch the vertex gadgets have to be used: one is not sufficient since $uw \notin E(G)$ and also two are not sufficient since in two steps one can only reach a vertex p_1^c for $c \in V(G)$ from p_1^v but no vertex $x_{s^*,1}^d$ for $d \in V(G)$. Hence, $\text{dist}_{G'}(u_v, u_w) \geq s/2 - 1 + 3 + s/2 - 1 = s + 1$, a contradiction. \square

3 Edge Triangle s -Club

In this section we settle the parameterized complexity of EDGE TRIANGLE s -CLUB with respect to the solution size k . Recall that a vertex set S is an edge- ℓ -triangle s -club if $G[S]$ contains a spanning subgraph $G' = (S, E')$ such that each edge in $E(G')$ is contained in at least ℓ triangles within G' and the diameter of G' is at most s . First, we show that EDGE TRIANGLE s -CLUB is FPT with respect to k when $\ell = 1$ irrespective of the value of s by providing a Turing kernel. To show this, it is sufficient to delete edges which are not part of a triangle. Afterwards, we prove W[1]-hardness of EDGE TRIANGLE s -CLUB with respect to k for all fixed $\ell \geq 2$.

3.1 Edge Triangle s -Club with $\ell = 1$

Now, we prove that EDGE TRIANGLE s -CLUB for $\ell = 1$ admits a Turing kernel with respect to k implying that the problem is FPT. To obtain the kernel we need the following reduction rule which removes edges which are in no triangle.

Reduction Rule 8. *Let (G, k) be an instance of EDGE TRIANGLE s -CLUB. Delete all edges from G which are not part of any triangle.*

It is clear that Reduction Rule 8 is correct and can be applied in polynomial time. The idea that after Reduction Rule 8 is applied, we can bound the size of the neighborhood of each vertex. Next, we prove that after the application of Reduction Rule 8 each edge with both endpoints in the closed neighborhood of a vertex is contained in a triangle.

Lemma 6. *Let (G, k) be an instance of EDGE TRIANGLE s -CLUB with $\ell = 1$ to which Reduction Rule 8 is applied. For each vertex $v \in V(G)$ each edge in $G[N[v]]$ is contained in at least one triangle in $G[N[v]]$.*

Proof. First, we consider edges of the form uv where u is a neighbor of v . Since Reduction Rule 8 is applied, there exists another vertex $w \in V(G)$ such that $G[\{u, v, w\}]$ is a triangle. Observe that $w \in N(v)$.

Second, each edge uw with $u, w \in N(v)$ is in a triangle with vertex v . \square

Next, we show that any instance such that sufficiently many vertices are close to some vertex v is a yes-instance.

Lemma 7. *Let (G, k) be an instance of EDGE TRIANGLE s -CLUB with $\ell = 1$ to which Reduction Rule 8 is applied. Then, (G, k) is a yes-instance if $|N_{\lfloor s/2 \rfloor}[v]| \geq k$ for some vertex $v \in V(G)$.*

Proof. By Lemma 6 each edge in $N_{\lfloor s/2 \rfloor}[v]$ is contained in at least one triangle since $N_{\lfloor s/2 \rfloor}[v] = \bigcup_{w \in N_{\lfloor s/2 \rfloor - 1}[v]} N[w]$. Furthermore, each vertex in $N_{\lfloor s/2 \rfloor}[v]$ has distance at most $\lfloor s/2 \rfloor$ to vertex v . Hence, $N_{\lfloor s/2 \rfloor}[v]$ is an s -club and by definition $|N_{\lfloor s/2 \rfloor}[v]| \geq k$. \square

Lemma 7 implies a Turing kernel for k which implies that the problem is fixed-parameter tractable.

Theorem 9. *EDGE TRIANGLE s -CLUB for $\ell = 1$ admits a k^2 -vertex Turing kernel if s is even and a k^3 -vertex Turing kernel if s is odd and $s \geq 3$.*

Proof. First, we apply Reduction Rule 8 exhaustively. Because of Lemma 7 we conclude that (G, k) is a trivial yes-instance, if $|N_{\lfloor s/2 \rfloor}[v]| \geq k$ for some $v \in V(G)$. Hence, in the following we can assume that $k > |N_{\lfloor s/2 \rfloor}[v]|$ for each vertex $v \in V(G)$.

First, we consider the case that s is even. Then $\lfloor s/2 \rfloor = s/2$ and we obtain that $N_s[v] \subseteq N_{s/2}[N_{s/2}[v]]$ for each $v \in V(G)$. Thus, $|N_s[v]| \leq k^2$.

Second, we consider the case that s is odd. Observe that we have $N_s[v] \subseteq N_{\lfloor s/2 \rfloor}[N_{\lfloor s/2 \rfloor}[N_{\lfloor s/2 \rfloor}[v]]]$ for each $v \in V(G)$. Thus, $|N_s[v]| \leq k^3$. \square

3.2 Edge Triangle s -Club for $\ell \geq 2$

Now we show W[1]-hardness for the remaining cases.

Theorem 10. *EDGE TRIANGLE s -CLUB is W[1]-hard for parameter k if $\ell \geq 2$.*

Next, we describe the construction of the reduction to prove Theorem 10. We reduce from CLIQUE. The idea is to construct one vertex gadget for each vertex of the CLIQUE instance and to add edges between two different vertex gadgets if and only if the two corresponding vertices are adjacent in such a way that all these edges are in exactly ℓ triangles. For an illustration of this construction see Fig. 2.

Construction 11. Let (G, k) be an instance of CLIQUE with $k \geq 3$. We construct an equivalent instance (G', k') of EDGE-TRIANGLE s -CLUB for some fixed $\ell \geq 2$ as follows. Let $\ell^* := \lceil \ell/2 \rceil$ and let $x := 6 \cdot \ell^*(s-1) + \lfloor \ell/2 \rfloor$. For each vertex $v \in V(G)$, we construct the following vertex gadget T^v . For better readability, all sub-indices of the vertices in T^v are considered modulo x . Our construction distinguishes even and odd values of ℓ . First, we describe the part of the construction which both cases have in common.




Fig. 2. Construction for Theorem 10 when $s = 3$ and $\ell = 2$ and G is a P_3 on $\{u, v, w\}$ with $uv \notin E(G)$. Only the gadgets A_v, B_w, A_w , and B_u are shown. For simplicity, no edges within A_v, B_w, A_w , and B_u are drawn and edges between B_w and A_w are only drawn if one endpoint is a_3^w, a_4^w, a_9^w , or a_{10}^w . Blue encircled vertices are neighbors of a_0^v , red encircled vertices have distance 2 to a_0^v , and black encircled vertices have distance 3 to a_0^v . Thus, a_0^v and $b_7^u = b_{0+1+3 \cdot 1 \cdot 2}^u = b_{0+\lfloor \ell/2 \rfloor + 3\ell^*(s-1)}^u$ have distance at least 4.

1. We add vertex sets $A_v := \{a_i^v \mid i \in [0, x]\}$ and $B_v := \{b_i^v \mid i \in [0, x]\}$ to G' .
2. We add the edges $a_i^v a_{i+j}^v$, and $b_i^v b_{i+j}^v$ for each $i \in [0, x]$ and each $j \in [-3\ell^*, 3\ell^*] \setminus \{0\}$ to G' .
3. We add the edge $a_i^v b_{i+j}^v$ for each $i \in [0, x]$ and each $j \in [-3\ell^*, 3\ell^*]$ to G' .

In other words, an edge $a_i^v b_j^v$ is added if the indices differ by at most $3\ell^*$. For even ℓ , this completes the construction of T^v . For odd ℓ , we extend T^v as follows:

- 0-1. We add the vertex set $C_v := \{c_i^v \mid i \in [0, x] \text{ and } i \equiv 0 \pmod{\ell^*}\}$ to G' . Note that C_v consists of exactly $6s - 5$ vertices.
- 0-2. We add the edges $c_i^v a_{i+j}^v$ and $c_i^v b_{i+j}^v$ for each $i \in [0, x]$ such that $i \equiv 0 \pmod{\ell^*}$ and each $j \in [-3\ell^*, 3\ell^*]$ to G' .
- 0-3. Also, we add the edge $c_i^v c_{i+j}^v$ to G' for each $i \in [0, x]$ such that $i \equiv 0 \pmod{\ell^*}$ and each $j \in [-3\ell^*, 3\ell^*] \setminus \{0\}$ to G' if the corresponding vertex c_{i+j}^v exists.

In other words, an edge between c_i^v and a_j^v, b_j^v , or c_j^v is added if the indices differ by at most $3\ell^*$. Now, for each edge $uv \in E(G)$, we add the following to G' :

- 0-4. We add the edges $a_i^v b_{i+j}^u$ and $a_i^u b_{i+j}^v$ for each $i \in [0, x]$ and $j \in [0, \lfloor \ell/2 \rfloor]$.
- 0-5. If ℓ is odd, we also add the edges $c_i^v b_{i+j}^u$ and $c_i^u b_{i+j}^v$ for each $i \in [0, x]$ such that $i \equiv 0 \pmod{\ell^*}$ and each $j \in [0, \lfloor \ell/2 \rfloor]$ to G' .

Observe that each vertex b_{i+j}^u is adjacent to *exactly* one vertex in C_v .

In other words, an edge between a_i^v or c_i^v and b_j^v is added if j exceeds i by at most $\lfloor \ell/2 \rfloor$. Finally, if ℓ is even, we set $k' := 2(x+1)k = (\ell(6s-5) + 2) \cdot k$, and if ℓ is odd, we set $k' := (2(x+1) + 6s-5)k = (\ell+2)(6s-5) \cdot k$.

Construction 11 has two key mechanisms: First, if $uv \notin E(G)$ then for each vertex $a \in A_v$ there is at least one vertex $b \in B_u$ such that $\text{dist}(a, b) > s$.

Second, each edge with one endpoint in A_v and one endpoint in B_u is contained in *exactly* ℓ triangles. Furthermore, if ℓ is odd, then this also holds for each edge with one endpoint in C_v and one in B_u . Consider an edge- ℓ -triangle s -club S and let $\tilde{G} = (S, \tilde{E})$ be a spanning subgraph of $G[S]$ with the maximal number of edges, such that each edge of \tilde{E} is contained in at least ℓ triangles in \tilde{G} and the diameter of \tilde{G} is s . As we will show, the two mechanics ensure that an edge with one endpoint in A_v (or C_v) and the other endpoint in B_u is contained in \tilde{E} if and only if S contains all vertices of A_v (and C_v) and B_u . We call this the *enforcement property*. Next, we formalize this property. To this end, we introduce the following notation. By E_{uv} we denote the set of all edges with one endpoint in A_v (or C_v if ℓ is odd) and the other endpoint in B_u .

Lemma 8. *Let S be an edge- ℓ -triangle s -club G' constructed in Construction 11. More precisely, let $\tilde{G} = (S, \tilde{E})$ be a maximal subgraph of $G[S]$ such that each edge in $E(\tilde{G})$ is contained in at least ℓ triangles within \tilde{G} and the diameter of \tilde{G} is at most s . Let $e \in E_{uv}$. Then $e \in E(\tilde{G})$ if and only if $A_v, B_u \subseteq S$ (and $C_v \subseteq S$, if ℓ is odd).*

Proof. Before we show the two implications, we prove the following cascading property of edge- ℓ -triangle s -clubs which contain at least one edge of E_{uv} .

Claim 4. If $a_i^v b_j^u \in E(\tilde{G})$ or $c_i^v b_j^u \in E(\tilde{G})$, then, $E_{uv} \subseteq E(\tilde{G})$.

Proof. First, we consider even values of ℓ . Note that $\ell^* = \lfloor \ell/2 \rfloor = \ell/2 = \lceil \ell/2 \rceil$. By construction we have

$$N[a_i^v] = \{a_{i+i'}^v, b_{i+i'}^v \mid i' \in [-3\ell/2, 3\ell/2]\} \cup \{b_{i+i'}^w \mid i' \in [0, \ell/2] \text{ and } vw \in E(G)\},$$

and similar

$$N[b_j^u] = \{a_{j+i'}^u, b_{j+i'}^u \mid i' \in [-3\ell/2, 3\ell/2]\} \cup \{a_{j-i'}^w \mid i' \in [0, \ell/2] \text{ and } uw \in E(G)\}.$$

Since $a_i^v b_j^u \in E(G')$ we obtain by Part 0-4. of Construction 11 that $j = i + z$ for some $z \in [0, \ell/2]$. Let $y := \ell/2 - z$ and observe that $N(a_i^v) \cap N(b_j^u) = \{a_{i+i'}^v \mid i' \in [-y, z] \setminus \{0\}\} \cup \{b_{j+i'}^u \mid i' \in [-z, y] \setminus \{0\}\}$. Thus, the edge $a_i^v b_j^u$ is contained in exactly ℓ triangles whose vertex sets are all contained in $G'[A_v \cup B_u]$. Since $a_i^v b_j^u \in E(\tilde{G})$, we thus conclude that all vertices and edges which form these ℓ triangles are contained in \tilde{G} . In particular, we obtain that $a_{i+1}^v b_{j+1}^u \in E(\tilde{G})$ and thus also that $a_{i+1}^v, b_{j+1}^u \in V(\tilde{G})$. Now, since $a_{i+1}^v b_{j+1}^u \in E(\tilde{G})$ we can repeat the above argumentation for the edge $a_{i+1}^v b_{j+1}^u$ and inductively for the edge $a_{i+q}^v b_{i+q}^u$ for all $q \in [x]$. We then have verified that $A_v \cup B_u \subseteq V(\tilde{G})$ and that each edge in E_{uv} is contained in $E(\tilde{G})$. Recall that A_v and B_u have size $x + 1$.

Second, we consider odd values of ℓ , that is $\ell = 2t + 1$ for some integer t . Note that $\lfloor \ell/2 \rfloor = t$ and that $\ell^* = \lceil \ell/2 \rceil = t + 1$. Furthermore, observe that vertex b_j^u has exactly one neighbor $c_{j+i'}^v$ in C_v for some $i' \in [0, t]$ such that $j + i' \bmod (t + 1) = 0$.

Hence, by construction we have

- $N[a_i^v] = N[c_i^v] = \{a_{i+i'}^v, b_{i+i'}^v \mid i' \in [-3(t+1), 3(t+1)]\} \cup \{c_{i+i'}^v \mid i' \in [-3(t+1), 3(t+1)] \text{ and } (i+i') \bmod (t+1) = 0\} \cup \{b_{i+i'}^w \mid i' \in [0, t] \text{ and } vw \in E(G)\}$,
 - $N[b_j^u] = \{a_{j+i'}^u, b_{j+i'}^u \mid i' \in [-3(t+1), 3(t+1)]\} \cup \{c_{j+i'}^u \mid i' \in [-3(t+1), 3(t+1)] \text{ and } (j+i') \bmod (t+1) = 0\} \cup \{a_{j-i'}^w \mid i' \in [0, t] \text{ and } uw \in E(G)\} \cup \{c_{j-i'}^w \mid i' \in [0, t] \text{ and } (j-i') \bmod (t+1) = 0 \text{ and } uw \in E(G)\}$,
- and

Since $a_i^v b_j^u \in E(G')$ we obtain by Part 0-4. of Construction 11 that $j = i + z$ for some $z \in [0, t]$. Now, let $y := t - z$ and let $c_{i'}^v$ be the unique neighbor of b_j^u in C_v . We conclude that $N(a_i^v) \cap N(b_j^u) = \{a_{i+j'}^v \mid j' \in [-y, z] \setminus \{0\}\} \cup \{b_{j+j'}^u \mid j' \in [-z, y] \setminus \{0\}\} \cup \{c_{i'}^v\}$.

Hence, both vertices have exactly $t + t + 1 = \ell$ common neighbors and thus $N(a_i^v) \cap N(b_j^u) \subseteq S$. By similar arguments a similar statement can be shown for the edge $c_i^v b_j^u$. Thus, the edges $a_i^v b_j^u$ and $c_i^v b_j^u$ are contained in exactly ℓ triangles whose vertex sets are contained in $G[A_v \cup C_v \cup B_u]$.

Since $a_i^v b_j^u \in E(\tilde{G})$ we thus conclude that all vertices and edges which form these ℓ triangles are contained in \tilde{G} . In particular, we obtain that $a_{i+1}^v b_{j+1}^u \in E(\tilde{G})$ and also that $a_{i+1}^v, b_{j+1}^u \in V(\tilde{G})$. Now, since $a_{i+1}^v b_{j+1}^u \in E(\tilde{G})$ we can repeat the above argumentation for the edge $a_{i+1}^v b_{j+1}^u$ and inductively for the edge $a_{i+q}^v b_{j+q}^u$ for all $q \in [x]$. We then have verified that $A_v \cup B_u \cup C_v \subseteq V(\tilde{G})$ and that each edge in E_{uv} is contained in $E(\tilde{G})$. ■

Now, we are ready to prove the two implications. The implication that if $e \in E(\tilde{G})$ then $A_v, B_u \subseteq S$ (and $C_v \subseteq S$, if ℓ is odd) directly follows from Claim 4. It remains to show the other implication. From Claim 4 we conclude that either $E_{uv} \subseteq E(\tilde{G})$ or $E_{uv} \cap E(\tilde{G}) = \emptyset$. If $E_{uv} \subseteq E(\tilde{G})$, then we are done, so assume towards a contradiction that $E_{uv} \cap E(\tilde{G}) = \emptyset$. Recall that $A_v \cup B_u \subseteq S$ (and also $C_v \subseteq S$ if ℓ is odd). Furthermore, recall that \tilde{G} is maximal, that is, there exists no spanning subgraph of $G[S]$ which has more edges than \tilde{G} . Hence, adding all edges in E_{uv} to \tilde{G} is still an edge- ℓ -triangle s -club, a contradiction to the maximality of \tilde{G} . □

Now, we prove the correctness of the reduction for Theorem 10.

Proof (of Theorem 10). We show that G contains a clique of size at least k if and only if G' contains an edge- ℓ -triangle s -club of size at least k' .

Let K be a clique of size k in G . Recall that T^v is the gadget of vertex $v \in V(G)$. We verify that $S := \{u \in V(T^v) \mid v \in K\}$ is an edge- ℓ -triangle s -club of size at least k' . More precisely, we show that $\tilde{G} := G[S]$ fulfills all properties of being an edge- ℓ -triangle s -club. Since $|T^v| = 2\ell^*(6s - 5) + 2$ if ℓ is even and $|T^v| = (2\ell^* + 1)(6s - 5)$ if ℓ is odd for each $v \in K$ and since $|K| \geq k$, we have $|S| = |V(\tilde{G})| \geq k'$. It remains to show that \tilde{G} is an edge- ℓ -triangle s -club.

Next, we show that \tilde{G} is an s -club.

First, we show that T^v is an s -club. Therefore, consider the vertex pair $\{a_i^v, a_j^v\}$ for some $v \in K$. Observe that $P := (a_i^v, a_{i+1}^v, \dots, a_{i+p}^v)$ for $i + p = j$ is a path of length p from a_i^v to a_j^v and that $Q := (a_i^v, a_{i-1}^v, \dots, a_{i-q}^v)$ for $i - q = j$

is a path of length q from a_i^v to a_j^v . Clearly, $p + q = x + 1$. Hence, $\min(p, q) \leq (x + 1)/2 \leq 3\ell^*(s - 1) + \lfloor \ell/2 \rfloor$. Without loss of generality, assume that the minimum is achieved by path P and assume that $p = \alpha \cdot (3\ell^*) + \beta$ for some $\alpha \in [s - 1]$ and some $\beta < 3\ell^*$. Recall that by Part 2 of Construction 11, $a_i^v, a_j^v \in E(G')$ if and only if $j' = i' + z$ for some $z \in [-3\ell^*, 3\ell^*] \setminus \{0\}$. Hence,

$$(a_i^v, a_{i+1 \cdot (3\ell^*)}^v, \dots, a_{i+\alpha \cdot (3\ell^*)}^v, a_{i+\alpha \cdot (3\ell^*)+\beta}^v)$$

is a path of length at most $(s - 1) + 1 = s$ from a_i^v to a_j^v .

These arguments also apply symmetrically to the vertex pairs $\{b_i^v, b_j^v\}$ and $\{a_i^v, b_j^v\}$ for each $v \in K$. Furthermore, if ℓ is odd, observe that the above argumentation can also be used to show that the vertex pairs $\{c_i^v, a_j^v\}$, $\{c_i^v, b_j^v\}$, and $\{c_i^v, c_j^v\}$ have distance at most s to each other.

Second, we show that a_i^v has distance at most s to b_j^u . Note that by Part 0-4. of Construction 11, a_i^v has neighbors $b_i^u, \dots, b_{i+\lfloor \ell/2 \rfloor}^u$ since $uv \in E(G)$. In the following, we assume that $j \neq i + z$ for all $z \in [0, \lfloor \ell/2 \rfloor]$. Consider the paths $P := (a_i^v, b_{i+\lfloor \ell/2 \rfloor}^u, b_{i+\lfloor \ell/2 \rfloor+1}^u, \dots, b_{i+\lfloor \ell/2 \rfloor+p}^u)$ for $i + \lfloor \ell/2 \rfloor + p = j$ of length $p + 1$ and $Q := (a_i^v, b_i^u, b_{i-1}^u, \dots, b_{i-q}^u)$ for $i - q = j$ of length $q + 1$. Observe that $(p + 1) + (q + 1) = (x + 3) - \lfloor \ell/2 \rfloor$. Thus, $p + q = (x + 1) - \lfloor \ell/2 \rfloor = 6\ell^*(s - 1) + 1$. Since p and q are integers we have $\min(p, q) \leq ((x + 1) - \lfloor \ell/2 \rfloor)/2 = 3\ell^*(s - 1)$.

Without loss of generality assume that the minimum is achieved by path P and assume that $p = \alpha \cdot (3\ell^*) + \beta$ for some $\alpha \in [s - 2]$ and some $\beta \leq 3\ell^*$. Recall that by Part 2 of Construction 11, we have $b_i^u, b_j^u \in E(G')$ if and only if $j' = i' + z$ for some $z \in [-3\ell^*, 3\ell^*] \setminus \{0\}$. Now, observe that

$$(a_i^v, b_{i+\lfloor \ell/2 \rfloor}^u, b_{i+\lfloor \ell/2 \rfloor+1 \cdot (3\ell^*)}^u, \dots, b_{i+\lfloor \ell/2 \rfloor+\alpha \cdot (3\ell^*)}^u, b_{i+\lfloor \ell/2 \rfloor+\alpha \cdot (3\ell^*)+\beta}^u)$$

is a path of length at most $1 + (s - 2) + 1 = s$ from a_i^v to b_j^u .

Furthermore, if ℓ is odd, observe that the above argumentation can also be used to show that the vertex pairs $\{c_i^v, b_j^u\}$ have distance at most s to each other by replacing a_i^v with c_i^v in the paths P and Q .

The fact that vertices a_i^v and a_j^u , and b_i^v and b_j^u , respectively, have distance at most s to each other can be proven similar as we showed that a_i^v and b_j^u have distance at most s by observing that a_i^v has distance 2 to each vertex a_{i+z}^u with $z \in [-3\ell^*, \lfloor \ell/2 \rfloor + 3\ell^*]$ since a_i^v has neighbors $b_i^u, \dots, b_{i+\lfloor \ell/2 \rfloor}^u$ and since by Part 2 of Construction 11, we have $b_i^u, a_j^u \in E(G')$ if and only if $j' = i' + z'$ for some $z' \in [-3\ell^*, 3\ell^*]$.

Furthermore, if ℓ is odd, observe that the above argumentation can also be used to show that the vertex pairs $\{c_i^v, a_j^u\}$ and $\{c_i^v, c_j^u\}$ have distance at most s to each other by replacing a_i^v with c_i^v and replacing a_j^u with c_j^u , respectively, in the paths P and Q .

Hence, \tilde{G} is indeed an s -club.

Next, we show that each edge in $E(\tilde{G})$ is contained in at least ℓ triangles which are contained in \tilde{G} .

Consider the edge $a_i^v a_{i+j}^v$ for some $j \in [-3\ell^*, 3\ell^*]$. Without loss of generality, assume that $j > 0$. By Part 2 Construction 11, both vertices are adjacent to each

vertex $a_{i+i'}^v$ with $i' \in [3\ell^*] \setminus \{j\}$. Hence, both vertices are in at least $3\ell^* - 1 \geq \ell$ triangles.

Furthermore, the statement can be shown analogously for the edges $b_i^v b_{i+j}^v$ (and $c_i^v c_{i+j}^v$ if ℓ is odd) for some $j \in [-3\ell^*, 3\ell^*]$.

Also, the statement can be shown analogously for the edge $a_i^v b_{i+j}^v$ for some $j \in [-3\ell^*, 3\ell^*]$. If $j \geq 0$, then a_{i+z}^v for each $z \in [3\ell^*]$ is a common neighbor of both vertices and thus the edge $a_i^v b_{i+j}^v$ is contained in at least ℓ triangles. The case $j < 0$ can be shown analogously.

For odd ℓ , the statement can be shown analogously for the edges $a_i^v c_{i+j}^v$, $b_i^v c_{i+j}^v$, and $c_i^v c_{i+j}^v$ for some $j \in [-3\ell^*, 3\ell^*]$.

The fact that the edges $a_i^v b_{i+j}^u$ and $c_i^v b_{i+j}^u$ are contained in exactly ℓ triangles follows from the proof of Lemma 8.

We conclude that each edge in $E(\tilde{G})$ is contained in at least ℓ triangles in \tilde{G} . Thus, S is indeed an edge- ℓ -triangle s -club of size k' .

Conversely, let S be an edge- ℓ -triangle s -club of size at least k' in G' . More precisely, let \tilde{G} be a maximal spanning subgraph of $G[S]$ which has diameter at most s and such that each edge in $E(\tilde{G})$ is contained in at least ℓ triangles in \tilde{G} . We show that \tilde{G} contains a clique of size at least k .

First, we show that for each vertex $x \in A_v \cup B_v \cup C_v$ there exists a vertex $y \in A_u \cup B_u \cup C_u$ such that $\text{dist}(x, y) \geq s + 1$ if $uv \notin E(G)$. For this, recall that by construction each two vertices with sub-indices i' and j' are not adjacent if their difference (modulo x) is larger than $3\ell^*$.

Claim 5. In G' we have $\text{dist}(x_i, y_j) \geq s + 1$ for each $i \in [0, x]$, $j := i + \lfloor \ell/2 \rfloor + 3\ell^*(s - 1)$, $x_i \in \{a_i^v, b_i^v, c_i^v\}$, and $y_j \in \{a_j^u, b_j^u, c_j^u\}$ if $uv \notin E(G)$.

Proof. There are two possible paths from x_i to $y_{i+\lfloor \ell/2 \rfloor + 3\ell^*(s-1)}$ with respect to the indices. First, there is a subsequence of the indices which is increasing $(i, i + 1, \dots, i + \lfloor \ell/2 \rfloor + 3\ell^*(s - 1))$. This path has length $\lfloor \ell/2 \rfloor + 3\ell^*(s - 1)$. Second, there is a subsequence of the indices which is decreasing $(i, i - 1, \dots, i - i')$, where $-i' = i + \lfloor \ell/2 \rfloor + 3\ell^*(s - 1)$. This path has length $3\ell^*(s - 1) + 1$.

Hence, each path from x_i to $y_{i+\lfloor \ell/2 \rfloor + 3\ell^*(s-1)}$ has to overcome at least $3\ell^*(s - 1) + 1$ indices. Observe that whenever an edge between A_p (or C_p) and B_q for $p, q \in V(G)$ with $pq \in E(G)$ is traversed, by construction the index can increase/decrease by at most $\lfloor \ell/2 \rfloor$. Now, we use the fact that $uv \notin E(G)$: There are no edges between the vertex gadgets T^v and T^u . Thus, at least two times such a traversal of at most $\lfloor \ell/2 \rfloor$ indices has to be done. Hence, the index i can increase or decrease by at most $2 \cdot \lfloor \ell/2 \rfloor + 3\ell^*(s - 2) < 3\ell^*(s - 1) + 1$ if at least 2 edge traversals between different vertex gadgets are necessary. Thus, both vertices have distance at least $s + 1$. ■

The following statement directly follows from Claim 5 and Lemma 8.

Claim 6. If $A_v \subseteq S$ (or $C_v \subseteq S$ if ℓ is odd) and $B_u \subseteq S$ then $uv \in E(G)$.

We now use Claims 5 and 6 to show that G contains a clique of size at least k . We distinguish the cases whether S contains only parts of one of the

gadgets $A_v, B_v,$ or C_v or whether S contains all vertices of the gadgets A_v (or $B_v,$ or C_v) completely.

First, assume that for some vertex $v \in V(G)$ we have $A_v \cap S \neq \emptyset$ and $A_v \not\subseteq S$. In the following, we show that S only contains vertices of gadget T^v and from gadgets T^u such that $uv \in E(G)$. Since $A_v \not\subseteq S$, we conclude that in \tilde{G} we have $N_{\tilde{G}}(A_v \cap S) \subseteq (B_v \cup C_v)$: Otherwise, vertex a_v has a neighbor $b_u \in B_u$ and by Lemma 8 we would obtain $A_u \subseteq S$, a contradiction to the assumption $A_v \not\subseteq S$. If $B_v \not\subseteq S$, then by Lemma 8 no vertex in B_v can have a neighbor a_i^w or c_i^w for some $w \neq v$. Hence, $S \cap T^v$ would be a connected component of size at most $3(x+1)$, a contradiction to the size of S since $k \geq 3$. Thus, we may assume that $B_v \subseteq S$. Observe that if $a_i^w \in S$ or $c_i^w \in S$ for some $w \in V(G)$ such that $vw \in E(\tilde{G})$, that is, also $vw \in E(G')$, then we have $A_w \subseteq S$ and $C_w \subseteq S$ by Lemma 8 since each vertex a_i^w and c_i^w has a neighbor in B_v . Let $W := \{w_1, \dots, w_t\}$ denote the set of vertices w_j such that $vw_j \in E(G)$ and $A_{w_j}, C_{w_j} \subseteq S$. If $w_x w_y \notin E(G)$ for some $x, y \in [t]$ with $x \neq y$, then $a_0^{w_x}$ and $a_{\lfloor \ell/2 \rfloor + 3\ell^*(s-1)}^{w_y}$ have distance at least $s+1$ by Claim 5. Thus $w_x w_y \in E(G)$ for each $x, y \in [t]$ with $x \neq y$.

Assume towards a contradiction that $a_i^p \in S$ or $c_i^p \in S$ for some $p \in V(G) \setminus w$ with $p \neq v$. Note that $pv \notin E(G)$ since otherwise $p \in W$ by the definition of W . Observe that since $B_v \subseteq S$ we also have $b_{i+\lfloor \ell/2 \rfloor + 3\ell^*(s-1)}^v \in S$. But since $pv \notin E(G)$ we obtain from Claim 5 that $\text{dist}(z_i, b_{i+\lfloor \ell/2 \rfloor + 3\ell^*(s-1)}^v) \geq s+1$ for $z_i = a_i$ or $z_i = c_i$, a contradiction. We conclude that S does not contain any vertex a_i^p or c_i^p with $p \neq v$ or $p \neq w_j$ for $j \in [t]$.

Next, assume towards a contradiction that $b_i^p \in S$ for some $p \in V(G)$ with $p \neq v$ and $p \notin W$. If $pv \notin E(G)$, then b_i^p and $b_{i+\lfloor \ell/2 \rfloor + 3\ell^*(s-1)}^v$ have distance at least $s+1$ again by Claim 5. Thus, we can assume that $pv \in E(G)$. Recall that $b_{i+\lfloor \ell/2 \rfloor + 3\ell^*(s-1)}^v \in S$. As defined by Claim 5, each shortest path from b_i^p to $b_{i+\lfloor \ell/2 \rfloor + 3\ell^*(s-1)}^v$ can swap at most once between different vertex gadgets. In this case, there is exactly one swap from T^p to T^v . From the above we know that $(A_p \cup C_p) \cap S = \emptyset$. Thus, each shortest path from b_i^p to $b_{i+\lfloor \ell/2 \rfloor + 3\ell^*(s-1)}^v$ uses at least one vertex in $A_v \cup C_v$. Since at least one edge with endpoint $A_v \cup C_v$ is contained in S , we conclude from Lemma 8 that $A_v \cup C_v \subseteq S$, a contradiction to the assumption $A_v \not\subseteq S$.

Hence, there is no vertex $p \neq v$ and $p \notin W$ such that $T^p \cap S \neq \emptyset$. In other words, S contains only vertices from the gadget T^v and from gadgets T^u with $vu \in E(G)$. Thus, $S \subseteq T^v \cup \bigcup_{j=1}^t T^{w_j}$. By definition of k' , we have $t \geq k-1$ and we conclude that G contains a clique of size at least k . The case that we have $B_v \cap S \neq \emptyset$ and $B_v \not\subseteq S$ or $C_v \cap S \neq \emptyset$ and $C_v \not\subseteq S$ for some vertex $v \in V(G)$ can be handled similarly.

Second, consider the case that for each set A_v with $A_v \cap S \neq \emptyset$ we have $A_v \subseteq S$, that for each set B_v with $B_v \cap S \neq \emptyset$ we have $B_v \subseteq S$, and that for each set C_v with $C_v \cap S \neq \emptyset$ we have $C_v \subseteq S$. Let $W_A := \{w_A^j \mid A_{w_j} \subseteq S\}$, $W_B := \{w_B^j \mid B_{w_j} \subseteq S\}$, and $W_C := \{w_C^j \mid C_{w_j} \subseteq S\}$. If $W_A = \emptyset$ or $W_B = \emptyset$ or $W_C = \emptyset$ then each connected component in $G'[S]$ has size at most $2(x+1) < k'$. Thus, we may assume that $W_A \neq \emptyset$, $W_B \neq \emptyset$, and $W_C \neq \emptyset$. By Claim 6, we have $w_A^i w_B^j \in E(G)$

for each $w_A^i \in W_A$ and $w_B^j \in W_B$ and also $w_C^i w_B^j \in E(G)$ for each $w_C^i \in W_C$ and $w_B^j \in W_B$. Furthermore, by Claim 5, we have $w_B^j w_B^{j'} \in E(G)$ for $w_B^j, w_B^{j'} \in W_B$, $w_A^i w_A^{i'} \in E(G)$ for $w_A^i, w_A^{i'} \in W_A$, $w_C^i w_C^{i'} \in E(G)$ for $w_C^i, w_C^{i'} \in W_C$, and also $w_A^i w_C^{i'} \in E(G)$ for $w_A^i \in W_A$ and $w_C^{i'} \in W_C$. Hence, we obtain that $\min(|W_A|, |W_B|, |W_C|) \geq k$ and thus G contains a clique of size k . \square

4 Seeded s -Club

In this section we study the parameterized complexity of SEEDED s -CLUB with respect to the standard parameter solution size k . Recall that in this problem we aim to find an s -club containing a given seed of vertices. Here, we assume that $|W| < k$ since otherwise the problem can be solved in polynomial time.

4.1 Tractable Cases

For clique seeds, we provide the following kernel. Note that here we present a kernel and not only a Turing kernel.

Theorem 12. *SEEDED s -CLUB admits a kernel with $\mathcal{O}(k^{2|W|+1})$ vertices if W is a clique.*

Note that this kernel has polynomial size if W has constant size. In the following, assume that $G[W]$ is a clique. To prove the kernel, we first remove all vertices with distance at least $s + 1$ to any vertex in W . Second, we show that if the remaining graph, that is, $N_s[W]$, is sufficiently large, then (G, W, k) is a trivial yes-instance.

Reduction Rule 13. *Let (G, W, k) be an instance of SEEDED s -CLUB. If G contains a vertex u such that $\text{dist}(u, w) \geq s + 1$ for some $w \in W$, then remove u .*

Clearly, Reduction Rule 13 is correct and can be applied in polynomial time. Next, we show that if the remaining graph is sufficiently large then (G, W, k) is a yes-instance of SEEDED s -CLUB.

Lemma 9. *An instance (G, W, k) of SEEDED s -CLUB with $|N_{s-1}[W]| \geq k^2$ is a yes-instance.*

To prove Lemma 9 for $s \geq 3$ we need the following technical lemmas.

Lemma 10. *An instance (G, W, k) of SEEDED s -CLUB with $s \geq 3$ is a yes-instance if $|N_{\lfloor (s+1)/2 \rfloor - 1}[W]| \geq k$.*

Proof. By definition $W \subseteq N_{\lfloor (s+1)/2 \rfloor - 1}[W]$ and $|N_{\lfloor (s+1)/2 \rfloor - 1}[W]| \geq k$. Thus, it remains to show that $N_{\lfloor (s+1)/2 \rfloor - 1}[W]$ is an s -club. Therefore, consider a pair of vertices $u, v \in N_{\lfloor (s+1)/2 \rfloor - 1}[W]$. Observe that by definition $\text{dist}(u, W) \leq \lfloor (s+1)/2 \rfloor - 1$ and $\text{dist}(v, W) \leq \lfloor (s+1)/2 \rfloor - 1$. Since W is a clique, we have $\text{dist}(u, v) \leq (\lfloor (s+1)/2 \rfloor - 1) + 1 + (\lfloor (s+1)/2 \rfloor - 1) \leq s$. Hence, the lemma follows. \square

Note that the assumption $s \geq 3$ in Lemma 10 is necessary to guarantee that $\lfloor (s+1)/2 \rfloor - 1 \geq 1$. Next, we show that if a vertex in $N_{\lfloor (s+1)/2 \rfloor - 1}(W)$ has many vertices close to it, then (G, W, k) is a yes-instance.

Lemma 11. *An instance (G, W, k) of SEEDED s -CLUB in which $s \geq 3$ and $|N_{\lfloor s/2 \rfloor}(v)| \geq k$ for some vertex $v \in N_{\lfloor (s+1)/2 \rfloor - 1}(W)$ is a yes-instance.*

Proof. Let v be a vertex as specified in the lemma. By definition of v , there exists a path $P := (q_0, q_1, \dots, q_{\lfloor (s+1)/2 \rfloor - 1})$ of length $\lfloor (s+1)/2 \rfloor - 1$ in G such that $q_{\lfloor (s+1)/2 \rfloor - 1} = v$, $w_0 \in W$, and $q_i \in N_i(q_0)$. We show that $S := N_{\lfloor s/2 \rfloor}(v) \cup W \cup P$ is an s -club of size k containing W . Clearly, $W \subseteq S$ and $|S| \geq k$. Thus, it remains to show that S is an s -club.

Consider a vertex $w \in W$. Vertex w has distance at most $i+1$ to vertex w_i . In particular, $\text{dist}(w, v) \leq \lfloor (s+1)/2 \rfloor$. Since each vertex $u \in N_{\lfloor s/2 \rfloor}(v)$ has distance at most $\lfloor s/2 \rfloor$ to v we obtain that $\text{dist}(w, u) \leq \lfloor (s+1)/2 \rfloor + \lfloor s/2 \rfloor = s$. By similar arguments we can also show that vertex w_i for $i \in [\lfloor (s+1)/2 \rfloor - 1]$ has distance at most s to each vertex in S .

Finally, consider two vertices $x, y \in N_{\lfloor s/2 \rfloor}(v)$. Note that $\text{dist}(x, v) \leq \lfloor s/2 \rfloor$ and also $\text{dist}(y, v) \leq \lfloor s/2 \rfloor$ and thus $\text{dist}(x, y) \leq s$.

Thus, S is indeed an s -club. \square

With those two lemmas we are now able to prove Lemma 9.

Proof (Proof of Lemma 9). First, we consider the case $s = 2$. It is sufficient to show that (G, W, k) is a yes-instance if $|N[w]| \geq k$ for some $w \in W$ since $|W| \leq k$ and since $N[W] \geq k^2$ by our assumption. Since all vertices in $N(w)$ have the common neighbor w , we conclude that $N[w]$ is a 2-club. Also, since W is a clique, we have $W \subseteq N[w]$. The size bound of $N_{s-1}[W]$ follows from $|N[w]| \geq k$. Thus, (G, W, k) is a yes-instance.

Second, we consider the case $s \geq 3$. Observe that

$$N_{s-1}[W] = N_{\lfloor (s+1)/2 \rfloor - 1}[W] \cup \bigcup_{v \in N_{\lfloor (s+1)/2 \rfloor - 1}[W]} N_{\lfloor s/2 \rfloor}(v).$$

By Lemma 10, (G, W, k) is a yes-instance if $|N_{\lfloor (s+1)/2 \rfloor - 1}[W]| \geq k$ and, by Lemma 11, (G, W, k) is a yes-instance if $|N_{\lfloor s/2 \rfloor}(v)| \geq k$ for some $v \in N_{\lfloor (s+1)/2 \rfloor - 1}(W)$. Thus, by the above equality we conclude that (G, W, k) is a yes-instance. \square

Finally, we bound the size of $N_s(W)$. There we assume that $|N_{s-1}[W]| < k^2$ by Lemma 9 and that Reduction Rule 13 is applied.

Lemma 12. *An instance (G, W, k) of SEEDED s -CLUB with $|N_s(W)| \geq k^{2|W|+1}$ which is reduced with respect to Reduction Rule 13 is a yes-instance.*

Proof. Since Reduction Rule 13 has been applied exhaustively, each vertex $p \in N_s(W)$ has distance exactly s to each vertex in W . In other words, for each vertex $w_\ell \in W$ there exists a vertex $u_{s-1}^\ell \in N_{s-1}(w_\ell)$ such that $pu_{s-1}^\ell \in E(G)$.

Note that $N_{s-1}(w_\ell) \subseteq N_{s-1}[W]$. Moreover, by Lemma 9 we may assume that $|N_{s-1}[W]| < k^2$. In particular: $|N_{s-1}(W)| < k^2$. Since $|N_s(W)| \geq k^{2|W|+1}$, by the pigeonhole principle there exists a set $\{u_{s-1}^1, u_{s-1}^2, \dots, u_{s-1}^{|W|}\}$ with $u_{s-1}^\ell \in N_{s-1}(w_\ell)$ for $\ell \in [|W|]$ such that the set $P := N_s(W) \cap \bigcap_{\ell \in [|W|]} N(u_{s-1}^\ell)$ has size at least k . The size bound follows from the observation that each $N_{s-1}(w_\ell)$ has size at most k^2 and we have exactly $|W|$ many of these sets. By the definition of vertex u_{s-1}^ℓ , there exists for each $i \in [s-2]$ a vertex $u_i^\ell \in N_i(w_\ell)$ such that $w_\ell, u_1^\ell, \dots, u_{s-1}^\ell$ is a path of length $s-1$ in G . We define the set $U := \{u_i^\ell \mid \ell \in [|W|], i \in [s-1]\}$. Next, we show that $Z := P \cup W \cup U$ induces an s -club.

First, observe that all vertices in P have distance at most 2 to each other since they have the common neighbor u_{s-1}^1 . Second, note that the vertices $w_\ell, u_1^\ell, \dots, u_{s-1}^\ell, p, u_{s-1}^j, \dots, u_1^j, w_j$ form a cycle with $2s+1$ vertices, for each $p \in P$ and each two indices $j, \ell \in [|W|]$. Each vertex in this cycle has distance at most s to each other vertex in that cycle. Hence, Z is indeed an s -club. \square

Recall that Lemma 9 showed that the number of vertices with distance at most $s-1$ to W is bounded by k^2 . Together with Lemma 12 now Theorem 12 is proven.

4.2 Intractable Cases

Now, we show hardness for some of the remaining cases.

Theorem 14. *Let H be a fixed graph. SEEDED s -CLUB is $W[1]$ -hard parameterized by k even if $G[W]$ is isomorphic to H , when*

- $s = 2$ and H contains at least two non-adjacent vertices, or if
- $s \geq 3$ and H contains at least two connected components.

Hardness for $s = 2$. First, we prove hardness for $s = 2$ when H contains at least one non-edge. For an illustration of Construction 15 we refer to Figure 3.

Construction 15. Let (G, k) be an instance of CLIQUE. We construct an equivalent instance (G', k') of SEEDED s -CLUB as follows. Initially, we add the set W to G' , and add edges such that $G'[W]$ is isomorphic to H . Since H is not a clique, there exist two vertices $u, v \in V(H)$ such that $uv \notin E(H)$. Let $R := W \setminus \{u, v\}$. Next, we add two copies G_u and G_v of G to G' , make u adjacent to each vertex in $V(G_u)$, and make v adjacent to each vertex in $V(G_v)$. We denote with x_u , and x_v the copies of x in G_u and G_v , respectively. Next, we add the edge $x_u x_v$ for each $x \in V(G)$. Furthermore, we add a new vertex p and make it adjacent to each vertex in W . Next, we add a new vertex u^* adjacent to p , each vertex in $V(G_u)$, and each vertex in R . Analogously, we add a new vertex v^* which is adjacent to p , each vertex in $V(G_v)$, and each vertex in R . Finally, we set $k' := 2k + |W| + 3$.

Now, we prove the correctness of Construction 15.




Fig. 3. Illustration of Construction 15.

Lemma 13. *For any graph H which is not a clique, SEEDED 2-CLUB parameterized by k is $W[1]$ -hard if $G'[W]$ is isomorphic to H .*

Proof. We prove that G contains a clique of size k if and only if G' contains a 2-club S containing W of size $k' = 2k + |W| + 3$.

Let K be a clique of size k in G and let K_u and K_v be the copies of K in G_u and G_v . We argue that $S := K_u \cup K_v \cup W \cup \{u^*, v^*, p\}$ is a 2-club of size at least k' containing W . Clearly, $|S| = k'$ and S contains W . Thus, it remains to show that $G[S]$ is a 2-club.

First, we show that each vertex in R has distance at most 2 to each vertex in S : All vertices in R have the common neighbors p , u^* , and v^* . Since u and v are neighbors of p , each vertex in $V(G_u)$ is a neighbor of u^* , and since each vertex in $V(G_v)$ is a neighbor of v^* we conclude that each vertex in R has distance at most 2 to any vertex in S .

Second, we show that each vertex in K_u has distance at most 2 to each vertex in $S \setminus R$. Observe that $\{u, u^*, x_v\} \cup K_u \subseteq N[x_u]$ for each vertex $x_u \in K_u$. Hence, x_u has distance at most 2 to p via u^* , v , and v^* via x_v , each vertex, and each vertex in K_v via the corresponding vertex in K_u . By symmetric arguments the statement also holds for each vertex in K_v .

Finally, each pair of vertices of $\{p, u, u^*, v, v^*\}$ has distance at most 2 to each other since $u, u^*, v, v^* \in N(p)$.

Thus, S is indeed a 2-club.

Conversely, suppose that G' contains a 2-club S of size at least $2k + |W| + 3$ which contains all vertices of W . Observe that we have $N(x_u) \cap N(v) = \{x_v\}$ for each vertex $x_u \in V(G_u)$, and symmetrically $N(x_v) \cap N(u) = \{x_u\}$ for each vertex $x_v \in V(G_v)$. Hence, $x_u \in S$ if and only if $x_v \in S$. Let $K_u := S \cap V(G_u)$. By definition of k' we obtain that $|K_u| \geq k$. Assume towards a contradiction that K_u contains a pair of nonadjacent vertices x_u and y_u . By the argumentation above we obtain $y_v \in S$. Now, observe that $N(x_u) = \{u, u^*, x_v\} \cup \bigcup \{z_u \mid xz \in E(G)\}$ and that $N(y_v) = \{v, v^*, y_u\} \cup \bigcup \{z_v \mid yz \in E(G)\}$. Since $xy \notin E(G)$ we thus

obtain $N(x_u) \cap N(y_v) = \emptyset$, a contradiction. Thus, G contains a clique of size k . \square

Hardness for seeds with at least two connected components and $s \geq 3$. Now, we show $W[1]$ -hardness for the case $s \geq 3$ when the seed contains at least two connected components. Fix a graph H with at least two connected components. We show $W[1]$ -hardness for $s \geq 3$ even if $G[W]$ is isomorphic to H .

Construction 16. Let (G, k) be an instance of CLIQUE. We construct an equivalent instance (G', k') of SEEDED s -CLUB as follows. Initially, we add the set W to G' , and add edges such that $G'[W]$ is isomorphic to H . Let D_1 be one connected component of $G'[W]$. By assumption, $D_2 := W \setminus D_1$ is not empty. Next, we add two copies G_1 and G_2 of G to G' . Then, we add edges to G' such that each vertex in D_1 is adjacent to each vertex in $V(G_1)$ and such that each vertex in D_2 is adjacent to each vertex in $V(G_2)$. Furthermore, we add a path (p_1, \dots, p_{s-1}) consisting of exactly $s - 1$ new vertices to G' , make p_1 adjacent to each $u \in D_1$, and make p_{s-1} adjacent to each $v \in D_2$. By $P := \{p_i \mid i \in [s - 1]\}$ we denote the set of these newly added vertices. Now, for each $x \in V(G)$ we do the following. Consider the copies $x_1 \in V(G_1)$ and $x_2 \in V(G_2)$ of vertex $x \in V(G)$. We add a path $(x_1, q_1^x, \dots, q_{s-2}^x, x_2)$ consisting of $s - 2$ new vertices to G' . By $Q_x := \{q_i^x \mid i \in [s - 2]\}$ we denote the set of the new internal path vertices. Finally, we set $k' := sk + |W| + s - 1$.

Now, we prove the correctness of Construction 16.

Lemma 14. *Let H be a fixed graph with at least two connected components. SEEDED s -CLUB parameterized by k is $W[1]$ -hard even if $G[W]$ is isomorphic to H .*

Proof. We show that G contains a clique of size k if and only if G' contains a W -seeded s -club of size at least $k' = sk + |W| + s - 1$.

Let K be a clique of size k in G . Furthermore, let K_1 and K_2 denote the copies of K in G_1 and G_2 , respectively. We show that $S := W \cup P \cup K_1 \cup K_2 \cup \bigcup_{x \in K} Q_x$ is a W -seeded s -club of size at least k' . Clearly, $|S| = k'$ and S contains W . Thus, it remains to verify that $G[S]$ is an s -club. Note that since each vertex in $V(G_1)$ is adjacent to each vertex in D_1 , each two vertices in D_1 have distance at most 2, and similarly each two vertices in $V(G_2)$ have distance at most 2. Analogously, we can show that each two vertices in D_2 and each two vertices in $V(G_2)$ have distance at most 2. Furthermore, the vertices $(x, p_1, p_2, \dots, p_{s-1}, y, u_2, q_{s-2}^u, \dots, q_1^u, u_1)$ for each vertex $x \in D_1$, each vertex $y \in D_2$, and each vertex $u \in K$ form a C_{2s+1} , a cycle with $2s + 1$ vertices. Observe that also the vertices $(u_1, q_1^u, \dots, q_{s-2}^u, u_2, v_2, q_{s-2}^v, \dots, q_1^v, v_1)$ form a C_{2s} for each two vertices $u, v \in K$. Since all remaining distances are covered by these two cycles, we conclude that S is indeed an s -club.

Conversely, suppose that G' contains a W -seeded s -club S of size at least k' . Let $Q'_v := \{v_1, q_1^v, \dots, q_{s-2}^v, v_2\}$ for each $v \in V(G)$. We show that $Q'_v \cap S \neq \emptyset$ if and only if $Q'_v \subseteq S$. Assume towards a contradiction, that $Q'_v \cap S \neq \emptyset$

for some $v \in V(G)$ such that $Q'_v \not\subseteq S$. If $v_1 \notin S$, and also $v_2 \notin S$, then no vertex in $S \cap Q'_v$ is connected to any vertex in $S \setminus Q'_v$. Hence, we can assume without loss of generality that $v_1 \in S$. Note that $N(D_2) = V(G_2) \cup \{p_{s-1}\}$. Furthermore, observe that $\text{dist}(v_1, p_{s-1}) = s$, that $\text{dist}(v_1, q_{s-2}^v) = s - 2$, and that $\text{dist}(v_1, q_{s-2}^u) \geq s - 1$ for each $u \in V(G) \setminus \{v\}$. Thus, the unique path of length at most s from v_1 to D_2 contains all vertices in Q'_v . Hence, $Q'_v \cap S \neq \emptyset$ if and only if $Q'_v \subseteq S$. By the definition of k' we may thus conclude that $Q'_v \subseteq S$ for at least k vertices $v \in V(G)$.

Now, assume towards a contradiction that $Q'_u \subseteq S$ and $Q'_v \subseteq S$ such that $uv \notin E(G)$. We consider the vertices v_1 and u_2 . Observe that by construction each path from v_1 to u_2 containing any vertex p_i has length at least $s + 1$. Hence, each shortest path from v_1 to u_2 contains the vertex set of Q'_w for some $w \in V(G)$. Since the path induced by each Q'_w has length $s - 1$, we conclude that $w = u$ or $w = v$. Assume without loss of generality that $w = v$. Hence, the $(s - 1)$ th vertex on the path from v_1 is v_2 . Since $uv \notin E(G)$ we have by construction that $u_2v_2 \notin E(G')$. Hence, $\text{dist}(v_1, u_2) \geq s + 1$, a contradiction to the fact that S is an s -club. Thus, $\{v \mid Q'_v \subseteq S\}$ is a clique of size at least k in G . \square

5 Conclusion

We provided a complexity dichotomy for VERTEX TRIANGLE s -CLUB and EDGE TRIANGLE s -CLUB for the standard parameter solution size k with respect to s and ℓ . Furthermore, we also provided a complexity dichotomy for SEEDED 2-CLUB for k in terms of the structure of $G[W]$. For SEEDED s -CLUB with $s \geq 3$ we provided an FPT-algorithm with respect to k when $G[W]$ is a clique and we showed W[1]-hardness for k when $G[W]$ contains at least 2 connected components. Hence, an immediate open question is the parameterized complexity of SEEDED s -CLUB for $s \geq 3$ when $G[W]$ is connected but not a clique. One aim should be to also provide a dichotomy for k for SEEDED s -CLUB with $s \geq 3$ for all possible structures of $G[W]$. It is particularly interesting to study seeds of constant size since this seems to be the most interesting case for applications.

For future work, it seems interesting to study the complexity of the considered variants of s -CLUB with respect to further parameters, for example with respect to structural parameters of the input graph G such as the treewidth of G . Additionally, the parameterized complexity of further robust variants of s -CLUB such as t -HEREDITARY s -CLUB [16,14] with respect to k remains open. It is also interesting to study other problems for detecting communities with seed constraints. One prominent example is s -PLEX. This problem is also NP-hard for $W \neq \emptyset$, since an algorithm for the case when $|W| = 1$ can be used as a black box to solve the unseeded variants. From a practical perspective, we plan to implement combinatorial algorithms for all three problem variants for the most important special case $s = 2$. Based on experience with previous implementations for 2-CLUB [10] and some of its robust variants [14] we are optimistic that these problems can be solved efficiently on sparse real-world instances.

References

1. Almeida, M.T., Brás, R.: The maximum l -triangle k -club problem: Complexity, properties, and algorithms. *Computers & Operations Research* **111**, 258–270 (2019)
2. Balasundaram, B., Butenko, S., Trukhanov, S.: Novel approaches for analyzing biological networks. *Journal of Combinatorial Optimization* **10**(1), 23–39 (2005)
3. Bodlaender, H.L., Downey, R.G., Fellows, M.R., Hermelin, D.: On problems without polynomial kernels. *Journal of Computer and System Sciences* **75**(8), 423–434 (2009)
4. Bourjolly, J., Laporte, G., Pesant, G.: An exact algorithm for the maximum k -club problem in an undirected graph. *European Journal of Operational Research* **138**(1), 21–28 (2002)
5. Carvalho, F.D., Almeida, M.T.: The triangle k -club problem. *Journal of Combinatorial Optimization* **33**(3), 814–846 (2017)
6. Chang, M., Hung, L., Lin, C., Su, P.: Finding large k -clubs in undirected graphs. *Computing* **95**(9), 739–758 (2013)
7. Cygan, M., Fomin, F.V., Kowalik, L., Lokshtanov, D., Marx, D., Pilipczuk, M., Pilipczuk, M., Saurabh, S.: *Parameterized Algorithms*. Springer (2015)
8. Downey, R.G., Fellows, M.R.: *Fundamentals of Parameterized Complexity*. Texts in Computer Science, Springer (2013)
9. Golovach, P.A., Heggenes, P., Kratsch, D., Rafiey, A.: Finding clubs in graph classes. *Discrete Applied Mathematics* **174**, 57–65 (2014)
10. Hartung, S., Komusiewicz, C., Nichterlein, A.: Parameterized algorithmics and computational experiments for finding 2-clubs. *Journal of Graph Algorithms and Applications* **19**(1), 155–190 (2015)
11. Hartung, S., Komusiewicz, C., Nichterlein, A., Suchý, O.: On structural parameterizations for the 2-club problem. *Discrete Applied Mathematics* **185**, 79–92 (2015)
12. Kanawati, R.: Seed-centric approaches for community detection in complex networks. In: *Proceedings of the 6th International Conference on Social Computing and Social Media (SCSM '14)*. Lecture Notes in Computer Science, vol. 8531, pp. 197–208. Springer (2014)
13. Komusiewicz, C.: Multivariate algorithmics for finding cohesive subnetworks. *Algorithms* **9**(1), 21 (2016)
14. Komusiewicz, C., Nichterlein, A., Niedermeier, R., Picker, M.: Exact algorithms for finding well-connected 2-clubs in sparse real-world graphs: Theory and experiments. *European Journal of Operational Research* **275**(3), 846–864 (2019)
15. Mokken, R.J., et al.: Cliques, clubs and clans. *Quality & Quantity* **13**(2), 161–173 (1979)
16. Pattillo, J., Youssef, N., Butenko, S.: On clique relaxation models in network analysis. *European Journal of Operational Research* **226**(1), 9–18 (2013)
17. Schäfer, A., Komusiewicz, C., Moser, H., Niedermeier, R.: Parameterized computational complexity of finding small-diameter subgraphs. *Optimization Letters* **6**(5), 883–891 (2012)
18. Veremyev, A., Boginski, V.: Identifying large robust network clusters via new compact formulations of maximum k -club problems. *European Journal of Operational Research* **218**(2), 316–326 (2012)
19. Whang, J.J., Gleich, D.F., Dhillon, I.S.: Overlapping community detection using seed set expansion. In: *Proceedings of the 22nd ACM International Conference on Information and Knowledge Management (CIKM '13)*. pp. 2099–2108. ACM (2013)

20. Yezerka, O., Pajouh, F.M., Butenko, S.: On biconnected and fragile subgraphs of low diameter. *European Journal of Operational Research* **263**(2), 390–400 (2017)