BSE-PROPERTIES OF SECOND DUAL OF BANACH ALGEBRAS

MARYAM AGHAKOOCHAKI AND ALI REJALI*

ABSTRACT. Let A be a commutative semisimple Arens regular unital Banach algebra. The correlation between the BSE-property of the Banach algebra A and its second duals are assessed. It is found and approved that if A is a BSE-algebra, then so is A^{**} . The opposite correlation will hold in certain conditions. The correlation of the BSE-norm property of the Banach algebra and its second dual are assessed and examined. It is revealed that, if A is a commutative Arens regular unital Banach algebra where A^{**} is semisimple, then, A and A^{**} are BSE-norm algebra.

1. INTRODUCTION

Let A be a Banach algebra. The researcher in [3] and [4], introduced the two Arens multiplications on A^{**} which convert it into a Banach algebra. A Banach algebra is an Arens regular if the Arens multiplications coincide with its second dual.

The Bochner-Schoenberg-Eberlein (BSE) is derived from the famous theorem proved in 1980 by Bochner and Schoenberg for the group of real numbers; [5] and [13]. The researcher in [8], revealed that if G is any locally compact abelian group, then the group algebra $L_1(G)$ is a BSE algebra. The researcher in [13],[15],[16] assessed the commutative Banach algebras that meet the Bochner-Schoenberg-Eberlein- type theorem and explained their properties. They are introduced and assessed in [14] the first and second types of BSE algebras. This concept is expanded in [9] and [11]. The concept of an n-BSE function and Banach algebras $C_{\text{BSE}(n)}(\Delta(A))$, for each $n \in \mathbb{N}$ is introduced in [15], where the correlation between these concepts and the BSE property is assessed. In 1992, a class of commutative Banach algebras was introduced by the same researchers in the form of the following equation:

$$\widehat{A} = C_{\rm BSE}(\Delta(A))$$

They proved that the commutative uniform C^* -algebras belong to this category; [15].

Key words and phrases. Banach algebra, BSE algebra, BSE-norm algebra, second dual of Banach algebra.

²⁰²⁰ Mathematics Subject Classification. D8, H51.

^{*}Corresponding auther.

The concept of BSE-norm algebra, where $L_1(G)$ is a BSE-norm algebra, for any locally compact group, Hausdorff, and abelian group G is devised in [16]. Another essential reference for BSE-norm algebras is [7]. The BSE property for some other Banach algebras, including the direct sum of Banach algebras is assessed in [10]. That the Lipschitz algebra $\text{Lip}_{\alpha}(K, A)$ is a BSEalgebra if and only if A is a BSE-algebra, where K is a compact metric space, A is a commutative unital semisimple Banach algebra, and $0 < \alpha \leq 1$ is proved in [1].

In this article, A is a commutative unital Banach algebra, where A is Arens regular and A^{**} is semisimple. That the $C_{BSE}(\Delta(A^{**})) \cong C_{BSE}(\Delta(A))$ will be proved in the article first, followed by proving that if A is a BSEalgebra, so is A^{**} . It will be concluded that if A^{**} is a BSE-algebra and $\widehat{A}|_{\Delta(A)} \cong \widehat{A^{**}}|_{\widehat{\Delta(A)}}$, then A is a BSE algebra.

The BSE- norm property for A^{**} and proving that A and A^{**} are BSEnorm algebras are assessed in this article.

2. Preliminaries

The basic terminologies and the related information on BSE algebras are extracted from [14], [15], and [16].

Let A be a commutative semisimple Banach algebra, and $\Delta(A)$ be the character space of A with the Gelfand topology. In this study, $\Delta(A)$ represents the set of all non-zero multiplicative linear functionals over A.

Assume that $C_b(\Delta(A))$ is the space consisting of all complex-valued continuous and bounded functions on $\Delta(A)$. A linear operator T on A is named a multiplier if for all $x, y \in A$, T(xy) = xT(y). The set of all multipliers on A will be expressed as M(A). It is obvious that M(A) is a Banach algebra, and if A is an unital Banach algebra, then $M(A) \cong A$. As observed in [12], for each $T \in M(A)$ there exists a unique bounded continuous function \hat{T} on $\Delta(A)$ expressed as:

$$\varphi(Tx) = \widehat{T}(\varphi)\varphi(x),$$

for all $x \in A$ and $\varphi \in \Delta(A)$. By setting $\{\widehat{T} : T \in M(A)\}$, the $\widehat{M(A)}$ is yield.

If the Banach algebra A is semisimple, then the Gelfand map $\Gamma_A : A \to \widehat{A}$, $f \mapsto \widehat{f}$, is injective or equivalently, and the following equation holds:

$$\bigcap_{\varphi \in \Delta(A)} \ker(\varphi) = \{0\}$$

A bounded complex-valued continuous function σ on $\Delta(A)$ is named a BSE function, if there exists a positive real number β in a sense that for every finite complex-number c_1, \dots, c_n , and the same many $\varphi_1, \dots, \varphi_n$ in $\Delta(A)$ the following inequality

$$|\sum_{j=1}^{n} c_j \sigma(\varphi_j)| \leq \beta \|\sum_{j=1}^{n} c_j \varphi_j\|_{A^*}$$

 $\mathbf{2}$

3

holds.

The set of all BSE-functions is expressed by $C_{BSE}(\Delta(A))$, where for each σ , the BSE-norm of σ , $\|\sigma\|_{BSE}$ is the infimum of all β s applied in the above inequality. That $(C_{BSE}(\Delta(A)), \|.\|_{BSE})$ is a semisimple Banach subalgebra of $C_b(\Delta(A))$ is in Lemma 1 proved in [14]. Algebra A is named a BSE algebra if it meets the following condition:

$$\widetilde{M(A)} \cong C_{\text{BSE}}(\Delta(A)).$$

If A is unital, then $\widehat{M(A)} \cong \widehat{A}|_{\Delta(A)}$, indicating that A is a BSE algebra if and only if $C_{BSE}(\Delta(A)) \cong \widehat{A}|_{\Delta(A)}$. The semisimple Banach algebra A is named a norm-BSE algebra if there exists some K > 0 in a sense that for each $a \in A$, the following holds:

$$||a||_A \le K ||\hat{a}||_{\text{BSE}}$$

Let A be a Banach algebra, $F, G \in A^{**}$, $f \in A^*$ and $x, y \in A$. 1) The first Arens multiplication is defined by:

$$F\Box G(f) = F(G.f)$$
$$G.f(x) = G(f.x)$$
$$f.x(y) = f(xy)$$

2) The second Arens multiplication on A^{**} is defined by:

$$F \Diamond G(f) = G(f.F)$$

$$f.F(x) = F(x.f)$$

$$x.f(y) = f(yx)$$

It is obvious that (A^{**}, \Box) and (A^{**}, \diamondsuit) are Banach algebras. If the binary operation \Box is the same as the action of \diamondsuit , then A is named the Arens regular Banach algebra. If A is a commutative Banach algebra, then A^{**} is a commutative Banach algebra if and only if A is an Arens regular Banach algebra; [6].

3. BSE Algebras and their second Dual

The structure of the BSE functions on $\Delta(A^{**})$ is characterization and the correlations between the BSE property of A and their second duals are assessed.

Let A be a commutative semisimple Banach algebra. Then (i) $C_{\text{BSE}}(\Delta(A))$ equals the set of all $\sigma \in C_b(\Delta(A))$ for which there exists at least one bounded net (a_α) in A with $\lim \widehat{a_\alpha}(\varphi) = \sigma(\varphi)$ for all $\varphi \in \Delta(A)$. (ii) $C_{\text{BSE}}(\Delta(A)) = C_b(\Delta(A)) \cap A^{**}$, where A^{**} is the second dual of A. Proved in [Theorem4, [14]].

Lemma 1. Let A be a commutative semisimple Banach algebra. Then

$$\overline{\langle \Delta(A) \rangle}^{w^*} = A^*$$

Proof. Assume that $B = \overline{\langle \Delta(A) \rangle}^{w^*}$ and $B \subsetneq A^*$. Define $\Theta : B \longrightarrow \mathbb{C}$ by $\Theta(b) = 0$ for all $b \in B$. Let $b_0 \in A^* \setminus B$. Then by referring to the Hahn–Banach theorem there exists a w^* - continuous linear function $\overline{\Theta} :$ $A^* \longrightarrow \mathbb{C}$ where $\overline{\Theta}(b_0) = 1$ and $\overline{\Theta}(b) = 0$ for all $b \in B$. Because (A^*, w^*) is a locally convex and $\overline{\Theta}$ is w^* -continuous, thus there exists $a \in A$ where $\overline{\Theta} = \hat{a}$. If $\varphi \in \Delta(A)$, the following is yield:

$$\varphi(a) = \hat{a}(\varphi) = \overline{\Theta}(\varphi) = 0$$

Because A is semisimple, a = 0. Thus,

$$\overline{\Theta}(b_0) = \hat{a}(b_0) = b_0(0) = 0$$

which is a contradiction; therefore $\overline{\langle \Delta(A) \rangle}^{w^*} = A^*$.

Lemma 2. Let A be a commutative Banach algebra and $\overline{\langle \Delta(A) \rangle}^{w^*} = A^*$. Then $\Delta(A^{**}) = \overline{\Delta(A)}^{w^*}$.

Proof. Provided in [6].

Corollary 3. Let A be a commutative semisimple unital Banach algebra. Then

$$\Delta(A^{**}) = \widehat{\Delta(A)}$$

Proof. Because A is unital, $\Delta(A)$ is w^* - compact, thus, $\widehat{\Delta(A)} = \overline{\Delta(A)}^w^*$. Then by applying Lemma 1 and Lemma 2, the equation $\Delta(A^{**}) = \widehat{\Delta(A)}$ is yield.

Remark 1. Let A be a commutative semisimple unital Banach algebra. Assume that $\varphi_1, \dots, \varphi_n \in \Delta(A)$ and $c_1, \dots, c_n \in \mathbb{C}$, then the following is yield:

$$\|\sum_{i=1}^{n} c_{i}\varphi_{i}\|_{A^{*}} = \|\sum_{i=1}^{n} c_{i}\widehat{\varphi}_{i}\|_{A^{***}}$$

Remark 2. Let A be a commutative Arens regular Banach algebra. If A^{**} is semisimple, then A is semisimple. Because $\widehat{\ker(\varphi)} \subseteq \ker(\widehat{\varphi})$ and $\widehat{\operatorname{Rad}(A)} \subseteq \operatorname{Rad}(A^{**})$. then A is semisimple, because A^{**} is semisimple, .

Theorem 4. Let A be a commutative Arens regular semisimple unital Banach algebra. Then the following statement holds:

$$C_{\rm BSE}(\Delta(A^{**})) \cong C_{\rm BSE}(\Delta(A))$$

These two as Banach algebras are isometric;

Proof. Because A is semisimple, by applying Corollary 3, the $\Delta(A^{**}) = \widehat{\Delta(A)}$ is yield, thus, $C_b(\Delta(A^{**})) \cong C_b(\Delta(A))$, and $A^{****} \mid_{\widehat{\Delta(A)}} \cong A^{**} \mid_{\Delta(A)}$. Consequently by applying [14, Theorem 4], $C_{BSE}(\Delta(A^{**})) \cong C_{BSE}(\Delta(A))$ is yield.

The correlation between the BSE-property of algebra A and it's second dual A^{**} is assessed as follows:

Lemma 5. Let A be a commutative unital Banach algebra such that A is Arens regular and A^{**} is semisimple. If A is a BSE algebra, so is A^{**} .

Proof. Let $\Sigma \in C_{BSE}(\Delta(A^{**}))$. Define $\sigma(\varphi) = \Sigma(\widehat{\varphi})$ for all $\varphi \in \Delta(A)$. Obviously, σ is well defined and $\sigma \in C_{BSE}(\Delta(A))$. By allowing $\varphi_1, \dots, \varphi_n \in \Delta(A)$ and $c_1, \dots, c_n \in \mathbb{C}$ the following is yield:

$$\sum_{j=1}^{n} c_j \sigma(\varphi_j) \mid = \mid \sum_{j=1}^{n} c_j \Sigma(\widehat{\varphi_j}) \mid$$
$$\leq M \mid\mid \sum_{j=1}^{n} c_j \widehat{\varphi_j} \mid\mid_{A^{***}}$$
$$= M \mid\mid \sum_{j=1}^{n} c_j \varphi_j \mid\mid_{A^{*}}$$

where M > 0. Because $C_{BSE}(\Delta(A)) = \widehat{A}|_{\Delta(A)}$, there exists $a \in A$ where $\sigma = \widehat{a}$.

Assume that $\Phi \in \Delta(A^{**})$, thus $\Phi = \hat{\varphi}$ for some $\varphi \in \Delta(A)$. Which yield:

$$\Sigma(\Phi) = \Sigma(\hat{\varphi}) = \sigma(\varphi)$$
$$= \hat{a}(\varphi)$$
$$= \hat{a}(\hat{\varphi})$$
$$= \hat{a}(\Phi)$$

where, $\Sigma = \widehat{\hat{a}}$, therefore $C_{\text{BSE}}(\Delta(A^{**})) = \widehat{A^{**}}|_{\widehat{\Delta(A)}}$.

Lemma 6. Let A be a commutative unital Banach algebra where A is Arens regular and A^{**} is semisimple. If A^{**} is a BSE algebra and $\widehat{A} \mid_{\Delta(A)} \cong \widehat{A^{**}} \mid_{\widehat{\Delta(A)}}$, then A is a BSE-algebra.

Proof. Assume that A^{**} is a BSE algebra, the following is yield:

$$C_{BSE}(\Delta(A)) \cong C_{BSE}(\Delta(A^{**}))$$
$$\cong \widehat{A^{**}} \mid_{\widehat{\Delta(A)}}$$
$$\cong \widehat{A} \mid_{\Delta(A)}$$

thus, A is a BSE- algebra.

At this stage, based on the established prerequisite the primary Theorem, is expressed as follows:

Theorem 7. Let A be a commutative unial Banach algebra, such that A is Arens regular and A^{**} is semisimple. Then the following statements are equivalent:

(i) A is a BSE algebra. (ii) A^{**} is a BSE algebra and $\widehat{A} \mid_{\Delta(A)} \cong \widehat{A^{**}} \mid_{\widehat{\Delta(A)}}$.

Example 1. Let (K, d) be a compact metric space and A be a unital commutative semisimple Banach algebra. Then A is a BSE-algebra if and only if Lip(K, A) is a BSE-algebra; [1]. In this context, the following statements are equivalent:

(i) A is a BSE algebra.
(ii) A^{**} is a BSE algebra and |_{Δ(A)}≅ Â^{**} |_{Â(A)}.
(iii) Lip(K, A) is a BSE-algebra.
(iv) Lip(K, A^{**}) is a BSE-algebra and |_{Δ(A)}≅ Â^{**} |_{Â(A)}.

Example 2. Let X be a metric space and A = Lip(X). Then A is unital and BSE algebra; [2]. By applying Lemma 5, $\text{Lip}(X)^{**}$ becomes a BSE algebra.

Example 3. Let A be a commutative reflexive Banach algebra. Then

$$C_{\rm BSE}(\Delta(A)) = A|_{\Delta(A)}$$

Moreover, A is unital, then A is a BSE algebra.

Proof. According to [Theorem 4, [14]] because A is reflexive, and the following is yield:

$$C_{\text{BSE}}(\Delta(A)) = A^{**} \mid_{\Delta(A)} \cap C_b(\Delta(A))$$
$$= \widehat{A} \mid_{\Delta(A)} \cap C_b(\Delta(A))$$
$$= \widehat{A} \mid_{\Delta(A)}$$

4. BSE-NORM ALGEBRA

The BSE-norm property of the second dual of Banach algebra is assessed and the correlation between the BSE-norm of the Banach algebra A and its second dual is assessed.

Theorem 8. Let A be a commutative unial Banach algebra such that A is Arens regular and A^{**} is semisimple. Then A^{**} is a BSE-norm algebra.

Proof. By applying the open mapping theorem there exists some M > 0where $||F||_{A^{**}} \leq M||F||_{\Delta(A)} ||_{A^{**}}$ for each $F \in A^{**}$, because the map $\Theta: A^{**} \longrightarrow C_b(\Delta(A))$ given by $\Theta(F) = F|_{\Delta(A)}$ is continuous, linear and

injective, thus:

$$\begin{aligned} \|\hat{F}\|_{\text{BSE}} &= \sup\{ \|\sum_{i=1}^{n} c_{i} \hat{F}(\Phi_{i}) \| \Phi_{i} \in \Delta(A^{**}), \|\sum_{i=1}^{n} c_{i} \Phi_{i}\|_{A^{***}} \leq 1 \} \\ &= \sup\{ \|\sum_{i=1}^{n} c_{i} \hat{F}(\hat{\varphi}_{i}) \| \varphi_{i} \in \Delta(A), \|\sum_{i=1}^{n} c_{i} \varphi_{i}\|_{A^{*}} \leq 1 \} \\ &= \sup\{ \|\sum_{i=1}^{n} c_{i} F(\varphi_{i}) \| \varphi_{i} \in \Delta(A), \|\sum_{i=1}^{n} c_{i} \varphi_{i}\|_{A^{*}} \leq 1 \} \\ &= \|F\|_{\Delta(A)} \|_{\text{BSE}} \end{aligned}$$

in addition:

$$\|F\|_{\Delta(A)} \|_{A^{**}} = \sup\{|F(\varphi)| : \varphi \in \Delta(A)\}$$

$$\leq \|F\|_{\Delta(A)} \|_{BSE}$$

Then by applying the open mapping theorem, the following is yield:

 $||F||_{A^{**}} \le M ||F||_{\Delta(A)} ||_{A^{**}} \le M ||F||_{\Delta(A)} ||_{BSE} = M ||\hat{F}||_{BSE}$

therefore, A^{**} is a BSE-norm algebra.

Theorem 9. Let A be a commutative unital Banach algebra, such that A is Arens regular and A^{**} is semisimple. Then A is a BSE-norm algebra.

Proof. Because A^{**} is a BSE-norm algebra, consequently, by definition there exists M > 0 where $||F||_{A^{**}} \leq M ||\widehat{F}||_{BSE}$ for each $F \in A^{**}$. If $a \in A$, the following is yield:

$$||a||_A = ||\hat{a}||_{A^{**}} \le M ||\hat{a}||_{BSE}$$

in addition

$$\begin{aligned} \|\widehat{\hat{a}}\|_{BSE} &= \sup\{|\sum_{i=1}^{n} c_{i} \Phi_{i}(\widehat{a})| \mid \Phi_{i} \in \Delta(A^{**}), \|\sum_{i=1}^{n} c_{i} \Phi_{i}\|_{A^{***}} \leq 1\} \\ &= \sup\{|\sum_{i=1}^{n} c_{i} \widehat{\varphi_{i}}(\widehat{a})| \mid \|\sum_{i=1}^{n} c_{i} \widehat{\varphi_{i}}\|_{A^{***}} \leq 1\} \\ &= \sup\{|\sum_{i=1}^{n} c_{i} \varphi_{i}(a)| \mid \|\sum_{i=1}^{n} c_{i} \varphi_{i}\|_{A^{*}} \leq 1\} \\ &= \|\widehat{a}\|_{BSE} \end{aligned}$$

therefore, $||a||_A \leq M ||\hat{a}||_{BSE}$. Then A is a BSE-norm algebra.

Corollary 10. Let A be a commutative unital Banach algebra, such that A is Arens regular and A^{**} is semisimple. Then A and A^{**} are BSE-norm algebras.

Example 4. Let A be a commutative semisimple reflexive Banach algebra where $\Delta(A)$ is a discrete space. Then A is a BSE-norm algebra.

7

Proof. By applying Example 3, $C_{BSE}(\Delta(A)) = \hat{A}|_{\Delta(A)}$, making $\hat{A}|_{\Delta(A)}$ is a closed BSE-norm space, hence $\Phi : A \longrightarrow \hat{A}|_{\Delta(A)}$ is an one-to-one continuous map where $\Phi(A) = \hat{A}|_{\Delta(A)}$. Then by applying the open mapping theorem it is revealed that there exists some M > 0 where $||a||_A \le M ||\hat{a}||_{BSE}$ for each $a \in A$, therefore A is a BSE-norm algebra.

Example 5. 1) If A is a commutative unital finite dimensional Banach algebra, then A is a BSE-norm algebra. 2) Let G be a compact group and $A = l_p(G)$. Then A is a BSE-norm algebra.

Proof. 1) Since A is finite dimensional, $A \cong C^n$, for some positive integer n under equivalent norms. Example 4 implies that A is a BSE-norm algebra. 2) Because $\Delta(A) = \hat{G}$ is a discrete space, A is a semisimple and reflexive algebra, thus by according to Example 4, A is a BSE-norm algebra. \Box

Example 6. Assume that G is a compact group and A is a reflexive semisimple Banach algebra where $\Delta(A)$ is discrete and $1 . Then <math>L_p(G, A)$ is a BSE-norm algebra.

Proof. The following can be written:

$$L_p(G, A)^{**} \cong L_q(G, A^*)^*$$
$$\cong L_p(G, A^{**}) \cong L_p(G, A)$$

Consequently $L_p(G, A)$ is reflexive, and

$$\Delta(L_p(G,A)) = \widehat{G} \times \Delta(A)$$

Because G is a compact group, and $\Delta(A)$ is discrete, it is concluded that $\Delta(L_p(G, A))$ is discrete. Therefore by applying Example 4, $L_p(G, A)$ is a BSE-norm algebra.

5. Acknowledgment

Appreciation is extended to Assistant Professor Hojjat Farzadfard for his valuable comments and suggestions regarding the manuscript.

References

- F. Abtahi, Z. Kamali, M. Toutounchi, The Bochner Schoenberg- Eberline type property for vector- valued Lipschitz algebras. J.Math. Anal. Appl. (2019).
- [2] F. Abtahi, Z. Kamali, M. Toutounchi, THE BSE concepts for vector- valued Lipschitz algebras. Commun. Pure Appl. Anal. (2020).
- [3] R. Arens, Operations induced in function classes, Monatsch. Math. 55 (1951), 1-19.
- [4] R. Arens, The adjoint of a bilinear operation, Proc. Amer. Math. Soc. 2 (1951), 839-848.
- [5] S. Bochner, A theorem on Fourier- Stieltjes integrals, Bull.Amer Math. Soc., 40(1934),271-276.
- [6] H.G. Dales, A.T.-M. Lau, The second duals of Beurling algebras, Mem. Amer. Math. Soc. 177 (836) (2005).

- [7] H. G. Dales and A. Ulger, Approximate identities and BSE norms for Banach function algebras. Fiels Institute, Toronto, 2014.
- [8] W. F. Eberlein, Characterizations of Fourier- Stieltjes transforms, Duke Math. J.,22(1955), 465- 468.
- [9] J.Inoue and S-E. Takahasi, On characterization of image of the Gelfand transform of commutative Banach algebras, Math. Nachr, 280 (2007), 105- 129.
- [10] Z. Kamali and M. Lashkarizadeh Bami, The multiplier algebras and BSE property of the direct sum of Banach algebras, Bull. Of Austral . Math. Soc., 88, No.2, (2013), 250-n258.
- [11] E. Kaniuth and A. Ulger, The Bochner Schoenberg- Eberline property for commutative Banach algebras, especially Fourier- Stieltjes algebras, Trans. Amer. Math. Soc., 362(2010), 4331- 4356.
- [12] R. Larsen, An Introduction to the Theory of Multipliers (Springer, New York, 1971).
- [13] I.J. Schoenberg, A remark on the preceding note by Bochner, Bull.Amer Math. Soc., 40(1934),277-278.
- [14] S. E.Takahasi and O.Hatori, Commutative Banach algebras Which satisfy a Bochner- Schonberg- Eberlein- type theorem, Proc.Amer. Math. Soc., 110(1990), 149-158.
- [15] S.E.Takahasi and O.Hatori, Commutative Banach algebras and BSE- inequalities, Math. Japon, 37(1992), 47- 52.
- [16] S.E.Takahasi and O.Hatori and K. Tanahashi, Commutative Banach algebras and BSE- norm, Math. Japon,46(1997), 59- 80.

M. Aghakoochaki Department of Pure Mathematis Faculty of Mathematics and Statistics University of Isfahan Isfahan, 81746-73441 Iran mkoochaki@sci.ui.ac.ir

A. Rejali Department of Pure Mathematis Faculty of Mathematics and Statistics University of Isfahan Isfahan, 81746-73441 Iran rejali@sci.ui.ac.ir