CHOW MOTIVES OF GENUS ONE FIBRATIONS

(*) DAIKI KAWABE

ABSTRACT. Let $f: X \to C$ be a genus 1 fibration from a smooth projective surface, i.e. its generic fiber is a regular genus 1 curve. Let $j: J \to C$ be the Jacobian fibration of f. In this paper, we prove that the Chow motives of X and J are isomorphic. As an application, combined with our concomitant work on motives of quasi-elliptic fibrations, we prove Kimura finite-dimensionality for smooth projective surfaces not of general type with geometric genus 0. This generalizes Bloch-Kas-Lieberman's result to arbitrary characteristic.

1. INTRODUCTION

1.1. Motivation. Let k be an algebraically closed field of arbitrary characteristic. Let $f : X \to C$ be a fibration from a smooth projective surface over k to a smooth projective curve. It is a proper, surjective, k-morphism such that $f_*\mathcal{O}_X \cong \mathcal{O}_C$. Let η be the generic point of C and X_η the generic fiber of f. We study the following:

(i) f is a genus 1 fibration if X_n is a regular genus 1 curve. More precisely,

 X_{η} is a *regular*, projective, geometrically-integral, curve with arithmetic genus 1.

- (ii) A genus 1 fibration f is elliptic if X_{η} is smooth, i.e. geometrically-regular.
- (iii) A genus 1 fibration f is quasi-elliptic if X_n is not smooth.

Note that quasi-elliptic surfaces exist only in characteristic 2 or 3 (Proposition 5.6). We say that a genus 1 fibration $f: X \to C$ is *minimal* if every birational morphism $f: X \to X'$ over C onto a smooth projective surface X' is an isomorphism.

From now on, we let $f: X \to C$ be a genus 1 fibration and assume f is minimal if not stated otherwise. Then X_{η} does not necessarily have a η -rational point, hence fmay have multiple fibers. To remedy this problem, we consider the Jacobian fibration $j: J \to C$ of f ([CD89]). Its generic fiber J_{η} is the regular compactification of the Jacobian variety of X_{η} . Then, J is a smooth projective surface, and j is a genus 1 fibration that has no multiple fibers. Clearly, it is easier to study J than X. It is known that there are relations between some invariants of X and J, e.g. the equalities of the i-th Betti numbers $b_i(X) = b_i(J)$ and the Picard numbers $\rho(X) = \rho(J)$ ([CD89]). For a smooth projective surface S over k, we denote by h(S) the Chow motive of S with \mathbb{Q} -coefficients, and by T(S) the kernel of the Albanese map $a_S : \operatorname{CH}_0(S)_{\mathbb{Z}}^0 \to \operatorname{Alb}_{S/k}(k)$.

In 1976, Bloch-Kas-Lieberman proved the following relation between X and J:

Proposition 1.1. ([BKL76, Prop. 4, p.138]). Let $f : X \to C$ be an elliptic fibration over \mathbb{C} and $j : J \to C$ the Jacobian fibration of f. If T(J) = 0, then T(X) = 0.

Date: March 9, 2024.

Key words and phrases. Chow motives, genus one fibrations, surfaces not of general type, Kimura-O'Sullivan finite-dimensionality.

^(*) Email address: daiki.kawabe.math@gmail.com, ORCID: 0000-0003-2766-7115

Conflict of interest: This work is supported by the JSPS KAKENHI Grant Number 18H03667 and Tohoku University Global Hagi Scholarship.

(*) DAIKI KAWABE

In 1992, K. Coombes proved the following relation between X and J:

Proposition 1.2. ([Coo92, Prop. 3.1, p.52]). Let k be an algebraically closed field. Let X be an Enriques surface over k with an elliptic fibration $f : X \to \mathbb{P}^1$, and $j : J \to \mathbb{P}^1$ the Jacobian fibraton of f. Then there is an isomorphism of Chow motives $h(X) \cong h(J)$.

The author was inspired by Propositions 1.1 and 1.2. In this paper, we generalize Proposition 1.2 for genus 1 fibrations defined over arbitrary algebraically closed fields.

S. Kimura and O'Sullivan independently introduced a notion of finite-dimensionality for Chow motives ([Kim05, Def. 3.7] and [And04, Ch. 12]). We call it Kimurafiniteness. They conjectured that every Chow motive is Kimura-finite. Moreover, for a surface S over \mathbb{C} with $p_g = 0$, Kimura proved the theorem that h(S) is Kimura-finite if and only if the Albanese map $a_S : CH_0(S)^0_{\mathbb{Z}} \to Alb_{S/\mathbb{C}}(\mathbb{C})$ is an isomorphism ([Kim05, Coro. 7.7]). By using his theorem, the result of Bloch-Kas-Lieberman [BKL76] (see Theorem 11.1) is equivalent to the following:

Theorem 1.3. (= Theorem 11.2). Let X be a smooth projective surface over \mathbb{C} . Assume that X has geometric genus 0 and Kodaira dimension < 2. Then h(X) is Kimura-finite.

In this paper, we generalize Theorem 1.3 to arbitrary characteristic.

1.2. Main theorems of this paper.

We prove two main theorems (Theorems 1.4 and 1.7). The first one is the following:

Theorem 1.4. (= Theorem 8.1). Let k be an arbitrary algebraically closed field. Let $f: X \to C$ be a minimal genus 1 fibration from a smooth projective surface over k, and $j: J \to C$ the Jacobian fibration of f. Then, there is an isomorphism

$$h(X) \cong h(J)$$

in the category $CH\mathcal{M}(k,\mathbb{Q})$ of Chow motives over k with \mathbb{Q} -coefficients.

Theorem 1.4 is a generalization of Proposition 1.2 to genus 1 fibrations. Here, we give a sketch of the proof of Theorem 1.4. Let us consider the Chow-Künneth decompositions of Murre [Mur90] (see Proposition 4.2) of h(X) and h(J), respectively

$$h(X) \cong \bigoplus_{i=0}^{4} h_i(X) \cong 1 \oplus h_1(X) \oplus h_2^{alg}(X) \oplus t_2(X) \oplus h_3(X) \oplus (\mathbb{L} \otimes \mathbb{L})$$
$$h(J) \cong \bigoplus_{i=0}^{4} h_i(J) \cong 1 \oplus h_1(J) \oplus h_2^{alg}(J) \oplus t_2(J) \oplus h_3(J) \oplus (\mathbb{L} \otimes \mathbb{L}).$$

Here, 1 is the unit motive, \mathbb{L} is the Lefschetz motive, and $h_2^{alg}(-)$ (resp. $t_2(-)$) is the algebraic (resp. transcendental) part of $h_2(-)$ ([KMP07]). Thus, it suffices to prove

$$h_i(X) \cong h_i(J)$$
 for $1 \le i \le 3$.

First, assume i = 1 or 3. For a smooth projective variety V over k, we denote by $(\operatorname{Pic}_{V/k}^0)_{red}$ (resp. $\operatorname{Alb}_{V/k}$) the Picard (resp. Albanese) variety of V. We prove the following key proposition:

Proposition 1.5. (= Proposition 7.16). There are isogenies of abelian k-varieties

$$(\operatorname{Pic}^{0}_{X/k})_{red} \sim_{isog} (\operatorname{Pic}^{0}_{J/k})_{red}, \quad \operatorname{Alb}_{X/k} \sim_{isog} \operatorname{Alb}_{J/k}.$$

Using Proposition 1.5, we prove $h_i(X) \cong h_i(J)$ for i = 1 or 3.

Finally, assume i = 2. By $\rho(X) = \rho(J)$, we get $h_2^{alg}(X) \cong \mathbb{L}^{\oplus \rho(X)} \cong h_2^{alg}(J)$. Thus, it remains to prove $t_2(X) \cong t_2(J)$. The outline of the proof is as follows. If f is quasi-elliptic, so is j. Using the result of the author ([Kaw22]), we get

$$t_2(X) = 0 = t_2(J).$$

Thus, it suffices to consider the case where f is elliptic. Let us consider the following functors between the category of Chow motives

$$\operatorname{CH}\mathcal{M}(\eta, \mathbb{Q}) \xleftarrow{i} \operatorname{CH}\mathcal{M}(C, \mathbb{Q}) \xrightarrow{F} \operatorname{CH}\mathcal{M}(k, \mathbb{Q}).$$

Here $\operatorname{CH}\mathcal{M}(C,\mathbb{Q})$ is the category of Chow motives over C ([CH00, Def. 2.8]). In particular, *i* is not fully-faithful. Then, we consider the following two-step process.

(i) We prove an isomorphism of Chow motives between the generic fibers

$$h(X_{\eta}) \cong h(J_{\eta})$$
 in $CH\mathcal{M}(\eta, \mathbb{Q}).$

(ii) We extend the isomorphism $h(X_n) \cong h(J_n)$ to the isomorphism

$$t_2(X) \cong t_2(J)$$
 in $\operatorname{CH}\mathcal{M}(k, \mathbb{Q}).$

Concerning (i), we prove

Theorem 1.6. (= Theorem 6.1). Let K be an arbitrary field. Let A be an abelian variety over K. Let T be a torsor over K for A. Then there is an isomorphism $h(T) \cong h(A)$ in the category $CH\mathcal{M}(K, \mathbb{Q})$ of Chow motives over K with \mathbb{Q} -coefficients.

The second main theorem of this paper is the following:

Theorem 1.7. (= Theorem 11.3). Let X be a smooth projective surface over an algebraically closed field k of characteristic $p \ge 0$. Assume that X has geometric genus 0 and Kodaira dimension < 2, that is, $p_g = 0$ and $\kappa < 2$. Then h(X) is Kimura-finite in the category $CH\mathcal{M}(k,\mathbb{Q})$ of Chow motives over k with \mathbb{Q} -coefficients.

Theorem 1.7 is a generalization of Theorem 1.3 to arbitrary characteristic. The outline of the proof of Theorem 1.7 is as follows. If $\kappa < 0$, the assertion is clear. Assume $\kappa = 0$ or 1. Then X has a genus 1 fibration $f: X \to C$ by the classification of surfaces. We take the Jacobian fibration $j: J \to C$ of f, and prove h(J) is Kimurafinite. By Theorem 1.4, we have $h(X) \cong h(J)$, and see h(X) is Kimura-finite.

1.3. Organization. This paper is organized as follows.

The main parts of this paper are Sections 8 (Theorem 1.4) and 11 (Theorem 1.7). In Section 2, we recall several basic objects in algebraic geometry. In Section 3, we collect and prove some facts about relative correspondences. In Section 4, we recall the theory of Chow motives, e.g. Chow-Künneth decompositions and transcendental motives. In Section 5, to prove Theorem 1.6, we prove several facts about torsors for commutative group varieties. We treat (not necessarily smooth) genus 1 curves.

In Section 6, we prove Theorem 1.6 $(h(T) \cong h(A))$ which is crucial for the proof of $t_2(X) \cong t_2(J)$. In Section 7, we recall relations between some invariants of a genus 1 fibration and the associated Jacobian fibration. Moreover, we prove $h_i(X) \cong h_i(J)$ for i = 1, 3 and $h_2^{alg}(X) \cong h_2^{alg}(J)$. This section is based on [CD89] and [CDLK].

In Section 8, using the results of Sections 2 - 7, we prove Theorem 1.4 $(h(X) \cong h(J))$. We extend $h(X_{\eta}) \cong h(J_{\eta})$ to $t_2(X) \cong t_2(J)$. In Section 9, we collect some properties of Kimura-finiteness. Recall that any curve has Kimura-finiteness and Kimura-finiteness is stable under direct sums, tensor products, and quotients (Proposition 9.3). In Section 10, we quickly review the classification of surfaces.

In Section 11, using the results of Sections 8 - 10, we prove Theorem 1.7.

1.4. Conventions and Terminology. We fix a base field k. By k-variety we mean a reduced separated k-scheme of finite type. Unless otherwise stated, we assume irreducibility for any k-variety. By k-curve (resp. k-surface) we mean a variety of dimension 1 (resp. dimension 2).

Let $i \in \mathbb{Z}_{\geq 0}$. For a k-scheme X, We denote by $\operatorname{CH}_i(X)$ (resp. $\operatorname{CH}^i(X)$) the Chow group of *i*-dimensional (resp. *i*-codimensional) cycles on X modulo rational equivalence with \mathbb{Q} -coefficients, and set $\operatorname{CH}(X) = \bigoplus_i \operatorname{CH}^i(X)$. For an irreducible k-variety X, we have $\operatorname{CH}_i(X) = \operatorname{CH}^{\dim(X)-i}(X)$. Let $f: X \to Y$ be a morphism of k-schemes. If f is proper, then $f_*: \operatorname{CH}_i(X) \to \operatorname{CH}_i(Y)$ denote the proper-pushfoward. If f is flat of relative dimension l, then $f^*: \operatorname{CH}_i(X) \to \operatorname{CH}_{i+l}(Y)$ denote the flat-pullback.

Let $\mathcal{V}(k)$ denote the category of smooth projective k-varieties. For $X, Y \in \mathcal{V}(k)$, set $\operatorname{Corr}^{r}(X, Y) := \bigoplus_{\alpha} \operatorname{CH}^{\dim(X_{\alpha})+r}(X_{\alpha} \times Y)$ where $X = \sqcup X_{\alpha}$, with X_{α} equidimensional.

- For X an irreducible variety over k, we use following notations: k(X): the function field of X X_M := X ×_{Spec(k)} Spec(M) for any extension M of k X(M) := Hom_{Sch(k)}(Spec(M), X) for an extension M of k
 For X a projective variety over k, we use following notations:
- $h^{i}(\mathcal{F}) = h^{i}(X, \mathcal{F}) = \dim_{k} H^{i}(X, \mathcal{F})$ for any coherent sheaf \mathcal{F} on X $\chi(\mathcal{F}) := \sum_{i} (-1)^{i} h^{i}(\mathcal{F})$ for any coherent sheaf \mathcal{F} on X $p_{a}(X) := (-1)^{\dim(X)}(\chi(\mathcal{O}_{X}) - 1)$: the arithmetic genus $q(X) := h^{1}(X, \mathcal{O}_{X})$: the irregularity ω_{X} : the dualizing sheaf of X
- For X a smooth projective variety over k, we use following notations: ω_X : the canonical sheaf of X

 K_X : the canonical divisor of X

 $P_m(X) := h^0(X, \omega_X^{\otimes m})$: the *m*-genus, for $m = 1, 2, \cdots$

- $p_g(X) := P_1(X)$: the geometric genus
- $b_i(X) := \dim_{\mathbb{Q}_l} H^i_{\text{\acute{e}t}}(X, \mathbb{Q}_l)$: the *i*-th Betti number for a prime number $l \neq \text{char}(k)$ $e(X) := \sum_i (-1)^i b_i(X)$: the topological Euler characteristic
- For S a smooth projective surface over k, we have $p_q(S) = h^0(S, \omega_S) = h^2(S, \mathcal{O}_S)$.
- For simplicity, we use following notations:
 - $X \cong Y$ X and Y are isomorphic.
 - $X \sim_{birat} Y$ X and Y are birationally equivalent as varieties
 - $A \sim_{isog} B$ A and B are isogenous as abelian varieties

In the rest of this paper, let k be an algebraically closed field of arbitrary characteristic if not stated otherwise, i.e. $k = \overline{k}$.

2. Picard schemes

In this section, we recall several basic objects in algebraic geometry.

• For a scheme X, we denote by Pic(X) the *Picard group* of X. Its elements are isomorphism classes of invertible sheaves on X. Then

$$\operatorname{Pic}(X) \cong H^1_{Zar}(X, \mathcal{O}_X^*) \cong H^1_{\acute{e}t}(X, \mathbb{G}_m).$$

The second isomorphism uses Hilbert's Theorem 90 ([Mil80, III. Prop. 4.9, p.124]).

• For a scheme X, we denote by Div(X) the group of *Cartier divisors* on X.

• Let S be a scheme and $f: X \to S$ an S-scheme. The relative Picard functor $\underline{\operatorname{Pic}}_{X/S}$ is defined by

$$\underline{\operatorname{Pic}}_{X/S} : (\operatorname{Sch}/S)^{\operatorname{op}} \to (\operatorname{Sets}) \; ; \; T \mapsto \operatorname{Pic}(X_T) / f_T^* \operatorname{Pic}(T)$$

where $f_T : X_T := X \times_S T \to T$ is the second projection. We denote by $\underline{\operatorname{Pic}}_{(X/S)(\acute{et})}$ (resp. $\underline{\operatorname{Pic}}_{(X/S)(fppf)}$) its associated sheaf in the étale (resp. fppf) topology.

Theorem 2.1. Let $f : X \to S$ be a projective flat morphism between Noetherian schemes with geometrically integral fibers. Then the functor $\underline{\operatorname{Pic}}_{(X/S)(\acute{e}t)}$ is represented by a group S-scheme $\operatorname{Pic}_{X/S}$ which is separated, locally of finite type over S. Moreover, $\operatorname{Pic}_{X/S}$ is a disjoint union of open quasi-projective subschemes over S.

Proof. See [G62V, Thm. 3.1, p.148] or [Kle05, Thm. 9.4.8, p.263].

The group scheme $\operatorname{Pic}_{X/S}$ is called the *Picard scheme* of X over S.

Theorem 2.2. Let $f: X \to S$ be a proper morphism between schemes. Assume S is a spectrum of a field k, and X is complete. Then the functor $\underline{\operatorname{Pic}}_{(X/k)(fppf)}$ is represented by a group scheme $\operatorname{Pic}_{X/k}$ which is separated, locally of finite type over k. Moreover $\operatorname{Pic}_{X/k}$ is a disjoint union of open quasi-projective subschemes.

Proof. See [Kle05, Thm. 9.4.18.3, p.276]. The heart is [Mur64, Thm. 2, p.42]. \Box

• Let G be a commutative group scheme over a field k, i.e. separated of locally finite type over k. Then G is separated. We denote by G^0 the connected component of the identity of G. Then G^0 is a commutative group k-scheme of finite type. We denote by G_{red} the reduced scheme associated with G.

Theorem 2.3. Let X be a proper scheme over a field k.

- (i) $\operatorname{Pic}_{X/k}^{0}$ is a group scheme which is a separated of finite type over k.
- (ii) If X is geometrically normal, then $\operatorname{Pic}_{X/k}^{0}$ is proper. So $(\operatorname{Pic}_{X/k}^{0})_{red}$ is an abelian variety, and is called the *Picard variety* of X.
- (iii) If $h^2(\mathcal{O}_X) = 0$, then $\operatorname{Pic}_{X/k}^0$ is smooth of dim $h^1(\mathcal{O}_X)$, so $\operatorname{Pic}_{X/k}^0 = (\operatorname{Pic}_{X/k}^0)_{red}$. If also X is geometrically-normal, $\operatorname{Pic}_{X/k}^0$ is an abelian variety of dim $h^1(\mathcal{O}_X)$.

Proof. (i) follows from Theorem 2.2 and the above argument.

(ii) See [G62VI, Thm. 2.1 (ii) p.231, Coro.3.2. p.236] or [Kle05, Rem. 9.5.6, p.277].
(iii) The smoothness is [G62VI, Prop. 2.10 (ii), p.236] or [Kle05, Prop. 9.5.19, p.285].
The dimension is [G62VI, Prop. 2.10 (iii), p.236] or [Kle05, Coro. 9.5.13, p.283].

 $|^r C$ an its irredu

• Let C be a proper scheme over a field k of dimension 1, $C = \bigsqcup_{i=1}^{r} C_i$ an its irreducible decomposition, and m_i the multiplicity of C_i . The *total degree* map of C is defined by

deg :
$$\operatorname{Pic}(C) \to \mathbb{Z}$$
; $\mathcal{L} \mapsto \chi(\mathcal{L}) - \chi(\mathcal{O}_C)$.

We set $\operatorname{Pic}^{0}(C) = \operatorname{Ker}(\operatorname{deg})$. The Jacobian variety of C is defined by

$$\operatorname{Jac}(C) := \operatorname{Pic}_{C/k}^0$$
.

Since $h^2(\mathcal{O}_C) = 0$, $\operatorname{Jac}(C)$ is smooth of dimension $h^1(\mathcal{O}_C)$ by Theorem 2.3 (iii). • Assume $k = \overline{k}$. Let $X \in \mathcal{V}(k)$. Then $\operatorname{Pic}(X) = (\operatorname{Pic}_{X/k})_{red}(k)$. The group

 $NS(X) := (Pic_{X/k})_{red}(k)/(Pic_{X/k}^0)_{red}(k)$ (resp. Num(X) := NS(X)/Torsion)

(*) DAIKI KAWABE

is called the Neron-Severi group of X (resp. the Picard lattice of X). Then

Proposition 2.4. ([Kle05, Coro. 9.6.17, p.298]). NS(X) is a finitely-generated abelian group. Its rank is called the *Picard number* of X and is denoted by $\rho(X)$.

• Let X be a geometrically-integral variety over a field k having a point $p_0 \in X(k)$. By [Ser60], there are an abelian k-variety $Alb_{X/k}$ and a k-morphism $alb_X : X \to Alb_{X/k}$ such that:

- (i) $alb_X(p_0) = 0;$
- (ii) for every k-morphism $g: X \to A$ of X into an abelian variety A, there is an unique k-homomorphism $g_*: alb_X \to A$ such that $g = g_* \circ alb_X$.

We call $Alb_{X/k}$ the Albanese variety of X and call alb_X the Albanese morphism of X. If X is smooth projective, $Alb_{X/k}$ is the dual abelian variety of $(Pic_{X/k}^0)_{red}$.

• Assume $k = \overline{k}$. Let $X \in \mathcal{V}(k)$ be an integral variety. Fix a point $p_0 \in X(k)$.

Let $\operatorname{CH}_0(X)^0_{\mathbb{Z}}$ be the Chow group of 0-cycles of degree 0 on X with \mathbb{Z} -coefficients. Then, there is a surjective homomorphism

$$a_X : \operatorname{CH}_0(X)^0_{\mathbb{Z}} \to \operatorname{Alb}_{X/k}(k) \; ; \; \sum_i n_i[p_i] \mapsto \sum_i n_i[\operatorname{alb}_X(p_i)].$$

This map is called the *Albanese map* of X. Moreover, its kernel is called the *Albanese Kernel* of X, and is denoted by T(X).

• The cohomological Brauer group of a scheme X is defined by $\operatorname{Br}(X) := H^2_{\acute{e}t}(X, \mathbb{G}_m)$. If K is a field, we set $\operatorname{Br}(K) := \operatorname{Br}(\operatorname{Spec}(K))$.

Proposition 2.5. Let $f : X \to S$ be a separated morphism of finite type between locally Noetherian schemes. Assume $f_*\mathcal{O}_X \cong \mathcal{O}_S$ holds universally, i.e. $f_{T*}\mathcal{O}_{X_T} \cong \mathcal{O}_T$ for any S-scheme T. Then, there is an exact sequence

$$0 \to \operatorname{Pic}(X_T)/\operatorname{Pic}(T) \xrightarrow{\alpha} \underline{\operatorname{Pic}}_{(X/S)(\acute{et})}(T) \xrightarrow{\delta} \operatorname{Br}(T)$$

for any S-scheme T. The map α is bijective if f_T has a section or if Br(T) = 0.

Proof. Let us consider the Leray spectral sequence $E_2^{p,q} = H^p(T, R^q f_{T*}\mathbb{G}_m) \Rightarrow H^{p+q}(X_T, \mathbb{G}_m)$. Then the exact sequence of terms of low degree is:

$$0 \to H^1(T, f_{T*}\mathbb{G}_m) \to H^1(X_T, \mathbb{G}_m) \to H^0(T, R^1 f_{T*}\mathbb{G}_m) \to H^2(T, f_{T*}\mathbb{G}_m) \to H^2(X_T, \mathbb{G}_m).$$

Since $f_*\mathcal{O}_X \cong \mathcal{O}_S$ holds universally, the above exact sequence becomes

$$0 \to \operatorname{Pic}(T) \to \operatorname{Pic}(X_T) \to \underline{\operatorname{Pic}}_{(X/S)(\acute{et})}(T) \to \operatorname{Br}(T) \to \operatorname{Br}(X_T).$$

Thus the assertion follows.

3. Correspondences

In this section, we recall and prove some facts about correspondences.

3.1. Correspondences over a field. Let k be a field and $X, Y \in \mathcal{V}(k)$.

Definition 3.1. A *correspondence* from X to Y is an element of $CH(X \times Y)$. For simplicity, we write $\alpha \in CH(X \times Y)$ as $\alpha : X \vdash Y$.

If $\alpha : X \vdash Y$, $\beta : Y \vdash Z$, the composition $\beta \circ \alpha : X \vdash Z$ is defined by

$$\beta \circ \alpha := p_{XZ*}(p_{XY}^*(\alpha) \cdot p_{YZ}^*(\beta)),$$

Here p_{XY} , p_{YZ} , p_{XZ} denote the projections from $X \times Y \times Z$ to $X \times Y$, $Y \times Z$, $X \times Z$.

For $\alpha: X \vdash Y$, define a homomorphism $\alpha_* : \operatorname{CH}(X) \to \operatorname{CH}(Y)$ by $\alpha_*(a) = p_{Y*}^{XY}(\alpha \cdot p_X^{XY*}(a))$, and a homomorphism $\alpha^* : \operatorname{CH}(Y) \to \operatorname{CH}(X)$ by $\alpha^*(b) = p_{X*}^{XY}(\alpha \cdot p_Y^{XY*}(b))$. A correspondence $\alpha: X \vdash Y$ has a transpose ${}^t\alpha: Y \vdash X$ defined by ${}^t\alpha = \tau_*(\alpha)$ where $\tau: X \times Y \to Y \times X$ reverses the factors. For any morphism $f: X \to Y$, we denote by

$$\Gamma_f: X \vdash Y$$

the graph of f. If $f = \delta_X : X \hookrightarrow X \times X$ is the diagonal embedding, set $\Delta_X := \Gamma_f$.

Lemma 3.2. (Lieberman's lemma). Let $\alpha : X \vdash Y$ and $\beta : X' \vdash Y'$. Let $f : X \to X'$ and $g : Y \to Y'$ be proper and flat morphisms. Then

$$(f \times g)_*(\alpha) = \Gamma_g \circ \alpha \circ {}^t\Gamma_f.$$

Proof. It follows from [Ful84, Prop. 16.1.1].

For any $X, T \in \mathcal{V}(k)$, let $X(T) := \operatorname{CH}(T \times X)$. For $\phi : X \vdash Y$, we define

$$\phi_T : X(T) \to Y(T) \; ; \; \alpha \mapsto \phi \circ \alpha.$$

Theorem 3.3. (Manin's identity principle) Let $\phi, \psi : X \vdash Y$. Then

(i) $\phi = \psi$ \iff (ii) $\phi_T = \psi_T$ for all $T \in \mathcal{V}(k)$ \iff (iii) $\phi_X = \psi_X$.

Proof. (i) \Rightarrow (ii) \Rightarrow (iii) are trivial. (iii) \Rightarrow (i) follows from taking $\alpha = \Delta_X$.

Remark 3.4. The Manin identity principle induces

$$\Gamma_f = \Gamma_g \text{ in } \operatorname{CH}(X \times Y) \iff (\operatorname{id}_T \times f)_* = (\operatorname{id}_T \times g)_* : \operatorname{CH}(T \times X) \to \operatorname{CH}(T \times Y) \ \forall T$$

Proposition 3.5. Let $X, Y \in \mathcal{V}(k)$. Let $\pi : X \to Y$ be a finite morphism.

- (i) Let d be the degree of π . Then $\Gamma_{\pi} \circ {}^t\Gamma_{\pi} = d \cdot \Delta_Y$ in $CH(Y \times Y)$.
- (ii) Let G be a finite group which acts freely on X, and let Y := X/G. Then ${}^t\Gamma_{\pi} \circ \Gamma_{\pi} = \sum_{\sigma \in G} \Gamma_{\sigma}$ in $\operatorname{CH}(X \times X)$.

Proof. The proof of (i) is similar to (ii), so it suffices to prove (ii). Let $CH(X)^G$ be the *G*-invariant subgroup. Then $CH(Y) \cong CH(X)^G$ by [Ful84, Exa. 1.7.6], so $\pi^*\pi_* = \sum_{\sigma \in G} \sigma_*$ in Aut(CH(X)), and hence ${}^t\Gamma_{\pi} \circ \Gamma_{\pi} = \sum_{\sigma \in G} \Gamma_{\sigma}$ by Remark 3.4. \Box

3.2. Relative correspondences and base changes.


To extend $h(X_{\eta}) \cong h(J_{\eta})$ to $t_2(X) \cong t_2(J)$, we prove some facts about relative correspondences. This subsection is based on [CH00] and [MNP13, Ch. 8].

Let B be a quasi-projective variety over a field k. Now, we explain some concepts. • Let $\mathcal{V}(B)$ be the category whose objects are pairs (X, f) with X a smooth quasiprojective k-variety and $f: X \to B$ a projective morphism.

- A morphism from (X, f) to (Y, g) is a morphism $h: X \to Y$ such that $g \circ h = f$.
- Let $(X, f), (Y, g) \in \mathcal{V}(B)$. Assume that Y is equidimensional. Set

 $\operatorname{Corr}_B^r(X,Y) := \operatorname{CH}_{\dim(Y)-r}(X \times_B Y) \text{ and } \operatorname{Corr}_B(X,Y) := \oplus_r \operatorname{Corr}_B^r(X,Y).$

• There are defined for Cartesian squares



If *i* is regular embedding of codimension *d*, the upper map induces $i^! : \operatorname{CH}_k(Y) \to \operatorname{CH}_{k-d}(X)$. We apply this construction using the following diagram with right-hand side Cartesian squares

$$\begin{array}{cccc} X \times_B Y & \xleftarrow{p_{XZ}} & X \times_B Y \times_B Z & \xrightarrow{\delta'} & (X \times_B Y) \times_k (Y \times_B Z) \\ & & & \downarrow & & \downarrow \\ & & & & \downarrow \\ & & & & Y & \xrightarrow{\delta} & & Y \times_k Y. \end{array}$$

Since Y is smooth, δ is a regular embedding and so the refined Gysin homomorphism $\delta^!$ is well-defined. For $\Gamma_1 \in \operatorname{Corr}_B^r(X, Y)$ and $\Gamma_2 \in \operatorname{Corr}_B^s(Y, Z)$, we define

$$\Gamma_2 \circ_B \Gamma_1 := (p_{XZ})_* ((\delta')! (\Gamma_1 \times_k \Gamma_2)) \in \operatorname{Corr}_B^{r+s}(X, Z).$$

The following two lemmas say that the composition of relative correspondences is compatible with some base change.

Lemma 3.6. Let $X, Y, Z \in \mathcal{V}(B)$. Let $i : U \hookrightarrow B$ be an open immersion. Let $i_{XY} : X \times_B Y \times_B U \to X \times_B Y$ be the projection, and similarly for i_{VW} and i_{UW} . Let $\Gamma_1 \in \operatorname{Corr}_B(X, Y)$ and $\Gamma_2 \in \operatorname{Corr}_B(Y, Z)$. Then in $\operatorname{Corr}_U(X, Z)$

 $((i_{YZ})^*\Gamma_2) \circ_U ((i_{XY})^*\Gamma_1) = (i_{XZ})^*(\Gamma_2 \circ_B \Gamma_1).$

Proof. Since i is an open immersion, there are the following Cartesian diagrams

where the morphisms i_{XZ} , p, q are open immersion. Then in $CH(X \times_B Y \times_B Z \times_B U)$

$$(\delta'')^! (i_{XY} \times_k i_{YZ})^* (\Gamma_1 \times_k \Gamma_2) = (\delta'')^! p^* (\Gamma_1 \times_k \Gamma_2) = q^* (\delta')^! (\Gamma_1 \times_k \Gamma_2)$$
(1)

Apply to $(p'_{XZ})_*$ to this formula to obtain:

$$\begin{aligned} ((i_{YZ})^*\Gamma_2) \circ_U ((i_{XY})^*\Gamma_1) &= (p'_{XZ})_* (\delta'')! ((i_{YZ})^*\Gamma_2 \times_k (i_{XY})^*\Gamma_1) \\ &= (p'_{XZ})_* (\delta'')! (i_{YZ} \times_k i_{XY})^* (\Gamma_2 \times_k \Gamma_1) \\ &= (p'_{XZ})_* q^* (\delta')! (\Gamma_1 \times_k \Gamma_2) \qquad \text{by (1)} \\ &= (i_{XZ})^* (p_{XZ})_* (\delta')! (\Gamma_1 \times_k \Gamma_2) \qquad \text{by base change theorem} \\ &= (i_{XZ})^* (\Gamma_1 \circ_B \Gamma_2). \end{aligned}$$

8

Lemma 3.7. ([MNP13, Lem. 8.1.6, p.108]). Let $X, Y, Z \in \mathcal{V}(B)$. Let $t : B \to B'$ be a k-morphism. Let $j_{XY} : X \times_B Y \to X \times_{B'} Y$ be the canonical morphism, and similarly for j_{XY} and j_{XZ} . Let $\Gamma_1 \in \operatorname{Corr}_B(X, Y), \Gamma_2 \in \operatorname{Corr}_B(Y, Z)$. Then in $\operatorname{Corr}_{B'}(X, Z)$

$$((j_{YW})_*\Gamma_2) \circ_{B'} ((j_{XY})_*\Gamma_1) = (j_{XZ})_*(\Gamma_2 \circ_B \Gamma_1).$$

Proof. Let us consider the following commutative diagrams

The remainder is similar to the proof of Lemma 3.6.

4. Chow motives

In this section, we review the theory of Chow motives.

4.1. The category of Chow motives. Let $\operatorname{CH}\mathcal{M}(k) = \operatorname{CH}\mathcal{M}(k, \mathbb{Q})$ be the category of Chow motives over a field k with \mathbb{Q} -coefficients. Objects in $\operatorname{CH}\mathcal{M}(k)$ are given by triples (X, p, m) where $X \in \mathcal{V}(k), p \in \operatorname{Corr}^0(X, X)$ is a projector (i.e. $p \circ p = p$), and $m \in \mathbb{Z}$. Morphisms are in $\operatorname{CH}\mathcal{M}(k)$ given by

$$\operatorname{Hom}_{\operatorname{CH}\mathcal{M}(k)}((X, p, m), (Y, p, n)) = q \circ \operatorname{Corr}^{n-m}(X, Y) \circ p.$$

Let $M = (X, p, m), N = (Y, q, n) \in CH\mathcal{M}(k)$. One can define a motive $M \otimes N := (X \times Y, \pi_X^* p \cdot \pi_Y^* q, m+n)$ where $\pi_X : (X \times Y) \times (X \times Y) \to X \times X$ be the projection, and similar for π_Y . Also, one can define $M \oplus N$. For simplicity, we give only the definition in case m = n. Then $M \oplus N := (X \sqcup Y, p \oplus q, m)$, and refer to [Kim05, Def. 2.9 (ii), p.178] for the general case.

We denote by $h(-): \mathcal{V}(k)^{\mathrm{op}} \to \mathrm{CH}\mathcal{M}(k)$ the contravariant functor which associates to any $X \in \mathcal{V}(k)^{\mathrm{op}}$ its Chow motive

$$h(X) = (X, \Delta_X, 0),$$

where $\Delta_X \in \operatorname{Corr}^0(X, X)$ is the diagonal, and to a morphism $f: X \to Y$ the correspondence $h(f) = {}^t\Gamma_f \in \operatorname{Corr}^0(Y, X)$. For $X, Y \in \mathcal{V}(k)$, one has $h(X \times Y) = h(X) \otimes h(Y)$.

Let $1 = (\operatorname{Spec}(k), \Delta_{\operatorname{Spec}(k)}, 0)$ be the unit motive and $\mathbb{L} = (\operatorname{Spec}(k), \Delta_{\operatorname{Spec}(k)}, -1)$ the Lefschetz motive. For an non-negative integer n, we let $\mathbb{L}^{\oplus n} := \mathbb{L} \oplus \cdots \oplus \mathbb{L}$ (*n*-times). Let H^* be a fixed Weil-cohomology theory. For $M = (X, p, m) \in \operatorname{CH}\mathcal{M}(k)$, one

define $\operatorname{CH}^{i}(M) := p_* \operatorname{CH}^{i+m}(X)$ and $H^i(M) := p_* H^{i+2m}(X)$.

 $\operatorname{CH}\mathcal{M}(k)$ is *pseudo-abelian*, i.e. every projector $f \in \operatorname{End}(M)$ has an image, and the canonical map $\operatorname{Im}(f) \oplus \operatorname{Im}(\operatorname{id} - f) \to M$ is an isomorphism. For $M = (X, p, m) \in \operatorname{CH}\mathcal{M}(k)$ and $f = p \circ f \circ p \in \operatorname{End}(M)$, one has $M = (X, p \circ f \circ p, m) \oplus (X, p - p \circ f \circ p, m)$.

4.2. Chow-Künneth decompositions.

Let k be an algebraically closed field and $X \in \mathcal{V}(k)$ a variety of dimension d.

Definition 4.1. ([Mur93]) We say that X admits a Chow-Künneth decomposition (CK-decomposition for short) if there exist $\pi_i(X) \in CH_d(X \times X)$ such that:

(i)
$$\Delta_X = \sum_{i=0}^{2d} \pi_i(X) \text{ in } \operatorname{CH}_d(X \times X)$$

(ii)
$$\pi_i(X) \circ \pi_j(X) = \begin{cases} \pi_i(X) & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$$

(iii) $cl_{X\times X}^d(\pi_i(X))$ is the (i, 2d-i)-th component of Δ_X in $H^{2d}(X\times X)$.

If π_i exist, we put $h_i(X) := (X, \pi_i(X), 0)$, and have $h(X) \cong \bigoplus_{i=0}^{2d} h_i(X)$ in $CH\mathcal{M}(k, \mathbb{Q})$. The Künneth components are unique in $H(X \times X)$ but $\pi_i(X)$ are not unique in $CH_d(X \times X)$ in general. For a surface, we will always use:

Proposition 4.2. ([Mur90]). Let $S \in \mathcal{V}(k)$ be a surface. Let $P \in S$ be a closed point. Then S admits a CK-decomposition $h(S) \cong \bigoplus_{i=0}^{4} h_i(S)$ such that:

- (i) $\pi_0 := (1/\deg(P))[S \times P]$ and $\pi_4 = (1/\deg(P))[P \times S];$
- (ii) There is a curve $C \subset S$ s.t. π_1 (resp. π_3) is supported on $S \times C$ (resp. $C \times S$);
- (iii) $\pi_2 := \Delta_S \pi_0 \pi_1 \pi_3 \pi_4;$
- (iv) $\pi_i = {}^t \pi_{4-i}$ for $0 \le i \le 4$.

Moreover, these projectors induce isomorphisms:

(i') $h_0(S) \cong 1$ and $h_4(S) \cong \mathbb{L} \otimes \mathbb{L}$;

(ii') $h_1(S) \cong h_1((\operatorname{Pic}^0_{S/k})_{red})$ and $h_3(S) \cong h_{2\dim(\operatorname{Alb}_{S/k})-1}(\operatorname{Alb}_{S/k}) \otimes \mathbb{L}^{2-\dim(\operatorname{Alb}_{S/k})}$.

Proposition 4.3. ([KMP07]). Let $S \in \mathcal{V}(k)$ be a surface. Let $D_i \in \mathrm{NS}(S)_{\mathbb{Q}}$ be an orthogonal basis. Then there is a unique splitting in $\mathrm{CH}_2(S \times S)$

$$\pi_2 = \pi_2^{alg} + \pi_2^{ti}$$

such that $\pi_2^{alg} := \sum_{i=1}^{\rho} 1/(D_i \cdot D_i)[D_i \times D_i]$, where $(D_i \cdot D_i)$ the intersection number. Moreover, the above splitting induces a decomposition in $CH\mathcal{M}(k, \mathbb{Q})$

$$h_2(S) \cong h_2^{alg}(S) \oplus t_2(S)$$

such that $h_2^{alg}(S) := (S, \pi_2^{alg}, 0) \cong \mathbb{L}^{\oplus \rho(S)}$ and $t_2(S) := (S, \pi_2^{tr}, 0)$. Finally, $\operatorname{CH}^*(h_2^{alg}(S)) = \operatorname{NS}(S)_{\mathbb{Q}}, \quad \operatorname{CH}^*(t_2(S)) = T(S)_{\mathbb{Q}}, \quad H^*(h_2^{alg}(S)) = H^2(S)_{alg}, \quad H^2(t_2(S)) = H^2(S)_{tr}.$

The motive $t_2(S)$ is called the *transcendental motive* of S.

4.3. Homomorphisms between transcendental motives.

Here we prove some results about homomorphisms between transcendental motives. Let k be an algebraically closed field. Let $X, Y \in \mathcal{V}(k)$ be surfaces. We let $\operatorname{CH}_2(X \times Y)_{\equiv}$: the subgroup of $\operatorname{CH}_2(X \times Y)$ generated by the classes supported on

subvarieties of the form $X \times N$ or $M \times Y$, with M a closed subvariety of X of

dimension < 2 and N a closed subvariety of Y of dimension < 2.

We define a homomorphism

$$\Phi_{X,Y} : \mathrm{CH}_2(X \times Y) \to \mathrm{Hom}_{\mathrm{CH}\mathcal{M}(k)}(t_2(X), t_2(Y))$$
$$\alpha \mapsto \pi_2^{tr}(Y) \circ \alpha \circ \pi_2^{tr}(X).$$

Theorem 4.4. ([KMP07, Thm. 7.4.3, p.165]). There is an isomorphism of groups $\operatorname{CH}_2(X \times Y)/\operatorname{CH}_2(X \times Y)_{\equiv} \cong \operatorname{Hom}_{\operatorname{CH}\mathcal{M}(k)}(t_2(X), t_2(Y)).$ For this subsection, we need the following lemma:

Lemma 4.5. Let $\alpha \in CH_2(X \times Y)$ and $\gamma \in CH_2(Y \times X)_{\equiv}$. Then

(i) $\gamma \circ \alpha \in \operatorname{CH}_2(X \times X)_{\equiv}$ and (ii) $\alpha \circ \gamma \in \operatorname{CH}_2(Y \times Y)_{\equiv}$.

Proof. The proof of (ii) is similar to (i), so it suffices to show (i). Without loss of generality, we may assume γ is irreducible and supported on $Y \times C$ with dim $(C) \leq 1$.

First, we assume dim(C) = 0. Let $p \in X$ be the closed point. For $\gamma = [Y \times p]$, then

$$\gamma \circ \alpha = [Y \times p] \circ \alpha = p_{YY*}^{YXY}(\alpha \times Y \cdot Y \times X \times p) = p_{YY*}^{YXY}(\alpha \times p) = [p_{Y*}^{YX}(\alpha) \times p].$$

Thus $\gamma \circ \alpha \in \operatorname{CH}_2(X \times X)_{\equiv}$.

Next, we assume $\dim(C) = 1$. Since γ is supported on $Y \times C$, there are a smooth irreducible curve C and a closed embedding $\iota : C \hookrightarrow X$ such that $\gamma = \Gamma_{\iota} \circ D$ in $\operatorname{CH}_2(Y \times X)$, where $\Gamma_{\iota} \in \operatorname{CH}_1(C \times X)$ is the graph of ι and $D \in \operatorname{CH}_2(Y \times C)$. Since the support of the second projection of Γ_{ι} has dimension ≤ 1 , the support of the second projection of $\gamma \circ \alpha$ has dimension ≤ 1 , and hence $\gamma \circ \alpha \in \operatorname{CH}_2(X \times X)_{\equiv}$.

Lemma 4.5 gives the following facts. The first one is the functorial relation for $\Phi_{X,Y}$:

Proposition 4.6. ([Ped12, p.62]). For surfaces $X, Y, Z \in \mathcal{V}(k)$,

$$\Psi_{Y,Z}(\beta) \circ \Psi_{X,Y}(\alpha) = \Psi_{X,Z}(\beta \circ \alpha) \text{ in } \operatorname{Hom}_{\operatorname{CH}\mathcal{M}(k)}(t_2(X), t_2(Z)).$$

Proof. For example, see [Kaw22].

Proposition 4.7. There is a bilinear homomorphism

$$\circ: \frac{\operatorname{CH}_2(X \times Y)}{\operatorname{CH}_2(X \times Y)_{\equiv}} \times \frac{\operatorname{CH}_2(Y \times Z)}{\operatorname{CH}_2(Y \times Z)_{\equiv}} \to \frac{\operatorname{CH}_2(X \times Z)}{\operatorname{CH}_2(X \times Z)_{\equiv}}$$
$$([\alpha], [\beta]) \mapsto [\beta] \circ [\alpha] := [\beta \circ \alpha].$$

Proof. By Lemma 4.5, the composition $[\beta] \circ [\alpha] := [\beta \circ \alpha] \in CH_2(X \times Z)/CH_2(X \times Z)_{\equiv}$ is well-defined. Thus, the assertion follows.

The main proposition of this subsection is:

Proposition 4.8. Let $X, Y \in \mathcal{V}(k)$ be surfaces. Assume there are elements $[\alpha] \in CH_2(X \times Y)/CH_2(X \times Y)_{\equiv}$ and $[\beta] \in CH_2(Y \times X)/CH_2(Y \times X)_{\equiv}$ such that:

(i) $[\Delta_Y] = [\alpha] \circ [\beta]$ in $\operatorname{CH}_2(Y \times Y)/\operatorname{CH}_2(Y \times Y)_{\equiv};$ (ii) $[\Delta_X] = [\beta] \circ [\alpha]$ in $\operatorname{CH}_2(X \times X)/\operatorname{CH}_2(X \times X)_{\equiv}.$

Then, there is an isomorphism $t_2(X) \cong t_2(Y)$ in $\operatorname{CH}\mathcal{M}(k,\mathbb{Q})$.

Proof. Assume (i). By Proposition 4.7, $[\Delta_Y] = [\alpha \circ \beta]$ in $\operatorname{CH}_2(Y \times Y)/\operatorname{CH}_2(Y \times Y)_{\equiv}$. By Theorem 4.4, in $\operatorname{Hom}_{\operatorname{CH}\mathcal{M}(k)}(t_2(Y), t_2(Y))$,

$$\Phi_{Y,Y}(\Delta_Y) = \Phi_{Y,Y}(\alpha \circ \beta) \tag{2}$$

Here, consider the two morphisms

 $\Phi_{X,Y}(\alpha) \in \operatorname{Hom}_{\operatorname{CH}\mathcal{M}(k)}(t_2(X), t_2(Y)) \quad \text{and} \quad \Phi_{Y,X}(\beta) \in \operatorname{Hom}_{\operatorname{CH}\mathcal{M}(k)}(t_2(Y), t_2(X)).$

In $\operatorname{Hom}_{\operatorname{CH}\mathcal{M}(k)}(t_2(Y), t_2(Y)),$

$$\Phi_{X,Y}(\alpha) \circ \Phi_{Y,X}(\beta) = \Phi_{Y,Y}(\alpha \circ \beta) \qquad \text{by Proposition 4.6} = \Phi_{Y,Y}(\Delta_Y) \qquad \text{by (2)} = \pi_2^{tr}(Y) \qquad \text{by } \pi_2^{tr}(Y) \circ \pi_2^{tr}(Y) = \pi_2^{tr}(Y) = \mathrm{id}_{t_2(Y)}.$$

Similarly, by (ii), we get $\Phi_{Y,X}(\beta) \circ \Phi_{X,Y}(\alpha) = \mathrm{id}_{t_2(X)}$ in $\mathrm{Hom}_{\mathrm{CH}\mathcal{M}(k)}(t_2(X), t_2(X))$. Therefore, we get $t_2(X) \cong t_2(Y)$ in $\mathrm{CH}\mathcal{M}(k, \mathbb{Q})$.

5. Torsors over group varieties

To prove Theorem 1.6, we prove several facts about torsors for commutative group varieties. This section is based on [LT58], [Sil86, X], [Mil80, III, §4]. In this section, let K be a field, \overline{K} an algebraic closure of K, and K^s a separable closure of K.

5.1. Torsors. Let A be a commutative group K-variety and T a smooth, not necessarily projective, geometrically-integral, K-variety. Assume A acts by the K-morphism

 $\mu: T \times A \to T.$

Definition 5.1. We say that T (or (T, μ)) is a *torsor* for A/K if

 $T\times A \to T\times T \ ; \ (p,P)\mapsto (p,\mu(p,P))$

is a K-isomorphism. Then the action of the group $A(\overline{K})$ on $T(\overline{K})$ is simply transitive.

For simplicity, write $\mu(p, P)$ as p + P. Here we define a subtraction map on T by

$$\nu: T \times T \to A,$$

 $\nu(p,q) = (\text{the unique } P \in A(\overline{K}) \text{ satisfying } \mu(p,P) = q).$

Then ν is a K-morphism of varieties¹. For simplicity, write $\nu(p,q)$ as q-p.

Two torsors (T, μ) , (T', μ') over K for A/K are isomorphic as (torsors) if there is an isomorphism $\theta : T \to T'$ of K-varieties that is A-equivariant. In other words, $\theta \circ \mu = \mu' \circ \theta \times id_A$. Note that any torsor is trivial after some base extension:

Proposition 5.2. Let T/K be a torsor for A/K. Let M/K be a field extension in K^s with $T(M) \neq \emptyset$. Choose $p_0 \in T(M)$ and define the *M*-morphism

 $\phi^{-1} = \phi_{p_0}^{-1} : A_M \to T_M ; P \mapsto p_0 + P$

Then ϕ^{-1} is an isomorphism of torsors and its inverse ϕ is

$$\phi: T_M \to A_M \; ; \; p \mapsto p - p_0$$

In particular, a torsor T is K-isomorphic to A if and only if $T(K) \neq \emptyset$.

Proof. Since T is a smooth variety, $T(K^s) \neq \emptyset$ by [Liu02, Prop. 2.20, p.93], so such M exists. Since the action μ is defined over K, for all $\sigma \in \text{Gal}(K^s/M), P \in A(K^s)$,

$$\phi^{-1}(P)^{\sigma} = (p_0 + P)^{\sigma} = p_0^{\sigma} + P^{\sigma} = p_0 + P^{\sigma} = \phi^{-1}(P^{\sigma}).$$

This shows that ϕ^{-1} is defined over M. Since $T \times A \cong T \times T$, the induced map $\phi^{-1}(\overline{K}) : A_M(\overline{K}) = A(\overline{K}) \to T(\overline{K}) = T_M(\overline{K})$ is an isomorphism of rational points, so ϕ^{-1} is an isomorphism of M-varieties. For any $P, Q \in A_M(\overline{K})$, we have

$$\phi^{-1}(P+Q) = p_0 + (P+Q) = (p_0 + P) + Q = \phi^{-1}(P) + Q.$$

¹because ν can be described as the composition of morphisms: $\nu = -_E \circ \phi \times \phi : T \times T \to E \times E \to E$

This shows that ϕ^{-1} is A-equivariant. Thus ϕ^{-1} is an isomorphism of torsors. Hence the rest of the assertions follow.

For a group G, let G_{tor} denote the torsion subgroup. To prove Thm. 1.6, we need:

Proposition 5.3. Let T/K be a torsor for a commutative group variety A/K. Then there are a finite Galois extension L/K and a point $p_0 \in T(L)$ such that

$$p_0 - p_0^{\sigma} \in A(L)_{tor}$$
 for all $\sigma \in \operatorname{Gal}(L/K)$

Proof. Fix a point $p \in T(K^s)$. Let n be an order of the element

$$\{a: \sigma \mapsto p - p^{\sigma}\} \in H^1(\operatorname{Gal}(K^s/K), A(K^s)).$$

The Kummer sequence $0 \to A(K^s)[n] \to A(K^s) \xrightarrow{n} A(K^s) \to 0$ gives an exact sequence $0 \to A(K^s)/nA(K^s) \to H^1(\operatorname{Gal}(K^s/K), A(K^s)[n]) \to H^1(\operatorname{Gal}(K^s/K), A(K^s))[n] \to 0.$ Then, there is an element $\{b\} \in H^1(\operatorname{Gal}(K^s/K), A(K^s)[n])$ such that $\{b\} = \{a\}$ in $H^1(\operatorname{Gal}(K^s/K), A(K^s)).$ So there is a point $P \in A(K^s)$ such that

$$b(\sigma) = a(\sigma) + P - P^{\sigma} \in A(K^s)$$
 for all $\sigma \in \operatorname{Gal}(K^s/K)$.

Namely,

$$(p - p^{\sigma}) + P - P^{\sigma} = b(\sigma) \in A(K^s)[n]$$
 for all $\sigma \in \operatorname{Gal}(K^s/K)$.
Set $p_0 := p + P \in T(K^s)$. Then, for all $\sigma \in \operatorname{Gal}(K^s/K)$,

 $p_0 - p_0^{\sigma} = (p+P) - (p+P)^{\sigma} = (p+P) - (p^{\sigma} + P^{\sigma}) = (p-p^{\sigma}) + P - P^{\sigma} = b(\sigma) \in A(K^s)[n].$ Since $p_0 \in C(K^s)$, there is a finite Galois extension L/K such that $p_0 \in T(L)$. Since $A(L)_{tor} = A(K^s)_{tor} \cap A(L)$, we get $p_0 - p_0^{\sigma} \in A(L)_{tor}$ for all $\sigma \in \text{Gal}(L/K)$. \Box

5.2. Genus one curves. In this paper, we use the following terminology:

Definition 5.4. Let C be a projective, *geometrically-integral*, curve over a field K.

- (i) C is a genus 1 curve if $p_a(C) = \dim H^1(X, \mathcal{O}_C) = 1$.
- (ii) C is an *elliptic curve* if it is a *smooth* genus 1 curve with $C(K) \neq \emptyset$. In other words, C is an abelian K-variety of dimension 1.

If C is a smooth genus 1-curve, then Jac(C) is an elliptic curve by Theorem 2.3 (iii).

Proposition 5.5. ([Sch10, Prop. 6.1, p.54]). Let C be a genus 1 curve over a field K. Then C is Gorenstein. Moreover, $\omega_C \cong \mathcal{O}_C$.

For a K-variety $g: X \to \operatorname{Spec}(K)$, let $X^{\#} := \{ x \in X \mid g \text{ is smooth at } x \}.$

Proposition 5.6. Assume $K = \overline{K}$. Let C be a non-smooth, genus 1 curve over K.

- (i) C has exactly one singular point.
- (ii) char(K) = 2 or 3.

Proof. (i) Let $\mu : \tilde{C} \to C$ be the normalization. As in [Har77, IV. Exercise 1.8] or [Liu02, p304], we have an exact sequence of coherent sheaves on C:

$$0 \to \mathcal{O}_C \to \mu_*\mathcal{O}_{\tilde{C}} \to \bigoplus_{c \in C} \mathcal{O}'_{C,p} / \mathcal{O}_{C,p} \to 0$$

where $\mathcal{O}'_{C,p}$ is the integral closure of $\mathcal{O}_{C,p}$. Then we have $p_a(C) = p_a(\tilde{C}) + \sum_{p \in C} \delta(p)$ where $\delta(p) := \text{length}(\mathcal{O}'_{C,p}/\mathcal{O}_{C,p})$. Now, for a closed point $p \in C$, $\delta(p) = 0$ if and only if p is a smooth point. By assumption, $p_a(C) = 1$, so $p_a(\tilde{C}) = 0$ and $\sum_{p \in C} \delta(p) = 1$. Thus, C has exactly one singular point.

(ii) By [Tat52, Coro. 1, p.404], $\sum_{p \in C} \delta(p)$ is an integer multiple of $(\operatorname{char}(K) - 1)/2$. By (i), we have $\sum_{p \in C} \delta(p) = 1$, so get $\operatorname{char}(K) = 2$ or 3.

Lemma 5.7. (Abel). Let C be a genus 1 curve over a field K. Fix $p_0 \in C^{\#}(\overline{K})$. Then, there is a bijection of sets

$$C^{\#}(\overline{K}) \to \operatorname{Pic}^{0}(C_{\overline{K}}) \; ; \; p \mapsto [p] - [p_0]$$

Proof. It follows from the Riemann-Roch for the Gorenstein curve $C_{\overline{K}}$.

Proposition 5.8. For a genus 1 curve C over a field K, $C^{\#}$ is a torsor for Jac(C)/K.

Proof. Fix $p_0 \in C^{\#}(\overline{K})$. By [MI21, Coro. 1.8.1 (4), p.13], there is a K^s -morphism

$$\phi: C_{K^s}^{\#} \to \operatorname{Jac}(C)_{K^s}.$$

By Lemma 5.7 and [Mil86, Thm. 8.1, p.192], the induced map $\phi(\overline{K}) : C^{\#}(\overline{K}) \to \operatorname{Jac}(C_{\overline{K}})$ is an isomorphism of rational points, so ϕ is an isomorphism of group varieties.

Define a map

$$\mu: C^{\#} \times \text{Jac}(C) \to C^{\#} \;\; ; \;\; (p, P) \mapsto \phi^{-1}(\phi(p) + P) = p + \phi^{-1}(P).$$

Then μ is a group action of $\operatorname{Jac}(C)$ on $C^{\#}$ over K^s .

(i) First, we prove μ is defined over K. For all $p \in C^{\#}(K^s)$, $P = [q] - [p_0] \in \operatorname{Jac}(C)(K^s)$, $\sigma \in \operatorname{Gal}(K^s/K)$,

$$\mu(p, P)^{\sigma} = (p + \phi^{-1}([q] - [p_0]))^{\sigma} = p^{\sigma} + q^{\sigma} = p^{\sigma} + \phi^{-1}([q^{\sigma}] - [p_0])$$
$$= p^{\sigma} + \phi^{-1}([q^{\sigma}] - [p_0^{\sigma}]) = \mu(p^{\sigma}, P^{\sigma}).$$

This shows that μ is defined over K.

(ii) Next, we prove $C^{\#} \times \operatorname{Jac}(C) \to C^{\#} \times C^{\#}$; $(p, P) \mapsto (p, \phi^{-1}(\phi(p) + P))$ is a K-isomorphism. Indeed, for all $p, q \in C^{\#}(\overline{K})$, one has $\mu(p, P) = q$ if and only if $\phi^{-1}(\phi(p) + P) = q$. Thus, the only choice for P is $P = \phi(q) - \phi(p)$. By (i) and (ii), $C^{\#}$ is a torsor for $\operatorname{Jac}(C)$.

The following facts will be used for the study of relations between f and j.

Proposition 5.9. Let C be a regular genus 1 curve over a field K. Let E be a regular compactification of Jac(C), i.e. it is a projective regular K-curve containing as a dense open subset Jac(C).

- (i) There is a separable field extension M/K with $C_M \cong E_M$. In particular, $C_{\overline{K}} \cong E_{\overline{K}}$. If $C^{\#}(K) \neq \emptyset$, then $C \cong E$.
- (ii) E is also a regular genus 1 curve over K.
- (iii) There is an isomorphism $\operatorname{Jac}(C) \cong \operatorname{Jac}(E)$.

Proof. (i) By Proposition 5.8, $C^{\#}$ is a torsor for $\operatorname{Jac}(C)$, so there is a finite separable field extension M/K such that $C_M^{\#} \cong \operatorname{Jac}(C)_M$. So $C_M \sim_{birat} E_M$. By assumption, C is regular. Since M/K is separable, C_M is also regular. Similarly, E_M is regular. Since C_M, E_M are regular proper curves, $C_M \cong E_M$. By base change, $C_{\overline{K}} \cong E_{\overline{K}}$. If $C^{\#}(K) \neq \emptyset$, then $C^{\#} \cong \operatorname{Jac}(C)$ by Proposition 5.2. Since M = K, we have $C \cong E$. (ii) By assumption, $p_a(C) = 1$. By [Liu02, Def. 3.19, p.279], $p_a(C_{\overline{K}}) = 1$. By (i),

 $C_{\overline{K}} \cong E_{\overline{K}}$. By the same fact, $p_a(E) = p_a(E_{\overline{K}}) = 1$.

(iii) If $C_{\overline{K}}$ is regular, the assertion is clear. Assume that $C_{\overline{K}}$ is non-regular.

(iii-i) First, we prove $\operatorname{Jac}(C) \cong E^{\#}$. By the definition of $E^{\#}$, we get $\operatorname{Jac}(C) \subset E^{\#}$. By (i), $p_a(E) = p_a(C) = 1$. By Proposition 5.6 (i), we have

$$|E(\overline{K}) \setminus E^{\#}(\overline{K})| = |C(\overline{K}) \setminus C^{\#}(\overline{K})| = 1.$$
(3)

Here, |-| denote the order of -. As in (i), $C_{\overline{K}}^{\#} \cong \operatorname{Jac}(C)_{\overline{K}}$ and $C_{\overline{K}} \cong E_{\overline{K}}$, so

$$|E(\overline{K}) \setminus \operatorname{Jac}(C)(\overline{K})| = |C(\overline{K}) \setminus C^{\#}(\overline{K})| = 1.$$
(4)

By (3) and (4), we have

$$|E(\overline{K}) \setminus \operatorname{Jac}(C)(\overline{K})| = |E(\overline{K}) \setminus E^{\#}(\overline{K})| = 1,$$

and hence we get $E^{\#} \cong \operatorname{Jac}(C)$.

(iii-ii) Next, we prove $\operatorname{Jac}(E) \cong E^{\#}$. By Proposition 5.8, $E^{\#}$ is a torsor for $\operatorname{Jac}(E)$. Since $E^{\#}(K) \neq \emptyset$, $E^{\#}$ is the trivial torsor for $\operatorname{Jac}(E)$. So $E^{\#} \cong \operatorname{Jac}(E)$.

Combining (iii-i) and (iii-ii), we get $\operatorname{Jac}(C) \cong E^{\#} \cong \operatorname{Jac}(E)$.

To prove Theorem 1.6, we need the following:

Proposition 5.10. Let A be an abelian variety over a field K of dimension d and $P_n \in A(K)$ a torsion point of order $n \in \mathbb{Z}_{>0}$.

- (i) Let $t: A \to A$ be the translation by P_n . Then $\Gamma_t = \Delta_A$ in $CH_d(A \times A)$.
- (ii) Let T/K be a torsor for A with $T(K) \neq \emptyset$. Let $t: T \to T$; $p \mapsto p + P_n$ be the action by P_n . Then $\Gamma_t = \Delta_T$ in $CH_d(T \times T)$.

Proof. (i) This is the same proof for $K = \mathbb{C}$ as in [JY17, Lem. 2.1, p.573]. Let $n: A \to A$ be the multiplication by n. Set $n_A := (id_A \times n) : A \times A \to A \times A$, and $\operatorname{CH}^d_s(A \times A) := \{\xi \in \operatorname{CH}^d(A \times A) \mid n^*_A(\xi) = n^{2d-s}\xi \text{ for all } n\}$. By [DM91, Coro. 2.21], we have

$$\operatorname{CH}^{d}(A \times A) \cong \bigoplus_{s=p''}^{p'} \operatorname{CH}^{d}_{s}(A \times A).$$

This induces an isomorphism

$$n_A^* : \operatorname{CH}^d(A \times A) \to \operatorname{CH}^d(A \times A)$$
 (5)

By $n \circ t = n$ in Isom(A), we have $n_A^* \Gamma_t = n_A^* \Delta_A$, so get $\Gamma_t = \Delta_A$ by (5).

(ii) Since $T(K) \neq \emptyset$, T is an abelian K-variety. Let $n : T \to T$ be the multiplication by n. Regarding t as an isomorphism of trivial torsors for T, we have $n \circ t = n$ in Isom(T). By the same argument of (i), $\Gamma_t = \Delta_T$.

6. Chow motives of torsors under Abelian varieties

The purpose of this section is to prove the following:

Theorem 6.1. Let K be an arbitrary field. Let A be an abelian variety over K of dimension d. Let T be a torsor over K under A.² Then there is an isomorphism

$$h(T) \cong h(A)$$

in the category $CH\mathcal{M}(K,\mathbb{Q})$ of Chow motives.

 $^{^{2}}$ First, the author only proved Theorem 6.1 for the case of smooth genus 1 curves under the guidance of Prof. Hanamura. Thanks to the referee's suggestion, he proved the above theorem.

Proof of Theorem 6.1.

By definition, we have to prove: there are elements $a \in CH_d(T \times A)$, $b \in CH_d(A \times T)$ such that

$$\begin{cases} a \circ b = \Delta_A & \text{in } \operatorname{CH}_d(A \times A) \\ b \circ a = \Delta_T & \text{in } \operatorname{CH}_d(T \times T) \end{cases}$$

Step 1. Construct correspondences on the varieties A and T. By Proposition 5.3, there are a finite Galois extension L/K and a point $p_0 \in T(L)$ such that

$$p_0 - p_0^{\sigma} \in A(L)_{tor}$$
 for all $\sigma \in \operatorname{Gal}(L/K)$.

Let n be the degree of L/K and let $G := \operatorname{Gal}(L/K)$. By definition of a torsor, there is an isomorphism of L-varieties

$$\phi = \phi_{p_0} : T_L \to A_L.$$

Let $p: T_L \to T$ and $q: A_L \to A$ be the projections. Let

$$\Gamma_{\phi} \in \operatorname{CH}_1(T_L \times_L A_L), \quad \Gamma_p \in \operatorname{CH}_1(T_L \times_K T), \text{ and } \Gamma_q \in \operatorname{CH}_1(A_L \times_K A).$$

be the graph of ϕ , p, and q, respectively. We define

$$a := (1/n) \ \Gamma_q \circ \Gamma_\phi \circ {}^t\Gamma_p \in \operatorname{CH}_1(T \times A)$$
$$b := (1/n) \ \Gamma_p \circ {}^t\Gamma_\phi \circ {}^t\Gamma_q \in \operatorname{CH}_1(A \times T)$$

Step 2. Translations. To prove $a \circ b = \Delta_A$ and $b \circ a = \Delta_T$, we prove an elementary lemma. For any $\sigma \in G$, let $\sigma : \operatorname{Spec}(L) \to \operatorname{Spec}(L)$ denote the induced morphism.

For any $\sigma \in G$, we define

$$\phi^{\sigma} \circ \phi^{-1} := (\mathrm{id}_A \times_K \sigma) \circ \phi \circ (\mathrm{id}_T \times_K \sigma^{-1}) \circ \phi^{-1} \in \mathrm{Isom}(A_L/L)$$
$$(\phi^{-1})^{\sigma} \circ \phi := (\mathrm{id}_T \times_K \sigma) \circ \phi^{-1} \circ (\mathrm{id}_A \times_K \sigma^{-1}) \circ \phi \in \mathrm{Isom}(T_L/L)$$

For short, we denote by $p_0 - p_0^{\sigma} : A_L \to A_L$ the translation by $p_0 - p_0^{\sigma} \in A(L)_{tor}$ and denote by $p_0^{\sigma} - p_0 : T_L \to T_L$ the action by $p_0^{\sigma} - p_0 \in A(L)_{tor}$.

Lemma 6.2. For any $\sigma \in G$, we have

(i)
$$\phi^{\sigma} \circ \phi^{-1} = p_0 - p_0^{\sigma}$$
 in $\text{Isom}(A_L/L)$. (ii) $(\phi^{-1})^{\sigma} \circ \phi = p_0^{\sigma} - p_0$ in $\text{Isom}(T_L/L)$.

Proof. (i) It suffices to prove $(\phi^{\sigma} \circ \phi^{-1})(K^s) = (p_0 - p_0^{\sigma})(K^s)$ in Isom $(A(K^s))$. The morphisms ϕ and ϕ^{-1} induce isomorphisms of K^s -rational points

$$\phi(K^s): T(K^s) \to A(K^s) \quad ; \quad p \mapsto p - p_0$$

$$\phi^{-1}(K^s): A(K^s) \to T(K^s) \quad ; \quad P \mapsto p_0 + P$$

Thus, $\phi^{\sigma} \circ \phi^{-1}$ induces an isomorphism of K^{s} -rational points

 $A(K^{s}) \ni P \xrightarrow{\phi^{-1}(K^{s})} p_{0} + P \xrightarrow{\sigma^{-1}} p_{0}^{\sigma^{-1}} + P^{\sigma^{-1}} \xrightarrow{\phi(K^{s})} (p_{0}^{\sigma^{-1}} + P^{\sigma^{-1}}) - p_{0} \xrightarrow{\sigma} (p_{0} + P) - p_{0}^{\sigma} \in A(K^{s})$ Let $O \in A$ be the origin. We have in $A(K^{s})$

$$(p_0 + P) - p_0^{\sigma} = (p_0 + P) - (p_0^{\sigma} + O) = (p_0 - p_0^{\sigma}) + P - O = (p_0 - p_0^{\sigma}) + P.$$

Therefore, we get $\phi^{\sigma} \circ \phi^{-1} = p_0 - p_0^{\sigma}$ in Isom (A_L/L) .

(ii) Similarly, $(\phi^{-1})^{\sigma} \circ \phi$ induces an isomorphism of K^s -rational points

$$T(K^s) \ni p \stackrel{\phi(K^s)}{\mapsto} p - p_0 \stackrel{\sigma^{-1}}{\mapsto} p^{\sigma^{-1}} - p_0^{\sigma^{-1}} \stackrel{\phi^{-1}(K^s)}{\mapsto} p_0 + (p^{\sigma^{-1}} - p_0^{\sigma^{-1}}) \stackrel{\sigma}{\mapsto} p_0^{\sigma} + (p - p_0^{\sigma}) \in T(K^s)$$

As the action $T \times A \to T$ is equal to $\phi^{-1} \circ +_A \circ \phi \times \operatorname{id}_A$, we have in $T(K^s)$ $p_0^{\sigma} + (p - p_0) = p_0 + \phi(p) = \phi^{-1}(\phi(p_0^{\sigma}) + \phi(p)) = \phi^{-1}(\phi(p) + \phi(p_0^{\sigma})) = p + \phi(p_0^{\sigma}) = p + (p_0^{\sigma} - p_0).$ Therefore, we get $(\phi^{-1})^{\sigma} \circ \phi = p_0^{\sigma} - p_0$ in $\operatorname{Isom}(T_L/L)$.

Step 3. Calculate the correspondences on the varieties A and T. First, we prove $a \circ b = \Delta_A$. Let

$$A_L \times_L A_L \xrightarrow{q' \times q'} A \times_K A_L \xrightarrow{q \times q} A \times A.$$

$$\begin{split} & \text{In CH}_d(A \times A), \\ & a \circ b = (1/n^2) \ (\Gamma_q \circ \Gamma_\phi \circ {}^t\Gamma_p) \circ (\Gamma_p \circ {}^t\Gamma_\phi \circ {}^t\Gamma_q) & \text{by } a = \Gamma_q \circ \Gamma_\phi \circ {}^t\Gamma_p \text{ and } b = \Gamma_p \circ {}^t\Gamma_\phi \circ {}^t\Gamma_q \\ & = (1/n^2) \ \Gamma_q \circ \Gamma_\phi \circ \sum_{\sigma^{-1} \in G} \Gamma_{\sigma^{-1}} \circ {}^t\Gamma_\phi \circ {}^t\Gamma_q & \text{by Proposition 3.5 (ii)} \\ & = (1/n^2) \sum_{\sigma^{-1} \in G} \Gamma_q \circ \Gamma_\sigma \circ \Gamma_\phi \circ \Gamma_{\sigma^{-1}} \circ {}^t\Gamma_\phi \circ {}^t\Gamma_q & \text{by } q \circ \sigma = q \\ & = (1/n^2) \sum_{\sigma \in G} \Gamma_q \circ \Gamma_{\sigma \circ \phi \circ \sigma^{-1} \circ \phi^{-1}} \circ {}^t\Gamma_q & \text{by Lemma 6.2 (i)} \\ & = (1/n^2) \sum_{\sigma \in G} (q \times q)_* (\Gamma_{p_0 - p_0^{\sigma}}) & \text{by Lieberman's lemma} \\ & = (1/n^2) \sum_{\sigma \in G} (q' \times q')_* (\Gamma_{p_0 - p_0^{\sigma}}) & \text{by Proposition 5.10 (i)} \\ & = \Delta_A & \text{by the proper push-forward of the cycle} \end{split}$$

Next, we prove $b \circ a = \Delta_T$. Indeed, T_L is an abelian *L*-variety since $T(L) \neq \emptyset$. Replacing $\phi^{\sigma} \circ \phi^{-1}$ with $(\phi^{-1})^{\sigma} \circ \phi$ (Lemma 6.2 (i) with (ii)) in the previous discussion, we get $b \circ a = \Delta_T$ in $\operatorname{CH}_d(T \times T)$. Hence, we get $h(T) \cong h(A)$ in $\operatorname{CH}\mathcal{M}(K, \mathbb{Q})$.

Remark 6.3. The statement of Theorem 6.1 is mentioned in [FV20, Rem. 1.6, p.62].

7. Genus one fibrations and Jacobian fibrations

In this section, let k be an algebraically closed field of arbitrary characteristic. The purpose of this section is to prove the following:

Theorem 7.1. Let $f: X \to C$ be a minimal genus 1 fibration over k and $j: J \to C$ the Jacobian fibration of f. Then there are isomorphisms of Chow motives

(i)
$$h_1(X) \cong h_1(J), h_3(X) \cong h_3(J).$$

(ii)
$$h_2^{uig}(X) \cong h_2^{uig}(J)$$
.

The results of this section are based on [CD89] and [CDLK].

7.1. Fibrations. First of all, we fix the terminologies, so begin with:

Definition 7.2. Let $X, Y \in \mathcal{V}(k)$. A morphism $f : X \to Y$ is a *fibration* if it is a proper surjective morphism such that $f_*\mathcal{O}_X \cong \mathcal{O}_Y$.

Proposition 7.3. For a fibration $f: X \to Y, f^* : \operatorname{Pic}(Y) \to \operatorname{Pic}(X)$ is injective.

Proof. Let $\mathcal{L} \in \operatorname{Pic}(Y)$ with $f^*\mathcal{L} \cong \mathcal{O}_X$. By projection formula,

$$\mathcal{L} \cong \mathcal{L} \otimes \mathcal{O}_Y \cong \mathcal{L} \otimes f_* \mathcal{O}_X \cong f_* (f^* \mathcal{L} \otimes \mathcal{O}_X) \cong f_* (f^* \mathcal{L}) \cong f_* \mathcal{O}_X \cong \mathcal{O}_Y.$$

Let $f: X \to C$ be a fibration from a surface to a curve. Then f is flat, and all fibers of f are connected (e.g. [Har77, III. Prop. 9.7 and Coro. 11.3]). We consider a fiber X_c of f over a closed point $c \in C$ as an effective Cartier divisor with the sheaf of ideals $\mathcal{O}_X(-X_c) = f^*(\mathcal{O}_C(-c))$. Since X is regular, we can identify X_c with corresponding Weil divisor, and write the fiber

$$X_c = \sum_{i=1}^r m_i E_i$$

as the finite sum of its irreducible components. The number m_i is called the *multiplicity* of the component E_i , the number $m_c := \gcd(m_1, \dots, m_r)$ is called *multiplicity* of X_c , and the fiber X_c is called *multiple* (resp. *non-multiple*) if $m_c > 1$ (resp. $m_c = 1$). For any fiber X_c , we denote by $\overline{X_c}$ the divisor $1/m_c \cdot X_c$. Then $X_c = m_c \overline{X_c}$.

Let S be a surface. Let D and D' be divisors on S. We denote by $(D \cdot D')$ the intersection number of D and D'. The divisor D is *numerically equivelent* to D' (for short $D \equiv D'$) if $(D \cdot C) = (D' \cdot C)$ for any curve C on S.

7.2. Genus one fibrations. Let k be an algebraically closed field. Let

$$f: X \to C$$

be a minimal genus 1 fibration from a surface over k. Since f is a fibration, it is proper and flat. By general properties of morphism of schemes, all geometric fibers are geometrically-connected and there is a dense open subset U of C such that an *elliptic* (resp. a quasi-elliptic) fibration f is smooth (resp. geometrically-integral) over U. Let Σ be the finite set of closed points $c \in C$ such that the scheme-theoretical fiber X_c is not-smooth (resp. not-integral) if f is elliptic (resp. f is quasi-elliptic). We call $X_c, c \in \Sigma$, the singular fibers of f.

The following formula is well-known and is very useful:

Theorem 7.4. (canonical bundle formula). Let $f : X \to C$ be a genus 1 fibration. Let $R^1 f_* \mathcal{O}_X = \mathcal{L} \oplus T$ be the decomposition, with L an invertible sheaf on C and T an \mathcal{O}_C -module of finite length. Then

$$\omega_X \cong f^*(\mathcal{L}^{-1} \otimes \omega_C) \otimes \mathcal{O}_X(\sum_{i=1}^r n_i \overline{X_{c_i}}),$$

where

- (i) $m_i \overline{X_{c_i}} = X_{c_i} \ (c_i \in C)$ are all the multiple fibers of f,
- (ii) $0 \le n_i < m_i$,
- (iii) $n_i = m_i 1$ if c_i is not supported in T, and
- (iv) $\deg(\mathcal{L}^{-1} \otimes \omega_C) = 2p_a(C) 2 + \chi(\mathcal{O}_X) + \operatorname{length}(T).$

Proof. For example, see [BM77, Thm. 2, p.27] or [Băd01, Thm. 7.15, p.100].

In the rest of this subsection, we refine the relative Mordell-Weil theorem for the proof of Proposition 1.5. Let $f: X \to C$ be a genus 1 fibration. For any point $c \in C$ (not necessarily closed), we denote by

$$r_c: \operatorname{Pic}(X) \to \operatorname{Pic}(X_c)$$

the homomorphism obtained by restriction of invertible sheaves on X to X_c . We set

$$\operatorname{Pic}(X)_0 := \operatorname{Ker}(\operatorname{deg} \circ r_\eta)$$
$$= \{ \mathcal{L} \in \operatorname{Pic}(X) \mid \operatorname{deg} \circ r_c(\mathcal{L}) = 0 \text{ for any } c \in C \} \subset \operatorname{Pic}(X)$$

Here, the second equality follows from the function $c \mapsto \chi(X_c, r_c(\mathcal{L}))$ is constant on C ([Mum70, Coro. (b), p.50]). Then $\operatorname{Pic}(X)_0 \supset (\operatorname{Pic}^0_{X/k})_{red}(k)$. We define

$$\operatorname{Pic}(X)_f := \operatorname{Ker}(r_\eta) \subset \operatorname{Pic}(X)_0$$
$$\mathcal{E}(C) := \operatorname{Pic}(X)_f / f^* \operatorname{Pic}(C) \subset \operatorname{Pic}(X) / f^* \operatorname{Pic}(C)$$

For a singular fiber $X_c = \sum_{i=1}^r n_i E_i \in \text{Div}(X)$ with $E_i \neq E_j$ for $i \neq j$, set $\text{Num}_c(X) := \sum_{i=1}^r \mathbb{Z}[E_i]$ in Num(X). To refine the MW theorem, we need:

Proposition 7.5. ([CD89, Prop. 5.2.1, p.293]). Let $f: X \to C$ be a genus 1 fibration.

- (i) $\mathcal{E}(C) = \bigoplus_{c \in \Sigma} ((\operatorname{Num}_c(X) / \mathbb{Z}[\overline{X_c}]) \oplus \mathbb{Z} / m_c \mathbb{Z})$, where Σ is the set of singular points of f.
- (ii) $\operatorname{Pic}(X)_f / f^* \operatorname{Pic}^0(C)$ is a finitely-generated abelian group.
- (iii) $\operatorname{Pic}(X)_0/\operatorname{Pic}(X)_f \cong \operatorname{Jac}(X_n)(K).$

Proof. (i) By the local exact sequence $\bigoplus_c \operatorname{CH}_1(X_c) \xrightarrow{i_*} \operatorname{CH}_1(X) \xrightarrow{i_{\eta}} \operatorname{CH}_0(X_{\eta}) \to 0$, we have $\operatorname{Pic}(X)_f = \sum_c (\sum_i \mathbb{Z}[E_i])$, where c runs over all closed points of C and E_i are irreducible components of X_c . Note

 $[D] \in f^* \operatorname{Pic}(C)$ if and only if $[D] = \sum n_c[X_c] \ (n_c \in \mathbb{Z}).$

For a closed point $c \in C$, let $\mathcal{E}_c(C)$ be the subgroup of $\mathcal{E}(C)$ generated by the images of irreducible components of X_c . In other words, if $\operatorname{Pic}_c(X)$ is the subgroup of $\operatorname{Pic}(X)$ generated by the images of irreducible components of X_c , then

$$\mathcal{E}_c(C) = \operatorname{Pic}_c(X) / (\operatorname{Pic}_c(X) \cap f^*\operatorname{Pic}(C)) \cong (\operatorname{Pic}_c(X) + f^*\operatorname{Pic}(C)) / f^*\operatorname{Pic}(C).$$

(i-i) We have $\mathcal{E}(C) = \bigoplus_c \mathcal{E}_c(C)$. Indeed, assume $D_c \in \text{Div}(X)$ are divisors supported on X_c , and $\sum D_c = 0$ in $\mathcal{E}(C)$. Then $\sum_c D_c \sim f^*(\delta)$ for some $\delta \in \text{Div}(C)$, namely $\sum_c D_c - f^*(\delta) = \text{div}(\phi)$ for some $\phi \in k(X)^{\times}$. Since $\text{div}(\phi)$ has support on fibers, there is a non-empty open set U of C such that ϕ is regular on $f^{-1}U$; noting $f_*\mathcal{O}_X \cong \mathcal{O}_C$ one has $\phi \in \mathcal{O}_X(f^{-1}U) = \mathcal{O}_C(U)$ so $\phi = f^*\psi$ for some $\psi \in k(C)^{\times}$. Therefore $\sum_c D_c = f^*(\delta + \text{div}(\psi))$; rewriting δ for $\delta + \text{div}(\psi)$, we have $\sum_c D_c = f^*(\delta)$ in Div(X). If $\delta = \sum n_c[c]$, one has $D_c = n_c[X_c]$ for each c, in particular $D_c = 0$ in $\mathcal{E}(C)$.

(i-ii) For a closed point $c \in C$, let E_1, \dots, E_r be the irreducible components of X_c . Then $\mathcal{E}_c(C) \cong \mathbb{Z}\{E_i\}/\mathbb{Z}[X_c]$, where $\mathbb{Z}\{E_i\}$ is the free abelian group with basis $\{E_i\}$. Indeed, there is a surjection $\mathbb{Z}\{E_i\} \to \mathcal{E}_c(C)$. If $D \in \mathbb{Z}\{E_i\}$ maps to zero in $\mathcal{E}(C)$, the argument in (i) shows $D = n_c[X_c]$ for $n_c \in \mathbb{Z}$.

(i-iii) Recall that X_c is a primitive generator of $\mathbb{Z} \cdot X_c$, so $X_c = m_c \overline{X_c}$. One has obviously a short exact sequence

$$0 \to \mathbb{Z}/m_c\mathbb{Z} \to \mathbb{Z}\{E_i\}/\mathbb{Z}[X_c] \to \mathbb{Z}\{E_i\}/\mathbb{Z}[\overline{X_c}] \to 0.$$

The last group is free of rank r-1, so there is a non-canonical splitting of this short exact sequence. Recall $\operatorname{Num}_c(X)$ is the subgroup of $\operatorname{Num}(X)$ generated by the classes of irreducible components of X_c .

(i-iv) There is an isomorphism $\mathbb{Z}\{E_i\}/\mathbb{Z}[\overline{X_c}] \to \operatorname{Num}_c(X)/\mathbb{Z}[\overline{X_c}]$. Indeed, there is a surjection $\mathbb{Z}\{E_i\} \to \operatorname{Num}_c(X)$, which induces a surjection $\mathbb{Z}\{E_i\} \to \operatorname{Num}_c(X)/\mathbb{Z}[\overline{X_c}]$. Assume $D = \sum n_i E_i$ goes to 0 by this homomorphism; then one has $D - n\overline{X_c} \equiv 0$, so $(D^2) = 0$ by Chow's moving lemma, so $D \in \mathbb{Z}\overline{X_c}$ by [Băd01, Coro. 2.6, p.19].

Combining (i-i)-(i-iv), we have $\mathcal{E}(C) = \bigoplus_c \mathcal{E}_c(C)$ with non-canonical isomorphisms

$$\mathcal{E}_c(C) \cong (\operatorname{Num}_c(X) / \mathbb{Z}[\overline{X_c}]) \oplus \mathbb{Z} / m_c \mathbb{Z}.$$

In particular, if X_c is integral, then $\mathcal{E}_c(C) = 0$.

(ii) For two points c, c' of C, one has $[c] - [c'] \in \operatorname{Pic}^0(C)$, thus $[X_c] - [X_{c'}] \in f^* \operatorname{Pic}^0(C)$.

First, assume f has a singular fiber. If we choose a singular point c', in the group $\operatorname{Pic}(X)_f/f^*\operatorname{Pic}^0(C)$, any smooth fiber $[X_c]$ equals the singular fiber $[X_{c'}]$. Since $\operatorname{Pic}(X)_f$ is generated by the irreducible components of all fibers, it follows that $\operatorname{Pic}(X)_f/f^*\operatorname{Pic}^0(C)$ is generated by the irreducible components of the singular fibers only. Next, assume f has no singular fibers. If we choose a smooth point c', in the group $\operatorname{Pic}(X)_f/f^*\operatorname{Pic}^0(C)$, any smooth fiber $[X_c]$ equals the smooth fiber $[X_{c'}]$. Thus $\operatorname{Pic}(X)_f/f^*\operatorname{Pic}^0(C)$ is generated by the smooth fiber only.

(iii) By Tsen's thm.([GS06, Thm. 6.2.8, p.143]), Br(K) = 0. By Proposition 2.5,

$$\operatorname{Pic}^{0}(X_{\eta}) \cong \operatorname{Jac}(X_{\eta})(K)$$
 (6)

By definition, $\operatorname{Pic}(X)_0 = \operatorname{Ker}(\operatorname{Pic}(X) \xrightarrow{r_\eta} \operatorname{Pic}(X_\eta) \xrightarrow{\operatorname{deg}} \mathbb{Z})$, and hence the restriction $r_\eta^0 : \operatorname{Pic}(X)_0 \to \operatorname{Pic}^0(X_\eta)$ is well-defined. By the restriction $r_\eta : \operatorname{Pic}(X) \to \operatorname{Pic}(X_\eta)$ is surjective, so is r_η^0 . Thus, there is an isomorphism of groups

$$\operatorname{Pic}(X)_0/\operatorname{Pic}(X)_f \cong \operatorname{Pic}^0(X_\eta) \tag{7}$$

Combining (6) and (7), we get an isomorphism $\operatorname{Pic}(X)_0/\operatorname{Pic}(X)_f \cong \operatorname{Jac}(X_\eta)(K)$. \Box

To state a refined MW theorem, we recall a notion: Let K/k be a primary field extension and A an abelian K-variety. We say that a couple $(\operatorname{Tr}_{K/k}(A), \tau)$ consisting of an abelian k-variety $\operatorname{Tr}_{K/k}(A)$ and a K-homomorphism $\tau : \operatorname{Tr}_{K/k}(A)_K \to A$ is a K/k-trace of A if τ has finite kernel and satisfies the following condition. For an abelian k-variety B and a K-homomorphism $\beta : B_K \to A$, there is a k-homomorphism $\beta' : B \to \operatorname{Tr}_{K/k}(A)$ such that $\beta = \tau \circ \beta'_K$. Indeed, it exists ([Lag59, Thm. 8, p.213]).

The following theorem is a refined version of the MW (or the Lang-Néron) theorem. We focus on the K/k-trace of $\operatorname{Jac}(X_{\eta})$ that is isogenous to a sub-abelian variety of $(\operatorname{Pic}^{0}_{X/k})_{red}$ for the proof of Proposition 1.5:

Theorem 7.6. Let $f: X \to C$ be a genus 1 fibration. Let T be an abelian k-variety of dimension ≤ 1 that is the K/k-trace $(\operatorname{Tr}_{K/k}(\operatorname{Jac}(X_{\eta})), \tau)$ of $\operatorname{Jac}(X_{\eta})$ if f is elliptic and is 0 if f is quasi-elliptic. Then

- (i) There is an isogeny of abelian k-varieties $(\operatorname{Pic}^{0}_{X/k})_{red} \sim_{isog} T \times \operatorname{Pic}^{0}_{C/k}$
- (ii) $T = 0 \iff b_1(X) = b_1(C)$

$$\iff \operatorname{Jac}(X_{\eta})(K)/\tau(T)(k) = \operatorname{Jac}(X_{\eta})(K)$$
 is f.g. (Lang-Néron)
Moreover, if these conditions are satisfied, then

$$\operatorname{rank}(\operatorname{Jac}(X_{\eta})(K)) = \rho(X) - 2 - \sum_{c \in C} (\#\operatorname{Irr}(X_c) - 1) \quad \text{(Shioda-Tate formula)}$$

Here, $Irr(X_c)$ is the set of the distinct irreducible components of the singular fiber X_c .

(iii) $T \neq 0 \iff b_1(X) = b_1(C) + 2$ $\iff \operatorname{Jac}(X_\eta)(K)$ is an not finitely-generated abelian group \iff there is an isogeny of elliptic K-curves

$$\psi: T \times_{\operatorname{Spec}(k)} \operatorname{Spec}(K) \to \operatorname{Jac}(X_{\eta}).$$

In particular, if f is quasi-elliptic, then $Jac(X_n)(K)$ is finitely-generated.

Proof. This proof is based on [CD89, Prop. 5.2.1 (v), p.293]. By Proposition 7.3, there is an injective morphism of group schemes $f^* : \operatorname{Pic}_{C/k} \hookrightarrow \operatorname{Pic}_{X/k}$. So we get an injection

$$f^* : \operatorname{Pic}^0_{C/k} = (\operatorname{Pic}^0_{C/k})_{red} \hookrightarrow (\operatorname{Pic}^0_{X/k})_{red}$$

By Theorem 2.3, $\operatorname{Pic}_{C/k}^{0}$ and $(\operatorname{Pic}_{X/k}^{0})_{red}$ are abelian k-varieties. By Poincare's reducibility theorem, there is an abelian k-subvariety $A \subset (\operatorname{Pic}_{X/k}^{0})_{red}$ such that

$$(\operatorname{Pic}^{0}_{X/k})_{red} \sim_{isog} A \times \operatorname{Pic}^{0}_{C/k}.$$
(8)

First, we prove that A satisfies assertions (i)-(iii). By (8), it remains to prove $\dim(A) \leq 1$ and A satisfies assertions (ii) and (iii).

Assume A = 0. Then $\operatorname{Pic}_{C/k}^0 \sim_{isog} (\operatorname{Pic}_{X/k}^0)_{red}$. So $\operatorname{Pic}_{C/k}^0(k) = (\operatorname{Pic}_{X/k}^0)_{red}(k)$. Now

$$\operatorname{Pic}(X)/f^*\operatorname{Pic}(C) = (\operatorname{Pic}_{X/k})_{red}(k)/\operatorname{Pic}_{C/k}(k)$$
$$(\operatorname{Pic}_{X/k})_{red}(k)/(\operatorname{Pic}_{X/k}^0)_{red}(k) = \operatorname{NS}(X)$$
$$(\operatorname{Pic}_{C/k})_{red}(k)/(\operatorname{Pic}_{C/k}^0)_{red}(k) = \mathbb{Z}.$$

Therefore,

$$\operatorname{Pic}(X)/f^*\operatorname{Pic}(C) \cong \operatorname{NS}(X)/\mathbb{Z}.$$

By Proposition 2.4, NS(X) is f.g. of rank $\rho(X)$. Thus, the group $\operatorname{Pic}(X)/f^*\operatorname{Pic}(C)$ is f.g. of rank $\rho(X) - 1$. Since $\operatorname{Pic}(X)_0 = \operatorname{Ker}(\operatorname{deg} \circ r_\eta : \operatorname{Pic}(X) \to \mathbb{Z})$, one has $\operatorname{Pic}(X)/\operatorname{Pic}(X)_0 \cong \operatorname{Im}(\operatorname{deg} \circ r_\eta) \cong \mathbb{Z}$. Hence

the group
$$\operatorname{Pic}(X)_0/f^*\operatorname{Pic}(C)$$
 is f.g. of rank $\rho(X) - 2$ (9)

On the other hand,

$$\operatorname{Jac}(X_{\eta})(K) = \frac{\operatorname{Pic}(X)_{0}}{\operatorname{Pic}(X)_{f}} = \frac{\operatorname{Pic}(X)_{0}/f^{*}\operatorname{Pic}(C)}{\operatorname{Pic}(X)_{f}/f^{*}\operatorname{Pic}(C)} = \frac{\operatorname{Pic}(X)_{0}/f^{*}\operatorname{Pic}(C)}{\bigoplus_{c}((\operatorname{Num}_{c}(X)/\mathbb{Z}[\overline{X_{c}}]) \oplus \mathbb{Z}/m_{c}\mathbb{Z})}$$
(10)

Here, the first quality uses Proposition 7.5 (iii), and the third Proposition 7.5 (i). By (9) and (10),

the group
$$\operatorname{Jac}(X_{\eta})(K)$$
 is f.g. of rank $\rho(X) - 2 - \sum_{c \in C} (\#\operatorname{Irr}(X_c) - 1)$

In particular, we see that

$$``A = 0 \Rightarrow \operatorname{Jac}(X_{\eta})(K) \text{ is f.g.''}$$
(11)

Next, assume $A \neq 0$. Then $A \times_{\text{Spec}(k)} \text{Pic}_{C/k}^0 \sim_{isog} \text{Pic}_{X/k}^0$. By the base extension K/k, we get a morphism of group K-varieties

 $(A \times_{\operatorname{Spec}(k)} \operatorname{Spec}(K)) \times_{\operatorname{Spec}(K)} (\operatorname{Pic}^{0}_{C/k} \times_{\operatorname{Spec}(k)} \operatorname{Spec}(K)) \to \operatorname{Pic}^{0}_{X/k} \times_{\operatorname{Spec}(k)} \operatorname{Spec}(K).$

Then, there is a morphism of group K-varieties

$$\phi: A \times_{\operatorname{Spec}(k)} \operatorname{Spec}(K) \to \operatorname{Pic}^{0}_{X/k} \times_{\operatorname{Spec}(k)} \operatorname{Spec}(K) \cong \operatorname{Pic}^{0}_{X \times_{\operatorname{Spec}(k)} \operatorname{Spec}(K)/K}.$$

Here, let $i: X_{\eta} = X \times_C \operatorname{Spec}(K) \to X \times_{\operatorname{Spec}(k)} \operatorname{Spec}(K)$ be the closed K-immersion. Then, we get a pull-back of group K-varieties

$$i^* : \operatorname{Pic}^0_{X \times_{\operatorname{Spec}(k)} \operatorname{Spec}(K)} \to \operatorname{Pic}^0_{X_{\eta}/K} = \operatorname{Jac}(X_{\eta}).$$

Hence, we obtain a morphism of group K-varieties

$$\psi := i^* \circ \phi : A_K = A \times_{\operatorname{Spec}(k)} \operatorname{Spec}(K) \to \operatorname{Jac}(X_\eta).$$

Here we need:

Lemma 7.7. Ker (ψ) is a finite group scheme.

Proof. Assume dim(Ker(ψ)) > 0. Then Ker(ψ)⁰ $\subset A_K$ is an abelian K-variety of positive dimension. Since C is geometrically connected, K/k is a primary field extension. By [Lag59, Thm. 5, p.26], there is an abelian k-variety $B \neq 0$ such that

$$\operatorname{Ker}(\psi)^0 \cong B_K. \tag{12}$$

Since the m-torsion subgroup B(k)[m] is isomorphic to $(\mathbb{Z}/m\mathbb{Z})^{2\dim(B)}$ for any integer m which is not divisible by char(k) (cf. [Mum70, Prop. (3), p.64]), we see that

$$B(k)$$
 is not f.g. (13)

On the other hand, since $\operatorname{Ker}(\psi)^0(K) = B(K)$ by (12), we have

$$B(k) = \operatorname{Ker}(\psi)^{0}(K) \cap B(k) \subset \operatorname{Ker}(\psi)(K) \cap A(k)$$

Since $\operatorname{Jac}(X_{\eta})(K) = \operatorname{Pic}(X)_0/\operatorname{Pic}(X)_f$ by Proposition 7.5 (iii), we have $\operatorname{Ker}(\psi)(K) = A(K) \cap \operatorname{Pic}(X)_f$, so we get

$$B(k) \subset \operatorname{Ker}(\psi)(K) \cap A(k) = A(k) \cap \operatorname{Pic}(X)_f.$$
(14)

Thus, it suffices to show that $A(k) \cap \operatorname{Pic}(X)_f$ is f.g. Indeed, by Proposition 7.5 (ii), $\operatorname{Pic}(X)_f/f^*\operatorname{Pic}^0(C)$ is f.g. By (8), $A(k) \cap f^*\operatorname{Pic}^0(C)$ is finite. By the inclusion

$$\frac{A(k) \cap \operatorname{Pic}(X)_f}{A(k) \cap f^* \operatorname{Pic}^0(C)} \subset \frac{\operatorname{Pic}(X)_f}{f^* \operatorname{Pic}^0(C)},$$

we see that $A(k) \cap \operatorname{Pic}(X)_f$ is f.g. By (14), we see that

$$B(k)$$
 is f.g. (15)

By (13) and (15), we obtain a contradiction. Hence $\text{Ker}(\psi)$ is a finite group scheme. \Box

Let us go back to the proof of Theorem 7.6. The morphism $\psi : A_K \to \text{Jac}(X_\eta)$ is either constant or surjective because $\dim(\text{Jac}(X_\eta)) = 1$. Since $\text{Ker}(\psi)$ is finite by Lemma 7.7, ψ is surjective, so ψ is isogeny. In particular,

$$``A \neq 0 \Rightarrow \operatorname{Jac}(X_{\eta}) \sim_{isog} A_K \Rightarrow \operatorname{Jac}(X_{\eta})(K) \text{ is not f.g.''}$$
(16)

Combining (11) and (16), we get $\dim(A) \leq 1$ and

"
$$A = 0 \iff \operatorname{Jac}(X_{\eta})(K)$$
 is f.g." and " $A \neq 0 \iff \operatorname{Jac}(X_{\eta})(K)$ is not f.g."

Now, by (8) and dim $((\text{Pic}_{-/k}^{0})_{red}) = 1/2 \cdot b_1(-)$ ([CD89, Prop. 0.7.4, p.69]), we get

$$\dim(A) = \dim((\operatorname{Pic}^{0}_{X/k})_{red}) - \dim((\operatorname{Pic}^{0}_{C/k})) = 1/2(b_1(X) - b_1(C)) \le 1.$$

In particular, we get

"
$$A = 0 \Leftrightarrow b_1(X) = b_1(C)$$
" and " $A \neq 0 \Leftrightarrow b_1(X) = b_1(C) + 2$ "

Thus, we see that A satisfies assertions (i)-(iii). In particular, if $A \neq 0$, then $\text{Jac}(X_{\eta})$ is an elliptic curve, so f is elliptic. On the contrary, f is quasi-elliptic, then A = 0, so $\text{Jac}(X_{\eta})(K)$ is finitely-generated.

Finally, we prove the existence of a k-isogeny $T \stackrel{isog}{\sim} A$. Assume T = 0. Then the map $\tau : T_K \to \text{Jac}(X_\eta)$ is not a K-isogeny since $\dim(T_K) < \dim(\text{Jac}(X_\eta))$, so we get A = 0 by assertion (iii) and the universal property of τ , and hence $T \stackrel{isog}{\sim} A$.

Next, assume $T \neq 0$. Then the map $\tau : T_K \to \operatorname{Jac}(X_\eta)$ is a K-isogeny since $\dim(T_K) = \dim(\operatorname{Jac}(X_\eta))$, so $\operatorname{Jac}(X_\eta)(K)$ is not finitely-generated. By assertion (iii), we have a K-isogeny $\psi : A_K \to \operatorname{Jac}(X_\eta)$, so we have a K-isogeny $T_K \overset{isog}{\sim} A_K$. Now K/k is primary. By [Lag59, Thm. 5, p.26], there is a k-isogeny of elliptic curves

$$T \stackrel{isog}{\sim} A$$

Since A satisfies assertions (i)-(iii), so is T.

7.3. Sections. Let $f: X \to C$ be a fibration from a surface to a curve.

Definition 7.8. A morphism $s: C \to X$ is a section of f if $f \circ s = id_C$.

We identify a section s of f with its image in X. This is a curve S such that $f|_S$ is an isomorphism, or equivalently $(S \cdot X_c) = 1$ for every closed fiber X_c of f. Let X(C)denote the set of sections of f. For any morphism $g: Z \to S$ of regular schemes, let

 $Z^{\#} := \{ z \in Z \mid g \text{ is smooth at } z \}.$

We will use the following properties of sections:

Lemma 7.9. Let $f: X \to C$ be a fibration from a surface to a curve. Let X_{η} be the generic fiber of f. Let K be the function field of C. Then

- (i) Giving a section s of f is equivalent to giving a K-rational point of X_{η} . Moreover, the image of s lies in $X^{\#}$ and corresponds to a point of $X_{\eta}^{\#}(K)$.
- (ii) If f has a section, then f has no multiple fibers.

Proof. (i) Let s be a section of f. By the base change $\operatorname{Spec}(K) \to C$, we get $s \times \operatorname{id}_K :$ $\operatorname{Spec}(K) \cong C \times_C \operatorname{Spec}(K) \hookrightarrow X_\eta$, so $s \times \operatorname{id}_K \in X_\eta(K)$. Conversely, let $\xi \in X_\eta(K)$. Take the closure S of ξ in X. Since $k(S) \cong k(C) \cong K$ and C is a normal k-curve, the proper birational map $f|_S : S \dashrightarrow C$ is an isomorphism, so $f|_S^{-1} \in X(C)$.

So, it remains to show for any $s \in X(C)$, any closed fiber X_c is smooth at x := s(c). Let $R := \mathcal{O}_{C,c}$, m its maximal ideal, and $f' : X' \to C'$ the base change of f by $\operatorname{Spec}(R) \to C$. By $X_c \cong X'_c$, we may assume $C = \operatorname{Spec}(R)$. Let $A := \mathcal{O}_{X,x}$ and m_A its maximal ideal. Then $\mathcal{O}_{X_c,x} = A/mA$ and its maximal ideal is $M := m_A/mA$. The image of m under the local homomorphism $R \to A \to R$ defined by f and s is equal to m. This implies that mA is not contained in m_A^2 . Thus the surjection $m_A/m_A^2 \to m_A/(m_A^2 + mA) = M/M^2$ is not injective, so $\dim(M/M^2) < \dim(m_A/m_A^2)$. As X is smooth at x, $\dim(m_A/m_A^2) = \dim(X) = 2$. Also $\dim(A/mA) = \dim(X_c) = 1$. Hence $\dim(A/mA) \leq \dim(M/M^2)$ is an equality, so X_c is smooth at x. As the above, $s \times \operatorname{id}_K \in X^{\#}_n(K)$.

(ii) If X_c is a multiple fiber of f, we can write $X_c = m_c D$ with $m_c \in \mathbb{Z}_{>1}$. If $S \subset X$ is a section of f, then $(S \cdot X_c) = m_c(S \cdot D) \ge m_c > 1$. So we obtain a contradiction. \Box

Let S be a Dedekind scheme and K its function field. Let G be a separated group K-scheme of finite type. Let X be a smooth group S-scheme of finite type.

Definition 7.10. X is Néron model of G if it satisfies the following conditions:

- (i) X is an S-model of G, i.e. $X_K \cong G$.
- (ii) For each smooth S-scheme Y and each K-morphism $\phi_K : Y_K \to X_K$, there is a unique S-morphism $\psi : Y \to X$ which extends ϕ_K .

7.4. Jacobian fibrations. The main object of this paper is the following.

Definition 7.11. A genus 1 fibration $f : X \to C$ is called *Jacobian* if it admits a section, i.e. $X(C) \neq \emptyset$. Equivalently, X_{η} has a η -rational point by Lemma 7.9 (i).

Now we associate to every genus 1 fibration a Jacobian genus 1 fibration that satisfies some good properties. Roughly speaking, it is easier to study j than f:

Proposition 7.12. ([CD89, Prop. 5.2.5, p.299]). Let $f : X \to C$ be a genus 1 fibration. Then there is a unique (up to *C*-isomorphism) Jacobian genus 1 fibration $j : J \to C$ such that the smooth locus $J^{\#}$ is *C*-isomorphic to the Néron model of $Jac(X_{\eta})$. In particular, *J* satisfies the following properties:

- (i) $J_{\eta}^{\#} \cong \operatorname{Jac}(X_{\eta}) \neq \emptyset$, where $J_{\eta}^{\#}$ is the generic fiber of $j: J^{\#} \to C$,
- (ii) the image of any section $C \to J$ lies in $J^{\#}$, and
- (iii) the natural map of sections $J(C) \to \operatorname{Jac}(X_n)(K)$ is bijective.

Proof. First, we prove the existence of J. Let $\eta \in C$ be the generic point, X_{η} the generic fiber of f, and J_{η} a regular compactification of $\operatorname{Jac}(X_{\eta})$ as in Proposition 5.9. Now, we have injections $J_{\eta} \hookrightarrow \mathbb{P}^n_K \hookrightarrow \mathbb{P}^n_k \times C$. Take schematic closure:

$$J \hookrightarrow \mathbb{P}^n_k \times C.$$

Then there is a regular projective scheme $J \to C$, flat over C and with the generic fiber J_{η} . After resolving singularities of J, blowing down (-1) curves in closed fibers, and replacing J by it, we obtain a genus 1 fibration $j : J \to C$. The uniqueness of J follows from the theory of minimal models. By the same argument as in [Art86, Prop. 2.15, p.218] or [LT16, Thm. 1.1], we see that $J^{\#}$ is C-isomorphic to the Néreon model of $Jac(X_{\eta})$, so get (i)-(iii).

We call $j: J \to C$ the Jacobian fibration of $f: X \to C$. First, we see:

Proposition 7.13. ([CD89, Prop. 5.3.2, p.303]). Every Jacobian genus 1 fibration $f: X \to C$ is *C*-isomorphic to the Jacobian fibration $j: J \to C$ of f.

Proof. By assumption, $X(C) \neq \emptyset$. By Lemma 7.9 (i), $X_{\eta}^{\#}(K) \neq \emptyset$. By Proposition 7.12, J_{η} is a regular compactification of $\operatorname{Jac}(X_{\eta})$. By Proposition 5.9 (i), we get $X_{\eta} \cong J_{\eta}$. This induces an isomorphism of function fields $k(X) \cong k(J)$. Then there is a birational map $J \dashrightarrow X$. Since X is minimal and $p_a(X_{\eta}) \geq 1$, any birational map to X is a birational morphism (cf. [Liu02, Coro. 3.24, p.422]), so we get $X \cong J$. \Box

Proposition 7.14. (canonical bundle formula). Let $j : J \to C$ be a Jacobian genus 1 fibration. Let $\mathcal{L}' := R^1 j_* \mathcal{O}_J$ be the invertible sheaf on C. Then

$$\omega_J \cong j^*(\mathcal{L}'^{-1} \otimes \omega_C), \ \ \deg(\mathcal{L}') = -\chi(\mathcal{O}_J).$$

Proof. By Lem. 7.9 (ii), j has no multiple fibers. Thus, it follows from Thm. 7.4. \Box

From now on, we prove relations between some invariants of f and j, so begin with:

Corollary 7.15. ([CD89, p.307]). Let $f : X \to C$ be a genus 1 fibration and $j : J \to C$ the Jacobian fibration of f. Then there is an isomorphism of group K-varieties

$$\operatorname{Jac}(X_{\eta}) \cong \operatorname{Jac}(J_{\eta}).$$

Proof. By Proposition 7.12, J_{η} is a regular compactification of $\text{Jac}(X_{\eta})$. Thus, it follows from Proposition 5.9 (iii).

The following proposition plays a key role in the proof of Theorem 7.1 (i).

Proposition 7.16. Let $f: X \to C$ be a genus 1 fibration and $j: J \to C$ the Jacobian fibration of f. Then there are isogenies of abelian k-varieties

$$(\operatorname{Pic}^{0}_{X/k})_{red} \sim_{isog} (\operatorname{Pic}^{0}_{J/k})_{red}, \quad \operatorname{Alb}_{X/k} \sim_{isog} \operatorname{Alb}_{J/k}.$$

Proof. It suffices to prove the existence of a k-isogeny $(\operatorname{Pic}^{0}_{X/k})_{red} \sim_{isog} (\operatorname{Pic}^{0}_{J/k})_{red}$. By Theorem 7.6, there are abelian k-varieties T, T' such that

$$(\operatorname{Pic}^{0}_{X/k})_{red} \sim_{isog} T \times \operatorname{Pic}^{0}_{C/k}, \qquad (\operatorname{Pic}^{0}_{J/k})_{red} \sim_{isog} T' \times \operatorname{Pic}^{0}_{C/k}$$

First, assume f is quasi-elliptic. By the construction of j, j is also quasi-elliptic. By our convention in Theorem 7.6, we have

$$T = T' = 0.$$

Thus, we get an isogeny of abelian k-varieties

$$(\operatorname{Pic}^{0}_{X/k})_{red} \sim_{isog} \operatorname{Pic}^{0}_{C/k} \sim_{isog} (\operatorname{Pic}^{0}_{J/k})_{red}.$$

Next, assume f is elliptic. By construction, j is also elliptic. By Theorem 7.6,

$$T = \operatorname{Tr}_{K/k}(\operatorname{Jac}(X_{\eta})), \qquad T' = \operatorname{Tr}_{K/k}(\operatorname{Jac}(J_{\eta}))$$

By Corollary 7.15, we have $\operatorname{Jac}(X_{\eta}) \cong \operatorname{Jac}(J_{\eta})$, so we get an isomorphism of k-abelian varieties

 $T \cong T'.$

Therefore, we get an isogeny of abelian k-varieties

$$(\operatorname{Pic}^{0}_{X/k})_{red} \sim_{isog} T \times_{\operatorname{Spec}(k)} \operatorname{Pic}^{0}_{C/k} \cong T' \times_{\operatorname{Spec}(k)} \operatorname{Pic}^{0}_{C/k} \sim_{isog} (\operatorname{Pic}^{0}_{J/k})_{red}.$$

Using Proposition 7.16, we prove the first main theorem of this section:

Theorem 7.17. Let $f: X \to C$ be a genus 1 fibration and $j: J \to C$ the Jacobian fibration of f. Then there are isomorphism of Chow motives

$$h_1(X) \cong h_1(J), \quad h_3(X) \cong h_3(J).$$

Proof. First, we prove $h_1(X) \cong h_1(J)$. By Proposition 7.16, there is an isogeny of abelian varieties

$$(\operatorname{Pic}^{0}_{X/k})_{red} \sim_{isog} (\operatorname{Pic}^{0}_{J/k})_{red}$$

Since isogenous abelian varieties have isomorphic Chow motives (cf. [DM91, Thm. 3.1]), we get an isomorphism

$$h_1((\operatorname{Pic}^0_{X/k})_{red}) \cong h_1((\operatorname{Pic}^0_{J/k})_{red}).$$
 (17)

On the other hand, by Proposition 4.2, we have an isomorphism of Chow motives

$$h_1(X) \cong h_1((\operatorname{Pic}^0_{X/k})_{red}).$$
 (18)

Combining (17) and (18), we get $h_1(X) \cong h_1(J)$. Similarly, we get $h_3(X) \cong h_3(J)$ (because $(\operatorname{Pic}^0_{X/k})_{red} \sim_{isog} \operatorname{Alb}_{X/k}$).

Remark 7.18. Taking the étale realizations of $h_i(X) \cong h_i(J)$ (i = 1, 3), we have isomorphisms $H^i_{\acute{e}t}(X, \mathbb{Q}_l) \cong H^i_{\acute{e}t}(J, \mathbb{Q}_l)$, so get $b_i(X) = b_i(J)$ (i = 1, 3).

To prove Thm. 7.1 (ii), we recall some known relations between f and j:

Proposition 7.19. ([CD89, Prop. 5.3.6, p.308]). Let $f : X \to C$ be a genus 1 fibration and $j : J \to C$ the Jacobian fibration of f. Then

- (i) $\chi(\mathcal{O}_X) = \chi(\mathcal{O}_J)$ and e(X) = e(J).
- (ii) $\rho(X) = \rho(J)$ and $b_i(X) = b_i(J)$ for every $i \ge 0$.

Proof. (i) See [CD89, Prop. 5.3.6, p.308]. The heart is [LLR04, Thm. 4.2, p.482].³ (ii) See [CD89, Coro. 5.3.5, p.310]. The proof of (ii) is used by (i). \Box

We prove the second main theorem of this section:

Theorem 7.20. Let $f: X \to C$ be a genus 1 fibration and $j: J \to C$ the Jacobian fibration of f. Then there is an isomorphism of Chow motives

$$h_2^{alg}(X) \cong h_2^{alg}(J)$$

Proof. For a surface S, $h_2^{alg}(S) \cong \mathbb{L}^{\rho(S)}$. By Proposition 7.19 (ii), $\rho(X) = \rho(J)$, so $h_2^{alg}(X) \cong \mathbb{L}^{\oplus \rho(X)} = \mathbb{L}^{\oplus \rho(J)} \cong h_2^{alg}(J)$.

8. Chow motives of genus one fibrations

In this section, we prove the first main theorem of this paper:

Theorem 8.1. Let k be an algebraically closed field of arbitrary characteristic. Let $f: X \to C$ be a minimal genus 1 fibration from a smooth projective surface over k, and $j: J \to C$ the Jacobian fibration of f. Then there is an isomorphism

$$h(X) \cong h(J)$$

in the category $CH\mathcal{M}(k,\mathbb{Q})$ of Chow motives.

Proof of Theorem 8.1. Let us consider the CK-decompositions of Murre (Proposition 4.2) and Kahn-Murre-Pedrini (Proposition 4.3) of h(X) and h(J), respectively:

$$h(X) \cong 1 \oplus h_1(X) \oplus h_2^{alg}(X) \oplus t_2(X) \oplus h_3(X) \oplus (\mathbb{L} \otimes \mathbb{L}).$$

$$h(J) \cong 1 \oplus h_1(J) \oplus h_2^{alg}(J) \oplus t_2(J) \oplus h_3(J) \oplus (\mathbb{L} \otimes \mathbb{L}).$$

By Theorem 7.17 and Theorem 7.20, we have

$$h_1(X) \cong h_1(J), \quad h_3(J) \cong h_3(J), \quad h_2^{alg}(X) \cong h_2^{alg}(J).$$

Therefore, to prove Theorem 8.1, it suffices to prove the following:

³Liu-Lorenzini-Raynaud showed that the singular fibers X_t and J_t are of the "same type" ([LLR04, Thm. 6.6, p.455]). This is one of the most profound results about the relations between f and j.

Theorem 8.2. Let $f: X \to C$ be a minimal genus 1 fibration and $j: J \to C$ the Jacobian fibration of f. There is an isomorphism of transcendental motives

$$t_2(X) \cong t_2(J)$$
 in $\operatorname{CH}\mathcal{M}(k, \mathbb{Q}).$

Proof of Theorem 8.2. By definition, f is either elliptic or quasi-elliptic.

8.1. The transcendental motives of the quasi-elliptic surfaces.

First, we assume f is quasi-elliptic. Then, j is also quasi-elliptic by the construction of j. Let us recall the result of the author:

Theorem 8.3. ([Kaw22]). Let $f: X \to C$ be a quasi-elliptic surface. Then $t_2(X) = 0$.

We apply Theorem 8.3 to f and j, and get

$$t_2(X) = 0 = t_2(J).$$

Thus, we complete the proof of Theorem 8.2 for the case where f is quasi-elliptic.

8.2. The transcendental motives of the elliptic surfaces.

Next, we assume f is elliptic. By Proposition 4.8, it suffices to prove: there are elements $[\alpha] \in CH_2(X \times J)/CH_2(X \times J)_{\equiv}, [\beta] \in CH_2(J \times X)/CH_2(J \times X)_{\equiv}$ such that

$$\begin{cases} [\alpha] \circ_k [\beta] = [\Delta_J] & \text{in } \operatorname{CH}_2(J \times J)/\operatorname{CH}_2(J \times J)_{\equiv} \\ [\beta] \circ_k [\alpha] = [\Delta_X] & \text{in } \operatorname{CH}_2(X \times X)/\operatorname{CH}_2(X \times X)_{\equiv} \end{cases}$$

Step 1. Construct correspondences on the elliptic surfaces.

We construct the correspondences $[\alpha]$ and $[\beta]$. Let $\eta \in C$ be the generic point. Let X_{η} and J_{η} be the generic fibers of $f: X \to C$ and $j: J \to C$, respectively. Since f is elliptic, X_{η} is a smooth genus 1 curve. By the construction of $J_{\eta}, J_{\eta} \cong \text{Jac}(X_{\eta})$. By Proposition 5.8, X_{η} is a torsor for J_{η} . By Theorem 6.1, we get

$$h(X_{\eta}) \cong h(J_{\eta})$$
 in $\operatorname{CH}\mathcal{M}(\eta, \mathbb{Q}).$

Thus, there are elements $a \in \operatorname{CH}_1(X_\eta \times_\eta J_\eta)$ and $b \in \operatorname{CH}_1(J_\eta \times_\eta X_\eta)$ such that

$$\begin{cases} a \circ_{\eta} b = \Delta_{J_{\eta}} & \text{in } \operatorname{CH}_{1}(J_{\eta} \times_{\eta} J_{\eta}) \\ b \circ_{\eta} a = \Delta_{X_{\eta}} & \text{in } \operatorname{CH}_{1}(X_{\eta} \times_{\eta} X_{\eta}) \end{cases}$$

In this proof, we let

$$r_{XJ}$$
: CH₂ $(X \times_C J) \to$ CH₁ $(X_\eta \times_\eta J_\eta)$

be the flat-pullback, and similar for r_{JX} and r_{JJ} . Let

$$\iota_{XJ}: \mathrm{CH}_2(X \times_C J) \to \mathrm{CH}_2(X \times J)$$

be the proper-pushfoward, and similar for ι_{JX} and ι_{JJ} .

Lemma 8.4. There are homomorphisms of groups

$$\operatorname{CH}_1(X_\eta \times_\eta J_\eta) \xleftarrow{\simeq}_{r_{XJ}} \frac{\operatorname{CH}_2(X \times_C J)}{\bigoplus_c \operatorname{CH}_2(X_c \times_c J_c)} \xrightarrow{\iota_{XJ}} \frac{\operatorname{CH}_2(X \times J)}{\operatorname{CH}_2(X \times J)_{\equiv}}.$$

Proof. The left isomorphism r_{XJ} follows from the local exact sequences

$$\oplus_c \mathrm{CH}_2(X_c \times_c J_c) \longrightarrow \mathrm{CH}_2(X \times J) \xrightarrow[r_{XJ}]{} \mathrm{CH}_1(X_\eta \times_\eta J_\eta) \to 0.$$

Thus, it remains to prove $\iota_{XJ}(\bigoplus_c \operatorname{CH}_2(X_c \times_c J_c)) \subset \operatorname{CH}_2(X \times J)_{\equiv}$. Indeed, let $z \in \bigoplus_c \operatorname{CH}_2(X_c \times_c J_c)$. Then $z = \sum_c \sum_{i,j} n_{c,i,j} [E_{c,i} \times_{k(c)} F_{c,j}]$ with $n_{c,i,j} \in \mathbb{Q}$. Here, $E_{c,i}$ and $F_{c,j}$ runs over all irreducible components of X_c and J_c , respectively. Thus

$$\iota_{XJ}(z) = \sum_{c} \sum_{i,j} n_{c,i,j} [E_{c,i} \times_k F_{c,j}] \in \operatorname{CH}_2(X \times_C J)_{\equiv}.$$

By Lemma 8.4, we can define

$$[\alpha] := \iota_{XJ}(r_{XJ}^{-1}(a)) \in \mathrm{CH}_2(X \times J)/\mathrm{CH}_2(X \times J, \mathbb{Q})_{\equiv}.$$

Similarly, we define

$$[\beta] := \iota_{JX}(r_{JX}^{-1}(b)) \in \mathrm{CH}_2(J \times X) / \mathrm{CH}_2(J \times X)_{\equiv}.$$

Step 2. Calculate the correspondences on the elliptic surfaces.

To compute the correspondences $[\alpha] \circ_k [\beta]$ and $[\beta] \circ_k [\alpha]$, we prove the following:

Lemma 8.5. Use above notations. Then

(i) For $[y] \in \operatorname{CH}_2(X \times_C J) / \oplus_c \operatorname{CH}_2(X_c \times_c J_c), [z] \in \operatorname{CH}_2(J \times_C X) / \oplus_c \operatorname{CH}_2(J_c \times_c X_c),$ $\iota_{XJ}([y]) \circ_k \iota_{JX}([z]) = \iota_{JJ}([y] \circ_C [z]) \text{ in } \operatorname{CH}_2(J \times J) / \operatorname{CH}_2(J \times J)_{\equiv}.$ (ii) For $d \in \operatorname{CH}_1(X_\eta \times_\eta J_\eta), e \in \operatorname{CH}_1(J_\eta \times_\eta X_\eta),$

$$r_{XJ}^{-1}(d) \circ_C r_{JX}^{-1}(e) = r_{JJ}^{-1}(d \circ_\eta e) \quad \text{in} \quad \operatorname{CH}_2(J \times_C J) / \oplus_c \operatorname{CH}_2(J_c \times_c J_c).$$

Proof. We prove (i). By Proposition 4.7, there is a bilinear homomorphism

$$\circ: \frac{\operatorname{CH}_2(J \times X)}{\operatorname{CH}_2(J \times X)_{\equiv}} \times \frac{\operatorname{CH}_2(X \times J)}{\operatorname{CH}_2(X \times J)_{\equiv}} \to \frac{\operatorname{CH}_2(J \times J)}{\operatorname{CH}_2(J \times J)_{\equiv}}$$
(19)
$$([\delta], [\gamma]) \mapsto [\gamma] \circ_k [\delta] := [\gamma \circ_k \delta]$$

Similarly, there is a bilinear homomorphism

$$\circ: \frac{\operatorname{CH}_2(J \times_C X)}{\bigoplus_c \operatorname{CH}_2(J_c \times X_c)} \times \frac{\operatorname{CH}_2(X \times_C J)}{\bigoplus_c \operatorname{CH}_2(X_c \times J_c)} \to \frac{\operatorname{CH}_2(J \times_C J)}{\bigoplus_c \operatorname{CH}_2(J_c \times J_c)}$$
(20)
$$([z], [y]) \mapsto [y] \circ_C [z] := [y \circ_C z]$$

Thus, in $\operatorname{CH}_2(J \times_C J) / \oplus_c \operatorname{CH}_2(J_c \times_c J_c)$,

$$\iota_{XJ}([y]) \circ_k \iota_{JX}([z]) = [\iota_{XJ}(y)] \circ_k [\iota_{JX}(z)] \stackrel{(19)}{=} [\iota_{XJ}(y) \circ_k \iota_{JJ}(z)]$$
$$= [\iota_{JJ}(y \circ_C z)] = \iota_{JJ}([y \circ_C z]) \stackrel{(20)}{=} \iota_{JJ}([y] \circ_C [z])$$

Here, the third equality uses Lemma 3.7 for B' = C and B = Spec(k). So, we get (i). The proof of (ii) is similar to (i) (Use Lemma 3.6 for U = Spec(k(C)) and B = C).

Now, we prove the following main proposition in this proof:

Proposition 8.6. Use above notations. Then in $CH_2(J \times J)/CH_2(J \times J)_{\equiv}$,

$$[\alpha] \circ_k [\beta] = [\Delta_J].$$

Proof. In
$$\operatorname{CH}_2(J \times J)/\operatorname{CH}_2(J \times J)_{\equiv}$$
,

$$[\alpha] \circ_k [\beta] = \iota_{XJ}(r_{XJ}^{-1}(a)) \circ_k \iota_{JX}(r_{JX}^{-1}(b)) \qquad \text{by definition}$$

$$= \iota_{JJ}(r_{XJ}^{-1}(a) \circ_C r_{JX}^{-1}(b)) \qquad \text{by Lemma 8.5 (i)}$$

$$= \iota_{JJ}(r_{JJ}^{-1}(a \circ_\eta b)) \qquad \text{by Lemma 8.5 (ii)}$$

$$= \iota_{JJ}(r_{JJ}^{-1}(\Delta_{J\eta})) \qquad \text{by } a \circ_\eta b = \Delta_{J\eta}$$

$$= [\Delta_J].$$

By Proposition 8.6, we get $[\alpha] \circ_k [\beta] = [\Delta_J]$ in $\operatorname{CH}_2(J \times J)/\operatorname{CH}_2(J \times J)_{\equiv}$. Similarly, $[\beta] \circ_k [\alpha] = [\Delta_X]$ in $\operatorname{CH}_2(X \times X)/\operatorname{CH}_2(X \times X)_{\equiv}$. Thus, we get

$$t_2(X) \cong t_2(J)$$
 in $\operatorname{CH}\mathcal{M}(k, \mathbb{Q})$.

Therefore, we complete the proof of Theorem 8.2 for the case where f is elliptic. Therefore, we complete the proof of Theorem 8.2, and hence of Theorem 8.1.

Remark 8.7. Taking the étale realization of $t_2(X) \cong t_2(J)$, we have an isomorphism $H^2_{\acute{e}t}(X, \mathbb{Q}_l)_{tr} \cong H^2_{\acute{e}t}(J, \mathbb{Q}_l)_{tr}$, so get $h^2_{\acute{e}t}(X, \mathbb{Q}_l)_{tr} = h^2_{\acute{e}t}(J, \mathbb{Q}_l)_{tr}$.

Remark 8.8. Thm. 8.2 is proved for the case where X is an Enriques surface with an elliptic fibration [Coo92] (see Prop. 1.2). Then we have $b_2(X) = 10$ by Def. 10.5, so $b_1(X) = 0$ by Prop. 10.4, and hence $(\operatorname{Pic}^0_{X/k})_{red} = 0$ by Prop. 2.4 (i). Thus, we get $h_1(X) = h_3(X) = 0$ by the argument as in the proof of Thm. 7.17.

Remark 8.9. As studied in [Huy16, Ch. 16] the case of elliptic K3 surfaces, it is expected that there is an equivalence between bounded derived categories of coherent sheaves $D^{b}(X) \cong D^{b}(J)$. This again induces $h(X) \cong h(J)$ (by [FV21] or [Huy19]).

9. KIMURA-FINITENESS

Here we collect some properties of Kimura-finiteness for the reader's convenience.

Definition 9.1. Let \mathcal{C} be a \mathbb{Q} -linear pseudo-abelian tensor category (e.g. $CH\mathcal{M}(k,\mathbb{Q})$)

- (i) An object A of C is evenly finite if $\wedge^n(A) = 0$ for n large enough.
- (ii) An object A of C is oddly finite if $\operatorname{Sym}^n(A) = 0$ for n large enough.
- (iii) An object A of C is Kimura-finite if there is a decomposition $A = A_+ \oplus A_$ such that A_+ is even and A_- is odd.

Conjecture 9.2. ([Kim05] or [And04, Ch. 12]). Every Chow motive is Kimura-finite.

For example, the motives 1 and \mathbb{L} are Kimura-finite.

Proposition 9.3. Let k be a field.

- (i) The motive of any smooth projective curve over k is Kimura-finite.
- (ii) The motive of any abelian variety is Kimura-finite.
- (iii) Let M and N be Kimura-finite dimensional motives. Then $M \oplus N$ and $M \otimes N$ are Kimura-finite.
- (iv) Let $\pi: V \to W$ be a dominant morphism of smooth projective varieties over k. If h(V) is Kimura-finite, so is h(W).
- (v) (Birational invariant) Let X and Y be smooth projective surfaces over k which are birationally equivalent. If h(X) is Kimura-finite, so is h(Y).

Proof. See [Kim05]. In particular, (v) follows from Manin's blow-up formula [Man68]. \Box

Remark 9.4. We only mention the case of surfaces. Let $k = \overline{k}$ and $S \in \mathcal{V}(k)$ a surface. Then $h_i(S)$ $(i \neq 2)$ and $h_2^{alg}(S)$ are Kimura-finite. Indeed, $h_0(S) \cong 1$, $h_4(S) \cong \mathbb{L}^{\otimes 2}$, and $h_2^{alg}(S) \cong \mathbb{L}^{\oplus \rho(S)}$. Thus, h_0 , h_4 , and h_2^{alg} are Kimura-finite. By Prop. 4.2, $h_1(S) \cong h_1((\operatorname{Pic}_{S/k}^0)_{red})$. By Prop. 9.3 (ii), $h_1((\operatorname{Pic}_{S/k}^0)_{red})$ is Kimura finite, so is h_1 . Similarly, h_3 is also Kimura-finite. However, the Kimura-finiteness of $t_2(S)$ is unknown in general. At least the following are known to have Kimura-finiteness:

- (i) varieties of dimension ≤ 3 rationally dominated by products of curves ([Via17, Ex. 3.15]). In particular, the K3 surfaces ([Par88], [ILP09]).
- (ii) K3 surfaces with Picard number 19, 20 ([Ped12]).
- (iii) some K3 surfaces obtained as complete intersections ([Lat18], [BL23]).
- (iv) many examples of surfaces with $p_g = 0$ over \mathbb{C} ([GP02], [PW15]).

In the proof of Theorem 1.7, we will use:

Proposition 9.5. ([Kim05]) Let k be a field and let $C, D \in \mathcal{V}(k)$ be curves. Let G be a finite group which acts freely on $C \times D$. Let X be a surface which is birational to $C \times D/G$. Then h(X) is Kimura-finite.

Proof. It follows from Proposition 9.3. Indeed, both h(C) and h(D) are Kimura-finite by (i). Then $h(C \times D) = h(C) \otimes h(D)$ is Kimura-finite by (iii), so $h(C \times D/G)$ is Kimura-finite by (iv), and hence h(X) is Kimura-finite by (v).

Proposition 9.6. ([KMP07, Coro. 7.6.11, p.181]). Let $S \in \mathcal{V}(\mathbb{C})$ be a surface. Then the following properties are equivalent:

- (i) $a_S : CH_0(S)^0_{\mathbb{Z}} \cong Alb_{S/\mathbb{C}}(\mathbb{C});$
- (ii) $p_g(S) = 0$ and h(S) is Kimura-finite in $CH\mathcal{M}(\mathbb{C}, \mathbb{Q})$;

(iii) $t_2(S) = 0$.

10. Classification of Algebraic surfaces

Here we quickly review the classification of surfaces for the reader's convenience. In this section, let k be an algebraically closed field and $S \in \mathcal{V}(k)$ a surface.

10.1. Kodaira dimension. We define the Kodaira dimension of S to be

$$\kappa(S) := \begin{cases} -\infty & \text{if } P_m(S) = h^0(S, \omega_S^{\otimes m}) = 0 \text{ for every } m \ge 1 \\ \text{tr.deg}_k(\bigoplus_{m \ge 0} H^0(S, \omega_S^{\otimes m})) - 1 & \text{otherwise} \end{cases}$$

Then $\kappa = -\infty$, 0, 1, or 2. Since P_m is a birational invariant, so is κ . A surface S is minimal if and only if S do contain smooth rational curves E satisfying $(E^2) = (E \cdot K_S) = -1$. If $\kappa(S) \ge 0$, then S is minimal if and only if K_S is nef, i.e. $(K_S \cdot C) \ge 0$ for every curve C. We denote by \equiv the numerical equivalence of divisors. Recall:

Theorem 10.1. For any surface $S \in \mathcal{V}(k)$, there is a birational morphism $f : S \to S'$ onto a minimal surface $S' \in \mathcal{V}(k)$ that satisfies one of the following properties:

- (i) $\kappa(S') = -\infty, S' \cong \mathbb{P}^2$ or S' is a minimal ruled surface;
- (ii) $\kappa(S') = 0, \ (K_{S'}^2) = 0, \ K_{S'} \equiv 0;$
- (iii) $\kappa(S') = 1, \ (K_{S'}^2) = 0, \ K_{S'} \not\equiv 0;$
- (iv) $\kappa(S') = 2, (K_{S'}^2) > 0.$

In particular, if $\kappa(S) \ge 0$, S' is unique.

10.2. Surfaces not of general type. We focus on surfaces with $\kappa < 2$.

Theorem 10.2. ([Băd01, Thm. 13.2, p.195]). Let $S \in \mathcal{V}(k)$ be a surface. $\kappa(S) = -\infty$ if and only if S is birationally ruled, i.e. it is birational to $\mathbb{P}^1 \times C$ for some curve C.

Proposition 10.3. (e.g. [Băd01, Thm. 5.1, p.72]). Let $S \in \mathcal{V}(k)$ be a surface. Then $10 - 8q + 12p_q = (K_S^2) + b_2 + 2\Delta$,

where $\Delta = 2q - b_1$. Also, $\Delta = 0$ if char(k) = 0, and $0 \le \Delta \le 2p_g$ if char(k) > 0.

Using Proposition 10.3, we get the following:

Proposition 10.4. Let $S \in \mathcal{V}(k)$ be a minimal surface with $(K_S^2) = 0$ and $p_g \leq 1$.

(i) $b_2 = 22$, $b_1 = 0$, $\chi = 2$, q = 0, $p_g = 1$, $\Delta = 0$. (ii) $b_2 = 14$, $b_1 = 2$, $\chi = 1$, q = 1, $p_g = 1$, $\Delta = 0$. (iii) $b_2 = 10$, $b_1 = 0$, $\chi = 1$, q = 0, $p_g = 0$, $\Delta = 0$. (iv) $b_2 = 10$, $b_1 = 0$, $\chi = 1$, q = 1, $p_g = 1$, $\Delta = 2$. (v) $b_2 = 6$, $b_1 = 4$, $\chi = 0$, q = 2, $p_g = 1$, $\Delta = 0$. (vi) $b_2 = 2$, $b_1 = 2$, $\chi = 0$, q = 1, $p_g = 0$, $\Delta = 0$. (vii) $b_2 = 2$, $b_1 = 2$, $\chi = 0$, q = 2, $p_g = 1$, $\Delta = 2$.

In particular, any minimal surface with $\kappa = 0$ belongs to the above list.

Definition 10.5. A minimal surface $S \in \mathcal{V}(k)$ with $\kappa(S) = 0$.

• S is called an *Enriques surface* if $b_2(S) = 10$.

Theorem 10.6. ([Băd01, Thm. 10.17]). Any Enriques surface has a genus 1 fibration.

Proposition 10.7. (e.g. [Băd01, Thm. 8.6, p.113]). Let $S \in \mathcal{V}(k)$ be a minimal surface with $\kappa(S) = 0$. Let $alb_S : S \to Alb_{S/k}$ be the Albanese morphism of S. If $b_1(S) = 2$, then the morphism alb_S gives rise to a fibration $a : S \to E$ onto an elliptic curve E, all of whose fibers are integral curves of arithmetic genus one.

Definition 10.8. Let $S \in \mathcal{V}(k)$ be a minimal surface with $\kappa(S) = 0$ and $b_2(S) = 2$. Let $a: S \to E$ be the Albanese fibration of S as in Proposition 10.7.

- (i) S is hyper-elliptic if the generic fiber of a is smooth.
- (ii) S is quasi hyper-elliptic if the generic fiber of a is non-smooth.

To prove the Kimura-finiteness of hyper-elliptic surfaces, we need the following:

Proposition 10.9. ([BM77, Thm. 4, p.35]). Let S be a hyper-elliptic surface. Then $S \cong E \times F/G$,

where E and F are elliptic curves, and G is a finite subgroup scheme of E.

As far as I know, the motive of the following object is not well-understood:

Theorem 10.10. (e.g. [Băd01, Thm. 9.9, p.129]). Let $S \in \mathcal{V}(k)$ be a minimal surface. If $\kappa(S) = 1$, then S has a genus 1 fibration.

To prove the second main theorem, we need:

Proposition 10.11. (e.g. [Băd01, Rem. 8.3, p.112]). Let $f : X \to C$ be a minimal genus 1 fibration. Use the same notations as in the canonical bundle formula. We set

$$\lambda(f) := 2p_a(C) - 2 + \chi(\mathcal{O}_X) + \operatorname{length}(T) + \sum_{i=1}^r n_i / m_i.$$

- (i) $\lambda(f) < 0$ if and only if $\kappa(X) = -\infty$.
- (ii) $\lambda(f) = 0$ if and only if $\kappa(X) = 0$.
- (iii) $\lambda(f) > 0$ if and only if $\kappa(X) = 1$.

11. Chow motives of surfaces not of general type with $p_g = 0$

In this section, we prove the second main theorem of this paper (Theorem 11.3). Let k be an algebraically closed field and let $X \in \mathcal{V}(k)$ be a surface.

Let us recall the result of Bloch-Kas-Lieberman:

Theorem 11.1. ([BKL76]). Assume that $k = \mathbb{C}$, $p_g = 0$, and $\kappa < 2$. Then

$$a_X : \mathrm{CH}_0(X)^0_{\mathbb{Z}} \cong \mathrm{Alb}_{X/\mathbb{C}}(\mathbb{C})$$

By Proposition 9.6, Theorem 11.1 is equivalent to the following:

Theorem 11.2. Assume that $k = \mathbb{C}$, $p_g = 0$, and $\kappa < 2$. Then h(X) is Kimura-finite.

In this paper, we generalize Theorem 11.2 to arbitrary characteristic:

Theorem 11.3. Let X be a smooth projective surface over an algebraically closed field k. If $p_q(X) = 0$ and $\kappa(X) < 2$, then h(X) is Kimura-finite in $CH\mathcal{M}(k, \mathbb{Q})$.

Proof of Theorem 11.3. The ideas of the proof are based on [BKL76, Prop. 4, p.138], [GP02, Coro. 2.12, p.187], and [Voi03, Thm. 11.10, p.313].

(i) we assume $\kappa(X) < 0$. By Theorem 10.2, we have $X \sim_{birat} \mathbb{P}^1 \times C$ for some k-curve C. We apply Proposition 9.5 to G = 0, and see that h(X) is Kimura-finite.

So we assume $\kappa(X) \ge 0$. By Proposition 9.3 (v), we may assume X is minimal. Since $0 \le \kappa < 2$, we have $(K^2) = 0$ by Theorem 10.1. Then the Noether formula $10 - 8q + 12p_q = (K^2) + b_2 + 2\Delta$ becomes

$$10 - 8q = b_2.$$

Since $b_2 \ge 0$, we must consider the following two cases:

(a) $q(X) = 0, \ b_2(X) = 10;$ (b) $q(X) = 1, \ b_2(X) = 2.$

Lemma 11.4. Let $S \in \mathcal{V}(k)$ be a minimal surface. If $p_g = 0$ and $0 \le \kappa < 2$, then S has a genus 1 fibration.

Proof. First, we assume $\kappa = 0$. If $\kappa = p_g = q = 0$, then S is an Enriques surface, so S has a genus 1 fibration by Theorem 10.6. If $\kappa = p_g = 0$ and q = 1, then $b_2 = 2$ as in (b). By Definition 10.8, S is hyper-elliptic or quasi-hyper elliptic, so S has a genus 1 fibration. Next, we assume $\kappa = 1$. Then S has a genus 1 fibration by Theorem 10.10.

By Lemma 11.4, we see that X has a genus 1 fibration

$$f: X \to C.$$

Since f is a fibration, $q(X) \ge q(C) = p_a(C)$. Since $\chi = 1 - q + p_g = 1 + q$, by (a) and (b), we must consider the following two cases:

(a) $\chi(\mathcal{O}_X) = 1$, $p_a(C) = 0$, $b_2(X) = 10$; (b) $\chi(\mathcal{O}_X) = 0$, $p_a(C) \le 1$, $b_2(X) = 2$.

Let $j: J \to C$ be the Jacobian fibration of f. By Theorem 8.1, we get an isomorphism $h(X) \cong h(J)$ in $\operatorname{CH}\mathcal{M}(k, \mathbb{Q})$.

Thus, it suffices to prove that h(J) is Kimura-finite (In fact, we only use $t_2(X) \cong t_2(J)$).

(ii) we assume (a). So $\chi(\mathcal{O}_X) = 1$ and $p_a(C) = 0$. By Proposition 7.19 (i), $\chi(\mathcal{O}_J) = \chi(\mathcal{O}_X) = 1$. Thus,

$$\lambda(j) := 2p_a(C) - 2 + \chi(\mathcal{O}_J) = 0 - 2 + 1 = -1.$$

We apply Proposition 10.11 to $\lambda(j) = -1$, and get

$$\kappa(J) < 0.$$

By (i), we see that h(J) is Kimura-finite.

- (iii) we assume (b). So $\chi(\mathcal{O}_X) = 0$, $p_a(C) \leq 1$, and $b_2(X) = 2$.
- (iii-i) We prove that J is a hyper elliptic surface or a quasi hyper-elliptic surface.
- First, we prove $\kappa(J) \leq 0$. By Proposition 7.19 (i), $\chi(\mathcal{O}_J) = \chi(\mathcal{O}_X) = 0$. Thus,

$$\lambda(j) := 2p_a(C) - 2 + \chi(\mathcal{O}_J) \le 2 - 2 + 0 = 0.$$

We apply Proposition 10.11 to $\lambda(j) \leq 0$, and get

$$\kappa(J) \le 0.$$

By (i), we only consider the case where $\kappa(J) = 0$.

Next, we prove $b_2(J) = 2$. Now, $b_2(X) = 2$. By Proposition 7.19 (ii), we get $b_2(J) = b_2(X) = 2$. By Definition 10.8, J is hyper-elliptic or quasi hyper-elliptic.

(iii-ii) We prove that M(J) is Kimura-finite.

First, we assume that J is quasi hyper-elliptic. By Definition 10.8, J has a quasielliptic fibration $j': J \to C'$. Using Theorem 8.3, we have

$$t_2(J) = 0.$$

Then $h(J) \cong \bigoplus_{i=0,\neq 2}^{4} h_i(J) \oplus h_2^{alg}(J)$. By Remark 9.4, $h_i(J)(i \neq 2)$ and $h_2^{alg}(J)$ are Kimura-finite. By Proposition 9.3 (iii), we see that h(J) is Kimura-finite.

Next, we assume that J is hyper-elliptic. By Proposition 10.9, there are elliptic curves E, F, and a finite subgroup scheme G of E such that

$$J \cong (E \times F)/G$$

By Proposition 9.5, h(J) is Kimura-finite. This completes the proof of Theorem 11.3.

Remark 11.5. Let $f: X \to C$ be a genus 1 fibration with $p_g(X) = 0$, and $j: J \to C$ the Jacobian fibration of f. By the above arguments, $\kappa(X) \ge \kappa(J)$ (if X is an Enriques surface, then J is rational by the argument as in (ii), so $\kappa(X) = 0$ and $\kappa(J) < \infty$). "Roughly speaking", it is easier to prove the Kimura-finiteness of surfaces with low Kodaira dimension than that of surfaces with high Kodaira dimension.

Remark 11.6. Surfaces with $p_g = 0$:

	$\kappa = 0$	$\kappa = 1$
q = 0	Enriques surface	canonical fibration
q = 1	Albanese fibration	canonical fibration or Albanese fibration
_		

Let X be a surface with $p_g = 0$, $\kappa = 1$, and q = 1. Bloch-Kas-Lieberman considered the Albanese fibration $a: X \to \text{Alb}_{X/k}$. The genus of the generic fiber of a is ≥ 1 .

On the other hand, we consider the canonical fibration $f: X \to C$. The genus of the generic fiber of f is equal to 1. "Roughly speaking", it is easier to treat genus 1 fibrations than genus $g \ge 2$ fibrations, in positive characteristics.

Acknowledgements. First and foremost, I would like to thank my Ph.D. supervisor, Prof. Masaki Hanamura for many helpful comments and suggestions. I am deeply grateful to Prof. Takao Yamazaki for his comments, and giving me an opportunity to talk about this paper at Algebraic Seminar in Tohoku University. I would like to thank Prof. Nobuo Tsuzuki for his generous support, and the referee for very useful comments and suggestions. I express my appreciation for the hospitality of Research Alliance Center for Mathematical Sciences, Tohoku University, where this work was done.

References

- [And04] Y. André, Une introduction aux motifs (Motifs purs, motifs mixtes, périodes), Panoramas et Synthèses, vol. 17, Société Mathématique de France, Paris (2004).
- [Art86] M. Artin, Néron models, in Arithmetic geometry, eds G. Cornell and J. H. Silverman, Springer (1986), 213-230.
- [Băd01] L. Bădescu, Algebraic surfaces, Universitext, Springer (2001).
- [BKL76] S. Bloch, A. Kas, and D. Lieberman, Zero cycles on surfaces with $p_g = 0$, Compositio Math. **33** (1976), 135-145.
- [BL23] M. Bolognesi and R. Laterveer, A 9-dimensional family of K3 surfaces with finite dimensional motive, (2023) arXiv:2310.11981
- [BM77] E. Bombieri and D. Mumford, Enriques' classification of surfaces in Char. p: II, in Complex analysis and algebraic geometry, Iwanami Shoten, Tokyo (1977), 23–42.
- [Coo92] K. Coombes, The K-cohomology of Enriques surfaces, Contemp. Math. 126 (1992), 47-57.
- [CH00] A. Corti and M. Hanamura, Motivic decomposition and intersection Chow group I, Duke Math. J. vol.103 (2000), 459-522.
- [CD89] F. Cossec and I. Dolgachev, *Enriques surfaces I*, Birkhäuser (1989).
- [CDLK] F. Cossec, I. Dolgachev, C. Liedtke, with Appendix by S. Kondō, Enriques surfaces I, New edition, Preprint.
- [DM91] C. Deninger and J. P. Murre, Motivic decomposition of abelian schemes and the Fourier transform, J. Reine Angew. Math. 422 (1991), 201-219.
- [FV20] L. Fu and C. Vial, Distinguished cycles on varieties with motive of abelian type and the section property, J. Algebraic Geometry 29 (2020), 53–107.
- [FV21] L. Fu and C. Vial, A motivic global Torelli theorem for isogenous K3 surfaces, Advances in Mathematics 383 (2021).
- [Ful84] W. Fulton, Intersection theory, Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge, vol.2, second edition, Springer, Berlin (1998).
- [GS06] P. Gille and T. Szamuely, *Central simple algebras and Galois cohomology*, Cambridge Studies in Advanced Mathematics, vol.101, Cambridge University Press, Cambridge (2006).
- [G62V] A. Grothendieck, Technique de descente et théorèmes d'existence en géométrie algébrique. V. Les schémas de Picard: théorèmes d'existence, Séminaire N. Bourbaki, 1962, exp. no 232, 143-161.

- [G62VI] A. Grothendieck, Technique de descente et théorèmes d'existence en géométrie algébrique. VI. Les schémas de Picard: propriétés générales, Séminaire N. Bourbaki, 1962, exp. no 236, 221-243.
- [GP02] V. Guletskii and C. Pedrini, *The Chow motive of the Godeaux surface*, in Algebraic Geometry, a Volume in Memory of Paolo Francia, Walter de Gruyter (2002), 180-195.
- [Har77] R. Hartshorne, Algebraic Geometry, Graduate Texts in Mathematics, vol. 52, Springer, New York (1977).
- [Huy16] D. Huybrechts, Lectures on K3 surfaces, Cambridge University Press, Cambridge 2016.
- [Huy19] D. Huybrechts, Motives of isogenous K3 surfaces, Comment. Math. Helvetici, vol. 94 (2019), no.3, 445-458
- [ILP09] C. Ingalls, A. Logan and O. Patashnick, Explicit coverings of families of elliptic surfaces by squares of curves, arXiv:2009.07807
- [JY17] Z. Jiang and Q. Yin, On the Chow ring of certain rational cohomology tori, C. R. Acad. Sci. Paris, Ser. I 355 (2017), 571-576.
- [KMP07] B. Kahn, J. P. Murre, and C. Pedrini, On the transcendental part of the motive of a surface, in Algebraic cycles and Motives Vol II, London Mathematical Society Lectures Notes Series, vol. 344, Cambridge University Press, Cambridge (2008), 143-202.
- [Kaw22] D. Kawabe, Chow motives of quasi elliptic surfaces, (2022) arXiv:2201.06152, to appear in Osaka Journal of Math.
- [Kim05] S. Kimura, Chow groups are finite-dimensional, in some sense, Math. Ann. 331 (2005), 173-205.
- [Kle05] S. Kleiman, The Picard scheme, in Fundamental algebraic geometry : Grothendieck's FGA explained, Mathematical Surveys and Monographs, vol.123, American Mathematical Society (2005).
- [Lag59] S. Lang, *Abelian varieties*, Interscience, New York (1959).
- [LT58] S. Lang and J. Tate, Principal homogeneous spaces over abelian varieties, Amer. J. Math. 80 (1958), 659-684.
- [Lat18] R. Laterveer, A family of cubic fourfolds with finite-dimensional motive, Journal of the Mathematical Society of Japan, vol. 70 (2018), no. 4, 1453-1473.
- [Liu02] Q. Liu, Algebraic geometry and arithmetic curves, Oxford University Press (2002).
- [LLR04] Q. Liu, D. Lorenzini, and M. Raynaud, Néron models, Lie algebras, and reduction of curves of genus one, Invent. Math. 157 (2004), 455–518. [Corrigendum, Invent. Math. 214 (2018), 593–604].
- [LT16] Q. Liu and J. Tong, Néron models of algebraic curves, Trans. Amer. Math. Soc. 368 (2016), no. 10, 7019–7043.
- [Man68] Y. I. Manin, Correspondences, motives and monoidal transformations, Math. USSR. Sbornik 6 (1968), 439-470.
- [Mil80] J. S. Milne, *Étale cohomology*, Princeton University Press, Princeton, NJ (1980).
- [Mil86] J. S. Milne, *Jacobian varieties*, in *Arithmetic geometry*, eds G. Cornell and J. H. Silverman, Springer (1986) 167-212.
- [MI21] M. Miyanishi and H. Ito, Algebraic surfaces in positive characteristics: purely inseparable phenomena in curves and surfaces, World Scientific (2021).
- [Mum70] D. Mumford, Abelian varieties, Tata Institute of Fundamental Research Studies in Mathematics, Oxford University Press (1970).
- [Mur64] J. P. Murre, On contravariant functors from the category of preschemes over a field into the category of abelian groups (with an application to the Picard functor), Publ. Math. IHES. 23 (1964), 5-43.
- [Mur90] J. P. Murre, On the motive of an algebraic surface, J. Reine angew. Math. **409** (1990), 190–204.
- [Mur93] J. P. Murre. On a conjectural filtration on the Chow groups of an algebraic variety. Part I: The general conjectures and some examples, Indag. Math. (N.S.), 4(2):177–188, 1993.
- [MNP13] J. P. Murre, J. Nagel, and C. Peters, *Lectures on the theory of pure motives*, University Lecture Series, vol. 61, American Mathematical Society (2013).
- [Par88] K. Paranjape, Abelian varieties associated to certain K3 surfaces, Compositio Math. vol. 68 (1988), no.1, 11-22.

(*) DAIKI KAWABE

- [Ped12] C. Pedrini, On the finite dimensionality of a K3 surface, Manuscripta Math. 138 (2012), 59-72.
- [PW15] C. Pedrini and C. Weibel, Some surfaces of general type for which Bloch's conjecture holds, in: Period Domains, Algebraic Cycles, and Arithmetic, Cambridge Univ. Press, (2015).
- [Sch10] S. Schröer, On fibrations whose geometric fibers are nonreduced, Nagoya Math. J. 200 (2010), 35–57.
- [Ser60] J.-P. Serre, *Morphismes universels et variété d'Albanese*, Séminaire Claude Chevalley, tome 4 (1958-1959), exp. no. 10, 1–22.
- [Sil86] J. Silverman, *The arithmetic of elliptic curves*, Graduate Texts in Mathematics, vol. 106, second edition, Springer (2009).
- [Tat52] J. Tate, Genus change in inseparable extensions of function fields, Proc. Amer. Math. Soc. 3 (1952), 400–406.
- [Via17] C. Vial, Remarks on motives of abelian type, Tohoku Math. J. 69 (2017), 195-220.
- [Voi03] C. Voisin, *Hodge theory and complex algebraic geometry II*, Cambridge Studies in Advanced Mathematics, vol. 77, Cambridge University Press, Cambridge (2003).