

EFFECTIVE COUNTING AND SPIRALLING OF LATTICE APPROXIMATES

NATHAN HUGHES

ABSTRACT. Given $d \geq 2$, we show that the number of approximates $\frac{1}{q}\mathbf{p} \in \mathbb{Q}^d$ of $\mathbf{x} \in \mathbb{R}^d$ satisfying $|q\mathbf{x} - \mathbf{p}| \leq cq^{-\frac{1}{d}}$ with denominator $1 \leq q < T$ decays to the asymptotic term $\text{cvol}_d(B_d(0, 1)) \log T$ with error of order $(\log T)^{-\frac{1}{2}} (\log \log T)^{\frac{3}{2}} (\log \log \log T)^{\frac{1}{2} + \epsilon}$ for almost all $\mathbf{x} \in \mathbb{R}^d$ and for any $\epsilon > 0$. Results with the same order are proven for primitive lattice approximates for all $d \geq 1$ and also for the case of linear forms and affine lattices. These results, especially in the primitive case for $d = 1$, are an improvement to the results of Schmidt.

1. INTRODUCTION

In this paper, we show effective results for counting and spiralling of lattice approximates - namely, we give error rates for these results in Diophantine approximation. For spiralling, these error rates are new results, and new proofs are provided for the counting results, different from those provided by Schmidt [17, 18]. In particular, for the primitive case with $d = 1$ (see theorem 6), our result improves the error rate given by Schmidt. We also give effective results for Diophantine approximation of systems of linear forms and affine lattices.

1.1. Approximation of Real Vectors. Recall the following corollary to Dirichlet's Approximation Theorem.

Theorem 1 (Dirichlet [4]). *For each $d \geq 1$, there exists a constant $C = C(d) > 0$ such that, for all $\mathbf{x} \in \mathbb{R}^d$, there exists infinitely many $(\mathbf{p}, q) \in \mathbb{Z}^d \times \mathbb{N}$ satisfying*

$$\|q\mathbf{x} - \mathbf{p}\| < C|q|^{-\frac{1}{d}}$$

Here, $\|\cdot\|$ can be any norm on \mathbb{R}^d , so we assume it is the Euclidean norm for simplicity. We might ask how these approximates are distributed with respect to \mathbf{x} . More concretely, fix $\mathbf{x} \in \mathbb{R}^d$, let \mathbb{S}^{d-1} be the unit sphere in \mathbb{R}^d , and let

$$\theta_{\mathbf{x}} : \mathbb{R}^d \times \mathbb{R} \rightarrow \mathbb{S}^{d-1}, \quad \theta_{\mathbf{x}}(\mathbf{p}, q) = \frac{q\mathbf{x} - \mathbf{p}}{\|q\mathbf{x} - \mathbf{p}\|}$$

Then $\theta_{\mathbf{x}}(\mathbf{p}, q)$ measures the direction of the approximate (\mathbf{p}, q) relative to \mathbf{x} ; this is referred to as *spiralling of lattice approximates* [1]. Given measurable $A \subset \mathbb{S}^{d-1}$, we can ask what proportion of the directions of approximates $\theta_{\mathbf{x}}(\mathbf{p}, q)$ lie in A , and they showed that it is proportional to the volume of A in \mathbb{S}^{d-1} for almost all $\mathbf{x} \in \mathbb{R}^d$.

Theorem 2 (Athreya, Ghosh, Tseng [1, Theorem 1.1]). *Let $d \geq 1$ and $A \subset \mathbb{S}^{d-1}$ be measurable. For $T > 1$ and $\mathbf{x} \in \mathbb{R}^d$, define the sets*

$$N(\mathbf{x}, T) = \#\left\{(\mathbf{p}, q) \in \mathbb{Z}^d \times \mathbb{N} \mid 0 < q \leq T, \|q\mathbf{x} - \mathbf{p}\| < C|q|^{-\frac{1}{d}}\right\}$$

$$N(\mathbf{x}, T, A) = \#\left\{(\mathbf{p}, q) \in N(\mathbf{x}, T) \mid \theta_{\mathbf{x}}(\mathbf{p}, q) \in A\right\}$$

Then for almost every $\mathbf{x} \in \mathbb{R}^d$,

$$\lim_{T \rightarrow \infty} \frac{N(\mathbf{x}, T, A)}{N(\mathbf{x}, T)} = \text{vol}_{\mathbb{S}^{d-1}}(A)$$

This result was shown using the Siegel transform of a lattice, which links Euclidean space to the space of unimodular lattices, allowing the introduction of ergodic techniques (such as Birkhoff's ergodic theorem and Moore's ergodicity theorem) and is another example of the link between homogeneous dynamics and number theory. The purpose of this paper is to give an effective version of theorem 2 and related problems in Diophantine approximation.

1.2. Linear Forms and Affine Lattices. The problem of Diophantine approximation can be generalised to higher dimensions as follows. Let $d, m, n \geq 1$ be natural numbers satisfying $d = m + n$. Given $M \in \text{Mat}_{m,n}(\mathbb{R})$ and $C > 0$, we can ask how many pairs $(\mathbf{p}, \mathbf{q}) \in \mathbb{Z}^m \times \mathbb{Z}^n$ satisfy

$$(1) \quad \|M\mathbf{q} - \mathbf{p}\| < C \|\mathbf{q}\|^{-\frac{n}{m}}$$

Define

$$\Lambda_M = \begin{pmatrix} \text{Id}_m & M \\ \mathbf{0} & \text{Id}_n \end{pmatrix}$$

Then we have the equality of sets

$$\Lambda_M \mathbb{Z}^d = \left\{ \begin{pmatrix} M\mathbf{q} - \mathbf{p} \\ \mathbf{q} \end{pmatrix} \mid \mathbf{p} \in \mathbb{Z}^m, \mathbf{q} \in \mathbb{Z}^n \right\}$$

Define the area

$$R_{T,c} = \left\{ (\mathbf{x}, \mathbf{y}) \in \mathbb{R}^m \times \mathbb{R}^n \mid \|\mathbf{x}\|^m \|\mathbf{y}\|^n \leq c, 1 \leq \|\mathbf{y}\| < T \right\} \subset \mathbb{R}^d$$

We calculate $\text{vol}_d(R_{T,c}) = cB_m C_n \log T$, where B_m is the m -dimensional volume of the unit m -ball and C_n is the surface area of the unit sphere \mathbb{S}^{n-1} . The number of solutions to (1) is then equal to $\#(\Lambda_M \mathbb{Z}^d \cap R_{T,c})$ for some $c > 0$. This reformulation of a problem of Diophantine approximation to a problem in homogeneous dynamics is known as Dani's correspondence (see [9, Section 5.2c] for more details). Let X_d be the set of all unimodular lattices in \mathbb{R}^d . Then we may identify X_d with the quotient group $\text{SL}_d(\mathbb{R})/\text{SL}_d(\mathbb{Z})$ and assign a left- $\text{SL}_d(\mathbb{R})$ -invariant Haar measure $\mu = \mu_d$ onto it, which is unique up to scalar multiplication, induced by the unique measure with the same properties on $\text{SL}_d(\mathbb{R})$. Asymptotically, we can count the number of points of a unimodular lattice $\Lambda \in X_d$ contained in the region $R_{T,c}$ accordingly.

Theorem 3 (Athreya, Parrish, Tseng [2, Theorem 1.5]). *For all $c > 0$ and for μ -a.e. unimodular lattice $\Lambda \in X_d$,*

$$\lim_{T \rightarrow \infty} \frac{\#(\Lambda \cap R_{T,c})}{\text{vol}_d(R_{T,c})} = 1$$

A generalisation of this result for affine unimodular lattices is also possible. Given $\Lambda \in X_d$ and a non-zero vector $\xi \in \mathbb{R}^d/\Lambda$, $\Lambda + \xi$ is an affine unimodular lattice and we denote by Y_d the set of all such affine unimodular lattices. Similarly to X_d , we may further identify the space Y_d with the quotient space $\mathrm{SL}_d(\mathbb{R}) \ltimes \mathbb{R}^d / \mathrm{SL}_d(\mathbb{Z}) \ltimes \mathbb{Z}^d$ and endow Y_d with a natural Haar measure $\nu = \nu_d$. We have the following theorem related to theorem 3.

Theorem 4 (Athreya, Parrish, Tseng [2, Theorem 1.6]). *For all $c > 0$ and for ν -a.e. unimodular lattice $\Lambda \in Y_d$,*

$$\lim_{T \rightarrow \infty} \frac{\#((\Lambda + \xi) \cap R_{T,c})}{\mathrm{vol}_d(R_{T,c})} = 1$$

1.3. Statement of Results. In the following theorems, $f(x) = o_\gamma(g(x))$ means that there exists a constant $C(\gamma) > 0$ and $x_0 \geq 0$ such that for all $x > x_0$, $|f(x)| < C(\gamma)(g(x))$. Additionally, define

$$P_{T,c} = \left\{ (\mathbf{v}_1, v_2) \in \mathbb{R}^d \times \mathbb{R} \mid \|\mathbf{v}_1\| v_2 \leq c, 1 < v_2 \leq T \right\} \subset \mathbb{R}^{d+1}$$

$$P_{T,c,A} = \left\{ (\mathbf{v}_1, v_2) \in P_{T,c} \mid \frac{\mathbf{v}_1}{\|\mathbf{v}_1\|} \in A \right\}$$

Our first result is an effective version of theorem 2 for $d \geq 3$.

Theorem 5. *Let $d \geq 3$ and $A \subset \mathbb{S}^{d-2}$ be measurable. Then for any $\epsilon > 0$, for μ -a.e. $\Lambda \in X_d$ and for sufficiently large T ,*

$$\frac{\#(\Lambda \cap P_{T,c,A})}{\#(\Lambda \cap P_{T,c})} = \mathrm{vol}_{\mathbb{S}^{d-2}}(A) + o_{A,c,\Lambda,d}((\log T)^{-\frac{1}{2}}(\log \log T)^{\frac{3}{2}}(\log \log \log T)^{\frac{1}{2}+\epsilon})$$

A similar spiralling result can be obtained for all $d \geq 2$ by restricting the lattice to only its primitive elements. Given $\Lambda \in X_d$, define Λ^{pr} to be the set of all points $\mathbf{v} \in \Lambda$ such that $\alpha \mathbf{v} \notin \Lambda$ for all non-zero $|\alpha| \neq 1$.

Theorem 6. *Let $d \geq 2$, and let $A \subset \mathbb{S}^{d-2}$ be measurable. Then for any $\epsilon > 0$, for μ -a.e. $\Lambda \in X_d$ and for sufficiently large T ,*

$$\frac{\#(\Lambda^{\mathrm{pr}} \cap P_{T,c,A})}{\#(\Lambda^{\mathrm{pr}} \cap P_{T,c})} = \mathrm{vol}_{\mathbb{S}^{d-2}}(A) + o_{A,c,\Lambda,d}((\log T)^{-\frac{1}{2}}(\log \log T)^{\frac{3}{2}}(\log \log \log T)^{\frac{1}{2}+\epsilon})$$

For linear forms, we have the following effective version of theorem 3.

Theorem 7. *Let $d, m, n \geq 1$ be natural numbers satisfying $d = m + n$. Then for all $\epsilon > 0$, for μ -a.e. $\Lambda \in X_d$, and for sufficiently large T ,*

$$\#(\Lambda \cap R_{T,c}) = cB_m C_n \log T + o_{c,\Lambda,d}((\log T)^{-\frac{1}{2}}(\log \log T)^{\frac{3}{2}}(\log \log \log T)^{\frac{1}{2}+\epsilon})$$

Finally, for affine lattices, we have the following effective version of theorem 4.

Theorem 8. *Let $d, m, n \geq 1$ be natural numbers satisfying $d = m + n$. Then for all $\epsilon > 0$, for ν -a.e. $\Lambda + \xi \in Y_d$ and for sufficiently large T ,*

$$\#((\Lambda + \xi) \cap R_{T,c}) = cB_m C_n \log T + o_{c,\Lambda,\xi,d}((\log T)^{-\frac{1}{2}}(\log \log T)^{\frac{3}{2}}(\log \log \log T)^{\frac{1}{2}+\epsilon})$$

1.4. Structure of Paper. In section 2, we provide necessary background on the homogeneous spaces of unimodular lattices and integrals of some functions on these spaces. We employ the use of a theorem of Gaposhkin (theorem 11) on the relationship between the second moment of an ergodic sum and the convergence of said sum, introduced in section 3. In most cases, Rogers' theorem (theorem 10) is suitable to calculate the second moment, but this theorem fails to hold on X_2 . Section 4 is dedicated to using a theorem of Kleinbock and Yu (theorem 14) to calculate the second moment in the primitive two-dimensional case. In section 5, the results for counting lattice approximates for systems of linear forms and affine lattices are given, the latter using a result of El-Baz, Marklof and Vinogradov (lemma 2).

2. THE HOMOGENEOUS SPACE OF UNIMODULAR LATTICES

2.1. Mean Value Theorems on the Space of Unimodular Lattices. Let $d \geq 2$, $G_d = \mathrm{SL}_d(\mathbb{R})$, $\Gamma_d = \mathrm{SL}_d(\mathbb{Z})$ and consider the quotient space G_d/Γ_d . This space can be identified with the homogeneous space X_d via the mapping $g\Gamma_d \mapsto g\mathbb{Z}^d$, where $g \in G_d$. As shown in section 1.2, a vector $\mathbf{x} \in \mathbb{R}^{d-1}$ can be identified with a lattice $\Lambda_{\mathbf{x}} \in X_d$.

For $d \geq 2$, let $f : \mathbb{R}^d \rightarrow \mathbb{R}$ be a Riemann-integrable with compact support and $\Lambda \in X_d$. The Siegel transform of f , denoted \hat{f} , is defined as

$$\hat{f}(\Lambda) = \sum_{\mathbf{v} \in \Lambda \setminus \{\mathbf{0}\}} f(\mathbf{v})$$

A theorem of Siegel shows sufficient conditions for a Siegel transform to be integrable and provides a formula to calculate said integral. Let $d\mathbf{x}$ denote the Lebesgue measure on \mathbb{R}^d .

Theorem 9 ([19, Siegel], see also [12]). *Let $f \in L^1(\mathbb{R}^d, d\mathbf{x})$. Then $\hat{f} \in L^1(X_d, \mu)$ and*

$$\int_{\mathbb{R}^d} f(\mathbf{x}) d\mathbf{x} = \int_{X_d} \hat{f}(\Lambda) d\mu(\Lambda)$$

For our purposes, observe that for any measurable set $B \subset \mathbb{R}^d \setminus \{\mathbf{0}\}$,

$$\int_{X_d} \hat{\mathbb{1}}_B(\Lambda) d\mu(\Lambda) = \int_{\mathbb{R}^d} \mathbb{1}_B(\mathbf{x}) d\mathbf{x} = \mathrm{vol}_d(B)$$

Theorem 9 was partially generalised by Rogers, providing an equation for more general functions on X_d , where $d \geq 3$.

Theorem 10 (Rogers [16, Theorem 4] [12]). *Let $d \geq 3$, and suppose $\rho : (\mathbb{R}^d)^k \rightarrow \mathbb{R}$ is non-negative and Borel measurable, for some $1 \leq k \leq d-1$. Then*

$$(2) \quad \begin{aligned} & \int_{X_d} \sum_{x_1, \dots, x_k \in \Lambda} \rho(x_1, \dots, x_k) d\mu(\Lambda) = \rho(\mathbf{0}) + \int_{\mathbb{R}^d} \cdots \int_{\mathbb{R}^d} \rho(x_1, \dots, x_k) dx_1 \cdots dx_k + \\ & \sum_{(\nu; \mu) \in \mathcal{S}} \sum_{q=1}^{\infty} \sum_{D \in \Phi(\mathcal{S}, q)} \left(\frac{e_1}{q} \cdots \frac{e_m}{q} \right)^d \int_{\mathbb{R}^d} \cdots \int_{\mathbb{R}^d} \rho \left(\sum_{i=1}^m \frac{d_{i1}}{q} x_i, \dots, \sum_{i=1}^m \frac{d_{ik}}{q} x_i \right) dx_1 \cdots dx_m \end{aligned}$$

where \mathcal{S} is the set of all partitions of $\{1, \dots, k\}$ into two increasing sequences (ν_1, \dots, ν_m) , $(\mu_1, \dots, \mu_{m-k})$ for some $1 \leq m \leq k-1$, where $\Phi((\nu; \mu), q)$ is the set of all $D \in \mathrm{Mat}_{m,k}(\mathbb{Z})$

with elements having highest common factor relatively prime to q , with

$$\begin{aligned} d_{i\nu_j} &= q\delta_{ij} \\ d_{i\mu_j} &= 0 \quad \text{if } \mu_j < \nu_i \end{aligned}$$

for $1 \leq i \leq m$, $1 \leq j \leq k - m$, and where $e_i = \gcd(\epsilon_i, q)$ for $1 \leq i \leq m$, for elementary divisors ϵ_i of the matrix $D \in \Phi((\nu; \mu), q)$ and δ_{ij} is the Kronecker delta function.

In particular, this theorem allows us to calculate the integral of \hat{f}^k over X_d , for any $d \geq 3$ and $1 \leq k \leq d - 1$.

2.2. Second Moments of Characteristic Functions on the Space of Unimodular Lattices. Suppose (X, μ) is a probability space and let ϕ be an ergodic transformation on X . For $f \in L^2(X, \mu)$, the n^{th} correlation coefficient is defined as

$$b_n(f, \phi) = \int_X (f \circ \phi^n) f \, d\mu$$

We will calculate the decay of these correlations and employ theorem 11 to obtain the rate of convergence of an ergodic sum. Notice that

$$\left\| \sum_{k=0}^{N-1} f \circ \phi^k \right\|_2^2 = Nb_0(f, \phi) + 2 \sum_{k=1}^{N-1} (N - k) b_k(f, \phi) \in O(N^2 b_N(f, \phi))$$

so it suffices to calculate the asymptotic behaviour in n of the second moment of an ergodic sum.

Lemma 1. *Let $d \geq 3$ and $B \subset \mathbb{R}^d \setminus \{0\}$ be measurable. Then*

$$\int_{X_d} \hat{\mathbb{1}}_B(\Lambda)^2 \, d\mu(\Lambda) \leq (\text{vol}_d(B))^2 + O_d(\text{vol}_d(B))$$

Proof. We apply theorem 10 with $k = 2$ and $\rho(x, y) = \mathbb{1}_B(x)\mathbb{1}_B(y)$. Note that, since $\mathbf{0} \notin B$,

$$\sum_{x, y \in \Lambda} \mathbb{1}_B(x)\mathbb{1}_B(y) = \left(\sum_{x \in \Lambda} \mathbb{1}_B(x) \right) \left(\sum_{y \in \Lambda} \mathbb{1}_B(y) \right) = \hat{\mathbb{1}}_B(\Lambda)^2$$

Since $k = 2$, we have $m = 1$ and $\mathcal{S} = \{(1; 2), (2; 1)\}$. For each element of \mathcal{S} , any matrix D is of size 1×2 with coprime entries. For any $q \in \mathbb{N}$, if $D \in \Phi((1; 2), q)$ then $D = (q, r)$ for some $r \in \mathbb{Z}$ with $\gcd(q, r) = 1$. To compute the elementary divisor of D , note that equivalent matrices have the same elementary divisors [15]. Since $\gcd(q, r) = 1$, D is equivalent to $(1, 0)$, so 1 is the elementary divisor of D . Suppose now that $D \in \Phi((2; 1), q)$. Writing $D = (d_{11}, d_{22})$, we have $d_{12} = q$ and $d_{11} = 0$, but $\gcd(\gcd(0, q), q) = q$, so $\Phi((2; 1), q)$ is empty. We can bound

equation 2 by

$$\begin{aligned}
\int_{X_d} \widehat{\mathbb{1}}_B(\Lambda)^2 d\mu(\Lambda) &= \left(\int_{\mathbb{R}^d} \mathbb{1}_B(x) dx \right)^2 + \sum_{q=1}^{\infty} \sum_{\substack{r \in \mathbb{Z} \\ \gcd(q,r)=1}} \left(\frac{1}{q} \right)^d \int_{\mathbb{R}^d} \mathbb{1}_B(x) \mathbb{1}_B\left(\frac{rx}{q}\right) dx \\
&= (\text{vol}_d(B))^2 + \sum_{q=1}^{\infty} \sum_{\substack{r \in \mathbb{Z} \\ \gcd(q,r)=1}} \int_{\mathbb{R}^d} \mathbb{1}_B(qx) \mathbb{1}_B(rx) dx \\
&\leq (\text{vol}_d(B))^2 + \sum_{q=1}^{\infty} \sum_{r \in \mathbb{Z} \setminus \{0\}} \int_{\mathbb{R}^d} \mathbb{1}_B(qx) \mathbb{1}_B(rx) dx
\end{aligned}$$

Finally, using Hölder's inequality,

$$\begin{aligned}
&\sum_{q=1}^{\infty} \sum_{r \in \mathbb{Z} \setminus \{0\}} \int_{\mathbb{R}^d} \mathbb{1}_B(qx) \mathbb{1}_B(rx) dx \\
&\leq \sum_{q=1}^{\infty} \left(\int_{\mathbb{R}^d} \mathbb{1}_B(qx) dx \right)^{\frac{1}{2}} \sum_{r \in \mathbb{Z} \setminus \{0\}} \left(\int_{\mathbb{R}^d} \mathbb{1}_B(rx) dx \right)^{\frac{1}{2}} \\
&= \sum_{q=1}^{\infty} \left(\frac{1}{q^d} \int_{\mathbb{R}^d} \mathbb{1}_B(x) dx \right)^{\frac{1}{2}} \sum_{r \in \mathbb{Z} \setminus \{0\}} \left(\frac{1}{|r|^d} \int_{\mathbb{R}^d} \mathbb{1}_B(x) dx \right)^{\frac{1}{2}} \\
&= 2\zeta\left(\frac{d}{2}\right)^2 \text{vol}_d(B)
\end{aligned}$$

□

3. EFFECTIVE ERGODIC THEOREMS

3.1. Effective Theorem for Siegel Transforms. Our method of proving theorems 5, 6, 7 and 8 uses the following result derived from a theorem of Gaposhkin.

Theorem 11 (Gaposhkin [6, Theorem 3(iv)], Kachurovskii [8, Theorem 15(ii)]). *Let $f \in L^2(X, \mu)$ and let ϕ be an ergodic transformation of X . If*

$$(3) \quad \int_X \left(\sum_{n=0}^{N-1} \left(f \circ \phi^n - \int_X f d\mu \right) \right)^2 d\mu \in O_{f,\phi}(N)$$

Then for any $\epsilon > 0$, and μ -a.e. $x \in X$,

$$(4) \quad \frac{1}{N} \sum_{n=0}^{N-1} f(\phi^n(x)) = \int_X f d\mu + o_{f,\phi}(N^{-\frac{1}{2}}(\log N)^{\frac{3}{2}}(\log \log N)^{\frac{1}{2}+\epsilon})$$

This theorem was originally stated by Gaposhkin for mean-square continuous wide-sense stationary random processes, and later stated by Kachurovskii for functions $f \in L^2(X, \mu)$ satisfying $\int_X f d\mu = 0$. In the context of homogeneous spaces, this theorem has been used by Kleinbock, Shi and Weiss to give error terms for the convergence of the ergodic integral

$$\frac{1}{T} \int_0^T \phi(g_t u(\vartheta) \Lambda) dt = \int_X \phi d\mu + o\left(T^{-\frac{1}{2}} \log^{\frac{3}{2}+\epsilon} T\right)$$

for any smooth, compactly supported functions on X_d for $d \geq 2$ and for almost every $\vartheta \in \text{Mat}_{m,n}(\mathbb{R})$ (to be contrasted with the measure used in theorem 7), leading to a generalisation of theorem 3 in the non-equal weight case [10].

Proof of theorem 5. For $t \in \mathbb{R}$, define

$$g_t = \begin{pmatrix} e^t \text{Id}_{d-1} & \mathbf{0} \\ \mathbf{0}^T & e^{-t(d-1)} \end{pmatrix}$$

Then $\{g_t\}_{t \in \mathbb{R}}$ defines a flow on X_d via left translation: $g_t \cdot g\Gamma_d = (g_t g)\Gamma_d$. By Moore's Ergodicity Theorem [14, Theorem 3] (see also [3, §3.2 Theorem 2.1]), since the closure of $\{g_t\}_{t \in \mathbb{R}}$ is non-compact in G_d , this flow is ergodic.

Let $T > 1$, $N \in \mathbb{N}$ and set $s = \frac{\log T}{d-1}$. For any distinct integers $n > m \geq 1$, $g_s^n P_{T,c} \cap g_s^m P_{T,c} = \emptyset$ and

$$\bigcup_{k=m}^n g_s^{-k} P_{T,c} = P_{T^{n+1},c} \setminus P_{T^m,c}$$

In particular, we have

$$\begin{aligned} \bigcup_{k=0}^{N-1} g_s^{-k} P_{T,c} &= P_{T^N,c} \\ \sum_{k=0}^{N-1} \widehat{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) &= \widehat{\mathbb{1}}_{P_{T^N,c}}(\Lambda) \end{aligned}$$

By Siegel's Mean Value Theorem (Theorem 9), the function $\widehat{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) - \text{vol}_d(P_{T,c})$ is integrable for all $k \in \mathbb{N}$, and moreover,

$$\int_{X_d} \widehat{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) - \text{vol}_d(P_{T,c}) d\mu = 0$$

Rogers' theorem (theorem 10) implies that $\left(\widehat{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) - \text{vol}_d(P_{T,c})\right)^2$ is integrable, so we can calculate

$$\begin{aligned} & \int_{X_d} \left(\sum_{k=0}^{N-1} \widehat{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) - N \text{vol}_d(P_{T,c}) \right)^2 d\mu \\ &= \int_{X_d} \left[\left(\sum_{k=0}^{N-1} \widehat{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) \right)^2 - 2N \text{vol}_d(P_{T,c}) \sum_{k=0}^{N-1} \widehat{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) + N^2 \text{vol}_d(P_{T,c})^2 \right] d\mu \\ &= \int_{X_d} \left(\sum_{k=0}^{N-1} \widehat{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) \right)^2 d\mu - N^2 \text{vol}_d(P_{T,c})^2 \end{aligned}$$

Lemma 1 implies that this can be bounded as

$$\begin{aligned} & \int_{X_d} \left(\sum_{k=0}^{N-1} \widehat{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) \right)^2 d\mu - N^2 \text{vol}_d(P_{T,c})^2 \\ & \leq \left(\int_{\mathbb{R}^d} \sum_{k=0}^{N-1} \mathbb{1}_{P_{T,c}}(\mathbf{x}) d\mathbf{x} \right)^2 + O_d(N \text{vol}_d(P_{T,c})) - N^2 \text{vol}_d(P_{T,c})^2 \\ & = O_{T,c,d}(N) \end{aligned}$$

This shows condition (3) in theorem 11 to be true, so for any $\epsilon > 0$,

$$(5) \quad \frac{1}{N} \sum_{k=0}^{N-1} \widehat{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) = \frac{1}{N} \widehat{\mathbb{1}}_{P_{T^N,c}}(\Lambda) = \text{vol}_d(P_{T,c}) + o_{T,c,\Lambda,N,d}(N^{-\frac{1}{2}} (\log N)^{\frac{3}{2}} (\log \log N)^{\frac{1}{2}+\epsilon})$$

Given any $\tau > 2$ there exists unique $T \in [2, 4)$ and $N \in \mathbb{N}$ with $\tau = T^N$. Write $\tau = T^N > 1$. Then for any $\epsilon > 0$,

$$(6) \quad \frac{1}{\log \tau} \widehat{\mathbb{1}}_{P_{\tau,c}}(\Lambda) = cB_d + o_{T,c,\Lambda,d}((\log \tau)^{-\frac{1}{2}} (\log \log \tau)^{\frac{3}{2}} (\log \log \log \tau)^{\frac{1}{2}+\epsilon})$$

Similarly, for any measurable $A \subset \mathbb{S}^{d-2}$,

$$\frac{1}{\log \tau} \widehat{\mathbb{1}}_{P_{\tau,c,A}}(\Lambda) = cB_d \text{vol}_{\mathbb{S}^{d-2}}(A) + o_{A,T,c,\Lambda,d}((\log \tau)^{-\frac{1}{2}} (\log \log \tau)^{\frac{3}{2}} (\log \log \log \tau)^{\frac{1}{2}+\epsilon})$$

By varying T , we conclude

$$(7) \quad \frac{\widehat{\mathbb{1}}_{P_{\tau,c,A}}(\Lambda)}{\widehat{\mathbb{1}}_{P_{\tau,c}}(\Lambda)} = \text{vol}_{\mathbb{S}^{d-2}}(A) + o_{A,c,\Lambda,d}((\log \tau)^{-\frac{1}{2}} (\log \log \tau)^{\frac{3}{2}} (\log \log \log \tau)^{\frac{1}{2}+\epsilon})$$

□

Remark 1. In 1960, Schmidt studied a similar counting problem for lattice points of a (not necessarily unimodular) lattice Λ contained in a family of nested finite measurable sets $\Phi \ni S$ which contain arbitrarily large volumes [17]. He found that in dimensions 3 and greater, for any non-decreasing $\psi : [0, \infty) \rightarrow (0, \infty)$ such that $\int_0^\infty \psi(s)^{-1} ds$ exists, that

$$\#(\Lambda \cap S) = \text{vol}_d(S) + O(\text{vol}_d(S)^{\frac{1}{2}} \log \text{vol}_d(S) \psi^{\frac{1}{2}}(\log \text{vol}_d(S)))$$

Noting that $\text{vol}_d(P_{\tau,c,A}) = c \text{vol}_d(B_d) \text{vol}_d(A) \log \tau$, we may rewrite (7) as

$$(8) \quad \#(\Lambda \cap P_{\tau,c,A}) = \text{vol}_d(P_{\tau,c,A}) + o_{\Lambda,c,d}((\log \tau)^{\frac{1}{2}} (\log \log \tau)^{\frac{3}{2}} (\log \log \tau)^{\frac{1}{2}+\epsilon})$$

Then setting ψ to be $f(s) = (\log s)^3 (\log \log s)^{1+2\epsilon}$ in Schmidt's theorem would give (8), but notice that $\int_1^\infty (f(s))^{-1} ds$ does not converge. The result of theorem 5 is therefore a slight improvement to Schmidt's result in the specific case of $\Phi = \{P_{T,c,A} \mid T > 1\}$ for any $c > 0$ and $A \subset \mathbb{S}^{d-2}$. Additionally, Schmidt found that, in two dimensions,

$$\#(\Lambda \cap S) = \text{vol}_2(S) + O(\text{vol}_2(S)^{\frac{1}{2}} \log^2 \text{vol}_2(S) \psi^{\frac{1}{2}}(\log \text{vol}_2(S)))$$

We will see in section 4 that, in the case of $\Phi = \{P_{T,c,A} \mid T > 1\}$, this can be improved to the same bound as in the higher-dimensional case.

Remark 2. As Gaposhkin states in his work, any number of iterated logarithms of the form $(N)^{-\frac{1}{2}} (\log N)^{\frac{3}{2}} \cdots (\log \cdots \log N)^{\frac{1}{2}+\epsilon}$ may be used in 5 [6, §3]. In fact, any function $\Psi(t)$ that satisfies

$$\int_1^\infty \frac{1}{t\Psi(t)} dt < \infty$$

can be used to derive an error term. In 4, the function

$$\Psi(t) = \begin{cases} 1 & 1 \leq t \leq e + \delta \\ (\log t)(\log \log t)^{1+\epsilon} & t > e + \delta \end{cases}$$

was used, for any $\delta > 0$. Given such a function Ψ , the effective term will be of the form

$$o\left(\sqrt{\frac{\Psi(T)}{T}} \log T\right)$$

(see Lemma 7 in [6]).

3.2. Effective Theorem for Primitive Siegel Transforms. Recall that the primitive lattice Λ^{pr} is the set of all points $\mathbf{v} \in \Lambda$ such that $\alpha \mathbf{v} \notin \Lambda$ for all non-zero $|\alpha| \neq 1$. Define the primitive Siegel transform of a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ by

$$\tilde{f} : X_d \rightarrow \mathbb{R}, \quad \tilde{f}(\Lambda) = \sum_{\mathbf{v} \in \Lambda^{\text{pr}}} f(\mathbf{v})$$

We are interested in obtaining similar theorems to those obtained above with this different transform. Siegel provided the following formula for the primitive Siegel transform.

Theorem 12 ([19, Siegel], see also [12]). *Let $f \in L^1(\mathbb{R}^d, d\mathbf{x})$. Then $\tilde{f} \in L^1(X_d, \mu)$ and*

$$\frac{1}{\zeta(d)} \int_{\mathbb{R}^d} f(\mathbf{x}) d\mathbf{x} = \int_{X_d} \tilde{f}(\Lambda) d\mu(\Lambda)$$

In the same paper that [16, Theorem 4] was proved, Rogers also provided a moment formula for the primitive Siegel transform.

Theorem 13 ([16, Theorem 5]). *Let $d \geq 3$ and suppose $f : X_d \rightarrow \mathbb{R}_{\geq 0}$ is measurable. Then*

$$\begin{aligned} \int_{X_d} \tilde{f}(\Lambda) d\mu(\Lambda) &= \left(\frac{1}{\zeta(d)} \int_{\mathbb{R}^d} f(\mathbf{x}) d\mathbf{x} \right)^2 + \frac{1}{\zeta(d)} \int_{\mathbb{R}^d} (f(\mathbf{x}))^2 d\mathbf{x} \\ &\quad + \frac{1}{\zeta(d)} \int_{\mathbb{R}^d} f(\mathbf{x}) f(-\mathbf{x}) d\mathbf{x} \end{aligned}$$

Proof of theorem 6 for $d \geq 3$. We have, for $s = \frac{\log T}{d-1}$, and using theorem 12,

$$\begin{aligned} &\int_{X_d} \left(\sum_{k=0}^{N-1} \tilde{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) - \frac{N \text{vol}(P_{T,c})}{\zeta(d)} \right)^2 d\mu \\ &= \int_{X_d} \left(\sum_{k=0}^{N-1} \tilde{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) \right)^2 d\mu - \frac{2N \text{vol}(P_{T,c})}{\zeta(d)} \int_{X_d} \sum_{k=0}^{N-1} \tilde{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) d\mu + \left(\frac{N \text{vol}(P_{T,c})}{\zeta(d)} \right)^2 \\ &= \int_{X_d} \left(\sum_{k=0}^{N-1} \tilde{\mathbb{1}}_{P_{T,c}}(g_s^k \Lambda) \right)^2 d\mu - \left(\frac{N \text{vol}(P_{T,c})}{\zeta(d)} \right)^2 \\ &= \int_{X_d} \left(\tilde{\mathbb{1}}_{P_{T^N,c}}(\Lambda) \right)^2 d\mu - \left(\frac{\text{vol}(P_{T^N,c})}{\zeta(d)} \right)^2 \end{aligned}$$

Using Theorem 13,

$$\begin{aligned} &\int_{X_d} \left(\tilde{\mathbb{1}}_{P_{T^N,c}}(\Lambda) \right)^2 d\mu - \left(\frac{\text{vol}(P_{T^N,c})}{\zeta(d)} \right)^2 \\ &= \frac{1}{\zeta(d)^2} \left(\int_{\mathbb{R}^d} \mathbb{1}_{P_{T^N,c}}(\mathbf{x}) d\mathbf{x} \right)^2 + \frac{1}{\zeta(d)} \int_{\mathbb{R}^d} (\mathbb{1}_{P_{T^N,c}}(\mathbf{x}))^2 d\mathbf{x} \\ &\quad + \frac{1}{\zeta(d)} \int_{\mathbb{R}^d} \mathbb{1}_{P_{T^N,c}}(\mathbf{x}) \mathbb{1}_{P_{T^N,c}}(-\mathbf{x}) d\mathbf{x} - \left(\frac{\text{vol}(P_{T^N,c})}{\zeta(d)} \right)^2 \\ &= \frac{\text{vol}(P_{T^N,c})^2}{\zeta(d)^2} + \frac{\text{vol}(P_{T^N,c})}{\zeta(d)} - \left(\frac{\text{vol}(P_{T^N,c})}{\zeta(d)} \right)^2 \\ &= \frac{\text{vol}(P_{T^N,c})}{\zeta(d)} \\ &= O_{T,c,d}(N) \end{aligned}$$

A similar argument to that of theorem 5 yields, for any $A \subset \mathbb{S}^{d-2}$ with measurable boundary,

$$(9) \quad \tilde{\mathbb{1}}_{P_{T,c,A}}(\Lambda) = cB_{d-1} \text{vol}(A) \log T + o_{A,c,\Lambda,d}((\log T)^{-\frac{1}{2}} (\log \log T)^{\frac{3}{2}} (\log \log \log T)^{\frac{1}{2}+\epsilon})$$

$$(10) \quad \frac{\tilde{\mathbb{1}}_{P_{T,c,A}}(\Lambda)}{\tilde{\mathbb{1}}_{P_{T,c,A}}(\Lambda)} = \text{vol}(A) + o_{A,c,\Lambda,d}((\log T)^{-\frac{1}{2}} (\log \log T)^{\frac{3}{2}} (\log \log \log T)^{\frac{1}{2}+\epsilon})$$

□

4. THE TWO-DIMENSIONAL CASE

4.1. Outline of Proof. We wish to find a bound, asymptotic in T , for the expression

$$\begin{aligned} & \int_{X_2} \left(\tilde{\mathbb{1}}_{P_{T,c}}(\Lambda) - \frac{\text{area}(P_{T,c})}{\zeta(2)} \right)^2 d\mu(\Lambda) \\ &= \left\| \tilde{\mathbb{1}}_{P_{T,c}} \right\|_2^2 - 2 \frac{\text{area}(P_{T,c})}{\zeta(2)} \int_{X_2} \tilde{\mathbb{1}}_{P_{T,c}}(\Lambda) d\mu(\Lambda) + \frac{\text{area}(P_{T,c})^2}{\zeta(2)^2} \\ &= \left\| \tilde{\mathbb{1}}_{P_{T,c}} \right\|_2^2 - \frac{4c^2}{\zeta(2)^2} \log(T) \end{aligned}$$

Rogers' theorems (theorems 10 and 13) cannot be used to calculate this expression in either the full or primitive case of the Siegel transform, since these theorems do not hold in the case $d = 2$. In its stead, we will use the following theorem.

Theorem 14 ([11, Theorem 2.1]). *Let \mathcal{S} be a measurable and bounded subset of \mathbb{R}^2 , let $f = \mathbb{1}_{\mathcal{S}}$, and let $-\mathcal{S} = \{\mathbf{x} \in \mathbb{R}^2 \mid -\mathbf{x} \in \mathcal{S}\}$. Then*

$$\left\| \tilde{f} \right\|_2^2 = \frac{1}{\zeta(2)} \left(\text{area}(\mathcal{S}) + \text{area}(\mathcal{S} \cap -\mathcal{S}) + \sum_{n \neq 0} \frac{\varphi(|n|)}{|n|} \iint_{\mathcal{S}} \left| \mathcal{I}_{(x,y)}^n(\mathcal{S}) \right| dx dy \right)$$

where

$$\left| \mathcal{I}_{(x,y)}^n(\mathcal{S}) \right| = \left\{ t \in \mathbb{R} \mid \frac{n}{x^2 + y^2}(-y, x) + t(x, y) \in \mathcal{S} \right\}.$$

Using this theorem, we will show that $\tilde{\mathbb{1}}_{P_{T,c}} \in L^2(X_2, \mu)$, in which case we may apply theorem 11. To calculate $\left\| \tilde{\mathbb{1}}_{P_{T,c}} \right\|_2^2$, we first find an expression for $\left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right|$ in terms of the y-coordinates of the intersections of the line $\left\{ \left(\frac{n}{x^2 + y^2}(-y, x) + t(x, y) \right) \mid t \in \mathbb{R} \right\}$ with $\partial P_{T,c}$. The integral $\iint_{P_{T,c}} \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| dx dy$ can be decomposed into a sum of integrals whose integrands are related to these intersection's y-coordinates. Finally, the sum $\sum_{n \neq 0} \iint_{P_{T,c}} \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| dx dy$ is calculated by finding a suitable power series for $\left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right|$ and using [7, Theorem 421] to find the asymptotic growth of the sum.

4.2. Setting up the Integrals. Let $P_{T,c} = \{(x, y) \in \mathbb{R}^2 \mid |x|y < c, 1 < y \leq T\}$. First, we wish to calculate, for any $n \in \mathbb{Z} \setminus \{0\}$, $T > 1$, $c > 0$ and $(x, y) \in P_{T,c}$, the value of

$$\left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| = \left| \left\{ t \in \mathbb{R} \mid \left(\frac{-ny}{x^2 + y^2} + tx, \frac{nx}{x^2 + y^2} + ty \right) \in P_{T,c} \right\} \right|$$

where $|\cdot|$ is the one-dimensional Lebesgue measure. For any $n \neq 0$, $(x, y) \in P_{T,c}$, let

$$L(n, x, y, t) = \left(\frac{-ny}{x^2 + y^2} + tx, \frac{nx}{x^2 + y^2} + ty \right), \quad L(n, x, y) = \left\{ L(n, x, y, t) \mid t \in \mathbb{R} \right\}$$

Then $L(n, x, y)$ is an infinite line parameterised by $t \in \mathbb{R}$ intersecting the point $\frac{n}{x^2+y^2}(-y, x)$ with gradient $\frac{y}{x}$ (or a vertical line when $x = 0$). Suppose $n > 0$. Then

$$\frac{(-n) \pm \sqrt{(-n)^2 \pm 4xyc}}{2x} = \frac{n \mp \sqrt{n^2 \pm 4xyc}}{-2x} = \frac{n \mp \sqrt{n^2 \mp 4(-x)yc}}{2(-x)}$$

Furthermore,

$$\left(\frac{-(-n)y}{x^2+y^2} + t(-x), \frac{(-n)(-x)}{x^2+y^2} + ty \right) = \left(-\left(\frac{-ny}{x^2+y^2} + tx \right), \frac{nx}{x^2+y^2} + ty \right)$$

so, by the symmetry of $P_{T,c}$ around the y -axis, $|\mathcal{I}_{(x,y)}^n(P_{T,c})| = |\mathcal{I}_{(x,y)}^{-n}(P_{T,c})|$. Therefore, we need only examine the case that $n > 0$.

Fix $n \in \mathbb{N}^+$ and $c > 0$. Let $Q_c = \{(x, y) \in \mathbb{R}^2 \mid |xy| \leq c\}$. The line L must intersect ∂Q_c at either 2 or 4 points (the case that L intersects at 2 points and is tangent at a third point is covered by the latter case). The y -coordinate of these intersections, for $x \neq 0$, is given by

$$\frac{n + (-1)^a \sqrt{n^2 + (-1)^b 4xyc}}{2x}$$

where $a, b \in \{0, 1\}$. Note that when $x = 0$, there are two intersections with y -coordinate $\pm \frac{cy}{n}$. For ease of notation, define

$$\begin{aligned} y_1 &= \frac{n - \sqrt{n^2 + 4xyc}}{2x} \\ y_2 &= \frac{n - \sqrt{n^2 - 4xyc}}{2x} \\ y_3 &= \frac{n + \sqrt{n^2 - 4xyc}}{2x} \\ y_4 &= \frac{n + \sqrt{n^2 + 4xyc}}{2x} \end{aligned}$$

For $x > 0$, and when they exist, these can be ordered as

$$y_1 < 0 < y_2 < y_3 < y_4$$

Let t_i be the value of t such that the y -coordinate of $L(n, x, y, t_i)$ is y_i . When $n < 2c$, for $t \in [t_1, t_4]$, $L(n, x, y, t) \in Q_c$. When $n \geq 2c$, for $t \in [t_1, t_2] \cup [t_3, t_4]$, $L(n, x, y, t) \in Q_c$. When $x < 0$,

$$y_3 < y_4 < y_1 < 0 < y_2.$$

Similarly, $L(n, x, y, t) \in Q_c$ when $t \in [t_3, t_2]$ and $n < 2c$ or when $t \in [t_3, t_4] \cup [t_1, t_2]$ and $n \geq 2c$. We now wish to find the intersections of $L(n, x, y)$ with $\partial P_{T,c}$. Notice that if $(x, y) \in P_{T,c}$ then $y > 1$, so any $y_i < 0$ will not correspond to an intersection of $L(n, x, y)$ with $\partial P_{T,c}$. In particular, $y_1 < 0$ for all x , so we need not consider it in these calculations. In view of the above, the line $L(n, x, y)$ can intersect $\partial P_{T,c}$ in the following ways:

- (1) through the lines $\{y = 1\}$ and $\{y = T\}$,
- (2) through the line $\{y = 1\}$ and intersecting once more at a point with y -coordinate $y_2 \in [1, T]$,

- (3) through the line $\{y = 1\}$ and intersecting once more at a point with y -coordinate $y_4 \in [1, T]$,
- (4) through the line $\{y = T\}$ and intersecting once more at a point with y -coordinate $y_3 \in [1, T]$,
- (5) intersecting twice at points with y -coordinates $y_3, y_4 \in [1, T]$,
- (6) through the line $\{y = 1\}$ and intersecting three more times at points with y -coordinates $y_2, y_3, y_4 \in [1, T]$,
- (7) through the lines $\{y = 1\}$, $\{y = T\}$ and intersecting twice more at points with y -coordinates $y_2, y_3 \in [1, T]$,
- (8) no intersection with $\partial P_{T,c}$.

Suppose we have n, x, y such that $L(n, x, y)$ intersects $\partial P_{T,c}$ at two points $(\alpha_1, \beta_1), (\alpha_2, \beta_2)$ with $\beta_2 > \beta_1$. Then

$$(11) \quad \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| = \left(\frac{\beta_2}{y} - \frac{nx}{yx^2 + y^3} \right) - \left(\frac{\beta_1}{y} - \frac{nx}{yx^2 + y^3} \right) = \frac{\beta_2 - \beta_1}{y}$$

Similarly, if $L(n, x, y)$ intersects $\partial P_{T,c}$ at (α_i, β_i) for $i = 1, 2, 3, 4$ with $\beta_1 < \beta_2 < \beta_3 < \beta_4$, then

$$\left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| = \frac{\beta_4 - \beta_3 + \beta_2 - \beta_1}{y}$$

For each of the 8 statements about intersection above, denote by B_i the set of all $(x, y) \in P_{T,c}$ such that $L(n, x, y)$ intersects $\partial P_{T,c}$ as in statement i above. The collection $\{B_i\}$ is a partition of $P_{T,c}$. The integral we wish to calculate can therefore be written as

$$(12) \quad \iint_{P_{T,c}} \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| dx dy = \sum_{i=1}^7 \iint_{B_i} \frac{1}{y} (\delta_{T,i}T + \delta_{y_4,i}y_4 - \delta_{y_3,i}y_3 + \delta_{y_2,i}y_2 - \delta_{1,i}) dx dy$$

where $\delta_{\alpha,i} = 1$ when statement i includes an intersection of type $\alpha \in \{1, y_2, y_3, y_4, T\}$ and is 0 otherwise. For example, $\delta_{T,1} = \delta_{T,4} = \delta_{T,7} = 1$, and $\delta_{\alpha,8} = 0$ for all intersection types. To ease the calculations to come, we may use the linearity of the integral to split (12) into a sum of five integrals, each with integrand of the form $\frac{\alpha}{y}$. The domains of these integrals are given by $A_\alpha = \bigcup_{i:\delta_{\alpha,i}=1} B_i$. Using the example of T as before, $A_T = B_1 \cup B_4 \cup B_7$. Therefore,

$$(13) \quad \begin{aligned} \iint_{P_{T,c}} \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| dx dy &= \iint_{A_T} \frac{T}{y} dx dy + \iint_{A_{y_4}} \frac{y_4}{y} dx dy - \iint_{A_{y_3}} \frac{y_3}{y} dx dy \\ &\quad + \iint_{A_{y_2}} \frac{y_2}{y} dx dy - \iint_{A_1} \frac{1}{y} dx dy. \end{aligned}$$

Notice also that A_α is the subset of $P_{T,c}$ such that $L(n, x, y)$ has an intersection with $\partial P_{T,c}$ of type α . Explicitly, these sets are:

- A_1 : all (x, y) such that $L(n, x, y)$ intersects the line segment $\{(x, 1) \mid |x| < c\} \subset \partial P_{T,c}$,
- A_{y_2} : all (x, y) such that y_2 exists and $1 < y_2 < T$,
- A_{y_3} : all (x, y) such that y_3 exists and $1 < y_3 < T$,
- A_{y_4} : all (x, y) such that y_4 exists and $1 < y_4 < T$,
- A_T : all (x, y) such that $L(n, x, y)$ intersects the line segment $\{(x, T) \mid |x| < \frac{c}{T}\} \subset \partial P_{T,c}$.




FIGURE 1. An example of a point $(a, b) \in P_{T,c}$ and a line $L(n, a, b)$. For this line, we have $y_1 < 1 < y_2 < y_3 < T < y_4$, so $(a, b) \in A_1 \cap A_{y_2} \cap A_{y_3} \cap A_T$.

Some cases will contain sub-cases; whether or not $n < 2c$ affects the boundary conditions for all sets except A_{y_4} . To help calculate these areas, we calculate that the derivatives of the y_i with respect to x are

$$\begin{aligned} \frac{\partial}{\partial x} y_1 &= \frac{\frac{n^2+2xyc}{\sqrt{n^2+4xyc}} - n}{2x^2} \geq 0 \\ \frac{\partial}{\partial x} y_2 &= \frac{\frac{n^2-2xyc}{\sqrt{n^2-4xyc}} - n}{2x^2} \geq 0 \\ \frac{\partial}{\partial x} y_3 &= -\frac{\frac{n^2-2xyc}{\sqrt{n^2-4xyc}} + n}{2x^2} \leq 0 \\ \frac{\partial}{\partial x} y_4 &= -\frac{\frac{n^2+2xyc}{\sqrt{n^2+4xyc}} + n}{2x^2} \leq 0 \end{aligned}$$

A_1 : Suppose $x > 0$. If $x > \frac{n^2}{4yc}$, then y_2, y_3 do not exist, therefore the line L intersects Q_c at two points, each point with y -coordinate $y_1 < 0 < y_4$. In this case, L intersects the line $\{(x, 1) \mid |x| < c\}$ when $y_4 \geq 1$, so $x < n + cy$. Note that the line $x = n + cy$ lies outside of $P_{T,c}$. When $0 < x \leq \frac{n^2}{4yc}$, both y_2 and y_3 exist. In this case, the line L leaves $P_{T,c}$ at y_2 and re-enters at y_3 . So L intersects $\{(x, 1) \mid |x| < c\}$ when $y_2 > 1$ or when $y_3 < 1$, hence $x > n - cy$. Finally, when $x < 0$, only $y_2 > 0$, so we require that

$y_2 > 1$, hence $x > n - cy$. If $n \leq 2c$, the integral is

$$\iint_{A_1} \frac{1}{y} dx dy = \int_1^T \int_{\max\{-\frac{c}{y}, n-cy\}}^{\frac{c}{y}} \frac{1}{y} dx dy$$

otherwise,

$$\iint_{A_1} \frac{1}{y} dx dy = \int_{\frac{n+\sqrt{n^2-4c^2}}{2c}}^T \int_{\max\{-\frac{c}{y}, n-cy\}}^{\frac{c}{y}} \frac{1}{y} dx dy$$

This integral is only valid for $n \leq \frac{c(T^2+1)}{T}$.




FIGURE 2. The area $A_1 \subset P_{T,c}$ such that any $(x, y) \in A_1$ has $L(x, n, y)$ intersecting $\partial P_{T,c} \cap \{y = 1\}$. The parameters $n = c = 1$ and $T = 10$ are used.

A_{y_2} : y_2 exists when $x < \frac{n^2}{4yc}$. Then

$$\begin{aligned} \frac{n - \sqrt{n^2 - 4xyc}}{2x} &= 1 \\ \sqrt{n^2 - 4xyc} &= n - 2x \\ n^2 - 4xyc &= n^2 - 4xn + 4x^2 \\ x &= n - cy \end{aligned}$$

Substituting this into the expression for y_2 gives

$$\frac{n - \sqrt{n^2 - 4x \left(\frac{n-x}{c}\right) c}}{2x} = \frac{n - |n - 2x|}{2x} = \begin{cases} 1 & x \leq \frac{n}{2} \\ 1 - \frac{n}{x} & x > \frac{n}{2} \end{cases}$$

The point with x -coordinate $\frac{n}{2}$ on the line $x = n - cy$ has y -coordinate $\frac{n}{2c}$, and $(\frac{n}{2}, \frac{n}{2c}) \in P_{T,c}$ if and only if $n = 2c$, corresponding to the point $(c, 1) \in \partial P_{T,c}$. The line $x = n - cy$ intersects $x = \frac{c}{y}$ at $y = \frac{n \pm \sqrt{n^2 - 4c^2}}{2c}$, which only exist when $n \geq 2c$.

$$\begin{aligned} \frac{n - \sqrt{n^2 - 4xyc}}{2x} &= T \\ \sqrt{n^2 - 4xyc} &= n - 2Tx \\ n^2 - 4xyc &= n^2 - 4Txn + 4T^2x^2 \\ x &= \frac{Tn - cy}{T^2} \end{aligned}$$

Substituting this into the expression for y_2 gives

$$\frac{n - \sqrt{n^2 - 4x \left(\frac{Tn - T^2x}{c} \right) c}}{2x} = \frac{n - |n - 2Tx|}{2x} = \begin{cases} T & x \leq \frac{n}{2T} \\ \frac{n}{x} - T & x \geq \frac{n}{2T} \end{cases}$$

The point with x -coordinate $\frac{n}{2T}$ on the line $x = \frac{Tn - cy}{T^2}$ has y -coordinate $\frac{nT}{2c}$. Since $\frac{\partial}{\partial x} y_2 \geq 0$, and for $n \leq 2c$, this integral is of the form

$$\begin{aligned} \iint_{A_{y_2}} \frac{n - \sqrt{n^2 - 4xyc}}{2xy} dx dy &= \int_1^{\frac{nT}{2c}} \int_{\max\{n - cy, -\frac{c}{y}\}}^{\frac{n^2}{4yc}} \frac{n - \sqrt{n^2 - 4xyc}}{2xy} dx dy \\ &\quad + \int_{\frac{nT}{2c}}^T \int_{-\frac{c}{y}}^{\frac{Tn - cy}{T^2}} \frac{n - \sqrt{n^2 - 4xyc}}{2xy} dx dy \end{aligned}$$

If $n > 2c$, then $(\frac{n}{2T}, \frac{nT}{2c}) \notin P_{T,c}$, so the integral is of the form

$$\iint_{A_{y_2}} \frac{n - \sqrt{n^2 - 4xyc}}{2xy} dx dy = \int_{\frac{n + \sqrt{n^2 - 4c^2}}{2c}}^T \int_{\max\{n - cy, -\frac{c}{y}\}}^{\frac{c}{y}} \frac{n - \sqrt{n^2 - 4xyc}}{2xy} dx dy$$

This integral is only valid for $n \leq \frac{c(T^2 + 1)}{T}$.




FIGURE 3. The area $A_{y_2} \subset P_{T,c}$ such that any $(x, y) \in A_{y_2}$ has $1 \leq y_2 \leq T$. The parameters $n = c = 1$ and $T = 10$ are used.

A_{y_3} : y_3 exists when $x \neq 0$, $x < \frac{n^2}{4yc}$ and $y_3 > 0$ only when $x > 0$. Then

$$\begin{aligned} \frac{n + \sqrt{n^2 - 4xyc}}{2x} &= 1 \\ \sqrt{n^2 - 4xyc} &= 2x - n \\ n^2 - 4xyc &= 4x^2 - 4xn + n^2 \\ x &= n - cy \end{aligned}$$

Substituting this into the expression for y_3 gives

$$\frac{n + \sqrt{n^2 - 4x \left(\frac{n-x}{c}\right) c}}{2x} = \frac{n + |n - 2x|}{2x} = \begin{cases} \frac{n}{x} - 1 & x \leq \frac{n}{2} \\ 1 & x > \frac{n}{2} \end{cases}$$

On the other hand,

$$\begin{aligned} \frac{n + \sqrt{n^2 - 4xyc}}{2x} &= T \\ \sqrt{n^2 - 4xyc} &= 2Tx - n \\ n^2 - 4xyc &= n^2 - 4Tnx + 4T^2x^2 \\ x &= \frac{Tn - cy}{T^2} \end{aligned}$$

and substituting this into the expression for y_3 gives

$$\frac{n + \sqrt{n^2 - 4x \left(\frac{Tn - T^2x}{c} \right) c}}{2x} = \frac{n + |n - 2Tx|}{2x} = \begin{cases} \frac{n}{x} - T & x \leq \frac{n}{2T} \\ T & x > \frac{n}{2T} \end{cases}$$

When $n \leq 2c$, this integral is of the form

$$\iint_{A_{y_3}} \frac{n + \sqrt{n^2 - 4xyc}}{2xy} dx dy = \int_1^{\frac{nT}{2c}} \int_{\frac{Tn-cy}{T^2}}^{\frac{n^2}{4yc}} \frac{n + \sqrt{n^2 - 4xyc}}{2xy} dx dy$$




FIGURE 4. The area $A_{y_3} \subset P_{T,c}$ such that any $(x, y) \in A_{y_3}$ has $1 \leq y_3 \leq T$. The parameters $n = c = 1$ and $T = 10$ are used.

When $n > 2c$, the line $x = \frac{Tn-cy}{T^2}$ intersects $x = \frac{c}{y}$ at $y = \frac{T(n - \sqrt{n^2 - 4c^2})}{2c}$, so the integral becomes

$$\iint_{A_{y_3}} \frac{n + \sqrt{n^2 - 4xyc}}{2xy} dx dy = \int_1^{\frac{T(n - \sqrt{n^2 - 4c^2})}{2c}} \int_{\frac{Tn-cy}{T^2}}^{\frac{c}{y}} \frac{n + \sqrt{n^2 - 4xyc}}{2xy} dx dy$$

This integral is only valid for $n \leq \frac{c(T^2+1)}{T}$.

A_{y_4} : y_4 is only positive when $x > 0$.

$$\begin{aligned}\frac{n + \sqrt{n^2 + 4xyc}}{2x} &= 1 \\ \sqrt{n^2 + 4xyc} &= 2x - n \\ n^2 + 4xyc &= 4x^2 - 4xn + n^2 \\ x &= n + cy\end{aligned}$$

For $c, n > 0$, this line does not intersect $P_{T,c}$.

$$\begin{aligned}\frac{n + \sqrt{n^2 + 4xyc}}{2x} &= T \\ \sqrt{n^2 + 4xyc} &= 2Tx - n \\ n^2 + 4xyc &= 4T^2x^2 - 4Txn + n^2 \\ x &= \frac{Tn + yc}{T^2}\end{aligned}$$

Substituting this into the expression for y_4 gives

$$\frac{n + \sqrt{n^2 + 4x \left(\frac{xT^2 - Tn}{c} \right) c}}{2x} = \frac{n + |n - 2xT|}{2x} = \begin{cases} T & x \geq \frac{n}{2T} \\ \frac{n}{x} - T & x < \frac{n}{2T} \end{cases}$$

The line $x = \frac{Tn+cy}{T^2}$ intersects the line $x = \frac{c}{y}$ at the point $(\frac{n+\sqrt{n^2+4c^2}}{2T}, \frac{T(-n+\sqrt{n^2+4c^2})}{2c})$. Therefore this integral is of the form

$$\iint_{A_{y_4}} \frac{n + \sqrt{n^2 + 4xyc}}{2xy} dx dy = \int_1^{\frac{T(-n+\sqrt{n^2+4c^2})}{2c}} \int_{\frac{nT+yc}{T^2}}^{\frac{c}{y}} \frac{n + \sqrt{n^2 + 4xyc}}{2xy} dx dy$$

This integral is only valid for $n \leq \frac{c(T^2-1)}{T}$.




FIGURE 5. The area $A_{y_4} \subset P_{T,c}$ such that any $(x, y) \in A_{y_4}$ has $1 \leq y_4 \leq T$. The parameters $n = c = 1$ and $T = 10$ are used.

A_T : If $n \leq 2c$, then the integral is

$$\iint_{A_T} \frac{T}{y} dx dy = \int_1^T \int_{\frac{Tn-cy}{T^2}}^{\min\{\frac{c}{y}, \frac{Tn+cy}{T^2}\}} \frac{T}{y} dx dy$$

otherwise,

$$\iint_{A_T} \frac{T}{y} dx dy = \int_1^{\frac{T(n-\sqrt{n^2-4c^2})}{2c}} \int_{\frac{Tn-cy}{T^2}}^{\min\{\frac{c}{y}, \frac{Tn+cy}{T^2}\}} \frac{T}{y} dx dy$$

This integral is only valid for $n \leq \frac{c(T^2+1)}{T}$.




FIGURE 6. The area $A_T \subset P_{T,c}$ such that any $(x, y) \in A_T$ has $L(n, x, y)$ intersecting $P_{T,c} \cap \{y = T\}$. The parameters $n = c = 1$ and $T = 10$ are used.

4.3. Calculating the Integrals. The following calculations have been verified using Wolfram Mathematica.

A_1 Case $n \leq 2c$

$$\begin{aligned} & \int_1^T \int_{\max\{-\frac{c}{y}, n-cy\}}^{\frac{c}{y}} \frac{1}{y} dx dy \\ &= -\frac{2c}{T} + \sqrt{n^2 + 4c^2} + n \log \left(\frac{\sqrt{n^2 + 4c^2} - n}{2c} \right) \\ &= O(1) \end{aligned}$$

Case $n > 2c$

$$\begin{aligned} & \int_{\frac{n+\sqrt{n^2-4c^2}}{2c}}^T \int_{\max\{-\frac{c}{y}, n-cy\}}^{\frac{c}{y}} \frac{1}{y} dx dy \\ &= -\frac{2c}{T} + \sqrt{n^2 + 4c^2} - \sqrt{n^2 - 4c^2} + n \log \left(\frac{\sqrt{n^2 + 4c^2} - n}{n - \sqrt{n^2 - 4c^2}} \right) \end{aligned}$$

A_{y_2} Case $n \leq 2c$

$$\begin{aligned}
& \int_1^{\frac{nT}{2c}} \int_{\max\{n-cy, -\frac{c}{y}\}}^{\frac{n^2}{4yc}} \frac{n - \sqrt{n^2 - 4xyc}}{2xy} dx dy + \int_{\frac{nT}{2c}}^T \int_{-\frac{c}{y}}^{\frac{Tn-cy}{T^2}} \frac{n - \sqrt{n^2 - 4xyc}}{2xy} dx dy \\
&= \sqrt{n^2 + 4c^2} - \sqrt{n^2 - 4c^2} + \frac{n}{2} \log \left(n + \sqrt{n^2 - 4c^2} \right)^2 - \frac{n}{2} \log \left(n + \sqrt{n^2 + 4c^2} \right)^2 \\
&\quad - \left(n + \sqrt{n^2 - 4c^2} - n \log \left(n + \sqrt{n^2 - 4c^2} \right) \right) \log \left(\frac{n + \sqrt{n^2 + 4c^2}}{n + \sqrt{n^2 - 4c^2}} \right) \\
&+ \left(\sqrt{n^2 + 4c^2} - \sqrt{n^2 - 4c^2} + n \log \left(\frac{n + \sqrt{n^2 - 4c^2}}{n + \sqrt{n^2 + 4c^2}} \right) \right) \log \left(\frac{T \left(\sqrt{n^2 + 4c^2} - n \right)}{2c} \right) \\
&= O(\log T)
\end{aligned}$$

Case $n > 2c$

$$\begin{aligned}
& \int_{\frac{n + \sqrt{n^2 - 4c^2}}{2c}}^T \int_{\max\{n-cy, -\frac{c}{y}\}}^{\frac{c}{y}} \frac{n - \sqrt{n^2 - 4xyc}}{2xy} dx dy \\
&= \left(n \log \left(n + \sqrt{n^2 - 4c^2} \right) - \sqrt{n^2 - 4c^2} - n \right) \log \left(\frac{n + \sqrt{n^2 + 4c^2}}{n + \sqrt{n^2 - 4c^2}} \right) \\
&\quad - \frac{n}{2} \log^2 \left(n + \sqrt{n^2 + 4c^2} \right) + \frac{n}{2} \log^2 \left(n + \sqrt{n^2 - 4c^2} \right) + \left(\sqrt{n^2 + 4c^2} - \sqrt{n^2 - 4c^2} \right) \\
&+ \left(n \log \left(\frac{n + \sqrt{n^2 - 4c^2}}{n + \sqrt{n^2 + 4c^2}} \right) + \sqrt{n^2 + 4c^2} - \sqrt{n^2 - 4c^2} \right) \left(\log T - \log \left(\frac{n + \sqrt{n^2 + 4c^2}}{2c} \right) \right)
\end{aligned}$$

A_{y_3} Case $n \leq 2c$

$$\begin{aligned}
& \int_1^{\frac{nT}{2c}} \int_{\frac{Tn-cy}{T^2}}^{\frac{n^2}{4yc}} \frac{n + \sqrt{n^2 - 4xyc}}{2xy} dx dy \\
&= \frac{n}{2} \log^2 \left(\frac{nT}{2c} \right) + O(\log T)
\end{aligned}$$

Case $n > 2c$

$$\begin{aligned}
& \int_1^{\frac{T(n - \sqrt{n^2 - 4c^2})}{2c}} \int_{\frac{Tn-cy}{T^2}}^{\frac{c}{y}} \frac{n + \sqrt{n^2 - 4xyc}}{2xy} dx dy \\
&= \left(n \log \left(n - \sqrt{n^2 - 4c^2} \right) + \sqrt{n^2 - 4c^2} - n \right) \log \left(\frac{T(n - \sqrt{n^2 - 4c^2})}{2c} \right) \\
&\quad - \frac{n}{2} \log^2 \left(n - \sqrt{n^2 - 4c^2} \right) + n - \sqrt{n^2 - 4c^2} + \frac{n}{2} \log^2 \left(\frac{2c}{T} \right) - \frac{2c}{T}
\end{aligned}$$

A_{y_4}

$$\begin{aligned}
& \int_1^{\frac{T(\sqrt{n^2+4c^2}-n)}{2c}} \int_{\frac{nT+yc}{T^2}}^{\frac{c}{y}} \frac{n + \sqrt{n^2 + 4xyc}}{2xy} dx dy \\
&= \left(n \log \left(\sqrt{n^2 + 4c^2} - n \right) + \sqrt{n^2 + 4c^2} - n \right) \log \left(\frac{T(\sqrt{n^2 + 4c^2} - n)}{2c} \right) \\
&\quad + n - \sqrt{n^2 + 4c^2} - \frac{1}{2} n \log^2 \left(\sqrt{n^2 + 4c^2} - n \right) + \frac{2c}{T} + \frac{n}{2} \log^2 \left(\frac{2c}{T} \right) \\
&= \frac{n}{2} \log^2 \left(\frac{2c}{T} \right) + O(\log T)
\end{aligned}$$

 A_T Case $n \leq 2c$

$$\begin{aligned}
& \int_1^T \int_{\frac{Tn-cy}{T^2}}^{\min\left\{\frac{c}{y}, \frac{Tn+cy}{T^2}\right\}} \frac{T}{y} dx dy \\
&= \sqrt{n^2 + 4c^2} - \frac{2c}{T} + n \log \left(\frac{\sqrt{n^2 + 4c^2} - n}{2c} \right) \\
&= O(1)
\end{aligned}$$

Case $n > 2c$

$$\begin{aligned}
& \int_1^{\frac{T(n-\sqrt{n^2-4c^2})}{2c}} \int_{\frac{Tn-cy}{T^2}}^{\min\left\{\frac{c}{y}, \frac{Tn+cy}{T^2}\right\}} \frac{T}{y} dx dy \\
&= \sqrt{n^2 + 4c^2} - \sqrt{n^2 - 4c^2} - \frac{2c}{T} + n \log \left(\frac{\sqrt{n^2 + 4c^2} - n}{n - \sqrt{n^2 - 4c^2}} \right)
\end{aligned}$$

We now calculate $\iint_{P_{T,c}} \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| dx dy$ using equation (13) and the calculations of the previous section.

When $n \leq 2c$ we can bound this integral by

$$\begin{aligned}
& O(1) + \frac{n}{2} \log^2 \left(\frac{2c}{T} \right) + O(\log T) - \frac{n}{2} \log^2 \left(\frac{nT}{2c} \right) + O(\log T) + O(\log T) - O(1) \\
&= O(\log T)
\end{aligned}$$

When $n > 2c$, the summand in this case is equal to

$$\begin{aligned} \log T \left(n \log \left(\frac{(\sqrt{n^2 + 4c^2} - n)(n + \sqrt{n^2 - 4c^2})}{(n + \sqrt{n^2 + 4c^2})(n - \sqrt{n^2 - 4c^2})} \right) + 2(\sqrt{n^2 + 4c^2} - \sqrt{n^2 - 4c^2}) \right) \\ + n(1 + \log(2c)) \log \left(\frac{(n + \sqrt{n^2 - 4c^2})(n - \sqrt{n^2 - 4c^2})}{(\sqrt{n^2 + 4c^2} - n)(n + \sqrt{n^2 + 4c^2})} \right) \\ + \sqrt{n^2 + 4c^2} \log \left(\frac{\sqrt{n^2 + 4c^2} - n}{\sqrt{n^2 + 4c^2} + n} \right) - \sqrt{n^2 - 4c^2} \log \left(\frac{n - \sqrt{n^2 - 4c^2}}{n + \sqrt{n^2 - 4c^2}} \right) + \frac{4c}{T} \\ + \frac{n}{2} \left(\log^2(\sqrt{n^2 + 4c^2} - n) + \log^2(\sqrt{n^2 + 4c^2} + n) \right. \\ \left. - \log^2(n + \sqrt{n^2 - 4c^2}) - \log^2(n - \sqrt{n^2 - 4c^2}) \right) \end{aligned}$$

Notice that

$$\frac{(n + \sqrt{n^2 - 4c^2})(n - \sqrt{n^2 - 4c^2})}{(\sqrt{n^2 + 4c^2} - n)(n + \sqrt{n^2 + 4c^2})} = 1$$

so the summand is equal to

$$\begin{aligned} \log T \left(n \log \left(\frac{(\sqrt{n^2 + 4c^2} - n)(n + \sqrt{n^2 - 4c^2})}{(n + \sqrt{n^2 + 4c^2})(n - \sqrt{n^2 - 4c^2})} \right) + 2(\sqrt{n^2 + 4c^2} - \sqrt{n^2 - 4c^2}) \right) \\ + \sqrt{n^2 + 4c^2} \log \left(\frac{\sqrt{n^2 + 4c^2} - n}{\sqrt{n^2 + 4c^2} + n} \right) - \sqrt{n^2 - 4c^2} \log \left(\frac{n - \sqrt{n^2 - 4c^2}}{n + \sqrt{n^2 - 4c^2}} \right) + \frac{4c}{T} \\ + \frac{n}{2} \left(\log^2(\sqrt{n^2 + 4c^2} - n) + \log^2(\sqrt{n^2 + 4c^2} + n) \right. \\ \left. - \log^2(n + \sqrt{n^2 - 4c^2}) - \log^2(n - \sqrt{n^2 - 4c^2}) \right) \end{aligned}$$

4.4. Calculating the Second Moment. Recall that

$$\left\| \tilde{\mathbb{I}}_{P_{T,c}} \right\|_2^2 = \frac{1}{\zeta(2)} \left(\text{area}(P_{T,c}) + \sum_{n \neq 0} \frac{\varphi(|n|)}{|n|} \iint_{P_{T,c}} |\mathcal{I}_{(x,y)}^n(P_{T,c})| \, dx \, dy \right)$$

Since $|\mathcal{I}_{(x,y)}^n(P_{T,c})| = |\mathcal{I}_{(x,y)}^{-n}(P_{T,c})|$, we can rewrite the sum as

$$\sum_{n \neq 0} \frac{\varphi(|n|)}{|n|} \iint_{P_{T,c}} |\mathcal{I}_{(x,y)}^n(P_{T,c})| \, dx \, dy = 2 \sum_{n \in \mathbb{N} \setminus \{0\}} \frac{\varphi(|n|)}{|n|} \iint_{P_{T,c}} |\mathcal{I}_{(x,y)}^n(P_{T,c})| \, dx \, dy$$

Therefore,

$$(14) \quad \left\| \tilde{\mathbb{1}}_{P_{T,c}} \right\|_2^2 = \frac{1}{\zeta(2)} \text{area}(P_{T,c}) + \frac{2}{\zeta(2)} \sum_{n \in \mathbb{N} \setminus \{0\}} \frac{\varphi(|n|)}{|n|} \iint_{P_{T,c}} \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| dx dy$$

We split the sum in (14) into the cases $n \leq 2c$ and $n > 2c$.

$$\begin{aligned} & \sum_{n \in \mathbb{N} \setminus \{0\}} \frac{\varphi(|n|)}{|n|} \iint_{P_{T,c}} \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| dx dy \\ = & \sum_{0 < n \leq 2c: n \in \mathbb{N}} \frac{\varphi(|n|)}{|n|} \iint_{P_{T,c}} \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| dx dy + \sum_{2c < n: n \in \mathbb{N}} \frac{\varphi(|n|)}{|n|} \iint_{P_{T,c}} \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| dx dy \end{aligned}$$

Then the first of these sums can be bounded as

$$\begin{aligned} & \sum_{0 < n \leq 2c: n \in \mathbb{N}} \frac{\varphi(|n|)}{|n|} \iint_{P_{T,c}} \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| dx dy \\ & \leq C + \sum_{n=1}^{[2c]} \frac{\varphi(n)}{n} O_c(\log T) \\ & = O_c(\log T) \end{aligned}$$

The second sum is as follows.

$$\begin{aligned} & \sum_{n=[2c]}^{[cT + \frac{c}{T}]} \frac{\varphi(n)}{|n|} \left[\log T \left(n \log \left(\frac{(\sqrt{n^2 + 4c^2} - n)(n + \sqrt{n^2 - 4c^2})}{(n + \sqrt{n^2 + 4c^2})(n - \sqrt{n^2 - 4c^2})} \right) \right. \right. \\ & \quad \left. \left. + \frac{4c}{T} + 2 \left(\sqrt{n^2 + 4c^2} - \sqrt{n^2 - 4c^2} \right) \right) + \sqrt{n^2 + 4c^2} \log \left(\frac{\sqrt{n^2 + 4c^2} - n}{n + \sqrt{n^2 + 4c^2}} \right) \right. \\ & \quad \left. + \sqrt{n^2 - 4c^2} \log \left(\frac{n + \sqrt{n^2 - 4c^2}}{n - \sqrt{n^2 - 4c^2}} \right) + \frac{n}{2} \left(\log^2 \left(\sqrt{n^2 + 4c^2} - n \right) + \log^2 \left(\sqrt{n^2 + 4c^2} + n \right) \right. \right. \\ & \quad \left. \left. - \log^2 \left(n + \sqrt{n^2 - 4c^2} \right) - \log^2 \left(n - \sqrt{n^2 - 4c^2} \right) \right) \right] - \epsilon_{y_4} \end{aligned}$$

Here, ϵ_{y_4} is equal to $\iint_{A_{y_4}} \frac{y_4}{y} dx dy$ evaluated at $n = [cT + \frac{c}{T}]$ and appears in the expression

since this particular integral does not validly contribute to the value of $\left\| \tilde{\mathbb{1}}_{P_{T,c}} \right\|_2^2$ for this value of n (as shown above, it is only valid for $n \leq \frac{c(T^2-1)}{T}$), hence we deduct its value from this sum. We may calculate, for $n = [cT + \frac{c}{T}]$, that

$$\lim_{T \rightarrow \infty} \iint_{A_{y_4}} \frac{y_4}{y} dx dy = 0$$

hence $\epsilon_{y_4} \rightarrow 0$. Since ϵ_{y_4} does not diverge as $T \rightarrow \infty$, we will ignore it when calculating the growth of the second sum.

4.5. Power Series for the Summands. We now wish to find suitable power series for each of the summands of the second sum which will allow us to calculate its asymptotic growth. Using

$$\begin{aligned}\sqrt{z+1} &= \sum_{k=0}^{\infty} \binom{\frac{1}{2}}{k} \frac{1}{z^{k-\frac{1}{2}}}, \quad x > 1 \\ \sqrt{z-1} &= \sum_{k=0}^{\infty} (-1)^k \binom{\frac{1}{2}}{k} \frac{1}{z^{k-\frac{1}{2}}}, \quad x > 1\end{aligned}$$

we calculate:

$$\begin{aligned}\sqrt{n^2 - 4c^2} &= 2c \cdot \sqrt{\frac{n^2}{4c^2} - 1} = \sum_{k=0}^{\infty} (-1)^k \binom{\frac{1}{2}}{k} \frac{4^k c^{2k}}{n^{2k-1}}, \quad n > 2c \\ \sqrt{n^2 + 4c^2} &= 2c \cdot \sqrt{\frac{n^2}{4c^2} + 1} = \sum_{k=0}^{\infty} \binom{\frac{1}{2}}{k} \frac{4^k c^{2k}}{n^{2k-1}}, \quad n > 2c\end{aligned}$$

To find a power series for the $\log(\sqrt{n^2 + 4c^2} - n)$, consider

$$\frac{\partial}{\partial n} \left(\log(\sqrt{n^2 + 4c^2} - n) - \log\left(\frac{2c^2}{n}\right) \right) = \frac{1}{n} - \frac{1}{\sqrt{n^2 + 4c^2}}$$

Recall that

$$(15) \quad \frac{1}{\sqrt{1-x^2}} = \sum_{k=0}^{\infty} \binom{2k}{k} \frac{1}{4^k} \frac{1}{x^{2k+1}}, \quad x > 1$$

$$(16) \quad \frac{1}{\sqrt{1+x^2}} = \sum_{k=0}^{\infty} \binom{2k}{k} \frac{(-1)^k}{4^k} \frac{1}{x^{2k+1}}, \quad x > 1$$

To calculate the power series for $n \log(\sqrt{n^2 + 4c^2} - n)$, notice that

$$\frac{\partial}{\partial n} \log\left(\frac{\sqrt{n^4 + 4c^2 n^2} - n^2}{2c^2}\right) = \frac{1}{n} - \frac{1}{\sqrt{n^2 + 4c^2}}$$

By equation (16),

$$\frac{1}{n} - \frac{1}{\sqrt{n^2 + 4c^2}} = \frac{2c^2}{n^3} - \frac{6c^4}{n^5} + \frac{20c^6}{n^7} - \frac{70c^8}{n^9} \dots$$

Integrating with respect to n yields

$$\log\left(\frac{\sqrt{n^4 + 4c^2 n^2} - n^2}{2c^2}\right) = \lim_{n \rightarrow \infty} \log\left(\frac{\sqrt{n^4 + 4c^2 n^2} - n^2}{2c^2}\right) - \frac{c^2}{n^2} + \frac{3c^4}{2n^4} - \frac{10c^6}{3n^6} + \frac{35c^8}{4n^8} - \dots$$

Since $\lim_{n \rightarrow \infty} \log \left(\frac{\sqrt{n^4 + 4c^2 n^2} - n^2}{2c^2} \right) = 0$, rearranging this equation and multiplying by n gives the power series

$$n \log \left(\sqrt{n^2 + 4c^2} - n \right) = n \log \left(\frac{2c^2}{n} \right) - \frac{c^2}{n} + \frac{3c^4}{2n^3} - \frac{10c^6}{3n^5} + \frac{35c^8}{4n^7} - \dots$$

Similarly, using (15) and (16),

$$n \log \left(n + \sqrt{n^2 - 4c^2} \right) = \log(2n) - \frac{c^2}{n} - \frac{3c^4}{2n^3} - \frac{10c^6}{3n^5} - \frac{35c^8}{4n^7} + \dots, \quad n > 2c$$

$$n \log \left(n + \sqrt{n^2 + 4c^2} \right) = \log(2n) + \frac{c^2}{n} - \frac{3c^4}{2n^3} + \frac{10c^6}{3n^5} - \frac{35c^8}{4n^7} + \dots, \quad n > 2c$$

$$n \log \left(n - \sqrt{n^2 - 4c^2} \right) = \log \left(\frac{2c^2}{n} \right) + \frac{c^2}{n} + \frac{3c^4}{2n^3} + \frac{10c^6}{3n^5} + \frac{35c^8}{4n^7} + \dots, \quad n > 2c$$

Using these equations, we find the following power series:

$$\begin{aligned} n \log \left(\frac{(\sqrt{n^2 + 4c^2} - n)(n + \sqrt{n^2 - 4c^2})}{(n + \sqrt{n^2 + 4c^2})(n - \sqrt{n^2 - 4c^2})} \right) &= -\frac{4c^2}{n} - \frac{40c^6}{3n^5} + \dots \\ \sqrt{n^2 + 4c^2} - \sqrt{n^2 - 4c^2} &= \frac{4c^2}{n} + \frac{8c^6}{n^5} + \dots \\ \sqrt{n^2 + 4c^2} \log \left(\frac{\sqrt{n^2 + 4c^2} - n}{n + \sqrt{n^2 + 4c^2}} \right) &= 2n(\log c - \log n) + \frac{4c^2(\log c - \log n - \frac{1}{2})}{n} + \dots \\ \sqrt{n^2 - 4c^2} \log \left(\frac{n + \sqrt{n^2 - 4c^2}}{n - \sqrt{n^2 - 4c^2}} \right) &= -2n(\log c - \log n) + \frac{4c^2(\log c - \log n - \frac{1}{2})}{n} + \dots \\ \frac{n}{2} \left(\log^2 \left(\sqrt{n^2 + 4c^2} - n \right) + \log^2 \left(\sqrt{n^2 + 4c^2} + n \right) \right. \\ \left. - \log^2 \left(n + \sqrt{n^2 - 4c^2} \right) - \log^2 \left(n - \sqrt{n^2 - 4c^2} \right) \right) &= \frac{4c^2(\log n - \log c)}{n} + \frac{80c^6 \log n}{3n^6} + \dots \end{aligned}$$

Substituting these into the sum gives the following bound on the second sum.

$$\begin{aligned} \sum_{n=[2c]}^{\lfloor cT + \frac{c}{T} \rfloor} \frac{\varphi(n)}{|n|} &\left[\log T \left\{ \left(-\frac{4c^2}{n} - \frac{40c^6}{3n^5} + \dots \right) + 2 \left(\frac{4c^2}{n} + \frac{8c^6}{n^5} + \dots \right) \right\} \right. \\ &+ \left(2n(\log c - \log n) + \frac{4c^2(\log c - \log n - \frac{1}{2})}{n} + \dots \right) \\ &+ \left(-2n(\log c - \log n) + \frac{4c^2(\log c - \log n - \frac{1}{2})}{n} + \dots \right) \\ &\left. + \frac{4c}{T} + \left(\frac{4c^2(\log n - \log c)}{n} + \frac{80c^6 \log n}{3n^6} + \dots \right) \right] - \epsilon_{y_4} \end{aligned}$$

$$\begin{aligned}
&= \sum_{n=\lceil 2c \rceil}^{\lfloor cT + \frac{c}{T} \rfloor} \frac{\varphi(n)}{n} \left[\frac{4c^2 \log T}{n} - \frac{4c^2 \log n}{n} + \frac{4c^2(\log c - 1)}{n} + \dots \right] - \epsilon_{y_4} \\
&= \sum_{n=\lceil 2c \rceil}^{\lfloor cT + \frac{c}{T} \rfloor} \frac{\varphi(n)}{n} \frac{4c^2}{n} \log \left(\frac{T}{n} \right) + O(\log T)
\end{aligned}$$

We will make use of the following theorem.

Theorem 15 ([7, Theorem 421]). *Let $\{c_i\}_{i=1}^{\infty}$ be a sequence of real numbers and $f : \mathbb{R} \rightarrow \mathbb{R}$. Suppose there exists $s \in \mathbb{R}$ such that $c_j = 0$ whenever $j < s$ and $f(t)$ has continuous derivative at all $t \geq s$. Then*

$$\sum_{n \leq x} c_n f(n) = \left(\sum_{n \leq x} c_n \right) f(x) - \int_s^x \left(\sum_{n \leq t} c_n \right) \cdot f'(t) dt$$

We also have the following:

Theorem 16 ([20, Chapter 3]).

$$\sum_{n=1}^N \frac{\varphi(n)}{n} = \frac{N}{\zeta(2)} + O\left((\log N)^{\frac{2}{3}} (\log \log N)^{\frac{4}{3}}\right)$$

As a consequence of theorem 16,

$$\sum_{n=\lceil 2c \rceil}^{\lfloor t \rfloor} \frac{\varphi(n)}{n} = \frac{\lfloor t \rfloor - \lceil 2c \rceil + 1}{\zeta(2)} + O\left((\log t)^{\frac{2}{3}} (\log \log t)^{\frac{4}{3}}\right)$$

Take $c_n = \frac{\varphi(n)}{n}$ for $n \geq [2c]$, $c_n = 0$ otherwise, and $f(n) = \frac{1}{n} \log\left(\frac{T}{n}\right)$. These satisfy the assumptions of theorem 15, therefore for large enough T ,

$$\begin{aligned}
& \sum_{n=[2c]}^{\lfloor cT + \frac{c}{T} \rfloor} \frac{\varphi(n)}{n} \frac{1}{n} \log\left(\frac{T}{n}\right) \\
&= \left[\frac{1}{\zeta(2)} \left(\lfloor cT + \frac{c}{T} \rfloor - [2c] \right) + O\left((\log T)^{\frac{2}{3}} (\log \log T)^{\frac{4}{3}} \right) \right] \frac{1}{\lfloor cT + \frac{c}{T} \rfloor} \log\left(\frac{T}{\lfloor cT + \frac{c}{T} \rfloor}\right) \\
&+ \int_{[2c]}^{\lfloor cT + \frac{c}{T} \rfloor} \left(\frac{\lfloor t \rfloor - [2c + 1]}{\zeta(2)} + O\left((\log t)^{\frac{2}{3}} (\log \log t)^{\frac{4}{3}} \right) \right) \frac{\log\left(\frac{T}{t}\right) + 1}{t^2} dt \\
&= \left(\frac{1}{\zeta(2)} - \frac{[2c]}{\zeta(2) \lfloor cT + \frac{c}{T} \rfloor} + \frac{O\left((\log T)^{\frac{2}{3}} (\log \log T)^{\frac{4}{3}} \right)}{\lfloor cT + \frac{c}{T} \rfloor} \right) \log\left(\frac{T}{\lfloor cT + \frac{c}{T} \rfloor}\right) \\
&+ \frac{1}{\zeta(2)} \int_{[2c]}^{\lfloor cT + \frac{c}{T} \rfloor} \frac{\lfloor t \rfloor}{t^2} \left(\log\left(\frac{T}{t}\right) + 1 \right) dt - \frac{[2c + 1]}{\zeta(2)} \int_{[2c]}^{\lfloor cT + \frac{c}{T} \rfloor} \frac{1}{t^2} \left(\log\left(\frac{T}{t}\right) + 1 \right) dt \\
&+ \frac{1}{\zeta(2)} \int_{[2c]}^{\lfloor cT + \frac{c}{T} \rfloor} \frac{O\left((\log t)^{\frac{2}{3}} (\log \log t)^{\frac{4}{3}} \right)}{t^2} \left(\log\left(\frac{T}{t}\right) + 1 \right) dt \\
&:= \frac{1}{\zeta(2)} \int_{[2c]}^{\lfloor cT + \frac{c}{T} \rfloor} \frac{\lfloor t \rfloor}{t^2} \log\left(\frac{T}{t}\right) dt + E'(T) \\
&:= \frac{1}{\zeta(2)} \int_1^T \frac{\log\left(\frac{T}{t}\right)}{t} dt + E''(T) + E'(T) \\
(17) \quad &= \frac{1}{2\zeta(2)} \log^2(T) + E''(T) + E'(T)
\end{aligned}$$

Let $E(T) := E''(T) + E'(T)$. Then equations (14) and (17) imply

$$\begin{aligned}
& \int_{X^2} \left(\widetilde{\mathbb{I}}_{P_{T,c}} - \frac{\text{area}(P_{T,c})}{\zeta(2)} \right)^2 d\mu \\
&= \left\| \widetilde{\mathbb{I}}_{P_{T,c}} \right\|_2^2 - \frac{4c^2}{\zeta(2)} \log^2(T) \\
&= \frac{1}{\zeta(2)} \left(\text{area}(P_{T,c}) + 2 \sum_{n \in \mathbb{N}} \frac{\varphi(|n|)}{|n|} \iint_{P_{T,c}} \left| \mathcal{I}_{(x,y)}^n(P_{T,c}) \right| dx dy \right) - \frac{4c^2}{\zeta(2)} \log^2(T) \\
&= O(\log T) + \frac{2}{\zeta(2)} \frac{4c^2}{2\zeta(2)} \log^2(T) + E(T) - \frac{4c^2}{\zeta(2)^2} \log^2(T) \\
&= O(\log(T) + E(T))
\end{aligned}$$

It remains to calculate the asymptotic growth of $E(T)$.

$$\begin{aligned}
E(T) &= \frac{1}{\zeta(2)} \left(\left(1 - \frac{\lfloor 2c \rfloor}{\lfloor cT + \frac{c}{T} \rfloor} \right) \log \left(\frac{T}{\lfloor cT + \frac{c}{T} \rfloor} \right) \right) + O \left(T^{-1} (\log T)^{\frac{2}{3}} (\log \log T)^{\frac{4}{3}} \right) \\
&\quad + \int_{\lfloor 2c \rfloor}^{\lfloor cT + \frac{c}{T} \rfloor} \left[O \left(\frac{\log \left(\frac{T}{t} \right)}{t^2} \right) + O \left(t^{-2} (\log t)^{\frac{2}{3}} (\log \log t)^{\frac{4}{3}} \right) \right. \\
&\quad \left. + O \left(t^{-2} (\log t)^{\frac{5}{3}} (\log \log t)^{\frac{4}{3}} \right) + O(t^{-2}) \right] dt \\
&= O(1) + O \left(\frac{1}{T} \right) + O \left(T^{-1} (\log T)^{\frac{2}{3}} (\log \log T)^{\frac{4}{3}} \right) + O \left(\frac{\log T}{T} \right) + O \left(\frac{(\log T)^{\frac{10}{3}}}{T} \right) \\
&= O(1)
\end{aligned}$$

Therefore $O(E(T)) + O(\log(T)) = O(\log T)$.

Since $\int_{X^2} \left(\tilde{\mathbb{I}}_{P_{TN,c}} - \frac{\text{area}(P_{TN,c})}{\zeta(2)} \right)^2 d\mu = O(N)$, a similar argument to theorem 5 yields, for any measurable $A \subset \mathbb{S}^0$,

$$\frac{\tilde{\mathbb{I}}_{P_{\tau,c,A}}(\Lambda)}{\tilde{\mathbb{I}}_{P_{\tau,c}}(\Lambda)} = \text{vol}(A) + o_{A,c,\Lambda,d}((\log \tau)^{-\frac{1}{2}} (\log \log \tau)^{\frac{3}{2}} (\log \log \log \tau)^{\frac{1}{2}+\epsilon})$$

Remark 3. Let $\mathcal{H}_1(0)$ be the stratum of all unit area translation surfaces with one marked point. Then $\mathcal{H}_1(0) \cong \text{SL}(2, \mathbb{R})/\text{SL}(2, \mathbb{Z})$. Given a unimodular lattice $\Lambda \subset \mathbb{C} \cong \mathbb{R}^2$, we may find a translation surface $(X, \omega) \in \mathcal{H}_1(0)$ for which Λ is its set of holonomy vectors. The number of saddle connections with holonomy vector lying in $P_{T,c}$ can then be counted using the result of section 4, giving an error term for theorem 1.1 of [2] in the case of the stratum $\mathcal{H}(0)$. See [13, 21] for more details on translation surfaces.

5. FURTHER RESULTS

5.1. Effective Counting for Linear Forms. Recall the definition of $R_{T,c}$ from section 1.3 and the calculation $\text{vol}_d(R_{T,c}) = cB_m C_n \log T$.

Proof of theorem 7. Define the flow $g_t : \mathbb{R} \rightarrow X_d$ by

$$g_t = \begin{pmatrix} e^{\frac{n}{m}t} \text{Id}_m & 0 \\ 0 & e^{-t} \text{Id}_n \end{pmatrix}$$

Then by Moore's Ergodicity Theorem [14, Theorem 3], this flow is ergodic on X_d . For $T \geq 1$ and for any $N \in \mathbb{N}$,

$$\sum_{i=0}^{N-1} \widehat{\mathbb{I}}_{R_{T,c}}(g_{\log T}^i \Lambda) = \widehat{\mathbb{I}}_{R_{TN,c}}(\Lambda)$$

By lemma 1,

$$\begin{aligned} \int_{X_d} \sum_{i=0}^{N-1} \widehat{\mathbb{1}}_{R_{T,c}}(g_{\log T}^i \Lambda) d\mu &\leq \left(\int_{\mathbb{R}^d} \mathbb{1}_{R_{T^N,c}}(\mathbf{x}) d\mathbf{x} \right)^2 + O_d(\text{vol}_d(R_{T^N,c})) \\ &\leq c^2 k^2 B_m^2 C_n^2 \log^2 T + O_d(\log T) \end{aligned}$$

Therefore,

$$\begin{aligned} &\int_{X_d} \left(\sum_{i=0}^{N-1} \widehat{\mathbb{1}}_{R_{T,c}}(g_{\log T}^i \Lambda) - \text{vol}_d(R_{T^N,c}) \right)^2 d\mu(\Lambda) \\ &= \int_{X_d} \left(\widehat{\mathbb{1}}_{R_{T^N,c}}(\Lambda) \right)^2 d\mu(\Lambda) - 2\text{vol}_d(R_{T,c}) \int_{X_d} \widehat{\mathbb{1}}_{R_{T^N,c}}(\Lambda) d\mu(\Lambda) + \text{vol}_d(R_{T^N,c})^2 \\ &\leq N^2 c^2 B_m^2 C_n^2 \log^2 T + O_d(\log T^N) - N^2 c^2 B_m^2 C_n^2 \log^2 T \\ &= O_{T,d}(N) \end{aligned}$$

The result follows by theorem 11 similarly to the proof of theorem 5. \square

5.2. Effective Counting for Affine Lattices. To calculate the second moment of a function over the space of affine lattices, we use the following lemma.

Lemma 2 ([5, Proposition 14]). *Let $d \geq 2$ be a natural number and $F \in L^1(\mathbb{R}^{2d})$. Then*

$$\int_{Y_d} \sum_{\mathbf{v} \neq \mathbf{w} \in \Lambda + \xi} F(\mathbf{v}, \mathbf{w}) d\nu(\Lambda, \nu) = \int_{\mathbb{R}^{2d}} F(\mathbf{x}) d\mathbf{x}$$

proof of theorem 8. Using theorems 9 and lemma 2, we deduce that, for any $f \in L^1(\mathbb{R}^d)$,

$$\int_{Y_d} \left(\widehat{f}(\Lambda + \xi) \right)^2 d\nu(\Lambda, \xi) = \left(\int_{\mathbb{R}^d} f(\mathbf{x}) d\mathbf{x} \right)^2 + \int_{\mathbb{R}^d} f(\mathbf{x})^2 d\mathbf{x}$$

By setting $f = \mathbb{1}_B$, we have

$$\int_{Y_d} \widehat{\mathbb{1}}_B(\Lambda + \xi) d\nu(\Lambda, \xi) = (\text{vol}_d(B))^2 + \text{vol}_d(B)$$

The result then follows using a similar argument to theorem 7. \square

REFERENCES

- [1] J. S. Athreya, A. Ghosh, and J. Tseng. Spiraling of approximations and spherical averages of Siegel transforms. *Journal of the London Mathematical Society*, 91(2):383–404, 2015.
- [2] J. S. Athreya, A. Parrish, and J. Tseng. Ergodic Theory and Diophantine Approximation for Translation Surfaces and Linear Forms. *Nonlinearity*, 29(8):2173–2190, Aug. 2016.
- [3] M. B. Bekka and M. Mayer. *Ergodic Theory and Topological Dynamics of Group Actions on Homogeneous Spaces*. Cambridge University Press, Cambridge, 2000.
- [4] L. Dirichlet. Verallgemeinerung eines Satzes aus der Lehre von den Kettenbrüchen nebst einigen Anwendungen auf die Theorie der Zahlen. *SB Preuss. Akad. Wiss.*, 1842:93–95, 1842.
- [5] D. El-Baz, J. Marklof, and I. Vinogradov. The Distribution of Directions in an Affine Lattice: Two-Point Correlations and Mixed Moments. *International Mathematics Research Notices*, 2015(5):1371–1400.

- [6] V. F. Gaposhkin. On the Dependence of the Convergence Rate in the Strong Law of Large Numbers for Stationary Processes on the Rate of Decay of the Correlation Function. *Theory of Probability & Its Applications*, 26(4):706–720, Jan. 1982.
- [7] G. H. Hardy, E. M. Wright, D. R. Heath-Brown, and J. H. Silverman. *An Introduction to the Theory of Numbers*. Posts & Telecom Press, Pékin, 2009.
- [8] A. Kachurovskii. The rate of convergence in ergodic theorems. *Russian Mathematical Surveys*, 51:653–703, 1996.
- [9] D. Y. Kleinbock, N. Shah, and A. Starkov. Dynamics of Subgroup Actions on Homogeneous Spaces of Lie Groups and Applications to Number Theory. In *Handbook of Dynamical Systems*, volume 1, chapter 11, pages 813–930. Elsevier, 2002.
- [10] D. Y. Kleinbock, R. Shi, and B. Weiss. Pointwise equidistribution with an error rate and with respect to unbounded functions. *Mathematische Annalen*, 367(1-2):857–879, Feb. 2017.
- [11] D. Y. Kleinbock and S. Yu. A dynamical Borel-Cantelli lemma via improvements to Dirichlet’s theorem. *Moscow Journal of Combinatorics and Number Theory*, 9(2):101–122, February 2020.
- [12] A. M. Macbeath and C. A. Rogers. Siegel’s Mean Value Theorem in the Geometry of Numbers. *Mathematical Proceedings of the Cambridge Philosophical Society*, 54(2):139–151, Apr. 1958.
- [13] H. Masur. Ergodic Theory of Translation Surfaces. In B. Hasselblatt and A. B. Katok, editors, *Handbook of Dynamical Systems*. N.H. North Holland : Elsevier, Amsterdam; New York, 2002.
- [14] C. C. Moore. Ergodicity of Flows on Homogeneous Spaces. *American Journal of Mathematics*, 88(1):154, Jan. 1966.
- [15] M. Newman. The Smith Normal Form. *Linear Algebra and its Applications*, 254(1-3):367–381, Mar. 1997.
- [16] C. A. Rogers. Mean Values over the Space of Lattices. *Acta Mathematica*, 94(0):249–287, 1955.
- [17] W. Schmidt. A Metrical Theorem In Geometry of Numbers. *Transactions of the American Mathematical Society*, 95(3):516–529, 1960.
- [18] W. Schmidt. A Metrical Theorem in Diophantine Approximation. *Canadian Journal of Mathematics*, 12:619–631, 1960.
- [19] C. L. Siegel. A Mean Value Theorem in Geometry of Numbers. *The Annals of Mathematics*, 46(2):340, Apr. 1945.
- [20] A. Walfisz. *Weylsche Exponentialsummen in der neueren Zahlentheorie*. VEB Deutscher Verlag der Wissenschaften, Berlin, 1963.
- [21] A. Zorich. Flat Surfaces. In *Frontiers in Number Theory, Physics, and Geometry I*, pages 437 – 583. Springer, Berlin, 2006.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF EXETER, EXETER, EX4 4QF, UK
 Email address: `nh477@exeter.ac.uk`