# arXiv:2201.06441v1 [math.FA] 17 Jan 2022

# ASYMPTOTIC ALMOST AUTOMORPHY FOR ALGEBRAS OF GENERALIZED FUNCTIONS

CHIKH BOUZAR AND MERYEM SLIMANI

ABSTRACT. Asymptotic almost automorphy is introduced and studied in the context of some algebras of generalized functions. We also give applications to neutral difference differential systems in the framework of such generalized functions.

### 1. INTRODUCTION

Bochner S. defined explicitly almost automorphic functions in the papers [2]-[4], where also some of their basic properties are given. In [3] he studied linear difference differential equations in the framework of such functions. It is well known that the concept of almost automorphy is strictly more general than the almost periodicity of H. Bohr [5], however the Stepanoff almost periodicity [26] and the Levitan almost periodicity [18] don't enter into the Bochner concept. Asymptotic almost periodicity of functions as a perturbation of almost periodic functions by functions vanishing at infinity is due to Fréchet M. in [16]. Asymptotic almost automorphy of classical functions is considered in [19], see also [12]. The almost periodicity and the asymptotic almost periodicity of Sobolev-Schwartz distributions, [25] and [22], are respectively considered by L. Schwartz in [22] and I. Cioranescu in [13], while almost automorphy and asymptotic almost automorphy in the setting of these distributions are respectively the subject of the recent works [11] and [12].

In view of the result [23] on the impossibility of the multiplication of distributions, see [21] for more details, algebras of generalized functions containing spaces of Sobolev-Schwartz type distributions have been studied, see [14], [15], [1] and [20]. The concepts of almost periodicity and asymptotic almost periodicity as well as almost automorphy in the context of such algebras of generalized functions are introduced, studied and applied in the papers [6], [7] [8], [9] and [10]. So, the paper first introduces and studies a class of asymptotically almost automorphic generalized functions, denoted by  $\mathcal{G}_{aaa}$ . In the sense of multiplication, not only  $\mathcal{G}_{aaa}$  is stable under multiplication and it contains the space of asymptotically almost automorphic distributions of [12], but moreover some nonlinear operations are performed within the algebra  $\mathcal{G}_{aaa}$ . As a by pass result, we give a Seeley type result on extention of functions in the context of the introduced generalized functions, this is needed in the proof of a fundamental result on the

<sup>2020</sup> Mathematics Subject Classification. 46F10. 34K14. 46F30.

*Key words and phrases.* Asymptotic almost automorphy, Generalized functions, Neutral difference differential equations.

uniqueness of decomposition of an asymptotically almost automorphic generalized function. The papers [8] and [9] can be considered as consequences of this work. The paper aims also, as in [16], to lift a Frechet existence result of asymptotically almost automorphic solutions of differential equations to the level of neutral difference differential systems in the framework of  $\mathcal{G}_{aaa}$ .

It is worth noting that the meaning of generalized functions is utilized differently by authors as distributions or ultradistributions, even as hyperfunctions, but in this work by generalized functions we mean in the sense of the works [14], [15], [1] and [20].

The paper is organized as follows: section two recalls definitions and some properties of asymptotically almost automorphic functions and asymptotically almost automorphic distributions as in [12]. Section three introduces asymptotically almost automorphic generalized functions and gives some of their important properties. The study of a Seeley type result on extensions of generalized functions is given in section four. In section six non-linear operations on asymptotically almost automorphic generalized function are studied. The last section is dedicated to linear neutral difference differential systems in the framework of asymptotically almost automorphic generalized functions.

# 2. Asymptotic almost automorphy of functions and distributions

Let  $\mathcal{C}_b$  denotes the space of bounded and continuous complex-valued functions defined on  $\mathbb{R}$ , endowed with the norm  $\|\cdot\|_{L^{\infty}(\mathbb{R})}$  of uniform convergence on  $\mathbb{R}$ , it is well-known that  $(\mathcal{C}_b, \|\cdot\|_{L^{\infty}(\mathbb{R})})$ is a Banach algebra.

A complex-valued function g defined and continuous on  $\mathbb{R}$  is called almost automorphic if for any sequence  $(s_m)_{m \in \mathbb{N}} \subset \mathbb{R}$ , one can extract a subsequence  $(s_{m_k})_k$  such that

$$\tilde{g}(x) := \lim_{k \to +\infty} g(x + s_{m_k})$$
 exists for every  $x \in \mathbb{R}$ ,

and

$$\lim_{k \to +\infty} \tilde{g} \left( x - s_{m_k} \right) = g \left( x \right) \text{ for every } x \in \mathbb{R}.$$

The space of almost automorphic functions on  $\mathbb{R}$  is denoted by  $\mathcal{C}_{aa}$ .

The space  $\mathcal{C}_{+,0}$  is the set of all bounded and continuous complex-valued functions defined on  $\mathbb{R}$  and vanishing at  $+\infty$ .

**Definition 1.** We say that a function  $f \in C_b$  is asymptotically almost automorphic, if there exist  $g \in C_{aa}$  and  $h \in C_{+,0}$  such that f = g + h on  $\mathbb{J} := [0, +\infty[$ . The space of asymptotically almost automorphic functions is denoted by  $C_{aaa}$ .

For a study of asymptotically almost automorphic functions and asymptotically almost automorphic distributions see [11] and [12] and the references list therein.

**Proposition 1.** The decomposition of an asymptotically almost automorphic function is unique on  $\mathbb{J}$ .

**Notation 1.** If  $f \in C_{aaa}$  and f = g + h on  $\mathbb{J}$ , where  $g \in C_{aa}$  and  $h \in C_{+,0}$ . Due to the uniqueness of the decomposition of f, the function g is said the principal term of f and the function h the corrective term of f, we denote them respectively by  $f_{aa}$  and  $f_{cor}$ . The notation  $f = (f_{aa} + f_{cor}) \in C_{aaa}$  means that  $f_{aa} \in C_{aa}$ ,  $f_{cor} \in C_{+,0}$  and  $f = f_{aa} + f_{cor}$  on  $\mathbb{J}$ .

Let  $\mathcal{E}(\mathbb{I})$  be the algebra of space of smooth functions on  $\mathbb{I} = \mathbb{R}$  or  $\mathbb{J}$ , and define the space

$$\mathcal{D}_{L^{p}}\left(\mathbb{I}\right) := \left\{\varphi \in \mathcal{E}(\mathbb{I}) : \forall j \in \mathbb{Z}_{+}, \varphi^{(j)} \in L^{p}\left(\mathbb{I}\right)\right\}, p \in \left[1, +\infty\right],$$

that we endow with the topology defined by the family of semi-norms

$$\left|\varphi\right|_{k,p,\mathbb{I}} := \sum_{j \le k} \left\|\varphi^{(j)}\right\|_{L^{p}(\mathbb{I})}, k \in \mathbb{Z}_{+}.$$

So,  $\mathcal{D}_{L^{p}}(\mathbb{I})$  is a Fréchet subalgebra of  $\mathcal{E}(\mathbb{I})$ . Denote  $\mathcal{B}(\mathbb{I}) := \mathcal{D}_{L^{\infty}}(\mathbb{I})$ .

**Remark 1.** We have  $\lim_{\substack{x\to 0\\>}} \varphi^{(j)}(0)$  exists for every  $j \in \mathbb{Z}_+$  when  $\varphi \in \mathcal{D}_{L^p}(\mathbb{J})$ .

The space of smooth almost automorphic functions  $\mathcal{B}_{aa}$  and smooth asymptotically almost automorphic functions  $\mathcal{B}_{aaa}$  are defined respectively by

$$\mathcal{B}_{aa} := \left\{ \varphi \in \mathcal{E} \left( \mathbb{R} \right) : \forall j \in \mathbb{Z}_+, \ \varphi^{(j)} \in \mathcal{C}_{aa} \right\}.$$
$$\mathcal{B}_{aaa} := \left\{ \varphi \in \mathcal{E} \left( \mathbb{R} \right) : \ \forall j \in \mathbb{Z}_+, \ \varphi^{(j)} \in \mathcal{C}_{aaa} \right\}$$

We endow  $\mathcal{B}_{aa}$  and  $\mathcal{B}_{aaa}$  with the topology induced by  $\mathcal{B} := \mathcal{D}_{L^{\infty}}(\mathbb{R})$ .

**Proposition 2.** (1) The space  $\mathcal{B}_{aaa}$  is a Fréchet subalgebra of  $\mathcal{B}$  stable by translation. (2)  $\mathcal{B}_{aaa} \times \mathcal{B}_{aa} \subset \mathcal{B}_{aaa}$ . (3)  $\mathcal{B}_{aaa} * L^1 \subset \mathcal{B}_{aaa}$ .

The space of  $L^p$ -distributions,  $p \in [1, +\infty]$ , denoted by  $\mathcal{D}'_{L^p}(\mathbb{R})$ , is the topological dual of  $\mathcal{D}_{L^q}(\mathbb{R})$ , where  $\frac{1}{p} + \frac{1}{q} = 1$ . Let  $\dot{\mathcal{B}}$  be the closure in  $\mathcal{B}$  of the space  $\mathcal{D} \subset \mathcal{E}(\mathbb{R})$  of functions with compact support. The topological dual of  $\dot{\mathcal{B}}$  is denoted by  $\mathcal{D}'_{L^1}(\mathbb{R})$ . The space of bounded distributions  $\mathcal{D}'_{L^\infty}(\mathbb{R})$  is denoted by  $\mathcal{B}'$ .

**Definition 2.** The space of almost automorphic distributions, denoted by  $\mathcal{B}'_{aa}$ , is the space of  $T \in \mathcal{B}'$  satisfying one of the following equivalent statements

- (1)  $T * \varphi \in \mathcal{C}_{aa}, \ \forall \varphi \in \mathcal{D}.$
- (2)  $\exists k \in \mathbb{Z}_+ \text{ and } g_j \in \mathcal{C}_{aa}, \ 0 \le j \le k, \text{ such that } T = \sum_{j=0}^k g_j^{(j)}.$

The space of bounded distributions vanishing at infinity, denoted by  $\mathcal{B}'_{+,0}$ , is the space of  $Q \in \mathcal{B}'$  satisfying

$$\lim_{\omega \to +\infty} \langle \tau_{\omega} Q, \varphi \rangle := \lim_{\omega \to +\infty} \langle Q, \tau_{-\omega} \varphi \rangle = 0, \ \forall \varphi \in \mathcal{D},$$

where  $\tau_{\omega}\varphi(\cdot) := \varphi(\cdot + \omega), \omega \in \mathbb{R}$ .

**Theorem 1.** The space of asymptotically almost automorphic distributions, denoted by  $\mathcal{B}'_{aaa}$ , is the space of  $T \in \mathcal{B}'$  satisfying one of the following equivalent statements

(1)  $\exists P \in \mathcal{B}'_{aa}, \exists Q \in \mathcal{B}'_{+,0} \text{ such that } T = P + Q \text{ on } \mathbb{J}.$ (2)  $T * \varphi \in \mathcal{C}_{aaa}, \forall \varphi \in \mathcal{D}.$ (3)  $\exists k \in \mathbb{Z}_+ \text{ and } f_j \in \mathcal{C}_{aaa}, 0 \leq j \leq k, \text{ such that } T = \sum_{j=0}^k f_j^{(j)}.$ (4)  $\exists (\theta_m)_{m \in \mathbb{N}} \subset \mathcal{B}_{aaa} \text{ such that } \lim_{m \to +\infty} \theta_m = T \text{ in } \mathcal{B}'.$ 

**Notation 2.** If  $T \in \mathcal{B}'_{aaa}$  and T = P + Q on  $\mathbb{J}$ , since this decomposition is unique by ([12], Proposition 12), the distribution P is called the principal term of T, the distribution Q is called the corrective term of T, we denote them respectively  $T_{aa}$  and  $T_{cor}$ . This is summarized by the notation  $T = (T_{aa} + T_{cor}) \in \mathcal{B}'_{aaa}$ .

## 3. Asymptotically almost automorphic generalized functions

We introduce and study an algebra of asymptotically almost automorphic generalized functions.

Let 
$$I := ]0, 1], (u_{\varepsilon})_{\varepsilon} \in (\mathcal{D}_{L^{p}}(\mathbb{I}))^{I}, p \in [1, +\infty], m \in \mathbb{Z} \text{ and } k \in \mathbb{Z}_{+}, \text{ then the notation}$$
  
 $|u_{\varepsilon}|_{k,p,\mathbb{I}} = O(\varepsilon^{m}), \varepsilon \to 0. \Leftrightarrow \exists c > 0, \exists \varepsilon_{0} \in I, \forall \varepsilon < \varepsilon_{0}, |u_{\varepsilon}|_{k,p,\mathbb{I}} \leq c\varepsilon^{m}.$ 

**Definition 3.** The algebra of asymptotically almost automorphic generalized functions is denoted and defined as the quotient algebra

$$\mathcal{G}_{aaa} := rac{\mathcal{M}_{aaa}}{\mathcal{N}_{aaa}}$$

where the space of moderate elements is denoted and defined by

$$\mathcal{M}_{aaa} := \left\{ \left( u_{\varepsilon} \right)_{\varepsilon} \in \left( \mathcal{B}_{aaa} \right)^{I} : \forall k \in \mathbb{Z}_{+}, \exists m \in \mathbb{Z}_{+}, \left| u_{\varepsilon} \right|_{k,\infty,\mathbb{R}} = O\left( \varepsilon^{-m} \right), \varepsilon \to 0 \right\},\$$

and the space of null elements by

$$\mathcal{N}_{aaa} := \left\{ (u_{\varepsilon})_{\varepsilon} \in (\mathcal{B}_{aaa})^{I} : \forall k \in \mathbb{Z}_{+}, \forall m \in \mathbb{Z}_{+}, |u_{\varepsilon}|_{k,\infty,\mathbb{R}} = O(\varepsilon^{m}), \varepsilon \to 0 \right\}.$$

**Example 1.** We have  $\mathcal{G}_{aap} \subsetneq \mathcal{G}_{aaa}$ , where  $\mathcal{G}_{aap}$  is the algebra of asymptotically almost periodic generalized functions of [8].

Some properties of  $\mathcal{M}_{aaa}$  and  $\mathcal{N}_{aaa}$  are given in the following results.

**Proposition 3.** (1) We have the null characterization of  $\mathcal{N}_{aaa}$ , i.e.

$$\mathcal{N}_{aaa} = \left\{ (u_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa} : \forall m \in \mathbb{Z}_{+}, \ |u_{\varepsilon}|_{0,\infty,\mathbb{R}} = O(\varepsilon^{m}), \varepsilon \to 0 \right\}$$

- (2) The space  $\mathcal{M}_{aaa}$  is an algebra stable under translation and derivation.
- (3) The space  $\mathcal{N}_{aaa}$  is an ideal of  $\mathcal{M}_{aaa}$ .

*Proof.* (1) The proof is based on the following Landau-Kolmogorov inequality

$$\left\| f^{(p)} \right\|_{L^{\infty}(\mathbb{R})} \leq 2\pi \left\| f \right\|_{L^{\infty}(\mathbb{R})}^{1-\frac{p}{n}} \left\| f^{(n)} \right\|_{L^{\infty}(\mathbb{R})}^{\frac{p}{n}}$$

where 0 and <math>f is of class  $\mathcal{C}^n(\mathbb{R})$ . Let  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$ , i.e.

(3.1) 
$$\forall k \in \mathbb{Z}_+, \ \exists m \in \mathbb{Z}_+, \ \exists c > 0, \ \exists \varepsilon_1 \in I, \ \forall \varepsilon < \varepsilon_1, \ |u_\varepsilon|_{k,\infty,\mathbb{R}} \le c\varepsilon^{-m},$$

and it satisfies the estimate of order zero, i.e.

(3.2) 
$$\forall m_2 \in \mathbb{Z}_+, \ \exists c_2 > 0, \ \exists \varepsilon_2 \in I, \ \forall \varepsilon < \varepsilon_2, \ |u_\varepsilon|_{0,\infty,\mathbb{R}} \le c_2 \varepsilon^{m_2}.$$

The Landau-Kolmogorov inequality for p = j, n = 2j, (3.1) and (3.2) give  $\forall k \in \mathbb{Z}_+$ ,

$$\begin{aligned} |u_{\varepsilon}|_{k,\infty,\mathbb{R}} &\leq \sum_{j\leq k} 2\pi \left\| u_{\varepsilon} \right\|_{L^{\infty}(\mathbb{R})}^{1-\frac{1}{2}} \left\| u_{\varepsilon}^{(2j)} \right\|_{L^{\infty}(\mathbb{R})}^{\frac{1}{2}}, \\ &\leq 2\pi \left( |u_{\varepsilon}|_{0,\infty,\mathbb{R}} \right)^{\frac{1}{2}} \sum_{l\leq 2k} \left\| u_{\varepsilon}^{(l)} \right\|_{L^{\infty}(\mathbb{R})}^{\frac{1}{2}}, \\ &\leq 2\pi c^{\frac{1}{2}} c_{2}^{\frac{1}{2}} \varepsilon^{-\frac{m}{2}} \varepsilon^{\frac{m_{2}}{2}}. \end{aligned}$$

Taking  $m_2 \in \mathbb{Z}_+$  such that  $m_0 = 2\left(-\frac{m}{2} + \frac{m_2}{2}\right) \in \mathbb{Z}_+$ , then we obtain  $\forall k \in \mathbb{Z}_+$ ,  $\forall m_0 \in \mathbb{Z}_+$ ,  $\exists c = 2\pi c^{\frac{1}{2}} c_2^{\frac{1}{2}} > 0$ ,  $\exists \varepsilon_0 = \inf(\varepsilon_1, \varepsilon_2) \in I$ ,  $\forall \varepsilon < \varepsilon_0$ ,  $|u_{\varepsilon}|_{k,\infty,\mathbb{R}} \leq c\varepsilon^{m_0}$ . Which gives  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aaa}$ .

(2) The stability under translation and derivation of the space  $\mathcal{M}_{aaa}$  is obvious. Let  $(u_{\varepsilon})_{\varepsilon}, (v_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$ , i.e. they satisfy (3.1) and for  $j \in \mathbb{Z}_+$ ,

$$\begin{aligned} \left\| (u_{\varepsilon}v_{\varepsilon})^{(j)} \right\|_{L^{\infty}(\mathbb{R})} &\leq \sum_{i\leq j} \frac{j!}{i!(j-i)!} \left\| u_{\varepsilon}^{(i)} \right\|_{L^{\infty}(\mathbb{R})} \left\| v_{\varepsilon}^{(j-i)} \right\|_{L^{\infty}(\mathbb{R})} \\ &\leq |u_{\varepsilon}|_{j,\infty,\mathbb{R}} \left| v_{\varepsilon} \right|_{j,\infty,\mathbb{R}} \sum_{i\leq j} \frac{j!}{i!(j-i)!}, \\ &\leq 2^{j}c_{1}c_{2}\varepsilon^{-m_{1}}\varepsilon^{-m_{2}}, \end{aligned}$$

consequently,  $\forall k \in \mathbb{Z}_+$ ,  $\exists m = (m_1 + m_2) \in \mathbb{Z}_+$ ,  $|u_{\varepsilon}v_{\varepsilon}|_{k,\infty,\mathbb{R}} = O(\varepsilon^{-m}), \varepsilon \to 0$ . Which shows that  $(u_{\varepsilon}v_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$ .

(3) Let  $(v_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$  and  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aaa}$ , i.e.  $(v_{\varepsilon})_{\varepsilon}$  satisfies (3.1) and  $(u_{\varepsilon})_{\varepsilon}$  satisfies

$$\forall k \in \mathbb{Z}_+, \ \forall m' \in \mathbb{Z}_+, \ \exists c' > 0, \ \exists \varepsilon_1' \in I, \ \forall \varepsilon < \varepsilon_1', \ |u_\varepsilon|_{k,\infty,\mathbb{R}} \le c' \varepsilon^{m'}.$$

Since the family of the norms  $|\cdot|_{k,\infty,\mathbb{R}}$  is compatible with the algebraic structure of  $\mathcal{B}$ , i.e.  $\forall k \in \mathbb{Z}_+, \exists c_k > 0$  such that

$$|u_{\varepsilon}v_{\varepsilon}|_{k,\infty,\mathbb{R}} \leq c_k |u_{\varepsilon}|_{k,\infty,\mathbb{R}} |v_{\varepsilon}|_{k,\infty,\mathbb{R}} \leq c_k cc' \varepsilon^{-m} \varepsilon^{m'}.$$

Take  $m' \in \mathbb{Z}_+$  such that  $m_0 = (-m + m') \in \mathbb{Z}_+$ , so we obtain  $\forall k \in \mathbb{Z}_+$ ,  $\forall m_0 \in \mathbb{Z}_+$ ,  $\exists C = c_k cc' > 0$ ,  $\exists \varepsilon_0 = \inf (\varepsilon_1, \varepsilon_1') \in I$ ,  $\forall \varepsilon < \varepsilon_0$ ,  $|u_{\varepsilon} v_{\varepsilon}|_{k,\infty,\mathbb{R}} \leq C \varepsilon^{m_0}$ , which implies  $(u_{\varepsilon} v_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aaa}$ .

Let  $\rho \in \mathcal{S}$  such that  $\int_{\mathbb{R}} \rho(x) dx = 1$  and  $\int_{\mathbb{R}} x^k \rho(x) dx = 0, \forall k \in \mathbb{N}$ . Set  $\rho_{\varepsilon}(\cdot) := \frac{1}{\varepsilon} \rho\left(\frac{\cdot}{\varepsilon}\right), \varepsilon > 0$ . Define the following maps

**Proposition 4.** The following diagram of linear embeddings

$$egin{array}{ccc} \mathcal{B}_{aaa} & \stackrel{\gamma_{aaa}}{\longrightarrow} & \mathcal{B}_{aaa}' \ \sigma_{aaa} \searrow & \downarrow \iota_{aaa} \ \mathcal{G}_{aaa} \end{array}$$

is commutative.

Proof. For  $f \in \mathcal{B}_{aaa}$ , we have  $\gamma_{aaa}(f) \in \mathcal{B}_{aaa} \subset \mathcal{C}_{aaa}$ . We conclude from ([12], Example 3 – (1)) that  $\gamma_{aaa}(f) \in \mathcal{B}'_{aaa}$ . Let  $T \in \mathcal{B}'_{aaa}$ , by the characterization of an asymptotically almost automorphic distribution  $\exists (f_i)_{i \leq m} \subset \mathcal{C}_{aaa}$  such that  $T = \sum_{i \leq m} f_i^{(i)}$ . If  $j \in \mathbb{Z}_+$ ,

$$\begin{aligned} \left| \left( T * \rho_{\varepsilon} \right)^{(j)} (x) \right| &\leq \sum_{i \leq m} \frac{1}{\varepsilon^{i+j}} \int_{\mathbb{R}} \left| f_i \left( x - \varepsilon y \right) \rho^{(i+j)} (y) \right| dy \\ &\leq \sum_{i \leq m} \frac{1}{\varepsilon^{i+j}} \| f_i \|_{L^{\infty}(\mathbb{R})} \int_{\mathbb{R}} \left| \rho^{(i+j)} (y) \right| dy, \end{aligned}$$

consequently, there exists  $c_{m,j} > 0$  such that

$$\left\| \left( T * \rho_{\varepsilon} \right)^{(j)} \right\|_{L^{\infty}(\mathbb{R})} \leq \frac{c_{m,j}}{\varepsilon^{m+j}},$$

hence,  $\forall k \in \mathbb{Z}_+, \exists m_1 \in \mathbb{Z}_+,$ 

$$|T * \rho_{\varepsilon}|_{k,\infty,\mathbb{R}} = O\left(\varepsilon^{-m_1}\right), \ \varepsilon \to 0,$$

which gives that  $(T * \rho_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$ . The linearity of  $\iota_{aaa}$  results from the fact that the convolution is linear. If  $(T * \rho_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aaa}$ , then

(3.4) 
$$\forall m' \in \mathbb{Z}_+, \ \exists c' > 0, \ \exists \varepsilon_1' \in I, \ \forall \varepsilon < \varepsilon_1', \ \|T * \rho_\varepsilon\|_{L^\infty(\mathbb{R})} \le c' \varepsilon^{m'}.$$

Due to regularization we have

$$\langle T, \psi \rangle = \lim_{\varepsilon \to 0} \int \left( T * \rho_{\varepsilon} \right) (x) \psi (x) \, dx, \psi \in \mathcal{D}_{L^1} (\mathbb{R}) \, .$$

From (3.4), it holds  $\forall m' \in \mathbb{Z}_+, \ \exists c' > 0, \ \exists \varepsilon_1' \in I, \forall \varepsilon < \varepsilon_1',$ 

$$\left| \int (T * \rho_{\varepsilon})(x)\psi(x)dx \right| \le \|\psi\|_{L^{1}(\mathbb{R})} c' \varepsilon^{m'},$$

consequently, when  $\varepsilon \to 0$  we obtain  $\langle T, \psi \rangle = 0$ ,  $\forall \psi \in \mathcal{D}_{L^1}(\mathbb{R})$ , therefore  $\iota_{aaa}$  is injective. Finally,

$$\iota_{aaa} \left( T^{(j)} \right) = \left( T^{(j)} * \rho_{\varepsilon} \right)_{\varepsilon} + \mathcal{N}_{aaa} = \left( T * \rho_{\varepsilon} \right)_{\varepsilon}^{(j)} + \mathcal{N}_{aaa} = \left( \iota_{aaa}(T) \right)^{(j)}, \ \forall j \in \mathbb{Z}_{+},$$

i.e. the embedding  $\iota_{aaa}$  commutes with derivatives.

Let  $f \in \mathcal{B}_{aaa}$ , we have to show that  $(f * \rho_{\varepsilon} - f)_{\varepsilon} \in \mathcal{N}_{aaa}$ . By Taylor's formula, for  $\theta \in ]0, 1[$ and  $m \in \mathbb{N}$ , we set for  $j \in \mathbb{Z}_+$ , that  $(f^{(j)} * \rho_{\varepsilon} - f^{(j)})(x)$  equals

$$\int_{\mathbb{R}} \sum_{l=1}^{m-1} \frac{\left(-\varepsilon y\right)^{l}}{l!} f^{(l+j)}\left(x\right) \rho\left(y\right) dy + \int_{\mathbb{R}} \frac{\left(-\varepsilon y\right)^{m}}{m!} f^{(m+j)}\left(x-\theta \varepsilon y\right) \rho\left(y\right) dy,$$

and then

$$\begin{split} \left\| f^{(j)} * \rho_{\varepsilon} - f^{(j)} \right\|_{L^{\infty}(\mathbb{R})} &\leq \frac{\varepsilon^{m}}{m!} \sup_{x \in \mathbb{R}} \int_{\mathbb{R}} \left| (-y)^{m} \right| \left| f^{(m+j)} \left( x - \theta \varepsilon y \right) \right| \left| \rho \left( y \right) \right| dy, \\ &\leq \frac{\varepsilon^{m}}{m!} \left\| f^{(m+j)} \right\|_{L^{\infty}(\mathbb{R})} \left\| y^{m} \rho \right\|_{L^{1}(\mathbb{R})}. \end{split}$$

Hence,  $\forall k \in \mathbb{Z}_+, \forall m \in \mathbb{N}$ ,

$$|f * \rho_{\varepsilon} - f|_{k,\infty,\mathbb{R}} \le \frac{\varepsilon^m}{m!} |f|_{m+k,\infty,\mathbb{R}} ||y^m \rho||_{L^1(\mathbb{R})},$$

which means that  $(f * \rho_{\varepsilon} - f)_{\varepsilon} \in \mathcal{N}_{aaa}$ .

Recall the algebra of almost automorphic generalized functions, see [10], denoted and defined by

$$\mathcal{G}_{aa} := rac{\mathcal{M}_{aa}}{\mathcal{N}_{aa}},$$

where

$$\mathcal{M}_{aa} := \left\{ (u_{\varepsilon})_{\varepsilon} \in (\mathcal{B}_{aa})^{I} : \forall k \in \mathbb{Z}_{+}, \exists m \in \mathbb{Z}_{+}, |u_{\varepsilon}|_{k,\infty,\mathbb{R}} = O\left(\varepsilon^{-m}\right), \varepsilon \to 0 \right\}.$$
$$\mathcal{N}_{aa} := \left\{ (u_{\varepsilon})_{\varepsilon} \in (\mathcal{B}_{aa})^{I} : \forall k \in \mathbb{Z}_{+}, \forall m \in \mathbb{Z}_{+}, |u_{\varepsilon}|_{k,\infty,\mathbb{R}} = O\left(\varepsilon^{m}\right), \varepsilon \to 0 \right\}.$$

The algebra of  $L^p$ -generalized functions on  $\mathbb{I}$ ,  $1 \leq p \leq +\infty$ , see [17], is defined as the quotient algebra

$$\mathcal{G}_{L^{p}}\left(\mathbb{I}
ight):=rac{\mathcal{M}_{L^{p}}\left(\mathbb{I}
ight)}{\mathcal{N}_{L^{p}}\left(\mathbb{I}
ight)},$$

where

$$\mathcal{M}_{L^{p}}(\mathbb{I}) := \left\{ (u_{\varepsilon})_{\varepsilon} \in (\mathcal{D}_{L^{p}}(\mathbb{I}))^{I} : \forall k \in \mathbb{Z}_{+}, \exists m > 0, \ |u_{\varepsilon}|_{k,p,\mathbb{I}} = O(\varepsilon^{-m}), \ \varepsilon \to 0 \right\}$$
$$\mathcal{N}_{L^{p}}(\mathbb{I}) := \left\{ (u_{\varepsilon})_{\varepsilon} \in (\mathcal{D}_{L^{p}}(\mathbb{I}))^{I} : \forall k \in \mathbb{Z}_{+}, \forall m > 0, \ |u_{\varepsilon}|_{k,p,\mathbb{I}} = O(\varepsilon^{m}), \ \varepsilon \to 0 \right\}.$$

Notation 3. Denote  $\mathcal{G}_{L^{\infty}}(\mathbb{I}) := \mathcal{G}_{\mathcal{B}}(\mathbb{I}), \mathcal{G}_{\mathcal{B}} := \mathcal{G}_{\mathcal{B}}(\mathbb{R}) \text{ and } \mathcal{G}_{L^{1}} := \mathcal{G}_{L^{1}}(\mathbb{R}).$ 

For  $\omega \in \mathbb{R}$ , the translate  $\tau_{\omega} \widetilde{u}$  of  $\widetilde{u} = [(u_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{\mathcal{B}}$  is defined by

$$\tau_{\omega}\widetilde{u} := \left[ \left( \tau_{\omega} u_{\varepsilon} \right)_{\varepsilon} \right]$$

For  $j \in \mathbb{Z}_+$ , the derivation  $\widetilde{u}^{(j)}$  of  $\widetilde{u}$  is defined by

$$\widetilde{u}^{(j)} := \left[ \left( u_{\varepsilon}^{(j)} \right)_{\varepsilon} \right].$$

Let  $\widetilde{v} = [(v_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{\mathcal{B}}$ , the product  $\widetilde{u} \times \widetilde{v}$  is defined by

$$\widetilde{u} \times \widetilde{v} := \left[ (u_{\varepsilon} v_{\varepsilon})_{\varepsilon} \right].$$

Let  $\widetilde{v} = [(v_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{L^1}$ , the convolution  $\widetilde{u} * \widetilde{v}$  is defined by

$$\widetilde{u} * \widetilde{v} := \left[ \left( u_{\varepsilon} * v_{\varepsilon} \right)_{\varepsilon} \right]$$

The following results lift the results of Proposition 2 to  $\mathcal{G}_{aaa}$ .

# **Proposition 5.** (1) $\mathcal{G}_{aaa}$ is a subalgebra of $\mathcal{G}_{\mathcal{B}}$ stable under translation and derivation. (2) $\mathcal{G}_{aaa} \times \mathcal{G}_{aa} \subset \mathcal{G}_{aaa}$ . (3) $\mathcal{G}_{aaa} * \mathcal{G}_{L^1} \subset \mathcal{G}_{aaa}$ .

- *Proof.* (1) From Proposition 3-(2), we deduce that  $\mathcal{G}_{aaa}$  is an algebra stable under translation and derivation. Let  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$ , satisfies (3.1) and as  $\mathcal{B}_{aaa} \subset \mathcal{B}$ , hence  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{\mathcal{B}}$ . If  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aaa}$ , in the same way we prove that  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{\mathcal{B}}$ .
  - (2) Let  $\widetilde{u} = [(u_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{aaa}$  and  $\widetilde{v} = [(v_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{aa}$ . As  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$  and  $(v_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aa}$ , so  $(u_{\varepsilon})_{\varepsilon}$  and  $(v_{\varepsilon})_{\varepsilon}$  satisfy the estimate (3.1). In view of Proposition 2-(2) it follows that  $\forall \varepsilon \in I, \ u_{\varepsilon}v_{\varepsilon} \in \mathcal{B}_{aaa}$ , and for every  $j \in \mathbb{Z}_{+}$ ,

$$\| (u_{\varepsilon}v_{\varepsilon})^{(j)} \|_{L^{\infty}(\mathbb{R})} \leq 2^{j} |u_{\varepsilon}|_{j,\infty,\mathbb{R}} |v_{\varepsilon}|_{j,\infty,\mathbb{R}} \leq 2^{j}c_{1}c_{2}\varepsilon^{-m_{1}}\varepsilon^{-m_{2}},$$

therefore,  $\forall k \in \mathbb{Z}_+$ ,  $\exists m = (m_1 + m_2) \in \mathbb{Z}_+$ ,  $|u_{\varepsilon}v_{\varepsilon}|_{k,\infty,\mathbb{R}} = O(\varepsilon^{-m})$ ,  $\varepsilon \to 0$ , i.e.  $(u_{\varepsilon}v_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$ . The product  $\widetilde{u} \times \widetilde{v}$  is independent on the representatives. Indeed, suppose that

 $(u'_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$  and  $(v'_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aa}$  are others representatives of  $\widetilde{u}$  and  $\widetilde{v}$  respectively, for  $j \in \mathbb{Z}_+$ ,

$$\begin{split} \left\| \left( u_{\varepsilon} v_{\varepsilon} - u'_{\varepsilon} v'_{\varepsilon} \right)^{(j)} \right\|_{L^{\infty}(\mathbb{R})} &= \left\| \left( u_{\varepsilon} v_{\varepsilon} - u'_{\varepsilon} v_{\varepsilon} + u'_{\varepsilon} v_{\varepsilon} - u'_{\varepsilon} v'_{\varepsilon} \right)^{(j)} \right\|_{L^{\infty}(\mathbb{R})}, \\ &\leq \left\| \left( (u_{\varepsilon} - u'_{\varepsilon}) v_{\varepsilon} \right)^{(j)} \right\|_{L^{\infty}(\mathbb{R})} + \left\| \left( u'_{\varepsilon} (v_{\varepsilon} - v'_{\varepsilon})^{(j)} \right\|_{L^{\infty}(\mathbb{R})}, \\ &\leq 2^{j} \left( \left| u_{\varepsilon} - u'_{\varepsilon} \right|_{j,\infty,\mathbb{R}} |v_{\varepsilon}|_{j,\infty,\mathbb{R}} + \left| u'_{\varepsilon} \right|_{j,\infty,\mathbb{R}} \left| v_{\varepsilon} - v'_{\varepsilon} \right|_{j,\infty,\mathbb{R}} \right), \end{split}$$

since  $(u_{\varepsilon} - u'_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aaa}, (v_{\varepsilon} - v'_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aa}$ , then  $\forall k \in \mathbb{Z}_{+}, \forall m \in \mathbb{Z}_{+}, |u_{\varepsilon}v_{\varepsilon} - u'_{\varepsilon}v'_{\varepsilon}|_{k,\infty,\mathbb{R}} = O(\varepsilon^{m})$ , as  $\varepsilon \to 0$ , which implies that  $(u_{\varepsilon}v_{\varepsilon} - u'_{\varepsilon}v'_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aaa}$ .

(3) Let  $\widetilde{u} = [(u_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{aaa}$  and  $\widetilde{v} = [(v_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{L^1}$ , so  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$  and  $(v_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{L^1}$ , that is  $(u_{\varepsilon})_{\varepsilon}$  satisfies the estimate (3.1), and  $(v_{\varepsilon})_{\varepsilon}$  satisfies

$$\forall k \in \mathbb{Z}_+, \ \exists m_1 > 0, \ \exists c > 0, \ \exists \varepsilon_0 \in I, \ \forall \varepsilon < \varepsilon_0, \ |v_\varepsilon|_{k,1,\mathbb{R}} \le c\varepsilon^{-m_1}$$

Proposition 2-(3) gives  $\forall \varepsilon \in I, u_{\varepsilon} * v_{\varepsilon} \in \mathcal{B}_{aaa}$ . Due to Young inequality we obtain

$$\|(u_{\varepsilon} * v_{\varepsilon})^{(j)}\|_{L^{\infty}(\mathbb{R})} \leq \|u_{\varepsilon}^{(j)}\|_{L^{\infty}(\mathbb{R})} \|v_{\varepsilon}\|_{L^{1}(\mathbb{R})}.$$

For every  $k \in \mathbb{Z}_+$ ,

$$|u_{\varepsilon} * v_{\varepsilon}|_{k,\infty,\mathbb{R}} \leq \sum_{j \leq k} \left\| u_{\varepsilon}^{(j)} \right\|_{L^{\infty}(\mathbb{R})} \left\| v_{\varepsilon} \right\|_{L^{1}(\mathbb{R})} \leq |u_{\varepsilon}|_{k,\infty,\mathbb{R}} \left| v_{\varepsilon} \right|_{0,1,\mathbb{R}},$$

consequently,  $\forall k \in \mathbb{Z}_+, \exists m \in \mathbb{Z}_+, |u_{\varepsilon} * v_{\varepsilon}|_{k,\infty,\mathbb{R}} = O(\varepsilon^{-m}), \varepsilon \to 0$ , so  $(u_{\varepsilon} * v_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$ . The convolution  $\widetilde{u} * \widetilde{v}$  does not depend on the representatives. Indeed, suppose that  $(u'_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$  and  $(v'_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{L^1}$  are others representatives of  $\widetilde{u}$  and  $\widetilde{v}$  respectively, for every  $k \in \mathbb{Z}_+$ ,

$$\begin{aligned} \left| u_{\varepsilon} * v_{\varepsilon} - u_{\varepsilon}^{'} * v_{\varepsilon}^{'} \right|_{k,\infty,\mathbb{R}\mathbb{R}} &= \left| u_{\varepsilon} * v_{\varepsilon} - u_{\varepsilon}^{'} * v_{\varepsilon} + u_{\varepsilon}^{'} * v_{\varepsilon} - u_{\varepsilon}^{'} * v_{\varepsilon}^{'} \right|_{k,\infty,\mathbb{R}}, \\ &\leq \left| u_{\varepsilon} - u_{\varepsilon}^{'} \right|_{k,\infty,\mathbb{R}} \left| v_{\varepsilon} \right|_{0,1,\mathbb{R}} + \left| u_{\varepsilon}^{'} \right|_{k,\infty,\mathbb{R}} \left| v_{\varepsilon} - v_{\varepsilon}^{'} \right|_{0,1,\mathbb{R}}, \\ &\text{as } \left( u_{\varepsilon} - u_{\varepsilon}^{'} \right)_{\varepsilon} \in \mathcal{N}_{aaa}, \left( v_{\varepsilon} - v_{\varepsilon}^{'} \right)_{\varepsilon} \in \mathcal{N}_{L^{1}}, \text{ then } \forall k \in \mathbb{Z}_{+}, \forall m \in \mathbb{Z}_{+}, \left| u_{\varepsilon} * v_{\varepsilon} - u_{\varepsilon}^{'} * v_{\varepsilon}^{'} \right|_{k,\infty,\mathbb{R}} = \\ &O(\varepsilon^{m}), \text{ as } \varepsilon \to 0, \text{ which means that } \left( u_{\varepsilon} * v_{\varepsilon} - u_{\varepsilon}^{'} * v_{\varepsilon}^{'} \right)_{\varepsilon} \in \mathcal{N}_{aaa}. \end{aligned}$$

# 4. A GENERALIZED SEELEY THEOREM

We give a result on the extension operators in the context of generalized functions. It is needed in the proof of the decomposition of an asymptotically almost automorphic generalized function. Let's first recall a technical Lemma, see [24].

**Lemma 1.** There are two sequences of real numbers  $(a_l)_{l \in \mathbb{Z}_+}$  and  $(b_l)_{l \in \mathbb{Z}_+}$  such that

(1)  $b_l < 0, \forall l \in \mathbb{Z}_+.$ (2)  $\sum_{l=0}^{+\infty} |a_l| |b_l|^n < +\infty, \forall n \in \mathbb{Z}_+.$ (3)  $\sum_{l=0}^{+\infty} a_l b_l^n = 1, \forall n \in \mathbb{Z}_+.$ (4)  $b_l \to -\infty, \ l \to +\infty.$ 

Define the space

$$\mathcal{B}_{+,0}\left(\mathbb{I}\right) := \left\{ \varphi \in \mathcal{B}(\mathbb{I}) : \forall j \in \mathbb{Z}_+, \ \lim_{x \to +\infty} \varphi^{(j)}(x) = 0 \right\}.$$

The algebra of bounded generalized functions vanishing at infinity on  $\mathbb{I}$  is defined by

$$\mathcal{G}_{+,0}\left(\mathbb{I}
ight):=rac{\mathcal{M}_{+,0}\left(\mathbb{I}
ight)}{\mathcal{N}_{+,0}\left(\mathbb{I}
ight)},$$

where

$$\mathcal{M}_{+,0}\left(\mathbb{I}\right) := \left\{ \left(u_{\varepsilon}\right)_{\varepsilon} \in \left(\mathcal{B}_{+,0}\left(\mathbb{I}\right)\right)^{I} : \forall k \in \mathbb{Z}_{+}, \ \exists m \in \mathbb{Z}_{+}, \ |u_{\varepsilon}|_{k,\infty,\mathbb{I}} = O(\varepsilon^{-m}), \ \varepsilon \to 0 \right\}.$$
$$\mathcal{N}_{+,0}\left(\mathbb{I}\right) := \left\{ \left(u_{\varepsilon}\right)_{\varepsilon} \in \left(\mathcal{B}_{+,0}\left(\mathbb{I}\right)\right)^{I} : \forall k \in \mathbb{Z}_{+}, \ \forall m \in \mathbb{Z}_{+}, \ |u_{\varepsilon}|_{k,\infty,\mathbb{I}} = O(\varepsilon^{m}), \ \varepsilon \to 0 \right\}.$$

**Theorem 2.** The linear extension operator  $\widetilde{E} : \mathcal{G}_{\mathcal{B}}(\mathbb{J}) \longrightarrow \mathcal{G}_{\mathcal{B}}(\mathbb{R}), \ \widetilde{u} = [(u_{\varepsilon})_{\varepsilon}] \longmapsto \widetilde{E}\widetilde{u} = [(Eu_{\varepsilon})_{\varepsilon}], where$ 

$$Eu_{\varepsilon}(x) := \begin{cases} u_{\varepsilon}(x), & \text{if } x \ge 0, \\ \sum_{l=0}^{+\infty} a_{l}u_{\varepsilon}(b_{l}x), & \text{if } x < 0, \end{cases}$$

is well defined and we have  $\widetilde{E}\widetilde{u}_{|\mathbb{J}} = \widetilde{u}$ . In particular,  $\forall \widetilde{u} \in \mathcal{G}_{+,0}(\mathbb{J}), \widetilde{E}\widetilde{u} \in \mathcal{G}_{+,0}(\mathbb{R})$ .

*Proof.* Let  $\widetilde{u} = [(u_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{\mathcal{B}}(\mathbb{J})$ , and  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{\mathcal{B}}(\mathbb{J})$  be a representative of  $\widetilde{u}$ . So  $\forall \varepsilon \in I$ ,  $Eu_{\varepsilon} \in \mathcal{B}(\mathbb{R})$  and  $Eu_{\varepsilon|\mathbb{J}} = u_{\varepsilon}$ . Indeed, if x < 0, then  $b_l x > 0$ ,  $\forall l \in \mathbb{Z}_+$  in view of Lemma 1-(1). Moreover according to Lemma 1-(2) and as  $u_{\varepsilon} \in \mathcal{B}(\mathbb{J}), \forall \varepsilon \in I$ , hence  $\forall n \in \mathbb{Z}_+, \forall \varepsilon \in I, \forall x < 0$ ,

(4.1) 
$$\left| \left( Eu_{\varepsilon} \right)^{(n)} (x) \right| \leq \left\| u_{\varepsilon}^{(n)} \right\|_{L^{\infty}(\mathbb{J})} \sum_{l=0}^{+\infty} |a_l| |b_l|^n < +\infty,$$

consequently,  $\forall n \in \mathbb{Z}_+$ , the series

$$(Eu_{\varepsilon})^{(n)}(x) = \sum_{l=0}^{+\infty} a_l b_l^n u_{\varepsilon}^{(n)}(b_l x), \varepsilon \in I,$$

absolutely converge. Furthermore due to Lemma 1-(3),

$$\lim_{\substack{x \to 0 \\ <}} (Eu_{\varepsilon})^{(n)}(x) = \sum_{l=0}^{+\infty} a_l b_l^n \lim_{\substack{x \to 0 \\ <}} u_{\varepsilon}^{(n)}(b_l x),$$
$$= u_{\varepsilon}^{(n)}(0) \sum_{l=0}^{+\infty} a_l b_l^n = u_{\varepsilon}^{(n)}(0).$$

so  $\forall \varepsilon \in I, Eu_{\varepsilon} \in \mathcal{E}(\mathbb{R})$ . As  $\forall \varepsilon \in I, u_{\varepsilon} \in \mathcal{B}(\mathbb{J})$  and by (4.1), it follows that  $\forall n \in \mathbb{Z}_+, \forall \varepsilon \in \mathcal{B}(\mathbb{J})$  $I, (Eu_{\varepsilon})^{(n)} \in L^{\infty}(\mathbb{R})$ . In order to show that  $(Eu_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{\mathcal{B}}(\mathbb{R})$ , we prove the following estimate

(4.2) 
$$\forall \varepsilon \in I, \ \forall k \in \mathbb{Z}_+, \ \exists C_k > 0, \ |Eu_\varepsilon|_{k,\infty,\mathbb{R}} \le C_k |u_\varepsilon|_{k,\infty,\mathbb{J}}.$$

...

Indeed, it is clear that

$$\left\| \left( E u_{\varepsilon} \right)^{(n)} \right\|_{L^{\infty}(\mathbb{J})} = \left\| u_{\varepsilon}^{(n)} \right\|_{L^{\infty}(\mathbb{J})},$$

and the estimate (4.1), gives

$$\left\| (Eu_{\varepsilon})^{(n)} \right\|_{L^{\infty}(\mathbb{R}\setminus\mathbb{J})} \leq \left\| u_{\varepsilon}^{(n)} \right\|_{L^{\infty}(\mathbb{J})} \sum_{l=0}^{+\infty} |a_{l}| |b_{l}|^{n},$$

therefore

$$\left\| (Eu_{\varepsilon})^{(n)} \right\|_{L^{\infty}(\mathbb{R})} \leq C_{n} \left\| u_{\varepsilon}^{(n)} \right\|_{L^{\infty}(\mathbb{J})},$$
  
where  $C_{n} = \max\left(1, \sum_{l=0}^{+\infty} |a_{l}| |b_{l}|^{n}\right)$ . So  $\forall k \in \mathbb{Z}_{+}, \ \exists C_{k} := \sum_{n \leq k} C_{n} < +\infty, \ \forall \varepsilon \in I, \ u_{\varepsilon} \in \mathcal{B}\left(\mathbb{J}\right),$   
(4.3)  
 $|Eu_{\varepsilon}|_{k,\infty,\mathbb{R}} \leq C_{k} |u_{\varepsilon}|_{k,\infty,\mathbb{J}},$ 

this implies  $(Eu_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{\mathcal{B}}(\mathbb{R})$ . The definition of  $\widetilde{E}\widetilde{u}$  is independent on representatives. Indeed, if  $(u_{\varepsilon})_{\varepsilon}$  and  $(v_{\varepsilon})_{\varepsilon}$  are representatives of  $\widetilde{u}$ , hence by (4.3),

$$\forall \varepsilon \in I, \ \forall k \in \mathbb{Z}_+, \ \exists C_k > 0, \ |Eu_\varepsilon - Ev_\varepsilon|_{k,\infty,\mathbb{R}} \le C_k |u_\varepsilon - v_\varepsilon|_{k,\infty,\mathbb{J}}.$$

As  $(u_{\varepsilon} - v_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{\mathcal{B}}(\mathbb{J})$  then  $\forall k \in \mathbb{Z}_+, \ \forall m > 0, \ |Eu_{\varepsilon} - Ev_{\varepsilon}|_{k \infty \mathbb{R}} = O(\varepsilon^m), \ \varepsilon \to 0$ , which shows that  $(Eu_{\varepsilon} - Ev_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{\mathcal{B}}(\mathbb{R})$ .

We have  $\widetilde{E}\widetilde{u}_{|\mathbb{J}} = \widetilde{u}$  in  $\mathcal{G}_{\mathcal{B}}(\mathbb{J})$ . Indeed, as  $\widetilde{E}\widetilde{u} = [(Eu_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{\mathcal{B}}(\mathbb{R})$  and  $\widetilde{u} = [(u_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{\mathcal{B}}(\mathbb{J})$ , therefore  $\widetilde{E}\widetilde{u}_{|\mathbb{J}} - \widetilde{u} := \left[ \left( Eu_{\varepsilon|\mathbb{J}} \right)_{\varepsilon} \right] - \widetilde{u} = \left[ \left( u_{\varepsilon} \right)_{\varepsilon} \right] - \widetilde{u} = \widetilde{u} - \widetilde{u} = 0$  in  $\mathcal{G}_{\mathcal{B}}\left( \mathbb{J} \right)$ .

If  $\widetilde{u} = [(u_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{+,0}(\mathbb{J}) \subset \mathcal{G}_{\mathcal{B}}(\mathbb{J})$ , then  $\widetilde{E}\widetilde{u} = [(Eu_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{\mathcal{B}}(\mathbb{R})$ . So  $\forall \varepsilon \in I, Eu_{\varepsilon} \in \mathcal{B}(\mathbb{R})$ . The fact that  $\forall \varepsilon \in I, Eu_{\varepsilon} = u_{\varepsilon}$  on  $\mathbb{J}$ , and  $\forall \varepsilon \in I, u_{\varepsilon} \in \mathcal{B}_{+,0}(\mathbb{J})$  implies  $\lim_{x \to +\infty} Eu_{\varepsilon}(x) = 0$ , i.e.  $\forall \varepsilon \in I, Eu_{\varepsilon} \in \mathcal{C}_{+,0}$ . By ([12], Proposition 5 – (5)), we obtain that  $\forall \varepsilon \in I, Eu_{\varepsilon} \in \mathcal{C}_{+,0} \cap \mathcal{B}(\mathbb{R}) =$  $\mathcal{B}_{+,0}(\mathbb{R})$ , it follows that  $E\widetilde{u} \in \mathcal{G}_{+,0}(\mathbb{R})$ . 

### CHIKH BOUZAR AND MERYEM SLIMANI

### 5. The decomposition

In this section we show that an asymptotically almost automorphic generalized function is uniquely decomposed as in the classical case.

**Theorem 3.** Let  $\widetilde{u} \in \mathcal{G}_{aaa}(\mathbb{R})$  then there exist  $\widetilde{v} \in \mathcal{G}_{aa}(\mathbb{R})$  and  $\widetilde{w} \in \mathcal{G}_{+,0}(\mathbb{R})$  such that  $\widetilde{u} = \widetilde{v} + \widetilde{w}$  on  $\mathbb{J}$ , and the decomposition is unique on  $\mathbb{J}$ .

Proof. Let  $\widetilde{u} = [(u_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{aaa}$ , so  $\forall \varepsilon \in I, \forall j \in \mathbb{Z}_+, u_{\varepsilon}^{(j)} \in \mathcal{C}_{aaa}$ . Then there exist  $v_{\varepsilon,j} \in \mathcal{C}_{aa}, w_{\varepsilon,j} \in \mathcal{C}_{+,0}$ , such that  $\forall j \in \mathbb{N}, u_{\varepsilon}^{(j)} = (v_{\varepsilon,j} + w_{\varepsilon,j}) \in \mathcal{C}_{aaa}$  on  $\mathbb{J}$ , and for  $j = 0, u_{\varepsilon} = v_{\varepsilon} + w_{\varepsilon}$  on  $\mathbb{J}$ . By ([12], Proposition 8), it holds that  $\forall j \in \mathbb{N}, v_{\varepsilon,j} = (v_{\varepsilon})^{(j)}$  on  $\mathbb{R}$  and  $w_{\varepsilon,j} = (w_{\varepsilon})^{(j)}$  on  $\mathbb{J}$ , which gives  $v_{\varepsilon} \in \mathcal{B}_{aa}$  and  $w_{\varepsilon} \in \mathcal{B}_{+,0}(\mathbb{J})$ . Let's show that  $(v_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aa}$ . As  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$ , therefore

(5.1) 
$$\forall k \in \mathbb{Z}_+, \ \exists m \in \mathbb{Z}_+, \ \exists c > 0, \ \exists \varepsilon_0 \in I, \ \forall \varepsilon < \varepsilon_0, \ |u_\varepsilon|_{k,\infty,\mathbb{R}} \le c\varepsilon^{-m}.$$

Due to ([12], Proposition 3 - (5)), we obtain

(5.2) 
$$\forall j \in \mathbb{Z}_+, \left\| v_{\varepsilon}^{(j)} \right\|_{L^{\infty}(\mathbb{R})} \le \left\| u_{\varepsilon}^{(j)} \right\|_{L^{\infty}(\mathbb{J})}$$

it follows that

(5.3) 
$$\forall k \in \mathbb{Z}_+, \exists m \in \mathbb{Z}_+, \exists c > 0, \exists \varepsilon_0 \in I, \forall \varepsilon < \varepsilon_0, |v_\varepsilon|_{k,\infty,\mathbb{R}} \le c\varepsilon^{-m}$$

this means that  $(v_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aa}$ . If  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aaa}$  then

(5.4) 
$$\forall k \in \mathbb{Z}_+, \forall m \in \mathbb{Z}_+, \exists c > 0, \exists \varepsilon_0 \in I, \forall \varepsilon < \varepsilon_0, |u_\varepsilon|_{k,\infty,\mathbb{R}} \le c\varepsilon^m,$$

and from (5.2) it holds  $(v_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aa}$ . Consequently,  $\tilde{v} = [(v_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{aa}$ . On the other hand, we have

(5.5) 
$$\forall j \in \mathbb{Z}_+, \left\| w_{\varepsilon}^{(j)} \right\|_{L^{\infty}(\mathbb{J})} \le \left\| u_{\varepsilon}^{(j)} \right\|_{L^{\infty}(\mathbb{J})} + \left\| v_{\varepsilon}^{(j)} \right\|_{L^{\infty}(\mathbb{J})}.$$

The estimates (5.1), (5.3) and (5.5) give

$$\forall k \in \mathbb{Z}_+, \ \exists m \in \mathbb{Z}_+, \ \exists c > 0, \ \exists \varepsilon_0 \in I, \ \forall \varepsilon < \varepsilon_0, |w_\varepsilon|_{k,\infty,\mathbb{J}} \le 2c\varepsilon^{-m}$$

hence  $(w_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{+,0}(\mathbb{J})$ . If  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aaa}$ , then  $(w_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{+,0}(\mathbb{J})$  follows from (5.4) and (5.5). Thus  $\widetilde{w} = [(w_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{+,0}(\mathbb{J})$ . By Theorem 2 extending  $\widetilde{w} \in \mathcal{G}_{+,0}(\mathbb{J})$  to  $\widetilde{E}\widetilde{w} \in \mathcal{G}_{+,0}(\mathbb{R})$  with  $\widetilde{E}\widetilde{w} = \widetilde{w}$  on  $\mathbb{J}$ . Finally,  $\widetilde{u} = \widetilde{v} + \widetilde{w}$  on  $\mathbb{J}$ .

If  $\tilde{u} \in \mathcal{G}_{aaa}$  has two decompositions, i.e.

$$\widetilde{u} = \widetilde{v}_i + \widetilde{w}_i$$
 on  $\mathbb{J}, i = 1, 2,$ 

where  $\widetilde{v}_i \in \mathcal{G}_{aa}$  and  $\widetilde{w}_i \in \mathcal{G}_{+,0} := \mathcal{G}_{+,0}(\mathbb{R})$ . Let  $(v_{\varepsilon,i})_{\varepsilon} \in \mathcal{M}_{aa}$  and  $(w_{\varepsilon,i})_{\varepsilon} \in \mathcal{M}_{+,0}$  be respectively representatives of  $\widetilde{v}_i$  and  $\widetilde{w}_i, i = 1, 2$ . So  $(v_{\varepsilon,1} - v_{\varepsilon,2})_{\varepsilon} + (w_{\varepsilon,1} - w_{\varepsilon,2})_{\varepsilon} \in \mathcal{N}_{\mathcal{B}}(\mathbb{J})$ , i.e.  $\forall k \in \mathbb{Z}_+, \forall m > 0, \exists c > 0, \exists \varepsilon_0 \in I, \forall \varepsilon < \varepsilon_0,$ 

(5.6) 
$$|v_{\varepsilon,1} - v_{\varepsilon,2} + w_{\varepsilon,1} - w_{\varepsilon,2}|_{k,\infty,\mathbb{J}} \le c\varepsilon^m.$$

Due to ([12], Proposition 9) as  $\forall \varepsilon \in I, v_{\varepsilon,i} \in \mathcal{B}_{aa}, i = 1, 2$ , for any real sequence  $(s_m)_{m \in \mathbb{N}}$ , such that  $s_m \to +\infty$  there exist  $(s_{m_{l(\varepsilon)}})_l$  a subsequence of  $(s_m)_{m \in \mathbb{N}}$  and  $g_{\varepsilon,i}, i = 1, 2$ , such that  $\forall x \in \mathbb{R}, \forall j \in \mathbb{Z}_+,$ 

$$g_{\varepsilon,i}^{(j)}(x) := \lim_{l \to +\infty} v_{\varepsilon,i}^{(j)} \left( x + s_{m_{l(\varepsilon)}} \right) \quad \text{and} \quad \lim_{l \to +\infty} g_{\varepsilon,i}^{(j)} \left( x - s_{m_{l(\varepsilon)}} \right) = v_{\varepsilon,i}^{(j)}(x).$$

Furthermore, as  $\forall \varepsilon \in I, w_{\varepsilon,i} \in \mathcal{B}_{+,0}, i = 1, 2,$ 

$$\lim_{l \to +\infty} w_{\varepsilon,i}^{(j)} \left( x + s_{m_{l(\varepsilon)}} \right) = 0, \forall x \in \mathbb{R}, \forall j \in \mathbb{Z}_{+}$$

By using (5.6) we have  $\forall m > 0, \exists c > 0, \exists \varepsilon_0 \in I, \forall \varepsilon < \varepsilon_0, \forall x \ge -s_{m_{l(\varepsilon)}}, \forall x \ge -$ 

$$\left| v_{\varepsilon,1}^{(j)} \left( x + s_{m_{l(\varepsilon)}} \right) - v_{\varepsilon,2}^{(j)} \left( x + s_{m_{l(\varepsilon)}} \right) + w_{\varepsilon,1}^{(j)} \left( x + s_{m_{l(\varepsilon)}} \right) - w_{\varepsilon,2}^{(j)} \left( x + s_{m_{l(\varepsilon)}} \right) \right| \le c\varepsilon^m,$$

so when  $l \to +\infty$  we obtain  $\forall m > 0, \exists c > 0, \exists \varepsilon_0 \in I, \forall \varepsilon < \varepsilon_0, \forall x \ge -s_{m_{l(\varepsilon)}}, \forall x$ 

$$\left|g_{\varepsilon,1}^{(j)}(x) - g_{\varepsilon,2}^{(j)}(x)\right| \le c\varepsilon^m$$

by taking the translate  $-s_{m_{l(\varepsilon)}}$  and let  $l \to +\infty$  we get  $\forall m > 0, \exists c > 0, \exists \varepsilon_0 \in I, \forall \varepsilon < \varepsilon_0, \forall x \ge 0, \forall x = x = x, x = x, x = x, x = x$ 

$$\left|v_{\varepsilon,1}^{(j)}(x) - v_{\varepsilon,2}^{(j)}(x)\right| \le c\varepsilon^m.$$

By ([12], Proposition 3 - (5)), it follows

(5.7) 
$$\forall k \in \mathbb{Z}_+, \forall m > 0, \exists c > 0, \exists \varepsilon_0 \in I, \forall \varepsilon < \varepsilon_0, |v_{\varepsilon,1} - v_{\varepsilon,2}|_{k,\infty,\mathbb{R}} \le c\varepsilon^m,$$

which shows that  $(v_{\varepsilon,1} - v_{\varepsilon,2})_{\varepsilon} \in \mathcal{N}_{\mathcal{B}}(\mathbb{R})$ , so  $\tilde{v}_1 = \tilde{v}_2$  in  $\mathcal{G}_{\mathcal{B}}(\mathbb{R})$ . From (5.6) and (5.7) it holds that  $(w_{\varepsilon,1} - w_{\varepsilon,2})_{\varepsilon} \in \mathcal{N}_{\mathcal{B}}(\mathbb{J})$ , i.e.  $\tilde{w}_1 = \tilde{w}_2$  on  $\mathbb{J}$ .

Notation 4. Let  $\tilde{u} \in \mathcal{G}_{aaa}$  and  $\tilde{u} = \tilde{v} + \tilde{w}$  on  $\mathbb{J}$ , where  $\tilde{v} \in \mathcal{G}_{aa}$  and  $\tilde{w} \in \mathcal{G}_{+,0}$ , then  $\tilde{v}$  and  $\tilde{w}$  are called respectively the principal term and the corrective term of  $\tilde{u}$  and we denote them respectively  $\tilde{u}_{aa}$  and  $\tilde{u}_{cor}$ . Also  $\tilde{u} = (\tilde{u}_{aa} + \tilde{u}_{cor}) \in \mathcal{G}_{aaa}$  means that  $\tilde{u}_{aa} \in \mathcal{G}_{aa}$ ,  $\tilde{u}_{cor} \in \mathcal{G}_{+,0}$  and  $\tilde{u} = \tilde{u}_{aa} + \tilde{u}_{cor}$  on  $\mathbb{J}$ .

## 6. Non linear operations

The algebra of tempered generalized functions on  $\mathbb{C}$  denoted by  $\mathcal{G}_{\tau}(\mathbb{C})$ , see [17] and [20] for more details, is the quotient algebra

$$\mathcal{G}_{\tau}(\mathbb{C}) := rac{\mathcal{M}_{\tau}(\mathbb{C})}{\mathcal{N}_{\tau}(\mathbb{C})},$$

where

$$\mathcal{M}_{\tau}(\mathbb{C}) := \left\{ \begin{array}{c} \left(f_{\varepsilon}\right)_{\varepsilon} \in \left(\mathcal{E}\left(\mathbb{R}^{2}\right)\right)^{I} : \forall j \in \mathbb{Z}_{+}^{2}, \exists m \in \mathbb{Z}_{+}, \\ \sup_{x \in \mathbb{R}^{2}} \left(1 + |x|\right)^{-m} \left|f_{\varepsilon}^{(j)}(x)\right| = O\left(\varepsilon^{-m}\right), \varepsilon \to 0 \end{array} \right\}.$$

CHIKH BOUZAR AND MERYEM SLIMANI

$$\mathcal{N}_{\tau}(\mathbb{C}) := \left\{ \begin{array}{l} (f_{\varepsilon})_{\varepsilon} \in (\mathcal{E}(\mathbb{R}^{2}))^{I} : \forall j \in \mathbb{Z}_{+}^{2}, \exists n \in \mathbb{Z}_{+}, \forall m \in \mathbb{Z}_{+}, \\ \sup_{x \in \mathbb{R}^{2}} (1 + |x|)^{-n} \left| f_{\varepsilon}^{(j)}(x) \right| = O(\varepsilon^{m}), \varepsilon \to 0 \end{array} \right\}.$$

**Example 2.** Any polynomial function is a tempered generalized function.

**Theorem 4.** Let  $\widetilde{u} = [(u_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{aaa}$  and  $\widetilde{F} = [(f_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{\tau}(\mathbb{C})$ , then  $\widetilde{F} \circ \widetilde{u} := [(f_{\varepsilon} \circ u_{\varepsilon})_{\varepsilon}]$ 

is a well-defined element of  $\mathcal{G}_{aaa}$ . The principal term and the corrective term of  $\widetilde{F} \circ \widetilde{u}$  are respectively  $\widetilde{F}(\widetilde{u}_{aa})$  and  $\widetilde{F}(\widetilde{u}_{aa} + \widetilde{u}_{cor}) - \widetilde{F}(\widetilde{u}_{aa})$ , where  $\widetilde{u} = \widetilde{u}_{aa} + \widetilde{u}_{cor}$  on  $\mathbb{J}$ .

*Proof.* Let  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$  and  $(f_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{\tau}(\mathbb{C})$ , by the classical Faà di Bruno formula, we have  $\forall j \in \mathbb{Z}_+$ ,

(6.1) 
$$\frac{(f_{\varepsilon} \circ u_{\varepsilon})^{(j)}(x)}{j!} = \sum_{\substack{l_1+2l_2+\dots+jl_j=j\\r=l_1+\dots+l_j}} \frac{f_{\varepsilon}^{(r)}(u_{\varepsilon}(x))}{l_1!\dots l_j!} \prod_{i=1}^j \left(\frac{u_{\varepsilon}^{(i)}(x)}{i!}\right)^{l_i}.$$

As  $\forall \varepsilon \in I$ ,  $\forall j \in \mathbb{Z}_+$ ,  $u_{\varepsilon}^{(j)} \in \mathcal{C}_{aaa}$  and  $f_{\varepsilon} \in \mathcal{E}(\mathbb{R})$ , it follows by ([12], Proposition 3 – (4)), that  $f_{\varepsilon}^{(r)}(u_{\varepsilon}) \in \mathcal{C}_{aaa}$ , and since  $\mathcal{C}_{aaa}$  is an algebra, then  $\forall \varepsilon \in I$ ,  $f_{\varepsilon} \circ u_{\varepsilon} \in \mathcal{B}_{aaa}$ . As  $(u_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$ , then  $\forall k \in \mathbb{Z}_+, \exists n_k \in \mathbb{Z}_+, \exists c_k > 0, \exists \varepsilon_k \in I, \forall \varepsilon < \varepsilon_k, |u_{\varepsilon}|_{k,\infty,\mathbb{R}} \leq c_k \varepsilon^{-n_k}$ .

The fact that  $(f_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{\tau}(\mathbb{C})$  gives

$$\forall j \in \mathbb{Z}_+, \exists N_j \in \mathbb{Z}_+, \exists C_j > 0, \exists \varepsilon'_j \in I, \forall \varepsilon < \varepsilon'_j, \left\| f_{\varepsilon}^{(j)}(u_{\varepsilon}) \right\|_{L^{\infty}(\mathbb{R})} \le C_j \varepsilon^{-N_j} \left\| 1 + u_{\varepsilon} \right\|_{L^{\infty}(\mathbb{R})}^{N_j}$$

Consequently, by (6.1) we obtain

$$\frac{\left\| (f_{\varepsilon} \circ u_{\varepsilon})^{(j)} \right\|_{L^{\infty}(\mathbb{R})}}{j!} \leq \sum_{\substack{l_1+2l_2+\dots+jl_j=j\\r=l_1+\dots+l_j}} \frac{C_r \varepsilon^{-N_r} \left\| 1+u_{\varepsilon} \right\|_{L^{\infty}(\mathbb{R})}^{N_r}}{l_1! \cdots l_j!} \prod_{i=1}^j \left( \frac{\left\| u_{\varepsilon}^{(i)} \right\|_{L^{\infty}(\mathbb{R})}}{i!} \right)^{l_i},$$

hence there exists c > 0,

$$\frac{\left\|\left(f_{\varepsilon} \circ u_{\varepsilon}\right)^{(j)}\right\|_{L^{\infty}(\mathbb{R})}}{j!} \leq \sum_{\substack{l_{1}+2l_{2}+\dots+jl_{j}=j\\r=l_{1}+\dots+l_{j}}} \frac{c\varepsilon^{-N_{r}(1+n_{0})}}{l_{1}!\dots l_{j}!} \prod_{i=1}^{j} \left(\frac{c_{i}\varepsilon^{-n_{i}}}{i!}\right)^{l_{i}},$$

$$\leq \sum_{\substack{l_{1}+2l_{2}+\dots+jl_{j}=j\\r=l_{1}+\dots+l_{j}}} \frac{c\varepsilon^{-(N_{r}(1+n_{0})+\sum_{i=1}^{j}n_{i}l_{i})}}{l_{1}!\dots l_{j}!} \prod_{i=1}^{j} \left(\frac{c_{i}}{i!}\right)^{l_{i}},$$

$$\leq C'\varepsilon^{-m},$$

14

where

Finally, with  $C = C' \sum_{j \leq k} j!$ , it holds

$$\forall k \in \mathbb{Z}_{+}, \exists m \in \mathbb{Z}_{+}, \exists C > 0, \exists \varepsilon^{"} = \inf_{1 \leq i \leq j \leq k} \left( \varepsilon_{i}, \varepsilon_{j}^{'} \right), \forall \varepsilon < \varepsilon^{"}, \left| f_{\varepsilon} \circ u_{\varepsilon} \right|_{k, \infty, \mathbb{R}} \leq C \varepsilon^{-m},$$

which means that  $(f_{\varepsilon} \circ u_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$ . This composition does not depend on the representatives. Indeed, suppose that  $(v_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$  and  $(g_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{\tau}(\mathbb{C})$  are others representatives of  $\widetilde{u}$  and  $\widetilde{F}$  respectively. Set  $(n_{\varepsilon})_{\varepsilon} := ((v_{\varepsilon})_{\varepsilon} - (u_{\varepsilon})_{\varepsilon}) \in \mathcal{N}_{aaa}$  and  $(m_{\varepsilon})_{\varepsilon} := ((f_{\varepsilon})_{\varepsilon} - (g_{\varepsilon})_{\varepsilon}) \in \mathcal{N}_{\tau}(\mathbb{C})$ . To show that  $(f_{\varepsilon} \circ u_{\varepsilon} - g_{\varepsilon} \circ v_{\varepsilon})_{\varepsilon} \in \mathcal{N}_{aaa}$ , since  $(f_{\varepsilon} \circ u_{\varepsilon} - g_{\varepsilon} \circ v_{\varepsilon})_{\varepsilon} \in \mathcal{M}_{aaa}$ , according to Proposition 3–(1), it is enough to prove that  $(f_{\varepsilon} \circ u_{\varepsilon} - g_{\varepsilon} \circ v_{\varepsilon})_{\varepsilon}$  satisfies (3.2). Indeed, we have

$$\begin{aligned} |f_{\varepsilon} \circ u_{\varepsilon} - g_{\varepsilon} \circ v_{\varepsilon}|_{0,\infty,\mathbb{R}} &\leq |f_{\varepsilon} \circ u_{\varepsilon} - f_{\varepsilon} \circ v_{\varepsilon}|_{0,\infty,\mathbb{R}} + |f_{\varepsilon} \circ v_{\varepsilon} - g_{\varepsilon} \circ v_{\varepsilon}|_{0,\infty,\mathbb{R}}, \\ &= |f_{\varepsilon}(u_{\varepsilon}) - f_{\varepsilon}(u_{\varepsilon} + n_{\varepsilon})|_{0,\infty,\mathbb{R}} + |m_{\varepsilon}(v_{\varepsilon})|_{0,\infty,\mathbb{R}}, \\ &\leq |n_{\varepsilon}|_{0,\infty,\mathbb{R}} \left| f_{\varepsilon}'(u_{\varepsilon}) \right|_{0,\infty,\mathbb{R}} + |m_{\varepsilon}(v_{\varepsilon})|_{0,\infty,\mathbb{R}}. \end{aligned}$$

It is clear that  $\forall k \in \mathbb{Z}_+$ ,  $|n_{\varepsilon}|_{0,\infty,\mathbb{R}} |f'_{\varepsilon}(u_{\varepsilon})|_{0,\infty,\mathbb{R}} = O(\varepsilon^k)$ ,  $\varepsilon \to 0$ , and also  $\forall l \in \mathbb{Z}_+$ ,  $|m_{\varepsilon}(v_{\varepsilon})|_{0,\infty,\mathbb{R}} = O(\varepsilon^l)$ ,  $\varepsilon \to 0$ . Therefore,  $\forall q \in \mathbb{Z}_+$ ,  $|f_{\varepsilon} \circ u_{\varepsilon} - g_{\varepsilon} \circ v_{\varepsilon}|_{0,\infty,\mathbb{R}} = O(\varepsilon^q)$ ,  $\varepsilon \to 0$ .

Let  $\widetilde{u} = (\widetilde{u}_{aa} + \widetilde{u}_{cor}) \in \mathcal{G}_{aaa}$ . As  $\widetilde{F} \circ \widetilde{u} = \widetilde{F}(\widetilde{u}_{aa}) + (\widetilde{F}(\widetilde{u}) - \widetilde{F}(\widetilde{u}_{aa}))$ , then  $\widetilde{F} \circ \widetilde{u} = \widetilde{F}(\widetilde{u}_{aa}) + (\widetilde{F}(\widetilde{u}_{aa} + \widetilde{u}_{cor}) - \widetilde{F}(\widetilde{u}_{aa}))$  on  $\mathbb{J}$ . In view of ([10], Proposition 9) we obtain  $\widetilde{F}(\widetilde{u}_{aa}) \in \mathcal{G}_{aa}$ . It remains to prove that  $\widetilde{F}(\widetilde{u}_{aa} + \widetilde{u}_{cor}) - \widetilde{F}(\widetilde{u}_{aa}) \in \mathcal{G}_{+,0}$ . Since  $\mathcal{G}_{aaa}$  and  $\mathcal{G}_{aa}$  are contained in  $\mathcal{G}_{\mathcal{B}}$  then  $\widetilde{F}(\widetilde{u}_{aa} + \widetilde{u}_{cor}) - \widetilde{F}(\widetilde{u}_{aa}) \in \mathcal{G}_{\mathcal{B}}$ . It suffices to show that  $\forall \varepsilon \in I, f_{\varepsilon}(u_{aa,\varepsilon} + u_{cor,\varepsilon}) - f_{\varepsilon}(u_{aa,\varepsilon}) \in \mathcal{B}_{+,0}$ , where  $(f_{\varepsilon})_{\varepsilon}, (u_{aa,\varepsilon})_{\varepsilon}$  and  $(u_{cor,\varepsilon})_{\varepsilon}$  are respective representatives of  $\widetilde{F}, \widetilde{u}_{aa}$  and  $\widetilde{u}_{cor}$ . The classical result on composition of asymptotically almost automorphic function with continuous function shows that the corrective term of  $f_{\varepsilon}(u_{aa,\varepsilon} + u_{cor,\varepsilon})$  is  $f_{\varepsilon}(u_{aa,\varepsilon} + u_{cor,\varepsilon}) - f_{\varepsilon}(u_{aa,\varepsilon})$  and the fact that  $\widetilde{F}(\widetilde{u}_{aa} + \widetilde{u}_{cor}) - \widetilde{F}(\widetilde{u}_{aa}) \in \mathcal{G}_{\mathcal{B}}$  gives  $\forall \varepsilon \in I, (f_{\varepsilon}(u_{aa,\varepsilon} + u_{cor,\varepsilon}) - f_{\varepsilon}(u_{aa,\varepsilon})) \in \mathcal{B}$ . By ([12], Proposition 5 - (5)), we have  $\forall \varepsilon \in I, f_{\varepsilon}(u_{aa,\varepsilon} + u_{cor,\varepsilon}) - f_{\varepsilon}(u_{aa,\varepsilon}) \in \mathcal{C}_{+,0} \cap \mathcal{B} = \mathcal{B}_{+,0}$ .

### 7. LINEAR NEUTRAL DIFFERENCE DIFFERENTIAL SYSTEMS

We consider linear neutral difference differential systems for the unknown vector function  $\widetilde{u} = (\widetilde{u}_1, \ldots, \widetilde{u}_n)$ ,

(7.1) 
$$L_{\omega}\widetilde{u} := \sum_{i=0}^{p} \sum_{j=0}^{q} \widetilde{A}_{ij} \left(\tau_{\omega_{j}}\widetilde{u}\right)^{(i)} + \widetilde{K} * \widetilde{u} = \widetilde{f},$$

where  $\widetilde{f} = \left(\widetilde{f}_1, \ldots, \widetilde{f}_n\right) \in \left(\mathcal{G}_{aaa}\right)^n$ ,  $\omega = \left(\omega_j\right)_{0 \le j \le q} \subset \mathbb{R}^q_+$  and for  $i \le p, j \le q$ ,  $\widetilde{A}_{ij} = \left(\widetilde{A}_{ij}^{rl}\right)_{1 \le r, l \le n}$ and  $\widetilde{K} = \left(\widetilde{K}^{rl}\right)_{1 \le r, l \le n}$  are square matrices of almost automorphic generalized functions and  $L^1$ -generalized functions respectively.

**Lemma 2.** If  $\widetilde{u} \in (\mathcal{G}_{aaa})^n$  then  $L_{\omega}\widetilde{u} \in (\mathcal{G}_{aaa})^n$ .

*Proof.* If  $\tilde{u} \in (\mathcal{G}_{aaa})^n$  then due to results of Proposition 5 we obtain that  $\forall i \leq p, j \leq q, \forall \omega = (\omega_j)_{0 \leq j \leq q} \subset \mathbb{R}^q_+$ ,

$$\left(\widetilde{A}_{ij}\left(\tau_{\omega_{j}}\widetilde{u}\right)^{(i)}\right)\in\left(\mathcal{G}_{aaa}\right)^{n}$$

and also

$$\widetilde{K} * \widetilde{u} \in \left(\mathcal{G}_{aaa}\right)^n,$$

 $\mathbf{SO}$ 

$$\left(\sum_{i=0}^{p}\sum_{j=0}^{q}\widetilde{A}_{ij}\left(\tau_{\omega_{j}}\widetilde{u}\right)^{(i)}+\widetilde{K}*\widetilde{u}\right)\in\left(\mathcal{G}_{aaa}\right)^{n},$$

i.e.  $L_{\omega}\widetilde{u} \in (\mathcal{G}_{aaa})^n$ .

**Definition 4.** A generalized function  $\widetilde{u} \in (\mathcal{G}_{\mathcal{B}})^n$  is said a generalized solution of (7.1) on  $\mathbb{J}$  if it satisfies

$$\left(\sum_{l=1}^{n}\sum_{i=1}^{p}\sum_{j=1}^{q}A_{ij,\varepsilon}^{rl}\left(\tau_{\omega_{j}}u_{l,\varepsilon}\right)^{(i)}+K_{\varepsilon}^{rl}*u_{l,\varepsilon}-f_{r,\varepsilon}\right)_{\varepsilon}\in\mathcal{N}_{\mathcal{B}}\left(\mathbb{J}\right), r=1,...,n_{\varepsilon}$$

where  $(u_{l,\varepsilon})_{\varepsilon}$ ,  $(A_{ij,\varepsilon}^{rl})_{\varepsilon}$ ,  $(K_{\varepsilon}^{rl})_{\varepsilon}$  and  $(f_{r,\varepsilon})_{\varepsilon}$ , r, l = 1, ..., n, are respective representatives of  $\tilde{u}_r, \tilde{A}_{ij}^{rl}, \tilde{K}^{rl}$  and  $\tilde{f}_r$ .

We give the main result of this section.

**Theorem 5.** Let  $\tilde{f} = (\tilde{f}_{aa} + \tilde{f}_{cor}) \in (\mathcal{G}_{aaa})^n$ , the equation (7.1) admits a generalized solution  $\tilde{u} \in (\mathcal{G}_{aaa})^n$  on  $\mathbb{J}$  if and only if there exist  $\tilde{v} \in (\mathcal{G}_{aa})^n$  and  $\tilde{w} \in (\mathcal{G}_{+,0})^n$  such that

(7.2) 
$$L_{\omega}\widetilde{v} = \widetilde{f}_{aa} \quad on \quad \mathbb{R},$$

and

(7.3) 
$$L_{\omega}\widetilde{w} = \widetilde{f}_{cor} \quad on \quad \mathbb{J}.$$

*Proof.* If  $\tilde{u} = (\tilde{u}_{aa} + \tilde{u}_{cor}) \in (\mathcal{G}_{aaa})^n$  is a generalized solution of (7.1), then the equations (7.1) explicitly are written as follows

$$\begin{cases} \sum_{l=1}^{n} \left( \sum_{i=0}^{p} \sum_{j=0}^{q} \widetilde{A}_{ij}^{1l} \left( \tau_{\omega_{j}} \widetilde{u}_{l} \right)^{(i)} + \widetilde{K}^{1l} * \widetilde{u}_{l} \right) = \widetilde{f}_{1}. \\ \vdots \\ \sum_{l=1}^{n} \left( \sum_{i=0}^{p} \sum_{j=0}^{q} \widetilde{A}_{ij}^{nl} \left( \tau_{\omega_{j}} \widetilde{u}_{l} \right)^{(i)} + \widetilde{K}^{nl} * \widetilde{u}_{l} \right) = \widetilde{f}_{n}. \end{cases}$$

Let  $(u_{aa,l,\varepsilon})_{\varepsilon}, (u_{cor,l,\varepsilon})_{\varepsilon}, (A_{ij,\varepsilon}^{rl})_{\varepsilon}, (K_{\varepsilon}^{rl})_{\varepsilon}, (f_{aa,l,\varepsilon})_{\varepsilon}$  and  $(f_{cor,l,\varepsilon})_{\varepsilon}$  be respective representatives of  $\widetilde{u}_{aa,l}, \ \widetilde{u}_{cor,l}, \widetilde{A}_{ij}^{rl}, \widetilde{K}^{rl}, \widetilde{f}_{aa,l}$  and  $\widetilde{f}_{cor,l}, 1 \leq l \leq n$ . For every  $r = 1, \ldots, n$  and  $\varepsilon \in I$ , denoting

$$S_{aa,r,\varepsilon} := \sum_{l=1}^{n} \left( \sum_{i=1}^{p} \sum_{j=1}^{q} A_{ij,\varepsilon}^{rl} \left( \tau_{\omega_j} u_{aa,l,\varepsilon} \right)^{(i)} + K_{\varepsilon}^{rl} * u_{aa,l,\varepsilon} \right),$$
  
$$S_{cor,r,\varepsilon} := \sum_{l=1}^{n} \left( \sum_{i=1}^{p} \sum_{j=1}^{q} A_{ij,\varepsilon}^{rl} \left( \tau_{\omega_j} u_{cor,l,\varepsilon} \right)^{(i)} + K_{\varepsilon}^{rl} * u_{cor,l,\varepsilon} \right),$$

we have  $\forall \varepsilon \in I, S_{aa,r,\varepsilon} \in \mathcal{B}_{aa}$  and  $S_{cor,r,\varepsilon} \in \mathcal{B}_{+,0}$ , and also  $[(S_{aa,r,\varepsilon})_{\varepsilon}] \in \mathcal{G}_{aa}$  and  $[(S_{cor,r,\varepsilon})_{\varepsilon}] \in \mathcal{G}_{+,0}$ . Since  $\widetilde{u}$  is a generalized solution of (7.1) on  $\mathbb{J}$ , it means that  $\forall k \in \mathbb{Z}_+, \forall m > 0, \exists c_k > 0, \exists c_k \in I, \forall \varepsilon < \varepsilon_k, \forall x \in \mathbb{J}, \forall \varepsilon < \varepsilon_k, \forall z \in \mathbb{J}, \forall z \in \mathbb{J$ 

(7.4) 
$$\sum_{j \le k} \left| S_{aa,r,\varepsilon}^{(j)}\left(x\right) - f_{aa,r,\varepsilon}^{(j)}\left(x\right) + S_{cor,r,\varepsilon}^{(j)}\left(x\right) - f_{cor,r,\varepsilon}^{(j)}\left(x\right) \right| \le c_k \varepsilon^m, r = 1, \dots, n.$$

For any real sequence  $(s_m)_{m\in\mathbb{N}}$ , such that  $s_m \to +\infty$  there exist  $(s_{m_{p(\varepsilon)}})_p$  a subsequence of  $(s_m)_{m\in\mathbb{N}}$ , such that taking the translate at  $s_{m_{p(\varepsilon)}} - s_{m_{q(\varepsilon)}}$  in the estimate (7.4) and let  $p, q \to +\infty$ , we obtain due to ([12], Proposition 9) that  $\forall k \in \mathbb{Z}_+, \forall m > 0, \exists c_k > 0, \exists \varepsilon_k \in I, \forall \varepsilon < \varepsilon_k, \forall x \in \mathbb{J}$ ,

$$\sum_{j \le k} \left| S_{aa,r,\varepsilon}^{(j)}\left(x\right) - f_{aa,r,\varepsilon}^{(j)}\left(x\right) \right| \le c_k \varepsilon^m, \forall r = 1, \dots, n,$$

consequently, by Proposition 3-(5) of [12], it holds  $\forall k \in \mathbb{Z}_+, \forall m > 0, \exists c_k > 0, \exists \varepsilon_k \in I, \forall \varepsilon < \varepsilon_k, \forall \varepsilon$ 

$$S_{aa,r,\varepsilon} - f_{aa,r,\varepsilon} |_{k,\infty,\mathbb{R}} \le c_k \varepsilon^m, \forall r = 1, \dots, n_k$$

i.e.

(7.5) 
$$(S_{aa,r,\varepsilon} - f_{aa,r,\varepsilon})_{\varepsilon} \in \mathcal{N}_{\mathcal{B}}(\mathbb{R}), \forall r = 1, \dots, n,$$

which means that  $\widetilde{u}_{aa} = (\widetilde{u}_{aa,1}, \ldots, \widetilde{u}_{aa,n})$  is a generalized solution of (7.2) on  $\mathbb{R}$ . By (7.4) and (7.5), we deduce

$$S_{cor,r,\varepsilon} - f_{cor,r,\varepsilon} \rangle_{\varepsilon} \in \mathcal{N}_{\mathcal{B}}\left(\mathbb{J}\right), \forall r = 1, \dots, n,$$

i.e.  $\widetilde{u}_{cor} = (\widetilde{u}_{cor,1}, \dots, \widetilde{u}_{cor,n})$  is a generalized solution of (7.3) on  $\mathbb{J}$ .

Conversely if there exist  $\widetilde{v} \in (\mathcal{G}_{aa})^n$  and  $\widetilde{w} \in (\mathcal{G}_{+,0})^n$  such that (7.2) and (7.3) hold, then we have  $\widetilde{u} := (\widetilde{v} + \widetilde{w}) \in (\mathcal{G}_{aaa})^n$  is a generalized solution of (7.1) on  $\mathbb{J}$ .

**Remark 2.** Theorem 5 generalizes Theorem 6 of [9] and Theorem 3 of [12].

As a particular case we consider linear systems of ordinary differential equations

(7.6) 
$$\widetilde{u} + \widetilde{A}\widetilde{u} = \widetilde{f}$$

where  $\widetilde{A}$  is a square matrix of almost automorphic generalized functions.

**Corollary 1.** Let  $\tilde{f} = (\tilde{f}_{aa} + \tilde{f}_{cor}) \in (\mathcal{G}_{aaa})^n$ , the system (7.6) admits a generalized solution  $\tilde{u} \in (\mathcal{G}_{aaa})^n$  on  $\mathbb{J}$  if and only if there exist  $\tilde{v} \in (\mathcal{G}_{aa})^n$  and  $\tilde{w} \in (\mathcal{G}_{+,0})^n$  such that

$$\widetilde{v} + \widetilde{A}\widetilde{v} = \widetilde{f}_{aa} \ on \ \mathbb{R}$$

and

 $\overset{\cdot}{\widetilde{w}} + \widetilde{A}\widetilde{w} = \widetilde{f}_{cor} \ on \ \mathbb{J}.$ 

Let  $\widetilde{u} = [(u_{\varepsilon})_{\varepsilon}] \in \mathcal{G}_{\mathcal{B}}, x_0 \in \mathbb{R}$  and define the primitive of  $\widetilde{u}$  by  $\widetilde{U} = [(U_{\varepsilon})_{\varepsilon}]$ , where

$$U_{\varepsilon}(x) := \int_{x_0}^x u_{\varepsilon}(t) dt, \varepsilon \in I.$$

**Corollary 2.** A generalized function  $\widetilde{U} = (\widetilde{U}_{aa} + \widetilde{U}_{cor}) \in \mathcal{G}_{\mathcal{B}}$  is a primitive of  $\widetilde{u} = (\widetilde{u}_{aa} + \widetilde{u}_{cor}) \in \mathcal{G}_{aaa}$  on  $\mathbb{J}$  if and only if

$$U_{aa}$$
 is a primitive of  $\widetilde{u}_{aa}$  on  $\mathbb{R}$ ,

and

$$U_{cor}$$
 is a primitive of  $\widetilde{u}_{cor}$  on  $\mathbb{J}$ .

### References

- Antonevich A. B., Radyno Ya. V., On a general method of constructing algebras of new generalized functions. Soviet. Math. Dokl., vol. 43:3, p. 680-684, (1991).
- Bochner S., Uniform convergence of monotone sequences of functions. Proc. Natl. Acad. Sci. USA, Vol. 47, p. 582-585, (1961).
- [3] Bochner S., A new approach to almost periodicity. Proc. Natl. Acad. Sci. USA, Vol. 48, No. 12, p. 2039-2043, (1962).
- Bochner S., Continuous mappings of almost automorphic and almost periodic functions. Proc. Natl. Acad. Sci. USA, Vol. 52, p. 907-910, (1964).
- [5] Bohr H., Almost periodic functions. Chelsea Publishing Company, (1947).
- [6] Bouzar C., Khalladi M.T., Almost periodic generalized functions. Novi Sad J. Math, Vol. 41, N<sup>0</sup>1, p. 33-42, (2011).
- [7] Bouzar C., Khalladi M.T., Linear differential equations in the algebra of almost periodic generalized functions. Rend. Sem. Univ. Politec. Torino, Vol. 70: 2, p. 111-120, (2012).
- [8] Bouzar C., Khalladi M. T., Asymptotically almost periodic generalized functions. Operator Theory : Advances and Applications, Vol. 231, p. 261-272, (2013).
- Bouzar C., Khalladi M. T., On asymptotically almost periodic generalized solutions of differential equations. Operator Theory : Advances and Applications, Vol. 231, p. 35-43, (2015).

- [10] Bouzar C., Khalladi M. T., Tchouar F. Z., Almost automorphic generalized functions. Novi Sad J. Math., Vol. 45, No. 1, p. 207-214, (2015).
- [11] Bouzar C., Tchouar F. Z., Almost automorphic distributions. Mediterr. J. Math, Vol. 14, No. 04, p. 1-13, (2017).
- [12] Bouzar C., Tchouar F. Z., Asymptotic almost automorphy of functions and distributions. Ural Mathematical Journal, Vol. 6, No. 1, p. 54-70, (2020).
- [13] Cioranescu I. Asymptotically almost periodic distributions. Appl. Anal., Vol. 34, No. 3–4. P. 251–259, (1990).
- [14] Colombeau J.-F., Elementary introduction to new generalized functions. North Holland. (1985).
- [15] Egorov Yu. V., A contribution to the theory of generalized functions, Russ. Math. Surveys, 45-5, p. 1-49, (1990).
- [16] Fréchet M., Les fonctions asymptotiquement presque périodiques. Revue Sci., 79, p. 341-354, (1941).
- [17] Garetto C., Topological structures in Colombeau algebras. Monatsh. Math., 146(3), p. 203–226, (2005).
- [18] Levitan B. M., Almost Periodic Functions. Gos. Izdat. Tekh-Teor. Lit. Moscow (in Russian), (1953).
- [19] N'Guérékata G. M., Some remarks on asymptotically almost automorphic functions. Riv. di Mat. della Universita di Parma (4) 13, p. 301-303, (1987).
- [20] M. Nedeljkov, S. Pilipović, D. Scarpalézos. The linear theory of Colombeau generalized functions, Longman, (1998).
- [21] Oberguggenberger M., Multiplication of distributions and applications to partial differential equations. Longman, (1992).
- [22] Schwartz L., Théorie des distributions. Hermann, 2<sup>ième</sup> Edition, (1966).
- [23] Schwartz L., Sur l'impossibilité de la multiplication des distributions. C. R. Acad. Sci., 239, 847-848, (1954).
- [24] Seeley R. T., Extension of  $C^{\infty}$  functions defined in a half space. Proc. Amer. Math. Soc., Vol. 15, No. 4, p. 625-626, (1964).
- [25] Sobolev S. L., Applications of functional analysis in mathematical physics. American Mathematical Society, (1963).
- [26] Stepanoff V. V., Sur quelques généralisations des fonctions presque périodiques. C. R. Acad. Sci. Paris, Vol. 181, p. 90-92, (1925).

LABORATORY OF MATHEMATICAL ANALYSIS AND APPLICATIONS. UNIVERSITÉ ORAN 1, AHMED BEN BELLA, 31000, ORAN, ALGERIA.

Email address: ch.bouzar@gmail.com

LABORATORY OF MATHEMATICAL ANALYSIS AND APPLICATIONS. UNIVERSITÉ ORAN 1, AHMED BEN BELLA, 31000, ORAN, ALGERIA.

Email address: meryemslimani@yahoo.com