

Singular Lagrangians, Constrained Hamiltonian Systems and Gauge Invariance: An Example of the Dirac–Bergmann Algorithm

J. David Brown

*Department of Physics, North Carolina State University,
2401 Stinson Drive, Raleigh, NC 27695*

E-mail: david_brown@ncsu.edu

ABSTRACT: The Dirac–Bergmann algorithm is a recipe for converting a theory with a singular Lagrangian into a constrained Hamiltonian system. Constrained Hamiltonian systems include gauge theories—general relativity, electromagnetism, Yang Mills, string theory, etc. The Dirac–Bergmann algorithm is elegant but at the same time rather complicated. It consists of a large number of logical steps linked together by a subtle chain of reasoning. Examples of the Dirac–Bergmann algorithm found in the literature are designed to isolate and illustrate just one or two of those logical steps. In this paper I analyze a finite–dimensional system that exhibits all of the major steps in the algorithm. The system includes primary and secondary constraints, first and second class constraints, restrictions on Lagrange multipliers, and both physical and gauge degrees of freedom. This relatively simple system provides a platform for discussing the Dirac conjecture, constructing Dirac brackets, and applying gauge conditions.

ARXIV EPRINT: [2201.06558](https://arxiv.org/abs/2201.06558)

Contents

1	Introduction	2
2	Lagrangian Analysis	4
3	Gauge Invariance	5
4	Primary Constraints and the Canonical Hamiltonian	7
5	Primary Hamiltonian and the Primary Action	8
6	Hamilton's Equations and the Initial Value Problem	9
7	Consistency Conditions, Secondary Constraints and Restrictions on the Lagrange Multipliers	10
8	Weak Equality and Lagrange Multiplier Analysis	12
9	Total Hamiltonian	13
10	First and Second Class Constraints	14
11	First Class Hamiltonian, Gauge Generators and the Dirac Conjecture	15
12	Extended Hamiltonian	17
13	Extended Action	18
14	Dirac Brackets and the Partially Reduced Hamiltonian	20
15	Partially Reduced Action	22
16	Gauge Conditions and the Fully Reduced Hamiltonian	24
17	Fully Reduced Action	26
18	Summary and Comments	27

1 Introduction

In the early 1950's Dirac and Bergmann independently developed the Hamiltonian formalism for systems with singular Lagrangians [1–10]. These systems, often called “constrained Hamiltonian systems”, include gauge theories. Gauge freedom is more clearly and more completely displayed in the Hamiltonian setting, with the generators of gauge transformations expressed as functions on phase space. Historically, the main motivation for casting gauge theories in Hamiltonian form was to facilitate their canonical quantization. Dirac and Bergmann were primarily motivated by the prospect of developing a quantum theory of gravity based on a Hamiltonian formulation of general relativity.

Textbook treatments of Lagrangian and Hamiltonian mechanics invariably assume that the Lagrangian $L(q, \dot{q})$ is nonsingular; that is, that the matrix of second derivatives of $L(q, \dot{q})$ with respect to the velocities is invertible. In classical mechanics, the nonsingular case appears to be sufficient to cover problems of physical interest. However, one might argue that textbooks avoid certain physically interesting problems simply because their Lagrangians are singular.

In field theory, the issue of singular Lagrangians cannot be avoided. Nearly every field theory of physical interest—electrodynamics, Yang–Mills theory, general relativity, relativistic string theory—has gauge freedom. The Lagrangians for these theories are singular.

The Dirac–Bergmann algorithm transforms a singular Lagrangian system into a Hamiltonian system. The formalism is elegant but at the same time rather complex. It consists of a large number of logical steps, linked together by a chain of reasoning that can be difficult to keep straight. Of course there are many examples in the literature in which the Dirac–Bergmann algorithm is applied, converting a singular Lagrangian into Hamiltonian form. But to my knowledge, all of these examples are designed to illustrate just one or two of the logical steps in the algorithm. The student of the subject is faced with the task of linking these examples together to create a complete picture of the algorithm.

For those who learn by example, what is needed is a single example that illustrates all of the major logical steps in the Dirac–Bergmann algorithm and shows how these steps are linked together. Such a “complete” example is not easy to identify because there is no obvious way to predict, starting with a particular Lagrangian, which of the steps in the algorithm will be needed.

The system analyzed in this paper is defined by the Lagrangian

$$L(q, \dot{q}) = \frac{1}{2} \left\{ (q_1 + \dot{q}_2 + \dot{q}_3)^2 + (\dot{q}_4 - \dot{q}_2)^2 + (q_1 + 2q_2)(q_1 + 2q_4) \right\} \quad (1.1)$$

where the dot denotes a time derivative. The matrix of second derivatives with respect to the velocities is

$$\frac{\partial^2 L}{\partial \dot{q}_i \partial \dot{q}_j} = \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & 2 & 1 & -1 \\ 0 & 1 & 1 & 0 \\ 0 & -1 & 0 & 1 \end{pmatrix}. \quad (1.2)$$

This matrix is singular; it has rank 2.

As we will see, the system defined by the Lagrangian (1.1) is relatively complete.¹ It contains both primary and secondary constraints, both first and second class constraints, and restrictions on the Lagrange multipliers. The first class constraints for this system are not all primary; this allows us to address the Dirac conjecture. The second class constraints can be eliminated by introducing Dirac brackets. Finally, this system contains both physical and gauge degrees of freedom. The gauge freedom can be eliminated with suitable gauge conditions.

One characteristic of any complete example such as (1.1) is that the configuration space, the space of q 's, must be at least four-dimensional. Here is why: The number of physical degrees of freedom is equal to the dimension of the configuration space, minus the number of first class constraints, minus half the number of second class constraints. If the example is to have at least one physical degree of freedom, at least two first class constraints (one primary and one secondary), and at least two second class constraints (the number of second class constraints must be even), then the configuration space must be at least four-dimensional.

The study of constrained Hamiltonian systems predates Dirac and Bergmann with earlier work by Rosenfeld [12, 13]. Like Rosenfeld, Bergmann and his collaborators [1, 2, 4, 6, 7, 14] were focused on field theories like general relativity that are covariant with respect to general four-dimensional coordinate transformations. Dirac took a more basic approach to the problem by considering a generic singular Lagrangian [3, 5, 8–10]. He developed the algorithm for the case of systems with a finite number of degrees of freedom. His view was that the generalization to field theory, with an infinite number of degrees of freedom, would be “merely a formal matter”.

In this paper I apply the Dirac–Bergmann algorithm to the Lagrangian (1.1), following closely the general treatment given by Henneaux and Teitelboim [11]. In turn, the account of Henneaux and Teitelboim closely follows Dirac’s 1964 *Lectures on Quantum Mechanics* [10]. Presentations of the Dirac–Bergmann algorithm can also be found in books by Hanson, Regge and Teitelboim [15], Sundermeyer [16], Rothe and Rothe [17], and Lusanna [18].

Throughout the paper I attempt to explain the reasoning behind the logical steps of the Dirac–Bergmann algorithm, but avoid general proofs. The reader is referred to references [11, 15–18] for more details.

We begin in Sec. 2 with a derivation of Lagrange’s equations for the singular Lagrangian (1.1). The general solution is derived, and in Sec. 3 we discuss the gauge freedom at the Lagrangian level. We begin construction of the Hamiltonian theory in Sec. 4 with a derivation of the primary constraints and canonical Hamiltonian. In Sec. 5 we introduce the primary Hamiltonian and the primary action. Section 6 is devoted to a discussion of the initial value problem and the need to go beyond the primary Hamiltonian. In Sec. 7 we apply Dirac’s consistency conditions to derive the secondary constraints and restrictions on the Lagrange multipliers. The concept of weak equality is introduced in Sec. 8, along with a formal analysis of the restrictions on the Lagrange multipliers. The total Hamiltonian

¹This example does not cover every contingency. In particular, it does not include redundant constraints [11].

is computed in Sec. 9, and in Sec. 10 we sort the constraints into first and second class. The first class Hamiltonian and gauge generators are identified in Sec. 11, where we also introduce the Dirac conjecture. In Secs. 12 and 13 we define the extended Hamiltonian and extended action. Dirac brackets are used in Sec. 14 to eliminate the second class constraints, which yields a partially reduced Hamiltonian. The corresponding partially reduced action is derived in Sec. 15. Gauge conditions are introduced in Sec. 16 and Dirac brackets are used to eliminate the constraints and gauge conditions. This yields a fully reduced Hamiltonian. The fully reduced action is derived in Sec. 17. Finally, Sec. 18 contains a short summary and some comments on the Dirac–Bergmann algorithm.

2 Lagrangian Analysis

The action is the integral of the Lagrangian (1.1):

$$S[q] = \int_0^T dt L(q, \dot{q}) . \quad (2.1)$$

The notation $S[q]$ indicates that S is a functional of the complete set of coordinates, $q_i = \{q_1, q_2, q_3, q_4\}$. The equations of motion are obtained by extremizing the action. For this example, we are not concerned with boundary conditions and integrate by parts freely. Lagrange’s equations are

$$0 = \frac{\delta S}{\delta q_1} = \dot{q}_2 + \dot{q}_3 + 2q_1 + q_2 + q_4 , \quad (2.2a)$$

$$0 = \frac{\delta S}{\delta q_2} = -2\ddot{q}_2 - \ddot{q}_3 + \ddot{q}_4 - \dot{q}_1 + q_1 + 2q_4 , \quad (2.2b)$$

$$0 = \frac{\delta S}{\delta q_3} = -\ddot{q}_2 - \ddot{q}_3 - \dot{q}_1 , \quad (2.2c)$$

$$0 = \frac{\delta S}{\delta q_4} = \ddot{q}_2 - \ddot{q}_4 + q_1 + 2q_2 . \quad (2.2d)$$

We can rewrite these as follows. First, add Eqs. (2.2 b) and (2.2 d), then subtract Eq. (2.2 c). This gives

$$q_1 + q_2 + q_4 = 0 . \quad (2.3a)$$

Next, subtract this result from Eq. (2.2 a) to obtain

$$\dot{q}_2 + \dot{q}_3 + q_1 = 0 . \quad (2.3b)$$

The time derivative of this equation yields Eq. (2.2 c). Finally, we find the result

$$\ddot{q}_4 - \ddot{q}_2 = q_2 - q_4 . \quad (2.3c)$$

by solving Eq. (2.3 a) for q_1 and using the equation of motion (2.2 d).

Equations (2.3) are equivalent to Lagrange’s equations (2.2). In particular, Eq. (2.2 a) is the sum of Eqs. (2.3 a) and (2.3 b); Eq. (2.2 b) is the sum of Eqs (2.3 a) and (2.3 c) with the time derivative of (2.3 b) subtracted; Eq. (2.2 c) is the negative of the time derivative of (2.3 b); Eq. (2.2 d) is obtained by subtracting (2.3 c) from (2.3 a).

The equations of motion for this simple linear system are easily solved. Note that the combination $q_4 - q_2$ is determined by (2.3 c) along with initial or boundary data; thus, we have

$$q_4(t) - q_2(t) = A \sin t + B \cos t , \quad (2.4)$$

where A and B are constants. Now Eq. (2.3 a) gives

$$q_1(t) + 2q_2(t) = -A \sin t - B \cos t . \quad (2.5)$$

If we knew $q_2(t)$, we could solve the previous two equations for $q_1(t)$ and $q_4(t)$, then integrate Eq. (2.3 b) to obtain $q_3(t)$. Clearly we do not have enough information to fully determine each of the q 's as functions of time. One of the q 's must remain undetermined. For example, let us choose q_2 arbitrarily by setting $q_2(t) = -\Psi(t)/2$ for some function $\Psi(t)$. We can then use the equations above to solve for q_1 , q_3 and q_4 :

$$q_1(t) = -A \sin t - B \cos t + \Psi(t) , \quad (2.6a)$$

$$q_2(t) = -\Psi(t)/2 , \quad (2.6b)$$

$$q_3(t) = -A \cos t + B \sin t + \Psi(t)/2 - \int_0^t ds \Psi(s) + C , \quad (2.6c)$$

$$q_4(t) = A \sin t + B \cos t - \Psi(t)/2 , \quad (2.6d)$$

where C is an integration constant. This is the general solution of the equations of motion.

3 Gauge Invariance

The undetermined function $\Psi(t)$ that appears in the general solution (2.6) can be freely specified. This is the gauge freedom of the theory. We can express the gauge freedom in another way: the Lagrangian (1.1) and the equations of motion (either (2.2) or (2.3)) are invariant under the replacements

$$q_1(t) \rightarrow q_1(t) + \Psi(t) , \quad (3.1a)$$

$$q_2(t) \rightarrow q_2(t) - \Psi(t)/2 , \quad (3.1b)$$

$$q_3(t) \rightarrow q_3(t) + \Psi(t)/2 - \int_0^t ds \Psi(s) , \quad (3.1c)$$

$$q_4(t) \rightarrow q_4(t) - \Psi(t)/2 , \quad (3.1d)$$

where $\Psi(t)$ is an arbitrary function of time.

Although each configuration of the system (that is, each set of q values) corresponds to a specific physical state of the system, the converse is not true. Because of the gauge freedom, there are many sets of q 's that describe one and the same physical state.

Let us examine the gauge freedom more closely, in anticipation of the Hamiltonian description of evolution. To begin, choose the gauge $\Psi(t) = 0$ and consider the general

solution (2.6). This solution describes the evolution of the system from initial data

$$q_1(0) = -B, \quad \dot{q}_1(0) = -A, \quad (3.2a)$$

$$q_2(0) = 0, \quad \dot{q}_2(0) = 0, \quad (3.2b)$$

$$q_3(0) = C - A, \quad \dot{q}_3(0) = B, \quad (3.2c)$$

$$q_4(0) = B, \quad \dot{q}_4(0) = A. \quad (3.2d)$$

The configuration at some arbitrary final time $t = T$ is

$$q_1(T) = -A \sin T - B \cos T, \quad (3.3a)$$

$$q_2(T) = 0, \quad (3.3b)$$

$$q_3(T) = C - A \cos T + B \sin T, \quad (3.3c)$$

$$q_4(T) = A \sin T + B \cos T. \quad (3.3d)$$

This configuration corresponds to a particular state of the physical system.

We can choose a different gauge in Eqs. (2.6). As long as the new gauge satisfies $\Psi(0) = \dot{\Psi}(0) = 0$, the solution will describe evolution from the same initial data (3.2). For example, with

$$\Psi(t) = \frac{(\pi^2 - 4)\epsilon}{8} [\cos(\pi t/T) - 1] + \frac{\pi\epsilon}{4} [\pi t/T - \sin(\pi t/T)] \quad (3.4)$$

where $\epsilon = \text{const}$, the configuration at $t = T$ is

$$q_1(T) = -A \sin T - B \cos T + \epsilon, \quad (3.5a)$$

$$q_2(T) = -\epsilon/2, \quad (3.5b)$$

$$q_3(T) = C - A \cos T + B \sin T + \epsilon/2, \quad (3.5c)$$

$$q_4(T) = A \sin T + B \cos T - \epsilon/2. \quad (3.5d)$$

The configurations (3.3) and (3.5) represent the same physical state of the system, since they evolve from the same initial data.

We can express this result more compactly as

$$\delta q_1 = \epsilon, \quad (3.6a)$$

$$\delta q_2 = -\epsilon/2, \quad (3.6b)$$

$$\delta q_3 = \epsilon/2, \quad (3.6c)$$

$$\delta q_4 = -\epsilon/2. \quad (3.6d)$$

Here, δq_i denotes the change in q_i at the generic time T , due to the change in gauge function $\Psi(t)$.

Here is another example. With

$$\Psi(t) = \frac{\pi^2\epsilon}{4T} [\cos(\pi t/T) - 1] + \frac{\pi\epsilon}{2T} [\pi t/T - \sin(\pi t/T)] \quad (3.7)$$

we obtain a configuration that differs from Eqs. (3.3) by

$$\delta q_1 = 0 , \tag{3.8a}$$

$$\delta q_2 = 0 , \tag{3.8b}$$

$$\delta q_3 = \epsilon , \tag{3.8c}$$

$$\delta q_4 = 0 . \tag{3.8d}$$

This configuration is also evolved from the initial data (3.2), and represents the same physical state as the configurations (3.3) and (3.5).

Although the gauge transformation (3.1) contains a single arbitrary function of time, the gauge invariance naturally splits into two types. The first consists of variations subject to $\delta q_2 = -\delta q_3 = \delta q_4 = -\delta q_1/2$. The second consists of arbitrary variations in δq_3 , with $\delta q_1 = \delta q_2 = \delta q_4 = 0$. This apparent ‘‘doubling’’ of the gauge freedom arises because the solution (3.1c) for $q_3(t)$ (unlike the other variables) includes the integral of $\Psi(t)$. There is enough freedom of choice in $\Psi(t)$ to allow variations in q_3 that are independent of the variations among the other variables. Both types of gauge transformations leave the physical state of the system unchanged.

The consequences of gauge invariance are most clearly expressed in the Hamiltonian formalism. The extended Hamiltonian defined in Sec. 12 includes phase space generators for both types of gauge transformations.

4 Primary Constraints and the Canonical Hamiltonian

We now begin construction of the Hamiltonian description of the system. The conjugate momenta are defined as usual by $p_i = \partial L/\partial \dot{q}_i$. For the Lagrangian (1.1), we have

$$p_1 = 0 \tag{4.1a}$$

$$p_2 = 2\dot{q}_2 + \dot{q}_3 - \dot{q}_4 + q_1 \tag{4.1b}$$

$$p_3 = \dot{q}_2 + \dot{q}_3 + q_1 \tag{4.1c}$$

$$p_4 = \dot{q}_4 - \dot{q}_2 \tag{4.1d}$$

Because the Lagrangian is singular, the matrix of second derivatives $\partial^2 L/\partial \dot{q}_i \partial \dot{q}_j$ is not invertible and we cannot solve Eqs. (4.1) for the velocities as functions of the coordinates and momenta. The definitions (4.1) yield two *primary constraints*,

$$\phi_1 \equiv p_1 = 0 , \tag{4.2a}$$

$$\phi_2 \equiv p_2 - p_3 + p_4 = 0 , \tag{4.2b}$$

that restrict the phase space variables p_i, q_i . We will denote these constraints collectively by ϕ_a , where $a = 1, 2$. Note that in this simple example, the primary constraints are independent of the q 's.

There is some freedom in choosing how the constraints are written. For example, we could replace ϕ_1 above with $\phi_1 \equiv p_1 - p_2 + p_3 - p_4 = 0$. However, we are not allowed to

choose $\phi_1 \equiv p_1^2 = 0$. This is because the constraints must satisfy a *regularity condition*: the Jacobian matrix formed from the derivatives of the constraints with respect to the p 's and q 's must have maximal rank on the constraint subspace [10, 11]. Roughly speaking, the constraints should have nonzero gradients.

The next step in constructing the Hamiltonian formalism is to compute the *canonical Hamiltonian*. The canonical Hamiltonian H_C is defined from the usual prescription by writing $p_i \dot{q}_i - L(q, \dot{q})$ in terms of p 's and q 's. Although we cannot solve for all of the \dot{q} 's in terms of p 's and q 's, it can be shown that the combination $p_i \dot{q}_i - L(q, \dot{q})$ depends only on the phase space variables [10, 11]. For our example problem the canonical Hamiltonian is

$$H_C = \frac{1}{2} \left[p_3^2 + p_4^2 - 2p_3 q_1 - (q_1 + 2q_2)(q_1 + 2q_4) \right]. \quad (4.3)$$

Note that H_C is ambiguous. For example, we could use the primary constraint (4.2 b) to replace the term $-2p_3 q_1$ with $-2(p_2 + p_4)q_1$.

5 Primary Hamiltonian and the Primary Action

The *primary Hamiltonian* H_P is obtained from the canonical Hamiltonian H_C by adding the primary constraints with Lagrange multipliers,

$$H_P = H_C + \lambda_a \phi_a, \quad (5.1)$$

where a sum over the repeated index a is implied. The *primary action* is built from the primary Hamiltonian in the usual way: $S_P[q, p, \lambda] = \int_0^T dt \{ p_i \dot{q}_i - H_P \}$. Explicitly, we have

$$S_P[q, p, \lambda] = \int_0^T dt \left\{ p_i \dot{q}_i - \frac{1}{2} \left[p_3^2 + p_4^2 - 2p_3 q_1 - (q_1 + 2q_2)(q_1 + 2q_4) \right] - \lambda_1 p_1 - \lambda_2 (p_2 - p_3 + p_4) \right\}. \quad (5.2)$$

The primary action is a functional of the complete set of phase space coordinates, q_i , p_i , as well as the Lagrange multipliers λ_1 and λ_2 .

The equations of motion are obtained by extremizing the primary action S_P . Extremization with respect to the momenta p_i gives

$$\dot{q}_1 = \lambda_1, \quad (5.3a)$$

$$\dot{q}_2 = \lambda_2, \quad (5.3b)$$

$$\dot{q}_3 = p_3 - q_1 - \lambda_2, \quad (5.3c)$$

$$\dot{q}_4 = p_4 + \lambda_2, \quad (5.3d)$$

while extremization with respect to the coordinates q_i yields

$$\dot{p}_1 = p_3 + q_1 + q_2 + q_4, \quad (5.3e)$$

$$\dot{p}_2 = q_1 + 2q_4, \quad (5.3f)$$

$$\dot{p}_3 = 0, \quad (5.3g)$$

$$\dot{p}_4 = q_1 + 2q_2. \quad (5.3h)$$

Extremizing the action S_P with respect to the Lagrange multipliers gives the constraints,

$$\phi_1 \equiv p_1 = 0 \quad , \quad (5.3i)$$

$$\phi_2 \equiv p_2 - p_3 + p_4 = 0 \quad . \quad (5.3j)$$

These equations of motion (5.3) are equivalent to Lagrange's equations (2.2). To show this, we first solve Eqs. (5.3 c,d,i,j) for the momenta to obtain

$$p_1 = 0 \quad , \quad (5.4a)$$

$$p_2 = \dot{q}_3 - \dot{q}_4 + q_1 + 2\lambda_2 \quad , \quad (5.4b)$$

$$p_3 = \dot{q}_3 + q_1 + \lambda_2 \quad , \quad (5.4c)$$

$$p_4 = \dot{q}_4 - \lambda_2 \quad . \quad (5.4d)$$

Using these results along with Eqs. (5.3 a,b) for the Lagrange multipliers, we find that Eqs. (5.3 e,f,g,h) agree precisely with Lagrange's equations (2.2).

6 Hamilton's Equations and the Initial Value Problem

At this point one might ask whether the task of expressing the singular system (1.1) in Hamiltonian form is complete. After all, the primary action (5.2) provides the correct equations of motion for the phase space variables q_i and p_i . In fact, we can obtain the time evolution of any phase space function F from $\dot{F} = [F, H_P]$ where H_P is the primary Hamiltonian and $[\cdot, \cdot]$ denotes Poisson brackets. Hamilton's equations for the coordinates and momenta, $\dot{q}_i = [q_i, H_P]$ and $\dot{p}_i = [p_i, H_P]$, coincide with Eqs. (5.3 a-h).

Our task of expressing the singular system in Hamiltonian form is not yet complete because we still need to interpret Hamilton's equations (5.3 a-h) as an initial value problem. That is, Hamilton's equations should determine the future history of the system solely from initial data. In contrast, the primary action (5.2) defines a boundary value problem in which the configuration variables, the q 's, are specified at initial and final times.

The key difference between the equations of motion $\delta S_P = 0$ and Hamilton's equations $\dot{F} = [F, H_P]$ is that the former include the primary constraints, Eqs. (5.3 i,j), whereas the later do not. Thus, the phase space trajectories that extremize the action S_P must lie entirely in the primary constraint surface. (The primary constraint "surface" is the subspace of phase space that satisfies the primary constraints.) In contrast, the trajectories obtained from Hamiltonian evolution $\dot{F} = [F, H_P]$ are defined throughout the entire phase space. Note that we cannot simply append the primary constraint equations to Hamilton's equations, because in that case the complete system would not be in Hamiltonian form.

Of course the physically allowed phase space trajectories must satisfy the primary constraints. With an initial value interpretation of Hamilton's equations, we can try to enforce the primary constraints with appropriate choices of initial data and Lagrange multipliers. In particular, we can choose initial data that lie on the primary constraint surface $\phi_1(0) \equiv p_1(0) = 0$ and $\phi_2(0) \equiv p_2(0) - p_3(0) + p_4(0) = 0$. But this is not enough, because the primary constraints are not necessarily satisfied at later times as the system evolves into the future.

We can describe the situation as follows. The trajectories that extremize the action S_P , the *physical trajectories*, do not necessarily fill the entire primary constraint surface. Instead they might span only a subspace of the primary constraint surface. If the initial data lie in the primary constraint surface but *outside* the subspace of physical trajectories, then the primary constraints will not be preserved as the data is evolved.

How should the initial data and Lagrange multipliers be restricted such that the primary constraints hold throughout the evolution? The primary constraints will hold for all time if they hold initially and their time derivatives (to all orders) also vanish initially. In the general case this leads to a hierarchy of restrictions on the initial data in the form of secondary, tertiary, etc. constraints.² It can also lead to restrictions on the Lagrange multipliers.

The higher order (secondary, tertiary, etc.) constraints and restrictions on the Lagrange multipliers are not new—imposing them does not change the content or predictions of the physical theory. This is because the higher order constraints and restrictions on the Lagrange multipliers are direct consequences of the equations of motion (5.3) that follow from the primary action (5.2). They are simply “hidden” in those equations. The process of identifying the higher order constraints and restrictions on Lagrange multipliers reveals these hidden conditions.

7 Consistency Conditions, Secondary Constraints and Restrictions on the Lagrange Multipliers

We can ensure that the primary constraints hold for all time by applying Dirac’s *consistency conditions* [10]. Begin by computing the time derivatives of the primary constraints with the primary Hamiltonian, $\dot{\phi}_a = [\phi_a, H_P]$. Now set these equal to zero:

$$[\phi_a, H_P] = 0 . \tag{7.1}$$

For each value of the index a , there are three possibilities.³ First, $[\phi_a, H_P]$ might vanish on the constraint surface $\phi_a = 0$, so that the consistency condition (7.1) reduces to the identity $0 = 0$. Second, $[\phi_a, H_P]$ could be a (non-constant) phase space function that is independent of the Lagrange multipliers. In this case Eq. (7.1) is a *secondary constraint*. Finally, $[\phi_a, H_P]$ might depend on the Lagrange multipliers. Then Eq. (7.1) fixes one of the Lagrange multipliers in terms of the phase space variables and the other Lagrange multipliers.

The secondary constraints that arise from this process must themselves satisfy the consistency conditions. This can lead to *tertiary constraints* and more restrictions on the Lagrange multipliers. In turn the tertiary constraints can lead to quaternary constraints, and so forth. We must continue to apply the consistency condition until the process naturally stops.

²Some authors use the term “secondary constraints” to refer to all higher-order constraints—that is, all constraints beyond the primary level.

³This assumes Lagrange’s equations are self-consistent. Otherwise, the consistency conditions could lead to a contradiction such as $1 = 0$.

For our example, the primary constraints are

$$\phi_1 \equiv p_1 = 0 , \quad (7.2a)$$

$$\phi_2 \equiv p_2 - p_3 + p_4 = 0 , \quad (7.2b)$$

and their time derivatives are

$$\dot{\phi}_1 = [\phi_1, H_P] = p_3 + q_1 + q_2 + q_4 , \quad (7.3a)$$

$$\dot{\phi}_2 = [\phi_2, H_P] = 2(q_1 + q_2 + q_4) . \quad (7.3b)$$

Thus, we find the secondary constraints

$$\psi_1 \equiv p_3 + q_1 + q_2 + q_4 = 0 , \quad (7.4a)$$

$$\psi_2 \equiv 2(q_1 + q_2 + q_4) = 0 . \quad (7.4b)$$

These will be denoted collectively by ψ_a .

Applying the consistency condition to the secondary constraints gives

$$\dot{\psi}_1 = [\psi_1, H_P] = p_4 + \lambda_1 + 2\lambda_2 = 0 , \quad (7.5a)$$

$$\dot{\psi}_2 = [\psi_2, H_P] = 2(p_4 + \lambda_1 + 2\lambda_2) = 0 . \quad (7.5b)$$

These equations restrict the Lagrange multipliers to satisfy

$$p_4 + \lambda_1 + 2\lambda_2 = 0 . \quad (7.6)$$

The process has now terminated. In this example there are no tertiary or higher-order constraints.

Recall from the previous section that our goal was to restrict the initial data and Lagrange multipliers such that the primary constraints vanish for all time under the Hamiltonian evolution defined by H_P . We achieve this by imposing the primary constraints at the initial time,

$$\phi_1(0) \equiv p_1(0) = 0 , \quad (7.7a)$$

$$\phi_2(0) \equiv p_2(0) - p_3(0) + p_4(0) = 0 , \quad (7.7b)$$

the secondary constraints at the initial time,

$$\psi_1(0) \equiv p_3(0) + q_1(0) + q_2(0) + q_4(0) = 0 , \quad (7.8a)$$

$$\psi_2(0) \equiv 2[q_1(0) + q_2(0) + q_4(0)] = 0 , \quad (7.8b)$$

and restricting the Lagrange multipliers to satisfy Eq. (7.6) for *all* time t .

Let's review the reasoning. From Eqs. (7.5), the restriction (7.6) on the Lagrange multipliers tells us that $\dot{\psi}_a = 0$ for all time. By Eqs. (7.8), ψ_a vanishes initially, so we see that ψ_a must vanish for all time. Now we use Eqs. (7.3) to conclude that $\dot{\phi}_a$ must vanish for all time. Since ϕ_a vanishes initially, by Eqs. (7.7), it follows that the primary constraints $\phi_a = 0$ must hold for all time t .

8 Weak Equality and Lagrange Multiplier Analysis

It will be useful to follow the general Dirac–Bergmann algorithm closely and carry out a formal analysis of the restriction (7.6) on the Lagrange multipliers [10, 11]. We begin with the concept of *weak equality*.

Let \mathcal{C}_A denote the complete set of (primary, secondary, tertiary, etc.) constraints. For our example,

$$\mathcal{C}_A \equiv \begin{pmatrix} \phi_1 \\ \phi_2 \\ \psi_1 \\ \psi_2 \end{pmatrix} = \begin{pmatrix} p_1 \\ p_2 - p_3 + p_4 \\ p_3 + q_1 + q_2 + q_4 \\ 2(q_1 + q_2 + q_4) \end{pmatrix} \quad (8.1)$$

where the index A runs from 1 to 4.

Two phase space functions F and G are *weakly equal* if they are equal when the (primary, secondary, tertiary, etc.) constraints hold. In other words, F and G are weakly equal if they coincide on the constraint surface, the subspace of phase space defined by $\mathcal{C}_A = 0$. Weak equality is written as $F \approx G$.

Functions F and G are *strongly equal* if they agree throughout phase space. Strong equality is written as $F = G$.

Now we turn to the formal analysis of the restriction (7.6) on the Lagrange multipliers. This restriction can be expressed as the weak equality $\dot{\mathcal{C}}_A \approx 0$. From Eqs. (7.3) and (7.5), we have

$$\dot{\mathcal{C}}_A = [\mathcal{C}, H_P] = \begin{pmatrix} p_3 + q_1 + q_2 + q_4 \\ 2(q_1 + q_2 + q_4) \\ p_4 + \lambda_1 + 2\lambda_2 \\ 2(p_4 + \lambda_2 + 2\lambda_2) \end{pmatrix} \approx \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix} \quad (8.2)$$

which simplifies to

$$\begin{pmatrix} 0 \\ 0 \\ p_4 \\ 2p_4 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 1 & 2 \\ 2 & 4 \end{pmatrix} \begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} \approx \begin{pmatrix} 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}. \quad (8.3)$$

This is a system of inhomogeneous linear equations for the Lagrange multipliers. A particular solution is

$$\begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} \Big|_{\text{particular}} = \begin{pmatrix} 0 \\ -p_4/2 \end{pmatrix}, \quad (8.4)$$

and the homogeneous solutions are

$$\begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} \Big|_{\text{homogeneous}} = \begin{pmatrix} 1 \\ -1/2 \end{pmatrix} \lambda, \quad (8.5)$$

where λ is arbitrary. The general solution is the sum of particular and homogeneous solutions:

$$\begin{pmatrix} \lambda_1 \\ \lambda_2 \end{pmatrix} \Big|_{\text{general}} = \begin{pmatrix} \lambda \\ -(\lambda + p_4)/2 \end{pmatrix}. \quad (8.6)$$

Thus, the restriction (7.6) on the Lagrange multipliers yields $\lambda_1 = \lambda$ and $\lambda_2 = -(\lambda + p_4)/2$, where λ is an arbitrary function of time.

9 Total Hamiltonian

The *total Hamiltonian* H_T is obtained from the primary Hamiltonian H_P by inserting the general solution for the Lagrange multipliers:

$$\begin{aligned} H_T &= H_P \Big|_{\lambda_1=\lambda, \lambda_2=-(\lambda+p_4)/2} = H_C + \lambda\phi_1 - (\lambda + p_4)\phi_2/2 \\ &= \frac{1}{2} [p_3^2 - p_2p_4 + p_3p_4 - 2p_3q_1 - (q_1 + 2q_2)(q_1 + 2q_4)] \\ &\quad + \lambda(p_1 - p_2/2 + p_3/2 - p_4/2) . \end{aligned} \quad (9.1)$$

Physical phase space trajectories are defined by the total Hamiltonian as the weak equality $\dot{F} \approx [F, H_T]$, with initial data that satisfy the complete set of constraints, $\mathcal{C}_A = 0$.

Hamilton's equations for the total Hamiltonian H_T are

$$\dot{q}_1 \approx \lambda , \quad (9.2a)$$

$$\dot{q}_2 \approx -p_4/2 - \lambda/2 , \quad (9.2b)$$

$$\dot{q}_3 \approx p_3 + p_4/2 - q_1 + \lambda/2 , \quad (9.2c)$$

$$\dot{q}_4 \approx p_3/2 - p_2/2 - \lambda/2 , \quad (9.2d)$$

and

$$\dot{p}_1 \approx p_3 + q_1 + q_2 + q_4 , \quad (9.2e)$$

$$\dot{p}_2 \approx q_1 + 2q_4 , \quad (9.2f)$$

$$\dot{p}_3 \approx 0 , \quad (9.2g)$$

$$\dot{p}_4 \approx q_1 + 2q_2 . \quad (9.2h)$$

Since these are weak equalities, we can use the constraints to simplify the results. Observe that the constraints $\mathcal{C}_A = 0$ imply $p_1 = p_3 = 0$, $p_2 + p_4 = 0$ and $q_1 + q_2 + q_4 = 0$. Therefore we can set p_1 and p_3 to zero, replace p_4 with $-p_2$, and replace q_4 with $-q_1 - q_2$. Then Eqs. (9.2d,e,g,h) are either redundant or vacuous, and the remaining equations are

$$\dot{q}_1 \approx \lambda , \quad (9.3a)$$

$$\dot{q}_2 \approx p_2/2 - \lambda/2 , \quad (9.3b)$$

$$\dot{q}_3 \approx -p_2/2 - q_1 + \lambda/2 , \quad (9.3c)$$

$$\dot{p}_2 \approx -q_1 - 2q_2 . \quad (9.3d)$$

These equations, along with the constraints $\mathcal{C}_A = 0$, give a complete description of the physical system.

Let's check the results. The Lagrange multiplier λ can be eliminated from Eqs. (9.3a,b) to give $\dot{q}_1 + 2\dot{q}_2 \approx p_2$. Now differentiate this equation and eliminate \dot{p}_2 with Eq. (9.3d) to obtain $\ddot{q}_1 + 2\ddot{q}_2 \approx -q_1 - 2q_2$. The constraints allow us to set $q_1 \approx -q_2 - q_4$, which gives

$$\ddot{q}_4 - \ddot{q}_2 \approx q_2 - q_4 . \quad (9.4)$$

This is Eq. (2.3 c), which follows directly from Lagrange's equations. The result (2.3 a) from Lagrange's equations is simply the secondary constraint $\mathcal{C}_4 \equiv 2(q_1 + q_2 + q_4) = 0$. Finally, the result (2.3 b) is obtained by summing Eqs. (9.3 b) and (9.3 c).

Recall that Eqs. (2.3 a–c) are equivalent to Lagrange's equations. Thus, we have verified that Hamilton's equations (9.2), along with the primary and secondary constraints $\mathcal{C}_A = 0$, are equivalent to Lagrange's equations.

10 First and Second Class Constraints

A first class function F is a phase space function that has weakly vanishing Poisson brackets with all primary and secondary constraints:

$$[F, \mathcal{C}_A] \approx 0 \iff F \text{ is first class.} \quad (10.1)$$

It can be shown that the Poisson brackets of any two first class functions is itself a first class function [10, 11].

The constraints themselves can be first class; constraints that are not first class are called second class. The constraints are separated into first and second class by examining the matrix of Poisson brackets:

$$[\mathcal{C}_A, \mathcal{C}_B] = \begin{pmatrix} 0 & 0 & -1 & -2 \\ 0 & 0 & -2 & -4 \\ 1 & 2 & 0 & 0 \\ 2 & 4 & 0 & 0 \end{pmatrix}. \quad (10.2)$$

The rank of this 4×4 matrix is 2, and its nullity is $4 - 2 = 2$. It follows that there are 2 independent eigenvectors with eigenvalues equal to zero; for example $u^A = (1, -1/2, 0, 0)$ and $v^A = (0, 0, 1, -1/2)$. Then there are two independent combinations of constraints that are first class, namely $u^A \mathcal{C}_A$ and $v^A \mathcal{C}_A$. (A sum over the repeated index A is implied.) The first class constraints are

$$\mathcal{C}_1^{(fc)} \equiv \phi_1 - \phi_2/2 = p_1 - p_2/2 + p_3/2 - p_4/2, \quad (10.3a)$$

$$\mathcal{C}_2^{(fc)} \equiv \psi_1 - \psi_2/2 = p_3. \quad (10.3b)$$

One can check that the first class conditions $[\mathcal{C}_1^{(fc)}, \mathcal{C}_B] = 0$ and $[\mathcal{C}_2^{(fc)}, \mathcal{C}_B] = 0$ hold. The most general first class constraint is a linear combination of $\mathcal{C}_1^{(fc)}$ and $\mathcal{C}_2^{(fc)}$.

There are two remaining linear combinations of constraints, which we take to be

$$\mathcal{C}_1^{(sc)} \equiv (\phi_1 + \phi_2)/3 = (p_1 + p_2 - p_3 + p_4)/3, \quad (10.4a)$$

$$\mathcal{C}_2^{(sc)} \equiv (\psi_1 + \psi_2)/3 = p_3/3 + q_1 + q_2 + q_4. \quad (10.4b)$$

These are the second class constraints. They have nonvanishing Poisson brackets with each other,

$$[\mathcal{C}_2^{(sc)}, \mathcal{C}_1^{(sc)}] = 1. \quad (10.5)$$

The most general second class constraint is a linear combination of $\mathcal{C}_1^{(sc)}$, $\mathcal{C}_2^{(sc)}$, $\mathcal{C}_1^{(fc)}$ and $\mathcal{C}_2^{(fc)}$, with nonzero coefficients on one or both of $\mathcal{C}_1^{(sc)}$ and $\mathcal{C}_2^{(sc)}$.

The splitting of constraints into first and second class is independent of the splitting into primary and secondary. In this example the first class constraints are mixtures of primary and secondary constraints. Likewise, the second class constraints are mixtures of primary and secondary constraints.

11 First Class Hamiltonian, Gauge Generators and the Dirac Conjecture

The total Hamiltonian (9.1) includes the product of an arbitrary Lagrange multiplier λ with the first class constraint $\mathcal{C}_1^{(fc)} \equiv \phi_1 - \phi_2/2$. We refer to $\mathcal{C}_1^{(fc)}$ as a *primary first class constraint*, since it is constructed entirely from primary constraints.

If we remove the primary first class constraint from the total Hamiltonian, what remains is the *first class Hamiltonian* H_{fc} . That is, the total Hamiltonian can be written as

$$H_T = H_{fc} + \lambda \mathcal{C}_1^{(fc)} , \quad (11.1)$$

where

$$H_{fc} = \frac{1}{2} \left[p_3^2 - p_2 p_4 + p_3 p_4 - 2p_3 q_1 - (q_1 + 2q_2)(q_1 + 2q_4) \right] \quad (11.2)$$

is the first class Hamiltonian. A common notation for H_{fc} , the notation used by Dirac [10], is H' .

We can check directly that the first class Hamiltonian (11.2) is a first class function. But this isn't necessary, because we know that the constraints are preserved under the time evolution defined by H_T . That is, $\dot{\mathcal{C}}_A = [\mathcal{C}_A, H_T] \approx 0$. Thus, the total Hamiltonian must be first class, $[H_T, \mathcal{C}_A] \approx 0$. Of course the primary first class constraint $\mathcal{C}_1^{(fc)}$ is first class. It then follows from the definition (11.1) that H_{fc} must also be a first class function.

The splitting (11.1) of the total Hamiltonian into the first class Hamiltonian and the primary first class constraint is not special to our example. This splitting will occur for any constrained Hamiltonian system [10, 11]. In general, H_T will include the products of every primary first class constraint with an arbitrary multiplier.

Primary first class constraints generate gauge transformations. Consider the change in a phase space function F generated by the primary first class constraint $\mathcal{C}_1^{(fc)}$,

$$\delta F = \delta\epsilon [F, \mathcal{C}_1^{(fc)}] . \quad (11.3)$$

This transformation does not change the physical state of the system. We can see this by considering F to be evaluated as a function of the q 's and p 's at some particular time t . At an infinitesimally later time $t + \delta t$, this function becomes $F(t + \delta t) = F(t) + [F, H_T] \delta t$. In terms of the first class Hamiltonian, we have

$$F(t + \delta t) = F(t) + \left\{ [F, H_{fc}] + \lambda [F, \mathcal{C}_1^{(fc)}] \right\} \delta t . \quad (11.4)$$

The Lagrange multiplier is arbitrary, so we can make a different choice during the time interval from t to $t + \delta t$, say, $\tilde{\lambda}$. Then the function F at time $t + \delta t$ will be

$$\tilde{F}(t + \delta t) = F(t) + \left\{ [F, H_{fc}] + \tilde{\lambda} [F, \mathcal{C}_1^{(fc)}] \right\} \delta t . \quad (11.5)$$

The physical state of the system at $t + \delta t$ should not depend on our choice of Lagrange multiplier, so $F(t + \delta t)$ and $\tilde{F}(t + \delta t)$ must represent the same physical state. The result (11.3) is obtained by subtracting Eq. (11.4) from Eq. (11.5) and defining $\delta F \equiv \tilde{F} - F$ and $\delta\epsilon \equiv (\tilde{\lambda} - \lambda)\delta t$.

For the phase space coordinates, the gauge transformation generated by the primary first class constraint $\mathcal{C}_1^{(fc)}$ is

$$\delta q_1 = \delta\epsilon , \tag{11.6a}$$

$$\delta q_2 = -\delta\epsilon/2 , \tag{11.6b}$$

$$\delta q_3 = \delta\epsilon/2 , \tag{11.6c}$$

$$\delta q_4 = -\delta\epsilon/2 . \tag{11.6d}$$

The transformations of the p 's all vanish. This result agrees with the gauge transformation from Eq. (3.6), with the change of notation $\epsilon \leftrightarrow \delta\epsilon$. Here, we denote the gauge parameter by $\delta\epsilon$ because the transformation is infinitesimal; in Eq. (3.6) we used ϵ because the transformation was finite. It is clear that the infinitesimal transformation (11.6) can be iterated to obtain the finite transformation (3.6).

In general, a gauge transformation is defined as a transformation $\delta F = \delta\epsilon[F, \mathcal{G}]$ that does not alter the physical state of the system. The function \mathcal{G} is the gauge generator. We have seen that the primary first class constraints generate gauge transformations. But not all gauge transformations are generated by primary first class constraints. In fact, it can be shown [10, 11] that the Poisson brackets between any primary first class constraint and the first class Hamiltonian is itself a first class constraint that generates a gauge transformation.⁴

For our example problem, the Poisson brackets of the primary first class constraint $\mathcal{C}_1^{(fc)}$ and the first class Hamiltonian H_{fc} is

$$[\mathcal{C}_1^{(fc)}, H_{fc}] = p_3 . \tag{11.7}$$

This is the secondary first class constraint, $\mathcal{C}_2^{(fc)} \equiv \psi_1 - \psi_2/2 = p_3$. Thus, we see that in this example both the primary and secondary first class constraints are generators of gauge transformations. Explicitly, the transformation $\delta F = \delta\epsilon[F, \mathcal{C}_2^{(fc)}]$ is

$$\delta q_1 = 0 , \tag{11.8a}$$

$$\delta q_2 = 0 , \tag{11.8b}$$

$$\delta q_3 = \delta\epsilon , \tag{11.8c}$$

$$\delta q_4 = 0 , \tag{11.8d}$$

with the transformations of the p 's all vanishing. We can iterate this infinitesimal gauge transformation (11.8) to obtain the finite transformation (3.8).

The ‘‘doubling’’ of the gauge freedom identified in Sec. 3 appears quite naturally in the Hamiltonian formalism. The two types of gauge transformation are generated by the two first class constraints, $\mathcal{C}_1^{(fc)}$ and $\mathcal{C}_2^{(fc)}$.

⁴For systems with more than one primary first class constraint, the Poisson brackets of any two primary first class constraints is also a first class constraint that generates a gauge transformation [10, 11].

The Dirac conjecture [10] says that *all* first class constraints (whether they are primary, secondary, etc., or a combination of primary, secondary, etc.) generate gauge transformations. This conjecture does not hold as a general theorem—there are known examples in which the transformation generated by a secondary first class constraint does not coincide with any invariance of the original Lagrangian system.⁵ Nevertheless, the Dirac conjecture is usually taken as an assumption. It appears that in practice, for systems of physical interest, all first class constraints generate gauge transformations.

12 Extended Hamiltonian

The Dirac conjecture tells us that all first class constraints generate gauge transformations and should be treated on an equal footing. The *extended* Hamiltonian H_E is defined by adding all first class constraints $\mathcal{C}_a^{(fc)}$ with Lagrange multipliers γ_a to the first class Hamiltonian:

$$H_E = H_{fc} + \gamma_a \mathcal{C}_a^{(fc)} . \quad (12.1)$$

(A sum over the index a is implied.)

For our example, the extended Hamiltonian is

$$H_E = \frac{1}{2} \left[p_3^2 - p_2 p_4 + p_3 p_4 - 2 p_3 q_1 - (q_1 + 2 q_2)(q_1 + 2 q_4) \right] + \gamma_1 (p_1 - p_2/2 + p_3/2 - p_4/2) + \gamma_2 p_3 . \quad (12.2)$$

The equations of motion $\dot{F} \approx [F, H_E]$ are

$$\dot{q}_1 \approx \gamma_1 , \quad (12.3a)$$

$$\dot{q}_2 \approx -p_4/2 - \gamma_1/2 , \quad (12.3b)$$

$$\dot{q}_3 \approx p_3 + p_4/2 - q_1 + \gamma_1/2 + \gamma_2 , \quad (12.3c)$$

$$\dot{q}_4 \approx p_3/2 - p_2/2 - \gamma_1/2 , \quad (12.3d)$$

and

$$\dot{p}_1 \approx p_3 + q_1 + q_2 + q_4 , \quad (12.3e)$$

$$\dot{p}_2 \approx q_1 + 2q_4 , \quad (12.3f)$$

$$\dot{p}_3 \approx 0 , \quad (12.3g)$$

$$\dot{p}_4 \approx q_1 + 2q_2 . \quad (12.3h)$$

Let's compare these results to the equations of motion (9.2) obtained from the total Hamiltonian H_T . There are just two differences. The first is trivial: the Lagrange multiplier λ in Eqs. (9.2) has changed names to γ_1 in Eqs. (12.3). The second difference is significant: the equation for \dot{q}_3 has an extra term γ_2 on the right-hand side. This is a new feature of the extended Hamiltonian. It makes explicit the fact that the gauge freedom allows q_3 to be changed arbitrarily, and independently, from the other variables.

⁵Counterexamples to the Dirac conjecture are discussed in Refs. [11, 19–25] and elsewhere. Proofs of the conjecture have been constructed by adopting various simplifying assumptions [11, 26, 27]. The status of the conjecture is a subtle issue; see for example Refs. [17, 28, 29].

We can check the equations of motion for H_E following the same reasoning that was applied to the equations of motion for H_T . First recall that the (first and second class) constraints imply $p_1 = p_3 = 0$, $p_4 = -p_2$ and $q_4 = -q_1 - q_2$. Then Eqs. (12.3 e) and (12.3 g) are vacuous, and Eqs. (12.3 f) and (12.3 h) are redundant. It also follows that with the constraints imposed, Eq. (12.3 d) is a consequence of Eqs. (12.3 a) and (12.3 b). The remaining equations are

$$\dot{q}_1 \approx \gamma_1 , \quad (12.4a)$$

$$\dot{q}_2 \approx p_2/2 - \gamma_1/2 , \quad (12.4b)$$

$$\dot{q}_3 \approx -p_2/2 - q_1 + \gamma_1/2 + \gamma_2 , \quad (12.4c)$$

$$\dot{p}_2 \approx -q_1 - 2q_2 . \quad (12.4d)$$

These agree with Eqs. (9.3), apart from the change of notation $\lambda \rightarrow \gamma_1$ and the extra term γ_2 on the right-hand side of the \dot{q}_3 equation.

By eliminating γ_1 , the equations of motion generated by the extended Hamiltonian H_E become

$$\dot{q}_1 + 2\dot{q}_2 \approx p_2 , \quad (12.5a)$$

$$\dot{q}_2 + \dot{q}_3 \approx -q_1 + \gamma_2 , \quad (12.5b)$$

$$\dot{p}_2 \approx -q_1 - 2q_2 . \quad (12.5c)$$

If we differentiate the first equation, combine with the second, and use the constraint $q_1 + q_2 + q_4 = 0$, we obtain $(\ddot{q}_4 - \ddot{q}_2) \approx -(q_4 - q_2)$. This is the expected result (2.3 c). In fact, the only difference between Hamilton's equations $\dot{F} \approx [F, H_E]$ and the results (2.3) (which are equivalent to Lagrange's equations) is the extra term γ_2 in Eq. (12.5 b) above. That term does not appear in the corresponding Lagrangian equation (2.3 b).

Note that we can use the first class constraints to simplify the extended Hamiltonian H_E . For example, using $C_2^{(fc)} = p_3$, we can set $p_3 = 0$ everywhere in Eq. (12.2), except of course in the term $\gamma_2 p_3$. The extended Hamiltonian becomes

$$H_E = \frac{1}{2} \left[-p_2 p_4 - (q_1 + 2q_2)(q_1 + 2q_4) \right] + \gamma_1 (p_1 - p_2/2 - p_4/2) + \gamma_2 p_3 . \quad (12.6)$$

This amounts to replacing the Lagrange multiplier γ_2 in Eq. (12.2) by

$$\gamma_2 \rightarrow \gamma_2 - p_3/2 - p_4/2 + q_1 - \gamma_1/2 . \quad (12.7)$$

This replacement does not change the physical content of the theory, since the Lagrange multiplier γ_2 is arbitrary.

13 Extended Action

The equations of motion for the extended theory can be derived from the action [11]

$$S_E[q, p, \gamma, \sigma] = \int_0^T dt \left\{ p_i \dot{q}_i - H_E - \sigma_1 C_1^{(sc)} - \sigma_2 C_2^{(sc)} \right\} , \quad (13.1)$$

which includes the second class constraints with Lagrange multipliers σ_a . Recall that the first class constraints $\mathcal{C}_a^{(fc)}$ are included in the extended Hamiltonian H_E with multipliers γ_a , so S_E includes all four constraints.

We can use either form of the extended Hamiltonian, Eq. (12.2) or (12.6), in the extended action. Let's use Eq. (12.2). Then the equations of motion that follow from extremizing S_E with respect to the momenta p_i are

$$\dot{q}_1 = \gamma_1 + \sigma_1/3, \quad (13.2a)$$

$$\dot{q}_2 = -p_4/2 - \gamma_1/2 + \sigma_1/3, \quad (13.2b)$$

$$\dot{q}_3 = p_3 + p_4/2 - q_1 + \gamma_1/2 + \gamma_2 - \sigma_1/3 + \sigma_2/3, \quad (13.2c)$$

$$\dot{q}_4 = p_3/2 - p_2/2 - \gamma_1/2 + \sigma_1/3. \quad (13.2d)$$

Extremizing S_E with respect to the coordinates yields

$$\dot{p}_1 = p_3 + q_1 + q_2 + q_4 - \sigma_2, \quad (13.2e)$$

$$\dot{p}_2 = q_1 + 2q_4 - \sigma_2, \quad (13.2f)$$

$$\dot{p}_3 = 0, \quad (13.2g)$$

$$\dot{p}_4 = q_1 + 2q_2 - \sigma_2, \quad (13.2h)$$

and the constraints

$$\mathcal{C}_1^{(fc)} \equiv p_1 - p_2/2 + p_3/2 - p_4/2 = 0, \quad (13.2i)$$

$$\mathcal{C}_2^{(fc)} \equiv p_3 = 0, \quad (13.2j)$$

$$\mathcal{C}_1^{(sc)} \equiv (p_1 + p_2 - p_3 + p_4)/3 = 0, \quad (13.2k)$$

$$\mathcal{C}_2^{(sc)} \equiv p_3/3 + q_1 + q_2 + q_4 = 0, \quad (13.2l)$$

follow from extremizing S_E with respect to the Lagrange multipliers γ_a and σ_a .

Let's check these equations of motion. The constraints imply $p_1 = p_3 = 0$, $p_4 = -p_2$ and $q_1 + q_2 + q_4 = 0$. Then the equation of motion (13.2e) gives $\sigma_2 \approx 0$. The constraints also imply $\dot{q}_1 + \dot{q}_2 + \dot{q}_4 = 0$. The sum of Eqs. (13.2a), (13.2b) and (13.2d) then yields $\sigma_1 \approx 0$. Now, if we set σ_1 and σ_2 to zero, the equations of motion (13.2a–h) agree precisely with Hamilton's equations (12.3) for the extended Hamiltonian H_E . In Sec. 12 we showed that the equations generated by H_E agree with Lagrange's equations apart from the extra term γ_2 in the equation for \dot{q}_3 . This term extends the original Lagrangian theory by making explicit the fact that the gauge freedom allows for independent transformations of q_3 .

Finally, we note that the extended action is invariant under the transformation defined by

$$\delta F = \epsilon_1[F, \mathcal{C}_1^{(fc)}] + \epsilon_2[F, \mathcal{C}_2^{(fc)}] \quad (13.3a)$$

for the phase space variables and

$$\delta\gamma_1 = \dot{\epsilon}_1, \quad (13.3b)$$

$$\delta\gamma_2 = \dot{\epsilon}_2 + \epsilon_1, \quad (13.3c)$$

$$\delta\sigma_1 = 0, \quad (13.3d)$$

$$\delta\sigma_2 = 0, \quad (13.3e)$$

for the Lagrange multipliers. Here, the gauge parameters ϵ_1 and ϵ_2 are functions of time. These equations express the gauge invariance at the level of the action S_E .

14 Dirac Brackets and the Partially Reduced Hamiltonian

We now return to the evolution defined by the extended Hamiltonian H_E of Eq. (12.2), and Hamilton's equations (12.3). To obtain a physically allowed trajectory, we must choose initial data that satisfy the four constraints $\mathcal{C}_a^{(fc)} = 0$ and $\mathcal{C}_a^{(sc)} = 0$. Apart from restricting the initial data, the second class constraints play no role in the formalism. It would be convenient if we could restrict the variables from the outset such that the second class constraints are automatically satisfied. For example, we could use $\mathcal{C}_1^{(sc)} = 0$ and $\mathcal{C}_2^{(sc)} = 0$ from Eqs. (13.2 k,l) to replace q_1 with $-q_2 - q_4 - p_3/3$ and replace p_2 with $-p_1 + p_3 - p_4$.

We are not allowed to apply the second class constraints in this way. For example, consider the Poisson brackets $[q_1, p_1] = 1$. If we were to replace q_1 with $-q_2 - q_4 - p_3/3$, we would find a different answer: $[-q_2 - q_4 - p_3/3, p_1] = 0$. The second class constraints cannot be imposed before Poisson brackets are computed.

Dirac devised a way to allow the second class constraints to be imposed from the outset by modifying the Poisson brackets [10]. The result is the Dirac brackets.

To construct Dirac brackets, we first compute the matrix of Poisson brackets among the second class constraints:

$$M_{ab} \equiv [\mathcal{C}_a^{(sc)}, \mathcal{C}_b^{(sc)}] = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}. \quad (14.1)$$

Let

$$M^{ab} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \quad (14.2)$$

denote the inverse of M_{ab} . Then the Dirac brackets $[F, G]^*$ of two phase space functions F and G are defined by

$$[F, G]^* \equiv [F, G] - [F, \mathcal{C}_a^{(sc)}] M^{ab} [\mathcal{C}_b^{(sc)}, G]. \quad (14.3)$$

Dirac brackets, like Poisson brackets, are antisymmetric and satisfy the Jacobi identity [10, 11].

Explicitly, the Dirac brackets among the coordinates are

$$[q_i, q_j]^* = \frac{1}{9} \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 \\ -1 & -1 & 0 & -1 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \quad (14.4)$$

and the Dirac brackets between the q 's and p 's are

$$[q_i, p_j]^* = \frac{1}{3} \begin{pmatrix} 2 & -1 & 0 & -1 \\ -1 & 2 & 0 & -1 \\ 1 & 1 & 3 & 1 \\ -1 & -1 & 0 & 2 \end{pmatrix} \quad (14.5)$$

For our example the Dirac brackets among the momenta all vanish: $[p_i, p_j]^* = 0$.

There are two key properties that make Dirac brackets relevant. First, the Dirac brackets agree weakly with Poisson brackets if one of the two functions is first class. Since the extended Hamiltonian is first class, we have $[F, H_E]^* \approx [F, H_E]$ for any F . It follows that we can write the equations of motion as

$$\dot{F} \approx [F, H_E]^* , \quad (14.6)$$

using Dirac brackets.

The second key property of the Dirac brackets is that they weakly vanish if one of the functions is a second class constraint: $[F, \mathcal{C}_a^{(sc)}]^* \approx 0$. This allows us to apply the second class constraints before computing brackets. For example, we can use either q_1 or $-q_2 - q_4 - p_3/3$ to compute Dirac brackets with p_1 :

$$[q_1, p_1]^* = [-q_2 - q_4 - p_3/3, p_1]^* = 2/3 . \quad (14.7)$$

With Dirac brackets, the second class constraints can be treated as strong equations and imposed before computing the equations of motion.

Let's use the second class constraints (13.2k,l) to eliminate q_1 and p_2 and write the extended Hamiltonian in terms of the smaller set of variables q_2, q_3, q_4, p_1, p_3 and p_4 . Setting

$$q_1 = -q_2 - q_4 - p_3/3 , \quad (14.8a)$$

$$p_2 = -p_1 + p_3 - p_4 , \quad (14.8b)$$

we have

$$H_{PR} = 7p_3^2/9 + \frac{1}{2} [p_4^2 + p_1p_4 + 2p_3(q_2 + q_4) + (q_2 - q_4)^2] + \gamma_1(3p_1/2) + \gamma_2p_3 . \quad (14.9)$$

This is the *partially reduced Hamiltonian*, obtained from the extended Hamiltonian by applying the second class constraints.

Of course the partially reduced Hamiltonian is not unique. We could use the second class constraints to eliminate some other pair of variables instead of q_1 and p_2 .

The equations of motion generated by the partially reduced Hamiltonian, $\dot{F} \approx [F, H_R]^*$, are

$$\dot{q}_2 \approx -p_1/6 - p_4/2 - \gamma_1/2 , \quad (14.10a)$$

$$\dot{q}_3 \approx 4p_3/3 + p_1/6 + p_4/2 + q_2 + q_4 + \gamma_1/2 + \gamma_2 , \quad (14.10b)$$

$$\dot{q}_4 \approx p_1/3 + p_4/2 - \gamma_1/2 , \quad (14.10c)$$

$$\dot{p}_1 \approx 2p_3/3 , \quad (14.10d)$$

$$\dot{p}_3 \approx 0 , \quad (14.10e)$$

$$\dot{p}_4 \approx -p_3/3 + q_2 - q_4 . \quad (14.10f)$$

We can also use H_{PR} and the Dirac brackets to compute \dot{q}_1 and \dot{p}_2 . The results are equivalent to those obtained by differentiating the right-hand sides of Eqs. (14.8) and using the equations of motion (14.10).

Let's check the equations of motion. With the second class constraints applied, the first class constraints imply $p_1 = p_3 = 0$ and $p_4 = -p_2$. Thus, Eqs. (14.10 d) and (14.10 e) are vacuous and the remaining equations become

$$\dot{q}_2 \approx p_2/2 - \gamma_1/2 , \quad (14.11a)$$

$$\dot{q}_3 \approx -p_2/2 + q_2 + q_4 + \gamma_1/2 + \gamma_2 , \quad (14.11b)$$

$$\dot{q}_4 \approx -p_2/2 - \gamma_1/2 , \quad (14.11c)$$

$$\dot{p}_2 \approx -q_2 + q_4 , \quad (14.11d)$$

Compare these to the independent equations (12.4) that follow from the extended Hamiltonian. Equations (12.4 b,c,d) agree with Eqs. (14.11 a,b,d) once we use $q_1 = -q_2 - q_4$. The final equation (12.4 a) is obtained by differentiating $q_1 = -q_2 - q_3$ in time and using Eqs.(14.11 a) and (14.11 b).

15 Partially Reduced Action

The partially reduced equations of motion (14.10) can be obtained from the extended action S_E by eliminating the superfluous variables. Note that the equations of motion obtained by varying S_E with respect to p_2 , q_1 , σ_1 and σ_2 are Eqs. (13.2 b,e,k,l), respectively. We can eliminate these variables by solving these equations and substituting the results into the action.⁶ The result is:

$$q_1 = -p_3/3 - q_2 - q_4 , \quad (15.1a)$$

$$p_2 = -p_1 + p_3 - p_4 , \quad (15.1b)$$

$$\sigma_1 = 3\dot{q}_2 + 3p_4/2 + 3\gamma_1/2 , \quad (15.1c)$$

$$\sigma_2 = -\dot{p}_1 + 2p_3/3 . \quad (15.1d)$$

Inserting these into the extended action (13.1), we find

$$\begin{aligned} & S_{PR}[q_2, q_3, q_4, p_1, p_3, p_4, \gamma_1, \gamma_2] \\ &= \int_0^T dt \{ -p_1\dot{p}_3/3 + (p_3 - p_4 - 2p_1)\dot{q}_2 + p_3\dot{q}_3 + (p_4 - p_1)\dot{q}_4 - H_{PR} \} . \quad (15.2) \end{aligned}$$

This is the partially reduced action.

⁶Any action can be reduced by using the equations of motion obtained by varying with respect some set of variables, solving those equations for the same set of variables, then substituting the results into the action. In general, it is not permissible to reduce an action by using the equations obtained by varying with respect to one set of variables but solving those equations for a different set of variables.

The equations of motion obtained from varying S_{PR} with respect to the phase space variables are

$$2\dot{p}_1 - \dot{p}_3 + \dot{p}_4 - p_3 - q_2 + q_4 = 0 , \quad (15.3a)$$

$$-\dot{p}_3 = 0 , \quad (15.3b)$$

$$\dot{p}_1 - \dot{p}_4 - p_3 + q_2 - q_4 = 0 , \quad (15.3c)$$

$$-2\dot{q}_2 - \dot{q}_4 - \dot{p}_3/3 - p_4/2 - 3\gamma_1/2 = 0 , \quad (15.3d)$$

$$\dot{q}_2 + \dot{q}_3 + \dot{p}_1/3 - 14p_3/9 - q_2 - q_4 - \gamma_2 = 0 , \quad (15.3e)$$

$$\dot{q}_2 + \dot{q}_4 - p_1/2 - p_4 = 0 , \quad (15.3f)$$

and the equations obtained by varying with respect to the Lagrange multipliers γ_1 and γ_2 are

$$-3p_1/2 = 0 , \quad (15.4a)$$

$$-p_3 = 0 . \quad (15.4b)$$

These are of course the first class constraints, reduced by using the second class constraints to eliminate q_1 and p_2 .

We can now solve the equations (15.3) for the time derivatives of q_2, q_3, q_4, p_1, p_3 and p_4 . The result coincides with the equations of motion (14.10) obtained from the partially reduced Hamiltonian H_{PR} and the Dirac brackets.

The partially reduced action S_{PR} is invariant under the transformation defined by

$$\delta F = \epsilon_1 [F, C_1^{(fc)}]^* + \epsilon_2 [F, C_2^{(fc)}]^* \quad (15.5a)$$

for the phase space variables and

$$\delta\gamma_1 = \dot{\epsilon}_1 , \quad (15.5b)$$

$$\delta\gamma_2 = \dot{\epsilon}_2 + \epsilon_1 , \quad (15.5c)$$

for the Lagrange multipliers. These equations express the gauge invariance of the theory at the level of the action principle with the second class constraints eliminated.

Finally, it is not too difficult to find a change of variables that will bring S_{PR} into "canonical form". For example, let

$$q_2 = Q_1 - P_2/9 , \quad (15.6a)$$

$$q_3 = Q_2 + P_2/9 , \quad (15.6b)$$

$$q_4 = Q_3 - P_2/9 , \quad (15.6c)$$

$$p_1 = (-P_1 + P_2 - P_3)/3 , \quad (15.6d)$$

$$p_3 = P_2 , \quad (15.6e)$$

$$p_4 = (-P_1 + P_2 + 2P_3)/3 , \quad (15.6f)$$

define a new set of variables Q_α, P_α for the secondary constraint surface. (The index α ranges over 1, 2 and 3.) The partially reduced action becomes

$$S_{PR}[Q, P, \gamma] = \int_0^T dt \left\{ P_\alpha \dot{Q}_\alpha - H_{PR} \right\} \quad (15.7)$$

with

$$H_{PR} = \frac{1}{18} [2P_1^2 + 12P_2^2 + 2P_3^2 - 4P_1P_2 + 5(P_2 - P_1)P_3 + 18P_2(Q_1 + Q_3) + 9(Q_1 - Q_3)^2] + \gamma_1(-P_1 + P_2 - P_3)/2 + \gamma_2P_2 . \quad (15.8)$$

The equations of motion $\delta S_{PR} = 0$ include $\dot{Q}_\alpha = [Q_\alpha, H_{PR}]$ and $\dot{P}_\alpha = [P_\alpha, H_{PR}]$, where $[\cdot, \cdot]$ are the usual Poisson brackets.

16 Gauge Conditions and the Fully Reduced Hamiltonian

Let's return to the theory described by the extended Hamiltonian, prior to the elimination of the second class constraints.

For our example problem, phase space is eight-dimensional. The physical trajectories fill the constraint "surface", which is the four-dimensional subspace where all first and second class constraints hold. Each point in the constraint surface can be mapped into a physically equivalent state by the gauge generators, namely, the first class constraints $\mathcal{C}_a^{(fc)}$. Since there are two independent gauge generators, each physical state of the system corresponds to a two-dimensional subspace of the constraint surface. The constraint surface is foliated by these two-dimensional slices, referred to as gauge "orbits".

We can select a single phase space point on each gauge orbit to represent the physical state. We do this by applying gauge conditions. In particular we will consider a *canonical gauge*⁷ which takes the form $\mathcal{G}_a(q, p) \approx 0$ with $a = 1, 2$. A good canonical gauge condition must not be gauge invariant, otherwise it would allow more than one point on the gauge orbit to represent the physical state of the system. To be precise, the matrix of Poisson brackets of gauge conditions and gauge generators, $[\mathcal{G}_a, C_b^{(fc)}]$, must be nonsingular [11].

As an example, let's choose

$$\mathcal{G}_1 = q_1 - q_2 , \quad (16.1a)$$

$$\mathcal{G}_2 = q_3 + p_4 . \quad (16.1b)$$

as our gauge conditions. This is a good gauge:

$$\det[\mathcal{G}_a, C_b^{(fc)}] = \begin{vmatrix} 3/2 & 0 \\ 1/2 & 1 \end{vmatrix} = 3/2 . \quad (16.2)$$

The matrix $[\mathcal{G}_a, C_b^{(fc)}]$ is nonsingular, as required.

The gauge conditions $\mathcal{G}_a = 0$, like the first and second class constraints, restrict the phase space variables. The full set of restrictions

$$\mathcal{C}_A^{(all)} = \{\mathcal{G}_1, \mathcal{G}_2, \mathcal{C}_1^{(fc)}, \mathcal{C}_2^{(fc)}, \mathcal{C}_1^{(sc)}, \mathcal{C}_2^{(sc)}\} \quad (16.3)$$

⁷Canonical gauges restrict the phase space variables. Noncanonical gauges [11] involve the Lagrange multipliers.

reduce the available phase space from eight dimensions to two dimensions. (Here, the index A ranges from 1 to 6.) Taken as a whole, the six conditions $\mathcal{C}_A^{(all)}$ are second class. We see this by computing the Poisson brackets

$$M_{AB} \equiv [\mathcal{C}_A^{(all)}, \mathcal{C}_B^{(all)}] = \frac{1}{6} \begin{pmatrix} 0 & 0 & 9 & 0 & 0 & 0 \\ 0 & 0 & 3 & 6 & -2 & -4 \\ -9 & -3 & 0 & 0 & 0 & 0 \\ 0 & -6 & 0 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 & 0 & -6 \\ 0 & 4 & 0 & 0 & 6 & 0 \end{pmatrix}. \quad (16.4)$$

This matrix has nonzero determinant, $\det(M) = 9/4$, which is the condition for the set of constraints and gauge conditions to be second class.

We can eliminate the constraints and gauge conditions by constructing Dirac brackets. The inverse of M_{AB} is

$$M^{AB} = \frac{1}{3} \begin{pmatrix} 0 & 0 & -2 & 1 & 0 & 0 \\ 0 & 0 & 0 & -3 & 0 & 0 \\ 2 & 0 & 0 & 0 & 0 & 0 \\ -1 & 3 & 0 & 0 & -2 & 1 \\ 0 & 0 & 0 & 2 & 0 & 3 \\ 0 & 0 & 0 & -1 & -3 & 0 \end{pmatrix}, \quad (16.5)$$

and the Dirac brackets are defined by

$$[F, G]^* = [F, G] - [F, \mathcal{C}_A^{(all)}] M^{AB} [\mathcal{C}_B^{(all)}, G]. \quad (16.6)$$

The Dirac brackets among the phase space variables are

$$[q_1, q_3]^* = [q_1, p_2]^* = -[q_1, p_4]^* = 1/3, \quad (16.7a)$$

$$[q_2, q_3]^* = [q_2, p_2]^* = -[q_2, p_4]^* = 1/3, \quad (16.7b)$$

$$[q_3, q_4]^* = -[q_4, p_2]^* = [q_4, p_4]^* = 2/3, \quad (16.7c)$$

with all other brackets vanishing.

The constraints can be solved in various ways and the results can be used freely, either before or after computing Dirac brackets. For example, the constraints imply

$$q_1 = -q_4/2, \quad (16.8a)$$

$$q_2 = -q_4/2, \quad (16.8b)$$

$$q_3 = p_2, \quad (16.8c)$$

$$p_1 = 0, \quad (16.8d)$$

$$p_3 = 0, \quad (16.8e)$$

$$p_4 = -p_2. \quad (16.8f)$$

We can use these to eliminate the variables q_1 , q_2 , q_3 , p_1 , p_3 and p_4 . Then the extended Hamiltonian H_E becomes the *fully reduced Hamiltonian*

$$H_{FR} = \frac{1}{2} \left[p_2^2 + \frac{9}{4} q_4^2 \right], \quad (16.9)$$

which depends only on q_4 and p_2 .

The Dirac brackets of the variables that remain are $[q_4, p_2]^* = -2/3$. Thus, the equations of motion become

$$\dot{q}_4 = [q_4, H_{FR}]^* = -2p_2/3 , \quad (16.10a)$$

$$\dot{p}_2 = [p_2, H_{FR}]^* = 3q_4/2 . \quad (16.10b)$$

These are the equations for a simple harmonic oscillator with solution

$$q_4(t) = \alpha \sin t + \beta \cos t , \quad (16.11a)$$

$$p_2(t) = -(3\alpha/2) \cos t + (3\beta/2) \sin t , \quad (16.11b)$$

where α and β are arbitrary constants.

With the gauge fixed the dynamics take place on the fully reduced phase space, the two-dimensional surface defined by the constraints and gauge conditions $\mathcal{C}_A^{(all)} = 0$. There are many different choices of coordinates for this surface. Instead of solving the constraints for q_4 and p_2 , we could solve them for q_2 and q_3 . In that case the fully reduced Hamiltonian is

$$H_{FR} = \frac{9}{2}q_2^2 + \frac{1}{2}q_3^2 \quad (16.12)$$

and Hamilton's equations are

$$\dot{q}_2 = [q_2, H_{FR}]^* = q_3/3 , \quad (16.13a)$$

$$\dot{q}_3 = [q_3, H_{FR}]^* = -3q_2 \quad (16.13b)$$

Again, this describes the simple harmonic oscillator.

We can choose other coordinates on the fully reduced phase space. For example, let $q_4 = Q + q_2$ and $p_2 = -P$, then use the constraints to eliminate q_1 , q_2 , q_3 , p_1 , p_3 and p_4 . The fully reduced Hamiltonian becomes

$$H_{FR} = \frac{1}{2}(Q^2 + P^2) . \quad (16.14)$$

The nonzero Dirac brackets are $[Q, P]^* = 1$, and the equations of motion are simply $\dot{Q} = P$ and $\dot{P} = -Q$.

In each case, the fully reduced theory exhibits the single physical degree of freedom that we expect.

17 Fully Reduced Action

The fully reduced equations of motion can be derived from the action that includes all of the constraints and gauge conditions. For lack of a better name, let's denote this action with the subscript "all":

$$S_{all} = \int_0^T dt \left\{ p_i \dot{q}_i - H_{fc} - \gamma_a \mathcal{C}_a^{(fc)} - \sigma_a \mathcal{C}_a^{(sc)} - \rho_a \mathcal{G}_a \right\} . \quad (17.1)$$

Now extremize S_{all} with respect to variations in $p_1, p_3, p_4, q_1, q_2, q_3$ and q_4 :

$$\dot{q}_1 = \gamma_1 + \sigma_1/3 , \quad (17.2a)$$

$$\dot{q}_3 = -q_1 + p_3 + p_4/2 + \gamma_1/2 + \gamma_2 - \sigma_1/3 + \sigma_2/3 , \quad (17.2b)$$

$$\dot{q}_4 = -p_2/2 + p_3/2 - \gamma_1/2 + \sigma_1/3 + \rho_2 , \quad (17.2c)$$

$$\dot{p}_1 = q_1 + q_2 + q_4 + p_3 - \sigma_2 - \rho_1 , \quad (17.2d)$$

$$\dot{p}_2 = q_1 + 2q_4 + \sigma_2 - \rho_1 , \quad (17.2e)$$

$$\dot{p}_3 = -\rho_2 . \quad (17.2f)$$

Also vary S_{all} with respect to the Lagrange multipliers to obtain the constraints $\mathcal{C}_A^{(all)} = 0$. The solution of the full set of equations, (17.2) and $\mathcal{C}_A^{(all)} = 0$, is given by Eqs. (16.8) along with

$$\gamma_1 = -p_2/3 - 2\dot{p}_3/2 + 2\dot{q}_1/3 - 2\dot{q}_4/3 , \quad (17.3a)$$

$$\gamma_2 = p_2 - 3q_4/4 + \dot{p}_1/6 + \dot{p}_2/6 + \dot{p}_3 + \dot{q}_3 + \dot{q}_4 , \quad (17.3b)$$

$$\sigma_1 = p_2 + 2\dot{p}_3 + \dot{q}_1 + 2\dot{q}_4 , \quad (17.3c)$$

$$\sigma_2 = 3q_4/4 - \dot{p}_1/2 - \dot{p}_2/2 , \quad (17.3d)$$

$$\rho_1 = -3q_4/4 - \dot{p}_1/2 + \dot{p}_2/2 , \quad (17.3e)$$

$$\rho_2 = -\dot{p}_3 . \quad (17.3f)$$

We now insert these results into S_{all} to obtain the *fully reduced action*

$$S_{FR}[q_4, p_2] = \int_0^T dt \left\{ -\frac{3}{2}p_2\dot{q}_4 - \frac{1}{2} \left[p_2^2 + \frac{9}{4}q_4^2 \right] \right\} , \quad (17.4)$$

which is a functional of q_4 and p_2 . The equations of motion $\delta S_{FR} = 0$ are

$$0 = \frac{\delta S_{FR}}{\delta q_4} = \frac{3}{2}\dot{p}_2 - \frac{9}{4}q_4 , \quad (17.5a)$$

$$0 = \frac{\delta S_{FR}}{\delta p_2} = -\frac{3}{2}\dot{q}_4 - p_2 . \quad (17.5b)$$

These are equivalent to Hamilton's equations (16.10) for the fully reduced Hamiltonian.

We can place S_{FR} into "canonical form" by defining new variables $P = -p_2$ and $Q = 3q_4/2$. Then

$$S_{FR}[Q, P] = \int_0^T dt \left\{ P\dot{Q} - \frac{1}{2}[P^2 + Q^2] \right\} , \quad (17.6)$$

which is the familiar action for the harmonic oscillator.

18 Summary and Comments

Here is the Dirac–Bergmann algorithm:

- Compute the conjugate momenta $p_i = \partial L / \partial \dot{q}_i$ and define the canonical Hamiltonian H_C as $p_i \dot{q}^i - L(q, \dot{q})$, written in terms of p 's and q 's.

- Identify the primary constraints. The primary Hamiltonian H_P is obtained from H_C by adding the primary constraints with Lagrange multipliers.
- Apply Dirac's consistency conditions to identify higher-order constraints and restrictions on the Lagrange multipliers.
- The total Hamiltonian H_T is found from H_P by incorporating the restrictions on the Lagrange multipliers.
- Separate the primary, secondary, and higher-order constraints into first and second class.
- The first class Hamiltonian H_{fc} is the part of H_T with the primary first class constraints removed.
- The extended Hamiltonian H_E is obtained from the first class Hamiltonian H_{fc} by adding all of the first class constraints with Lagrange multipliers.
- The partially reduced Hamiltonian H_{PR} is found from H_E by using Dirac brackets to eliminate the second class constraints.
- Gauge freedom is removed by assigning gauge conditions. The fully reduced Hamiltonian H_{FR} is obtained from H_E by using Dirac brackets to impose all constraints and gauge conditions.

The theory defined by the singular Lagrangian (1.1) provides a relatively complete example of each step in the algorithm.

One reason the Dirac–Bergmann algorithm can be confusing is that typical examples are chosen for simplicity, allowing some of the logical steps to be skipped. This causes the distinction between Hamiltonians to become blurred. For example, if there are no restrictions on the Lagrange multipliers, then the primary Hamiltonian H_P and the total Hamiltonian H_T coincide. Likewise, if there are no secondary (or higher-order) first class constraints, then the total Hamiltonian H_T and the extended Hamiltonian H_E coincide. Also, for theories with no second class constraints and no gauge conditions imposed, Dirac brackets and the reduction process are not needed.

Acknowledgments

I would like to thank Claudio Bunster and Marc Henneaux for sparking my interest in this subject many years ago. I also thank Ashiqul Islam Dip for helpful discussions.

References

- [1] P. Bergmann, *Non-Linear Field Theories*, *Physical Review* **75** (1949) 680.
- [2] P. Bergmann and J. Brunings, *Non-Linear Field Theories II. Canonical Equations and Quantization*, *Rev. Mod. Phys.* **21** (1949) 480.

- [3] P. Dirac, *Generalized Hamiltonian dynamics*, *Canadian Journal of Mathematics* **2** (1950) 129.
- [4] P. Bergmann, R. Penfield, R. Schiller, and H. Zatzkis, *The Hamiltonian of the General Theory of Relativity with Electromagnetic Field*, *Physical Review* **80** (1950) 81.
- [5] P. Dirac, *The Hamiltonian form of field dynamics*, *Canadian Journal of Mathematics* **3** (1951) 1.
- [6] J. Anderson and P. Bergmann, *Constraints in Covariant Field Theories*, *Physical Review* **83** (1951) 1018.
- [7] P. Bergmann and I. Goldberg, *Dirac bracket transformations in phase space*, *Physical Review* **98** (1955) 531.
- [8] P. Dirac, *Generalized Hamiltonian dynamics*, *Proc. Roy. Soc. (London)* **246** (1958) 326.
- [9] P. Dirac, *The theory of gravitation in Hamiltonian form*, *Proc. Roy. Soc. (London)* **246** (1958) 333.
- [10] P. Dirac, *Lectures on Quantum Mechanics*. Belfer Graduate School of Sciences, Yeshiva University, New York, 1964.
- [11] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems*. Princeton University Press, Princeton, New Jersey, 1992.
- [12] L. Rosenfeld, *Zur Quantelung der Wellenfelder*, *Annalen der Physik* **397** (1930) 113.
- [13] L. Rosenfeld, *On the quantization of wave fields*, *Eur. Phys. J. H* **42** (2017) 63.
- [14] R. Penfeld, *Hamiltonians without Parametrization*, *Physical Review* **84** (1951) 737.
- [15] A. Hanson, T. Regge, and C. Teitelboim, *Constrained Hamiltonian Systems*. Accademia Nazionale dei Lincei, Roma, 1976.
- [16] K. Sundermeyer, *Constrained Dynamics*. Springer-Verlag, Berlin, 1982.
- [17] H. Rothe and K. Rothe, *Classical and Quantum Dynamics of Constrained Hamiltonian Systems*. World Scientific, Singapore, 2010.
- [18] L. Lusanna, *Non-Inertial Frames and Dirac Observables in Relativity*. Cambridge University Press, 6, 2019.
- [19] R. Cawley, *Determination of the Hamiltonian in the Presence of Constraints*, *Phys. Rev. Lett.* **42** (1979) 413.
- [20] Z. Li, *On the invalidity of a conjecture of dirac*, *Chinese Physics Letters* **10** (feb, 1993) 68–70.
- [21] Y.-L. Wang, Z.-P. Wang, and K. Wang, *Gauge Symmetries and the Dirac Conjecture*, *Int. J. Theor. Phys.* **48** (2009) 1894–1904.
- [22] A. Frenkel, *Comment on Cawley’s counterexample to a conjecture of Dirac*, *Phys. Rev. D* **21** (1980) 2986–2987.
- [23] R. Sugano and T. Kimura, *Comments on Constraints of Gauge Theories*, *J. Phys. A* **16** (1983) 4417.
- [24] B.-C. Wu, *Dirac’s conjecture*, *Int. J. Theor. Phys.* **33** (1994) 1529–1533.
- [25] O. Miskovic and J. Zanelli, *Dynamical structure of irregular constrained systems*, *J. Math. Phys.* **44** (2003) 3876–3887, [[hep-th/0302033](#)].
- [26] M. E. V. Costa, H. O. Girotti, and T. J. M. Simoes, *Dynamics of Gauge Systems and Dirac’s Conjecture*, *Phys. Rev. D* **32** (1985) 405.

- [27] A. Cabo and D. Louis-Martinez, *On Dirac's conjecture for Hamiltonian systems with first and second class constraints*, *Phys. Rev. D* **42** (1990) 2726–2735.
- [28] H. Rothe and K. Rothe, *Gauge identities and the Dirac conjecture*, *Annals Phys.* **313** (2004) 479–496, [[hep-th/0406044](#)].
- [29] J. Pons, *On Dirac's incomplete analysis of gauge transformations*, *Stud. Hist. Phil. Sci. B* **36** (2005) 491–518, [[physics/0409076](#)].