

# THE $K(\pi, 1)$ -CONJECTURE IMPLIES THE CENTER CONJECTURE FOR ARTIN GROUPS

KASIA JANKIEWICZ AND KEVIN SCHREVE

**ABSTRACT.** In this note, we prove that the  $K(\pi, 1)$ -conjecture for Artin groups implies the center conjecture for Artin groups. Specifically, every Artin group without a spherical factor that satisfies the  $K(\pi, 1)$ -conjecture has a trivial center.

## 1. INTRODUCTION

A Coxeter system  $(W, S)$  consists of a group  $W$  and a generating set  $S$  where  $W$  is given by a presentation

$$W = \langle s \in S \mid s^2 = (st)^{m_{st}} = 1 \rangle,$$

where  $m_{st} \in \{2, 3, \dots\} \cup \{\infty\}$ . The associated Artin group  $A$  is given by the presentation

$$A = \langle s \in S \mid \underbrace{sts \cdots}_{m_{st} \text{ terms}} = \underbrace{tst \cdots}_{m_{st} \text{ terms}} \rangle.$$

An Artin group  $A$  is *spherical* if the corresponding Coxeter group is finite, and otherwise  $A$  is *infinite type*. The Coxeter diagram  $\Gamma_S$  is a graph with vertices corresponding to  $S$  and where two vertices are joined by an edge if and only if  $m_{st} > 2$ . If  $m_{st} \geq 4$  we label the edge with  $m_{st}$ . A *special subgroup* of  $A$  is a subgroup generated by some subset of  $S$ . Each special subgroup is itself an Artin group [vdL83]. Each Artin group with standard generating set admits (a possibly trivial) decomposition  $A = A_{T_1} \times \cdots \times A_{T_n}$  where each  $T_i \subseteq S$  defines a connected component of the Coxeter graph  $\Gamma_S$ . An Artin group  $A$  is *irreducible* if its Coxeter diagram is connected. We say  $A_{T_i}$  is a *spherical factor* of  $A$  if  $A_{T_i}$  is spherical. Every irreducible spherical Artin group has an infinite cyclic center [Del72, BS72]. Conjecturally, those are the only irreducible Artin groups with nontrivial center.

**Conjecture 1** (The Center Conjecture). Every Artin group without a spherical factor has trivial center.

The center conjecture holds for FC-type Artin groups and 2-dimensional Artin groups [GP12]. Charney and Morris-Wright have shown the center conjecture holds for Artin groups whose defining graphs are not stars of a single vertex [CMW19]. Godelle and Paris further showed that if all Artin groups with  $m_{st} \neq \infty$  for all  $s, t \in S$  satisfy the center conjecture, then all Artin groups satisfy the center conjecture [GP12].

The FC-type and 2-dimensional Artin groups also satisfy the  $K(\pi, 1)$ -conjecture [CD95].

---

*Date:* January 19, 2022.

*2020 Mathematics Subject Classification.* 20F36, 20F65.

*Key words and phrases.* Artin groups, center, the  $K(\pi, 1)$ -conjecture.

The first author was supported by the NSF grant DMS-2203307.




FIGURE 1. The Coxeter diagram of the smallest mysterious Artin group according to McCammond.

**Conjecture 2** (The  $K(\pi, 1)$ -conjecture). The orbit space  $\mathcal{H}(W)/W$  of a complexified hyperplane arrangement associated to a Coxeter system  $(W, S)$  is a  $K(\pi, 1)$  for the Artin group  $A$  associated to  $W$ .

For a precise definition of  $\mathcal{H}(W)$  and more background, see e.g. the survey paper [Par14]. It is known that the fundamental group of  $\mathcal{H}(W)/W$  is equal to  $A$ , so the conjecture is about the asphericity of  $\mathcal{H}(W)/W$ . In this note, we prove the following:

**Theorem 3.** Every Artin group without a spherical factor that satisfies the  $K(\pi, 1)$ -conjecture has trivial center.

In fact, we only need the following consequence of the  $K(\pi, 1)$ -conjecture: an Artin group  $A$  which satisfies the  $K(\pi, 1)$ -conjecture has finite cohomological dimension which is realized by a spherical subgroup, i.e.  $\text{cd}(A) = \text{cd}(A_T) = |T|$  for some spherical subset  $T$ . See Theorem 12 for the more general statement of our main theorem.

For example, in [McC17] McCammond mentions that the center conjecture is unknown for the Artin group  $A$  with the Coxeter diagram as in Figure 1. The group  $A$  is given by the presentation

$$\langle a, b, c, d \mid aba = bab, bcb = bcb, cdc = dcd, dad = ada, bdb = dbd, ac = ca \rangle.$$

The  $K(\pi, 1)$ -conjecture holds for  $A$  by a theorem of Charney [Cha04]. That allows us to answer the question of McCammond.

**Corollary 4.** The Artin group with the Coxeter diagram as in Figure 1 has trivial center.

Another class of Artin groups which satisfy the  $K(\pi, 1)$ -conjecture are the *locally reducible* Artin groups, where all irreducible spherical subgroups are of rank  $\leq 2$  [Cha00]. There are many of these with  $m_{st} \neq \infty$  for each  $s, t \in S$ , and as far as we know the center conjecture was open here.

**Corollary 5.** Every locally reducible Artin group without spherical factor has trivial center.

## 2. REPRESENTATIONS OF ARTIN GROUPS IN MAPPING CLASS GROUPS

An Artin group  $A$  with standard generating set  $S$  has *small type* if  $m_{st} \in \{2, 3\}$  for all  $s, t \in S$ . In this section we recall a representation of small type Artin groups in mapping class groups, due to Crisp-Paris [CP01], and analyze where certain elements of  $A$  are mapped.

Let  $\Sigma$  be a surface. A *multicurve* is a collection of pairwise disjoint simple closed curves on  $\Sigma$ . We say that two multicurves are *disjoint* if their isotopy classes have disjoint representatives. A *multitwist* about a multicurve  $\gamma$  is a product of non-trivial powers of Dehn




FIGURE 2. An example of the surface  $\Sigma_S$  (right) corresponding to the Artin group  $A$  with the standard generating set  $S$  whose Coxeter graph  $\Gamma$  is illustrated (left). For each vertex  $s$  in  $\Gamma$ , there is a corresponding rectangle in the right picture, which is glued along the dotted sides to an annulus  $An_s$ . Its meridian  $\gamma_s$  has the same color as  $s$ .

twists about simple closed curves in  $\gamma$  (the powers can be different for different curves). We shall need the following lemma about commuting multitwists about multicurves.

**Lemma 6.** Let  $\gamma$  and  $\gamma'$  be essential multicurves on  $\Sigma$  that do not share a simple closed curve. Let  $T_\gamma$  and  $T_{\gamma'}$  be the associated multitwists along  $\gamma$  and  $\gamma'$ . Then  $T_\gamma$  and  $T_{\gamma'}$  commute if and only if  $\gamma$  and  $\gamma'$  are disjoint.

*Proof.* The if direction is clear, so suppose  $T_\gamma$  and  $T_{\gamma'}$  commute. Then  $T_\gamma^N$  and  $T_{\gamma'}^N$  commute for all  $N$ . By a theorem of Koberda [Kob12], the group generated by large powers of Dehn twists of all curves in  $\gamma \cup \gamma'$  is a right-angled Artin group. Therefore, we have two words  $w$  and  $w'$  in a RAAG, where  $w$  and  $w'$  are nontrivial powers of commuting generators. It follows from the normal form for RAAG's that these commute exactly when each generator in  $w$  commutes with each generator of  $w'$  [HM95]. Therefore, the curves in  $\gamma$  can be isotoped to be disjoint from the curves in  $\gamma'$ .  $\square$

Let  $A_S$  be a small type Artin group with standard generating set  $S$ . We can build an associated surface

$$\Sigma_S = \bigcup_{s \in S} An_s$$

where each  $An_s$  is an annulus. We denote the meridian of  $An_s$  by  $\gamma_s$ . If  $m_{st} = 3$ , then we arrange that the annuli  $An_s, An_t$  intersect in a single square so that  $\gamma_s, \gamma_t$  intersect transversely at one point, and any triple intersection of annuli is empty. If  $m_{st} = 2$ , then  $An_s, An_t$  will be disjoint. See Figure 2 for an example. There is a representation of  $A_S$  in the mapping class group of  $\Sigma_S$  where each generator  $s \in S$  is mapped to the Dehn twist about the simple closed curve  $\gamma_s$  (see Proposition 9). Full details can be found in [CP01]. Each subset  $T$  of  $S$  has an associated subsurface  $\Sigma_T$  of  $\Sigma_A$ . By construction, the subsurface  $\Sigma_T$  for an irreducible spherical subset  $T \subseteq S$  and the induced homomorphism  $\rho_T : A_T \rightarrow \text{Mod}(\Sigma_T)$  are exactly the Perron-Vannier representation of small type spherical Artin groups [PV96] (see also [JS20]). For every spherical subset  $T$ , the generator of the center of the group  $A_T$  is denoted by  $z_T$ . By [LP01, Prop 2.12]  $\rho(z_T^4)$  is a multitwist about the boundary of  $\Sigma_T$ . We denote that multicurve by  $\Gamma_T$ .

Let  $K_S$  be the graph that is the union  $\bigcup_{s \in S} \gamma_s \subseteq \Sigma_S$ . By construction, there is a deformation retraction  $r : \Sigma_S \rightarrow K_S$ . Thus  $H_1(\Sigma_S) = H_1(K_S)$ , and in particular if  $S$  is spherical then  $H_1(\Sigma_S) = \mathbb{Z}^S$ . For any  $S' \subset S$ , the map  $H_1(\Sigma_{S'}) \rightarrow H_1(\Sigma_S)$  induced by the inclusion  $\Sigma_{S'} \hookrightarrow \Sigma_S$  is injective and  $H_1(\Sigma_{S'}) \subsetneq H_1(\Sigma_S)$ .

Every closed path in a graph is homotopic to a cycle (i.e. a closed path without backtracks). In particular, every homotopy class of a simple closed curve in  $\Sigma_S$  can be realized as a cycle in  $K_S$ . We will now view all the simple closed curves in  $\Sigma_S$  as cycles in  $K_S$ . In particular, we view components of  $\gamma_T$  for any spherical subset  $T \subseteq S$  as cycles in  $K_S$ .

**Lemma 7.** Let  $T \subseteq S$  be an irreducible spherical subset, and  $s \in S - T$ . Then  $\gamma_s$  intersects  $\gamma_T$  if and only if  $\gamma_s$  intersects  $\gamma_t$  for some  $t \in T$ .

*Proof.* If  $\gamma_s$  does not intersect any  $\gamma_t$  for  $t \in T$ , then  $\rho(s)$  commutes with  $\rho(t)$  for all  $t \in T$ . Then  $\rho(s)$  must also commute with  $\rho(z_T^2)$ , and by Lemma 6  $\gamma_s$  and  $\gamma_T$  are disjoint.

Now suppose that  $\gamma_s$  intersects  $\gamma_t$  for some  $t \in T$ . By construction  $\gamma_s, \gamma_t$  intersect exactly once. Suppose  $\gamma_s$  can be isotoped in  $\Sigma_S$  to be disjoint from  $\partial \Sigma_T$ . Then  $\gamma_s \subseteq \Sigma_T$ , since  $\gamma_s, \gamma_t$  still must intersect. In particular,  $[\gamma_s] \in H_1(\Sigma_T) = H_1(K_T)$ . This is a contradiction, since  $H_1(K_T) \subsetneq H_1(K_{T \cup \{s\}})$ .  $\square$

**Lemma 8.** Let  $T_1, T_2 \subseteq S$  be two disjoint, irreducible, spherical subsets. Then any component of  $\gamma_{T_1}$  and any component of  $\gamma_{T_2}$  are non-isotopic and disjoint in  $\Sigma_S$ .

*Proof.* Since  $T_1, T_2 \subseteq S$  are disjoint, by construction, the subgraphs  $K_{T_1}, K_{T_2} \subseteq K_S$  are disjoint. Every connected component of  $\gamma_{T_i}$  can be realized as a cycle contained in  $K_{T_i}$ . Any two disjoint cycles in a graph are non-isotopic. The conclusion follows.  $\square$

**Proposition 9** ([CP01]). For every small type Artin group  $A$  with the standard generating set  $S$ , there exists a surface with boundary  $\Sigma_S$  and a homomorphism  $\rho : A \rightarrow \text{Mod}(\Sigma_S)$  where

- (a) for each  $s \in S$ ,  $\rho(s)$  is the Dehn twist about a simple closed curve  $\gamma_s$ ,
- (b) the simple closed curves  $\gamma_s, \gamma_t$  are disjoint  $\iff m_{st} = 2$ ,
- (c) the simple closed curves  $\gamma_s, \gamma_t$  intersect exactly once  $\iff m_{st} = 3$ .

Moreover,

- (d) for every irreducible spherical subset  $T \subseteq S$ ,  $\rho(z_T^2)$  is the multitwist about a multicurve  $\gamma_T$  which is the boundary of the subsurface  $\Sigma_T$ , and
- (e) for every irreducible spherical  $T \subseteq S$  and  $s \in S - T$ , the simple closed curve  $\gamma_s$  and the multicurve  $\Gamma_T$  are disjoint if and only if  $[s, t] = 1$  for all  $t \in T$ .

*Proof.* The fact that  $\rho$  is a homomorphism follows from standard relations between Dehn twists, see [CP01, Prop 4]. The parts (a), (b), (c) follow from [CP01] as well. Part (d) follows from [LP01] (see discussion above). Finally part (e) is a consequence of Lemma 7 and Lemma 6.  $\square$

Let  $A$  be an Artin group with standard generating set  $S$ . We say  $A$  is *free-of-infinity* if  $m_{st} < \infty$  for all  $s, t \in S$ .

**Proposition 10** ([CP01]). Let  $A$  be a free-of-infinity Artin group. Then there exists a small type Artin group  $\tilde{A}$  with standard generating set  $\tilde{S}$  and a homomorphism  $\phi : A \rightarrow \tilde{A}$  such that

- there exists a partition  $\bigsqcup_{s \in S} I(s)$  of  $\tilde{S}$  such that the elements of  $I(s)$  pairwise commute

$$\text{and } \phi(s) = \prod_{r \in I(s)} r,$$

- $m_{st} = 2$  if and only if every element of  $I(s)$  and every element of  $I(t)$  commute, and
- if  $m_{st} \geq 3$  then the subgroup generated by  $I(s) \cup I(t)$  is a direct product of braid groups on  $m_{st}$  strands.

Let  $\rho \circ \phi : A \rightarrow \text{Mod}(\Sigma_{\tilde{A}})$  be the composition of the homomorphism  $\phi$  with the homomorphism  $\rho : \tilde{A} \rightarrow \text{Mod}(\Sigma_{\tilde{A}})$  from Proposition 9. Then

- (a) for each  $s \in S$ ,  $\rho \circ \phi(s)$  is a multitwist about a multicurve  $\gamma_s = \bigcup_{r \in I(s)} \gamma_r$ ,
- (b)  $m_{st} = 2$  if and only if every component of  $\gamma_s$  and every component of  $\gamma_t$  are disjoint,
- (c) for every spherical subset  $T \subset S$ ,  $\rho(z_T^2)$  is the multitwists about a multicurve  $\gamma_T$ , and
- (d) for every  $T \subseteq S$  and  $s \in S - T$ , the multicurve  $\gamma_s$  and the multicurve  $\gamma_T$  are disjoint if and only if  $[s, t] = 1$  for all  $t \in T$ .

*Proof.* The homomorphism  $\phi$  is described in [Cri99] and also in [CP01]. Parts (a) and (b) follow directly from the construction. Part (c) is proven in [JS20, Lem 6.1]. Part (d) follows from Lemma 7 and Lemma 6.  $\square$

### 3. THE MAIN THEOREM

We will need the following lemma.

**Lemma 11.** Let  $A_S$  be an Artin group which splits as a product  $A_S = A_U \times A_V$  where  $A_U$  is the maximal spherical factor. Suppose that  $\text{cd } A_S < \infty$ . Then  $\text{cd } A_S = \text{cd } A_U + \text{cd } A_V$ .

*Proof.* By [Bie81, Thm 5.5] a group  $G = N \times Q$  has  $\text{cd } G = \text{cd } N + \text{cd } Q$  provided that

- $\text{cd } Q < \infty$ , and
- $N$  is of type FP and  $H^n(N, \mathbb{Z}N)$  is free for  $n = \text{cd } N$ .

Clearly  $\text{cd } A_V < \infty$  since  $\text{cd } A_S < \infty$ . Since  $A_U$  is a spherical Artin group,  $A_U$  has type FP. By [Squ94, Thm B] (see also [Bes99])  $A_U$  is a duality group, so  $H^n(A_U, \mathbb{Z}A_U)$  is free. The conclusion follows.  $\square$

**Theorem 12.** Let  $A_S$  be an Artin group of infinite type with the standard generating set  $S$  such that  $A_S$  has no spherical factors. If  $\text{cd } A = \text{cd } A_T = |T|$  for some spherical subset  $T \subseteq S$ , then  $A_S$  has trivial center. In particular if  $A_S$  satisfies the  $K(\pi, 1)$ -conjecture, then  $A_S$  has trivial center.

*Proof for free-of-infinity case.* First suppose that  $A_S$  is free-of-infinity. Let  $T \subseteq S$  be a maximal spherical subset such that  $\text{cd } A_S = \text{cd } A_T$ . Let  $T_1 \sqcup T_2 \sqcup \cdots \sqcup T_n$  be the decomposition of  $T$  into irreducible spherical subsets inducing the decomposition  $A_T = A_{T_1} \times \cdots \times A_{T_n}$ . Since  $A_S$  has no spherical factors for each  $i = 1, \dots, n$  there exists  $s_i \in S - T$  such that  $[s_i, z_{T_i}] \neq 1$  as otherwise  $A_{T_i}$  would be a spherical factor of  $A_S$ . In particular, for each  $i = 1, \dots, n$ , there exists  $t_i \in T_i$  such that  $[s_i, t_i] \neq 1$ .

Consider the representation of  $\rho : A_S \rightarrow \text{Mod}(\Sigma_S)$  from Proposition 10. By Proposition 10,  $\rho(s_i)$  and  $\rho(z_{T_i})$  are the Dehn twists about multicurves  $\gamma_{s_i}$  and  $\gamma_{T_i}$  respectively, where  $\gamma_{s_i}$  and  $\gamma_{T_i}$  intersect.

Suppose that  $A_S$  has nontrivial center and let  $y \in Z(A_S)$  with  $y \neq e$ . Note that  $y$  has infinite order since  $A_S$  is torsion-free, as  $\text{cd } A_S < \infty$ . If  $y^k \notin A_T$  for any  $k \neq 0$ , then  $\langle A_T, y \rangle \simeq A_T \times \mathbb{Z}$  is a subgroup of  $\text{cd } A + 1$ , which is a contradiction. Thus there exists  $k \in \mathbb{N}$  such that  $y^k \in A_T$ . Then  $y^k \in Z(A_T)$ , i.e.  $y^m = \prod_{i=1}^n z_{T_i}^{m_i}$  for some  $m > 0$  and at least one of  $m_1, \dots, m_n$ , say  $m_1$ , is non-zero. By Lemma 8,  $\rho(y^m)$  is a multitwist about a multicurve  $\gamma = \sqcup \gamma_{T_i}$  in  $\Sigma$  where the union is taken over all  $i$  such that  $m_i \neq 0$ . In particular,  $\gamma$  intersects  $\gamma_{s_1}$ . By Lemma 6,  $[\rho(y^m), \rho(s_1)] \neq 1$ . Thus  $[y, s_1] \neq 1$ . This contradicts the fact that  $y$  is a central element of  $A$ .  $\square$

*Proof for general case.* The general case is induction on the cardinality of  $S$ . Suppose  $\text{cd } A_S = \text{cd } A_T$  where  $T \subseteq S$  is a spherical subset. Suppose there exist generators  $v, w \in S$  such that  $m_{vw} = \infty$ . The group  $A_S$  splits as an amalgamated product  $A_{S \setminus \{v\}} *_{A_{S \setminus \{v, w\}}} A_{S \setminus \{w\}}$ . Since  $T$  cannot contain both  $v$  and  $w$ , we have  $T \subseteq S \setminus \{v\}$  or  $T \subseteq S \setminus \{w\}$ . Without loss of generality we assume that  $T \subseteq S \setminus \{v\}$ . It follows that  $\text{cd } A_{S \setminus \{v\}} = \text{cd } A_T$ , as  $\text{cd } A_{S \setminus \{v\}} \leq \text{cd } A_S$ . If  $A_{S \setminus \{v\}}$  has no spherical factor, then by induction  $A_{S \setminus \{v\}}$  has trivial center. By [GP12, Lem 3.2] the center of the amalgamated product  $A$  is also trivial.

Now suppose that  $A_{S \setminus \{v\}}$  has a nontrivial spherical factor. Let

$$A_{U_1} \times \cdots \times A_{U_p} \times A_{V_1} \times \cdots \times A_{V_q}$$

be the decomposition of  $A_{S \setminus \{v\}}$  into irreducible factors where each  $A_{U_i}$  is spherical and each  $A_{V_j}$  has infinite type. Let  $A_V = A_{V_1} \times \cdots \times A_{V_q}$ . By maximality  $U_i \subseteq T$  for all  $i = 1, \dots, p$ . Let  $T' = V \cap T$ . Then  $\text{cd } A_V = \text{cd } A_{T'}$ . Indeed by Lemma 11 ,

$$\text{cd } A_V = \text{cd } A_{S \setminus \{v\}} - \sum_{i=1}^p \text{cd } A_{U_i} = \text{cd } A_T - \sum_{i=1}^p \text{cd } A_{U_i} = A_{T'}.$$

By the inductive assumption  $Z(A_V) = \{1\}$ , and thus  $Z(A_{S \setminus \{v\}}) \subseteq \langle z_{U_1} \rangle \times \cdots \times \langle z_{U_n} \rangle$ .

Since  $A_S$  does not have a spherical factor, for every  $i = 1, \dots, n$  we have  $[v, z_{U_i}] \neq 1$ . In particular, each set  $U_i$  contains a standard generator  $u_i$  such that  $m_{vu_i} \geq 3$ . Since  $Z(A_{S \setminus \{v\}}) \subseteq A_T$  and by [GP12, Lem 3.2]  $Z(A) \subseteq Z(A_{S \setminus \{v\}})$ , it suffices to prove that  $v$  does not commute with any nontrivial element of  $Z(A_T) = \langle z_{T_1}, \dots, z_{T_n} \rangle$ . By maximality of  $T$ ,  $A_{T \cup \{v\}}$  is not spherical. By the discussion above,  $A_{T \cup \{v\}}$  is irreducible, and in particular it has no spherical factors. If  $A_{T \cup \{v\}}$  is free-of-infinity, we are done.

We now assume that  $A_{T \cup \{v\}}$  is not free-of-infinity. Consider the quotient homomorphism  $\phi : A_{T \cup \{v\}} \rightarrow A_{\overline{T \cup \{v\}}}$ , where for every  $t \in T$  such that  $m_{tv} = \infty$  the corresponding generators  $\bar{t}, \bar{v} \in \overline{T \cup \{v\}}$  have  $m_{\bar{t}\bar{v}} = 7$ . The group  $A_{\overline{T \cup \{v\}}}$  is irreducible. The only irreducible spherical Artin group containing label 7 is the dihedral Artin group. If  $A_{\overline{T \cup \{v\}}}$  is the dihedral Artin group, then  $A_{T \cup \{v\}} = F_2$  and so  $\text{cd } A_S = 1$ , i.e.  $A_S = F(S)$ . Then clearly,  $A_S$  has trivial center. Otherwise  $A_{\overline{T \cup \{v\}}}$  is irreducible and has infinite type. Also  $\text{cd } A_{\overline{T \cup \{v\}}} = \text{cd } A_{\overline{T}}$ . By the free-of-infinity case,  $[\bar{v}, \bar{y}] \neq 1$  for any nontrivial  $\bar{y} \in \langle z_{\overline{T_1}}, \dots, z_{\overline{T_n}} \rangle$ , as otherwise  $y$  would be a central element of  $A_{\overline{T \cup \{v\}}}$ . Thus  $[v, y] \neq 1$  for any nontrivial  $y \in \langle z_{T_1}, \dots, z_{T_n} \rangle$ . This completes the proof.  $\square$

## REFERENCES

- [Bes99] Mladen Bestvina, *Non-positively curved aspects of Artin groups of finite type*, Geom. Topol. **3** (1999), 269–302. MR1714913
- [Bie81] Robert Bieri, *Homological dimension of discrete groups*, Second, Queen Mary College Department of Pure Mathematics, London, 1981.

- [BS72] Egbert Brieskorn and Kyoji Saito, *Artin-Gruppen und Coxeter-Gruppen*, Invent. Math. **17** (1972), 245–271. MR323910
- [CD95] Ruth Charney and Michael W. Davis, *The  $K(\pi, 1)$ -problem for hyperplane complements associated to infinite reflection groups*, J. Amer. Math. Soc. **8** (1995), no. 3, 597–627. MR1303028
- [Cha00] Ruth Charney, *The Tits conjecture for locally reducible Artin groups*, Internat. J. Algebra Comput. **10** (2000), no. 6, 783–797. MR1809385
- [Cha04] ———, *The Deligne complex for the four-strand braid group*, Trans. Amer. Math. Soc. **356** (2004), no. 10, 3881–3897. MR2058510
- [CMW19] Ruth Charney and Rose Morris-Wright, *Artin groups of infinite type: trivial centers and acylindrical hyperbolicity*, Proc. Amer. Math. Soc. **147** (2019), no. 9, 3675–3689. MR3993762
- [CP01] John Crisp and Luis Paris, *The solution to a conjecture of Tits on the subgroup generated by the squares of the generators of an Artin group*, Invent. Math. **145** (2001), no. 1, 19–36. MR1839284
- [Cri99] John Crisp, *Injective maps between Artin groups*, Geometric group theory down under (Canberra, 1996), 1999, pp. 119–137. MR1714842
- [Del72] Pierre Deligne, *Les immeubles des groupes de tresses généralisés*, Invent. Math. **17** (1972), 273–302. MR0422673 (54 #10659)
- [GP12] Eddy Godelle and Luis Paris, *Basic questions on Artin-Tits groups*, Configuration spaces, 2012, pp. 299–311. MR3203644
- [HM95] Susan Hermiller and John Meier, *Algorithms and geometry for graph products of groups*, J. Algebra **171** (1995), no. 1, 230–257.
- [JS20] Kasia Jankiewicz and Kevin Schreve, *Right-angled Artin subgroups of Artin groups*, 2020. J. Lond. Math. Soc., to appear.
- [Kob12] Thomas Koberda, *Right-angled Artin groups and a generalized isomorphism problem for finitely generated subgroups of mapping class groups*, Geom. Funct. Anal. **22** (2012), no. 6, 1541–1590. MR3000498
- [LP01] Catherine Labruère and Luis Paris, *Presentations for the punctured mapping class groups in terms of Artin groups*, Algebr. Geom. Topol. **1** (2001), 73–114. MR1805936
- [McC17] Jon McCammond, *The mysterious geometry of Artin groups*, Winter Braids Lect. Notes **4** (2017), no. Winter Braids VII (Caen, 2017), Exp. No. 1, 30. MR3922033
- [Par14] Luis Paris,  *$K(\pi, 1)$  conjecture for Artin groups*, Ann. Fac. Sci. Toulouse Math. (6) **23** (2014), no. 2, 361–415. MR3205598
- [PV96] B. Perron and J. P. Vannier, *Groupe de monodromie géométrique des singularités simples*, Math. Ann. **306** (1996), no. 2, 231–245. MR1411346
- [Squ94] Craig C. Squier, *The homological algebra of Artin groups*, Math. Scand. **75** (1994), no. 1, 5–43. MR1308935
- [vdL83] Harm van der Lek, *Extended Artin groups*, Singularities, Part 2 (Arcata, Calif., 1981), 1983, pp. 117–121. MR713240

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, SANTA CRUZ, CA 95064  
*Email address:* kasia@ucsc.edu

DEPARTMENT OF MATHEMATICS, LOUISIANA STATE UNIVERSITY, BATON ROUGE, LA 70806  
*Email address:* kschreve@lsu.edu