

High dimensional AdS black hole and Phase transition in Einstein-bumblebee gravity

Chikun Ding^{1,2,3,*}, Yu Shi¹, Jun Chen¹, Yuebing Zhou¹, and Changqing Liu²

¹*Department of Physics, Huaihua University, Huaihua, 418008, P. R. China*

²*Department of Physics, Hunan University of Humanities, Science and Technology, Loudi, Hunan 417000, P. R. China*

³*Key Laboratory of Low Dimensional Quantum Structures and Quantum Control of Ministry of Education, and Synergetic Innovation Center for Quantum Effects and Applications, Hunan Normal University, Changsha, Hunan 410081, P. R. China*

Abstract

In this paper we obtain an exact high dimensional anti-de Sitter (AdS) black hole solution in Einstein-bumblebee gravity theory. This AdS black hole can only exist with a linear functional potential of the bumblebee field. We find that the Smarr formula and the first law of black hole thermodynamics can still be constructed in this Lorentz symmetry breaking black hole spacetime as long as its temperature, entropy and volume are slightly modified. We find also that there exist two kinds of phase transition: small-large black hole phase transition and Hawking-Page phase transition, like those of Schwarzschild AdS black hole. After Lorentz symmetry breaking, the black hole mass at divergent point of heat capacity becomes small, and the Gibbs free energy of the meta-stable large black hole is also smaller, showing that the large stable black hole can be more easily formed.

PACS numbers: 04.50.Kd, 04.20.Jb, 04.70.Dy

I. INTRODUCTION

Local Lorentz invariance (LI) is the most fundamental spacetime symmetry of the special/general relativity (SR/GR) and the standard model (SM) of particle physics [1]. LI means that the equations of motion for particles and fields are invariant under all Lorentz transformations(global for SR, local for GR) and diffeomorphism transformation for a particle. However, LI should not be an exact principle at very high energy scales [2]. GR and SM are the best theories describing the four fundamental forces and there are no known experimental conflicts. Nonetheless, they are fundamentally different in that SM is a quantum theory disregarding the gravitational effects of all particles, while GR is a classical geometrical theory ignoring all the quantum features of particles. For particles with energies on the order of 10^{30} eV (above the Planck scale),

*Corresponding author; Electronic address: dingchikun@163.com

the gravitational interactions estimated by GR are very powerful, and so one shouldn't ignore it [3]. Thus, on the very high energy scales, one has to reconsider combining the SM with GR in a unified theory, i.e., "quantum gravity". An effective field theory combining GR and SM on low energy scales is called standard model extension (SME), which couples the SM to GR, and involves extra items embracing information about the violations of Lorentz invariance(Lorentz violation, LV) happening on the Planck scale [4].

Studying the Lorentz violation (LV) is a useful approach toward investigating the foundations of modern physics. Besides SME, string theory [5], noncommutative field theories [6], spacetime-varying fields [7], loop quantum gravity theory [8], brane world scenarios [9], massive gravity [10] and Einstein-aether theory [11] are other proposals of Lorentz violation. The SME is experimentally accessible for studying possible observable signals of violation the particle Lorentz symmetry. In this model, the local Lorentz violating is provided by a spontaneous symmetry breaking potential due to self-interacting tensor fields, whose vacuum expectation value (VEV) yields to background tensor fields. A specific theory is the bumblebee field B^μ , an self-interacting tensor field has a nonzero VEV b^μ which defines a privileged direction in spacetime and spontaneously breaks the local Lorentz symmetry.

There are many forms of the bumblebee potential. One is the smooth functional bumblebee potential is $V(X) = kX^2/2$ which has a minimum when the bumblebee field equals its VEV, i.e., $V = 0$ and $V'(X) = kX = 0$ if $X = 0$, where k is a constant [12] and $X = B^\mu B_\mu \pm b^2$. With this potential, the static black hole solutions were found by Bertolami *et al.* [13] and by Casana *et al.* [14]; the rotating black hole solutions were built by Ding *et al.* [15] and by Jha *et al.* [16]. An other form of bumblebee potential is a linear multiplier function $V(X) = \lambda X/2$, where λ is a Lagrange-multiplier. With this potential, the Schwarzschild-anti-de Sitter-like and Schwarzschild-de Sitter-like black hole solutions were found by Maluf *et al.* [17].

The quantum field theory in a spacetime with a horizon exhibits thermal behavior which must be thought of as an emergent phenomena [18]. A black hole event horizon has the conceptions of temperature and entropy [19], and the first law of thermodynamics $dU = TdS - PdV$ can be built, where P is the radial pressure of the matter, U is the internal energy and V is the volume of the black hole. There also has transfer of heat energy and the conception of phase transition [20]. Then the gravity system can be mapped to a thermodynamic system and studying on black hole thermodynamics can help to construct a quantum theory of gravity.

We will study the high dimensional black hole solutions and its thermodynamics and phase transition with a negative cosmological constant in the theory of Einstein- bumblebee gravity. The rest paper is organized as follows. In Sec. II we give the background for the Einstein-Gauss-Bonnet-bumblebee theory and derive the

black hole solution by solving the gravitational field equations. In Sec. III, we study its thermodynamical properties. In Sec. IV, we study the phase transitions and find some effects of the Lorentz breaking constant ℓ . Sec. V is for a summary.

II. HIGH DIMENSIONAL ADS BLACK HOLE IN EINSTEIN-BUMBLEBEE GRAVITY

In the bumblebee gravity model, one introduces the bumblebee vector field B_μ which has a nonzero vacuum expectation value, to lead a spontaneous Lorentz symmetry breaking in the gravitational sector via a given potential. In the higher $D \geq 4$ dimensional spacetime, the action of Einstein-bumblebee gravity is [21],

$$\mathcal{S} = \int d^D x \sqrt{-g} \left[\frac{R - 2\Lambda}{2\kappa} + \frac{\varrho}{2\kappa} B^\mu B^\nu R_{\mu\nu} - \frac{1}{4} B^{\mu\nu} B_{\mu\nu} - V(B_\mu B^\mu \mp b^2) + \mathcal{L}_M \right], \quad (2.1)$$

where R is Ricci scalar, Λ is the negative cosmological constant. $\kappa = 8\pi G_D/c^4$, where G_D is the D -dimensional gravitational constant and has the relation to Newton's constant G as $G_D = G\Omega_{D-2}/4\pi$ [22], $\Omega_{D-2} = 2\sqrt{\pi}^{D-1}/\Gamma[(D-1)/2]$ is the area of a unit $(D-2)$ -sphere. Here and hereafter, we take $G_D = 1$ and $c = 1$ for convenience.

The coupling constant ϱ dominates the non-minimal gravity interaction to bumblebee field B_μ . The term \mathcal{L}_M represents possible interactions with matter or external currents. The constant b is a real positive constant. The potential $V(B_\mu B^\mu \mp b^2)$ triggers Lorentz and/or CPT (charge, parity and time) violation. It gives a nonzero vacuum expectation value (VEV) for bumblebee field B_μ indicating that the vacuum of this model obtains a prior direction in the spacetime. This potential has a minimum at $B^\mu B_\mu \pm b^2 = 0$ and $V'(b_\mu b^\mu) = 0$ to ensure the destroying the $U(1)$ symmetry, where the field B_μ acquires a nonzero VEV, $\langle B^\mu \rangle = b^\mu$. Another vector b^μ is a function of the spacetime coordinates and has a constant value $b_\mu b^\mu = \mp b^2$, where \pm signs mean that b^μ is timelike or spacelike, respectively. The bumblebee field strength is

$$B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu. \quad (2.2)$$

This antisymmetry of $B_{\mu\nu}$ implies the constraint [12]

$$\nabla^\mu \nabla^\nu B_{\mu\nu} = 0. \quad (2.3)$$

Varying the action (2.1) with respect to the metric yields the gravitational field equations

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu}^B + \kappa T_{\mu\nu}^M, \quad (2.4)$$

where $G_{\mu\nu} = R_{\mu\nu} - g_{\mu\nu}R/2$, and the bumblebee energy momentum tensor $T_{\mu\nu}^B$ is

$$\begin{aligned} T_{\mu\nu}^B &= B_{\mu\alpha}B_{\nu}^{\alpha} - \frac{1}{4}g_{\mu\nu}B^{\alpha\beta}B_{\alpha\beta} - g_{\mu\nu}V + 2B_{\mu}B_{\nu}V' \\ &+ \frac{\varrho}{\kappa} \left[\frac{1}{2}g_{\mu\nu}B^{\alpha}B^{\beta}R_{\alpha\beta} - B_{\mu}B^{\alpha}R_{\alpha\nu} - B_{\nu}B^{\alpha}R_{\alpha\mu} \right. \\ &\left. + \frac{1}{2}\nabla_{\alpha}\nabla_{\mu}(B^{\alpha}B_{\nu}) + \frac{1}{2}\nabla_{\alpha}\nabla_{\nu}(B^{\alpha}B_{\mu}) - \frac{1}{2}\nabla^2(B^{\mu}B_{\nu}) - \frac{1}{2}g_{\mu\nu}\nabla_{\alpha}\nabla_{\beta}(B^{\alpha}B^{\beta}) \right]. \end{aligned} \quad (2.5)$$

The prime denotes differentiation with respect to the argument,

$$V' = \frac{\partial V(x)}{\partial x} \Big|_{x=B^{\mu}B_{\mu}\pm b^2}. \quad (2.6)$$

Varying instead with respect to the the bumblebee field generates the bumblebee equations of motion (supposing that there is no coupling between the bumblebee field and \mathcal{L}_M),

$$\nabla^{\mu}B_{\mu\nu} = 2V'B_{\nu} - \frac{\varrho}{\kappa}B^{\mu}R_{\mu\nu}. \quad (2.7)$$

The contracted Bianchi identities ($\nabla^{\mu}G_{\mu\nu} = 0$) lead to conservation of the total energy-momentum tensor

$$\nabla^{\mu}T_{\mu\nu} = \nabla^{\mu}(T_{\mu\nu}^B + T_{\mu\nu}^M - \Lambda g_{\mu\nu}) = 0. \quad (2.8)$$

We suppose that there is no matter field and the bumblebee field is frozen at its VEV like in Refs [13, 14], i.e., it is

$$B_{\mu} = b_{\mu}. \quad (2.9)$$

In order to generate a black hole solution in the case of non-zero cosmological constant Λ , we should apply a linear function form of the potential [17]

$$V = \frac{\lambda}{2}(B_{\mu}B^{\mu} - b^2), \quad (2.10)$$

where λ is a non-zero constant and deserves as a Lagrange-multiplier field. This potential is $V = 0$ under the condition (2.9) and its derivative is $V' = \lambda/2$ which can modify the Einstein equations. However, this additional degree of freedom of the λ field is auxiliary due to that it is restricted by the equation (2.25). Then the first two terms in Eq. (2.5) are like those of the electromagnetic field, the only distinctiveness are the coupling items to Ricci tensor and this λ term. Under this condition, Eq. (2.4) leads to gravitational field equations [21]

$$G_{\mu\nu} = \kappa(\lambda b_{\mu}b_{\nu} + b_{\mu\alpha}b_{\nu}^{\alpha} - \frac{1}{4}g_{\mu\nu}b^{\alpha\beta}b_{\alpha\beta}) + \varrho \left(\frac{1}{2}g_{\mu\nu}b^{\alpha}b^{\beta}R_{\alpha\beta} - b_{\mu}b^{\alpha}R_{\alpha\nu} - b_{\nu}b^{\alpha}R_{\alpha\mu} \right) + \bar{B}_{\mu\nu}, \quad (2.11)$$

with

$$\bar{B}_{\mu\nu} = \frac{\rho}{2} \left[\nabla_\alpha \nabla_\mu (b^\alpha b_\nu) + \nabla_\alpha \nabla_\nu (b^\alpha b_\mu) - \nabla^2 (b_\mu b_\nu) - g_{\mu\nu} \nabla_\alpha \nabla_\beta (b^\alpha b^\beta) \right]. \quad (2.12)$$

The static spherically symmetric black hole metric in the D dimensional spacetime has the form

$$ds^2 = -e^{2\phi(r)} dt^2 + e^{2\psi(r)} dr^2 + r^2 d\Omega_{D-2}^2, \quad (2.13)$$

where Ω_{D-2} is a standard $D - 2$ sphere. In the present study, we pay attention to that the bumblebee field has a radial vacuum energy expectation because that the spacetime curvature has a strong radial variation, on the contrary that the temporal changes are very slow. So the bumblebee field is supposed to be spacelike as that

$$b_\mu = (0, be^{\psi(r)}, 0, 0, \dots, 0), \quad (2.14)$$

where b is a positive constant. Then the bumblebee field strength is

$$b_{\mu\nu} = \partial_\mu b_\nu - \partial_\nu b_\mu, \quad (2.15)$$

whose components are all zero. And their divergences are all zero, i.e.,

$$\nabla^\mu b_{\mu\nu} = 0. \quad (2.16)$$

From the bumblebee field motion equation (2.7), we have the projection of the Ricci tensor along the bumblebee field is

$$b^\mu R_{\mu\nu} = \frac{\kappa\lambda}{\rho} b_\nu. \quad (2.17)$$

As to gravitational field equation (2.11), one can obtain the following three component equations

$$(D - 2)(1 + \ell)[2r\psi' - (D - 3)] + e^{2\psi} [(D - 2)(D - 3) - 2\Lambda r^2] = 0, \quad (2.18)$$

$$2\ell r^2(\phi'' + \phi'^2 - \phi'\psi') - 2\ell(D - 2)r(\psi' + \phi') - 2(D - 2)r\phi' + e^{2\psi} [(D - 2)(D - 3) + 2\kappa\lambda b^2 r^2 - 2\Lambda r^2] - (1 + \ell)(D - 2)(D - 3) = 0, \quad (2.19)$$

$$(1 + \ell) \left[r^2(\phi'' + \phi'^2 - \phi'\psi') + \frac{(D - 3)(D - 4)}{2} + (D - 3)r(\phi' - \psi') \right] + e^{-2\psi} \left[\Lambda r^2 - \frac{(D - 3)(D - 4)}{2} \right] = 0, \quad (2.20)$$

where we have redefined the Lorentz-violating parameter $\ell = \rho b^2$ and, the prime ' is the derivative with respect to the corresponding argument, respectively. From the Eq. (2.18), one can obtain a metric function as

$$e^{2\psi} = \frac{1 + \ell}{f(r)}, \quad (2.21)$$

where

$$f(r) = 1 - \frac{16\pi M}{(D-2)\Omega_{D-2}r^{D-3}} - \frac{2\Lambda}{(D-1)(D-2)}r^2, \quad (2.22)$$

and M is the mass of the black hole. In order to recover the 4-dimensional Schwarzschild-like solution [14] in the cases of $\Lambda = 0$, then we let another function to

$$e^{2\phi} = f(r). \quad (2.23)$$

It is easy to prove that the above results of $e^{2\phi} = f(r)$, $e^{2\psi} = (1 + \ell)/f(r)$ can meet the gravitational Eq. (2.20). Then the bumblebee field is

$$b_\mu = (0, b\sqrt{(1 + \ell)/f(r)}, 0, 0, \dots, 0). \quad (2.24)$$

With the bumblebee field motion equation (2.17) or the gravitational Eq. (2.19), a solution of this type will be possible if and only if

$$\Lambda = \frac{(D-2)\kappa\lambda}{2\varrho}(1 + \ell), \quad (2.25)$$

which is the constraint on the additional field λ coming from the potential (2.10). So λ isn't a new freedom and we can define the effective cosmological constant Λ_e as following to absorb it,

$$\Lambda_e = \frac{(D-2)\kappa\lambda}{2\varrho}. \quad (2.26)$$

Then $\Lambda = (1 + \ell)\Lambda_e$ and, the present black hole metric is

$$ds^2 = -f(r)dt^2 + \frac{1 + \ell}{f(r)}dr^2 + r^2 d\Omega_{D-2}^2, \quad (2.27)$$

with the metric function $f(r)$ as

$$f(r) = 1 - \frac{16\pi M}{(D-2)\Omega_{D-2}r^{D-3}} - \frac{2(1 + \ell)\Lambda_e}{(D-1)(D-2)}r^2. \quad (2.28)$$

This metric can't be asymptotic to the AdS spacetime due to the factor $(1 + \ell)$ before g_{rr} forbidding this limit. The black hole horizon locates at the largest real root of the following equation

$$1 - \frac{16\pi M}{(D-2)\Omega_{D-2}r_h^{D-3}} - \frac{2(1 + \ell)\Lambda_e}{(D-1)(D-2)}r_h^2 = 0. \quad (2.29)$$

It shows that the bumblebee field affects the location of the event horizons, contrary to the Schwarzschild-like solution in which the horizon radius is the same as of the Schwarzschild black hole.

III. THERMODYNAMICS OF EINSTEIN-BUMBLEBEE ADS BLACK HOLE

A black hole is not only a gravity system, but also a special thermodynamic system due to that its surface gravity κ and horizon area A are the close similarity to the temperature T and entropy S , $T = \kappa/2\pi$, $S = A/4$. Four laws of black hole thermodynamics were established in Ref. [23]. LV can modify the geometry of a black hole, so in this kind of black hole, can the thermodynamical laws still hold? In this section we study the Smarr formula and the first law of this LV black hole.

In this D -dimensional spacetime, the Killing vector is $\xi^a = (1, 0, 0, 0, \dots, 0)$. In order to construct a Komar integral relation for non-zero cosmological constant, one should introduce Killing potential ω^{ab} [24] which can be obtained according to relation $\xi^b = \nabla_a \omega^{ab}$. For the static Killing vector ξ^a , we have

$$\omega^{rt} = -\omega^{tr} = \frac{r}{D-1}. \quad (3.1)$$

We can integrate the Killing equation $\nabla_b(\nabla^b \xi^a) = -R_b^a \xi^b$ over a hypersurface S ,

$$\int_{\partial S} \nabla^b \xi^a d\Sigma_{ab} = - \int_S R_b^a \xi^b d\Sigma_a, \quad (3.2)$$

where $d\Sigma_{ab}$ and $d\Sigma_a$ are the surface elements of ∂S and S , respectively, and ∂S is a 2-surface bounding S .

With the metric (2.13) in mind, we have the relation

$$R_b^a \xi^b = \frac{2}{D-2}(\Lambda_e, 0, 0, 0, \dots, 0) = \frac{2\Lambda_e}{D-2}\xi^a, \quad (3.3)$$

then the Eq. (3.2) can be rewritten as

$$\frac{D-2}{8\pi} \int_{\partial S} \left(\nabla^b \xi^a + \frac{2}{D-2} \Lambda_e \omega^{ab} \right) d\Sigma_{ab} = 0, \quad (3.4)$$

which is called the Komar integral relation. The closed 2-surface ∂S has two parts, horizon ∂S_h and infinite ∂S_∞ , so Eq. (3.4) can be rewritten as

$$\frac{D-2}{8\pi} \int_{\partial S_\infty} \left(\nabla^b \xi^a + \frac{2}{D-2} \Lambda_e \omega^{ab} \right) d\Sigma_{ab} = \frac{D-2}{8\pi} \int_{\partial S_h} \left(\nabla^b \xi^a + \frac{2}{D-2} \Lambda_e \omega^{ab} \right) d\Sigma_{ab}. \quad (3.5)$$

This integral can give the Smarr formula

$$(D-3)M = (D-2)TS - 2PV, \quad (3.6)$$

where the Hawking temperature [25] and the pressure are

$$T = \frac{\sqrt{f'(r_h)g'(r_h)}}{4\pi} = \frac{1}{4\pi\sqrt{1+\ell r_h}} \left[(D-3) - \frac{2}{D-2}(1+\ell)\Lambda_e r_h^2 \right], \quad P = -\frac{\Lambda_e}{8\pi}, \quad (3.7)$$

and the entropy and the volume are,

$$S = \int \frac{dM}{T} = \frac{\sqrt{1+\ell}}{4} \Omega_{D-2} r_h^{D-2}, \quad V = \left(\frac{\partial M}{\partial P} \right)_S = (1+\ell) \frac{\Omega_{D-2}}{D-1} r_h^{D-1}. \quad (3.8)$$

One can see that the Smarr formula can still be constructed in this LV black hole spacetime as long as its temperature, entropy and volume are slightly modified. Using Hamiltonian perturbation method [24], it is easy to prove there exists the first law of black hole thermodynamics as following

$$dM = TdS + VdP, \quad (3.9)$$

which still also holds in this LV black hole spacetime.

IV. PHASE TRANSITION

A phase transition is a discontinuous change in the properties of a substance, as its environment is changed only infinitesimally. An isolated Schwarzschild black hole is thermodynamically unstable due to its negative heat capacity. However for a Schwarzschild AdS black hole, besides the negative heat phase, a phase of positive heat capacity will occur. At a given temperature and pressure, the stable phase is always the one with the lower Gibbs free energy. In an AdS space, the Gibbs energy of the thermal radiation phase is zero [26]. So the negative free energy of the positive heat capacity black hole phase is stable phase. In this section, we study this two phase transitions.

From the Eq. (2.29), we can express the mass M as

$$M = \frac{(D-2)\Omega_{D-2}}{16\pi} r_h^{D-3} + PV. \quad (4.1)$$

M can be interpreted as an enthalpy [24] $H = U + PV$, so the black hole internal energy U is

$$U = \frac{(D-2)\Omega_{D-2}}{16\pi} r_h^{D-3}. \quad (4.2)$$

Then the first law (3.9) can be rewritten as following

$$dU = TdS - PdV, \quad (4.3)$$

which is the same as the fundamental thermodynamical equation of a simple gas system. The temperature (3.7) can be rewritten as

$$T = \frac{1}{4\pi\sqrt{1+\ell} r_h} \left[(D-3) + \frac{16\pi P}{D-2} (1+\ell) r_h^2 \right], \quad (4.4)$$




FIG. 1: Black hole temperature and heat capacity C_P at constant pressure P vs. black hole mass M with $D = 4$ and different coupling constant ℓ . The temperature has a minimum value and the the heat capacity is divergent at the point of black hole mass $M = M_d$, where $M_d = 1/\sqrt{18(1+\ell)\pi P}$.

which is shown in Fig. 1 when $D = 4$. There exist a minimum temperature

$$T_{min} = 2\sqrt{\frac{(D-3)P}{(D-2)\pi}}, \quad (4.5)$$

when $r_h = \sqrt{(D-2)(D-3)/[16(1+\ell)\pi P]}$. When $T > T_{min}$, there are two possible black hole masses [20] which can be in equilibrium with thermal radiation, i.e., $M > M_d$ or $M < M_d$. Fig. 1 shows that the temperature of the large black hole ($M > M_d$) increases when the LV coupling constant ℓ becomes bigger. At a fixed pressure P , the black hole heat capacity C_P is

$$C_P = \left(\frac{\partial H}{\partial T}\right)_P = \frac{(D-2)\Omega_{D-2}}{4} r_h^{D-2} \frac{16\pi(1+\ell)Pr_h^2 + (D-2)(D-3)}{16\pi(1+\ell)Pr_h^2 - (D-2)(D-3)}, \quad (4.6)$$

which is also shown in Fig. 1 when $D = 4$. It is easy to see that the heat capacity at constant pressure is divergent at the black hole mass M_d ,

$$M_d = \frac{(D-2)^2\Omega_{D-2}}{8\pi(D-1)} \left[\frac{(D-2)(D-3)}{16(1+\ell)\pi P} \right]^{(D-3)/2}. \quad (4.7)$$

At this divergent point, the black hole entropy is

$$S_d = 4^{(1-D)}\Omega_{D-2}\sqrt{1+\ell}^{(3-D)} \left[\frac{(D-2)(D-3)}{\pi P} \right]^{(D-2)/2}, \quad (4.8)$$

which is shown in Fig. 2 when $D = 4$. From the Fig. 1, one can see that the small black hole mass has negative specific heat, so it is unstable and will decay. While the large black hole mass has positive heat capacity and therefore it is thermally stable. The phase transition between these two phases is called small-large black hole phase transition.

There is another kind of phase transition called Hawking-Page(HP) phase transition, which is between a metastable large black hole phase and stable large black hole phase in an anti-de Sitter spacetime. Witten [27]




FIG. 2: Black hole entropy S vs. black hole temperature T with $D = 4$ and different coupling constant ℓ . They all have both branches, the above one is corresponding to the large black hole $M > M_d$, the below one to the small black hole $M < M_d$. At the temperature $T = T_{min}$, the entropy is $S_d = 1/8\sqrt{1 + \ell P}$.

explained that this HP phase transition is a confinement/deconfinement phase transition in the AdS/CFT correspondence. It can also be understood as a solid/liquid phase transition in black hole chemistry [28]. HP phase transition occurs when Gibbs free energy is zero. The black hole Gibbs free energy can be obtained

$$G = H - TS = \frac{\Omega_{D-2}}{8\pi(D-1)} \left[(D-2) - 2\pi\sqrt{1 + \ell T r_h} \right] r_h^{D-3}, \quad (4.9)$$

which is shown in Fig. 3 when $D = 4$. It can be seen that there are two branches, the above parts corresponding to small black hole phase whose free energy are positive for any temperature; the below ones corresponding to metastable large black hole phase when $G > 0$ and stable large black hole phase when $G < 0$. When the temperature $T < T_{min}$, the spacetime has only one phase—thermal radiation, whose free energy is zero. At the temperature $T = T_{min}$, its free energy has a maximum value G_{max} ,

$$G_{max} = \frac{\Omega_{D-2}}{(D-1)\pi} 2^{(3-2D)} \left[\frac{(D-2)(D-3)}{(1+\ell)\pi P} \right]^{(D-3)/2}. \quad (4.10)$$

The temperature at Hawking-Page phase transition point can be obtained when $G = 0$,

$$T_{HP} = 2\sqrt{\frac{(D-2)P}{(D-1)\pi}}. \quad (4.11)$$

At this point $r_h = \sqrt{(D-1)(D-2)/[16(1+\ell)\pi P]}$ and the black hole mass is,

$$M_{HP} = \frac{(D-2)\Omega_{D-2}}{8\pi} \left[\frac{(D-1)(D-2)}{16(1+\ell)\pi P} \right]^{(D-3)/2}, \quad (4.12)$$

its entropy is

$$S_{HP} = 4^{(1-D)}\Omega_{D-2}\sqrt{1+\ell}^{(3-D)} \left[\frac{(D-1)(D-2)}{\pi P} \right]^{(D-2)/2}. \quad (4.13)$$




FIG. 3: Black hole Gibbs free energy vs. temperature with $D = 4$ and different coupling constant ℓ . They vanish at the point $T_{HP} = \sqrt{8P/3\pi}$. They all have both branches, the above one is corresponding to the small black hole $M < M_d$, the below one to the large black hole $M > M_d$. At the temperature $T = T_{min}$, the free energy is $G_d = 1/12\sqrt{2(1+\ell)\pi P}$.

TABLE I: Quantities of heat capacity, entropy, temperature and Gibbs free energy for three phases of the AdS black hole in Einstein-bumblebee gravity.

	Black hole mass	Heat capacity	Entropy	Temperature	Free energy
Unstable small black hole	$0 < M < M_d$	$C_p < 0$	$S < S_d$	$T > T_{min}$	$G > 0$
Metastable large black hole	$M_d < M < M_{HP}$	$C_p > 0$	$S_d < S < S_{HP}$	$T_{min} < T < T_{HP}$	$0 < G < G_d$
Stable large black hole	$M > M_{HP}$	$C_p > 0$	$S > S_{HP}$	$T > T_{HP}$	$G < 0$

We summarize these three phases in Tab. I. From the figures 1 to 3 and the table I, one can see the following picture for the formation of a AdS black hole: i), in a AdS space, the thermal radiation with zero Gibbs free energy has a very high temperature, accidentally, a tiny mass black hole begins to form due to quantum fluctuation; ii), the tiny black hole begins to grow, its temperature decreases quickly, its free energy increases slowly; iii), when the black hole mass grows to M_d , its free energy approaches to a maximum, its temperature approaches to a minimum, and the small-large black hole phase transition occurs; iv), the black hole mass continues to grow, its positive free energy decreases, its temperature increases slowly and it is a metastable black hole; v), when the mass grows to M_{HP} , the HP phase transition occurs; vi), finally, the stable large black hole has formed.

Now we consider some relationships between the both kinds of phase transition. The first are for the temperature and entropy,

$$\frac{T_{HP}}{T_{min}} = \frac{D-2}{\sqrt{(D-1)(D-3)}}, \quad \frac{S_{HP}}{S_d} = \left(\frac{D-1}{D-3}\right)^{(D-2)/2}, \quad (4.14)$$

which are only dependant on the dimension number D and the same as those in Refs. [26, 29]. The sec-

ond relationship is between the D dimensional Hawking-Page temperature T_{HP} and the $D + 1$ dimensional minimum temperature T_{min} ,

$$T_{HP}(D) = T_{min}(D + 1), \quad (4.15)$$

which is the same as that in Ref. [26]. This relation is also independent of the pressure and the bumblebee coupling constant, and is like the AdS/CFT correspondence. T_{min} is a physical quantity in the bulk, then T_{HP} can be treated as the dual physical quantity on the boundary.

V. SUMMARY

In this paper, we have derived the high dimensional AdS black hole solution in Einstein-bumblebee gravity theory. It can exist only under the condition that the bumblebee potential has a linear functional form with a Lagrange-multiplier field λ . This additional field is strictly constrained by the bumblebee motion equation and can be absorbed by the definition of an effective cosmological constant Λ_e . Like Schwarzschild-like black hole [14], it can't be asymptotically to anti-de Sitter spacetime, so it is a Schwarzschild-AdS-like black hole solution [17]. Contrary to Schwarzschild-like black hole, the bumblebee field will affect the locations of the black hole horizon.

By using black hole chemistry, we have studied the thermodynamics and phase transitions of this high dimensional Schwarzschild-AdS-like black hole. With Komar integral method, we find that the Smarr formula can exist as long as its temperature, entropy and volume are slightly modified. With Hamiltonian perturbation method, there still exist the first law of thermodynamics.

Its temperature has a minimum value T_{min} , corresponding to the black hole mass M_d . Small black hole with mass $0 < M < M_d$ has a negative heat capacity C_p , while the large black hole with mass $M > M_d$ has a positive one. At the point M_d , the C_p is divergent. The LV constant ℓ decreases the mass M_d , showing that the phase transition from a small black hole to large one will more easily occur.

Its Gibbs energy has a maximum value G_{max} at the point of mass M_d and has a zero point $G = 0$ at the point of mass M_{HP} . Small black hole with mass $0 < M < M_d$ and large black hole with mass $M_d < M < M_{HP}$ have a positive Gibbs free energy, while the large black hole with mass $M > M_d$ has a negative one. The LV constant ℓ decreases the Gibbs free energy of the meta-stable large black hole with the mass $M_d < M < M_{HP}$, showing that the HP phase transition from a meta-stable large black hole to large stable one will be more easily to occur.

The dualities found in Ref. [26] still holds when the Lorentz symmetry is spontaneously broken: the ratio of temperature between small/large black hole phase transition and HP phase transition T_{HP}/T_{min} , and the equivalence of the temperature of D -dimensional HP phase transition to the temperature of $D+1$ dimensional small/large black hole phase transition $T_{HP}(D) = T_{min}(D+1)$.

Acknowledgments

This work was supported by the Scientific Research Fund of the Hunan Provincial Education Department under No. 19A257 and No. 19A260, the National Natural Science Foundation (NNSFC) of China (grant No. 11247013), Hunan Provincial Natural Science Foundation of China grant No. 2015JJ2085.

-
- [1] A. Ashtekar and V. Petkov, *Springer handbook of spacetime*P485, Springer Dordrecht, (2014).
 - [2] D. Mattingly, Living Rev. Rel. **8**, 5 (2005).
 - [3] G. Amelino-Camelia, Liv. Rev. Rel. **16**, 5 (2013).
 - [4] V. A. Kostelecký, Phys. Rev. D **69**, 105009 (2004).
 - [5] V. A. Kostelecký and S. Samuel, Phys. Rev. D **39**, 683 (1989).
 - [6] S. M. Carroll, J. A. Harvey, V. A. Kostelecky, C. D. Lane, and T. Okamoto, Phys. Rev. Lett. **87**, 141601 (2001); I. Mocioiu, M. Pospelov, and R. Roiban, Phys. Lett. B **489**, 390 (2000); A. F. Ferrari, M. Gomes, J. R. Nascimento, E. Passos, A. Yu. Petrov, and A. J. da Silva, Phys. Lett. B **652**, 174 (2007).
 - [7] O. Bertolami, R. Lehnert, R. Potting, and A. Ribeiro, Phys. Rev. D **69**, 083513 (2004); V. A. Kostelecký, R. Lehnert, and M. Perry, Phys. Rev. D **68**, 123511 (2003); R. Jackiw and S.-Y. Pi, Phys. Rev. D **68**, 104012 (2003).
 - [8] R. Gambini and J. Pullin, Phys. Rev. D **59**, 124021 (1999); J. R. Ellis, N. E. Mavromatos, and D. V. Nanopoulos, Gen. Relativ. Gravit. **32**, 127 (2000).
 - [9] C. P. Burgess, J. Cline, E. Filotas, J. Matias, and G.D. Moore, J. High Energy Phys. **03** (2002) 043; A. R. Frey, J. High Energy Phys. **04** (2003) 012; J. Cline and L. Valcárcel, J. High Energy Phys. **03** (2004) 032.
 - [10] A. Fernando, T. Clark, Gen. Relativ. Grav. **46**, 1834 (2014).
 - [11] T. Jacobson and D. Mattingly, Phys. Rev. D **64**, 024028 (2001); T. Jacobson, Proc. Sci. QG-PH (2007) 020.
 - [12] R. Bluhm, Shu-Hong Fung and V. A. Kostelecký, Phys. Rev. D **77**, 065020 (2008).
 - [13] O. Bertolami and J. Páramos, Phys. Rev. D **72**, 044001 (2005).
 - [14] R. Casana and A. Cavalcante, Phys. Rev. D **97**, 104001 (2018).
 - [15] C. Ding, C. Liu, R. Casana and A. Cavalcate, Eur. Phys. C **80**, 178 (2020).
 - [16] S. K. Jha and A. Rahaman, arXiv: 2011.14916, (2021).
 - [17] R. V. Maluf and J. C. S. Neves, Phys. Rev. D **103**, 044002 (2021).
 - [18] T. Padmanabhan, *Gravitation: Foundations and Frontiers*, Cambridge University Press, (2010).
 - [19] S.W. Hawking, Commun. Math. Phys. **43**, 199 (1975); J. Bekenstein, Lett. Nuovo Cimento **4**, 737 (1972); J. D. Bekenstein, Phys. Rev. D **7**, 2333 (1973).

- [20] S. W. Hawking and D. N. Page, *Cummum. Math. Phys.* **87**, 577 (1983).
- [21] C. Ding and X. Chen, *Chin. Phys. C* **45**, 025106 (2021).
- [22] D. G. Boulware and S. Deser, *Phys. Rev. Lett.* **55**, 2656 (1985).
- [23] J. M. Bardeen, B. Carter, and S. Hawking, *Commun. Math. Phys.* **31**, 161 (1973).
- [24] D. Kastor, S. Ray and J. Traschen, *Class. Quantum Grav.* **26**, 195011 (2009).
- [25] C. Ding and J. Jing, *Class. Quantum Grav.* **25**, 145015 (2008).
- [26] Shao-Wen Wei and Yu-Xiao Liu, *Phys. Rev. D* **102**, 104011 (2020).
- [27] E. Witten, Anti-de Sitter space, *Adv. Theor. Math. Phys.* **2**, 505 (1998).
- [28] D. Kubiznak and R. B. Mann, *Can. J. Phys.* **93**, 999 (2015).
- [29] A. Belhaj, A. El Balali, W. El Hadri and E. Torrente-Lujan, *Phys. Lett. B* **811**, 135871 (2020).