

Generalized Lévy-Leblond equation with external potentials and effects of rest energy for non-relativistic Fermi fields

Fuad M. Saradzhev ¹

Centre for Science, Athabasca University, Athabasca, Alberta, Canada

Abstract

The generalized Lévy-Leblond equation (GLL) is used to study the bound state problem for a non-relativistic Fermi field in pseudoscalar external potentials. Two spherically symmetrical external potentials, a pseudoscalar spherical well of finite depth and a pseudoscalar Coulomb potential, are considered. It is shown that the rest energy of the Fermi field affects non-trivially the bound state spectrum. The existence of bound states, their number and energies all depend on the value of the rest energy.

¹fuads@athabascau.ca

1 Introduction

Non-relativistic particles (fields), defined through the theory of representations of the Galilei group, are known to acquire spin and magnetic moment [1], the properties which were originally believed to be purely relativistic. The dynamics of the spin 1/2 particles is governed by the Lévy-Leblond (LL) equation, the non-relativistic version of the Dirac equation. In [1], it was derived in the Dirac way by linearization of the Schrödinger equation.

Another way of deriving the LL equation is by making use of a $(4+1)$ -dimensional covariant formulation of Galilean covariance [2] based on an extended space-time approach [3],[4]. A formulation of Galilean covariance in one higher dimension allows us to incorporate into the theory the notion of rest (internal) energy: one of the Casimir invariants of the Galilei group in $(4+1)$ -dimensions includes the rest energy operator. Reduction to $(3+1)$ -dimensions by eliminating the additional, fifth coordinate results in the generalized LL equation with a rest energy contribution [5]. A generalized LL equation was also discussed in [6].

It is often stated that the rest energy is a reference point for the energy of non-relativistic systems and the only effect of taking the rest energy into account is a shift of all energy levels by its value. This is equivalent to a change in a phase of wave function. In this paper, we aim to show that the rest energy of non-relativistic Fermi particles can have a non-trivial impact on their dynamics. We study the GLL equation in external, pseudoscalar potentials to determine how the rest energy affects the bound state spectrum.

Our paper is organized as follows. In Sect.2, we consider a free non-relativistic Fermi field. We start with the Schrödinger equation with the rest energy term, and we show that its linearization leads to the GLL equation. This provides an alternative derivation of the GLL equation in the same way as the LL equation was derived. A rest mass associated with the rest energy is introduced, and the component form of the GLL equation is given.

The covariance of the LL equation under reflection of space coordinates was proven in [7]. We construct a representation of the parity transformation, and we demonstrate that the rest energy term violates the covariance of the GLL equation. Only one component of the non-relativistic Fermi field can be characterized by a definite parity, while the Fermi field itself does not have it.

We establish a two-step transition from the Dirac equation to the GLL equation. Taking the non-relativistic limit of the Dirac equation yields the LL equation [1], and an additional transformation is required to get its generalized version.

In Sect.3, we consider a non-relativistic Fermi field in external potentials. We find that if the potentials are pseudoscalar, then the rest energy cannot be removed from the GLL equation. We derive a Schrödinger type equation for the independent component of the Fermi field, its effective potential exhibiting the rest energy contributions and the parity violation.

We study in details two spherically symmetric external potentials: a pseudoscalar spherical well of finite depth and a pseudoscalar Coulomb potential. In the first case, the effective potential is also a well which depth depends on the value of the rest mass. We determine the spectrum of the bound S-states, and we find the minimum value of the rest energy below which no bound states exist.

In the case of pseudoscalar Coulomb potential created by an external charge, the effective potential includes an interaction with a background field which magnitude is determined by the rest mass. We construct an additional constant of motion with the rest mass contribution, and we determine how it changes the structure of angular parts of eigenfunctions. We find the energy spectrum of the system in the approximation when the external charge is very large. We show that the spectrum is affected by the orientation of the particle's spin with respect to the direction of the background field.

We conclude with Discussion in Sect.4.

2 GLL equation for a free non-relativistic Fermi field

The Schrödinger equation for a free non-relativistic Fermi field $\psi_+(\mathbf{r}, t)$ with inertial mass m and rest energy E_0 reads

$$i\hbar \frac{\partial \psi_+(\mathbf{r}, t)}{\partial t} = \left(-\hbar^2 \frac{\Delta}{2m} + E_0 \right) \psi_+(\mathbf{r}, t), \quad (1)$$

where $\mathbf{r} = (x^1 = x, x^2 = y, x^3 = z)$, $\Delta \equiv \partial_a^2$, $\partial_a \equiv \frac{\partial}{\partial x^a}$, $a = 1, 2, 3$, and the subscript "+" indicates that we restrict ourselves to the positive inertial mass sector.

By re-defining the wave function $\psi_+(\mathbf{r}, t)$ as

$$\psi_+(\mathbf{r}, t) \rightarrow \tilde{\psi}_+(\mathbf{r}, t) \equiv e^{\frac{i}{\hbar} E_0 t} \psi_+(\mathbf{r}, t), \quad (2)$$

we can remove the rest energy term from the Schrödinger equation:

$$i\hbar \frac{\partial \tilde{\psi}_+(\mathbf{r}, t)}{\partial t} = -\hbar^2 \frac{\Delta}{2m} \tilde{\psi}_+(\mathbf{r}, t). \quad (3)$$

The LL equation is a non-relativistic equation obtained as a result of linearization of the Schrödinger equation (3) without the rest energy term. Let us follow the similar procedure to get a linearized version of Eq.(1) with the rest energy term included. We represent the linearized equation as

$$\left(\Gamma_1 \frac{\partial}{\partial t} + \Gamma^a \partial_a + \Gamma_2 \right) \psi_+(\mathbf{r}, t) = 0, \quad (4)$$

where Γ_1, Γ^a , and Γ_2 are (4×4) -matrices to be determined from the requirement that the solutions of Eq.(4) should obey Eq.(1) as well. This means that there should exist

some operator $\Gamma'_1 \frac{\partial}{\partial t} + \Gamma'^a \partial_a + \Gamma'_2$, such that

$$\left(\Gamma'_1 \frac{\partial}{\partial t} + \Gamma'^a \partial_a + \Gamma'_2 \right) \left(\Gamma_1 \frac{\partial}{\partial t} + \Gamma^b \partial_b + \Gamma_2 \right) = S, \quad (5)$$

where

$$S \equiv i\hbar \frac{\partial}{\partial t} + \hbar^2 \frac{\Delta}{2m} - E_0 \quad (6)$$

is the Schrödinger operator. This yields the set of conditions

$$\begin{aligned} \Gamma'_1 \Gamma_1 &= 0, & \Gamma'_1 \Gamma_2 + \Gamma'_2 \Gamma_1 &= i\hbar, \\ \Gamma'_1 \Gamma^a + \Gamma'^a \Gamma_1 &= 0, & \Gamma'^a \Gamma_2 + \Gamma'_2 \Gamma^a &= 0, \\ \Gamma'^a \Gamma^b + \Gamma'^b \Gamma^a &= \frac{\hbar^2}{m} \delta^{ab}, & \Gamma'_2 \Gamma_2 &= -E_0. \end{aligned} \quad (7)$$

These conditions differ from the ones given by Lévy-Leblond only in the last equation in (7) where we can see the rest energy contribution. To bring them to the form given in [1], we replace Γ_2 and Γ'_2 by

$$\begin{aligned} \tilde{\Gamma}_2 &= \Gamma_2 + i\sqrt{E_0}I, \\ \tilde{\Gamma}'_2 &= \Gamma'_2 + i\sqrt{E_0}I, \end{aligned}$$

where I is the identity matrix. Then

$$\tilde{\Gamma}'_2 \tilde{\Gamma}_2 = i\sqrt{E_0} (\tilde{\Gamma}'_2 + \tilde{\Gamma}_2),$$

so the product $\tilde{\Gamma}'_2 \tilde{\Gamma}_2$ becomes independent of the rest energy if we take $\tilde{\Gamma}'_2 = -\tilde{\Gamma}_2$.

To avoid the appearance of the rest energy terms in other equations of (7), we also take $\Gamma'_1 = -\Gamma_1$ and $\Gamma'^a = -\Gamma^a$. This brings Eq.(7) to the form

$$\begin{aligned} \Gamma_1^2 &= 0, & \Gamma_1 \tilde{\Gamma}_2 + \tilde{\Gamma}_2 \Gamma_1 &= -i\hbar, \\ \Gamma_1 \Gamma^a + \Gamma^a \Gamma_1 &= 0, & \Gamma^a \tilde{\Gamma}_2 + \tilde{\Gamma}_2 \Gamma^a &= 0, \\ \Gamma^a \Gamma^b + \Gamma^b \Gamma^a &= -\frac{\hbar^2}{m} \delta^{ab}, & \tilde{\Gamma}_2^2 &= 0. \end{aligned} \quad (8)$$

Representing Γ_1 , Γ^a and $\tilde{\Gamma}_2$ as

$$\Gamma_1 = -\frac{\hbar}{c\sqrt{2m}}\gamma^4, \quad \Gamma^a = -\frac{\hbar}{\sqrt{2m}}\gamma^a, \quad \tilde{\Gamma}_2 = ic\sqrt{\frac{m}{2}}\gamma^5,$$

where c is the speed of light, we rewrite these conditions as the algebra

$$\gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2g^{\mu\nu}, \quad (9)$$

where $\mu, \nu = 1, \dots, 5$ and

$$g_{\mu\nu} = \begin{pmatrix} -\mathbf{1}_{3 \times 3} & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}.$$

In what follows, we will use the following representation for the γ -matrices [8]:

$$\gamma^a = \begin{pmatrix} 0 & i\sigma^a \\ i\sigma^a & 0 \end{pmatrix}, \quad \gamma^4 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -1 & -1 \end{pmatrix}, \quad \gamma^5 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & -1 \\ 1 & -1 \end{pmatrix}. \quad (10)$$

The linearized Eq.(4) becomes

$$\left(i\hbar\gamma^{\bar{\mu}}\partial_{\bar{\mu}} + mc\gamma^5 - \sqrt{2mE_0}I \right) \psi_+(\mathbf{r}, t) = 0, \quad (11)$$

where $\bar{\mu}$ runs from 1 to 4, and $\partial_4 \equiv \frac{1}{c} \frac{\partial}{\partial t}$. Herein, $\sqrt{2mE_0}$ can be interpreted as a momentum k corresponding to the rest energy, so that $E_0 = k^2/(2m)$. We can also introduce a rest mass m_0 associated with the rest energy E_0 as $m_0 \equiv E_0/c^2$. This gives us the following relation between the parameters k and c :

$$k = \sqrt{2mm_0}c.$$

Eqs.(11) takes the compact form

$$(i\hbar\gamma^{\bar{\mu}}\partial_{\bar{\mu}} - kI_+) \psi_+(\mathbf{r}, t) = 0, \quad (12)$$

where

$$I_+ \equiv I - \frac{mc}{k}\gamma^5.$$

We will refer to this equation as the generalized LL (GLL) equation for a free non-relativistic Fermi field. For $k = 0$, it reduces to the LL equation.

The representation (10) is not unique. There are infinitely many representations of the algebra (9) which are equivalent to each other. Indeed, Eq.(12) is invariant with respect to unitary transformations

$$\gamma^{\bar{\mu}} \rightarrow U\gamma^{\bar{\mu}}U^{-1}, \quad \gamma^5 \rightarrow U\gamma^5U^{-1}, \quad \psi_+ \rightarrow U\psi_+, \quad (13)$$

where U is a unitary (4×4) -matrix. The standard Dirac representation of the γ -matrices used in [1] can be obtained from Eq.(10) by making use of the transformation (13) with

$$U = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}.$$

As in the case of the Schrödinger equation, the rest energy term can be removed from the GLL equation as well. To show this, let us introduce the new field

$$\psi_+(\mathbf{r}, t) \rightarrow \tilde{\psi}_+(\mathbf{r}, t) \equiv \left(I - \frac{\mu}{\sqrt{2}}\gamma^4 \right) e^{\frac{i}{\hbar}E_0t} \psi_+(\mathbf{r}, t),$$

where μ is a parameter to be determined. The GLL equation given by Eq.(12) becomes

$$\left[i\hbar\gamma^4\partial_4 + i\hbar\left(I - \frac{\mu}{\sqrt{2}}\gamma^4\right)\gamma^a\partial_a - kI_+ \left(I + \frac{\mu}{\sqrt{2}}\gamma^4\right) + \frac{1}{c}E_0\gamma^4 \right] \tilde{\psi}_+(\mathbf{r}, t) = 0.$$

Multiplying both sides of this equation from the left by $\left(I + \frac{\mu}{\sqrt{2}}\gamma^4\right)$, we bring it to the form

$$\left[i\hbar\gamma^{\bar{\mu}}\partial_{\bar{\mu}} + mc\gamma^5 + (\mu mc\sqrt{2} - k)I + \frac{1}{2mc}(\mu mc\sqrt{2} - k)^2\gamma^4 \right] \tilde{\psi}_+(\mathbf{r}, t) = 0. \quad (14)$$

If we take

$$\mu = \frac{k}{mc\sqrt{2}} = \sqrt{\frac{m_0}{m}}, \quad (15)$$

then the last two terms in Eq.(14) vanish, so the field

$$\tilde{\psi}_+(\mathbf{r}, t) = \left(I - \sqrt{\frac{m_0}{2m}}\gamma^4\right) e^{\frac{i}{\hbar}E_0t}\psi_+(\mathbf{r}, t), \quad (16)$$

obeys the original LL equation without the rest energy term:

$$(i\hbar\gamma^{\bar{\mu}}\partial_{\bar{\mu}} + mc\gamma^5)\tilde{\psi}_+(\mathbf{r}, t) = 0. \quad (17)$$

In addition to the factor $\exp\{\frac{i}{\hbar}E_0t\}$ which we had before in Eq.(2), we get here the factor $(1 - \sqrt{\frac{m_0}{2m}}\gamma^4)$ that depends on the ratio of the rest and inertial masses and mixes the components of the original field $\psi_+(\mathbf{r}, t)$.

Representing the field $\psi_+(\mathbf{r}, t)$ as

$$\psi_+(\mathbf{r}, t) = \begin{pmatrix} \psi_{1,+}(\mathbf{r}, t) \\ \psi_{2,+}(\mathbf{r}, t) \end{pmatrix},$$

where $\psi_{1,+}(\mathbf{r}, t)$, $\psi_{2,+}(\mathbf{r}, t)$ are 2-component fields, and introducing their linear combinations

$$\begin{aligned} \eta_{1,+}(\mathbf{r}, t) &\equiv \psi_{1,+}(\mathbf{r}, t) + \psi_{2,+}(\mathbf{r}, t) \\ \eta_{2,+}(\mathbf{r}, t) &\equiv \psi_{1,+}(\mathbf{r}, t) - \psi_{2,+}(\mathbf{r}, t) \end{aligned}$$

we can rewrite the GLL equation as a system of two equations:

$$\begin{aligned} i\hbar\partial_4\eta_{1,+} + p_-\eta_{2,+} &= 0, \\ p_+\eta_{1,+} - mc\eta_{2,+} &= 0. \end{aligned} \quad (18)$$

where

$$p_{\pm} \equiv \frac{1}{\sqrt{2}}(\hbar\sigma^a\partial_a \pm k),$$

Only one of the components is dynamically independent. Its time evolution determines the time evolution of another component as well. Eliminating, for instance, $\eta_{2,+}(\mathbf{r}, t)$ in favor of $\eta_{1,+}(\mathbf{r}, t)$, this brings us back to the Schrödinger equation (1):

$$i\hbar\partial_4\eta_{1,+} + \frac{p_-p_+}{mc}\eta_{1,+} = 0 \quad (19)$$

with $p_-p_+ = \frac{1}{2}(\hbar^2\Delta - k^2)$. The GLL equation for a free non-relativistic Fermi field can be therefore reduced to the Schrödinger equation for its independent component. In this sense, the GLL equation and the Schrödinger equation are equivalent to each other, and we can use any of these two equations as a starting point in the study of free non-relativistic Fermi fields.

2.1 Parity

Let us determine if the GLL equation is covariant under reflection of space coordinates:

$$\mathbf{r}' = -\mathbf{r}, \quad t' = t. \quad (20)$$

The covariance means that in the primed frame the GLL equation has the same form as in the unprimed one,

$$(i\hbar\gamma^{\bar{\mu}}\partial_{\bar{\mu}}' - kI_+) \psi_+'(\mathbf{r}', t) = 0, \quad (21)$$

where

$$\psi_+'(\mathbf{r}', t) = S(P)\psi_+(\mathbf{r}, t) \quad (22)$$

and $S(P)$ is a representation of the parity transformation (20). It is assumed that $(S(P))^2 = 1$.

To find $S(P)$, we substitute (22) into the GLL equation in the primed frame and multiply it by $S^{-1}(P)$. Since

$$\partial_a' = -\partial_a, \quad \partial_4' = \partial_4,$$

this gives us

$$\begin{aligned} & \{-i\hbar S^{-1}(P)\gamma^a S(P)\partial_a + i\hbar S^{-1}(P)\gamma^4 S(P)\partial_4 \\ & \quad - kI + mcS^{-1}(P)\gamma^5 S(P)\}\psi_+(\mathbf{x}, t) = 0. \end{aligned} \quad (23)$$

Let us consider first the case when k is equal to zero. Then Eq.(23) coincides with the corresponding equation in the unprimed frame if

$$S^{-1}(P)\gamma^a S(P) = \gamma^a, \quad S^{-1}(P)\gamma^4 S(P) = -\gamma^4, \quad S^{-1}(P)\gamma^5 S(P) = -\gamma^5 \quad (24)$$

(see Appendix A). This is satisfied by

$$S(P) = e^{i\delta\frac{1}{2}(\gamma^5\gamma^4 - \gamma^4\gamma^5)}, \quad (25)$$

where $e^{i\delta} = \pm 1$. This phase is related to the intrinsic parity of the field.

In the component form, the relation (22) between the primed and unprimed fields becomes

$$\begin{aligned}\psi'_{1,+}(\mathbf{r}, t) &= e^{i\delta}\psi_{2,+}(-\mathbf{r}, t), \\ \psi'_{2,+}(\mathbf{r}, t) &= e^{i\delta}\psi_{1,+}(-\mathbf{r}, t).\end{aligned}$$

If the field $\psi_+(\mathbf{r}, t)$ has a definite parity, then

$$\psi'_{1,+}(\mathbf{r}, t) = \psi_{1,+}(\mathbf{r}, t), \quad \psi'_{2,+}(\mathbf{r}, t) = \psi_{2,+}(\mathbf{r}, t). \quad (26)$$

This yields

$$\eta_{1,+}(\mathbf{r}, t) = e^{i\delta}\eta_{1,+}(-\mathbf{r}, t), \quad (27)$$

$$\eta_{2,+}(\mathbf{r}, t) = -e^{i\delta}\eta_{2,+}(-\mathbf{r}, t), \quad (28)$$

i.e. the spatial parity of the dependent component is opposite to the spatial parity of the independent one.

For a non-zero k , the equation (23) with $S(P)$ given by Eq.(25) takes the form

$$(i\hbar\gamma^{\bar{\mu}}\partial_{\bar{\mu}} - kI_+ + 2kI)\psi_+(\mathbf{r}, t) = 0.$$

The additional term $2kI$ related to the rest energy violates the covariance of the GLL equation under the parity transformation. In this case, the relations (26) are no longer valid.

However, the independent component $\eta_{1,+}(\mathbf{r}, t)$ can still be characterized by a definite parity, positive or negative, while it is not true for $\eta_{2,+}(\mathbf{r}, t)$ (and for the overall field $\psi_{1,+}(\mathbf{r}, t)$). The component $\eta_{1,+}(\mathbf{r}, t)$ satisfies the Schrödinger equation (19) which obviously remains invariant in form under the parity transformation. So, Eq.(27) is still valid. Using the relation between the components $\eta_{1,+}(\mathbf{r}, t)$ and $\eta_{2,+}(\mathbf{r}, t)$ (see the second equation in (18)), we get

$$\eta_{2,+}(\mathbf{r}, t) = e^{i\delta} \left[-\eta_{2,+}(-\mathbf{r}, t) + 2\mu\eta_{1,+}(-\mathbf{r}, t) \right]$$

which is a generalization of equation (28) for the case of non-zero k . It shows that $\eta_{2,+}(\mathbf{r}, t)$ is not invariant under parity. The parity transformation changes the dependent component into a combination of the dependent and independent ones.

2.2 Connection with the Dirac equation

The rest energy of a free relativistic Fermi field is incorporated nontrivially in its total or relativistic energy. Its rest mass m_0 is at the same time a measure of its inertia, i.e. of the tendency of the field to resist changes in velocity. Thus, m_0 is its inertial mass as well [9].




Figure 1: A schematic illustration of the connection between the relativistic Dirac equation and two non-relativistic equations, LL and GLL: (a) represents the non-relativistic limit $c \rightarrow \infty$; (b) stands for a two-way transformation between the LL and GLL equations.

For non-relativistic Fermi fields, the rest energy contribution decouples from the kinetic energy and becomes an additive factor. In this case, the rest energy is independent from inertia, and the rest mass m_0 is not in general the same as the inertial mass m [10]. The masses m_0 and m are equal to each other only if the theory of non-relativistic Fermi fields is considered to be the limit of the relativistic one when velocities are much smaller than the speed of light.

In the limit $c \rightarrow \infty$, the rest energy becomes infinitely large in value and should be subtracted from the total energy before the non-relativistic limit of the Dirac equation is taken. This is achieved by re-defining the Dirac wave function $\psi_D(\mathbf{r}, t)$ in the way similar to Eq.(2):

$$\psi_D(\mathbf{r}, t) \rightarrow \tilde{\psi}_D(\mathbf{r}, t) \equiv e^{\frac{i}{\hbar} m_0 c^2 t} \psi_D(\mathbf{r}, t). \quad (29)$$

As a result, the non-relativistic limit of the Dirac equation is the LL equation without any rest energy contribution. It is sometimes stated that the rest energy is a purely relativistic effect and has no non-relativistic limit [11].

The transition from the Dirac equation to the GLL equation requires two steps (see Figure 1). First, the non-relativistic limit of the Dirac equation is taken (step (a)). Then, within the non-relativistic formulation, a final amount of rest energy E_0 is introduced by transforming the Fermi fields (step (b)) as

$$\tilde{\psi}_+(\mathbf{r}, t) \rightarrow \psi_+(\mathbf{r}, t) = \left(I + \sqrt{\frac{m_0}{2m}} \gamma^4 \right) e^{-\frac{i}{\hbar} E_0 t} \tilde{\psi}_+(\mathbf{r}, t). \quad (30)$$

This transformation is the inverse of the transformation given by Eq.(16).

3 GLL equation with external potentials

A general form of the external potential can be represented by a 4×4 matrix-valued function $V(\mathbf{r}, t)$, so that the GLL equation becomes

$$(i\hbar\gamma^{\bar{\mu}}\partial_{\bar{\mu}} - kI_+ - V)\psi_+(\mathbf{r}, t) = 0. \quad (31)$$

Using again the field $\tilde{\psi}_+(\mathbf{r}, t)$ given by Eq.(16), we convert Eq.(31) into

$$(i\hbar\gamma^{\bar{\mu}}\partial_{\bar{\mu}} + mc\gamma^5 - \tilde{V})\tilde{\psi}_+(\mathbf{r}, t) = 0, \quad (32)$$

where

$$\tilde{V} \equiv V + \frac{1}{\sqrt{2}}\mu\{\gamma^4, V\} + \frac{1}{2}\mu^2\gamma^4\{\gamma^4, V\} \quad (33)$$

and $\{A, B\} = AB + BA$ is the anticommutator of two operators A and B .

If the external potential obeys the condition

$$\{\gamma^4, V\} = 0, \quad (34)$$

then $\tilde{V} = V$ and the rest energy term is removed from the GLL equation in the way similar to the case of free non-relativistic Fermi fields.

An example of such external potentials is a potential produced by an electromagnetic field $A_{\bar{\mu}}$. It can be introduced by replacing $\partial_{\bar{\mu}}$ with $D_{\bar{\mu}} \equiv \partial_{\bar{\mu}} - \frac{i}{\hbar} \frac{e}{c} A_{\bar{\mu}}$, where e is the charge of the Fermi field. This yields

$$V = -\frac{e}{c}\gamma^{\bar{\mu}}A_{\bar{\mu}}.$$

Since the matrices $\gamma^{\bar{\mu}}$ anti-commute with γ^4 , the condition (34) is satisfied and the rest energy E_0 does not show itself in the solutions of Eq.(32).

The external potentials that include the matrix γ^5 do not obey the condition (34). The simplest example is the potential of the form

$$V = \frac{1}{c}\gamma^5\varphi, \quad (35)$$

where $\varphi(\mathbf{r}, t)$ is a single-valued function that does not have a matrix structure and behaves like a scalar under the Galilean transformations. The transformed potential

$$\tilde{V} = V + \sqrt{2}\frac{\mu}{c}\varphi + \frac{\mu^2}{c}\gamma^4\varphi$$

exhibits the rest energy contribution in its last two terms. It means that for this type of potentials the rest energy will contribute non-trivially to the bound state spectrum of the GLL equation.

Applying the parity transformation to the GLL equation with the external potential given by Eq.(35) in the same way as we did it above to the free GLL equation, we get the following transformation for the function $\varphi(\mathbf{r}, t)$:

$$\varphi(\mathbf{r}, t) \rightarrow \varphi'(\mathbf{r}, t) = \varphi(-\mathbf{r}, t), \quad (36)$$

i.e. $\varphi(\mathbf{r}, t)$ behaves like a scalar under space reflection as well, while V is a pseudoscalar. As before in the case without external potentials, the covariance of the GLL equation is violated for non-zero values of k .

In the component form, Eq.(31) becomes

$$\begin{aligned} i\hbar\partial_4\eta_{1,+} + p_-\eta_{2,+} &= 0, \\ cp_+\eta_{1,+} - (mc^2 - \varphi)\eta_{2,+} &= 0. \end{aligned} \quad (37)$$

Assuming that φ is a smooth function and eliminating $\eta_{2,+}(\mathbf{r}, t)$, this yields for the independent component $\eta_{1,+}(\mathbf{r}, t)$ a Schrödinger-type equation with contributions from the external potential:

$$\begin{aligned} i\hbar\partial_4\eta_{1,+} + \left[\frac{c}{2(mc^2 - \varphi)}(\hbar^2\Delta - k^2) + \frac{1}{2} \frac{c\hbar^2\sigma^a\partial_a\varphi}{(mc^2 - \varphi)^2}\sigma^b\partial_b \right. \\ \left. + \frac{k}{2} \frac{c\hbar\sigma^a\partial_a\varphi}{(mc^2 - \varphi)^2} \right] \eta_{1,+} = 0. \end{aligned} \quad (38)$$

This equation is not invariant under the parity transformation unless the potential φ is constant in space. The last term in Eq.(38) changes its sign under $\mathbf{r} \rightarrow -\mathbf{r}$, while other terms remain unchanged. This means that in smooth pseudoscalar potentials of the form given by Eq.(35) even the independent component $\eta_{1,+}(\mathbf{r}, t)$ cannot be characterized by a definite parity.

3.1 Spherically symmetric pseudoscalar potential

Let us assume that the potential (35) is time-independent and spherically symmetric, i.e. $\varphi = \varphi(r)$, where $r = |\mathbf{r}|$. Representing $\eta_{1,+}(\mathbf{r}, t)$ as

$$\eta_{1,+}(\mathbf{r}, t) = \eta_{1,+}(\mathbf{r})e^{-\frac{i}{\hbar}Et}$$

and introducing the dimensionless variables

$$\boldsymbol{\xi} \equiv \frac{mc}{\hbar}\mathbf{r}, \quad \mathcal{E} \equiv \frac{E}{mc^2}, \quad \bar{\varphi} \equiv \frac{\varphi}{mc^2},$$

we rewrite Eq.(38) as the eigenvalue equation

$$\mathcal{H}\chi(\mathbf{r}) = \mathcal{E}\chi(\mathbf{r}) \quad (39)$$

with

$$\chi(\mathbf{r}) \equiv \frac{1}{\sqrt{1-\bar{\varphi}}}\eta_{1,+}(\mathbf{r}),$$

and the Hamiltonian density \mathcal{H} given by

$$\begin{aligned} \mathcal{H} \equiv & \frac{1}{2} \frac{1}{1-\bar{\varphi}} (\mathbf{p}_\xi^2 + 2\mu^2) + \frac{3}{8} \frac{1}{(1-\bar{\varphi})^3} (\nabla_\xi \bar{\varphi})^2 + \frac{1}{4} \frac{1}{(1-\bar{\varphi})^2} \nabla_\xi^2 \bar{\varphi} \\ & + \frac{1}{2} \frac{1}{(1-\bar{\varphi})^2} \boldsymbol{\sigma} [\nabla_\xi \bar{\varphi} \times \mathbf{p}_\xi] - \frac{\mu}{\sqrt{2}} \frac{(\boldsymbol{\sigma} \nabla_\xi \bar{\varphi})}{(1-\bar{\varphi})^2}, \end{aligned} \quad (40)$$

where $\nabla_\xi \equiv (\hbar/mc)\nabla = (\hbar/mc)(\partial_1, \partial_2, \partial_3)$, and $\mathbf{p}_\xi = -i\nabla_\xi$ is the momentum operator.

We observe here two well-known terms: a Darwin type term $(1/4(1-\bar{\varphi})^2)\nabla_\xi^2 \bar{\varphi}$ and the spin-orbit coupling,

$$\frac{1}{2} \frac{1}{(1-\bar{\varphi})^2} \boldsymbol{\sigma} [\nabla_\xi \bar{\varphi} \times \mathbf{p}_\xi] = \frac{1}{(1-\bar{\varphi})^2} \frac{1}{\xi} \frac{d\bar{\varphi}}{d\xi} (\mathbf{S}\mathbf{L}_\xi),$$

where $\mathbf{L}_\xi = [\boldsymbol{\xi} \times \mathbf{p}_\xi]$ and $\mathbf{S} = (1/2)\boldsymbol{\sigma}$ are the angular momentum and spin operators, respectively.

We also see terms representing the rest energy contribution. The term $\mu^2/(1-\bar{\varphi})$ is present for all types of the potential $\bar{\varphi}$ and reduces to a shift of the energy spectrum as a whole by μ^2 if the potential is taken equal to zero. The term

$$-\frac{\mu}{\sqrt{2}} \frac{\boldsymbol{\sigma} \nabla_\xi \bar{\varphi}}{(1-\bar{\varphi})^2} = \boldsymbol{\sigma} \mathbf{B}_{\text{eff}} \quad (41)$$

describes the interaction of the non-relativistic Fermi field with an effective background field

$$\mathbf{B}_{\text{eff}} \equiv -\frac{\mu}{\sqrt{2}} \frac{1}{(1-\bar{\varphi})^2} \frac{d\bar{\varphi}}{d\xi} \mathbf{n} \quad (42)$$

where $\mathbf{n} \equiv \boldsymbol{\xi}/\xi$, and results, as we mentioned above, in the violation of parity. This interaction introduces non-central forces, i.e. forces depending on the direction of $\boldsymbol{\xi}$ with respect to \mathbf{S} . The term (41) depends linearly on $(\boldsymbol{\sigma}\mathbf{n})$ and violates the time reversal symmetry as well.

As in the case of the Dirac equation, the angular momentum and spin are not constants of motion, and it is the total angular momentum $\mathbf{J} = \mathbf{L}_\xi + \mathbf{S}$ that is conserved, i.e. $[\mathbf{J}, \mathcal{H}]_- = 0$. For the Dirac equation with spherically symmetric potentials, there is another constant of motion which in the non-relativistic limit takes the form [12]

$$\mathcal{K} = 1 + \boldsymbol{\sigma}\mathbf{L}_\xi. \quad (43)$$

Evaluating its commutator with the Hamiltonian \mathcal{H} , we obtain

$$[\mathcal{K}, \mathcal{H}]_- = -2(\boldsymbol{\sigma}\mathbf{B}_{\text{eff}})\mathcal{K}. \quad (44)$$

This is another effect of rest energy: due to the presence of the background field, \mathcal{K} is not conserved in the case of the GLL equation.

3.2 Pseudoscalar spherical well of finite depth

For the spherical well potential

$$\bar{\varphi}(\xi) = \begin{cases} -\bar{\varphi}_0 & \text{for } \xi < \xi_0 \\ 0 & \text{for } \xi > \xi_0, \end{cases} \quad (45)$$

where $\bar{\varphi}_0 > 0$, the parity violating term (41) as well as the spin-orbit coupling and Darwin type terms vanish both inside and outside the well.

The Hamiltonian density given by Eq.(40) becomes

$$\mathcal{H} = \begin{cases} \frac{1}{2} \frac{1}{1+\bar{\varphi}_0} (\mathbf{p}_\xi^2 + 2\mu^2) & \text{for } \xi < \xi_0 \\ \frac{1}{2} (\mathbf{p}_\xi^2 + 2\mu^2) & \text{for } \xi > \xi_0. \end{cases} \quad (46)$$

The set of operators commuting with \mathcal{H} is then $\mathbf{L}^2, \mathbf{L}_3$, so the solutions of the eigenvalue equation (39) have the form

$$\chi_{lM} = \frac{1}{\xi} R_l Y_l^M, \quad (47)$$

where $(1/\xi)R_l$ and Y_l^M are their radial and angular parts, respectively, Y_l^M being spherical harmonics.

The quantum numbers l, M are defined as

$$\begin{aligned} \mathbf{L}^2 \chi_{lM} &= l(l+1) \chi_{lM}, \\ \mathbf{L}_3 \chi_{lM} &= M \chi_{lM}. \end{aligned}$$

Substituting the ansatz (47) into the eigenvalue equation (39) and using the expression for \mathbf{p}_ξ^2 in spherical polar coordinates,

$$\mathbf{p}_\xi^2 = -\frac{\partial^2}{\partial \xi^2} - \frac{2}{\xi} \frac{\partial}{\partial \xi} + \frac{1}{\xi^2} \mathbf{L}_\xi^2,$$

we bring the radial part of the eigenvalue equation to the form

$$\frac{d^2 R_l}{d\xi^2} + \left(a^2 - \frac{l(l+1)}{\xi^2} \right) R_l = 0 \quad \text{for } \xi < \xi_0 \quad (48)$$

and

$$\frac{d^2 R_l}{d\xi^2} + \left(-b^2 - \frac{l(l+1)}{\xi^2} \right) R_l = 0 \quad \text{for } \xi > \xi_0, \quad (49)$$

where we have introduced the abbreviations

$$a^2 \equiv 2(1 + \bar{\varphi}_0) \left[\frac{\mu^2 \bar{\varphi}_0}{1 + \bar{\varphi}_0} - |\bar{\mathcal{E}}| \right], \quad b^2 \equiv 2|\bar{\mathcal{E}}|, \quad \bar{\mathcal{E}} \equiv \mathcal{E} - \mu^2. \quad (50)$$




Figure 2: *The effective potential well of the radial Schrödinger equation derived from the GLL equation with the spherical well potential.*

We recognize in Eqs.(48) and (49) the radial part of the Schrödinger equation with the potential illustrated in Figure 2.

We are interested in the bound state solutions for which $|\bar{\mathcal{E}}| < \mu^2\bar{\varphi}_0/(1 + \bar{\varphi}_0)$. The solutions are [15],[16]

$$R_l(\xi) = \begin{cases} N_- j_l(a\xi) & \text{for } \xi < \xi_0 \\ N_+ h_l^{(1)}(ib\xi) & \text{for } \xi > \xi_0, \end{cases} \quad (51)$$

where N_-, N_+ are normalization constants, while j_l and $h_l^{(1)}$ are spherical Bessel and Hankel functions, respectively. At $\xi = \xi_0$, both $R_l(\xi)$ and its first derivative should be continuous. The normalization constants can be eliminated from the continuity relation of the logarithmic derivative. This yields the condition

$$ib \frac{h_l^{(1)'}(ib\xi_0)}{h_l^{(1)}(ib\xi_0)} = a \frac{j_l'(a\xi_0)}{j_l(a\xi_0)}, \quad (52)$$

where the primes denote differentiations to the respective arguments. This is the transcendental equation for the eigenvalue $\bar{\mathcal{E}}$. The quantum number M does not affect the eigenvalues, so each energy level is $(2l + 1)$ -fold degenerate.

For $l = 0$, i.e. for S-states, the condition (52) reduces to

$$\rho \cot(\bar{\xi}_0 \rho) = -\sqrt{\frac{1 - \rho^2}{1 + \bar{\varphi}_0}}, \quad (53)$$

where

$$\rho \equiv \sqrt{1 - \frac{1 + \bar{\varphi}_0}{\mu^2 \bar{\varphi}_0} |\bar{\mathcal{E}}|}, \quad \bar{\xi}_0 \equiv \sqrt{2\mu^2 \bar{\varphi}_0} \xi_0.$$

The bound states spectrum is given by

$$|\bar{\mathcal{E}}| = \frac{\mu^2 \bar{\varphi}_0}{1 + \bar{\varphi}_0} (1 - \rho^2), \quad 0 < \rho \leq 1. \quad (54)$$

	1	2	3	4
$\frac{\mu^2 \bar{\varphi}_0}{1+\bar{\varphi}_0}$	0.0089	0.0133	0.0089	0.0094
ρ	0.5184	0.4420 0.9768	0.2985 0.6370 0.9882	0.3853 0.8843

Figure 3: The values of depth of the effective potential well and the solutions for ρ for various sets of parameters $(\bar{\varphi}_0, \xi_0, \mu^2)$: (1) $\bar{\varphi}_0 = 8, \xi_0 = 10, \mu^2 = 0.01$; (2) $\bar{\varphi}_0 = 8, \xi_0 = 10, \mu^2 = 0.015$; (3) $\bar{\varphi}_0 = 8, \xi_0 = 20, \mu^2 = 0.01$; (4) $\bar{\varphi}_0 = 15, \xi_0 = 10, \mu^2 = 0.01$.




Figure 4: A sketch of the effective potential well with its respective S -states energy levels for the sets of parameters from Figure 3.

The values of ρ satisfying Eq.(53) can be found graphically [15]. The number of bound states depends on the depth and width of the original well potential given by Eq.(45), i.e. on $\bar{\varphi}_0$ and ξ_0 , as well as on the rest energy represented by the parameter μ^2 . As we can see from Figures 3 and 4, the larger μ^2 , the deeper the effective potential well and the more energy levels in it. The number of energy levels also increases with increasing either $\bar{\varphi}_0$ or ξ_0 or both of them.

It can be seen graphically that only when

$$(a^2 + b^2)\bar{\xi}_0^2 = \frac{2\mu^2\bar{\xi}_0^2\bar{\varphi}_0}{1 + \bar{\varphi}_0}(1 + \bar{\varphi}_0\rho^2) \geq \frac{\pi^2}{4},$$

or

$$\mu^2 \geq \mu_{\min}^2 \equiv \frac{\pi}{4\bar{\varphi}_0\xi_0},$$

can Eq.(53) have a solution. For given values of $\bar{\varphi}_0$ and ξ_0 , μ_{\min}^2 is the minimum non-zero value of μ^2 to achieve the bound S-states. In dimensional variables, this corresponds to the following minimum value of the rest energy:

$$E_{0,\min} \equiv \frac{\pi mc^3\hbar}{4\varphi_0 r_0},$$

where $r_0 = (\hbar/mc)\xi_0$.

For $\mu^2 = 0$, the effective potential well and bound states disappear. Let us demonstrate this for S-states. In this case, the radial part of the eigenvalue equation becomes

$$\frac{d^2 R_0}{d\xi^2} + 2\mathcal{E}(1 + \bar{\varphi}_0)R_0 = 0 \quad \text{for} \quad \xi < \xi_0$$

and

$$\frac{d^2 R_0}{d\xi^2} + 2\mathcal{E}R_0 = 0 \quad \text{for} \quad \xi > \xi_0,$$

where $\mathcal{E} < 0$. The solution satisfying the boundary condition $R_0(0) = 0$ and decreasing exponentially for large values of ξ is

$$R_0(\xi) = \begin{cases} N_- (e^{-\sqrt{2(1+\bar{\varphi}_0)|\mathcal{E}|\xi}} - e^{\sqrt{2(1+\bar{\varphi}_0)|\mathcal{E}|\xi}}) & \text{for} \quad \xi < \xi_0 \\ N_+ e^{-\sqrt{2|\mathcal{E}|\xi}} & \text{for} \quad \xi > \xi_0. \end{cases}$$

Imposing the condition of continuity of R_0 and its derivative at $\xi = \xi_0$ and eliminating the normalization constants we get the equation for the bound state eigenvalues:

$$e^{2\sqrt{2(1+\bar{\varphi}_0)|\mathcal{E}|\xi_0}} = \frac{1 - \sqrt{1 + \bar{\varphi}_0}}{1 + \sqrt{1 + \bar{\varphi}_0}}.$$

Since its right-hand side is negative, the equation has no solution.

3.3 Pseudoscalar Coulomb potential

The non-central forces mix the radial and angular parts of eigenfunctions and make explicit calculations for a general form of spherically symmetric pseudoscalar potential rather complicated. In analyzing the effects of the spin orientation it is convenient to use the helicity operator $\mathbf{S}\mathbf{n}$.

The helicity is rotational invariant and commutes with total angular momentum, $[\mathbf{S}\mathbf{n}, \mathbf{J}]_- = 0$. Its commutator with the Hamiltonian \mathcal{H} is

$$[\mathbf{S}\mathbf{n}, \mathcal{H}]_- = \frac{1}{1-\bar{\varphi}} \frac{1}{\xi} \left[\frac{1}{(1-\bar{\varphi})} \frac{d\bar{\varphi}}{d\xi} - \frac{1}{\xi} \right] (\mathbf{S}\mathbf{n})\mathcal{K}. \quad (55)$$

Let us define the operator

$$\tilde{\mathcal{K}} \equiv \mathcal{K} + \tau(\mathbf{S}\mathbf{n}), \quad (56)$$

where τ is a constant. If we impose the condition

$$[\tilde{\mathcal{K}}, \mathcal{H}]_- = 0, \quad (57)$$

then $\tilde{\mathcal{K}}$ becomes an additional constant of motion. Using the commutators (44) and (55), we rewrite this condition as

$$\frac{1}{1-\bar{\varphi}} \left[2\mu\sqrt{2} + \frac{\tau}{\xi} \right] \frac{d\bar{\varphi}}{d\xi} - \frac{\tau}{\xi^2} = 0.$$

It is solved by

$$\bar{\varphi}(\xi) = -\frac{\bar{q}^2}{\xi}, \quad (58)$$

provided $\tau = 2\mu\bar{q}^2\sqrt{2}$, where

$$\bar{q}^2 \equiv Z \frac{e^2}{c\hbar}$$

is a dimensionless parameter, and Ze is the magnitude of the external charge creating the potential.

The operator $\tilde{\mathcal{K}}$ is similar to the Biedenharn-Temple operator for the Dirac-Coulomb problem [17],[18]; see also [19]. It can be brought to the original form

$$\tilde{\mathcal{K}} = 1 + \boldsymbol{\sigma}\tilde{\mathbf{L}}_\xi$$

(compare to Eq.(43)), if a "non-integer" orbital angular momentum is introduced as

$$\tilde{\mathbf{L}}_\xi \equiv \mathbf{L}_\xi + \mu\bar{q}^2\sqrt{2}\mathbf{n}.$$

With the potential (58), the Hamiltonian density \mathcal{H} given by Eq.(40) takes the form

$$\begin{aligned} \mathcal{H} \equiv & -\frac{1}{2} \frac{1}{\xi + \bar{q}^2} \left(\xi \frac{\partial^2}{\partial \xi^2} + 2 \frac{\partial}{\partial \xi} \right) + \frac{1}{2} \frac{1}{\xi(\xi + \bar{q}^2)} (\tilde{\mathcal{K}}^2 - \tilde{\mathcal{K}} - 2\bar{q}^4\mu^2) \\ & + \frac{\xi}{\xi + \bar{q}^2} \mu^2 + \frac{3}{8} \frac{\bar{q}^4}{\xi(\xi + \bar{q}^2)^3} + \frac{1}{2} \frac{\bar{q}^2}{\xi(\xi + \bar{q}^2)^2} (\tilde{\mathcal{K}} - 1). \end{aligned} \quad (59)$$

Its angular dependence is expressed in terms of $\tilde{\mathcal{K}}$, and this means that the separation of the radial and angular parts of eigenfunctions is still possible.

3.3.1 Angular eigenfunctions

Let us first discuss the structure of eigenfunctions in the case $\mu = 0$. Then the set of operators commuting with \mathcal{H} is $\mathbf{J}^2, \mathbf{J}_3, \mathcal{K}$, so the solutions of the eigenvalue equation (39) have the form

$$\chi_{jM}^\kappa = \frac{1}{\xi} R_\kappa \mathcal{Y}_{jM}^\kappa, \quad (60)$$

$(1/\xi)R_\kappa$ and \mathcal{Y}_{jM}^κ being their radial and angular parts, respectively. The quantum numbers j, M, κ are defined as

$$\begin{aligned} \mathbf{J}^2 \chi_{jM}^\kappa &= j(j+1) \chi_{jM}^\kappa, \\ \mathbf{J}_3 \chi_{jM}^\kappa &= M \chi_{jM}^\kappa, \\ \mathcal{K} \chi_{jM}^\kappa &= -\kappa \chi_{jM}^\kappa. \end{aligned}$$

Furthermore, \mathbf{S}^2 and \mathbf{L}_ξ^2 are also constants of motion but they are not reflected in the ansatz (60) since \mathbf{S}^2 always has eigenvalue 3/4, while \mathbf{L}_ξ^2 is expressed in terms of \mathcal{K} : $\mathbf{L}_\xi^2 = \mathcal{K}^2 - \mathcal{K}$. In particular, the eigenfunctions (60) have a definite value of orbital angular momentum quantum number l as well, and this value is fixed uniquely by the parity.

The minimum value of $|\kappa|$ is 1, so the eigenfunctions are divided into two sets corresponding to positive and negative values of κ [13],[14]. For $\kappa < 0$, $\kappa = -(l+1)$, and for odd values of $|\kappa|$ the eigenfunctions have positive parity, while for even values of $|\kappa|$ the parity is negative. For $\kappa > 0$, $\kappa = l$, and the situation with the parity is opposite: the parity of the eigenfunctions is positive for even values of $|\kappa|$ and negative for odd ones. For both sets, $|\kappa| = j + 1/2$. The sign of κ determines whether the orthogonal projection of $\boldsymbol{\sigma}$ on \mathbf{J} is parallel ($\kappa < 0$) or antiparallel ($\kappa > 0$) to \mathbf{J} .

The normalized angular parts of the eigenfunctions are

$$\mathcal{Y}_{jM}^{-|\kappa|} = \sqrt{\frac{j+M}{2j}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} Y_{j-\frac{1}{2}}^{M-\frac{1}{2}} + \sqrt{\frac{j-M}{2j}} \begin{pmatrix} 0 \\ 1 \end{pmatrix} Y_{j-\frac{1}{2}}^{M+\frac{1}{2}} \quad (61)$$

for $\kappa < 0$ and

$$\mathcal{Y}_{jM}^{|\kappa|} = -\sqrt{\frac{j-M+1}{2j+2}} \begin{pmatrix} 1 \\ 0 \end{pmatrix} Y_{j+\frac{1}{2}}^{M-\frac{1}{2}} + \sqrt{\frac{j+M+1}{2j+2}} \begin{pmatrix} 0 \\ 1 \end{pmatrix} Y_{j+\frac{1}{2}}^{M+\frac{1}{2}} \quad (62)$$

for $\kappa > 0$, where $\begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ are "spin-up" and "spin-down" column matrices, and Y_j^M are spherical harmonics.

The angular parts are related as follows

$$(\boldsymbol{\sigma} \mathbf{n}) \mathcal{Y}_{jM}^{\pm|\kappa|} = -\mathcal{Y}_{jM}^{\mp|\kappa|}, \quad (63)$$

i.e. the eigenfunctions of opposite parities are connected by the helicity operator.

For non-zero values of μ , the set of operators commuting with \mathcal{H} becomes $\mathbf{J}^2, \mathbf{J}_3, \tilde{\mathcal{K}}$, and we use the following ansatz for eigenfunctions:

$$\chi_{jM}^{\tilde{\kappa}} = \frac{1}{\xi} R_{\tilde{\kappa}} \mathcal{Y}_{jM}^{\tilde{\kappa}}, \quad (64)$$

where the quantum numbers $j, M, \tilde{\kappa}$ are defined as

$$\begin{aligned} \mathbf{J}^2 \chi_{jM}^{\tilde{\kappa}} &= j(j+1) \chi_{jM}^{\tilde{\kappa}}, \\ \mathbf{J}_3 \chi_{jM}^{\tilde{\kappa}} &= M \chi_{jM}^{\tilde{\kappa}}, \\ \tilde{\mathcal{K}} \chi_{jM}^{\tilde{\kappa}} &= -\tilde{\kappa} \chi_{jM}^{\tilde{\kappa}}. \end{aligned}$$

Since the parity is violated, \mathbf{L}_ξ^2 is not a constant of motion, and the eigenfunctions (64) are not characterized by a definite value of l .

The helicity operator anti-commutes with \mathcal{K} , $(\mathbf{S}\mathbf{n})\mathcal{K} + \mathcal{K}(\mathbf{S}\mathbf{n}) = 0$. This leads us to the following expression for $\tilde{\mathcal{K}}^2$:

$$\tilde{\mathcal{K}}^2 = \mathcal{K}^2 + 2\mu^2 \bar{q}^4.$$

As before in the case of $\mu = 0$, the eigenfunctions are divided into two sets corresponding to positive and negative (irrational) values of $\tilde{\kappa}$:

$$\tilde{\kappa} = \pm \sqrt{\kappa^2 + 2\mu^2 \bar{q}^4} = \pm \sqrt{\left(j + \frac{1}{2}\right)^2 + 2\mu^2 \bar{q}^4}.$$

The minimum value of $|\tilde{\kappa}|$ is $\sqrt{1 + 2\mu^2 \bar{q}^4}$. The sign of $\tilde{\kappa}$ is determined by the spin orientation with respect to both \mathbf{J} and \mathbf{n} : $\tilde{\kappa} > 0$ if $(\boldsymbol{\sigma}\tilde{\mathbf{J}}) < 1/2$ and $\tilde{\kappa} < 0$ if $(\boldsymbol{\sigma}\tilde{\mathbf{J}}) > 1/2$, where

$$\tilde{\mathbf{J}} \equiv \mathbf{J} + \mu \bar{q}^2 \sqrt{2} \mathbf{n}$$

is the irrational total angular momentum.

The angular parts of the eigenfunctions are constructed as linear combinations of the ones for $\mu = 0$:

$$\mathcal{Y}_{jM}^{-|\tilde{\kappa}|} = \alpha_j \mathcal{Y}_{jM}^{-|\kappa|} - \beta_j \mathcal{Y}_{jM}^{|\kappa|} \quad \text{for} \quad \tilde{\kappa} < 0 \quad (65)$$

and

$$\mathcal{Y}_{jM}^{|\tilde{\kappa}|} = \alpha_j \mathcal{Y}_{jM}^{|\kappa|} + \beta_j \mathcal{Y}_{jM}^{-|\kappa|} \quad \text{for} \quad \tilde{\kappa} > 0, \quad (66)$$

where the coefficients α_j and β_j obey the conditions

$$\begin{aligned} (|\tilde{\kappa}| - |\kappa|)\alpha_j - \mu \bar{q}^2 \sqrt{2} \beta_j &= 0, \\ (|\tilde{\kappa}| + |\kappa|)\beta_j - \mu \bar{q}^2 \sqrt{2} \alpha_j &= 0, \end{aligned} \quad (67)$$

and $\mathcal{Y}_{jM}^{-|\kappa|}$ and $\mathcal{Y}_{jM}^{|\kappa|}$ are given by Eqs.(61) and (62).

The normalization conditions for the eigenfunctions (65) and (66) yield

$$\alpha_j^2 + \beta_j^2 = 1. \quad (68)$$

Solving Eqs.(67) and (68), we get

$$\alpha_j \equiv \frac{1}{\sqrt{2}} \left(1 + \frac{|\kappa|}{|\tilde{\kappa}|} \right)^{1/2}, \quad \beta_j \equiv \frac{1}{\sqrt{2}} \left(1 - \frac{|\kappa|}{|\tilde{\kappa}|} \right)^{1/2}.$$

The angular parts of the eigenfunctions from the two sets are related as

$$(\boldsymbol{\sigma n})\mathcal{Y}_{jM}^{\pm|\tilde{\kappa}|} = \mp \frac{1}{|\tilde{\kappa}|} \mu \bar{q}^2 \sqrt{2} \mathcal{Y}_{jM}^{\pm|\tilde{\kappa}|} - \frac{|\kappa|}{|\tilde{\kappa}|} \mathcal{Y}_{jM}^{\mp|\tilde{\kappa}|}.$$

These relations reduce to those in Eq.(63) when either μ or the external charge vanish.

The angular dependence of wave functions determines the conditions when the transitions between given states are possible. For the case $\mu = 0$, these conditions are parity selection rules which forbid transitions between any two states of opposite parities. The violation of parity for non-zero values of μ enlarges the number of possible transitions.

This has an immediate impact on the properties of the system. If the transitions are caused by a rotation, then the allowed ones are those for which the matrix elements of the components of the total angular momentum are non-zero (see Appendix B). In the representation in which \mathbf{J}^2 and J_3 are diagonal, the matrix elements of J_3 do not depend on our choice of μ . All matrix elements for transitions between the states of different j and M vanish

For the matrix elements of J_1 and J_2 , the situation is different. For $\mu = 0$, the transitions between two states within the set of positive (or negative) κ states are allowed if the states have the same value of j , while the values of M differ by one unit:

$$\Delta j = 0, \quad \Delta M = \pm 1.$$

The transitions between the sets are allowed if both j and M change by one unit:

$$\Delta j = \pm 1, \quad \Delta M = \pm 1.$$

The effects of non-zero values of μ are two-fold. The selection rules for both type of transitions become

$$\Delta j = 0, \pm 1, \quad \Delta M = \pm 1,$$

i.e. the transitions, which were forbidden in the case $\mu = 0$, are now allowed. In addition, the matrix elements become μ -dependent.

3.3.2 Radial eigenfunctions

Substituting the ansatz (64) into the eigenvalue equation (39), we get its radial part in the form of the Schrödinger equation

$$\frac{d^2 R_{\tilde{\kappa}}}{d\xi^2} + 2 \left(\bar{\mathcal{E}} - V_{\text{eff}}^{\pm} \right) R_{\tilde{\kappa}} = 0 \quad (69)$$

with the energy-dependent effective potentials

$$\begin{aligned} V_{\text{eff}}^{\pm} &= \left[-\mathcal{E}\bar{q}^2 \pm \frac{1}{2\bar{q}^2} \left(|\tilde{\kappa}| \mp \frac{1}{2} \right) \right] \frac{1}{\xi} + \frac{j(j+1)}{2\xi^2} \\ &+ \frac{3}{8} \frac{1}{(\xi + \bar{q}^2)^2} \mp \frac{1}{2} \frac{1}{\bar{q}^2(\xi + \bar{q}^2)} \left(|\tilde{\kappa}| \mp \frac{1}{2} \right), \end{aligned}$$

the upper signs referring to the case $\tilde{\kappa} > 0$ and the lower ones to the case $\tilde{\kappa} < 0$.

The energy-dependent effective potential appears here in the same way as in the case of the Dirac equation reduced to the Pauli-Schrödinger one [20],[21]. Relativistic wave equations with energy-dependent potential were studied in [22]-[27].

Let us assume that the external charge is very large, $\bar{q}^2 \gg 1/\mu$. Then, expanding $|\tilde{\kappa}|$ in powers of $1/\bar{q}^2$, we can represent it as

$$|\tilde{\kappa}| = \mu\sqrt{2}\bar{q}^2 + \left[\sqrt{\kappa^2 + 2\mu^2\bar{q}^4} - \mu\sqrt{2}\bar{q}^2 \right],$$

where the expression in the square brackets only includes terms of the power $1/\bar{q}^2$ and higher. This allows us to rewrite the potentials V_{eff}^{\pm} as

$$V_{\text{eff}}^{\pm} = V_0^{\pm} + V_{\text{pert}}^{\pm},$$

where

$$V_0^{\pm} \equiv \frac{C_1^{\pm}}{\xi} + \frac{j(j+1)}{2\xi^2}$$

with

$$C_1^{\pm} \equiv -\mathcal{E}\bar{q}^2 \pm \frac{\mu}{\sqrt{2}},$$

and

$$V_{\text{pert}}^{\pm} = \pm \left[\frac{1}{2\bar{q}^2} \left(|\tilde{\kappa}| \mp \frac{1}{2} \right) - \frac{\mu}{\sqrt{2}} \right] \frac{1}{\xi} + \frac{3}{8} \frac{1}{(\xi + \bar{q}^2)^2} \mp \frac{1}{2} \frac{1}{\bar{q}^2(\xi + \bar{q}^2)} \left(|\tilde{\kappa}| \mp \frac{1}{2} \right)$$

are perturbations.

With V_{pert}^{\pm} neglected, Eq.(69) becomes exactly solvable. In this approximation, the solution in arbitrary normalization is

$$R_{\tilde{\kappa}} = e^{-b_{\pm}\xi} \xi^{\lambda} {}_1F_1 \left(\lambda + \frac{C_1^{\pm}}{b_{\pm}}, 2\lambda; 2b_{\pm}\xi \right), \quad (70)$$

where $\lambda \equiv |\kappa| + 1/2$, while b_{\pm} for the positive and negative $\tilde{\kappa}$ bound states are defined in the same way as b in Eq.(50). The function $R_{\tilde{\kappa}}$ vanishes for $\xi \rightarrow \infty$ and becomes normalizable if the confluent series ${}_1F_1$ breaks off. This happens when

$$\lambda + \frac{C_1^{\pm}}{b_{\pm}} = -n, \quad n = 0, 1, 2, \dots \quad (71)$$

This relation indicates that C_1^{\pm} should be negative. The condition $C_1^{\pm} < 0$, in turn, yields energy intervals where bound states exist:

$$|\bar{\mathcal{E}}| < \bar{\mathcal{E}}_- \equiv \mu^2 + \frac{\mu}{\bar{q}^2 \sqrt{2}} \quad (72)$$

for the negative $\tilde{\kappa}$ bound states, and

$$|\bar{\mathcal{E}}| < \bar{\mathcal{E}}_+ \equiv \mu^2 - \frac{\mu}{\bar{q}^2 \sqrt{2}} \quad (73)$$

for the positive $\tilde{\kappa}$ bound states.

Eq.(71) leads to the energy levels

$$\bar{\mathcal{E}}_{n,j}^{\pm} \equiv -\frac{(\lambda + n)^2}{2\bar{q}^4} \left[\sqrt{1 + \frac{2\bar{q}^4 \bar{\mathcal{E}}_{\pm}}{(\lambda + n)^2}} - 1 \right]^2, \quad (74)$$

the subscripts indicating that the energy levels depend on two quantum numbers, n and j . The energy dependence of the effective potential modifies the scalar product [26]. For a given value of $\tilde{\kappa}$, the orthonormality condition for the radial parts of eigenfunctions reads (see Appendix C)

$$\begin{aligned} (R_{n,\tilde{\kappa}}^*, R_{n',\tilde{\kappa}}) &\equiv \int_0^{\infty} d\xi R_{n,\tilde{\kappa}}^* \left(1 - \frac{\partial V_{\text{eff}}}{\partial \mathcal{E}} \right) R_{n',\tilde{\kappa}} \\ &= \int_0^{\infty} d\xi R_{n,\tilde{\kappa}}^* \left(1 + \frac{\bar{q}^2}{\xi} \right) R_{n',\tilde{\kappa}} = \delta_{nn'}. \end{aligned} \quad (75)$$

For the energy levels with $n = 0$ and $j = 1/2$, the normalized radial parts of their eigenfunctions are (see Figure 5)

$$R_{0,1/2}^{\pm} = 4\sqrt{2} |\bar{\mathcal{E}}_{0,1/2}^{\pm}| \left(3 + 2\bar{q}^2 \sqrt{2|\bar{\mathcal{E}}_{0,1/2}^{\pm}|} \right)^{-1/2} \exp\{-\sqrt{2|\bar{\mathcal{E}}_{0,1/2}^{\pm}|} \xi\} \xi^{3/2}.$$

The values of $|\bar{\mathcal{E}}_{0,1/2}^{\pm}|$ represent the energies required to destroy the bond between the Fermi field and the charge q . We have used here a slightly different notation for the radial parts of eigenfunctions: $R_{n,j}^+$ and $R_{n,j}^-$ for positive and negative values of $\tilde{\kappa}$, respectively, instead of $R_{n,\tilde{\kappa}}$ for all possible values of $\tilde{\kappa}$.

The potentials V_0^{\pm} (see Figure 6) have the form of Kratzer's diatomic molecule potential. They are made up of the Coulomb potential, which varies as $(-1/\xi)$, and




Figure 5: The radial part of eigenfunction for the positive $\tilde{\kappa}$ bound states with $n = 0$, $j = \frac{1}{2}$ and $\tilde{q}^2 = 100$. The larger μ , the closer the Fermi particle to the location of the charge q . The radial part of eigenfunction for the negative $\tilde{\kappa}$ bound states has a similar shape.




Figure 6: The potential V_0^+ for the positive $\tilde{\kappa}$ bound states with $n = 0$, $j = \frac{1}{2}$ and $\tilde{q}^2 = 100$. The larger μ , the deeper the minimum. The potential V_0^- for the negative $\tilde{\kappa}$ bound states has a similar shape.

the centripetal one, which varies as $1/\xi^2$. The centripetal potential dominates at small values of ξ . As a result, the potentials V_0^\pm are repulsive at short distances for $j \geq 1/2$ and are more repulsive the greater the value of j .

Figure 6 shows that the Fermi field and the charge q attract each other at large values of ξ , pass through a state of minimum energy and begin to repel at small distances. The minima are

$$V_0^\pm(\xi_{0,\pm}) = -\frac{1}{2} \frac{\bar{q}^4 (\bar{\mathcal{E}}_\pm - |\bar{\mathcal{E}}|)^2}{j(j+1)}$$

at

$$\xi_{0,\pm} = \frac{j(j+1)}{\bar{q}^2 (\bar{\mathcal{E}}_\pm - |\bar{\mathcal{E}}|)}.$$

For the same value of $\bar{\mathcal{E}}$ and the same parameter set (μ^2, \bar{q}^2) , the minima of the potentials for $j > 1/2$ are farther from the location of the charge q than the ones for $j = 1/2$. In dimensional variables, the location of the minima is at

$$r_{0,\pm} = r_b \frac{j(j+1)}{Z(\bar{\mathcal{E}}_\pm - |\bar{\mathcal{E}}|)},$$

where $r_b \equiv \frac{\hbar^2}{me^2}$. In the case of electron, r_b is the Bohr radius, $r_b = 5.29 \times 10^{-9}$ cm.

If we expand the potentials about their minima as

$$V_0^\pm(\xi) = V_0^\pm(\xi_{0,\pm}) + \frac{1}{2} \omega_\pm^2 (\xi - \xi_{0,\pm})^2 + \dots,$$

this yields the classical frequencies for small harmonic vibrations of the Fermi field

$$\omega_\pm = \frac{\bar{q}^4 (\bar{\mathcal{E}}_\pm - |\bar{\mathcal{E}}|)^2}{[j(j+1)]^{3/2}}. \quad (76)$$

The dimensional frequencies are

$$w_\pm \equiv \frac{mc^2}{\hbar} \omega_\pm = \nu_0 Z^2 \frac{(\bar{\mathcal{E}}_\pm - |\bar{\mathcal{E}}|)^2}{[j(j+1)]^{3/2}},$$

where $\nu_0 \equiv \frac{me^4}{\hbar^3}$ is the Coulomb units for frequency [28]. For electron, $\nu_0 = 4.13 \times 10^{16}$ Hz.

For the bound states corresponding to the energy levels given by Eq.(74), the frequencies take the form

$$w_{n,j}^\pm = \nu_0 \frac{Z^2}{\bar{q}^4} \frac{2(\lambda+n)^2}{[j(j+1)]^{3/2}} |\bar{\mathcal{E}}_{n,j}^\pm|. \quad (77)$$

For $\bar{q}^2 = 100$, the values of $w_{0,\frac{1}{2}}^+$ are $1.64\nu_0 Z^2 \times 10^{-4}$ for $\mu = 0.5$ and $6.73\nu_0 Z^2 \times 10^{-4}$ for $\mu = 1$.

For $\mu^2 = 0$, $\tilde{\mathcal{K}}$ reduces to \mathcal{K} , and we still get two sets of eigenfunctions corresponding now to positive and negative values of κ . Their radial parts are still given by Eq.(70) if we take $\mu^2 = 0$ in it. However, there are no bound states in this case. The bound states do not exist because the parameters C_1^\pm in Eq.(70) become positive.

The first order corrections to the energies $|\bar{\mathcal{E}}_{0,1/2}^\pm|$ are

$$\left(R_{0,\frac{1}{2}}^{*,\pm}, V_{\text{pert}}^\pm R_{0,\frac{1}{2}}^\pm \right) = \int_0^\infty d\xi R_{0,\frac{1}{2}}^{*,\pm} V_{\text{pert}}^\pm R_{0,\frac{1}{2}}^\pm \left(1 + \frac{\bar{q}^2}{\xi} \right).$$

Performing the integration, we get

$$\begin{aligned} \left(R_{0,\frac{1}{2}}^{*,\pm}, V_{\text{pert}}^\pm R_{0,\frac{1}{2}}^\pm \right) &= \frac{2b_{0,\pm}}{3 + 2\bar{q}^2 b_{0,\pm}} \left\{ \mp \frac{\mu}{\sqrt{2}} (1 + \bar{q}^2 b_{0,\pm}) \pm \frac{1}{2} |\tilde{\kappa}| b_{0,\pm} + \frac{1}{8} b_{0,\pm} \right. \\ &\quad \left. + \frac{3}{4} b_{0,\pm}^2 \left[2\bar{q}^4 b_{0,\pm} e^{2b_{0,\pm}\bar{q}^2} E_1(2b_{0,\pm}\bar{q}^2) - \bar{q}^2 \right] \right\}, \end{aligned} \quad (78)$$

where $b_{0,\pm} = \sqrt{2|\bar{\mathcal{E}}_{0,1/2}^\pm|}$, and E_1 is the exponential integral

$$E_1(z) \equiv \int_1^\infty dt \frac{e^{-zt}}{t}.$$

For $\bar{q}^2 \gg 1$, $E_1(2b_{0,\pm}\bar{q}^2)$ can be approximated as

$$E_1(2b_{0,\pm}\bar{q}^2) \approx \frac{1}{2b_{0,\pm}\bar{q}^2} e^{-2b_{0,\pm}\bar{q}^2}.$$

This brings Eq.(78) into the form

$$\left(R_{0,\frac{1}{2}}^{*,\pm}, V_{\text{pert}}^\pm R_{0,\frac{1}{2}}^\pm \right) = \frac{4|\bar{\mathcal{E}}_{0,1/2}^\pm|}{b_{0,\pm}(3 + 2\bar{q}^2 b_{0,\pm})} \left[\mp \frac{\mu}{\sqrt{2}} (1 + \bar{q}^2 b_{0,\pm}) \pm \frac{1}{2} |\tilde{\kappa}| b_{0,\pm} + \frac{1}{8} b_{0,\pm} \right]. \quad (79)$$

For $\bar{q}^2 = 100$ and $\mu = 0.5$, the first order correction to $|\bar{\mathcal{E}}_{0,1/2}^+|$ is 1.1% of its value. For $\mu = 1$, it is even smaller: 0.54%. For the first order correction to $|\bar{\mathcal{E}}_{0,1/2}^-|$, we get 1.75% and 0.90%, respectively. This confirms the validity of the large \bar{q}^2 approximation for the $j = \frac{1}{2}$ ground states. The larger \bar{q}^2 or μ^2 , the more exact the approximation is.

4 Discussion

The non-relativistic counterpart of the Dirac equation should not be considered necessarily as its non-relativistic limit. With the rest energy included, the GLL equation leads to a different physical picture some elements of which cannot be seen in the framework of the LL equation. We demonstrated this for a non-relativistic Fermi field in pseudoscalar external potentials.

One of the new elements is the fact that the existence of bound states depends on the rest energy of the Fermi particle. The particle entering the region of a pseudoscalar spherical potential well is confined to this well if the rest energy differs from zero and exceeds a minimum value. This minimum is determined by the depth and width of the well. The larger the rest energy, the more bound states we get. Otherwise, the particle is not "captured" by the well. If it is in a bound state and its rest energy goes below the minimum value, this destroys the bound state, and the particle leaves the well.

The picture becomes richer and includes more new elements if we consider the motion of the Fermi particle in a pseudoscalar Coulomb potential. The particle experiences a rest energy dependent background field. This violates the parity and modifies the symmetry of the system which in turn results in restructuring of eigenfunctions. The bound state spectrum depends on the orientation of spin with respect to both the background field and the total angular momentum and consists of two sets corresponding to positive and negative values of the modified constant of motion.

Appendix A. Parity transformation

As we can see from Eq.(23), for the case $k = 0$, there are in general two ways to make the GLL equation in the primed frame the same as the one in the unprimed frame. The first way is to choose the parity transformation to satisfy the conditions

$$S^{-1}(P)\gamma^a S(P) = -\gamma^a, \quad S^{-1}(P)\gamma^4 S(P) = \gamma^4, \quad S^{-1}(P)\gamma^5 S(P) = \gamma^5.$$

These are equivalent to

$$\{\gamma^a, S(P)\} = 0, \quad [\gamma^4, S(P)]_- = 0, \quad [\gamma^5, S(P)]_- = 0.$$

However, these conditions only have the trivial solution $S(P) = 0$. Indeed, a general form of $S(P)$ anti-commuting with γ^a and commuting with γ^4 and γ^5 is a linear combination of all possible products of odd numbers of γ^4 and γ^5 . Such products are either equal to zero like

$$\gamma^4\gamma^5\gamma^5 = 0, \quad \gamma^5\gamma^4\gamma^4 = 0,$$

or reduce to γ^4 or γ^5 like

$$\gamma^4\gamma^5\gamma^4 = 2\gamma^4, \quad \gamma^5\gamma^4\gamma^5 = 2\gamma^5.$$

So, a general form of $S(P)$ is

$$S(P) = d_1\gamma^4 + d_2\gamma^5.$$

It anti-commutes with γ^a for any values of d_1 and d_2 . Its commutators with γ^4 and γ^5 become

$$[\gamma^4, S(P)]_- = d_2(\gamma^4\gamma^5 - \gamma^5\gamma^4), \quad [\gamma^5, S(P)]_- = -d_1(\gamma^4\gamma^5 - \gamma^5\gamma^4).$$

This yields $d_1 = d_2 = 0$, so no parity transformation can be found in this way.

The second way is to choose the conditions given by Eq.(24). Their equivalent form is

$$[\gamma^a, S(P)]_- = 0, \quad \{\gamma^4, S(P)\} = 0, \quad \{\gamma^5, S(P)\} = 0.$$

A general form of $S(P)$ commuting with γ^a and anti-commuting with γ^4 and γ^5 is a linear combination of all possible products of even numbers of γ^4 and γ^5 . Such products are either equal to zero or reduce to $\gamma^4\gamma^5$ or $\gamma^5\gamma^4$. Representing $S(P)$ as

$$S(P) = d_1\gamma^5\gamma^4 + d_2\gamma^4\gamma^5,$$

we find that $S(P)$ commutes with γ^a for any values of d_1 and d_2 , while its anti-commutators with γ^4 and γ^5 become

$$\{\gamma^4, S(P)\} = 2(d_1 + d_2)\gamma^4, \quad \{\gamma^5, S(P)\} = 2(d_1 + d_2)\gamma^5.$$

This yields $d_2 = -d_1$. From the condition $(S(P))^2 = 1$ we then find (up to the intrinsic parity) that $d_1 = 1/2$.

Appendix B. Matrix elements of J_1 , J_2 and J_3

Let us determine the matrix elements for the components of the total angular momentum in the case of pseudoscalar Coulomb potential. The spin operator components act on "spin-up" and "spin-down" columns, while the angular momentum components act on spherical harmonics as follows

$$\begin{aligned} L_3 Y_j^M &= M Y_j^M, \\ L_{\pm} Y_j^M &= -P_{j,\pm M} Y_j^{M\pm 1}, \end{aligned}$$

where

$$P_{j,M} \equiv \sqrt{j(j+1) - M(M+1)}$$

and

$$L_{\pm} = L_1 \pm i L_2.$$

Using the angular parts of the eigenfunctions given by Eqs.(61) and (62), we get the angular parts of the matrix elements for J_3 in the case $\mu = 0$:

$$\begin{aligned} (\mathcal{Y}_{j'M'}^{*,-|\kappa'|}, J_3 \mathcal{Y}_{jM}^{-|\kappa|}) &= (\mathcal{Y}_{j'M'}^{*,|\kappa'|}, J_3 \mathcal{Y}_{jM}^{|\kappa|}) = M \delta_{j'j} \delta_{M'M}, \\ (\mathcal{Y}_{j'M'}^{*,-|\kappa'|}, J_3 \mathcal{Y}_{jM}^{|\kappa|}) &= (\mathcal{Y}_{j'M'}^{*,|\kappa'|}, J_3 \mathcal{Y}_{jM}^{-|\kappa|}) = 0, \end{aligned}$$

where the products $(\mathcal{Y}_{j'M'}^{*,\kappa'}, J_a \mathcal{Y}_{jM}^{\kappa})$ are defined as

$$(\mathcal{Y}_{j'M'}^{*,\kappa'}, J_a \mathcal{Y}_{jM}^{\kappa}) \equiv \int_0^{2\pi} d\phi \int_0^{\pi} d\theta \sin\theta \mathcal{Y}_{j'M'}^{*,\kappa'} J_a \mathcal{Y}_{jM}^{\kappa},$$

so the matrix elements of J_3 which correspond to transitions between states of different j or M as well as states with opposite parities are zero.

The corresponding matrix elements for J_1 and J_2 are

$$\begin{aligned} (\mathcal{Y}_{j'M'}^{*,-|\kappa'|}, J_1 \mathcal{Y}_{jM}^{-|\kappa|}) &= -\frac{j-1}{2j} \delta_{j'j} (P_{j,M} \delta_{M',M+1} + P_{j,-M} \delta_{M',M-1}), \\ (\mathcal{Y}_{j'M'}^{*,|\kappa'|}, J_1 \mathcal{Y}_{jM}^{|\kappa|}) &= -\frac{j+2}{2j+2} \delta_{j'j} (P_{j,M} \delta_{M',M+1} + P_{j,-M} \delta_{M',M-1}), \\ (\mathcal{Y}_{j'M'}^{*,-|\kappa'|}, J_1 \mathcal{Y}_{jM}^{|\kappa|}) &= \frac{1}{2j+2} \delta_{j',j+1} (Q_{j+1,M} \delta_{M',M+1} - Q_{j+1,-M} \delta_{M',M-1}), \\ (\mathcal{Y}_{j'M'}^{*,|\kappa'|}, J_1 \mathcal{Y}_{jM}^{-|\kappa|}) &= -\frac{1}{2j} \delta_{j',j-1} (Q_{j-1,-M} \delta_{M',M+1} - Q_{j-1,M} \delta_{M',M-1}). \end{aligned}$$

and

$$\begin{aligned} (\mathcal{Y}_{j'M'}^{*,-|\kappa'|}, J_2 \mathcal{Y}_{jM}^{-|\kappa|}) &= i \frac{j-1}{2j} \delta_{j'j} (P_{j,M} \delta_{M',M+1} - P_{j,-M} \delta_{M',M-1}), \\ (\mathcal{Y}_{j'M'}^{*,|\kappa'|}, J_2 \mathcal{Y}_{jM}^{|\kappa|}) &= i \frac{j+2}{2j+2} \delta_{j'j} (P_{j,M} \delta_{M',M+1} - P_{j,-M} \delta_{M',M-1}), \\ (\mathcal{Y}_{j'M'}^{*,-|\kappa'|}, J_2 \mathcal{Y}_{jM}^{|\kappa|}) &= -i \frac{1}{2j+2} \delta_{j',j+1} (Q_{j+1,M} \delta_{M',M+1} + Q_{j+1,-M} \delta_{M',M-1}), \\ (\mathcal{Y}_{j'M'}^{*,|\kappa'|}, J_2 \mathcal{Y}_{jM}^{-|\kappa|}) &= i \frac{1}{2j} \delta_{j',j-1} (Q_{j-1,-M} \delta_{M',M+1} + Q_{j-1,M} \delta_{M',M-1}), \end{aligned}$$

respectively, where

$$Q_{j,M} \equiv \sqrt{\frac{j+M}{j-M}} P_{j,M}.$$

Using next the relations given by Eqs.(65) and (66), we evaluate the angular parts of the matrix elements for the total angular momentum components in the case of non-zero values of μ . The matrix elements for J_3 have the same structure as in the case $\mu = 0$:

$$\begin{aligned} (\mathcal{Y}_{j'M'}^{*,+}, J_3 \mathcal{Y}_{jM}^-) &= (\mathcal{Y}_{j'M'}^{*,+}, J_3 \mathcal{Y}_{jM}^+) = M \delta_{j'j} \delta_{M'M}, \\ (\mathcal{Y}_{j'M'}^{*,+}, J_3 \mathcal{Y}_{jM}^+) &= (\mathcal{Y}_{j'M'}^{*,+}, J_3 \mathcal{Y}_{jM}^-) = 0, \end{aligned}$$

while the matrix elements for J_1 and J_2 become

$$\begin{aligned} (\mathcal{Y}_{j'M'}^{*,s'}, J_1 \mathcal{Y}_{jM}^s) &= A_j^{s's} \delta_{j'j} (P_{j,M} \delta_{M',M+1} + P_{j,-M} \delta_{M',M-1}) \\ &+ B_j^{s's} \delta_{j',j+1} (Q_{j+1,M} \delta_{M',M+1} - Q_{j+1,-M} \delta_{M',M-1}) \\ &+ C_j^{s's} \delta_{j',j-1} (Q_{j-1,-M} \delta_{M',M+1} - Q_{j-1,M} \delta_{M',M-1}) \end{aligned}$$

and

$$\begin{aligned} (\mathcal{Y}_{j'M'}^{*,s'}, J_2 \mathcal{Y}_{jM}^s) &= -i A_j^{s's} \delta_{j'j} (P_{j,M} \delta_{M',M+1} - P_{j,-M} \delta_{M',M-1}) \\ &- i B_j^{s's} \delta_{j',j+1} (Q_{j+1,M} \delta_{M',M+1} + Q_{j+1,-M} \delta_{M',M-1}) \\ &- i C_j^{s's} \delta_{j',j-1} (Q_{j-1,-M} \delta_{M',M+1} + Q_{j-1,M} \delta_{M',M-1}), \end{aligned}$$

where $s', s = (+, -)$,

$$\begin{aligned} A_j^{--} &\equiv -\frac{1}{2} \left(\alpha_j^2 \frac{j-1}{j} + \beta_j^2 \frac{j+2}{j+1} \right), \\ A_j^{++} &\equiv -\frac{1}{2} \left(\alpha_j^2 \frac{j+2}{j+1} + \beta_j^2 \frac{j-1}{j} \right), \\ A_j^{-+} &= A_j^{+-} \equiv \frac{1}{2} \alpha_j \beta_j \frac{2j+1}{j(j+1)}, \end{aligned}$$

and

$$\begin{aligned} B_j^{--} &\equiv -\alpha_{j+1} \beta_j \frac{1}{2j+2}, & C_j^{--} &\equiv \beta_{j-1} \alpha_j \frac{1}{2j}, \\ B_j^{++} &\equiv \beta_{j+1} \alpha_j \frac{1}{2j+2}, & C_j^{++} &\equiv -\alpha_{j-1} \beta_j \frac{1}{2j}, \\ B_j^{-+} &\equiv \alpha_{j+1} \alpha_j \frac{1}{2j+2}, & C_j^{-+} &\equiv \beta_{j-1} \beta_j \frac{1}{2j}, \\ B_j^{+-} &\equiv -\beta_{j+1} \beta_j \frac{1}{2j+2}, & C_j^{+-} &\equiv -\alpha_{j-1} \alpha_j \frac{1}{2j}. \end{aligned}$$

Appendix C. Proof of orthonormality condition

Let us first show that, for a given value of j , the radial parts of eigenfunctions corresponding to different values of quantum number n ,

$$R_{n,j}^{\pm}(\xi) = N_{n,j}^{\pm} e^{-b_{n,\pm}\xi} \xi^{\lambda} {}_1F_1(-n, 2\lambda; 2b_{n,\pm}\xi),$$

with $b_{n,\pm} \equiv \sqrt{2|\bar{\mathcal{E}}_{n,j}^{\pm}|}$, and $N_{n,j}^{\pm}$ being the normalization constants, are orthogonal. Introducing the integral [28]

$$J_{p'}^{sp}(f, f'; g, g') \equiv \int_0^{\infty} dz e^{-\frac{1}{2}(g+g')z} z^{p'-1+s} {}_1F_1(f, p'; gz) {}_1F_1(f', p' - p; g'z),$$

we can rewrite the matrix elements $(R_{n,j}^{\star,\pm}, R_{n',j}^{\pm})$ as

$$(R_{n,j}^{\star,\pm}, R_{n',j}^{\pm}) = N_{n,j}^{\pm} N_{n',j}^{\pm} \left[J_{2\lambda}^{10}(-n, -n'; 2b_{n,\pm}, 2b_{n',\pm}) + \bar{q}^2 J_{2\lambda}^{00}(-n, -n'; 2b_{n,\pm}, 2b_{n',\pm}) \right],$$

where $n \neq n'$.

The integrals $J_{p'}^{10}$ and $J_{p'}^{00}$ are related as follows:

$$J_{p'}^{10}(f, f'; g, g') = \frac{4}{g^2 - g'^2} \left[\frac{1}{2} p'(g - g') - fg + f'g' \right] J_{p'}^{00}(f, f'; g, g').$$

In our case, the relation takes the form

$$\begin{aligned} J_{2\lambda}^{10}(-n, -n'; 2b_{n,\pm}, 2b_{n',\pm}) &= \frac{2}{b_{n,\pm}^2 - b_{n',\pm}^2} [(\lambda + n)b_{n,\pm} - (\lambda + n')b_{n',\pm}] \\ &\times J_{2\lambda}^{00}(-n, -n'; 2b_{n,\pm}, 2b_{n',\pm}). \end{aligned}$$

The combination $(\lambda + n)b_{n,\pm} - (\lambda + n')b_{n',\pm}$ is determined by the energy spectrum. Using Eqs.(71) and (74), we get

$$(\lambda + n)b_{n,\pm} - (\lambda + n')b_{n',\pm} = -\frac{\bar{q}^2}{2} (b_{n,\pm}^2 - b_{n',\pm}^2).$$

This gives us

$$J_{2\lambda}^{10}(-n, -n'; 2b_{n,\pm}, 2b_{n',\pm}) + \bar{q}^2 J_{2\lambda}^{00}(-n, -n'; 2b_{n,\pm}, 2b_{n',\pm}) = 0,$$

so that $(R_{n,j}^{\star,\pm}, R_{n',j}^{\pm})$ vanish.

Next, to find the normalization constants, we consider the case $n = n'$. For the ground states $n = n' = 0$, ${}_1F_1(0, 2\lambda; 2b_{0,\pm}\xi) = 1$, and the orthonormality condition (75) becomes

$$(N_{0,j}^{\pm})^2 \int_0^{\infty} d\xi e^{-2b_{0,\pm}\xi} \xi^{2\lambda} \left(1 + \frac{\bar{q}^2}{\xi} \right) = 1.$$

This yields

$$N_{0,j}^{\pm} = (2b_{0,\pm})^{j+1} \left[\frac{1}{(2j+1)!} \frac{b_{0,\pm}}{j+1+b_{0,\pm}\bar{q}^2} \right]^{1/2}.$$

For $n = n' \geq 1$, we use the integral

$$J_n(\nu, p, g) \equiv \int_0^{\infty} dz e^{-gz} z^{\nu-1} [{}_1F_1(-n, p; gz)]^2$$

which can be evaluated as [28]

$$J_n(\nu, p, g) = \frac{\Gamma(\nu)n!}{g^{\nu}p(p+1)\cdots(p+n-1)} \times \left\{ 1 + \sum_{s=0}^{n-1} \frac{n(n-1)\cdots(n-s)(p-\nu-s-1)(p-\nu-s)\cdots(p-\nu+s)}{[(s+1)!]^2 p(p+1)\cdots(p+s)} \right\}.$$

We are interested in two particular cases: $p = \nu$ and $p = \nu - 1$. The values of the integral in these two cases are

$$J_n(\nu, \nu, g) = \frac{\Gamma(\nu)n!}{g^{\nu}\nu(\nu+1)\cdots(\nu+n-1)}$$

and

$$J_n(\nu, \nu-1, g) = \frac{\Gamma(\nu)n!}{g^{\nu}(\nu-1)\nu\cdots(\nu+n-2)} \left[1 + \frac{2n}{\nu-1} \right],$$

respectively. Then the orthonormality condition

$$(N_{n,j}^{\pm})^2 \left[J_n(2\lambda+1, 2\lambda, 2b_{n,\pm}) + \bar{q}^2 J_n(2\lambda, 2\lambda, 2b_{n,\pm}) \right] = 1$$

gives us

$$N_{n,j}^{\pm} = \frac{(2b_{n,\pm})^{j+\frac{3}{2}}}{\Gamma(2j+2)} \left[\frac{\Gamma(2j+n+2)}{2n!(j+n+1+\bar{q}^2 b_{n,\pm})} \right]^{1/2}, \quad n \geq 1.$$

References

- [1] J.M. Lévy-Leblond, *Comm. Math. Phys.* 6 (1967) 286.
- [2] M. de Montigny, F.C. Khanna, A.E. Santana, E.S. Santos, J.D.M. Vianna, *Ann. Phys. (N.Y.)* 277 (1999) 144;
M. de Montigny, F.C. Khanna, A.E. Santana, E.S. Santos, J.D.M. Vianna, *J. Phys. A : Math. Gen.* 33 (2000) L273;
M. de Montigny, F.C. Khanna, A.E. Santana, E.S. Santos, *J. Phys. A : Math. Gen.* 34 (2001) 8901;
M. de Montigny, F.C. Khanna, A.E. Santana, *Int. J. Theor. Phys.* 42 (2003) 649;
M. de Montigny, F.C. Khanna, A.E. Santana, *J. Phys. A: Math. Gen.* 36 (2003) 2009;
L. Abreu, M. de Montigny, F.C. Khanna, A.E. Santana, *Ann. Phys. (N.Y.)* 308 (2003) 244;
E.S. Santos, M. de Montigny, F.C. Khanna, A.E. Santana, *J. Phys. A: Math. Gen.* 37 (2004) 9771;
E.S. Santos, M. de Montigny, F.C. Khanna, *Ann. Phys. (NY)* 320 (2005) 21.
- [3] Y. Takahashi, *Fortschr. Phys.* 36 (1988) 63;
Y. Takahashi, *Fortschr. Phys.* 36 (1988) 83.
- [4] M. Omote, S. Kamefuchi, T. Takahashi, Y. Ohnuki, *Fortschr. Phys.* 37 (1989) 933.
- [5] F.M. Saradzhev, *Phys. Rev. D* 84 (2011) 125025.
- [6] R. Huegele, Z.E. Musielak, J.L. Fry, *Adv. Studies Theor. Phys.* 7 (2013) 825.
- [7] A. Horzela, E. Kapuścik, *Electromagnetic Phenomena*, 3 (2003) 63.
- [8] M. de Montigny, F. Khanna, F.M. Saradzhev, *Ann. Phys. (NY)* 323 (2008) 1191.
- [9] See, for instance, M. Jammer, *Concepts of mass in contemporary physics and philosophy*, Princeton University Press, 1999.
- [10] R. Jackiw, V.P. Nair, *Phys. Lett. B* 551 (2003) 166;
- [11] B. Thaller, *The Dirac equation*, Springer, 1992.
- [12] B.R. Desai, *Quantum mechanics with basic field theory*, Cambridge University Press, 2010.

- [13] F. Gross, Relativistic quantum mechanics and field theory, John Wiley and Sons, 1993.
- [14] J.J. Sakurai, Advanced Quantum Mechanics, Addison-Wesley, 1967.
- [15] S. Flügge, Practical quantum mechanics, Springer-verlag, 1974.
- [16] A.Z. Capri, Nonrelativistic quantum mechanics, World Scientific, 2002.
- [17] G. Temple, The general principles of quantum mechanics, Methuen, New York, 1948;
P.C. Martin, R.J. Glauber, Phys. Rev. 109 (1958) 1307.
- [18] L.C. Biedenharn, Phys. Rev. 126 (1962) 845;
L.C. Biedenharn, N.V.V.J. Swamy, Phys. Rev. 133 (1964) B1353.
- [19] F. Bloore, P.A. Horváthy, J. Math. Phys. 33 (1992) 1869;
P.A. Horváthy, A.J. Macfarlane, J.W. van Holten, Phys. Lett. B 486 (2000) 346;
S. Haouat, L. Chetouani, J. Phys. A 40 (2007) 10541.
- [20] W. Pauli, Z. Phys. 43 (1927) 601.
- [21] H.A. Bethe, E.E. Salpeter, Quantum mechanics of one and two electron systems, Springer, 1957.
- [22] H. Snyder, J. Weinberg, Phys. Rev. 57 (1940) 307;
L.I. Schiff, H. Snyder, J. Weinberg, Phys. Rev. 57 (1940) 315.
- [23] A.M. Green, Nucl. Phys. 33 (1962) 218.
- [24] H. Leutwyler, J. Stern, Ann. Phys. (NY) 112 (1978) 24.
- [25] I.T. Todorov, Phys. Rev. (1971) 2351;
V.A. Rizov, H. Sazdjian, I.T. Todorov, Ann. Phys. (NY) 165 (1985) 59.
- [26] G.P. Lepage, Phys. Rev. A16 (1977) 863;
H. Sazdjian, J. Math. Phys. 29 (1988) 1620.
- [27] J. Formánek, R.J. Lombard, J. Mares, Czech. J. Phys. 54 (2004) 289;
R.J. Lombard, J. Mares, C. Volpe, J. Phys. G34 (2007) 1879;
R. Yekken, R.J. Lombard, J. Phys. A43 (2010) 125301, and references therein.
- [28] L.D. Landau, E.M. Lifshitz, Quantum Mechanics, Pergamon Press, 1991.