

PARALLEL TRANSPORT FOR HIGGS BUNDLES OVER p -ADIC CURVES

DAXIN XU, WITH AN APPENDIX BY TONGMU HE, DAXIN XU

ABSTRACT. Faltings conjectured that under the p -adic Simpson correspondence, finite dimensional p -adic representations of the geometric étale fundamental group of a smooth proper p -adic curve X are equivalent to semi-stable Higgs bundles of degree zero over X . In this article, we establish an equivalence between these representations and Higgs bundles, whose underlying bundles potentially admit a strongly semi-stable reduction of degree zero. We show that these Higgs bundles are semi-stable of degree zero and investigate some evidence for Faltings' conjecture.

CONTENTS

1. Introduction	1
2. Higgs crystals and Higgs bundles	6
3. Small Higgs bundles with strongly semi-stable reduction of degree zero	17
4. Review on the p -adic Simpson correspondence	19
5. From \mathbf{C} -representations of the geometric fundamental group to Higgs bundles	25
6. Parallel transport for Higgs bundles	29
Appendix A. Twisted inverse image for Higgs bundles over curves, d'après Faltings	35
References	42

1. INTRODUCTION

1.1.1. In his seminal work [30], Simpson established a correspondence between finite dimensional complex linear representations of the topological fundamental group of a connected projective complex manifold and semi-stable Higgs bundles with vanishing Chern classes. This result generalizes the Narasimhan–Seshadri correspondence [23] between unitary irreducible representations of the fundamental group of a compact Riemann surface of genus ≥ 2 and stable vector bundles of degree zero.

Inspired by these works, Faltings developed an analogue theory between (generalized) p -adic local systems and Higgs bundles over p -adic smooth proper varieties, called *the p -adic Simpson correspondence* [13]. Faltings' short paper has been systematically studied by Abbes–Gros and Tsuji [4], each introducing their own new approach and laying the foundations of this theory.

Simultaneously and independently, Deninger and Werner developed a p -adic analogue of the Narasimhan–Seshadri correspondence: they associated functorially to every vector bundle over a p -adic smooth projective curve whose reduction is strongly semi-stable of degree zero a p -adic representation of the geometric fundamental group of the curve [10]. The author showed that the construction of Deninger–Werner is compatible with Faltings' theory in the case where the Higgs field is trivial [36]. These results have been generalized to the case of higher dimensional p -adic varieties by Deninger–Werner [11] and Würthen [35].

Based on Scholze's work [28], Liu and Zhu [21] developed a p -adic Riemann–Hilbert correspondence for \mathbb{Q}_p -local systems on a smooth rigid analytic variety over a finite extension of \mathbb{Q}_p , which allows them to associate nilpotent Higgs bundles to these arithmetic local systems. This work is recently extended to the

logarithmic case by Diaol-Lan-Liu-Zhu [12]. Y. Wang compared Faltings' approach in the good reduction case with the work of Liu-Zhu in his thesis [34].

Faltings imbedded the category of p -adic representations of the geometric fundamental group into the much larger category of so called "generalized representations" that he expected to be equivalent to the category of all Higgs bundles. The fundamental problem is then to understand which Higgs bundles arise from true p -adic representations. He formulated, at least for curves, the hope that true p -adic representations correspond to semi-stable Higgs bundles of slope zero. Faltings sketched a strategy of two steps to establish the equivalence between generalized representations and Higgs bundles. First, one establishes an equivalence between the subcategories of objects that are p -adically close from the trivial objects, that he qualifies as being "small". Second, one extends the equivalence to all objects by descent, as any generalized representation becomes small over a finite étale covering. The first step has been developed by Abbes-Gros and Tsuji [4] in any dimension. The second step, sketched by Faltings only for curves, remained mysterious.

In this article, we develop for curves an alternative descent argument to the second step of Faltings strategy. We treat only true p -adic representations, avoiding the difficult issue of descending generalized representation. Our construction provides a characterization of the Higgs bundles associated to true p -adic representations that we compare to Faltings' expected description.

Let K be a finite extension of \mathbb{Q}_p , \overline{K} an algebraic closure of K , \mathbf{C} (resp. \mathfrak{o}) the p -adic completion of \overline{K} (resp. $\mathcal{O}_{\overline{K}}$), X a geometrically connected, smooth proper curve over K and \overline{x} a geometric point of X . Roughly speaking, we establish an equivalence between the category $\mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C})$ of continuous finite dimensional \mathbf{C} -representations of $\pi_1(X_{\overline{K}}, \overline{x})$ and a category of Higgs bundles over $X_{\mathbf{C}}$, whose underlying bundles admit a strongly semi-stable reduction of degree zero as in Deninger-Werner's theory, after taking a twisted inverse image over a finite étale cover of $X_{\overline{K}}$. This equivalence is compatible with the p -adic Simpson correspondence when restricted to small objects. We prove that these Higgs bundles are semi-stable of degree zero. It is conjectured that the converse holds. We discuss these results in more details in the following.

1.1.2. We first briefly explain the p -adic Simpson correspondence.

We set $S = \mathrm{Spec}(\mathcal{O}_K)$. Let X be a projective S -scheme with semi-stable reduction¹ and geometrically connected generic fiber $X_{\overline{K}}$ and \overline{x} a geometric point of $X_{\overline{K}}$. The p -adic Simpson correspondence applies to certain objects more general than those of $\mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C})$, called *generalized representations*. They are certain modules of the Faltings ringed topos and locally can be expressed as semi-linear representations of $\pi_1(X_{\overline{K}}, \overline{x})$ over finite projective modules over a certain p -adic ring equipped with a continuous action of $\pi_1(X_{\overline{K}}, \overline{x})$. A generalized representation is called *small*, if locally, the associated representation admits a basis, whose $\pi_1(X_{\overline{K}}, \overline{x})$ -action is trivial modulo p^β for some $\beta > \frac{2}{p-1}$ (definition 4.1.9). The category $\mathbf{GRep}(X)$ of *generalized representations* of X contains $\mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C})$ as a full subcategory and we denote by $\mathbf{GRep}_{\mathrm{small}}(X)$ its full subcategory of small objects.

On the other hand, a Higgs bundle over $X_{\mathbf{C}}$ is a pair (M, θ) consisting of a vector bundle M over $X_{\mathbf{C}}$ together with a Higgs field $\theta : M \rightarrow M \otimes_{\mathcal{O}_{X_{\mathbf{C}}}} \Omega_{X_{\mathbf{C}}}^1(-1)$ such that $\theta \wedge \theta = 0$, where (-1) denotes the inverse of the Tate twist. We say a Higgs bundle (M, θ) over $X_{\mathbf{C}}$ is *small* if it admits an integral model (M°, θ) over $X_\circ = X \otimes_{\mathcal{O}_K} \mathfrak{o}$ such that $\theta(M^\circ) \subset p^\alpha M^\circ \otimes_{\mathcal{O}_X} \tilde{\Omega}_{X/S}^1$ for some $\alpha > \frac{1}{p-1}$. Here, $\tilde{\Omega}_{X/S}^1$ denotes the \mathcal{O}_X -module of logarithmic differentials for the logarithmic structures \mathcal{M}_X and \mathcal{M}_S defined by the special fibers of X and S respectively. We denote by $\mathrm{HB}(X_{\mathbf{C}})$ the category of Higgs bundles over $X_{\mathbf{C}}$ and by $\mathrm{HB}_{\mathrm{small}}(X_{\mathbf{C}})$ its full subcategory of small Higgs bundles.

Set $\mathcal{O}_K^b = \varprojlim_{x \rightarrow x^p} \mathcal{O}_{\overline{K}}/p\mathcal{O}_{\overline{K}}$ and $A_{\mathrm{inf}} = W(\mathcal{O}_K^b)$ and let $\theta : A_{\mathrm{inf}} \rightarrow \mathfrak{o}$ be Fontaine's theta map. Let \check{X} be the p -adic completion of X_\circ . We take a family of log-smooth liftings $\mathcal{X} = (\mathcal{X}_n)_{n \geq 1}$ of \check{X} over $(A_{\mathrm{inf}}/\ker(\theta)^n)_{n \geq 1}$ with respect to logarithmic structures extending by \mathcal{M}_X and \mathcal{M}_S (see § 2.1.9 for more details). Based on the works of Faltings [13], Abbes-Gros [4], Tsuji [4, 33], the lifting \mathcal{X} induces an equivalence of categories:

$$(1.1.2.1) \quad \mathbf{H}_{\mathcal{X}} : \mathbf{GRep}_{\mathrm{small}}(X) \simeq \mathrm{HB}_{\mathrm{small}}(X_{\mathbf{C}}),$$

¹We restrict ourself to the projective with semistable reduction case for the simplicity of presentation. The p -adic Simpson correspondence applies to more general settings.

called the *p -adic Simpson correspondence*.

1.1.3. In the following, we assume moreover that X has *relative dimension one* over S . In this article, we construct a functor (5.2.2.1) compatible with the p -adic Simpson correspondence $\mathbf{H}_{\mathcal{X}}$:

$$(1.1.3.1) \quad \mathbb{H}_{\mathcal{X}, \text{Exp}} : \mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C}) \rightarrow \text{HB}(X_{\mathbf{C}}).$$

The construction relies on a section $\text{Exp} : \mathbf{C} \rightarrow 1 + \mathfrak{m}$ (A.0.0.2) of the p -adic logarithmic homomorphism $\log : 1 + \mathfrak{m} \rightarrow \mathbf{C}$, $x \mapsto \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(x-1)^n}{n}$. For small \mathbf{C} -representations, the above functor is explicitly defined by gluing a local version of the p -adic Simpson correspondence ([4] § II.13, II.14).

To extend the construction for all \mathbf{C} -representations, an important tool is the *twisted inverse image functor* for Higgs bundles, sketched in Faltings' paper [13]. We present this functor in a more general setting in the appendix, jointly with T. He. Let Y be a smooth projective S -curve with semi-stable reduction and $f : Y \rightarrow X$ a proper S -morphism, which is finite étale over X_K , and \mathcal{Y} a family of log-smooth liftings of \check{Y} over $(A_{\text{inf}}/\ker(\theta)^n)_{n \geq 1}$. There exists a *twisted inverse image functor* associated to these data (A.3.0.2):

$$(1.1.3.2) \quad f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^{\circ} : \text{HB}(X_{\mathbf{C}}) \rightarrow \text{HB}(Y_{\mathbf{C}}),$$

which is different from the usual inverse image functor by a twist. Its restriction to small Higgs bundles is compatible with the natural inverse image functor of generalized representations via p -adic Simpson correspondences $\mathbf{H}_{\mathcal{X}}$ and $\mathbf{H}_{\mathcal{Y}}$. The twisted inverse image functor and the étale descent for Higgs bundles allow us to define the functor $\mathbb{H}_{\mathcal{X}, \text{Exp}}$.

The same argument also extends $\mathbf{H}_{\mathcal{X}}$ to a functor $\mathbf{H}_{\mathcal{X}, \text{Exp}}$ from generalized representations over X to Higgs bundles over $X_{\mathbf{C}}$, which is the easy direction in the second step of Faltings' approach (1.1.1). However, the difficulty occurs in the other direction, which requires the descent of certain generalized representations. In this article, we describe the essential image of $\mathbb{H}_{\mathcal{X}, \text{Exp}}$, related to Deninger–Werner's theory [10], and then construct a quasi-inverse of $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ by another method (1.1.11), which avoids this difficulty.

1.1.4. In summary, we have the following commutative diagram

$$(1.1.4.1) \quad \begin{array}{ccc} \mathbf{GRep}_{\text{small}}(X) & \xrightarrow[\sim]{\mathbf{H}_{\mathcal{X}}} & \text{HB}_{\text{small}}(X_{\mathbf{C}}) \\ \downarrow & & \downarrow \\ \mathbf{GRep}(X) & \xrightarrow[\text{---}]{\mathbf{H}_{\mathcal{X}, \text{Exp}}} & \text{HB}(X_{\mathbf{C}}) \\ \uparrow & \nearrow \mathbb{H}_{\mathcal{X}, \text{Exp}} & \uparrow \\ \mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C}) & \xleftarrow{\mathbb{V}^{\text{DW}}} & \text{VB}^{\text{DW}}(X_{\mathbf{C}}) \end{array}$$

Let us briefly explain how Deninger–Werner's functor $\mathbb{V}^{\text{DW}} : \text{VB}^{\text{DW}}(X_{\mathbf{C}}) \rightarrow \mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C})$ [10] fits into this picture. Recall that a vector bundle over a smooth proper curve C of characteristic $p > 0$ is *strongly semi-stable* if its inverse images by all the power of the absolute Frobenius F_C of C are semi-stable. A vector bundle M over $X_{\mathbf{C}}$ is called *Deninger–Werner* if its restriction to the normalisation of each irreducible component of $X_{\overline{K}}$ is strongly semi-stable of degree zero. A vector bundle over $X_{\mathbf{C}}$ is *of Deninger–Werner*, if it admits a Deninger–Werner vector bundle over $X_{\mathbf{C}}$ as an integral model. For every integer $n \geq 1$, the reduction modulo p^n of a Deninger–Werner vector bundle over $X_{\mathbf{C}}$ can be trivialized by a proper morphism of S -curves $Y \rightarrow X$, which is finite étale over X_K (after taking a finite extension of K). By parallel transport, Deninger and Werner constructed a functor \mathbb{V}^{DW} from the category $\text{VB}^{\text{DW}}(X_{\mathbf{C}})$ of Deninger–Werner vector bundles to $\mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C})$. In the above diagram, the functor $\text{VB}^{\text{DW}}(X_{\mathbf{C}}) \rightarrow \text{HB}(X_{\mathbf{C}})$ is given by equipping each vector bundle with the trivial Higgs field, and factors through $\text{HB}_{\text{small}}(X_{\mathbf{C}})$. The compatibility between \mathbb{V}^{DW} and $\mathbf{H}_{\mathcal{X}}$ is proved in the author's thesis [36].

1.1.5. In the following, we characterize the essential image of $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ (1.1.3.1) in terms of certain Higgs bundles related to Deninger–Werner vector bundles.

We denote by $\mathrm{HB}^{\mathrm{DW}}(X_{\circ})$ the category of pairs (M, θ) consisting of a Deninger–Werner vector bundle M on X_{\circ} together with a small Higgs field θ on M , and by $\mathrm{HB}^{\mathrm{DW}}(X_{\mathbf{C}})$ the full subcategory of $\mathrm{HB}(X_{\mathbf{C}})$ consisting of Higgs bundles which admits an integral model in $\mathrm{HB}^{\mathrm{DW}}(X_{\circ})$. A Higgs bundle over $X_{\mathbf{C}}$ is called *potentially Deninger–Werner*, if it belongs to $\mathrm{HB}^{\mathrm{DW}}(Y_{\mathbf{C}})$ after taking a twisted inverse image functor (1.1.3.2). We denote by $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$ the category of potentially Deninger–Werner Higgs bundles over $X_{\mathbf{C}}$. Here is our main result.

Theorem 1.1.6 (5.2.5, 6.0.1). (i) *The functor $\mathbb{H}_{\mathcal{X}, \mathrm{Exp}}$ factors through the category $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$:*

$$(1.1.6.1) \quad \mathbb{H}_{\mathcal{X}, \mathrm{Exp}} : \mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C}) \rightarrow \mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}}).$$

(ii) *There exists a quasi-inverse $\mathbb{V}_{\mathcal{X}, \mathrm{Exp}}$ of $\mathbb{H}_{\mathcal{X}, \mathrm{Exp}}$:*

$$(1.1.6.2) \quad \mathbb{V}_{\mathcal{X}, \mathrm{Exp}} : \mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}}) \xrightarrow{\sim} \mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C}).$$

Higgs bundles of $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$ satisfy following properties:

Theorem 1.1.7 (§ 5.3, 6.4.3). (i) [Faltings] *Every object of $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$ is semi-stable of degree zero.*

(ii) *Every Higgs line bundle of degree zero over $X_{\mathbf{C}}$ belongs to $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$.*

(iii) *The category $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$ is abelian and is closed under extension.*

(iv) *When the genus of $X_{\mathbf{C}}$ is zero or one, every semi-stable Higgs bundle of degree zero belongs to $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$.*

In ([13] § 5), Faltings claimed that Higgs bundles over $X_{\mathbf{C}}$, associated to $\mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C})$, are semi-stable of degree zero and gave a brief sketch of the proof. Moreover, he expressed the hope that all semi-stable Higgs bundles of degree zero are obtained in this way. Our proofs of theorems 1.1.6(i), 1.1.7(i) are inspired by Faltings' sketched argument. Based on theorem 1.1.6, we can reformulate Faltings' conjecture as follow.

Conjecture 1.1.8 (Faltings). *Every semi-stable Higgs bundle of degree zero over $X_{\mathbf{C}}$ belongs to $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$.*

Moreover, we can compare the cohomologies of two sides in this correspondence by considering extensions:

Proposition 1.1.9. *Let (M, θ) be a Higgs bundle of $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$ and $V = \mathbb{V}_{\mathcal{X}, \mathrm{Exp}}(M, \theta)$ the associated \mathbf{C} -representation. There exists a canonical \mathbf{C} -linear isomorphism:*

$$(1.1.9.1) \quad \mathrm{H}^1(X_{\overline{K}, \mathrm{\acute{e}t}}, V) \simeq \mathrm{H}^1(X_{\mathbf{C}}, M \xrightarrow{\theta} M \otimes_{\mathcal{O}_{X_{\mathbf{C}}}} \Omega_{X_{\mathbf{C}}}^1(-1)).$$

Remark 1.1.10. (i) A Higgs bundle $(M, 0)$ with zero Higgs field belongs to $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$ if and only if M potentially belongs to $\mathrm{VB}^{\mathrm{DW}}(X_{\mathbf{C}})$. And the functor $\mathbb{V}_{\mathcal{X}, \mathrm{Exp}}$ is compatible with \mathbb{V}^{DW} (proposition 6.4.1).

(ii) In the complex Simpson correspondence [30], Higgs bundles, fixed by the natural \mathbb{G}_m -action on Higgs fields, correspond to variations of Hodge structures. It is an important question to characterize these Higgs bundles in the p -adic theory. Note that Higgs bundles fixed by the \mathbb{G}_m -action are nilpotent and the associated generalized representations are of Hodge–Tate in the sense of [3].

1.1.11. Let us briefly explain the construction of the functor $\mathbb{V}_{\mathcal{X}, \mathrm{Exp}}$, which is inspired by \mathbb{V}^{DW} .

Recall (1.1.4) that, for every $n \geq 1$, the reduction modulo p^n of a Deninger–Werner vector bundle over X_{\circ} can be trivialized by a proper morphism of S -curves $f : Y \rightarrow X$ such that f_K is finite étale. We extend this trivializing property to $\mathrm{HB}^{\mathrm{DW}}(X_{\circ})$ (see theorem 3.2.2 for a precise statement in terms of Higgs crystals). Via Tsuji's approach to the p -adic Simpson correspondence ([4] Chapter IV), we obtain a similar trivializing property for generalized representations over X associated to $\mathrm{HB}^{\mathrm{DW}}(X_{\mathbf{C}})$. This allows us to apply a parallel transport functor, constructed by the author ([36] § 8), from these generalized representations over X to $\mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C})$. Combined with these constructions, we obtain a functor

$$\mathbb{V}_{\mathcal{X}} : \mathrm{HB}^{\mathrm{DW}}(X_{\mathbf{C}}) \rightarrow \mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C}).$$

Using the descent for sheaves of finite étale topos, we extend $\mathbb{V}_{\mathcal{X}}$ to the functor $\mathbb{V}_{\mathcal{X}, \text{Exp}}$.

When $\mathbb{V}_{\mathcal{X}, \text{Exp}}$ (resp. $\mathbb{H}_{\mathcal{X}, \text{Exp}}$) restricts to small Higgs bundles (resp. small \mathbf{C} -representations), we show that it is compatible with the p -adic Simpson correspondence $\mathbf{H}_{\mathcal{X}}$ (and is independent of the choice of Exp). This assertion for $\mathbb{V}_{\mathcal{X}, \text{Exp}}$ relies on a recent work of T. He on the cohomological descent in Faltings topos [17] (another approach to this result is also obtained by Abbes–Gros [2] proposition 4.6.30). A key ingredient in T. He’s proof is the arc-descent of perfectoid algebras due to Bhatt–Scholze [7]. Then we deduce that $\mathbb{V}_{\mathcal{X}, \text{Exp}}$ is a quasi-inverse of $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ via the p -adic Simpson correspondence.

Finally, we prove assertions (i-iii) of theorem 1.1.7 using similar properties for Deninger–Werner vector bundles [10]. Assertion (iv) follows from (iii), as any Higgs bundle over an elliptic curve can be written as a successive extension of Higgs line bundles.

1.1.12. We briefly go over the structure of this article. Section 2 contains a brief review of Tsuji’s interpretation of small Higgs bundles as Higgs crystals over a site and some complements. Section 3 is devoted to the trivializing property of Deninger–Werner Higgs bundles (1.1.11). In section 4, we briefly review Tsuji’s approach to p -adic Simpson correspondence and a local description as well. In section 5, we present the construction of $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ and prove theorem 1.1.6(i) and theorem 1.1.7. Section 6 is devoted to the construction of $\mathbb{V}_{\mathcal{X}, \text{Exp}}$ and the proof of theorem 1.1.6(ii). In the appendix, jointly with T. He, we define the twisted inverse image functor (1.1.3.2) in a more general setting.

Acknowledgement. The author is grateful to Ahmed Abbes for valuable discussions and his comments on this paper. The author would like to thank Takeshi Tsuji for sharing his note on the p -adic Simpson correspondence. He would like to thank Tongmu He, Annette Werner and Xinwen Zhu for useful discussions. The author is partially supported by National Natural Science Foundation of China (No. 11688101).

1.2. Notations and conventions. In this article, \mathbb{N} denotes the set of positive integers.

1.2.1. Let K be a complete discrete valuation ring of characteristic zero, with algebraically closed residue field k of characteristic $p > 0$, \mathcal{O}_K the valuation ring of K , \overline{K} an algebraic closure of K , $\mathcal{O}_{\overline{K}}$ the integral closure of \mathcal{O}_K in \overline{K} , \mathfrak{o} the p -adic completion of $\mathcal{O}_{\overline{K}}$ and \mathbf{C} the fraction field of \mathfrak{o} . We choose a compatible system $(p^{\frac{1}{n}})_{n \geq 1}$ of n -th roots of p in $\mathcal{O}_{\overline{K}}$. For any rational number $\varepsilon > 0$, we set $p^\varepsilon = (p^{\frac{1}{n}})^{n\varepsilon}$, where n is a positive integer such that $n\varepsilon$ is an integer, and $\mathfrak{o}_\varepsilon = \mathfrak{o}/p^\varepsilon \mathfrak{o}$.

We set $S = \text{Spec}(\mathcal{O}_K)$, $\overline{S} = \text{Spec}(\mathcal{O}_{\overline{K}})$ and equip S with the log structure \mathcal{M}_S defined by the special fiber of S . We denote by s , η and $\overline{\eta}$ the special point, generic point and geometrically generic point of S respectively. Let A_{inf} be $W(\mathcal{O}_{\overline{K}}^\flat)$, $\underline{p} = (p, p^{\frac{1}{p}}, \dots) \in (\mathcal{O}_{\overline{K}}^\flat)^\flat$, $\xi = p - [\underline{p}]$ a generator of $\ker(\theta : A_{\text{inf}} \rightarrow \mathfrak{o})$ and B_{dR}^+ the completion of $A_{\text{inf}}[\frac{1}{p}]$ with respect to $\ker(\theta[\frac{1}{p}])$.

Given an S -scheme X , we set $\underline{X} = X \times_S \overline{S}$. We say an S -scheme of finite type X has a *semi-stable reduction* if, étale locally, X is isomorphic to

$$(1.2.1.1) \quad \text{Spec}(\mathcal{O}_K[t_1, \dots, t_n]/(t_1 \cdots t_m - \pi)),$$

where π is a uniformizer of \mathcal{O}_K , and $n \geq m \geq 1$ are integers.

Let X be a scheme (resp. formal scheme, resp. a rigid analytic space over \mathbf{C}) such that \mathcal{O}_X is coherent. Then, we say a coherent \mathcal{O}_X -module is a *vector bundle* if it is locally free of finite type. We denote by $\text{Coh}(X)$ (resp. $\text{VB}(X)$) the category of coherent modules (resp. vector bundles) over X .

1.2.2. A *curve over a field F* means a geometrically connected, separated scheme of finite type over F of dimension one.

Let $T = \text{Spec}(R)$ be the spectrum of a valuation ring R of height one with fraction field F , and \overline{F} an algebraic closure of F . We denote by τ (resp. t) the generic point (resp. special) point of T . Given a proper F -scheme X , an *T -model of X* is a flat, proper T -scheme of finite presentation Y with generic fiber $Y_\tau \simeq X$.

A *T -curve* is a flat, proper T -scheme of finite presentation X of relative dimension one whose generic fiber is a smooth curve over F .

By an abuse of language, we say an \overline{S} -curve X has a semi-stable reduction if there exists a finite extension K' of K , an $S' = \text{Spec}(\mathcal{O}_{K'})$ -curve with semi-stable reduction Y such that $X \simeq Y \times_{S'} \overline{S}$. By semi-stable reduction theorem, an \overline{S} -model X of a smooth proper curve $X_{\overline{\eta}}$ over \overline{K} is dominated by another semi-stable \overline{S} -model X' of $X_{\overline{\eta}}$.

A morphism $\varphi : Y \rightarrow X$ of T -curves is called τ -cover (resp. *generic τ -cover*), if φ is proper of finite presentation and $\varphi_{\tau} : Y_{\tau} \rightarrow X_{\tau}$ is finite étale (resp. finite). Let $\varphi : Y \rightarrow X$ be a τ -cover. We say φ is a *Galois τ -cover*, if φ_{τ} is a Galois cover with automorphism group G , and the action of G on Y_{τ} extends to a X -action on Y .

1.2.3. Almost mathematics. In this article, we consider the category of almost \mathfrak{o} -modules with respect to the maximal ideal \mathfrak{m} of \mathfrak{o} . We denote by $(-)_\#$ the functor

$$(1.2.3.1) \quad \mathbf{Mod}(\mathfrak{o}) \rightarrow \mathbf{Mod}(\mathfrak{o}), \quad M \mapsto M_\# = \text{Hom}_{\mathfrak{o}}(\mathfrak{m}, M).$$

Recall that it sends almost isomorphisms to isomorphisms.

Let \mathcal{A} be an abelian category, $\text{End}(\text{id}_{\mathcal{A}})$ the ring of endomorphisms of the identity functor and $\varphi : \mathfrak{o} \rightarrow \text{End}(\text{id}_{\mathcal{A}})$ a homomorphism. For every object M of \mathcal{A} and $\gamma \in \mathfrak{o}$, $\varphi(\gamma)$ defines an endomorphism $\mu_{\gamma}(M)$ of M . We say M is almost-zero if it is annihilated by every element of \mathfrak{m} . We denote by $\alpha\text{-}\mathcal{A}$ the quotient of \mathcal{A} by the thick full subcategory of almost zero objects. We denote by α the canonical functor

$$\alpha : \mathcal{A} \rightarrow \alpha\text{-}\mathcal{A}, \quad M \mapsto M^{\alpha}.$$

We say a morphism f of \mathcal{A} is an almost isomorphism if $\alpha(f)$ is an isomorphism.

1.2.4. Let T be a topos. The projective systems of objects of T indexed by the ordered set of positive integers \mathbb{N} forms a topos, that we denote by $T^{\mathbb{N}^{\circ}}$.

Given a ring $\check{A} = (A_n)_{n \geq 1}$ of $T^{\mathbb{N}^{\circ}}$, we say an \check{A} -module $M = (M_n)_{n \geq 1}$ is *adic* if for every integers $1 \leq i < j$, the morphism $M_j \otimes_{A_j} A_i \rightarrow M_i$, induced by $M_j \rightarrow M_i$, is an isomorphism.

2. HIGGS CRYSTALS AND HIGGS BUNDLES

In this section, we briefly review the notion of *Higgs crystal* introduced by Tsuji ([4] IV.2), which provides a site-theoretic description of small Higgs bundles. We work with the base $(A_{\text{inf}}, (\xi))$ (c.f. [4] IV.1 Notation).

2.1. Higgs envelope. We first review the notion of Higgs envelope of algebras.

Definition 2.1.1 ([4] IV.2.1.1). (1) We denote by \mathcal{A} the category of A_{inf} -algebras A with a decreasing filtration $\{F^n A\}_{n \geq 0}$ of ideals such that $F^0 A = A$ and $\xi^n A \subset F^n A$ for $n \geq 0$. A morphism in \mathcal{A} is a homomorphism of A_{inf} -algebras compatible with the filtrations.

(2) For $r \in \mathbb{N}_{>0}$ (resp. $r = \infty$), we denote by \mathcal{A}^r the full subcategory of \mathcal{A} consisting of $(A, F^n A)$ satisfying the following conditions:

- (i) The rings A and $A/F^n A$ ($n \in \mathbb{N}_{>0}$) are p -torsion free.
- (ii) The ring A is ξ -torsion free.
- (iii) For every $s \in \mathbb{N} \cap [0, r]$ (resp. $s \in \mathbb{N}$), we have $pF^s A \subset \xi^s A$ (resp. $F^s A = \xi^s A$).
- (iv) The inverse image of $F^n A$ under $\xi : A \rightarrow A$ is $F^{n-1} A$ for every $n \in \mathbb{N}_{>0}$.

(3) We denote by \mathcal{A}_p^r the full subcategory of $\mathcal{A}_{\text{alg}}^r$ consisting of $(A, F^n A) \in \text{Ob} \mathcal{A}_{\text{alg}}^r$ satisfying the following condition: The rings A and $A/F^n A$ ($n \in \mathbb{N}_{>0}$) are p -adically complete and separated.

(4) For $r \in \mathbb{N}_{>0} \cup \{\infty\}$, we define \mathcal{A}^r to be the full subcategory of \mathcal{A}_p^r consisting of $(A, F^n A) \in \text{Ob} \mathcal{A}_p^r$ satisfying the following condition: The natural homomorphism $A \rightarrow \varprojlim_n A/F^n A$ is an isomorphism.

Definition 2.1.2 ([4] IV.2.15). (1) We denote by $\mathcal{A}_{p,\bullet}$ the category of inverse systemes of A_{inf} -algebras $(A_N)_{N \geq 1}$ such that A_N are p -adically complete and separated, $\xi^N A_N = 0$ and the transition maps $A_{N+1} \rightarrow A_N$ are surjective.

(2) For $r \in \mathbb{N}_{>0}$ (resp. $r = \infty$), we denote by \mathcal{A}_{\bullet}^r the full subcategory of $\mathcal{A}_{p,\bullet}$ consisting of (A_N) satisfying the following conditions ([4] IV.2.1.5):

- (i) The rings A_N ($N \in \mathbb{N}_{>0}$) are p -torsion free.
- (ii) For every $s \in \mathbb{N} \cap [0, r]$ (resp. $s \in \mathbb{N}$) and $N \in \mathbb{N}_{>0}$, we have $pF^s A_N \subset \xi^s A_N$ (resp. $F^s A_N = \xi^s A_N$), where $F^n A_N$ ($n \in \mathbb{N}$) is the decreasing filtration of A_N by ideals defined by $F^0 A_N = A_N$, $F^n A_N = \text{Ker}(\pi : A_N \rightarrow A_n)$ ($1 \leq n \leq N$) and $F^n A_N = 0$ ($n > N$).
- (iii) The kernel of the homomorphism $A_N \rightarrow A_N; x \mapsto \xi x$ is $F^{N-1} A_N$ for every $N \in \mathbb{N}_{>0}$.

For $N \geq n$, we set $F^n A_N = \text{Ker}(A_N \rightarrow A_n)$. Then the functor $\mathcal{A}_{p,\bullet} \rightarrow \mathcal{A}$, defined by $(A_N) \mapsto (\varprojlim_N A_N, \varprojlim_N F^n A_N)$ is fully faithful ([4] IV.2.1.7). The projective limit functor induces an equivalence of categories $\mathcal{A}_{\bullet}^r \xrightarrow{\sim} \mathcal{A}^r$ ([4] IV.2.1.7). In summary, we have the following diagram of fully faithful functors:

$$\begin{array}{ccccccc}
 \mathcal{A}_{\bullet}^r & \xrightarrow{\hspace{2cm}} & & \mathcal{A}_{p,\bullet} & & & \\
 \downarrow \sim & & & \downarrow & & & \\
 \mathcal{A}^r & \longrightarrow & \mathcal{A}_p^r & \longrightarrow & \mathcal{A}_{\text{alg}}^r & \longrightarrow & \mathcal{A}
 \end{array}$$

The functors in the second row have left adjoint functors ([4] IV.2.1.2). Then so is the inclusion functor in the first row. We denote the left adjoint functor by $D_{\text{Hig}}^r : \mathcal{A} \rightarrow \mathcal{A}^r$ (resp. $D_{\text{Hig}}^r : \mathcal{A}_{p,\bullet} \rightarrow \mathcal{A}_{\bullet}^r$) and call it *Higgs envelope (of level r)*.

2.1.3. We present an example of Higgs envelope. Let $A = (A_N)$ be an object of \mathcal{A}_{\bullet}^r . If $r \in \mathbb{N}_{>0}$, for integers $N \geq 1, m, n \geq 0$, we set

$$A_N^{(m)r,n} = p^{-n_1} \xi^{n-n_2} F^{n_2} A_N \subset A_N, \quad \text{and} \quad A_N^{(m)r} = \sum_{n \geq 0} A_N^{(m)r,n},$$

where $n_1 = \min\{\lfloor \frac{n-m}{r} \rfloor, 0\}$ and $n_2 = n - m - n_1 r$. If $r = \infty$, we set $A_N^{(m)r} = A_N$.

Let d be an integer ≥ 1 , and define an object $B = (B_N)$ of $\mathcal{A}_{p,\bullet}$ over A by

$$B_1 = A_1, \quad B_N = A_N \{T_1, \dots, T_N\} = \varprojlim_m A_N / p^m A_N [T_1, \dots, T_d], \quad (N \geq 2),$$

$$B_2 \rightarrow B_1, T_i \mapsto 0, \quad B_{N+1} \rightarrow B_N, T_i \mapsto T_i.$$

We define the (A_{inf}/ξ^N) -submodule $A_N[W_1, \dots, W_d]_r$ of $A_N[W_1, \dots, W_d]$ to be

$$\bigoplus_{I \in \mathbb{N}^d} A_N^{(|I|)r} \underline{W}^I,$$

which is an A_N -subalgebra ([4] IV.2.3.13(1)). We define $A_N\{W_1, \dots, W_d\}_r$ to be the p -adic completion of $A_N[W_1, \dots, W_d]_r$ and we obtain an object $(A_N\{W_1, \dots, W_d\}_r)$ of $\mathcal{A}_{p,\bullet}$, denoted by $A\{W_1, \dots, W_d\}_r$.

By ([4] 2.3.15), $A\{W_1, \dots, W_d\}_r$ is actually an object of \mathcal{A}_{\bullet}^r and $D_{\text{Hig}}^r(B)$ with the adjunction morphism $B \rightarrow D_{\text{Hig}}^r(B)$ is isomorphic to the following homomorphism in $\mathcal{A}_{p,\bullet}$ over A :

$$(2.1.3.1) \quad B_N = A_N\{T_1, \dots, T_d\} \rightarrow A_N\{W_1, \dots, W_d\}_r, \quad T_i \mapsto \xi W_i, \quad N \geq 2.$$

In the following, we mainly use the description (2.1.3.1) in the case $r = \infty$, where $A_N\{W_1, \dots, W_d\}_{\infty}$ is the p -adic completion of the polynomial algebra $A_N[W_1, \dots, W_N]$. We will present a simple proof of (2.1.3.1) in this case.

2.1.4. Let $(C, F^n C)$ be an object of \mathcal{A} such that $F^1 C$ is finitely generated by $(\xi, a_i)_{i=1}^m$, $F^n C = \xi^n C$ for $n \geq 2$ and that $(\xi, a_i, p)_{i=1}^m$ is a regular sequence. We denote by $C_0 = C[\frac{F^1 C}{\xi}]$ the dilatation of the dilatation of the ideal $F^1 C$ with respect to ξ ([37] 3.3). We set $\widehat{C}_0 = \varprojlim_{n \geq 1} C_0 / p^n C_0$ and $D = \varprojlim_{n \geq 1} \widehat{C}_0 / \xi^n \widehat{C}_0$.

Then the description (2.1.3.1) in the case $r = \infty$ follows from the following lemma.

Lemma 2.1.5. *With the above assumption, $(D, \xi^n D)$ belongs to \mathcal{A}^{∞} and is isomorphic to Higgs envelope $D_{\text{Hig}}^{\infty}(C, F^n C)$.*

Proof. In view of the proof of ([4] IV.2.1.2), it suffices to show that $(C_0, \xi^n C_0)$ belongs to $\mathcal{A}_{\text{alg}}^\infty$ and is isomorphic to the image of $(C, F^n C)$ via the left adjoint of $\mathcal{A}_{\text{alg}}^\infty \rightarrow \mathcal{A}$. We have an isomorphism of C -algebras:

$$C_0 \simeq C[x_1, \dots, x_m] / (a_i - \xi x_i)_{i=1}^m.$$

As C is ξ -torsion free, then C_0 is a sub-ring of $C[\frac{1}{\xi}]$ and is also ξ -torsion free. Since $(\xi, a_i, p)_{i=1}^m$ is a regular sequence of C , then the quotient $C_0 / \xi C_0 \simeq (C / F^1 C)[x_1, \dots, x_m]$ is p -torsion free, i.e. (ξ, p) is a regular sequence of C_0 . Then we deduce that for every $N \geq 1$, $C_0 / \xi^N C_0$ is p -torsion free. Therefore, $(C_0, \xi^n C_0)$ is an object of $\mathcal{A}_{\text{alg}}^\infty$.

Let $(E, F^n E)$ be an object of $\mathcal{A}_{\text{alg}}^\infty$ and $f : (C, F^n C) \rightarrow (E, F^n E)$ a homomorphism compatible with filtrations. Since E is ξ -torsion free and $f(F^1 C) \subset \xi E$, f induces a homomorphism of C -algebras $g : C_0 \rightarrow E$ compatible with filtrations. This finishes the proof. \square

2.1.6. Let $X \rightarrow \text{Spf}(\mathbb{Z}_p)$ be a p -adic formal scheme (i.e. $p\mathcal{O}_X$ is an ideal of definition) over \mathbb{Z}_p . For $m \geq 1$, we denote by X_m the reduction modulo p^m of X . A *fine log structure* \mathcal{M} on X is a family of fine log structures \mathcal{M}_m on X_m and exact closed immersions $(X_m, \mathcal{M}_m) \rightarrow (X_{m+1}, \mathcal{M}_{m+1})$ ($m \geq 1$) extending the closed immersion $X_m \rightarrow X_{m+1}$ ([4] IV.2.2). We define a *p -adic fine log formal scheme* to be a p -adic formal scheme endowed with a fine log structure. A morphism of p -adic fine log formal schemes $f : (X, \mathcal{M}) \rightarrow (Y, \mathcal{N})$ is a family of morphisms of fine log schemes $f_m : (X_m, \mathcal{M}_m) \rightarrow (Y_m, \mathcal{N}_m)$ ($m \in \mathbb{N}$) compatible with the exact closed immersions $(X_m, \mathcal{M}_m) \rightarrow (X_{m+1}, \mathcal{M}_{m+1})$ and $(Y_m, \mathcal{N}_m) \rightarrow (Y_{m+1}, \mathcal{N}_{m+1})$. We say f is *smooth* (resp. *étale*, resp. an *exact closed immersion*, resp. a *closed immersion*, resp. *strict*, resp. *integral*) if for every $m \geq 1$, f_m is smooth (resp. étale, ...).

Given a p -adic fine log formal scheme X , the étale site $X_{\text{ét}}$ is defined by the category of p -adic fine formal schemes strict étale over X , equipped with étale topology. We define the sheaf of rings \mathcal{O}_X (resp. sheaf of monoids \mathcal{M}_X) on $X_{\text{ét}}$ by $\Gamma(U, \mathcal{O}_X) = \varprojlim_m \Gamma(U_m, \mathcal{O}_{U_m})$ (resp. $\Gamma(U, \mathcal{M}_X) = \varprojlim_m \Gamma(U_m, \mathcal{M}_{U_m})$). We refer to *loc.cit* for more details on p -adic fine log formal schemes.

2.1.7. Next, we review the notion of *Higgs envelope* of p -adic fine log formal schemes following *loc.cit*.

We set $\Sigma_N = \text{Spf}(\mathbb{A}_{\text{inf}}/\xi^N)$, equipped with the trivial log structure. We refer to ([4] IV.2.2.1) for the definition of category \mathcal{C} (over $(\Sigma_N)_{N \geq 1}$) and its full subcategory \mathcal{C}^r for $r \in \mathbb{N} \cup \{\infty\}$. Recall that an object of \mathcal{C} is a sequence of morphisms $\mathcal{Y} = (\mathcal{Y}_1 \rightarrow \mathcal{Y}_2 \rightarrow \dots \rightarrow \mathcal{Y}_N \rightarrow \dots)_{N \geq 1}$ of p -adic fine log formal schemes \mathcal{Y}_N over Σ_N , compatible with $\Sigma_N \rightarrow \Sigma_{N+1}$ such that:

- the morphism $\mathcal{Y}_1 \rightarrow \mathcal{Y}_2$ is an immersion, and for $N \geq 2$, the morphism $\mathcal{Y}_N \rightarrow \mathcal{Y}_{N+1}$ is an exact closed immersion;
- for $N \geq 2$, the morphism of the modulo p reduction of $\mathcal{Y}_N \rightarrow \mathcal{Y}_{N+1}$ is a nilpotent immersion.

An object $\mathcal{Y} = (\mathcal{Y}_N)$ of \mathcal{C} belongs to \mathcal{C}^∞ , if it satisfies moreover the following conditions:

- $\mathcal{Y}_1 \rightarrow \mathcal{Y}_2$ is an exact closed immersion.
- The morphism $p : \mathcal{O}_{\mathcal{Y}_N} \rightarrow \mathcal{O}_{\mathcal{Y}_N}$ is injective for all $N \geq 1$.
- For every $N \geq 1$, $\mathcal{O}_{\mathcal{Y}_N}$ is killed by ξ^N and $\text{Ker}(\mathcal{O}_{\mathcal{Y}_N} \rightarrow \mathcal{O}_{\mathcal{Y}_n}) = \xi^n \mathcal{O}_{\mathcal{Y}_N}$ for $1 \leq n \leq N$.
- $\text{Ker}(\xi : \mathcal{O}_{\mathcal{Y}_N} \rightarrow \mathcal{O}_{\mathcal{Y}_N})$ is $\xi^{N-1} \mathcal{O}_{\mathcal{Y}_N}$.

Given an object $\mathcal{Y} = (\mathcal{Y}_N)_{N \geq 1}$ of \mathcal{C} and an immersion $X \rightarrow \mathcal{Y}_1$ over Σ_1 , we also denote by $(X \rightarrow \mathcal{Y})$ the object $(X \rightarrow \mathcal{Y}_2 \rightarrow \mathcal{Y}_3 \rightarrow \dots)$ of \mathcal{C} . If a caligraphic letter $\mathcal{Y} = (\mathcal{Y}_N)_{N \geq 1}$ denotes an object of \mathcal{C} , we use the correspondence capital letter Y to denote \mathcal{Y}_1 .

For $N \geq 1$, we denote by $\Sigma_{N,S}$ the p -adic formal scheme Σ_N , equipped with the fine log structure extended by \mathcal{M}_S on S (c.f. [4] II.9.6), and by Σ_S the object $(\Sigma_{N,S})_{N \geq 1}$ of \mathcal{C} . Note that, Σ_S belongs to \mathcal{C}^r , for every $r \in \mathbb{N} \cup \{\infty\}$.

Let $f = (f_N)_{N \geq 1} : \mathcal{Y} = (\mathcal{Y}_N) \rightarrow \mathcal{Y}' = (\mathcal{Y}'_N)$ be a morphism of \mathcal{C} . We say f is smooth (resp. étale, resp. integral) if f_N is smooth (resp. étale, resp. integral) for every $N \geq 1$. We say f is *Cartesian* if the morphism $\mathcal{Y}'_N \rightarrow \mathcal{Y}'_{N+1} \times_{\mathcal{Y}_{N+1}} \mathcal{Y}_N$ induced by f is an isomorphism for every $N \geq 1$.

2.1.8. For $r \in \mathbb{N} \cup \{\infty\}$, the natural inclusion $\mathcal{C}^r \rightarrow \mathcal{C}$ admits a right adjoint functor $D_{\text{HIG}}^r : \mathcal{C} \rightarrow \mathcal{C}^r$, called *Higgs envelope of level r* ([4] IV.2.2.2, IV.2.2.9). When $r = \infty$, we simply call it Higgs envelope.

If \mathcal{Y} is an object of \mathcal{C} such that \mathcal{Y}_N are affine and $\mathcal{Y}_1 \rightarrow \mathcal{Y}_2$ is an exact closed immersion, the Higgs envelope of level r of \mathcal{Y} is constructed by the functor $D_{\text{HIG}}^r : \mathcal{A}_{p,\bullet} \rightarrow \mathcal{A}_{\bullet}^r$ applying to $(\Gamma(\mathcal{Y}_N, \mathcal{O}_{\mathcal{Y}_N}))_{N \geq 1}$, and the log structure induced by $(\mathcal{M}_{\mathcal{Y}_N})_{N \geq 1}$ ([4] IV.2.2.14). We refer to ([4] IV.2.2) for more details.

2.1.9. Let $X \rightarrow \Sigma_{1,S}$ be a smooth integral morphism of p -adic fine log formal schemes and $\mathcal{X} = (\mathcal{X}_N)_{N \geq 1} \rightarrow \Sigma_S$ a smooth Cartesian morphism lifting of X (i.e. $\mathcal{X}_1 = X$). Then \mathcal{X} belongs to \mathcal{C}^r for every $r \in \mathbb{N} \cup \{\infty\}$.

Let s be an integer ≥ 1 . We denote by \mathcal{X}^{s+1} the product of $(s+1)$ -copies of \mathcal{X} over Σ_S in \mathcal{C} and $p(s) : \mathcal{D}(s) = (\mathcal{D}(s)_N)_{N \geq 1} \rightarrow (X \rightarrow \mathcal{X}^{s+1})$ the Higgs envelope of level r of the diagonal embedding $(X \rightarrow \mathcal{X}^{s+1})$. We set $\mathcal{D}(s) = \mathcal{D}(s)_1$. We denote by $q_i : \mathcal{D}(1) \rightarrow \mathcal{X}$ (resp. $p_{ij} : \mathcal{D}(2) \rightarrow \mathcal{D}(1)$) be the projection in the i -th component for $i = 1, 2$ (resp. i, j -th component for $1 \leq i < j \leq 3$) and by $\Delta_s : \mathcal{X} \rightarrow \mathcal{D}(s)$ the morphism induced by the diagonal immersion $\mathcal{X} \rightarrow \mathcal{X}^{s+1}$.

Suppose that \mathcal{X} is affine and that there exist $x_i = (x_{i,N}) \in \varprojlim_N \Gamma(\mathcal{X}_N, \mathcal{M}_{\mathcal{X}_N})$ ($1 \leq i \leq d$) ([4] IV.2.2) such that $\{d \log(x_{i,N})\}_{1 \leq i \leq d}$ is a basis of $\Omega_{\mathcal{X}_N/\Sigma_{N,S}}^1$ for $N \geq 1$. By ([4] IV.2.3.17), for $i \in \{1, \dots, d\}$ and $N \geq 2$, there exists a unique element $u_{i,N} \in 1 + F^1 \mathcal{O}_{\mathcal{D}(1)_N}$, such that

$$p(1)_N^*(1 \otimes x_{i,N}) = p(1)_N^*(x_{i,N} \otimes 1)u_{i,N} \in \mathcal{O}_{\mathcal{D}(1)_N},$$

and that there exists canonical isomorphisms of $(\mathcal{O}_{\mathcal{X}_N})_{N \geq 1}$ -algebras in $\mathcal{A}_{\bullet}^\infty$

$$(2.1.9.1) \quad \mathcal{O}_{\mathcal{X}_N}\{W_1, \dots, W_d\}_r \xrightarrow{\sim} q_{j*}(\mathcal{O}_{\mathcal{D}(1)_N}), \quad W_i \mapsto \frac{u_{i,N+1}-1}{\xi}, \quad 1 \leq i \leq d, \quad j = 1, 2.$$

2.2. Higgs crystals.

2.2.1. Let Y be a p -adic fine log formal scheme over $\Sigma_{1,S}$. For $r \in \mathbb{N}_{>0} \cup \{\infty\}$, we denote by $(Y/\Sigma_S)_{\text{HIG}}^r$ (resp. $(Y/\Sigma_S)_{\text{HIG}}^r$) the Higgs site (resp. Higgs topos) defined in ([4] IV.3.1.4). Recall ([4] IV.3.1.1) that an object of $(Y/\Sigma_S)_{\text{HIG}}^r$ is a pair (\mathcal{T}, z) consisting of an object $\mathcal{T} = (\mathcal{T}_N)_{N \geq 1}$ of $\mathcal{C}_{/\Sigma_S}^r$ and a $\Sigma_{1,S}$ -morphism $z : T(= \mathcal{T}_1) \rightarrow Y$. A morphism $(\mathcal{T}', z') \rightarrow (\mathcal{T}, z)$ is a morphism $u : \mathcal{T} \rightarrow \mathcal{T}'$ of $\mathcal{C}_{/\Sigma_S}^r$ such that $z \circ u_1 = z'$.

For an object (\mathcal{T}, z) of $(Y/\Sigma_S)_{\text{HIG}}^r$, the set of coverings $\text{Cov}((\mathcal{T}, z))$ is defined to be

$$\left\{ (u_\alpha : (\mathcal{T}_\alpha, z_\alpha) \rightarrow (\mathcal{T}, z))_{\alpha \in A} \mid \begin{array}{l} \text{(i) } u_\alpha \text{ is strict étale and Cartesian for all } \alpha \in A \\ \text{(ii) } \bigcup_{\alpha \in A} u_{\alpha,1}(T_\alpha) = T \end{array} \right\}$$

The functor $(\mathcal{T}, z) \mapsto \Gamma(T, \mathcal{O}_T)$ defines a sheaf of rings, denoted by $\overline{\mathcal{O}}_{Y/\Sigma_S}^2$ on $(Y/\Sigma_S)_{\text{HIG}}^r$. For each rational number $\alpha \in \mathbb{Q}_{>0}$, we set $\overline{\mathcal{O}}_{Y/\Sigma_S, \alpha} = \overline{\mathcal{O}}_{Y/\Sigma_S}/p^\alpha \overline{\mathcal{O}}_{Y/\Sigma_S}$ and $\overline{\mathcal{O}}_{Y/\Sigma_S, \mathbb{Q}} = \overline{\mathcal{O}}_{Y/\Sigma_S}[\frac{1}{p}]$.

An $\overline{\mathcal{O}}_{Y/\Sigma_S}$ -module \mathcal{M} is equivalent to a collection of data $\{\mathcal{M}_{(\mathcal{T}, z)}, \tau_u\}$, where for each object (\mathcal{T}, z) of $(Y/\Sigma_S)_{\text{HIG}}^r$, $\mathcal{M}_{(\mathcal{T}, z)}$ is an \mathcal{O}_T -module of $T_{\text{ét}}$ and for each morphism $u : (\mathcal{T}', z') \rightarrow (\mathcal{T}, z)$ of $(Y/\Sigma_S)_{\text{HIG}}^r$, τ_u is an $\mathcal{O}_{T'}$ -linear morphism

$$\tau_u : u_1^*(\mathcal{M}_{(\mathcal{T}, z)}) \rightarrow \mathcal{M}_{(\mathcal{T}', z')},$$

satisfying following conditions:

- (i) for any morphism $u : (\mathcal{T}', z') \rightarrow (\mathcal{T}, z)$ such that the underlying morphism $u : \mathcal{T}' \rightarrow \mathcal{T}$ in \mathcal{C} is strict étale and Cartesian, the morphism τ_u is an isomorphism;
- (ii) for any composable morphisms $(\mathcal{T}'', z'') \xrightarrow{v} (\mathcal{T}', z') \xrightarrow{u} (\mathcal{T}, z)$, we have $\tau_v \circ v_1^*(\tau_u) = \tau_{uv}$.

A *Higgs isocrystal of level r* is an $\overline{\mathcal{O}}_{Y/\Sigma_S, \mathbb{Q}}$ -module \mathcal{M} on $(Y/\Sigma_S)_{\text{HIG}}^r$ such that for every morphism u of $(Y/\Sigma_S)_{\text{HIG}}^r$, τ_u is an isomorphism. A *Higgs crystal* is an $\overline{\mathcal{O}}_{Y/\Sigma_S}$ -module \mathcal{M} on $(Y/\Sigma_S)_{\text{HIG}}^\infty$ such that for every morphism u of $(Y/\Sigma_S)_{\text{HIG}}^\infty$, τ_u is an isomorphism³.

²This sheaf of rings is denoted by $\mathcal{O}_{Y/\Sigma_S, 1}$ in [4] IV.3.1.4

³Our definition of (iso-)crystals is slightly different to that of [4] IV.3.3, where certain finiteness condition is imposed.

2.2.2. To introduce a finiteness condition on Higgs (iso-)crystals in a simple way, we assume X is an S -scheme with semi-stable reduction in the following of this section. Let \mathcal{M}_X be the log structure defined by its special fiber and \check{X} the p -adic completion of $X \otimes_{\mathcal{O}_K} \mathfrak{o}$, equipped with the fine log structure induced by \mathcal{M}_X . Then \check{X} is smooth over $\Sigma_{1,S}$.

We denote by \mathbf{Q} ([4] III.10.5) the full subcategory of $\mathbf{\acute{E}t}/_X$ étale schemes over X consisting of affine schemes U such that there exists a fine and saturated chart $M \rightarrow \Gamma(U, \mathcal{M}_X)$ that induces an isomorphism

$$M \xrightarrow{\sim} \Gamma(U, \mathcal{M}_X) / \Gamma(U, \mathcal{O}_U^\times).$$

A Higgs isocrystal of level r (resp. a Higgs crystal) \mathcal{M} is called *finite*, if for any object U of \mathbf{Q} and any smooth Cartesian lifting \mathcal{U} of \check{U} over Σ_S (regarded as an object of $(\check{X}/\Sigma_S)_{\text{HIG}}^r$), the $\mathcal{O}_{\check{U}}[\frac{1}{p}]$ -module (resp. $\mathcal{O}_{\check{U}}$ -module) $\mathcal{M}_{\mathcal{U}}$ is finitely generated and projective (resp. coherent and $\mathcal{M}_{\mathcal{U}}[\frac{1}{p}]$ is projective). In particular, when \check{X} admits a smooth Cartesian lifting \mathcal{X} over Σ_S , this condition is equivalent to the fact that $\mathcal{M}_{\mathcal{X}}$ is finitely generated and locally projective (resp. coherent and $\mathcal{M}_{\mathcal{X}}[\frac{1}{p}]$ is locally projective).

We denote by $\text{HC}_{\mathbb{Q}_p}^r(\check{X}/\Sigma_S)$ (resp. $\text{HC}_{\mathbb{Z}_p}^r(\check{X}/\Sigma_S)$) the category of Higgs isocrystals on $(\check{X}/\Sigma_S)_{\text{HIG}}^r$ (resp. Higgs crystals on $(\check{X}/\Sigma_S)_{\text{HIG}}^\infty$) and by $\text{HC}_{\mathbb{Q}_p, \text{fin}}^r(\check{X}/\Sigma_S)$ (resp. $\text{HC}_{\mathbb{Z}_p, \text{fin}}^r(\check{X}/\Sigma_S)$) the full subcategory consisting of objects which are finite.

The canonical functor $\text{HC}_{\mathbb{Z}_p, \text{fin}}^r(\check{X}/\Sigma_S)_{\mathbb{Q}} \rightarrow \text{HC}_{\mathbb{Q}_p, \text{fin}}^r(\check{X}/\Sigma_S)$ is fully faithful ([4] IV.3.5.1).

Remark 2.2.3. Under the assumption of § 2.2.2, the above finiteness condition is stronger than the condition in ([4] IV.3.3.1(ii) and IV.3.3.2(i,iii)). Hence we can apply results of ([4] IV) to our setting.

2.2.4. Let $\mathcal{X} \rightarrow \Sigma_S$ be a smooth Cartesian lifting of \check{X} in \mathcal{C} . We keep the notation in § 2.1.9.

We denote by $\text{HS}_{\mathbb{Q}_p}^r(\check{X}, \mathcal{X}/\Sigma_S)$ (resp. $\text{HS}_{\mathbb{Z}_p}^r(\check{X}, \mathcal{X}/\Sigma_S)$) the category of pairs (M, ε) consisting of an $\mathcal{O}_{\check{X}}[\frac{1}{p}]$ -module (resp. $\mathcal{O}_{\check{X}}$ -module) M together with an $\mathcal{O}_{D(1)}$ -linear isomorphism $\varepsilon : q_2^*(M) \xrightarrow{\sim} q_1^*(M)$, called *Higgs stratification*, such that: (a) $\Delta_1^*(\varepsilon) = \text{id}_M$; (b) $p_{12}^*(\varepsilon) \circ p_{23}^*(\varepsilon) = p_{13}^*(\varepsilon)$.

By a standard argument ([4] IV.3.4.4), the lifting \mathcal{X} induces equivalences of categories between (2.2.4.1)

$$\text{HC}_{\mathbb{Q}_p}^r(\check{X}/\Sigma_S) \xrightarrow{\sim} \text{HS}_{\mathbb{Q}_p}^r(\check{X}, \mathcal{X}/\Sigma_S) \quad (\text{resp. } \text{HC}_{\mathbb{Z}_p}^r(\check{X}/\Sigma_S) \xrightarrow{\sim} \text{HS}_{\mathbb{Z}_p}^r(\check{X}, \mathcal{X}/\Sigma_S)), \quad \mathcal{M} \mapsto (\mathcal{M}_{\mathcal{X}}, \varepsilon),$$

where ε is constructed by the composition $q_2^*(\mathcal{M}_{\mathcal{X}}) \simeq \mathcal{M}_{D(1)} \simeq q_1^*(\mathcal{M}_{\mathcal{X}})$.

We denote by $\text{HS}_{\mathbb{Q}_p, \text{fin}}^r(\check{X}, \mathcal{X}/\Sigma_S)$ (resp. $\text{HS}_{\mathbb{Z}_p, \text{fin}}^r(\check{X}, \mathcal{X}/\Sigma_S)$) the full subcategory of $\text{HS}_{\mathbb{Q}_p}^r(\check{X}, \mathcal{X}/\Sigma_S)$ (resp. $\text{HS}_{\mathbb{Z}_p}^r(\check{X}, \mathcal{X}/\Sigma_S)$) corresponding to finite Higgs (iso-)crystals (of level r).

2.2.5. We denote by $\tilde{\Omega}_{X/S}^1$ the module of log differentials $\Omega_{(X, \mathcal{M}_X)/(S, \mathcal{M}_S)}^1$ and by $\Omega_{\check{X}/\Sigma_{1,S}}^1$ the p -adic completion of $\tilde{\Omega}_{X/S}^1 \otimes_{\mathcal{O}_K} \mathcal{O}_{\overline{K}}$ over \check{X} . Given an $\mathcal{O}_{\check{X}}$ -module M and an $\mathcal{O}_{\check{X}}$ -linear morphism $\theta : M \rightarrow M \otimes_{\mathcal{O}_{\check{X}}} \xi^{-1} \Omega_{\check{X}/\Sigma_{1,S}}^1$, we can define an $\mathcal{O}_{\check{X}}$ -linear morphism, for $q \geq 1$:

$$\theta^q : M \otimes_{\mathcal{O}_{\check{X}}} \xi^{-q} \Omega_{\check{X}/\Sigma_{1,S}}^q \rightarrow M \otimes_{\mathcal{O}_{\check{X}}} \xi^{-q-1} \Omega_{\check{X}/\Sigma_{1,S}}^{q+1}, \quad x \otimes \xi^{-q} \omega \mapsto \theta(x) \wedge \xi^{-q} \omega.$$

If $\theta^1 \circ \theta = 0$, we say θ is a *Higgs field* and (M, θ) is a *Higgs module*.

Since X admits a semi-stable reduction over S , there exists a strict étale morphism $U \rightarrow \check{X}$ and $(t_i)_{i=1}^d \in \Gamma(U, \mathcal{M}_U)$ such that $\{d \log(t_i)\}_{i=1}^d$ is a basis of $\Omega_{U/\Sigma_{1,S}}^1$. Let θ_i ($1 \leq i \leq d$) be the endomorphism of M defined by $\theta(x) = \sum_{1 \leq i \leq d} \xi^{-1} \theta_i(x) \otimes d \log(t_i)$. The condition $\theta^1 \circ \theta = 0$ is equivalent to the fact that θ_i and θ_j commutes to each other.

Suppose M is a coherent $\mathcal{O}_U[\frac{1}{p}]$ -module (resp. a p -torsion free coherent \mathcal{O}_U -module). We endow M with the p -adic topology and for $x \in \Gamma(U, M)$ and $\underline{m} = (m_i) \in \mathbb{N}^d$, we set $\theta_{\underline{m}}(x) = \prod_{1 \leq i \leq d} \theta_i^{m_i}(x)$. Consider the following convergent condition $(\text{Conv})_r$ (resp. integral condition (Int)) on M ([4] IV.3.4.12):

$(\text{Conv})_r$: $p^{-[\underline{m}|r^{-1}]} \frac{1}{\underline{m}!} \theta_{\underline{m}}(x)$ converges to 0 as $|\underline{m}| \rightarrow \infty$, where $|\underline{m}|r^{-1} = 0$ if $r = \infty$.

(Int) : we have $\frac{1}{\underline{m}!} \theta_{\underline{m}}(x) \in M$.

Definition 2.2.6. (i) We denote by $\mathrm{HM}_{\mathbb{Q}_p}(\check{X}/\Sigma_{1,S})$ (resp. $\mathrm{HM}_{\mathbb{Z}_p}(\check{X}/\Sigma_{1,S})$) the category of pairs (M, θ) consisting of an $\mathcal{O}_{\check{X}}[\frac{1}{p}]$ -module (resp. $\mathcal{O}_{\check{X}}$ -module) together with a Higgs field θ , and by $\mathrm{HB}_{\mathbb{Q}_p}(\check{X}/\Sigma_{1,S})$ the full subcategory of $\mathrm{HM}_{\mathbb{Q}_p}(\check{X}/\Sigma_{1,S})$ whose underlying $\mathcal{O}_{\check{X}}[\frac{1}{p}]$ -module is locally projective of finite type.

(ii) We denote by $\mathrm{HB}_{\mathbb{Q}_p, \text{small}}^r(\check{X}/\Sigma_{1,S})$ (resp. $\mathrm{HB}_{\mathbb{Z}_p, \text{conv}}^r(\check{X}/\Sigma_{1,S})$) the full subcategory of $\mathrm{HM}_{\mathbb{Q}_p}(\check{X}/\Sigma_{1,S})$ (resp. $\mathrm{HM}_{\mathbb{Z}_p}(\check{X}/\Sigma_{1,S})$) consisting of (M, θ) , where M is a finitely generated locally projective $\mathcal{O}_{\check{X}}[\frac{1}{p}]$ -module (resp. a p -torsion free coherent $\mathcal{O}_{\check{X}}$ -module such that $M[\frac{1}{p}]$ is locally projective) and θ satisfies the following condition: There exist a strict étale covering $(U_\alpha \rightarrow \check{X})_{\alpha \in A}$ and $(t_{i,\alpha})_{i=1}^d \in \Gamma(U_\alpha, \mathcal{M}_{U_\alpha})$ such that

- $d \log t_{i,\alpha}$ is a basis of $\Omega_{U_\alpha/\Sigma_{1,S}}^1$ for every $N \geq 1$,
- U_α is affine
- the pair $(\Gamma(U_\alpha, M), \Gamma(U_\alpha, \theta))$ satisfies $(\text{Conv})_r$ (resp. (Int)) with respect to $t_{i,\alpha}$.

Proposition 2.2.7 ([4] IV.3.4.16). *Let \mathcal{X} be a smooth Cartesian lifting of \check{X} over Σ_S .*

(i) *The lifting \mathcal{X} induces a canonical functor:*

(2.2.7.1)

$$\mathrm{HS}_{\mathbb{Q}_p}^r(\check{X}, \mathcal{X}/\Sigma_S) \rightarrow \mathrm{HM}_{\mathbb{Q}_p}^r(\check{X}/\Sigma_{1,S}) \quad (\text{resp. } \mathrm{HS}_{\mathbb{Z}_p}(\check{X}, \mathcal{X}/\Sigma_S) \rightarrow \mathrm{HM}_{\mathbb{Z}_p}(\check{X}/\Sigma_{1,S})), \quad (M, \varepsilon) \mapsto (M, \theta).$$

(ii) *The functor (2.2.7.1) induces an equivalence of categories $\iota_{\mathcal{X}}$ between $\mathrm{HB}_{\mathbb{Q}_p, \text{small}}^r(\check{X}/\Sigma_S)$ (resp. $\mathrm{HB}_{\mathbb{Z}_p, \text{conv}}^r(\check{X}/\Sigma_S)$) and $\mathrm{HC}_{\mathbb{Q}_p, \text{fin}}^r(\check{X}/\Sigma_S)$ (resp. full subcategory of $\mathrm{HC}_{\mathbb{Z}_p, \text{fin}}(\check{X}/\Sigma_S)$ such that $\mathcal{M}_{\mathcal{X}}$ is p -torsion free).*

We briefly review the construction of functor (2.2.7.1). Let $\mathcal{D}(1)_N^1$ be the first infinitesimal neighborhood of $\Delta_{1,N} : \mathcal{X}_N \rightarrow \mathcal{D}(1)_N$. Then there exists a canonical isomorphism ([4] (IV.2.4.7))

$$\mathrm{Ker}(\mathcal{O}_{\mathcal{D}(1)_N^1} \rightarrow \mathcal{O}_{\mathcal{X}_N}) \xrightarrow{\sim} \xi^{-1} \Omega_{\mathcal{X}_N/\Sigma_{N,S}}^1.$$

Given (M, ε) as above, ε induces an isomorphism $\varepsilon^1 : \mathcal{O}_{D(1)^1} \otimes M \rightarrow M \otimes \mathcal{O}_{D(1)^1}$ and θ is defined by

$$\theta : M \rightarrow \xi^{-1} M \otimes_{\mathcal{O}_{\check{X}}} \Omega_{\check{X}/\Sigma_{1,S}}^1, \quad x \mapsto \varepsilon^1(1 \otimes x) - x \otimes 1.$$

2.2.8. We have a canonical isomorphism of rigid-analytic spaces $X_{\mathbb{C}}^{\text{an}} \simeq \check{X}^{\text{rig}}$. Let $\check{X}_{\text{ad}}^{\text{rig}}$ be the rigid space \check{X}^{rig} , equipped with the admissible topology and

$$\rho_{\check{X}} : (\check{X}_{\text{ad}}^{\text{rig}}, \mathcal{O}_{\check{X}^{\text{rig}}}) \rightarrow (\check{X}, \mathcal{O}_{\check{X}})$$

the morphism of ringed topoi ([1] 4.7.5.2). We have the following commutative diagram:

$$(2.2.8.1) \quad \begin{array}{ccc} \mathrm{Coh}(\mathcal{O}_{X_{\mathbb{C}}}) & \xrightarrow[\sim]{\widehat{}} & \mathrm{Coh}(\mathcal{O}_{\check{X}}) \\ \downarrow & & \downarrow \rho_{\check{X}}^* \\ \mathrm{Coh}(\mathcal{O}_{X_{\mathbb{C}}}) & \xrightarrow[\sim]{\text{an}} & \mathrm{Coh}(\mathcal{O}_{X_{\mathbb{C}}^{\text{an}}}) \end{array}$$

where horizontal functors are the p -adic completion functor and analytification functor and induce equivalences of categories, the vertical functor is essentially surjective and is fully faithful up to isogeny.

Proposition 2.2.9. (i) *The analytification functor induces an equivalence of categories $\mathrm{VB}(X_{\mathbb{C}}) \xrightarrow{\sim} \mathrm{VB}(X_{\mathbb{C}}^{\text{an}})$.*

(ii) *The functor $\rho_{\check{X}}^*$ induces an equivalence between the category $\mathrm{LP}(\mathcal{O}_{\check{X}}[\frac{1}{p}])$ of locally projective $\mathcal{O}_{\check{X}}[\frac{1}{p}]$ -modules of finite type and $\mathrm{VB}(X_{\mathbb{C}}^{\text{an}})$.*

Proof. Assertion (i) follows from ([36] proposition 4.7).

(ii) By ([36] 5.15), $\rho_{\check{X}}^*$ sends $\mathrm{LP}(\mathcal{O}_{\check{X}}[\frac{1}{p}])$ to $\mathrm{VB}(X_{\mathbb{C}}^{\text{an}})$. It remains to show it is essentially surjective.

Let E be a vector bundle over $X_{\mathbb{C}}$ and \mathcal{E} a torsion-free coherent $\mathcal{O}_{X_{\mathbb{C}}}$ -module with generic fiber E . For any affine open subscheme $U = \mathrm{Spec}(R)$ of $X_{\mathbb{C}}$ such that $U \cap X_s$ is non-empty, the $R[\frac{1}{p}]$ -module $\mathcal{E}(U)[\frac{1}{p}]$ is finite projective. Then the $\widehat{R}[\frac{1}{p}]$ -module $\widehat{\mathcal{E}(U)}[\frac{1}{p}] \simeq \mathcal{E}(U)[\frac{1}{p}] \otimes_R \widehat{R}$ is also finite projective and $\widehat{\mathcal{E}}[\frac{1}{p}]$ is therefore locally projective of finite type. Then the essential surjectivity follows. \square

Definition 2.2.10 ([13]). (i) Let M be a coherent $\mathcal{O}_{\check{X}}$ -module and θ a Higgs field on M .

For $\alpha \in \mathbb{Q}_{>0}$, we say θ is α -small if $\theta(M) \subset p^\alpha M \otimes_{\mathcal{O}_{\check{X}}} \xi^{-1} \Omega_{\check{X}/\Sigma_{1,S}}^1$.

We say θ is *small* if it is α -small for some $\alpha > \frac{1}{p-1}$.

(ii) Let \mathcal{N} be a locally projective $\mathcal{O}_{\check{X}}[\frac{1}{p}]$ -module and θ a Higgs field on \mathcal{N} .

We say θ is *small* if there exists a coherent sub- $\mathcal{O}_{\check{X}}$ -module \mathcal{N}° of \mathcal{N} such that it generates \mathcal{N} over $\mathcal{O}_{\check{X}}[\frac{1}{p}]$ and that the restriction of θ on \mathcal{N}° is a small in the sense of (i).

Given a p -torsion free coherent $\mathcal{O}_{\check{X}}$ -module M , a small Higgs fields on M satisfies condition (Int).

Given a locally projective $\mathcal{O}_{\check{X}}[\frac{1}{p}]$ -module \mathcal{N} of finite type, θ is small if and only if (\mathcal{N}, θ) belongs to $\text{HB}_{\mathbb{Q}_p, \text{small}}^r(\check{X}/\Sigma_{1,S})$ for some $r \in \mathbb{N}$ ([4] IV.3.6.6).

We denote by $\text{HB}(X_{\mathbb{C}})$ (resp. $\text{HB}(X_{\mathbb{C}}^{\text{an}})$) the category of pairs (M, θ) of a vector bundle M and a Higgs field $\theta : M \rightarrow M \otimes_{\mathcal{O}_{X_{\mathbb{C}}}} \xi^{-1} \Omega_{X_{\mathbb{C}}/\mathbb{C}}^1$ (resp. $\theta : M \rightarrow M \otimes_{\mathcal{O}_{X_{\mathbb{C}}^{\text{an}}}} \xi^{-1} \Omega_{X_{\mathbb{C}}^{\text{an}}/\mathbb{C}}^1$). By proposition 2.2.9, we have equivalences of categories

$$(2.2.10.1) \quad \text{HB}_{\mathbb{Q}_p}(\check{X}/\Sigma_{1,S}) \xrightarrow{\sim} \text{HB}(X_{\mathbb{C}}^{\text{an}}), \quad \text{HB}(X_{\mathbb{C}}) \xrightarrow{\sim} \text{HB}(X_{\mathbb{C}}^{\text{an}}).$$

We say a Higgs bundle (M, θ) of $\text{HB}(X_{\mathbb{C}}^{\text{an}})$ (resp. $\text{HB}(X_{\mathbb{C}})$) is *small (with respect to \check{X})* if there exists a small Higgs bundle of $\text{HB}_{\mathbb{Q}_p}(\check{X}/\Sigma_{1,S})$ associated to (M, θ) via the above equivalences.

Corollary 2.2.11. *Let \mathcal{M} be an object of $\text{HC}_{\mathbb{Z}_p, \text{fin}}(\check{X}/\Sigma_S)$ such that $M = \mathcal{M}_{\mathcal{X}}$ is p -torsion free and ε (resp. θ) the associated Higgs stratification (resp. field) on M . If θ is α -small and set $\beta = \alpha - \frac{1}{p-1}$, then for any local sections $a \in \mathcal{O}_{D(1)}, x \in M$, we have*

$$(2.2.11.1) \quad \varepsilon(a \otimes x) \equiv x \otimes a \pmod{p^\beta}.$$

In particular, if $M/p^\beta M$ is moreover trivial, then there exists an isomorphism of Higgs crystals modulo p^β :

$$\mathcal{M}/p^\beta \mathcal{M} \simeq (\overline{\mathcal{O}}_{\check{X}/\Sigma_S, \beta})^{\oplus r}.$$

Proof. Let $U \rightarrow \check{X}$ be a strict étale morphism as in 2.2.5. With the notations of 2.1.9 and 2.2.5, the Higgs stratification ε can be written, for local section $m \in M$, as:

$$\varepsilon(1 \otimes x) = \sum_{\underline{m} \in \mathbb{N}^d} \frac{\theta_{\underline{m}}(x)}{\underline{m}!} \otimes \prod_{1 \leq i \leq d} \left(\frac{u_{i,2} - 1}{\xi} \right)^{m_i}.$$

Since θ is α -small, we deduce that $\frac{\theta_{\underline{m}}(x)}{\underline{m}!} \equiv 0 \pmod{p^\beta}$. Then the congruence (2.2.11.1) follows.

The second assertion follows from the equivalence (2.2.4.1). \square

2.3. Inverse image of Higgs crystals. In this subsection, we discuss the inverse image functor for Higgs crystals and its realization in terms of modules with Higgs stratification (2.2.4) and Higgs modules.

2.3.1. Let K' be a finite extension of K in \overline{K} and $S' = \text{Spec}(\mathcal{O}_{K'})$. With the notations of § 2.1.7, there exists a canonical morphism $\Sigma_{S'} \rightarrow \Sigma_S$ in \mathcal{C} .

Let X' be a projective S' -scheme with semi-stable reduction, $\mathcal{M}_{X'}$ the log structure defined by its special fiber and \check{X}' the p -adic completion of $X' \otimes_{\mathcal{O}_{K'}} \mathfrak{o}$, equipped with the fine log structure induced $\mathcal{M}_{X'}$. Let $f : X' \rightarrow X_{S'} = X \times_S S'$ be an S' -morphism, which is compatible with logarithmic structures. We denote by $\check{f} : \check{X}' \rightarrow \check{X}$ the morphism of p -adic fine log formal schemes over $\Sigma_{1,S}$ induced by f . For $r', r \in \mathbb{N}_{>0} \cup \{\infty\}$ such that $r' \geq r$, the morphism \check{f} induces a morphism of topoi ([4] IV.3.1.5)

$$\check{f}_{\text{HIG}} : (\check{X}'/\Sigma_{S'})^{r', \sim} \rightarrow (\check{X}/\Sigma_S)^{r, \sim},$$

such that $\check{f}_{\text{HIG}}^{-1}(\overline{\mathcal{O}}_{\check{X}/\Sigma_S}) = \overline{\mathcal{O}}_{\check{X}'/\Sigma_{S'}}$. Its inverse image functors preserve Higgs (iso-)crystals:

$$\check{f}_{\text{HIG}}^* : \text{HC}_{\mathbb{Q}_p}^r(\check{X}/\Sigma_S) \rightarrow \text{HC}_{\mathbb{Q}_p}^{r'}(\check{X}'/\Sigma_{S'}), \quad \text{HC}_{\mathbb{Z}_p}(\check{X}/\Sigma_S) \rightarrow \text{HC}_{\mathbb{Z}_p}(\check{X}'/\Sigma_{S'}).$$

If a Higgs crystal \mathcal{M} is finite, then so is its inverse image $\check{f}_{\text{HIG}}^*(\mathcal{M})$. When $S' = S$ and $X' = X$, the above functor, defined for Higgs isocrystals of different levels, is fully faithful ([4] IV.3.5.3).

In the following, we focus on the inverse image for Higgs (iso-)crystals (i.e. $r' = r = \infty$). We ignore the superscript ∞ from the notations $\text{HS}_{\mathbb{Q}_p}^\infty$ and $\text{HB}_{\mathbb{Q}_p}^\infty$.

Let \mathcal{X} (resp. \mathcal{X}') be a smooth Cartesian lifting of \check{X} over Σ_S (resp. \check{X}' over $\Sigma_{S'}$) and $\mathcal{D}(s)$ (resp. $\mathcal{D}'(s)$) the Higgs envelope of $X \rightarrow \mathcal{X}^{s+1}$ (resp. $X' \rightarrow \mathcal{X}'^{s+1}$). By (2.2.4.1) and 2.2.7(ii), for $\bullet \in \{\mathbb{Z}_p, \mathbb{Q}_p\}$, we obtain twisted inverse image functors for modules with stratifications and convergent Higgs bundles

$$(2.3.1.1) \quad \check{f}_{\text{HIG}, \mathcal{X}', \mathcal{X}}^* : \text{HS}_\bullet(\check{X}, \mathcal{X}/\Sigma_S) \rightarrow \text{HS}_\bullet(\check{X}', \mathcal{X}'/\Sigma_{S'}), \quad \text{HB}_{\bullet, \text{conv}}(\check{X}/\Sigma_{1,S}) \rightarrow \text{HB}_{\bullet, \text{conv}}(\check{X}'/\Sigma_{1,S'}).$$

Lemma 2.3.2. *Suppose $\tilde{f} : \mathcal{X}' \rightarrow \mathcal{X}$ is a lifting of $f : \check{X}' \rightarrow \check{X}$ over Σ_S (which locally exists). Then it induces canonical morphisms $\mathcal{D}'(s) \rightarrow \mathcal{D}(s)$ compatible with natural projections and the diagonal embedding $\mathcal{X} \rightarrow \mathcal{D}(s)$ (c.f. §2.1.9).*

Proof. We have a commutative diagram

$$\begin{array}{ccccc} \mathcal{D}'(s) & \longrightarrow & \check{X}' & \xrightarrow{\check{f}} & \check{X} \\ \downarrow & & \Delta \downarrow & & \Delta \downarrow \\ \mathcal{D}(s) & \longrightarrow & \mathcal{X}'^{s+1} & \xrightarrow{\tilde{f}^{s+1}} & \mathcal{X}^{s+1} \end{array}$$

Then by the universal property of Higgs envelope, we deduces a canonical morphism $\mathcal{D}'(s) \rightarrow \mathcal{D}(s)$ above \tilde{f}^{s+1} . The compatibility can be verified in a similar way. \square

2.3.3. Keep the assumption of lemma 2.3.2. Then, for $\bullet \in \{\mathbb{Z}_p, \mathbb{Q}_p\}$, the morphism \tilde{f} induces functors:

$$(2.3.3.1) \quad \tilde{f}^* : \text{HS}_\bullet(\check{X}, \mathcal{X}/\Sigma_S) \rightarrow \text{HS}_\bullet(\check{X}', \mathcal{X}'/\Sigma_{S'}), \quad (M, \varepsilon) \mapsto (\check{f}^* M, \varepsilon \otimes_{\mathcal{O}_{D(1)}} \mathcal{O}_{D(1)'})$$

$$(2.3.3.2) \quad \tilde{f}^* : \text{HB}_{\bullet, \text{conv}}(\check{X}/\Sigma_{1,S}) \rightarrow \text{HB}_{\bullet, \text{conv}}(\check{X}'/\Sigma_{1,S'}), \quad (M, \theta) \mapsto \check{f}^*(M, \theta),$$

which are compatible with the twisted inverse image functor (2.3.1.1) via functors (2.2.4.1) and (2.2.7.1). The restriction of \tilde{f}^* to underlying $\mathcal{O}_{\check{X}}$ -modules coincides with the usual inverse image functor \check{f}^* .

Proposition 2.3.4. (i) *Suppose \tilde{f}' is another lifting of \check{f} . There exists a natural isomorphism of functors $\alpha_{\tilde{f}, \tilde{f}'} : \tilde{f}'^* \xrightarrow{\sim} \tilde{f}^*$ (2.3.3.1).*

(ii) *The isomorphisms $\alpha_{\tilde{f}, \tilde{f}'}$ satisfy a cocycle condition for three liftings of \check{f} .*

(iii) *By gluing local constructions, we obtain inverse image functors independent of the choice lifting \tilde{f} :*

$$(2.3.4.1) \quad \check{f}_{\mathcal{X}', \mathcal{X}}^* : \text{HS}_\bullet(\check{X}, \mathcal{X}/\Sigma_S) \rightarrow \text{HS}_\bullet(\check{X}', \mathcal{X}'/\Sigma_{S'}), \quad \text{HB}_{\bullet, \text{conv}}(\check{X}/\Sigma_{1,S}) \rightarrow \text{HB}_{\bullet, \text{conv}}(\check{X}'/\Sigma_{1,S'}).$$

The above functor is isomorphic to the functor (2.3.1.1).

Proof. By proposition 2.2.7, it suffices to prove the proposition for the categories HS .

(i) By the universal property of Higgs envelope, the canonical morphism $(\tilde{f}, \tilde{f}') : \mathcal{X}' \rightarrow \mathcal{X}^2$ induces a canonical morphism over Σ_S :

$$(2.3.4.2) \quad \alpha : \mathcal{X}' \rightarrow \mathcal{D}(1).$$

Given an object (M, ε) of $\text{HS}_\bullet(\check{X}, \mathcal{X}/\Sigma_S)$, then we obtain an isomorphism

$$\alpha^*(\varepsilon) : \tilde{f}'^*(M) \xrightarrow{\sim} \tilde{f}^*(M).$$

It remains to show that this isomorphism is compatible with $\mathcal{O}_{D'(1)}$ -stratifications. Indeed, we need to show the following diagram commutes:

$$\begin{array}{ccc} \mathcal{O}_{D(1)'} \otimes_{\mathcal{O}_{\tilde{X}'}} \tilde{f}'^*(M) & \xrightarrow{\tilde{f}'^*(\varepsilon)} & \tilde{f}'^*(M) \otimes_{\mathcal{O}_{\tilde{X}'}} \mathcal{O}_{D(1)'} \\ \mathcal{O}_{D(1)'} \otimes \alpha \downarrow & & \downarrow \alpha \otimes \mathcal{O}_{D(1)'} \\ \mathcal{O}_{D(1)'} \otimes_{\mathcal{O}_{\tilde{X}'}} \tilde{f}^*(M) & \xrightarrow{\tilde{f}^*(\varepsilon)} & \tilde{f}^*(M) \otimes_{\mathcal{O}_{\tilde{X}'}} \mathcal{O}_{D(1)'} \end{array}$$

The morphism $(\tilde{f}', \tilde{f}) : \mathcal{X}'^2 \rightarrow \mathcal{X}^2$ induces a canonical morphism $(\tilde{f}', \tilde{f}) : \mathcal{D}'(1) \rightarrow \mathcal{D}(1)$, which coincides following two compositions:

$$\mathcal{D}'(1) \xrightarrow{(\tilde{f}', \tilde{f}), \alpha \circ q_1} \mathcal{D}(1) \times_{\mathcal{D}} \mathcal{D}(1) \simeq \mathcal{D}(2) \xrightarrow{q_{13}} \mathcal{D}(1), \quad \mathcal{D}'(1) \xrightarrow{\alpha \circ q_2, (\tilde{f}, \tilde{f})} \mathcal{D}(1) \times_{\mathcal{D}} \mathcal{D}(1) \simeq \mathcal{D}(2) \xrightarrow{q_{13}} \mathcal{D}(1).$$

By the cocycle condition of ε , we deduce that

$$(\alpha \otimes \mathcal{O}_{D(1)'}) \circ \tilde{f}'^*(\varepsilon) \simeq (\tilde{f}', \tilde{f})^*(\varepsilon) \simeq \tilde{f}^*(\varepsilon) \circ (\mathcal{O}_{D(1)'} \otimes \alpha).$$

Hence the above diagram is commutative. In this way, we construct a natural isomorphism $\alpha_{\tilde{f}, \tilde{f}'} : \tilde{f}'^* \xrightarrow{\sim} \tilde{f}^*$ and finish the proof of assertion (i).

Assertion (ii) follows from the cocycle condition for stratifications.

(iii) By gluing local constructions, we obtain functors (2.3.4.1). In view of the equivalences between Higgs crystals and modules with Higgs stratification, we deduce the second claim. \square

Remark 2.3.5. We present a local description of the isomorphism $\alpha^*(\varepsilon)$. Suppose \mathcal{X} is affine and there exist $t_i = (t_{i,N}) \in \varprojlim_N \Gamma(\mathcal{X}_N, \mathcal{M}_{\mathcal{X}_N})$ ($1 \leq i \leq d$) such that $\{d \log(t_{i,N})\}_{1 \leq i \leq d}$ is a basis of $\Omega_{\mathcal{X}_N/\Sigma_N, S}^1$ for $N \geq 1$. We keep the notation of 2.1.9. Then the homomorphism $\mathcal{O}_{D(1)} \rightarrow \mathcal{O}_{\tilde{X}'}$ (2.3.4.2) sends

$$\frac{u_{i,2} - 1}{\xi} \mapsto \frac{\tilde{f}'^*(t_{i,2}) \tilde{f}^*(t_{i,2})^{-1} - 1}{\xi},$$

where $\tilde{f}'^*(t_{i,2}) \tilde{f}^*(t_{i,2})^{-1}$ denotes an element $v_{i,2} \in 1 + \xi \mathcal{O}_{\mathcal{X}'_2}$ such that $\tilde{f}'^*(t_{i,2}) = v_{i,2} \tilde{f}^*(t_{i,2})$. Let (M, ε) be an object of $\text{HS}_{\bullet}(X, \mathcal{X}/\Sigma_S)$ and θ the associated Higgs field. The $\mathcal{O}_{D(1)}$ -stratification ε on M is locally defined, for $x \in M$, by

$$\varepsilon(1 \otimes x) = \sum_{\underline{m} \in \mathbb{N}^d} \frac{\theta_{\underline{m}}(x)}{\underline{m}!} \otimes \prod_{i=1}^d \left(\frac{u_{i,2} - 1}{\xi} \right)^{m_i}.$$

Note that this formula p -adically converges as θ satisfies $(\text{Conv})_r$ (resp. (Int)) (2.2.5).

Then the isomorphism $\alpha^*(\varepsilon) : \tilde{f}'^*(M) \xrightarrow{\sim} \tilde{f}^*(M)$ is locally given by:

$$(2.3.5.1) \quad \alpha^*(\varepsilon)(\check{f}^*(x)) = \sum_{\underline{m} \in \mathbb{N}^d} \check{f}^* \left(\frac{\theta_{\underline{m}}(x)}{\underline{m}!} \right) \otimes \prod_{i=1}^d \left(\frac{\tilde{f}'^*(t_{i,2}) \tilde{f}^*(t_{i,2})^{-1} - 1}{\xi} \right)^{m_i}.$$

Corollary 2.3.6. *Let (M, θ) be a Higgs bundle of $\text{HB}_{\mathbb{Z}_p, \text{conv}}(\tilde{X}/\Sigma_{1,S})$.*

(i) *If the morphism $df : f^* \tilde{\Omega}_{X/S}^1 \rightarrow \tilde{\Omega}_{X'/S'}^1$ of log differentials (2.2.5) is divisible by p^n , then the Higgs field of $\check{f}_{\mathcal{X}', \mathcal{X}}^*(M, \theta)$ is trivial modulo p^n .*

(ii) *If the Higgs field θ is α -small for $\alpha > \frac{1}{p-1}$, for $\beta = \alpha - \frac{1}{p-1}$, the reduction modulo p^β of the underlying bundle of $\check{f}_{\mathcal{X}', \mathcal{X}}^*(M, \theta)$ is canonically isomorphic to that of $\check{f}^*(M)$.*

Proof. (i) The assertion being local property, we may assume that there exists a lifting $\tilde{f} : \mathcal{X}' \rightarrow \mathcal{X}$ of \check{f} . Then the assertion follows from the formula (2.3.3.2).

(ii) If a local lifting \tilde{f} of \check{f} exists, the underlying bundle of $\check{f}_{\check{X}', \check{X}}^*(M, \theta)$ is isomorphic to $\check{f}^*(M)$. These local constructions are glued by the formula (2.3.5.1), which is congruent to the identity modulo p^β . Then the assertion follows. \square

Remark 2.3.7. The above construction also applies to the inverse image functoriality of Cartier transform of Ogus-Vologodsky in characteristic $p > 0$ and its lifting modulo p^n via Oyama topoi [24, 25].

2.4. Twisted inverse image for small Higgs bundles via exponential. We present an alternative construction of the twisted inverse image functor for small Higgs bundles along a finite morphism of smooth proper p -adic curves. This approach will be generalized to all Higgs bundles over p -adic curves in appendix A.

2.4.1. We briefly review the notion of *spectral cover* over a curve following [6]. Let C be a smooth projective curve over \mathbf{C} (1.2.2) and L a line bundle over C . We denote by $\text{HB}(C, L)$ the category of pairs (M, θ) consisting of a vector bundle M and an \mathcal{O}_C -linear map $\theta : M \rightarrow M \otimes L$. Let (M, θ) be such a pair of rank r . For $0 \leq i \leq r$, the map θ induces a canonical morphism $\wedge^i M \otimes \wedge^i M^\vee \rightarrow L^{\otimes i}$ and we denote by $c_i(\theta)$ (or simply c_i , if there is no confusion) the image of $(-1)^i \text{id}_{\wedge^i M}$ in $\Gamma(C, L)$ via this map. Then θ satisfies the *characteristic polynomial*:

$$\theta^r + c_1 \theta^{r-1} + \cdots + c_r = 0 \quad \in \text{Hom}(M, M \otimes L^{\otimes r}).$$

The above construction underlies to the Hitchin map h from the moduli stack of pairs (M, θ) of $\text{HB}(C, L)$ of rank r , to the Hitchin base $B := \oplus_{i=1}^r \Gamma(C, L^{\otimes i})$, sending (M, θ) to $(c_i(\theta))_{i=1}^d$ [20]. The spectral cover, reviewed below, shows that the Hitchin map is surjective.

For a closed point $c = (c_i)_{i=1}^r$ of the Hitchin base B , we denote by \mathcal{I}_c the ideal sheaf of $\text{Sym}_{\mathcal{O}_C}(L)$ locally generated by

$$(2.4.1.1) \quad \{(a, c_1 \cdot a, \dots, c_r \cdot a) \mid a \in L^{\otimes r}\} \subset L^{\otimes r} \oplus \cdots \oplus L \oplus \mathcal{O}_C,$$

and we set

$$\mathcal{A}_c = \text{Sym}_{\mathcal{O}_C}(L) / \mathcal{I}_c.$$

The finite morphism $\pi_c : C_c := \text{Spec}_{\mathcal{O}_C}(\mathcal{A}_c) \rightarrow C$ (resp. the algebra \mathcal{A}_c) is called *spectral cover* of c . The direct image functor $\pi_{c,*}$ induces an equivalence between invertible \mathcal{A}_c -modules and pairs (M, θ) of $\text{HB}(C, L)$ such that $h(M, \theta) = c$ ([6] proposition 3.6). For such a pair (M, θ) , we also denote the ideal sheaf and spectral cover associated to $h(M, \theta)$ by $\mathcal{I}_\theta, \mathcal{A}_\theta$ (resp. $\pi_\theta : C_\theta \rightarrow C$).

We usually work with $L = \xi^{-1} \Omega_C^1$, where a pair (M, θ) of $\text{HB}(X, L)$ is the same as a Higgs bundle.

In the following, let X be a projective S -curve with semi-stable reduction (1.2.2) and \check{X} the p -adic completion of $X_\circ = X \otimes_{\mathcal{O}_K} \mathfrak{o}$, equipped with the fine log structure induced by \mathcal{M}_X . We simply write $\Omega_{\check{X}/\Sigma_{1,S}}^1$ as $\Omega_{\check{X}}^1$ and we denote its dual by $T_{\check{X}}$.

Here is a criterion for the smallness of a Higgs bundle in terms of its image in the Hitchin base.

Proposition 2.4.2. *A Higgs bundle (M, θ) of rank r over $X_{\mathbf{C}}$ is small if and only if there exists a rational number $\alpha > \frac{1}{p-1}$ such that $c_i(\theta) \in p^{\alpha i} \Gamma(\check{X}, \xi^{-i} \Omega_{\check{X}}^{\otimes i}) \subset \Gamma(X_{\mathbf{C}}, \xi^{-i} \Omega_{X_{\mathbf{C}}}^{\otimes i})$ for every $1 \leq i \leq r$.*

Proof. Assume (M, θ) is small. There exists a coherent $\mathcal{O}_{\check{X}}$ -module \mathcal{M} with generic fiber M and a rational number $\alpha > \frac{1}{p-1}$ such that $\theta(\mathcal{M}) \subset p^\alpha \mathcal{M} \otimes_{\mathcal{O}_{\check{X}}} \xi^{-1} \Omega_{\check{X}}^1$. For each $i \geq 1$, $\wedge^i \theta$ sends $\wedge^i \mathcal{M}$ to $p^{\alpha i} \wedge^i \mathcal{M} \otimes_{\mathcal{O}_{\check{X}}} \xi^{-i} \Omega_{\check{X}}^{\otimes i}$. Then we deduce $c_i(\theta) \in p^{\alpha i} \Gamma(\check{X}, \xi^{-i} \Omega_{\check{X}}^{\otimes i})$.

On the other hand, assume $c_i(\theta) \in p^{\alpha i} \Gamma(\check{X}, \xi^{-i} \Omega_{\check{X}}^{\otimes i})$ for every $1 \leq i \leq r$. We take local sections m of M , $d \log(x)$ of $\Omega_{\check{X}}^1$ and its dual ∂ of $T_{\check{X}}$. And we write

$$\theta(m) = \theta_\partial(m) \otimes \xi^{-1} d \log(x).$$

Locally, let \mathcal{N} be the $\mathcal{O}_{\check{X}}$ -submodule of M generated by $\{m, \theta_\partial(m), \dots, (\theta_\partial)^{r-1}(m)\}$. In view of the characteristic polynomial of θ and the assumption, for $n \geq r$, the section $(\theta_\partial)^n(m)$ belongs to $p^{\alpha(n-r)} \mathcal{N}$. We take

a positive integer s such that $\alpha > \frac{1}{s} + \frac{1}{p-1}$. Then we deduce that $p^{-s^{-1}n}(\theta_\partial)^n(m)/n!$ tends to zero, when n tends to infinity, i.e. (M, θ) satisfies $(\text{Conv})_s$ (2.2.5). By ([4] IV.3.6.4), the Higgs field θ is small. \square

2.4.3. Let (M, θ) be a small Higgs bundle over $X_{\mathbf{C}}$ and $\alpha \in \mathbb{Q}_{>\frac{1}{p-1}}$ a rational number as in proposition 2.4.2. By *loc.cit.*, we can define an ideal sheaf \mathcal{J}_θ of $\text{Sym}_{\mathcal{O}_{\tilde{X}}}^\bullet(p^{-\alpha}\xi T_{\tilde{X}})$ by a similar formula of (2.4.1.1) with coefficients $\{c_i(\theta)\}_{i=1}^r$. We denote by $\hat{\mathcal{A}}_\theta$ the finite, locally free $\mathcal{O}_{\tilde{X}}$ -algebra

$$\hat{\mathcal{A}}_\theta = \text{Sym}_{\mathcal{O}_{\tilde{X}}}^\bullet(p^{-\alpha}\xi T_{\tilde{X}})/\mathcal{J}_\theta.$$

As $\text{Spf}_{\mathcal{O}_{\tilde{X}}}(\hat{\mathcal{A}}_\theta)$ is projective over \check{S} , we have a canonical isomorphism $(\text{Spf}_{\mathcal{O}_{\tilde{X}}}(\hat{\mathcal{A}}_\theta))^{\text{rig}} \simeq (\text{Spec}_{\mathcal{O}_{X_{\mathbf{C}}}}(\mathcal{A}_\theta))^{\text{an}}$.

Let K' be a finite extension of K , Y a projective $S' (= \text{Spec}(\mathcal{O}_{K'}))$ -curve with semi-stable reduction, $f : Y \rightarrow X_{S'}$ a generic η' -cover (1.2.2) and $\check{f} : \check{Y} \rightarrow \check{X}$ be the associated morphism of p -adic fine log formal schemes. Let \mathcal{Y} be a smooth lifting of \check{Y} over $\Sigma_{S'}$. The obstruction of lifting \check{f} to $\Sigma_{2,S}$ defines an element o_f of $H^1(\check{Y}, \check{f}^*(\xi T_{\check{X}}))$. We consider the following exponential map:

$$(2.4.3.1) \quad H^1(\check{Y}, \check{f}^*(\xi T_{\check{X}})) \rightarrow p^\alpha H^1(\check{Y}, \check{f}^*\hat{\mathcal{A}}_\theta) \xrightarrow{\exp} H^1(\check{Y}, (\check{f}^*\hat{\mathcal{A}}_\theta)^\times). \\ \varphi = (\varphi_{ij}) \mapsto \exp(\varphi) = (\exp(\varphi_{ij})).$$

The second map is well-defined due to $\alpha > \frac{1}{p-1}$. By applying the above homomorphism to o_f , we obtain an invertible $\check{f}^*\hat{\mathcal{A}}_\theta$ -module $\mathcal{L}_{f,\theta}$.

For $(f^*(M), f^*(\theta)) \in \text{HB}(Y_{\mathbf{C}}, \xi^{-1}f^*(\Omega_{X_{\mathbf{C}}}^1))$, we have a canonical homomorphism:

$$(2.4.3.2) \quad H^1(\check{Y}, (\check{f}^*\hat{\mathcal{A}}_\theta)^\times) \rightarrow H^1(Y_{\mathbf{C}}^{\text{an}}, (\mathcal{A}_{f^*(\theta)})^{\text{an}, \times}) \simeq H^1(Y_{\mathbf{C}}, \mathcal{A}_{f^*(\theta)}^\times).$$

We denote abusively by $\mathcal{L}_{f,\theta}$ the invertible $(\mathcal{A}_{f^*(\theta)})^{\text{an}}$ -module (resp. $\mathcal{A}_{f^*(\theta)}$ -module) associated to $\mathcal{L}_{f,\theta}$.

Via $f_{\mathbf{C}}^*(\Omega_{X_{\mathbf{C}}}^1) \rightarrow \Omega_{Y_{\mathbf{C}}}^1$, we associate to (M, θ) a Higgs bundle $f_{\mathcal{Y},\mathcal{X}}^\circ(M, \theta) := f^*(M, \theta) \otimes_{\mathcal{A}_{f^*(\theta)}} \mathcal{L}_{f,\theta}$ over $Y_{\mathbf{C}}$, called *twisted inverse image* of (M, θ) . By proposition 2.4.2, $f_{\mathcal{Y},\mathcal{X}}^\circ(M, \theta)$ is a small Higgs bundle.

We compare $f_{\mathcal{Y},\mathcal{X}}^\circ$ with the twisted inverse image functor $\check{f}_{\mathcal{Y},\mathcal{X}}^*$ defined via Higgs crystals (2.3.4.1).

Proposition 2.4.4. *Via equivalences (2.2.10.1), there exists a canonical isomorphism of Higgs bundles*

$$f_{\mathcal{Y},\mathcal{X}}^\circ(M, \theta) \xrightarrow{\sim} \check{f}_{\mathcal{Y},\mathcal{X}}^*(M, \theta).$$

Proof. We first make an explicit description of $\mathcal{L}_{f,\theta}$. Let $\{U_i\}_{i \in I}$ be an affine open coverings of \check{Y} and $\mathcal{U}_i \rightarrow \mathcal{Y}$ the associated open formal subschemes in \mathcal{C} . We take a family of local liftings $\tilde{f}_i : \mathcal{U}_i \rightarrow \mathcal{X}$ of \check{f} .

The difference between \tilde{f}_i and \tilde{f}_j (modulo ξ^2) on the intersection $\mathcal{U}_{ij} = \mathcal{U}_i \cap \mathcal{U}_j$ defines a class

$$\varphi_{ij} \in \text{Hom}(\check{f}^*(\Omega_{\check{X}}^1)|_{\mathcal{U}_{ij}}, \xi \mathcal{O}_{\mathcal{U}_{ij,2}}).$$

The data $\varphi = (\varphi_{ij})$ defines a class in $H^1(\check{Y}, \xi^{-1}\check{f}^*(T_{\check{X}}))$. The invertible $\check{f}^*(\hat{\mathcal{A}}_\theta)$ -module $\mathcal{L}_{f,\theta}$ is equivalent to the family $\{\check{f}^*(\hat{\mathcal{A}}_\theta)|_{\mathcal{U}_i}\}_{i \in I}$ equipped with the gluing data $\{\exp(\varphi_{ij})\}_{i,j \in I}$.

Then the $\check{f}^*(\hat{\mathcal{A}}_\theta)$ -module $f_{\mathcal{Y},\mathcal{X}}^\circ(M, \theta)$ is equivalent to the family $\{\check{f}^*(M)|_{\mathcal{U}_i}\}_{i \in I}$ equipped with the gluing data $\{\Phi_{ij} = \text{id} \otimes \exp(\varphi_{ij})\}_{i,j \in I}$.

We may assume that \mathcal{X} is affine and there exists $x = (x_N) \in \varprojlim_N \Gamma(\mathcal{U}_{ij,N}, \mathcal{M}_{\mathcal{U}_{ij,N}})$ such that $d \log(x_N)$ is a basis of $\Omega_{\mathcal{U}_{ij,N}/\Sigma_{N,S}}^1$ for $N \geq 1$ as in 2.3.5. Then we have

$$\varphi_{ij}(d \log(x_1)) = \tilde{f}_i^*(x_2)\tilde{f}_j^*(x_2)^{-1} - 1.$$

As an isomorphism of $\mathcal{O}_{\mathcal{U}_{ij}}$ -modules, the transition morphism $\Phi_{ij} : \check{f}^*(M)|_{\mathcal{U}_{ij}} \xrightarrow{\sim} \check{f}^*(M)|_{\mathcal{U}_{ij}}$ is given by:

$$(2.4.4.1) \quad \check{f}^*(m) \mapsto \sum_{n \geq 0} \check{f}^*(\theta_{\partial^n}(m)) \otimes \frac{1}{n!} \left(\frac{\tilde{f}_i^*(x_2)\tilde{f}_j^*(x_2)^{-1} - 1}{\xi} \right)^n.$$

The above isomorphism coincides with the transition map of $\check{f}_{\mathcal{Y},\mathcal{X}}^*(M, \theta)$ (2.3.5.1) and finishes the proof. \square

2.4.5. Let us investigate the twisted inverse image functor for Higgs line bundles. Let α be a rational number $> \frac{1}{p-1}$ and $\theta \in p^\alpha \Gamma(\check{X}, \xi^{-1} \Omega_{\check{X}}^1)$. Then \mathcal{A}_θ is isomorphic to \mathcal{O}_{X_C} and an invertible \mathcal{A}_θ -module is equivalent to a line bundle together with the Higgs field θ . The tensor product of invertible \mathcal{A}_θ -modules is the same as the tensor product of underlying line bundles together with the Higgs field θ .

In view of formula (2.4.4.1) applied to $(\mathcal{O}_{\check{X}}, \theta)$, the special fiber of the line bundle $\mathcal{L}_{f, \theta}$ is trivial.

3. SMALL HIGGS BUNDLES WITH STRONGLY SEMI-STABLE REDUCTION OF DEGREE ZERO

In the following, we assume K is an algebraic extension of \mathbb{Q}_p with algebraically closed residue field k .

3.1. Review on Deninger–Werner’s construction. In this subsection, we briefly review some basic definitions and results in [10].

Definition 3.1.1. (i) Let C be a smooth proper curve over k and $F_C : C \rightarrow C$ the absolute Frobenius of C . We say a vector bundle E over C is *strongly semi-stable* if $(F_C^n)^*(E)$ is semi-stable for all $n \geq 0$.

(ii) Let C be a proper curve over k and $\pi : \tilde{C} \rightarrow C$ the normalization of the reduced subscheme of C . We say a vector bundle E over C is *strongly semi-stable of degree zero* if $\pi^*(E)$ is strongly semi-stable of degree zero on each irreducible connected component of \tilde{C} .

Definition 3.1.2. (i) Let X be an \overline{S} -model of a smooth proper curve $X_{\overline{\eta}}$ over \overline{K} (1.2.2) and \check{X} the p -adic completion of X . We denote by $\text{VB}^{\text{DW}}(\check{X})$ the full subcategory of vector bundles E over \check{X} whose special fiber E_s is a strongly semi-stable bundle of degree zero over X_s .

(ii) Let Y be a smooth proper \overline{K} -curve. We denote by $\text{VB}^{\text{DW}}(Y_{\mathbb{C}}^{\text{an}})$ the full subcategory of vector bundles E over $Y_{\mathbb{C}}^{\text{an}}$ such that there exists an \overline{S} -model X of Y and an object \mathcal{E} of $\text{VB}^{\text{DW}}(\check{X})$ with rigid generic fiber E . We also denote by $\text{VB}^{\text{DW}}(Y_{\mathbb{C}})$ the essential image of $\text{VB}^{\text{DW}}(Y_{\mathbb{C}}^{\text{an}})$ via equivalence $\text{VB}(Y_{\mathbb{C}}) \simeq \text{VB}(Y_{\mathbb{C}}^{\text{an}})$.

We denote by $\text{VB}^{\text{pDW}}(Y_{\mathbb{C}})$ the full subcategory of $\text{VB}(Y_{\mathbb{C}})$ consisting of vector bundles E , which admits a finite étale cover $g : Z \rightarrow Y$ over \overline{K} such that $g_{\mathbb{C}}^*(E)$ belongs to $\text{VB}^{\text{DW}}(Z_{\mathbb{C}})$.

Theorem 3.1.3 ([10] theorems 16, 17 and [32] théorém 2.4). *Let X be an \overline{S} -model of a smooth proper \overline{K} -curve. A vector bundle E over \check{X} belongs to $\text{VB}^{\text{DW}}(\check{X})$ if and only if, for each $n \geq 1$, there exists an $\overline{\eta}$ -cover (1.2.2) $\varphi : X' \rightarrow X$ such that $\varphi_n^*(E_n)$ is free of finite type, where φ_n (resp. E_n) denotes the reduction modulo p^n .*

Let \overline{x} be a geometric point of $X_{\overline{\eta}}$. Using parallel transport, Deninger and Werner constructed functors:

$$(3.1.3.1) \quad \mathbb{V}_X^{\text{DW}} : \text{VB}^{\text{DW}}(\check{X}) \rightarrow \mathbf{Rep}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathfrak{o}), \quad \mathbb{V}_{X_{\mathbb{C}}}^{\text{DW}} : \text{VB}^{\text{pDW}}(X_{\mathbb{C}}) \rightarrow \mathbf{Rep}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathbb{C}).$$

3.2. Covers trivializing small Higgs bundles with strongly semi-stable degree zero reduction.

3.2.1. Let X be a projective S -curve with semi-stable reduction and \mathcal{M}_X be the log structure on X defined by its special fiber X_s . We denote by \check{X} the p -adic completion of $X \times_S \overline{S}$, equipped with the fine log structure induced by \mathcal{M}_X . We fix a smooth Cartesian lifting \mathcal{X} of \check{X} over Σ_S , which always exists.

We denote by $\text{HB}_{\mathbb{Z}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$ the full subcategory of $\text{HB}_{\mathbb{Z}_p, \text{conv}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$ (2.2.6) consisting of Higgs modules (M, θ) such that M belongs to $\text{VB}^{\text{DW}}(\check{X})$ and that θ is small (2.2.10).

Our main result in this section is the following:

Theorem 3.2.2. *Let M be a vector bundle of rank r over \check{X} , θ a small Higgs field on M , and \mathcal{M} the associated Higgs crystal on $(\check{X}/\Sigma_S)_{\text{HIG}}^\infty$. The following conditions are equivalence:*

(i) (M, θ) belongs to $\text{HB}_{\mathbb{Z}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$.

(ii) *For every integer $n \geq 1$, there exists a finite extension K' of K , a projective $S' (= \text{Spec}(\mathcal{O}_{K'}))$ -curve Y with semi-stable reduction and an η' -cover $f : Y \rightarrow X_{S'}$ satisfying the following property: if $\check{f} : \check{Y} \rightarrow \check{X}$ denotes the p -adic completion of $f \times_{S'} \overline{S}$ and the associated morphism of p -adic fine log formal schemes, then there exists an isomorphism of reduction modulo p^n of Higgs crystals*

$$(3.2.2.1) \quad \check{f}_{\text{HIG}}^*(\mathcal{M})_n \simeq (\overline{\mathcal{O}}_{\check{Y}/\Sigma_{S',n}})^{\oplus r}.$$

The above theorem generalizes the description of $\text{VB}^{\text{DW}}(\check{X})$ in theorem 3.1.3.

We will use the following argument of Faltings on η -covers of X whose associated inverse image of log differentials is divisible by a power of p .

Proposition 3.2.3 (Faltings, [13] Proof of theorem 6). *Let n be an integer ≥ 1 . There exists a finite extension K' of K , a proper $S' (= \text{Spec}(\mathcal{O}_{K'}))$ -curve Y with semi-stable reduction, equipped with the log structure \mathcal{M}_Y defined by its special fiber and an (S, \mathcal{M}_S) -morphism $f : (Y, \mathcal{M}_Y) \rightarrow (X, \mathcal{M}_X)$ such that the image of the morphism of log differentials $df_{\log} : f^*(\tilde{\Omega}_{X/S}^1) \rightarrow \tilde{\Omega}_{Y/S'}^1$ is divisible by p^n .*

Proof. Let J_K be the Jacobian of X_K and J its Néron model, which is isomorphic to the connected component of the relative Picard $\text{Pic}_{X/S}^0$ ([8] 9.5 theorem 1). After taking an extension of K and choosing an S -point of X , we take a closed immersion $X_K \rightarrow J_K$ defined by this point. The multiplication by p^n on J_K induces an étale cover $f_K : X_{n,K} \rightarrow X_K$. After replacing K by a finite extension, there exist an S -model X_n with semi-stable reduction of $X_{n,K}$ and a model $f : X_n \rightarrow X$ of f_K .

If $J_{n,K}$ denotes the Jacobian of $X_{n,K}$ and J_n its Néron model, f_K induces a morphism $g_K : J_{n,K} \rightarrow J_K$ by the Albanese property and g_K factors through $p^n : J_K \rightarrow J_K$. Moreover, by the universal property of Néron models, g_K induces a morphism $g : J_n \rightarrow J$, which factors through p^n . Now we consider the following diagram of global sections of differentials:

$$\begin{array}{ccc}
 \Gamma(X, \tilde{\Omega}_{X/S}^1) & \xrightarrow{f^*} & \Gamma(X_n, \tilde{\Omega}_{X_n/S}^1) \\
 \downarrow & & \downarrow \\
 \Gamma(X_K, \Omega_{X_K/K}^1) & \xrightarrow{f_K^*} & \Gamma(X_{n,K}, \Omega_{X_{n,K}/K}^1) \\
 \downarrow \wr & & \downarrow \wr \\
 \Gamma(J_K, \Omega_{J_K/K}^1) & \xrightarrow{g_K^*} & \Gamma(J_{n,K}, \Omega_{J_{n,K}/K}^1) \\
 \downarrow & & \downarrow \\
 \Gamma(J, \Omega_{J/S}^1) & \xrightarrow{g^*} & \Gamma(J_n, \Omega_{J_n/S}^1)
 \end{array}$$

Note that $\tilde{\Omega}_{X/S}^1$ is isomorphic to the dualizing sheaf $\omega_{X/S}$. The isomorphisms in the above diagram are induced by $H^1(X, \mathcal{O}_X) \simeq \text{Lie } J$ and the coherent duality.

Since g factors through p^n , the image of g^* is divisible by p^n and so is f^* . Moreover, the sub- \mathcal{O}_{X_n} -module of $\tilde{\Omega}_{X_n/S}^1$ generated by global sections contains $p^{n_0} \tilde{\Omega}_{X_n/S}^1$ for some integer n_0 . Hence the image of $df : f^*(\tilde{\Omega}_{X/S}^1) \rightarrow \tilde{\Omega}_{X_n/S}^1$ is divisible by p^{n-n_0} . This finishes the proof. \square

Proof of theorem 3.2.2. (ii) \Rightarrow (i): Let \mathcal{Y} be a smooth Cartesian lifting of \check{Y} over $\Sigma_{S'}$. By corollary 2.3.6(ii), $\check{f}^*(M)_s$ is isomorphic to the special fiber of the underlying bundle of $\check{f}_{\mathcal{Y}, \mathcal{X}}^*(M, \theta)$, which is trivial by (3.2.2.1). Then we conclude that M_s is strongly semi-stable of degree zero by [10] theorem 18.

(i) \Rightarrow (ii): We first take an η' -cover $f : Y \rightarrow X_{S'}$ as in proposition 3.2.3 such that the image of df is divisible by p^{n+1} and a smooth Cartesian lifting \mathcal{Y} of \check{Y} over $\Sigma_{S'}$. By corollary 2.3.6(i), the Higgs field of $\check{f}_{\mathcal{Y}, \mathcal{X}}^*(M, \theta)$ is $(n+1)$ -small. In view of corollary 2.3.6(ii), the special fiber of its underlying vector bundle $(\mathcal{M}_{\mathcal{Y}})_s$ is isomorphic to $\check{f}_s^*(M_s)$. Since M_s is strongly semi-stable of degree zero, then so is $\check{f}_s^*(M_s)$ in view of [10] theorem 18.

By theorem 3.1.3 and the semi-stable reduction theorem, there exists a finite extension K'' of K' a projective $S'' (= \text{Spec}(\mathcal{O}_{K''}))$ -curve Z with semi-stable reduction and an η'' -cover $g : Z \rightarrow Y_{S''}$ satisfying following properties: If \check{g} denotes the p -adic completion of $g \times_{S''} \bar{S}$, then $\check{g}^*(\mathcal{M}_{\mathcal{Y}})_{n+1}$ is trivial.

In the following, we show that the functor $(\check{g} \circ \check{f})_{\text{HIG}}^*$ trivializes Higgs crystal \mathcal{M} modulo p^n .

We fix a smooth Cartesian lifting \mathcal{Z} of \check{Z} over $\Sigma_{S''}$. The Higgs fields of $\check{f}_{\mathcal{Y},\mathcal{X}}^*(M, \theta)$ and of $(\check{g} \circ \check{f})_{\mathcal{Z},\mathcal{X}}^*(M, \theta)$ are $(n+1)$ -small. By applying corollary 2.3.6(ii) to \check{g} , we obtain a canonical isomorphism of Higgs modules modulo p^n in $\mathrm{HM}(\check{Z}, \mathcal{Z}/\Sigma_{S''})$ (2.3.4.1):

$$(\mathcal{O}_{\check{Z}_n}^{\oplus r}, 0) \xrightarrow{\sim} \left((\check{g} \circ \check{f})_{\mathcal{Z},\mathcal{X}}^*(M, \theta) \right)_n.$$

Then we conclude the assertion from corollary 2.2.11. \square

4. REVIEW ON THE p -ADIC SIMPSON CORRESPONDENCE

4.1. Faltings topos.

4.1.1. Let $Y \rightarrow X$ be a morphism of coherent schemes (“coherent” means “quasi-compact and quasi-separated”). We denote by $E_{Y \rightarrow X}$ the category of morphisms of schemes $(V \rightarrow U)$ above $(Y \rightarrow X)$, i.e. commutative diagrams

$$\begin{array}{ccc} V & \longrightarrow & U \\ \downarrow & & \downarrow \\ Y & \longrightarrow & X \end{array}$$

such that $U \rightarrow X$ is étale and $V \rightarrow Y \times_X U$ is finite étale. We endow $E_{Y \rightarrow X}$ with the topology generated by the following types of families of morphisms

- (v) $\{(V_m \rightarrow U) \rightarrow (V \rightarrow U)\}_{m \in M}$, where $\{V_m \rightarrow V\}_{m \in M}$ is a finite étale covering,
- (c) $\{(V \times_U U_n \rightarrow U_n) \rightarrow (V \rightarrow U)\}_{n \in N}$, where $\{U_n \rightarrow U\}_{n \in N}$ is an étale covering.

We denote by $\tilde{E}_{Y \rightarrow X}$ the topos of sheaves of sets on $E_{Y \rightarrow X}$ and call it *Faltings topos* associated to Y .

The presheaf $\overline{\mathcal{B}}_{Y \rightarrow X}$ on $E_{Y \rightarrow X}$ defined by

$$\overline{\mathcal{B}}_{Y \rightarrow X}(V \rightarrow U) = \Gamma(U^V, \mathcal{O}_{U^V}),$$

where U^V is the integral closure of U in V ([38] 0BAK), is a sheaf of rings ([17] proposition 7.6). For $n \geq 1$, we set $\overline{\mathcal{B}}_{Y \rightarrow X, n} = \overline{\mathcal{B}}_{Y \rightarrow X} / p^n \overline{\mathcal{B}}_{Y \rightarrow X}$.

We denote by $\tilde{\overline{\mathcal{B}}}_{Y \rightarrow X}$ the sheaf of rings of $\tilde{E}_{Y \rightarrow X}^{\mathbb{N}^0}$ (§ 1.2.4), defined by the projective system $(\overline{\mathcal{B}}_{Y \rightarrow X, n})_{n \geq 1}$.

Proposition 4.1.2 ([4] VI.5.10). *A sheaf F on $E_{Y \rightarrow X}$ is equivalent to the datum of a sheaf F_U of $U_{Y, \text{fét}}$, with $U_Y = U \times_X Y$ for every object U of $\hat{\mathbf{E}t}_X$ and a morphism $F_U \rightarrow (f_Y)_{\text{fét},*}(F_{U'})$ for every morphism $f : U' \rightarrow U$ of $\hat{\mathbf{E}t}_X$ satisfying the following compatibility condition: for every covering $(f_i : U_i \rightarrow U)_{i \in I}$ of $\hat{\mathbf{E}t}_X$, if we set $U_{ij} = U_i \times_U U_j$ and $f_{ij} : U_{ij} \rightarrow U$ the canonical morphism, then the following sequence of morphisms of sheaves of $U_{Y, \text{fét}}$ is exact:*

$$F_U \rightarrow \prod_{i \in I} (f_i)_{\text{fét},*}(F_{U_i}) \rightrightarrows \prod_{i, j \in I} (f_{ij, \bar{\eta}})_{\text{fét},*}(F_{U_{ij}}).$$

Corollary 4.1.3. *Let $Y = \bigsqcup_{i \in I} Y_i$ be the decomposition of Y into finitely many connected components. Then we have a canonical isomorphism of topoi:*

$$\tilde{E}_{Y \rightarrow X} \simeq \bigsqcup_{i \in I} \tilde{E}_{Y_i \rightarrow X}.$$

Proof. For each scheme U étale over X , we have a functorial decomposition $U_Y \simeq \bigsqcup_{i \in I} U_{Y_i}$ and the associated topoi $U_{Y, \text{fét}} \simeq \bigsqcup_{i \in I} U_{Y_i, \text{fét}}$. Then we conclude the corollary from proposition 4.1.2. \square

4.1.4. We briefly review the cohomological descent in Faltings topoi. We assume that X is a coherent \overline{S} -scheme, Y is a coherent $\overline{\eta}$ -scheme and moreover that in the following diagram:

$$\begin{array}{ccccc} Y & \longrightarrow & X^Y & \longrightarrow & X \\ \downarrow & & \square & & \downarrow \\ \overline{\eta} & \longrightarrow & \overline{S} & & \end{array}$$

where X^Y denotes the integral closure of X in Y ([38] 0BAK), the square is Cartesian.

Let $X_\bullet \rightarrow X$ be a proper hypercovering of \overline{S} -schemes, and $\varepsilon : \widetilde{E}_{Y_\bullet \rightarrow X_\bullet} \rightarrow \widetilde{E}_{Y \rightarrow X}$ the augmentation of simplicial topoi with $Y_\bullet = X_\bullet \times_X Y$. For any $\overline{\mathcal{B}}_{Y \rightarrow X}$ -module \mathcal{N} , we have a canonical morphism:

$$(4.1.4.1) \quad \mathcal{N} \rightarrow R\varepsilon_*(\varepsilon^* \mathcal{N}).$$

We say \mathcal{N} satisfies the *cohomological descent (in Faltings topoi)*, if for any proper hypercovering $X_\bullet \rightarrow X$ of \overline{S} -schemes, the above morphism is an almost isomorphism.

Recently, T. He [17] showed that the cohomological descent holds for $\mathcal{N} = \mathbb{L} \otimes_{\mathbb{Z}} \overline{\mathcal{B}}_{Y \rightarrow X}$, with a finite locally constant abelian sheaf \mathbb{L} of $\widetilde{E}_{Y \rightarrow X}$. A key ingredient of his proof is the arc-descent for perfectoid algebras due to Bhatt–Scholze [7]. In particular, we obtain the following corollary:

Corollary 4.1.5. *Assume $Y = X_{\overline{\eta}}$. For any integer $n \geq 1$, a locally free $\overline{\mathcal{B}}_{Y \rightarrow X, n}$ -module M of finite type satisfies the cohomological descent.*

Proof. Let $\{(V_i \rightarrow U_i) \rightarrow (X_{\overline{\eta}} \rightarrow X)\}_{i \in I}$ be a covering such that $M|_{(V_i \rightarrow U_i)}$ is free of finite type. If $(V \rightarrow U) \rightarrow (X_{\overline{\eta}} \rightarrow X)$ is an element of this covering, then $U \rightarrow X$ is étale and $V \rightarrow U_{\overline{\eta}}$ is finite étale. We may assume that V, U are coherent. By ([38] 035K), the generic fiber of U^V is isomorphic to V . The cohomological descent ([17] corollaries 8.14, 8.18) holds for the restriction of M to the localization $E_{Y \rightarrow X}/(V \rightarrow U) \simeq E_{V \rightarrow U}$. As the assertion is local on $E_{Y \rightarrow X}$, the corollary follows. \square

4.1.6. In the following, let X be a flat, normal, separated S -scheme (resp. \overline{S} -scheme) of finite presentation such that $\underline{X} = X \times_S \overline{S}$ (§1.2.1) (resp. X) is normal and $Y = X_{\overline{\eta}}$. We will use both of two settings in this article. Then \underline{X} (resp. X) is locally irreducible (cf. [4] III.3.1). Indeed, as \underline{X} (resp. X) is flat over \overline{S} , its generic points are the generic points of $X_{\overline{\eta}}$, which are finite (cf. [4] III.3.2(ii)).

For simplicity, we denote the Faltings site $E_{X_{\overline{\eta}} \rightarrow X}$ (resp. the structure sheaf $\overline{\mathcal{B}}_{X_{\overline{\eta}} \rightarrow X}$) by E_X or E (resp. $\overline{\mathcal{B}}_X$ or $\overline{\mathcal{B}}$), if there is no confusion. The Faltings topoi \widetilde{E}_X admits a closed sub-topoi $\widetilde{E}_{X, s}$ ([4] IV 9.3.5). For each $n \geq 1$, the ring $\overline{\mathcal{B}}_{X, n}$ belongs to $\widetilde{E}_{X, s}$.

The functor $\mathbf{F}\acute{\mathbf{e}}\mathbf{t}_{X_{\overline{\eta}}} \rightarrow E$, defined by $V \mapsto (V \rightarrow Y)$, is continuous and left exact and induces a morphism of topoi $\beta : \widetilde{E}_X \rightarrow X_{\overline{\eta}, \text{fét}}$. Moreover, it induces morphisms of ringed topoi ([36] 7.13)

$$(4.1.6.1) \quad \beta_{X, n} : (\widetilde{E}_{X, s}, \overline{\mathcal{B}}_n) \rightarrow (X_{\overline{\eta}, \text{fét}}, \mathfrak{o}_n), \quad \forall n \geq 1, \quad \check{\beta}_X : (\widetilde{E}_X^{\text{No}}, \check{\overline{\mathcal{B}}}_X) \rightarrow (X_{\overline{\eta}, \text{fét}}^{\text{No}}, \check{\mathfrak{o}}),$$

where \mathfrak{o}_n is the constant sheaf of $X_{\overline{\eta}, \text{fét}}$ with value \mathfrak{o}_n , and $\check{\mathfrak{o}} = (\mathfrak{o}_n)_{n \geq 1}$.

The functor $\mathbf{E}\mathbf{t}_X \rightarrow E$, defined by $U \mapsto (U_{\overline{\eta}} \rightarrow U)$, is continuous and left exact and induces a morphism of topoi $\sigma : \widetilde{E}_X \rightarrow X_{\text{ét}}$. Moreover, it induces morphisms of ringed topoi ([4] III.9.9, III.9.10)

$$(4.1.6.2) \quad \sigma_{X, n} : (\widetilde{E}_{X, s}, \overline{\mathcal{B}}_{X, n}) \rightarrow (X_{s, \text{ét}}, \mathcal{O}_{\underline{X}_n}), \quad \forall n \geq 1, \quad \check{\sigma}_X : (\widetilde{E}_X^{\text{No}}, \check{\overline{\mathcal{B}}}_X) \rightarrow (X_{s, \text{ét}}^{\text{No}}, \check{\mathcal{O}}_{\underline{X}}),$$

where $\mathcal{O}_{\underline{X}_n}$ is the étale sheaf of $X_{s, \text{ét}}$ associated to the coherent sheaf $\mathcal{O}_{\underline{X}_n}$ of $X_{s, \text{zar}}$ and $\check{\mathcal{O}}_{\underline{X}} = (\mathcal{O}_{\underline{X}_n})_{n \geq 1}$.

When X is an S -scheme, the canonical functor $E_X \rightarrow E_{\underline{X}}$, defined by $(V \rightarrow U) \mapsto (V \rightarrow \underline{U})$, induces a morphism of topoi

$$(4.1.6.3) \quad h : \widetilde{E}_{\underline{X}} \rightarrow \widetilde{E}_X.$$

We have $h_*(\overline{\mathcal{B}}_{\underline{X}}) = \overline{\mathcal{B}}_X$ and the above morphism is ringed by $\overline{\mathcal{B}}_{\underline{X}}$ and $\overline{\mathcal{B}}_X$.

Lemma 4.1.7. (i) For any \mathfrak{o}_n -module \mathbb{L} of $X_{\overline{\eta}, \text{fét}}$, the following canonical morphism is an isomorphism:

$$\beta_{X,n}^*(\mathbb{L}) \xrightarrow{\sim} h_*(\beta_{\underline{X},n}^*(\mathbb{L})).$$

(ii) For any $\mathcal{O}_{\underline{X}_n}$ -module M of $X_{s,\text{ét}}$, the following canonical morphism is an isomorphism:

$$\sigma_{X,n}^*(M) \xrightarrow{\sim} h_*(\sigma_{\underline{X},n}^*(M)).$$

Proof. (i) By ([4] VI.8.9 and VI.10.9), both two sheaves are isomorphic to the sheaf associated to the presheaf:

$$\{U \mapsto f_{\overline{\eta}, \text{fét}}^*(\mathbb{L}) \otimes_{\mathfrak{o}_n} (\overline{\mathcal{B}}_{X,n})|_U, \quad f: U \rightarrow X \in \hat{\mathbf{E}}\mathbf{t}_X\},$$

where \mathfrak{o}_n is the constant sheaf of $U_{\overline{\eta}, \text{fét}}$. Then the assertion follows.

Assertion (ii) can be proved in a similar way using ([4] VI.5.34(ii)). \square

4.1.8. In the following, we assume X is an S -scheme of finite type with semi-stable reduction, equipped with log structure \mathcal{M}_X defined by the special fiber. Then \underline{X} is normal.

Let $U = \text{Spec}(R) \rightarrow X$ be an object of \mathbf{Q} (2.2.2) and $\overline{y} \rightarrow U_{\overline{\eta}}$ a geometric point such that the image y of \overline{y} in U_{η} is a generic point and that $\overline{\mathcal{K}} := \Gamma(\overline{y}, \mathcal{O}_{\overline{y}})$ is an algebraic closure of the residue field \mathcal{K} of U_{η} at y . We define \mathcal{K}^{ur} to be the union of all finite extensions of \mathcal{K} contained in $\overline{\mathcal{K}}$ such that the integral closures of R in \mathcal{L} are étale over $R[\frac{1}{p}]$. Let \overline{R} be the integral closure of R in \mathcal{K}^{ur} and $\widehat{\overline{R}}$ the p -adic completion of \overline{R} .

Let $U_{\overline{\eta}}^{\overline{y}}$ be the connected component of $U_{\overline{\eta}}$ containing \overline{y} . The morphism $\overline{y} \rightarrow U_{\overline{\eta}}$ induces an extension $\overline{K} \rightarrow \overline{\mathcal{K}}$ above $K \rightarrow \mathcal{K}$. We denote by $G_{(U,\overline{y})}$ the Galois group $\text{Gal}(\mathcal{K}^{\text{ur}}/\mathcal{K})$, and by $\Delta_{(U,\overline{y})}$ the kernel of $G_{(U,\overline{y})} \rightarrow \text{Gal}(\overline{K}/K)$, which is canonically isomorphic to $\pi_1(U_{\overline{\eta}}^{\overline{y}}, \overline{y})$. Let $\mathbf{B}_{\Delta_{(U,\overline{y})}}$ be the classifying topos of the profinite group $\Delta_{(U,\overline{y})}$. Under the equivalence: $\nu_{\overline{y}} : (U_{\overline{\eta}}^{\overline{y}})_{\text{fét}} \xrightarrow{\sim} \mathbf{B}_{\Delta_{(U,\overline{y})}}$ defined by the fiber functor $F \mapsto F_{\overline{y}}$, we have a canonical isomorphism ([4] III.8.15)

$$\nu_{\overline{y}}(\overline{\mathcal{B}}_U|(U_{\overline{\eta}}^{\overline{y}})_{\text{fét}}) \xrightarrow{\sim} \widehat{\overline{R}}.$$

Moreover, the action $\Delta_{(U,\overline{y})}$ on $\widehat{\overline{R}}$ is continuous with respect to the p -adic topology.

For each pair (U, \overline{y}) as above, we associate to adic $\overline{\mathcal{B}}$ -modules of finite type (1.2.4) continuous $\widehat{\overline{R}}$ -representations of $\Delta_{(U,\overline{y})}$ via $\nu_{\overline{y}}$. In the following, we collect some notions about representations:

Definition 4.1.9 ([4] II.13.1, II.13.2). Let G be a topological group, A an \mathfrak{o} -algebra that is complete and separated for the p -adic topology, equipped with a continuous G -action (via homomorphisms of \mathfrak{o} -algebras).

(i) A continuous A -representation of G is an A -module M of finite type, equipped with the p -adic topology and a continuous A -semi-linear action of G .

(ii) Let M be a continuous A -representation of G and α a rational number > 0 .

We say M is α -small if M is a free A -module of finite rank having a basis over A consisting of elements which are G -invariant modulo p^{α} .

We say M is small if M is α -small for a rational number $\alpha > \frac{2}{p-1}$.

(iii) A continuous $A[\frac{1}{p}]$ -representation of G is an $A[\frac{1}{p}]$ -module N of finite type, equipped with the p -adic topology and a continuous $A[\frac{1}{p}]$ -semi-linear action of G .

(iv) We say a continuous $A[\frac{1}{p}]$ -representation N of G is small, if N is moreover projective, and there exists a rational number $\alpha > \frac{2}{p-1}$, a sub- A -module of finite type N° which is G -invariant and generates N over $A[\frac{1}{p}]$, and a finite number of generators of N° that are G -invariant modulo p^{α} .

We denote by $\mathbf{Rep}(G, A)$ (resp. $\mathbf{Rep}(G, A[\frac{1}{p}])$) the category of continuous A -representations (resp. $A[\frac{1}{p}]$ -representations) of G and by $\mathbf{Rep}_{\text{small}}(G, A)$ (resp. $\mathbf{Rep}_{\text{small}}(G, A[\frac{1}{p}])$) the full subcategory of $\mathbf{Rep}(G, A)$ (resp. $\mathbf{Rep}(G, A[\frac{1}{p}])$) consisting of small representations.

4.1.10. We denote by $\mathbf{LocSys}(X_{\overline{\eta}, \text{fét}}, \mathfrak{o}_n)$ (resp. $\mathbf{LocSys}(X_{\overline{\eta}, \text{fét}}, \check{\mathfrak{o}})$) the category of locally free of finite type \mathfrak{o}_n -modules on $X_{\text{fét}}$ (resp. adic $\check{\mathfrak{o}}$ -modules $M = (M_n)_{n \geq 1}$ of finite type such that each M_n belongs to $\mathbf{LocSys}(X_{\overline{\eta}, \text{fét}}^{\mathbb{N}^\circ}, \mathfrak{o}_n)$). Let \overline{x} be a geometric point of $X_{\overline{\eta}}$. The fiber functor $\nu_{\overline{x}}$ at \overline{x} induces fully faithful functors and an equivalence of categories (c.f. [36] 3.29):

$$(4.1.10.1) \quad \mathbf{LocSys}(X_{\overline{\eta}, \text{fét}}, \mathfrak{o}_n) \rightarrow \mathbf{Rep}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathfrak{o}_n), \quad \mathbf{LocSys}(X_{\overline{\eta}, \text{fét}}^{\mathbb{N}^\circ}, \check{\mathfrak{o}}) \rightarrow \mathbf{Rep}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathfrak{o}),$$

$$(4.1.10.2) \quad \mathbf{LocSys}(X_{\overline{\eta}, \text{fét}}^{\mathbb{N}^\circ}, \check{\mathfrak{o}})_{\mathbb{Q}} \xrightarrow{\sim} \mathbf{Rep}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathbb{C}).$$

The essential images of (4.1.10.1) are representations whose underlying \mathfrak{o}_n -modules, \mathfrak{o} -modules are free.

Let $\rho : X_{\overline{\eta}, \text{ét}} \rightarrow X_{\overline{\eta}, \text{fét}}$ be the canonical morphism of topoi. The adjunction morphism $\text{id} \rightarrow \rho_* \rho^*$ is an isomorphism. The inverse image of ρ induces a fully faithful functor:

$$(4.1.10.3) \quad \rho^* : \mathbf{Mod}(X_{\overline{\eta}, \text{fét}}, \mathfrak{o}_n) \rightarrow \mathbf{Mod}(X_{\overline{\eta}, \text{ét}}, \mathfrak{o}_n), \quad \forall n \geq 1.$$

4.2. p -adic Simpson correspondence via Higgs crystals.

4.2.1. We keep the assumption and notation of 4.1.8. We denote by \overline{R}^\flat the perfection $\varprojlim_{x \mapsto x^p} \overline{R}/p\overline{R}$ of $\overline{R}/p\overline{R}$ and by $\theta : W(\overline{R}^\flat) \rightarrow \widehat{\overline{R}}$ the canonical homomorphism defined by Fontaine, which is surjective ([4] II.9.5). The element ξ (§1.2.1) generates the ideal $\text{Ker}(\theta)$.

For $N \geq 1$, we set $A_N(\overline{R}) = W(\overline{R}^\flat)/\xi^N W(\overline{R}^\flat)$, which is also p -adically complete and separated. There exists a canonical and continuous action of $G_{(U, \overline{y})}$ on $A_N(\overline{R})$ ([4] IV.5.1.2).

We denote by \overline{U} the p -adic formal scheme $\overline{U} = \text{Spf}(\widehat{\overline{R}})$ and for $N \geq 1$, $D_N(\overline{U})$ the p -adic formal scheme $\text{Spf}(A_N(\overline{R}))$. They are equipped with fine and saturated log structures induced by \mathcal{M}_U and there is a canonical isomorphism $\overline{U} \xrightarrow{\sim} D_1(\overline{U})$ (cf. [4] IV.5.1.4).

The inductive system $D(\overline{U}) = (D_N(\overline{U}))_{N \geq 1}$ together with a canonical morphism $z_{\overline{U}} : \overline{U} \rightarrow \check{U}$ form an object of $(\check{X}/\Sigma_S)_{\text{HIG}}^r$ for $r \in \mathbb{N} \cup \{\infty\}$ (cf. [4] IV.5.2). Then $\Delta_{(U, \overline{y})}$ acts on $(D(\overline{U}), z_{\overline{U}})$.

4.2.2. We denote by $\text{LPM}(\widehat{\overline{R}})$ the category of finitely generated $\widehat{\overline{R}}$ -modules M such that $M[\frac{1}{p}]$ is projective over $\widehat{\overline{R}}[\frac{1}{p}]$, and by $\mathbf{Rep}^{\text{LPM}}(\Delta_{(U, \overline{y})}, \widehat{\overline{R}})$ (resp. $\mathbf{Rep}^{\text{PM}}(\Delta_{(U, \overline{y})}, \widehat{\overline{R}}[\frac{1}{p}])$) the full subcategory of $\mathbf{Rep}(\Delta_{(U, \overline{y})}, \widehat{\overline{R}})$ (resp. $\mathbf{Rep}(\Delta_{(U, \overline{y})}, \widehat{\overline{R}}[\frac{1}{p}])$) whose underlying module belongs to $\text{LPM}(\widehat{\overline{R}})$ (resp. is finitely generated and projective over $\widehat{\overline{R}}[\frac{1}{p}]$).

For an object U of \mathbf{Q} , a geometric point \overline{y} of $U_{\overline{\eta}}$ as in 4.1.8 and an object \mathcal{M} of $\text{HC}_{\mathbb{Z}_p, \text{fin}}(\check{X}/\Sigma_S)$ (resp. $\text{HC}_{\mathbb{Q}_p, \text{fin}}^r(\check{X}/\Sigma_S)$), we associate an object of $\text{LPM}(\widehat{\overline{R}})$ (resp. a projective $\widehat{\overline{R}}[\frac{1}{p}]$ -module):

$$T_{(U, \overline{y})}(\mathcal{M}) = \Gamma((D(\overline{U}), z_{\overline{U}}), \mathcal{M}), \quad (\text{resp. } V_{(U, \overline{y})}^r(\mathcal{M}) = \Gamma((D(\overline{U}), z_{\overline{U}}), \mathcal{M})).$$

The action of $\Delta_{(U, \overline{y})}$ on $(D(\overline{U}), z_{\overline{U}})$ provides a semi-linear action of $\Delta_{(U, \overline{y})}$ on $T_{(U, \overline{y})}(\mathcal{M})$. Then we obtain the following functors ([4] IV.5.2):

$$(4.2.2.1) \quad T_{(U, \overline{y})} : \text{HC}_{\mathbb{Z}_p, \text{fin}}(\check{X}/\Sigma_S) \rightarrow \mathbf{Rep}^{\text{LPM}}(\Delta_{(U, \overline{y})}, \widehat{\overline{R}}), \quad \mathcal{M} \mapsto \Gamma((D(\overline{U}), z_{\overline{U}}), \mathcal{M}),$$

$$(4.2.2.2) \quad V_{(U, \overline{y})}^r : \text{HC}_{\mathbb{Q}_p, \text{fin}}^r(\check{X}/\Sigma_S) \rightarrow \mathbf{Rep}^{\text{PM}}(\Delta_{(U, \overline{y})}, \widehat{\overline{R}}[\frac{1}{p}]), \quad \mathcal{M} \mapsto \Gamma((D(\overline{U}), z_{\overline{U}}), \mathcal{M}).$$

Let $f : U' \rightarrow U$ be a morphism of \mathbf{Q} and \overline{y}' a geometric generic point of $U'_{\overline{\eta}}$ above \overline{y} . It induces a homomorphism $\Delta_{(U', \overline{y}')} \rightarrow \Delta_{(U, \overline{y})}$ and a morphism $D(\overline{U}') \rightarrow D(\overline{U})$ of $(\check{X}/\Sigma_S)_{\text{HIG}}^r$ compatible with $\Delta_{(U, \overline{y})}$ -actions. Moreover, we deduce a $\widehat{\overline{R}}$ -linear and $\Delta_{(U, \overline{y})}$ -equivariant morphism:

$$T_{(U, \overline{y})}(\mathcal{M}) \rightarrow T_{(U', \overline{y}')}(\mathcal{M}) \quad (\text{resp. } V_{(U, \overline{y})}^r(\mathcal{M}) \rightarrow V_{(U', \overline{y}')}^r(\mathcal{M})).$$

Since \mathcal{M} is a crystal, we deduce the following \widehat{R}' -linear $\Delta_{(U', \overline{y}')}$ -equivariant isomorphism:

$$(4.2.2.3) \quad T_{(U, \overline{y})}(\mathcal{M}) \otimes_{\widehat{R}} \widehat{R}' \xrightarrow{\sim} T_{(U', \overline{y}')}(\mathcal{M}) \quad (\text{resp. } V_{(U, \overline{y})}^r(\mathcal{M}) \otimes_{\widehat{R}} \widehat{R}' \xrightarrow{\sim} V_{(U', \overline{y}')}^r(\mathcal{M})).$$

4.2.3. The above construction globalizes to a functor ([4] IV.6.4.5):

$$(4.2.3.1) \quad T : \mathrm{HC}_{\mathbb{Z}_p, \mathrm{fin}}(\check{X}/\Sigma_S) \rightarrow \mathbf{Mod}(\check{\mathcal{B}}), \quad \mathcal{M} \mapsto (T_{(U, \overline{y})}(\mathcal{M})_n)_{n \geq 1, U \in \mathbf{Q}, \overline{y} \in \mathrm{Con}(U_{\overline{y}})}.$$

It is fully faithful up to isogeny ([4] theorem IV.6.4.9).

We denote by $\mathrm{HC}_{\mathbb{Z}_p, \mathrm{fin}}^r(\check{X}/\Sigma_S)$ the category whose object is a triple consisting of an object \mathcal{M} of $\mathrm{HC}_{\mathbb{Z}_p, \mathrm{fin}}^r(\check{X}/\Sigma_S)$, an object \mathcal{M}° of $\mathrm{HC}_{\mathbb{Z}_p, \mathrm{fin}}(\check{X}/\Sigma_S)$ and an isomorphism $\iota_{\mathcal{M}} : \mathcal{M}_{\mathbb{Q}}^\circ \simeq \mathcal{M}|_{(\check{X}/\Sigma_S)_{\mathrm{HIG}}^\infty}$. A morphism is defined in a natural way (c.f. [4] definition IV.3.5.4).

The functor T induces a fully faithful functor:

$$(4.2.3.2) \quad V^r : \mathrm{HC}_{\mathbb{Z}_p, \mathrm{fin}}^r(\check{X}/\Sigma_S)_{\mathbb{Q}} \rightarrow \mathbf{Mod}(\check{\mathcal{B}})_{\mathbb{Q}}, \quad (\mathcal{M}, \mathcal{M}^\circ) \mapsto T(\mathcal{M}^\circ)_{\mathbb{Q}},$$

which is compatible with the functor $V_{(U, \overline{y})}^r$.

Proposition 4.2.4 ([4] IV.6.4.6). *The functors T, V^r are compatible with the inverse image functor of Higgs crystals and the inverse image functor of Faltings ringed topoi.*

4.2.5. The functor $T_{(U, \overline{y})}$ (resp. $V_{(U, \overline{y})}^r$) can be reinterpreted as an admissible isomorphism for a “period ring”. Following ([4] IV.5.2), we denote by $\mathcal{D}_{\check{X}, \mathcal{X}}^r(\overline{U})$ the object of \mathcal{C}^r defined by the Higgs envelope (§ 2.1.8)

$$\mathcal{D}_{\check{X}, \mathcal{X}}^r(\overline{U}) = D_{\mathrm{HIG}}^r(\overline{U} \rightarrow \mathcal{X} \times_{\Sigma_S} D(\overline{U})),$$

which is affine. We define the object $\mathcal{A}_{\check{X}, \mathcal{X}}^r(\overline{R}) = (\mathcal{A}_{\check{X}, \mathcal{X}, N}^r(\overline{R}))$ of \mathcal{A}_\bullet^r (§ 2.1.1) by

$$\mathcal{A}_{\check{X}, \mathcal{X}, N}^r(\overline{R}) = \Gamma(\mathcal{D}_{\check{X}, \mathcal{X}, N}^r(\overline{U}), \mathcal{O}_{\mathcal{D}_{\check{X}, \mathcal{X}, N}^r(\overline{U})}^r).$$

We ignore the notation \check{X}, \mathcal{X} from $\mathcal{A}_{\check{X}, \mathcal{X}, N}^r$ if there is no confusion and we focus on \mathcal{A}_1^r .

Let $\underline{U}^{\overline{y}} = \mathrm{Spec}(R_1)$ be the connected component (or equivalently, irreducible component) of $\underline{U} = U \times_S \overline{S}$ containing \overline{y} and $\check{U}^{\overline{y}} = \mathrm{Spf}(\widehat{R}_1)$ its p -adic completion. The projections $\mathcal{D}^r(\overline{U}) \rightarrow D(\overline{U})$ and $\mathcal{D}^r \rightarrow \mathcal{X}$ induce compatible homomorphisms $\widehat{R} \rightarrow \mathcal{A}_1^r(\overline{R})$ and $\widehat{R}_1 \rightarrow \mathcal{A}_1^r(\overline{R})$. We refer to ([4] IV.5.2.5) for a description of \widehat{R} -algebra structure of $\mathcal{A}_1^r(\overline{R})$. The ring $\mathcal{A}_1^r(\overline{R})$ is equipped with a continuous action of $\Delta_{(U, \overline{y})}$ with respect to the p -adic topology ([4] IV.5.2.6) and a Higgs field

$$\theta_{\mathcal{A}} : \mathcal{A}_1^r(\overline{R}) \rightarrow \xi^{-1} \mathcal{A}_1^r(\overline{R}) \otimes_R \widetilde{\Omega}_{R/\mathcal{O}_K}^1.$$

Let \mathcal{M} be a Higgs isocrystal of $\mathrm{HC}_{\mathbb{Z}_p, \mathrm{fin}}^r(\check{X}/\Sigma_S)$ (resp. a Higgs crystal of $\mathrm{HC}_{\mathbb{Z}_p, \mathrm{fin}}(\check{X}/\Sigma_S)$). We set $M = \Gamma(\check{U}^{\overline{y}}, \mathcal{M}_{\mathcal{X}})$ and $\theta : M \rightarrow \xi^{-1} M \otimes_R \widetilde{\Omega}_{R/\mathcal{O}_K}^1$ the associated Higgs field on M . We set $V(\mathcal{M})$ to be $V_{(U, \overline{y})}^r(\mathcal{M})$ (resp. $T_{(U, \overline{y})}(\mathcal{M})$). Then there exists a canonical $\Delta_{(U, \overline{y})}$ -equivariant $\mathcal{A}_1^r(\overline{R})$ -linear isomorphism compatible with Higgs fields ([4] IV.5.2.12):

$$(4.2.5.1) \quad \mathcal{A}_1^r(\overline{R}) \otimes_{\widehat{R}} V(\mathcal{M}) \xrightarrow{\sim} \mathcal{A}_1^r(\overline{R}) \otimes_{\widehat{R}_1} M,$$

where the Higgs field on $V(\mathcal{M})$ is trivial and $\Delta_{(U, \overline{y})}$ -action on M is also trivial.

Moreover, Tsuji compares the algebra $\mathcal{A}_1^r(\overline{R})$ with the Higgs-Tate algebra introduced by Abbes–Gros ([4] II.10). More precisely, let $\mathcal{C}_{\mathcal{X}_2, (U, \overline{y})}$ be the Higgs-Tate \widehat{R} -algebra defined by \mathcal{X}_2 over $\Sigma_{2, S}$ associated to (U, \overline{y}) ([4] II.10.5) and by $\widehat{\mathcal{C}}_{\mathcal{X}_2, (U, \overline{y})}$ its p -adic completion. There exists a canonical $\Delta_{(U, \overline{y})}$ -equivariant isomorphism compatible with Higgs fields ([4] (IV.5.4.3)):

$$(4.2.5.2) \quad \widehat{\mathcal{C}}_{\mathcal{X}_2, (U, \overline{y})} \xrightarrow{\sim} \mathcal{A}_{\check{X}, \mathcal{X}, 1}^\infty(\overline{R}).$$

For $r \geq 1$, there exists an injection of \widehat{R} -homomorphism from $\mathcal{A}_1^r(\overline{R})$ to $\widehat{\mathcal{C}}_{\mathcal{X}_2, (U, \overline{y})}^{(1/r)}$ the p -adic completion of the Higgs-Tate algebra of thickness $1/r$ ([4] II.12.1) whose cokernel is annihilated by p ([4] (IV.5.4.4)).

4.2.6. The Higgs-Tate algebra admits a globalization as a $\check{\mathcal{B}}_X$ -algebra $\check{\mathcal{C}}_{\mathcal{X}_2}^{(s)}$ of $\widetilde{E}_X^{\mathbb{N}^\circ}$ for $s \in \mathbb{Q}$ ([4] III.10.31). For a pair of an adic $\check{\mathcal{B}}_{X, \mathbb{Q}}$ -module of finite type \mathcal{N} and a Higgs bundle (M, θ) of $\mathrm{HB}_{\mathbb{Q}_p}(\check{X}/\Sigma_{1, S})$ (2.2.6), Abbes and Gros introduced the definition of \mathcal{N} and (M, θ) are associated by an admissible condition associated to $\{\check{\mathcal{C}}_{\mathcal{X}_2}^{(s)}\}_{s \in \mathbb{Q}_{>0}}$ ([4] III.12.11). This allows them to establish an equivalence between certain Higgs bundles and $\check{\mathcal{B}}_{X, \mathbb{Q}}$ -modules satisfying this admissible condition.

Their approach to the p -adic Simpson correspondence is compatible Tsuji's approach via $\iota_{\mathcal{X}}$ (proposition 2.2.7(ii)). More precisely, in view of § 4.2.5, we have:

Proposition 4.2.7. *Let $(\mathcal{M}, \mathcal{M}^\circ)$ be an object of $\mathrm{HC}_{\mathbb{Z}_p, \mathrm{fin}}^r(\check{X}/\Sigma_S)$. Then the Higgs bundle $\iota_{\mathcal{X}}^{-1}(\mathcal{M})$ and the $\check{\mathcal{B}}_{X, \mathbb{Q}}$ -module $T(\mathcal{M}^\circ)_{\mathbb{Q}} (= V^r(\mathcal{M}, \mathcal{M}^\circ))$ are associated.*

4.3. A local version of the p -adic Simpson correspondence. In this subsection, we assume that id_X belongs \mathbf{Q} and fix a geometric generic point \overline{y} of $X_{\overline{y}}$ as in 4.1.8. In particular, X is affine. We review a local description of the p -adic Simpson correspondence on X following ([13], [4] II.13, II.14).

4.3.1. We denote by $\mathrm{HB}_{\mathbb{Z}_p, \mathrm{small}}^{\mathrm{free}}(\widehat{R}_1)$ the category of pairs (M, θ) consisting of a free \widehat{R}_1 -module M of finite rank together with a small Higgs field $\theta : M \rightarrow \xi^{-1}M \otimes_R \widetilde{\Omega}_{R/\sigma_K}^1$ (2.2.10). We note $\Delta_{(X, \overline{y})}$ simply by Δ and refer to ([4] II.6.10) for the definition of Δ_∞ . There exist equivalences of categories ([4] II.13.11, [4] II.14.4):

$$(4.3.1.1) \quad \mathrm{HB}_{\mathbb{Z}_p, \mathrm{small}}^{\mathrm{free}}(\widehat{R}_1) \xrightarrow{\sim} \mathbf{Rep}_{\mathrm{small}}(\Delta_\infty, \widehat{R}_1) \xrightarrow{\sim} \mathbf{Rep}_{\mathrm{small}}(\Delta, \widehat{R}),$$

where the first equivalence preserves the underlying \widehat{R}_1 -modules and the second one is defined by $M \mapsto M \otimes_{\widehat{R}_1} \widehat{R}$. A Higgs bundle (M, θ) of $\mathrm{HB}_{\mathbb{Z}_p, \mathrm{small}}^{\mathrm{free}}(\widehat{R}_1)$ and the corresponding small \widehat{R} -representation of Δ are associated in the sense of 4.2.6 ([4] II.13.16).

If $\mathrm{HB}_{\mathbb{Z}_p, \mathrm{small}}^{\mathrm{free}}(\check{X}/\Sigma_{1, S})$ denotes the full subcategory of $\mathrm{HB}_{\mathbb{Z}_p, \mathrm{conv}}(\check{X}/\Sigma_{1, S})$ (2.2.10) consisting of objects whose underlying $\mathcal{O}_{\check{X}}$ -module is free and θ is small, then there exists a canonical functor $\mathrm{HB}_{\mathbb{Z}_p, \mathrm{small}}^{\mathrm{free}}(\check{X}/\Sigma_{1, S}) \rightarrow \mathrm{HB}_{\mathbb{Z}_p, \mathrm{small}}^{\mathrm{free}}(\widehat{R}_1)$ defined by restriction to $\check{U}^{\overline{y}}$. We denote the composition of above functors by

$$(4.3.1.2) \quad \mathbf{V} : \mathrm{HB}_{\mathbb{Z}_p, \mathrm{small}}^{\mathrm{free}}(\check{X}/\Sigma_{1, S}) \rightarrow \mathbf{Rep}_{\mathrm{small}}(\Delta, \widehat{R}).$$

Proposition 4.3.2. *Let \mathcal{X} be a smooth Cartesian lifting of \check{X} over Σ_S . Then the following diagram is commutative up to isomorphisms (2.2.7(ii))*

$$\begin{array}{ccc} \mathrm{HB}_{\mathbb{Z}_p, \mathrm{small}}^{\mathrm{free}}(\check{X}/\Sigma_{1, S}) & \xrightarrow{\mathbf{V}} & \mathbf{Rep}^{\mathrm{LPM}}(\Delta, \widehat{R}) \\ & \searrow \iota_{\mathcal{X}} & \nearrow T_{(X, \overline{y})} \\ & \mathrm{HC}_{\mathbb{Z}_p, \mathrm{fin}}(\check{X}/\Sigma_S) & \end{array}$$

Proof. Let (M, θ) be an object of $\mathrm{HB}_{\mathbb{Z}_p, \mathrm{small}}^{\mathrm{free}}(\check{X}/\Sigma_{1, S})$ and \mathcal{M} the associated Higgs crystal on $(\check{X}/\Sigma_S)_{\mathrm{HIG}}^\infty$. We denote abusively by (M, θ) the restriction of (M, θ) to $\mathrm{HB}_{\mathbb{Z}_p, \mathrm{small}}^{\mathrm{free}}(\widehat{R}_1)$. By proposition 4.2.7 and ([4] II.13.16), we have the following $\widehat{\mathcal{C}}_{\mathcal{X}_2, (X, \overline{y})}$ -linear isomorphism

$$\widehat{\mathcal{C}}_{\mathcal{X}_2, (X, \overline{y})} \otimes_{\widehat{R}} T_{(X, \overline{y})}(\mathcal{M}) \xrightarrow{\sim} \widehat{\mathcal{C}}_{\mathcal{X}_2, (X, \overline{y})} \otimes_{\widehat{R}_1} (M, \theta),$$

compatible with Higgs fields. We compare it with the isomorphism (4.2.5.1) via (4.2.5.2). Then the assertion follows from taking Higgs field invariants and the fact that $(\widehat{\mathcal{C}}_{\mathcal{X}_2, (X, \overline{y})})^{\theta=0} = \widehat{R}$ ([4] IV.5.2.10). \square

5. FROM \mathbf{C} -REPRESENTATIONS OF THE GEOMETRIC FUNDAMENTAL GROUP TO HIGGS BUNDLES

Let X be a projective S -curve with semi-stable reduction, \check{X} the p -adic completion of $X_{\mathfrak{o}} = X \otimes_{\mathcal{O}_K} \mathfrak{o}$, equipped with the log structure induced by \mathcal{M}_X on X . Let \bar{x} be a geometric point of $X_{\bar{\eta}}$ and $\pi_1(X_{\bar{\eta}}, \bar{x})$ the étale fundamental group. We fix a smooth Cartesian lifting \mathcal{X} of \check{X} to Σ_S .

The p -adic logarithmic homomorphism $\log : 1 + \mathfrak{m} \rightarrow \mathbf{C}$, $x \mapsto \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(x-1)^n}{n}$ admits a section $\exp : y \mapsto \sum_{n=0}^{\infty} \frac{y^n}{n!}$ on the open unit ball of radius $p^{-\frac{1}{p-1}}$ of \mathbf{C} . We fix a section $\text{Exp} : \mathbf{C} \rightarrow 1 + \mathfrak{m}$ of \log , extending \exp (A.0.0.2).

In this section, we globalise the construction in §4.3 and then apply the descent for the p -adic Simpson correspondence over a curve to construct a functor (depending on \mathcal{X} and Exp):

$$\mathbb{H}_{\mathcal{X}, \text{Exp}} : \mathbf{Rep}(\pi_1(X_{\bar{\eta}}, \bar{x}), \mathbf{C}) \rightarrow \text{HB}(X_{\mathbf{C}}).$$

We also provide a description of the essential image of $\mathbb{H}_{\mathcal{X}, \text{Exp}}$.

5.1. Higgs bundles associated to small \mathfrak{o} -representations.

Proposition 5.1.1. *Let X be an S -scheme with semi-stable reduction and geometrically connected generic fiber, and \mathcal{X} a smooth Cartesian lifting of \check{X} over Σ_S .*

(i) *The lifting \mathcal{X} induces a functor (2.2.6, 4.1.9)*

$$\mathbb{H}_{\mathcal{X}} : \mathbf{Rep}_{\text{small}}(\pi_1(X_{\bar{\eta}}, \bar{x}), \mathfrak{o}) \rightarrow \text{HB}_{\mathbb{Z}_p, \text{conv}}(\check{X}/\Sigma_{1,S}).$$

(ii) *If an \mathfrak{o} -representation V is α -small for some $\alpha > \frac{2}{p-1}$, then $\mathbb{H}_{\mathcal{X}}(V)$ is trivial modulo $p^{\alpha - \frac{2}{p-1}}$. In particular, when X has relative dimension one over S , the functor $\mathbb{H}_{\mathcal{X}}$ factors through the full subcategory $\text{HB}_{\mathbb{Z}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$ of $\text{HB}_{\mathbb{Z}_p, \text{conv}}(\check{X}/\Sigma_{1,S})$ (3.2.1).*

Proof. (i) Let V be a α -small \mathfrak{o} -representation of $\pi_1(X_{\bar{\eta}}, \bar{x})$ for $\alpha > \frac{2}{p-1}$. We set $\beta = \alpha - \frac{1}{p-1}$ and take a rational number $s > 0$ such that $\beta > s + \frac{1}{p-1}$. Let U be an object of \mathbf{Q} and \bar{y} a geometric generic point of $U_{\bar{\eta}}$ as in 4.1.8.

By applying equivalences (4.3.1.1) to the α -small representation $V \otimes_{\mathfrak{o}} \widehat{R}$ of $\Delta_{(U, \bar{y})}$, there exists a Higgs bundle $M_{(U, \bar{y})} = (V \otimes_{\mathfrak{o}} \widehat{R}_1, \theta_{(U, \bar{y})})$ of $\text{HB}_{\mathbb{Z}_p, \text{small}}^{\text{free}}(\widehat{R}_1)$ and a canonical $\Delta_{(U, \bar{y})}$ -equivariant $\widehat{\mathcal{C}}_{\mathcal{X}_2, (U, \bar{y})}^{(s)}$ -linear isomorphism compatible with Higgs fields ([4] II.13.16):

$$(5.1.1.1) \quad M_{(U, \bar{y})} \otimes_{\widehat{R}_1} \widehat{\mathcal{C}}_{\mathcal{X}_2, (U, \bar{y})}^{(s)} \xrightarrow{\sim} V \otimes_{\mathfrak{o}} \widehat{\mathcal{C}}_{\mathcal{X}_2, (U, \bar{y})}^{(s)},$$

where $M_{(U, \bar{y})}$ is equipped with the trivial $\Delta_{(U, \bar{y})}$ -action, V is equipped with the trivial Higgs field. Moreover, $\theta_{(U, \bar{y})}$ is β -small and the above isomorphism is congruent to the identity map modulo $p^{\beta - \frac{1}{p-1}}$ in view of the proof of ([4] II.13.15)⁴.

If $g : U' \rightarrow U$ is a morphism of \mathbf{Q} and \bar{y}' a geometric generic point of $U'_{\bar{\eta}}$ above \bar{y} , it induces a $\Delta_{(U', \bar{y}')} -$ equivariant homomorphism $\widehat{\mathcal{C}}_{\mathcal{X}_2, (U, \bar{y})}^{(s)} \rightarrow \widehat{\mathcal{C}}_{\mathcal{X}_2, (U', \bar{y}')}^{(s)}$ via $\Delta_{(U', \bar{y}')} \rightarrow \Delta_{(U, \bar{y})}$. We deduce the following $\Delta_{(U', \bar{y}')} -$ equivariant isomorphism from (5.1.1.1):

$$M_{(U, \bar{y})} \otimes_{\widehat{R}_1} \widehat{\mathcal{C}}_{\mathcal{X}_2, (U', \bar{y}')}^{(s)} \xrightarrow{\sim} M_{(U', \bar{y}')} \otimes_{\widehat{R}'_1} \widehat{\mathcal{C}}_{\mathcal{X}_2, (U', \bar{y}')}^{(s)}.$$

By taking $\Delta_{(U', \bar{y}')} -$ invariant and ([4] II.11.8), we obtain an isomorphism of Higgs bundles

$$(5.1.1.2) \quad c_g : (M_{(U, \bar{y})}, \theta_{(U, \bar{y})}) \otimes_{\widehat{R}_1} \widehat{R}'_1 \xrightarrow{\sim} (M_{(U', \bar{y}')}, \theta_{(U', \bar{y}')}).$$

The transition isomorphisms $\{c_g\}$ satisfy a cocycle condition. Then the data $(M_{(U, \bar{y})}, \theta_{(U, \bar{y})}; c_g)$ descent to a locally free $\mathcal{O}_{\check{X}}$ -module M of rank r and a β -small Higgs field $\theta : M \rightarrow \xi^{-1} M \otimes_{\mathcal{O}_{\check{X}}} \Omega_{\check{X}/\Sigma_{1,S}}^1$. The construction of $V \mapsto (M, \theta)$ is clearly functorial.

⁴In *loc.cit.*, the isomorphism (5.1.1.1) is represented by a matrix E ([4] II.13.15.4), and the claim that E is congruent to id modulo $p^{\beta - r}$ should be “modulo $p^{\beta - \frac{1}{p-1}}$ ”.

(ii) Since θ is β -small, it remains to show the assertion for underlying vector bundle.

The isomorphism (5.1.1.1) coincides with the identity modulo $p^{\alpha - \frac{2}{p-1}}$ and so is its Δ -invariants. Then we deduce that the transition isomorphism (5.1.1.2) is the identity modulo $p^{\alpha - \frac{2}{p-1}}$. This finishes the proof. \square

The functor $\mathbb{H}_{\mathcal{X}}$ is compatible with the functor T (4.2.3.1). More precisely, we have:

Corollary 5.1.2. *The composition (2.2.7(ii), (4.2.3.1))*

$$T \circ \iota_{\mathcal{X}} \circ \mathbb{H}_{\mathcal{X}} : \mathbf{Rep}_{\text{small}}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathfrak{o}) \rightarrow \mathbf{HB}_{\mathbb{Z}_p, \text{conv}}(\check{X}/\Sigma_{1,S}) \rightarrow \mathbf{HC}_{\mathbb{Z}_p, \text{fin}}(\check{X}/\Sigma_S) \rightarrow \mathbf{Mod}(\check{\mathcal{B}}_X)$$

is isomorphic to the canonical functor $V \mapsto \check{\beta}_X^*(V)$ (4.1.6.1), (4.1.10.1).

Proof. It suffices to show that for each pair (U, \overline{y}) , the composition of the above functor with $T_{(U, \overline{y})} \circ \iota_{\mathcal{X}} \circ \mathbb{H}_{\mathcal{X}}$ is isomorphic to $V \mapsto V \otimes_{\mathfrak{o}} \widehat{R}$. We may assume that $U = X$ belongs to \mathbf{Q} . In this case, the composition $\mathbf{V} \circ \mathbb{H}_{\mathcal{X}}$ (4.3.1.2) is isomorphic to the functor $V \mapsto V \otimes_{\mathfrak{o}} \widehat{R}$. Then the assertion follows from proposition 4.3.2. \square

5.1.3. In the following of this section, we keep the assumption in the beginning of § 5. We denote by $\mathbf{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$ the essential image of $\mathbf{HB}_{\mathbb{Z}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$ in $\mathbf{HB}_{\mathbb{Q}_p}(\check{X}/\Sigma_{1,S})$. By passing to categories up to isogeny, the functor $\mathbb{H}_{\mathcal{X}}$ extends to the following functor

$$(5.1.3.1) \quad \mathbb{H}_{\mathcal{X}} : \mathbf{Rep}_{\text{small}}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathbf{C}) \rightarrow \mathbf{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S}),$$

that we still denote by $\mathbb{H}_{\mathcal{X}}$. We also consider $\mathbf{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{Y}/\Sigma_{1,S})$ as a full subcategory of $\mathbf{HB}(Y_{\mathbf{C}})$ via equivalences (2.2.10.1).

Proposition 5.1.4. *Let V be a small \mathbf{C} -representation of $\pi_1(X_{\overline{\eta}}, \overline{x})$. Then the Higgs bundle $\mathbb{H}_{\mathcal{X}}(V)$ and the $\check{\mathcal{B}}_{X, \mathbf{Q}}$ -module $\check{\beta}_{X, \mathbf{Q}}^*(V)$ are associated (4.2.6).*

Proof. Let V° be an \mathfrak{o} -representaiton lattice of V . There exists an integer $r \geq 1$ and an object $(\mathcal{M}, \mathcal{M}^{\circ})$ of $\mathbf{HC}_{\mathbb{Z}_p, \text{fin}}^r(\check{X}/\Sigma_S)$ (4.2.3) such that $\mathcal{M}^{\circ} \simeq \iota_{\mathcal{X}}(\mathbb{H}_{\mathcal{X}}(V^{\circ}))$. Then the assertion follows from proposition 4.2.7 and corollary 5.1.2. \square

Lemma 5.1.5. *Let K' be a finite extension of K , Y a projective $S' = \text{Spec}(\mathcal{O}_{K'})$ -curve with semi-stable reduction, $f : Y \rightarrow X_{S'}$ an η' -cover and \mathcal{Y} a smooth Cartesian lifting of \check{Y} over $\Sigma_{S'}$.*

(i) *The twisted inverse image functor $\check{f}_{\mathcal{Y}, \mathcal{X}}^*$ (2.3.4.1) sends $\mathbf{HB}_{\mathbb{Z}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$ (resp. $\mathbf{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$) to $\mathbf{HB}_{\mathbb{Z}_p, \text{small}}^{\text{DW}}(\check{Y}/\Sigma_{1,S'})$ (resp. $\mathbf{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{Y}/\Sigma_{1,S'})$).*

(ii) *The following diagram is commutative up to isomorphisms:*

$$\begin{array}{ccc} \mathbf{Rep}_{\text{small}}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathbf{C}) & \xrightarrow{\mathbb{H}_{\mathcal{X}}} & \mathbf{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S}) \\ \downarrow & & \downarrow \check{f}_{\mathcal{Y}, \mathcal{X}}^* \\ \mathbf{Rep}_{\text{small}}(\pi_1(Y_{\overline{\eta}}, \overline{y}), \mathbf{C}) & \xrightarrow{\mathbb{H}_{\mathcal{Y}}} & \mathbf{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{Y}/\Sigma_{1,S'}) \end{array}$$

where the left vertical arrow is induced by the canonical morphism $\pi_1(Y_{\overline{\eta}}, \overline{y}) \rightarrow \pi_1(X_{\overline{\eta}}, \overline{x})$.

Proof. (i) The assertion follows from corollary 2.3.6(ii) and the fact that strongly semi-stable vector bundles of degree zero are preserved by inverse image.

(ii) The composition $T \circ \iota_{\mathcal{X}}$ is compatible with twisted inverse image functor of Higgs bundles and inverse image functor in Faltings ringed topos (proposition 4.2.4). Applying $T \circ \iota_{\mathcal{X}}$ to the above diagram, the assertion follows from corollary 5.1.2 and full faithfulness of $T_{\mathbf{Q}} : \mathbf{HC}_{\mathbb{Z}_p, \text{fin}}(\check{X}/\Sigma_S)_{\mathbf{Q}} \rightarrow \mathbf{Mod}(\overline{\mathcal{B}}_X)_{\mathbf{Q}}$ ([4] theorem IV.6.4.9). \square

5.2. Construction of $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ via descent. To extend the construction of $\mathbb{H}_{\mathcal{X}}$ to all \mathbf{C} -representations, we need the twisted inverse image functor for Higgs bundles, that we summarize in the following.

5.2.1. Let K' be a finite extension of K , Y a projective $S' = \text{Spec}(\mathcal{O}_{K'})$ -curve with semi-stable reduction, \mathcal{Y} a smooth Cartesian lifting of \check{Y} over $\Sigma_{1, S'}$, and $f : Y \rightarrow X_{S'}$ an η' -cover. Let $\text{Exp} : \mathbf{C} \rightarrow 1 + \mathfrak{m}$ be a section of \log as in the beginning of this section.

Let (M, θ) be a Higgs bundle over $X_{\mathbf{C}}$ and $\mathcal{A}_{f^*(\theta)}$ the spectral cover of the Higgs field $f^*(\theta)$ over $Y_{\mathbf{C}}$ (2.4.1). The obstruction to lift $\check{f} : \check{Y} \rightarrow \check{X}$ to $\Sigma_{2, S}$ defines a Higgs bundle $\mathcal{L}_{f, \theta}^{\text{Exp}} \in H^1(Y_{\mathbf{C}}, \mathcal{A}_{f^*(\theta)}^{\times})$ (A.3.0.1). The twisted inverse image functor is defined by (A.3.0.2)

$$f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^{\circ} : \text{HB}(X_{\mathbf{C}}) \rightarrow \text{HB}(Y_{\mathbf{C}}), \quad (M, \theta) \mapsto f_{\mathbf{C}}^*(M, \theta) \otimes_{\mathcal{A}_{f^*(\theta)}} \mathcal{L}_{f, \theta}^{\text{Exp}}.$$

When it restricts to small Higgs bundles, it is compatible with the functor $\check{f}_{\mathcal{Y}, \mathcal{X}}^*$ (2.3.4.1) and is independent of the choice of Exp (see proposition A.3.1).

5.2.2. We construct a functor

$$(5.2.2.1) \quad \mathbb{H}_{\mathcal{X}, \text{Exp}} : \mathbf{Rep}(\pi_1(X_{\bar{\eta}}, \bar{x}), \mathbf{C}) \rightarrow \text{HB}(X_{\mathbf{C}}).$$

For each \mathbf{C} -representation V of $\mathbf{Rep}(\pi_1(X_{\bar{\eta}}, \bar{x}), \mathbf{C})$, there exists a Galois étale cover $\pi : Y_{\bar{\eta}} \rightarrow X_{\bar{\eta}}$ and a geometric point \bar{y} above \bar{x} such that the restriction of V to $\pi_1(Y_{\bar{\eta}}, \bar{y})$ is small. By semi-stable reduction theorem, there exists a finite extension K' of K , a projective $S' = \text{Spec}(\mathcal{O}_{K'})$ -curve Y with semi-stable reduction and geometric generic fiber $Y_{\bar{\eta}}$ and an η' -cover $f : Y \rightarrow X_{S'}$ such that $f_{\bar{\eta}} = \pi$. We may assume f is a Galois η' -cover ([36] proposition 5.11(iii)) and we take a smooth Cartesian lifting \mathcal{Y} of \check{Y} over $\Sigma_{S'}$.

Since V is a small representation of $\pi_1(Y_{\bar{\eta}}, \bar{y})$, we obtain a small Higgs bundle $\mathbb{H}_{\mathcal{Y}}(V)$ over $Y_{\mathbf{C}}$ (5.1.3.1), whose Higgs field is denoted by ψ . The automorphism group $\text{Aut}(Y_{\bar{\eta}}/X_{\bar{\eta}})$, that we denote by G , extends to an action on Y above $X_{S'}$. By proposition 4.2.4, the G -action on V induces a G -action on $\mathbb{H}_{\mathcal{Y}}(V)$:

$$(5.2.2.2) \quad \{\varphi_g : g_{\mathcal{Y}, \mathcal{Y}, \text{Exp}}^{\circ}(\mathbb{H}_{\mathcal{Y}}(V)) \xrightarrow{\sim} \mathbb{H}_{\mathcal{Y}}(V) \mid g \in G\}, \text{ such that } \varphi_g \circ g^{\circ}(\varphi_{g'}) = \varphi_{g'g}.$$

In particular, we deduce that $g^*(c_i(\psi)) = c_i(\psi) \in \Gamma(Y_{\mathbf{C}}, \xi^{-i} \Omega_{Y_{\mathbf{C}}}^{\otimes i})$ for each i and $g \in G$. By descent, we obtain a point $\theta = \{c_i\}_{i=1}^r$ of the Hitchin base $\oplus_{i=1}^r \Gamma(X_{\mathbf{C}}, \xi^{-i} \Omega_{X_{\mathbf{C}}}^{\otimes i})$ of $X_{\mathbf{C}}$ such that $f_{\mathbf{C}}^*(c_i) = c_i(\psi)$. Since $f \circ g = f$, we have (A.2.6.1)

$$\mathcal{L}_{f, \theta}^{\text{Exp}} \simeq \mathcal{L}_{f \circ g, \theta}^{\text{Exp}} \simeq \mathcal{L}_{g, f^*(\theta)}^{\text{Exp}} \otimes g^*(\mathcal{L}_{f, \theta}^{\text{Exp}}).$$

Then the data (5.2.2.2) defines a usual descent data on $\mathbb{H}_{\mathcal{Y}}(V) \otimes_{\mathcal{A}_{f^*(\theta)}} (\mathcal{L}_{f, \theta}^{\text{Exp}})^{-1}$:

$$\{\phi_g : g_{\mathbf{C}}^*(\mathbb{H}_{\mathcal{Y}}(V) \otimes_{\mathcal{A}_{f^*(\theta)}} (\mathcal{L}_{f, \theta}^{\text{Exp}})^{-1}) \xrightarrow{\sim} \mathbb{H}_{\mathcal{Y}}(V) \otimes_{\mathcal{A}_{f^*(\theta)}} (\mathcal{L}_{f, \theta}^{\text{Exp}})^{-1} \mid g \in G\}, \text{ such that } \phi_g \circ g_{\mathbf{C}}^*(\phi_{g'}) = \phi_{g'g}.$$

By descent, we obtain a Higgs bundle $\mathbb{H}_{\mathcal{X}, \text{Exp}}(V)$ on $X_{\mathbf{C}}$ such that $f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^{\circ}(\mathbb{H}_{\mathcal{X}, \text{Exp}}(V)) \simeq \mathbb{H}_{\mathcal{Y}}(V)$.

Proposition 5.2.3. (i) *The construction $V \mapsto \mathbb{H}_{\mathcal{X}, \text{Exp}}(V)$ is independent of the choice of the Galois η' -cover $f : Y \rightarrow X_{S'}$ and of the smooth Cartesian lifting \mathcal{Y} over $\Sigma_{S'}$.*

(ii) *The functor $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ is well-defined.*

(iii) *The functor $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ is exact and commutes with duals.*

Proof. (i) Let K'' be a finite extension of K , Z an $S'' = \text{Spec}(\mathcal{O}_{K''})$ -curve with semi-stable reduction, $g : Z \rightarrow X_{S''}$ a Galois η'' -cover and \bar{z} a geometric point of $Z_{\bar{\eta}}$ above \bar{x} such that V is small as a \mathbf{C} -representation of $\pi_1(Z_{\bar{\eta}}, \bar{z})$. To prove the assertion, we may assume that g dominates f . Then the assertion follows from lemma 5.1.5 and proposition A.2.1(iv).

(ii) Let $u : V \rightarrow V'$ be a morphism of $\mathbf{Rep}(\pi_1(X_{\bar{\eta}}, \bar{x}), \mathbf{C})$. By (i), we may choose a Galois η' -cover $f : Y \rightarrow X_{S'}$ as in §5.2.2 such that V and V' are both small as \mathbf{C} -representation of $\pi_1(Y_{\bar{\eta}}, \bar{y})$. Then the functoriality of $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ follows from that of $\mathbb{H}_{\mathcal{Y}}$ and of $f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^{\circ}$.

(iii) The functor $\mathbb{H}_{\mathcal{X}}$ clearly satisfies these properties. The assertion follows from the fact that functors $\mathbb{H}_{\mathcal{X}}$ and $f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^{\circ}$ satisfy the corresponding properties (proposition A.2.1). \square

To describe the essential image of $\mathbb{H}_{\mathcal{X}, \text{Exp}}$, we introduce the following categories:

Definition 5.2.4. We denote by $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$ the full subcategory of $\text{HB}(X_{\mathbf{C}})$ consisting of Higgs bundles (M, θ) such that there exists a finite extension K' of K , a projective $S' = \text{Spec}(\mathcal{O}_{K'})$ -curve Y with semi-stable reduction, a Galois η' -cover $f : Y \rightarrow X_{S'}$ and a smooth Cartesian lifting \mathcal{Y} of \check{Y} over $\Sigma_{S'}$ such that $f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^{\circ}(M, \theta)$ belongs to the essential image of $\text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{Y}/\Sigma_{1, S'})$ in $\text{HB}(Y_{\mathbf{C}})$.

By ([36], corollaire 5.8), the twisted inverse image functor $f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^{\circ}$ sends $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$ to $\text{HB}_{\mathcal{Y}, \text{Exp}}^{\text{pDW}}(Y_{\mathbf{C}})$. In view of proposition 5.1.1(ii), lemma 5.1.5 and the construction of $\mathbb{H}_{\mathcal{X}, \text{Exp}}$, we have:

Corollary 5.2.5. (i) *The functor $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ factors through the full subcategory $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$:*

$$(5.2.5.1) \quad \mathbb{H}_{\mathcal{X}, \text{Exp}} : \mathbf{Rep}(\pi_1(X_{\check{\eta}}, \bar{x}), \mathbf{C}) \rightarrow \text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}}).$$

(ii) *With the assumption and notation of 5.1.5, the following diagram is commutative up to isomorphisms:*

$$\begin{array}{ccc} \mathbf{Rep}(\pi_1(X_{\check{\eta}}, \bar{x}), \mathbf{C}) & \xrightarrow{\mathbb{H}_{\mathcal{X}, \text{Exp}}} & \text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}}) \\ \downarrow & & \downarrow f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^{\circ} \\ \mathbf{Rep}(\pi_1(Y_{\check{\eta}}, \bar{y}), \mathbf{C}) & \xrightarrow{\mathbb{H}_{\mathcal{Y}, \text{Exp}}} & \text{HB}_{\mathcal{Y}, \text{Exp}}^{\text{pDW}}(Y_{\mathbf{C}}) \end{array}$$

5.3. Semistability of $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$.

Proposition 5.3.1. (i) *Every Higgs bundle of $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$ is semi-stable of degree zero.*

(ii) *The category $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$ is closed under extension and duals.*

Proof. (i) Since Deninger–Werner vector bundles are semi-stable of degree zero ([10] theorem 13), we deduce that every object of $\text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1, S})$ is a semi-stable Higgs bundle of degree zero over $X_{\mathbf{C}}$. Given a Higgs bundle (M, θ) of $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$, its twisted inverse image along an η -cover is semi-stable of degree zero. By proposition A.2.1(iii), we conclude that (M, θ) is also semi-stable of degree zero.

(ii) The corresponding results hold for Deninger–Werner vector bundles ([10] proposition 9, theorem 11). We deduce corresponding results for $\text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1, S})$. Then the assertion follows from proposition A.3.1. \square

Proposition 5.3.2. (i) *Every Higgs line bundle (L, θ) of degree zero over $X_{\mathbf{C}}$ belongs to $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$.*

(ii) *When the genus of $X_{\mathbf{C}}$ is 0, 1, every semi-stable Higgs bundle of degree zero belongs to $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$.*

Proof. (i) By proposition A.2.1, we may assume θ is small and L is of degree zero after applying a twisted inverse image. Recall that a line bundle of $\text{Pic}_{X/S}^0(\mathfrak{o})$ is of Deninger–Werner over \check{X} ([10] claim of theorem 12). Since the cokernel of the inclusion

$$\text{Pic}_{X/S}^0(\mathfrak{o}) \hookrightarrow \text{Pic}_{X_{\eta}/K}^0(\mathbf{C})$$

is torsion ([9] theorem 4.1), there exists an integer $N \geq 1$ such that $L^{\otimes N} \in \text{Pic}_{X/S}^0(\mathfrak{o})$. By applying the argument of proposition 3.2.3, there exists a finite extension K' of K , an $S' (= \text{Spec}(\mathcal{O}_{K'}))$ -curve Y with semi-stable reduction and an η' -cover $f : Y \rightarrow X_{S'}$ such that the usual inverse image $f_{\mathbf{C}}^*(L) \simeq L^{\otimes N}$. We take a smooth Cartesian lifting \mathcal{Y} of \check{Y} over $\Sigma_{S'}$.

Line bundle $\mathcal{L}_{f, \theta}$ admits an integral model with trivial special fiber and belongs to the image of $\text{Pic}_{X/S}^0(\mathfrak{o})$ (2.4.5). Then so is the underlying bundle $L^{\otimes N} \otimes_{\mathcal{O}_{X_{\mathbf{C}}}} \mathcal{L}_{f, \theta}$ of $f_{\mathcal{Y}, \mathcal{X}}^{\circ}(L, \theta)$. This finishes the proof.

(ii) The assertion is clear if the genus is zero.

When $X_{\mathbf{C}}$ is an elliptic curve, every stable Higgs bundle of degree zero has rank one ([15] proposition 4.3, [5]). Every semi-stable Higgs bundle of degree zero is isomorphic to a successive extension of Higgs line bundle of degree zero. Then the assertion follows from assertion (i) and proposition 5.3.1(ii). \square

Remark 5.3.3. (i) Assertion (i) (resp. (ii)) of proposition 5.3.2 is also treated by Heuer [18] (resp. Heuer–Mann–Werner [19]) by a different method in a general context.

(ii) Recall that any Deninger–Werner vector bundle over a curve is semi-stable of degree zero. When the genus $X_{\mathbf{C}}$ is ≥ 2 , there exist semi-stable Higgs bundles of degree zero whose underlying bundles are not semi-stable and does not belong to $\mathrm{VB}^{\mathrm{DW}}(X_{\mathbf{C}})$. Let L be a square root of $\Omega_{X_{\mathbf{C}}}^1$, which has degree $g - 1 > 0$. The isomorphism $\theta : L \xrightarrow{\sim} L^{-1} \otimes \Omega_{X_{\mathbf{C}}}^1$ defines a Higgs field on $L \oplus L^{-1}$ and this Higgs bundle is stable while its underlying bundle is not semi-stable. It is expected that $(L \oplus L^{-1}, \theta)$ belongs to $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$ (Conjecture 1.1.8).

6. PARALLEL TRANSPORT FOR HIGGS BUNDLES

In this section, we will apply the parallel transport construction to Higgs bundles of $\mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}})$ (definition 5.2.4) and prove the following result.

Theorem 6.0.1. (i) *There exists a canonical functor*

$$(6.0.1.1) \quad \mathbb{V}_{\mathcal{X}, \mathrm{Exp}} : \mathrm{HB}_{\mathcal{X}, \mathrm{Exp}}^{\mathrm{pDW}}(X_{\mathbf{C}}^{\mathrm{an}}) \rightarrow \mathbf{Rep}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathbf{C}).$$

(ii) *The functors $\mathbb{V}_{\mathcal{X}, \mathrm{Exp}}$ and $\mathbb{H}_{\mathcal{X}, \mathrm{Exp}}$ (5.2.5.1) are quasi-inverse to each others.*

6.1. Parallel transport via Faltings topoi. In [36] § 8, we construct the parallel transport functor for certain modules in Faltings topos inspired by Deninger–Werner’s construction [10]. In this subsection, we present another construction of this functor via h-descent for étale sheaves and we apply this functor to modules associated to Higgs bundles of $\mathrm{HB}_{\mathbb{Z}_p, \mathrm{small}}^{\mathrm{DW}}(\check{X}/\Sigma_{1,S})$ (§ 3.2.1).

In this subsection, let X be an \overline{S} -model of a smooth proper \overline{K} -variety (§1.2.2), $(\tilde{E}_X, \overline{\mathcal{B}}_X)$ the Faltings ringed topos associated to the pair $(X_{\overline{\eta}} \rightarrow X)$. If Y is another \overline{S} -model and $\varphi : Y \rightarrow X$ an \overline{S} -morphism, we denote the functorial morphism of Faltings ringed topoi associated to φ by:

$$(6.1.0.1) \quad \Phi : (\tilde{E}_Y, \overline{\mathcal{B}}_Y) \rightarrow (\tilde{E}_X, \overline{\mathcal{B}}_X).$$

Definition 6.1.1 ([36] définition 8.5). (i) We say a $\overline{\mathcal{B}}_{X,n}$ -module M_n is *potentially free of finite type* if it is of finite type and there exists a proper \overline{S} -morphism $\varphi : Y \rightarrow X$ such that $\varphi_{\overline{\eta}}$ is finite étale and that the inverse image $\Phi^*(M_n)$ is isomorphic to a free $\overline{\mathcal{B}}_{Y,n}$ of finite type.

(ii) We say a $\overline{\mathcal{B}}_X$ -module $M = (M_n)_{n \geq 1}$ is *potentially free of finite type* if it is adic of finite type (1.2.4) and for each $n \geq 1$, the $\overline{\mathcal{B}}_{X,n}$ -module M_n is potentially free of finite type.

We denote by $\mathbf{Mod}^{\mathrm{pltf}}(\overline{\mathcal{B}}_{X,n})$ (resp. $\mathbf{Mod}^{\mathrm{pltf}}(\check{\overline{\mathcal{B}}}_X)$) the full subcategory of $\mathbf{Mod}(\overline{\mathcal{B}}_{X,n})$ (resp. $\mathbf{Mod}(\check{\overline{\mathcal{B}}}_X)$) consisting of potentially free of finite type objects, and by $\mathbf{Mod}_{\mathrm{cohd}}^{\mathrm{pltf}}(\overline{\mathcal{B}}_{X,n})$ (resp. $\mathbf{Mod}_{\mathrm{cohd}}^{\mathrm{pltf}}(\check{\overline{\mathcal{B}}}_X)$) the full subcategory of $\mathbf{Mod}^{\mathrm{pltf}}(\overline{\mathcal{B}}_{X,n})$ (resp. $\mathbf{Mod}^{\mathrm{pltf}}(\check{\overline{\mathcal{B}}}_X)$) consisting of $\overline{\mathcal{B}}_{X,n}$ -modules satisfying cohomological descent (§4.1.4) (resp. the full subcategory of $\mathbf{Mod}^{\mathrm{pltf}}(\overline{\mathcal{B}}_{X,n})$ consisting of $\overline{\mathcal{B}}$ -modules $M = (M_n)_{n \geq 1}$ such that each M_n belongs to $\mathbf{Mod}_{\mathrm{cohd}}^{\mathrm{pltf}}(\overline{\mathcal{B}}_{X,n})$).

The inverse image of morphisms of ringed topoi $\beta_n : (\tilde{E}_X, \overline{\mathcal{B}}_{X,n}) \rightarrow (X_{\overline{\eta}, \mathrm{fét}}, \mathfrak{o}_n)$, $\check{\beta} : (\tilde{E}_X^{\mathrm{N}^\circ}, \check{\overline{\mathcal{B}}}_X) \rightarrow (X_{\overline{\eta}, \mathrm{fét}}^{\mathrm{N}^\circ}, \check{\mathfrak{o}})$ induce functors (4.1.6, 4.1.10):

$$(6.1.1.1) \quad \beta_n^* : \mathbf{LocSys}(X_{\overline{\eta}, \mathrm{fét}}, \mathfrak{o}_n) \rightarrow \mathbf{Mod}_{\mathrm{cohd}}^{\mathrm{pltf}}(\overline{\mathcal{B}}_{X,n}), \quad \check{\beta}^* : \mathbf{LocSys}(X_{\overline{\eta}, \mathrm{fét}}^{\mathrm{N}^\circ}, \check{\mathfrak{o}}) \rightarrow \mathbf{Mod}_{\mathrm{cohd}}^{\mathrm{pltf}}(\check{\overline{\mathcal{B}}}_X).$$

Indeed, if \mathbb{L} is an object of $\mathbf{LocSys}(X_{\overline{\eta}, \mathrm{fét}}, \mathfrak{o}_n)$ and Z is a finite étale cover of $X_{\overline{\eta}}$ trivializing \mathbb{L} , then the integral closure $X^Z \rightarrow X$ of X in Z is an $\overline{\eta}$ -cover. Then the morphism of Faltings ringed topoi, induced by $X^Z \rightarrow X$, trivializes $\beta_n^*(\mathbb{L})$ (c.f. [36] proposition 8.7 for more details). Moreover, $\beta_n^*(\mathbb{L})$ satisfies the cohomological descent by corollary 4.1.5.

6.1.2. In the following, we construct in another direction, the parallel transport functor:

$$(6.1.2.1) \quad \mathbb{L}_n : \mathbf{Mod}^{\text{pltf}}(\overline{\mathcal{B}}_{X,n}) \rightarrow \mathbf{Mod}(X_{\overline{\eta},\text{fét}}, \mathfrak{o}_n).$$

Let \mathcal{N} be an object of $\mathbf{Mod}^{\text{pltf}}(\overline{\mathcal{B}}_{X,n})$ and $\varphi : Y \rightarrow X$ an $\overline{\eta}$ -cover trivializing \mathcal{N} as in 6.1.1. We consider the hypercovering $\varepsilon_\bullet : Y_\bullet \rightarrow X$, where for $m \geq 0$, $Y_m \rightarrow X$ is defined by the $(m+1)$ -th self product of Y over X . Let $\varepsilon_{\bullet, \overline{\eta}} : (Y_{\bullet, \overline{\eta}})_{\text{ét}} \rightarrow X_{\overline{\eta}, \text{ét}}$ be the augmentation of simplicial étale topoi. In the following, we apply h-descent to ε_\bullet to construct $\mathbb{L}_n(\mathcal{N})$.

For each integer $m \geq 1$, the morphism of Faltings topoi associated to $\varepsilon_m : Y_m \rightarrow X$ trivializes \mathcal{N} . By lemma 4.1.3, we have a canonical decomposition of the cohomology $\varepsilon_m^*(\mathcal{N})$ in terms of the connected components of $Y_{m, \overline{\eta}}$:

$$(6.1.2.2) \quad \Gamma(\tilde{E}_{Y_m}, \varepsilon_m^*(\mathcal{N}))_\# \simeq \bigoplus_{Z \in \text{Con}(Y_{m, \overline{\eta}})} \Gamma(\tilde{E}_{Z \rightarrow Y_m}, \varepsilon_m^*(\mathcal{N}))_\#,$$

where the functor $\# : \mathbf{Mod}(\mathfrak{o}) \rightarrow \mathbf{Mod}(\mathfrak{o})$ is defined by $M \mapsto \text{Hom}_{\mathfrak{o}}(\mathfrak{m}, M)$ (1.2.3.1).

By Faltings' comparison theorem ([17] theorem 1.16), each component of (6.1.2.2) is isomorphic to $\Gamma(Z_{\text{ét}}, \mathfrak{o}_n^{\oplus r})_\# \simeq (\mathfrak{o}_n^{\oplus r})_\#$, where r is the rank of \mathcal{N} . We denote abusively by $\Gamma(\tilde{E}_{Y_m}, \varepsilon_m^*(\mathcal{N}))_\#$ the \mathfrak{o}_n -module of $(Y_{m, \overline{\eta}})_{\text{ét}}$, defined by the constant \mathfrak{o}_n -module $\Gamma(\tilde{E}_{Z \rightarrow Y_m}, \varepsilon_m^*(\mathcal{N}))_\#$ in each connected component Z of $Y_{m, \overline{\eta}}$.

For $m' > m$, every morphism $\phi : Y_{m'} \rightarrow Y_m$ in the hypercovering $Y_\bullet \rightarrow X$ induces a canonical morphism:

$$(6.1.2.3) \quad \Phi^* : \Gamma(\tilde{E}_{Y_m}, \varepsilon_m^*(\mathcal{N}))_\# \rightarrow \Gamma(\tilde{E}_{Y_{m'}}, \varepsilon_{m'}^*(\mathcal{N}))_\#.$$

If Z is a connected component of $Y_{m', \overline{\eta}}$, then Φ^* induces an isomorphism

$$\Gamma(\tilde{E}_{\phi(Z) \rightarrow Y_m}, \varepsilon_m^*(\mathcal{N}))_\# \xrightarrow{\sim} \Gamma(\tilde{E}_{Z \rightarrow Y_{m'}}, \varepsilon_{m'}^*(\mathcal{N}))_\#$$

Therefore, Φ^* induces an isomorphism of \mathfrak{o}_n -modules on $(Y_{m', \overline{\eta}})_{\text{ét}}$:

$$(6.1.2.4) \quad \phi_\#^*(\Gamma(\tilde{E}_{Y_m}, \varepsilon_m^*(\mathcal{N}))_\#) \xrightarrow{\sim} \Gamma(\tilde{E}_{Y_{m'}}, \varepsilon_{m'}^*(\mathcal{N}))_\#.$$

In summary, we obtain a descent datum of étale sheaves $\Gamma(\tilde{E}_{Y_\bullet}, \varepsilon_\bullet^*(\mathcal{N}))_\# \in (Y_{\bullet, \overline{\eta}})_{\text{ét}}$ with respect to the hypercovering $Y_{\bullet, \overline{\eta}} \rightarrow X_{\overline{\eta}}$. By h-descent ([38] 0GF0), this descent datum gives rise to an étale \mathfrak{o}_n -module on $X_{\overline{\eta}}$, denoted by $\mathbb{L}_n(\mathcal{N})$.

Proposition 6.1.3. (i) *The construction $\mathcal{N} \mapsto \mathbb{L}_n(\mathcal{N})$ defines a functor*

$$\mathbb{L}_n : \mathbf{Mod}^{\text{pltf}}(\overline{\mathcal{B}}_{X,n}) \rightarrow \mathbf{Mod}(X_{\overline{\eta}, \text{fét}}, \mathfrak{o}_n)$$

(ii) *The functor \mathbb{L}_n factors through the full subcategory $\mathbf{Mod}(X_{\overline{\eta}, \text{fét}}, \mathfrak{o}_n)$ (4.1.10.3).*

Proof. (i) We first show that $\mathcal{N} \mapsto \mathbb{L}_n(\mathcal{N})$ is independent of the choice of the η' -cover $f : Y \rightarrow X_{S'}$. Let $Z \rightarrow X$ be another $\overline{\eta}$ -cover trivializing \mathcal{N} . After replacing Z by $Z \times_X Y$, we may assume that $Z \rightarrow X$ factors through $\pi : Z \rightarrow Y$. The inverse image of the descent data $\Gamma(\tilde{E}_{Y_\bullet}, \varepsilon_\bullet^*(\mathcal{N}))_\#$ defined by Y_\bullet via $\pi_\bullet : Z_\bullet \rightarrow Y_\bullet$ is isomorphic to the descent data $\Gamma(\tilde{E}_{Z_\bullet}, \varepsilon_\bullet^*(\mathcal{N}))_\#$ defined by Z_\bullet . Then the claim follows.

Given a morphism $u : \mathcal{N} \rightarrow \mathcal{N}'$ of $\mathbf{Mod}^{\text{pltf}}(\overline{\mathcal{B}}_{X,n})$, we may take an $\overline{\eta}$ -cover $Y \rightarrow Z$ trivializing both \mathcal{N} and \mathcal{N}' . Then the functoriality of \mathbb{L}_n follows.

Assertion (ii) follows from the fact that $\mathbb{L}_n(\mathcal{N})$ is trivialized by a finite étale cover $Y_{\overline{\eta}} \rightarrow X_{\overline{\eta}}$. \square

By a similar argument of [36] proposition 8.16, we conclude the following proposition:

Proposition 6.1.4. *Let \mathbb{L} be an object of $\mathbf{LocSys}(X_{\overline{\eta}, \text{fét}}, \mathfrak{o}_n)$. Then, there exists a canonical and functorial almost isomorphism of \mathfrak{o}_n -modules of $X_{\overline{\eta}, \text{fét}}$:*

$$\mathbb{L}_n(\beta_n^*(\mathbb{L})) \xrightarrow{\sim} \mathbb{L}.$$

Proposition 6.1.5. *The functors β_n^* and \mathbb{L}_n induce equivalences of categories quasi-inverse to each other up to almost isomorphisms (1.2.1):*

$$(6.1.5.1) \quad (\beta_n^*)^\alpha : \alpha\text{-}\mathbf{LocSys}(X_{\overline{\eta}, \text{fét}}, \mathfrak{o}_n) \rightleftarrows \alpha\text{-}\mathbf{Mod}_{\text{coh}}^{\text{pltf}}(\overline{\mathcal{B}}_{X,n}) : (\mathbb{L}_n)^\alpha.$$

Remark 6.1.6. In a recent preprint of Mann–Werner [22], the authors obtained a similar result between local systems with integral models and certain trivializable modules over a proper adic space of finite type over \mathbf{C} via v -descent.

The above proposition follows from proposition 6.1.4 and the following lemma.

Lemma 6.1.7. *Let \mathcal{N} be an object of $\mathbf{Mod}_{\text{cohd}}^{\text{pltf}}(\overline{\mathcal{B}}_{X,n})$. Then, there exists a canonical isomorphism of $(\overline{\mathcal{B}}_{X,n})^\alpha$ -modules:*

$$(6.1.7.1) \quad (\beta_n^*(\mathbb{L}_n(\mathcal{N})))^\alpha \xrightarrow{\sim} \mathcal{N}^\alpha.$$

Proof. We simply write $\mathbb{L}_n(\mathcal{N})$ as \mathbb{L} . In view of the construction in § 6.1.2, we have an isomorphism of $(\overline{\mathcal{B}}_{Y_\bullet,n})$ -modules:

$$\varepsilon^*(\mathbb{L} \otimes_{\mathfrak{o}_n} \overline{\mathcal{B}}_{X,n}) \simeq \Gamma(\tilde{E}_{Y_\bullet}, \varepsilon^*(\mathcal{N}))_\# \otimes_{\mathfrak{o}_n} \overline{\mathcal{B}}_{Y_\bullet,n}.$$

Since $\varepsilon^*(\mathcal{N})$ is isomorphic to a free $(\overline{\mathcal{B}}_{Y_\bullet,n})$ -module of finite type, the following canonical morphisms are almost isomorphisms by Faltings' comparison theorem ([17] theorem 1.16)

$$\Gamma(\tilde{E}_{Y_\bullet}, \varepsilon^*(\mathcal{N})) \otimes_{\mathfrak{o}_n} \overline{\mathcal{B}}_{Y_\bullet,n} \rightarrow \Gamma(\tilde{E}_{Y_\bullet}, \varepsilon^*(\mathcal{N}))_\# \otimes_{\mathfrak{o}_n} \overline{\mathcal{B}}_{Y_\bullet,n}, \quad \Gamma(\tilde{E}_{Y_\bullet}, \varepsilon^*(\mathcal{N})) \otimes_{\mathfrak{o}_n} \overline{\mathcal{B}}_{Y_\bullet,n} \rightarrow \varepsilon^*(\mathcal{N}).$$

Then, these almost isomorphisms fit into the following diagram

$$(6.1.7.2) \quad \begin{array}{ccc} \mathbb{L} \otimes_{\mathfrak{o}_n} \overline{\mathcal{B}}_{X,n} & & \mathcal{N} \\ \approx \downarrow & & \downarrow \approx \\ \mathrm{R}\varepsilon_*(\varepsilon^*(\beta_n^*(\mathbb{L}_n(\mathcal{N})))) & \xleftarrow{\sim} \mathrm{R}\varepsilon_*(\Gamma(\tilde{E}_{Y_\bullet}, \varepsilon^*(\mathcal{N})) \otimes_{\mathfrak{o}_n} \overline{\mathcal{B}}_{Y_\bullet,n}) \xrightarrow{\sim} & \mathrm{R}\varepsilon_*(\varepsilon^*(\mathcal{N})) \end{array}$$

Two horizontal arrows are almost isomorphisms due to the cohomological descent. Then we conclude the isomorphism of $(\overline{\mathcal{B}}_{X,n})^\alpha$ -modules (6.1.7.1). \square

6.1.8. For $m \geq n \geq 1$, the functors \mathbb{L}_m and \mathbb{L}_n are compatible up to almost isomorphisms via reduction ([36] lemma 8.13). By considering the projective system $(\mathbb{L}_n)_{n \geq 1}$, we obtain following functor ([36] 8.14.1):

$$(6.1.8.1) \quad \mathbb{L} : \mathbf{Mod}^{\text{pltf}}(\check{\overline{\mathcal{B}}}) \rightarrow \mathbf{Mod}(X_{\overline{\eta}, \text{fét}}^{\text{N}^\circ}, \check{\mathfrak{o}}), \quad M = (M_n)_{n \geq 1} \mapsto (\mathfrak{m} \otimes_{\mathfrak{o}} \mathbb{L}_n(M_n))_{n \geq 1}.$$

By passing to categories up to isogeny, the functor $\mathbb{L}_{\mathbb{Q}}$ factors through the full subcategory $\mathbf{LocSys}(X_{\overline{\eta}, \text{fét}}^{\text{N}^\circ}, \check{\mathfrak{o}})_{\mathbb{Q}}$, which is equivalent to $\mathbf{Rep}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathbf{C})$ (4.1.10.2) (c.f. [36] 8.14):

$$\mathbb{L}_{\mathbb{Q}} : \mathbf{Mod}^{\text{pltf}}(\check{\overline{\mathcal{B}}})_{\mathbb{Q}} \rightarrow \mathbf{Rep}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathbf{C}).$$

6.2. Parallel transport for Deninger–Werner Higgs bundles. In the following of this section, we keep the assumption and notation of §5.

Theorem 6.2.1. *Let (M, θ) be an object of $\mathrm{HB}_{\mathbb{Z}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$, \mathcal{M} the associated Higgs crystal and $\check{h} : (\tilde{E}_{\underline{X}}^{\text{N}^\circ}, \check{\overline{\mathcal{B}}}_{\underline{X}}) \rightarrow (\tilde{E}_X^{\text{N}^\circ}, \check{\overline{\mathcal{B}}}_X)$ the canonical morphism of ringed topoi (4.1.6.3).*

(i) *Then $\check{h}^*(T(\mathcal{M}))$ is potentially free of finite type. In particular, this allows us to define the parallel transport functor $\mathbb{V}_{\mathcal{X}}$ as the composition (4.2.3.1):*

$$(6.2.1.1) \quad \mathbb{V}_{\mathcal{X}} : \mathrm{HB}_{\mathbb{Z}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S}) \xrightarrow{\iota_{\mathcal{X}}} \mathrm{HC}_{\mathbb{Z}_p, \text{fin}}(\check{X}/\Sigma_S) \xrightarrow{\check{h}^* \circ T} \mathbf{Mod}^{\text{pltf}}(\check{\overline{\mathcal{B}}}_{\underline{X}}) \xrightarrow{\mathbb{L}} \mathbf{Mod}(X_{\overline{\eta}, \text{fét}}^{\text{N}^\circ}, \check{\mathfrak{o}}).$$

(ii) *The $\check{\overline{\mathcal{B}}}_{\underline{X}}$ -module $\check{h}^*(T(\mathcal{M}))$ satisfies the cohomological descent.*

(iii) *There exists a canonical isomorphism of $(\check{\overline{\mathcal{B}}}_X)^\alpha$ -modules:*

$$(6.2.1.2) \quad (\check{\beta}_X^*(\mathbb{V}_{\mathcal{X}}(M, \theta)))^\alpha \simeq (T(\mathcal{M}))^\alpha.$$

Proof. (i) Let n be an integer ≥ 1 and $\varphi : Y \rightarrow X_{S'}$ be an η' -cover trivializing \mathcal{M}_n as in theorem 3.2.2. Since the functor T is compatible with the inverse image functor of Higgs crystals and Φ^* (6.1.0.1) (proposition 4.2.4), then assertion (i) follows.

Assertion (ii) follows from corollary 4.1.5 and the following lemma.

Lemma 6.2.2. *The $\overline{\mathcal{B}}_{X,n}$ -module $T(\mathcal{M})_n$ is locally free of finite type.*

Proof. By ([4] II.5.17), there exists an étale covering $\{U_i \rightarrow X\}_{i \in I}$ of \mathbf{Q} such that $M|_{U_i}$ is a trivial vector bundle of rank r .

Let \overline{y} be a geometric generic point of $U_{i,\overline{\eta}}$, $U_{i,\overline{\eta}}^{\overline{y}}$ the connected component of $U_{i,\overline{\eta}}$ containing \overline{y} and we take again the notation of §4.3 for the pair (U_i, \overline{y}) . In view of equivalences (4.3.1.1), there exists a trivialization of the underlying $\overline{R}/p^n \overline{R}$ -module of $T(\mathcal{M})_{n,U}|_{(U_{i,\overline{\eta}}^{\overline{y}})_{\text{fét}}}$ (c.f. proposition 4.1.2):

$$\nu_{\overline{y}}(T(\mathcal{M})_{n,U}|_{(U_{i,\overline{\eta}}^{\overline{y}})_{\text{fét}}}) \simeq (\overline{R}/p^n \overline{R})^{\oplus r}.$$

Moreover, there exists a finite étale Galois cover $V_i^{\overline{y}} \rightarrow U_{i,\overline{\eta}}^{\overline{y}}$ such that the action of $\Delta_{(U_i, \overline{y})} = \pi_1(U_{i,\overline{\eta}}^{\overline{y}}, \overline{y})$ on the above module factors through $\text{Gal}(V_i^{\overline{y}}/U_{i,\overline{\eta}}^{\overline{y}})$. In view of (4.2.2.3) and ([4] III.10.5), the above isomorphism induces a trivialization of $\overline{\mathcal{B}}_{V_i^{\overline{y}} \rightarrow U_i, n}$ -modules of $\tilde{E}_{V_i^{\overline{y}} \rightarrow U_i}$:

$$T(\mathcal{M})_n|_{(V_i^{\overline{y}} \rightarrow U_i)} \simeq (\overline{\mathcal{B}}_{V_i^{\overline{y}} \rightarrow U_i, n})^{\oplus r}.$$

Since $\{(V_i^{\overline{y}} \rightarrow U_i) \rightarrow (X_{\overline{\eta}} \rightarrow X)\}_{i \in I, \overline{y} \in \text{Con}(U_{i,\overline{\eta}})}$ forms a covering in Faltings site, the lemma follows. \square

(iii) By assertion (ii) and lemma 6.1.7, we conclude the following $(\check{\mathcal{B}}_{\underline{X}})^{\alpha}$ -linear isomorphism:

$$(6.2.2.1) \quad (\check{\beta}_{\underline{X}}^*(\mathbb{V}_{\mathcal{X}}(M, \theta)))^{\alpha} \simeq (\check{h}^*(T(\mathcal{M})))^{\alpha}.$$

On the one hand, the following canonical morphism is an isomorphism in view of lemma 4.1.7(ii)

$$\check{\beta}_{\underline{X}}^*(\mathbb{V}_{\mathcal{X}}(M, \theta)) \xrightarrow{\sim} \check{h}_*(\check{\beta}_{\underline{X}}^*(\mathbb{V}_{\mathcal{X}}(M, \theta))).$$

On the other hand, by lemma 6.2.2, the following canonical morphism is an isomorphism:

$$T(\mathcal{M}) \xrightarrow{\sim} \check{h}_* \check{h}^*(T(\mathcal{M})).$$

Then the assertion follows from applying \check{h}_* to the isomorphism (6.2.2.1). \square

By passing to categories up to isogeny and § 6.1.8, we deduce from $\mathbb{V}_{\mathcal{X}}$ the following functor, that we abusively denote by $\mathbb{V}_{\mathcal{X}}$:

$$(6.2.2.2) \quad \mathbb{V}_{\mathcal{X}} : \text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S}) \rightarrow \mathbf{Rep}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathbf{C}).$$

Corollary 6.2.3. *Let (M, θ) be a Higgs bundle of $\text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$. Then $\check{\beta}_{X, \mathbf{Q}}^*(\mathbb{V}_{\mathcal{X}}(M, \theta))$ and (M, θ) are associated (4.2.6).*

Proof. It follows from propositions 4.2.7, 6.1.4 and theorem 6.2.1(iii). \square

Corollary 6.2.4. *Let K' be a finite extension of K , Y a projective $S' = \text{Spec}(\mathcal{O}_{K'})$ -curve with semi-stable reduction, $f : Y \rightarrow X_{S'}$ an η' -cover and \mathcal{Y} a smooth Cartesian lifting of \check{Y} over $\Sigma_{S'}$. The following diagram is commutative up to isomorphisms:*

$$\begin{array}{ccc} \text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S}) & \xrightarrow{\mathbb{V}_{\mathcal{X}}} & \mathbf{Rep}(\pi_1(X_{\overline{\eta}}, \overline{x}), \mathbf{C}) \\ \downarrow \check{f}_{\mathcal{Y}, \mathcal{X}}^* & & \downarrow \\ \text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{Y}/\Sigma_{1,S'}) & \xrightarrow{\mathbb{V}_{\mathcal{Y}}} & \mathbf{Rep}(\pi_1(Y_{\overline{\eta}}, \overline{y}), \mathbf{C}) \end{array}$$

where the right vertical arrow is defined by the canonical morphism $\pi_1(Y_{\overline{\eta}}, \overline{y}) \rightarrow \pi_1(X_{\overline{\eta}}, \overline{x})$.

Proof. By Faltings' comparison theorem ([17] theorem 1.16), the following functor is fully faithful (c.f. [36] proposition 8.8):

$$(6.2.4.1) \quad \check{\beta}_{\mathbb{Q}}^* : \mathbf{Rep}(\pi_1(X_{\bar{\eta}}, \bar{x}), \mathbf{C}) \simeq \mathbf{LocSys}(X_{\bar{\eta}, \text{fét}}, \check{\mathfrak{o}})_{\mathbb{Q}} \rightarrow \mathbf{Mod}(\check{\mathcal{B}}_{\underline{X}})_{\mathbb{Q}}.$$

Composing the above functor in the above diagram, the assertion follows from corollary 6.2.3 and the fact that $f_{\mathcal{Y}, \mathcal{X}}^*$ and the inverse image functor of Faltings topos are compatible with the p -adic Simpson correspondence (proposition 4.2.4). \square

6.3. Construction of $\mathbb{V}_{\mathcal{X}, \text{Exp}}$ via descent.

6.3.1. In this subsection, we construct the functor (6.0.1.1):

$$(6.3.1.1) \quad \mathbb{V}_{\mathcal{X}, \text{Exp}} : \text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}}) \rightarrow \mathbf{Rep}(\pi_1(X_{\bar{\eta}}, \bar{x}), \mathbf{C}).$$

For each Higgs bundle (M, θ) of $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$, there exists a finite extension K' of K , a projective $S' = \text{Spec}(\mathcal{O}_{K'})$ -curve Y with semi-stable reduction and geometric generic fiber $Y_{\bar{\eta}}$, an η' -cover $f : Y \rightarrow X_{S'}$ and a smooth Cartesian lift \mathcal{Y} of \check{Y} over $\Sigma_{S'}$ such that $f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^{\circ}(M, \theta)$ belongs to $\text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{Y}/\Sigma_{1, S'})$. We may assume f is a Galois η' -cover ([36] proposition 5.11(iii)) and we set $G = \text{Aut}(Y_{\bar{\eta}}/X_{\bar{\eta}})$. By considering $\pi_1(Y_{\bar{\eta}}, \bar{y})$ -representation $V := \mathbb{V}_{\mathcal{Y}}(f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^{\circ}(M, \theta))$ as an object of $\mathbf{LocSys}(X_{\bar{\eta}, \text{fét}}^{\text{No}}, \check{\mathfrak{o}})_{\mathbb{Q}}$, it is equipped with a collection of G -action data induced by that of $f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^{\circ}(M, \theta)$:

$$\phi_g : g^*(V) \xrightarrow{\sim} V, \text{ such that } \phi_g \circ g^*(\phi_{g'}) = \phi_{g'g}, \quad \forall g, g' \in G.$$

By the following lemma and descent, we obtain a \mathbf{C} -representation of $\pi_1(X_{\bar{\eta}}, \bar{x})$, that we denote by $\mathbb{V}_{\mathcal{X}, \text{Exp}}(M, \theta)$.

Lemma 6.3.2. *There exists a finite free $\pi_1(Y_{\bar{\eta}}, \bar{y})$ -invariant \mathfrak{o} -lattice Γ of V such that each ϕ_g induces an isomorphism $g^*(\Gamma) \xrightarrow{\sim} \Gamma$.*

Proof. Let $\tilde{\Gamma}$ be a finite free $\pi_1(Y_{\bar{\eta}}, \bar{y})$ -invariant \mathfrak{o} -lattice of V . We set $\Gamma = \sum_{g \in G} \phi_g(g^*(\tilde{\Gamma}))$ in V . As Γ is finitely generated torsion-free \mathfrak{o} -modules, it is also a finite free $\pi_1(Y_{\bar{\eta}}, \bar{y})$ -invariant \mathfrak{o} -lattice of V . For each $g, h \in G$, the action of ϕ_h on the component $\phi_g(g^*(\tilde{\Gamma}))$ fits into the following diagram:

$$h^*g^*(\tilde{\Gamma}) \xrightarrow{h^*(\phi_g)} h^*(\phi_g(g^*(\tilde{\Gamma}))) \xrightarrow{\phi_h} \phi_h(h^*(\phi_g(g^*(\tilde{\Gamma})))) = \phi_{gh}((gh)^*\tilde{\Gamma}) \subset \Gamma,$$

whose composition is ϕ_{gh} . Hence ϕ_h preserves Γ and is surjective. On the other hand, ϕ_h is clearly injective and the lemma follows. \square

Theorem 6.0.1 follows from the following proposition:

Proposition 6.3.3. (i) *The construction $(M, \theta) \mapsto \mathbb{V}_{\mathcal{X}, \text{Exp}}(M, \theta)$ is independent of the choice of the η' -cover $f : Y \rightarrow X_{S'}$ and of the smooth Cartesian lifting \mathcal{Y} over $\Sigma_{S'}$.*

(ii) *The functor $\mathbb{V}_{\mathcal{X}, \text{Exp}}$ is well-defined.*

(iii) *The functor $\mathbb{V}_{\mathcal{X}, \text{Exp}}$ is compatible with twisted inverse images of Higgs bundles and the restriction of representations as in corollary 6.2.4.*

(iv) *The functors $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ and $\mathbb{V}_{\mathcal{X}, \text{Exp}}$ are quasi-inverse to each others.*

Proof. (i) Let K'' be a finite extension of K , Z an $S'' = \text{Spec}(\mathcal{O}_{K''})$ -curve with semi-stable reduction, $g : Z \rightarrow X_{S''}$ an η'' -cover and \mathcal{Z} a smooth Cartesian lifting of \check{Z} over $\Sigma_{S''}$ such that $f_{\mathcal{Z}, \mathcal{X}, \text{Exp}}^{\circ}(M, \theta)$ belongs to $\text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{Z}/\Sigma_{1, S''})$. To prove the assertion, we may assume that g dominates f . Then the assertion follows from corollary 6.2.4 and proposition A.2.1.

(ii) Let $u : (M, \theta) \rightarrow (M', \theta')$ be a morphism of $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$. By (i), we may choose an η' -cover $f : Y \rightarrow X_{S'}$ as in §6.3.1 and a smooth Cartesian lift \mathcal{Y} of \check{Y} over Σ_S such that both $f^{\circ}(M, \theta)$ and $f^{\circ}(M', \theta')$ belong to $\text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{Y}/\Sigma_{1, S})$. Then the functoriality of $\mathbb{V}_{\mathcal{X}, \text{Exp}}$ follows from that of $\mathbb{V}_{\mathcal{Y}}$ and of descent for \mathbf{C} -representations.

(iii) The assertion follows from corollary 6.2.4 and the construction of $\mathbb{V}_{\mathcal{X}, \text{Exp}}$.

(iv) We ignore \mathcal{X}, Exp from the notations $\mathbb{V}_{\mathcal{X}, \text{Exp}}, \mathbb{H}_{\mathcal{X}, \text{Exp}}$ for simplicity. We show that for each object V of $\mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C})$ (resp. (M, θ) of $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$), there exist functorial isomorphisms

$$(6.3.3.1) \quad V \xrightarrow{\sim} \mathbb{V}(\mathbb{H}(V)), \quad (M, \theta) \xrightarrow{\sim} \mathbb{H}(\mathbb{V}(M, \theta)).$$

By (iii), corollary 5.2.5(ii) and descents, it suffices to prove above isomorphisms under the assumption that V is small and that (M, θ) belongs to $\text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$ and $\mathbb{V}(M, \theta)$ is small.

On the one hand, let V be a small \mathbf{C} -representation of $\pi_1(X_{\overline{K}}, \overline{x})$. In view of corollary 5.1.2(i), proposition 6.1.4 and theorem 6.2.1(i), we have a functorial isomorphism of $\check{\mathcal{B}}_{X, \mathbb{Q}}$ -module:

$$\check{\beta}_{X, \mathbb{Q}}^*(V) \xrightarrow{\sim} \check{\beta}_{X, \mathbb{Q}}^*(\mathbb{V}_{\mathcal{X}} \mathbb{H}_{\mathcal{X}}(V)).$$

In view of the full faithfulness of functor (6.2.4.1), the first isomorphism of (6.3.3.1) follows.

On the other hand, let (M, θ) be a Higgs bundle of $\text{HB}_{\mathbb{Q}_p, \text{small}}^{\text{DW}}(\check{X}/\Sigma_{1,S})$ such that $\mathbb{H}(M, \theta)$ is α -small for some $\alpha > \frac{2}{p-1}$. By proposition 5.1.4 and corollary 6.2.3, the following pairs are associated:

$$\left((M, \theta), \check{\beta}_{X, \mathbb{Q}}^*(\mathbb{V}(M, \theta)) \right), \quad \left(\mathbb{H}(\mathbb{V}(M, \theta)), \check{\beta}_{X, \mathbb{Q}}^*(\mathbb{V}(M, \theta)) \right).$$

Then we deduce a functorial isomorphism $(M, \theta) \xrightarrow{\sim} \mathbb{H}(\mathbb{V}(M, \theta))$ via the p -adic Simpson correspondence. \square

6.4. Comparison with Deninger–Werner’s theory. We first clarify the relationship between $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$ and Deninger–Werner’s vector bundles.

Proposition 6.4.1. *Let M be a vector bundle over $X_{\mathbf{C}}$. The following properties are equivalent:*

- (i) *The Higgs bundle $(M, 0)$ belongs to $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$.*
- (ii) *There exists a \mathbf{C} -representation V of $\pi_1(X_{\overline{K}}, \overline{x})$ and a $\check{\mathcal{B}}_{X, \mathbb{Q}}$ -linear isomorphism $\check{\beta}^*(V) \simeq \check{\sigma}^*(M)$ (i.e. M is Weil–Tate in the sense of [36] definition 10.3).*
- (iii) *The vector bundle M belongs to $\text{VB}^{\text{pDW}}(X_{\mathbf{C}})$ (3.1.2).*

Proof. The Higgs bundle $(M, 0)$ is clearly small. The equivalence between (i) and (ii) follows from corollary 6.2.3 and ([36] proposition 11.7).

When the twisted inverse image functor applies to Higgs bundles with trivial Higgs field, it coincides with the usual inverse image functor. Hence, condition (i) is equivalent to condition (iii). \square

Corollary 6.4.2. *Via the equivalence in proposition 6.4.1, Deninger–Werner’s functor $\mathbb{V}_{X_{\mathbf{C}}}^{\text{DW}}$ (3.1.3.1) is compatible with functor $\mathbb{V}_{\mathcal{X}, \text{Exp}}$.*

Proof. Deninger–Werner’s construction can be reinterpreted as a $\check{\mathcal{B}}_{\mathbb{Q}}$ -admissibility and is therefore compatible with the p -adic Simpson correspondence ([36] théorème 1.10, proposition 11.7). Then the assertion follows from corollary 6.2.3. \square

It is expected that the category $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$ is equivalent to the category of semi-stable Higgs bundles of degree zero over $X_{\mathbf{C}}$ (see conjecture 1.1.8). The latter is an abelian category. The same property holds for $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$.

Corollary 6.4.3. *The categories $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$ and $\text{VB}^{\text{pDW}}(X_{\mathbf{C}})$ are abelian.*

Proof. Since $\mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C})$ is abelian and $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ is an exact equivalence, the assertion for $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$ follows. The assertion for $\text{VB}^{\text{pDW}}(X_{\mathbf{C}})$ follows from that of $\text{HB}_{\mathcal{X}, \text{Exp}}^{\text{pDW}}(X_{\mathbf{C}})$ and proposition 6.4.1. \square

6.4.4. Proof of proposition 1.1.9. The equivalence $\mathbb{H}_{\mathcal{X}, \text{Exp}}$ induces an isomorphism of extension classes:

$$\text{Ext}_{\mathbf{Rep}(\pi_1(X_{\overline{K}}, \overline{x}), \mathbf{C})}^1(V, \mathbf{C}) \xrightarrow{\sim} \text{Ext}_{\text{HB}(X_{\mathbf{C}})}^1((M, \theta), (\mathcal{O}_{X_{\mathbf{C}}}, 0)).$$

The left hand side is isomorphic to $H_{\text{ét}}^1(X_{\overline{K}}, V)$. And the right hand side is calculated by the first Higgs cohomology. Then we deduce an isomorphism (1.1.9.1):

$$H_{\text{ét}}^1(X_{\overline{K}}, V) \simeq H^1(X_{\mathbf{C}}, M \xrightarrow{\theta} M \otimes_{\mathcal{O}_{X_{\mathbf{C}}}} \xi^{-1} \Omega_{X_{\mathbf{C}}}^1). \quad \square$$

6.4.5. When the Higgs bundle $(M, 0)$ has the trivial Higgs field, the above isomorphism degenerates to a decomposition

$$H_{\text{ét}}^1(X_{\overline{K}}, V) \simeq H^1(X_{\mathbf{C}}, M) \oplus H^0(X_{\mathbf{C}}, M \otimes_{\mathcal{O}_{X_{\mathbf{C}}}} \xi^{-1} \Omega_{X_{\mathbf{C}}}^1).$$

The first component $H^1(X_{\mathbf{C}}, M)$ represents the extension classes of vector bundles.

APPENDIX A. TWISTED INVERSE IMAGE FOR HIGGS BUNDLES OVER CURVES, D'APRÈS FALTINGS

Tongmu He, Daxin Xu

In this appendix, we define the twisted inverse image functor for Higgs bundles along a finite morphism between smooth proper p -adic curves. We present its construction in the following setting, which is more general than the original definition of Faltings [13].

Let $f : Y \rightarrow X$ be a finite morphism between smooth proper \mathbf{C} -curves, and $\mathfrak{Y}, \mathfrak{X}$ two smooth liftings of Y, X over B_{dR}^+/ξ^2 respectively. The p -adic logarithmic map $\log : 1 + \mathfrak{m} \rightarrow \mathbf{C}$, defined by $x \mapsto \sum_{n=1}^{\infty} (-1)^{n+1} \frac{(x-1)^n}{n}$, gives rise to a short exact sequence:

$$(A.0.0.1) \quad 0 \rightarrow \mu_{p^\infty} \rightarrow 1 + \mathfrak{m} \xrightarrow{\log} \mathbf{C} \rightarrow 0.$$

For any $\varepsilon \in \mathbb{Q}$, we denote by B_ε° (resp. B_ε) the open (resp. closed) ball $\{x \in \mathbf{C} \mid |x| < \varepsilon\}$ (resp. $|x| \leq \varepsilon$) of radius ε . We set $r_p = p^{-\frac{1}{p-1}}$. The exponential map $\exp : B_{r_p}^\circ \rightarrow 1 + \mathfrak{m}$, defined by $x \mapsto \sum_{n \geq 0} \frac{x^n}{n!}$, is a section of \log on $B_{r_p}^\circ$. Since $(1 + \mathfrak{m}, \times)$ is divisible and is therefore an injective abelian group, we can choose a homomorphism, called *exponential map*

$$(A.0.0.2) \quad \text{Exp} : \mathbf{C} \rightarrow 1 + \mathfrak{m},$$

satisfying: $\log \circ \text{Exp} = \text{id}$ and $\text{Exp}|_{B_{r_p}^\circ} = \exp$.

With above data, we will define the twisted inverse image functor between Higgs bundles:

$$(A.0.0.3) \quad f_{\mathfrak{Y}, \mathfrak{X}, \text{Exp}}^\circ : \text{HB}(X) \rightarrow \text{HB}(Y).$$

A.1. Splitting of the p -adic logarithm for commutative smooth algebraic groups. We review the p -adic logarithm for commutative smooth algebraic groups, following Fargues [14]. Then we define a functorial splitting of the p -adic logarithm, which is a key ingredient in the construction of $f_{\mathfrak{Y}, \mathfrak{X}, \text{Exp}}^\circ$.

A.1.1. Let G be a commutative smooth algebraic group over \mathbf{C} , $\text{Lie } G$ its Lie algebra, G^{rig} the associated rigid-analytic group over \mathbf{C} and G^{an} the associated analytic group in the sense of Berkovich (c.f. [14] §1). If V is a \mathbf{C} -vector space, we denote by $\mathbb{G}_a^{\text{rig}} \otimes V$ the rigid analytic group, which represents the functor $X \mapsto (\Gamma(X, \mathcal{O}_X) \otimes V, +)$. After choosing a basis of V , it is isomorphic to a sum of copies of $\mathbb{G}_a^{\text{rig}}$.

For a rigid-analytic space (resp. analytic space in the sense of Berkovich) X , we denote by $|X|$ the set of points of X (resp. the topological space of X). We can identify $G(\mathbf{C})$ with the set $|G^{\text{rig}}|$.

By [14] théorème 1.2, there exists an open analytic subgroup U_G^{an} of G^{an} such that:

$$U_G^{\text{an}} = \{g \in |G^{\text{an}}| \mid \lim_{n \rightarrow \infty} p^n g = 0\}.$$

We denote by U_G the associated rigid-analytic subgroup of G^{rig} . There exists a unique étale morphism of rigid-analytic groups

$$\log_G : U_G \rightarrow \mathbb{G}_a^{\text{rig}} \otimes \text{Lie } G,$$

inducing the identity map on Lie algebras and an isomorphism on a neighborhood of the identity element (c.f. [14] proposition 9). When G admits a smooth model over \mathfrak{o} , we present an explicit description of this local isomorphism in §A.1.4. Let $G[p^\infty]$ be the p -divisible group defined by p -power torsions of G . We obtain a short exact sequence of rigid analytic groups (see [14] théorème 1.2 and the following proposition):

$$(A.1.1.1) \quad 0 \rightarrow G[p^\infty] \rightarrow U_G \xrightarrow{\log_G} \mathbb{G}_a^{\text{rig}} \otimes \text{Lie } G \rightarrow 0.$$

When $G = \mathbb{G}_m$, $|U_G| = 1 + \mathfrak{m}$ and the underlying points of the above sequence recovers (A.0.0.1).

Proposition A.1.2. (i) *The rigid-analytic group U_G is p -divisible in the sense that $p : U_G \rightarrow U_G$ is finite, faithfully flat and satisfies $\forall g \in U_G^{\text{an}}, \lim_{n \rightarrow \infty} p^n g = 0$ for the topology of the Berkovich space U_G^{an} (c.f. [14] définition 2). In particular, the morphism \log_G is surjective.*

(ii) *The homomorphism Exp (A.0.0.2) induces a functorial homomorphism in G :*

$$(A.1.2.1) \quad \text{Exp}_G : \text{Lie } G \rightarrow |U_G|,$$

which is a section of $|\log_G|$ (A.1.1.1) on underlying points.

Proof. (i) By Rosenlicht's theorem, the group G is an extension of by an abelian variety by a linear algebraic group. Then, we see that $p : G \rightarrow G$ is finite and faithfully flat, and the corresponding properties hold for G^{rig} and U_G . The last property follows from the definition of U_G .

The second assertion follows from ([14] proposition 16).

(ii) The p -adic logarithm induces an isomorphism if G is unipotent. We may assume that G is isomorphic to an extension $0 \rightarrow T \rightarrow G \rightarrow A \rightarrow 0$ of an abelian variety by a torus.

Let $\Lambda = T_p G$ be the Tate module of G . By [14] théorème 3.3, there exists a unique \mathbf{C} -linear morphism

$$f : \text{Lie } G \rightarrow \Lambda \otimes_{\mathbb{Z}_p} \mathbf{C}(-1)$$

and a diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & G[p^\infty] & \longrightarrow & U_G & \xrightarrow{\log_G} & \mathbb{G}_a^{\text{rig}} \otimes \text{Lie } G \longrightarrow 0 \\ & & \downarrow \sim & & \downarrow & & \downarrow f \\ 0 & \longrightarrow & \Lambda(-1) \otimes \mu_{p^\infty} & \longrightarrow & \Lambda(-1) \otimes \widehat{\mathbb{G}}_m^{\text{rig}} & \xrightarrow{\text{id} \otimes \log_{\mathbb{G}_m}} & \mathbb{G}_a^{\text{rig}} \otimes (\Lambda \otimes_{\mathbb{Z}_p} \mathbf{C}(-1)) \longrightarrow 0 \end{array}$$

where $\widehat{\mathbb{G}}_m$ denotes the p -adic completion of \mathbb{G}_m over \mathfrak{o} , and the top extension is isomorphic to the pullback of the lower one by f . If we consider the underlying points of the above sequence, we obtain

$$\begin{array}{ccccccc} 0 & \longrightarrow & G[p^\infty] & \longrightarrow & |U_G| & \xrightarrow{|\log_G|} & \text{Lie } G \longrightarrow 0 \\ & & \downarrow \sim & & \downarrow & \square & \downarrow f \\ 0 & \longrightarrow & \Lambda(-1) \otimes_{\mathbb{Z}_p} \mu_{p^\infty} & \longrightarrow & \Lambda(-1) \otimes_{\mathbb{Z}_p} (1 + \mathfrak{m}) & \xrightarrow{\text{id} \otimes \log} & \Lambda \otimes_{\mathbb{Z}_p} \mathbf{C}(-1) \longrightarrow 0 \\ & & & & & \nwarrow \text{id} \otimes \text{Exp} & \end{array}$$

where the right square is Cartesian. Then the section $\text{Exp} : \mathbf{C} \rightarrow 1 + \mathfrak{m}$ of \log induces an exponential map $\text{Exp}_G : \text{Lie } G \rightarrow |U_G|$. In view of the construction, the map Exp_G is functorial on G . \square

Remark A.1.3. Fargues [14] showed that the functor $\mathcal{G} \mapsto (T_p \mathcal{G}, \text{Lie } \mathcal{G}, f)$ induces an equivalence between the category of p -divisible rigid-analytic groups over \mathbf{C} and the category of triples (T, W, f) consisting of a free \mathbb{Z}_p -module T of finite rank, a \mathbf{C} -vector space W and a \mathbf{C} -linear morphism $f : W \rightarrow T \otimes_{\mathbb{Z}_p} \mathbf{C}(-1)$. Moreover, Scholze and Weinstein showed that the full subcategory of these linear algebra data (T, W, f) such that f is injective, is equivalent to the category of p -divisible groups over \mathfrak{o} [29].

A.1.4. Let $\mathcal{G} = \mathrm{Spf}(R)$ be a formally smooth formal group of dimension d over $\mathrm{Spf}(\mathfrak{o})$. Its generic fiber $\mathcal{G}^{\mathrm{rig}}$ is a rigid-analytic group. After taking an isomorphism $R \simeq \mathfrak{o}[[x_1, \dots, x_d]]$, the rigid-analytic space $\mathcal{G}^{\mathrm{rig}}$ is isomorphic to the open unit ball of dimension d . For $\varepsilon \in \mathbb{Q}_{<1}$, the open (resp. closed) ball of radius ε forms a rigid-analytic subgroup of $\mathcal{G}^{\mathrm{rig}}$, that we denote by $\mathcal{G}_\varepsilon^{\circ, \mathrm{rig}}$ (resp. $\mathcal{G}_\varepsilon^{\mathrm{rig}}$). These subgroups are independent of the choice of formal coordinates of \mathcal{G} (see [14] exemple 1 (7)). We have an admissible covering:

$$\mathcal{G}^{\mathrm{rig}} = \bigcup_{\varepsilon} \mathcal{G}_\varepsilon^{\mathrm{rig}} = \bigcup_{\varepsilon} \mathcal{G}_\varepsilon^{\circ, \mathrm{rig}}.$$

In this case, the p -adic logarithm of $\mathcal{G}^{\mathrm{rig}}$ is given by the logarithm of the formal group \mathcal{G} ([31] §2.4) and is defined on $\mathcal{G}^{\mathrm{rig}}$ ([14] exemple 5 (4)).

Suppose that there exists a smooth group scheme \mathcal{G} over \mathfrak{o} of relative dimension d with generic fiber G . Its Lie algebra $\mathrm{Lie} \mathcal{G}$ is an \mathfrak{o} -lattice of $\mathrm{Lie} G$. We denote abusively by $\mathrm{Lie} \mathcal{G}$ (resp. $\mathrm{Lie} G$) the associated affine space over \mathfrak{o} (resp. \mathbf{C}), equipped with the additive group structure.

Let \mathcal{G} (resp. $\mathrm{Lie} \mathcal{G}$) be the formally smooth formal group defined by the completion of \mathcal{G} (resp. $\mathrm{Lie} \mathcal{G}$) with respect to the augmentation ideal of the unit section over \mathfrak{o} . With the above notations, we have $\mathcal{G}^{\mathrm{rig}} \subset U_G$ and $\mathrm{Lie} \mathcal{G}^{\mathrm{rig}} \subset \mathbb{G}_a^{\mathrm{rig}} \otimes \mathrm{Lie} G$. Recall $r_p = p^{-\frac{1}{p-1}}$. The restriction of the logarithm map \log_G to $\mathcal{G}^{\mathrm{rig}}$

$$\log_G : \mathcal{G}^{\mathrm{rig}} \rightarrow \mathrm{Lie} G \otimes \mathbb{G}_a^{\mathrm{rig}}$$

induces an isomorphism between $\mathcal{G}_{r_p}^{\circ, \mathrm{rig}}$ and $\mathrm{Lie} \mathcal{G}_{r_p}^{\circ, \mathrm{rig}}$ ([31] §2.4). We denote the inverse of this map by

$$(A.1.4.1) \quad \exp_G : \mathrm{Lie} \mathcal{G}_{r_p}^{\circ, \mathrm{rig}} \xrightarrow{\sim} \mathcal{G}_{r_p}^{\circ, \mathrm{rig}} \subset U_G.$$

For $\alpha \in \mathbb{Q}_{>0}$ and $\varepsilon = p^{-\alpha}$, the sets of points $|\mathrm{Lie} \mathcal{G}_\varepsilon^{\mathrm{rig}}| \subset |\mathbb{G}_a^{\mathrm{rig}} \otimes \mathrm{Lie} G|$ identify with $p^\alpha \mathrm{Lie} \mathcal{G} \subset \mathrm{Lie} G$. For $\alpha \in \mathbb{Q}_{>\frac{1}{p-1}}$, the above homomorphism on underlying classical points

$$(A.1.4.2) \quad |\exp_G| : p^\alpha \mathrm{Lie} \mathcal{G} \rightarrow |U_G|$$

is compatible with Exp_G (A.1.2.1). Moreover, if we regard $|\exp_G|$ as a homomorphism of p -adic Lie groups, its differential map coincides with the identity map on $\mathrm{Lie} G$.

A.2. Construction of $f_{\mathfrak{Y}, \mathfrak{X}, \mathrm{Exp}}^\circ$. We keep the notation in the beginning of appendix A. Let (M, θ) be a Higgs bundle over X , and $\mathcal{A}_{f^*(\theta)}$ the spectral cover of $f^*(\theta) : f^*(M) \rightarrow f^*(M) \otimes \xi^{-1} f^*(\Omega_X^1)$ over Y (2.4.1). We denote by \mathcal{P} the relative Picard functor of $\mathrm{Spec}_{\mathcal{O}_{X, \mathbf{C}}}(\mathcal{A}_{f^*(\theta)})$ over \mathbf{C} , which is represented by a commutative smooth algebraic group over \mathbf{C} ([26] corollaire 1.5.2, 2.3.2). We denote the exponential map for \mathcal{P} (A.1.2.1) by

$$\mathrm{Exp}_{\mathcal{A}_{f^*(\theta)}} : \mathrm{Lie} \mathcal{P} \simeq H^1(X_{\mathbf{C}}, \mathcal{A}_{f^*(\theta)}) \rightarrow \mathcal{P}(\mathbf{C}) = H^1(X_{\mathbf{C}}, \mathcal{A}_{f^*(\theta)}^\times).$$

The obstruction of lifting f to B_{dR}^+/ξ^2 defines an element o_f of $H^1(Y, \xi f^*(T_X))$, where T_X denotes the tangent sheaf of X over \mathbf{C} . We denote by $\mathcal{L}_{f, \theta}^{\mathrm{Exp}}$ the image of o_f in the following composition:

$$(A.2.0.1) \quad H^1(Y, \xi f^*(T_X)) \rightarrow H^1(Y, \mathcal{A}_{f^*(\theta)}) \xrightarrow{\mathrm{Exp}_{\mathcal{A}_{f^*(\theta)}}} H^1(Y, \mathcal{A}_{f^*(\theta)}^\times),$$

and by $(L_{f, \theta}^{\mathrm{Exp}}, \theta_f)$ the Higgs bundle over Y associated to $\mathcal{L}_{f, \theta}^{\mathrm{Exp}}$.

Proposition A.2.1. (i) *The association $(M, \theta) \mapsto f^*(M, \theta) \otimes_{\mathcal{A}_{f^*(\theta)}} \mathcal{L}_{f, \theta}^{\mathrm{Exp}}$ defines a functor (2.4.1)*

$$(A.2.1.1) \quad f_{\mathfrak{Y}, \mathfrak{X}, \mathrm{Exp}}^\circ : \mathrm{HB}(X) \rightarrow \mathrm{HB}(Y, \xi^{-1} f^*(\Omega_X^1)),$$

also denoted by f° if there is no confusion. Via the composition $\mathrm{HB}(Y, \xi^{-1} f^(\Omega_X^1)) \rightarrow \mathrm{HB}(Y)$ induced by $f^*(\Omega_X^1) \rightarrow \Omega_Y^1$, we obtain the twisted inverse image functor (A.0.0.3).*

(ii) *The functor f° (resp. functor (A.0.0.3)) is exact and commutes with duals.*

(iii) *Given a Higgs bundle (M, θ) over X , the degrees of $f^*(M, \theta)$ and of $f^\circ(M, \theta)$ are the same.*

(iv) Let $g : Z \rightarrow Y$ be a finite étale morphism of smooth proper \mathbf{C} -curves and \mathfrak{Z} a smooth lifting of Z to B_{dR}^+/ξ^2 . Then there exists a canonical isomorphism of functors (A.0.0.3):

$$(f \circ g)_{\mathfrak{Z}, \mathfrak{X}, \text{Exp}}^{\circ} \simeq g_{\mathfrak{Z}, \mathfrak{Y}, \text{Exp}}^{\circ} \circ f_{\mathfrak{Y}, \mathfrak{X}, \text{Exp}}^{\circ}.$$

A.2.2. We first investigate the duality of twisted inverse image. Let (M, θ) be a Higgs bundle over X , $(N, \vartheta) = (f^*(M), f^*(\theta))$ and \mathcal{N} the invertible \mathcal{A}_{ϑ} associated to (N, ϑ) .

The canonical morphism defined, for any local sections ε of N^{\vee} , m of N and ∂ of T_X , by

$$(A.2.2.1) \quad \begin{aligned} N^{\vee} = \mathcal{H}om_{\mathcal{O}_Y}(N, \mathcal{O}_Y) &\rightarrow \mathcal{N}^{\vee} = \mathcal{H}om_{\mathcal{A}_{\vartheta}}(\mathcal{N}, \mathcal{A}_{\vartheta}), \\ \varepsilon &\mapsto (m \mapsto (\varepsilon(m), \varepsilon(\vartheta_{f^* \partial}(m)) \otimes \xi f^*(\partial), \dots,)) \end{aligned}$$

is an isomorphism of vector bundles over Y . Let $\vartheta^{\vee} : N^{\vee} \rightarrow N^{\vee} \otimes \xi^{-1} f^*(\Omega_X^1)$ be the morphism defined by $\vartheta^{\vee}(\varepsilon) = -\varepsilon(\vartheta(m))$. If we take a basis of N locally on Y , then the matrix representation of $-\vartheta^{\vee}$ with respect to this basis is the transpose of that of ϑ . Hence ϑ and $-\vartheta^{\vee}$ have the same characteristic polynomial, and we obtain an isomorphism $\mathcal{A}_{\vartheta} \simeq \mathcal{A}_{-\vartheta^{\vee}}$. Equipped with $-\vartheta^{\vee}$ on the left hand side, the above isomorphism is \mathcal{A}_{ϑ} -linear.

We claim that there exists a canonical isomorphism of invertible \mathcal{A}_{ϑ} -modules:

$$(A.2.2.2) \quad (\mathcal{L}_{f, \theta}^{\text{Exp}})^{\vee} \simeq (L_{f, \theta^{\vee}}^{\text{Exp}}, -\theta_f^{\vee}).$$

Indeed, the isomorphism $T_X \rightarrow T_X, \partial \mapsto -\partial$ induces isomorphisms:

$$\mathcal{A}_{\vartheta^{\vee}} \xrightarrow{\sim} \mathcal{A}_{-\vartheta^{\vee}}, \quad \iota : H^1(Y, \mathcal{A}_{\vartheta^{\vee}}^{\times}) \xrightarrow{\sim} H^1(Y, \mathcal{A}_{-\vartheta^{\vee}}^{\times}),$$

where the latter isomorphism sends a Higgs bundle (N, φ) to $(N, -\varphi)$. The left hand side of (A.2.2.2) is identified with the $\text{Exp}(-o_f)$ in $H^1(Y, \mathcal{A}_{-\vartheta^{\vee}}^{\times})$ via $H^1(Y, \mathcal{A}_{-\vartheta^{\vee}}^{\times}) \simeq H^1(Y, \mathcal{A}_{\vartheta}^{\times})$. Then it corresponds to $\iota(\text{Exp}(o_f))$ via ι . Hence, we obtain the right hand side of (A.2.2.2) by (A.2.2.1) and the claim follows.

By (A.2.2.1) and (A.2.2.2), we obtain the following canonical isomorphism between duals:

$$\begin{aligned} f^{\circ}(M^{\vee}, \theta^{\vee}) = (N^{\vee}, \vartheta^{\vee}) \otimes_{\mathcal{A}_{\vartheta^{\vee}}} \mathcal{L}_{f, \theta^{\vee}}^{\text{Exp}} &\simeq \iota \left((N^{\vee}, -\vartheta^{\vee}) \otimes_{\mathcal{A}_{-\vartheta^{\vee}}} (L_{f, \theta^{\vee}}^{\text{Exp}}, -\theta_f^{\vee}) \right) \\ &\simeq \iota \left((\mathcal{N} \otimes_{\mathcal{A}_{\vartheta}} \mathcal{L}_{f, \theta}^{\text{Exp}})^{\vee} \right) = \left(f^{\circ}(M, \theta) \right)^{\vee}. \end{aligned}$$

Hence, the twisted inverse image is compatible with the dual of Higgs bundles.

A.2.3. Let $0 \rightarrow (M'', \theta'') \rightarrow (M, \theta) \rightarrow (M', \theta') \rightarrow 0$ be a short exact sequence of Higgs bundles over X . Since θ and θ' are compatible, the action of \mathcal{A}_{θ} on M induces an action on M' , which is compatible with that of $\text{Sym}_{\mathcal{O}_X}(\xi T_X)$ via θ' . Then we obtain a surjective homomorphism: $\mathcal{A}_{\theta} \twoheadrightarrow \mathcal{A}_{\theta'}$, whose kernel is denoted by I . Via this quotient, we have a canonical isomorphism of Higgs modules:

$$(A.2.3.1) \quad (M, \theta) \otimes_{\mathcal{A}_{\theta}} \mathcal{A}_{\theta'} \xrightarrow{\sim} (M', \theta').$$

On the other hand, by taking dual, we obtain a surjective homomorphism $\mathcal{A}_{\theta} \rightarrow \mathcal{A}_{\theta''}$. We claim that the composition:

$$(A.2.3.2) \quad I \hookrightarrow \mathcal{A}_{\theta} \twoheadrightarrow \mathcal{A}_{\theta''}$$

is an isomorphism of \mathcal{A}_{θ} -modules. To prove it, we may work locally on X and take a base of T_X over \mathcal{O}_X . If $F_{\theta}(x) = \sum_{i=0}^r c_i(\theta) x^{r-i}$ denotes the characteristic polynomial of θ , we have a local isomorphism $\mathcal{A}_{\theta} \simeq \mathcal{O}_X[x]/(F_{\theta}(x))$. Since we have a decomposition $F_{\theta}(x) = F_{\theta'}(x)F_{\theta''}(x)$, then the claim follows.

Let $0 \rightarrow (N'', \vartheta'') \rightarrow (N, \vartheta) \rightarrow (N', \vartheta') \rightarrow 0$ be the inverse image of the above sequence to Y via f^* . In view of the construction, we have canonical isomorphisms of $\mathcal{A}_{\theta'}$ -modules (resp. $\mathcal{A}_{\theta''}$ -modules):

$$\mathcal{L}_{f, \theta}^{\text{Exp}} \otimes_{\mathcal{A}_{\theta}} \mathcal{A}_{\theta'} \simeq \mathcal{L}_{f, \theta'}^{\text{Exp}}, \quad \mathcal{L}_{f, \theta}^{\text{Exp}} \otimes_{\mathcal{A}_{\theta}} \mathcal{A}_{\theta''} \simeq \mathcal{L}_{f, \theta''}^{\text{Exp}}.$$

By applying $f^\circ(M, \theta) \otimes_{\mathcal{A}_\theta} -$ to the short exact sequence $0 \rightarrow I \rightarrow \mathcal{A}_\theta \rightarrow \mathcal{A}_{\theta'} \rightarrow 0$, we deduce the following short exact sequence:

$$0 \rightarrow f^\circ(M'', \theta'') \rightarrow f^\circ(M, \theta) \rightarrow f^\circ(M', \theta') \rightarrow 0.$$

In this way, we obtain the construction of $f^\circ(u)$ for a morphism u of $\text{HB}(X)$, whose cokernel is still a Higgs bundle. We also show that f° is exact.

If $u : (M_1, \theta_1) \rightarrow (M_2, \theta_2)$ is a monomorphism of Higgs bundles whose quotient is a torsion \mathcal{O}_X -modules. A simple calculation shows that the characteristic polynomials of θ_1 and of θ_2 coincide. Since f is flat, this observation allows us to define a monomorphism $f^\circ(u) : f^\circ(M_1, \theta_1) \rightarrow f^\circ(M_2, \theta_2)$.

Since $\dim X = 1$, a coherent \mathcal{O}_X -module can be decomposed as a direct sum of a vector bundle over X and a torsion \mathcal{O}_X -module. From previous constructions, we establish the functoriality of f° .

Remark A.2.4. For a torsion quasi-coherent \mathcal{O}_X -module M together with a Higgs field θ , we can canonically define the twisted inverse image $f^\circ(M, \theta)$ by its usual inverse image $f^*(M, \theta)$. Together with the above construction, we can slightly extend f° to the category of coherent \mathcal{O}_X -modules with a Higgs field.

A.2.5. *Proof of proposition A.2.1(iii).* Let Bun_n be the moduli stack of vector bundles of rank n on X , $T^* \text{Bun}_n$ the moduli stack of Higgs bundles of rank n , $\mathcal{P}\text{ic}(\mathcal{A}_\theta)$ the Picard stack of $\text{Spec}_{\mathcal{O}_X}(\mathcal{A}_\theta)$. Let \mathcal{P} be the rigidified Picard functor for a suitable rigidification ([26] §2.1), which is representable by a smooth group scheme over \mathbf{C} . We have a quotient $r : \mathcal{P} \rightarrow \mathcal{P}$, whose kernel is connected ([26] proposition 2.4.3) and a canonical morphism $\mathcal{P} \rightarrow \mathcal{P}\text{ic}(\mathcal{A}_\theta)$ sending (\mathcal{L}, φ) to \mathcal{L} .

Let (M, θ) be a Higgs bundle of rank r on X . We show that $f^\circ(M, \theta)$ has the same degree as $f^*(M, \theta)$. Let $c = (c_i(\theta))_{i=1}^r \in B = \oplus_{i=1}^r \Gamma(X, \Omega_X^{\otimes i})$ be the image of (M, θ) under the Hitchin map. We consider the following diagram

$$\begin{array}{ccccccc} \mathcal{P} & \longleftarrow & \mathcal{P} & \longrightarrow & \mathcal{P}\text{ic}(\mathcal{A}_\theta) & \longrightarrow & T^* \text{Bun}_n \longrightarrow \text{Bun}_n \\ & & & & \downarrow & & \downarrow h \\ & & & & c & \longrightarrow & B \end{array}$$

where the right vertical arrow is the Hitchin map and the square is a fibered product. The composition $\mathcal{P} \rightarrow \text{Bun}_n$ sends a pair (\mathcal{M}, α) of an invertible \mathcal{A}_θ -module \mathcal{M} together with a rigidification α to the vector bundle $\pi_{\theta,*}(\mathcal{M})$, where $\pi_\theta : X_\theta \rightarrow X$ denotes the spectral cover.

By construction, $\mathcal{L}_{f,\theta}^{\text{Exp}}$ lies in the connected component of \mathcal{P} . Then we deduce that $f^*(M, \theta)$ and $f^\circ(M, \theta)$ (resp. their liftings to \mathcal{P}) lie in the same connected component of \mathcal{P} (resp. \mathcal{P}). Hence their underlying vector bundles lie in the same connected component of Bun_n , i.e. they have the same degree. \square

A.2.6. *Proof of proposition A.2.1(iv).* Let $o_{f \circ g}, o_g$ and o_f be the obstruction of lifting $f \circ g, g$ and f to B_{dR}^+/ξ^2 . Then we have:

$$o_{f \circ g} = o_g + g^*(o_f) \in H^1(Z, \xi(f \circ g)^*(T_X)).$$

Let (M, θ) be a Higgs bundle over X . By the functoriality of Exp_G , we deduce from above identity a canonical isomorphism of invertible $\mathcal{A}_{(f \circ g)^*(\theta)}$ -modules

$$(A.2.6.1) \quad \mathcal{L}_{f \circ g, \theta}^{\text{Exp}} \simeq \mathcal{L}_{g, f^*(\theta)}^{\text{Exp}} \otimes g^*(\mathcal{L}_{f, \theta}^{\text{Exp}}).$$

Then the assertion follows. \square

A.3. Comparison with inverse image of Higgs crystals. Let K' be a finite extension of K , Y a projective S' -curve with semi-stable reduction, $f : Y \rightarrow X_{S'}$ a generic η' -cover (1.2.2) and $\check{f} : \check{Y} \rightarrow \check{X}$ the associated morphism of p -adic fine log formal schemes. We fix a lifting \mathcal{X} (resp. \mathcal{Y}) of \check{X} over Σ_S (resp. \check{Y} over $\Sigma_{S'}$). The obstruction of lifting \check{f} to Σ_S defines an element o_f of $H^1(\check{Y}, \xi \check{f}^*(T_{\check{X}}))$.

As in (A.2.0.1), we denote by $\mathcal{L}_{f, \theta}^{\text{Exp}}$ the image of o_f in the following composition:

$$(A.3.0.1) \quad H^1(\check{Y}, \xi \check{f}^*(T_{\check{X}})) \rightarrow H^1(Y_{\mathbf{C}}, \xi f_{\mathbf{C}}^*(T_{X_{\mathbf{C}}})) \rightarrow H^1(Y_{\mathbf{C}}, \mathcal{A}_{f^*(\theta)}) \xrightarrow{\text{Exp}_{\mathcal{A}_{f^*(\theta)}}} H^1(Y_{\mathbf{C}}, \mathcal{A}_{f^*(\theta)}^\times).$$

Via $f_{\mathbf{C}}^*(\Omega_{X_{\mathbf{C}}}^1) \rightarrow \Omega_{Y_{\mathbf{C}}}^1$, we can define the twisted inverse image functor

$$(A.3.0.2) \quad f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^\circ : \text{HB}(X_{\mathbf{C}}) \rightarrow \text{HB}(Y_{\mathbf{C}}), \quad (M, \theta) \mapsto f_{\mathbf{C}}^*(M, \theta) \otimes_{\mathcal{A}_{f^*(\theta)}} \mathcal{L}_{f, \theta}^{\text{Exp}}.$$

One can verify proposition A.2.1 for $f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^\circ$ in a similar way. When the liftings \mathcal{X}_2 and \mathcal{Y}_2 are algebraic, the above functor is clearly compatible with functor (A.0.0.3) defined by $\mathcal{X}_2 \otimes_{\text{A}_{\text{inf}}} \text{B}_{\text{dR}}^+$, $\mathcal{Y}_2 \otimes_{\text{A}_{\text{inf}}} \text{B}_{\text{dR}}^+$.

Proposition A.3.1. *The functor $f_{\mathcal{Y}, \mathcal{X}, \text{Exp}}^\circ$ is compatible with the functor $f_{\mathcal{Y}, \mathcal{X}}^\circ$ (2.4.3), when it restricts to the small Higgs bundles.*

A.3.2. To prove above proposition, we need to study the exponential maps on rigidified Picard schemes.

Let \mathcal{B} be a finite locally free $\mathcal{O}_{X_{\mathbf{C}}}$ -algebra. We denote by \mathcal{P} the relative Picard functor of $\text{Spec}_{\mathcal{O}_{X_{\mathbf{C}}}}(\mathcal{B})$ over \mathbf{C} , which is represented by a commutative smooth algebraic group over \mathbf{C} ([26] corollaire 1.5.2, 2.3.2). We denote the exponential map for \mathcal{P} (A.1.2.1) by

$$(A.3.2.1) \quad \text{Exp}_{\mathcal{B}} : \text{Lie } \mathcal{P} \simeq H^1(X_{\mathbf{C}}, \mathcal{B}) \rightarrow \mathcal{P}(\mathbf{C}) = H^1(X_{\mathbf{C}}, \mathcal{B}^\times).$$

On the other hand, let \mathcal{B} be a finite locally free \mathcal{O}_{X_\circ} -algebra and $\widehat{\mathcal{B}}$ the p -adic completion of \mathcal{B} over \check{X} . For $\alpha \in \mathbb{Q}_{> \frac{1}{p-1}}$, we have the following exponential map

$$(A.3.2.2) \quad \exp_{\mathcal{B}} : p^\alpha H^1(\check{X}, \widehat{\mathcal{B}}) \rightarrow H^1(\check{X}, \widehat{\mathcal{B}}^\times), \quad \varphi = (\varphi_{\lambda, \mu}) \mapsto \exp(\varphi) = (\exp(\varphi_{\lambda, \mu})),$$

which is well-defined due to $\alpha > \frac{1}{p-1}$.

We use a similar construction of $\exp_{\mathcal{B}}$ to define the twisted inverse image functor for small Higgs bundles over a curve in § 2.4. We will generalize this functor for all Higgs bundles over a curve via $\text{Exp}_{\mathcal{B}_{\mathbf{C}}}$ next subsection. To compare two constructions, we need the following result.

Proposition A.3.3. *The exponential maps $\text{Exp}_{\mathcal{B}_{\mathbf{C}}}$ and $\exp_{\mathcal{B}}$ are compatible.*

A.3.4. *Proof of proposition A.3.1.* Let (M, θ) be a small Higgs bundle over $X_{\mathbf{C}}$. We take again notations of § 2.4. The left square is clearly commutative and the commutativity of the right square follows from proposition A.3.3:

$$\begin{array}{ccccc} H^1(\check{Y}, \check{f}^*(T_{\check{X}})) & \longrightarrow & p^\alpha H^1(\check{Y}, \check{f}^*(\widehat{\mathcal{A}}_\theta)) & \xrightarrow{\exp} & H^1(\check{Y}, \check{f}^*(\widehat{\mathcal{A}}_\theta)) \\ \downarrow & & \downarrow & & \downarrow \\ H^1(Y, f^*(T_{X_{\mathbf{C}}})) & \hookrightarrow & H^1(Y_{\mathbf{C}}, \mathcal{A}_{f^*(\theta)}^\times) & \xrightarrow{\text{Exp}} & H^1(Y, \mathcal{A}_{f^*(\theta)}^\times). \end{array}$$

Hence the invertible $\check{f}^*(\widehat{\mathcal{A}}_\theta)$ -module $\mathcal{L}_{f, \theta}$ (§ 2.4.3) is sent to $\mathcal{L}_{f, \theta}^{\text{Exp}}$ via the right vertical arrow. Then the assertion follows. \square

A.3.5. We prove above proposition in the following. To do it, we need to investigate the integral structure of the relative Picard functor \mathcal{P} , whose representability exists after taking certain rigidification.

Let $i : D = \sqcup \text{Spec}(\mathfrak{o}) \rightarrow X_\circ$ be a divisor such that the following maps are injective:

$$\Gamma(X_{\mathbf{C}}, \mathcal{B}_{\mathbf{C}}) \rightarrow \Gamma(D_{\mathbf{C}}, i^*(\mathcal{B}_{\mathbf{C}})), \quad \Gamma(X_s, \mathcal{B}_s) \rightarrow \Gamma(D_s, i^*(\mathcal{B}_s)).$$

Then (i, D) is a rigidification of the relative Picard functor $\text{Pic}_{\mathcal{B}/\mathfrak{o}}$ ([26] corollaire 2.2.2). Such a rigidification for \mathcal{B} always exists ([26] proposition 2.2.3).

The rigidified Picard functor \mathcal{P} ([26] §2.1), associated to D and \mathcal{B} , is defined, for any \mathfrak{o} -scheme T by

$$(A.3.5.1) \quad \mathcal{P}(T) = \{(\mathcal{L}, \alpha) \mid \mathcal{L}, \mathcal{B}_T\text{-line bundle, } \alpha : i_T^* \mathcal{L} \xrightarrow{\sim} i_T^* \mathcal{B}_T\} / \text{isom}.$$

Note that any automorphism of an object of $\mathcal{P}(T)$ is the identity.

The functor \mathcal{P} is represented by a smooth algebraic space locally of finite presentation over \mathfrak{o} ([26] théorème 2.3.1 et corollaire 2.3.2).

Let T be an affine \mathfrak{o} -scheme. We set $I_T = \text{Ker}(\mathcal{B}_T \rightarrow i_{T,*} i_T^*(\mathcal{B}_T))$, $J_T = \text{Ker}(\mathcal{B}_T^* \rightarrow i_{T,*} i_T^*(\mathcal{B}_T^*))$. By considering the long exact sequence of $0 \rightarrow I_T \rightarrow \mathcal{B}_T \rightarrow i_{T,*} i_T^*(\mathcal{B}_T) \rightarrow 0$ and ([26] corollaire 2.2.2), we deduce $H^0(X_T, I_T) = 0$, and short exact sequences:

$$(A.3.5.2) \quad 0 \rightarrow H^0(T, i_T^* \mathcal{B}_T) / H^0(X_T, \mathcal{B}_T) \rightarrow H^1(X_T, I_T) \rightarrow H^1(X_T, \mathcal{B}_T) \rightarrow 0,$$

$$(A.3.5.3) \quad 0 \rightarrow H^0(T, i_T^* \mathcal{B}_T^*) / H^0(X_T, \mathcal{B}_T^*) \rightarrow H^1(X_T, J_T) \rightarrow H^1(X_T, \mathcal{B}_T^*) \rightarrow 0.$$

By ([26] proposition 2.4.1), we deduce the following isomorphism:

$$H^1(X_T, J_T) \simeq \mathcal{P}(T), \quad H^1(X_T, I_T) \simeq \text{Ker}(\mathcal{P}(T \otimes_{\mathfrak{o}} \mathfrak{o}[\varepsilon]/(\varepsilon^2)) \rightarrow \mathcal{P}(T)).$$

We denote by $\text{Lie } \mathcal{P}$ the Lie algebra of \mathcal{P} , defined by $\text{Lie } \mathcal{P}(T) = \text{Ker}(\mathcal{P}(T \otimes_{\mathfrak{o}} \mathfrak{o}[\varepsilon]/(\varepsilon^2)) \rightarrow \mathcal{P}(T))$.

A.3.6. The functors $\mathcal{P}_{\mathbf{C}}$ is also represented by a commutative smooth algebraic group over \mathbf{C} , the quotient $r : \mathcal{P}_{\mathbf{C}} \rightarrow \mathcal{P}$ is smooth and surjective ([26] proposition 2.4.3). Its kernel is a connected smooth affine group scheme over \mathbf{C} and is isomorphic to the functor defined, for \mathbf{C} -scheme T , by

$$T \mapsto H^0(T, (i_T^* \mathcal{B}_T)^{\times}) / H^0(X_T, (\mathcal{B}_T)^{\times}).$$

Its Lie algebra $\text{Lie Ker}(r)$ is isomorphic to $\Gamma(D_{\mathbf{C}}, i_{\mathbf{C}}^*(\mathcal{B}_{\mathbf{C}})) / \Gamma(X_{\mathbf{C}}, \mathcal{B}_{\mathbf{C}})$ and the inclusion $\text{Lie Ker}(r) \rightarrow \text{Lie } \mathcal{P}_{\mathbf{C}}$ coincides with the first map of (A.3.5.2) with $T = \text{Spec}(\mathbf{C})$.

A.3.7. Let \mathcal{E} be the scheme-theoretic closure of the unit section of $\mathcal{P}_{\mathbf{C}}$ in \mathcal{P} ([26] §3.2 c)). Then \mathcal{E} is a sheaf of groups (for the fppf topology) and the quotient $\mathcal{G} = \mathcal{P} / \mathcal{E}$ is represented by a smooth group scheme over \mathfrak{o} ([26] proposition 3.3.5).

Since $\mathcal{E}^0 \simeq 0$ ([26] proposition 5.2), we have $\text{Lie } \mathcal{E} = 0$. Then we deduce an isomorphism $\text{Lie } \mathcal{P} \xrightarrow{\sim} \text{Lie } \mathcal{G}$, which is isomorphic to an affine space over \mathfrak{o} , equipped with the additive group structure.

Let α be a positive rational number. With the notation of § A.1.4, the rigid space $\text{Lie } \mathcal{G}_{p^{-\alpha}}^{\text{rig}}$ admits a formal model defined as below. Let \mathcal{J} be the ideal sheaf of $\text{Lie } \mathcal{G}$ defined by the unit section over \mathfrak{o} and $\text{Lie } \mathcal{G}^{\alpha} = \text{Spec}_{\mathcal{O}_{\text{Lie } \mathcal{G}}}(\mathcal{O}_{\text{Lie } \mathcal{G}}[\frac{\mathcal{J}}{p^{\alpha}}])$ the dilatation of $\text{Lie } \mathcal{G}$ of $(\mathcal{J}, p^{\alpha})$ with respect to p^{α} . Then the rigid generic fiber of the p -adic completion $\widehat{\text{Lie } \mathcal{G}^{\alpha}}$ of $\text{Lie } \mathcal{G}^{\alpha}$ is isomorphic to $\text{Lie } \mathcal{G}_{p^{-\alpha}}^{\text{rig}}$.

Let n be an integer ≥ 1 and A an \mathfrak{o}_n -algebra, $\text{Lie } \mathcal{G}(A)$ is an \mathfrak{o} -module. The image of $\text{Lie } \mathcal{G}^{\alpha}(A) \rightarrow \text{Lie } \mathcal{G}(A)$ is contained in $p^{\alpha} \text{Lie } \mathcal{G}(A)$. This allows us to define a homomorphism by the exponential

$$(A.3.7.1) \quad \begin{aligned} \exp^{\alpha} : \text{Lie } \mathcal{G}^{\alpha}(A) &\rightarrow p^{\alpha} \text{Lie } \mathcal{G}(A) \simeq p^{\alpha} H^1(X_A, I_A) \rightarrow \mathcal{P}(A) \simeq H^1(X_A, I_A^{\times}), \\ \varphi = (\varphi_{\lambda, \mu}) &\mapsto \exp(\varphi) = (\exp(\varphi_{\lambda, \mu})). \end{aligned}$$

Note that the above exponential is actually a finite sum due to $\alpha > \frac{1}{p-1}$ and is therefore well-defined.

Together with the quotient $\mathcal{P} \rightarrow \mathcal{G}$, the above morphism induces a homomorphism of group \mathfrak{o}_n -schemes:

$$\exp_n^{\alpha} : (\text{Lie } \mathcal{G}^{\alpha})_n \rightarrow \mathcal{G}_n.$$

In view of (A.3.7.1), for $\beta = \alpha - \frac{1}{p-1} > 0$, the reduction modulo p^{β} of the above morphism factors through the unit section of \mathcal{G}_{β} . Let \mathcal{J} be the ideal sheaf of \mathcal{G} generated by p^{β} and the ideal sheaf of the unit section of \mathcal{G} over \mathfrak{o} . Then there exists an integer k_n such that the above morphism factors through:

$$\exp_n^{\alpha} : (\text{Lie } \mathcal{G}^{\alpha})_n \rightarrow \text{Spec}(\mathcal{O}_{\mathcal{G}} / \mathcal{J}^{k_n}).$$

With the notation of A.1.4, we have $\mathcal{G} \simeq \text{Spf}(\varprojlim_n \mathcal{O}_{\mathcal{G}} / \mathcal{J}^{k_n})$. Then the above homomorphisms induce a homomorphism of formal group schemes and moreover of rigid-analytic groups:

$$(A.3.7.2) \quad \widehat{\exp^{\alpha}} : \widehat{\text{Lie } \mathcal{G}^{\alpha}} \rightarrow \mathcal{G}, \quad \widehat{\exp^{\alpha}}^{\text{rig}} : \text{Lie } \mathcal{G}_{p^{-\alpha}}^{\text{rig}} \rightarrow \mathcal{G}^{\text{rig}}.$$

Lemma A.3.8. For $\alpha \in \mathbb{Q}_{> \frac{1}{p-1}}$, the homomorphism $|\widehat{\exp^{\alpha}}^{\text{rig}}| : p^{\alpha} \text{Lie } \mathcal{G} \rightarrow |\mathcal{G}|$ on the underlying points is compatible with $|\exp_{\mathcal{G}}|$ (A.1.4.2) and $\text{Exp}_{\mathcal{G}_{\mathbf{C}}}$ (A.1.2.1) as well.

Proof. It suffices to prove the assertion for $|\exp_{\mathcal{G}}|$. We consider $|\widehat{\exp^{\alpha}}^{\text{rig}}|$ as a homomorphism of p -adic Lie groups. In view of the construction (A.3.7.1), we deduce that its differential is the identity $\text{id}_{\text{Lie } \mathcal{G}_C}$. By ([27] proposition 18.17), we deduce that $|\widehat{\exp^{\alpha}}^{\text{rig}}|$ coincides with the exponential map $|\exp_{\mathcal{G}}|$ on an open neighborhood U of 0. For any element x of $p^{\alpha} \text{Lie } \mathcal{G}$, there exists an integer n such that $p^n x \in U$. We have

$$p^n \log_G(\widehat{\exp^{\alpha}}^{\text{rig}}(x)) = \log_G(\widehat{\exp^{\alpha}}^{\text{rig}}(p^n x)) = \log_G(\exp_{\mathcal{G}}(p^n x)) = p^n x.$$

Since $\times p^n$ is injective on $\text{Lie } \mathcal{G}$, the assertion follows from the isomorphism (A.1.4.1). \square

A.3.9. We denote by \mathcal{P}' (resp. \mathcal{G}') the subfunctor of \mathcal{P} (resp. \mathcal{G}) such that $\mathcal{P}'(R) = \{s \in \mathcal{P}(R) \mid s_k = e \in \mathcal{P}(R \otimes_{\circ} k)\}$ (resp. $\mathcal{G}'(R) = \{s \in \mathcal{G}(R) \mid s_k = e \in \mathcal{G}(R \otimes_{\circ} k)\}$), where e denotes the unit section. Since $\mathcal{E}(\mathfrak{o}) \simeq \mathcal{E}(k)$ ([26] théorème 6.4.1 (2)) and $\mathcal{G} = \mathcal{P}/\mathcal{E}$ for the étale topology, the canonical morphism $\mathcal{P}'(\mathfrak{o}) \rightarrow \mathcal{G}'(\mathfrak{o})$ is an isomorphism.

Let \hat{I} (resp. \hat{J}) be the p -adic completion of I (resp. \hat{J}). We have an inclusion $\varprojlim_n \mathcal{P}'(\mathfrak{o}_n) \subset H^1(\check{X}, \hat{J}) \simeq \varprojlim_n \mathcal{P}(\mathfrak{o}_n)$. In view of the algebraicity of formal line bundles, we have a canonical isomorphism $\mathcal{P}'(\mathfrak{o}) \simeq \varprojlim_n \mathcal{P}'(\mathfrak{o}_n)$. As subsets of $\mathcal{G}(\mathfrak{o})$, we have inclusion $\mathcal{G}(\mathfrak{o}) \subset \mathcal{G}'(\mathfrak{o}) \simeq \mathcal{P}'(\mathfrak{o})$.

By lemma A.3.8 and taking projective limit of (A.3.7.1), we deduce that the homomorphism $|\widehat{\exp^{\alpha}}^{\text{rig}}|$ (A.3.7.2) (and therefore $\text{Exp}_{\mathcal{P}_C} = \text{Exp}_{\mathcal{G}_C}$) is compatible with the composition

$$(A.3.9.1) \quad p^{\alpha} H^1(\check{X}, \hat{I}) \rightarrow \mathcal{G}(\mathfrak{o}) \subset H^1(\check{X}, \hat{J}), \quad \varphi = (\varphi_{\lambda, \mu}) \mapsto \exp(\varphi) = (\exp(\varphi_{\lambda, \mu})).$$

Since Exp_G is functorial (proposition A.1.2), we conclude that $\text{Exp}_{\mathcal{B}_C}$ and $\exp_{\mathcal{B}}$ are compatible, and this finishes the proof of proposition A.3.3. \square

REFERENCES

- [1] A. Abbes, *Éléments de Géométrie Rigide: Volume I. Construction et étude géométrique des espaces rigides*. Springer-Verlag, (2010).
- [2] A. Abbes, M. Gros, *Les suites spectrales de Hodge-Tate*, preprint (2020), <https://arxiv.org/abs/2003.04714>.
- [3] A. Abbes, M. Gros, *Systèmes locaux de Hodge-Tate*, in preparation.
- [4] A. Abbes, M. Gros, T. Tsuji, *The p -adic Simpson correspondence*, Annals of Mathematics Studies, Vol. 193 (2016).
- [5] M. F. Atiyah, *Vector bundles over an elliptic curve*, Proc. London Math. Soc., Third series, 7 (1957), 414-452.
- [6] A. Beauville, M.S. Narasimhan, S. Ramanan, *Spectral curves and the generalised theta divisor*. Journal für die Reine und Angewandte Mathematik, 398 (1989), 169–179.
- [7] B. Bhatt, P. Scholze, *Prisms and prismatic cohomology*. <https://arxiv.org/abs/1905.08229>, 2019.
- [8] S. Bosch, W. Lütkebohmert and M. Raynaud, *Néron Models*, (Springer, Berlin, 1990).
- [9] R. Coleman, *Reciprocity laws on curves*, Comp. Math. 72 (1989) 205–235.
- [10] C. Deninger, A. Werner, *Vector bundles on p -adic curves and parallel transport*. Ann. Sci. Éc. Norm. Supér., Ser. 4, 38(4): 553–597, 2005.
- [11] C. Deninger, A. Werner, *Parallel transport for vector bundles on p -adic varieties*. J. Algebraic Geom., 29: 1–52, 2020.
- [12] H. Diao, K-W. Lan, R. Liu, and X. Zhu, *Logarithmic Riemann–Hilbert correspondences for rigid varieties*, arXiv:1803.05786.
- [13] G. Faltings, *A p -adic Simpson correspondence*, Adv. Math. 198 (2005), 847-862.
- [14] L. Fargues, *Groupes analytiques rigides p -divisibles*. Math. Ann. 374, 723–791 (2019).
- [15] E. Franco, O. Garcia-Prada, P. Newstead, *Higgs bundles over elliptic curves*, Illinois J. Math. Volume 58, Number 1 (2014), 43-96.
- [16] A. Grothendieck, J. Dieudonné, *Éléments de Géométrie Algébrique IV, Étude locale des schémas et des morphismes de schémas*, Pub. Math. IHES 20 (1964), 24 (1965), 28 (1966), 32 (1967).
- [17] T. He, *Cohomological descent for Faltings’ p -adic Hodge theory and applications*, preprint (2021), <https://arxiv.org/abs/2104.12645v2>.
- [18] B. Heuer, *Line bundles on rigid spaces in the v -topology*, preprint (2020), <https://arxiv.org/abs/2012.07918>.
- [19] B. Heuer, L. Mann, A. Werner, *The p -adic Corlette-Simpson correspondence for abeloids*, preprint (2021), <https://arxiv.org/abs/2107.09403>.
- [20] N. J. Hitchin, *Stable bundles and integrable systems*. Duke Mathematical Journal, 54 (1987), 91-114.
- [21] R. Liu, X. Zhu, *Rigidity and a Riemann-Hilbert correspondence for p -adic local systems*, Invent. Math., 207(1): 291–343, 2017.

- [22] L. Mann, A. Werner, *Local systems on diamonds and p -adic vector bundles*, preprint (2020), <https://arxiv.org/abs/2005.06855>.
- [23] M. S. Narasimhan and C. S. Seshadri, *Stable and unitary vector bundles on a compact Riemann surface*, Ann. of Math. (2) 82 (1965), 540–567.
- [24] A. Ogus, V. Vologodsky, *Nonabelian Hodge theory in characteristic p* , Publ. Math. IHÉS 106.1 (2007), 1–138.
- [25] H. Oyama, *PD Higgs crystals and Higgs cohomology in characteristic p* , J. Algebraic Geom. 26 (2017), no. 4, 735–802.
- [26] M. Raynaud, *Specialisation du foncteur de Picard*. Publ. Math. IHES 38, 27–76 (1970).
- [27] P. Schneider, *p -adic Lie groups*. Vol. 344. Springer Science and Business Media, 2011.
- [28] P. Scholze, *p -adic Hodge theory for rigid analytic varieties*. Forum Math. Pi, 1:e1, 2013.
- [29] P. Scholze and J. Weinstein, *Moduli of p -divisible groups*, Camb. J. Math. 1 (2013), no. 2, 145–237.
- [30] C. Simpson, *Higgs bundles and local systems*, Publ. Math. IHÉS, 75 (1992), 5–95.
- [31] J. T. Tate, *p -divisible groups*. In Proc. Conf. Local Fields (Driebergen, 1966), pp. 158–183. Springer, Berlin (1967)
- [32] J. Tong, *Application d’Albanese pour les courbes et contractions*; Math. Annalen 338 (2007), 405–420.
- [33] T. Tsuji, *Notes on the local p -adic Simpson correspondence*, Math. Ann. 371 (2018), no. 1–2, 795–881.
- [34] Y. Wang, *A p -adic Simpson correspondence for rigid analytic varieties*, preprint (2021), <https://arxiv.org/abs/2101.00263>.
- [35] M. Würthen, *Vector bundles with numerically flat reduction on rigid analytic varieties and p -adic local systems*, International Mathematics Research Notices, (2022), <https://doi.org/10.1093/imrn/rnab214>.
- [36] D. Xu, *Transport parallèle et correspondance de Simpson p -adique*, Forum Math. Sigma (2017), Vol. 5, e13, 127 pp.
- [37] D. Xu, *Lifting the Cartier transform of Ogus-Vologodsky modulo p^n* , Mém. Soc. Math. Fr. (N.S.) No. 163 (2019), vi+133 pp.
- [38] The Stacks project authors. The stacks project. <https://stacks.math.columbia.edu>, 2022.

DAXIN XU, MORNINGSIDES CENTER OF MATHEMATICS AND HUA LOO-KENG KEY LABORATORY OF MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, CHINESE ACADEMY OF SCIENCES, BEIJING 100190, CHINA;
 SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING, 100049, CHINA.
E-mail address: daxin.xu@amss.ac.cn
 TONGMU HE, INSTITUT DES HAUTES ETUDES SCIENTIFIQUES, 35 ROUTE DE CHARTRES, 91440 BURES-SUR-YVETTE, FRANCE
E-mail address: hetm15@ihes.fr