

MAGNETIC SLOWDOWN OF TOPOLOGICAL EDGE STATES

G. BAL, S. BECKER, AND A. DROUOT

ABSTRACT. We study the propagation of wavepackets along curved interfaces between topological, magnetic materials. Our Hamiltonian is a massive Dirac operator with a magnetic potential. We construct semiclassical wavepackets propagating along the curved interface as adiabatic modulations of straight edge states under constant magnetic fields. While in the magnetic-free case, the wavepackets propagate coherently at speed one, here they experience slowdown, dispersion, and Aharonov–Bohm effects. Several numerical simulations illustrate our results.

1. INTRODUCTION

This paper analyzes wavepackets propagating along an interface between two topologically distinct materials, in the presence of an external magnetic field. It extends constructions carried out in [BBD⁺21] for magnetic-free models. We represent here the electron dynamics via a two-dimensional Dirac equation:

$$(\varepsilon D_t + \mathcal{D})\Psi(t, x) = 0, \quad (t, x) \in \mathbb{R} \times \mathbb{R}^2. \quad (1.1)$$

In (1.1), \mathcal{D} denotes a Dirac operator with sign-changing mass and magnetic field:

$$\mathcal{D} = (\varepsilon D_1 - A_1(x))\sigma_1 + (\varepsilon D_2 - A_2(x))\sigma_2 + \kappa(x)\sigma_3, \quad \text{where:}$$

- $\varepsilon > 0$ is a small semiclassical parameter and $\varepsilon D_t = -i\varepsilon\partial_t$, $\varepsilon D_j = -i\varepsilon\partial_j$ denote the self-adjoint semiclassical derivatives;
- $A = (A_1, A_2)^t \in C^\infty(\mathbb{R}^2, \mathbb{R}^2)$ is a magnetic potential with $\nabla A \in C_b^\infty$ (i.e. smooth with all derivatives uniformly bounded), inducing the magnetic field $B = \partial_1 A_2 - \partial_2 A_1$;
- $\kappa \in C^\infty(\mathbb{R}^2, \mathbb{R})$ has varying sign and satisfies $\nabla \kappa \in C_b^\infty$;
- $\sigma_1, \sigma_2, \sigma_3$ are the standard 2×2 Pauli matrices.

The sign of the domain wall κ characterizes the topological phase of the material. Specifically, under the transversality condition

$$\kappa(y) = 0 \quad \Rightarrow \quad \nabla \kappa(y) \neq 0, \quad (1.2)$$

the interface $\Gamma = \kappa^{-1}(0)$ separates regions of distinct local topology [Bal19a, BBD⁺21]. For analytic reasons, we consider here a uniform version of (1.2):

$$\inf\{|\nabla \kappa(x)| : x \in \Gamma\} > 0.$$

The bulk-edge correspondence for Dirac operators [Bal19a] predicts that a tubular neighborhood of $\Gamma = \kappa^{-1}(0)$ supports asymmetric currents, hence (some analogue of) edge states for small ε . For vanishing magnetic potentials, we constructed in [BBD⁺21] long-lived solutions to (1.1). These were confined and propagating at speed one along Γ . We referred to

them as dynamical edge states, since we derived them as time-dependent adiabatic modulations of straight edge states. This paper extends the construction to the magnetic case. A non-zero magnetic field induces several new phenomena for dynamical edge states:

- (i) A systematic slowdown, see Figure 1;
- (ii) A large phase-shift, generating a Aharonov–Bohm effect when Γ is a loop;
- (iii) In general, a mesoscopic dispersion along Γ .

We set a few notations in §1.1, state a simplified main result in §1.2, and detail the effects (i)–(iii) in §1.3.




FIGURE 1. Snapshots of the evolution of Gaussian wavepackets propagating along a straight interface $\kappa(x) = x_2$, under a constant magnetic field $B = 0, 0.5, 1, 1.5, 2$ (from right to left), computed numerically. The packets slow down in stronger fields.

1.1. Notations. We first define the normal and tangent vector fields to the level sets of κ :

$$n(x) = \frac{\nabla \kappa(x)}{|\nabla \kappa(x)|}, \quad \tau(x) = Jn(x), \quad J = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

Given $y_0 \in \Gamma$, we let y_t be the solution of the ODE

$$\dot{y}_t = c(y_t)\tau(y_t), \quad c(y) := \frac{|\nabla \kappa(y)|}{\sqrt{|\nabla \kappa(y)|^2 + B(y)^2}}, \quad (1.3)$$

where $B = \nabla \times A$ is the magnetic field. Note that $\partial_t \kappa(y_t) = 0$ since $\nabla \kappa(x) \cdot \tau(x) = 0$, hence $y_t \in \Gamma$ for any $t \in \mathbb{R}$. We then define the quantities

$$B_t = B(y_t), \quad n_t = \frac{\nabla \kappa(y_t)}{|\nabla \kappa(y_t)|}, \quad r_t = |\nabla \kappa(y_t)|, \quad \rho_t = \sqrt{r_t^2 + B_t^2}, \quad \gamma_t = \frac{B_t}{\rho_t}.$$

To capture the local geometry of the interface near y_t , we define two smoothly varying angles φ_t and θ_t (and the corresponding clockwise rotation R_{θ_t}) by

$$\cos \varphi_t = \frac{r_t}{\rho_t} := c_t, \quad \sin \varphi_t = \frac{B_t}{\rho_t} := s_t, \quad R_{\theta_t} = \begin{bmatrix} \cos \theta_t & \sin \theta_t \\ -\sin \theta_t & \cos \theta_t \end{bmatrix}, \quad (1.4)$$

see Figure 2; we assume φ_0 and θ_0 belong to $[0, 2\pi)$ for concreteness. With these notations, $R_{\theta_t} n_t = e_2$ and $\dot{y}_t = c_t \tau_t$ with $\tau_t = Jn_t$. Differentiating the equation defining θ_t , we observe that $\dot{n}_t + \dot{\theta}_t \tau_t = 0$. Since $\dot{y}_t = c_t \tau_t$, we deduce that $\dot{\theta}_t = -c_t(\tau \cdot \partial_\tau n)(y_t)$. In particular, $|\dot{\theta}_t| = c_t K_t$ where K_t is the curvature of the curve Γ at y_t .

We also introduce the unitary pullback operator \mathcal{R}_{θ_t} by R_{θ_t} as $\mathcal{R}_{\theta_t} g(z) = g(R_{\theta_t} z)$. Associated to θ_t and φ_t are two spinorial rotations,

$$U_{2, \varphi_t} = e^{-i \frac{\varphi_t}{2} \sigma_2}, \quad U_{3, \theta_t} = e^{-i \frac{\theta_t}{2} \sigma_3}.$$




FIGURE 2. Geometry of curved interface.

1.2. Simplified main result. For the sake of simplicity, we consider a restricted setup: the domain wall satisfies a geometric condition and the magnetic field is constant. Our main result, 6.2, takes then a clearer form.

The aforementioned assumptions are:

- $\kappa \in C^\infty(\mathbb{R}^2, \mathbb{R})$ with $\nabla \kappa \in C_b^\infty$ satisfies

$$y \in \kappa^{-1}(0) \quad \Rightarrow \quad |\nabla \kappa(y)| = 1, \quad \Delta \kappa(y) = 0; \quad (1.5)$$

- The magnetic field B is constant.

While (1.5) is analytically restrictive, it is not geometrically restrictive: any one-dimensional submanifold of \mathbb{R}^2 is the zero set of a function κ satisfying (1.5). For instance, for the straight edge $\Gamma = \mathbb{R}e_2$, we can choose $\kappa(y) = y_2$ while for the unit circle $\Gamma = \mathbb{S}^1$ we can choose $\kappa(y) = \ln |y|$ (near $|y| = 1$). When the two above conditions hold, the quantities $r_t, B_t, \rho_t, \gamma_t, \varphi_t, c_t$ and s_t do not depend on t , and we omit the subscript t .

The most elementary setup with these two conditions consists of $\kappa(x) = x_2$ and $A = -Bx_2e_1$. The corresponding Dirac operator is:

$$\not{D}_{0,B} := (\varepsilon D_1 + Bx_2)\sigma_1 + \varepsilon D_2\sigma_2 + x_2\sigma_3. \quad (1.6)$$

We remark that the magnetic potential vanishes along the interface $\mathbb{R}e_1$ and is parallel to it. The equation $(\varepsilon D_t + \not{D}_{0,B})\Psi = 0$ admits an explicit family of non-dispersive wavepacket solutions that propagate along Γ at speed $c = (1 + B^2)^{-1/2}$:

$$\Psi_{0,B}(x) := \frac{1}{\sqrt{\varepsilon}} \psi_{0,B} \left(\frac{x - ct e_1}{\sqrt{\varepsilon}} \right), \quad \psi_{0,B}(z) := \int_{\mathbb{R}} e^{i\xi z_1 - \frac{1}{2}\rho(z_2 + \gamma\xi)^2} \hat{f}(\xi) d\xi \cdot U_{2,\varphi} \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad (1.7)$$

where \hat{f} is any Schwartz-class function, $\rho = \sqrt{1 + B^2}$, $\gamma = \frac{B}{1+B^2}$, and $\varphi = \arctan B$.

We extend this statement to interfaces tilted by an angle $\theta \in \mathbb{R}$. Let $\not{D}_{\theta,B}$ be the Dirac operator with domain wall and magnetic potential

$$\kappa(x) = n \cdot x, \quad A(x) = B\kappa(x)\tau, \quad n = \begin{bmatrix} -\sin \theta \\ \cos \theta \end{bmatrix}, \quad \tau = - \begin{bmatrix} \cos \theta \\ \sin \theta \end{bmatrix}.$$

The equation $(\varepsilon D_t + \mathcal{D}_{\theta,B})\Psi = 0$ is unitarily equivalent to the case $\theta = 0$ and admits a family of solutions constructed from $\Psi_{0,B}$:

$$\Psi_{\theta,B}(x) := \frac{1}{\sqrt{\varepsilon}} \psi_{\theta,B} \left(\frac{x - ct\tau}{\sqrt{\varepsilon}} \right), \quad \psi_{\theta,B}(z) := U_{3,\theta} \mathcal{R}_{\theta} \psi_{0,B}(z).$$

Our main result produces approximate solutions to $(\varepsilon D_t + \mathcal{D})\Psi = 0$ as modulations of $\psi_{\theta,B}$. To state it, we need the distribution g_t on \mathbb{R}^2 defined by:

$$g_t(z) := \int_{\mathbb{R}^2} e^{i\zeta z + i\gamma(\theta_t - \theta_0)\zeta_1^2} d\zeta = \frac{e^{-i\pi/4}}{|2\pi\gamma(\theta_t - \theta_0)|^{1/2}} e^{i\frac{-z_1^2}{4\gamma(\theta_t - \theta_0)}} \cdot \delta_0(z_2). \quad (1.8)$$

The second equality is valid for $\theta_t \neq \theta_0$ and should be replaced by a Dirac mass when $\theta_t = \theta_0$. Our simplified theorem for constant magnetic fields reads then as follows.

Theorem 1.1. *Assume that κ satisfies the condition (1.5) and that B is constant. For any $T > 0$, the equation $(\varepsilon D_t + \mathcal{D})\Psi = 0$ admits solutions that satisfy, uniformly for $t \in [-T, T]$ and $x \in \mathbb{R}^2$:*

$$\Psi(t, x) = \frac{e^{\frac{i}{\varepsilon}\chi(t,x)}}{\sqrt{\varepsilon}} \psi \left(t, \frac{x - y_t}{\sqrt{\varepsilon}} \right) + O_{L^2}(\varepsilon^{1/2}), \quad \text{where } \psi(t, z) := (\mathcal{R}_{\theta_t} g_t) * \psi_{\theta_t,B}(z), \quad (1.9)$$

$$\text{and } \chi(t, x) = \int_0^t \dot{y}_s \cdot A(y_s) ds + A(y_t) \cdot (x - y_t) + (x - y_t) \cdot (\nabla A(y_t)^\top - B n_t \tau_t^\top) (x - y_t).$$

The leading order term in (1.9):

- (i) Propagates at speed $c = (1 + B^2)^{-1/2}$ along Γ , in the prescribed direction τ .
- (ii) Is semiclassically localized at the phase-space point (y_t, A_t) with $A_t = A(y_t)$. This is where the eigenvalues of the symbol of \mathcal{D} are degenerate. While the rapid oscillations generated by A_t can be locally gauged away, they cannot be globally neglected when Γ is a closed loop: this is the Aharonov–Bohm effect.
- (iii) Disperses along Γ at rate prescribed by the difference $\theta_t - \theta_0$:

$$|\psi(t, z)| \leq \frac{C}{1 + |\theta_t - \theta_0|^{1/2}} \sup_{z \in \mathbb{R}^2} |\psi(0, z)|. \quad (1.10)$$

In particular, the leading part in (1.9) is controlled by $\varepsilon^{-1/2} |\theta_t - \theta_0|^{-1/2}$. This is a relatively weak dispersion as it comes from dispersion of the wave envelop $\mathbb{R}_{\theta_t} g_t \star \psi_{\theta_t,B}$ rather than dispersive relations of plane waves. It produces effects for times of order one, in contrast with dispersion in e.g. the semiclassical Schrödinger equation which arises at time ε .

We detail these three effects in §1.3. Theorem 6.2 will extend the result of Theorem 1.1 to cover varying magnetic fields, general domain walls and longer times of validity. The corresponding wavepackets will more generally have a variable speed, given by the ODE (1.3); a phase with properties identical to (ii) above; and a more complicated rate of dispersion.

1.3. Effects of the magnetic field. We comment here on the structure of the wavepacket (1.7), when B is constant and κ satisfies (1.5); and detail how things change when these conditions are relaxed (see Theorem 6.2).




FIGURE 3. Aharonov–Bohm effect for the domain wall $\kappa(y) = \ln |y|$ (corresponding to a circle interface). The plot shows the rapid evolution of the phase of the first spinor component as the wavepacket performs a single revolution around the circle. See §7.3 for details.

Turning on a magnetic field systematically slows down the propagation. In the general setup of Theorem 6.2, the wavepackets move at speed $(1 + B_t^2)^{-1/2}$, which is smaller than 1 whenever the magnetic field does not vanish.

The wavepacket is semiclassically localized at (y_t, A_t) . This point lies in the crossing set of the semiclassical symbol

$$\mathcal{D}(x, \xi) = (\xi_1 - A_1(x))\sigma_1 + (\xi_2 - A_2(x))\sigma_2 + \kappa(x)\sigma_3;$$

that is, the eigenvalues of $\mathcal{D}(y_t, A_t)$ are repeated. Hence, (y_t, A_t) is an exotic semiclassical trajectory, in the sense that it is not among those predicted by standard propagation of singularity, such as [DH72].

The semiclassical action $\dot{y}_t \cdot A_t$ generates a large phase-shift $e^{\frac{i}{\varepsilon} \int_0^t \dot{y}_s \cdot A_s ds}$ that can be locally – but not globally – gauged away. When Γ is a loop, after a full revolution the phase shift




FIGURE 4. Snapshots of the evolution of a wavepacket around the unit disc (with $\kappa(y) = \ln |y|$ and $\dot{\theta} = c$) in magnetic fields $B = 1/2, 1/\sqrt{2}, 1$ (top row) and $B = 2, 3, 4$ (bottom row). We observe gradual dispersion consistent with our predicted dispersion rate $\gamma|\theta_t - \theta_0| = \gamma ct = B(1 + B^2)^{-3/2}t$, see (1.11). For equal distance of propagation the dispersion is maximal when $B = 1$ while for equal time of propagation the dispersion is strongest when $B = 1/\sqrt{2}$.

relates to the magnetic flux $\Phi = \int_{\Gamma} A$: it is $e^{\frac{i}{\varepsilon}\Phi}$, see Figure 3 for the case of the circle. This is the Aharonov–Bohm effect.

The envelop of the wavepacket typically disperses along Γ . Under the geometric condition (1.5), the rate of change of dispersion is $|\theta_t - \theta_0|^{-1/2}$. Indeed, writing ψ in terms of g_t and $\psi_{0,B}$, and using the formula (1.8), we have for $|\theta_t - \theta_0| \geq 1$:

$$\begin{aligned} \sup_{z \in \mathbb{R}^2} |\psi(t, z)| &= \sup_{z \in \mathbb{R}^2} |g_t * \psi_{0,B}(z)| \leq \frac{1}{\sqrt{2\pi\gamma|\theta_t - \theta_0|}} \sup_{z_2 \in \mathbb{R}^2} \int_{\mathbb{R}} |\psi(y_1, z_2)| dy_1 \\ &\leq \frac{C}{\sqrt{\gamma|\theta_t - \theta_0|}} = \frac{C'}{\sqrt{\gamma|\theta_t - \theta_0|}} \sup_{z \in \mathbb{R}^2} |\psi(0, z)|. \end{aligned} \quad (1.11)$$

This yields (1.10). While the rate of change of dispersion $|\theta_t - \theta_0|^{-1/2}$ is bounded above if Γ is asymptotically flat, it can be as small as $t^{-1/2}$ when Γ is a loop or a spiral. In these cases, the wavepacket loses coherence over long times as displayed in Figure 4; it should be noted however that larger magnetic fields do not necessarily give rise to larger dispersion. In the general setup of Theorem 6.2, the rate of dispersion ν_t takes a more complicated form: see Lemma 5.3, (5.3) and (3.21). We comment that dispersion can have long-time effects. When the dispersion is strongest, $\nu_t \sim t$ and our construction holds up to times $T \ll \varepsilon^{-1/8}$; when it is weakest, $\nu_t = O(1)$ and we recover the time of validity $T \ll \varepsilon^{-1/2}$ of [BBD⁺21].

When B or $|\nabla\kappa|$ vary along Γ (that is, outside the setup of Theorem 1.1), an additional effect emerges: time-dependent anisotropic compression/stretching in the normal and tangent directions of Γ . We refer the reader to the formula (5.6) and Theorem 6.2. The compression factors, $\sqrt{\rho_t}$ (in the normal direction) and c_0^2/c_t^2 (in the tangent direction), remain bounded above and below in the limit $t \rightarrow \infty$. This contrasts with the parameter ν_t that controls the dispersion, which can grow like t .

1.4. Strategy of proof. Conceptually speaking, our approach consists in constructing successive transformations of \mathcal{D} that bring us closer to the flat Dirac operator $\mathcal{D}_{0,B}$ of (1.6). The main steps are as follows:

1. In §2, we conjugate \mathcal{D} by a gauge transform $e^{i\chi(t,x)/\varepsilon}$, cooked up so that the resulting magnetic potential $A - \nabla\chi$ vanishes along Γ and is tangent to the level sets of κ . These are shared features with the flat Dirac operator $\mathcal{D}_{0,B}$. The expansion of the gauge term $e^{i\chi(t,x)/\varepsilon}$ near y_t produces the large oscillatory phase of (1.9).
2. In §3.1 we look for solutions to the gauge-modified Dirac equation $(\varepsilon D_t + \tilde{\mathcal{D}})\tilde{\Psi} = 0$ as semiclassical wavepackets localized at $(y_t, 0)$. A formal Taylor expansion produces a hierarchy of equations for the envelops.
3. In §3.2, we perform a series of spatial and spinorial rotations on the leading equation. The first two rotations (which already appear in [BBD⁺21]) flatten the interface; the last is magnetically induced and is among the new ingredients. The result is the leading equation that one would get starting from $\mathcal{D}_{0,B}$.
4. Up to a partial shifted Fourier transform, the leading operator takes the same form as in the absence of magnetic fields [BBD⁺21]. In §4, we compute explicitly its kernel and prove stability estimates – later needed for the subleading transport equation.
5. In §5 we explicitly integrate the transport equation. In Fourier variables, the solutions have a quadratic phase with Hessian $\theta_t - \theta_0$ (in the setup of §1.2). In physical space, this transfers to dispersion at rate $|\theta_t - \theta_0|^{-1/2}$.

6. Higher-order approximations are constructed iteratively in §6. Combining them with the unitarity of $e^{-it\mathcal{D}}$ and a Duhamel argument, we obtain our main result, Theorem 6.2. It gives an approximate solution to $(\varepsilon D_t + \mathcal{D})\Psi = 0$ that propagates along Γ , slowed down by the magnetic field, and explicitly expressed through the above transformations.

We then illustrate our findings with a series of numerical simulations in §7.

1.5. Related literature. For systems of semiclassical PDEs, the symbols that govern the macroscopic transport are the eigenvalues of the (matrix-valued) symbol. A symbolic diagonalization argument shows that their Hamiltonian flow governs the leading-order dynamics. When the eigenvalues of the symbol are degenerate, the classical equations of motion break down. The situation studied here is among the simplest such cases. Our analysis shows that the phase-space crossing set $\{(x, A(x)), x \in \Gamma\}$ support wavepackets with the dynamics (1.3). As mentioned above, (1.3) is an exotic semiclassical trajectory not predicted by the standard results on propagation of singularities [DH72]. In other setups, wavepackets may e.g. start away from the crossing set, reach it, and undergo a Landau–Zener transition; see e.g. [Hag94, HJ98, FKG03, CdV04].

From a physical point of view, our motivation stems from the ubiquity of \mathcal{D} in the field of topological phases of matter [Wit16, MM21], and in particular one-particle models of topological insulators and topological superconductors [Vol89, Ber13, PSB16, Bal19a, Bal19b], which generically come with conical points [Dro21b]. The domain wall $\kappa(x)$ models the interface between two topologically distinct insulating phases [FLTW16, Dro19b, DW20]. This in turn generates an asymmetric transport along the interface Γ by a principle called the bulk-interface correspondence; see e.g. [EG02, GP13, PSB16, Dro19a, Bal20, Bal21, Dro21a, BM22]. The wavepackets analyzed here encode this asymmetry; see [BBD⁺21, §1.4] for a discussion when $A = 0$. The operator \mathcal{D} also emerges in the effective analysis of graphene and its pseudomagnetic (strained) analogues, see for instance [GRW21].

Note that the magnetic field is essential in the integer quantum Hall effect [TKNdN82, ASS94, BvESB94], which was the first observed example of topologically non-trivial state of matter. There, the insulating gaps are obtained from the degenerate Landau levels associated to a constant magnetic field. This contrasts with the situation considered here: the non-trivial topology imposed by the domain wall is stable against magnetic contributions; see [Bal21]. This means that the magnetic field does not influence the existence of edge states. It however affects their quantitative features, see §1.3.

Our results in Theorems 1.1 and 6.2 concern weakly dispersive wavepackets with a macroscopic center y_t propagating along $\Gamma = \kappa^{-1}(0)$. Changing the metric (i.e. replacing D by $(D_j + \Gamma_j(x))g^{jk}(x)\sigma_k$) would likely preserve this structure. However, adding an electric potential $V(x)$ to \mathcal{D} modifies the energy landscape, hence the interface $\Gamma = \kappa^{-1}(0)$ between the two topological media cannot properly support wavepackets. These are likely to propagate instead within a thicker strip close to Γ , with splitting according to Landau–Zener transition rules.

Acknowledgments. This work is a sequel to [BBD⁺21], which started during a 2020 AIM workshop, Mathematics of topological insulators. The authors thank the organizers: Daniel Freed, Gian Michele Graf, Rafe Mazzeo and Michael Weinstein. In addition, the authors

are very grateful to Clotilde Fermanian Kammerer, Jianfeng Lu, and Alexander Watson for initial discussions. The authors acknowledge support from the NSF grants DMS-2118608 and DMS-2054589 (AD), DMS-1908736 and EFMA-1641100 (GB).

2. GAUGE TRANSFORMATION

This section constructs a gauge function that reduces \tilde{D} to an operator with magnetic properties closer to those of the model operator (1.6).

2.1. Equivalent tangent magnetic potential. We construct first a magnetic potential \tilde{A} on \mathbb{R}^2 such that $\nabla \times \tilde{A} = B$ near Γ , and \tilde{A} vanishes along $\Gamma = \kappa^{-1}(0)$ and is carried by the vector field τ . In particular, the operator

$$\tilde{D} = (\varepsilon D_1 - \tilde{A}_1(x))\sigma_1 + (\varepsilon D_2 - \tilde{A}_2(x))\sigma_2 + \kappa(x)\sigma_3 \quad (2.1)$$

will share many of the characteristics of the model (1.6): the magnetic potential is tangent to Γ and vanishes along Γ . Moreover, from magnetic equivalence between A and \tilde{A} , D and \tilde{D} are locally conjugated. We use below the notation (with Euclidean distance d)

$$U_\eta = \{x \in \mathbb{R}^2 : d(x, \Gamma) < \eta\}.$$

Lemma 2.1. *There exist $\eta > 0$ and a function $\beta \in C^\infty(\mathbb{R}^2, \mathbb{R})$ vanishing on $\mathbb{R}^2 \setminus U_{2\eta}$ such that, defining $\tilde{A} := \beta\kappa\tau$, we have*

$$\nabla \times (A - \tilde{A}) = \nabla \times (A - \beta\kappa\tau) = 0 \quad \text{on } U_\eta; \quad \beta = \frac{B}{|\nabla\kappa|} \quad \text{on } \Gamma. \quad (2.2)$$

Proof. We first note that it suffices to construct β on U_η and extend it to \mathbb{R}^2 as a smooth function with support in $U_{2\eta}$. For f a smooth function, we observe that

$$\nabla \times (f\tau) = \nabla f \times \tau + f\nabla \times \tau = \partial_n f + f\nabla \times \tau, \quad (2.3)$$

where we used $\nabla f \times \tau = \partial_1 f \tau_2 - \partial_2 f \tau_1 = \partial_n f$. Let $B = \nabla \times A$. To find β such that $B = \nabla \times (\beta\kappa\tau)$, we first solve

$$\partial_n f + f\nabla \times \tau = B, \quad f|_\Gamma = 0. \quad (2.4)$$

We note that the coefficients of ∂_n are in C_b^∞ . In particular, the flow $e^{s\partial_n}(x)$ is defined for all times. We now define the map $\Phi : \Gamma \times \mathbb{R} \rightarrow \mathbb{R}^2$ by

$$\Phi(x, s) = e^{s\partial_n}(x).$$

If f solves (2.4), then $\tilde{f} = f \circ \Phi$ solves

$$\partial_s \tilde{f} + \tilde{f} \cdot (\nabla \times \tau) \circ \Phi = B \circ \Phi, \quad \tilde{f}|_\Gamma = 0.$$

This equation clearly admits a solution $\tilde{f} \in C^\infty(\Gamma \times \mathbb{R}, \mathbb{R})$. Moreover, note that ∂_n is transverse to Γ ; that $|\nabla\kappa|$ is uniformly bounded above and below; and that $\kappa \in C_b^\infty$, Φ is a diffeomorphism from $\Gamma \times (-\delta, \delta)$ to its range, which contains a neighborhood of the form U_η . Therefore, \tilde{f} induces the solution $f = \tilde{f} \circ \Phi^{-1} \in C^\infty(U_\eta, \mathbb{R})$ of (2.4).

Because f vanishes on Γ and κ vanishes transversely on Γ (with $|\nabla\kappa|$ bounded below), after potentially reducing η we can write $f = \beta\kappa$ for some smooth function β on U_η . From the equation (2.4) and the identity (2.3), we conclude that

$$\nabla \times A = B = \partial_n f + f\nabla \times \tau = \nabla \times (f\tau) = \nabla \times (\beta\kappa\tau).$$

Moreover, using that f and κ vanish on Γ , we have along Γ :

$$B = \partial_n f = \partial_n(\beta\kappa) = \beta\partial_n\kappa = \beta|\nabla\kappa|.$$

This completes the proof. \square

Thanks to (2.2), A and \tilde{A} give rise to the same magnetic field on U_η , hence the difference $A - \tilde{A}$ is locally a gradient field:

Lemma 2.2. *Let $\eta > 0$ given by Lemma 2.1 and $U \subset U_\eta$ be a simply connected set. For any $(y_0, \eta_0) \in U \times \mathbb{R}$, there exists a unique $\chi_0 \in C^\infty(U, \mathbb{R})$ such that*

$$\tilde{A} = A - \nabla\chi_0 \quad \text{in } U; \quad \text{and} \quad \chi_0(y_0) = \eta_0.$$

Proof of Lemma 2.2. Thanks to (2.2), we have $\nabla \times (A - \beta\kappa\tau) = 0$ on U_η , hence on U . Since U is simply connected, by Poincaré's lemma there exists a unique $\chi_0 \in C^\infty(U, \mathbb{R})$ such that $\nabla\chi_0 = A - \beta\kappa\tau$ and $\chi_0(y_0) = \eta_0$. \square

It follows from Lemma 2.2 that the operators \mathcal{D} and $\tilde{\mathcal{D}}$ defined in (2.1) are locally conjugate: on open sets U produced by Lemma 2.2, we have

$$e^{-i\chi_0/\varepsilon} \mathcal{D} e^{i\chi_0/\varepsilon} = \tilde{\mathcal{D}}. \quad (2.5)$$

2.2. Global gauge. When Γ is simply connected, we can pick $U = U_\eta$ in Lemma 2.2, and the conjugation relation (2.5) holds on a full neighborhood of Γ . This however fails when Γ is a loop: χ_0 is only defined on part of Γ . We circumvent this obstacle by using instead a time-dependent gauge that follows the center of mass y_t of our wavepacket, i.e. defined on a set of the form

$$\Omega_\delta = \{(t, x), t \in \mathbb{R}, |x - y_t| < \delta\}.$$

Proposition 2.3. *Let \tilde{A} be as defined in Lemma 2.1. There exist $\delta > 0$ and $\chi \in C^\infty(\mathbb{R} \times \mathbb{R}^2, \mathbb{R})$ with support in $\Omega_{2\delta}$ with $\chi(0, y_0) = 0$ and such that for $(t, x) \in \Omega_\delta$:*

$$\tilde{A}(x) = A(x) - \nabla\chi(t, x), \quad \partial_t\chi(t, x) = 0. \quad (2.6)$$

Proof. As in the proof of Lemma 2.1, it suffices to construct χ on Ω_δ and to extend it to \mathbb{R}^2 as a smooth function with support in $\Omega_{2\delta}$. Without loss of generality, we can assume that Γ is connected. If Γ is also simply connected, then we simply take $\delta = \eta$ and $\chi(t, x) = \chi_0(x)$. If Γ is not simply connected, then it is a loop; in particular it is compact.

Fix $t \in \mathbb{R}$. According to Lemma 2.2, there exists $\delta_t > 0$ and a smooth function $\chi(t, \cdot)$ defined on the ball $B(y_t, \delta_t)$ such that

$$\tilde{A}(x) = A(x) - \nabla\chi(t, x), \quad x \in B(y_t, \delta_t); \quad \chi(t, y_t) = \int_0^t \dot{y}_s \cdot A(y_s) ds. \quad (2.7)$$

Since Γ is compact, we can pick δ_t independent of t ; we write below $\delta = \delta_t$. Varying t , we obtain a uniquely defined function χ on Ω_δ . We take time-derivative of both identities in (2.7). The first one yields $\nabla\partial_t\chi(t, x) = 0$ for $x \in B(y_t, \delta)$. The second one gives

$$\partial_t\chi(t, y_t) = \partial_t(\chi(t, y_t)) - \dot{y}_t \cdot \nabla\chi(t, y_t) = \dot{y}_t \cdot A(y_t) - \dot{y}_t \cdot A(y_t) = 0,$$

where we used $\nabla\chi = \beta\kappa\tau - A$ hence $\nabla\chi(t, y_t) = A(y_t)$. From these identities, we deduce that $\partial_t\chi(t, x) = 0$ for $x \in B(y_t, \delta)$; in particular $\partial_t\chi$ is smooth. Moreover, $\nabla\chi = A - \tilde{A}$ is also smooth. This implies that χ is smooth, which completes the proof. \square

Thanks to (2.6), we have the relation

$$e^{-i\chi/\varepsilon}(\varepsilon D_t + \tilde{D})e^{i\chi/\varepsilon} = \varepsilon D_t + \tilde{D} + R, \quad \text{where:} \quad (2.8)$$

$$R = \partial_t \chi + (\partial_1 \chi - A_1 + \tilde{A}_1)\sigma_1 + (\partial_2 \chi - A_2 + \tilde{A}_2)\sigma_2. \quad (2.9)$$

It follows from (2.8) that Ψ solves (1.1) if and only if $\tilde{\Psi} = e^{-i\chi/\varepsilon}\Psi$ solves

$$(\varepsilon D_t + \tilde{D} + R)\tilde{\Psi} = 0. \quad (2.10)$$

While R looks like a leading-order term, it will effectively be of order ε^∞ because it vanishes on Ω_δ – a domain where our wavepacket is concentrated. We will eventually treat it as a small source term in §6.

2.3. Gauge expansion. We conclude this section with an expansion of χ near (t, y_t) , which will serve to prove Theorem 1.1.

Lemma 2.4. *We have:*

$$\chi(t, y_t) = \int_0^t \dot{y}_s \cdot A_s ds; \quad \nabla \chi(t, y_t) = A_t; \quad \nabla^2 \chi(t, y_t) = \nabla A^\top(y_t) - B_t n_t \tau_t^\top. \quad (2.11)$$

Proof. The first formula of (2.11) comes from the equation (2.7) defining χ . The second one follows from

$$\nabla \chi(t, y_t) = A(y_t) - \beta(y_t) \kappa(y_t) \tau(y_t) = A_t.$$

In order to see the last one, we notice that

$$\begin{aligned} \nabla^2 \chi(t, y_t) &= \nabla \cdot (\nabla^\top \chi)(y_t) = \nabla (A^\top - \beta \kappa \tau^\top)(y_t) \\ &= \nabla A^\top(y_t) - \beta(y_t) \nabla \kappa(y_t) \tau^\top(y_t) = \nabla A^\top(y_t) - B_t n_t \tau_t^\top, \end{aligned}$$

where we used $\beta_t = B_t/r_t$ on Γ , see (2.2). This completes the proof. \square

3. LOCAL NORMAL FORM

We now start our construction of approximate solutions to $(\varepsilon D_t + \tilde{D})\tilde{\Psi} = 0$. This section describes a series of transformations that allows us to explicitly invert an operator that governs the leading dynamics: a rescaling of $\tilde{\Psi}$ in natural coordinates, spatial and spinorial rotations, and finally a shifted partial Fourier transform.

3.1. Spatial rescaling. We first write the wavepacket in natural coordinates:

$$\psi(t, z) = \varepsilon^{\frac{1}{2}} S \tilde{\Psi}(t, z),$$

with the scaling transformation defined as

$$Sf(z) = f(y_t + \sqrt{\varepsilon}z), \quad S^{-1}f(x) = f\left(\frac{x - y_t}{\sqrt{\varepsilon}}\right). \quad (3.1)$$

We observe that $\varepsilon^{\frac{1}{2}}S$ is an isometry on $L^2(\mathbb{R}^2)$, while formally,

$$Sf(z) = \sum_{\alpha} \frac{\varepsilon^{\frac{|\alpha|}{2}} z^\alpha}{\alpha!} \partial^\alpha f(y_t) = \sum_{j \geq 0} \varepsilon^{\frac{j}{2}} S_j f(z), \quad S_j f(z) := \sum_{|\alpha|=j} \frac{z^\alpha}{\alpha!} \partial^\alpha f(y_t). \quad (3.2)$$

This scaling emerges from the following consideration. For $\kappa(x) = x_2$, the wavepacket is confined by a harmonic-like oscillator $\varepsilon D_2 \sigma_2 + x_2 \sigma_3$ and hence comes with a natural

scale $\sqrt{\varepsilon}x_2$. This scale is then imposed to all directions when κ admits more sophisticated variations, i.e. when Γ is curved.

We then verify that $\psi(t, z)$ solves

$$L\psi = 0; \quad L = \varepsilon^{-\frac{1}{2}}S(\varepsilon D_t + \tilde{D})S^{-1} = \varepsilon^{\frac{1}{2}}D_t - \dot{y}_t \cdot D + (D + \varepsilon^{-\frac{1}{2}}Sh(z)) \cdot \sigma \quad (3.3)$$

with $D = (D_1, D_2, 0)^t$, $\sigma = (\sigma_1, \sigma_2, \sigma_3)^t$, and $h = (-\tilde{A}_1, -\tilde{A}_2, \kappa)^t = (-\beta\kappa\tau, \kappa)^t$. We still use \dot{y}_t for $(\dot{y}_t, 0)^t \in \mathbb{R}^3$. Using (3.2), we find formally $L = \sum_{j \geq 0} \varepsilon^{\frac{j}{2}} L_j$, where we observe that $L_{-1} = 0$ since \tilde{A} and hence h vanish on $\Gamma = \kappa^{-1}(0)$, and

$$L_0 = -\dot{y}_t \cdot D + D \cdot \sigma + S_1 h \cdot \sigma \quad (3.4)$$

$$L_1 = D_t + S_2 h \cdot \sigma \quad (3.5)$$

$$L_j = S_{j+1} h \cdot \sigma = \sum_{|\alpha|=j+1} \frac{1}{\alpha!} z^\alpha \partial^\alpha h(y_t) \cdot \sigma, \quad j \geq 2. \quad (3.6)$$

Our next objective is to transform L_0 in an appropriate basis so that its infinite dimensional kernel and its inverse on the orthogonal complement may be written explicitly.

3.2. Rotations. We now introduce spatial and spinorial rotations already mentioned in the introduction. We define, for $1 \leq j \leq 3$ and $\theta \in \mathbb{R}$, the spinor rotation acting on \mathbb{C}^2 :

$$U_{j,\theta} = e^{-i\frac{\theta}{2}\sigma_j} = \cos\left(\frac{\theta}{2}\right) - i \sin\left(\frac{\theta}{2}\right) \sigma_j.$$

Associated to it is a standard three-dimensional rotation acting on \mathbb{R}^3 of angle θ about the j -axis (with usual orientation) $\tilde{R}_{j,\theta}$ such that for $c \in \mathbb{R}^3$:

$$\tilde{R}_{j,\theta} c \cdot \sigma = c \cdot U_{j,\theta}^* \sigma U_{j,\theta}, \quad U_{j,\theta}^* \sigma U_{j,\theta} =: \tilde{R}_{j,\theta}^* \sigma.$$

Note that

$$U_{k,\theta}^* \sigma_j U_{k,\theta} = \delta_{j \neq k} \cos \theta \sigma_j + \epsilon_{jkl} \sin \theta \sigma_l + \delta_{jk} \sigma_j.$$

From this, we deduce two formulas that we will use later:

$$\tilde{R}_{2,\varphi_t} = \begin{bmatrix} \cos \varphi_t & 0 & -\sin \varphi_t \\ 0 & 1 & 0 \\ \sin \varphi_t & 0 & \cos \varphi_t \end{bmatrix} = \begin{bmatrix} c_t & 0 & -s_t \\ 0 & 1 & 0 \\ s_t & 0 & c_t \end{bmatrix}, \quad \tilde{R}_{3,\theta_t} = \begin{bmatrix} R_{\theta_t} & 0 \\ 0 & 1 \end{bmatrix}, \quad (3.7)$$

where R_{θ_t} is the spatial rotation defined in (1.4). We finally define the L^2 -unitary transform related to the spatial rotations

$$\mathcal{R}_{\theta_t} f = f \circ R_{\theta_t}$$

and the operator

$$U_t = U_{2,\varphi_t} U_{3,\theta_t} \mathcal{R}_{\theta_t}. \quad (3.8)$$

When $\varphi_t \equiv 0$ (i.e., when $B = 0$), this transformation already appears in [BBD⁺21]. We now compute the operators $U_t^* L_j U_t$. We start with preliminary relations:

Lemma 3.1. *We find:*

$$U_t^* (-\dot{y}_t \cdot D) U_t = c_t D_1, \quad U_t^* D \cdot \sigma U_t = (c_t \sigma_1 + s_t \sigma_3) D_1 + \sigma_2 D_2, \quad (3.9)$$

$$U_t^* D_t U_t = D_t - \frac{\dot{\varphi}_t}{2} \sigma_2 - \frac{\dot{\theta}_t}{2} (-s_t \sigma_1 + c_t \sigma_3) - \dot{\theta}_t (z_1 D_2 - z_2 D_1). \quad (3.10)$$

Moreover $U_t^* S_j h \cdot \sigma U_t$ is a multiplication operator by a time-dependent homogeneous polynomial of degree j in z ; and for $j = 1, 2$:

$$U_t^* S_1 h \cdot \sigma U_t = \rho_t z_2 \sigma_3, \quad U_t^* S_2 h \cdot \sigma U_t = -r_t c_t z_2 \langle z, \nabla \beta(y_t) \rangle + q_t(z) \cdot \sigma \quad (3.11)$$

where $q_t \cdot \sigma$ is a matrix-valued quadratic form in z carried by σ_2, σ_3 .

Proof. 1. Space differentiation. We first observe that

$$\mathcal{R}_{\theta_t}^* D \mathcal{R}_{\theta_t} = \tilde{R}_{3,\theta_t}^* D, \quad \mathcal{R}_{\theta_t}^* D \cdot v \mathcal{R}_{\theta_t} = D \cdot \tilde{R}_{3,\theta_t} v. \quad (3.12)$$

Applied to $v = -\dot{y}_t$ with $R_{\theta_t} \dot{y}_t = -c_t e_1$ by construction, we find $-\tilde{R}_{3,\theta_t} \dot{y}_t \cdot D = c_t D_1$. Since spinorial rotations commute with scalars, we obtain the first relation in (3.9).

We now apply (3.12) to $v = \sigma$ and find

$$(\mathcal{R}_{\theta_t} U_{3,\theta_t})^* D \cdot \sigma \mathcal{R}_{\theta_t} U_{3,\theta_t} = D \cdot \sigma.$$

Therefore, using (3.7), we find

$$U_t^* D \cdot \sigma U_t = U_{2,\varphi_t}^* D \cdot \sigma U_{2,\varphi_t} = D \cdot \tilde{R}_{2,\varphi_t}^* \sigma = (c_t \sigma_1 + s_t \sigma_3) D_1 + \sigma_2 D_2.$$

This proves the second identity in (3.9).

2. Time differentiation. We note that

$$\mathcal{R}_{\theta_t}^* D_t \mathcal{R}_{\theta_t} = D_t - \dot{\theta}_t J z \cdot D_z, \quad U_{3,\theta_t}^* D_t U_{3,\theta_t} = D_t - \frac{\dot{\theta}_t}{2} \sigma_3, \quad U_{2,\varphi_t}^* D_t U_{2,\varphi_t} = D_t - \frac{\dot{\varphi}_t}{2} \sigma_2.$$

We deduce that

$$\begin{aligned} U_t^* D_t U_t &= U_{2,\varphi_t}^* \left(D_t - \frac{\dot{\theta}_t}{2} \sigma_3 \right) U_{2,\varphi_t} - \dot{\theta}_t (z_1 D_2 - z_2 D_1) \\ &= D_t - \frac{\dot{\varphi}_t}{2} \sigma_2 - \frac{\dot{\theta}_t}{2} (-s_t \sigma_1 + c_t \sigma_3) - \dot{\theta}_t (z_1 D_2 - z_2 D_1). \end{aligned}$$

This proves (3.10).

3. Multiplicative operators. We observe that $\mathcal{R}_{\theta_t}^* S h \mathcal{R}_{\theta_t} = h(y_t + \sqrt{\varepsilon} R_{\theta_t} z)$ as a multiplication operator so that

$$U_t^* S h \cdot \sigma U_t = \tilde{R}_{2,\varphi_t}^* \tilde{R}_{3,\theta_t} h(y_t + \sqrt{\varepsilon} R_{\theta_t} z) \cdot \sigma = \sum_{j \geq 1} \varepsilon^{\frac{j}{2}} \sum_{|\alpha|=j} z^\alpha \nu_\alpha \cdot \sigma,$$

where we have defined the 3-vectors $\nu_\alpha = \frac{1}{\alpha!} \tilde{R}_{2,\varphi_t} \tilde{R}_{3,\theta_t} (R_{-\theta_t} \nabla)^\alpha h(y_t)$. Indeed,

$$h(y_t + \sqrt{\varepsilon} R_\theta z) = \sum_{j \geq 1} \frac{1}{j!} (\sqrt{\varepsilon} z \cdot R_{-\theta} \nabla)^j h(y_t) = \sum_{j \geq 1} \varepsilon^{\frac{j}{2}} \sum_{|\alpha|=j} \frac{z^\alpha}{\alpha!} (R_{-\theta} \nabla)^\alpha h(y_t)$$

using multinomial coefficients. Therefore, $S_j h \cdot \sigma$ is a multiplicative operator by a homogeneous polynomial of degree j depending on t ; composing with rotations preserves this feature: $U_t^* S_j h \cdot \sigma U_t$ is also a homogeneous polynomial of degree j .

We now focus on $j = 1$. Define $\tilde{h} = (-\beta \tau, 1)^t$ so that $h = \kappa \tilde{h}$. Since $\kappa(y_t) = 0$, we have $S_1 h = \tilde{h}(y_t) \cdot S_1 \kappa$. Since spinorial rotations commute with the scalar $S_1 \kappa$,

$$U_t^* S_1 h \cdot \sigma U_t = \mathcal{R}_{\theta_t}^* S_1 \kappa \mathcal{R}_{\theta_t} U_t^* \tilde{h}(y_t) \cdot \sigma U_t. \quad (3.13)$$

We now compute separately $\mathcal{R}_{\theta_t}^* S_1 \kappa \mathcal{R}_{\theta_t}$ and $U_t^* \tilde{h}(y_t) \cdot \sigma U_t$ (note that \mathcal{R}_{θ_t} does not technically affect that last term since $\tilde{h}(y_t)$ does not depend on z). We have:

$$\mathcal{R}_{\theta_t}^* S_1 \kappa \cdot \sigma \mathcal{R}_{\theta_t} = \mathcal{R}_{\theta_t}^* z \cdot \nabla \kappa(y_t) \mathcal{R}_{\theta_t} = z \cdot R_{\theta_t} \nabla \kappa(y_t) = r_t z_2. \quad (3.14)$$

Moreover, since $B_t = \beta_t r_t$, we have $\tilde{h}(y_t) = (-r_t^{-1} B_t \tau_t, 1)^t$ hence $R_{3,\theta_t} \tilde{h} = (r_t^{-1} B_t, 0, 1)$. We deduce

$$U_{3,\theta_t}^* \tilde{h}(y_t) \cdot \sigma U_{3,\theta_t} = R_{3,\theta_t} \tilde{h}(y_t) \cdot \sigma = \frac{B_t \sigma_1}{r_t} + \sigma_3 = \frac{\rho_t}{r_t} (s_t \sigma_1 + c_t \sigma_3).$$

By (3.7), $U_{2,\varphi_t}^* (s_t \sigma_1 + c_t \sigma_3) U_{2,\varphi_t} = \sigma_3$, which is the main motivation for the definition of U_{2,φ_t} . Hence,

$$U_t^* \tilde{h}(y_t) \cdot \sigma U_t = U_{2,\varphi_t}^* U_{3,\theta_t}^* \tilde{h}(y_t) \cdot \sigma U_{3,\theta_t} U_{2,\varphi_t} = \frac{\rho_t}{r_t} \sigma_3. \quad (3.15)$$

Multiplying (3.14) and (3.15) and going back to (3.13), we conclude that:

$$U_t^* S_1 h \cdot \sigma U_t = \rho_t z_2 \sigma_3.$$

We go on with the case $j = 2$. We compute $U_t^* S_2 h \cdot \sigma U_t$. We recall that $h = \kappa \tilde{h}$, therefore (using again $\kappa(y_t) = 0$):

$$S_2 h = 2(S_1 \kappa)(S_1 \tilde{h}) + (S_2 \kappa) \tilde{h}(y_t). \quad (3.16)$$

We start with the contribution of $(S_2 \kappa) \tilde{h}(y_t)$, i.e. the term $U_t^* (S_2 \kappa) \tilde{h}(y_t) \cdot \sigma U_t$. Using that $S_2 \kappa$ is a scalar and $\tilde{h} \cdot \sigma$ does not depend on z we have

$$U_t^* (S_2 \kappa) \tilde{h}(y_t) U_t = \mathcal{R}_{\theta_t}^* (S_2 \kappa) \mathcal{R}_{\theta_t} U_t^* \tilde{h}(y_t) \cdot \sigma U_t = \mathcal{R}_{\theta_t}^* (S_2 \kappa) \mathcal{R}_{\theta_t} \frac{\rho_t}{r_t} \sigma_3, \quad (3.17)$$

where we used (3.15) in the last equality. This term is carried by σ_3 .

We now focus on $U_t^* (S_1 \kappa) (S_1 \tilde{h}(y_t) \cdot \sigma) U_t$. Using (3.14) and that spinorial rotations commute with scalars, we obtain:

$$U_t^* (S_1 \kappa) (S_1 \tilde{h}) \cdot \sigma U_t = \mathcal{R}_{\theta_t}^* (S_1 \kappa) \mathcal{R}_{\theta_t} U_t (S_1 \tilde{h}) \cdot \sigma U_t = r_t z_2 U_t (S_1 \tilde{h}) \cdot \sigma U_t. \quad (3.18)$$

It remains to compute $U_t (S_1 \tilde{h}) \cdot \sigma U_t$. We recall that $\tilde{h} = (-\beta \tau, 1)^t$. Moreover, τ has unit norm: $\langle \tau, \tau \rangle = 1$. Taking derivatives of this expression shows that $\partial_j \tau$ is normal to τ ; we write $\partial_j \tau = \alpha_j n$ below. We deduce that

$$\begin{aligned} (S_1 \tilde{h}) \cdot \sigma &= - \sum_{j,k=1}^2 z_j \partial_j (\beta \tau_k) \sigma_k = - \sum_{j,k=1}^2 z_j (\partial_j \beta) \tau_k \sigma_k - \beta \sum_{j,k=1}^2 z_j \partial_j \tau_k \sigma_k \\ &= -(S_1 \beta) \tau \cdot \sigma - \beta \sum_{j,k=1}^2 z_j \alpha_j n_k \sigma_k = -(S_1 \beta) \tau \cdot \sigma - \langle z, \alpha \rangle n \cdot \sigma. \end{aligned}$$

We observe that:

$$U_{3,\theta_t}^* \tau \cdot \sigma U_{3,\theta_t} = R_{3,\theta_t} \tau \cdot \sigma = -\sigma_1, \quad U_{2,\varphi_t}^* \sigma_1 U_{2,\varphi_t} = R_{2,\varphi_t} e_1 \cdot \sigma = c_t \sigma_1 - s_t \sigma_3;$$

$$U_{3,\theta_t}^* n \cdot \sigma U_{3,\theta_t} = R_{3,\theta_t} n \cdot \sigma = \sigma_2, \quad U_{2,\varphi_t}^* \sigma_2 U_{2,\varphi_t} = \sigma_2.$$

Therefore, the following equality is valid modulo terms carried by σ_2, σ_3 :

$$U_t (S_1 \tilde{h}) \cdot \sigma U_t = \mathcal{R}_{\theta_t}^* S_1 \beta \mathcal{R}_{\theta_t} c_t \sigma_1 = c_t \langle z, R_{\theta_t} \nabla \beta(y_t) \rangle \sigma_1.$$

Going back to (3.16), (3.17) and (3.18), we conclude that – again with equality valid modulo terms carried by σ_2, σ_3 :

$$U_t^* S_2 h \cdot \sigma U_t = U_t^* (S_1 \kappa) (S_1 \tilde{h}) \cdot \sigma U_t = -c_t r_t z_2 \langle z, R_{\theta_t} \nabla \beta(y_t) \rangle \sigma_1.$$

This completes the proof of (3.11), hence of the lemma. \square

Thanks to Lemma 3.1, we can describe $U_t^* L_j U_t$. For $j \geq 2$, $U_t^* L_j U_t$ is a multiplication operator by a homogeneous polynomial of degree j . For $j = 0, 1$, we have explicit expressions:

$$U_t^* L_0 U_t = c_t(1 + \sigma_1)D_1 + \sigma_2 D_2 + (\rho_t z_2 + s_t D_1)\sigma_3; \quad (3.19)$$

$$U_t^* L_1 U_t = D_t - \dot{\theta}_t(z_1 D_2 - z_2 D_1) + \left(\frac{\dot{\theta}_t s_t}{2} + j_t z_1 z_2 + k_t z_2^2 \right) \sigma_1 + E_1, \quad (3.20)$$

where E_1 is a multiplication operator by a polynomial of degree two and carried by σ_2 and σ_3 (we will see later that it does not contribute to leading order). The coefficients j_t and k_t are explicitly given in terms of β :

$$\begin{bmatrix} j_t \\ k_t \end{bmatrix} = r_t c_t R_{\theta_t} \nabla \beta(y_t) = r_t c_t \begin{bmatrix} -\partial_\tau \beta(y_t) \\ \partial_n \beta(y_t) \end{bmatrix}.$$

We end this section with a few important relations:

$$j_t = -r_t \dot{\beta}_t; \quad j_t \gamma_t = \frac{d \ln c_t}{dt}; \quad k_t = \frac{c_t}{2} \left(\partial_n B(y_t) - B_t \frac{\Delta \kappa(y_t)}{r_t} \right). \quad (3.21)$$

Proof of (3.21). We start with the first identity in (3.21). We recall that $\dot{y}_t = c_t \tau_t$. Since ∂_τ is tangent to Γ and $\beta = B/|\nabla \kappa|$ along Γ :

$$j_t = -r_t c_t \partial_\tau \beta(y_t) = -r_t c_t \tau_t \cdot \nabla \left(\frac{B}{|\nabla \kappa|} \right) (y_t) = -r_t \dot{y}_t \cdot \nabla \left(\frac{B}{|\nabla \kappa|} \right) (y_t) = -r_t \dot{\beta}_t. \quad (3.22)$$

Regarding the second identity: using (3.22) and $\beta_t = B_t/r_t$ on Γ , we obtain:

$$j_t \gamma_t = -\frac{r_t B_t}{r_t^2 + B_t^2} \dot{\beta}_t = -\frac{\beta_t \dot{\beta}_t}{1 + \beta_t^2} = -\frac{1}{2} \frac{d}{dt} \ln(1 + \beta_t^2) = -\frac{1}{2} \frac{d}{dt} \ln \frac{1}{c_t^2} = \frac{d \ln c_t}{dt}.$$

For the last identity, we first note that $|\nabla \kappa| = \partial_n \kappa$. Since $f = \beta \kappa$ solves (2.4):

$$B = \partial_n f + f \nabla \times \tau = \partial_n(\beta \kappa) + \beta \kappa \nabla \times \tau = \kappa \partial_n \beta + \beta \partial_n \kappa + \beta \kappa \nabla \times \tau.$$

Therefore, since κ vanishes on Γ , we deduce that on Γ :

$$\partial_n B = 2 \partial_n \kappa \partial_n \beta + \beta (\partial_n^2 \kappa + \partial_n \kappa \nabla \times \tau) = 2 |\nabla \kappa| \partial_n \beta + B \left(\frac{\partial_n^2 \kappa}{|\nabla \kappa|} + \nabla \times \tau \right). \quad (3.23)$$

Moreover, using $|\nabla \kappa| = \partial_n \tau$ and $\nabla f \times J \nabla g = \nabla f \cdot \nabla g$:

$$\nabla \times \tau = \frac{\nabla \times J \nabla \kappa}{|\nabla \kappa|} + \nabla \frac{1}{|\nabla \kappa|} \cdot J \nabla \kappa = \frac{\Delta \kappa}{|\nabla \kappa|} - \frac{\partial_n^2 \kappa}{|\nabla \kappa|}. \quad (3.24)$$

Plugging (3.24) into (3.23) and using the defining equation $\kappa_t = r_t c_t \partial_n \beta(y_t)$ yield (3.21). \square

3.3. Shifted Fourier transform. We finally introduce the L^2 - unitary transformation

$$\mathcal{V}_t a(z) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{iz_1 \xi} a(\xi, z_2 + \gamma_t \xi) d\xi. \quad (3.25)$$

Its inverse is given explicitly by

$$\mathcal{V}_t^* a(\xi, \zeta) = \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} e^{-iz_1 \xi} a(z_1, \zeta - \gamma_t \xi) dz_1.$$

Above, ξ is the one-dimensional dual Fourier variable to z_1 while the one-dimensional variable ζ is a shift of z_2 given by $\zeta = z_2 + \gamma_t \xi$. This shifted partial Fourier transform, which reduces to a partial Fourier transform in the first variable $z_1 \rightarrow \xi$ when $B = 0$ and introduces an additional shift $\gamma_t \xi$ to z_2 otherwise is reminiscent of the parametrization of Landau level eigenfunctions in a Landau gauge when $B \neq 0$. The construction of our wavepackets involves many functions of the form $a(t, \xi, \zeta) = f(t, \xi) \phi(\zeta)$, with f describing the wavepacket profile along Γ while ϕ describes the wavepacket profile across Γ .

The global transformation we need to consider is the composition of U_t introduced in the preceding section and \mathcal{V}_t given above. We thus set $\mathfrak{U}_t = U_t \mathcal{V}_t$. The operator in this new set of variables is $T := \mathfrak{U}_t^* L \mathfrak{U}_t$ with L defined in (3.3). We similarly define $T_j = \mathfrak{U}_t^* L_j \mathfrak{U}_t$ for $j \geq 0$. We will also be using $D_\xi = -i\partial_\xi$ and $D_\zeta = -i\partial_\zeta$.

In T_1 , it will turn out that the terms carried by σ_2 or σ_3 and the terms that are odd in (D_ζ, ζ) do not appear to leading order in our expansion. Therefore, we use the notation

$$P_1 \equiv P_2$$

if $P_1 - P_2$ is a differential operator made of odd terms in (ζ, D_ζ) and terms carried by σ_2, σ_3 .

Lemma 3.2. *With $\mathfrak{U}_t = U_t \mathcal{V}_t$ and $T_j = \mathfrak{U}_t^* L_j \mathfrak{U}_t$, we have:*

$$T_0 = c_t(1 + \sigma_1)\xi + D_\zeta \sigma_2 + \rho_t \zeta \sigma_3, \quad (3.26)$$

$$T_1 \equiv D_t + \dot{\theta}_t \gamma_t (D_\zeta^2 - \xi^2) + \left(\frac{\dot{\theta}_t s_t}{2} + j_t \gamma_t (D_\xi \xi - D_\zeta \zeta) + k_t (\zeta^2 + \gamma_t^2 \xi^2) \right) \sigma_1. \quad (3.27)$$

Moreover T_j is a linear combination of differential operators of the form $\xi^k \zeta^\ell D_\xi^m D_\zeta^{j+1-k-\ell-m}$, with $k, \ell \geq 0$ and coefficients bounded with respect to t .

Proof. We note that \mathcal{V}_t satisfies the canonical relations

$$\mathcal{V}_t^* (z_2 + \gamma_t D_1) \mathcal{V}_t = \zeta, \quad \mathcal{V}_t^* D_2 \mathcal{V}_t = D_\zeta, \quad \mathcal{V}_t^* D_1 \mathcal{V}_t = \xi, \quad \mathcal{V}_t^* z_1 \mathcal{V}_t = -(D_\xi + \gamma_t D_\zeta). \quad (3.28)$$

We deduce (3.26) for $T_0 = \mathfrak{U}_t^* L_0 \mathfrak{U}_t$ from the expression (3.19). Thanks to (3.28) we obtain $\mathcal{V}_t^* z_2 \mathcal{V}_t = \zeta - \gamma_t \xi$, hence using the notation \equiv :

$$\begin{aligned} \mathcal{V}_t^* z_1 z_2 \mathcal{V}_t &= -(D_\xi + \gamma_t D_\zeta)(\zeta - \gamma_t \xi) \equiv \gamma_t (D_\xi \xi - D_\zeta \zeta) \\ \mathcal{V}_t^* z_2^2 \mathcal{V}_t &= (\zeta - \gamma_t \xi)^2 \equiv \zeta^2 + \gamma_t^2 \xi^2 \\ \mathcal{V}_t^* (z_1 D_2 - z_2 D_1) \mathcal{V}_t &= -(D_\xi + \gamma_t D_\zeta) D_\zeta - (\zeta - \gamma_t \xi) \xi \equiv -\gamma_t (D_\zeta^2 - \xi^2). \end{aligned}$$

Moreover, we observe that

$$\mathcal{V}_t^* D_t \mathcal{V}_t = D_t + \dot{\gamma}_t \xi D_\zeta \equiv D_t.$$

We deduce from (3.20) the expression for T_1 :

$$T_1 \equiv D_t + \dot{\theta}_t \gamma_t (D_\zeta^2 - \xi^2) + \left(\frac{\dot{\theta}_t s_t}{2} + j_t \gamma_t (D_\xi \xi - D_\zeta \zeta) + k_t (\zeta^2 + \gamma_t^2 \xi^2) \right) \sigma_1.$$

Finally, since L_j is multiplication operator by a homogeneous polynomial of degree $j + 1$, T_j is a linear combination of differential operators of the form $\xi^k \zeta^\ell D_\xi^m D_\zeta^{j+1-k-\ell-m}$, with coefficients bounded with respect to t . This completes the proof. \square

3.4. Conjugation and asymptotic expansion. Introducing $a = \mathfrak{U}_t^* \psi$ and recalling that $T = \mathfrak{U}_t^* L \mathfrak{U}_t$, equation (3.3) is equivalent to $Ta = 0$. With L_j given in (3.4)–(3.6) and $T_j = \mathfrak{U}_t^* L_j \mathfrak{U}_t$, we decompose a as $a = \sum_{j \geq 0} \varepsilon^{\frac{j}{2}} a_j$ to reduce the equation $Ta = 0$ to the triangular system

$$T_0 a_0 = 0, \quad T_1 a_0 + T_0 a_1 = 0, \quad \sum_{k=0}^j T_{j-k} a_k = 0, \quad j \geq 2. \quad (3.29)$$

In §4 we identify the kernel of T_0 and we produce explicit formula for its inverse on the orthogonal complement. We describe how to solve the subleading equation $T_1 a_0 + T_0 a_1 = 0$ in §5. We finally present the higher-order asymptotic expansions and corresponding error estimates in §6.

4. INVERSION OF LEADING OPERATOR

To study the kernel of the operator

$$T_0 = c_t(1 + \sigma_1)\xi + D_\zeta \sigma_2 + \rho_t \zeta \sigma_3,$$

and invert it on the orthogonal complement, we first bring it to a normal form. Thanks to the change of variables $(\xi, \zeta) \mapsto (\rho_t^{1/2} c_t^{-1} \xi, \rho_t^{-1/2} \zeta)$, we can assume that $\rho_t = c_t = 1$. Moreover, with $Q = Q^* = \frac{1}{\sqrt{2}}(\sigma_1 + \sigma_3)$, we have

$$Q \sigma_1 Q = \sigma_3, \quad Q \sigma_2 Q = -\sigma_2, \quad Q \sigma_3 Q = \sigma_1.$$

It suffices then to work with the model operator

$$H = \xi(1 + \sigma_3) + \sigma_1 \zeta - D_\zeta \sigma_2 = \begin{pmatrix} 2\xi & \mathfrak{a}_\zeta \\ \mathfrak{a}_\zeta^* & 0 \end{pmatrix}, \quad \mathfrak{a}_\zeta = \partial_\zeta + \zeta,$$

instead of T_0 . Multiplication by 2ξ may also be written as $\mathfrak{a}_\xi + \mathfrak{a}_\xi^*$ with $\mathfrak{a}_\xi = \partial_\xi + \xi$.

4.1. Functional setting. To study the operator H , we use the standard basis of Hermite functions on $L^2(\mathbb{R})$ given by

$$h_0(\zeta) = \pi^{-\frac{1}{4}} e^{-\frac{1}{2}\zeta^2}, \quad h_n(\zeta) = \frac{1}{2^{\frac{n}{2}} \sqrt{n!}} (\mathfrak{a}_\zeta^*)^n h_0(\zeta), \quad n \geq 1.$$

They satisfy the relations

$$\mathfrak{a}_\zeta h_0 = 0; \quad \mathfrak{a}_\zeta h_n = \sqrt{2n} h_{n-1}, \quad n \geq 1; \quad \mathfrak{a}_\zeta^* h_n = \sqrt{2n+2} h_{n+1}, \quad n \geq 0.$$

With the decomposition $L^2(\mathbb{R}) \ni \psi(\zeta) = \sum_{n \geq 0} \psi_n h_n(\zeta)$, we may then define

$$\begin{aligned} \mathbf{a}_\zeta^{-1} \psi(\zeta) &= \sum_{n \geq 1} \frac{1}{\sqrt{2n}} \psi_{n-1} h_n(\zeta), \quad \psi \in L^2(\mathbb{R}) \\ (\mathbf{a}_\zeta^*)^{-1} \psi(\zeta) &= \sum_{n \geq 0} \frac{1}{\sqrt{2n+2}} \psi_{n+1} h_n(\zeta), \quad \psi \in (\text{Ker } \mathbf{a}_\zeta)^\perp = \overline{\text{Ran } \mathbf{a}_\zeta^*} = \{\psi \in L^2(\mathbb{R}); \psi_0 = 0\}. \end{aligned} \quad (4.1)$$

A natural scale of Hilbert spaces for $p \in \mathbb{N}$ associated to this decomposition is $\mathcal{S}_p(\mathbb{R}) = \{\psi \in L^2(\mathbb{R}); (\mathbf{a}_\zeta^*)^p \psi \in L^2(\mathbb{R})\}$ endowed with the norm

$$\|\psi\|_p^2 = \|(\mathbf{a}_\zeta^*)^p \psi\|_0^2 \cong \|\psi\|_0^2 + \|D_\zeta^p \psi\|_0^2 + \|\zeta^p \psi\|_0^2,$$

where $\|\cdot\|_0$ is the usual $L^2(\mathbb{R})$ -norm and where $a \cong b$ when there exists $C_p > 0$ such that $C_p^{-1}a \leq b \leq C_p a$. That the two above expressions for the norm are equivalent may be obtained by induction from the classical result

$$\|\mathbf{a}_\zeta^* \psi\|_0^2 = \|D_\zeta \psi\|_0^2 + \|\zeta \psi\|_0^2 + \|\psi\|_0^2$$

for $\psi \in \mathcal{S}_1(\mathbb{R})$. Note that $\mathcal{S}_0(\mathbb{R}) \equiv L^2(\mathbb{R})$.

With the decomposition $L^2(\mathbb{R}^2) \ni \psi(\xi, \zeta) = \sum_{m,n \geq 0} \psi_{mn} h_m(\xi) h_n(\zeta)$ in two-dimensional spaces, we similarly define $\mathcal{S}_p(\mathbb{R}^2) = \{\psi \in L^2(\mathbb{R}^2); (\mathbf{a}_\xi^*)^p \psi \in L^2(\mathbb{R}^2) \text{ and } (\mathbf{a}_\zeta^*)^p \psi \in L^2(\mathbb{R}^2)\}$ endowed with the norms

$$\|\psi\|_p^2 = \sum_{j=0}^p \|(\mathbf{a}_\xi^*)^j (\mathbf{a}_\zeta^*)^{p-j} \psi\|_0^2 \cong \|(1 + |\xi|^p + |\zeta|^p) \psi\|_0^2 + \|D_\xi^p \psi\|_0^2 + \|D_\zeta^p \psi\|_0^2.$$

For vector-valued functions, we also define the spaces $\mathcal{S}_p(\mathbb{R}^d, \mathbb{C}^q)$ component-wise. Below we simply write \mathcal{S}_p when the domain \mathbb{R}^d and range \mathbb{C}^q are clear and $\|\cdot\|_p$ the associated norm.

4.2. Inversion of model operator. As the lemma below demonstrates, the kernel $N \subset L^2(\mathbb{R}^2, \mathbb{C}^2)$ of H and its orthogonal complement are given by

$$\begin{aligned} N &= \left\{ \psi(\xi, \zeta) = f(\xi) h_0(\zeta) \begin{bmatrix} 0 \\ 1 \end{bmatrix}, f \in L^2(\mathbb{R}) \right\}, \\ N^\perp &= \left\{ \psi \in L^2(\mathbb{R}^2, \mathbb{C}^2), (\psi_2(\xi, \cdot), h_0(\cdot))_2 = 0, \forall \xi \in \mathbb{R} \right\}, \end{aligned}$$

where $(\cdot, \cdot)_2$ is the standard $L^2(\mathbb{R})$ -inner product in the second variable ζ and $\psi = (\psi_1, \psi_2)^t$.

Lemma 4.1. *For $p \in \mathbb{N}$, we define the bounded linear operator H^{-1} from $N^\perp \cap \mathcal{S}_{p+1}$ to $N^\perp \cap \mathcal{S}_p$ as*

$$H^{-1} = \begin{bmatrix} 0 & (\mathbf{a}_\zeta^*)^{-1} \\ \mathbf{a}_\zeta^{-1} & -\mathbf{a}_\zeta^{-1} 2\xi (\mathbf{a}_\zeta^*)^{-1} \end{bmatrix}.$$

All solutions $\psi \in \mathcal{S}_p$ of the equation $H\psi = g$ for $g \in N^\perp \cap \mathcal{S}_{p+1}$ are of the form $\psi = H^{-1}g + \psi_0$ with ψ_0 arbitrary in $N \cap \mathcal{S}_p$. In particular $\text{Ker } H = N$ on \mathcal{S}_0 .

Proof. The system $H\psi = g$ is equivalent to $\mathbf{a}_\zeta^* \psi_1 = g_2$ and $\mathbf{a}_\zeta \psi_2 + 2\xi \psi_1 = g_1$. Using (4.1), we find for $(g_2, h_0(\zeta))_2 = 0$, i.e., $g \in N^\perp$, that $\psi_1 = (\mathbf{a}_\zeta^*)^{-1} g_2$ and then $\psi_2 = \mathbf{a}_\zeta^{-1} (g_1 - 2\xi \psi_1)$ and hence the above result. We then note that \mathbf{a}_ζ^{-1} has range in N^\perp .

That $\psi \in \mathcal{S}_p$ when $g \in \mathcal{S}_{p+1}$ follows directly from the fact that the following operators are bounded: $\mathbf{a}_\zeta^{-1} : \mathcal{S}_{p+1} \rightarrow \mathcal{S}_{p+1} \cap N^\perp$, $(\mathbf{a}_\zeta^*)^{-1} : \mathcal{S}_{p+1} \cap N^\perp \rightarrow \mathcal{S}_{p+1}$, as well as $\xi : \mathcal{S}_{p+1} \rightarrow \mathcal{S}_p$.

Any function $\psi \in \mathcal{S}_p$ solution of $H\psi = g$ may be decomposed as $\psi_0 + \psi_1$ with $\psi_0 \in N$ and $\psi_1 \in N^\perp$. Since $H\psi_0 = 0$ and hence $\psi_1 = H^{-1}g$, ψ_0 is arbitrary in $\text{Ker } H = N$. \square

4.3. Microscopic balance. Our geometric assumptions impose that $C^{-1} \geq \rho_t \geq C > 0$ and that $0 < c_0 \leq c_t < 1$ for constants C and c_0 independent of $t \geq 0$.

For a fixed time t , we now solve the equation $T_0 a = b$. We introduce

$$\phi_t(\zeta) := \left(\frac{\rho_t}{4\pi}\right)^{\frac{1}{4}} e^{-\frac{\rho_t}{2}\zeta^2} \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad N_t = \text{Ker } T_0 = \{f(\xi)\phi_t(\zeta); f \in \mathcal{S}_0(\mathbb{R}; \mathbb{C})\}. \quad (4.2)$$

The normalization implies that $\|\phi_t\|_{L^2(\mathbb{R}, \mathbb{C}^2)} = 1$. That N_t is the kernel of T_0 comes from Lemma 4.1 and the change of variables given in the last section. We still denote by $\mathcal{S}_p(\mathbb{R}^2, \mathbb{C}^2)$ the spaces of functions in the (ξ, ζ) variables, which are equivalent to the corresponding spaces in the variables $(\rho_t^{-1}c_t\xi, \rho_t^{1/2}\zeta)$ by assumption on (c_t, ρ_t) .

The leading order equation is then solved as follows.

Lemma 4.2. *Let $b \in N_t^\perp \cap \mathcal{S}_{p+1}$. The equation $T_0 a = b$ admits a unique solution $T_0^{-1}b := a \in N_t^\perp \cap \mathcal{S}_p$ with inverse operator given explicitly by*

$$T_0^{-1} = Q \begin{bmatrix} 0 & (\rho_t\zeta - \partial_\zeta)^{-1} \\ (\rho_t\zeta + \partial_\zeta)^{-1} & -2\xi c_t(\rho_t\zeta + \partial_\zeta)^{-1}(\rho_t\zeta - \partial_\zeta)^{-1} \end{bmatrix} Q, \quad Q = \frac{1}{\sqrt{2}}(\sigma_1 + \sigma_3).$$

All solutions of that equation in \mathcal{S}_p are of the form $a = T_0^{-1}b + f(\xi)\phi(\zeta)$ for arbitrary $f(\xi) \in \mathcal{S}_p(\mathbb{R}, \mathbb{C})$.

Proof. This is a corollary of Lemma 4.1 and the invertible transform from T_0 to H . The operator T_0 is thus invertible on the orthogonal complement of $\phi_t(\zeta)$ with $T_0 a = b$ solvable if and only if $(b, \phi_t)_2 = 0$ in which case all solutions are given by $a = T_0^{-1}b + f(\xi)\phi_t$ for f arbitrary in \mathcal{S}_p . \square

The solution to the leading equation $T_0 a_0 = 0$ is therefore given by

$$a_0(t, \xi, \zeta) = f_0(t, \xi)\phi_t(\zeta) \quad (4.3)$$

with $f_0(t, \xi)$ arbitrary at this level.

5. TRANSPORT EQUATION

Lemma 4.2 states that the subleading equation in (3.29), $T_0 a_1 = -T_1 a_0$, admits a solution if and only if $T_1 a_0(t, \cdot) \in N_t^\perp$ for every t . From the expressions (4.3) of a_0 and (4.2) of N_t and $\phi_t(\zeta)$, we deduce that $T_0 a_1 = -T_1 a_0$ is solvable if and only if $\mathcal{T}f_0 = 0$, where

$$\mathcal{T}f(t, \xi) := \int_{\mathbb{R}} \phi_t(\zeta) \cdot T_1[f(t, \xi)\phi_t(\zeta)] d\zeta. \quad (5.1)$$

In this section we provide an explicit expression for \mathcal{T} and for the solutions to $\mathcal{T}f_0 = 0$ and more generally $\mathcal{T}f_0 = g$. We also provide a functional setting to analyze the map $g \rightarrow f_0$ when the dispersion is strongest.

5.1. Derivation of the transport operator.

Lemma 5.1. *We have the identity*

$$\mathcal{T} = D_t - \frac{k_t}{2\rho_t} - j_t\gamma_t \frac{\xi D_\xi + D_\xi \xi}{2} - \left(\dot{\theta}_t \gamma_t + k_t \gamma_t^2 \right) \xi^2. \quad (5.2)$$

Proof. We first observe that

$$\left\langle \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \sigma_j \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\rangle = -2\delta_{1j}.$$

Therefore, the terms in T_1 that are carried by σ_2, σ_3 do not contribute to \mathcal{T} . Likewise, we observe that

$$\int_{\mathbb{R}} \phi(\zeta) \cdot \zeta \phi(\zeta) d\zeta = \int_{\mathbb{R}} \phi(\zeta) \cdot D_\zeta \phi(\zeta) d\zeta = 0.$$

Hence, for the purpose of computing \mathcal{T} , we can ignore terms in T_1 carried by σ_2 or σ_3 , and terms linear in (ζ, D_ζ) . In other words, we can replace T_1 in (5.1) by the right hand side of (3.27):

$$D_t + \dot{\theta}_t \gamma_t (D_\zeta^2 - \xi^2) + \left(\frac{\dot{\theta}_t s_t}{2} + j_t \gamma_t (D_\xi \xi - D_\zeta \zeta) + k_t (\zeta^2 + \gamma_t^2 \xi^2) \right) \sigma_1.$$

We note moreover that

$$D_\zeta \zeta \phi = i(\rho_t \zeta^2 - 1)\phi, \quad D_\zeta^2 \phi(\zeta) = (\rho_t - \rho_t^2 \zeta^2)\phi(\zeta), \quad D_t \phi(\zeta) = i \frac{\dot{\rho}_t}{2} \zeta^2 \phi(\zeta) - i \frac{\dot{\rho}_t}{4\rho_t} \phi(\zeta).$$

Therefore, we deduce that

$$\begin{aligned} \mathcal{T} &= \left(\frac{\rho_t}{\pi} \right)^{\frac{1}{2}} \int_{\mathbb{R}} e^{-\rho_t \zeta^2} \left(D_t + i \frac{\dot{\rho}_t}{2} \zeta^2 - i \frac{1}{4} \frac{\dot{\rho}_t}{\rho_t} + \dot{\theta}_t \gamma_t (\rho_t - \rho_t^2 \zeta^2 - \xi^2) \right. \\ &\quad \left. - \left(\frac{\dot{\theta}_t s_t}{2} + j_t \gamma_t (D_\xi \xi - i \rho_t \zeta^2 + i) + k_t (\zeta^2 + \gamma_t^2 \xi^2) \right) \right) d\zeta. \end{aligned}$$

Furthermore, we have

$$\left(\frac{\rho_t}{\pi} \right)^{\frac{1}{2}} \int_{\mathbb{R}} e^{-\rho_t \zeta^2} d\zeta = 1, \quad \left(\frac{\rho_t}{\pi} \right)^{\frac{1}{2}} \int_{\mathbb{R}} \zeta^2 e^{-\rho_t \zeta^2} d\zeta = \frac{1}{2\rho_t}.$$

Hence, after performing the integration and realizing that the coefficients involving $\dot{\rho}_t$ cancel out, we obtain the formula

$$\begin{aligned} \mathcal{T} &= D_t + \dot{\theta}_t \gamma_t \left(\frac{\rho_t}{2} - \xi^2 \right) - \frac{\dot{\theta}_t s_t}{2} - j_t \gamma_t \left(D_\xi \xi + \frac{i}{2} \right) - k_t \left(\frac{1}{2\rho_t} + \gamma_t^2 \xi^2 \right) \\ &= D_t - \dot{\theta}_t \gamma_t \xi^2 - j_t \gamma_t \frac{\xi D_\xi + D_\xi \xi}{2} - k_t \left(\frac{1}{2\rho_t} + \gamma_t^2 \xi^2 \right) \\ &= D_t - \frac{k_t}{2\rho_t} - j_t \gamma_t \frac{\xi D_\xi + D_\xi \xi}{2} - \left(\dot{\theta}_t \gamma_t + k_t \gamma_t^2 \right) \xi^2, \end{aligned}$$

where in the second line we used $\gamma_t \rho_t = s_t$. □

5.2. Solving the transport equation. To produce an explicit solution of the transport equation $\mathcal{T}f = 0$ we define $(\lambda_t, \mu_t, \nu_t)$ such that

$$\lambda_t = \int_0^t \frac{k_s}{2\rho_s} ds, \quad \nu_t = 2 \int_0^t \frac{c_0^2}{c_s^2} \left(\dot{\theta}_s \gamma_s + k_s \gamma_s^2 \right) ds, \quad e^{\mu_t} = \frac{c_t}{c_0}. \quad (5.3)$$

Lemma 5.2. *The solution to $\mathcal{T}f = 0$ is given by*

$$f(t, \xi) = \exp \left(i\lambda_t + i\frac{\nu_t}{2} (e^{\mu_t} \xi)^2 \right) e^{\frac{\mu_t}{2}} f(0, e^{\mu_t} \xi). \quad (5.4)$$

Proof. 1. We recall that the self-adjoint operator $\frac{1}{2}(\xi D_\xi + D_\xi \xi)$ generates the semigroup of $(L^2\text{-unitary})$ dilations; that is, for F independent of t :

$$\left(D_\mu - \frac{\xi D_\xi + D_\xi \xi}{2} \right) U_\mu F = 0, \quad U_\mu F(\xi) = e^{\frac{\mu}{2}} F(e^\mu \xi). \quad (5.5)$$

We note that $e^{\mu_0} = 1$; moreover, thanks to (3.21), we have $\dot{\mu}_t = \partial_t \ln(c_t) = j_t \gamma_t$. Therefore, by the chain rule and (5.5), we have

$$U_{\mu_t}^{-1} \left(D_t - j_t \gamma_t \frac{\xi D_\xi + D_\xi \xi}{2} \right) U_{\mu_t} = D_t.$$

2. We deduce, using $U_{\mu_t}^{-1} \xi^2 U_{\mu_t} = e^{-2\mu_t} \xi^2$, that

$$U_{\mu_t}^{-1} \mathcal{T} U_{\mu_t} = D_t - \frac{k_t}{2\rho_t} - e^{-2\mu_t} \left(\dot{\theta}_t \gamma_t + k_t \gamma_t^2 \right) \xi^2 = D_t - \dot{\lambda}_t - \frac{1}{2} \dot{\nu}_t \xi^2 = e^{i\lambda_t + i\nu_t \frac{\xi^2}{2}} D_t e^{-i\lambda_t - i\nu_t \frac{\xi^2}{2}},$$

where we used the relations (5.3) for λ_t and ν_t . For the formula (5.5) for U_μ , we deduce that the solution to $\mathcal{T}f = 0 = \mathcal{T}U_{\mu_t} e^{i\lambda_t + i\frac{\nu_t}{2}\xi^2} g$ is $D_t g = 0$ and hence

$$f(t, \cdot) = U_{\mu_t} e^{i\lambda_t + i\frac{\nu_t}{2}\xi^2} f(0, \cdot).$$

Since $e^{\mu_t} = c_t/c_0$, we conclude that

$$f(t, \xi) = e^{\frac{\mu_t}{2}} \exp \left(i\lambda_t + i\frac{\nu_t}{2} (e^{\mu_t} \xi)^2 \right) f(0, e^{\mu_t} \xi).$$

This completes the proof. \square

5.3. Dispersive estimate. In this section, we study the L^∞ -decay of the leading order solution of (3.3), $\psi_0 = \mathfrak{U}_t a_0$, where a_0 takes the form prescribed by (4.3) and (5.4):

$$a_0 = e^{i\lambda_t} \tilde{a}_0 \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad \tilde{a}_0(t, \xi, \zeta) = \left(\frac{\rho_t}{4\pi} \right)^{\frac{1}{4}} e^{\frac{\mu_t}{2}} \exp \left(i\frac{\nu_t}{2} (e^{\mu_t} \xi)^2 - \frac{\rho_t}{2} \zeta^2 \right) \hat{f}(e^{\mu_t} \xi), \quad (5.6)$$

where $\hat{f}(\xi)$ denotes the Fourier transform of a function $f(z_1)$.

Lemma 5.3. (i) *With $\psi_0 = \mathfrak{U}_t a_0$ and a_0 given by (5.6), we have whenever $B_t \neq 0$ or $\nu_t \neq 0$:*

$$\psi_0(t, z) = e^{i\lambda_t} U_t (G_t *_1 f)(e^{-\mu_t} z_1, e^{\mu_t} z_2) \begin{bmatrix} 1 \\ -1 \end{bmatrix},$$

where $*_1$ denotes convolution with respect to the first variable and

$$G_t(e^{-\mu_t} z_1, e^{\mu_t} z_2) = \left(\frac{\rho_t}{4\pi} \right)^{\frac{1}{4}} \left(\frac{e^{\mu_t}}{Q_t} \right)^{\frac{1}{2}} e^{-\frac{1}{2}\rho_t z_2^2} e^{-\frac{1}{2}Q_t^{-1}(z_1 + is_t z_2)^2}, \quad Q_t = s_t \gamma_t - ie^{2\mu_t} \nu_t. \quad (5.7)$$

We recall that $s_t = \gamma_t \rho_t$.

(ii) In particular, there exists $C > 0$ such that as long as $\nu_t \neq 0$:

$$\sup_{z \in \mathbb{R}^2} |\psi_0(t, z)| \leq C \min \left(\frac{\|f\|_{L^1}}{|\nu_t|^{1/2}}, \|\hat{f}\|_{L^1} \right). \quad (5.8)$$

Proof. We recall that $\mathfrak{U}_t = U_t \mathcal{V}_t$ and we first write a formula for $\mathcal{V}_t \tilde{a}_0$. We have:

$$\mathcal{V}_t \tilde{a}_0(z) = \left(\frac{\rho_t}{4\pi} \right)^{\frac{1}{4}} e^{\frac{\mu_t}{2}} \int_{\mathbb{R}} e^{iz_1 \xi} \exp \left(i \frac{\nu_t}{2} (e^{\mu_t} \xi)^2 - \frac{\rho_t}{2} (z_2 + \gamma_t \xi)^2 \right) \hat{f}(e^{\mu_t} \xi) \frac{d\xi}{\sqrt{2\pi}}.$$

Hence,

$$\begin{aligned} \mathcal{V}_t \tilde{a}_0(e^{\mu_t} z_1, \frac{z_2}{e^{\mu_t}}) &= \left(\frac{\rho_t}{4\pi} \right)^{\frac{1}{4}} \int_{\mathbb{R}} e^{ie^{\mu_t} z_1 \xi} \exp \left(i \frac{\nu_t}{2} (e^{\mu_t} \xi)^2 - \frac{\rho_t e^{-2\mu_t}}{2} (z_2 + \gamma_t e^{\mu_t} \xi)^2 \right) \hat{f}(e^{\mu_t} \xi) \frac{e^{\frac{\mu_t}{2}} d\xi}{\sqrt{2\pi}} \\ &= \left(\frac{\rho_t}{4\pi} \right)^{\frac{1}{4}} \int_{\mathbb{R}} e^{iz_1 \xi} \exp \left(i \frac{\nu_t}{2} \xi^2 - \frac{\rho_t e^{-2\mu_t}}{2} (z_2 + \gamma_t \xi)^2 \right) \hat{f}(\xi) \frac{e^{-\frac{\mu_t}{2}} d\xi}{\sqrt{2\pi}}. \end{aligned} \quad (5.9)$$

The left-hand side involves the inverse Fourier transform in z_1 of a product and may therefore be written as the convolution

$$\mathcal{V}_t \tilde{a}_0(e^{\mu_t} z_1, z_2) = (f *_1 G_t)(z_1, e^{\mu_t} z_2) \quad (5.10)$$

where the Gaussian G_t is given by

$$G_t(z) = \left(\frac{\rho_t}{4\pi} \right)^{\frac{1}{4}} e^{\frac{-\mu_t}{2}} \int_{\mathbb{R}} e^{iz_1 \xi} \exp \left(i \frac{\nu_t}{2} \xi^2 - \frac{\rho_t e^{-2\mu_t}}{2} (z_2 + \gamma_t \xi)^2 \right) \frac{d\xi}{\sqrt{2\pi}}, \quad (5.11)$$

so that

$$\begin{aligned} G_t(z_1, e^{\mu_t} z_2) &= \left(\frac{\rho_t}{4\pi} \right)^{\frac{1}{4}} e^{\frac{-\mu_t}{2}} e^{-\frac{1}{2} \rho_t z_2^2} \int_{\mathbb{R}} e^{-\frac{1}{2} \tilde{Q}_t \xi^2} e^{i\xi(z_1 + i\rho_t \gamma_t e^{-\mu_t} z_2)} \frac{d\xi}{\sqrt{2\pi}}, \quad \tilde{Q}_t = \rho_t \gamma_t^2 e^{-2\mu_t} - i\nu_t \\ &= \left(\frac{\rho_t}{4\pi} \right)^{\frac{1}{4}} e^{\frac{-\mu_t}{2}} e^{-\frac{1}{2} \rho_t z_2^2} \tilde{Q}_t^{-\frac{1}{2}} e^{-\frac{1}{2} \tilde{Q}_t^{-1} (z_1 + i\rho_t \gamma_t e^{-\mu_t} z_2)^2}, \end{aligned} \quad (5.12)$$

as the Fourier transform of a Gaussian function when $\tilde{Q}_t \neq 0$. Since $Q_t = e^{2\mu_t} \tilde{Q}_t$, we find (5.7). Then (5.6) follows from applying U_t to $\mathcal{V}_t a_0$ using (5.6) and (5.10).

To prove the second part of the lemma, we first take L^∞ -norms on both sides of (5.10) and we apply Young's $L^1 - L^\infty$ convolution inequality (in the variable z_1). This produces:

$$\sup_{z \in \mathbb{R}} |\mathcal{V}_t \tilde{a}_0(z)| = \sup_{z_2 \in \mathbb{R}} \sup_{z_1 \in \mathbb{R}} |f *_1 G_t(z)| \leq \sup_{z_2 \in \mathbb{R}} \|f\|_{L^1} \sup_{z_1 \in \mathbb{R}} |G_t(z)| = \|f\|_{L^1} \sup_{z \in \mathbb{R}} |G_t(z)|. \quad (5.13)$$

The Gaussian $|G_t|$ attains its maximum at $z = 0$ because the real part of the quadratic form in z in (5.7) satisfies

$$\Re \left(\rho_t z_2^2 + \frac{1}{Q_t} (z_1 + i s_t z_2)^2 \right) = \frac{\rho_t}{(s_t \gamma_t)^2 + \tilde{\nu}_t^2} (\gamma_t z_1 - \tilde{\nu}_t z_2)^2 \geq 0, \quad \tilde{\nu}_t = e^{2\mu_t} \nu_t,$$

as we verify by an elementary computation using $\rho_t \gamma_t = s_t$. Using (5.13), we find

$$\sup_{z \in \mathbb{R}} |\mathcal{V}_t \tilde{a}_0(z)| \leq \|f\|_{L^1} |G_t(0)| = \left(\frac{\rho_t}{4\pi} \right)^{\frac{1}{4}} \left(\frac{e^{\mu_t}}{|Q_t|} \right)^{\frac{1}{2}} \|f\|_{L^1}.$$

From (5.9), we also obtain that $\sup_{z \in \mathbb{R}} |\mathcal{V}_t \tilde{a}_0(z)| \leq C \|\hat{f}\|_{L^1}$. To end up with (5.8), we observe that U_t preserves L^∞ -norms, that $|Q_t| \geq |\nu_t|$ and that c_t is bounded above and below for all times t . \square

When $Q_t = 0$, we have $\nu_t = 0$, $c_t = 1$, $\rho_t = r_t$ and the formulas (5.10) and (5.11) remain valid. But instead of being a Gaussian, G_t is now a multiple of the Dirac mass, and

$$G_t(z) = c_0^{1/2} \left(\frac{r_t}{4\pi} \right)^{1/4} \exp \left(-\frac{r_t}{2} z_2^2 \right) \delta_0(z_1), \quad \psi_0(t, z) = c_0^{1/2} \left(\frac{r_t}{4\pi} \right)^{1/4} U_t e^{-\frac{r_t}{2} z_2^2} f(z_1).$$

5.4. Dispersion-dependent functional setting. The solution of $\mathcal{T}f = 0$ with given initial condition admits an explicit expression as we saw in Lemma 5.2. In the analysis of the asymptotic expansion of $\psi(t, z)$ in powers of ε , we need to solve transport equations of the form $\mathcal{T}f = g$ with time-dependent source terms. To quantify the stability of the inverse transport operators and that of other relevant transforms, we introduce the following functional setting.

We recall the functional spaces \mathcal{S}_p were defined in section 4.1. To handle the time-dependence for $t \in [0, T]$ of the wavepackets, we introduce the spaces for $p \in \mathbb{N}$ and $k = \lfloor \frac{p}{2} \rfloor \in \mathbb{N}$ (i.e., $p = 2k$ or $p = 2k + 1$) defined for an interval $I \subset \mathbb{R}$ by

$$\mathbb{S}_p(I, \mathbb{R}^d, \mathbb{C}^q) = \cap_{r=0}^k C^r(I; \mathcal{S}_{p-2r}(\mathbb{R}^d, \mathbb{C}^q))$$

with norm given by the sum of the natural norms for the above spaces (see (5.16) below). The spaces are constructed so that any derivative in time corresponds to a loss of order 2 in the remaining variable.

To quantify the effects of dispersion, we define

$$\langle \nu \rangle_T = 1 + \sup_{0 \leq t \leq T} |\nu_t| \quad \text{and} \quad \varsigma = \langle \nu \rangle_T^{-\frac{1}{2}}. \quad (5.14)$$

We saw in Lemma 5.2 that dispersion resulted in a multiplication operator of the form $\phi_t(\xi) = e^{i\frac{1}{2}\nu(t)\xi^2}$. The operator $D_t\phi_t = \phi_t(D_t + \frac{1}{2}\nu'(t)\xi^2)$. This explains why the spaces \mathbb{S}_p are constructed so that both ∂_t and ξ^2 map \mathbb{S}_{p+2} to \mathbb{S}_p .

Similarly, $\partial_\xi\phi_t = \phi_t(\partial_\xi + i\nu(t)\xi)$, so that (the operator of multiplication by) ϕ_t is large as an operator on \mathbb{S}_1 when $\langle \nu \rangle_T$ is. Yet clearly, ϕ_t^2 is comparable to ϕ_t in the same sense. Since the construction of our wavepackets requires repeated application of operators of the form ϕ_t , we introduce scaled metrics on \mathcal{S}_p and \mathbb{S}_p so that application of ϕ_t results in a bounded operation independent of $\langle \nu \rangle_T$.

This is simply achieved by replacing $\mathbf{a}_\xi^* = -\partial_\xi + \xi$ by $\mathbf{a}_{\xi\varsigma}^* = -\varsigma\partial_\xi + \varsigma^{-1}\xi$ and endowing the spaces $\mathcal{S}_p(\mathbb{R})$ and $\mathbb{S}_p(\mathbb{R}^2)$ respectively with the norms

$$\|\psi\|_{p\varsigma}^2 = \|(\mathbf{a}_{\xi\varsigma}^*)^p \psi\|_0^2 \quad \text{and} \quad \|\psi\|_{p\varsigma}^2 = \sum_{j=0}^p \|(\mathbf{a}_{\xi\varsigma}^*)^j (\mathbf{a}_\xi^*)^{p-j} \psi\|_0^2. \quad (5.15)$$

We call $\mathcal{S}_{p\varsigma}$ the spaces \mathcal{S}_p endowed with these dispersion-scaled norms.

The space $\mathbb{S}_p(I, \mathbb{R}^d, \mathbb{C})$ are similarly endowed for $d = 1, 2$ with the norms (and called $\mathbb{S}_{p\varsigma}$)

$$\|\psi\|_{p\varsigma}^2 = \sup_{t \in I} \sum_{r=0}^{\lfloor \frac{p}{2} \rfloor} \|(\mathbf{a}_{\xi\varsigma}^*)^{p-2r} \partial_t^r \psi\|_0^2 \quad \text{and} \quad \|\psi\|_{p\varsigma}^2 = \sup_{t \in I} \sum_{r=0}^{\lfloor \frac{p}{2} \rfloor} \sum_{j=0}^{p-2r} \|(\mathbf{a}_{\xi\varsigma}^*)^j (\mathbf{a}_\xi^*)^{p-2r-j} \partial_t^r \psi\|_0^2. \quad (5.16)$$

Spaces of vector-valued functions are similarly constructed componentwise.

Note that $\varsigma^p \|f\|_p \lesssim \|f\|_{p\varsigma} \lesssim \varsigma^{-p} \|f\|_p$. Here and below, we use the notation $a \lesssim b$ to mean the existence of a ς -independent constant C such that $a \leq Cb$. Here is a number of dispersion-dependent estimates we will be using.

Lemma 5.4. *All operator bounds below are meant to be ς -independent bounds.*

Any operator $B \in \{\varsigma D_\xi, \varsigma^{-1}\xi, D_\zeta, \zeta\}$ is bounded from $\mathcal{S}_{p+1,\varsigma}$ to $\mathcal{S}_{p\varsigma}$ and from $\mathbb{S}_{p+1,\varsigma}$ to $\mathbb{S}_{p\varsigma}$. The operator ∂_t is bounded from $\mathbb{S}_{p+2,\varsigma}$ to $\mathbb{S}_{p\varsigma}$.

Let $\phi(\xi) = e^{i\frac{1}{2}\nu\xi^2}$ with $|\nu| \lesssim \langle\nu\rangle_T$. Then the operator of multiplication by $\phi(\xi)$ is bounded from $\mathcal{S}_{p\varsigma}$ to itself.

Let $\phi(t, \xi) = e^{i\frac{1}{2}\nu(t)\xi^2}$ with $\sup_{t \in \mathbb{I}} |\nu(t)| \lesssim \langle\nu\rangle_T$ and $\sup_{t \in \mathbb{I}} |\nu^{(j)}(t)| \leq C_j$ for $j \geq 1$. Then the operator of multiplication by $\phi(t, \xi)$ is bounded from $\mathbb{S}_{p\varsigma}$ to itself.

Proof. The first statement comes from the construction of the spaces since $\mathfrak{a}_{\xi\varsigma}^*$ controls $\varsigma\partial_\xi$ and $\varsigma^{-1}\xi$ in the sense that

$$\|\mathfrak{a}_{\xi\varsigma}^* \psi\|_0^2 = \varsigma^2 \|\partial_\xi \psi\|_0^2 + \varsigma^{-2} \|\xi \psi\|_0^2 + \|\psi\|_0^2.$$

One obtains from (5.16) for $p \geq 2$ that $\|\partial_t \psi\|_{p\varsigma}^2 \lesssim \|\psi\|_{p-2,\varsigma}^2$ and hence the bound on ∂_t .

Consider now the operator of multiplication by $\phi(\xi)$ in one dimension $d = 1$. We wish to show that

$$\|\phi \psi\|_{p\varsigma}^2 = \|(\mathfrak{a}_{\xi\varsigma}^*)^p (\phi \psi)\|_0^2 \lesssim \|(\mathfrak{a}_{\xi\varsigma}^*)^p \psi\|_0^2 = \|\psi\|_{p\varsigma}^2.$$

This holds when $p = 0$. Assume it holds for $p - 1 \geq 0$. Then

$$\|\phi \psi\|_{p\varsigma}^2 = \|(\mathfrak{a}_{\xi\varsigma}^*)^{p-1} (\mathfrak{a}_{\xi\varsigma}^*) (\phi \psi)\|_0^2 = \|(\mathfrak{a}_{\xi\varsigma}^*)^{p-1} (\phi \psi_1)\|_0^2 \lesssim \|\psi_1\|_{p-1,\varsigma}^2$$

by induction hypothesis, where $\psi_1(\xi) = (\mathfrak{a}_{\xi\varsigma}^* - i\varsigma\nu\xi)\psi$. By construction of the functional spaces and the above result for the operator $\varsigma^{-1}\xi$ knowing that $\varsigma|\nu| \lesssim \varsigma^{-1}$, we find $\|\psi_1\|_{p-1,\varsigma}^2 \lesssim \|\psi\|_{p,\varsigma}^2$ and the result is proved. The same proof applies in two dimensions $d = 2$ as well using the norm for $\mathcal{S}_p(\mathbb{R}^2)$ in (5.15) and that the commutator $[\mathfrak{a}_\zeta^*, \phi] = 0$.

The proof in the time-dependent setting uses that

$$\partial_t^r (\phi \psi) = \phi \left(\partial_t + i\frac{1}{2}\nu'(t)\xi^2 \right)^r \psi = \phi \psi_r, \quad \psi_r = \left(\sum_{j=0}^r \nu_j(t) \xi^{2(r-j)} \partial_t^j \right) \psi$$

for smooth and bounded functions $\nu_j(t)$ independent of $\langle\nu\rangle_T$ by assumption on $\nu(t)$. Assume dimension $d = 1$ as $d = 2$ is treated similarly. Using that $\xi^{2(r-j)} \partial_t^j$ maps $\mathbb{S}_{p\varsigma}$ to $\mathbb{S}_{p-2r,\varsigma}$ for each $0 \leq j \leq r$ and the bounds proved above in the time-independent setting, we find

$$\|(\mathfrak{a}_{\xi\varsigma}^*)^{p-2r} \partial_t^r \phi \psi\|_0 = \|(\mathfrak{a}_{\xi\varsigma}^*)^{p-2r} \phi \psi_r\|_0 \lesssim \|\psi_r\|_{p-2r,\varsigma} \lesssim \|\psi\|_{p\varsigma}.$$

The explicit expression of the norms in (5.16) and the above estimate conclude the proof of the lemma. \square

5.5. Stability of the transport and other operators. We then have the following stability result for the transport equation $\mathcal{T}f = g$. We recall that $\varsigma = \langle\nu\rangle_T^{-\frac{1}{2}}$.

Lemma 5.5. *Let $p \in \mathbb{N}$. The solution $f_0(t, \xi)$ of $\mathcal{T}f_0 = g$ on $[0, T] \times \mathbb{R}$ with initial condition $f_0(0, \xi) = \hat{f}(\xi) \in \mathcal{S}_{p\varsigma}(\mathbb{R}, \mathbb{C})$ and source term $g \in \mathbb{S}_{p\varsigma}(\mathbb{R}, \mathbb{C})$ satisfies the estimate*

$$\|f_0\|_{p\varsigma} \leq C_p (\|\hat{f}\|_{p\varsigma} + \langle T \rangle \|g\|_{p\varsigma})$$

where $\langle T \rangle = 1 + T$ and C_p is independent of T .

Proof. We first adapt Lemma 5.2 to handle volume sources and define

$$\lambda_{s,t} = \int_s^t \frac{k_\tau}{2\rho_\tau} d\tau, \quad \tilde{\nu}_{s,t} = 2 \int_s^t \frac{c_\tau^2}{c_\tau^2} (\dot{\theta}_\tau \gamma_\tau + k_\tau \gamma_\tau^2) d\tau, \quad e^{\mu_{s,t}} = \frac{c_t}{c_s},$$

to obtain that the solution to $\mathcal{T}f = 0$ with initial condition $f(s, \cdot)$ is given by

$$f(t, \xi) = \exp \left(i\lambda_{s,t} + i\frac{\tilde{\nu}_{s,t}}{2} \xi^2 \right) e^{\mu_{s,t}/2} f(s, e^{\mu_{s,t}} \xi). \quad (5.17)$$

By an application of the Duhamel principle, the solution to $\mathcal{T}f = g$ for $0 \leq t \leq T$ with $f(0, \xi) = 0$ is thus given explicitly by

$$f(t, \xi) = i \int_0^t \exp \left(i\lambda_{s,t} + i\frac{1}{2} \tilde{\nu}_{s,t} \xi^2 \right) e^{\mu_{s,t}/2} g(s, e^{\mu_{s,t}} \xi) ds.$$

Consider the solution in (5.17) at a fixed time s with the explicit time-dependence $\lambda(t) = \lambda_{s,t}$, $e^{\mu(t)} = e^{\mu_{s,t}}$ and $\nu(t) = \tilde{\nu}_{s,t}$ to simplify notation. We prove the lemma for the operator

$$f(t, \xi) \rightarrow e^{i\lambda(t)} e^{\frac{\mu(t)}{2}} e^{i\frac{1}{2}\nu(t)\xi^2} f(t, e^{\mu(t)} \xi).$$

The term $e^{i\lambda(t)}$ generates a smooth in time modulation. Since $\mu(t)$ is smooth and bounded above and below by positive constants independent of t , $e^{\mu(t)/2}$ is also smooth. The operator $f(t, \xi) \rightarrow e^{\frac{\mu(t)}{2}} e^{i\lambda(t)} f(t, \xi)$ is therefore bounded in the norms $\|\cdot\|_{p\zeta}$.

All other time dependent coefficients are smooth and uniformly bounded in time independent of T except for $\nu(t)$ that may grow linearly with $\nu'(t)$ uniformly bounded. Consider the operator $f(t, \xi) \mapsto h(t, \xi) = f(t, e^{\mu(t)} \xi)$ with $\hat{f} \in \mathbb{S}_{p\zeta}$. Plugging $h(t, \xi)$ into the definition of the norms (5.16), we directly obtain that $\|h\|_{p\zeta} \lesssim \|f\|_{p\zeta}$.

The final transformation (with $\nu(t)$ replaced by $\nu(t)e^{-2\mu(t)}$)

$$f(t, \xi) \mapsto h(t, \xi) = f(t, \xi) e^{i\frac{1}{2}\nu(t)\xi^2}$$

was analyzed in Lemma 5.4. This concludes the analysis of the map from $\hat{f}(\xi)$ to $f_0(t, \xi)$.

The volume source term $g(t, \xi)$ is treated similarly by the Duhamel principle, with an additional possible integration in time that provides the extra multiplication by $\langle T \rangle$. This concludes the proof of the lemma. \square

We conclude this section with a summary of the operators we introduced to construct approximations of the Dirac equation and some estimates they satisfy.

The terms in T_0 and T_1 that contribute to the construction of the leading term $a_0(t, \xi, \zeta)$ in the formal expansion $a = \sum_{j \geq 0} \varepsilon^{\frac{j}{2}} a_j$ were given in Lemma 3.2. Constructing higher-order terms a_j and proving convergence results require estimates on the operators T_j , which are constructed as in the proof of Lemma 3.2 and given explicitly by

$$T_1 = D_t + \mathcal{A} + \tilde{T}_1, \quad T_j = \tilde{T}_j, \quad j \geq 2,$$

with $\mathfrak{U}_t^* D_t \mathfrak{U}_t = D_t + \mathcal{A}$ where

$$\mathcal{A} := -\frac{1}{2}(\dot{\varphi}_t \sigma_2 + \dot{\theta}_t(-s_t \sigma_1 + c_t \sigma_3) - \dot{\theta}_t(-D_\xi D_\zeta - \zeta \xi + \gamma_t(\xi^2 - D_\zeta^2))) + \dot{\gamma}_t \xi D_\zeta,$$

and for $j \geq 1$,

$$\tilde{T}_j = \sum_{|\alpha|=j+1} \mathcal{V}_t^* z^\alpha \mathcal{V}_t \nu_\alpha \cdot \sigma, \quad \mathbb{C}^3 \ni \nu_\alpha = \frac{1}{\alpha!} \tilde{R}_{2,\varphi_t} \tilde{R}_{3,\theta_t} (R_{-\theta_t} \nabla)^\alpha h(y_t).$$

We verify for completeness that $\mathcal{V}_t^* z^\alpha \mathcal{V}_t = (\zeta - \gamma_t \xi)^{\alpha_2} (-1)^{\alpha_1} (D_\zeta + \gamma_t D_\xi)^{\alpha_1}$.

We recall that $(j_t, k_t) = (\nu_{111}, \nu_{021})$ while $\nu_{201} = 0$ was important to obtain an explicit expression for the transport solution since the transport operator involves no term of the form D_ξ^2 . We now summarize the estimates we will be needing.

Lemma 5.6. *We may write $T_0^{-1} = T_{01}^{-1} + T_{02}^{-1}$ and $T_1 = T_{11} + T_{12}$ such that the following operators are bounded with $\langle \nu \rangle_T$ -independent bounds:*

$$\begin{aligned} \zeta^{-1} T_{01}^{-1} : & \quad N_t^\perp \cap \mathbb{S}_{p+1,\zeta} \rightarrow \mathbb{S}_{p\zeta}, & T_{02}^{-1} : & \quad N_t^\perp \cap \mathbb{S}_{p\zeta} \rightarrow \mathbb{S}_{p\zeta}, \\ \zeta T_{11} : & \quad \mathbb{S}_{p+1,\zeta} \rightarrow \mathbb{S}_{p\zeta}, & T_{12} : & \quad \mathbb{S}_{p+2,\zeta} \rightarrow \mathbb{S}_{p\zeta}, \\ \zeta^{j+1} T_j : & \quad \mathbb{S}_{p+j+1,\zeta} \rightarrow \mathbb{S}_{p\zeta}, \quad j \geq 2, & \langle T \rangle^{-1} \mathcal{T}^{-1} : & \quad \mathbb{S}_{p\zeta} \rightarrow \mathbb{S}_{p\zeta}. \end{aligned}$$

Proof. From Lemma 4.2 and the decomposition

$$T_{01}^{-1} = Q \begin{bmatrix} 0 & 0 \\ 0 & -2\xi c_t(\rho_t \zeta + \partial_\zeta)^{-1}(\rho_t \zeta - \partial_\zeta)^{-1} \end{bmatrix} Q, \quad T_{02}^{-1} = T_0^{-1} - T_{01}^{-1},$$

and Lemma 5.4, we obtain the above first two bounds.

We define $T_1 = T_{11} + T_{12}$ with T_{11} the contribution that is linear in D_ξ while T_{12} accounts for the rest (no contribution in T_1 is quadratic in D_ξ). The above corresponding bounds then follow from Lemma 5.4 for quadratic expressions in D_ξ , D_ζ , ξ , and ζ . The same lemma is used to bound T_j for $j \geq 2$. The final estimate is a repeat of Lemma 5.5. \square

6. ASYMPTOTIC EXPANSION AND ERROR ESTIMATES

We recall that our objective is to construct approximations of solutions $\Psi(t, x)$ of the Dirac equation (1.1). Our first step was to perform a gauge transformation $\tilde{\Psi}(t, x) = e^{-i\chi(t,x)/\varepsilon} \Psi(t, x)$ replacing the Dirac operator \not{D} by $\tilde{\not{D}} + R$ in (2.10). Since R in (2.9) is a negligible perturbation to arbitrary order in ε , our second step was to look for wavepackets $\tilde{\Psi}(t, x)$ in the kernel of $\varepsilon D_t + \tilde{\not{D}}$.

Wavepackets in natural coordinates $\psi(t, z) = \varepsilon^{\frac{1}{2}} S \tilde{\Psi}(t, z)$ with the scaling S defined in (3.1) then solve $L\psi = 0$ in (3.3). Further transformations resulted in the definition of $a(\xi, \zeta) = \mathfrak{U}_t^* \psi(\xi, \zeta)$, where $\mathfrak{U}_t = U_t \mathcal{V}_t$ with U_t defined in (3.8) and \mathcal{V}_t in (3.25). The problem $L\psi = 0$ is then equivalent to $Ta = 0$ for $T = \mathfrak{U}_t^* L \mathfrak{U}_t$. Writing $L = \sum_{j \geq 0} \varepsilon^{\frac{j}{2}} L_j$ with L_j presented in (3.4)-(3.6), we have a corresponding expansion $T = \sum_{j \geq 0} \varepsilon^{\frac{j}{2}} T_j$ with $T_j = \mathfrak{U}_t^* L_j \mathfrak{U}_t$.

6.1. Construction of the asymptotic wavepacket. Using the notation recalled above, we now construct approximations $a^J = \sum_{j=0}^J \varepsilon^{\frac{j}{2}} a_j$ of $a(t, \xi, \zeta)$ solution of $Ta = 0$. Plugging the expansion for a^J in the equation $Ta^J = 0$, using $T = \sum_{j \geq 0} \varepsilon^{\frac{j}{2}} T_j$, and equating like

powers of ε gives the sequence of equations (3.29), which we recall here:

$$\sum_{k=0}^j T_k a_{j-k} = 0, \quad 0 \leq j \leq J. \quad (6.1)$$

We solve these equations in turn.

The leading order equation $T_0 a_0 = 0$ combined with Lemma 4.2 shows that $a_0(t, \xi, \zeta) = f_0(t, \xi) \phi_t(\zeta)$ with ϕ_t defined in (4.2). The compatibility condition for the next equation $T_0 a_1 = -T_1 a_0$ implies $\mathcal{T} f_0 = 0$ with the transport operator \mathcal{T} defined in Lemma 5.2. For a fixed initial condition $f_0(0, \xi) = \hat{f}(\xi)$, Lemma 5.5 provides a unique solution $f_0(t, \xi)$ of $\mathcal{T} f_0 = 0$ and a leading term $a_0(t, \xi, \zeta) = f_0(t, \xi) \phi_t(\zeta)$.

Consider next the construction of a_1 , solution to $T_0 a_1 + T_1 a_0 = 0$. It is given according to Lemma 4.2 by

$$a_1 = -T_0^{-1} T_1 a_0 + f_1 \phi_t$$

with f_1 arbitrary at this stage. We will consider regularity properties in a lemma below.

We now extend the construction to higher-order approximations. Let $j \geq 2$ and assume a_k for $0 \leq k \leq j-1$ constructed except for f_{j-1} . In order to define a_j , we impose the compatibility condition

$$\left(\sum_{k=1}^j T_k a_{j-k}, \phi_t \right)_2 = 0 \quad \text{or equivalently} \quad \mathcal{T} f_{j-1} = g_{j-1}, \quad g_{j-1} = - \left(\sum_{k=2}^j T_k a_{j-k}, \phi_t \right)_2.$$

We recall that $(\cdot, \cdot)_2$ is the standard inner product in $L^2(\mathbb{R}, \mathbb{C}^2)$ in the ζ variable. By lemma 5.5, this is a well-posed transport equation for f_{j-1} . Then, by lemma 4.2,

$$a_j = T_0^{-1} \left(- \sum_{k=1}^j T_k a_{j-k} \right) + f_j \phi_t.$$

This constructs a_j for $0 \leq j \leq J$ iteratively and we set $f_J = 0$ for concreteness. This concludes the construction of the approximation

$$a^J = \sum_{j=0}^J \varepsilon^{\frac{j}{2}} a_j$$

of formal order $\varepsilon^{\frac{J+1}{2}}$ of a solution of $Ta = 0$.

Note that the initial conditions for $a_j(0, \xi, \zeta)$ for $j \geq 1$ are defined implicitly by the above construction. Only $\hat{f}(\xi)$ in the initial condition $a_0(0, \xi, \zeta) = \hat{f}(\xi) \phi_t(\zeta)$ is prescribed. Our construction aims to propagate wavepackets that belong to an appropriate (non-dispersive) branch of continuous spectrum of $\tilde{\mathcal{D}}$. The initial condition for a^J ensures that the latter belongs to that branch with sufficient accuracy.

The terms of the above expansion satisfy the following estimate:

Lemma 6.1. *Let a_j be constructed as above for $0 \leq j \leq J$ (with $f_J = 0$) and for $p \in \mathbb{N}$, let $\hat{f} \in \mathcal{S}_{p+3J}$. Then we have the estimates for $0 \leq j \leq J-1$,*

$$\|a_j\|_p \leq C_{pJ} (\langle T \rangle \langle \nu \rangle_T^3)^j \langle \nu \rangle_T^p \|\hat{f}\|_{p+3j}, \quad \|a_J\|_p \leq C_{pJ} (\langle T \rangle \langle \nu \rangle_T^3)^{J-1} \langle \nu \rangle_T^{p+1} \|\hat{f}\|_{p+3J},$$

with constants C_{pJ} independent of T and $\langle \nu \rangle_T$.

Proof. Consider $0 < j < J$. The estimate for a_J is a bit different since $f_J = 0$.

We first observe that by construction of the terms a_j for $1 \leq j \leq J-1$,

$$a_j = -T_0^{-1} \sum_{k=0}^{j-1} T_{j-k} a_k - \phi_t(\zeta) \mathcal{T}^{-1} \sum_{k=0}^{j-1} (T_{j+1-k} a_k, \phi_t)_2. \quad (6.2)$$

We wish to prove by induction that

$$\|a_j\|_{p\varsigma} \lesssim (\langle T \rangle \varsigma^{-3})^j \|a_0\|_{p+3j, \varsigma}, \quad \|a_J\|_{p\varsigma} \lesssim (\langle T \rangle \varsigma^{-3})^{J-1} (\varsigma \|a_0\|_{p+3J, \varsigma} + \|a_0\|_{p+3J-1, \varsigma}). \quad (6.3)$$

Using that $\|a_0\|_{p\varsigma} = \|f_0 \phi_t\|_{p\varsigma} \lesssim \|\hat{f}\|_{p\varsigma}$ thanks to Lemma 5.5, then $\varsigma^p \|f\|_p \lesssim \|f\|_{p\varsigma} \lesssim \varsigma^{-p} \|f\|_p$ and $\varsigma = \langle \nu \rangle_T^{-\frac{1}{2}}$ thus provide the results stated in the lemma. It remains to verify (6.3).

From (6.2) for $j = 1$, we obtain $a_1 = -T_0^{-1} T_1 a_0 - \phi_t \mathcal{T}^{-1} (T_2 a_0, \phi_t)_2$ so that using the result of Lemma 5.6, we find the second term to be the least regular and

$$\|a_1\|_{p\varsigma} \lesssim \langle T \rangle \varsigma^{-3} \|a_0\|_{p\varsigma+3}$$

so that (6.3) holds when $j = 1$. Assume it holds for $j-1 \geq 1$. Then using (6.2) and Lemma 5.6, we find

$$\|a_j\|_{p\varsigma} \lesssim \langle T \rangle \sum_{k=0}^{j-1} \varsigma^{-(j+2-k)} \|a_k\|_{p+j+2-k, \varsigma} \lesssim \langle T \rangle \sum_{k=0}^{j-1} \varsigma^{-(j+2-k)} (\langle T \rangle \varsigma^{-3})^k \|a_0\|_{p+2k+j+2, \varsigma},$$

which is largest when $k = j-1$ and provides the sought estimate when $0 \leq j < J$. It remains to consider the term

$$a_J = -T_0^{-1} \sum_{k=0}^{J-1} T_{J-k} a_k = -T_0^{-1} \Pi_t \sum_{k=0}^{J-1} T_{J-k} a_k = -T_0^{-1} \Pi_t T_1 a_{J-1} - T_0^{-1} \Pi_t \sum_{k=0}^{J-2} T_{J-k} a_k,$$

by construction of the wavepackets, where Π_t projects onto N_t^\perp . This operator has a smooth Schwartz kernel in (t, ζ, ζ') that is independent of ξ . Therefore Π_t is bounded from $\mathbb{S}_{p\varsigma}$ to itself with ς -independent bound.

Since T_0^{-1} is bounded from $N^\perp \cap \mathbb{S}_{p+1, \varsigma}$ to $\mathbb{S}_{p\varsigma}$, we find for $0 \leq k \leq J-2$

$$\|T_0^{-1} \Pi_t T_{J-k} a_k\|_{p\varsigma} \lesssim \|a_k\|_{p+2+J-k, \varsigma} \lesssim (\langle T \rangle \varsigma^{-3})^k \|a_0\|_{p+2+J+2k, \varsigma} \lesssim (\langle T \rangle \varsigma^{-3})^{J-1} \|a_0\|_{p+3J-1, \varsigma}$$

since $2 + J + 2k \leq 3J - 1$.

It remains to consider $T_0^{-1} \Pi_t T_1 a_{J-1}$. We decompose $T_0^{-1} \Pi_t T_1 = A + B$ with $A = T_{01}^{-1} \Pi_t T_{11} + T_{02}^{-1} \Pi_t T_{12}$ with bound $\langle T \rangle$ from $\mathbb{S}_{p+2, \varsigma} \rightarrow \mathbb{S}_{p\varsigma}$ and $B = T_{01}^{-1} \Pi_t T_{12} + T_{02}^{-1} \Pi_t T_{11}$ with bound $\langle T \rangle \varsigma$ from $\mathbb{S}_{p+3, \varsigma} \rightarrow \mathbb{S}_{p\varsigma}$ as per Lemma 5.6. This implies

$$\|a_J\|_{p\varsigma} \lesssim (\langle T \rangle \varsigma^{-3})^{J-1} (\varsigma \|a_0\|_{p+3J, \varsigma} + \|a_0\|_{p+3J-1, \varsigma})$$

as was to be shown. Thus (6.3) holds and this concludes the proof of the lemma. \square

6.2. Main approximation result. We can now state our main result.

Theorem 6.2. *Let Ψ be the solution on $[0, T] \times \mathbb{R}^2$ of $(\varepsilon D_t + \not{D})\Psi = 0$ with initial condition $\Psi(0, x) = \Psi^J(0, x)$ where $\Psi^J(t, x) = e^{\frac{i}{\varepsilon} \chi(t, x)} (\varepsilon^{-\frac{1}{2}} S^{-1} \mathcal{U}_t a^J)(t, x)$ with a^J constructed in section 6.1 based on an initial condition $\hat{f}(\xi) \in \mathcal{S}_{3J+2}(\mathbb{R}, \mathbb{C})$.*

Then there exists $C_J > 0$ independent of T such that for every $\varepsilon \in (0, 1]$,

$$\sup_{t \in [0, T]} \|\Psi - \Psi^J\|_{L^2(\mathbb{R}^2, \mathbb{C}^2)} \leq C_J (\langle T \rangle \langle \nu \rangle_T^3)^J \varepsilon^{\frac{J}{2}}.$$

We recall that $\langle \nu \rangle_T$ is defined in (5.14) and $\langle T \rangle = 1 + T$. In particular, the approximation error is controlled so long as $\varepsilon^{\frac{1}{2}} \langle T \rangle \langle \nu \rangle_T^3 \ll 1$. When $\langle \nu \rangle_T$ is uniformly bounded, errors are controlled up to times $T \ll \varepsilon^{-\frac{1}{2}}$ as in [BBD⁺21]. However, when $\langle \nu \rangle_T$ is of order T , then errors are controlled up to times $T \ll \varepsilon^{-\frac{1}{8}}$.

Note that such results are qualitatively reasonable: in the presence of dispersion, it becomes more difficult to control spatial moments (necessitated by the Taylor expansion of the coefficients h) of the solution.

The rest of the section is devoted to the proof of this theorem. It is based on a similar approximation result in the local variables $z = (z_1, z_2)$, which we state as a result of independent interest.

Proposition 6.3. *Let ψ be the solution on $[0, T] \times \mathbb{R}^2$ of $L\psi = 0$ with initial condition $\psi(0, z) = \psi^J(0, z)$ where $\psi^J(t, z) = \mathfrak{U}_t a^J(t, z)$ with a^J constructed in section 6.1 based on an initial condition $\hat{f}(\xi) \in \mathcal{S}_{3J+2}(\mathbb{R}, \mathbb{C})$.*

Then there exists $C_J > 0$ independent of T such that for every $\varepsilon \in (0, 1]$,

$$\sup_{t \in [0, T]} \|\psi - \psi^J\|_{L^2(\mathbb{R}^2, \mathbb{C}^2)} \leq C_J (\langle T \rangle \langle \nu \rangle_T^3)^J \varepsilon^{\frac{J}{2}}.$$

Proof. Solutions of the local problems $L\psi = 0$ and $Ta = 0$ are equivalent via the relation $\psi = \mathfrak{U}_t a$. We now show that $\psi^J = \sum_{j=0}^J \varepsilon^{\frac{j}{2}} \psi_j$ with $\psi_j = \mathfrak{U}_t a_j$ approximately solves $L\psi = 0$ when $J \geq 1$. Define for $J \geq 1$ the remainder operator

$$L_{\geq J} = \varepsilon^{-\frac{J}{2}} \left(L - \sum_{j=0}^{J-1} \varepsilon^{\frac{j}{2}} L_j \right)$$

and the reduced operator $\tilde{L}_{\geq J} = L_{\geq J} - \delta_{1,J} D_t$ involving only the coefficients h .

We deduce from (6.1) that $\sum_{k=0}^j L_k \psi_{j-k} = 0$ for $0 \leq j \leq J$ and hence

$$L\psi^J = \sum_{j=0}^J \varepsilon^{\frac{j}{2}} L\psi_j = \sum_{j=0}^J \left(\sum_{k=0}^{J-j} \varepsilon^{\frac{j+k}{2}} L_k \psi_j + \varepsilon^{\frac{J+1}{2}} L_{\geq J+1-j} \psi_j \right) = \varepsilon^{\frac{J+1}{2}} \sum_{j=0}^J L_{\geq J+1-j} \psi_j,$$

since

$$\sum_{j=0}^J \sum_{k=0}^{J-j} \varepsilon^{\frac{j+k}{2}} L_k \psi_j = \sum_{j=0}^J \sum_{l=j}^J \varepsilon^{\frac{l}{2}} L_{l-j} \psi_j = \sum_{l=0}^J \varepsilon^{\frac{l}{2}} \sum_{j=0}^l L_{l-j} \psi_j = 0.$$

We thus obtain

$$L\psi^J = \varepsilon^{\frac{J+1}{2}} \left(D_t \psi_J + \sum_{j=0}^J \tilde{L}_{\geq J+1-j} \psi_j \right) =: r_J. \quad (6.4)$$

We now derive a uniform bound in time of order $\varepsilon^{\frac{J+1}{2}}$ in $L^2(\mathbb{R}^2, \mathbb{C}^2)$ for r_J .

We observe that $\mathcal{V}_t \mathcal{S}_p(\mathbb{R}^d; \mathbb{C}^q) \mathcal{V}_t^* \cong \mathcal{S}_p(\mathbb{R}^d; \mathbb{C}^q)$ as spaces of functions (of z for $d = 2$). In other words, $\|g\|_p \cong \|\mathcal{V}_t^* g\|_p$ define equivalent norms for each p . The reason is that the above

spaces based on Hermite functions are invariant by conjugation by Fourier transforms as well as invariant under invertible linear transforms $(\xi, z_2) \mapsto (\xi, z_2 + \gamma_t \xi)$ of the base variables (uniformly in time since γ_t is bounded). Since γ_t is smooth, we also obtain the equivalence of the norms of $\mathcal{V}_t \mathbb{S}_p(\mathbb{R}^d; \mathbb{C}^q) \mathcal{V}_t^* \cong \mathbb{S}_p(\mathbb{R}^d; \mathbb{C}^q)$. Note that there is no meaningful notion of anisotropic space $\mathbb{S}_{p\kappa}$ in the variable z .

The regularity results of lemma 6.1 therefore apply to $\psi_j = \mathfrak{U}_t a_j$ since conjugation by U_t also preserves norms as the rotation angles θ_t and φ_t are smooth in t . As a consequence,

$$\begin{aligned} \|\psi_J\|_2 &\lesssim \langle T \rangle^{J-1} \langle \nu \rangle_T^{3J} \|\hat{f}\|_{3J+2}, \\ \|\psi_j\|_{J-j+2} &\lesssim (\langle T \rangle \langle \nu \rangle_T^3)^j \langle \nu \rangle_T^{J-j+2} \|\hat{f}\|_{J+2j+2} \lesssim \langle T \rangle^{J-1} \langle \nu \rangle_T^{3J} \|\hat{f}\|_{3J+2} \end{aligned} \quad (6.5)$$

for $0 \leq j \leq J-1$.

Since D_t is bounded from \mathbb{S}_{p+2} to \mathbb{S}_p , we obtain that $\|D_t \psi_J\|_0 \lesssim \langle T \rangle^{J-1} \|\hat{f}\|_{3J+2}$.

Using Taylor expansions of the coefficients for $1 \leq j \leq J$,

$$h(y_t + \sqrt{\varepsilon} z) = \sum_{|\alpha| \leq j} \frac{1}{\alpha!} \partial^\alpha h(y_t) \varepsilon^{\frac{|\alpha|}{2}} z^\alpha + \varepsilon^{\frac{j+1}{2}} \sum_{|\beta|=j+1} z^\beta R_\beta(y_t + \sqrt{\varepsilon} z),$$

with smooth functions R_β by regularity assumptions on κ and A , we observe that $\tilde{L}_{\geq j}$ has the same regularity properties as \tilde{L}_j . In particular, it maps \mathbb{S}_{p+j+1} to \mathbb{S}_p . Therefore, for $0 \leq j \leq J$, $\|\tilde{L}_{\geq J+1-j} \psi_j\|_0 \lesssim \|\psi_j\|_{J+2-j} \lesssim \langle T \rangle^{J-1} \langle \nu \rangle_T^{3J} \|\hat{f}\|_{3J+2}$ thanks to (6.5).

By definition (6.4), this shows that

$$\|r_J\|_0 \lesssim \varepsilon^{\frac{J+1}{2}} \langle T \rangle^{J-1} \langle \nu \rangle_T^{3J} \|\hat{f}\|_{3J+2}. \quad (6.6)$$

Define $\tilde{L} = L - \varepsilon^{\frac{1}{2}} D_t$, which we verify is self adjoint [Tha92]. We thus observe that $L(\psi^J - \psi) = r^J$ is equivalent to

$$(D_t + \varepsilon^{-\frac{1}{2}} \tilde{L})(\psi^J - \psi) = \varepsilon^{-\frac{1}{2}} r_J.$$

By unitarity of the above Dirac operator [Tha92], we obtain an error on $\|\psi^J - \psi\|_0$ of order $\langle T \rangle \varepsilon^{-\frac{1}{2}} \|r_J\|_0 \lesssim \varepsilon^{\frac{J}{2}} \langle T \rangle^J \langle \nu \rangle_T^{3J}$. This concludes the proof of the proposition. \square

Proof. (Theorem 6.2) Let $\tilde{\Psi}$ be the solution of $(\varepsilon D_t + \tilde{D})\tilde{\Psi} = 0$ with initial conditions $\tilde{\Psi}(0, x) = \tilde{\Psi}^J(t, x) = (\varepsilon^{-\frac{1}{2}} S^{-1} \mathfrak{U}_t a^J)(t, x)$. Since $\tilde{\Psi} = \varepsilon^{-\frac{1}{2}} S^{-1} \psi$ and $\tilde{\Psi}^J = \varepsilon^{-\frac{1}{2}} S^{-1} \psi^J$ for $\varepsilon^{-\frac{1}{2}} S^{-1}$ an L^2 -isometry, we directly deduce from Proposition 6.3 that

$$\sup_{t \in [0, T]} \|\tilde{\Psi} - \tilde{\Psi}^J\|_{L^2(\mathbb{R}^2, \mathbb{C}^2)} \leq C_J (\langle T \rangle \langle \nu \rangle_T^3)^J \varepsilon^{\frac{J}{2}}.$$

By construction, $\Psi^J(t, x) = e^{\frac{i}{\varepsilon} \chi(t, x)} \tilde{\Psi}^J(t, x) = e^{\frac{i}{\varepsilon} \chi(t, x)} \varepsilon^{-\frac{1}{2}} S^{-1} \psi^J(t, x)$ so that

$$(\varepsilon D_t + \tilde{D})\Psi^J = e^{\frac{i}{\varepsilon} \chi(t, x)} (\varepsilon D_t + \tilde{D} + R) \tilde{\Psi}^J = e^{\frac{i}{\varepsilon} \chi(t, x)} \varepsilon^{-\frac{1}{2}} S^{-1} [\varepsilon^{\frac{1}{2}} r_J + (\varepsilon^{\frac{1}{2}} S R) \psi^J],$$

where we used $\varepsilon D_t + \tilde{D} = S^{-1} L \varepsilon^{\frac{1}{2}} S$ and where r_J defined in (6.4) and estimated in (6.6).

By unitarity for $(\varepsilon D_t + \tilde{D})$, the statement of the theorem follows if we show that $S R \psi^J(t, z) = R(t, y_t + \sqrt{\varepsilon} z) \psi^J(t, z)$ satisfy a bound similar to r_J as in (6.6).

Since R vanishes on Ω_δ and is uniformly bounded on $\mathbb{R} \times \mathbb{R}^2$ by construction, we find that for every $p \geq 0$,

$$\|(1 + |z|^2)^{\frac{p}{2}} R(t, y_t + \sqrt{\varepsilon} z)\|_\infty \lesssim \varepsilon^{\frac{p}{2}}.$$

Therefore, thanks to (6.5),

$$\begin{aligned} \|R(t, y_t + \sqrt{\varepsilon} z) \psi^J\|_0 &\lesssim \sum_{j=0}^J \varepsilon^{\frac{j}{2}} \|R(t, y_t + \sqrt{\varepsilon} z) \psi_j\|_0 \lesssim \sum_{j=0}^J \varepsilon^{\frac{j+1}{2}} \|(1 + |z|^2)^{\frac{j-j+1}{2}} \psi_j\|_0 \\ &\lesssim \varepsilon^{\frac{J+1}{2}} \sum_{j=0}^J \|\psi_j\|_{J-j+1} \lesssim \varepsilon^{\frac{J+1}{2}} \langle T \rangle^{J-1} \langle \nu \rangle_T^{3J} \|\hat{f}\|_{3J+2}. \end{aligned}$$

Thus, $SR\psi^J$ satisfies the same estimate (6.6) as r_J . Since \mathcal{D} is self-adjoint, we conclude the proof of Theorem 6.2 by the same unitarity principle as in Proposition 6.3. \square

Proof. (Theorem 1.1) The above result with $J = 1$ also provides a proof of Theorem 1.1 when the gauge transformation is based on $\chi(t, x)$. Indeed, in this simplified setting with B constant and $\Delta\kappa = 0$, we observe that $a_0(t, \xi, \zeta)$ is given by (5.6) with $\lambda_t = 0$, $e^{\mu_t} = 1$, and $\tilde{\nu}_t = \nu_t = 2\gamma(\theta_t - \theta_0)$ with $\gamma = \frac{B}{1+B^2}$. This provides the expression for the kernel $g_t(z)$ in (1.8), which quantifies the dispersive effects while $\psi_{\theta_t, B}(z)$ captures all other effects in $\mathcal{U}_t a_0(z)$ as one readily verifies.

An additional error in $\Psi^1(t, x) = e^{\frac{i}{\varepsilon}\chi(t, x)} \varepsilon^{-\frac{1}{2}} S^{-1} \psi^1(t, x)$ comes from replacing $\chi(t, x)$ by its quadratic expansion $\chi_2(t, x)$ (called $\chi(t, x)$ in Theorem 1.1) in (1.9). In the z variables, we find

$$e^{i\frac{1}{\varepsilon}\chi(y_t + \sqrt{\varepsilon} z)} = e^{i\frac{1}{\varepsilon}\chi_2(y_t + \sqrt{\varepsilon} z)} + O(\varepsilon^{\frac{1}{2}})|z|^3.$$

This term is multiplied by $\psi^1 = \psi_0 + \sqrt{\varepsilon}\psi_1$ and yields an error of order $\varepsilon^{\frac{1}{2}}$ in the L^2 -sense for $(e^{\frac{i}{\varepsilon}\chi(t, x)} - e^{\frac{i}{\varepsilon}\chi_2(t, x)})\varepsilon^{-\frac{1}{2}} S^{-1} \psi^1(t, x)$ when $\psi_1 \in \mathbb{S}_0$ and when $|z|^3 \psi_0 \in \mathbb{S}_0$ as well. Both bounds hold as soon as $\psi_0 \in \mathbb{S}_3$, and hence when $\hat{f} \in \mathcal{S}_5$ as required for $J = 1$ in Theorem 6.2 with a bound

$$\| [e^{i\frac{1}{\varepsilon}\chi(y_t + \sqrt{\varepsilon} z)} - e^{i\frac{1}{\varepsilon}\chi_2(y_t + \sqrt{\varepsilon} z)}] \psi^1 \|_{\mathbb{S}_0} \leq C\varepsilon^{\frac{1}{2}}.$$

We therefore obtain an overall bound on $\Psi - e^{\frac{i}{\varepsilon}\chi_2(t, x)} \varepsilon^{-\frac{1}{2}} S^{-1} \psi^1(t, x)$ growing as $C\varepsilon^{\frac{1}{2}}$ on an interval $[0, T]$ with T fixed. Since $\sqrt{\varepsilon}\psi_1$ is also of order $\varepsilon^{\frac{1}{2}}$, then so is $\Psi - e^{\frac{i}{\varepsilon}\chi_2(t, x)} \varepsilon^{-\frac{1}{2}} S^{-1} \psi_0(t, x)$. The construction performed on $[0, T]$ clearly holds on $[-T, 0]$ as well. This completes the proof of Theorem 1.1. \square

7. NUMERICAL SIMULATIONS

In this section, we illustrate numerically the main effects of the magnetic field and the curved interface on the propagating wavepackets: (i) slowdown; (ii) Aharonov–Bohm phase-shift; (iii) dispersion.

The leading term in the asymptotic expansion provided by Theorem 6.2 is given by

$$\Psi_0(t, x) = \varepsilon^{-\frac{1}{2}} e^{\frac{i}{\varepsilon}\chi(t, x)} (\mathcal{U}_t a_0) \left(t, \frac{x - y_t}{\sqrt{\varepsilon}} \right), \quad \text{where:} \quad (7.1)$$

- $\mathcal{U}_t a_0(t, z) = U_t \mathcal{V}_t a_0(t, z)$ with U_t a spatial and spinorial rotation that does not quantitatively affect the amplitude landscape of the wavepacket;

- \mathcal{V}_t is a Fourier-like transform given explicitly in (3.25).
- $a_0(t, \xi, \zeta) = f_0(t, \xi)\phi_t(\zeta)$ with $f_0(t, \xi)$ given respectively by (5.4) and (4.2).

For concreteness, we work below with the Gaussian initial condition $f_0(0, \xi) = e^{-\frac{1}{2}\sigma\xi^2}$ for $\sigma > 0$. Thanks to (5.4) and (5.12), we obtain

$$\mathcal{V}_t a_0(t, z) = \left(\frac{\rho_t}{4\pi}\right)^{\frac{1}{4}} e^{\frac{1}{2}\mu_t} (Q_\sigma)^{-\frac{1}{2}} e^{i\lambda_t} e^{-\frac{1}{2}\rho_t z_2^2} e^{-\frac{1}{2}Q_\sigma^{-1}(z_1 + is_t z_2)^2} \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \quad (7.2)$$

with $Q_\sigma = e^{2\mu_t}\sigma + s_t\gamma_t - ie^{2\mu_t}\nu_t$. The various constants that appear in these formulas are collected as follows

$$\begin{aligned} r_t &= |\nabla\kappa(y_t)|, \quad \rho_t = \sqrt{r_t^2 + B_t^2}, \quad c_t = \frac{r_t}{\rho_t}, \quad s_t = \frac{B_t}{\rho_t}, \quad \gamma_t = \frac{B_t}{\rho_t^2}, \quad \dot{j}_t\gamma_t = \frac{d\ln c_t}{dt}, \\ k_t &= \frac{c_t}{2} \left(\partial_n B(y_t) - B_t \frac{\Delta\kappa(y_t)}{r_t} \right), \quad \lambda_t = \int_0^t \frac{k_s}{2\rho_s} ds, \quad e^{\mu_t} = \frac{c_t}{c_0}, \quad \nu_t = 2 \int_0^t \frac{c_0^2}{c_s^2} (\dot{\theta}_s \gamma_s + k_s \gamma_s^2) ds, \end{aligned}$$

and $n(y_t)$ and $\tau(y_t)$ are the normal and tangent vectors to Γ at y_t .

The rate of change of dispersion $\dot{\nu}_t$ is, up to the multiplicative constant $2\gamma_t e^{-2\mu_t}$ given by the two contributions $\dot{\theta}_t + \gamma_t k_t$, which we write explicitly as

$$\dot{\theta}_t + \gamma_t k_t = c_t \left(\epsilon_t K_t + \frac{1}{2} \frac{B \partial_n B - B^2 r_t^{-1} \Delta\kappa}{B^2 + r_t^2}(y_t) \right),$$

with $\epsilon_t = \pm 1$ when $\{\mp\kappa > 0\}$ is convex in a neighborhood of y_t . This provides an expression to assess how the geometry of κ and that of B combine to amplify or suppress dispersion.

7.1. Magnetic slowdown in constant B-field. The most visible impact of the magnetic field is the slowdown of the wavepacket: it propagates at speed $c_t = (1 + B_t^2)^{-\frac{1}{2}}$, which is strictly less than 1 whenever $B_t \neq 0$. This is confirmed by the results shown in Figure 5.

In this simulation, a constant magnetic field for a straight interface given by $\kappa(x) = x_2$ is modeled by $\tilde{A} = A = -Bx_2 e_1 = Bx_2 \tau$ (with therefore $\chi = 0$ and $\beta = B$). This implies $\dot{j}_t = k_t = 0$ so that $e^{\mu_t} = 1$ and $\nu_t = 0$. The terms $\rho = \sqrt{1 + B^2}$ and $\gamma = \frac{B}{1+B^2}$ are constant. The wavepacket velocity is constant and given by

$$\dot{y}_t = -\frac{1}{\sqrt{1+B^2}} e_1 = -c e_1.$$

The wavepacket in (7.2) takes the simpler form

$$\mathcal{V}_t a_0(t, z) = \left(\frac{\rho}{4\pi}\right)^{\frac{1}{4}} \left(\frac{1}{\sigma + \rho\gamma^2}\right)^{\frac{1}{2}} e^{-\frac{1}{2}\left(\rho z_2^2 + \frac{1}{\sigma + \rho\gamma^2}(z_1 + is_t z_2)^2\right)} \begin{bmatrix} 1 \\ -1 \end{bmatrix}.$$

The oscillations of the spinor components are clearly visible in Figure 5.

7.2. Aharonov–Bohm effect for a circular interface. This effect emerges in the phase $\chi(t, x)/\varepsilon$ in (7.1) when Γ is a loop. Once y_t completes an exact full rotation around Γ , the wavepacket acquires the phase shift

$$\frac{\chi(t, y_t)}{\varepsilon} = \frac{1}{\varepsilon} \int_0^t \dot{y}_s A(y_s) ds = \frac{1}{\varepsilon} \int_\Gamma A, \quad (7.3)$$

see (2.11). This is a gauge-independent quantity involving the magnetic flux $\Phi = \int_\Gamma A$ enclosed by Γ , and of order $1/\varepsilon$.




FIGURE 5. $\varepsilon = 0.05$. Snapshots of real part (left figures) and imaginary part (right figures) of first (top figures) and second (bottom figures) component of the wavepacket with straight interface $\kappa(x) = x_2$. The wavepacket is propagating towards the reader, for constant magnetic fields $B = 1.5, 0.75, 0$ (from left to right in each individual figure).

Consider a magnetic vector potential with flux $\Phi > 0$ given in polar coordinates by

$$A(r, \theta) = \frac{\Phi}{2\pi r} e_\theta$$

and a circular interface Γ given by $|x| = R > 0$. Note that $B = \nabla \times A = \Phi \delta_0$ vanishes away from the origin and in particular in the vicinity of Γ . However, the wavepacket still feels a magnetic effect: after a full revolution around Γ , it acquires according to (7.3) a measurable phase-shift Φ/ε that cannot be gauged away. This is the Aharonov–Bohm effect.

7.3. Dispersive and phase effects in closed interfaces. The coefficient Q_σ in (7.2) controls the dispersion. The only term there that can grow with t is

$$\nu_t e^{2\mu t} = 2 \int_0^t \frac{c_t^2}{c_s^2} \left(\dot{\theta}_s \gamma_s + k_s \gamma_s^2 \right) ds.$$

Consider now a circle of radius R and a choice of domain wall $\kappa(x) = \frac{|x|^m - R^m}{mR^{m-1}}$ for $m > 0$ with $|\nabla \kappa| = 1 = r$ and $\Delta \kappa = m|x|^{-1}$ equal to mR^{-1} on Γ . Assume B constant so that all




FIGURE 6. Aharonov–Bohm effect with $\varepsilon = 3/40$, $\kappa(x) = \frac{|x|^2-1}{2}$ and $A(x) = \frac{\Phi}{2\pi|x|}$. The left and right panels correspond to $\Phi = 0$ and $\Phi = 2\pi$, respectively. The bottom panels are plots of $t \mapsto (\cos \varphi_t, \sin \varphi_t, t)$, where φ_t is the phase of the top spinor component. The case $\Phi = 2\pi$ induces $1/\varepsilon \simeq 13$ revolutions of the phase as the wavepacket travels once around the circle.




FIGURE 7. $\varepsilon = 0.05$. Snapshots showing one revolution of wavepacket, starting at 4 o'clock, on circular edge in different constant magnetic fields $B = 1/\sqrt{2}$ (left, enhanced spreading) and $B = 3/2$ (right, reduced spreading) with common interface $\kappa(x) = \frac{|x|^2-1}{2}$.

coefficients are independent of time and given by

$$c = \frac{1}{\sqrt{1+B^2}}, \quad k = -\frac{cBm}{2R}, \quad \dot{\theta} = \frac{c}{R}, \quad \dot{\theta} + \gamma k = \frac{c}{R} \left(1 - \frac{mB^2}{2(1+B^2)} \right).$$

We thus observe that $\nu_t = 2t\gamma(\dot{\theta} + \gamma k)$ grows linearly in time provided that $\dot{\theta} + \gamma k \neq 0$. In this case, the resulting wavepacket decreases like $t^{-1/2}$, in a way depending on B (higher magnetic fields, however, do not necessarily enhance dispersion). This was predicted in Lemma 5.3 and (7.2) and is numerically confirmed in Figure 7.

In the other hand, when $m = \frac{2(1+B^2)}{B^2}$, the resulting domain wall prevents dispersion; see Figure 8. This can be of interest in application where one wants to slow down propagation without losing on coherence.




FIGURE 8. Homogeneous magnetic field strength $B = 1$ with $\kappa(x) = \frac{|x|^4 - 1}{4}$. Snapshots showing an almost dispersion-free propagation around the circle.

In this scenario we can explicitly construct the local gauge χ . We have $n = e_r$ and $\tau = -e_\theta$, by using the defining relations $\partial_\tau \chi = A_\tau$ on Γ followed by $\partial_n \chi = A_n$ across it. We have

$$A = Bx_1 e_2 = \frac{Br}{2} ((1 + \cos 2\theta)e_\theta + \sin 2\theta e_r)$$

so that integrating along Γ , then across Γ , we obtain

$$\chi(R, \theta) = \frac{BR^2}{2} \left(\theta + \frac{\sin 2\theta}{2} \right), \quad \chi(r, \theta) = BR^2 \frac{\theta}{2} + Br^2 \frac{\sin 2\theta}{4}.$$

This shows that $\chi(r, \theta)$ is not globally defined as a continuous function on \mathbb{R}^2 : the term $BR^2 \frac{\theta}{2}$ jumps after each revolution. The increment πBR^2 is the magnetic flux: we retrieve a Aharonov–Bohm effect.

7.4. Dispersive and (limited) compression effect in varying magnetic fields. We consider here the setting of a flat interface (with $\kappa(x) = x_2$) but with a varying magnetic field. We then observe compression and dispersive effects consistent with (7.2).

In Figure 9, the intensity of the magnetic field increases as the wavepacket propagates along Γ . We thus expect an increase in ρ_t and as a result a compression of the wavepacket. This is confirmed by the numerical simulations of Figure 9.

We next consider the setting of a magnetic field that increases transversely to Γ : $B(x) = 1 + 4B_2 x_2$. Set $A = \tilde{A} = -(1 + 2B_2 x_2)x_2 e_1$, resulting in $\beta = 1 + 2B_2 x_2$; in particular $\partial_n \beta = 2B_2$ and $k = k_t = cB_2$ is constant. This shows that ν_t grows linearly with time. The wavepacket decays like $t^{-1/2}$, as confirmed numerically in Figure 10.

The amplitude drop generated by dispersion is, however, reversible. For instance, the magnetic field $B(x) = 1 + 4 \cos(2\pi x_1/15)x_2$ generates time-dependent oscillations in ν_t : this




FIGURE 9. Snapshots of wavepacket, starting on the right, for a potential $A_1(x) = -B_0 x_2(1 - \tanh(x_1 - 2))$ with $B_0 = 0, 1, 2$ from bottom to top, which corresponds to a magnetic field $B(x) = B_0(1 - \tanh(x_1 - 2))$, on a straight interface $\kappa(x) = x_2$. Right figure with $B_0 = 1$.




FIGURE 10. $\varepsilon = 0.075$, $B = 1 + 4B_2 x_2$ with $B_2 = 0, 0.5, 1$ and $\kappa(x) = x_2$. The left figure shows the snapshots of the three wavepackets, starting on the right, and on the right, we see the power-law decay in the non-constant magnetic field. The right figure shows the L^∞ -decay of the amplitudes.

coefficient is proportional to $\sin(2\pi t/15)$. There is no dispersion for $t \in 15\mathbb{Z}$, as shown in Figure 11.




FIGURE 11. $\varepsilon = 0.1$, snapshots of a wavepacket, starting on the right, for a straight interface $\kappa(x) = x_2$ with $B(x) = 1 + 4 \cos(\frac{2\pi x_1}{15})x_2$, a periodic modulation of the field in Figure 10, sampled over half a period of the cosine. The reversible amplitude drop caused by dispersion is clearly visible.

REFERENCES

- [ASS94] Joseph E. Avron, Ruedi Seiler, and Barry Simon, *Charge deficiency, charge transport and comparison of dimensions*, Comm. Math. Phys. **159** (1994), no. 2, 399–422.
- [Bal19a] Guillaume Bal, *Continuous bulk and interface description of topological insulators*, Journal of Mathematical Physics **60** (2019), no. 8, 081506.
- [Bal19b] Guillaume Bal, *Topological protection of perturbed edge states*, Communications in Mathematical Sciences **17** (2019), no. 1, 193–225.
- [Bal20] Guillaume Bal, *Topological invariants for interface modes* (2020), available at [arXiv:1906.08345](https://arxiv.org/abs/1906.08345).
- [Bal21] Guillaume Bal, *Topological charge conservation for continuous insulators* (2021), available at [arXiv:2106.08480](https://arxiv.org/abs/2106.08480).
- [BBD⁺21] Guillaume Bal, Simon Becker, Alexis Drouot, Clotilde Fermanian Kammerer, Jianfeng Lu, and Alexander Watson, *Edge state dynamics along curved interfaces* (2021), available at [arXiv:2106.00729](https://arxiv.org/abs/2106.00729).
- [Ber13] Bogdan Andrei Bernevig, *Topological insulators and topological superconductors*, Princeton University Press, 2013.
- [BM22] Guillaume Bal and Daniel Massatt, *Multiscale invariants of floquet topological insulators*, To appear in SIAM Multiscale Modeling and Simulation (2022), available at [arXiv:2101.06330](https://arxiv.org/abs/2101.06330).
- [BvESB94] Jean Bellissard, Andreas van Elst, and Hermann Schulz-Baldes, *The noncommutative geometry of the quantum Hall effect*, Journal of Mathematical Physics **35** (1994), no. 10, 5373–5451.
- [CdV04] Yves Colin de Verdière, *The level crossing problem in semi-classical analysis. II. The Hermitian case*, Annales de l’Institut Fourier **54** (2004), no. 5, 1423–1441.
- [DH72] J.J. Duistermaat and L. Hörmander, *Fourier integral operators. II.*, Acta Math. **128** (1972), no. 3-4, 183–269.
- [Dro19a] Alexis Drouot, *The bulk-edge correspondence for continuous honeycomb lattices*, Communication in Partial Differential Equations **44** (2019), no. 12, 1406–1430.
- [Dro19b] Alexis Drouot, *Characterization of edge states in perturbed honeycomb structures*, Pure and Applied Analysis **1** (2019), no. 3, 385–445.
- [Dro21a] Alexis Drouot, *Microlocal analysis of the bulk-edge correspondence*, Communications in Mathematical Physics **383** (2021), 2069–2112.
- [Dro21b] Alexis Drouot, *Ubiquity of conical points in topological insulators*, Journal de l’Ecole Polytechnique **8** (2021), 507–532.
- [DW20] Alexis Drouot and MI Weinstein, *Edge states and the valley Hall effect*, Advances in Mathematics **368** (2020), 107142.
- [EG02] Peter Elbau and Gian-Michele Graf, *Equality of bulk and edge Hall conductance revisited*, Communications in mathematical physics **229** (2002), no. 3, 415–432.

- [FKG03] Clotilde Fermanian Kammerer and Patrick Gérard, *A Landau-Zener formula for non-degenerated involutive codimension three crossings*, Ann. Henri Poincaré **4** (2003), 123–168.
- [FLTW16] Charles L Fefferman, James P Lee-Thorp, and Michael I Weinstein, *Edge states in honeycomb structures*, Annals of PDE **2** (2016), no. 2, 12.
- [GP13] Gian-Michele Graf and Marcello Porta, *Bulk-edge correspondence for two-dimensional topological insulators*, Communications in Mathematical Physics **324** (2013), no. 3, 851–895.
- [GRW21] J. Guglielmon, M. C. Rechtsman, and M. I. Weinstein, *Landau levels in strained two-dimensional photonic crystals*, Phys. Rev. A **103** (2021Jan), 013505.
- [Hag94] George A. Hagedorn, *Molecular propagation through electron energy level crossings*, Memoirs of the AMS **111** (1994), no. 536.
- [HJ98] George A Hagedorn and Alain (F-CNRS-T) Joye, *Landau–Zener transitions through small electronic eigenvalue gaps in the Born–Oppenheimer approximation.*, Ann. Inst. H. Poincaré Phys. Théor. **68** (1998), no. 1, 85–134.
- [MM21] Roderich Moessner and Joel E Moore, *Topological phases of matter*, Cambridge University Press, 2021.
- [PSB16] Emil Prodan and Hermann Schulz-Baldes, *Bulk and boundary invariants for complex topological insulators*, Springer verlag, Berlin, 2016.
- [Tha92] Bernd Thaller, *The Dirac equation*, Texts and monographs in physics, Springer-Verlag, 1992.
- [TKNdN82] D. J. Thouless, M. Kohmoto, M. P. Nightingale, and M. den Nijs, *Quantized Hall Conductance in a Two-Dimensional Periodic Potential*, Phys. Rev. Lett. **49** (1982Aug), 405–408.
- [Vol89] Grigory E. Volovik, *Nonlinear phenomena in condensed matter: Universe in a Helium droplet*, 1989.
- [Wit16] Edward Witten, *Three lectures on topological phases of matter*, Nuovo Cimento Rivista Serie **39** (2016), no. 7, 313–370.

(Guillaume Bal) UNIVERSITY OF CHICAGO, USA.

Email address: guillaumbal@uchicago.edu

(Simon Becker) COURANT INSTITUTE FOR MATHEMATICAL SCIENCES, NEW YORK CITY, USA.

Email address: simon.becker@cims.nyu.edu

(Alexis Drouot) UNIVERSITY OF WASHINGTON, USA.

Email address: adrouot@uw.edu