

RANDOM β -TRANSFORMATION ON FAT SIERPINSKI GASKETTINGYU ZHANG¹, KARMA DAJANI² AND WENXIA LI³

ABSTRACT. We consider the iterated function system (IFS)

$$f_{\vec{q}}(\vec{z}) = \frac{\vec{z} + \vec{q}}{\beta}, \vec{q} \in \{(0, 0), (1, 0), (0, 1)\}.$$

As is well known, for $\beta = 2$ the attractor, S_β , is a fractal called the Sierpiński gasket(or sieve) and for $\beta > 2$ it is also a fractal. Our goal is to study greedy, lazy and random β -transformations on the attractor for this IFS with $1 < \beta < 2$. For $1 < \beta \leq 3/2$, S_β is a triangle and it is shown that the greedy transformation T_β and the lazy transformation L_β are isomorphic and they both admit an absolutely continuous invariant measure. We show that all β -expansions of a point \vec{z} in S_β can be generated by a random map K_β defined on $\{0, 1\}^\mathbb{N} \times \{0, 1, 2\}^\mathbb{N} \times S_\beta$ and K_β has a unique invariant measure of maximal entropy when $1 < \beta \leq \beta_*$, where $\beta_* \approx 1.4656$ is the root of $x^3 - x^2 - 1 = 0$. We also show existence of a K_β -invariant probability measure, absolutely continuous with respect to $m_1 \otimes m_2 \otimes \lambda_2$, where m_1, m_2 are product measures on $\{0, 1\}^\mathbb{N}, \{0, 1, 2\}^\mathbb{N}$, respectively, and λ_2 is the normalized Lebesgue measure on S_β . For $3/2 < \beta \leq \beta^*$, where $\beta^* \approx 1.5437$ is the root of $x^3 - 2x^2 + 2x = 2$, there are radial holes in S_β . In this case, K_β is defined on $\{0, 1\}^\mathbb{N} \times S_\beta$. We also show that it has a unique invariant measure of maximal entropy.

MSC2020: 37H12, 37A44, 37A05, 11K55**Keywords:** fat Sierpinski gasket, random β -transformation, absolutely continuous invariant measures, measures of maximal entropy.

1. INTRODUCTION

Let $\beta > 1$ and consider the *iterated function system*(IFS):

$$f_{\vec{q}_0}(\vec{z}) = \frac{\vec{z} + \vec{q}_0}{\beta}, f_{\vec{q}_1}(\vec{z}) = \frac{\vec{z} + \vec{q}_1}{\beta}, f_{\vec{q}_2}(\vec{z}) = \frac{\vec{z} + \vec{q}_2}{\beta},$$

where the coordinates of the three points $\vec{q}_0, \vec{q}_1, \vec{q}_2$ are $(0, 0), (1, 0), (0, 1)$, respectively. It is well known that there exists a unique non-empty compact set $S_\beta \subset \mathbb{R}^2$ such that $S_\beta = \cup_{i=0}^2 f_{\vec{q}_i}(S_\beta)$; see [12] for further details. The *attractor* for the IFS, S_β , is a Sierpinski gasket. Denote the convex hull of S_β by Δ which is a triangle with vertices at $(0, 0), (\frac{1}{\beta-1}, 0)$ and $(0, \frac{1}{\beta-1})$. For every point $\vec{z} \in S_\beta$, there exists a

1,3: Department of Mathematics, Shanghai Key Laboratory of PMMP, East China Normal University, Shanghai 200241, People's Republic of China.

2: Department of Mathematics, Utrecht University, Fac Wiskunde en informatica and MRI, Budapestlaan 6, P.O. Box 80.000, 3508 TA Utrecht, the Netherlands.

Email addresses: tinguyzhang@ecnu.edu.cn (T. Zhang), k.dajani@uu.nl (K. Dajani), wxli@math.ecnu.edu.cn (W. Li).

sequence $(a_i)_{i=1}^\infty \in \{\vec{q}_0, \vec{q}_1, \vec{q}_2\}^\mathbb{N}$ such that

$$\vec{z} = \lim_{n \rightarrow \infty} f_{a_1} \circ \cdots \circ f_{a_n}(\vec{q}_0) = \sum_{i=1}^{\infty} \frac{a_i}{\beta^i}.$$

We call $(a_i)_{i=1}^\infty$ a *coding* of \vec{z} and $\sum_{i=1}^\infty a_i \beta^{-i}$ a *representation* of \vec{z} in base β .

Let $i \in \{0, 1, 2\}$. For $\beta > 2$, the images $f_{\vec{q}_i}(\Delta)$ are disjoint. In the case the IFS $\{f_{\vec{q}_i}\}$ satisfies the strong separation condition and each point in S_β has a unique coding. For $\beta = 2$, the sets $f_{\vec{q}_i}(\Delta)$ overlap only at the vertices. Therefore only countably many points in S_β have two codings, and all other points have a unique coding.

When $\beta \in (1, 2)$, we call S_β a *fat Sierpinski gasket*. For $1 < \beta \leq 3/2$, we have a non-empty triple overlap $C_{012} := f_{\vec{q}_0}(\Delta) \cap f_{\vec{q}_1}(\Delta) \cap f_{\vec{q}_2}(\Delta)$ (see Figure 1), and $S_\beta = \Delta$. In this case Lebesgue almost every point in S_β has a continuum of codings (see [19, Theorem 3.5]). For $3/2 < \beta < 2$, there are holes in S_β as well as overlaps (see Figure 3 for example), which make its structure more complex. In [6], Broomhead et al. described two special types of structures: those in which holes are radially distributed and those that are totally self-similar. We are more interested in S_β with the former characteristics. For more results of the Hausdorff dimension of the attractors, see [16, 15, 18, 17, 14].

In this article, we shall consider the random transformation on the two-dimensional Sierpinski gasket S_β . The main motivation of this consideration is from the random β -expansion in an interval. In [9], Dajani and Kraaikamp introduced a random β -transformation K_β on $\{0, 1\}^\mathbb{N} \times [0, [\beta]/(\beta - 1)]$ associated with the ‘greedy’ map and the ‘lazy’ map. They showed that the greedy expansion is isomorphic to the lazy expansion (see [8]). Dajani and de Vries [10] showed that K_β has a unique measure of maximal entropy. In [11] they also proved the existence and uniqueness of a K_β -invariant measure, absolutely continuous with respect to $m_p \otimes \lambda$, where m_p is the Bernoulli measure on $\{0, 1\}^\mathbb{N}$ and λ is the normalized Lebesgue measure on $[0, [\beta]/(\beta - 1)]$. Inspired by these results, we consider transformations which are defined on fat Sierpinski triangles with overlapping structures.

The rest of the article is organized as follows. We suppose $\beta \in (1, 3/2]$ in sections 3, 4 and 5. In section 2, we focus on some basic concepts and notations needed in the rest of the paper. In section 3, we introduce the greedy and lazy transformations on S_β and prove that they are isomorphic. We also prove the existence of invariant measures that are absolutely continuous with respect to Lebesgue measure (acim) for these two transformations. In section 4, we give the definition of the random transformation K_β on $\{0, 1\}^\mathbb{N} \times \{0, 1, 2\}^\mathbb{N} \times S_\beta$ and prove that it has a unique invariant measure of maximal entropy for $1 < \beta \leq \beta_*$, where $\beta_* \approx 1.4656$ is the root of $x^3 - x^2 - 1 = 0$. In section 5, we give a position dependent random map R on S_β . With two skew product transformations, we establish a connection between R and K_β , and finally prove that K_β has an invariant measure of the form $m_1 \otimes m_2 \otimes \mu_\beta$, where m_1 is the product measure on $\{0, 1\}^\mathbb{N}$ with weights $\{p, 1 - p\}$, m_2 is the product measure on $\{0, 1, 2\}^\mathbb{N}$ with weights $\{s, t, 1 - s - t\}$, and μ_β is R -invariant and absolutely continuous with respect to λ_2 , the normalized Lebesgue measure on S_β . In section 6, we modify the definition of the random transformation K_β for $3/2 < \beta \leq \beta^*$ and prove that it also has a unique invariant measure of maximal entropy, where $\beta^* \approx 1.5437$ is the root of $x^3 - 2x^2 + 2x = 2$.

2. PRELIMINARY

Given $1 < \beta < 2$, recall that the fat Sierpinski gasket S_β is the self-similar set in \mathbb{R}^2 generated by the IFS

$$(2.1) \quad f_{\vec{q}_0}(\vec{z}) = \frac{\vec{z} + \vec{q}_0}{\beta}, f_{\vec{q}_1}(\vec{z}) = \frac{\vec{z} + \vec{q}_1}{\beta}, f_{\vec{q}_2}(\vec{z}) = \frac{\vec{z} + \vec{q}_2}{\beta},$$

where $\vec{q}_0, \vec{q}_1, \vec{q}_2$ are $(0, 0), (1, 0), (0, 1)$, respectively. For every point $\vec{z} \in S_\beta$, there exists a sequence $(a_i)_{i=1}^\infty \in \{\vec{q}_0, \vec{q}_1, \vec{q}_2\}^\mathbb{N}$ such that

$$\vec{z} = \sum_{i=1}^{\infty} \frac{a_i}{\beta^i}.$$

Recall that Δ is the convex hull of S_β which is an isosceles right triangle.

Consider the ordering of points on the plane. We write $(x_1, y_1) < (x_2, y_2)$ if $x_1 + y_1 < x_2 + y_2$, or $x_1 + y_1 = x_2 + y_2$ and $y_1 < y_2$. Notice that $\vec{q}_0 < \vec{q}_1 < \vec{q}_2$.

Let $\Omega = \{0, 1\}^\mathbb{N}$ with the product σ -algebra \mathcal{A} and $\Upsilon = \{0, 1, 2\}^\mathbb{N}$ with the product σ -algebra \mathcal{B} . Define metrics d and ρ on Ω and Υ respectively by

$$d((\omega_i), (\omega'_i)) = 2^{-\min\{k: \omega_k \neq \omega'_k\}}$$

and

$$\rho((v_i), (v'_i)) = 3^{-\min\{k: v_k \neq v'_k\}}.$$

Throughout the article, the lexicographical ordering on Ω , Υ and $\{\vec{q}_0, \vec{q}_1, \vec{q}_2\}^\mathbb{N}$ are all denoted by \prec and \preceq . More precisely, for two sequences $(c_i), (d_i)$ we write $(c_i) \prec (d_i)$ if $c_1 < d_1$, or there exists $k \geq 2$ such that $c_i = d_i$ for all $1 \leq i < k$ and $c_k < d_k$. Similarly, we write $(c_i) \preceq (d_i)$ if $(c_i) \prec (d_i)$ or $(c_i) = (d_i)$.

We will use the following concepts and properties.

Definition 2.1. [20] A topological space X is called *separable* if it has a countable dense set. A topological space is called *completely metrizable* if its topology is induced by a complete metric. A *Polish space* is a separable, completely metrizable topological space.

There are some elementary observations.

- (i) The real line \mathbb{R} with the usual topology is Polish.
- (ii) Any countable discrete space is Polish. In particular, $\{0, 1, 2\}$ with the discrete topology is Polish.
- (iii) The product of countably many Polish spaces is Polish. In particular, $\Omega = \{0, 1\}^\mathbb{N}$ and $\Upsilon = \{0, 1, 2\}^\mathbb{N}$ are Polish.

Theorem 2.2. [20] *Let X, Y be Polish spaces, A a Borel subset of X , and $f : A \rightarrow Y$ a one-to-one Borel map. Then $f(A)$ is Borel.*

Definition 2.3. [7] A *dynamical system* is a quadruple (X, \mathcal{F}, μ, Q) , where X is a non-empty set, \mathcal{F} is a σ -algebra on X , μ is a probability measure on (X, \mathcal{F}) and $Q : X \rightarrow X$ is a surjective μ -measure preserving transformation.

If (X, \mathcal{F}, μ, Q) is a dynamical system, and $x \in X$, we call the sequence

$$x, Qx, \dots, Q^n x, \dots$$

the Q -orbit of x .

Definition 2.4. [7] Two dynamical systems (X, \mathcal{F}, μ, Q) and (Y, \mathcal{G}, ν, U) are *isomorphic* if there exist measurable sets $N \subset X$ and $M \subset Y$ with $\mu(N) = 0 = \nu(M)$ and $Q(X \setminus N) \subset X \setminus N, U(Y \setminus M) \subset Y \setminus M$, and finally if there exists a measurable map $\psi : X \setminus N \rightarrow Y \setminus M$ such that (i)–(iv) are satisfied for the systems restricted to $X \setminus N$ and $Y \setminus M$:

- (i) ψ is one-to-one and onto,
- (ii) ψ is measurable, i.e., $\psi^{-1}(G) \in \mathcal{F}$ for all $G \in \mathcal{G}$,
- (iii) ψ preserves the measures, i.e., $\nu(G) = \mu(\psi^{-1}(G))$,
- (iv) ψ preserves the dynamics of Q and U , i.e., $\psi \circ Q = U \circ \psi$,

The map ψ is called an *isomorphism*.

Proposition 2.5. [7] *Entropy is isomorphism invariant.*

3. GREEDY AND LAZY TRANSFORMATION FOR $1 < \beta \leq 3/2$

Let $1 < \beta \leq 3/2$, then $S_\beta = \Delta$. Divide S_β into the following sets according to the overlapping structure of $f_{\vec{q}_i}(S_\beta)$ (see Figure 1).

$$\begin{aligned}
 E_0 &= [0, \frac{1}{\beta}) \times [0, \frac{1}{\beta}), \\
 E_1 &= \{(x, y) : 0 \leq y < \frac{1}{\beta}, \frac{1}{\beta(\beta-1)} < x+y \leq \frac{1}{\beta-1}\}, \\
 E_2 &= \{(x, y) : 0 \leq x < \frac{1}{\beta}, \frac{1}{\beta(\beta-1)} < x+y \leq \frac{1}{\beta-1}\}, \\
 C_{01} &= \{(x, y) : x \geq \frac{1}{\beta}, 0 \leq y < \frac{1}{\beta}, x+y \leq \frac{1}{\beta(\beta-1)}\}, \\
 C_{12} &= \{(x, y) : x \geq \frac{1}{\beta}, y \geq \frac{1}{\beta}, \frac{1}{\beta(\beta-1)} < x+y \leq \frac{1}{\beta-1}\}, \\
 C_{02} &= \{(x, y) : 0 \leq x < \frac{1}{\beta}, y \geq \frac{1}{\beta}, x+y \leq \frac{1}{\beta(\beta-1)}\}, \\
 C_{012} &= \{(x, y) : x \geq \frac{1}{\beta}, y \geq \frac{1}{\beta}, x+y \leq \frac{1}{\beta(\beta-1)}\}.
 \end{aligned}
 \tag{3.1}$$

Notice that C_{012} is a single point set if $\beta = 3/2$. Let


$$C = C_{01} \cup C_{12} \cup C_{02}, \quad E = \cup_{i=0}^2 E_i.$$

Definition 3.1. The greedy transformation T_β from S_β into S_β is given by

$$T_\beta(\vec{z}) = \begin{cases} \beta\vec{z}, & \text{if } \vec{z} \in E_0, \\ \beta\vec{z} - \vec{q}_1, & \text{if } \vec{z} \in C_{01} \cup E_1, \\ \beta\vec{z} - \vec{q}_2, & \text{if } \vec{z} \in C_{012} \cup C_{12} \cup C_{02} \cup E_2. \end{cases}
 \tag{3.2}$$

The lazy transformation L_β from S_β into S_β is given by

$$L_\beta(\vec{z}) = \begin{cases} \beta\vec{z}, & \text{if } \vec{z} \in C_{012} \cup C_{01} \cup C_{02} \cup E_0, \\ \beta\vec{z} - \vec{q}_1, & \text{if } \vec{z} \in C_{12} \cup E_1, \\ \beta\vec{z} - \vec{q}_2, & \text{if } \vec{z} \in E_2. \end{cases}
 \tag{3.3}$$

FIGURE 1. S_β for $1 < \beta \leq \frac{3}{2}$

Notice that

$$\begin{aligned} T_\beta|_{E_i} &= L_\beta|_{E_i} = f_{\vec{q}_i}^{-1} \text{ for } i = \{0, 1, 2\}, \\ T_\beta|_{C_{ij}} &= f_{\vec{q}_j}^{-1}, L_\beta|_{C_{ij}} = f_{\vec{q}_i}^{-1} \text{ for } ij \in \{01, 12, 02\}, \\ T_\beta|_{C_{012}} &= f_{\vec{q}_2}^{-1}, L_\beta|_{C_{012}} = f_{\vec{q}_0}^{-1}, \end{aligned}$$

and

$$\begin{aligned} f_{\vec{q}_0}(S_\beta) &= E_0 \cup C_{01} \cup C_{02} \cup C_{012}, \\ f_{\vec{q}_1}(S_\beta) &= E_1 \cup C_{01} \cup C_{12} \cup C_{012}, \\ f_{\vec{q}_2}(S_\beta) &= E_2 \cup C_{12} \cup C_{02} \cup C_{012}. \end{aligned}$$

Then it is easy to verify that T_β and L_β are well defined.

Denote the Borel σ -algebra on S_β by \mathcal{S} . Let μ be an arbitrary T_β -invariant probability measure on (S_β, \mathcal{S}) . Let $\psi : S_\beta \rightarrow S_\beta$ be given by

$$\psi(\vec{z}) = \begin{bmatrix} 1 & 0 \\ -1 & -1 \end{bmatrix} \vec{z} + \begin{bmatrix} 0 \\ \frac{1}{\beta-1} \end{bmatrix}.$$

Theorem 3.2. *For $\beta \in (1, 3/2]$, the systems $(S_\beta, \mathcal{S}, \mu, T_\beta)$ and $(S_\beta, \mathcal{S}, \nu, L_\beta)$ are isomorphic, where $\nu = \mu \circ \psi^{-1}$.*

Proof. Denote $\begin{bmatrix} x' \\ y' \end{bmatrix} = \psi\left(\begin{bmatrix} x \\ y \end{bmatrix}\right)$. Then

$$x' = x, \quad y' = \frac{1}{\beta-1} - (x+y), \quad x' + y' = \frac{1}{\beta-1} - y.$$

Therefore, we can obtain

$$\begin{aligned}
\psi(E_0) &= \psi\{(x, y) : 0 \leq x < \frac{1}{\beta}, 0 \leq y < \frac{1}{\beta}\} \\
&= \{(x', y') : 0 \leq x' < \frac{1}{\beta}, \frac{1}{\beta-1} - \frac{1}{\beta} < x' + y' \leq \frac{1}{\beta-1}\} \\
&= \{(x', y') : 0 \leq x' < \frac{1}{\beta}, \frac{1}{\beta(\beta-1)} < x' + y' \leq \frac{1}{\beta-1}\} \\
&= E_2.
\end{aligned}$$

By doing the same calculation, we can also obtain

$$\psi(E_1) = E_1, \psi(C_{01}) = C_{12}, \psi(C_{02}) = C_{02}, \psi(C_{012}) = C_{012}.$$

Notice that $\psi^{-1} = \psi$ and $\psi(S_\beta) = S_\beta$, then we can get $\psi(E_2) = E_0$ and $\psi(C_{12}) = C_{01}$. Now we show $\psi \circ T_\beta = L_\beta \circ \psi$. For $(x, y) \in E_0$,

$$\psi \circ T_\beta \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = \psi \left(\begin{bmatrix} \beta x \\ \beta y \end{bmatrix} \right) = \begin{bmatrix} \beta x \\ \frac{1}{\beta-1} - \beta x - \beta y \end{bmatrix}.$$

Since $\psi((x, y)) \in E_2$, then

$$L_\beta \circ \psi \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) = \beta \begin{bmatrix} x \\ \frac{1}{\beta-1} - x - y \end{bmatrix} - \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} \beta x \\ \frac{\beta}{\beta-1} - \beta x - \beta y - 1 \end{bmatrix} = \psi \circ T_\beta \left(\begin{bmatrix} x \\ y \end{bmatrix} \right).$$

Similarly, we have

$$\begin{aligned}
\psi \circ T_\beta \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) &= \begin{bmatrix} \beta x - 1 \\ \frac{\beta}{\beta-1} - \beta x - \beta y \end{bmatrix} = L_\beta \circ \psi \left(\begin{bmatrix} x \\ y \end{bmatrix} \right), \text{ for } (x, y) \in C_{01} \cup E_1, \\
\psi \circ T_\beta \left(\begin{bmatrix} x \\ y \end{bmatrix} \right) &= \begin{bmatrix} \beta x \\ \frac{\beta}{\beta-1} - \beta x - \beta y \end{bmatrix} = L_\beta \circ \psi \left(\begin{bmatrix} x \\ y \end{bmatrix} \right), \text{ for } (x, y) \in C_{012} \cup C_{12} \cup C_{02} \cup E_2.
\end{aligned}$$

Since $\psi : S_\beta \rightarrow S_\beta$ is a bijection, then it follows from $\nu = \mu \circ \psi^{-1}$ that ψ is an isomorphism. \square

We end this section by recalling some definitions and results in [4] due to Boryarsky, A. and Góra, P., but rephrased to our setting. Let S be a bounded region in \mathbb{R}^N ($N \geq 2$ is an integer) and let $\mathcal{P} = \{S_1, S_2, \dots, S_m\}$ be a partition of S , where $m > 0$ is an integer. Let τ be a transformation from S into S . We say τ is *piecewise C^2 and expanding* with respect to \mathcal{P} if:

- (a) each S_i is a bounded closed domain having a piecewise C^2 boundary of finite $(N-1)$ -dimensional measure;
- (b) $\tau_i = \tau|_{S_i}$ is a C^2 , 1-1 transformation from $\text{int}(S_i)$ onto its image and can be extended as a C^2 transformation onto $S_i, i = 1, 2, \dots, m$;
- (c) there exists $0 < c < 1$ such that for any $i = 1, 2, \dots, m$,

$$\|D\tau_i^{-1}\| < c,$$

where $D\tau_i^{-1}$ is the derivative matrix of τ_i^{-1} and $\|\cdot\|$ is the euclidean matrix norm, i.e., $\|A\| = (\sum_{i=1}^m \sum_{j=1}^n a_{ij}^2)^{\frac{1}{2}}$ for $m \times n$ order matrix $A = (a_{ij})$.

Assume that the faces of ∂S_i meet at angles bounded uniformly away from 0. Fix $1 \leq i \leq m$. Let F denote the set of singular points of ∂S_i . At any $x \in F$ we

construct the largest cone having a vertex at x and which lies completely in S_i . Let $\theta(x)$ denote the angle subtended at the vertex of this cone. Then define

$$(3.4) \quad \gamma(S_i) = \min_{x \in F} \theta(x).$$

Since the faces of ∂S_i meet at angles bounded uniformly away from 0, $\gamma(S_i) > 0$. Let $\alpha(S_i) = \pi/2 + \gamma(S_i)$ and

$$(3.5) \quad a(S_i) = |\cos(\alpha(S_i))|.$$

Assume $a = \min\{a(S_i) : i = 1, \dots, m\}$.

Corollary 3.3. [4, Corollary 1] *Let $\tau : S \rightarrow S, S \subset \mathbb{R}^N$, be piecewise C^2 and such that some iterate τ^k satisfies $c(1 + 1/a) < 1$ (c and a corresponds to τ^k), then τ admits an absolutely continuous invariant measure (acim).*

Using the above corollary, we can prove the following theorem.

Theorem 3.4. *Let T_β be defined as in (3.2). Then T_β admits an acim. And so does L_β which is defined as in (3.3).*

Proof. Since $\beta > 1$, $\lim_{n \rightarrow \infty} \beta^n = \infty$. We can choose an integer k such that

$$\beta^k > 2(2 + \sqrt{2}).$$

Use the partition $\mathcal{P} : \{S_1, S_2, S_3\}$, where

$$S_1 = E_0, S_2 = C_{01} \cup E_1, S_3 = C_{012} \cup C_{12} \cup C_{02} \cup E_2.$$

Consider the iterate T_β^k and the corresponding partition $\vee_{i=0}^{k-1} T_\beta^{-i}(\mathcal{P})$. For $P_i \in \vee_{i=0}^{k-1} T_\beta^{-i}(\mathcal{P})$, let $T_{\beta,i}^k = T_\beta^k|_{P_i}$. Since the derivative matrix of $(T_{\beta,i}^k)^{-1}$ is $\begin{bmatrix} \frac{1}{\beta^k} & 0 \\ 0 & \frac{1}{\beta^k} \end{bmatrix}$, then the Euclidean matrix norm,

$$\|D(T_{\beta,i}^k)^{-1}\| = \frac{\sqrt{2}}{\beta^k} < \frac{2\sqrt{2}}{\beta^k} := c < 1.$$

For the partition $\vee_{i=0}^{k-1} T_\beta^{-i}(\mathcal{P})$, we have $a = \sqrt{2}/2$, which implies

$$c(1 + \frac{1}{a}) = \frac{2\sqrt{2}}{\beta^k}(1 + \sqrt{2}) < \frac{2\sqrt{2}}{2(2 + \sqrt{2})}(1 + \sqrt{2}) = 1.$$

By Corollary 3.3, T_β admits an acim.

For L_β , we use the partition $\mathcal{Q} = \{C_{012} \cup C_{01} \cup C_{02} \cup E_0, C_{12} \cup E_1, E_2\}$. In a similar way we can also prove that L_β has an acim. \square

4. RANDOM TRANSFORMATION FOR $1 < \beta \leq 3/2$

4.1. K_β and some properties. From Definition 3.1, one can see that on E_i both the greedy and lazy maps are identical which means that they assign the same digits. On C_{ij} , the greedy map assigns the digit \vec{q}_j , while the lazy map assigns the digit \vec{q}_i . On C_{012} , the digit \vec{q}_1 cannot be assigned by the two maps. Now, in order to define a random transformation, we need to randomly select the map used in the switch regions C_{ij} and C_{012} . When \vec{z} belongs to C_{ij} we flip a 2-sided coin to decide which map will be applied to \vec{z} , while we flip a 3-sided ‘coin’ when \vec{z} belongs to C_{012} .

Recall $\Omega = \{0, 1\}^\mathbb{N}$ and $\Upsilon = \{0, 1, 2\}^\mathbb{N}$. Let $\sigma : \Omega \rightarrow \Omega$ and $\sigma' : \Upsilon \rightarrow \Upsilon$ be the left shifts. Define $K_\beta : \Omega \times \Upsilon \times S_\beta \rightarrow \Omega \times \Upsilon \times S_\beta$ by

$$K_\beta(\omega, v, \vec{z}) = \begin{cases} (\omega, v, \beta\vec{z} - \vec{q}_i), & \text{if } \vec{z} \in E_i, i = 0, 1, 2, \\ (\sigma\omega, v, \beta\vec{z} - \vec{q}_i), & \text{if } \omega_1 = 0 \text{ and } \vec{z} \in C_{ij}, ij \in \{01, 12, 02\}, \\ (\sigma\omega, v, \beta\vec{z} - \vec{q}_j), & \text{if } \omega_1 = 1 \text{ and } \vec{z} \in C_{ij}, ij \in \{01, 12, 02\}, \\ (\omega, \sigma'v, \beta\vec{z} - \vec{q}_i), & \text{if } \vec{z} \in C_{012} \text{ and } v_1 = i \in \{0, 1, 2\}. \end{cases}$$

The digits are given by

$$d_1 = d_1(\omega, v, \vec{z}) = \begin{cases} \vec{q}_i, & \text{if } \vec{z} \in E_i, i = 0, 1, 2, \\ \text{or } (\omega, v, \vec{z}) \in \Omega \times \{v_1 = i\} \times C_{012}, \\ \text{or } (\omega, v, \vec{z}) \in \{\omega_1 = 0\} \times \Upsilon \times C_{ij}, ij \in \{01, 12, 02\}, \\ \vec{q}_j, & \text{if } (\omega, v, \vec{z}) \in \{\omega_1 = 1\} \times \Upsilon \times C_{ij}, ij \in \{01, 12, 02\}. \end{cases}$$

Then

$$K_\beta(\omega, v, \vec{z}) = \begin{cases} (\omega, v, \beta\vec{z} - d_1), & \text{if } \vec{z} \in E_0 \cup E_1 \cup E_2, \\ (\sigma\omega, v, \beta\vec{z} - d_1), & \text{if } \vec{z} \in C_{01} \cup C_{12} \cup C_{02}, \\ (\omega, \sigma'v, \beta\vec{z} - d_1), & \text{if } \vec{z} \in C_{012}. \end{cases}$$

Set $d_n = d_n(\omega, v, \vec{z}) = d_1(K_\beta^{n-1}(\omega, v, \vec{z}))$, and $\pi_3 : \Omega \times \Upsilon \times S_\beta \rightarrow S_\beta$ be the canonical projection onto the third coordinate. Then

$$\pi_3(K_\beta^n(\omega, v, \vec{z})) = \beta^n \vec{z} - \beta^{n-1} d_1 - \dots - \beta d_{n-1} - d_n,$$

and rewriting yields

$$\vec{z} = \frac{d_1}{\beta} + \frac{d_2}{\beta^2} + \dots + \frac{d_n}{\beta^n} + \frac{\pi_3(K_\beta^n(\omega, v, \vec{z}))}{\beta^n}.$$

Since $\pi_3(K_\beta^n(\omega, v, \vec{z})) \in S_\beta$ and S_β is a bounded set in \mathbb{R}^2 , it follows that

$$\|\vec{z} - \sum_{i=1}^n \frac{d_i}{\beta^i}\|_1 = \frac{\|\pi_3(K_\beta^n(\omega, v, \vec{z}))\|_1}{\beta^n} \rightarrow 0,$$

where $\|\cdot\|_1$ denotes the L_1 norm, i.e. the sum of the absolute values of the vector elements. This shows that for all $\omega \in \Omega$, $v \in \Upsilon$ and for all $\vec{z} \in S_\beta$ one has

$$\vec{z} = \sum_{i=1}^{\infty} \frac{d_i}{\beta^i} = \sum_{i=1}^{\infty} \frac{d_i(\omega, v, \vec{z})}{\beta^i}.$$

For each point $\vec{z} \in S_\beta$, consider the set

$$D_{\vec{z}} = \{(d_1(\omega, v, \vec{z}), d_2(\omega, v, \vec{z}), \dots) : \omega \in \Omega, v \in \Upsilon\}.$$

Recall that \prec and \preceq are the lexicographical ordering on Ω , Υ and $\{\vec{q}_0, \vec{q}_1, \vec{q}_2\}^{\mathbb{N}}$. The following theorem shows how the lexicographical ordering on Ω and Υ affect the ordering of the elements in $D_{\vec{z}}$.

Theorem 4.1. *Suppose $\omega, \omega' \in \Omega, v, v' \in \Upsilon$ are such that $\omega \prec \omega'$ and $v \prec v'$. Then for $\vec{z} \in S_\beta$,*

$$(d_1(\omega, v, \vec{z}), d_2(\omega, v, \vec{z}), \dots) \preceq (d_1(\omega', v', \vec{z}), d_2(\omega', v', \vec{z}), \dots).$$

Proof. Let $m := \inf\{i : \omega_i < \omega'_i\}$, $n := \inf\{i : v_i < v'_i\}$. Then we have $\omega_m < \omega'_m$ and $v_n < v'_n$. Denote by t_i the time of the i^{th} visit to the region $\Omega \times \Upsilon \times C$ of the orbit of (ω, v, \vec{z}) under K_β . Denote by s_j the time of the j^{th} visit to the region $\Omega \times \Upsilon \times C_{012}$ of the orbit of (ω, v, \vec{z}) under K_β . One can see that $K_\beta^{t_i}(\omega, v, \vec{z})$ hits $\Omega \times \Upsilon \times C$ for the i^{th} time and $K_\beta^{s_j}(\omega, v, \vec{z})$ hits $\Omega \times \Upsilon \times C_{012}$ for the j^{th} time.

Let $l = \min\{t_m, s_n\}$. Then $\pi_3(K_\beta^l(\omega, v, \vec{z})) = \pi_3(K_\beta^l(\omega', v', \vec{z}))$ for $i = 0, \dots, l$. It follows that $d_i(\omega, v, \vec{z}) = d_i(\omega', v', \vec{z})$ for $i = 0, \dots, l$.

If $l = \infty$, then $d_i(\omega, v, \vec{z}) = d_i(\omega', v', \vec{z})$ for all i . If $l < +\infty$, then $K_\beta^l(\omega, v, \vec{z}) = K_\beta^l(\omega', v', \vec{z})$ hits $\Omega \times \Upsilon \times C$ for the m^{th} time or $\Omega \times \Upsilon \times C_{012}$ for the n^{th} time. Since $\omega_m < \omega'_m$ and $v_n < v'_n$, then

$$d_{l+1}(\omega, v, \vec{z}) = d_1(K_\beta^l(\omega, v, \vec{z})) < d_1(K_\beta^l(\omega', v', \vec{z})) = d_{l+1}(\omega', v', \vec{z}).$$

□

Now we show that any representation of \vec{z} can be generated from the map K_β by choosing appropriate $\omega \in \Omega$ and $v \in \Upsilon$. We need the following lemma.

Lemma 4.2. *Let $\beta \in (1, 3/2]$. Let $(x, y) \in S_\beta$ and $(x, y) = \sum_{i=1}^\infty a_i \beta^{-i}$ with $a_i \in \{\vec{q}_0, \vec{q}_1, \vec{q}_2\}$ be a representation of (x, y) in base β . One has,*

- (i) *If $(x, y) \in E_i$ for some $i \in \{0, 1, 2\}$, then $a_1 = \vec{q}_i$;*
- (ii) *If $(x, y) \in C_{ij}$ for some $ij \in \{01, 12, 02\}$, then $a_1 \in \{\vec{q}_i, \vec{q}_j\}$;*
- (iii) *If $(x, y) \in C_{012}$, then $a_1 \in \{\vec{q}_0, \vec{q}_1, \vec{q}_2\}$.*

Proof. (i) Suppose $a_1 \neq \vec{q}_0$. From $a_1 \in \{\vec{q}_1, \vec{q}_2\}$ and $(x, y) = \sum_{i=1}^\infty a_i \beta^{-i}$ we have

$$x \geq \frac{1}{\beta} \text{ or } y \geq \frac{1}{\beta}.$$

Then $(x, y) \notin E_0$.

Suppose $a_1 \neq \vec{q}_1$. If $a_1 = \vec{q}_0$, then

$$x + y = \sum_{i=2}^\infty \frac{\|a_i\|_1}{\beta^i} \leq \frac{1}{\beta(\beta-1)}.$$

If $a_1 = \vec{q}_2$, then $y \geq \frac{1}{\beta}$. In both cases, $(x, y) \notin E_1$.

Suppose $a_1 \neq \vec{q}_2$. By a similar proof we have $(x, y) \notin E_2$.

(ii) Suppose $a_1 \notin \{\vec{q}_i, \vec{q}_j\}$. If $a_1 = \vec{q}_2$, then $y \geq \frac{1}{\beta}$, which implies $(x, y) \notin C_{01}$. If $a_1 = \vec{q}_0$, then $x + y \leq \frac{1}{\beta(\beta-1)}$, which implies $(x, y) \notin C_{12}$. If $a_1 = \vec{q}_1$, then $x \geq \frac{1}{\beta}$, which implies $(x, y) \notin C_{02}$.

(iii) holds trivially. □

Theorem 4.3. *For $\beta \in (1, 3/2]$, let $\vec{z} \in S_\beta$ and $\vec{z} = \sum_{i=1}^\infty a_i \beta^{-i}$ with $a_i \in \{\vec{q}_0, \vec{q}_1, \vec{q}_2\}$ be a representation of \vec{z} in base β . Then there exists an $\omega \in \Omega$ and an $v \in \Upsilon$ such that $a_i = d_i(\omega, v, \vec{z})$.*

Proof. Let $\vec{z}_n = \sum_{i=1}^\infty \frac{a_i + n - 1}{\beta^i}$. Notice that $\vec{z}_1 = \vec{z}$. Denote by $k_n(\vec{z})$ and $l_n(\vec{z})$ the number of times we flip a 2-sided coin and a 3-sided coin, respectively, which are

$$k_n(\vec{z}) = \sum_{i=1}^n \mathbf{1}_C(\vec{z}_n), \quad l_n(\vec{z}) = \sum_{i=1}^n \mathbf{1}_{C_{012}}(\vec{z}_n).$$

- If $\vec{z} \in E_i$, then $k_1(\vec{z}) = l_1(\vec{z}) = 0$. It follows from Lemma 4.2 that $a_1 = \vec{q}_i$. Let $\Omega_1 = \Omega$, $\Upsilon_1 = \Upsilon$.

- If $\vec{z} \in C_{ij}, i, j \in \{01, 12, 02\}$, then $k_1(\vec{z}) = 1, l_1(\vec{z}) = 0$. Let $\Upsilon_1 = \Upsilon$. It follows from Lemma 4.2 that $a_1 \in \{\vec{q}_i, \vec{q}_j\}$.
 - If $a_1 = \vec{q}_i$, we let $\Omega_1 = \{\omega \in \Omega : \omega_1 = 0\}$.
 - If $a_1 = \vec{q}_j$, we let $\Omega_1 = \{\omega \in \Omega : \omega_1 = 1\}$.
- If $\vec{z} \in C_{012}$, then $k_1(\vec{z}) = 0, l_1(\vec{z}) = 1$ and by Lemma 4.2, $a_1 = \vec{q}_i, i \in \{0, 1, 2\}$. We let $\Omega_1 = \Omega, \Upsilon_1 = \{v \in \Upsilon : v_1 = i\}$.

Obviously Ω_1 and Υ_1 are cylinders of length $k_1(\vec{z})$ and $l_1(\vec{z})$ respectively, and $d_1(\omega, v, \vec{z}) = a_1$ for all $\omega \in \Omega_1, v \in \Upsilon_1$. By a cylinder of length 0 we mean the whole space.

Now suppose we have obtained cylinders $\Omega_n \subseteq \dots \subseteq \Omega_1 \subseteq \Omega$ and $\Upsilon_n \subseteq \dots \subseteq \Upsilon_1 \subseteq \Upsilon$. Ω_n and Υ_n are cylinders of length $k_n(\vec{z})$ and $l_n(\vec{z})$ respectively. For all $\omega \in \Omega_n, v \in \Upsilon_n$, we have $d_1(\omega, v, \vec{z}) = a_1, \dots, d_n(\omega, v, \vec{z}) = a_n$ and $\vec{z}_{n+1} = \pi_3(K_\beta^n(\omega, v, \vec{z}))$.

- If $\vec{z}_{n+1} \in E_i$, then $k_{n+1}(\vec{z}) = k_n(\vec{z}), l_{n+1}(\vec{z}) = l_n(\vec{z})$ and by Lemma 4.2, $a_{n+1} = \vec{q}_i$. We let $\Omega_{n+1} = \Omega_n, \Upsilon_{n+1} = \Upsilon_n$.
- If $\vec{z}_{n+1} \in C_{ij}, i, j \in \{01, 12, 02\}$, then $k_{n+1}(\vec{z}) = k_n(\vec{z}) + 1, l_{n+1}(\vec{z}) = l_n(\vec{z})$. We let $\Upsilon_{n+1} = \Upsilon_n$. By Lemma 4.2, $a_{n+1} \in \{\vec{q}_i, \vec{q}_j\}$.
 - If $a_{n+1} = \vec{q}_i$, we let $\Omega_{n+1} = \{\omega \in \Omega_n : \omega_{k_{n+1}} = 0\}$.
 - If $a_{n+1} = \vec{q}_j$, we let $\Omega_{n+1} = \{\omega \in \Omega_n : \omega_{k_{n+1}} = 1\}$.
- If $\vec{z}_{n+1} \in C_{012}$, then $k_{n+1}(\vec{z}) = k_n(\vec{z}), l_{n+1}(\vec{z}) = l_n(\vec{z}) + 1$. By Lemma 4.2, $a_{n+1} = \vec{q}_i, i \in \{0, 1, 2\}$. Let $\Omega_{n+1} = \Omega_n$ and $\Upsilon_{n+1} = \{v \in \Upsilon_n : v_{l_{n+1}} = i\}$.

We can see that Ω_{n+1} is a cylinder of length $k_{n+1}(\vec{z})$ and Υ_{n+1} is a cylinder of length $l_{n+1}(\vec{z})$. For all $\omega \in \Omega_{n+1}, v \in \Upsilon_{n+1}$,

$$d_1(\omega, v, \vec{z}) = a_1, \dots, d_{n+1}(\omega, v, \vec{z}) = a_{n+1}.$$

If the map K_β hits $\Omega \times \Upsilon \times C$ infinitely many times, then $k_n(\vec{z}) \rightarrow \infty$ and $\cap \Omega_n$ consists of a single point. If this happens only finitely many times, then the set $\{k_n(\vec{z}) : n \in \mathbb{N}\}$ is finite and $\cap \Omega_n$ is a cylinder set. For $\cap \Upsilon_n$, it has the same property according to how many times K_β hits $\Omega \times \Upsilon \times C_{012}$.

In all cases $\cap \Omega_n$ and $\cap \Upsilon_n$ are non-empty, and for $\omega \in \cap \Omega_n, v \in \cap \Upsilon_n$, we have $d_j(\omega, v, \vec{z}) = a_j$ for all $j \geq 1$. \square

4.2. Unique K_β -invariant measure of maximal entropy for $\beta \in (1, \beta_*)$. Equip Υ with the uniform product measure \mathbb{P} and recall that σ' is the left shift on Υ . On the set $\Omega \times \Upsilon \times S_\beta$ we consider the product σ -algebra $\mathcal{A} \times \mathcal{B} \times \mathcal{S}$. Define the function $\rho_1 : \Omega \times \Upsilon \times S_\beta \rightarrow \{\vec{q}_0, \vec{q}_1, \vec{q}_2\}^\mathbb{N}$ by

$$\rho_1(\omega, v, \vec{z}) = (d_1(\omega, v, \vec{z}), d_2(\omega, v, \vec{z}), \dots).$$

Define the function $\rho_2 : \{\vec{q}_0, \vec{q}_1, \vec{q}_2\}^\mathbb{N} \rightarrow \Upsilon$ by

$$\rho_2(\vec{q}_{b_1}, \vec{q}_{b_2}, \vec{q}_{b_3}, \dots) = (b_1, b_2, b_3, \dots).$$

Denote by $\varphi = \rho_2 \circ \rho_1$ which is the function from $\Omega \times \Upsilon \times S_\beta$ to Υ . Then $\varphi \circ K_\beta = \sigma' \circ \varphi$, and φ is surjective from Theorem 4.3.

It is easily seen that φ is measurable. In fact, the inverse image of the cylinder set with the first digit fixed is measurable in $\Omega \times \Upsilon \times S_\beta$:

$$\begin{aligned}\varphi^{-1}(\{(b_1, b_2, \dots) \in \Upsilon : b_1 = 0\}) &= (\Omega \times \Upsilon \times E_0) \cup (\{\omega \in \Omega : \omega_1 = 0\} \times \Upsilon \times (C_{01} \cup C_{02})) \\ &\quad \cup (\Omega \times \{v \in \Upsilon : v_1 = 0\} \times C_{012}), \\ \varphi^{-1}(\{(b_1, b_2, \dots) \in \Upsilon : b_1 = 1\}) &= (\Omega \times \Upsilon \times E_1) \cup (\{\omega \in \Omega : \omega_1 = 1\} \times \Upsilon \times C_{01}) \\ &\quad \cup (\{\omega \in \Omega : \omega_1 = 0\} \times \Upsilon \times C_{12}) \cup (\Omega \times \{v \in \Upsilon : v_1 = 1\} \times C_{012}), \\ \varphi^{-1}(\{(b_1, b_2, \dots) \in \Upsilon : b_1 = 2\}) &= (\Omega \times \Upsilon \times E_2) \cup (\{\omega \in \Omega : \omega_1 = 1\} \times \Upsilon \times (C_{02} \cup C_{12})) \\ &\quad \cup (\Omega \times \{v \in \Upsilon : v_1 = 2\} \times C_{012}).\end{aligned}$$

To show that φ is an isomorphism, let

$$\begin{aligned}Z_1 &= \{(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times S_\beta : K_\beta^n(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times C \text{ infinitely often}\}, \\ Z_2 &= \{(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times S_\beta : K_\beta^n(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times C_{012} \text{ infinitely often}\}, \\ D_1 &= \{(b_1, b_2, \dots) \in \Upsilon : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \in C \text{ for infinitely many } j\text{'s}\}, \\ D_2 &= \{(b_1, b_2, \dots) \in \Upsilon : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \in C_{012} \text{ for infinitely many } j\text{'s}\}.\end{aligned}$$

Notice that

$$Z_1 = \cap_{n=1}^{\infty} \cup_{m=n}^{\infty} K_\beta^{-m}(\Omega \times \Upsilon \times C)$$

and

$$Z_2 = \cap_{n=1}^{\infty} \cup_{m=n}^{\infty} K_\beta^{-m}(\Omega \times \Upsilon \times C_{012}),$$

which imply that Z_1 and Z_2 are Borel sets in $\Omega \times \Upsilon \times S_\beta$. Let $Z = Z_1 \cap Z_2$, $D = D_1 \cap D_2$, then we have $K_\beta^{-1}(Z) = Z$, $(\sigma')^{-1}(D) = D$ and $\varphi(Z) = D$. Let $\varphi' = \varphi|_Z$.

Lemma 4.4. *The map $\varphi' : Z \rightarrow D$ is a bimeasurable bijection.*

Proof. For any sequence $(b_1, b_2, \dots) \in D$, we can obtain a point $\vec{z} = \sum_{i=1}^{\infty} \vec{q}_{b_i} \beta^{-i}$. To determine ω and v , we could define

$$\begin{aligned}r_1 &= \min\{j \geq 1 : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \in C\}, \quad r_k = \min\{j > r_{k-1} : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \in C\}, \\ s_1 &= \min\{j \geq 1 : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \in C_{012}\}, \quad s_k = \min\{j > s_{k-1} : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \in C_{012}\}.\end{aligned}$$

- If $\sum_{i=1}^{\infty} \vec{q}_{b_{r_k+i-1}} \beta^{-i} \in C_{ij}$, $ij \in \{01, 12, 02\}$, then $b_{r_k} \in \{i, j\}$ by Lemma 4.2.
 - If $b_{r_k} = i$, we let $\omega_k = 0$.
 - If $b_{r_k} = j$, we let $\omega_k = 1$.
- If $\sum_{i=1}^{\infty} \vec{q}_{b_{s_k+i-1}} \beta^{-i} \in C_{012}$, then $b_{s_k} \in \{0, 1, 2\}$ by Lemma 4.2. If $b_{s_k} = i$, we let $v_k = i$, $i \in \{0, 1, 2\}$.

Notice that for any $N > 0$, there exist $r, s > N$ such that

$$\sum_{i=1}^{\infty} \vec{q}_{b_{r+i-1}} \beta^{-i} \in C \text{ and } \sum_{i=1}^{\infty} \vec{q}_{b_{s+i-1}} \beta^{-i} \in C_{012},$$

which implies that $K_\beta^n(\omega, v, \vec{z})$ hits both $\Omega \times \Upsilon \times C$ and $\Omega \times \Upsilon \times C_{012}$ infinitely often. Then the infinite sequences $\omega = (\omega_1, \omega_2, \omega_3, \dots) \in \Omega$ and $v = (v_1, v_2, v_3, \dots) \in \Upsilon$

can be uniquely determined. Therefore, we can define the inverse of φ' . Let $(\varphi')^{-1} : D \rightarrow Z$ be

$$(\varphi')^{-1}((b_1, b_2, \dots)) = (\omega, v, \sum_{i=1}^{\infty} \frac{\vec{q}_{b_i}}{\beta^i}).$$

If $(\omega, v, \vec{z}) = (\omega', v', \vec{z}')$ then $\varphi'(\omega, v, \vec{z}) = \varphi'(\omega', v', \vec{z}')$. Since $Z = Z_1 \cap Z_2$ is a Borel set in $\Omega \times \Upsilon \times S_\beta$, then we have that $(\varphi')^{-1}$ is measurable by Theorem 2.2. Hence φ' is a bimeasurable bijection. \square

Let $\beta_* \approx 1.4656$ be the root of $x^3 - x^2 - 1 = 0$.

Lemma 4.5. *If $1 < \beta < \beta_*$, then $\mathbb{P}(D) = 1$.*

Proof. Let us first prove $\mathbb{P}(D_2) = 1$. For any sequence $(b_1, b_2, \dots) \in \Upsilon$ and $n \geq 4$, define

$$\vec{z}_n = (x_n, y_n) = \frac{\vec{q}_1}{\beta} + \frac{\vec{q}_2}{\beta^2} + \frac{\vec{q}_2}{\beta^3} + \frac{\vec{q}_0}{\beta^4} + \dots + \frac{\vec{q}_0}{\beta^n} + \frac{\vec{q}_{b_1}}{\beta^{n+1}} + \frac{\vec{q}_{b_2}}{\beta^{n+2}} + \dots.$$

Then we have $x_n \geq \frac{1}{\beta}$, and

$$y_n \geq \frac{1}{\beta^2} + \frac{1}{\beta^3},$$

$$x_n + y_n \leq \sum_{i=1}^3 \frac{1}{\beta^i} + \sum_{i=1}^{\infty} \frac{1}{\beta^{n+i}}.$$

If $1 < \beta < \beta_*$, then

$$\frac{1}{\beta^2} + \frac{1}{\beta^3} \geq \frac{1}{\beta},$$

and

$$\frac{1}{\beta} + \frac{1}{\beta^2} + \frac{1}{\beta^3} < \frac{1}{\beta(\beta-1)}.$$

Since $\sum_{i=1}^{\infty} \frac{1}{\beta^{n+i}} = \frac{1}{\beta^n(\beta-1)} \rightarrow 0$ as $n \rightarrow \infty$, there exists an integer $N > 3$ such that for any $n \geq N$,

$$y_n \geq \frac{1}{\beta},$$

$$x_n + y_n \leq \frac{1}{\beta(\beta-1)}.$$

It follows that $(x_n, y_n) \in C_{012}$ for any $n \geq N$.

Let

$$\mathbf{b} = 122 \underbrace{00 \dots 0}_{N-3 \text{ times}}.$$

Let

$$D' = \{(b_1, b_2, \dots) \in \Upsilon : b_j b_{j+1} \dots b_{j+N-1} = \mathbf{b} \text{ for infinitely many } j\}'s\}$$

then $D' \subset D_2$. Now we show that $\mathbb{P}(D') = 1$. Notice that $D' = \cap_{n=1}^{\infty} \cup_{m=n}^{\infty} D_m$, where $D_m = \{(b_1, b_2, \dots) \in \Upsilon : b_m b_{m+1} \dots b_{m+N-1} = \mathbf{b}\}$. Let $B_n = \cup_{m=n}^{\infty} D_m$. If $(b_1, b_2, \dots) \in \Upsilon \setminus B_n$, then we have that for any $j \geq n$, $b_j b_{j+1} \dots b_{j+N-1} \neq \mathbf{b}$. Clearly,

$$\Upsilon \setminus B_n \subseteq B' := \{(b_1, b_2, \dots) : b_{n+kN} \dots b_{n+(k+1)N-1} \neq \mathbf{b}, k = 0, 1, 2, \dots\}.$$

Since $\mathbb{P}(\Upsilon \setminus B_n) \leq \mathbb{P}(B') = \lim_{k \rightarrow \infty} (1 - 1/3^N)^k = 0$, then $\mathbb{P}(B_n) = 1$. It follows from $D' = \cap_{n=1}^{\infty} B_n$ and $B_1 \supseteq B_2 \supseteq \dots$ that $\mathbb{P}(D') = \lim_{n \rightarrow \infty} \mathbb{P}(B_n) = 1$. Then we get $\mathbb{P}(D_2) = 1$.

Similarly, define

$$\vec{z}_l = (x_l, y_l) = \frac{\vec{q}_1}{\beta} + \frac{\vec{q}_0}{\beta^2} + \dots + \frac{\vec{q}_0}{\beta^l} + \frac{\vec{q}_{b_1}}{\beta^{l+1}} + \frac{\vec{q}_{b_2}}{\beta^{l+2}} + \dots.$$

Then we have $x_l \geq \frac{1}{\beta}$, and

$$0 \leq y_l \leq \sum_{i=1}^{\infty} \frac{1}{\beta^{l+i}} = \frac{1}{\beta^l(\beta-1)},$$

$$\frac{1}{\beta} \leq x_l + y_l \leq \frac{1}{\beta} + \sum_{i=1}^{\infty} \frac{1}{\beta^{l+i}} = \frac{1}{\beta} + \frac{1}{\beta^l(\beta-1)}.$$

Since $\lim_{l \rightarrow \infty} \frac{1}{\beta^l(\beta-1)} = 0$, then there exists $L > 0$ such that for any $l \geq L$,

$$0 \leq y_l < \frac{1}{\beta}, x_l + y_l \leq \frac{1}{\beta(\beta-1)}.$$

It follows that $(x_l, y_l) \in C_{01}$ for any $l \geq L$. Let

$$(4.1) \quad D'' = \{(b_1, b_2, \dots) \in \Upsilon : b_j b_{j+1} \dots b_{j+L-1} = 1 \underbrace{00 \dots 0}_{L-1 \text{ times}} \text{ for infinitely many } j\},$$

then $D'' \subset D_1$. Since $\mathbb{P}(D'') = 1$, we have $\mathbb{P}(D_1) = 1$. Therefore, $\mathbb{P}(D) = 1$. \square

The following theorem can be obtained from Lemmas 4.4 and 4.5.

Theorem 4.6. *For $\beta \in (1, \beta_*]$, the dynamical systems $(\Omega \times \Upsilon \times S_\beta, \mathcal{A} \times \mathcal{B} \times \mathcal{S}, \nu_\beta, K_\beta)$ and $(\Upsilon, \mathcal{B}, \mathbb{P}, \sigma')$ are isomorphic, where $\nu_\beta(A) = \mathbb{P}(\varphi(Z \cap A))$.*

Remark 4.7. (i) Notice that if we change \mathbb{P} to the general product measure m_2 on Υ , Lemma 4.5 and Theorem 4.6 remain true.

(ii) Since \mathbb{P} is the unique measure of maximal entropy on Υ , the above theorem implies that any other K_β -invariant measure with support Z has entropy strictly less than $\log 3$. We now investigate the entropy of K_β -invariant measure μ for which $\mu(Z^c) > 0$.

Divide Z^c into three Borel sets as follows:

$$\begin{aligned} Z^c &= (Z_1 \cap Z_2)^c \\ &= Z_1^c \cup Z_2^c \\ &= (Z_1^c \setminus Z_2^c) \cup (Z_2^c \setminus Z_1^c) \cup (Z_1^c \cap Z_2^c) \\ &:= Z_3 \cup Z_4 \cup Z_5, \end{aligned}$$

where

$$\begin{aligned}
Z_3 &= (Z_1^c \setminus Z_2^c) \\
&= \{(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times S_\beta : K_\beta^n(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times C \text{ for finitely many } n\text{'s} \\
&\quad \text{and } K_\beta^n(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times C_{012} \text{ infinitely often}\}, \\
Z_4 &= (Z_2^c \setminus Z_1^c) \\
&= \{(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times S_\beta : K_\beta^n(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times C_{012} \text{ for finitely many } n\text{'s}, \\
&\quad \text{and } K_\beta^n(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times C \text{ infinitely often}\}, \\
Z_5 &= (Z_1^c \cap Z_2^c) \\
&= \{(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times S_\beta : K_\beta^n(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times C \text{ for finitely many } n\text{'s}, \\
&\quad \text{and } K_\beta^n(\omega, v, \vec{z}) \in \Omega \times \Upsilon \times C_{012} \text{ for finitely many } n\text{'s}\}.
\end{aligned}$$

We first prove the following lemma.

Lemma 4.8. *Let $\beta \in (1, \beta_*)$. Let μ_3 be a K_β -invariant measure for which $\mu_3(Z_3) = 1$. Then $h_{\mu_3}(K_\beta) < \log 3$. Similarly, let μ_4 and μ_5 be K_β -invariant measures for which $\mu_4(Z_4) = \mu_5(Z_5) = 1$. Then $h_{\mu_4}(K_\beta), h_{\mu_5}(K_\beta) < \log 3$ also holds.*

Proof. Let

$$\begin{aligned}
H_3 &= \{ \vec{z} = \sum_{i=1}^{\infty} \frac{\vec{q}_{b_i}}{\beta^i} \in S_\beta : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \text{ belongs to } C_{012} \text{ for infinitely many } j\text{'s}, \\
&\quad \text{and never belongs to } C \}.
\end{aligned}$$

Then $\Omega \times \Upsilon \times H_3 \subseteq K_\beta^{-1}(\Omega \times \Upsilon \times H_3)$ and $\cup_{i=0}^{\infty} K_\beta^{-i}(\Omega \times \Upsilon \times H_3) = Z_3$. It follows that $\mu_3(Z_3) = \lim_{i \rightarrow \infty} \mu_3(K_\beta^{-i}(\Omega \times \Upsilon \times H_3)) = 1$. Since μ_3 is K_β -invariant, then

$$\mu_3(\Omega \times \Upsilon \times H_3) = \mu_3(K_\beta^{-1}(\Omega \times \Upsilon \times H_3)) = \mu_3(K_\beta^{-2}(\Omega \times \Upsilon \times H_3)) = \cdots = 1.$$

Thus it is enough to study the entropy with respect to μ_3 of the map K_β restricted to $\Omega \times \Upsilon \times H_3$. Let π_1, π_2, π_3 be the canonical projection onto the three coordinates respectively. Notice that the action of the transformation K_β on the first coordinate is an identity, which implies that K_β is essentially a product transformation $I_\Omega \times K'_\beta$, where $K'_\beta = (\pi_2 \circ K_\beta) \times (\pi_3 \circ K_\beta)$ on $\Upsilon \times H_3$ and I_Ω is the identity on Ω . Since $(v, \vec{z}) \in \Upsilon \times S_\beta$ and $\omega \in \Omega$ are independent, it follows from $h_\mu(I_\Omega) = 0$ for any measure μ on (Ω, \mathcal{A}) and that

$$h_{\mu_3}(K_\beta) = h_{\mu'_3}(K'_\beta),$$

where $\mu'_3(B \times H) = \mu_3(\Omega \times B \times H)$ for $B \in \mathcal{B}, H \in (H_3 \cap \mathcal{S})$.

Let

$$\begin{aligned}
D_3 &= \{(b_1, b_2, \dots) \in \Upsilon : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \text{ belongs to } C_{012} \text{ for infinitely many } j\text{'s}, \\
&\quad \text{and never belongs to } C \}.
\end{aligned}$$

Define a map ϕ from $(\Upsilon \times H_3, \mathcal{B} \times (H_3 \cap \mathcal{S}), \mu'_3, K'_\beta)$ to $(D_3, D_3 \cap \mathcal{B}, \mu'_3 \circ \phi^{-1}, \sigma')$ as

$$\phi(v, \vec{z}) = \rho_2(\rho_1(0^\infty, v, \vec{z})).$$

Since $\vec{z} \in H_3$, then ϕ is well defined and bijective. ϕ is measurable and the inverse is also measurable by Theorem 2.2. Finally, ϕ preserves the measure and $\phi \circ K'_\beta = \sigma' \circ \phi$. Then ϕ is an isomorphism and it follows that

$$h_{\mu_3}(K_\beta) = h_{\mu'_3}(K'_\beta) = h_{\mu'_3 \circ \phi^{-1}}(\sigma') \leq h_{\mathbb{P}}(\sigma') = \log 3.$$

Since \mathbb{P} is the unique measure of maximal entropy on D_3 , to show $h_{\mu_3}(K_\beta) < \log 3$, it is enough to prove that $\mu'_3 \circ \phi^{-1} \neq \mathbb{P}$. This is done by contradiction. If $\mu'_3 \circ \phi^{-1} = \mathbb{P}$, then

$$\mathbb{P}(D_3) = \mu'_3(\Upsilon \times H_3) = \mu_3(\Omega \times \Upsilon \times H_3) = 1.$$

Since $D_3 \subset (D'')^c$, where D'' is defined as in (4.1), then $\mathbb{P}((D'')^c) = 1$, which is a contradiction to $\mathbb{P}(D'') = 1$. Therefore,

$$h_{\mu_3}(K_\beta) < \log 3.$$

Let

$$\begin{aligned} H_4 &= \left\{ \vec{z} = \sum_{i=1}^{\infty} \frac{\vec{q}_{b_i}}{\beta^i} \in S_\beta : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \text{ belongs to } C \text{ for infinitely many } j\text{'s,} \right. \\ &\quad \left. \text{and never belongs to } C_{012} \right\}, \\ H_5 &= \left\{ \vec{z} = \sum_{i=1}^{\infty} \frac{\vec{q}_{b_i}}{\beta^i} \in S_\beta : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \text{ never belongs to } C \right. \\ &\quad \left. \text{and never belongs to } C_{012} \right\}, \\ &= \{ \vec{z} \in S_\beta : \vec{z} \text{ has a unique } \beta\text{-expansion} \}. \end{aligned}$$

Then it follows that

$$\mu_4(\Omega \times \Upsilon \times H_4) = \mu_5(\Omega \times \Upsilon \times H_5) = 1.$$

We can also obtain that $h_{\mu_4}(K_\beta), h_{\mu_5}(K_\beta) < \log 3$ using the similar method. \square

From the above lemma we can obtain the upper bound of the entropy of K_β -invariant measure for which Z^c has positive measure.

Lemma 4.9. *Let $\beta \in (1, 3/2]$. Let μ be a K_β -invariant measure for which $\mu(Z^c) > 0$. Then $h_\mu(K_\beta) < \log 3$.*

Proof. Notice that Z, Z_3, Z_4 and Z_5 are pairwise disjoint and the union is $\Omega \times \Upsilon \times S_\beta$. Since Z, Z_3, Z_4 and Z_5 are K_β -invariant, then there exist K_β -invariant probability measures μ_{12}, μ_3, μ_4 and μ_5 concentrated on Z, Z_3, Z_4 and Z_5 , respectively, such that

$$\mu = (1 - \alpha_3 - \alpha_4 - \alpha_5)\mu_{12} + \alpha_3\mu_3 + \alpha_4\mu_4 + \alpha_5\mu_5,$$

where $0 \leq \alpha_3, \alpha_4, \alpha_5 \leq 1$ and $0 < \alpha_3 + \alpha_4 + \alpha_5 \leq 1$. Then

$$h_\mu(K_\beta) = (1 - \alpha_3 - \alpha_4 - \alpha_5)h_{\mu_{12}}(K_\beta) + \alpha_3h_{\mu_3}(K_\beta) + \alpha_4h_{\mu_4}(K_\beta) + \alpha_5h_{\mu_5}(K_\beta).$$

Since $h_{\mu_{12}}(K_\beta) \leq \log 3$ by Remark 4.7 and $h_{\mu_3}(K_\beta), h_{\mu_4}(K_\beta), h_{\mu_5}(K_\beta) < \log 3$ by Lemma 4.8, then the result follows. \square

Now we obtain the main result in this section.

Theorem 4.10. *Let $\beta \in (1, \beta_*)$. The measure $\nu_\beta(A) = \mathbb{P}(\varphi(Z \cap A))$ is the unique K_β -invariant measure of maximal entropy.*

5. AN ABSOLUTELY CONTINUOUS INVARIANT MEASURE FOR K_β WHEN $1 < \beta \leq \frac{3}{2}$

The analysis is similar in the two cases: $\beta \in (1, 3/2)$ and $\beta = 3/2$. Without special statement we assume $\beta \in (1, 3/2)$ in this section. A brief description of the case of $\beta = 3/2$ is given in the remarks. Recall the product measure m_1 on Ω with weights $\{p, 1-p\}$ and m_2 on Υ with weights $\{s, t, 1-s-t\}$. Consider the measure space $(S_\beta, \mathcal{S}, \lambda_2)$, where λ_2 is the normalized Lebesgue measure. In this section we will prove that K_β has an invariant measure of the form $m_1 \otimes m_2 \otimes \mu_\beta$, where μ_β is absolutely continuous with respect to λ_2 . We will show the result by several steps.

Step 1: a position dependent random transformation R .

Bahsoun and Góra [3], gave a sufficient condition for the existence of an absolutely continuous invariant measure for a random map with position dependent probabilities on a bounded domain of \mathbb{R}^N . We take some of their results a little further.

For $k = 1, \dots, K$, let $\tau_k : S_\beta \rightarrow S_\beta$ be piecewise one-to-one and C^2 , non-singular transformations on a common partition \mathcal{P} of S_β : $\mathcal{P} = \{S_1, \dots, S_q\}$ and $\tau_{k,i} = \tau_k|_{S_i}$, $i = 1, \dots, q$. Let $p_k : S_\beta \rightarrow [0, 1]$ be piecewise C^1 functions such that $\sum_{k=1}^K p_k = 1$. Denote by $R = \{\tau_1, \dots, \tau_K; p_1(\vec{z}), \dots, p_K(\vec{z})\}$ the position dependent random map, i.e., $R(\vec{z}) = \tau_k(\vec{z})$ with probability $p_k(\vec{z})$. Define the transition function for R as follows:

$$\mathbf{P}(\vec{z}, A) = \sum_{k=1}^K p_k(\vec{z}) \mathbb{1}_A(\tau_k(\vec{z})),$$

where A is any measurable set and $\mathbb{1}_A$ denotes the indicator function of the set A .

The iteration of R is denoted by $R^n := \{\tau_{k_1 k_2 \dots k_n}; p_{k_1 k_2 \dots k_n}\}$, $k_1 k_2 \dots k_n \in \{1, 2, \dots, K\}^n$, where

$$\tau_{k_1 k_2 \dots k_n}(\vec{z}) = \tau_{k_n} \circ \tau_{k_{n-1}} \circ \dots \circ \tau_{k_1}(\vec{z})$$

and

$$p_{k_1 k_2 \dots k_n}(\vec{z}) = p_{k_n}(\tau_{k_{n-1}} \circ \dots \circ \tau_{k_1}(\vec{z})) \cdot p_{k_{n-1}}(\tau_{k_{n-2}} \circ \dots \circ \tau_{k_1}(\vec{z})) \dots p_{k_1}(\vec{z}).$$

The transition function \mathbf{P} induces an operator \mathbf{P}_* on the set of probability measure on (S_β, \mathcal{S}) defined by

$$\mathbf{P}_* \mu(A) = \int \mathbf{P}(\vec{z}, A) d\mu(\vec{z}) = \sum_{k=1}^K \int_{\tau_k^{-1}(A)} p_k(\vec{z}) d\mu(\vec{z}) = \sum_{k=1}^K \sum_{i=1}^q \int_{\tau_{k,i}^{-1}(A)} p_k(\vec{z}) d\mu(\vec{z}).$$

We say that the measure μ is R -invariant iff $\mathbf{P}_* \mu = \mu$.

If μ has density f with respect to λ_2 , then $\mathbf{P}_* \mu$ has also a density which we denote by $P_R f$, i.e.,

$$\int_A P_R f(\vec{z}) d\lambda_2(\vec{z}) = \sum_{k=1}^K \sum_{i=1}^q \int_{\tau_{k,i}^{-1}(A)} p_k(\vec{z}) f(\vec{z}) d\lambda_2(\vec{z}).$$

We call P_R the Perron-Frobenius operator of the random map R and it has very useful properties[3]:

- (i) P_R is linear;
- (ii) P_R is non-negative;
- (iii) $P_R f = f \iff \mu = f \cdot \lambda_2$ is R -invariant;

- (iv) $\|P_R f\|_1 \leq \|f\|_1$, where $\|\cdot\|_1$ denotes the L^1 norm;
- (v) $P_{R \circ T} = P_R \circ P_T$. In particular, $P_R^N = P_{R^N}$.

Let each S_i be a bounded closed domain having a piecewise C^2 boundary of finite 1-dimensional measure. Assume that the faces of ∂S_i meet at angles bounded uniformly away from 0 and the probabilities $p_k(\vec{z})$ are piecewise C^1 functions on the partition \mathcal{P} . We assume:

Condition(A):

$$\max_{1 \leq i \leq q} \sum_{k=1}^K p_k(\vec{z}) \|D\tau_{k,i}^{-1}(\tau_{k,i}(\vec{z}))\| < c < 1,$$

where $D\tau_{k,i}^{-1}(\vec{z})$ is the derivative matrix of $\tau_{k,i}^{-1}$ at \vec{z} .

Using the multidimensional notion of variation [13]:

$$V(f) = \int_{\mathbb{R}^N} \|Df\| d\lambda_N = \sup \left\{ \int_{\mathbb{R}^N} f \operatorname{div}(g) d\lambda_N : g = (g_1, \dots, g_N) \in C_0^1(\mathbb{R}^N, \mathbb{R}^N) \right\},$$

where $f \in L_1(\mathbb{R}^N)$ has bounded support, Df denotes the gradient of f in the distributional sense, $\operatorname{div}(g) = \nabla \cdot g = \frac{\partial g_1}{\partial x_1} + \frac{\partial g_2}{\partial x_2} + \dots + \frac{\partial g_N}{\partial x_N}$ is the divergence operator, and $C_0^1(\mathbb{R}^N, \mathbb{R}^N)$ is the space of continuously differentiable functions from \mathbb{R}^N into \mathbb{R}^N having compact support. Consider the Banach space [13, Remark 1.12],

$$BV(S_\beta) = \{f \in L_1(S_\beta) : V(f) < +\infty\},$$

with the norm $\|f\|_{BV} = \|f\|_{L_1} + V(f)$.

Let $a(S_i)$ be defined as in (3.5) and $\gamma(S_i)$ defined as in (3.4). Now we start at points $y \in F$, where the minimal angle $\gamma(S_i)$ is attained, defining L_y to be central rays of the largest regular cones contained in S_i . Then we extend this field of segments to C^1 field, making L_y short enough to avoid overlapping. Let $\delta(y)$ be the length of $L_y, y \in \partial S_i$. By the compactness of ∂S_i we have

$$\delta(S_i) = \inf_{y \in \partial(S_i)} \delta(y) > 0.$$

Let \vec{z} be a point in ∂S_i and $J_{k,i}$ the Jacobian of $\tau_{k,i}$ at \vec{z} .

We recall the following theorem.

Theorem 5.1. [3, Theorem 6.3] *If R is a random map which satisfies Condition (A), then*

$$V(P_R f) \leq c(1 + 1/a)V(f) + (M + \frac{c}{a\delta})\|f\|_1,$$

where $a = \min\{a(S_i) : i = 1, \dots, q\} > 0, \delta = \min\{\delta(S_i) : i = 1, \dots, q\} > 0, M_{k,i} = \sup_{\vec{z} \in S_i} (Dp_k(\vec{z}) - \frac{DJ_{k,i}}{J_{k,i}} p_k(\vec{z}))$ and $M = \sum_{k=1}^K \max_{1 \leq i \leq q} M_{k,i}$.

Now, let R be a random map which is given by $\{\tau_1, \dots, \tau_6; p_1(\vec{z}), \dots, p_6(\vec{z})\}$ where

$$\begin{aligned}\tau_1(\vec{z}) &= \begin{cases} \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in E_i, i \in \{0, 1, 2\} \\ \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\} \\ \beta\vec{z}, & \text{if } \vec{z} \in C_{012}, \end{cases} \\ \tau_2(\vec{z}) &= \begin{cases} \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in E_i, i \in \{0, 1, 2\} \\ \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\} \\ \beta\vec{z} - \vec{q}_1, & \text{if } \vec{z} \in C_{012}, \end{cases} \\ \tau_3(\vec{z}) &= \begin{cases} \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in E_i, i \in \{0, 1, 2\} \\ \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\} \\ \beta\vec{z} - \vec{q}_2, & \text{if } \vec{z} \in C_{012}, \end{cases} \\ \tau_4(\vec{z}) &= \begin{cases} \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in E_i, i \in \{0, 1, 2\} \\ \beta\vec{z} - \vec{q}_j, & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\} \\ \beta\vec{z}, & \text{if } \vec{z} \in C_{012}, \end{cases} \\ \tau_5(\vec{z}) &= \begin{cases} \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in E_i, i \in \{0, 1, 2\} \\ \beta\vec{z} - \vec{q}_j, & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\} \\ \beta\vec{z} - \vec{q}_1, & \text{if } \vec{z} \in C_{012}, \end{cases} \\ \tau_6(\vec{z}) &= \begin{cases} \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in E_i, i \in \{0, 1, 2\} \\ \beta\vec{z} - \vec{q}_j, & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\} \\ \beta\vec{z} - \vec{q}_2, & \text{if } \vec{z} \in C_{012}, \end{cases}\end{aligned}$$

The probabilities are defined as follows.

$$(5.1) \quad \begin{aligned} p_1(\vec{z}) &= p \cdot s, & p_4(\vec{z}) &= (1-p) \cdot s, \\ p_2(\vec{z}) &= p \cdot t, & p_5(\vec{z}) &= (1-p) \cdot t, \\ p_3(\vec{z}) &= p \cdot (1-s-t), & p_6(\vec{z}) &= (1-p) \cdot (1-s-t). \end{aligned}$$

We have the following lemma.

Lemma 5.2. *For any $\vec{z} \in S_\beta$, $\sum_{k_1 k_2 \dots k_n \in \{1, \dots, 6\}^n} p_{k_1 k_2 \dots k_n}(\vec{z}) = 1, n \in \mathbb{Z}$.*

Proof. We prove this lemma by induction. For $n = 1$, $p_1(\vec{z}) + \dots + p_6(\vec{z}) = 1$. Assume it is true for $n = m$, i.e. for any $\vec{z} \in S_\beta$,

$$\sum_{k_1 k_2 \dots k_m \in \{1, \dots, 6\}^m} p_{k_1 k_2 \dots k_m}(\vec{z}) = 1.$$

For $n = m + 1$,

$$\begin{aligned}
& \sum_{k_1 k_2 \dots k_{m+1} \in \{1, \dots, 6\}^{m+1}} p_{k_1 k_2 \dots k_{m+1}}(\vec{z}) \\
&= \sum_{k_1 k_2 \dots k_{m+1} \in \{1, \dots, 6\}^{m+1}} p_{k_{m+1}}(\tau_{k_m} \circ \dots \circ \tau_{k_1}(\vec{z})) \cdot p_{k_m}(\tau_{k_{m-1}} \circ \dots \circ \tau_{k_1}(\vec{z})) \cdots p_{k_1}(\vec{z}) \\
&= \sum_{1 k_2 \dots k_{m+1} \in \{1, \dots, 6\}^{m+1}} p_{k_{m+1}}(\tau_{k_m} \circ \dots \circ \tau_{k_2}(\tau_1(\vec{z}))) \cdots p_{k_2}(\tau_1(\vec{z})) \cdot p_1(\vec{z}) \\
&\quad + \sum_{2 k_2 \dots k_{m+1} \in \{1, \dots, 6\}^{m+1}} p_{k_{m+1}}(\tau_{k_m} \circ \dots \circ \tau_{k_2}(\tau_2(\vec{z}))) \cdots p_{k_2}(\tau_2(\vec{z})) \cdot p_2(\vec{z}) \\
&\quad + \cdots + \sum_{6 k_2 \dots k_{m+1} \in \{1, \dots, 6\}^{m+1}} p_{k_{m+1}}(\tau_{k_m} \circ \dots \circ \tau_{k_2}(\tau_6(\vec{z}))) \cdots p_{k_2}(\tau_6(\vec{z})) \cdot p_6(\vec{z}) \\
&= p_1(\vec{z}) \sum_{k_2 \dots k_{m+1} \in \{1, \dots, 6\}^m} p_{k_2 \dots k_{m+1}}(\tau_1(\vec{z})) + p_2(\vec{z}) \sum_{k_2 \dots k_{m+1} \in \{1, \dots, 6\}^m} p_{k_2 \dots k_{m+1}}(\tau_2(\vec{z})) \\
&\quad + \cdots + p_6(\vec{z}) \sum_{k_2 \dots k_{m+1} \in \{1, \dots, 6\}^m} p_{k_2 \dots k_{m+1}}(\tau_6(\vec{z})) \\
&= p_1(\vec{z}) + \cdots + p_6(\vec{z}) \\
&= 1.
\end{aligned}$$

□

Now we can prove the existence of an acim for R .

Theorem 5.3. *Let $R = \{\tau_1, \dots, \tau_6; p_1(\vec{z}), \dots, p_6(\vec{z})\}$, then R admits an acim.*

Proof. Denote the partition (3.1) by \mathcal{P} with

$$S_1 = E_0, S_2 = E_1, S_3 = E_2, S_4 = C_{01}, S_5 = C_{12}, S_6 = C_{02}, S_7 = C_{012}.$$

Consider the iteration of the random map, R^n , the corresponding partition is $\vee_{i=0}^{n-1} R^{-i} \mathcal{P}$, where

$$R^{-i} \mathcal{P} = \vee_{k_1 k_2 \dots k_i \in \{1, \dots, 6\}^i} \tau_{k_1 k_2 \dots k_i}^{-1} \mathcal{P}.$$

For a set $P_i \in \vee_{i=0}^{n-1} R^{-i} \mathcal{P}$ and a sequence $k_1 \dots k_n \in \{1, \dots, 6\}^n$, let $\tau_{k_1 \dots k_n, i} = \tau_{k_1 \dots k_n}|_{P_i}$ and $M_{k_1 \dots k_n, i} = \sup_{\vec{z} \in P_i} (Dp_{k_1 \dots k_n}(\vec{z}) - \frac{DJ_{k_1 \dots k_n, i}}{J_{k_1 \dots k_n, i}} p_{k_1 \dots k_n, i}(\vec{z}))$, where $J_{k_1 \dots k_n, i}$ is the Jacobian of $\tau_{k_1 \dots k_n, i}$. Let

$$M_n = \sum_{k_1 \dots k_n \in \{0, 1, 2\}^n} \max_{P_i \in \vee_{i=0}^{n-1} R^{-i} \mathcal{P}} M_{k_1 \dots k_n, i} \quad \text{and} \quad \delta_n = \min_{P_i \in \vee_{i=0}^{n-1} R^{-i} \mathcal{P}} \delta(P_i).$$

For any set $P_i \in \vee_{i=0}^{n-1} R^{-i} \mathcal{P}$, the derivative matrix of $\tau_{k_1 k_2 \dots k_n}^{-1}$ is equal to

$$\begin{bmatrix} \frac{1}{\beta^n} & 0 \\ 0 & \frac{1}{\beta^n} \end{bmatrix}.$$

Using Lemma 5.2 we have

$$\max_{P_i \in \vee_{i=0}^{n-1} R^{-i} \mathcal{P}} \sum_{k_1 k_2 \dots k_n \in \{1, \dots, 6\}^n} p_{k_1 k_2 \dots k_n}(\vec{z}) \|D(\tau_{k_1 k_2 \dots k_n}|_{P_i})^{-1}\| = \frac{\sqrt{2}}{\beta^n} < \frac{2\sqrt{2}}{\beta^n} := c_n.$$

For the partition $\vee_{i=0}^{n-1} R^{-i}\mathcal{P}$, we have $a_n = \sqrt{2}/2$. Let

$$r_n = c_n(1 + \frac{1}{a_n}) = \frac{2\sqrt{2} + 4}{\beta^n}, \quad R_n = M_n + \frac{c_n}{a_n \delta_n}.$$

We can find $l > \log(2\sqrt{2} + 4)/\log \beta$ such that $r_l < 1$. Fix this l and let $C_1 = \max\{r_1, r_2, \dots, r_{l-1}\}$, $C_2 = \max\{R_1, R_2, \dots, R_{l-1}\}$. For any integer n , we have $n = jl + i$, where $0 \leq i \leq l-1$. Notice that $P_{R^n} = (P_{R^l})^j P_{R^i}$. Apply Theorem 5.1 on R^l , then we get

$$\begin{aligned} V(P_{R^n} f) &= V P_{R^l}^j (P_{R^i} f) \\ &\leq r_l \cdot V P_{R^l}^{j-1} (P_{R^i} f) + R_l \|f\|_1 \\ &\leq r_l \cdot (r_l \cdot V P_{R^l}^{j-2} (P_{R^i} f) + R_l \|f\|_1) + R_l \|f\|_1 \\ &\dots \\ &\leq r_l^j V(P_{R^i} f) + (r_l^{j-1} + r_l^{j-2} + \dots + r_l + 1) R_l \|f\|_1 \\ &\leq r_l^j (C_1 V(f) + C_2 \|f\|_1) + (r_l^{j-1} + r_l^{j-2} + \dots + r_l + 1) R_l \|f\|_1 \\ &= C_1 r_l^j V(f) + (C_2 r_l^j + r_l^{j-1} + r_l^{j-2} + \dots + r_l + 1) R_l \|f\|_1 \\ &\leq C_1 r_l^j V(f) + (C_2 + \frac{1}{1-r_l}) R_l \|f\|_1. \end{aligned}$$

By definition of the norm $\|\cdot\|_{BV}$,

$$\begin{aligned} \|P_{R^n} f\|_{BV} &= \|P_{R^n} f\|_1 + V(P_{R^n} f) \\ &\leq \|f\|_1 + C_1 r_l^j V(f) + (C_2 + \frac{1}{1-r_l}) R_l \|f\|_1. \end{aligned}$$

Then the result follows by the technique in [4, Theorem 1]. We write some details for completeness. From the above inequality it follows that the set $\{P_R^n \mathbf{1}\}_{n \geq l}$ is uniformly bounded, where $\mathbf{1}$ is the constant function equal to 1 on S_β . Hence P_R has a nontrivial fixed point $\mathbf{1}^*$ which is the density of an acim by the Kakutani-Yoshida Theorem. \square

Step 2: for the skew product transformation R' on $S_\beta \times [0, 1)$.

Let $(I, \mathcal{B}(I), \lambda_1)$ be the unit interval $I = [0, 1)$, with $\mathcal{B}(I)$ the Borel σ -algebra on I and λ_1 being Lebesgue measure on $(I, \mathcal{B}(I))$. Let $Y = S_\beta \times I$ and the set J_k be given by $J_k = \{(\vec{z}, w) : \sum_{i < k} p_i(\vec{z}) \leq w < \sum_{i \leq k} p_i(\vec{z})\}$. Define maps $\varphi_k : J_k \rightarrow I$ by

$$\varphi_k(\vec{z}, w) = \frac{1}{p_k(\vec{z})} w - \frac{\sum_{r=1}^{k-1} p_r(\vec{z})}{p_k(\vec{z})}.$$

Define the skew product transformation $R' : S_\beta \times I \rightarrow S_\beta \times I$ by

$$R'(\vec{z}, w) = (\tau_k(x), \varphi_k(\vec{z}, w))$$

for $(\vec{z}, w) \in J_k$.

Since $p_k(\vec{z})$ is defined as in (5.1), then we have

$$\begin{aligned}\varphi_1(\vec{z}, w) &= \frac{w}{ps}, & \varphi_4(\vec{z}, w) &= \frac{w-p}{(1-p)s}, \\ \varphi_2(\vec{z}, w) &= \frac{w-ps}{pt}, & \varphi_5(\vec{z}, w) &= \frac{w-p-(1-p)s}{(1-p)t}, \\ \varphi_3(\vec{z}, w) &= \frac{w-ps-pt}{p(1-s-t)}, & \varphi_6(\vec{z}, w) &= \frac{w-p-(1-p)s-(1-p)t}{(1-p)(1-s-t)}.\end{aligned}$$

We denote $p_k(\vec{z})$ and $\varphi_k(\vec{z}, w)$ by p_k and $\varphi_k(w)$, respectively, since each $p_k(\vec{z})$ is a constant. Therefore,

$$R'(\vec{z}, w) = \begin{cases} (\tau_1(\vec{z}), \varphi_1(w)), & \text{if } w \in [0, ps), \\ (\tau_2(\vec{z}), \varphi_2(w)), & \text{if } w \in [ps, ps+pt), \\ (\tau_3(\vec{z}), \varphi_3(w)), & \text{if } w \in [ps+pt, p), \\ (\tau_4(\vec{z}), \varphi_4(w)), & \text{if } w \in [p, p+(1-p)s), \\ (\tau_5(\vec{z}), \varphi_5(w)), & \text{if } w \in [p+(1-p)s, p+(1-p)s+(1-p)t), \\ (\tau_6(\vec{z}), \varphi_6(w)), & \text{if } w \in [p+(1-p)s+(1-p)t, 1). \end{cases}$$

Denote by μ_β an acim for the position dependent random transformation $R = \{\tau_1, \dots, \tau_6; p_1, \dots, p_6\}$, which means μ_β is R -invariant and absolutely continuous with respect to Lebesgue measure λ_2 in \mathbb{R}^2 . We start by recalling Lemma 3.2 in [1].

Lemma 5.4. μ_β is invariant for the random map R if and only if $\mu_\beta \otimes \lambda_1$ is invariant for the skew product R' .

Step 3: for the skew product transformation R_β on $\Omega \times \Upsilon \times S_\beta$.

Define the skew product transformation R_β on $\Omega \times \Upsilon \times S_\beta$ as follows:

$$R_\beta(\omega, v, \vec{z}) = \begin{cases} (\sigma\omega, \sigma'v, \beta\vec{z} - \vec{q}_i), & \text{if } \vec{z} \in E_i, i \in \{0, 1, 2\} \\ (\sigma\omega, \sigma'v, \beta\vec{z} - \vec{q}_i), & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\} \text{ and } \omega_1 = 0 \\ (\sigma\omega, \sigma'v, \beta\vec{z} - \vec{q}_j), & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\} \text{ and } \omega_1 = 1 \\ (\sigma\omega, \sigma'v, \beta\vec{z} - \vec{q}_i), & \text{if } \vec{z} \in C_{012}, v_1 = i, i \in \{0, 1, 2\}. \end{cases}$$

Lemma 5.5. $(S_\beta \times I, \mathcal{S} \times \mathcal{B}(I), \mu_\beta \otimes \lambda_1, R')$ and $(\Omega \times \Upsilon \times S_\beta, \mathcal{A} \times \mathcal{B} \times \mathcal{S}, m_1 \otimes m_2 \otimes \mu_\beta, R_\beta)$ are isomorphic.

Proof. Let $\pi_2 : S_\beta \times I \rightarrow I$ be the canonical projection onto the second coordinate. Consider the map $\varphi = \pi_2 \circ R'$ on $(I, \mathcal{B}(I), \lambda_1)$. One can see that $\varphi(w) = \varphi_k(w)$ for $w \in I_k$, where $I_1 = [0, p_1)$ and $I_k = \left[\sum_{i=1}^{k-1} p_i, \sum_{i=1}^k p_i \right)$ for $2 \leq k \leq 6$. Define

$$l(w) = \frac{1}{p_k} \quad \text{and} \quad h(w) = \frac{\sum_{i=1}^{k-1} p_i}{p_k}$$

for $w \in I_k$. It follows that $\varphi(w) = l(w) \cdot w - h(w)$. Let

$$l_n = l_n(w) := l(\varphi^{n-1}(w)) \quad \text{and} \quad h_n = h_n(w) := h(\varphi^{n-1}(w)).$$

For $w \in [0, 1)$, we can write the generalized Lüroth series (GLS) of w , which is

$$w = \frac{h_1}{l_1} + \frac{h_2}{l_1 l_2} + \dots + \frac{h_n}{l_1 \dots l_n} + \dots.$$

Consider the system $\{\{0, 1, 2, 3, 4, 5\}^{\mathbb{N}}, \mathcal{C}, m, \sigma''\}$, where \mathcal{C} is the product σ -algebra, σ'' is the left shift and m is the product measure with weights $\{p_1, \dots, p_6\}$ as (5.1). Let $\phi_1 : I \rightarrow \{0, 1, 2, 3, 4, 5\}^{\mathbb{N}}$ be given by

$$\phi_1 : w = \sum_{n=1}^{\infty} \frac{h_n}{l_1 l_2 \dots l_n} \mapsto (\gamma_1, \gamma_2, \dots),$$

where $\gamma_n = \gamma_n(w)$, $n \geq 1$ is defines as follows:

$$\gamma_n := \gamma_n(w) = k - 1 \iff \varphi^{n-1}(w) \in I_k,$$

for $k \in \{1, 2, 3, 4, 5, 6\}$. It is known that φ preserves the Lebesgue measure λ_1 and ϕ_1 is an isomorphism between the two dynamical systems $(I, \mathcal{B}(I), \lambda_1, \varphi)$ and $\{\{0, 1, 2, 3, 4, 5\}^{\mathbb{N}}, \mathcal{C}, m, \sigma''\}$. See [2] for more details.

Next we give a map ϕ_2 from $\{\{0, 1, 2, 3, 4, 5\}^{\mathbb{N}}, \mathcal{C}, m, \sigma''\}$ to $\{\Omega \times \Upsilon, \mathcal{A} \times \mathcal{B}, m_1 \otimes m_2, \sigma \times \sigma'\}$. Let $h_1 : \{0, 1, 2, 3, 4, 5\} \rightarrow \{0, 1\}$ and $h_2 : \{0, 1, 2, 3, 4, 5\} \rightarrow \{0, 1, 2\}$ be given by

$$h_1(x) = \begin{cases} 0, & \text{if } x = 0, 1, 2, \\ 1, & \text{if } x = 3, 4, 5, \end{cases}, \quad h_2(x) = \begin{cases} 0, & \text{if } x = 0, 3, \\ 1, & \text{if } x = 1, 4, \\ 2, & \text{if } x = 2, 5. \end{cases}$$

Define $\phi_2 : \{0, 1, 2, 3, 4, 5\}^{\mathbb{N}} \rightarrow \Omega \times \Upsilon$ by $\phi_2(\gamma) = (\omega, v)$, where

$$\begin{aligned} \omega &= (h_1(\gamma_1), h_1(\gamma_2), h_1(\gamma_3), \dots) := \tilde{h}_1(\gamma), \\ v &= (h_2(\gamma_1), h_2(\gamma_2), h_2(\gamma_3), \dots) := \tilde{h}_2(\gamma). \end{aligned}$$

One can see that ϕ_2 maps a cylinder of rank n in $\{0, 1, 2, 3, 4, 5\}^{\mathbb{N}}$ to the product of two cylinders of the same rank n in $\Omega \times \Upsilon$. It follows that ϕ_2 is a bimeasurable bijection. From the definition of the product measure, we can get the measure preservingness on cylinders. Finally, it is easy to see that $\phi_2 \circ \sigma'' = (\sigma \times \sigma') \circ \phi_2$. Therefore, ϕ_2 is an isomorphism.

Now let $\phi : S_\beta \times I \rightarrow \Omega \times \Upsilon \times S_\beta$ be given by

$$\phi(\vec{z}, w) = (\tilde{h}_1(\phi_1(w)), \tilde{h}_2(\phi_1(w)), \vec{z}).$$

In fact, $\phi = \iota \circ (I_{S_\beta} \times (\phi_2 \circ \phi_1))$, where I_{S_β} is the identity map on S_β and $\iota(\vec{z}, \omega, v) = (\omega, v, \vec{z})$ is a transformation that only changes the order of coordinates. Since $\phi_2 \circ \phi_1$ preserves the dynamics of $\pi_2 \circ R$ and $\sigma \times \sigma'$, i.e.,

$$(\phi_2 \circ \phi_1) \circ (\pi_2 \circ R) = (\sigma \times \sigma') \circ (\phi_2 \circ \phi_1),$$

we have that $\phi \circ R' = R_\beta \circ \phi$. Therefore, the result follows. \square

Step 4: for the random transformation K_β on $\Omega \times \Upsilon \times S_\beta$.

Define a skew product transformation R_β as follows:

$$R_\beta(\omega, v, \vec{z}) = \begin{cases} (\sigma\omega, \sigma'v, \tau_1(\vec{z})), & \text{if } \omega_1 = 0, v_1 = 0, \\ (\sigma\omega, \sigma'v, \tau_2(\vec{z})), & \text{if } \omega_1 = 0, v_1 = 1, \\ (\sigma\omega, \sigma'v, \tau_3(\vec{z})), & \text{if } \omega_1 = 0, v_1 = 2, \\ (\sigma\omega, \sigma'v, \tau_4(\vec{z})), & \text{if } \omega_1 = 1, v_1 = 0, \\ (\sigma\omega, \sigma'v, \tau_5(\vec{z})), & \text{if } \omega_1 = 1, v_1 = 1, \\ (\sigma\omega, \sigma'v, \tau_6(\vec{z})), & \text{if } \omega_1 = 1, v_1 = 2, \end{cases}$$

Let μ be an arbitrary probability measure on S_β . We will show that any product measure of the form $m_1 \otimes m_2 \otimes \mu$ is K_β -invariant if and only if it is R_β -invariant.

Lemma 5.6. $m_1 \otimes m_2 \otimes \mu \circ K_\beta^{-1} = m_1 \otimes m_2 \otimes \mu \circ R_\beta^{-1} = m_1 \otimes m_2 \otimes \nu$, where

$$\begin{aligned} \nu = & ps \cdot \mu \circ \tau_1^{-1} + pt \cdot \mu \circ \tau_2^{-1} + p(1-s-t) \cdot \mu \circ \tau_3^{-1} \\ & + (1-p) \cdot s \cdot \mu \circ \tau_4^{-1} + (1-p) \cdot t \cdot \mu \circ \tau_5^{-1} + (1-p) \cdot (1-s-t) \cdot \mu \circ \tau_6^{-1}. \end{aligned}$$

Proof. Denote by C_1 and C_2 arbitrary cylinders in Ω and Υ , respectively. Let S be a closed set in S_β . It suffices to verify that the measures coincide on sets of the form $C_1 \times C_2 \times S$, because the collection of these sets forms a generating π -system. Let $[i, C_1] = \{\omega_1 = i\} \cap \sigma^{-1}(C_1)$ for $i = 0, 1$ and $[i, C_2] = \{v_1 = i\} \cap (\sigma')^{-1}(C_2)$ for $i = 0, 1, 2$. Notice that

$$\begin{aligned} \tau_1(S) \cap E &= \tau_2(S) \cap E = \tau_3(S) \cap E = \tau_4(S) \cap E = \tau_5(S) \cap E = \tau_6(S) \cap E, \\ \tau_1(S) \cap C &= \tau_2(S) \cap C = \tau_3(S) \cap C, \\ \tau_4(S) \cap C &= \tau_5(S) \cap C = \tau_6(S) \cap C, \\ \tau_1(S) \cap C_{012} &= \tau_4(S) \cap C_{012}, \\ \tau_2(S) \cap C_{012} &= \tau_5(S) \cap C_{012}, \\ \tau_3(S) \cap C_{012} &= \tau_6(S) \cap C_{012}. \end{aligned}$$

We can get

$$\begin{aligned} K_\beta^{-1}(C_1 \times C_2 \times S) = & C_1 \times C_2 \times (\tau_1(S) \cap E) \\ & \cup [0, C_1] \times C_2 \times (\tau_1(S) \cap C) \cup [1, C_1] \times C_2 \times (\tau_4(S) \cap C) \\ & \cup C_1 \times [0, C_2] \times (\tau_1(S) \cap C_{012}) \cup C_1 \times [1, C_2] \times (\tau_2(S) \cap C_{012}) \\ & \cup C_1 \times [2, C_2] \times (\tau_3(S) \cap C_{012}) \end{aligned}$$

Hence,

$$\begin{aligned} & m_1 \otimes m_2 \otimes \mu \circ K_\beta^{-1}(C_1 \times C_2 \times S) \\ = & m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_1(S) \cap E) \\ & + p \cdot m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_1(S) \cap C) + (1-p) \cdot m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_4(S) \cap C) \\ & + s \cdot m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_1(S) \cap C_{012}) + t \cdot m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_2(S) \cap C_{012}) \\ & + (1-s-t) \cdot m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_3(S) \cap C_{012}) \\ = & ps \cdot m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_1(S)) + pt \cdot m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_2(S)) \\ & + p(1-s-t) \cdot m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_3(S)) + (1-p)s \cdot m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_4(S)) \\ & + (1-p)t \cdot m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_5(S)) + (1-p)(1-s-t) \cdot m_1(C_1) \cdot m_2(C_2) \cdot \mu(\tau_6(S)) \\ = & m_1 \otimes m_2 \otimes \nu(C_1 \times C_2 \times S). \end{aligned}$$

On the other hand,

$$\begin{aligned} & R_\beta^{-1}(C_1 \times C_2 \times S) \\ = & [0, C_1] \times [0, C_2] \times \tau_1(S) \cup [0, C_1] \times [1, C_2] \times \tau_2(S) \\ & \cup [0, C_1] \times [2, C_2] \times \tau_3(S) \cup [1, C_1] \times [0, C_2] \times \tau_4(S) \\ & \cup [1, C_1] \times [1, C_2] \times \tau_5(S) \cup [1, C_1] \times [2, C_2] \times \tau_6(S). \end{aligned}$$

Therefore, we complete the proof. \square

Now we give the main result in this section.

Theorem 5.7. *Let $\beta \in (1, 3/2)$. Then K_β has an invariant measure of the form $m_1 \otimes m_2 \otimes \mu_\beta$, where μ_β is absolutely continuous with respect to λ_2 .*

Proof. By Theorem 5.3, Lemma 5.4, Lemma 5.5, and Lemma 5.6, we complete the proof. \square

Remark 5.8. When $\beta = 3/2$, $C_{012} = \{(\frac{2}{3}, \frac{2}{3})\}$ is a point. We modify the definition of R, R', R_β , and give relevant conclusions.

(i) Let $R = \{\tau_1, \tau_2; p_1(\vec{z}), p_2(\vec{z})\}$ be a position dependent random transformation on S_β , where

$$\tau_1(\vec{z}) = \begin{cases} \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in E_i, i \in \{0, 1, 2\}, \\ \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\}, \\ \beta\vec{z}, & \text{if } \vec{z} = (\frac{2}{3}, \frac{2}{3}), \end{cases}$$

$$\tau_2(\vec{z}) = \begin{cases} \beta\vec{z} - \vec{q}_i, & \text{if } \vec{z} \in E_i, i \in \{0, 1, 2\}, \\ \beta\vec{z} - \vec{q}_j, & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\}, \\ \beta\vec{z}, & \text{if } \vec{z} = (\frac{2}{3}, \frac{2}{3}), \end{cases}$$

and $p_1(\vec{z}) = p_2(\vec{z}) = 1/2$ for $\vec{z} \in S_\beta$. Similar to Theorem 5.3, it is not difficult to prove that R has an acim μ_β .

(ii) By Lemma 3.2 in [1], $\mu_\beta \otimes \lambda_1$ is invariant for the skew product R' , where

$$R'(\vec{z}, w) = \begin{cases} (\tau_1(\vec{z}), \frac{w}{p}), & \text{if } w \in [0, p), \\ (\tau_2(\vec{z}), \frac{w-p}{1-p}), & \text{if } w \in [p, 1). \end{cases}$$

(iii) Define the *skew product transformation* R_β on $\Omega \times S_\beta$ as follows:

$$R_\beta(\omega, \vec{z}) = \begin{cases} (\sigma\omega, \beta\vec{z} - \vec{q}_i), & \text{if } \vec{z} \in E_i, i \in \{0, 1, 2\}, \\ (\sigma\omega, \beta\vec{z} - \vec{q}_i), & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\}, \omega_1 = 0, \\ (\sigma\omega, \beta\vec{z} - \vec{q}_j), & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\}, \omega_1 = 1, \\ (\sigma\omega, \beta\vec{z}), & \text{if } \vec{z} = (\frac{2}{3}, \frac{2}{3}). \end{cases}$$

Then we have that the dynamical systems $(S_\beta \times I, \mathcal{S} \times \mathcal{B}(I), \mu_\beta \otimes \lambda_1, R')$ and $(\Omega \times S_\beta, \mathcal{A} \times \mathcal{S}, m_1 \otimes \mu_\beta, R_\beta)$ are isomorphic. The proof is similar and easier than that of Lemma 5.5.

(iv) Let μ be an arbitrary probability measure on S_β and let $\tilde{K}_\beta : \Omega \times S_\beta \rightarrow \Omega \times S_\beta$ be given by

$$\tilde{K}_\beta(\omega, \vec{z}) = \begin{cases} (\omega, \beta\vec{z} - \vec{q}_i), & \text{if } \vec{z} \in E_i, i \in \{0, 1, 2\} \\ (\sigma\omega, \beta\vec{z} - \vec{q}_i), & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\}, \omega_1 = 0 \\ (\sigma\omega, \beta\vec{z} - \vec{q}_j), & \text{if } \vec{z} \in C_{ij}, ij \in \{01, 02, 12\}, \omega_1 = 1 \\ (\omega, \beta\vec{z}), & \text{if } \vec{z} = (\frac{2}{3}, \frac{2}{3}). \end{cases}$$

It is easy to check that

$$m_1 \otimes \mu \circ \tilde{K}_\beta^{-1} = m_1 \otimes \mu \circ R_\beta^{-1} = m_1 \otimes \nu,$$

where

$$\nu = p \cdot \mu \circ \tau_1^{-1} + (1-p) \cdot \mu \circ \tau_2^{-1}$$

by using the same method of calculation in Lemma 5.6. Therefore, it follows from (i-iv) that \tilde{K}_β has an invariant measure of the form $m_1 \otimes \mu_\beta$.

Next we give another method to prove that R_β has an acim by using Corollary 3.3. Denote $I_0 = [0, p)$, $I_1 = [p, 1]$, $J_0 = [0, s)$, $J_1 = [s, s+t)$, $J_2 = [s+t, 1]$. Let

$$\tilde{R}(\vec{z}, w, v) = \begin{cases} (\beta\vec{z} - \vec{q}_i, \frac{w}{p}, \frac{v}{s}) & \text{if } \vec{z} \in E_i \times [0, p) \times [0, s), i \in \{0, 1, 2\}, \\ (\beta\vec{z} - \vec{q}_i, \frac{w-p}{1-p}, \frac{v}{s}) & \text{if } \vec{z} \in E_i \times [p, 1] \times [0, s), i \in \{0, 1, 2\}, \\ (\beta\vec{z} - \vec{q}_i, \frac{w}{p}, \frac{v-s}{t}) & \text{if } \vec{z} \in E_i \times [0, p) \times [s, s+t), i \in \{0, 1, 2\}, \\ (\beta\vec{z} - \vec{q}_i, \frac{w-p}{1-p}, \frac{v-s}{t}) & \text{if } \vec{z} \in E_i \times [p, 1] \times [s, s+t), i \in \{0, 1, 2\}, \\ (\beta\vec{z} - \vec{q}_i, \frac{w}{p}, \frac{v-s-t}{1-s-t}) & \text{if } \vec{z} \in E_i \times [0, p) \times [s+t, 1], i \in \{0, 1, 2\}, \\ (\beta\vec{z} - \vec{q}_i, \frac{w-p}{1-p}, \frac{v-s-t}{1-s-t}) & \text{if } \vec{z} \in E_i \times [p, 1] \times [s+t, 1], i \in \{0, 1, 2\}, \\ (\beta\vec{z} - \vec{q}_i, \frac{w}{p}, \frac{v}{s}) & \text{if } \vec{z} \in C_{ij} \times [0, p) \times [0, s), ij \in \{01, 02, 12\}, \\ (\beta\vec{z} - \vec{q}_j, \frac{w-p}{1-p}, \frac{v}{s}) & \text{if } \vec{z} \in C_{ij} \times [p, 1] \times [0, s), ij \in \{01, 02, 12\}, \\ (\beta\vec{z} - \vec{q}_i, \frac{w}{p}, \frac{v-s}{t}) & \text{if } \vec{z} \in C_{ij} \times [0, p) \times [s, s+t), ij \in \{01, 02, 12\}, \\ (\beta\vec{z} - \vec{q}_j, \frac{w-p}{1-p}, \frac{v-s}{t}) & \text{if } \vec{z} \in C_{ij} \times [p, 1] \times [s, s+t), ij \in \{01, 02, 12\}, \\ (\beta\vec{z} - \vec{q}_i, \frac{w}{p}, \frac{v-s-t}{1-s-t}) & \text{if } \vec{z} \in C_{ij} \times [0, p) \times [s+t, 1], ij \in \{01, 02, 12\}, \\ (\beta\vec{z} - \vec{q}_j, \frac{w-p}{1-p}, \frac{v-s-t}{1-s-t}) & \text{if } \vec{z} \in C_{ij} \times [p, 1] \times [s+t, 1], ij \in \{01, 02, 12\}, \\ (\beta\vec{z}, \frac{w}{p}, \frac{v}{s}) & \text{if } \vec{z} \in C_{012} \times [0, p) \times [0, s), \\ (\beta\vec{z}, \frac{w-p}{1-p}, \frac{v}{s}) & \text{if } \vec{z} \in C_{012} \times [p, 1] \times [0, s), \\ (\beta\vec{z}, \frac{w}{p}, \frac{v-s}{t}) & \text{if } \vec{z} \in C_{012} \times [0, p) \times [s, s+t), \\ (\beta\vec{z}, \frac{w-p}{1-p}, \frac{v-s}{t}) & \text{if } \vec{z} \in C_{012} \times [p, 1] \times [s, s+t), \\ (\beta\vec{z}, \frac{w}{p}, \frac{v-s-t}{1-s-t}) & \text{if } \vec{z} \in C_{012} \times [0, p) \times [s+t, 1], \\ (\beta\vec{z}, \frac{w-p}{1-p}, \frac{v-s-t}{1-s-t}) & \text{if } \vec{z} \in C_{012} \times [p, 1] \times [s+t, 1]. \end{cases}$$

We can get the following theorem.

Lemma 5.9. *Let $\beta \in (1, 3/2)$, then \tilde{R} admits an acim.*

Proof. Use the partition $\mathcal{P} : \{E_i \times I_m \times J_n, C_{ij} \times I_m \times J_n, C_{012} \times I_m \times J_n\}, i = 0, 1, 2; m = 0, 1; n = 0, 1, 2$.

Consider the iterate \tilde{R}^k and the corresponding partition $\vee_{i=0}^{k-1} \tilde{R}^{-i}(\mathcal{P})$. For $P_i \in \vee_{i=0}^{k-1} \tilde{R}^{-i}(\mathcal{P})$, let $\tilde{R}_i^k = \tilde{R}^k|_{P_i}$. Since the derivative matrix of $(\tilde{R}_i^k)^{-1}$ is

$$\begin{bmatrix} \frac{1}{\beta^k} & 0 & 0 & 0 \\ 0 & \frac{1}{\beta^k} & 0 & 0 \\ 0 & 0 & p^l(1-p)^{k-l} & 0 \\ 0 & 0 & 0 & s^i t^j (1-s-t)^{k-i-j} \end{bmatrix},$$

where $l, i, j \in \{0, 1, \dots, k\}$ and $0 \leq i+j \leq k$, then the Euclidean matrix norm,

$$\|D(\tilde{R}_i^k)^{-1}\| = \sqrt{\frac{2}{\beta^{2k}} + p^{2l}(1-p)^{2(k-l)} + s^{2i}t^{2j}(1-s-t)^{2(k-i-j)}}.$$

Since $\beta > 1$ and $0 < p, s, t, s+t < 1$, then $\lim_{n \rightarrow \infty} \|D(\tilde{R}_i^k)^{-1}\| = 0$ for any $P_i \in \vee_{i=0}^{k-1} \tilde{R}^{-i}(\mathcal{P})$. It means that we can choose an integer k such that

$$\|D(\tilde{R}_i^k)^{-1}\| < c$$

for any $0 < c < 1$. For the partition $\vee_{i=0}^{k-1} \tilde{R}^{-i}(\mathcal{P})$, a is a positive constant which is not larger than 1 since any set in the partition is a polyhedra in \mathbb{R}^4 . Let $c < \frac{a}{a+1}$, then we have

$$c(1 + \frac{1}{a}) < 1.$$

By Corollary 3.3, \tilde{R} admits an acim. □

Theorem 5.10. R_β admits an acim.

Proof. Set

$$l'(x) = \begin{cases} \frac{1}{p}, & \text{if } x \in [0, p), \\ \frac{1}{1-p}, & \text{if } x \in [p, 1], \end{cases} \quad h'(x) = \begin{cases} 0, & \text{if } x \in [0, p), \\ \frac{p}{1-p}, & \text{if } x \in [p, 1]. \end{cases}$$

Let $\varphi'(x) = l'(x) \cdot x - h'(x)$ and

$$l'_n = l'_n(x) := l'(\varphi'^{n-1}(x)), \quad h'_n = h'_n(x) := h'(\varphi'^{n-1}(x)).$$

Define the map $\phi_1 : [0, 1] \rightarrow \Omega$ as follows:

$$\phi_1 : x = \sum_{n=1}^{\infty} \frac{h'_i}{l'_1 l'_2 \dots l'_i} \mapsto (\omega_1, \omega_2, \dots),$$

where $\omega_n = \omega_n(x) = k - 1 \iff \varphi'^{n-1}(x) \in I_k$ for $I_1 = [0, p)$ and $I_2 = [p, 1]$.

Similarly, set

$$l''(x) = \begin{cases} \frac{1}{s}, & \text{if } x \in [0, s), \\ \frac{1}{t}, & \text{if } x \in [s, s+t), \\ \frac{1}{1-s-t}, & \text{if } x \in [s+t, 1], \end{cases} \quad h''(x) = \begin{cases} 0, & \text{if } x \in [0, s), \\ \frac{s}{t}, & \text{if } x \in [s, s+t), \\ \frac{s+t}{1-s-t}, & \text{if } x \in [s+t, 1]. \end{cases}$$

and

$$l''_n = l''_n(x) := l''(\varphi''^{n-1}(x)), \quad h''_n = h''_n(x) := h''(\varphi''^{n-1}(x)),$$

where $\varphi''(x) = l''(x) \cdot x - h''(x)$. Let $\phi_2 : [0, 1] \rightarrow \Upsilon$ be given by

$$\phi_2 : x = \sum_{n=1}^{\infty} \frac{h''_i}{l''_1 l''_2 \dots l''_i} \mapsto (v_1, v_2, \dots),$$

where $v_n = v_n(x) = k - 1 \iff \varphi''^{n-1}(x) \in J_k$ for $J_1 = [0, s)$, $J_2 = [s, s+t)$ and $J_3 = [s+t, 1]$.

Consider the map $\phi = \iota \circ (I_{S_\beta} \times \phi_1 \times \phi_2)$ from $(S_\beta \times [0, 1] \times [0, 1], \mathcal{S} \times \mathcal{B}[0, 1] \times \mathcal{B}[0, 1], \nu, \tilde{R})$ to $(\Omega \times \Upsilon \times S_\beta, \mathcal{A} \times \mathcal{B} \times \mathcal{S}, \mu, R_\beta)$, where $\iota(\vec{z}, \omega, v) = (\omega, v, \vec{z})$ and I_{S_β} is the identity map on S_β . It is easy to check that ϕ is an isomorphism by the definition, and the result follows from Lemma 5.9. □

6. RANDOM TRANSFORMATION K_β FOR $\frac{3}{2} < \beta \leq \beta^*$

6.1. A special class of S_β . For $3/2 < \beta < 2$, we should notice that there are holes in the attractor. Broomhead et al.[6] discussed a special structure of those holes: they are all centered on three radial lines originating from the center of the attractor and extending to the three vertices. Since the Sierpinski triangle they discussed can be converted to S_β through affine transformation, we can obtain the corresponding range of β . Their set-up is as follows. Let S_λ be the attractor of an IFS

$$\{g_i(\vec{z}) = \lambda \vec{z} + (1 - \lambda) \vec{p}_i\}, \quad i = 0, 1, 2,$$

where $\vec{p}_i = \frac{2}{3}(\cos \frac{2\pi i}{3}, \sin \frac{2\pi i}{3})$, which are the vertices of an equilateral triangle Δ' . Denote the central hole by H' , i.e. $H' = \Delta' \setminus \cup_{i=0}^2 g_i(\Delta')$. Recall the following result in [6, Proposition 3.7].

Proposition 6.1. *Let $\lambda_* \approx 0.6478$ be the appropriate root of*

$$x^3 - x^2 + x = \frac{1}{2}.$$

Then S_λ has a non-empty interior if $\lambda \in [\lambda_, \frac{2}{3})$, and moreover, each hole has the form $g_i^n(H')$, see Figure 2.*




FIGURE 2. S_λ for $\lambda_* \leq \lambda < \frac{2}{3}$

Notice that the IFS given by (2.1) can also be written as

$$f_{\vec{q}_i}(\vec{z}) = \frac{1}{\beta}\vec{z} + (1 - \frac{1}{\beta})\frac{\vec{q}_i}{\beta - 1}, i = 0, 1, 2.$$

Since there is an invertible affine transformation l from S_λ to S_β , then the two attractors are affinely equivalent when $\lambda = 1/\beta$. Therefore we have the following proposition.


Proposition 6.2. *Let $\beta^* \approx 1.5437$ be the root of*

$$x^3 - 2x^2 + 2x = 2.$$

Then S_β has a non-empty interior if $\beta \in (\frac{3}{2}, \beta^]$, and each hole has the form $f_{\vec{q}_i}^n(H)$, see Figure 3.*

Proof. Let

$$l(\vec{z}) = \frac{1}{\beta - 1} \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ -1/2 & -\sqrt{3}/2 \end{bmatrix} \vec{z} + \frac{1}{\beta - 1} \begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}.$$

FIGURE 3. S_β for $\frac{3}{2} < \beta \leq \beta^*$

Then we have $l(\vec{p}_i) = \frac{1}{\beta-1}\vec{q}_i, i \in \{0, 1, 2\}$, which implies

$$(6.1) \quad l(\Delta') = \Delta.$$

Since

$$\begin{aligned} f_{\vec{q}_i} \circ l(\vec{z}) &= \frac{1}{\beta} \left(\frac{1}{\beta-1} \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ -1/2 & -\sqrt{3}/2 \end{bmatrix} \vec{z} + \frac{1}{\beta-1} \begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix} \right) + \left(1 - \frac{1}{\beta}\right) \frac{\vec{q}_i}{\beta-1} \\ &= \frac{1}{\beta(\beta-1)} \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ -1/2 & -\sqrt{3}/2 \end{bmatrix} \vec{z} + \frac{1}{\beta(\beta-1)} \begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix} + \left(1 - \frac{1}{\beta}\right) l(\vec{p}_i), \end{aligned}$$

and

$$\begin{aligned} l \circ g_i(\vec{z}) &= \frac{1}{\beta-1} \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ -1/2 & -\sqrt{3}/2 \end{bmatrix} (\lambda \vec{z} + (1-\lambda)\vec{p}_i) + \frac{1}{\beta-1} \begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix} \\ &= \frac{\lambda}{\beta-1} \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ -1/2 & -\sqrt{3}/2 \end{bmatrix} \vec{z} + \frac{1-\lambda}{\beta-1} \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ -1/2 & -\sqrt{3}/2 \end{bmatrix} \vec{p}_i + \frac{1}{\beta-1} \begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix} \\ &= \frac{\lambda}{\beta-1} \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ -1/2 & -\sqrt{3}/2 \end{bmatrix} \vec{z} + (1-\lambda) \left(l(\vec{p}_i) - \frac{1}{\beta-1} \begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix} \right) + \frac{1}{\beta-1} \begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix} \\ &= \frac{\lambda}{\beta-1} \begin{bmatrix} -1/2 & \sqrt{3}/2 \\ -1/2 & -\sqrt{3}/2 \end{bmatrix} \vec{z} + (1-\lambda) l(\vec{p}_i) + \frac{\lambda}{\beta-1} \begin{bmatrix} 1/3 \\ 1/3 \end{bmatrix}, \end{aligned}$$

then $f_{\vec{q}_i} \circ l = l \circ g_i$ by $\lambda = 1/\beta$. From this, we can get

$$(6.2) \quad l(H') = l(\Delta') \setminus \cup_{i=0}^2 l(g_i(\Delta')) = \Delta \setminus \cup_{i=0}^2 f_{\vec{q}_i}(\Delta) = H.$$

If $\beta \in (\frac{3}{2}, \beta^*]$, then $\lambda \in [\lambda_*, \frac{2}{3})$. By Proposition 6.1 (see [6, Proposition 3.7] for details), we have that for $n \geq 2$,

$$(1) \quad g_i g_j^{n-1}(H') \subset g_j(\Delta');$$

$$(2) \ g_i g_j^{n-1}(H') \cap g_j g_i^{n-1}(H') = \emptyset.$$

Then it follows from (6.1) and (6.2) that

$$\begin{aligned} (1) \ f_{\tilde{q}_i} f_{\tilde{q}_j}^{n-1}(H) &= f_{\tilde{q}_i} f_{\tilde{q}_j}^{n-1}(l(H')) = f_{\tilde{q}_i} f_{\tilde{q}_j}^{n-2}(l \circ g_i(H')) = l(g_i g_j^{n-1}(H')) \\ &\subset l \circ g_j(\Delta') = f_{\tilde{q}_j} \circ l(\Delta') = f_{\tilde{q}_j}(\Delta), \\ (2) \ f_{\tilde{q}_i} f_{\tilde{q}_j}^{n-1}(H) \cap f_{\tilde{q}_j} f_{\tilde{q}_i}^{n-1}(H) &= l(g_i g_j^{n-1}(H') \cap g_j g_i^{n-1}(H')) = \emptyset. \end{aligned}$$

□

6.2. Definition of K_β for $\beta \in (3/2, \beta^*]$. Because of the disappearance of the triple overlap, we only need one two-sided coin to be tossed to determine the transformation on the point. However, the partition of S_β is complicated by the presence of holes.

First we partition Δ_1 into \tilde{E}_i and \tilde{C}_{ij} (see Figure 4), where

$$\begin{aligned} \tilde{E}_0 &= ([0, \frac{1}{\beta}) \times [0, \frac{1}{\beta})) \setminus H, \\ \tilde{E}_1 &= \{(x, y) : 0 \leq y < \frac{1}{\beta}, \frac{1}{\beta(\beta-1)} < x+y \leq \frac{1}{\beta-1}\} \setminus H, \\ \tilde{E}_2 &= \{(x, y) : 0 \leq x < \frac{1}{\beta}, \frac{1}{\beta(\beta-1)} < x+y \leq \frac{1}{\beta-1}\} \setminus H, \\ \tilde{C}_{01} &= \{(x, y) : x \geq \frac{1}{\beta}, y \geq 0, x+y \leq \frac{1}{\beta(\beta-1)}\}, \\ \tilde{C}_{12} &= \{(x, y) : x \geq \frac{1}{\beta}, y \geq \frac{1}{\beta}, x+y \leq \frac{1}{\beta-1}\}, \\ \tilde{C}_{02} &= \{(x, y) : x \geq 0, y \geq \frac{1}{\beta}, x+y \leq \frac{1}{\beta(\beta-1)}\}. \end{aligned}$$

Here $H = \{(x, y) : x < \frac{1}{\beta}, y < \frac{1}{\beta}, x+y > \frac{1}{\beta(\beta-1)}\}$.


Then we have the following lemma.

Lemma 6.3. *For $\beta \in (3/2, \beta^*]$. Let $(x, y) \in S_\beta$ and $(x, y) = \sum_{i=1}^{\infty} \frac{a_i}{\beta^i}$ with $a_i \in \{\tilde{q}_0, \tilde{q}_1, \tilde{q}_2\}$ be a representation of (x, y) in base β . one has*

- (i) *If $(x, y) \in \tilde{E}_i$ for some $i \in \{0, 1, 2\}$, then $a_1 = \tilde{q}_i$,*
- (ii) *If $(x, y) \in \tilde{C}_{ij}$ for some $ij \in \{01, 12, 02\}$, then $a_1 \in \{\tilde{q}_i, \tilde{q}_j\}$.*

Proof. The proof is similar with Lemma 4.2. □

From the discussion in Proposition 6.2, we know that all holes $f_{\tilde{q}_i}^n(H)$ in the attractor S_β are in \tilde{E}_i (see Figure 3). And the ‘holes’ $f_{\tilde{q}_i} f_{\tilde{q}_j}^n(H) (i \neq j)$ that should exist in \tilde{C}_i are covered by $f_{\tilde{q}_j}(S_\beta)$, i.e. $f_{\tilde{q}_i} f_{\tilde{q}_j}^n(H) (i \neq j) \subset S_\beta$, which leaves only one choice \tilde{q}_j for the first digit of the expansions of the points in $f_{\tilde{q}_i} f_{\tilde{q}_j}^n(H)$. Now we can partition the Sierpinski carpet S_β into equality regions E_i and switch regions

FIGURE 4. Δ_1 for $\frac{3}{2} < \beta \leq \beta^*$

C_{ij} .

$$\begin{aligned}
E_0 &= \tilde{E}_0 \bigcup \bigcup_{n=1}^{\infty} f_{\vec{q}_1} f_{\vec{q}_0}^n(H) \bigcup \bigcup_{n=1}^{\infty} f_{\vec{q}_2} f_{\vec{q}_0}^n(H) \setminus \bigcup_{n=0}^{\infty} f_{\vec{q}_0}^n(H), \\
E_1 &= \tilde{E}_1 \bigcup \bigcup_{n=1}^{\infty} f_{\vec{q}_0} f_{\vec{q}_1}^n(H) \bigcup \bigcup_{n=1}^{\infty} f_{\vec{q}_2} f_{\vec{q}_1}^n(H) \setminus \bigcup_{n=0}^{\infty} f_{\vec{q}_1}^n(H), \\
E_2 &= \tilde{E}_2 \bigcup \bigcup_{n=1}^{\infty} f_{\vec{q}_1} f_{\vec{q}_2}^n(H) \bigcup \bigcup_{n=1}^{\infty} f_{\vec{q}_0} f_{\vec{q}_2}^n(H) \setminus \bigcup_{n=0}^{\infty} f_{\vec{q}_2}^n(H), \\
C_{01} &= \tilde{C}_{01} \setminus (\bigcup_{n=1}^{\infty} f_{\vec{q}_0} f_{\vec{q}_1}^n(H) \bigcup \bigcup_{n=1}^{\infty} f_{\vec{q}_1} f_{\vec{q}_0}^n(H)), \\
C_{12} &= \tilde{C}_{12} \setminus (\bigcup_{n=1}^{\infty} f_{\vec{q}_2} f_{\vec{q}_1}^n(H) \bigcup \bigcup_{n=1}^{\infty} f_{\vec{q}_1} f_{\vec{q}_2}^n(H)), \\
C_{02} &= \tilde{C}_{02} \setminus (\bigcup_{n=1}^{\infty} f_{\vec{q}_0} f_{\vec{q}_2}^n(H) \bigcup \bigcup_{n=1}^{\infty} f_{\vec{q}_2} f_{\vec{q}_0}^n(H)).
\end{aligned}$$

The selection of the digits in the expansion becomes more accurate.

Lemma 6.4. For $\beta \in (3/2, \beta^*]$. Let $\vec{z} \in S_\beta$ and $\vec{z} = \sum_{i=1}^{\infty} \frac{a_i}{\beta^i}$ with $a_i \in \{\vec{q}_0, \vec{q}_1, \vec{q}_2\}$ be a representation of \vec{z} in base β . one has

- (i) If $\vec{z} \in E_i$ for some $i \in \{0, 1, 2\}$, then $a_1 = \vec{q}_i$,
- (ii) If $\vec{z} \in C_{ij}$ for some $ij \in \{01, 12, 02\}$, then $a_1 \in \{\vec{q}_i, \vec{q}_j\}$.

Proof. It is enough to prove that $a_1 = \vec{q}_j$ for $\vec{z} \in f_{\vec{q}_i} f_{\vec{q}_j}^n(H)$.

Suppose $a_1 \neq \vec{q}_j$. Since $f_{\vec{q}_i} f_{\vec{q}_j}^n(H) \subset \tilde{C}_{ij}$, it follows from Lemma 6.3 that $a_1 = \vec{q}_i$. Then

$$f_{\vec{q}_i}^{-1}(\vec{z}) = \beta \vec{z} - \vec{q}_i = \sum_{i=1}^{\infty} \frac{a_{i+1}}{\beta^i} \in S_\beta,$$

which is a contradiction to the fact that $f_{\vec{q}_i}^{-1}(\vec{z}) \in f_{\vec{q}_j}^n(H)$ which is a hole. \square

Let

$$C = C_{01} \cup C_{12} \cup C_{02}, \quad E = E_0 \cup E_1 \cup E_2.$$

Recall $\Omega = \{0, 1\}^{\mathbb{N}}$ with the product σ -algebra \mathcal{A} and $\sigma : \Omega \rightarrow \Omega$ is the left shift, and define $K_\beta : \Omega \times S_\beta \rightarrow \Omega \times S_\beta$ by

$$K_\beta(\omega, \vec{z}) = \begin{cases} (\omega, \beta\vec{z} - \vec{q}_i), & \text{if } \vec{z} \in E_i, i = 0, 1, 2, \\ (\sigma\omega, \beta\vec{z} - \vec{q}_i), & \text{if } \omega_1 = 0 \text{ and } \vec{z} \in C_{ij}, ij \in \{01, 12, 02\}, \\ (\sigma\omega, \beta\vec{z} - \vec{q}_j), & \text{if } \omega_1 = 1 \text{ and } \vec{z} \in C_{ij}, ij \in \{01, 12, 02\}. \end{cases}$$

The digits are given by

$$d_1 = d_1(\omega, \vec{z}) = \begin{cases} \vec{q}_i, & \text{if } \vec{z} \in E_i, i = 0, 1, 2, \\ \text{or } (\omega, \vec{z}) \in \{\omega_1 = 0\} \times C_{ij}, ij \in \{01, 12, 02\}, \\ \vec{q}_j, & \text{if } (\omega, \vec{z}) \in \{\omega_1 = 1\} \times C_{ij}, ij \in \{01, 12, 02\}. \end{cases}$$

Then

$$K_\beta(\omega, \vec{z}) = \begin{cases} (\omega, \beta\vec{z} - d_1), & \text{if } \vec{z} \in E, \\ (\sigma\omega, \beta\vec{z} - d_1), & \text{if } \vec{z} \in C. \end{cases}$$

Set $d_n = d_n(\omega, \vec{z}) = d_1(K_\beta^{n-1}(\omega, \vec{z}))$, and $\pi_2 : \Omega \times S_\beta \rightarrow S_\beta$ be the canonical projection onto the second coordinate, then

$$\|\vec{z} - \sum_{i=1}^n \frac{d_i}{\beta^i}\|_1 = \frac{\|\pi_2(K_\beta^n(\omega, \vec{z}))\|_1}{\beta_n}.$$

Since $\pi_2(K_\beta^n(\omega, \vec{z})) \in \Delta$ and Δ is a bounded set in \mathbb{R}^2 , it follows that for all $\omega \in \Omega$ and for all $\vec{z} \in S_\beta$ one has

$$\vec{z} = \sum_{i=1}^{\infty} \frac{d_i}{\beta^i} = \sum_{i=1}^{\infty} \frac{d_i(\omega, \vec{z})}{\beta^i}.$$

One can see that the random map K_β for $\beta \in (3/2, \beta^*]$ satisfies similar properties as that for $\beta \in (1, 3/2]$. For each point $\vec{z} \in S_\beta$, consider the set

$$D_{\vec{z}} = \{(d_1(\omega, \vec{z}), d_2(\omega, \vec{z}), \dots) : \omega \in \Omega\}.$$

Theorem 6.5. *Suppose $\omega, \omega' \in \Omega$ are such that $\omega \prec \omega'$. Then*

$$(d_1(\omega, \vec{z}), d_2(\omega, \vec{z}), \dots) \preceq (d_1(\omega', \vec{z}), d_2(\omega', \vec{z}), \dots).$$

Proof. This result is easier to prove than Theorem 4.1 using the same method. \square

Theorem 6.6. *For $\beta \in (3/2, \beta^*]$. Let $\vec{z} \in S_\beta$ and $\vec{z} = \sum_{i=1}^{\infty} \frac{a_i}{\beta^i}$ with $a_i \in \{\vec{q}_0, \vec{q}_1, \vec{q}_2\}$ be a representation of \vec{z} in base β . Then there exists an $\omega \in \Omega$ such that $a_i = d_i(\omega, \vec{z})$.*

Proof. Using Lemma 6.4, the proof is similar with Theorem 4.3. \square

6.3. Unique K_β -invariant measure of maximal entropy. On the set $\Omega \times S_\beta$ we consider the product σ -algebra $\mathcal{A} \times \mathcal{S}$. Recall the function $\rho_2 : \{\vec{q}_0, \vec{q}_1, \vec{q}_2\}^{\mathbb{N}} \rightarrow \Upsilon$ which is given by

$$\rho_2(\vec{q}_{b_1}, \vec{q}_{b_2}, \vec{q}_{b_3}, \dots) = (b_1, b_2, b_3, \dots).$$

Define the function $\rho : \Omega \times S_\beta \rightarrow \{\vec{q}_0, \vec{q}_1, \vec{q}_2\}^{\mathbb{N}}$ by

$$\rho_1(\omega, \vec{z}) = (d_1(\omega, \vec{z}), d_2(\omega, \vec{z}), \dots).$$

Denote by $\varphi = \rho_2 \circ \rho_1$ the function from $\Omega \times S_\beta$ to Υ . It is easy to check that φ is measurable and $\varphi \circ K_\beta = \sigma' \circ \varphi$. It follows from Theorem 6.6 that φ is surjective.

Let

$$Z = \{(\omega, \vec{z}) \in \Omega \times S_\beta : K_\beta^n(\omega, \vec{z}) \in \Omega \times C \text{ infinitely often}\},$$

$$D = \{(b_1, b_2, \dots) \in \Upsilon : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \in C \text{ for infinitely many } j\text{'s}\}.$$

Then $\varphi(Z) = D$, $K_\beta^{-1}(Z) = Z$ and $(\sigma')^{-1}(\Upsilon) = \Upsilon$. Let φ' be the restriction of the map φ on Z .

Lemma 6.7. *The map $\varphi' : Z \rightarrow D$ is a bimeasurable bijection.*

Proof. For any sequence $(b_1, b_2, \dots) \in D$, we can get $\vec{z} = \sum_{i=1}^{\infty} \frac{\vec{q}_{b_i}}{\beta^i}$. To get ω , define

$$r_1 = \min\{j \geq 1 : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \in C\}, \quad r_k = \min\{j > r_{k-1} : \sum_{i=1}^{\infty} \frac{\vec{q}_{b_{j+i-1}}}{\beta^i} \in C\}.$$

If $b_{r_k} \in \{0, 2\}$, let $\omega_k = b_{r_k}/2$. If $b_{r_k} = 1$ and $\sum_{i=1}^{\infty} \frac{\vec{q}_{b_{r_k+i-1}}}{\beta^i} \in C_0$, let $\omega_k = 1$. If $b_{r_k} = 1$ and $\sum_{i=1}^{\infty} \frac{\vec{q}_{b_{r_k+i-1}}}{\beta^i} \in C_1$, let $\omega_k = 0$. Define $(\varphi')^{-1} : \Upsilon' \rightarrow Z$ by

$$(\varphi')^{-1}((b_1, b_2, \dots)) = (\omega, \vec{z}).$$

$(\varphi')^{-1}$ is measurable by Theorem 2.2, and is the inverse of φ' . \square

Lemma 6.8. $\mathbb{P}(D) = 1$, where \mathbb{P} is the uniform product measure.

Proof. The proof is similar with Theorem 4.5. For any sequence $(b_1, b_2, \dots) \in \Upsilon$ and $l \geq 3$, define

$$\vec{z}_l = (x_l, y_l) = \frac{\vec{q}_1}{\beta} + \frac{\vec{q}_2}{\beta^3} + \frac{\vec{q}_{b_1}}{\beta^{l+1}} + \frac{\vec{q}_{b_2}}{\beta^{l+2}} + \dots$$

Since $\beta \in [3/2, \beta^*)$ and $\lim_{n \rightarrow 0} \frac{1}{\beta^n(\beta-1)} = 0$, there exists $L > 0$ such that for any $l \geq L$,

$$x_l \geq \frac{1}{\beta}, \quad y_l \geq \frac{1}{\beta^3},$$

$$x_l + y_l \leq \frac{1}{\beta} + \frac{1}{\beta^3} + \frac{1}{\beta^n(\beta-1)} \leq \frac{1}{\beta(\beta-1)}.$$

It follows that $(x_l, y_l) \in \tilde{C}_{01}$. Notice that for any point $(x, y) \in \cup_{n=1}^{\infty} f_{\vec{q}_0}^n f_{\vec{q}_1}^n(H) \cup \cup_{n=1}^{\infty} f_{\vec{q}_1}^n f_{\vec{q}_0}^n(H)$, we have $y < 1/\beta^3$, which implies that $(x_l, y_l) \in C_{01} \subset C$ for any $l \geq L$. Let

$$D' = \{(b_1, b_2, \dots) \in \Upsilon : b_j b_{j+1} \dots b_{j+L-1} = 1 \underbrace{00 \dots 0}_{N-1 \text{ times}} \text{ for infinitely many } j\text{'s}\},$$

then $D' \subset D$. Since $\mathbb{P}(D') = 1$, we have $\mathbb{P}(D) = 1$. \square

Now, consider the K_β -invariant measure ν_β defined on $\mathcal{A} \times \mathcal{S}$ by $\nu_\beta(A) = \mathbb{P}(\varphi(Z \cap A))$. We have the following theorem.

Theorem 6.9. *Let $3/2 < \beta \leq \beta^*$. The dynamical systems $(\Omega \times S_\beta, \mathcal{A} \times \mathcal{S}, \nu_\beta, K_\beta)$ and $(\Upsilon, \mathcal{B}, \mathbb{P}, \sigma')$ are isomorphic. Moreover, the measure ν_β is the unique K_β -invariant measure of maximal entropy.*

Proof. It follows from Lemmas 6.7 and 6.8 that $\varphi : (\Omega \times S_\beta, \mathcal{A} \times \mathcal{S}, \nu_\beta, K_\beta) \rightarrow (\Upsilon, \mathcal{B}, \mathbb{P}, \sigma')$ is an isomorphism, which implies that $h_{\nu_\beta}(K_\beta) = h_{\mathbb{P}}(\sigma') = \log 3$ and any other K_β -invariant measure with support Z has entropy strictly less than $\log 3$. Let μ be a K_β -invariant measure for which $\mu(Z^c) > 0$. By a similar discussion with that of Lemma 4.9, we have $h_\mu(K_\beta) < \log 3$. \square

ACKNOWLEDGEMENTS

The first and third author were supported by National Natural Science Foundation of China (NSFC) No. 12071148 and Science and Technology Commission of Shanghai Municipality (STCSM) No. 18dz2271000. The first author was also supported by China Scholarship Council No.202006140156 during her visit to Utrecht University.

REFERENCES

- [1] Bahsoun W, Bose C and Quas A. *Deterministic representation for position dependent random maps*. Discrete Conti. Dyn. Syst., 22(3):529-540, 2012.
- [2] Barrionuevo J, Burton R, Dajani K and Kraaikamp C. *Ergodic properties of generalized Lüroth series*. Acta Arithmetica, 74: 311-327, 1996.
- [3] Bahsoun W and Góra P. *Position dependent random maps in one and higher dimensions*. Studia Mathematica, 166(3): 271-286, 2005.
- [4] Boyarsky A and Góra P. *Absolutely continuous invariant measures for piecewise expanding C^2 transformations in \mathbb{R}^N* . Israel Jour. Math., Vol. 67, No. 3, 1989, 272-286.
- [5] Boyarsky A and Góra P. *Laws of chaos*. Birkhauser, 1997.
- [6] Broomhead D, Montaldi J and Sidorov N. *Golden Gasket: Variations on the Sierpiński Sieve*. Nonlinearity, 17: 455-1480, 2004.
- [7] Dajani K, and Kraaikamp C. *Ergodic Theory of Numbers. 1st ed. Vol. 29*. Mathematical Association of America, 2002.
- [8] Dajani K and Kraaikamp C. *From greedy to lazy expansions and their driving dynamics*. Expo. Math., 20: 315-327, 2002.
- [9] Dajani K and Kraaikamp C. *Random β -expansions*. Ergod. Th. and Dynam. Sys., 23: 461-479, 2003.
- [10] Dajani K and de Vries M. *Measures of maximal entropy for random β -expansions*. J. Eur. Math. Soc., 7: 51-68, 2005.
- [11] Dajani K and de Vries M. *Invariant densities for random β -expansions*. J. Eur. Math. Soc., 9: 157-176, 2007.
- [12] Falconer K. *Fractal Geometry: Mathematical Foundations and Applications, 3rd Edition*. John Wiley and Sons, Ltd., 2014.
- [13] Giusti E. *Minimal Surfaces and Functions of Bounded Variation*. Birkhauser, 1984.
- [14] Hochman M. *On self-similar sets with overlaps and inverse theorems for entropy in \mathbb{R}^d* . Mem. Am. Math. Soc., to appear, (arXiv:1503.09043).
- [15] Hasselblatt B and Plante D Jr. *On the interior of 'fat' Sierpiński triangles*. Exp. Math., 23: 285-309, 2014.
- [16] Kong D and Li W. *Critical base for the unique codings of fat Sierpiński gasket*. Nonlinearity, 33(9): 4484-4511, 2020.
- [17] Jordan T and Pollicott M. *Properties of measures supported on fat Sierpinski carpets*. Ergod. Theor. Dynam. Syst., 26: 739-54, 2006.
- [18] Simon K and Solomyak B. *On the dimension of self-similar sets*. Fractals, 10: 59-65, 2002.
- [19] Sidorov N. *Combinatorics of linear iterated function systems with overlaps*. Nonlinearity, 20: 1299-312, 2007.
- [20] Srivastava S M. *A course on Borel sets*. Graduate Texts in Mathematics, 180, Springer, New York, 1998.