

p -ADIC SIMPSON CORRESPONDENCE VIA PRISMATIC CRYSTALS

YU MIN AND YUPENG WANG

ABSTRACT. Let \mathfrak{X} be a smooth p -adic formal scheme over \mathcal{O}_K with adic generic fiber X . We obtain a global equivalence between the category $\mathrm{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$ of rational Hodge–Tate crystals on the absolute prismatic site $(\mathfrak{X})_{\Delta}$ and the category $\mathrm{HIG}_*^{\mathrm{nil}}(X)$ of enhanced Higgs bundles on X . Along the way, we construct an inverse Simpson functor from $\mathrm{HIG}_*^{\mathrm{nil}}(X)$ to the category $\mathrm{Vect}(X_{\mathrm{pro\acute{e}t}}, \widehat{\mathcal{O}}_X)$ of generalised representations on X , which turns out to be fully faithful.

CONTENTS

1. Introduction	1
2. Recollection of the decompletion theory of Diao–Lan–Liu–Zhu	10
3. A p -adic Simpson correspondence for rigid analytic varieties over K	15
4. Local description of Hodge–Tate crystals	23
5. Hodge–Tate crystals as generalised representations	40
6. Inverse Simpson functor for enhanced Higgs bundles	42
7. The Sen operator	50
References	58

1. INTRODUCTION

1.1. Overview. In their groundbreaking work [BS22], Bhatt and Scholze introduced the prismatic cohomology, which is “motivic” in the sense that it can recover most existing p -adic cohomology theories (e.g. crystalline cohomology, de Rham cohomology and étale cohomology). Recently, the coefficient theory of the prismatic cohomology has caught a lot of attention. For example, one can recover étale \mathbb{Z}_p -local systems on the generic fibers of bounded p -adic formal schemes \mathfrak{X} by studying the *Laurent F -crystals* on the absolute prismatic site $(\mathfrak{X})_{\Delta}$ of \mathfrak{X} (cf. [Wu21], [BS21], [MW21a]). When \mathfrak{X} is smooth over \mathcal{O}_K , the ring of integers of a p -adic field K , there exists a nice equivalence between the category of crystalline \mathbb{Z}_p -local systems on its generic fiber and that of (*analytic*) *prismatic F -crystals* on $(\mathfrak{X})_{\Delta}$ (cf. [ALB19], [BS21], [DL21], [DLMS22], [GR22]). If \mathfrak{X} is smooth over $\widehat{\mathcal{O}}_{\widehat{K}}$, then the theory of prismatic crystals provides a q -deformation of local p -adic Simpson correspondence (cf. [GLSQ20], [MT20], [GLSQ22], etc.).

In this paper, we will focus on the theory of *Hodge–Tate prismatic crystal*. Note that in the geometric setting, Hodge–Tate crystals on the relative prismatic site can be **locally** understood as certain Higgs bundles (cf. [Tian21]). This phenomenon also appears in characteristic p and one can understand Hodge–Tate crystals as nilpotent Higgs bundles for “good” smooth scheme over \mathbb{F}_p (cf. [Og22]). In the arithmetic setting, Hodge–Tate crystals on the absolute prismatic site of \mathcal{O}_K can be understood as certain semi-linear \widehat{K} -representations of $\mathrm{Gal}(\overline{K}/K)$ and closely related with classical Sen theory (cf. [MW21b], [Gao22], [BL22a], [GMW22], [AHLB22], etc.).

Continuing our work [MW21b], we are going to combine the two aspects, i.e. the geometric side and the arithmetic side, of Hodge–Tate crystals mentioned above. More precisely, for a smooth p -adic formal scheme \mathfrak{X} over \mathcal{O}_K with generic fiber X , we will study the category $\mathrm{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$ of rational Hodge–Tate crystal on $(\mathfrak{X})_{\Delta}$ and investigate how the arithmetic part and the geometric part interact with each other. In fact, we will establish a *global* equivalence between the category of rational Hodge–Tate crystals and the category of enhanced Higgs bundles, which can be understood

2020 *Mathematics Subject Classification.* Primary 14G22, 14F30, 14G45.

Key words and phrases. rational Hodge–Tate crystal, enhanced Higgs bundle, p -adic Simpson correspondence, Sen theory.

as a hybrid of Higgs bundles and (arithmetic) Sen operators. Along the way, we will obtain a fully faithful functor from the category of enhanced Higgs bundles on X to the category of generalised representations on $X_{\text{proét}}$, which we call the *inverse Simpson functor*.

1.2. Main result. We freely use notations introduced in §1.4. In particular, K is a complete discretely valued field of mixed characteristic $(0, p)$ with ring of integers \mathcal{O}_K and perfect residue field κ and $C = \widehat{K}$ is a fixed p -complete algebraic closure of K .

1.2.1. Statement of main result. We first introduce some notions that will be considered in this paper. Let $\text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$ (resp. $\text{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$) be the category of rational Hodge–Tate crystals on the **absolute** prismatic site $(\mathfrak{X})_{\Delta}$ (resp, the absolute prismatic site of perfect prisms) of \mathfrak{X} (cf. Definition 1.24). Let $\text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ denote the category of generalised representations (i.e. locally finite free $\widehat{\mathcal{O}}_X$ -modules) on $X_{\text{proét}}$ (cf. Example 1.23) and $\text{HIG}_*^{\text{nil}}(X)$ be the category of enhanced Higgs bundles on X ; that is, Higgs bundles defined on X with a Sen operator satisfying certain conditions (cf. Definition 6.2). Let $\text{HIG}_{G_K}(X_C)$ denote the category of G_K -Higgs bundles on X_C ; that is, Higgs bundles on X_C together with compatible G_K -actions (cf. Definition 3.1).

Then our main theorem is as follows.

Theorem 1.1 (Theorem 6.4). *Assume \mathfrak{X} is a smooth p -adic formal scheme over \mathcal{O}_K with rigid analytic generic fiber X . Then there exists an equivalence of categories*

$$\rho : \text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \simeq \text{HIG}_*^{\text{nil}}(X),$$

which preserves ranks, tensor products and dualities. Moreover, this equivalence fits into the following commutative diagram of fully faithful functors:

$$(1.1) \quad \begin{array}{ccccc} \text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) & \xrightarrow[\simeq]{\rho} & \text{HIG}_*^{\text{nil}}(X) & & \\ \downarrow \text{Res} & & \swarrow F_S & \searrow F & \\ \text{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) & \xrightarrow{\simeq} & \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X) & \xrightarrow{\simeq} & \text{HIG}_{G_K}(X_C), \end{array}$$

where Res is induced by inclusion of sites $(\mathfrak{X})_{\Delta}^{\text{perf}} \subset (\mathfrak{X})_{\Delta}$, F will be defined in Construction 6.13 and all arrows with “ \simeq ” are equivalences of categories. In particular, we obtain a fully faithful inverse Simpson functor

$$F_S : \text{HIG}_*^{\text{nil}}(X) \rightarrow \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X).$$

Remark 1.2. The relation between Hodge–Tate crystals and Higgs bundles is not new: indeed, if one fix a transversal prism (A, I) and consider small affine formal scheme $\mathfrak{X} = \text{Spf}(R)$ over A/I , then there exists an equivalence between the category of topologically nilpotent Higgs bundles over \mathfrak{X} and the category of Hodge–Tate crystals on the **relative** prismatic site $(\mathfrak{X}/(A, I))_{\Delta}$ of \mathfrak{X} (cf. [Tian21, Thm. 4.10]). Our improvement is that if we consider the **absolute** prismatic site of \mathfrak{X} and rational Hodge–Tate crystals, then the Higgs bundles coming from rational Hodge–Tate crystals can be equipped with an additional operator so that we can establish a global theory. We emphasize that both inverting p and considering the absolute prismatic site are essential in our construction.

Remark 1.3. In the classical theory of Simpson correspondence over the field of complex numbers \mathbb{C} , a nilpotent Higgs bundle $(\mathcal{H}, \theta_{\mathcal{H}})$ over a smooth projective variety X may not induce a representation of the fundamental group $\pi_1(X(\mathbb{C}))$ or a \mathbb{G}_m -action on $(\mathcal{H}, \theta_{\mathcal{H}})$. However, in the setting of Theorem 1.1, if a nilpotent Higgs bundle $(\mathcal{H}, \theta_{\mathcal{H}})$ is enhanced, then it will induce a generalised representation and a G_K -action on its base-change to C , the field of p -adic complex numbers. We point out that the naive base-change “ $(\mathcal{H}, \theta_{\mathcal{H}}) \otimes_K C$ ” never carries the “induced G_K -action” because Tate twists appear in the definition of Higgs bundles and C does not contain p -adic analogue of $2\pi i$.

Remark 1.4. Theorem 1.1 sheds light on studying p -adic Simpson correspondence via prismatic crystals. In p -adic Simpson correspondence, constructing representations from certain Higgs bundles seems much harder than the other direction (for example, see [Xu22] for the curve case). Our inverse Simpson functor F_S in Theorem 1.1 provides an example in this direction by working with enhanced

Higgs bundles. Moreover, our method is also compatible with the works [Fal05], [AGT16], [Wang21] when \mathfrak{X} is smooth over \mathcal{O}_C . One can check in this case, locally the diagram in Theorem 1.1 can recover Faltings' local Simpson functor in [Fal05] and by using the period sheaf \mathcal{OC}^\dagger in [Wang21] instead of \mathcal{OC} in Theorem 1.5, our approach can show that the local construction in [Tian21] glues if one considers the categories of small Higgs bundles and small Hodge–Tate crystals and both categories are equivalent to the category of small generalised representations under the liftable assumption of \mathfrak{X} to A_{inf}/ξ^2 .

Next, we are going to explain our strategy of proving Theorem 1.1. We will first introduce the key tools used in the globalisation process and then talk about the local constructions and complete the strategy of the proof.

1.2.2. p -adic Simpson correspondence and Hodge–Tate crystals on perfect site. Similar to [Tian21], we can get a local correspondence between Hodge–Tate crystals and Higgs bundles in the absolute case. The main difficulty of getting a global one then lies in comparing local constructions. Our strategy is to compare local data in a bigger category through a fully faithful functor. The key ingredient is the p -adic Simpson correspondence.

Based on the previous work of Scholze [Sch12, Sch13], Liu–Zhu [LZ17] assigned to each étale \mathbb{Q}_p -local system on a smooth rigid analytic space over K a nilpotent Higgs field by using decompletion theory in [KL16] and got the rigidity of de Rham local systems. Their method also works in the logarithmic case [DLLZ18]. We will work on their arenas and prove the following result, in which X is not required to be a rigid generic fiber of a smooth p -adic formal scheme.

Theorem 1.5 (Theorem 3.3). *Let X be a smooth rigid space over K and $\nu : X_{\text{proét}} \rightarrow X_{C,\text{ét}}$ be the natural projection of sites. For any generalised representation $\mathcal{L} \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$, if we put $\Theta_{\mathcal{L}} := \text{id}_{\mathcal{L}} \otimes \Theta$, then the rule*

$$\mathcal{L} \mapsto (\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})}) := (\nu_*(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{OC}), \nu_*(\Theta_{\mathcal{L}}))$$

induces a rank-preserving equivalence from the category $\text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ of generalised representations on $X_{\text{proét}}$ to the category $\text{HIG}_{G_K}(X_C)$ of G_K -Higgs bundles on $X_{C,\text{ét}}$, which preserves tensor products and dualities. Moreover, the following assertions are true:

- (1) *For any $i \geq 1$, the higher direct image $R^i \nu_*(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{OC}) = 0$.*
- (2) *Let $\Theta_{\mathcal{H}(\mathcal{L})} := \theta_{\mathcal{H}(\mathcal{L})} \otimes \text{id}_{\mathcal{OC}} + \text{id}_{\mathcal{H}(\mathcal{L})} \otimes \Theta$. Then there exists a natural isomorphism*

$$(\mathcal{H} \otimes_{\mathcal{O}_{X_C}} \mathcal{OC}|_{X_C}, \Theta_{\mathcal{H}(\mathcal{L})}) \xrightarrow{\cong} (\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{OC}, \Theta_{\mathcal{L}})|_{X_C}$$

of Higgs fields.

- (3) *Let $(\text{HIG}(\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})}))$ denote the Higgs complex induced by $(\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})})$. Then there exists a natural quasi-isomorphism*

$$\text{R}\Gamma(X_{C,\text{proét}}, \mathcal{L}) \cong \text{R}\Gamma(X_{C,\text{ét}}, (\text{HIG}(\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})}))$$

which is compatible with G_K -actions. As a consequence, we get a quasi-isomorphism

$$\text{R}\Gamma(X_{\text{proét}}, \mathcal{L}) \cong \text{R}\Gamma(G_K, \text{R}\Gamma(X_{C,\text{ét}}, (\text{HIG}(\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})}))).$$

- (4) *Let $X' \rightarrow X$ be a smooth morphism of rigid space over K . Then the equivalence in (3) is compatible with pull-back along f . In other words, for any $\mathcal{L} \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ with corresponding $(\mathcal{H}, \theta_{\mathcal{H}}) \in \text{HIG}_{G_K}(X_C)$, we have*

$$(\mathcal{H}(f^*\mathcal{L}), \theta_{\mathcal{H}(f^*\mathcal{L})}) \cong (f^*\mathcal{H}, f^*\theta_{\mathcal{H}}).$$

An immediate corollary is the following.

Corollary 1.6 (Corollary 3.7). *Let d be the dimension of X over K .*

- (1) *Assume X is quasi-compact. Then $\text{R}\Gamma(X_{C,\text{proét}}, \mathcal{L})$ is concentrated in degree $[0, 2d]$ and $\text{R}\Gamma(X_{\text{proét}}, \mathcal{L})$ is concentrated in degree $[0, 2d + 1]$.*
- (2) *If moreover X is proper, then $\text{R}\Gamma(X_{C,\text{proét}}, \mathcal{L})$ is a perfect complex of C -representations of G_K and $\text{R}\Gamma(X_{\text{proét}}, \mathcal{L})$ is a perfect complex of K -vector spaces.*

Remark 1.7. By almost étale descent, one can replace C in Theorem 1.5 and Corollary 1.6 by its any perfectoid sub-field containing ζ_p^∞ . By analytic-étale comparison, one can use the Zariski site instead of étale site above.

Remark 1.8. (1) When X is affine, Theorem 1.5 was achieved by Tsuji [Tsu18, Thm. 15.2] by choosing a certain integral model (with log structure) of X , which is not necessary in our approach.

(2) For generalised representations \mathcal{L} coming from \mathbb{Q}_p -local systems \mathbb{L} (i.e. $\mathcal{L} = \widehat{\mathcal{O}}_X \otimes_{\mathbb{Q}_p} \mathbb{L}$), Theorem 1.5 was proved by Liu–Zhu as [LZ17, Thm. 2.1] by using decompletion theory for certain overconvergent period rings “ $\widetilde{\mathbf{B}}^\dagger$ ”. Our proof was inspired by theirs and works for any generalised representations by using decompletion theory developed in [DLLZ18] (cf. §2).

(3) Note that when X admits a good reduction \mathfrak{X} over $W(\kappa)$, the base-change along $W(\kappa) \rightarrow \mathbf{A}_{\text{inf}}$ induces a lifting $\widetilde{\mathfrak{X}}$ of $\mathfrak{X}_C := \mathfrak{X} \times_{W(\kappa)} \mathcal{O}_C$. So for a small generalised representations \mathcal{L} , the work of Abbes–Gros–Tsuji [AGT16] produces also a Higgs bundle over X_C with G_K -action. Theorem 1.5 should be compatible with theirs. Note that when X has semi-stable reduction \mathfrak{X} over $W(\kappa)$ merely, then the base-change of \mathfrak{X} to \mathbf{A}_{inf} is not a lifting of \mathfrak{X}_C as log schemes. Indeed, we do not know whether a lifting of \mathfrak{X}_C to \mathbf{A}_{inf} (as log scheme) exists except the curve case and the affine small case. So it seems not easy to apply results in [AGT16] directly in this case.

(4) When X is either abeloid or X is curve and \mathcal{L} is a line bundle, Theorem 1.5 can be also deduced from [Heu22a] and [HMW22] by noting that HTlog in loc.cit. is G_K -equivariant. We thanks Ben Heuer for pointing out this to us.

Before explaining how to apply Theorem 1.5, let us exhibit the relation between rational Hodge–Tate crystals and generalised representations. We again assume X is a rigid generic fiber of a smooth p -adic formal scheme over \mathcal{O}_K . The key observation is that both $X_{\text{proét}}$ (or X_v) and $(\mathfrak{X})_{\Delta}^{\text{perf}}$ have basis consisting of perfectoid algebras. Using this, we can prove the following étale comparison theorem:

Theorem 1.9 (Theorem 5.1). *There exists a natural equivalence of categories*

$$L : \text{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \rightarrow \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X),$$

which preserves ranks, tensor products and dualities, such that for any rational Hodge–Tate crystal \mathbb{M} on $(\mathfrak{X})_{\Delta}^{\text{perf}}$, we have a quasi-isomorphism

$$\text{R}\Gamma((\mathfrak{X})_{\Delta}^{\text{perf}}, \mathbb{M}) \cong \text{R}\Gamma(X_{\text{proét}}, L(\mathbb{M})),$$

which is functorial in \mathbb{M} .

Remark 1.10. When $\mathfrak{X} = \text{Spf}(\mathcal{O}_K)$, Theorem 1.9 reduces to [MW21b, Thm. 1.6]; that is, there exists an equivalence

$$\text{Rep}_{G_K}(C) \simeq \text{Vect}((\mathcal{O}_K)_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$$

between the category of semi-linear C -representations of G_K and the category of rational Hodge–Tate crystals on $(\mathcal{O}_K)_{\Delta}^{\text{perf}}$. Similar result also holds true for de Rham crystals (cf. [GMW22]). We will discuss this topic in higher dimensional case in a forthcoming paper with Hui Gao.

Due to Theorem 1.9, the inclusion of sites $(\mathfrak{X})_{\Delta}^{\text{perf}} \subset (\mathfrak{X})_{\Delta}$ induces a natural functor

$$\text{Res} : \text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \rightarrow \text{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \simeq \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X).$$

We will see that this functor is fully faithful (cf. Proposition 6.11) and then use it to compare local constructions that we are going to explain right now.

1.2.3. Local constructions. Assume $\mathfrak{X} = \text{Spf}(R^+)$ is small affine. By (p -completely) faithfully flat descent and [BS21, Prop. 2.7], one can study (rational) Hodge–Tate crystals in terms of stratifications by choosing a certain cover $(\mathfrak{S}(R^+), (E))$ (depending on the choice of charts on R^+) of final objects of $\text{Shv}((R^+)_{\Delta})$. In this case, parts of Theorem 1.5 upgrade to the integral level and one can give an explicit description of the functor Res . In this case, $\text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ is equivalent to the category $\text{Rep}_{\Gamma(\overline{K}/K)}(\widehat{R}_{C,\infty})$ of semi-linear $\widehat{R}_{C,\infty}$ -representations of $\Gamma(\overline{K}/K) \cong (\oplus_{i=1}^d \mathbb{Z}_p \gamma_i) \rtimes G_K$ (cf. Notation 2.19).

Theorem 1.11. *Keep notations as above and fix a framing of \mathfrak{X} .*

- (1) [Theorem 4.3] *The evaluation at $(\mathfrak{S}(R^+), (E))$ induces an equivalence of categories*

$$\rho : \text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}) \rightarrow \text{HIG}_*^{\text{nil}}(R^+) \text{ (resp. } \rho : \text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \rightarrow \text{HIG}_*^{\text{nil}}(R)),$$

which preserve ranks, tensor products and dualities, such that for any Hodge–Tate crystal (resp. rational Hodge–Tate crystal) \mathbb{M} with associated enhanced Higgs module over R^+ (resp. R), there exists a quasi-isomorphism

$$\text{R}\Gamma((R^+)_{\Delta}, \mathbb{M}) \cong \text{HIG}(H, \theta_H, \phi_H),$$

which is functorial in \mathbb{M} .

- (2) [Theorem 6.5] *For any $(H, \theta_H, \phi_H) \in \text{HIG}_*^{\text{nil}}(R)$ corresponding the rational Hodge–Tate crystal \mathbb{M} with induced $\widehat{R}_{C,\infty}$ -representation $\text{Res}(\mathbb{M})$ of $\Gamma(\overline{K}/K)$, we have $\text{Res}(\mathbb{M}) = H \otimes_R \widehat{R}_{C,\infty}$ such that for any $1 \leq i \leq d$, any $g \in G_K$ and any $x \in H$,*

$$\gamma_i(x) = \exp(-(\zeta_p - 1)\lambda\theta_i)(x) \text{ and } g(x) = (1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_H}{E'(\pi)}}(x).$$

Remark 1.12. Note that any Hodge–Tate crystal $\mathbb{M} \in \text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta})$ can be naturally viewed as a Hodge–Tate crystal on the relative prismatic site $(R^+ / (\mathfrak{S}, (E)))_{\Delta}$. One can apply [Tian21, Thm. 4.10] to get a topologically nilpotent Higgs field over R^+ , which turns out to be the underlying Higgs field (H, θ_H) of $\rho(\mathbb{M})$.

Remark 1.13. When $\mathcal{O}_K = W(\kappa)$, the first part of Theorem 1.11 was also obtained by Bhatt–Lurie [BL22b] by using the local splitting of the Hodge–Tate structure map $\text{WCart}_{\mathfrak{X}}^{\text{HT}} \rightarrow \mathfrak{X}$. Up to now, it is still a problem to achieve a global theory as their method needs a global “Frobenius endomorphism” (cf. [BL22b, §9]), which is a very restrictive condition. However, our Theorem 1.1 suggests it is reasonable to ask whether the “generic fiber” of the Hodge–Tate structure map $\text{WCart}_{\mathfrak{X}}^{\text{HT}}[\frac{1}{p}] \rightarrow \mathfrak{X}[\frac{1}{p}]$ admits a global splitting.

Inspired by Theorem 1.11 (2), one can assign to each enhanced Higgs bundle $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$ over X (which is necessary to be affine) a G_K -Higgs bundle over X_C by letting $g \in G_K$ act via the cocycle $(1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_H}{E'(\pi)}}$ and hence get the global functor F in Theorem 1.1. To prove F is fully faithful (cf. Proposition 6.11), we need to relate prismatic theory to classical Sen theory, which will be discussed in §1.2.4¹.

Now since everything can be described explicitly, one can check directly that Theorem 1.1 holds true in the affine small case. In particular, Diagram (1.1) is commutative in this case. To complete the proof of Theorem 1.1, we need to check that for rational Hodge–Tate crystals, the local equivalence ρ in Theorem 1.11 (1) is independent of the choice of charts on R^+ . Since F is fully faithful, it is enough to show $F \circ \rho$ is independent of the choice of charts. But this is obvious now as all functors except ρ in Diagram (1.1) are globally defined and Diagram (1.1) is commutative in affine case.

1.2.4. Sen operators. Finally, we explain how our work is related with Sen theory. Recall that for any semi-linear C -representation V of G_K , by considering the cyclotomic extension, Sen constructed an endomorphism Θ_V , which is called *Sen operator*, defined over K satisfying certain properties. His result was generalised by Berger–Colmez [BC16] to any extension of K whose Galois group is a p -adic Lie group of arbitrary dimension and recently by Pan [Pan20] to the geometric setting. On the other hand, still considering the cyclotomic extension, the method of Sen was also generalised by Shimizu [Shi18] and Petrov [Pet20] to higher dimensional rigid spaces: for any quasi-compact smooth rigid space X over K and any G_K -equivariant vector bundle \mathcal{E} over $X_{C,\text{ét}}$, they constructed an $\mathcal{O}_{\mathcal{X}}$ -vector bundle over the ringed space $\mathcal{X} = (X, \mathcal{O}_{\mathcal{X}} = \mathcal{O}_X \otimes_K K_{\text{cyc}})$ together with a Sen operator satisfying certain properties. This construction is compatible with that of Sen and is functorial in \mathcal{E} (see Proposition 7.2 for the explicit statement).

Now by Theorem 1.5 together with the above construction, we can assign to each generalised representation \mathcal{L} over $X_{\text{proét}}$ a Higgs bundle with a Sen operator $(\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})}, \phi_{\mathcal{H}(\mathcal{L})})$ over \mathcal{X} . We

¹There is another way to show the full faithfulness here; See Proposition 2.27 in <https://arxiv.org/pdf/2201.08030v1.pdf> for details.

also call $\phi_{\mathcal{H}(\mathcal{L})}$ the *Sen operator of \mathcal{L}* and denote it by $\phi_{\mathcal{L}}$. After base change to X_C , this will give rise to an *arithmetic Higgs bundle* over X_C (cf. Definition 7.14, 7.15). In other words, there is a functor

$$D : \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X) \rightarrow \text{HIG}^{\text{arith}}(X_C)$$

which is closely related with the cyclotomic extension. Before moving on, we raise the following question.

Question 1.14. Which arithmetic Higgs bundles come from generalised representations under the functor D ?

In fact, it is not clear to us how to describe the essential image of the functor D in general. When $\mathfrak{X} = \text{Spf}(\mathcal{O}_K)$, Fontaine gave a complete answer to Question 1.14 after classifying all semi-linear C -representations of G_K (cf. [Fon04, Thm. 2.14]) by working with a certain algebraic group. But his method looks very difficult to be generalised to the higher dimensional case. We will give a partial answer to this question provided by the theory of Hodge–Tate crystals at the end of this subsection.

Now we move back to investigate the relation between prismatic theory and Sen theory. Note that we have a local equivalence between rational Hodge–Tate crystals and enhanced Higgs bundles at the moment. The latter can give rise to arithmetic Higgs bundles as follows.

Example 1.15 (Construction 7.22 (1)). Assume $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$ is an enhanced Higgs bundle over X . Then $(\mathcal{H} \otimes_{\mathcal{O}_X} \mathcal{O}_{X_C}, -(\zeta_p - 1)\lambda\theta_{\mathcal{H}}, -\frac{\phi_{\mathcal{H}}}{E'(\pi)})$ is an arithmetic Higgs bundle over X_C , where λ is a certain unit in \mathcal{O}_C . We denote the induced functor by $F_{\infty} : \text{HIG}_{*}^{\text{nil}}(X) \rightarrow \text{HIG}^{\text{arith}}(X_C)$.

Given a rational Hodge–Tate crystal, we now have two ways to construct arithmetic Higgs bundles (locally): one is the cyclotomic way through the functor D , the other is the Kummer way via Example 1.15. It turns out that the two constructions coincide with each other.

Theorem 1.16 (Theorem 7.12, Hui Gao). *Let \mathbb{M} be a rational Hodge–Tate crystal on $(\mathfrak{X})_{\Delta}$ where $\mathfrak{X} = \text{Spf}(R^+)$ is small, with induced enhanced Higgs bundle $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$ and generalised representation \mathcal{L} . Then the Sen operator $\phi_{\mathcal{L}}$ of \mathcal{L} is exactly $-\frac{\phi_{\mathcal{H}}}{E'(\pi)}$.*

Remark 1.17. Theorem 1.16 was conjectured by the authors [MW21b, Conjecture 1.8] and was proved by Hui Gao [Gao22] by using ideas from [BC16] and the theory of (ϕ, τ) -modules. Gao informed us that the method in [Gao22] also works in the higher dimensional case and explained the proof to us. We thank Hui Gao for allowing us to include Theorem 1.16 here. When $R^+ = W(\kappa)$ is absolutely unramified extension of \mathbb{Z}_p , the result was also obtained in [BL22a, §3.8].

We will prove Theorem 1.16 by following the strategy of [Gao22]; that is, we will apply the theory of locally analytic vectors of Berger–Colmez [BC16] to compare Sen operators obtained from cyclotomic extension and Kummer extension. As a consequence, we also conclude the full faithfulness of $F : \text{HIG}_{*}^{\text{nil}}(X) \rightarrow \text{HIG}_{G_K}(X_C)$ which will complete the proof of Theorem 1.1.

Now given Theorem 1.1, Theorem 1.16 can be globalised. Let F_{cyc} denote the composition of functors $\text{HIG}_{*}^{\text{nil}}(X) \rightarrow \text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \rightarrow \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X) \xrightarrow{D} \text{HIG}^{\text{arith}}(X_C)$. This means there is an equivalence of functors $F_{\text{cyc}} \simeq F_{\infty} : \text{HIG}_{*}^{\text{nil}}(X) \rightarrow \text{HIG}^{\text{arith}}(X_C)$. As an application, we get a partial answer to Question 1.14.

Theorem 1.18 (Corollary 7.24). *If an arithmetic Higgs bundle is induced by an enhanced Higgs bundle under the functor F_{∞} , then it comes from a generalised representation over $X_{\text{proét}}$.*

1.3. Organization. We review the decompletion theory of [DLLZ18] in §2 and use it to establish the p -adic Simpson correspondence (i.e. Theorem 1.5) in §3. In §4, we construct the equivalence ρ in Theorem 1.1 in the small affine case. In §5, we prove Theorem 1.9. The §6 is devoted to proving Theorem 1.1 by leaving the full faithfulness of F to §7. Finally, we show that F is fully faithful by using Sen theory in §7 and prove Theorem 1.18 as an application.

1.4. Conventions and Notations. Let K be a complete discretely valued field of mixed characteristic $(0, p)$ with ring of integers \mathcal{O}_K and perfect residue field κ . We fix an algebraic closure \overline{K} of K and let C denote the p -adic completion of \overline{K} . For any Galois extension L/K in \overline{K} , we denote by $\text{Gal}(L/K)$ the corresponding Galois group. We fix a uniformizer $\pi \in \mathcal{O}_K$ and denote its minimal polynomial over $W(\kappa)$ by $E(u)$. Then $E(u) \in \mathfrak{S} := W(\kappa)[[u]]$.

Let $\{\zeta_{p^n}\}_{n \geq 0}$ (resp. $\{\pi^{\frac{1}{p^n}}\}_{n \geq 0}$) be a compatible sequence $\{\zeta_{p^n}\}_{n \geq 0}$ of primitive p^n -th units (resp. of p^n -th roots of π) in \overline{K} and let $\epsilon = (1, \zeta_p, \dots)$ (resp. $\pi^b = (\pi, \pi^{\frac{1}{p}}, \dots)$) be the induced element in C^b , the tilting of C . Let $K_{\text{cyc}} := \bigcup_{n \geq 0} K(\zeta_{p^n})$ (resp. $K_\infty = \bigcup_{n \geq 0} K(\pi^{\frac{1}{p^n}})$, resp. $K_{\text{cyc}, \infty} = K_{\text{cyc}, \infty}$) with the p -adic completion \widehat{K}_{cyc} (resp. \widehat{K}_∞ , resp. $\widehat{K}_{\text{cyc}, \infty}$) in C . Then \widehat{K}_{cyc} , \widehat{K}_∞ and $\widehat{K}_{\text{cyc}, \infty}$ are all perfectoid fields. Let $\chi : G_K \rightarrow \text{Gal}(K_{\text{cyc}}/K) \rightarrow \mathbb{Z}_p^\times$ be the p -adic cyclotomic character and let $c : G_K \rightarrow \text{Gal}(K_{\text{cyc}, \infty}/K) \rightarrow \mathbb{Z}_p$ be the function such that for any $g \in G_K$, $g(\pi^b) = \epsilon^{c(g)} \pi^b$. Let $t = \log[\epsilon]$ be the p -adic analogue of “ $2\pi i$ ”.

Convention 1.19. In this paper, we always assume $K_{\text{cyc}} \cap K_\infty = K$. This is always the case for $p \geq 3$ and for $p = 2$ when $K = K(\zeta_4)$ (cf. [Liu08, Lem. 5.1.2]). In general, for $p = 2$, the assumption is satisfied for a suitable choice of π (cf. [Wang17, Lem. 2.1]). Under this assumption, we have

$$\text{Gal}(K_{\text{cyc}, \infty}/K) \cong \mathbb{Z}_p \tau \rtimes \text{Gal}(K_{\text{cyc}}/K),$$

where $\tau \in \text{Gal}(K_{\text{cyc}, \infty}/K_{\text{cyc}})$ such that $\tau(\pi^b) = \epsilon \pi^b$ (i.e. $c(\tau) = 1$). Then for any $\gamma \in \text{Gal}(K_{\text{cyc}, \infty}/K_\infty) \cong \text{Gal}(K_{\text{cyc}}/K)$, we have $\gamma \tau \gamma^{-1} = \tau^{\chi(\gamma)}$. Moreover, one can identify $\text{Gal}(K_{\text{cyc}, \infty}/K)$ with an open subgroup of $\{(\begin{smallmatrix} a & b \\ c & d \end{smallmatrix}) \in \text{GL}_2(\mathbb{Z}_p) \mid c = 0, d = 1\}$ via the homomorphism $X : \text{Gal}(K_{\text{cyc}, \infty}/K) \rightarrow \text{GL}_2(\mathbb{Z}_p)$ sending each $g \in \text{Gal}(K_{\text{cyc}, \infty}/K)$ to $X(g) = \begin{pmatrix} \chi(g) & c(g) \\ 0 & 1 \end{pmatrix}$.

Put $\mathbf{A}_{\text{inf}} = W(\mathcal{O}_C^b)$ and $\xi = \frac{[\epsilon]-1}{[\epsilon]^{\frac{1}{p}}-1}$. Then ξ is a generator of Fontaine’s map $\theta : \mathbf{A}_{\text{inf}} \rightarrow \mathcal{O}_C$ and $(\mathbf{A}_{\text{inf}}, (\xi))$ is a prism. We equip \mathfrak{S} with a δ -structure such that the induced Frobenius on \mathfrak{S} carries u to u^p . Then $(\mathfrak{S}, (E))$ is also a prism and the morphism $\mathfrak{S} \rightarrow \mathbf{A}_{\text{inf}}$ sending u to $[\pi^b]$ defines a morphism $(\mathfrak{S}, (E)) \rightarrow (\mathbf{A}_{\text{inf}}, (\xi))$ of prisms in $(\mathcal{O}_K)_\Delta$.

Let Y, Y_1, \dots, Y_d be a free variable. For any $n \geq 0$, we define

$$\binom{Y}{n} := \frac{Y(Y-1) \cdots (Y-n+1)}{n!}$$

and

$$Y^{[n]} = \frac{Y^{[n]}}{n!}.$$

For any $\underline{n} = (n_1, \dots, n_d) \in \mathbb{N}^d$, we then define

$$\binom{\underline{Y}}{\underline{n}} = \prod_{i=1}^d \binom{Y_i}{n_i}$$

and

$$\underline{Y}^{\underline{n}} = \prod_{i=1}^d Y_i^{[n_i]}.$$

Put $\underline{0} = (0, 0, \dots, 0) \in \mathbb{Z}^d$ and for any $1 \leq i \leq d$, put $\underline{1}_i = (0, \dots, 1, \dots, 0)$ with 1 appearing exactly in the i -th component. For any $\underline{m} = (m_1, \dots, m_d), \underline{n} = (n_1, \dots, n_d) \in \mathbb{N}^d$, define

$$\binom{\underline{m} + \underline{n}}{\underline{n}} = \prod_{i=1}^d \binom{m_i + n_i}{n_i}$$

and

$$|\underline{n}| = n_1 + n_2 + \cdots + n_d.$$

We will also use the language of stratifications to study certain vector bundles on sites.

Convention 1.20. Let S^\bullet be a cosimplicial ring with face maps p_i ’s and degeneracy maps σ_i ’s; that is, p_i is induced by the injection $[n] \rightarrow [n+1] \setminus \{i\}$ and that σ_i is induced by the surjection

$[n+1] \rightarrow [n]$ satisfying $\sigma_i^{-1}(i) = \{i, i+1\}$. By a *stratification* with respect to S^\bullet , we mean a finite projective S^0 -module together with an S^1 -linear isomorphism

$$\varepsilon : M \otimes_{S^0, p_0} S^1 \xrightarrow{\cong} M \otimes_{S^0, p_1} S^1.$$

Sometimes, we also say ε is a stratification on M .

Let $q_i : S^0 \rightarrow S^n$ denote the morphism induced by the inclusion $[0] \xrightarrow{0 \rightarrow i} [n]$ of simplices. We say a stratification (M, ε) satisfies the *cocycle condition*, if the following assertions are true:

- (1) $p_2^*(\varepsilon) \circ p_0^*(\varepsilon) = p_1^*(\varepsilon) : M \otimes_{S^0, q_2} S^2 \rightarrow M \otimes_{S^0, q_0} S^2$.
- (2) $\sigma_0^*(\varepsilon) = \text{id}_M : M \rightarrow M$.

We denote by $\text{Strat}(S^\bullet)$ the category of stratifications with respect to S^\bullet satisfying the cocycle condition.

We also consider representations of several topological groups over certain topological rings.

Convention 1.21. Let R be a topological ring with a continuous action of a topological group G . By a *representation of G over R of rank l* , we mean a finite projective R -module M of rank l together with a semi-linear continuous action of G ; that is, for any $r \in R$, $m \in M$ and $g \in G$,

$$g(rm) = g(r)g(m).$$

Sometimes, we also call M an *R -representation of G of rank l* . We say an R -representation M of G is *free*, if M is finite free over R . We denote by $\text{Rep}_G(R)$ the category of representations of G over R and denote by $\text{Rep}_G^{\text{free}}(R)$ the full subcategory of free representations.

Assume G acts on R trivially and that M is an topological R module equipped with a continuous action of G . We denote by $\text{R}\Gamma(G, M)$ the continuous group cohomology of M , which is computed by the complex $C(G^\bullet, M)$. Here and from now on, we always denote by $C(G, M)$ the group of continuous functions from G to M . For any closed normal subgroup $H < G$ with quotient group G/H , in general, there is no Hochschild spectral sequence for $\text{R}\Gamma(G, M)$. However, if $G \rightarrow G/H$ admits a continuous cross-section (which is not necessary an homomorphism), then Hochschild–Serre spectral sequence holds true (See the proof of [Ked16, Lem 3.3], [Laz65, Chap. V (3.2), pp. 193] for more details). In this paper, we always deal with the case for H open or $G \cong H \rtimes G/H$ and hence can use Hochschild–Serre spectral sequence freely.

Let $X = \text{Spa}(R, R^+)$ be a smooth affinoid space of dimension d over K . By a *toric chart* on X , we mean an étale morphism

$$\square : X \rightarrow \mathbb{G}_m^d := \text{Spa}(K\langle T_1^{\pm 1}, \dots, T_d^{\pm 1} \rangle, \mathcal{O}_K\langle T_1^{\pm 1}, \dots, T_d^{\pm 1} \rangle).$$

Note that toric charts always exists étale locally on X . Let $\mathfrak{X} = \text{Spf}(R)$ be a smooth² p -adic formal scheme of dimension d over \mathcal{O}_K . We say it is *small*, if there exists a *framing* on \mathfrak{X} ; that is, an étale morphism

$$\square : \mathfrak{X} \rightarrow \text{Spf}(\mathcal{O}_K\langle T_1^{\pm 1}, \dots, T_d^{\pm 1} \rangle).$$

Note that in this case, \square induces a toric chart on the rigid generic fibre of \mathfrak{X} . Clearly, framings always exist étale locally on \mathfrak{X} . However, it is worth pointing out that framings even exist Zariski locally on \mathfrak{X} by [Bha16, Lem. 4.9]. By abuse of notations, we still denote by \square the induced morphism on the rings of coordinates.

We call an étale morphism of rigid spaces *standard étale*, if it is a composition of rational localisations and finite étale morphisms.

Convention 1.22. Let \mathcal{C} be a site and \mathbb{A} be a sheaf of rings on \mathcal{C} . By an \mathbb{A} -*crystal* on \mathcal{C} , we mean a sheaf \mathbb{M} of \mathbb{A} -modules such that

- (1) For any object $C \in \mathcal{C}$, $\mathbb{M}(C)$ is a finite projective $\mathbb{A}(C)$ -module of rank r .
- (2) For any arrow $C_1 \rightarrow C_2$ in \mathcal{C} , it induces a canonical isomorphism

$$\mathbb{M}(C_1) \otimes_{\mathbb{A}(C_1)} \mathbb{A}(C_2) \xrightarrow{\cong} \mathbb{M}(C_2)$$

of $\mathbb{A}(C_2)$ -modules.

We denote by $\text{Vect}(\mathcal{C}, \mathbb{A})$ the category of \mathbb{A} -crystals on \mathcal{C} .

²In this paper, smooth p -adic formal schemes are always assumed to be separated and of topologically finite type.

Example 1.23. By [KL16, Thm. 3.5.8], a $\widehat{\mathcal{O}}_X$ -crystal on $X_{\text{proét}}$ is exactly a *generalised representation* (i.e. a locally finite free $\widehat{\mathcal{O}}_X$ -module) on $X_{\text{proét}}$. For this reason, by abuse of notations, we denote by $\text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ the category of generalised representation on $X_{\text{proét}}$. When $X = \text{Spf}(\mathcal{O}_K)$, this is exactly the category of continuous C -representations of G_K .

The main object we will study in this paper is (rational) Hodge–Tate crystals.

Definition 1.24. Let \mathfrak{X} be a smooth p -adic formal scheme over \mathcal{O}_K . A *Hodge–Tate crystal* (resp. a rational Hodge–Tate crystal) is an $\overline{\mathcal{O}}_{\Delta}$ -crystal (resp. $\overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]$) on $(\mathfrak{X})_{\Delta}$. The category of Hodge–Tate crystals (resp. rational Hodge–Tate crystals) on $(\mathfrak{X})_{\Delta}$ is denoted by $\text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta})$ (resp. $\text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$). We will also consider the rational Hodge–Tate crystals (i.e. $\overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]$ -crystals) on the sub-site of perfect prism $(\mathfrak{X})_{\Delta}^{\text{perf}}$ and denote the corresponding category by $\text{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$.

When $\mathfrak{X} = \text{Spf}(R^+)$ is small affine, we also denote $\text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta})$ by $\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta})$, and similarly define $\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$ and $\text{Vect}((R^+)_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$.

1.5. Acknowledgement. The authors would like to thank Bhargav Bhatt, Ben Heuer, Ruochuan Liu and Takeshi Tsuji for their interests and valuable comments on the early draft of this paper. The authors also thank Hui Gao for allowing them to include his result (i.e. Theorem 1.16) in this paper. The work was finished when the first author stayed in Morningside Center of Mathematics as a postdoc and both authors thank the institute for providing the opportunity to cooperation. The first author was supported by China Postdoctoral Science Foundation E1900503.

2. RECOLLECTION OF THE DECOMPLETION THEORY OF DIAO–LAN–LIU–ZHU

In this section, we recall the decompletion theory formulated in [DLLZ18] and provide some examples, which will be used in the next sections.

2.1. Decompletion systems in [DLLZ18, Appendix A]. We give a quick review of decompletion theory developed in [DLLZ18, Appendix A] in this subsection.

Let $\{A_i\}_{i \in I}$ be a direct system of topological rings with a small filtered index category I , \widehat{A}_∞ be a complete topological ring with compatible homomorphisms $A_i \rightarrow \widehat{A}_\infty$ such that the induced map $\varinjlim_i A_i \rightarrow \widehat{A}_\infty$ has dense image, and Γ be a topological group acting continuously and compatibly on A_i 's and \widehat{A}_∞ .

Definition 2.1 ([DLLZ18, Def. A.1.2]). We call the triple $(\{A_i\}_{i \in I}, \widehat{A}_\infty, \Gamma)$ a *decompletion system* (resp. *weak decompletion system*) if the following two conditions hold:

- (1) For any (resp. free) \widehat{A}_∞ -representation L_∞ of Γ , there exists some $i \in I$, some (resp. free) A_i -representation of Γ , and some Γ -equivariant continuous A_i -linear morphism $\iota_i : L_i \rightarrow L_\infty$ which induces an isomorphism $L_i \otimes_{A_i} \widehat{A}_\infty \xrightarrow{\iota_i \otimes \text{id}} L_\infty$ of representations of Γ over \widehat{A}_∞ . We call such a pair (L_i, ι_i) a *model* of L_∞ over A_i .
- (2) For each model L_i over A_i , there exists some $i_0 \geq i$ such that for any $i' \geq i_0$, the model $(L_{i'}, \iota_{i'}) := (L_i \otimes_{A_i} A_{i'}, \iota_i \otimes \text{id})$ is *good* in the sense that $\iota_{i'}$ induces a quasi-isomorphism $\text{R}\Gamma(\Gamma, L_{i'}) \rightarrow \text{R}\Gamma(\Gamma, L_\infty)$.

Remark 2.2 ([DLLZ18, Rem. A.1.3]). If $(\{A_i\}_{i \in I}, \widehat{A}_\infty, \Gamma)$ is a (weak) decompletion system, then the natural functor $\varinjlim_i \text{Rep}_\Gamma(A_i) \rightarrow \text{Rep}_\Gamma(\widehat{A}_\infty)$ (resp. $\varinjlim_i \text{Rep}_\Gamma^{\text{free}}(A_i) \rightarrow \text{Rep}_\Gamma^{\text{free}}(\widehat{A}_\infty)$) is an equivalence and for any two models $(L_{i,1}, \iota_{i,1})$ and $(L_{i,2}, \iota_{i,2})$ of L_∞ over A_i , there exists some $i' \geq i$ such that $(L_{i,1} \otimes_{A_i} A_{i'}, \iota_{i,1} \otimes \text{id}) \cong (L_{i,2} \otimes_{A_i} A_{i'}, \iota_{i,2} \otimes \text{id})$.

The main results in [DLLZ18, Appendix A] give some sufficient conditions to clarify whether a triple $(\{A_i\}_{i \in I}, \widehat{A}_\infty, \Gamma)$ is a (weak) decompletion system. Recall a complex (C^\bullet, d) of Banach modules over a Banach ring A is called *uniformly strict exact* with respect to some $c \geq 0$, if for any cocycle $f \in C^s$, there exists a cochain $g \in C^{s-1}$ satisfying $|g| \leq c|f|$ such that $f = d(g)$.

Definition 2.3 ([DLLZ18, Def. A.1.6]). Let $(\{A_i\}_{i \in I}, \widehat{A}_\infty, \Gamma)$ be a triple as above. Assume that $A_i \rightarrow \widehat{A}_\infty$ are closed embeddings and that Γ is profinite. We say $(\{A_i\}_{i \in I}, \widehat{A}_\infty, \Gamma)$ is *weakly decompleting* if there exists a norm $|\cdot|$ on \widehat{A}_∞ making it a Banach ring (and hence making A_i 's Banach subrings) and an inverse system $\{\Gamma_i\}_{i \in I}$ of closed normal subgroup converging to 1 such that the projection $\Gamma \rightarrow \Gamma/\Gamma_i$ admits a continuous cross-section (which is not necessary a homomorphism) for each $i \in I$, satisfying the following conditions:

- (1) The action of Γ on \widehat{A}_∞ is isometric.
- (2) For each i , the projection $\widehat{A}_\infty \rightarrow \widehat{A}_\infty/A_i$ admits an isometric section of Banach A_i -modules.
- (3) There exists some $c > 0$ such that the complex $C(\Gamma_i^\bullet, \widehat{A}_\infty/A_i)$ is uniformly strict exact with respect to c , where we equip $C(\Gamma_i^s, \widehat{A}_\infty/A_i)$ with the supreme norm $|f| := \sup_{\gamma \in \Gamma_i^s} |f(\gamma)|$ for each s .

Then the first main result in [DLLZ18, Appendix A] is

Theorem 2.4 ([DLLZ18, Thm. A.1.8]). *A weakly decompleting triple is a weak decompletion system.*

Recall a Banach ring A is *stably uniform* if it is either the underlying ring of a stably uniform Huber pair in the sense of [SW20, Def. 5.2.4] or a stably uniform adic Banach ring in the sense of [KL15, Rem. 2.8.5] over a nonarchimedean field.

Definition 2.5 ([DLLZ18, Def. A.1.9]). Let $(\{A_i\}_{i \in I}, \widehat{A}_\infty, \Gamma)$ be a triple as above. Assume that $A_i \rightarrow \widehat{A}_\infty$ are closed embeddings and that Γ is profinite. We say $(\{A_i\}_{i \in I}, \widehat{A}_\infty, \Gamma)$ is *stably decompleting* if the following conditions are true:

- (1) A_i 's and \widehat{A}_∞ are stably uniform over a nonarchimedean field.

- (2) Each rational subset U of $\mathrm{Spa}(A_i, A_i^\circ)$ is stablized by some open normal subgroup Γ_U of Γ ; and the pull-back of $(\{A_j\}_{j \geq i}, \widehat{A}_\infty, \Gamma_U)$ to each such U is weakly decompleting.

Then the second main result in [DLLZ18, Appendix] is

Theorem 2.6 ([DLLZ18, Thm. A.1.10]). *A stably decompleting triple is a decompletion system.*

We give some examples in the next subsection.

2.2. Examples of decompletion system.

2.2.1. Generalised arithmetic tower. Assume $d \geq 0$ and define $\Gamma_{\mathrm{geo}} = \bigoplus_{i=1}^d \mathbb{Z}_p \gamma_i$, which is a finite free \mathbb{Z}_p -module with basis $\gamma_1, \dots, \gamma_d$. Let $\Gamma := \Gamma_{\mathrm{geo}} \rtimes \mathrm{Gal}(K_{\mathrm{cyc}}/K)$ such that for any $g \in \mathrm{Gal}(K_{\mathrm{cyc}}/K)$ and any $1 \leq i \leq d$, $g\gamma_i g^{-1} = \gamma_i^{\chi(g)}$.

Let (R, R^+) be an affinoid Tate algebra over K . For any $n \geq 1$, let $R_n^+ := R \otimes_{\mathcal{O}_K} \mathcal{O}_{K(\zeta_{p^n})}$, $R_n := R \otimes_K K(\zeta_{p^n})$, and $\Gamma_n := \Gamma_{\mathrm{geo}}^{p^n} \rtimes \mathrm{Gal}(K_{\mathrm{cyc}}/K(\zeta_{p^n}))$. Put $\widehat{R}_\infty^+ := R^+ \widehat{\otimes}_{\mathcal{O}_K} \mathcal{O}_{K_{\mathrm{cyc}}} \widehat{R}_\infty := R \widehat{\otimes}_K K_{\mathrm{cyc}}$. Then Γ acts on R_n 's and \widehat{R}_∞ compatibly and continuously via the quotient map $\Gamma \rightarrow \mathrm{Gal}(K_{\mathrm{cyc}}/K)$.

Lemma 2.7. *The triple $(\{R_n\}_{n \geq 0}, \widehat{R}_\infty, \Gamma)$ is stably decompleting.*

Proof. By proceeding as in the proof of [DLLZ18, Prop. 2.1.1], we are reduced to showing that for sufficiently large l and for any $n \geq 0$, $H^n(\Gamma_l, \widehat{R}_\infty^+/R_l^+)$ is killed by p^2 . Note that by [BMS18, Lem. 7.3], one may compute $\mathrm{R}\Gamma(\Gamma_{\mathrm{geo}}, \widehat{R}_\infty^+/R_l^+)$ by using Koszul complex $K(\gamma_1 - 1, \dots, \gamma_d - 1; \widehat{R}_\infty^+/R_l^+)$:

$$\widehat{R}_\infty^+/R_l^+ \xrightarrow{(\gamma_1 - 1, \dots, \gamma_d - 1)} (\widehat{R}_\infty^+/R_l^+)^d \rightarrow \dots \rightarrow \widehat{R}_\infty^+/R_l^+.$$

Since Γ_{geo} acts on $\widehat{R}_\infty^+/R_l^+$ trivially, $\mathrm{R}\Gamma(\Gamma_{\mathrm{geo}}, \widehat{R}_\infty^+/R_l^+)$ is computed by $K(0, \dots, 0; \widehat{R}_\infty^+/R_l^+)$. By using Hochschild–Serre spectral sequence, in order to conclude our result, we have to show that $\mathrm{R}\Gamma(\mathrm{Gal}(K_{\mathrm{cyc}}/K(\zeta_{p^l})), \widehat{R}_\infty^+/R_l^+)$ is killed by p^2 , which was confirmed in the proof of [DLLZ18, Prop. 2.1.1]. \square

Theorem 2.8. *The triple $(\{R_n\}_{n \geq 0}, \widehat{R}_\infty, \Gamma)$ is a decompletion system.*

Proof. This follows from Theorem 2.6 together with Lemma 2.7. \square

Remark 2.9. In [Shi18], Shimizu essentially showed that the triple $(\{R_n\}_{n \geq 0}, \widehat{R}_\infty, \Gamma)$ is a weak decompletion system by using the formalism of Tate–Sen theory developed in [BC08]. When $d = 0$, the result was first proved by Sen in [Sen80].

Proposition 2.10. *For any \widehat{R}_∞ -representation M of Γ , the subgroup Γ_{geo} acts on M quasi-unipotently.*

Proof. By Theorem 2.8, there exists an $n \geq 1$ and a representation N of Γ over R_n such that $M \cong N \otimes_{R_n} \widehat{R}_\infty$. Since $\Gamma' := \Gamma_{\mathrm{geo}} \rtimes \mathrm{Gal}(K_{\mathrm{cyc}}/K(\zeta_{p^n}))$ acts on R_n trivially. We see that N is a linear representation of Γ' . Then the result follows from the same proof of [LZ17, Lem. 2.15]. \square

Remark 2.11. Proposition 2.10 is optimal in the sense that one can not expect that Γ_{geo} acts on any \widehat{R}_∞ -representation M of Γ unipotently in general. For example, we may assume $M = R_{\widehat{K}_{\mathrm{cyc}}} e$ and $\Gamma = \mathbb{Z}_p \gamma \rtimes \mathrm{Gal}(K_{\mathrm{cyc}}/K)$ such that γ acts e via the scalar ζ_{p^n} for some fixed $n \geq 1$ and $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ acts on e trivially. Then $M \in \mathrm{Rep}_\Gamma(R_{\widehat{K}_{\mathrm{cyc}}})$ on which the Γ_{geo} -action is not unipotent.

2.2.2. Toric tower. Let $X = \mathrm{Spa}(R, R^+)$ be a smooth affinoid space of dimension d over K which admits a toric chart \square . Let $X_{\widehat{K}_{\mathrm{cyc}}} := \mathrm{Spa}(R_{\widehat{K}_{\mathrm{cyc}}}, R_{\widehat{K}_{\mathrm{cyc}}}^+)$ denote the base-change of X along $K \hookrightarrow \widehat{K}_{\mathrm{cyc}}$.

Notation 2.12. For any $n \geq 0$, let $R_n^+ := R^+ \widehat{\otimes}_{\mathcal{O}_K \langle T_1^{\pm 1}, \dots, T_d^{\pm 1} \rangle, \square} \mathcal{O}_{K(\zeta_{p^n})} \langle T_1^{\pm \frac{1}{p^n}}, \dots, T_d^{\pm \frac{1}{p^n}} \rangle$, $R_\infty^+ = \varprojlim_n R_n^+$, and $\widehat{R}_\infty^+ = (\varprojlim_n R_n^+)_p^\wedge$ be the p -adic completion of R_∞^+ . Put $R_n = R_n^+[\frac{1}{p}]$ and $\widehat{R}_\infty := \widehat{R}_\infty^+[\frac{1}{p}]$. Then both of them are stably uniform adic Banach rings. Let X_n (resp. X_∞) be the base-change (resp. the perfectoid space corresponding to the base-change) of X along

$$\mathrm{Spa}(K(\zeta_{p^n}) \langle T_1^{\pm \frac{1}{p^n}}, \dots, T_d^{\pm \frac{1}{p^n}} \rangle, \mathcal{O}_{K(\zeta_{p^n})} \langle T_1^{\pm \frac{1}{p^n}}, \dots, T_d^{\pm \frac{1}{p^n}} \rangle) \rightarrow \mathbb{G}_m^d$$

$$(\text{resp. } \text{Spa}(\widehat{K}_{\text{cyc}}\langle T_1^{\pm\frac{1}{p^\infty}}, \dots, T_d^{\pm\frac{1}{p^\infty}} \rangle, \mathcal{O}_{\widehat{K}_{\text{cyc}}} \langle T_1^{\pm\frac{1}{p^\infty}}, \dots, T_d^{\pm\frac{1}{p^\infty}} \rangle \rightarrow \mathbb{G}_m^d).$$

Then R_n (resp. \widehat{R}_∞) is the ring of regular functions on X_n (resp. X_∞). For any $n \geq 0$, the natural map $X_\infty \rightarrow X_n$ is a Galois cover with Galois group Γ_n . In particular, we denote $\Gamma := \Gamma_0$. The map $X_\infty \rightarrow X_{\widehat{K}_{\text{cyc}}}$ is also a Galois cover with Galois group

$$(2.1) \quad \Gamma_{\text{geo}} \cong \mathbb{Z}_p \gamma_1 \oplus \dots \oplus \mathbb{Z}_p \gamma_d$$

where for any $1 \leq i, j \leq d$ and any $n \geq 0$, $\gamma_i(T_j^{\frac{1}{p^n}}) = \zeta_{p^n}^{\delta_{ij}} T_j^{\frac{1}{p^n}}$ and δ_{ij} denotes Kronecker's delta. Then for any $n \geq 0$, we have an exact sequence

$$(2.2) \quad 1 \rightarrow \Gamma_{\text{geo}}^{p^n} \rightarrow \Gamma_n \rightarrow \text{Gal}(K_{\text{cyc}}/K(\zeta_{p^n})) \rightarrow 1,$$

which splits and induces an isomorphism $\Gamma_n \cong \Gamma_{\text{geo}}^{p^n} \rtimes \text{Gal}(K_{\text{cyc}}/K(\zeta_{p^n}))$ such that for any $g \in \text{Gal}(K_{\text{cyc}}/K(\zeta_{p^n}))$ and any $1 \leq i \leq d$, $g\gamma_i^{p^n}g^{-1} = \gamma_i^{p^n\chi(g)}$.

Lemma 2.13. *The triple $(\{R_n\}_{n \geq 0}, \widehat{R}_\infty, \Gamma)$ is stably decompleting.*

Proof. It suffices to show that any $(\{R_n\}_{n \geq 0}, \widehat{R}_\infty, \Gamma)$ is weakly decompleting, as its pullbacks to rational localizations of X_n satisfy the same assumptions. We remark that $\Gamma \rightarrow \Gamma/\Gamma_n$ admits a continuous cross-section as the target is finite.

The condition (1) of Definition 2.3 is satisfied trivially. For condition (2), by noting that for any $n \geq 0$, \widehat{R}_∞ admits a Γ -equivariant decomposition

$$(2.3) \quad \widehat{R}_\infty = \bigoplus_{(\alpha_1, \dots, \alpha_d) \in (\mathbb{N}[\frac{1}{p}] \cap [0, 1))^d} R_{\widehat{K}_{\text{cyc}}, n} T_1^{\frac{1}{p^n}\alpha_1} \dots T_d^{\frac{1}{p^n}\alpha_d},$$

where $R_{\widehat{K}_{\text{cyc}}, n} = R_{\widehat{K}_{\text{cyc}}, n}^+[\frac{1}{p}]$ and $R_{\widehat{K}_{\text{cyc}}, n}^+ = R_n^+ \widehat{\otimes}_{\mathcal{O}_{K(\zeta_{p^n})}} \mathcal{O}_{\widehat{K}_{\text{cyc}}}$, it suffices to check $R_{\widehat{K}_{\text{cyc}}, n} \rightarrow R_{\widehat{K}_{\text{cyc}}, n}/R_n$ admits an isometric section as Banach R_n -modules, which can be verified as in the proof of [LZ17, Prop. A.2.1.1].

It remains to check condition (3) of Definition 2.3. For this purpose, it is enough to show that $\text{R}\Gamma(\Gamma_n, \widehat{R}_\infty^+/R_n^+)$ is killed by p^2 for $n \gg 0$. Similar to decomposition (2.3), we see that

$$\widehat{R}_\infty^+/R_n^+ = R_{\widehat{K}_{\text{cyc}}, n}^+/R_n^+ \oplus \bigoplus_{\underline{0} \neq (\alpha_1, \dots, \alpha_d) \in (\mathbb{N}[\frac{1}{p}] \cap [0, 1))^d} R_{\widehat{K}_{\text{cyc}}, n}^+ T_1^{\frac{1}{p^n}\alpha_1} \dots T_d^{\frac{1}{p^n}\alpha_d}.$$

By [BMS18, Lem. 7.3], $\text{R}\Gamma(\Gamma_{\text{geo}}^{p^n}, R_{\widehat{K}_{\text{cyc}}, n}^+ T_1^{\frac{1}{p^n}\alpha_1} \dots R_{\widehat{K}_{\text{cyc}}, n}^+ T_d^{\frac{1}{p^n}\alpha_d})$ can be computed via the Koszul complex $\text{K}(\gamma_1^{p^n} - 1, \dots, \gamma_d^{p^n} - 1; R_{\widehat{K}_{\text{cyc}}, n}^+ T_1^{\frac{1}{p^n}\alpha_1} \dots R_{\widehat{K}_{\text{cyc}}, n}^+ T_d^{\frac{1}{p^n}\alpha_d})$:

$$R_{\widehat{K}_{\text{cyc}}, n}^+ T_1^{\frac{1}{p^n}\alpha_1} \dots T_d^{\frac{1}{p^n}\alpha_d} \xrightarrow{(\gamma_1^{p^n} - 1, \dots, \gamma_d^{p^n} - 1)} (R_{\widehat{K}_{\text{cyc}}, n}^+ T_1^{\frac{1}{p^n}\alpha_1} \dots T_d^{\frac{1}{p^n}\alpha_d})^d \rightarrow \dots \rightarrow R_{\widehat{K}_{\text{cyc}}, n}^+ T_1^{\frac{1}{p^n}\alpha_1} \dots T_d^{\frac{1}{p^n}\alpha_d}.$$

Then for $\underline{\alpha} \neq \underline{0}$, we argue as in the proof of [Sch13, Lem. 6.18] to conclude that

$$\text{R}\Gamma(\Gamma_{\text{geo}}^{p^n}, R_{\widehat{K}_{\text{cyc}}, n}^+ T_1^{\frac{1}{p^n}\alpha_1} \dots R_{\widehat{K}_{\text{cyc}}, n}^+ T_d^{\frac{1}{p^n}\alpha_d})$$

is concentrated in degree ≥ 1 and is annihilated by $\zeta_p - 1$. Therefore $\text{R}\Gamma(\Gamma_n, R_{\widehat{K}_{\text{cyc}}, n}^+ T_1^{\frac{1}{p^n}\alpha_1} \dots T_d^{\frac{1}{p^n}\alpha_d})$ is also concentrated in degree ≥ 1 and is annihilated by $\zeta_p - 1$ and hence so is

$$\text{R}\Gamma(\Gamma_n, \bigoplus_{\underline{0} \neq (\alpha_1, \dots, \alpha_d) \in (\mathbb{N}[\frac{1}{p}] \cap [0, 1))^d} R_{\widehat{K}_{\text{cyc}}, n}^+ T_1^{\frac{1}{p^n}\alpha_1} \dots T_d^{\frac{1}{p^n}\alpha_d}).$$

To see $\text{R}\Gamma(\Gamma_n, R_{\widehat{K}_{\text{cyc}}, n}^+/R_n^+)$ is killed by p^2 for $n \gg 0$, we are reduced to the proof of Lemma 2.7.

Finally, we conclude that $\text{R}\Gamma(\Gamma_n, \widehat{R}_\infty^+/R_n^+)$ is killed by p^2 for $n \gg 0$ and then complete the proof. \square

Theorem 2.14. *The triple $(\{R_n\}_{n \geq 0}, \widehat{R}_\infty, \Gamma)$ is a decompletion system.*

Proof. This follows from Theorem 2.6 together with Lemma 2.13. \square

Notation 2.15. Let $\text{Rep}_{\Gamma}^{\text{uni}}(R_{\widehat{K}_{\text{cyc}}})$ be the full subcategory of $\text{Rep}_{\Gamma}(R_{\widehat{K}_{\text{cyc}}})$ of representations on which Γ_{geo} acts unipotently.

The following result is a direct consequence of Theorem 2.14.

Proposition 2.16. *There exists an equivalence of categories*

$$\mathrm{Rep}_\Gamma^{\mathrm{uni}}(R_{\widehat{K}_{\mathrm{cyc}}}) \rightarrow \mathrm{Rep}_\Gamma(\widehat{R}_\infty)$$

via base-change such that for any $M \in \mathrm{Rep}_\Gamma^{\mathrm{uni}}(R_{\widehat{K}_{\mathrm{cyc}}})$ with associated $M_\infty \in \mathrm{Rep}_\Gamma(\widehat{R}_\infty)$, there exist quasi-isomorphisms

$$\mathrm{R}\Gamma(\Gamma_{\mathrm{geo}}, M) \rightarrow \mathrm{R}\Gamma(\Gamma_{\mathrm{geo}}, M_\infty)$$

and

$$\mathrm{R}\Gamma(\Gamma, M) \rightarrow \mathrm{R}\Gamma(\Gamma, M_\infty).$$

The equivalence is compatible with the standard étale localisation of X .

Proof. Let $M \in \mathrm{Rep}_\Gamma^{\mathrm{uni}}(R_{\widehat{K}_{\mathrm{cyc}}})$ with $M_\infty = M \otimes_{R_{\widehat{K}_{\mathrm{cyc}}}} \widehat{R}_\infty$. Using decomposition (2.3) for $n = 0$, we obtain a Γ -equivariant decomposition

$$(2.4) \quad M_\infty = \widehat{\bigoplus_{(\alpha_1, \dots, \alpha_d) \in (\mathbb{N}[\frac{1}{p}] \cap [0,1))^d}} MT_1^{\alpha_1} \cdots T_d^{\alpha_d}.$$

Since Γ_{geo} acts on M unipotently, if $\alpha_i \neq 0$ for some $1 \leq i \leq d$, we deduce that $\gamma_i - 1$ acts on $MT_1^{\alpha_1} \cdots T_d^{\alpha_d}$ invertibly. Therefore, we see that

$$\mathrm{R}\Gamma(\Gamma_{\mathrm{geo}}, \widehat{\bigoplus_{\emptyset \neq (\alpha_1, \dots, \alpha_d) \in (\mathbb{N}[\frac{1}{p}] \cap [0,1))^d}} MT_1^{\alpha_1} \cdots T_d^{\alpha_d}) = 0,$$

which implies that the natural map

$$\mathrm{R}\Gamma(\Gamma_{\mathrm{geo}}, M) \rightarrow \mathrm{R}\Gamma(\Gamma_{\mathrm{geo}}, M_\infty)$$

is a quasi-isomorphism and hence so is

$$\mathrm{R}\Gamma(\Gamma, M) \rightarrow \mathrm{R}\Gamma(\Gamma, M_\infty).$$

In particular, we see the functor $\mathrm{Rep}_\Gamma^{\mathrm{uni}}(R_{\widehat{K}_{\mathrm{cyc}}}) \rightarrow \mathrm{Rep}_\Gamma(\widehat{R}_\infty)$ is fully faithful.

We have to check the essential surjectivity of the above functor. Fix an $M_\infty \in \mathrm{Rep}_\Gamma(\widehat{R}_\infty)$. By Theorem 2.14, it admits a good model M_n over R_n for some $n \gg 0$. Then the argument in proof of [LZ17, Lem. 2.15] shows that Γ_{geo} acts on M_n and hence $M_{\widehat{K}_{\mathrm{cyc}}, n} := M_n \otimes_{R_n} R_{\widehat{K}_{\mathrm{cyc}}, n}$ quasi-unipotently. Now, the arguments in the paragraph below [LZ17, Lem. 2.15] applies. But for the convenience of reader, we repeat the details as follows:

Since Γ_{geo} acts on $M_{\widehat{K}_{\mathrm{cyc}}, n}$ quasi-unipotently, we have a decomposition

$$M_{\widehat{K}_{\mathrm{cyc}}, n} = \bigoplus_\tau M_{\widehat{K}_{\mathrm{cyc}}, n, \tau}$$

where τ 's are finite characters of Γ_{geo} and

$$M_{\widehat{K}_{\mathrm{cyc}}, n, \tau} = \{x \in M_{\widehat{K}_{\mathrm{cyc}}, n} \mid (\gamma - \tau(\gamma))^m(x) = 0 \text{ for } m \gg 0, \forall \gamma \in \Gamma_{\mathrm{geo}}\}$$

denotes the corresponding generalised eigenspaces. Each $M_{\widehat{K}_{\mathrm{cyc}}, n, \tau}$ is finite projective over $R_{\widehat{K}_{\mathrm{cyc}}}$ as $M_{\widehat{K}_{\mathrm{cyc}}, n}$ is. After enlarging n if necessary, we may assume the orders of all characters appearing in the above decomposition divide p^n . Then for each τ , there exists a $T_1^{\alpha_1} \cdots T_d^{\alpha_d} \in R_{\widehat{K}_{\mathrm{cyc}}, n}$ on which Γ_{geo} acts via τ . Denote by $M := M_{\widehat{K}_{\mathrm{cyc}}, n, 1}$ the generalised eigenspace corresponding the trivial character. Then M is stable by the action of Γ and the natural map $M \otimes_{R_{\widehat{K}_{\mathrm{cyc}}}} R_{\widehat{K}_{\mathrm{cyc}}, n} \rightarrow M_{\widehat{K}_{\mathrm{cyc}}, n}$ is surjective and hence an isomorphism. Then the essential surjectivity follows as

$$M \otimes_{R_{\widehat{K}_{\mathrm{cyc}}}} \widehat{R}_\infty \cong M_{\widehat{K}_{\mathrm{cyc}}, n} \otimes_{R_{\widehat{K}_{\mathrm{cyc}}}} \widehat{R}_\infty \cong M_\infty$$

and Γ_{geo} acts on M unipotently.

Finally, we complete the proof by noting that all constructions above are compatible with standard étale localisations of X . \square

Remark 2.17. Proposition 2.10 shows that for any $M \in \text{Rep}_\Gamma(R_{\widehat{K}_{\text{cyc}}})$, Γ_{geo} acts quasi-unipotently and Remark 2.11 shows that the inclusion $\text{Rep}_\Gamma^{\text{uni}}(R_{\widehat{K}_{\text{cyc}}}) \hookrightarrow \text{Rep}_\Gamma(R_{\widehat{K}_{\text{cyc}}})$ is not an equivalence. For any $M \in \text{Rep}_\Gamma(R_{\widehat{K}_{\text{cyc}}})$, we describe its image via the composition

$$\square^{\text{uni}} : \text{Rep}_\Gamma(R_{\widehat{K}_{\text{cyc}}}) \rightarrow \text{Rep}_\Gamma(\widehat{R}_\infty) \xrightarrow{\sim} \text{Rep}_\Gamma^{\text{uni}}(R_{\widehat{K}_{\text{cyc}}}),$$

where the first arrow is induced by base-change, as follows:

Since Γ_{geo} acts on M quasi-unipotently, M admits a decomposition

$$M = \bigoplus_\tau M_\tau$$

where τ 's are finite characters on Γ_{geo} and M_τ is the corresponding generalised eigenspace as above. Let $\underline{T}^\tau \in \widehat{R}_\infty$ be of the form $T_1^{\alpha_1} \cdots T_d^{\alpha_d}$ on which Γ_{geo} acts via τ^{-1} . Then

$$M' := \bigoplus_\tau M_\tau \underline{T}^\tau \subset M \otimes_{R_{\widehat{K}_{\text{cyc}}}} \widehat{R}_\infty$$

such that Γ_{geo} acts on M^{uni} unipotently and that

$$M' \otimes_{R_{\widehat{K}_{\text{cyc}}}} \widehat{R}_\infty = M \otimes_{R_{\widehat{K}_{\text{cyc}}}} \widehat{R}_\infty.$$

It is easy to see that $M^{\text{uni}} := M'$ is the image of M under \square^{uni} .

Remark 2.18. Theorem 2.14 and Proposition 2.16 were proved in [Tsu18, §14] by working with a suitable integral model (with some certain log structure) of X .

Notation 2.19. For any p -adically complete field F containing \widehat{K}_{cyc} , let $R_{F,n}$ (resp. $\widehat{R}_{F,\infty}$) denote the base-change of $R_{\widehat{K}_{\text{cyc}},n}$ (resp. \widehat{R}_∞) along the inclusion $\widehat{K}_{\text{cyc}} \rightarrow F$. Let L/K be a Galois extension containing K_{cyc} in \overline{K} and let $\Gamma(L/K)$ be the Galois group of the cover $X_{\widehat{L},\infty} = \text{Spa}(\widehat{R}_{\widehat{L},\infty}, \widehat{R}_{\widehat{L},\infty}) \rightarrow X$. Then we have

$$\Gamma(L/K) \cong \Gamma_{\text{geo}} \rtimes \text{Gal}(L/K)$$

such that for any $1 \leq i \leq d$ and any $g \in \text{Gal}(L/K)$, $g\gamma_i g^{-1} = \gamma_i^{\chi(g)}$. Clearly, $\Gamma(K_{\text{cyc}}/K) = \Gamma$.

Remark 2.20. Note that for any Galois extension L/K containing K_{cyc} in \overline{K} , $\text{Gal}(L/K_{\text{cyc}})$ is a closed normal subgroup of $\Gamma(L/K)$ with corresponding quotient group $\Gamma \cong \Gamma(L/K)/\text{Gal}(L/K_{\text{cyc}})$. Then by Faltings' almost purity theorem, taking $\text{Gal}(L/K_{\text{cyc}})$ -invariants induces an equivalence of categories

$$\text{Rep}_{\Gamma(L/K)}^*(R_{\widehat{L}}) \simeq \text{Rep}_\Gamma^*(R_{\widehat{K}_{\text{cyc}}}) \quad (\text{resp. } \text{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L},\infty}) \simeq \text{Rep}_\Gamma(\widehat{R}_\infty))$$

for $*$ = \emptyset or uni. Then Proposition 2.10 implies that for any $M \in \text{Rep}_{\Gamma(L/K)}(R_{\widehat{L}})$, Γ_{geo} acts on M quasi-unipotently while Proposition 2.16 implies an equivalence of categories

$$\text{Rep}_{\Gamma(L/K)}^{\text{uni}}(R_{\widehat{L}}) \simeq \text{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L},\infty}).$$

For these reasons, from now on, we will not distinguish the category $\text{Rep}_{\Gamma(L/K)}^*(R_{\widehat{L}})$ (resp. $\text{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L},\infty})$) with the category $\text{Rep}_\Gamma^*(R_{\widehat{K}_{\text{cyc}}})$ (resp. $\text{Rep}_\Gamma(\widehat{R}_\infty)$), where $*$ = \emptyset or uni.

A similar but much easier argument for the proof of Lemma 2.13 also shows the following.

Lemma 2.21. *The triple $(\{R_{F,n}\}, \widehat{R}_{F,\infty}, \Gamma_{\text{geo}})$ is stably decompleting.*

We leave its proof to readers as we will not use this lemma in this paper.

Theorem 2.22. *The triple $(\{R_{F,n}\}, \widehat{R}_{F,\infty}, \Gamma_{\text{geo}})$ is a decomposition system.*

Proof. Just combine Theorem 2.6 with Lemma 2.21. □

Remark 2.23. (1) Theorem 2.22 was proved essentially in [AGT16] and [Tsu18].

(2) Unlike Proposition 2.16, an $\widehat{R}_{L,\infty}$ -representation of Γ_{geo} may not admits a model over R_L . However, if M is small, which means that the Γ_{geo} -action on M is “close” to the identity, then it admits a good model over R_L . See [AGT16, Chap. II.14], [Tsu18, §12] or [Wang21, §3] for details.

3. A p -ADIC SIMPSON CORRESPONDENCE FOR RIGID ANALYTIC VARIETIES OVER K

In this section, we construct a p -adic Simpson correspondence for generalised representations on $X_{\text{proét}}$ for rigid analytic varieties X of dimension d over K by working on the arenas of [LZ17] and [DLLZ18]. The main ingredient is a period sheaf \mathcal{OC} which was firstly studied in [Hy89] in the local case, was defined as the graded piece $\text{Gr}^0 \mathcal{OB}_{\text{dR}}$ of the de Rham period sheaf on $X_{\text{proét}}$ in [Sch13, §6], and was used to study Simpson correspondence in [LZ17]. It is a sheaf of $\widehat{\mathcal{O}}_X$ -algebras on $X_{\text{proét}}$ with a universal Higgs field

$$\Theta : \mathcal{OC} \rightarrow \mathcal{OC} \otimes_{\mathcal{O}_X} \Omega_X^1(-1)$$

such that the induced Higgs complex $\text{HIG}(\mathcal{OC}, \Theta)$ is an resolution of $\widehat{\mathcal{O}}_X$. As remarked in [LZ17, Rem. 2.1], let

$$0 \rightarrow \widehat{\mathcal{O}}_X \rightarrow \mathcal{E} \rightarrow \widehat{\mathcal{O}}_X \otimes_{\mathcal{O}_X} \Omega_X^1(-1) \rightarrow 0$$

denote the Faltings' extension (which can be found in [Sch13, Cor. 6.14] up to a Tate twist). Then

$$\mathcal{OC} = \varinjlim_n \text{Sym}^n \mathcal{E}$$

with the translation morphisms sending each local section $x_1 \otimes \cdots \otimes x_n$ of $\text{Sym}^n \mathcal{E}$ to $1 \otimes x_1 \otimes \cdots \otimes x_n$ and the universal Higgs field Θ is induced by sending each local section $x_1 \otimes \cdots \otimes x_n$ of $\text{Sym}^n \mathcal{E}$ to the local section $-\sum_{i=1}^n x_1 \otimes \cdots \otimes x_{i-1} \otimes x_{i+1} \otimes \cdots \otimes x_n \otimes \bar{x}_i$ of $\text{Sym}^{n-1} \mathcal{E} \otimes_{\mathcal{O}_X} \Omega_X^1(-1)$ ³, where \bar{x}_i denotes the image of x_i under the projection $\mathcal{E} \rightarrow \widehat{\mathcal{O}}_X \otimes_{\mathcal{O}_X} \Omega_X^1(-1)$. See [Hy89] for more details.

Definition 3.1. Let L/K be an algebraic Galois extension containing μ_{p^∞} with Galois group $\text{Gal}(L/K)$ and \widehat{L} be the completion of L in C . By a $\text{Gal}(L/K)$ -Higgs bundle of rank l on $X_{\widehat{L}, \text{ét}}$, we mean a Higgs bundle $(\mathcal{H}, \theta_{\mathcal{H}})$ with $\theta_{\mathcal{H}}$ nilpotent of rank l on $X_{\widehat{L}, \text{ét}}$ together with a $\text{Gal}(L/K)$ -action on \mathcal{H} such that the morphism

$$\mathcal{H} \xrightarrow{\theta_{\mathcal{H}}} \mathcal{H} \otimes_{\mathcal{O}_X} \Omega_X^1(-1)$$

is $\text{Gal}(L/K)$ -equivariant. We denote by $\text{HIG}_{\text{Gal}(L/K)}(X_{\widehat{L}})$ the category of $\text{Gal}(L/K)$ -Higgs bundles on $X_{\widehat{L}, \text{ét}}$. In particular, when $L = \overline{K}$ (and hence $\widehat{L} = C$), we denote $\text{HIG}_{\text{Gal}(\overline{K}/K)}(X_{\widehat{K}})$ by $\text{HIG}_{G_K}(X_C)$.

In particular, when $X = \text{Spa}(R, R^+)$ is smooth affinoid, one can define $\text{Gal}(L/K)$ -Higgs modules (H, θ_H) of rank l over $R_{\widehat{L}}$ similarly and denote the category of those (H, θ_H) by $\text{HIG}_{\text{Gal}(L/K)}(R_{\widehat{L}})$.

Remark 3.2. We claim that the nilpotency condition on $\theta_{\mathcal{H}}$ in Definition 3.1 is not necessary. To see this, we may assume $X = \text{Spa}(R, R^+)$ admits a toric chart and assume $(\mathcal{H}, \theta_{\mathcal{H}})$ is induced by a $\text{Gal}(K_{\text{cyc}}/K)$ -Higgs module (H, θ_H) over $R_{\widehat{K}_{\text{cyc}}}$. By Theorem 2.8, we may further assume (H, θ_H) is defined over $R_{K(\zeta_{p^n})}$ for some $n \geq 0$. Write $\Theta_H = \sum_{i=1}^d \theta_i \otimes \frac{d \log T_i}{t}$. Then for any $x \in H$ and any $g \in \text{Gal}(K_{\text{cyc}}/K)$, we have

$$g(\theta_H(x)) = g\left(\sum_{i=1}^d \theta_i(x) \otimes \frac{d \log T_i}{t}\right) = \chi(g)^{-1} \sum_{i=1}^d g(\theta_i(x)) \otimes \frac{d \log T_i}{t}$$

and that

$$\theta_H(g(x)) = \sum_{i=1}^d \theta_i(g(x)) \otimes \frac{d \log T_i}{t}.$$

So we get $g\theta_i g^{-1} = \chi(g)\theta_i$. Then the proof of [LZ17, Lem. 2.15] applies.

Then our main result can be stated as follows:

Theorem 3.3. Let $\nu : X_{\text{proét}} \rightarrow X_{C, \text{ét}}$ be the natural projection of sites. For any generalised representation $\mathcal{L} \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$, if we put $\Theta_{\mathcal{L}} := \text{id}_{\mathcal{L}} \otimes \Theta$, then the rule

$$\mathcal{L} \mapsto (\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})}) := (\nu_*(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{OC}), \nu_*(\Theta_{\mathcal{L}}))$$

induces a rank-preserving equivalence from the category $\text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ of generalised representations on $X_{\text{proét}}$ to the category $\text{HIG}_{G_K}(X_C)$ of G_K -Higgs bundles on $X_{C, \text{ét}}$, which preserves tensor products and dualities. Moreover, the following assertions are true:

³Our Higgs field Θ differs from that in [Hy89] by a sign but is compatible with $\text{Gr}^0 \nabla$ in [Sch13, §6].

- (1) For any $i \geq 1$, the higher direct image $R^i \nu_* (\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}_{\mathbb{C}}) = 0$.
- (2) Let $\Theta_{\mathcal{H}(\mathcal{L})} := \theta_{\mathcal{H}(\mathcal{L})} \otimes \text{id}_{\mathcal{O}_{\mathbb{C}}} + \text{id}_{\mathcal{H}(\mathcal{L})} \otimes \Theta$. Then there exists a natural isomorphism

$$(\mathcal{H} \otimes_{\mathcal{O}_{X_C}} \mathcal{O}_{\mathbb{C}}|_{X_C}, \Theta_{\mathcal{H}(\mathcal{L})}) \xrightarrow{\cong} (\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}_{\mathbb{C}}, \Theta_{\mathcal{L}})|_{X_C}$$

of Higgs fields.

- (3) Let $(\text{HIG}(\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})}))$ denote the Higgs complex induced by $(\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})})$. Then there exists a natural quasi-isomorphism

$$\text{R}\Gamma(X_{C, \text{proét}}, \mathcal{L}) \cong \text{R}\Gamma(X_{C, \text{ét}}, (\text{HIG}(\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})}))$$

which is compatible with G_K -actions. As a consequence, we get a quasi-isomorphism

$$\text{R}\Gamma(X_{\text{proét}}, \mathcal{L}) \cong \text{R}\Gamma(G_K, \text{R}\Gamma(X_{C, \text{ét}}, (\text{HIG}(\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})}))).$$

- (4) Let $X' \rightarrow X$ be a smooth morphism of rigid analytic varieties over K . Then the equivalence in (3) is compatible with pull-back along f . In other words, for any $\mathcal{L} \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ with corresponding $(\mathcal{H}, \theta_{\mathcal{H}}) \in \text{HIG}_{G_K}(X_C)$, we have

$$(\mathcal{H}(f^* \mathcal{L}), \theta_{\mathcal{H}(f^* \mathcal{L})}) \cong (f^* \mathcal{H}, f^* \theta_{\mathcal{H}}).$$

Remark 3.4. Let L/K be a Galois extension of K in \overline{K} containing K_{cyc} . By Faltings' almost purity theorem (and almost étale descent), the scalar extension

$$\text{HIG}_{\text{Gal}(L/K)}(X_{\widehat{L}}) \rightarrow \text{HIG}_{G_K}(X_C)$$

is indeed an equivalence of categories whose quasi-inverse is induced by taking $\text{Gal}(\overline{K}/L)$ -invariants. For the same reason, one can replace C and G_K by \widehat{L} and $\text{Gal}(L/K)$ respectively in the statement of Theorem 3.3 and then conclude an equivalence of categories

$$\text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X) \simeq \text{HIG}_{\text{Gal}(L/K)}(X_{\widehat{L}})$$

such that the corresponding assertions (1), (2) and (3) hold true, and are compatible with Theorem 3.3 via the base-change along $\widehat{L} \rightarrow C$.

Remark 3.5. By analytic-étale comparison (cf. [FvdP04, Prop. 8.2.3]), we may replace étale site $X_{C, \text{ét}}$ by analytic site $X_{C, \text{an}}$ in the statement of Theorem 3.3 such that all results still hold true. The same remark also applies to Remark 3.4.

- Remark 3.6.**
- (1) When X is affine, Theorem 3.3 was also achieved by Tsuji [Tsu18, Thm 15.2] by choosing a certain integral model of X (and considering a certain log structure on the chosen model), which is not necessary in our approach.
 - (2) When $\mathcal{L} = \mathbb{L} \otimes_{\mathbb{Z}} \widehat{\mathcal{O}}_X$ is induced by a \mathbb{Q}_p -local system on $X_{\text{ét}}$, Theorem 3.3 reduces to [LZ17, Thm 2.1] by studying decompletion theory for relative analogue of the overconvergent period ring $\widetilde{\mathbf{B}}^{\dagger}$. Our proof is inspired of the work in [LZ17] and improve theirs to any generalised representations by using decompletion theory in the previous section.
 - (3) Theorem 3.3 can be also deduced from [Heu22a] when X is a curve and [HMW22] when X is abeloid by noticing that the morphism “HTlog” in loc.cit. is G_K -equivariant.

Before moving on, let us give a immediate corollary of Theorem 3.3.

Corollary 3.7. *Let d be the dimension of X over K .*

- (1) *Assume X is quasi-compact. Then $\text{R}\Gamma(X_{C, \text{proét}}, \mathcal{L})$ is concentrated in degree $[0, 2d]$ and $\text{R}\Gamma(X_{\text{proét}}, \mathcal{L})$ is concentrated in degree $[0, 2d + 1]$.*
- (2) *If moreover X is proper, then $\text{R}\Gamma(X_{C, \text{proét}}, \mathcal{L})$ is a perfect complex of C -representations of G_K and $\text{R}\Gamma(X_{\text{proét}}, \mathcal{L})$ is a perfect complex of K -vector spaces.*

Proof. It suffices to prove Item (1) as Item (2) follows from the properness of X .

Since X is quasi-compact and locally noetherian, it is a noetherian space. Then we obtain that $\text{R}\Gamma(X_{C, \text{proét}}, \mathcal{L})$ is concentrated in degree $[0, 2d]$ by combining Remark 3.5 with Grothendieck's vanishing theorem [Gro57, Thm 3.6.5].

It remains to show that $\text{R}\Gamma(X_{\text{proét}}, \mathcal{L})$ is concentrated in degree $[0, 2d + 1]$. Due to Remark 3.4, one may replace \widehat{K}_{cyc} and $\text{Gal}(K_{\text{cyc}}/K)$ instead of C and G_K in Theorem 3.3 and deduce that

$R\Gamma(X_{\widehat{K}_{\text{cyc}, \text{proét}}, \mathcal{L}})$ is concentrated in degree $[0, 2d]$. Then we conclude by using Theorem 3.3 (3) by noting that $\text{Gal}(K_{\text{cyc}}/K)$ is of cohomological dimension 1. \square

3.1. A local version of Theorem 3.3. This subsection is devoted to a local version of Theorem 3.3. More precisely, we assume $X = \text{Spa}(R, R^+)$ is smooth affinoid of dimension d over K , which admits a toric chart, and keep the notations in Notation 2.12 and Notation 2.19.

The following lemma is well-known:

Lemma 3.8. *The evaluation at $X_{\widehat{L}, \infty}$ induces an equivalence from the category $\text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ to the category $\text{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L}, \infty})$ of representations of $\Gamma(L/K)$ over $\widehat{R}_{\widehat{L}, \infty}$, which preserves tensor products and dualities. Moreover, for any $\mathcal{L} \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$, there exists a natural quasi-isomorphism*

$$R\Gamma(X_{\text{proét}}, \mathcal{L}) \simeq R\Gamma(\Gamma(L/K), \mathcal{L}(X_{\widehat{L}, \infty})).$$

Proof. Note that we may regard generalised representations as $\widehat{\mathcal{O}}_X$ -crystals on $X_{\text{proét}, \text{aff}, \text{perf}}$. Since $X_{\widehat{L}, \infty}$ is a cover of X with Galois group $\Gamma(L/K)$, it is a cover of the final object of $\text{Shv}(X_{\text{proét}})$. If we denote the corresponding Čech nerve by $X_{\widehat{L}, \infty}^{\bullet/X}$, then for any $n \geq 0$, $X_{\widehat{L}, \infty}^{n/X}$ is the self-product of $n + 1$ copies of $X_{\widehat{L}, \infty}$ over X and is isomorphic to

$$X_{\widehat{L}, \infty} \times \underline{\Gamma(L/K)^n} = \text{Spa}(C(\Gamma(L/K)^n, \widehat{R}_{\widehat{L}, \infty}), C(\Gamma(L/K)^n, \widehat{R}_{\widehat{L}, \infty}^+))$$

as argued in the proof of [Sch13, Lem. 5.6], where $C(\Gamma(L/K)^n, \widehat{R}_{\widehat{L}, \infty}^+)$ denotes the continuous functions from $\Gamma(L/K)^n$ to $\widehat{R}_{\widehat{L}, \infty}^+$. In particular, we have an isomorphism of cosimplicial rings

$$\widehat{\mathcal{O}}_X(X_{\widehat{L}, \infty}^{\bullet/X}) \cong C(\Gamma(L/K)^\bullet, \widehat{R}_{\widehat{L}, \infty}).$$

By [KL15, Thm. 9.2.15], we get an equivalence of categories

$$\text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X) \simeq \text{Strat}(\widehat{\mathcal{O}}_X(X_{\widehat{L}, \infty}^{\bullet/X})) \simeq \text{Strat}(C(\Gamma(L/K)^\bullet, \widehat{R}_{\widehat{L}, \infty}))$$

Then the desired equivalence follows from the Galois descent. It follows from the standard linear algebra that the equivalence above preserves tensor products and dualities.

It remains to prove the “moreover” part. Using [KL15, Thm. 9.2.15] again, we get isomorphisms

$$\mathcal{L}(X_{\widehat{L}, \infty}^{\bullet/X}) \cong \mathcal{L}(X_{\widehat{L}, \infty}) \otimes_{\widehat{R}_{\widehat{L}, \infty}} C(\Gamma(L/K)^\bullet, \widehat{R}_{\widehat{L}, \infty}) \cong C(\Gamma(L/K)^\bullet, \mathcal{L}(X_{\widehat{L}, \infty})).$$

Since $R\Gamma(\Gamma(L/K), \mathcal{L}(X_{\widehat{L}, \infty}))$ is computed by $C(\Gamma(L/K)^\bullet, \mathcal{L}(X_{\widehat{L}, \infty}))$, we are reduced to showing that $R\Gamma(X_{\text{proét}}, \mathcal{L})$ can be computed by the Čech-Alexander complex $\mathcal{L}(X_{\widehat{L}, \infty}^{\bullet/X})$. However, this follows from [LZ17, Prop. 2.3] together with the Čech-to-derived spectral sequence. \square

According to Lemma 3.8, we may work with \widehat{R}_∞ -representations of Γ instead of generalised representations on $X_{\text{proét}}$ in this subsection. We also need the following local description of \mathcal{OC} .

Lemma 3.9. *Let L/K be a Galois extension in \overline{K} containing K_{cyc} . There is an isomorphism of sheaves*

$$\widehat{\mathcal{O}}_X[Y_1, \dots, Y_d]_{|X_{\widehat{L}, \infty}} \rightarrow \mathcal{OC}_{|X_{\widehat{L}, \infty}}$$

on $X_{\text{proét}}/X_{\widehat{L}, \infty}$ such that for any $1 \leq i \leq d$, $Y_i = t^{-1} \log(\frac{[T_i]^\flat}{T_i})$ and via this isomorphism the Higgs field $\Theta = \sum_{i=1}^d -\frac{\partial}{\partial Y_i} \otimes \frac{d \log T_i}{t}$. Moreover, the action of $\Gamma(L/K) \cong \Gamma_{\text{geo}} \rtimes \text{Gal}(L/K)$ on Y_i 's is determined such that for any $1 \leq i, j \leq d$ and any $g \in \text{Gal}(L/K)$, we have $\gamma_i(Y_j) = Y_j + \delta_{ij}$ and $g(Y_j) = \chi(g)^{-1} Y_j$.

Proof. Let $X_i := T_i - [T_i]^\flat$ as in [Sch13, Prop. 6.10]. Then the desired isomorphism follows from [Sch13, Cor. 6.15] by noting that $\frac{X_i}{\xi}$ and Y_i differ from a unit in $\text{Gr}^0 \mathcal{O}\mathbb{B}_{\text{dR}}$. The expression of Θ

follows from the definition of the connection ∇ on $\mathcal{O}\mathbb{B}_{\text{dR}}^4$. Finally, the $\Gamma(L/K)$ -action on Y_i 's can be deduced similarly as in the proof of [Sch13, Lem. 6.17]. \square

Notation 3.10. Let $C_{L,\infty} := \mathcal{O}\mathbb{C}(X_{\widehat{L},\infty})$ with induced Higgs field Θ . Then Lemma 3.9 provides a $\Gamma(L/K)$ -equivariant isomorphism of Higgs fields

$$\iota : (\widehat{R}_{\widehat{L},\infty}[Y_1, \dots, Y_d], \sum_{i=1}^d -\frac{\partial}{\partial Y_i} \otimes \frac{d\log T_i}{t}) \xrightarrow{\cong} (C_{L,\infty}, \Theta),$$

where the Γ -action on $\widehat{R}_{\widehat{L},\infty}[Y_1, \dots, Y_d]$ is determined as in Lemma 3.9. Then we obtain a $\Gamma(L/K)$ -equivariant inclusion of Higgs fields $(R_{\widehat{L}}[Y_1, \dots, Y_d], \sum_{i=1}^d -\frac{\partial}{\partial Y_i} \otimes \frac{d\log T_i}{t}) \subset (C_{L,\infty}, \Theta)$ via ι .

Then the main result in this subsection is

Theorem 3.11. *Keep notations as above.*

- (1) For any $M_\infty \in \text{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L},\infty})$, let $H(M_\infty) := (M_\infty \otimes_{\widehat{R}_{\widehat{L},\infty}} C_{L,\infty})^{\Gamma_{\text{geo}}}$ and $\theta_{H(M_\infty)}$ be the restriction of $\Theta_{M_\infty} := \text{id}_{M_\infty} \otimes \Theta$ to $H(M_\infty)$. Then $(H(M_\infty), \theta_{H(M_\infty)})$ defines a $\text{Gal}(L/K)$ -Higgs modules over $R_{\widehat{L}}$. Moreover, for any $i \geq 1$, we have $H^i(\Gamma_{\text{geo}}, M_\infty \otimes_{\widehat{R}_{\widehat{L},\infty}} C_{L,\infty}) = 0$.
- (2) For any $(H, \theta_H) \in \text{HIG}_{\text{Gal}(L/K)}(R_{\widehat{L}})$, let $\Theta_H := \theta_H \otimes \text{id}_{C_{L,\infty}} + \text{id}_H \otimes \Theta$. Then $M_\infty(H, \theta_H) := (H \otimes_{R_{\widehat{L}}} C_{L,\infty})^{\Theta_H=0}$ is an $\widehat{R}_{\widehat{L},\infty}$ -representation of $\Gamma(L/K)$.
- (3) The functor $M_\infty \mapsto (H(M_\infty), \theta_{H(M_\infty)})$ induces an equivalence of categories

$$\text{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L},\infty}) \rightarrow \text{HIG}_{\text{Gal}(L/K)}(R_{\widehat{L}}),$$

whose quasi-inverse is given by the functor $(H, \theta_H) \mapsto M_\infty(H, \theta_H)$. The equivalence preserves ranks, tensor products and dualities.

- (4) For any $M_\infty \in \text{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L},\infty})$ with the associated $\text{Gal}(L/K)$ -Higgs module $(H, \theta_H) \in \text{HIG}_{\text{Gal}(L/K)}(R_{\widehat{L}})$. Then there exists a $\Gamma(L/K)$ -equivariant isomorphism of Higgs fields

$$(H \otimes_{R_{\widehat{L}}} C_{L,\infty}, \Theta_H) \cong (M_\infty \otimes_{\widehat{R}_{\widehat{L},\infty}} C_{L,\infty}, \Theta_{M_\infty})$$

which induces an $\text{Gal}(L/K)$ -equivariant quasi-isomorphism

$$\text{R}\Gamma(\Gamma_{\text{geo}}, M_\infty) \cong \text{HIG}(H, \theta_H),$$

where $\text{HIG}(H, \theta_H)$ denotes the Higgs field induced by (H, θ_H) . As a consequence, we have a quasi-isomorphism

$$\text{R}\Gamma(\Gamma(L/K), M_\infty) \cong \text{R}\Gamma(\text{Gal}(L/K), \text{HIG}(H, \theta_H)).$$

- (5) All results above are compatible with standard étale localisation of X .

Proof. (1) Let M be the representation of $\Gamma(L/K)$ over $R_{\widehat{L}}$ corresponding to $\mathcal{L}(X_{\widehat{L},\infty})$ in the sense of Proposition 2.16 (and Remark 2.20). In other words, it satisfies the following conditions:

- (a) $M_\infty \cong M \otimes_{R_{\widehat{L}}} \widehat{R}_{\widehat{L},\infty}$ as representations of $\Gamma(L/K)$ over $\widehat{R}_{\widehat{L},\infty}$.
- (b) Γ_{geo} acts on M unipotently and induces quasi-isomorphisms

$$H^i(\Gamma_{\text{geo}}, M) \cong H^i(\Gamma_{\text{geo}}, \mathcal{L}(X_{\widehat{L},\infty})).$$

In particular, we have

$$M \otimes_{R_{\widehat{L}}} C_{L,\infty} \cong M_\infty \otimes_{\widehat{R}_{\widehat{L},\infty}} C_{L,\infty}.$$

Using this to replace [LZ17, Prop. 2.8], we may conclude by the argument in the paragraph below [LZ17, Prop. 2.10]. But for the further use, we provide more details here.

Consider the $\Gamma(L/K)$ -equivariant inclusion

$$R_{\widehat{L}}[Y_1, \dots, Y_d] \subset \widehat{R}_{\widehat{L},\infty}[Y_1, \dots, Y_d] = \mathcal{O}\mathbb{C}(X_{\widehat{L},\infty}).$$

⁴Roughly speaking, if we regard $\mathcal{O}\mathbb{B}_{\text{dR}}$ as $\mathcal{O}_X \otimes_K \mathbb{B}_{\text{dR}}$, then ∇ is induced by that on \mathcal{O}_X via the scalar extension $K \rightarrow \mathbb{B}_{\text{dR}}$. See [Sch13, §6] for details. Using $\lim_{n \rightarrow +\infty} \frac{\gamma_i^{p^n} - 1}{p^n}(Y_j) = \delta_{ij}$, one deduce that the Higgs field is induced by the action of Lie algebra $\text{Lie}(\Gamma_{\text{geo}})$ of Γ_{geo} on Y_i 's. More precisely, it is indeed given by $\Theta = \sum_{i=1}^d -\log \gamma_i \otimes \frac{\log T_i}{t}$.

Then by condition (2) and the argument in the paragraph after [Sch13, Lem. 6.17], for any $i \geq 0$, we get an isomorphism

$$H^i(\Gamma_{\text{geo}}, M \otimes_{R_{\widehat{L}}} R_{\widehat{L}}[Y_1, \dots, Y_d]) \cong H^i(\Gamma_{\text{geo}}, (\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}_{\mathbb{C}})(X_{\widehat{L}, \infty})).$$

Using condition (1) above together [LZ17, Lem. 2.10], it follows from Hochschild–Serre spectral sequence that

$$H^i(\Gamma_{\text{geo}}, M \otimes_{R_{\widehat{L}}} R_{\widehat{L}}[Y_1, \dots, Y_d]) = 0$$

for any $i \geq 1$, which settles the “moreover” part of (1).

Since $\gamma_i^{-1} - 1$ acts on M nilpotently, for any $x \in M$,

$$\prod_{i=1}^d \gamma_i^{-Y_i} x := \sum_{n_1, \dots, n_d \geq 0} \left(\prod_{i=1}^d (\gamma_i^{-1} - 1)^{n_i}(x) \right) \binom{Y_1}{n_1} \cdots \binom{Y_d}{n_d}$$

is well-defined in $M \otimes_{R_{\widehat{L}}} R_{\widehat{L}}[Y_1, \dots, Y_d]$. By chasing the proof of [LZ17, Lem. 2.10], we see that

$$(3.1) \quad H(M_{\infty}) = H^0(\Gamma_{\text{geo}}, M \otimes_{R_{\widehat{L}}} R_{\widehat{L}}[Y_1, \dots, Y_d]) = \prod_{i=1}^d \gamma_i^{-Y_i}(M) := \left\{ \prod_{i=0}^d \gamma_i^{-Y_i} x \mid x \in M \right\}$$

is a finite projective $R_{\widehat{L}}$ -module as desired. Using the expression of Θ in Lemma 3.9, we see that

$$(3.2) \quad \theta_{H(M_{\infty})} = \sum_{i=1}^d \log \gamma_i \otimes \frac{d \log T_i}{t}.$$

Since for any $g \in \text{Gal}(L/K)$, $g\gamma_i g^{-1} = \gamma_i^{\chi(g)}$. We see that $g \log \gamma_i g^{-1} = \chi(g) \log \gamma_i$. Therefore, we obtain that $(H(M_{\infty}), \theta_{H(M_{\infty})}) \in \text{HIG}_{\text{Gal}(L/K)}(R_{\widehat{L}})$.

(2) We write $\theta_H = \sum_{i=1}^d \theta_i \otimes \frac{d \log T_i}{t}$ with nilpotent $R_{\widehat{L}}$ -linear endomorphisms θ_i 's of H which commute with each other such that for any $g \in \text{Gal}(L/K)$, $g\theta_i g^{-1} = \chi(g)\theta_i$. Note that for any $x \in H \otimes_{R_{\widehat{K}_{\text{cyc}}}} \widehat{R}_{\widehat{K}_{\text{cyc}}, \infty}[Y_1, \dots, Y_d]$, it is uniquely written as

$$x = \sum_{\underline{n}=(n_1, \dots, n_d) \in \mathbb{N}^d} h_{\underline{n}} Y_1^{[n_1]} \cdots Y_d^{[n_d]}$$

for finitely many $h_{\underline{n}}$'s in $H \otimes_{R_{\widehat{K}_{\text{cyc}}}} \widehat{R}_{\widehat{K}_{\text{cyc}}, \infty}$, where $Y^{[n]}$ denotes the polynomial $\frac{Y^n}{n!}$ for all $n \geq 0$. Using the description of Θ in Lemma 3.9, we see that

$$\begin{aligned} \Theta_H(x) &= \sum_{\underline{n}=(n_1, \dots, n_d) \in \mathbb{N}^d} (\theta_i(h_{\underline{n}}) Y_1^{[n_1]} \cdots Y_d^{[n_d]} - h_{\underline{n}} Y_1^{[n_1]} \cdots Y_i^{[n_i-1]} \cdots Y_d^{[n_d]}) \otimes \frac{d \log T_i}{t} \\ &= \sum_{\underline{n}=(n_1, \dots, n_d) \in \mathbb{N}^d} (\theta_i(h_{\underline{n}}) - h_{\underline{n}+\underline{1}_i}) Y_1^{[n_1]} \cdots Y_d^{[n_d]} \otimes \frac{d \log T_i}{t}. \end{aligned}$$

Therefore, we deduce that $\Theta_H(x) = 0$ if and only if for any $\underline{n} \in \mathbb{N}^d$ and any $1 \leq i \leq d$,

$$h_{\underline{n}} = \theta_i(h_{\underline{n}-\underline{1}_i}).$$

By iteration, we obtain that for any $\underline{n} = (n_1, \dots, n_d) \in \mathbb{N}^d$,

$$h_{\underline{n}} = \theta_1^{n_1} \cdots \theta_d^{n_d}(h_{\underline{0}}).$$

Since θ_i 's are nilpotent, we see that $h_{\underline{n}} = 0$ for $|\underline{n}| \gg 0$ and that

$$\prod_{i=1}^d \exp(\theta_i Y_i)(h) := \sum_{\underline{n}=(n_1, \dots, n_d) \in \mathbb{N}^d} \theta_1^{n_1} \cdots \theta_d^{n_d}(h) Y_1^{[n_1]} \cdots Y_d^{[n_d]}$$

is well defined for any $h \in H \otimes_{R_{\widehat{K}_{\text{cyc}}}} \widehat{R}_{\widehat{K}_{\text{cyc}}, \infty}$. So we obtain that

$$(3.3) \quad M_{\infty}(H, \theta_H) = \prod_{i=1}^d \exp(\theta_i Y_i)(H \otimes_{R_{\widehat{K}_{\text{cyc}}}} \widehat{R}_{\widehat{K}_{\text{cyc}}, \infty}) = \left\{ \prod_{i=1}^d \exp(\theta_i Y_i)(h) \mid h \in H \otimes_{R_{\widehat{K}_{\text{cyc}}}} \widehat{R}_{\widehat{K}_{\text{cyc}}, \infty} \right\},$$

which is finite projective over $\widehat{R}_{\widehat{K}_{\text{cyc}}, \infty}$. By Lemma 3.9, the Γ_{geo} -action on M is induced such that for any $1 \leq i \leq d$,

$$(3.4) \quad \gamma_i = \exp(\theta_i)$$

Then for any $g \in \text{Gal}(L/K)$, $g\gamma_i g^{-1} = \gamma_i^{\chi(g)}$ (as $g\theta_i g^{-1} = \chi(g)\theta_i$). In other words, $M_\infty(H, \theta_H)$ belongs to $\text{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L}, \infty})$.

(3) Fix an $M_\infty \in \text{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L}, \infty})$ and let M be as in the proof of (1). Since Γ_{geo} acts on M unipotently, we see that $\prod_{i=1}^d \gamma_i^{-Y_i}$ is a well-defined isomorphism of $M \otimes_{R_{\widehat{L}}} R_{\widehat{L}}[Y_1, \dots, Y_d]$. So we see that

$$M \otimes_{R_{\widehat{L}}} R_{\widehat{L}}[Y_1, \dots, Y_d] \cong H(M_\infty) \otimes_{R_{\widehat{L}}} R_{\widehat{L}}[Y_1, \dots, Y_d]$$

and hence that

$$M_\infty \otimes_{\widehat{R}_{\widehat{L}, \infty}} \widehat{R}_{\widehat{L}, \infty}[Y_1, \dots, Y_d] \cong H(M_\infty) \otimes_{R_{\widehat{L}}} \widehat{R}_{\widehat{L}, \infty}[Y_1, \dots, Y_d].$$

Using (3.1) and (3.2), we see above two isomorphisms are compatible with both Higgs fields and $\Gamma(L/K)$ -actions. In particular, we see that $M_\infty = M_\infty(H(M_\infty), \theta_{H(M_\infty)})$.

Similarly, fix an $(H, \theta_H) \in \text{HIG}_{\text{Gal}(L/K)}(R_{\widehat{L}})$ and write $\theta_H = \sum_{i=1}^d \theta_i \otimes \frac{d\log T_i}{t}$ as in the proof of (2). Since θ_i 's are nilpotent, we see that $\prod_{i=1}^d \exp(\theta_i Y_i)$ is a well-defined isomorphism of $H \otimes_{R_{\widehat{L}}} \widehat{R}_{\widehat{L}, \infty}[Y_1, \dots, Y_d]$. So we get an isomorphism

$$H \otimes_{R_{\widehat{L}}} \widehat{R}_{\widehat{L}, \infty}[Y_1, \dots, Y_d] \cong M_\infty(H, \theta_H) \otimes_{\widehat{R}_{\widehat{L}, \infty}} \widehat{R}_{\widehat{L}, \infty}[Y_1, \dots, Y_d].$$

By (3.3) and (3.4), the above isomorphism is compatible with both Higgs fields and $\Gamma(L/K)$ -actions. So we deduce that $(H(M_\infty(H, \theta_H)), \theta_{H(M_\infty(H, \theta_H))}) = (H, \theta_H)$.

Therefore, we see functors defined in (1) and (2) induce a rank-preserving equivalence between $\text{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L}, \infty})$ and $\text{HIG}_{\text{Gal}(L/K)}(R_{\widehat{L}})$. By standard linear algebra, we see above constructions are compatible with tensor products and dualities, which completes the proof of (3).

(4) The first part has been established in (3). To complete the proof, we are reduced to showing that

$$\text{R}\Gamma(\Gamma_{\text{geo}}, M_\infty) \simeq \text{HIG}(H, \theta_H).$$

Since Higgs complex $\text{HIG}(C_{L, \infty}, \Theta)$ induced by $(C_{L, \infty}, \Theta)$ is a resolution of $\widehat{R}_{\widehat{L}, \infty}$, we get quasi-isomorphisms

$$\text{R}\Gamma(\Gamma_{\text{geo}}, M_\infty) \cong \text{R}\Gamma(\Gamma_{\text{geo}}, \text{HIG}(M_\infty \otimes_{\widehat{R}_{\widehat{L}, \infty}} C_{L, \infty}, \Theta_{M_\infty})) \cong \text{R}\Gamma(\Gamma_{\text{geo}}, \text{HIG}(H \otimes_{R_{\widehat{L}}} C_{L, \infty}, \Theta_H)).$$

By (1), for any $i \geq 1$ and any $j \geq 0$, we have

$$\text{H}^i(\Gamma_{\text{geo}}, H \otimes_{R_{\widehat{L}}} C_{L, \infty} \otimes_R \Omega_R^j(-j)) = 0.$$

Using spectral sequence, we get a quasi-isomorphism

$$\text{R}\Gamma(\Gamma_{\text{geo}}, \text{HIG}(H \otimes_{R_{\widehat{L}}} C_{L, \infty}, \Theta_H)) \simeq \text{HIG}(H, \theta_H),$$

and hence that

$$\text{R}\Gamma(\Gamma_{\text{geo}}, M_\infty) \simeq \text{HIG}(H, \theta_H)$$

as desired. This completes the proof of (4).

(5) Since the equivalence in Proposition 2.16 is compatible with standard étale localisation, this follows by noting all constructions above are compatible with standard étale localisation. \square

Remark 3.12. For any $M_\infty \in \text{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L}, \infty})$, let $M \in \text{Rep}_{\Gamma(L/K)}^{\text{uni}}(R_{\widehat{L}})$ be as in the proof of Theorem 3.11 (1). Then one can check that $(M, \theta_M := \sum_{i=1}^d \log \gamma_i \otimes \frac{d\log T_i}{t})$ with the restricted $\text{Gal}(L/K)$ -action gives rise to an object in $\text{HIG}_{\text{Gal}(L/K)}(R_{\widehat{L}})$. Let $(H(M_\infty), \theta_{H(M_\infty)})$ be as in (3.1) and (3.2). It is easy to see that the map $\prod_{i=1}^d \gamma_i^{-Y_i} x \mapsto x$ gives rise to an isomorphism

$$(H(M_\infty), \theta_{H(M_\infty)}) \cong (M, \theta_M)$$

of objects in $\text{HIG}_{\text{Gal}(L/K)}(R_{\widehat{L}})$. This phenomenon was discovered in [LZ17, Lem. 2.11].

Corollary 3.13. *Keep notations as above. There exists an equivalence of categories*

$$\mathrm{Vect}(X_{\mathrm{pro\acute{e}t}}, \widehat{\mathcal{O}}_X) \simeq \mathrm{HIG}_{\mathrm{Gal}(L/K)}(X_{\widehat{L}})$$

such that for any generalised representation \mathcal{L} on $X_{\mathrm{pro\acute{e}t}}$ with associated $\mathrm{Gal}(L/K)$ -Higgs bundle $(\mathcal{H}, \theta_{\mathcal{H}})$, there exists a quasi-isomorphism

$$\mathrm{R}\Gamma(X_{\mathrm{pro\acute{e}t}}, \mathcal{L}) \cong \mathrm{R}\Gamma(\mathrm{Gal}(L/K), \mathrm{R}\Gamma(X_{\widehat{L}, \acute{e}t}, \mathrm{HIG}(\mathcal{H}, \theta_{\mathcal{H}}))).$$

Proof. The desired equivalence is induced by the composition

$$\mathrm{Vect}(X_{\mathrm{pro\acute{e}t}}, \widehat{\mathcal{O}}_X) \rightarrow \mathrm{Rep}_{\Gamma(L/K)}(\widehat{R}_{\widehat{L}, \infty}) \rightarrow \mathrm{HIG}_{\mathrm{Gal}(L/K)}(R_{\widehat{L}}) \xleftarrow{\Gamma(X_{\widehat{L}}, -)} \mathrm{HIG}_{\mathrm{Gal}(L/K)}(X_{\widehat{L}, \acute{e}t}),$$

where the first two arrows are induced by Lemma 3.8 and Theorem 3.11 (3) while the last arrow is induced by taking global sections $\Gamma(X_{\widehat{L}}, -)$, which is an equivalence as X is affinoid. Finally, since X is affinoid, we get

$$\mathrm{R}\Gamma(X_{\widehat{L}, \acute{e}t}, \mathrm{HIG}(\mathcal{H}, \theta_{\mathcal{H}})) \simeq \mathrm{HIG}(H, \theta_H),$$

where (H, θ_H) is the $\mathrm{Gal}(L/K)$ -Higgs module induced by $(\mathcal{H}, \theta_{\mathcal{H}})$. So the desired quasi-isomorphism follows from Lemma 3.8 combined with Theorem 3.11 (4). \square

Remark 3.14. We will see in the next subsection (cf. Corollary 3.17) that the functor

$$\mathrm{Vect}(X_{\mathrm{pro\acute{e}t}}, \widehat{\mathcal{O}}_X) \rightarrow \mathrm{HIG}_{\mathrm{Gal}(L/K)}(X_{\widehat{L}})$$

is actually induced by $\mathcal{L} \mapsto (\nu_*(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}), \nu_*(\theta_{\mathcal{L}}))$ as stated in Theorem 3.3. So it does **NOT** depend on the choice of toric chart on X . So one can glue the local constructions together to get a global equivalence.

3.2. Proof of Theorem 3.3. In this subsection, we focus on the proof of Theorem 3.3.

We first show that $\mathrm{R}^i \nu_*(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C})$ is a vector bundle on $X_{C, \acute{e}t}$ for $i = 0$ and vanishes for $i \geq 1$. Note that it is the sheafification of the presheaf

$$(Y \in X_{C, \acute{e}t}) \mapsto \mathrm{H}^i(X_{\mathrm{pro\acute{e}t}}/Y, \mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}).$$

By [LZ17, Cor. 2.6], we are reduced to showing that for any affinoid $Y = \mathrm{Spa}(R, R^+) \in X_{\acute{e}t}$ admitting a toric chart, $\mathrm{H}^0(X_{\mathrm{pro\acute{e}t}}/Y_C, \mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C})$ is a finite projective $R_{\widehat{L}}$ -module, and that for any standard étale localisation $Z = \mathrm{Spa}(S, S^+) \rightarrow Y$,

$$(3.5) \quad \mathrm{H}^0(X_{\mathrm{pro\acute{e}t}}/Z_C, \mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}) = \mathrm{H}^0(X_{\mathrm{pro\acute{e}t}}/Y_C, \mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}) \otimes_{R_C} S_C$$

and for any $i \geq 1$,

$$(3.6) \quad \mathrm{H}^i(X_{\mathrm{pro\acute{e}t}}/Z_C, \mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}) = 0.$$

We proceed as in [LZ17, §2.3].

Lemma 3.15. *For any affinoid perfectoid $U \in X_{\mathrm{pro\acute{e}t}}$ and for any $i \geq 1$,*

$$\mathrm{H}^i(X_{\mathrm{pro\acute{e}t}}/U, \mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}) = 0.$$

Proof. Since U is affinoid, the pro-étale site $U_{\mathrm{pro\acute{e}t}}$ is coherent. In particular, taking cohomology commutes with taking direct limits. Since $\mathcal{O}\mathbb{C} = \varinjlim_n \mathrm{Sym}_{\widehat{\mathcal{O}}_X}^n \mathcal{E}$, the result follows from [LZ17, Prop. 2.3]. \square

Lemma 3.16. *For any $i \geq 0$, there exists a natural isomorphism*

$$\mathrm{H}^i(\Gamma_{\mathrm{geo}}, (\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C})(Y_{C, \infty})) \simeq \mathrm{H}^i(X_{\mathrm{pro\acute{e}t}}/Y_C, \mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}),$$

where $Y_{C, \infty}$ is the analogue of $X_{C, \infty}$ for Y replacing X (cf. 2.12).

Proof. Lemma 3.15 together with Čech-to-derived spectral sequence implies that $\mathrm{R}\Gamma(X_{\mathrm{pro\acute{e}t}}/Y_C, \mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C})$ can be computed by Čech–Alexander complex $(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C})(Y_{C, \infty}^{\bullet/Y_C})$, where $Y_{C, \infty}^{\bullet/Y_C}$ denotes the self product of $n + 1$ copies of $Y_{C, \infty}$ over Y_C . By a similar argument in the proof of Lemma 3.8, we see that

$$(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C})(Y_{C, \infty}^{\bullet/Y_C}) \cong C(\Gamma_{\mathrm{geo}}, (\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C})(Y_{C, \infty})).$$

Then the result follows. \square

Now, by Theorem 3.3 (1) and (5), we conclude that $H^0(X_{\text{proét}}/Y_C, \mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C})$ is a finite projective R_C -module such that (3.5) and (3.6) hold true. This shows that the functor

$$\text{Vect}(X, \widehat{\mathcal{O}}_X) \rightarrow \text{HIG}_{G_K}(X_C)$$

sending each generalised representation \mathcal{L} to

$$(\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})}) := (\nu_*(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}), \nu_*(\Theta_{\mathcal{L}}))$$

is a well-defined rank-preserving functor such that Theorem 3.3 (1) is true. Moreover, we also deduce that

Corollary 3.17. *Assume X admits a toric chart. Then the functor $\text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X) \rightarrow \text{HIG}_{G_K}(X_C)$ introduced in Corollary 3.13 is independent of the choice of toric chart.*

Proof. By Lemma 3.16, we see that $\mathcal{H}(\mathcal{L})(X_{C,\infty}) = H^0(\Gamma_{\text{geo}}, (\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C})(X_{C,\infty}))$. Then the result follows from the construction of the functor in Corollary 3.13. \square

Thanks to above corollary, we can show that the functor $\mathcal{L} \mapsto (\mathcal{H}(\mathcal{L}), \theta_{\mathcal{H}(\mathcal{L})})$ is indeed an equivalence. For this purpose, we construct its quasi-inverse as follows:

Let $\{X_i \rightarrow X\}_{i \in I}$ be an étale covering such that each $X_i \rightarrow X$ is standard étale. For any $(\mathcal{H}, \theta_{\mathcal{H}}) \in \text{HIG}_{G_K}(X_{C,\text{ét}})$, let $(\mathcal{H}_i, \theta_{\mathcal{H}_i})$ be its restriction to X_i . Then we get canonical comparison isomorphisms

$$\iota_{ij} : (\mathcal{H}_i, \theta_{\mathcal{H}_i})|_{X_i \times_X X_j} \xrightarrow{\cong} (\mathcal{H}_j, \theta_{\mathcal{H}_j})|_{X_i \times_X X_j}.$$

Let $\mathcal{L}_i \in \text{Vect}(X_{i,\text{proét}}, \widehat{\mathcal{O}}_{X_i})$ be the generalised representation on $X_{i,\text{proét}}$ corresponding to $(\mathcal{H}_i, \theta_{\mathcal{H}_i})$ in the sense of Corollary 3.13. By Corollary 3.17, ι_{ij} 's induce canonical isomorphisms of

$$\mathcal{L}_i|_{X_i \times_X X_j} \xrightarrow{\cong} \mathcal{L}_j|_{X_i \times_X X_j}.$$

So \mathcal{L}_i 's glue and achieve an $\mathcal{L}(\mathcal{H}, \theta_{\mathcal{H}}) \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$. It is easy to check that the functor $(\mathcal{H}, \theta_{\mathcal{H}}) \mapsto \mathcal{L}(\mathcal{H}, \theta_{\mathcal{H}})$ is the desired quasi-inverse. By Theorem 3.11 (3), the above constructions preserve tensor products and dualities.

To complete the proof of Theorem 3.3, it remains to verify items (2), (3) and (4).

For (2), let $\mathcal{L} \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ with associated $(\mathcal{H}, \theta_{\mathcal{H}}) \in \text{HIG}_{G_K}(X_C)$. By adjunction formula, we get a morphism

$$(\mathcal{H} \otimes_{\mathcal{O}_{X_C}} \mathcal{O}\mathbb{C}|_{X_C}, \Theta_{\mathcal{H}}) \rightarrow (\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{L})|_{X_C}, \Theta_{\mathcal{L}})$$

and have to check this is an isomorphism, which reduces to Theorem 3.11 (3) as the problem is local.

For (3), let $\mathcal{L} \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ and $(\mathcal{H}, \theta_{\mathcal{H}}) \in \text{HIG}_{G_K}(X_C)$ be as above. By noting that the universal Higgs field $(\mathcal{O}\mathbb{C}, \Theta)$ induces a quasi-isomorphism $\widehat{\mathcal{O}}_X \xrightarrow{\cong} \text{HIG}(\mathcal{O}\mathbb{C}, \Theta)$, we get a quasi-isomorphism

$$\mathcal{L} \simeq \text{HIG}(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}, \Theta_{\mathcal{L}})$$

and hence quasi-isomorphisms

$$\begin{aligned} \text{R}\Gamma(X_{C,\text{proét}}, \mathcal{L}) &\simeq \text{R}\Gamma(X_{C,\text{proét}}, \text{HIG}(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}, \Theta_{\mathcal{L}})) \\ &\simeq \text{R}\Gamma(X_{C,\text{ét}}, \text{R}\nu_*(\text{HIG}(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}, \Theta_{\mathcal{L}}))) \\ &\simeq \text{R}\Gamma(X_{C,\text{ét}}, \nu_*(\text{HIG}(\mathcal{L} \otimes_{\widehat{\mathcal{O}}_X} \mathcal{O}\mathbb{C}, \Theta_{\mathcal{L}}))) \quad \text{by (1)} \\ &\simeq \text{R}\Gamma(X_{C,\text{ét}}, \nu_*(\text{HIG}(\mathcal{H} \otimes_{\mathcal{O}_X} \mathcal{O}\mathbb{C}, \Theta_{\mathcal{H}}))) \quad \text{by (2)} \\ &\simeq \text{R}\Gamma(X_{C,\text{ét}}, \text{HIG}(\mathcal{H}, \theta_{\mathcal{H}})), \end{aligned}$$

where the last quasi-isomorphism is deduced by noting that $(\nu_*(\mathcal{O}\mathbb{C}), \nu_*(\Theta)) \cong (\mathcal{O}_X, 0)$.

For (4), let $\mathcal{L} \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ and $(\mathcal{H}, \theta_{\mathcal{H}}) \in \text{HIG}_{G_K}(X_C)$ be as above. Since there exists a natural morphism $(f^*\mathcal{O}\mathbb{C}_X, f^*\Theta_X) \rightarrow (\mathcal{O}\mathbb{C}_{X'}, \Theta_{X'})$ of Higgs fields (by taking graded pieces of [Sch13,

Prop. 8.5] after inverting t), we get isomorphisms

$$\begin{aligned}
& (f^*\mathcal{H} \otimes_{\mathcal{O}_{X'_C}} \mathcal{O}\mathbb{C}_{|X'_C}, \Theta_{f^*\mathcal{H}}) \\
& \cong (f^*\mathcal{H} \otimes_{\mathcal{O}_{X'_C}} \mathcal{O}\mathbb{C}_{|X'_C}, f^*\Theta_{\mathcal{H}}) \quad (\text{as } f^*\Theta_{\mathcal{H}} = \Theta_{f^*\mathcal{H}}) \\
& \rightarrow (f^*\mathcal{L} \otimes_{\widehat{\mathcal{O}}_{X'}}, \mathcal{O}\mathbb{C}_{|X'_C}, f^*\Theta_{\mathcal{L}}) \quad (\text{by (2) via base-change along } f^*\mathcal{O}\mathbb{C}_X \rightarrow \mathcal{O}\mathbb{C}_{X'}) \\
& \cong (f^*\mathcal{L} \otimes_{\widehat{\mathcal{O}}_{X'}}, \mathcal{O}\mathbb{C}_{|X'_C}, \Theta_{f^*\mathcal{L}}) \quad (\text{as } f^*\Theta_{\mathcal{L}} = \Theta_{f^*\mathcal{L}}).
\end{aligned}$$

By taking kernels of Higgs fields, we see that

$$f^*\mathcal{L} \cong \mathcal{L}(f^*\mathcal{H}, f^*\theta_{\mathcal{H}}).$$

By applying $\nu_{X',*} : X'_{\text{proét}} \rightarrow X'_{C,\text{ét}}$ to $(f^*\mathcal{H} \otimes_{\mathcal{O}_{X'_C}} \mathcal{O}\mathbb{C}_{|X'_C}, \Theta_{f^*\mathcal{H}}) \cong (f^*\mathcal{L} \otimes_{\widehat{\mathcal{O}}_{X'}}, \mathcal{O}\mathbb{C}_{|X'_C}, \Theta_{f^*\mathcal{L}})$, we get

$$(\mathcal{H}(f^*\mathcal{L}), \theta_{\mathcal{H}(f^*\mathcal{L})}) \cong (f^*\mathcal{H}, f^*\theta_{\mathcal{H}}).$$

Then Item (4) follows.

Therefore, we complete the proof of Theorem 3.3.

4. LOCAL DESCRIPTION OF HODGE–TATE CRYSTALS

In this section, we study the category of (rational) Hodge–Tate crystals on $(\mathfrak{X})_{\Delta}$ for $\mathfrak{X} = \text{Spf}(R^+)$ small affine with a fixed chart

$$\square := \mathcal{O}_K \langle T_1^{\pm 1}, \dots, T_d^{\pm 1} \rangle \rightarrow R^+$$

as defined in §1.4 and denote by $X = \text{Spa}(R, R^+)$ the rigid generic fibre of \mathfrak{X} . For the sake of simplicity, we denote the prismatic site $(\mathfrak{X})_{\Delta}$ by $(R^+)_{\Delta}$, denote by $(R^+)_{\Delta}^{\text{perf}}$ the sub-site of all perfect prisms, and denote by $(R^+ / (\mathfrak{S}, (E)))_{\Delta}$ the relative prismatic site with the base prism $(\mathfrak{S}, (E))$. Let R_C^+ be the base-change of R^+ along $\mathcal{O}_K \rightarrow \mathcal{O}_C$.

To state our main result, we make the following definition:

Definition 4.1. By an *enhanced Higgs module* of rank l over R^+ , we mean a triple (H, θ_H, ϕ_H) such that

- (1) H is a finite projective module over R^+ of rank l and $\theta_H : H \rightarrow H \otimes_{R^+} \widehat{\Omega}_{R^+/\mathcal{O}_K}^1 \{-1\}$ defines a nilpotent Higgs field on H . We denote by $\text{HIG}(H, \theta_H)$ the induced Higgs complex.
- (2) $\phi_H \in \text{End}_{R^+}(H)$ such that
 - (a) $\lim_{n \rightarrow +\infty} \prod_{i=0}^{n-1} (\phi_H + iE'(\pi)) = 0$ with respect to the p -adic topology on H , and that
 - (b) $[\phi_H, \theta_H] = -E'(\pi)\theta_H$; that is, ϕ_H induces an endomorphism of $\text{HIG}(H, \theta_H)$ as follows:

$$\begin{array}{ccc}
(4.1) \quad H & \xrightarrow{\theta_H} H \otimes_{R^+} \widehat{\Omega}_{R^+/\mathcal{O}_K}^1 \{-1\} \rightarrow \cdots \rightarrow H \otimes_{R^+} \widehat{\Omega}_{R^+/\mathcal{O}_K}^d \{-d\} \\
\phi_H \downarrow & \phi_H + E'(\pi)\text{id}_H \downarrow & \phi_H + dE'(\pi)\text{id}_H \downarrow \\
H & \xrightarrow{\theta_H} H \otimes_{R^+} \widehat{\Omega}_{R^+/\mathcal{O}_K}^1 \{-1\} \rightarrow \cdots \rightarrow H \otimes_{R^+} \widehat{\Omega}_{R^+/\mathcal{O}_K}^d \{-d\}.
\end{array}$$

We denote by $\text{HIG}(H, \theta_H, \phi_H)$ the total complex of (4.1). In other words, $\text{HIG}(H, \theta_H, \phi_H)$ is the fibre of $\text{HIG}(H, \theta_H)$:

$$\text{HIG}(H, \theta_H, \phi_H) = \text{fib}(\text{HIG}(H, \theta_H) \xrightarrow{\phi_H} \text{HIG}(H, \theta_H)).$$

We denote by $\text{HIG}_*^{\text{nil}}(R^+)$ the category of enhanced Higgs modules over R^+ . One can similarly define the category $\text{HIG}_*^{\text{nil}}(R)$ of enhanced Higgs modules over R by replacing R^+ and $\widehat{\Omega}_{R^+/\mathcal{O}_K}^i$ by R and $\Omega_{R/K}^i$ respectively in the above sentences.

Remark 4.2. In Definition 4.1, the nilpotency condition on θ_H is not necessary. In fact, if we write $\theta_H = \sum_{i=1}^d \theta_i \otimes \frac{d \log T_i}{E(u)}$, then the condition $[\phi_H, \theta_H] = -E'(\pi)\theta_H$ amounts to that for any $1 \leq i \leq d$, $[\phi_H, \theta_i] = -E'(\pi)\theta_i$. So the Lie sub-algebra \mathfrak{g} of $\text{End}_{R^+}(H)$ generated by ϕ_H and θ_i 's is solvable with $[\mathfrak{g}, \mathfrak{g}]$ generated by θ_i 's. Then the desired nilpotency follows from standard Lie theory (cf. [Hum72, §3.3 Cor. A]) via embedding R^+ into $\text{Frac}(R^+)$, the fractional field of R^+ .

In this subsection, we want to prove the following result.

Theorem 4.3. *The evaluation at the Breuil–Kisin prism $(\mathfrak{S}(R^+), (E))$ attached to the fixed framing \square induces an equivalence of categories*

$$\rho_{\square} : \text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}) \rightarrow \text{HIG}_{*}^{\text{nil}}(R^+) \text{ (resp. } \rho_{\square} : \text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \rightarrow \text{HIG}_{*}^{\text{nil}}(R)),$$

which preserve ranks, tensor products and dualities, such that for any Hodge–Tate crystal (resp. rational Hodge–Tate crystal) \mathbb{M} with associated enhanced Higgs module over R^+ (resp. R), there exists a quasi-isomorphism

$$\text{R}\Gamma((R^+)_{\Delta}, \mathbb{M}) \cong \text{HIG}(H, \theta_H, \phi_H),$$

which is functorial in \mathbb{M} .

Proof. This follows from Proposition 4.23 and Proposition 4.26 proved below. \square

Remark 4.4. Since $(\mathfrak{S}(R^+), (E))$ depends on the framing on R^+ , the equivalence $\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}) \rightarrow \text{HIG}_{*}^{\text{nil}}(R^+)$ also depends on the given framing. However, we will see in Corollary 6.15 that after inverting p , the equivalence $\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \rightarrow \text{HIG}_{*}^{\text{nil}}(R)$ is indeed independent of the choice of framing by using p -adic Simpson correspondence studied in the previous section. So there is a global theory on the rational level (cf. Theorem 6.4). When $\mathcal{O}_K = W(\kappa)$ is unramified, Theorem 4.3 was also obtained by Bhatt–Lurie in [BL22b] in a stacky way by regarding the prismaticization of \mathfrak{X} as a classifying space of some certain group scheme. However, up to now, it is still a problem to achieve a global theory as their method need a global “Frobenius endomorphism” of \mathfrak{X} . See [BL22b, §9] for details.

Remark 4.5. Note that there exists an obvious functor $\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}) \rightarrow \text{Vect}((R^+ / (\mathfrak{S}, (E)))_{\Delta}, \overline{\mathcal{O}}_{\Delta})$ induced by restriction along the inclusion $(R^+ / (\mathfrak{S}, (E)))_{\Delta} \subset (R^+)_{\Delta}$. By abuse of notations, for each $\mathbb{M} \in \text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta})$ (with associated enhanced Higgs module (H, θ_H, ϕ_H)), we also denote by \mathbb{M} its image under the above functor. Using Proposition 4.13 below, it is easy to see that the induced topologically nilpotent Higgs module associated to $\mathbb{M} \in \text{Vect}((R^+ / (\mathfrak{S}, (E)))_{\Delta}, \overline{\mathcal{O}}_{\Delta})$ via the equivalences in [Tian21, Thm. 4.10] is exactly (H, θ_H) and there exists a quasi-isomorphism

$$\text{R}\Gamma((R^+ / (\mathfrak{S}, (E)))_{\Delta}, \mathbb{M}) = \text{HIG}(H, \theta_H).$$

Remark 4.6. The equivalence $\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}) \rightarrow \text{HIG}_{*}^{\text{nil}}(R^+)$ in Theorem 4.3 still holds true in a more general setting: Indeed, one can define quasi-coherent Hodge–Tate crystal as a sheaf \mathbb{M} of $\overline{\mathcal{O}}_{\Delta}$ -modules on $(R)_{\Delta}$ with p -complete evaluations satisfying Convention 1.22 (2) for p -complete tensor products and define quasi-coherent enhanced Higgs modules as p -complete R -modules H with Higgs fields θ_H and R -linear endomorphism ϕ_H satisfying Definition 4.1 (2). Then the same proof of Theorem 4.3 also yields an equivalence between the category of quasi-coherent Hodge–Tate crystals and the category quasi-coherent enhanced Higgs modules by using (p -completely) faithfully flat descent for general (p -complete) modules (rather than finite projective ones).

We will prove Theorem 4.3 by establishing the desired equivalence and then comparing cohomologies separately in sequels.

4.1. Preliminaries.

Construction 4.7. Let $\mathfrak{S}\langle \underline{T}^{\pm 1} \rangle$ be the (p, E) -adic completion of $\mathfrak{S}[\underline{T}^{\pm 1}]$ which is endowed with a δ -structure over \mathfrak{S} such that $\delta(T_i) = 0$ for all i . Then $(\mathfrak{S}\langle \underline{T}^{\pm 1} \rangle, (E))$ is a prism. By étaleness of \square , it induces a smooth lifting $\mathfrak{S}(R^+)$ of R^+ over \mathfrak{S} and a (p, E) -adically étale morphism $\mathfrak{S}\langle \underline{T}^{\pm 1} \rangle \rightarrow \mathfrak{S}(R^+)$. By [BS22, Lem. 2.18], there exists a unique δ -structure on $\mathfrak{S}(R^+)$ which is compatible with the one on $\mathfrak{S}\langle \underline{T}^{\pm 1} \rangle$ and makes $(\mathfrak{S}(R^+), (E))$ a prism in $(R^+)_{\Delta}$. One can similarly define $(\mathbf{A}_{\text{inf}}\langle \underline{T}^{\pm 1} \rangle, (\xi))$ and $(\mathbf{A}_{\text{inf}}(R^+), (\xi))$ with δ -structures compatible with those on $(\mathfrak{S}\langle \underline{T}^{\pm 1} \rangle, (E))$ and $(\mathfrak{S}(R^+), (E))$ via the morphism $\mathfrak{S} \rightarrow \mathbf{A}_{\text{inf}}$ such that $\mathbf{A}_{\text{inf}}(R^+)$ is a lifting of R_C^+ over \mathbf{A}_{inf} . Let

$$\widehat{R}_{C, \infty}^+ := R_C^+ \widehat{\otimes}_{\mathcal{O}_C\langle T_1^{\pm 1}, \dots, T_d^{\pm 1} \rangle} \mathcal{O}_C\langle T_1^{\pm \frac{1}{p^{\infty}}}, \dots, T_d^{\pm \frac{1}{p^{\infty}}} \rangle.$$

It is a perfectoid ring over R with the induced perfect prism $(\mathbf{A}_{\text{inf}}(\widehat{R}_{C, \infty}^+), (\xi))$ in the sense of [BS22, Thm. 3.10], which is the perfection of $(\mathbf{A}_{\text{inf}}(R^+), (\xi))$. We also adapt notations in 2.12 and Notation 2.19.

Remark 4.8. Note that for any étale localisation $\iota : R^+ \rightarrow S^+$, there exists a unique morphism of prisms $(\mathfrak{S}(R^+), (E)) \rightarrow (\mathfrak{S}(S^+), (E))$ lifting ι , following from Lemma [BS22, Lem. 2.18]. Using this, we deduce from the constructions in this section that the equivalence and quasi-isomorphism in Theorem 4.3 are compatible with étale localisations. More precisely, let $\mathbb{M} \in \text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta})$ be a Hodge–Tate crystal with corresponding enhanced Higgs module (H, θ_H, ϕ_H) . Then the enhanced Higgs module associated to the restriction of \mathbb{M} to $(S^+)_{\Delta}$ is exactly $(H \otimes_{R^+} S^+, \theta_H \otimes \text{id}_{S^+}, \phi_H \otimes \text{id}_{S^+})$ and there exists a quasi-isomorphism

$$\text{RG}((S^+)_{\Delta}, \mathbb{M}) \cong \text{HIG}(H, \theta_H, \phi_H) \otimes_{R^+} S^+.$$

Lemma 4.9. (1) Both $(\mathfrak{S}(R^+), (E))$, $(\mathbf{A}_{\text{inf}}(R^+), (\xi))$ and $(\mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+), (\xi))$ are covers of the final object of the topos $\text{Shv}((R^+)_{\Delta})$. As a consequence, these prisms are also covers of the final object of $\text{Shv}((R^+ / (\mathfrak{S}(E)))_{\Delta})$.

(2) The $(\mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+), (\xi))$ is a cover of the final object of the topos $\text{Shv}((R^+)_{\Delta}^{\text{perf}})$.

Proof. Since there exist morphisms of prisms $(\mathfrak{S}(R^+), (E)) \rightarrow (\mathbf{A}_{\text{inf}}(R^+), (\xi)) \rightarrow (\mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+), (\xi))$. So it is enough to show $(\mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+), (\xi))$ cover the final objects of both $\text{Shv}((R^+)_{\Delta})$ and $\text{Shv}((R^+)_{\Delta}^{\text{perf}})$.

For the first topos, let (A, I) be any bounded prism in $(R^+)_{\Delta}$. Then $A/I \widehat{\otimes}_{R^+} \widehat{R}_{C,\infty}^+$ is a quasi-syntomic cover of A/I . By [BS22, Prop. 7.11 (1)], there exists a cover (B, IB) of (A, I) such that $A/I \rightarrow B/IB$ factors through $A/I \rightarrow A/I \widehat{\otimes}_{R^+} \widehat{R}_{C,\infty}^+$. In particular, B/I is an $\widehat{R}_{C,\infty}^+$ -algebra. Since $\widehat{R}_{C,\infty}^+$ is perfectoid, by deformation theory, we get a morphism $(\mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+), (\xi)) \rightarrow (B, IB)$ as desired.

For the second topos, we conclude from a similar argument as above by using [BS22, Prop. 7.11 (2)] instead of [BS22, Prop. 7.11 (1)]. \square

Notation 4.10. Let $(\mathfrak{S}(R^+)^{\bullet}, (E))$ (resp. $(\mathfrak{S}(R^+)_{\text{geo}}^{\bullet}, (E))$, resp. $(\mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+)^{\bullet}, (E))$) be the Čech nerve of $\mathfrak{S}(R^+)$ (resp. $\mathfrak{S}(R^+)_{\text{geo}}$, resp. $\mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+)$) in $(R^+)_{\Delta}$ (resp. $(R^+ / (\mathfrak{S}(E)))_{\Delta}$, resp. $(R^+)_{\Delta}^{\text{perf}}$). Here, E denotes $E(u_0)$; that is, the corresponding $E(u)$ of the first component in each degree of the Čech nerves.

Lemma 4.11. We have isomorphisms of cosimplicial rings (with obvious degeneracy morphisms p_i and face morphisms σ_i):

- (1) $\mathfrak{S}(R^+)^{\bullet} = \mathfrak{S}(R^+) \widehat{\otimes}^{(\bullet+1)} \left\{ \frac{u_0 - u_i}{E(u_0)}, \frac{T_{1,0} - T_{1,i}}{E(u_0)T_{1,0}}, \dots, \frac{T_{d,0} - T_{d,i}}{E(u_0)T_{d,0}} \mid 1 \leq i \leq \bullet \right\}_{\delta}^{\wedge}$, where for any $n \geq 0$, $\mathfrak{S}(R^+) \widehat{\otimes}^{(n+1)}$ denotes the (p, E) -complete tensor product of $n+1$ copies of $\mathfrak{S}(R^+)$ over $W(\kappa)$ and for any $0 \leq i \leq n$, $u_i, T_{1,i}, \dots, T_{d,i}$ denote the corresponding u, T_1, \dots, T_d of the $(i+1)$ -factor.
- (2) $\mathfrak{S}(R^+)_{\text{geo}}^{\bullet} = \mathfrak{S}(R^+) \widehat{\otimes}^{(\bullet+1)} \left\{ \frac{T_{1,0} - T_{1,i}}{E(u_0)T_{1,0}}, \dots, \frac{T_{d,0} - T_{d,i}}{E(u_0)T_{d,0}} \mid 1 \leq i \leq \bullet \right\}_{\delta}^{\wedge}$, where for any $n \geq 0$, $\mathfrak{S}(R^+) \widehat{\otimes}^{(\bullet+1)}$ denotes the (p, E) -complete tensor product of $n+1$ copies of $\mathfrak{S}(R^+)$ over \mathfrak{S} with $T_{1,i}$'s as above.
- (3) $\overline{\mathcal{O}}_{\Delta}[\frac{1}{p}](\mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+)^{\bullet}) = C(\Gamma(\overline{K}/K)^{\bullet}, \widehat{R}_{C,\infty})$, where for any $n \geq 0$, $C(\Gamma(\overline{K}/K)^{\bullet}, \widehat{R}_{C,\infty})$ denotes the ring of continuous functions from $\Gamma(\overline{K}/K)^n$ to $\widehat{R}_{C,\infty}^+$ (cf. §1.4).

Proof. (1) For any $n \geq 0$, $(\mathfrak{S}(R^+)^n, (E))$ is the initial object of the category of prisms in $(R)_{\Delta}$ which are the targets of $n+1$ arrows from $(\mathfrak{S}(R^+), (E))$. Note that for any prism (A, I) in the such category, there exists a unique morphism $\mathfrak{S}(R^+) \widehat{\otimes}^{(n+1)} \rightarrow A$ such that the reductions of u_i 's and $T_{s,i}$'s modulo I are π and T_s respectively for all $0 \leq i \leq n$ and any $1 \leq s \leq d$. So it suffices to show that $\mathfrak{S}(R^+) \widehat{\otimes}^{(n+1)} \left\{ \frac{u_0 - u_i}{E(u_0)}, \frac{T_{1,0} - T_{1,i}}{E(u_0)T_{1,0}}, \dots, \frac{T_{d,0} - T_{d,i}}{E(u_0)T_{d,0}} \mid 1 \leq i \leq \bullet \right\}_{(p,E)}^{\wedge}$ is well-defined, which follows from [BS22, Prop. 3.13].

(2) This follows from a similar argument used in (1). Also see [Tian21, §3.3].

(3) By virtues of [BS22, Thm. 3.10], for any $n \geq 0$, $\mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+)^n / (E)$ is the initial object of the category of perfectoid algebras over R^+ which are the targets of $n+1$ arrows from $\widehat{R}_{C,\infty}^+$. Note that, for any perfectoid ring S in the such category, there exists $n+1$ morphisms of perfectoid spaces $\text{Spa}(S[\frac{1}{p}], S[\frac{1}{p}]^{\circ})$ to $X_{C,\infty} = \text{Spa}(\widehat{R}_{C,\infty}, \widehat{R}_{C,\infty}^+)$ over $X = \text{Spa}(R, R^+)$. So we see that

$(\mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+)^n/(E))[\frac{1}{p}]$ is the ring of regular functions on the $(n+1)$ -folds self-product of $X_{C,\infty}$ over X , which turns out to be $C(\Gamma(\overline{K}/K)^n, \widehat{R}_{C,\infty})$ as shown in the proof of Lemma 3.8. \square

Remark 4.12. Keep notations as in the proof of 4.11 (3). Then one can show that $\mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+)^n/(E)$ is almost isomorphic to $C(\Gamma(\overline{K}/K)^n, \widehat{R}_{C,\infty}^+)$ with respect to \mathfrak{m}_C in the sense of [Sch12, §4]. More precisely, we can show that the kernel and cokernel of the canonical morphism

$$\iota_n : \mathbf{A}_{\text{inf}}(\widehat{R}_{C,\infty}^+)^n/(E) \rightarrow C(\Gamma(\overline{K}/K)^n, \widehat{R}_{C,\infty}^+)$$

are both killed by \mathfrak{m}_C . Indeed, since ι_n becomes isomorphic after inverting p , we can conclude by using the structure theorem of perfectoid rings (cf. [Bha, Prop. 3.2]).

Proposition 4.13. Let $X_i := \frac{u_0 - u_i}{E(u_0)}$ and $Y_{s,i} = \frac{T_{s,0} - T_{s,i}}{E(u_0)T_{s,0}}$ for any $1 \leq s \leq d$ and any $i \geq 1$. We identify u_0 and $T_{s,0}$'s of the reduction of first factor of $\mathfrak{S}(R)_{\text{geo}}^\bullet$ (resp. $\mathfrak{S}(R^+)^\bullet$) with T_s 's of $\mathfrak{S}(R^+)$.

(1) We have isomorphisms of cosimplicial rings:

$$\overline{\mathcal{O}}_{\Delta}(\mathfrak{S}(R^+)_{\text{geo}}^\bullet) \cong R^+\{Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^\wedge$$

and

$$\overline{\mathcal{O}}_{\Delta}[\frac{1}{p}](\mathfrak{S}(R^+)_{\text{geo}}^\bullet) \cong R\{Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^\wedge,$$

where $R^+\{Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^\wedge$ is the p -adic completion of the free polynomial ring over R^+ generated by the reductions of $Y_{s,i}$'s modulo E with $1 \leq i \leq \bullet$ and

$$R\{Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^\wedge = R^+\{Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^\wedge[\frac{1}{p}]$$

such that the face maps are given by

$$(4.2) \quad p_i(Y_{s,j}) = \begin{cases} Y_{s,j+1} - Y_{s,j}, & i = 0 \\ Y_{s,j+1}, & i \leq j \\ Y_{s,j}, & i > j; \end{cases}$$

while the degeneracy maps are given by

$$(4.3) \quad \sigma_i(Y_{s,j}) = \begin{cases} 0, & i = 0, j = 1 \\ Y_{s,j-1}, & i < j, (i, j) \neq (0, 1) \\ Y_{s,j}, & i \geq j. \end{cases}$$

(2) We have isomorphisms of cosimplicial rings:

$$\overline{\mathcal{O}}_{\Delta}(\mathfrak{S}(R^+)_{\text{geo}}^\bullet) \cong R^+\{X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^\wedge$$

and

$$\overline{\mathcal{O}}_{\Delta}[\frac{1}{p}](\mathfrak{S}(R^+)_{\text{geo}}^\bullet) \cong R\{X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^\wedge,$$

where $R^+\{X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^\wedge$ is the p -adic completion of the free polynomial ring over R^+ generated by the reductions of X_i 's and $Y_{s,i}$'s modulo E with $1 \leq i \leq \bullet$ and

$$R\{X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^\wedge = R^+\{X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^\wedge[\frac{1}{p}]$$

such that the face maps are given by

$$(4.4) \quad \begin{aligned} p_i(X_j) &= \begin{cases} (X_{j+1} - X_1)(1 - E'(\pi)X_1)^{-1}, & i = 0 \\ X_{j+1}, & i \leq j \\ X_j, & i > j; \end{cases} \\ p_i(Y_{s,j}) &= \begin{cases} (Y_{s,j+1} - Y_{s,1})(1 - E'(\pi)X_1)^{-1}, & i = 0 \\ Y_{s,j+1}, & i \leq j \\ Y_{s,j}, & i > j; \end{cases} \end{aligned}$$

while the degeneracy maps are given by

$$(4.5) \quad \begin{aligned} \sigma_i(X_j) &= \begin{cases} 0, & i = 0, j = 1 \\ X_{j-1}, & i < j, (i, j) \neq (0, 1) \\ X_j, & i \geq j; \end{cases} \\ \sigma_i(Y_{s,j}) &= \begin{cases} 0, & i = 0, j = 1 \\ Y_{s,j-1}, & i < j, (i, j) \neq (0, 1) \\ Y_{s,j}, & i \geq j. \end{cases} \end{aligned}$$

Proof. We only need to show the $\overline{\mathcal{O}}_{\Delta}$ case while the $\overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]$ case follows from inverting p .

(1) This is [Tian21, Cor. 4.5].

(2) For any $n \geq 0$, since $\delta(u_i) = 0$ for any $0 \leq i \leq n$, the desired isomorphism

$$\overline{\mathcal{O}}_{\Delta}(\mathfrak{S}(R^+)^\bullet) \cong R^+ \{X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^\wedge$$

follows from the same calculations in the proof of [Tian21, Prop. 4.3 and Cor. 4.5]. It remains to check that (4.4) and (4.5) hold true. Almost all formulae follow from the definitions of p_i 's and σ_i 's directly (cf. Convention 1.20) while the only difficulty appears in checking (4.4) for p_0 . We show the details as follows:

Since $p_0(u_i) = u_{i+1}$ and $p_0(T_{s,i}) = T_{s,i+1}$, we see that

$$p_0(X_j) = \frac{u_1 - u_{j+1}}{E(u_1)} = (X_{j+1} - X_1) \frac{E(u_0)}{E(u_1)}$$

and that

$$p_0(Y_{s,j}) = \frac{T_{s,1} - T_{s,j+1}}{E(u_1)T_{s,1}} = (Y_{s,j+1} - Y_{s,j}) \frac{E(u_0)T_{s,0}}{E(u_1)T_{s,1}}.$$

Then the desired formulae can be obtained by noting that

$$\frac{T_{s,0}}{T_{s,1}} = (1 - E(u_0)Y_{s,1}) \equiv 1 \pmod{E}$$

and that

$$(4.6) \quad \begin{aligned} \frac{E(u_0)}{E(u_1)} &= \left(1 - \frac{E(u_0) - E(u_1)}{E(u_0)}\right)^{-1} \\ &= \left(1 - \sum_{i \geq 1}^{\deg E} \frac{E^{(i)}(u_1)(u_0 - u_1)^i}{i!E(u_0)}\right)^{-1} \quad (\text{by Taylor's expansion}) \\ &\equiv (1 - E'(\pi)X_1)^{-1} \pmod{E} \quad (\text{as } u_0 - u_1 = E(u_0)X_1). \end{aligned}$$

□

Example 4.14. Let $R^+ = \mathcal{O}_K$ and hence $d = 0$ in Proposition 4.13. Let \mathfrak{S}^\bullet be the underlying cosimplicial ring of Čech nerve of $(\mathfrak{S}, (E)) \in (\mathcal{O}_K)_\Delta$. Then we have

$$\overline{\mathcal{O}}_{\Delta}(\mathfrak{S}^\bullet, (E)) = \mathcal{O}_K \{X_1, \dots, X_\bullet\}^\wedge$$

with face and degeneracy maps given by (4.4) and (4.5) (for X_i 's), respectively.

Corollary 4.15. Let \mathcal{K} (resp. \mathcal{K}_g , resp. \mathcal{K}_a) be the kernel of $\sigma_0 : \overline{\mathcal{O}}_{\Delta}(\mathfrak{S}(R^+)^\bullet, (E)) \rightarrow \overline{\mathcal{O}}_{\Delta}(\mathfrak{S}(R^+), (E))$ (resp. $\sigma_0 : \overline{\mathcal{O}}_{\Delta}(\mathfrak{S}(R^+)^\bullet_{\text{geo}}, (E)) \rightarrow \overline{\mathcal{O}}_{\Delta}(\mathfrak{S}(R^+), (E))$, resp. $\sigma_0 : \overline{\mathcal{O}}_{\Delta}(\mathfrak{S}^\bullet, (E)) \rightarrow \overline{\mathcal{O}}_{\Delta}(\mathfrak{S}, (E))$). Then

- (1) \mathcal{K} is the closed pd-ideal generated by $\{X_1^{[n_0]}Y_{1,1}^{[n_1]} \cdots Y_{d,1}^{[n_d]} \mid n_0 + \cdots + n_d \geq 1\}$. In particular, by letting $d = 0$, we see that \mathcal{K}_a is the closed pd-ideal generated by $\{X_1^{[n]} \mid n \geq 1\}$.
- (2) \mathcal{K}_g is the closed pd-ideal generated by $\{Y_{1,1}^{[n_1]} \cdots Y_{d,1}^{[n_d]} \mid n_1 + \cdots + n_d \geq 1\}$.
- (3) For any $r \geq 1$, let $\mathcal{K}^{[r]}$ (resp. $\mathcal{K}_g^{[r]}$, resp. $\mathcal{K}_a^{[r]}$) be the r -th closed pd-power of \mathcal{K} (resp. \mathcal{K}_g , resp. \mathcal{K}_a). Then the obvious morphisms $\mathfrak{S}^\bullet \rightarrow \mathfrak{S}(R^+)^\bullet \rightarrow \mathfrak{S}(R^+)^\bullet_{\text{geo}}$ (induce a morphism ideals $\mathcal{K}_a \rightarrow \mathcal{K} \rightarrow \mathcal{K}_g$ and then) induce a short exact sequence R^+ -modules:

$$(4.7) \quad 0 \rightarrow \mathcal{K}_a/\mathcal{K}_a^{[2]} \otimes_{\mathcal{O}_K} R^+ \rightarrow \mathcal{K}/\mathcal{K}^{[2]} \rightarrow \mathcal{K}_g/\mathcal{K}_g^{[2]} \rightarrow 0$$

Similar results hold true for replacing $\overline{\mathcal{O}}_\Delta$ by $\overline{\mathcal{O}}_\Delta[\frac{1}{p}]$ after inverting p .

Proof. It suffices to consider the $\overline{\mathcal{O}}_\Delta$ case as the $\overline{\mathcal{O}}_\Delta[\frac{1}{p}]$ case follows from the flat base-change “ $-\otimes_{\mathbb{Z}_p}$ ”.

(1) By Proposition 4.13 (2), especially (4.5), we see that the closed ideal \mathcal{J} generated by

$$\{X_1^{[n_0]}Y_{1,1}^{[n_1]} \cdots Y_{d,1}^{[n_d]} \mid n_0 + \cdots + n_d \geq 1\}$$

is contained in \mathcal{K} . So $\sigma_0 : \overline{\mathcal{O}}_\Delta(\mathfrak{S}(R^+)^1) \rightarrow \overline{\mathcal{O}}_\Delta(\mathfrak{S}(R^+)) = R^+$ factors through the quotient $\mathfrak{S}(R^+)^1/\mathcal{J}$. It is enough to show that $\mathfrak{S}(R^+)^1/\mathcal{J} \cong R^+$, which can be easily deduced as follows:

Let \mathcal{J}_0 be the ideal of the pd-polynomial ring $R^+[X_1, Y_{1,1}, \dots, Y_{d,1}]_{\text{pd}}$ over R^+ generated by $X_1, Y_{1,1}, \dots, Y_{d,1}$. By taking (p, E) -adic completion along the exact sequence

$$0 \rightarrow \mathcal{J}_0 \rightarrow R^+[X_1, Y_{1,1}, \dots, Y_{d,1}]_{\text{pd}} \rightarrow R^+ \rightarrow 0,$$

we get the desired isomorphism $\overline{\mathcal{O}}_\Delta(\mathfrak{S}(R^+)^1)/\mathcal{J} \cong R^+$ by noting that all rings involved are (p, E) -torsion free and R^+ is itself (p, E) -complete.

(2) This follows from the same argument used in the proof of (1).

(3) By (1) and (2), we see that for any $r \geq 1$, $\mathcal{K}^{[r]}$ (resp. $\mathcal{K}_g^{[r]}$, resp. $\mathcal{K}_a^{[r]}$) is the closed ideal generated by

$$\{X_1^{[n_0]}Y_{1,1}^{[n_1]} \cdots Y_{d,1}^{[n_d]} \mid n_0 + \cdots + n_d \geq r\} \text{ (resp. } \{Y_{1,1}^{[n_1]} \cdots Y_{d,1}^{[n_d]} \mid n_1 + \cdots + n_d \geq r\}, \text{ resp. } \{X_1^{[n]} \mid n \geq r\}).$$

So, we deduce that

$$\mathcal{K}/\mathcal{K}^{[2]} \cong R^+ \cdot X_1 \oplus \bigoplus_{i=1}^d R^+ \cdot Y_i \text{ (resp. } \mathcal{K}_g/\mathcal{K}_g^{[2]} \cong \bigoplus_{i=1}^d R^+ \cdot Y_i, \text{ resp. } \mathcal{K}_a/\mathcal{K}_a^{[2]} \cong \mathcal{O}_K \cdot X_1),$$

which implies the desired exact sequence immediately. \square

Remark 4.16. Consider $\overline{\mathcal{O}}_\Delta$ in Corollary 4.15. By identifying X_1 with $\frac{du}{E(u)}$, we see that

$$\mathcal{K}_a/\mathcal{K}_a^{[2]} \cong \widehat{\Omega}_{\mathfrak{S}/W(\kappa)}^1\{-1\} \otimes_{\mathfrak{S}} \mathcal{O}_K \cong \mathcal{O}_K \cdot \frac{du}{E(u)}.$$

By the definition of $Y_{s,i}$'s, we see that

$$\mathcal{K}_g/\mathcal{K}_g^{[2]} \cong \widehat{\Omega}_{R^+/\mathcal{O}_K}^1\{-1\}$$

by identifying $Y_{s,i}$ with $\frac{d \log T_i}{E(u)}$, where $\widehat{\Omega}_{R^+/\mathcal{O}_K}^1$ denotes the module of continuous differentials and $\{-1\}$ denotes the Breuil–Kisin twist. In particular, the exact sequence splits (non-canonically) and induces an isomorphism

$$(4.8) \quad \mathcal{K}/\mathcal{K}^{[2]} \cong \mathcal{K}_a/\mathcal{K}_a^{[2]} \otimes_{\mathcal{O}_K} R^+ \oplus \mathcal{K}_g/\mathcal{K}_g^{[2]} \cong R^+ \cdot \frac{du}{E(u)} \oplus \widehat{\Omega}_{R^+/\mathcal{O}_K}^1\{-1\}.$$

A similar remark applies to $\overline{\mathcal{O}}_\Delta[\frac{1}{p}]$.

4.2. Hodge–Tate crystals as enhanced Higgs modules. Now, we establish the desired equivalence in Theorem 4.3.

Convention 4.17. For simplicity, we put $a = E'(\pi)$.

Convention 4.18. Let $A^{?,\bullet}$ be the cosimplicial ring $R^?\{X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}$ for $? \in \{\emptyset, +\}$ with face and degeneracy morphisms given by (4.4) and (4.5), respectively. By Proposition 4.13, we have

$$A^\bullet \cong \overline{\mathcal{O}}_\Delta[\frac{1}{p}](\mathfrak{S}(R^+)^{\bullet}, (E))$$

and

$$A^{+,\bullet} \cong \overline{\mathcal{O}}_\Delta(\mathfrak{S}(R^+)^{\bullet}, (E)).$$

Let $\text{Strat}(A^{?,\bullet})$ denote the category of stratifications with respect to $A^{?,\bullet}$ satisfying the cocycle condition (cf. Convention 1.20). Note that $A^{?,0} = R^?$.

We start with the following lemma:

Lemma 4.19. *There exists a canonical equivalence from $\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta})$ (resp. $\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$) to $\text{Strat}(A^{?,\bullet})$ for $? = +$ (resp. $? = \emptyset$) by evaluating (rational) Hodge–Tate crystals along the cosimplicial prism $(\mathfrak{S}(R^+)^\bullet, (E))$.*

Proof. By Lemma 4.9, $(\mathfrak{S}(R^+), (E))$ is a cover of final object of $\text{Shv}((R^+)_{\Delta})$. So the result follows from [BS21, Prop. 2.7]. \square

Therefore, in order to establish the desired equivalence in Theorem 4.3, it suffices to construct an equivalence between the categories $\mathfrak{S}(A^{?,\bullet})$ and $\text{HIG}_*^{\text{nil}}(R^?)$.

Let $(H, \varepsilon) \in \text{Strat}(A^{?,\bullet})$ be any fixed stratification satisfying the cocycle condition. By Proposition 4.13, via the embedding $H \xrightarrow{x \mapsto x \otimes 1} H \otimes_{R^?, p_0} A^{?,1}$, there exists a collection $\{\phi_{i,\underline{n}}\}_{i \geq 0, \underline{n} \in \mathbb{N}^d}$ of $R^?$ -linear endomorphisms of H such that for any $x \in H$,

$$\varepsilon(x) = \sum_{i \geq 0, \underline{n} \in \mathbb{N}^d} \phi_{i,\underline{n}}(x) X_1^{[i]} \underline{Y}_1^{[\underline{n}]}.$$

Note that in order to make ε well-defined, we require

$$\lim_{i+|\underline{n}| \rightarrow +\infty} \phi_{i,\underline{n}} = 0$$

with respect to the p -adic topology on H . By (4.4) and (4.5), we see that

$$\begin{aligned} & (4.9) \\ & p_2^*(\varepsilon) \circ p_0^*(\varepsilon)(x) \\ &= p_2^*(\varepsilon) \left(\sum_{j \geq 0, \underline{n} \in \mathbb{N}^d} \phi_{j,\underline{n}}(x) (1 - aX_1)^{-j-|\underline{n}|} (X_2 - X_1)^{[j]} (\underline{Y}_2 - \underline{Y}_1)^{[\underline{n}]} \right) \\ &= \sum_{i,j \geq 0, \underline{l}, \underline{n} \in \mathbb{N}^d} \phi_{i,\underline{l}}(\phi_{j,\underline{n}}(x)) (1 - aX_1)^{-j-|\underline{n}|} X_1^{[i]} (X_2 - X_1)^{[j]} \underline{Y}_1^{[\underline{l}]} (\underline{Y}_2 - \underline{Y}_1)^{[\underline{n}]} \\ &= \sum_{i,j,k \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{i,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x)) (1 - aX_1)^{-j-k-|\underline{m}|-|\underline{n}|} (-1)^{j+|\underline{m}|} X_1^{[i]} X_1^{[j]} X_2^{[k]} \underline{Y}_1^{[\underline{l}]} \underline{Y}_1^{[\underline{m}]} \underline{Y}_2^{[\underline{n}]} \\ &= \sum_{i,j,k \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{i,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x)) (1 - aX_1)^{-j-k-|\underline{m}|-|\underline{n}|} (-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X_1^{[i+j]} \underline{Y}_1^{[\underline{l}+\underline{m}]} X_2^{[k]} \underline{Y}_2^{[\underline{n}]}, \end{aligned}$$

that

$$(4.10) \quad p_1^*(\varepsilon)(x) = \sum_{k \geq 0, \underline{n} \in \mathbb{N}^d} \phi_{k,\underline{n}}(x) X_2^{[k]} \underline{Y}_2^{[\underline{n}]},$$

and that

$$(4.11) \quad \sigma_0^*(\varepsilon)(x) = \phi_{0,\underline{0}}(x).$$

Therefore, we have

Lemma 4.20. *For any stratification (H, ε) with respect to $A^{?,\bullet}$, if we write*

$$\varepsilon(x) = \sum_{i \geq 0, \underline{n} \in \mathbb{N}^d} \phi_{i,\underline{n}}(x) X_1^{[i]} \underline{Y}_1^{[\underline{n}]}$$

for any $x \in H$ with $\phi_{i,\underline{n}} \rightarrow 0$ as $i + |\underline{n}| \rightarrow +\infty$, then (H, ε) satisfies the cocycle condition if and only if the following conditions are satisfied:

- (1) $\phi_{0,\underline{0}} = \text{id}_H$, and
- (2) For any $k \geq 0, \underline{n} \in \mathbb{N}^d$ and for any $x \in H$, we have

$$\phi_{k,\underline{n}}(x) = \sum_{i,j \geq 0, \underline{l}, \underline{m} \in \mathbb{N}^d} \phi_{i,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x)) (1 - aX)^{-j-k-|\underline{m}|-|\underline{n}|} (-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]}.$$

Notation 4.21. For any $(H, \varepsilon) \in \text{Strat}(A^{?,\bullet})$ with $\phi_{i,\underline{n}}$'s as above. Define $\phi_H = \phi_{1,\underline{0}}$. For any $1 \leq i \leq d$, put $\theta_i = \phi_{0,\underline{1}_i}$ and for $? = +$ (resp. \emptyset), define

$$\theta_H : H \rightarrow H \otimes_{R^+} \widehat{\Omega}_{R^+/O_K}^1 \{-1\} \text{ (resp. } \theta_H : H \rightarrow H \otimes_R \Omega_{R/K}^1 \{-1\})$$

by $\theta_H = \sum_{i=1}^d \theta_i \otimes \frac{\text{dlog} T_i}{E(u)}$.

The following lemma is the key ingredient to establish the desired equivalence.

Lemma 4.22. *Let $\phi_{i,\underline{n}}$'s be endomorphisms of H with $\phi_{0,\underline{0}} = \text{id}_H$. Let ϕ_H , θ_H and θ_i 's be as in Notation 4.21. Then the following assertions are equivalent:*

(1) *For any $k \geq 0$, $\underline{n} \in \mathbb{N}^d$ and for any $x \in H$, we have*

(4.12)

$$\phi_{k,\underline{n}}(x) = \sum_{i,j \geq 0, \underline{l}, \underline{m} \in \mathbb{N}^d} \phi_{i,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]}.$$

(2) *For any $1 \leq i, j \leq d$, we have $[\phi_H, \theta_i] = -a\theta_i$ and $[\theta_i, \theta_j] = 0$ such that for any $k \geq 0$ and $\underline{n} \in \mathbb{N}^d$,*

$$(4.13) \quad \phi_{k,\underline{n}}(x) = \prod_{s=1}^d \theta_i^{n_i} \cdot \prod_{i=0}^{k-1} (\phi_H + ia)(x) = \prod_{i=0}^{k-1} (\phi_H + (i + |\underline{n}|)a) \cdot \prod_{i=1}^d \theta_i^{n_i}(x).$$

Moreover, if one of the above conditions is satisfied, then θ_i 's are nilpotent and

$$\lim_{k+|\underline{n}| \rightarrow +\infty} \phi_{k,\underline{n}} = 0 \Leftrightarrow \lim_{k \rightarrow +\infty} \prod_{i=0}^{k-1} (\phi_H + ia) = 0.$$

Proof. First, we assume (1) holds true for any $k \geq 0$ and any $\underline{n} \in \mathbb{N}^d$ and then verify (2) as follows:

By letting $\underline{Y} = \underline{0}$ in both sides of (4.12), we get

$$\phi_{k,\underline{n}}(x) = \sum_{i,j \geq 0} \phi_{i,\underline{0}}(\phi_{j+k,\underline{n}}(x))(1-aX)^{-j-k-|\underline{n}|}(-1)^j \binom{i+j}{i} X^{[i+j]}.$$

Compare the coefficients of X in both sides of the above formula and then we get

$$(k + |\underline{n}|)a\phi_{k,\underline{n}}(x) - \phi_{1+k,\underline{n}}(x) + \phi_{1,\underline{0}}(\phi_{k,\underline{n}}(x)) = 0,$$

which implies by iteration that for any $k \geq 0$ and $\underline{n} \in \mathbb{N}^d$,

$$(4.14) \quad \phi_{k,\underline{n}}(x) = (\phi_{1,\underline{0}} + (k-1+|\underline{n}|)a)\phi_{k-1,\underline{n}}(x) = \prod_{i=0}^{k-1} (\phi_H + (i + |\underline{n}|)a)\phi_{0,\underline{n}}(x).$$

By letting $X = 0$ in both sides of (4.12), we get

$$\phi_{k,\underline{n}}(x) = \sum_{\underline{l}, \underline{m} \in \mathbb{N}^d} \phi_{0,\underline{l}}(\phi_{k,\underline{m}+\underline{n}}(x))(-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} \underline{Y}^{[\underline{l}+\underline{m}]}.$$

Compare the coefficients of Y_i in both sides of the above formula and then we get

$$\phi_{0,\underline{1}_i}(\phi_{k,\underline{n}}(x)) - \phi_{k,\underline{1}_i+\underline{n}}(x) = 0.$$

In particular, by letting $k = 0$ and $\underline{n} = \underline{1} - i + \underline{1}_j$ in the above formula, we see that

$$\theta_i \theta_j = \phi_{0,\underline{1}_i+\underline{1}_j} = \theta_j \theta_i.$$

In other words, θ_i 's commute with each other. Also, by iteration (for all $1 \leq i \leq d$) that for any $k \geq 0$ and $\underline{n} \in \mathbb{N}^d$,

$$(4.15) \quad \phi_{k,\underline{n}}(x) = \phi_{0,\underline{1}_i}(\phi_{k,\underline{n}-\underline{1}_i}(x)) = \prod_{i=1}^d \theta_i^{n_i}(\phi_{k,\underline{0}}(x)).$$

Now, (4.13) follows from (4.14) and (4.15). In particular, by letting $k = 1$ and $\underline{n} = \underline{1}_i$, we see that $[\phi_H, \theta_i] = -a\theta_i$ as desired.

Now assume (2) is true. By noting that the constant term of the right hand side of (4.12) is exactly $\phi_{k,\underline{n}}(x)$, we are reduced to showing that the right hand side of (4.12) is independent of X and \underline{Y} .

First, we show that the right hand side of (4.12) is independent of X . In other words, if we put

$$\begin{aligned}
(\dagger) &= \partial_X \left(\sum_{i,j \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{i,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x)) (1-aX)^{-j-k-|\underline{m}|-|\underline{n}|} (-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \right) \\
&= \sum_{i,j \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} (j+k+|\underline{m}|+|\underline{n}|) a \phi_{i,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x)) \\
&\quad \cdot (1-aX)^{-j-k-1-|\underline{m}|-|\underline{n}|} (-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad + \sum_{i,j \geq 0, i+j \geq 1, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{i,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x)) (1-aX)^{-j-k-|\underline{m}|-|\underline{n}|} (-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j-1]} \underline{Y}^{[\underline{l}+\underline{m}]},
\end{aligned}$$

then we have to show that $(\dagger) = 0$. Note that

$$\{(i, j) \mid i, j \geq 0, i+j \geq 1\} = \{(i, j) \mid i, j-1 \geq 0\} \cup \{(i, 0) \mid i \geq 1\}.$$

We have

$$\begin{aligned}
(\dagger) &= \sum_{i,j \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} (j+k+|\underline{m}|+|\underline{n}|) a \phi_{i,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x)) \\
&\quad \cdot (1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|} (-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad - \sum_{i,j \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{i,\underline{l}}(\phi_{j+1+k,\underline{m}+\underline{n}}(x)) (1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|} (-1)^{j+|\underline{m}|} \binom{i+j+1}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad + \sum_{i \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{k,\underline{m}+\underline{n}}(x)) (1-aX)^{-k-|\underline{m}|-|\underline{n}|} (-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i]} \underline{Y}^{[\underline{l}+\underline{m}]}.
\end{aligned}$$

By (4.13), we can use $\phi_{j+1+k,\underline{m}+\underline{n}}(x) - \phi_H(\phi_{j+k,\underline{m}+\underline{n}}(x))$ instead of $(j+k+|\underline{m}|+|\underline{n}|)a(\phi_{j+k,\underline{m}+\underline{n}}(x))$ in above formula and conclude that

$$\begin{aligned}
(\dagger) &= \sum_{i,j \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} -\phi_{i,\underline{l}}(\phi_H(\phi_{j+k,\underline{m}+\underline{n}}(x))) (1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|} (-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad - \sum_{i,j \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{i,\underline{l}}(\phi_{j+1+k,\underline{m}+\underline{n}}(x)) (1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|} (-1)^{j+|\underline{m}|} \binom{i+j}{i-1} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad + \sum_{i \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{k,\underline{m}+\underline{n}}(x)) (1-aX)^{-k-|\underline{m}|-|\underline{n}|} (-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i]} \underline{Y}^{[\underline{l}+\underline{m}]}.
\end{aligned}$$

By (4.13) again, we can use $\phi_{i+1,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x)) - ia\phi_{i,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x))$ instead of $\phi_{i,\underline{l}}(\phi_H(\phi_{j+k,\underline{m}+\underline{n}}(x)))$ in above formula and obtain that

$$\begin{aligned}
(\dagger) &= \sum_{i,j \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} ia\phi_{i,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad - \sum_{i,j \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad - \sum_{i,j \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i,\underline{l}}(\phi_{j+1+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+j}{i-1} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad + \sum_{i \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{k,\underline{m}+\underline{n}}(x))(1-aX)^{-k-|\underline{m}|-|\underline{n}|}(-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&= \sum_{i,j \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} aX\phi_{i,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i-1+j}{i-1} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i-1+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad - \sum_{i,j \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad - \sum_{i,j \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i,\underline{l}}(\phi_{j+1+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+j}{i-1} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad + \sum_{i \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{k,\underline{m}+\underline{n}}(x))(1-aX)^{-k-|\underline{m}|-|\underline{n}|}(-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i]} \underline{Y}^{[\underline{l}+\underline{m}]},
\end{aligned}$$

where we use $i \binom{i+j}{i} X^{[i+j]} = X \binom{i-1+j}{i-1} X^{[i-1+j]}$ to get the second equality. In the most right hand side of the above formula, replacing $i-1$ by i in the first and third summands and using $\binom{j-1}{-1} = \binom{j}{-1} = 0$, we see that

$$\begin{aligned}
(\dagger) &= \sum_{i,j \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} aX\phi_{i+1,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad - \sum_{i,j \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad - \sum_{i,j \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{j+1+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+1+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+1+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad + \sum_{i \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{k,\underline{m}+\underline{n}}(x))(1-aX)^{-k-|\underline{m}|-|\underline{n}|}(-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&= - \sum_{i,j \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{j+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad - \sum_{i,j \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{j+1+k,\underline{m}+\underline{n}}(x))(1-aX)^{-j-1-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+1+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+1+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&\quad + \sum_{i \geq 0, \underline{l}, \underline{m}, \in \mathbb{N}^d} \phi_{i+1,\underline{l}}(\phi_{k,\underline{m}+\underline{n}}(x))(1-aX)^{-k-|\underline{m}|-|\underline{n}|}(-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i]} \underline{Y}^{[\underline{l}+\underline{m}]}.
\end{aligned}$$

Finally, replacing $j + 1$ by j in the second term of the most right hand side of above formula, we see that

$$\begin{aligned}
(\dagger) &= - \sum_{i,j \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{i+1, \underline{l}}(\phi_{j+k, \underline{m}+\underline{n}}(x))(1-aX)^{-j-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&+ \sum_{i \geq 0, j \geq 1, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{i+1, \underline{l}}(\phi_{j+k, \underline{m}+\underline{n}}(x))(1-aX)^{-j-k-|\underline{m}|-|\underline{n}|}(-1)^{j+|\underline{m}|} \binom{i+j}{i} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i+j]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&+ \sum_{i \geq 0, \underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{i+1, \underline{l}}(\phi_{k, \underline{m}+\underline{n}}(x))(1-aX)^{-k-|\underline{m}|-|\underline{n}|}(-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} X^{[i]} \underline{Y}^{[\underline{l}+\underline{m}]} \\
&= 0
\end{aligned}$$

as desired, which shows that the right hand side of (4.12) is independent of X . In particular, to deduce (4.12), it is enough to show that for any $k \geq 0$ and any $\underline{n} \in \mathbb{N}^d$, we have

$$(4.16) \quad \phi_{k, \underline{n}}(x) = \sum_{\underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{0, \underline{l}}(\phi_{k, \underline{m}+\underline{n}}(x))(-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} \underline{Y}^{[\underline{l}+\underline{m}]}.$$

We consider a special case for $k = 0$ in (4.16). In other words, we want to show that for any $\underline{n} \in \mathbb{N}^d$,

$$(4.17) \quad \phi_{0, \underline{n}}(x) = \sum_{\underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{0, \underline{l}}(\phi_{0, \underline{m}+\underline{n}}(x))(-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} \underline{Y}^{[\underline{l}+\underline{m}]}.$$

To do so, it suffices to show that for free variables Z_1, \dots, Z_d ,

$$\sum_{\underline{n} \in \mathbb{N}^d} \phi_{0, \underline{n}}(x) \underline{Z}^{[\underline{n}]} = \sum_{\underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{0, \underline{l}}(\phi_{0, \underline{m}+\underline{n}}(x))(-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} \underline{Y}^{[\underline{l}+\underline{m}]} \underline{Z}^{[\underline{n}]}.$$

Since θ_i 's commute with each other and $\phi_{0, \underline{n}} = \prod_{i=1}^d \theta_i^{n_i}$ by (4.13), we see that

$$\sum_{\underline{n} \in \mathbb{N}^d} \phi_{0, \underline{n}}(x) \underline{Z}^{[\underline{n}]} = \exp\left(\sum_{i=1}^d \theta_i Z_i\right)(x)$$

and that

$$\begin{aligned}
&\sum_{\underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{0, \underline{l}}(\phi_{0, \underline{m}+\underline{n}}(x))(-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} \underline{Y}^{[\underline{l}+\underline{m}]} \underline{Z}^{[\underline{n}]} \\
&= \sum_{\underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \prod_{i=1}^d \theta_i^{l_i+m_i+n_i}(x) (-1)^{|\underline{m}|} \underline{Y}^{[\underline{l}]} \underline{Y}^{[\underline{m}]} \underline{Z}^{[\underline{n}]} \\
&= \sum_{\underline{l} \in \mathbb{N}^d} \prod_{i=1}^d \theta_i^{l_i} \underline{Y}^{[\underline{l}]} \sum_{\underline{m} \in \mathbb{N}^d} \prod_{i=1}^d \theta_i^{m_i} (-\underline{Y})^{[\underline{m}]} \sum_{\underline{n} \in \mathbb{N}^d} \prod_{i=1}^d \theta_i^{n_i}(x) \underline{Z}^{[\underline{n}]} \\
&= \exp\left(\sum_{i=1}^d \theta_i Y_i\right)(x) \exp\left(-\sum_{i=1}^d \theta_i Y_i\right) \exp\left(\sum_{i=1}^d \theta_i Z_i\right)(x) \\
&= \exp\left(\sum_{i=1}^d \theta_i Z_i\right)(x).
\end{aligned}$$

Therefore, the special case (4.17) holds true. In particular, replacing x by $\phi_{k, \underline{0}}(x)$ in (4.17), we get

$$(4.18) \quad \phi_{0, \underline{n}}(\phi_{k, \underline{0}}(x)) = \sum_{\underline{l}, \underline{m}, \underline{n} \in \mathbb{N}^d} \phi_{0, \underline{l}}(\phi_{0, \underline{m}+\underline{n}}(\phi_{k, \underline{0}}(x)))(-1)^{|\underline{m}|} \binom{\underline{l}+\underline{m}}{\underline{l}} \underline{Y}^{[\underline{l}+\underline{m}]}.$$

By using (4.13), especially $\phi_{0, \underline{n}}(\phi_{k, \underline{0}}(x)) = \phi_{k, \underline{n}}(x)$, we see that (4.18) is exactly (4.16) as desired. So we conclude that (2) implies (1).

To complete the proof, it remains to show the “moreover” part. The nilpotency of θ_i ’s follows from the same argument use in Remark 4.2. Since $\phi_{k,\underline{n}} = \phi_{0,\underline{n}} \circ \phi_{k,0}$ for any $(k, \underline{n}) \in \mathbb{N} \times \mathbb{N}^d$, we conclude from the nilpotency of θ_i ’s that

$$\lim_{k+|\underline{n}| \rightarrow +\infty} \phi_{k,\underline{n}} = 0 \Leftrightarrow \lim_{k \rightarrow +\infty} \prod_{i=0}^{k-1} (\phi_H + ia) = 0.$$

□

Proposition 4.23. *The evaluation at $(\mathfrak{S}(R^+), (E))$ induces equivalences*

$$\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}) \rightarrow \text{Strat}(A^+, \bullet) \rightarrow \text{HIG}_*^{\text{nil}}(R^+) \text{ (resp. } \text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \rightarrow \text{Strat}(A^{\bullet}) \rightarrow \text{HIG}_*^{\text{nil}}(R))$$

of categories, which preserves ranks, tensor products and dualities. Moreover, for any $\mathbb{M} \in \text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta})$ (resp. $\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$) with the corresponding stratification $(H, \varepsilon) \in \text{Strat}(A^?, \bullet)$ for $? = +$ (resp. \emptyset), let ϕ_H, θ_H and θ_i ’s be as in Notation 4.21 and then we have

- (1) *the enhanced Higgs module induced by \mathbb{M} is (H, θ_H, ϕ_H) , and*
- (2) *the stratification ε on H is determined such that for any $x \in H$,*

(4.19)

$$\begin{aligned} \varepsilon(x) &= \exp\left(\sum_{i=1}^d \theta_i Y_i\right) (1 - aX_1)^{-\frac{\phi_H}{a}}(x) := \sum_{k \geq 0, \underline{n} \in \mathbb{N}^d} \prod_{i=1}^d \theta_i^{n_i} \left(\prod_{j=0}^{k-1} (\phi_H + jE'(\pi))(x)\right) X_1^{[k]} \underline{Y}_1^{[\underline{n}]} \\ &= (1 - aX_1)^{-\frac{\phi_H}{a}} \prod_{i=1}^d \exp((1 - aX_1)^{-1} \theta_i Y_i)(x) := \sum_{k \geq 0, \underline{n} \in \mathbb{N}^d} \prod_{j=0}^{k-1} (\phi_H + (j + |\underline{n}|)E'(\pi)) \left(\prod_{i=1}^d \theta_i^{n_i}(x)\right) X_1^{[k]} \underline{Y}_1^{[\underline{n}]} \end{aligned}$$

Proof. The “moreover” part of Lemma 4.22 shows that (H, θ_H, ϕ_H) is an enhanced Higgs module over $R^?$. Then the proposition follows from Lemma 4.19 and Lemma 4.22 immediately as all constructions involved preserve ranks, tensor products and dualities. □

Remark 4.24. We give an intrinsic construction of (H, θ_H, ϕ_H) from a stratification $(H, \varepsilon) \in \text{Strat}(A^?, \bullet)$ for $? = +$ (resp. \emptyset) as follows:

Let $\mathcal{K}, \mathcal{K}_g$ and \mathcal{K}_a be as in Corollary 4.15 (and Remark 4.16). Noticing that $\sigma_0^*(\varepsilon) = \text{id}_H$, we deduce that for any $x \in H$,

$$\varepsilon(x) - \text{id}_H(x) \in H \otimes_{R^?, p_1} A^1 \cdot \mathcal{K}.$$

Modulo $\mathcal{K}^{[2]}$, we get an $R^?$ -linear morphism

$$\overline{(\varepsilon - \text{id}_H)} : H \rightarrow H \otimes_{R^?} \mathcal{K}/\mathcal{K}^{[2]}.$$

Let $\theta : H \rightarrow H \otimes_{R^+} \widehat{\Omega}_{R^+/\mathcal{O}_K}^1\{-1\}$ (resp. $\theta : H \rightarrow H \otimes_R \Omega_{R/K}^1\{-1\}$) be the $R^?$ -linear morphism induced by the projection $\mathcal{K}/\mathcal{K}^{[2]} \rightarrow \mathcal{K}_g/\mathcal{K}_g^{[2]}$ in Corollary 4.15 (3). Using the decomposition (4.8), we see that $\overline{(\varepsilon - \text{id}_H)}$ is of the form

$$(\phi, \theta) : H \rightarrow H \oplus H \otimes_{R^+} \widehat{\Omega}_{R^+/\mathcal{O}_K}^1\{-1\} \text{ (resp. } (\phi, \theta) : H \rightarrow H \oplus H \otimes_R \Omega_{R/K}^1\{-1\}).$$

By (4.19), we see that $\phi = \phi_H$ and $\theta = \theta_H$.

Example 4.25 (Breuil–Kisin Twist). Let $\overline{\mathcal{O}}_{\Delta}\{n\} := \mathcal{I}_{\Delta}^n/\mathcal{I}_{\Delta}^{n+1}$ for any $n \in \mathbb{Z}$, which is known as the n -th Breuil–Kisin twist of \mathcal{O}_{Δ} . Then we see that

$$\overline{\mathcal{O}}_{\Delta}\{n\}(\mathfrak{S}(R^+), (E)) = R^+ \cdot E(u)^n.$$

Note that by Taylor’s expansion, we have

$$p_0(E(u_0)^n) = E(u_1)^n = E(u_0 - E(u_0)X_1)^n \equiv (1 - E'(u_0)X_1)^n E(u_0)^n \pmod{E(u_0)^{n+1}}.$$

By chasing constructions above, we see that the enhanced Higgs bundle associated to $\overline{\mathcal{O}}_{\Delta}\{n\}$ is $(R^+, 0, -nE'(\pi)\text{id}_H)$. In general, for any (rational) Hodge–Tate crystal \mathbb{M} on $(R^+)_{\Delta}$ with the associated enhanced Higgs module (H, θ_H, ϕ_H) , the enhanced Higgs module induced by $\mathbb{M}\{n\} := \mathbb{M} \otimes_{\overline{\mathcal{O}}_{\Delta}} \overline{\mathcal{O}}_{\Delta}\{n\}$ is $(H, \theta_H, \phi_H - nE'(\pi)\text{id}_H)$.

4.3. Prismatic cohomology vs Higgs cohomology: Local case. Now, we want to show the following result:

Proposition 4.26. *For any (rational) Hodge–Tate crystal \mathbb{M} on $(R^+)_{\Delta}$ with associated enhanced Higgs module (H, θ_H, ϕ_H) , there exists a quasi-isomorphism*

$$R\Gamma((R^+)_{\Delta}, \mathbb{M}) \simeq \mathrm{HIG}(H, \theta_H, \phi_H).$$

Remark 4.27. Note that one can understand $R\Gamma((R^+/\langle \mathfrak{S}, (E) \rangle)_{\Delta})$ as a perfect complex of (rational) Hodge–Tate crystals on $(\mathcal{O}_K)_{\Delta}$ in the sense of [BL22a]. We may compute $R\Gamma((R^+)_{\Delta}, \mathbb{M})$ via “ $R\Gamma((\mathcal{O}_K)_{\Delta}, R\Gamma((R^+/\langle \mathfrak{S}, (E) \rangle)_{\Delta}, \mathbb{M}))$ ”. Recall that $R\Gamma((R^+/\langle \mathfrak{S}, (E) \rangle)_{\Delta}, \mathbb{M})$ can be computed via $\mathrm{HIG}(H, \theta_H)$ (cf. [Tian21, Thm. 4.10]) and that for any (rational) Hodge–Tate crystal \mathbb{M} over $(\mathcal{O}_K)_{\Delta}$ with the induced “enhanced Higgs module” (H, ϕ_H) , $R\Gamma((\mathcal{O}_K)_{\Delta}, \mathbb{M})$ can be computed by the complex $[H \xrightarrow{\phi_H} H]$ (cf. [MW21b, Thm. 3.20]). So it is reasonable for the existence of the desired quasi-isomorphism between $R\Gamma((R^+)_{\Delta}, \mathbb{M})$ and $\mathrm{HIG}(H, \theta_H, \phi_H)$. Indeed, we will show Proposition 4.26 by combining calculations in [Tian21] with [MW21b] together.

For the sake of simplicity, we still assume $a = E'(\pi)$ as in Convention 4.17. We first show that one can compute $R\Gamma((R^+)_{\Delta}, \mathbb{M})$ via Čech–Alexander method.

Lemma 4.28. *Let \mathfrak{X} be any bounded p -adic formal scheme and let \mathcal{C} be either $(\mathfrak{X})_{\Delta}$ or $(\mathfrak{X})_{\Delta}^{\mathrm{perf}}$. For any (rational) Hodge–Tate crystal \mathbb{M} on \mathcal{C} and for any $\mathcal{U} = (A, I) \in \mathcal{C}$, we have $H^i(\mathcal{C}/\mathcal{U}, \mathbb{M}) = 0$ for any $i \geq 1$.*

Proof. For any $\mathcal{V} = (B, IB) \in (R)_{\Delta}$ which is a cover of \mathcal{U} in \mathcal{C} , we denote by $(B^{\bullet}, IB^{\bullet})$ the Čech nerve induced by \mathcal{V} . Since \mathbb{M} is a (rational) Hodge–Tate crystal on \mathcal{C} , we have a canonical isomorphism of cosimplicial B^{\bullet} -modules

$$\mathbb{M}(B^{\bullet}, IB^{\bullet}) \cong \mathbb{M}(A, I) \otimes_{A/I} B^{\bullet}/IB^{\bullet}.$$

Since \mathcal{V} is a cover of \mathcal{U} , by (p, I) -adically faithfully flat descent, we see that $H^i(\mathbb{M}(B^{\bullet}, IB^{\bullet})) = 0$ for any $i \geq 0$. So the lemma follows from the Čech-to-derived spectral sequence. \square

Now, we still keep the assumption on the smallness of R^+ .

Lemma 4.29. *For any (rational) Hodge–Tate crystal \mathbb{M} on $(R^+)_{\Delta}$, there exists a quasi-isomorphism*

$$R\Gamma((R^+)_{\Delta}, \mathbb{M}) \cong \mathbb{M}(\mathfrak{S}(R^+)^{\bullet}, (E)).$$

Proof. This follows from Lemma 4.9 together with Lemma 4.28 by using Čech-to-derived spectral sequence. \square

Construction 4.30. Let \mathbb{M} be a (rational) Hodge–Tate crystal on $(R^+)_{\Delta}$ with induced enhanced Higgs complex (H, θ_H, ϕ_H) . Then there exists a canonical isomorphism

$$H \otimes_{R^+, q_0} A^{?, \bullet} \cong \mathbb{M}(\mathfrak{S}(R^+)^{\bullet}, (E)).$$

By virtues of Proposition 4.13 (2), we may write

$$H \otimes_{R^+, q_0} A^{?, \bullet} = H\{X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\mathrm{pd}}^{\wedge}.$$

$$B^{\bullet} = R^+\{Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\mathrm{pd}}^{\wedge}.$$

We regard $A^{?, \bullet}$ as a B^m -algebra via identifying $Y_{s,i}$ 's⁵. Let $\widehat{\Omega}_{B^m}^q$ be the module of continuous q -forms of B^m over \mathcal{O}_K . By Proposition 4.13, for any $n \geq 0$, we have

$$R^+\{X_i, Y_{s,j} \mid 1 \leq i \leq n, 1 \leq j \leq m\}_{\mathrm{pd}}^{\wedge} \widehat{\otimes}_{B^m} \widehat{\Omega}_{B^m}^q \cong \widehat{\Omega}_{R^+\{X_i, Y_{s,j} \mid 1 \leq i \leq n, 1 \leq j \leq m\}_{\mathrm{pd}}^{\wedge} / R^+\{X_i \mid 1 \leq i \leq n\}_{\mathrm{pd}}^{\wedge}}^q.$$

For any $m, n, q \geq 0$, define

$$C^{m, \bullet}\{-q\} := H\{X_i, Y_{1,j}, \dots, Y_{d,j} \mid 1 \leq i \leq n, 1 \leq j \leq m\}_{\mathrm{pd}}^{\wedge} \widehat{\otimes}_{A^{+, m}} \widehat{\Omega}_{A^{+, m}}^q\{-q\}.$$

We then define a structure of “Higgs complex of bicosimplicial objects” on $C^{\bullet, \bullet}\{-\bullet\}$ as follows:

⁵We warn that the induced morphism $B^{\bullet} \rightarrow A^{?, \bullet}$ does not preserve the cosimplicial structures.

- (1) For any $0 \leq i \leq n+1$, define $p_i^1 : C^{n,\bullet}\{-\bullet\} \rightarrow C^{n+1,\bullet}\{-\bullet\}$ such that it acts via $(1 - aX_1)^{-1}$ on $E(u)$ ⁶ for $i = 0$ and the identity for $i \geq 1$, acts via (4.4) on X_k 's, acts on $Y_{s,j}$'s via the scalar $(1 - aX_1)^{-1}$ for $i = 0$ and as the identity for $i \geq 1$, and acts on $x \in H$ via $(1 - aX_1)^{-\frac{\phi_H}{a}}$ for $i = 0$ and via id_H for $i \geq 1$.
- (2) For any $0 \leq i \leq n$, define $\sigma_i^1 : C^{n+1,\bullet}\{-\bullet\} \rightarrow C^{n,\bullet}\{-\bullet\}$ such that it acts trivially on $E(u)$, acts via (4.5) on X_i 's and acts on $Y_{s,j}$'s and $x \in H$ as the identity.
- (3) For any $0 \leq j \leq m+1$, define $p_j^2 : C^{\bullet,m}\{-\bullet\} \rightarrow C^{\bullet,m+1}\{-\bullet\}$ such that it acts trivially on $E(u)$, acts via the identity on X_i 's, via (4.2) on $Y_{s,k}$'s, and acts on $x \in H$ via $\exp(\sum_{i=1}^d \theta_i Y_{i,1})$ for $i = 0$ and via the identity for $i \geq 1$.
- (4) For any $0 \leq j \leq m$, define $\sigma_j^2 : C^{\bullet,m+1}\{-\bullet\} \rightarrow C^{\bullet,m}\{-\bullet\}$ such that it acts trivially on $E(u)$, X_i 's and $x \in H$, and acts via (4.3) on $Y_{s,k}$'s.
- (5) For any $q \geq 0$, define $d_q^{\bullet,\bullet} : C^{\bullet,\bullet}\{-q\} \rightarrow C^{\bullet,\bullet}\{-q-1\}$ such that for any $x \in H\{X_i, Y_{s,j} \mid 1 \leq i \leq \bullet, 1 \leq j \leq \bullet\}_{\text{pd}}^\wedge$ and $\omega \in \widehat{\Omega}_{B,\bullet}^q\{-q\}$, it carries $x \otimes \omega$ to $\theta_H(x) \wedge \omega + x \otimes d\omega$, where θ_H extends to $H\{X_i, Y_{s,j} \mid 1 \leq i \leq \bullet, 1 \leq j \leq \bullet\}_{\text{pd}}^\wedge$ by linear extension and $d : \widehat{\Omega}_{B,\bullet}^q\{-q\} \rightarrow \widehat{\Omega}_{B,\bullet}^{q+1}\{-q-1\}$ is defined in the paragraph after [Tian21, Lem. 4.13] (which is denoted by d_R in loc.cit.).

Then we have the following lemma:

Lemma 4.31. *The $(C^{\bullet,\bullet}\{-\bullet\}, p_i^1, \sigma_i^1, p_i^2, \sigma_i^2, d_q^{\bullet,\bullet})$ defines a complex of bicosimplicial $R^?$ -modules. In other words, for any $q \geq 0$, $(C^{\bullet,\bullet}\{-q\}, p_i^1, \sigma_i^1, p_i^2, \sigma_i^2,)$ is a bicosimplicial $R^?$ -module and $d_q^{\bullet,\bullet} : C^{\bullet,\bullet}\{-q\} \rightarrow C^{\bullet,\bullet}\{-q-1\}$ preserves bicosimplicial structures such that $d_{q+1}^{\bullet,\bullet} \circ d_q^{\bullet,\bullet} = 0$.*

Proof. By [Tian21, Lem. 4.14], we see that for any $n, q \geq 0$, $(C^{n,\bullet}\{-q\}, p_i^2, \sigma_i^2)$ defines a cosimplicial $R^?$ -module such that $\{d_q^{n,\bullet} : C^{n,\bullet}\{-q\} \rightarrow C^{n,\bullet}\{-q-1\}\}_{q \geq 0}$ is a complex of cosimplicial $R^?$ -modules.

Next, we show that for any $m, q \geq 0$, $(C^{\bullet,m}\{-q\}, p_i^1, \sigma_i^1)$ also defines a cosimplicial $R^?$ -module such that $\{d_q^{\bullet,m} : C^{\bullet,m}\{-q\} \rightarrow C^{\bullet,m}\{-q-1\}\}_{q \geq 0}$ is a complex of cosimplicial $R^?$ -modules. It is easy to check the first criterion. Using Example 4.25, we only need to check that $(C^{\bullet,m}\{-q\}, p_i^1, \sigma_i^1)$ defines a cosimplicial $R^?$ -module for $q = 0$. But in this case, $(C^{\bullet,m}\{-0\}, p_i^1, \sigma_i^1)$ is the cosimplicial $R^?$ -module induced by evaluating \mathbb{M} at the fibre product $(\mathfrak{S}(R^+)_{\text{geo}}^m, (E)) \times_{(\mathfrak{S}, (E))} (\mathfrak{S}^n, (E))$ ⁷ of $(\mathfrak{S}(R^+)_{\text{geo}}^m, (E))$ and $(\mathfrak{S}^n, (E))$ over the first factor of the latter, where $(\mathfrak{S}^n, (E))$ is defined in Example 4.14. To show the second criterion, we have to check that $d_q^{\bullet,m}$ preserves cosimplicial structures. We only show that it commutes with p_0^1 as the rest is much easier and can be deduced in a similar way. Now, for any $x \in H$ and $\omega \in \widehat{\Omega}_{R^+\{X_i, Y_{s,j} \mid 1 \leq i \leq n, 1 \leq j \leq m\}_{\text{pd}}^\wedge / R^+\{X_i \mid 1 \leq i \leq n\}_{\text{pd}}^\wedge}^q\{-q\}$, we see that

$$\begin{aligned} p_0^1(d_q^{n,m}(x \otimes \omega)) &= p_0^1\left(\sum_{i=1}^d \theta_i(x) \otimes \frac{d \log T_i}{E(u)} \wedge \omega + x \otimes d\omega\right) \\ &= \sum_{i=1}^d (1 - aX_1)^{-\frac{\phi_H}{a}-1} \theta_i(x) \otimes \frac{d \log T_i}{E(u)} \wedge p_0^1(\omega) + (1 - aX_1)^{-\frac{\phi_H}{a}}(x) \otimes p_0^1(d\omega), \end{aligned}$$

where the second equality follows as $p_0^1(E(u)) = (1 - aX_1)E(u)$ (cf. Example 4.25). On the other hand, we have

$$\begin{aligned} d_q^{n,m}(p_0^1(x \otimes \omega)) &= d_q^{n,m}((1 - aX_1)^{-\frac{\phi_H}{a}}(x) \otimes p_0^1(\omega)) \\ &= \sum_{i=1}^d \theta_i((1 - aX_1)^{-\frac{\phi_H}{a}}(x)) \otimes \frac{d \log T_i}{E(u)} \wedge p_0^1(\omega) + (1 - aX_1)^{-\frac{\phi_H}{a}}(x) \otimes dp_0^1(\omega). \end{aligned}$$

By noting that d commutes with p_0^1 , we are reducing to show that for any $1 \leq i \leq d$,

$$(1 - aX_1)^{-\frac{\phi_H}{a}-1} \theta_i(x) = \theta_i((1 - aX_1)^{-\frac{\phi_H}{a}}(x)).$$

But this follows from (4.13) directly.

⁶Here, we view $E(u)$ as a basis of the Breuil–Kisin twist $R^? \{1\}$. See Example 4.25.

⁷The existence of the fibre product $(\mathfrak{S}(R^+)_{\text{geo}}^m, (E)) \times_{(\mathfrak{S}, (E))} (\mathfrak{S}^n, (E))$ can be deduced by checking that its underlying δ -ring is $\mathfrak{S}(R^+)_{\text{geo}}^m \widehat{\otimes}_{\mathfrak{S}} \mathfrak{S}^n$.

It remains to show for each $q \geq 0$, $(C^{\bullet, \bullet}\{-q\}, p_i^1, \sigma_i^1, p_i^2, \sigma_i^2)$ is a bicosimplicial $R^?$ -module. In other words, we need to show p_i^1 's and σ_i^1 's commute with p_i^2 's and σ_i^2 's. We only show p_0^1 commutes with p_0^2 's as the rest can be checked similarly. For any $x \in H$, we have

$$\begin{aligned} p_0^1(p_0^2(x)) &= p_0^1(\exp(\sum_{i=1}^d \theta_i Y_{i,1})x) \\ &= (1 - aX_1)^{-\frac{\phi_H}{a}}(\exp(\sum_{i=1}^d (1 - aX_1)^{-1} \theta_i Y_{i,1})x) \\ &= \exp(\sum_{i=1}^d \theta_i Y_{i,1})((1 - aX_1)^{-\frac{\phi_H}{a}}(x)) \quad (\text{by (4.19)}) \\ &= p_0^2(p_0^1(x)). \end{aligned}$$

For any $k \geq 1$, we have that

$$p_0^1(p_0^2(X_k)) = p_0^1(X_k) = (X_{k+1} - X_1)(1 - aX_1)^{-1} = p_0^2(p_0^1(X_k)),$$

and that for any $1 \leq s \leq d$,

$$p_0^1(p_0^2(Y_{s,k})) = p_0^1(Y_{s,k+1} - Y_{s,1}) = (Y_{s,k+1} - Y_{s,1})(1 - aX_1)^{-1} = p_0^2(Y_{s,k}(1 - aX_1)^{-1}) = p_0^2(p_0^1(Y_{s,k})),$$

as desired. The proof is complete. \square

The $(C^{\bullet, \bullet}\{-\bullet\}, p_i^1, \sigma_i^1, p_i^2, \sigma_i^2)$ is related with $\mathbb{M}(\mathfrak{S}(R)^{\bullet}, (E))$ in the following sense:

Lemma 4.32. *Let $(C^{\bullet, \bullet}, p_i^1, \sigma_i^1, p_i^2, \sigma_i^2)$ denote the bisimplicial $R^?$ -module of the restriction of the complex $(C^{\bullet, \bullet}\{-\bullet\}, p_i^1, \sigma_i^1, p_i^2, \sigma_i^2, d_q^{\bullet, \bullet})$ at $q = 0$. Let $(D^{\bullet}, p_i^D, \sigma_i^D)$ be the diagonal of this bicomplex object (cf. [Wei04, §8.5]). Then we have an isomorphism of cosimplicial $R^?$ -modules $D^{\bullet} = \mathbb{M}(\mathfrak{S}(R^+), (E))$.*

Proof. We identify $\mathbb{M}(\mathfrak{S}(R^+), (E))$ with $H\{X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq \bullet\}_{\text{pd}}^{\wedge}$ as in Construction 4.30 and denote the induced face and degeneracy morphisms by p_i^H 's and σ_i^H 's. Then by definition of $C^{\bullet, \bullet}$, we see that for any $n \geq 0$,

$$D^n = C^{n,n} = H\{X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n\}_{\text{pd}}^{\wedge}.$$

For any $f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n)x \in D^n$, we have that

$$\begin{aligned} & p_0^D(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n)x) \\ &= p_0^1(p_0^2(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n)x)) \\ &= p_0^1(f(X_i, Y_{1,i+1} - Y_{1,1}, \dots, Y_{d,i+1} - Y_{d,1} \mid 1 \leq i \leq n) \exp(\sum_{k=1}^d \theta_k Y_k)x) \\ &= f((X_{i+1} - X_1)(1 - aX_1)^{-1}, (Y_{1,i+1} - Y_{1,1})(1 - aX_1)^{-1}, \dots, (Y_{d,i+1} - Y_{d,1})(1 - aX_1)^{-1} \mid 1 \leq i \leq n) \\ & \quad \cdot (1 - aX_1)^{-\frac{\phi_H}{a}} \exp(\sum_{k=1}^d (1 - aX_1)^{-1} \theta_k Y_k)x \\ &= p_0^H(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n)x) \quad (\text{by (4.19)}) \end{aligned}$$

and that for any $1 \leq k \leq n+1$,

$$\begin{aligned}
& p_k^D(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n)x) \\
&= p_k^1(p_k^2(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n)x)) \\
&= p_k^1(f(X_i, Y_{1,j}, \dots, Y_{d,j} \mid 1 \leq i \leq n, 1 \leq j \leq n+1, j \neq k) \exp(\sum_{l=1}^d \theta_l Y_{l,1})x) \\
&= f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n+1, i \neq k) \exp(\sum_{l=1}^d \theta_l Y_{l,1})x \\
&= p_k^H(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n)x).
\end{aligned}$$

Similarly, for any $f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n+1)x \in D^n$, we have that

$$\begin{aligned}
& \sigma_0^D(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n+1)x) \\
&= \sigma_0^1(\sigma_0^2(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n+1)x)) \\
&= \sigma_0^1(f(X_i, 0, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n)x) \\
&= f(0, X_i, 0, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n)x \\
&= \sigma_0^H(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n+1)x),
\end{aligned}$$

and that for any $1 \leq k \leq n$, after letting X'_i (resp. $Y'_{s,i}$) be X_i (resp. $Y_{s,i}$) for $1 \leq i \leq k$ and be X_{i-1} (resp. $Y_{s,i-1}$) for $k+1 \leq i \leq n+1$,

$$\begin{aligned}
& \sigma_k^D(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n+1)x) \\
&= \sigma_k^1(\sigma_k^2(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n+1)x)) \\
&= \sigma_0^1(f(X_i, Y'_{1,i}, \dots, Y'_{d,i} \mid 1 \leq i \leq n+1)x) \\
&= f(X'_i, Y'_{1,i}, \dots, Y'_{d,i} \mid 1 \leq i \leq n+1)x \\
&= \sigma_k^H(f(X_i, Y_{1,i}, \dots, Y_{d,i} \mid 1 \leq i \leq n+1)x).
\end{aligned}$$

These imply the desired isomorphism of simplicial $R^?$ -modules and hence complete the proof. \square

Corollary 4.33. *Let $\text{Tot}(C^\bullet)$ denote be the total complex induced by the bi-cosimplicial $R^?$ -module $C^{\bullet,\bullet}$. Then we have a quasi-isomorphism*

$$\mathbb{M}(\mathfrak{S}(R^+)^{\bullet}, (E)) \simeq \text{Tot}(C^{\bullet,\bullet}).$$

Proof. This is well-known as the cosimplicial Eilenberg–Zilber theorem and the shuffle coproduct formula (cf. [GM02, Appendix]). \square

Lemma 4.34. *Let $\text{Tot}(C^{\bullet,\bullet}\{-\bullet\})$ (resp. $\text{Tot}(C^{\bullet,0}\{-\bullet\})$) be the total complex of*

$$(C^{\bullet,\bullet}\{-\bullet\}, p_i^1, \sigma_i^1, p_i^2, \sigma_i^2, d_q^{\bullet,\bullet}) \text{ (resp. } (C^{\bullet,0}\{-\bullet\}, p_i^1, \sigma_i^1, d_q^{\bullet,0})).$$

Then the natural inclusions $C^{\bullet,\bullet} \hookrightarrow C^{\bullet,\bullet}\{-\bullet\} \hookleftarrow C^{\bullet,0}\{-\bullet\}$ induce quasi-isomorphisms

$$\text{Tot}(C^{\bullet,\bullet}) \simeq \text{Tot}(C^{\bullet,\bullet}\{-\bullet\}) \simeq \text{Tot}(C^{\bullet,0}\{-\bullet\}).$$

Proof. Let \mathfrak{S}^\bullet be as in Example 4.14. Note that [Tian21, Lem. 4.13] gives rise to a quasi-isomorphism

$$C^{0,0}\{-\bullet\} \simeq \text{Tot}(C^{0,\bullet}\{-\bullet\}).$$

By Construction 4.30, for any $n \geq 0$, we get an isomorphism of complexes of simplicial $R^?$ -morphisms

$$C^{0,\bullet}\{-\bullet\} \hat{\otimes}_{\mathcal{O}_K} \mathfrak{S}^n/(E) \cong C^{n,\bullet}\{-\bullet\}.$$

Define $T^{n,\bullet} := \text{Tot}(C^{n,\bullet}\{-\bullet\})$ for all n . Then using faithful flatness of $\mathcal{O}_K \rightarrow \mathfrak{S}^n/(E)$, we get quasi-isomorphisms

$$C^{n,0}\{-\bullet\} \simeq \text{Tot}(C^{n,\bullet}\{-\bullet\}) = T^{n,0}$$

for all n . By Lemma 4.31, we see that $\{C^{n,0} \simeq T^{n,0}\}_{n \geq 0}$ is compatible with p_i^1 's and σ_i^1 's. So we deduce that

$$\text{Tot}(C^{\bullet,0}\{-\bullet\}) \simeq \text{Tot}(T^{\bullet,\bullet}) = \text{Tot}(C^{\bullet,\bullet}\{-\bullet\})$$

as desired.

On the other hand, by [BdJ11, Lem. 2.17], for any $q \geq 1$, $C^{0,\bullet}\{-q\}$ is homotopic to zero and hence so is $C^{n,\bullet}\{-q\}$ for each $n \geq 0$. So we get a quasi-isomorphism

$$C^{n,\bullet} \simeq T^{n,\bullet}$$

for each n , which is compatible with p_i^1 's and σ_i^1 's, again following from Lemma 4.31. Therefore, we obtain a quasi-isomorphism

$$\mathrm{Tot}(C^{\bullet,\bullet}) \simeq \mathrm{Tot}(T^{\bullet,\bullet}) = \mathrm{Tot}(C^{\bullet,\bullet}\{-\bullet\})$$

as desired. The proof is complete. \square

Construction 4.35. Let $F(X, Y) := \frac{(1-aX)^{-\frac{Y}{a}} - 1}{X} = X + \sum_{n \geq 1} \prod_{i=1}^n (Y + ia) X^{[n+1]}$. Then for any $q \geq 0$, as $\lim_{n \rightarrow +\infty} \prod_{i=0}^{n-1} (\phi_H + ia) = 0$, we see that $F(X, \phi_H + qa)$ is well-defined such that the following diagram

$$\begin{array}{ccc} H \otimes_{R^+} \widehat{\Omega}_{R^+/\mathcal{O}_K}^1 \{-q\} & \xrightarrow{\phi_H + qa} & H \otimes_{R^+} \widehat{\Omega}_{R^+/\mathcal{O}_K}^1 \{-q\} \\ \parallel & & \downarrow F(X_1, \phi_H + qa) \\ C^{0,0} \{-q\} & \xrightarrow{p_0^1 - p_1^1} & C^{1,0} \{-q\} \end{array}$$

commutes and hence we get a morphism of complexes

$$\rho_q : [H \otimes_{R^+} \widehat{\Omega}_{R^+/\mathcal{O}_K}^1 \{-q\} \rightarrow H \otimes_{R^+} \widehat{\Omega}_{R^+/\mathcal{O}_K}^1 \{-q\}] \rightarrow C^{\bullet,0} \{-q\}.$$

Since $[\phi_H, \theta_H] = -a\theta_H$, we see that ρ_\bullet induces a morphism of complexes

$$\rho : \mathrm{HIG}(H, \theta_H, \phi_H) \rightarrow \mathrm{Tot}(C^{\bullet,0}\{-\bullet\}).$$

Lemma 4.36. *The morphism ρ in Construction 4.35 is a quasi-isomorphism.*

Proof. Use spectral sequence for bicomplexes, we are reduced to showing that each ρ_q is a quasi-isomorphism. But this follows from [MW21b, Thm. 3.20] (by replacing \mathcal{O}_K there by $R^?$). \square

Now, we are prepared to prove Proposition 4.26:

Proof of Proposition 4.26: This follows from quasi-isomorphisms

$$\begin{aligned} \mathrm{R}\Gamma((R^+)_{\Delta}, \mathbb{M}) &\simeq \mathbb{M}(\mathfrak{S}(R^+)^\bullet, (E)) \quad (\text{by Lemma 4.29}) \\ &\simeq \mathrm{Tot}(C^{\bullet,\bullet}) \quad (\text{by Corollary 4.33}) \\ &\simeq \mathrm{Tot}(C^{\bullet,0}\{-\bullet\}) \quad (\text{by Lemma 4.34}) \\ &\simeq \mathrm{HIG}(H, \theta_H, \phi_H) \quad (\text{by Lemma 4.36}). \end{aligned}$$

Then we can conclude by noting that all constructions above are functorial in \mathbb{M} . \square

5. HODGE–TATE CRYSTALS AS GENERALISED REPRESENTATIONS

In this section, we work with smooth p -adic formal schemes over \mathcal{O}_K with rigid analytic generic fiber X to prove the following result.

Theorem 5.1. *There exists a natural equivalence of categories*

$$L : \text{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \rightarrow \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X),$$

which preserves ranks, tensor products and dualities, such that for any rational Hodge–Tate crystal \mathbb{M} on $(\mathfrak{X})_{\Delta}^{\text{perf}}$, we have a quasi-isomorphism

$$\text{R}\Gamma((\mathfrak{X})_{\Delta}^{\text{perf}}, \mathbb{M}) \cong \text{R}\Gamma(X_{\text{proét}}, L(\mathbb{M})),$$

which is functorial in \mathbb{M} .

Remark 5.2. By [KL16, Thm 3.5.8 and Cor. 3.5.9], one may replace $X_{\text{proét}}$ by X_v in Theorem 5.1.

Now, we focus on the proof of Theorem 5.1. We first construct the desired equivalence. Let $X_{v,\text{aff,perf}}$ be the site of affinoid perfectoid spaces over X with the v -topology. By [KL16, Thm. 3.5.8], we have an equivalence of categories

$$\text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X) \simeq \text{Vect}(X_{v,\text{aff,perf}}, \mathcal{O}_X).$$

We then construct the desired functor L as follows.

Construction 5.3. For any $U = \text{Spa}(S, S^+) \in X_{v,\text{aff,perf}}$, we can assign it to a unique perfect prism $\mathcal{A}_U := (\mathbf{A}_{\text{inf}}(S^+), \text{Ker}(\theta : \mathbf{A}_{\text{inf}}(S^+) \rightarrow S^+)) \in (\mathfrak{X})_{\Delta}^{\text{perf}}$ by using [BS22, Thm. 3.10]. Via this functor, we have

$$\widehat{\mathcal{O}}_X(U) = S = \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}](\mathcal{A}_U),$$

and then get a natural functor

$$L : \text{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \rightarrow \text{Vect}(X_{v,\text{aff,perf}}, \mathcal{O}_X)$$

such that for any $\mathbb{M} \in \text{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$ and for any affinoid perfectoid space $U \in X_{\text{proét}}$, we have

$$L(\mathbb{M})(U) = \mathbb{M}(\mathcal{A}_U).$$

Note that by [KL16, Thm 3.5.8], the presheaf $L(\mathbb{M})$ is indeed a sheaf.

To see L is an equivalence, we have to construct the quasi-inverse L^{-1} of L .

Construction 5.4. For any $\mathcal{A} = (A, I) \in (\mathfrak{X})_{\Delta}^{\text{perf}}$, we can assign it to an affinoid perfectoid space $U_{\mathcal{A}}$ over X by setting

$$U_{\mathcal{A}} := \text{Spa}((A/I)[\frac{1}{p}], (A/I)[\frac{1}{p}]^+)$$

by using [BS22, Thm. 3.10] again, where $(A/I)[\frac{1}{p}]^+$ denotes the p -adic completion of the integral closure of A/I in $(A/I)[\frac{1}{p}]$. Via this functor, we have

$$\mathcal{O}_X(U_{\mathcal{A}}) = (A/I)[\frac{1}{p}] = \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}](\mathcal{A}),$$

and then get a natural functor

$$L^{-1} : \text{Vect}(X_{v,\text{aff,perf}}, \mathcal{O}_X) \rightarrow \text{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$$

such that for any $\mathcal{L} \in \text{Vect}(X_{v,\text{aff,perf}}, \mathcal{O}_X)$ and for any perfect prism $\mathcal{A} \in (\mathfrak{X})_{\Delta}^{\text{perf}}$, we have

$$L^{-1}(\mathcal{L})(\mathcal{A}) = \mathcal{L}(U_{\mathcal{A}}).$$

It is easy to see that L^{-1} is the quasi-inverse of L as desired.

Remark 5.5. A similar construction of L also appeared in the work of Morrow–Tsuji [MT20]. In loc.cit., they worked with smooth formal schemes \mathfrak{X} over \mathcal{O}_C and established an equivalence between the category of prismatic crystals and the category of relative Breuil–Kisin–Fargues modules on $X_{\text{proét}}$.

Remark 5.6. The functors $U \mapsto \mathcal{A}_U$ and $\mathcal{A} \mapsto U_{\mathcal{A}}$ in Construction 5.3 and Construction 5.4 are not the quasi-inverse of each other. In fact, we have almost isomorphisms (with respect to the ideal $\sqrt{(p)}$)

$$\mathcal{O}_X^+(U) \rightarrow \mathcal{O}_X^+(U_{\mathcal{A}_U})$$

and

$$\overline{\mathcal{O}}_{\Delta}(\mathcal{A}) \rightarrow \overline{\mathcal{O}}_{\Delta}(\mathcal{A}_{U_{\mathcal{A}}}).$$

In order to complete the proof of Theorem 5.1, it remains to show that

$$\mathrm{R}\Gamma((\mathfrak{X})_{\Delta}^{\text{perf}}, \mathbb{M}) \simeq \mathrm{R}\Gamma(X_{\text{proét}}, L(\mathbb{M})).$$

To do so, let us fix an étale covering $\{\mathfrak{X}_i = \mathrm{Spf}(R_i^+) \rightarrow \mathfrak{X}\}_{i \in I}$ of small affine \mathfrak{X}_i 's. Let $\mathcal{A}_i := (\mathbf{A}_{\text{inf}}(\widehat{R}_{i,C,\infty}^+), (\xi))$ be as in Construction 4.7 with $X_{i,C,\infty} = \mathrm{Spa}(\widehat{R}_{i,C,\infty}, \widehat{R}_{i,C,\infty}^+)$. Then $\{X_{i,C,\infty} \rightarrow X\}_{i \in I}$ is a pro-étale covering of X . We claim that $\{\mathcal{A}_i\}_{i \in I}$ forms a cover of the final object of $\mathrm{Shv}((\mathfrak{X})_{\Delta}^{\text{perf}})$.

Indeed, for any $(A, I) \in (\mathfrak{X})_{\Delta}^{\text{perf}}$, using the separatedness of \mathfrak{X} , we see that $\mathrm{Spf}(A/I) \times_{\mathfrak{X}} \mathfrak{X}_i$ is affine and thus denote the corresponding ring of regular functions by \overline{A}_i . Then $\mathrm{Spf}(\overline{A}_i) \rightarrow \mathrm{Spf}(A/I)$ is an étale morphism. In particular, by Elkik's algebraization theorem and [CS19, Corollary 2.1.6], \overline{A}_i is a perfectoid ring. By deformation theory, \overline{A}_i admits a unique lifting A_i over A . Then [BS22, Lem. 2.18] shows that there exists a unique δ -structure on A_i which is compatible with the one on A such that (A_i, IA_i) is a perfect prism in $(\mathfrak{X}_i)_{\Delta}^{\text{perf}}$. As there exist finitely many i 's such that $\{(A, I) \rightarrow (A_i, IA_i)\}$ forms a cover of (A, I) , the claim now follows from Lemma 4.9 (2).

Note that \mathcal{A}_i corresponds to $X_{i,C,\infty}$ via the functors in Construction 5.3 and Construction 5.4.

Lemma 5.7. *For any finite subset $J = \{j_1, \dots, j_r\} \subset I$, the fibre product $\mathcal{A}_J = \mathcal{A}_{j_1} \times \dots \times \mathcal{A}_{j_r}$ exists in $(\mathfrak{X})_{\Delta}^{\text{perf}}$ such that*

$$X_{j_1,C,\infty} \times_X \dots \times_X X_{j_r,C,\infty} = U_{\mathcal{A}_J}.$$

Proof. Note that \mathcal{A}_J should be the initial object of the category of prisms $\mathcal{B} = (B, J) \in (\mathfrak{X})_{\Delta}^{\text{perf}}$ together with morphisms $\mathcal{A}_j \rightarrow \mathcal{B}$ for all $j \in J$. For any such a prism \mathcal{B} , the structure morphism $\mathrm{Spf}(B/J) \rightarrow \mathfrak{X}$ then factors through $\mathfrak{X}_J \rightarrow \mathfrak{X}$, where $\mathfrak{X}_J = \mathrm{Spf}(R_J^+) = \mathfrak{X}_{j_1} \times_{\mathfrak{X}} \dots \times_{\mathfrak{X}} \mathfrak{X}_{j_r}$ (as \mathfrak{X} is separated). Put $\widehat{R}_{j,C,\infty,J}^+ := \widehat{R}_{j,C,\infty}^+ \widehat{\otimes}_{R_j^+} R_J^+$. Then it is perfectoid (which plays the role of “ $\widehat{R}_{C,\infty}^+$ ” for R_J^+ instead of R_j^+ with respect to the induced framing from R_i^+) and hence induces a perfect prism $\mathcal{A}_{j,J} := (A_{j,J}, IA_{j,J}) \in (R_J^+)_{\Delta}^{\text{perf}}$ by [BS22, Thm. 3.10]. By construction, the given morphism $\mathcal{A}_i \rightarrow \mathcal{B}$ has to factor over $\mathcal{A}_j \rightarrow \mathcal{A}_{j,J}$ for each $j \in J$. To conclude, it is enough to show the fibre product $\mathcal{A}_{j_1,J} \times \dots \times \mathcal{A}_{j_r,J}$ exists in $(R_J^+)_{\Delta}^{\text{perf}}$ as it is exactly \mathcal{A}_J as desired. Using [BS22, Thm. 3.10], it is enough to show that the category of perfectoid rings over

$$S^+ := \widehat{R}_{j_1,C,\infty,J}^+ \widehat{\otimes}_{R_j^+} \dots \widehat{\otimes}_{R_j^+} \widehat{R}_{j_r,C,\infty,J}^+$$

has an initial object S_J^+ , which follows from [BS22, Cor. 7.3] as S^+ is quasi-regular semi-perfectoid. So we conclude the existence of \mathcal{A}_J . Moreover, by constructions above, $U_{\mathcal{A}_J} = \mathrm{Spa}(S_J[\frac{1}{p}], S_J[\frac{1}{p}]^+)$ is the initial object of the category of perfectoid spaces Y over X which admits a fixed morphism $Y \rightarrow X_{J,C,\infty}$ for any $j \in J$. This shows that $U_{\mathcal{A}_J} = X_{j_1,C,\infty} \times_X \dots \times_X X_{j_r,C,\infty}$ as desired. \square

Thanks to Lemma 5.7 above, for any $\mathbb{M} \in \mathrm{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$, by Construction 5.3 and Construction 5.4, we have the same Čech complex

$$\check{C}^\bullet(\{\mathcal{A}_i\}_{i \in I}, \mathbb{M}) = \check{C}^\bullet(\{X_{i,C,\infty}\}_{i \in I}, L(\mathbb{M})).$$

By Lemma 4.28 and [LZ17, Prop. 2.3], using Čech-to-derived spectral sequence, we get a quasi-isomorphism

$$\mathrm{R}\Gamma((\mathfrak{X})_{\Delta}^{\mathrm{perf}}, \mathbb{M}) \simeq \mathrm{R}\Gamma(X_{\mathrm{pro\acute{e}t}}, \mathrm{L}(\mathbb{M}))$$

as desired, which completes the proof of Theorem 5.1.

Example 5.8. Assume $\mathfrak{X} = \mathrm{Spf}(R^+)$ is small with respect to the framing \square (cf. §1.4). Then one can give a direct proof of Theorem 5.1 as follows:

Keep notations in Construction 4.7 and let $X_{C,\infty} = \mathrm{Spa}(\widehat{R}_{C,\infty}, \widehat{R}_{C,\infty}^+)$. Then $X_{C,\infty} \in X_{\mathrm{pro\acute{e}t}}$ is affinoid perfectoid and is a Galois cover of X with Galois group $\Gamma(\overline{K}/K)$ (cf. Notation 2.19). Note that $\mathcal{A}_{X_{C,\infty}} = (\mathbf{A}_{\mathrm{inf}}(\widehat{R}_{C,\infty}^+), (\xi))$, which is a cover of the final object of $\mathrm{Shv}((\mathfrak{X})_{\Delta}^{\mathrm{perf}})$, following from Lemma 4.9 (2). Let $X_{C,\infty}^{\bullet+1/X}$ (resp. $\mathcal{A}_{X_{C,\infty}}^{\bullet+1}$) be the induced Čech nerve of $X_{C,\infty}$ (resp. $\mathcal{A}_{X_{C,\infty}}$). Then the proof of Lemma 3.8 combined with Lemma 4.11 (3) shows that

$$\widehat{\mathcal{O}}_X(X_{C,\infty}^{\bullet+1/X}) = \mathrm{C}(\Gamma(\overline{K}/K)^{\bullet}, \widehat{R}_{C,\infty}) = \overline{\mathcal{O}}_{\Delta}(\mathcal{A}_{X_{C,\infty}}^{\bullet+1}).$$

Note that [BS21, Prop. 2.7] also applies to $(\mathfrak{X})_{\Delta}^{\mathrm{perf}}$. We obtain a natural equivalence

$$\mathrm{Vect}((\mathfrak{X})_{\Delta}^{\mathrm{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \simeq \mathrm{Strat}(\mathrm{C}(\Gamma(\overline{K}/K), \widehat{R}_{C,\infty})) \simeq \mathrm{Rep}_{\Gamma(\overline{K}/K)}(\widehat{R}_{C,\infty}).$$

Then we show that L is an equivalence by applying Lemma 3.8. Clearly, the above construction preserves ranks, tensor products and dualities.

By Lemma 4.28 and Čech-to-derived spectral sequence, for any rational Hodge–Tate crystal \mathbb{M} on $(\mathfrak{X})_{\Delta}^{\mathrm{perf}}$ with induced $\widehat{R}_{C,\infty}^+$ -representation M of $\Gamma(\overline{K}/K)$, we have quasi-isomorphism

$$\iota_{\square} : \mathrm{R}\Gamma((\mathfrak{X})_{\Delta}^{\mathrm{perf}}, \mathbb{M}) \simeq \mathrm{R}\Gamma(\Gamma(\overline{K}/K), M) \simeq \mathrm{R}\Gamma(X_{\mathrm{pro\acute{e}t}}, \mathrm{L}(\mathbb{M})),$$

where the second quasi-isomorphism follows from Lemma 3.8 again. We complete the proof.

At the end of this example, we show that the quasi-isomorphism ι_{\square} is actually independent of the choice of the framing \square on \mathfrak{X} . Indeed, let \square_1 and \square_2 be two framings on \mathfrak{X} with corresponding $X_{C,\infty}^{\square_1}$ and $X_{C,\infty}^{\square_2}$, respectively. Let $U_{i,j}$ be the fibre products of $(i+1)$ copies of $X_{C,\infty}^{\square_1}$ and $(j+1)$ copies of $X_{C,\infty}^{\square_2}$ over X . Then each $U_{i,j}$ is affinoid perfectoid over X and then we get an isomorphism of bicomplexes R -modules

$$\mathbb{M}(\mathcal{A}_{U_{\bullet,\bullet}}) \cong \mathrm{L}(\mathbb{M})(U_{\bullet,\bullet}).$$

By noting that $U_{\bullet,0}$ (resp. $U_{0,\bullet}$) is the Čech nerve associated to $X_{C,\infty}^{\square_1}$ (resp. $X_{C,\infty}^{\square_2}$), we have a natural commutative diagram

$$\begin{array}{ccccc} \mathbb{M}(\mathcal{A}_{X_{C,\infty}^{\square_1}}^{\bullet+1/X}) & \longrightarrow & \mathbb{M}(\mathcal{A}_{U_{\bullet,\bullet}}) & \longleftarrow & \mathbb{M}(\mathcal{A}_{X_{C,\infty}^{\square_2}}^{\bullet+1/X}) \\ \downarrow & & \downarrow & & \downarrow \\ \mathrm{L}(\mathbb{M})(U_{\bullet,0}) & \longrightarrow & \mathrm{L}(\mathbb{M})(U_{\bullet,\bullet}) & \longleftarrow & \mathrm{L}(\mathbb{M})(U_{0,\bullet}). \end{array}$$

Thanks to [LZ17, Prop. 2.3] and Lemma 4.28, all horizontal arrows induce quasi-isomorphisms of corresponding total complexes and hence so is the middle vertical arrow (as the left and the right arrows induce ι_{\square_1} and ι_{\square_2} on totalizations). Therefore, we get a natural comparison between ι_{\square_1} and ι_{\square_2} .

6. INVERSE SIMPSON FUNCTOR FOR ENHANCED HIGGS BUNDLES

We still work with smooth p -adic formal schemes over \mathcal{O}_K in this section.

Notation 6.1. Let λ be the image of $\frac{\xi}{E([\pi^b])}$ under the surjection $\theta : \mathbf{A}_{\text{inf}} \rightarrow \mathcal{O}_C$. Clearly, λ belongs to $\mathcal{O}_{\widehat{K}_{\text{cyc}, \infty}}$ and is indeed a unit. For any $g \in G_K$, we see that

$$\begin{aligned}
 (6.1) \quad g(\lambda) &= \theta(g(\frac{[\epsilon] - 1}{E([\pi^b])([\epsilon^{\frac{1}{p}}] - 1)})) \\
 &= \theta(\frac{[\epsilon]^{\chi(g)} - 1}{[\epsilon] - 1} \frac{[\epsilon^{\frac{1}{p}}] - 1}{[\epsilon^{\frac{1}{p}}]^{\chi(g)} - 1} \frac{\xi}{E([\pi^b])} \frac{E([\pi^b])}{E([\epsilon]^{c(g)}[\pi^b])}) \\
 &= \chi(g) \frac{\zeta_p - 1}{\zeta_p^{\chi(g)} - 1} \lambda \theta(\frac{E([\pi^b])}{E([\pi^b]) + E'([\pi^b])[\pi^b]([\epsilon]^{c(g)} - 1)}) \\
 &= \chi(g) \frac{\zeta_p - 1}{\zeta_p^{\chi(g)} - 1} \lambda (1 + \pi E'(\pi) \theta(\frac{[\epsilon]^{c(g)} - 1}{[\epsilon] - 1} \frac{[\epsilon] - 1}{E([\pi^b])}))^{-1} \\
 &= \chi(g) \frac{\zeta_p - 1}{\zeta_p^{\chi(g)} - 1} \lambda (1 + \pi E'(\pi) (\zeta_p - 1) \lambda c(g))^{-1},
 \end{aligned}$$

where $c : G_K \rightarrow \mathbb{Z}_p$ is defined by $g(\pi^b) = \epsilon^{c(g)} \pi^b$ as in §1.4.

Definition 6.2. By an *enhanced Higgs bundle* of rank l on $\mathfrak{X}_{\text{ét}}$, we mean a triple $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$ consisting of

- (1) a locally finite free $\mathcal{O}_{\mathfrak{X}}$ -module \mathcal{H} of rank l on \mathfrak{X} together with a nilpotent Higgs field $\theta_{\mathcal{H}}$ on \mathcal{H} ; denote by $\text{HIG}(\mathcal{H}, \theta_{\mathcal{H}})$ the induced Higgs complex of $(\mathcal{H}, \theta_{\mathcal{H}})$;
- (2) an $\mathcal{O}_{\mathfrak{X}}[\frac{1}{p}]$ -linear endomorphism $\phi_{\mathcal{H}}$ of $\phi_{\mathcal{H}}$ satisfying
 - (a) $\lim_{n \rightarrow +\infty} \prod_{i=0}^{n-1} (\phi_{\mathcal{H}} + iE'(\pi)) = 0$ for the p -adic topology,
 - (b) $[\phi_{\mathcal{H}}, \theta_{\mathcal{H}}] = -E'(\pi)\theta_{\mathcal{H}}$; that is, $\phi_{\mathcal{H}}$ makes the following diagram

$$(6.2) \quad \begin{array}{ccccccc}
 \mathcal{H} & \xrightarrow{\theta_{\mathcal{H}}} & H \otimes_{\mathcal{O}_{\mathfrak{X}}} \widehat{\Omega}_{\mathfrak{X}}^1\{-1\} & \rightarrow \cdots \rightarrow & \mathcal{H} \otimes_{\mathcal{O}_{\mathfrak{X}}} \widehat{\Omega}_{\mathfrak{X}}^d\{-d\} \\
 \phi_{\mathcal{H}} \downarrow & & \phi_{\mathcal{H}} + E'(\pi)\text{id}_{\mathcal{H}} \downarrow & & \phi_{\mathcal{H}} + dE'(\pi)\text{id}_{\mathcal{H}} \downarrow \\
 \mathcal{H} & \xrightarrow{\theta_{\mathcal{H}}} & \mathcal{H} \otimes_{\mathcal{O}_{\mathfrak{X}}} \widehat{\Omega}_{\mathcal{O}_{\mathfrak{X}}}^1\{-1\} & \rightarrow \cdots \rightarrow & \mathcal{H} \otimes_{\mathcal{O}_{\mathfrak{X}}} \widehat{\Omega}_{\mathfrak{X}}^d\{-d\}
 \end{array}$$

commute. We denote by $\text{HIG}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$ the total complex of the bicomplex (6.2) and hence

$$\text{HIG}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) = \text{fib}(\text{HIG}(\mathcal{H}, \theta_{\mathcal{H}}) \xrightarrow{\phi_{\mathcal{H}}} \text{HIG}(\mathcal{H}, \theta_{\mathcal{H}})).$$

We denote by $\text{HIG}_{*}^{\text{nil}}(\mathfrak{X})$ the category of enhanced Higgs bundles on $\mathfrak{X}_{\text{ét}}$. One can similarly define the category $\text{HIG}_{*}^{\text{nil}}(X)$ of enhanced Higgs bundles on $X_{\text{ét}}$.

Remark 6.3. The nilpotency of $\theta_{\mathcal{H}}$ is not necessary in Definition (cf. Remark 4.2).

This section is devoted to proving the following theorem.

Theorem 6.4. *Assume \mathfrak{X} is a smooth p -adic formal scheme over \mathcal{O}_K with rigid analytic generic fiber X . Then there exists an equivalence of categories*

$$\rho : \text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \simeq \text{HIG}_{*}^{\text{nil}}(X),$$

which preserves ranks, tensor products and dualities. Moreover, this equivalence fits into the following commutative diagram of functors:

$$(6.3) \quad \begin{array}{ccc}
 \text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) & \xrightarrow{\rho} & \text{HIG}_{*}^{\text{nil}}(X) \\
 \downarrow \text{Res} & \swarrow F_S & \downarrow F \\
 \text{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) & \xrightarrow{\simeq} \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X) \xrightarrow{\simeq} & \text{HIG}_{G_K}(X_C),
 \end{array}$$

where Res is induced by inclusion of sites $(\mathfrak{X})_{\Delta}^{\text{perf}} \subset (\mathfrak{X})_{\Delta}$, F will be defined in Construction 6.13. All arrows in Diagram 6.3 are fully faithful functors while the bottom arrows are given in Theorem

5.1 and **Theorem 3.3**. In particular, we obtain the fully faithful inverse Simpson functor

$$F_S : \mathrm{HIG}_*^{\mathrm{nil}}(X) \rightarrow \mathrm{Vect}(X_{\mathrm{pro\acute{e}t}}, \widehat{\mathcal{O}}_X)$$

defined as the composite of the functor F and the equivalence $\mathrm{HIG}_{G_K}(X_C) \xrightarrow{\sim} \mathrm{Vect}(X_{\mathrm{pro\acute{e}t}}, \widehat{\mathcal{O}}_X)$.

The equivalence $\rho : \mathrm{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \simeq \mathrm{HIG}_*^{\mathrm{nil}}(X)$ in **Theorem 6.4** has been established when $\mathfrak{X} = \mathrm{Spf}(R^+)$ is small affine and upgrades to the integral level in this case (cf. **Theorem 4.3**). However, the constructions there depend on the choice of framing on R^+ and look difficult to glue together. The problem can be solved by showing that both F and Res are fully faithful making **Diagram (6.3)** commute locally. Then by some standard argument, the local equivalence $\rho_{\square} : \mathrm{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \simeq \mathrm{HIG}_*^{\mathrm{nil}}(R)$ in **Theorem 4.3** is independent of the choice of framing and hence glue to a global equivalence. We will follow this strategy in sequels.

6.1. Local version of Theorem 6.4 and the construction of F . In this subsection, we assume $\mathfrak{X} = \mathrm{Spf}(R^+)$ is small affine and keep notations in **Notation 2.19**. Then by **Example 5.8** and **Lemma 3.8**, we have equivalences of categories

$$\mathrm{Vect}((\mathfrak{X})_{\Delta}^{\mathrm{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \simeq \mathrm{Rep}_{\Gamma(\overline{K}/K)}(\widehat{R}_{C,\infty}) \simeq \mathrm{Vect}(X_{\mathrm{pro\acute{e}t}}, \widehat{\mathcal{O}}_X).$$

Recall we also have the equivalence

$$\rho_{\square} : \mathrm{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \simeq \mathrm{HIG}_*^{\mathrm{nil}}(R).$$

The purpose in this subsection is to give an explicit description of Res as a functor

$$V : \mathrm{HIG}_*^{\mathrm{nil}}(R) \rightarrow \mathrm{Rep}_{\Gamma(\overline{K}/K)}(\widehat{R}_{C,\infty}),$$

and then describe F by using the local Simpson correspondence i.e. **Theorem 3.11**.

Lemma 6.5. *For any $(H, \theta_H, \phi_H) \in \mathrm{HIG}_*^{\mathrm{nil}}(R)$ with induced $\widehat{R}_{C,\infty}$ -representation $V := V(H, \theta_H, \phi_H)$ of $\Gamma(\overline{K}/K)$, we have $V = H \otimes_R \widehat{R}_{C,\infty}$ such that for any $1 \leq i \leq d$, any $g \in G_K$ and any $x \in H$,*

$$\gamma_i(x) = \exp(-(\zeta_p - 1)\lambda\theta_i)(x) \text{ and } g(x) = (1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_H}{E'(\pi)}}(x).$$

Proof. Let A^\bullet be the cosimplicial ring in **Convention 4.18**. By **Lemma 4.19** and **Example 5.8**, it suffices to compare stratifications induced by (H, θ_H, ϕ_H) and V with respect to A^\bullet and $C(\Gamma(\overline{K}/K)^\bullet, \widehat{R}_{C,\infty})$, respectively. Note that the natural map

$$\mathrm{Strat}(A^\bullet) \rightarrow \mathrm{Strat}(C(\Gamma(\overline{K}/K)^\bullet, \widehat{R}_{C,\infty}))$$

is induced by the morphism of cosimplicial rings $\mathfrak{S}(R)^\bullet \rightarrow \mathbf{A}_{\mathrm{inf}}(\widehat{R}_{C,\infty}^+)^\bullet$. By **Lemma 4.11** (3) and **Proposition 4.13**, we have to determine the functions in $C(\Gamma(\overline{K}/K), \widehat{R}_{C,\infty})$ induced by $X_1, Y_{1,1}, \dots, Y_{d,1}$.

Lemma 6.6. *For any $\underline{n} = (n_1, \dots, n_d) \in \mathbb{N}^d$, any $1 \leq s \leq d$, and any $g \in G_K$, as functions in $C(\Gamma(\overline{K}/K), \widehat{R}_{C,\infty})$, we have*

$$X_1(\gamma_1^{n_1} \cdots \gamma_d^{n_d} g) = -\pi(\zeta_p - 1)\lambda c(g)$$

and that

$$Y_{s,1}(\gamma_1^{n_1} \cdots \gamma_d^{n_d} g) = -n_s(\zeta_p - 1)\lambda.$$

Proof. Recall $X_1 = \frac{u_0 - u_1}{E(u_0)}$ and $Y_{s,1} = \frac{T_{s,1} - T_{s,0}}{E(u_0)T_{s,0}}$. By the proof of **Lemma 4.11** (3), as functions in $C(\Gamma(\overline{K}/K), \mathbf{A}_{\mathrm{inf}}(\widehat{R}_{C,\infty}^+))$, u_0 (resp. $T_{s,0}$) is the constant function with the value $[\pi^b]$ and u_1 (resp. $T_{s,1}$) the evaluation function on $\Gamma(\overline{K}/K)$ at $[\pi^b]$ (resp. $[T_s^b]$), where $T_s^b = (T_s, T_s^{\frac{1}{p}}, \dots) \in \widehat{R}_{C,\infty}^{+b}$. Therefore, we conclude that

$$X_1(\gamma_1^{n_1} \cdots \gamma_d^{n_d} g) = \theta\left(\frac{[\pi^b](1 - [\epsilon]^{c(g)})}{E([\pi^b])}\right) = -\pi(\zeta_p - 1)\lambda c(g)$$

and that

$$Y_{s,1}(\gamma_1^{n_1} \cdots \gamma_d^{n_d} g) = \theta\left(\frac{[T_s^b](1 - [\epsilon]^{n_s})}{E([\pi^b])[T_s^b]}\right) = -n_s(\zeta_p - 1)\lambda.$$

as desired. \square

Now, we continue the proof of Proposition 6.5. By Proposition 4.23, especially (4.19), we deduced from Lemma 6.6 that for any $x \in H$,

$$(6.4) \quad \gamma_1^{n_1} \cdots \gamma_d^{n_d} g(x) = \exp\left(\sum_{i=1}^d -n_i(\zeta_p - 1)\lambda\theta_i\right)(1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_H}{E'(\pi)}}(x).$$

This completes the proof. \square

Remark 6.7. Since θ_H and ϕ_H are both defined over R , we see that $H_C := H \otimes_R R_C$ is itself stable under the action of $\Gamma(\overline{K}/K)$. Noting that γ_i acts on H_C unipotently, we see that H_C is the R_C -representation in $\text{Rep}_{\Gamma(\overline{K}/K)}^{\text{uni}}(R_C)$ associated to $V(H, \theta_H, \phi_H)$ under the equivalence in Proposition 2.16 (and Remark 2.20).

Now we can describe the functor F in the case for $\mathfrak{X} = \text{Spf}(R^+)$.

Construction 6.8. Define $F : \text{HIG}_*^{\text{nil}}(R) \rightarrow \text{HIG}_{G_K}(R_C)$ as the composition

$$\text{HIG}_*^{\text{nil}}(R) \xrightarrow{V} \text{Rep}_{\Gamma(\overline{K}/K)}(\widehat{R}_{C,\infty}) \xrightarrow{\text{Theorem 3.11}} \text{HIG}_{G_K}(R_C).$$

In particular, it makes the Diagram 6.3 commute.

Lemma 6.9. For any $(H, \theta_H, \phi_H) \in \text{HIG}_*^{\text{nil}}(R)$, we have

$$F(H, \theta_H, \phi_H) = (H \otimes_R R_C, -(\zeta_p - 1)\lambda\theta_H)$$

with the G_K -action such that for any $x \in H$ and any $g \in G_K$,

$$g(x) = (1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_H}{E'(\pi)}}(x).$$

Proof. This follows from Remark 3.12 and Remark 6.7 immediately. \square

Remark 6.10. Note that $\lambda \in \mathcal{O}_{\widehat{K}_{\text{cyc},\infty}}$. Thanks to Remark 2.20 and Remark 3.4, one can use \widehat{L} and $\text{Gal}(L/K)$ instead of C and G_K in Lemma 6.5, Remark 6.7 and Lemma 6.9 for any Galois extension L/K in \overline{K} containing $K_{\text{cyc},\infty}$.

Proposition 6.11. The functors F , V and Res are both fully faithful.

Proof. We only need to show V is fully faithful. Granting this, we get the full faithfulness of F by Theorem 3.11 and then the full faithfulness of Res as Diagram (6.3) commutes in this case. Moreover, by étale descent, we can further assume H is finite free over R . Thanks to Remark 6.10, we may work with $\widehat{K}_{\text{cyc},\infty}$ and $\Gamma(K_{\text{cyc},\infty}/K)$ instead of C and $\Gamma(\overline{K}/K)$ in the following arguments. For simplicity, we put $L = K_{\text{cyc},\infty}$.

Let $H_{\widehat{L}} \in \text{Rep}_{\Gamma(L/K)}(R_{\widehat{L}})$ be as in Remark 6.7. By Proposition 2.16, if we put $V = V(H, \theta_H, \phi_H)$, then we have a quasi-isomorphism

$$\text{R}\Gamma(\Gamma(L/K), H_{\widehat{L}}) \simeq \text{R}\Gamma(\Gamma(L/K), V).$$

We claim that $\Gamma(L/K) \cong \Gamma_{\text{geo}} \rtimes \text{Gal}(L/K)$ satisfied Axiom (1)-(4) of Tate–Sen theory for $R_{\widehat{L}}$ formulated in [Por22, §5.1]. Indeed, since Γ_{geo} acts on $R_{\widehat{L}}$ trivially, we only need to check $\text{Gal}(L/K)$ satisfies desired axioms. Note that R^+ is a topologically free \mathcal{O}_K -module by lifting κ -basis of $R^+/\pi R^+$ over κ . By equipping $R_{\widehat{L}}$ with the supreme norm induced by the corresponding \mathcal{O}_K -basis of R^+ , we are reduced to checking that the desired axioms are satisfied for \widehat{L} , which reduces to the example below [Por22, Cor. 5.3]. In particular, by [Por22, Cor. 5.3], we get quasi-isomorphisms

$$\text{R}\Gamma(\Gamma(L/K), H_{\widehat{L}}) \simeq \text{R}\Gamma(\Gamma(L/K), H_{\widehat{L}}^{\text{la}}) \simeq \text{R}\Gamma(\text{Lie}(\Gamma(L/K)), H_{\widehat{L}}^{\text{la}})^{\Gamma(L/K)},$$

where $H_{\widehat{L}}^{\text{la}}$ denotes the locally analytic vectors in $H_{\widehat{L}}$ with respect to the action of $\Gamma(L/K)$, which turns out to be $H \otimes_R R_{\widehat{L}}^{\text{la}}$ as $\Gamma(L/K)$ acts on H analytically (cf. (6.4)).

So we are reduced to showing that

$$(6.5) \quad H^{\phi_H=0, \theta_1=\dots=\theta_d=0} = H^0(\text{Lie}(\Gamma(L/K)), H \otimes_R R_{\widehat{L}}^{\text{la}})^{\Gamma(L/K)}.$$

Recall $\Gamma(L/K) \cong (\mathbb{Z}_p\gamma_1 \oplus \cdots \oplus \mathbb{Z}_p\gamma_d \oplus \mathbb{Z}_p\tau) \rtimes \Gamma(K_{\text{cyc}}/K)$ and let γ be a topological generator of $\text{Gal}(K_{\text{cyc}}/K(\zeta_p))$. Let ∇_i (resp. ∇_τ , resp. ∇_γ) be the operator in $\text{Lie}(\Gamma(L/K))$ corresponding to $\log\gamma_i$ (resp. $\log\tau$, resp. $\log\gamma$). Then we have

$$\text{Lie}(\Gamma(L/K)) \cong \bigoplus_{i=1}^d \mathbb{Q}_p \nabla_i \oplus \mathbb{Q}_p \nabla_\tau \oplus \mathbb{Q}_p \nabla_\gamma$$

such that ∇_i 's and ∇_τ commute with each other and $[\nabla_\gamma, \nabla_*] = \nabla_*$ for $*$ in $\{1, \dots, d, \tau\}$. Note that $\lambda \in \widehat{L}^{\text{la}} \subset R_{\widehat{L}}^{\text{la}}$ by (6.1). It is easy to see that

$$\nabla_\tau(\lambda) = -\pi E'(\pi)(\zeta_p - 1)\lambda^2 \text{ and } \nabla_\gamma(\lambda) = \lambda.$$

We claim that

$$H^{\phi_H=0, \theta_1=\dots=\theta_d=0} \subset H^0(\text{Lie}(\Gamma(L/K)), H \otimes_R R_{\widehat{L}}^{\text{la}})^{\Gamma(L/K)}.$$

Indeed, by (6.4), the claim is an immediate consequence of that for any $x \in H$,

$$-\pi\lambda(\zeta_p - 1)\phi_H(x) = \lim_{m \rightarrow +\infty} \frac{\tau^{p^m} - 1}{p^m}(x)$$

and

$$-\lambda(\zeta_p - 1)\theta_i(x) = \lim_{m \rightarrow +\infty} \frac{\gamma_i^{p^m} - 1}{p^m}(x).$$

Moreover, the above argument shows that for $x \in H$, we have

$$\begin{aligned} \nabla_i(x) &= -\lambda(\zeta_p - 1)\theta_i, \\ \nabla_\tau(x) &= -\pi\lambda(\zeta_p - 1)\phi_H; \\ \nabla_\gamma(x) &= 0. \end{aligned}$$

It remains to prove that

$$((H \otimes_R R_{\widehat{L}}^{\text{la}})^{\nabla_1=\dots=\nabla_d=0, \nabla_\tau=\nabla_\gamma=0})^{\Gamma(L/K)} \subset H^{\phi_H=0, \theta_1=\dots=\theta_d=0}.$$

For this, we have to apply some results in §7 (whose proofs certainly do not rely on this proposition).

Let ∇_{Sen} be the operator in the proof of Theorem 7.11. Then we have

$$((H \otimes_R R_{\widehat{L}}^{\text{la}})^{\nabla_1=\dots=\nabla_d=0, \nabla_\tau=\nabla_\gamma=0})^{\Gamma(L/K)} \subset ((H \otimes_R R_{\widehat{L}}^{\text{la}})^{\nabla_1=\dots=\nabla_d=0, \nabla_{\text{Sen}}=0})^{\Gamma(L/K)}.$$

The right hand side is equal to

$$\begin{aligned} ((H \otimes_R R_{\widehat{L}}^{\text{la}})^{\theta_1=\dots=\theta_d=0, \phi_H=0})^{\text{Gal}(L/K)} &= (((H \otimes_R R_{\widehat{L}}^{\text{la}})^{\theta_1=\dots=\theta_d=0, \phi_H=0})^{\text{Gal}(L/K_\infty)})^{\text{Gal}(L/K_{\text{cyc}})} \\ &= ((H \otimes_R R_{K_\infty})^{\theta_1=\dots=\theta_d=0, \phi_H=0})^{\text{Gal}(L/K_{\text{cyc}})} \\ &= (H^{\theta_1=\dots=\theta_d=0, \phi_H=0} \otimes_R R_{K_\infty})^{\text{Gal}(L/K_{\text{cyc}})} \end{aligned}$$

where the second equality is due to $R_{\widehat{L}}^{\text{la}, \text{Gal}(L/K_\infty)=1} = R_{K_\infty}$ by Proposition 7.8 and the third equality is due to that $R \rightarrow R_{K_\infty}$ is faithfully flat.

Write $(H^{\theta_1=\dots=\theta_d=0, \phi_H=0} \otimes_R R_{K_\infty})^{\text{Gal}(L/K_{\text{cyc}})}$ as $\bigcup_m (H^{\theta_1=\dots=\theta_d=0, \phi_H=0} \otimes_R R_{K(\pi_m)})^{\text{Gal}(K(\pi_m, \zeta_{p^m})/K(\zeta_{p^m}))}$.

For each m , we see that

$$(H^{\theta_1=\dots=\theta_d=0, \phi_H=0} \otimes_R R_{K(\pi_m, \zeta_{p^m})})^{\text{Gal}(K(\pi_m, \zeta_{p^m})/K(\zeta_{p^m}))} = H^{\theta_1=\dots=\theta_d=0, \phi_H=0} \otimes_R R_{K(\zeta_{p^m})}$$

by Galois descent. Then

$$(H^{\theta_1=\dots=\theta_d=0, \phi_H=0} \otimes_R R_{K(\pi_m)})^{\text{Gal}(K(\pi_m, \zeta_{p^m})/K(\zeta_{p^m}))} = H^{\theta_1=\dots=\theta_d=0, \phi_H=0} \otimes_R R_{K(\pi_m)} \cap H^{\theta_1=\dots=\theta_d=0, \phi_H=0} \otimes_R R_{K(\zeta_{p^m})}$$

which is exactly $H^{\theta_1=\dots=\theta_d=0, \phi_H=0}$. Hence we obtain that

$$((H \otimes_R R_{\widehat{L}}^{\text{la}})^{\nabla_1=\dots=\nabla_d=0, \nabla_{\text{Sen}}=0})^{\Gamma(L/K)} = H^{\theta_1=\dots=\theta_d=0, \phi_H=0}.$$

So we are done. \square

Remark 6.12. The idea of using the cohomology of Lie algebra to prove the above proposition is due to Hui Gao.

Construction 6.13. Assume \mathfrak{X} is a smooth p -adic formal scheme over \mathcal{O}_K with generic fiber X . For any enhanced Higgs bundle $(\mathcal{H}, \theta_H, \phi_H) \in \text{HIG}_*^{\text{nil}}(X)$, define $F(\mathcal{H}, \theta_H, \phi_H) \in \text{HIG}_{G_K}(X_C)$ as follows:

- (1) The underlying Higgs bundle on $X_{C, \text{ét}}$ is $(\mathcal{H} \otimes_{\mathcal{O}_{\mathfrak{X}}} \mathcal{O}_{X_C}, -(\zeta_p - 1)\lambda\theta_H)$.

(2) For any local section x of \mathcal{H} and for any $g \in G_K$, we have

$$(6.6) \quad g(x) = (1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(x).$$

Proposition 6.14. *The functor $F : \text{HIG}_*^{\text{nil}}(X) \rightarrow \text{HIG}_{G_K}(X_C)$ is a well-defined fully faithful functor, which preserves ranks, tensor products and dualities.*

Proof. We first show that F is well-defined. More precisely, we have to check that (6.6) induces a G_K -action on $\mathcal{H} \otimes_{\mathcal{O}_{\mathbb{X}}} \mathcal{O}_{X_C}$ and the Higgs field $-(\zeta_p - 1)\lambda\theta_H$ is G_K -equivariant.

For any $g_1, g_2 \in G_K$ and any local section x , we have

$$\begin{aligned} g_1(g_2(x)) &= g_1((1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g_2))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(x)) \\ &= (1 + \pi E'(\pi)(\zeta_p^{\chi(g_1)} - 1)g_1(\lambda)c(g_2))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(g_1(x)) \quad (\text{as } \phi_{\mathcal{H}} \text{ is defined over } \mathcal{H}) \\ &= (1 + \chi(g_1)c(g_2)\frac{\pi E'(\pi)(\zeta_p - 1)\lambda}{1 + E'(\pi)(\zeta_p - 1)\lambda c(g_1)})^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(g_1(x)) \quad (\text{by (6.1)}) \\ &= ((1 + \chi(g_1)c(g_2)\frac{\pi E'(\pi)(\zeta_p - 1)\lambda}{1 + E'(\pi)(\zeta_p - 1)\lambda c(g_1)})(1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g_1)))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(x) \\ &= (1 + \pi E'(\pi)(\zeta_p - 1)(\lambda c(g_1) + \chi(g_1)c(g_2)))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(x) \\ &= (1 + \pi E'(\pi)(\zeta_p - 1)c(g_1g_2))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(x) \\ &= (g_1g_2)(x). \end{aligned}$$

So (6.6) is a well-defined G_K -action on $F(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$.

It remains to check that $-(\zeta_p - 1)\lambda\theta_{\mathcal{H}}$ is G_K -equivariant. For any $g \in G_K$ and any local section x , if we write $\theta_{\mathcal{H}}(x) = \sum_{i=1}^d \theta_i(x) \otimes \frac{\text{dlog} T_i}{t}$, then we have

$$\begin{aligned} &g(-(\zeta_p - 1)\lambda\theta_{\mathcal{H}}(x)) \\ &= -(\zeta_p^{\chi(g)} - 1)g(\lambda)g(\sum_{i=1}^d \theta_i(x) \otimes \frac{\text{dlog} T_i}{t}) \\ &= -\chi(g)^{-1}(\zeta_p^{\chi(g)} - 1)g(\lambda) \sum_{i=1}^d g(\theta_i(x)) \otimes \frac{\text{dlog} T_i}{t} \\ &= -\chi(g)^{-1}(\zeta_p^{\chi(g)} - 1)g(\lambda) \sum_{i=1}^d (1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(\theta_i(x)) \otimes \frac{\text{dlog} T_i}{t} \\ &= -(\zeta_p - 1)\lambda(1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-1} \sum_{i=1}^d (1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(\theta_i(x)) \otimes \frac{\text{dlog} T_i}{t} \quad (\text{by (6.1)}) \\ &= -(\zeta_p - 1)\lambda \sum_{i=1}^d (1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}-1}(\theta_i(x)) \otimes \frac{\text{dlog} T_i}{t} \end{aligned}$$

and that

$$\begin{aligned} &-(\zeta_p - 1)\lambda\theta_{\mathcal{H}}(g(x)) \\ &= -(\zeta_p - 1)\lambda\theta_{\mathcal{H}}((1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(x)) \\ &= -(\zeta_p - 1)\lambda \sum_{i=1}^d \theta_i((1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(x)) \otimes \frac{\text{dlog} T_i}{t}. \end{aligned}$$

Since $[\phi_{\mathcal{H}}, \theta_{\mathcal{H}}] = -E'(\pi)\theta_{\mathcal{H}}$ (and hence $[\phi_{\mathcal{H}}, \theta_i] = -E'(\pi)\theta_i$ for all i), by (4.13), we see that

$$(1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}-1}(\theta_i(x)) = \theta_i((1 + \pi E'(\pi)(\zeta_p - 1)\lambda c(g))^{-\frac{\phi_{\mathcal{H}}}{E'(\pi)}}(x)),$$

which shows that $-(\zeta_p - 1)\lambda\theta_{\mathcal{H}}$ is compatible with G_K -action. So we conclude that F is well-defined. By its construction, we see that F preserves ranks, tensor products and dualities.

It remains show that F is fully faithful. It suffices to show that there is an isomorphism

$$H^0(X, \text{HIG}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})) \cong H^0(X_C, \text{HIG}(F(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})))^{G_K},$$

where $\text{HIG}(F(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})) = \text{HIG}(\mathcal{H} \otimes_{\mathcal{O}_{\mathfrak{X}}} \mathcal{O}_{X_C}, -(\zeta_p - 1)\lambda\theta_H)$ denotes the Higgs complex induced by $F(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$.

Let $\{\mathfrak{X}_i \rightarrow \mathfrak{X}\}_{i \in I}$ be a covering of \mathfrak{X} by small affines $\mathfrak{X}_i = \text{Spf}(R_i^+)$ and for any $i, j \in I$, let $\mathfrak{X}_{ij} = \mathfrak{X}_i \times_{\mathfrak{X}} \mathfrak{X}_j = \text{Spf}(R_{ij}^+)$. Let X_i be the generic fibers of \mathfrak{X}_i and $X_{i,C}$ be its base-change to C . Similarly define X_{ij} and $X_{ij,C}$. By noting that

$$H^0(X, \text{HIG}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})) = \text{Ker}\left(\prod_{i \in I} H^0(X_i, \text{HIG}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})|_{X_i}) \rightarrow \prod_{i,j \in I} H^0(X_{ij}, \text{HIG}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})|_{X_{ij}})\right)$$

and that

$$\begin{aligned} & H^0(X_C, \text{HIG}(F(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})))^{G_K} \\ &= \text{Ker}\left(\prod_{i \in I} H^0(X_{i,C}, \text{HIG}(F(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}))|_{X_{i,C}})^{G_K} \rightarrow \prod_{i,j \in I} H^0(X_{ij,C}, \text{HIG}(F(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}))|_{X_{ij,C}})^{G_K}\right), \end{aligned}$$

we are reduced to showing that for $* = i$ or ij ,

$$H^0(X_*, \text{HIG}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})|_{X_*}) \cong H^0(X_{*,C}, \text{HIG}(F(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}))|_{X_{*,C}})^{G_K}.$$

Then Proposition 6.11 applies (as each \mathfrak{X}_* is small affine). We are done. \square

Corollary 6.15. *Assume $\mathfrak{X} = \text{Spf}(R^+)$ is small affine. Then the equivalence*

$$\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \simeq \text{HIG}_*^{\text{nil}}(R)$$

constructed in Theorem 4.3 is independent of the framing on \mathfrak{X} .

Proof. Let ρ_1 and ρ_2 be equivalences from $\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$ to $\text{HIG}_*^{\text{nil}}(R)$ corresponding to the framings \square_1 and \square_2 on R^+ in the sense of Theorem 4.3. Note that in this case, Theorem 6.4 holds true and in particular, the diagram (6.3) commutes. So we see that

$$F \circ \rho_1 = F \circ \rho_2,$$

which coincides with the composition

$$\text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \xrightarrow{\text{Res}} \text{Vect}((R^+)_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \xrightarrow{\sim} \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X) \xrightarrow{\sim} \text{HIG}_{G_K}(X_C),$$

where $X = \text{Spa}(R, R^+)$ denotes the generic fiber of \mathfrak{X} . We denote this composition by T . It is clearly independent of the choice of framings on R^+ .

By Proposition 6.11, both F and T are fully faithful. Therefore, for any $\mathbb{M} \in \text{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$, we have

$$\text{Hom}(\mathbb{M}, \mathbb{M}) = \text{Hom}(T(\mathbb{M}), T(\mathbb{M})) = \text{Hom}(F(\rho_1(\mathbb{M})), F(\rho_2(\mathbb{M}))) = \text{Hom}(\rho_1(\mathbb{M}), \rho_2(\mathbb{M})).$$

So the identity $\text{id}_{\mathbb{M}}$ provides a canonical isomorphism between $\rho_1(\mathbb{M})$ and $\rho_2(\mathbb{M})$. We win. \square

6.2. Proof of Theorem 6.4. Now, we are going to prove Theorem 6.4 by noting that both $\text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$ and $\text{HIG}_*^{\text{nil}}(X)$ are indeed étale stacks.

We fix a covering $\{\mathfrak{X}_i \rightarrow \mathfrak{X}\}_{i \in I}$ of \mathfrak{X} by small affine $\mathfrak{X}_i = \text{Spf}(R_i^+)$ and let $\mathfrak{X}_{ij} = \mathfrak{X}_i \times_{\mathfrak{X}} \mathfrak{X}_j = \text{Spf}(R_{ij}^+)$ for all $i, j \in I$. For any $\mathbb{M} \in \text{Vect}((\mathfrak{X}_i)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$, denote by \mathbb{M}_i its restriction to $(\mathfrak{X}_i)_{\Delta}$ for any $i \in I$. Then we get canonical isomorphisms $\iota_{ij} : \mathbb{M}_i|_{(\mathfrak{X}_{ij})_{\Delta}} \xrightarrow{\cong} \mathbb{M}_j|_{(\mathfrak{X}_{ij})_{\Delta}}$ satisfying the cocycle condition.

Now applying Theorem 4.3, each \mathbb{M}_i induces an enhanced Higgs bundle $(\mathcal{H}_i, \theta_{\mathcal{H}_i}, \phi_{\mathcal{H}_i})$ in $\text{HIG}_*^{\text{nil}}(X_i)$. By Corollary 6.15, the isomorphisms ι_{ij} 's induce isomorphisms

$$\rho_{ij} : (\mathcal{H}_i, \theta_{\mathcal{H}_i}, \phi_{\mathcal{H}_i})|_{X_{ij}} \xrightarrow{\cong} (\mathcal{H}_j, \theta_{\mathcal{H}_j}, \phi_{\mathcal{H}_j})|_{X_{ij}}$$

satisfying the cocycle condition. Therefore, we get an enhanced Higgs bundle $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) \in \text{HIG}_*^{\text{nil}}(X)$ whose restriction to X_i coincides with $(\mathcal{H}_i, \theta_{\mathcal{H}_i}, \phi_{\mathcal{H}_i})$. This induces a functor

$$\rho : \text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \rightarrow \text{HIG}_*^{\text{nil}}(X).$$

Conversely, we can similarly assign each enhanced Higgs bundle $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) \in \mathrm{HIG}_*^{\mathrm{nil}}(X)$ to a rational Hodge–Tate crystal $\mathbb{M} \in \mathrm{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$ and hence get a functor

$$\rho^{-1} : \mathrm{HIG}_*^{\mathrm{nil}}(X) \rightarrow \mathrm{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]).$$

By Theorem 4.3, we see from the construction above that ρ and ρ^{-1} are the quasi-inverse of each other.

Finally, let $T : \mathrm{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \rightarrow \mathrm{HIG}_{G_K}(X_C)$ be the composition

$$\mathrm{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \xrightarrow{\mathrm{Res}} \mathrm{Vect}((\mathfrak{X})_{\Delta}^{\mathrm{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \xrightarrow{\sim} \mathrm{Vect}(X_{\mathrm{pro\acute{e}t}}, \widehat{\mathcal{O}}_X) \xrightarrow{\sim} \mathrm{HIG}_{G_K}(X_C).$$

Since $F \circ \rho = T$ after restricting to \mathfrak{X}_i ’s and \mathfrak{X}_{ij} ’s, we have $F(\rho(\mathbb{M}_i)) = T(\mathbb{M}_i)$ and $F(\rho(\iota_{ij})) = T(\iota_{ij})$ for all $i, j \in I$. Hence the diagram (6.3) commutes as desired.

Remark 6.16. If we start with a smooth p -adic formal scheme \mathfrak{X} over \mathcal{O}_C , under some deformation condition on \mathfrak{X} , Theorem 6.4 also holds true for “small Hodge–Tate crystals” and “small Higgs bundles”, by using the overconvergent period sheaf \mathcal{OC}^{\dagger} constructed in [Wang21] to replace \mathcal{OC} . Here, “small Hodge–Tate” crystals should be understood as rational Hodge–Tate crystals admitting $\overline{\mathcal{O}}_{\Delta}$ -lattices which are “close” to $\overline{\mathcal{O}}_{\Delta}^l$ for some $l \geq 0$. Similar remark apply to “small Higgs bundles”. A similar but much more stronger result is also obtained by Tsuji and we appreciate him for telling us this.

7. THE SEN OPERATOR

Let $\mathfrak{X} = \mathrm{Spf}(R^+)$ be a small smooth p -adic formal scheme over \mathcal{O}_K . Given a rational Hodge–Tate crystal $\mathbb{M} \in \mathrm{Vect}((R^+)_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$, there is an associated enhanced Higgs bundle (H, θ_H, ϕ_H) by Proposition 4.23. The goal of this section is to prove that the linear operator ϕ_H is essentially the classical Sen operator in the cyclotomic case (after extending scalar). This result has been used in §6 to prove a global version of Proposition 4.23 in the p -inverted case and construct a global inverse Simpson functor at the same time.

When $R^+ = \mathcal{O}_K$, the coincidence of the operators is our [MW21b, Conjecture 1.4], which is proved later in [Gao22]. The basic idea to show this compatibility here is due to Hui Gao and was used in his previous work [Gao22] using the theory of locally analytic vectors in [BC16]. Hui Gao explained to us that his strategy also works in the relative case. The results in §7.2 are due to Gao, and we thank him for allowing us to include them here. For the basics of the theory of locally analytic vectors, we refer to [BC16, Section 2] and [Gao22, Section 2].

7.1. Sen theory in the cyclotomic case. Recall that in [Sen80], Sen established equivalences of the following categories

$$\mathrm{Rep}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(K_{\mathrm{cyc}}) \rightarrow \mathrm{Rep}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(\widehat{K}_{\mathrm{cyc}}) (\simeq \mathrm{Rep}_{G_K}(C))$$

which preserves ranks, tensor products and dualities. Using this, he constructed a faithful functor from $\mathrm{Rep}_{G_K}(C)$ to $\mathrm{Mod}_*(C)$, the category of pairs (V, ϕ_V) consisting of a finite dimensional C -space V and an endomorphism $\phi_V \in \mathrm{End}_C(V)$ such that for any $V \in \mathrm{Rep}_{G_K}(C)$ with the corresponding pair (W, ϕ_W) , there exists a G_K -equivariant isomorphism

$$\mathrm{R}\Gamma(G_K, V) \otimes_K C \simeq [W \xrightarrow{\phi_W} W].$$

We briefly review the construction of Sen. For any $V \in \mathrm{Rep}_{G_K}(C)$, one can regard it as a representation in $\mathrm{Rep}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(\widehat{K}_{\mathrm{cyc}})$ and denote by V_0 the corresponding K_{cyc} -representation of $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$. Then ϕ_V is the unique endomorphism of V_0 such that for any $v \in V_0$, there exists an open subgroup H_{v_0} of $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ such that for any $\gamma \in H_{v_0}$,

$$(7.1) \quad \gamma(v_0) = \exp(\phi_V \log \chi(\gamma))(v_0).$$

Note that $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ is a p -adic Lie group. It is easy to see that $V_0 = V^{\mathrm{la}}$, the subspace of V consisting of locally analytic vectors with respect to the action of $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$. Then ϕ_V is the generator of the Lie algebra of $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$. See [BC16] for more discussions.

The result of Sen can be generalised to the geometric case. Assume $X = \mathrm{Spa}(R, R^+)$ is smooth over K and admits a toric chart. In [Shi18], Shimizu showed that for any $V \in \mathrm{Rep}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}^{\mathrm{free}}(R_{\widehat{K}_{\mathrm{cyc}}})$ with associated $V_0 \in \varinjlim_n \mathrm{Rep}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}^{\mathrm{free}}(R_{K(\zeta_{p^n})})$ (cf. Theorem 2.8 and Remark 2.2), there exists a unique endomorphism ϕ_V of V_0 such that (7.1) is still true for any $v_0 \in V_0$ and some open subgroup $H_{v_0} \subset \mathrm{Gal}(K_{\mathrm{cyc}}/K)$. His result was generalised to any smooth quasi-compact rigid analytic spaces X over K by Petrov in [Pet20]. Let us fix some notations.

Notation 7.1. For any smooth quasi-compact rigid analytic space X over K , let \mathcal{X} be the ringed space $(X, \mathcal{X} := \mathcal{O}_X \otimes_K K_{\mathrm{cyc}})$. Note that there exists an obvious way to assign each vector bundle on \mathcal{X} (with a continuous $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ -action) to a vector bundle on $X_{\widehat{K}_{\mathrm{cyc}}}$ (with a continuous $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ -action.). Let $\mathrm{Vect}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(\mathcal{X})$ denote the category of vector bundles on \mathcal{X} with continuous $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ -actions.

Proposition 7.2 ([Pet20, Prop. 3.2]). *There exists an equivalence of categories*

$$\mathrm{Vect}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(\mathcal{X}) \simeq \mathrm{Vect}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(X_{\widehat{K}_{\mathrm{cyc}}})$$

which preserves ranks, tensor products and dualities. Here $\mathrm{Vect}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(X_{\widehat{K}_{\mathrm{cyc}}})$ denotes the category of vector bundles on $X_{\widehat{K}_{\mathrm{cyc}}}$ with continuous $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ -actions. Moreover, for any $\mathcal{E} \in \mathrm{Vect}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(\mathcal{X})$, there exists a unique endomorphism $\phi_{\mathcal{E}}$ of \mathcal{E} such that for any affinoid $U = \mathrm{Spa}(R, R^+) \subset X$ and for any section $x \in \mathcal{E}(U)$, there exists an open subgroup $H_x \subset \mathrm{Gal}(K_{\mathrm{cyc}}/K)$ such that for any $\gamma \in H_x$,

$$(7.2) \quad \gamma(x) = \exp(\phi_{\mathcal{E}} \log \chi(\gamma))(x).$$

Now let's assume $X = \mathrm{Spa}(R, R^+)$ admits a toric chart over $\mathrm{Spa}(K, \mathcal{O}_K)$. Following Sen's strategy, Shimizu uses finite vectors (i.e. vectors whose $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ -orbit is a finite set) to construct the decompletion of representations in $\mathrm{Rep}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}^{\mathrm{free}}(R_{\widehat{K}_{\mathrm{cyc}}})$. We are now going to show finite vectors coincide with locally analytic vectors in this case.

Theorem 7.3. *For any finite free $R_{\widehat{K}_{\mathrm{cyc}}}$ -representation W of $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$, we have $W^{\mathrm{fin}} = W^{\mathrm{la}}$, where W^{fin} denotes the subset of W consisting of elements whose $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ -orbit is finite while W^{la} denotes the subset consisting of locally analytic vectors in the sense of [BC16].*

Proof. The proof is similar to that of [BC16, Thm. 3.2]. We first deal with the trivial case, i.e. $W = \widehat{K}_{\mathrm{cyc}}$. It is easy to see $R_{K_{\mathrm{cyc}}} \subset R_{\widehat{K}_{\mathrm{cyc}}}^{\mathrm{la}}$. For the other direction, let $R_n : R_{\widehat{K}_{\mathrm{cyc}}} \rightarrow R_{K(\zeta_{p^n})}$ be the normalised Tate trace such that for any $x \in R_{\widehat{K}_{\mathrm{cyc}}}$, $\lim_{n \rightarrow +\infty} R_n(x) = x$. Suppose $x \in R_{\widehat{K}_{\mathrm{cyc}}}^{\mathrm{Gal}(K_{\mathrm{cyc}}/K)\text{-la}}$ is $\mathrm{Gal}(K_{\mathrm{cyc}}/K(\zeta_{p^m}))$ -analytic for some m . Then for any $k \geq 1$, we have that $R_{m+k}(x)$ is also $\mathrm{Gal}(K_{\mathrm{cyc}}/K(\zeta_{p^m}))$ -analytic and fixed by $\mathrm{Gal}(K_{\mathrm{cyc}}/K(\zeta_{p^{m+k}}))$. Therefore $R_{m+k}(x)$ is also fixed by $\mathrm{Gal}(K_{\mathrm{cyc}}/K(\zeta_{p^m}))$. This shows $x = \lim_{k \rightarrow +\infty} R_{m+k}(x)$ is in $R_{K(\zeta_{p^m})}$.

Now we come to the general case. By the description of the Galois action in Proposition 7.2, it is easy to see that $W^{\mathrm{fin}} \subset W^{\mathrm{la}}$. For the converse direction, let's choose a basis $\{e_1, \dots, e_d\}$ of W^{fin} . Then $W^{\mathrm{la}} = \bigoplus_{i=1}^d R_{\widehat{K}_{\mathrm{cyc}}}^{\mathrm{la}} \cdot e_i = \bigoplus_{i=1}^d R_{K_{\mathrm{cyc}}} \cdot e_i = W^{\mathrm{fin}}$ by [BC16, Prop. 2.3]. \square

Under the equivalence $\mathrm{Rep}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}^{\mathrm{free}}(R_{\widehat{K}_{\mathrm{cyc}}}) \simeq \mathrm{Rep}_{\mathrm{Gal}(K_{\mathrm{cyc},\infty}/K)}^{\mathrm{free}}(R_{\widehat{K}_{\mathrm{cyc},\infty}})$, we interpret the decompletion result in [Shi18], [Pet20] in terms of locally analytic vectors.

Theorem 7.4. *Let $W \in \mathrm{Rep}_{R_{\widehat{K}_{\mathrm{cyc},\infty}}}^{\mathrm{free}}(\mathrm{Gal}(K_{\mathrm{cyc},\infty}/K))$. Define*

$$D_{\mathrm{Sen}, K_{\mathrm{cyc}}}(W) := (W^{\mathrm{Gal}(K_{\mathrm{cyc},\infty}/K_{\mathrm{cyc}})})^{\gamma\text{-la}},$$

where by γ -la, we simply mean the $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ -locally analytic vectors. Then $D_{\mathrm{Sen}, K_{\mathrm{cyc}}}(W)$ is a finite free $R_{K_{\mathrm{cyc}}}$ -module such that $D_{\mathrm{Sen}, K_{\mathrm{cyc}}}(W) \otimes_{R_{K_{\mathrm{cyc}}}} R_{\widehat{K}_{\mathrm{cyc},\infty}} = W$. Moreover there is a linear operator

$$\nabla_{\gamma} : D_{\mathrm{Sen}, K_{\mathrm{cyc}}}(W) \rightarrow D_{\mathrm{Sen}, K_{\mathrm{cyc}}}(W)$$

which is called the Sen operator and can be defined as $\frac{\log(g)}{\log(\chi(g))}$ for $g \in \mathrm{Gal}(K_{\mathrm{cyc},\infty}/K_{\infty}) \cong \mathrm{Gal}(K_{\mathrm{cyc}}/K)$ which is close enough to 1.

7.2. Sen theory in the Kummer case. Following the strategy of [Gao22], we now give a Sen theory with respect to the Kummer tower, which turns out to be closely related to the prismatic theory.

Notation 7.5. (1) Let $K_m := K(\pi^{p^{-\frac{1}{m}}}, \zeta_{p^m})$ and $\hat{G}_m := \mathrm{Gal}(K_{\mathrm{cyc},\infty}/K_m)$ for any $m \geq 0$. In particular, put $\hat{G} = \mathrm{Gal}(K_{\mathrm{cyc},\infty}/K)$

(2) Given any \mathbb{Q}_p -Banach representation W of \hat{G} , let

$$W^{\tau\text{-la}, \gamma^{p^m}=1} := W^{\mathrm{Gal}(K_{\mathrm{cyc},\infty}/K_{\mathrm{cyc}})\text{-la}} \cap W^{\mathrm{Gal}(K_{\mathrm{cyc},\infty}/K_{\infty}(\zeta_{p^m}))=1}$$

for any $m \geq 0$, where $\tau \in \mathrm{Gal}(K_{\mathrm{cyc},\infty}/K_{\mathrm{cyc}})$ is described as in Convention 1.19.

Construction 7.6 ([BC16, §4.4]).

- (1) Since $c : G_K \rightarrow \mathbb{Z}_p$ represents a cocycle in $H^1(G_K, C(1)) = \{0\}$, there exists $\alpha \in C$ such that $c(g) = g(\alpha)\chi(g) - \alpha$. This shows $g(\alpha) = \frac{\alpha}{\chi(g)} + \frac{c(g)}{\chi(g)}$ and in particular $\alpha \in \widehat{K}_{\mathrm{cyc},\infty}^{\hat{G}\text{-la}}$.
- (2) Similarly as in the beginning of [BC16, §4.2], for any $n \geq 0$, let $\alpha_n \in K_{\mathrm{cyc},\infty}$ such that $\|\alpha - \alpha_n\| \leq p^{-n}$. Then there exists some $r(n) \gg 0$ such that if $m \geq r(n)$, then $\|\alpha - \alpha_n\|_{\hat{G}} = \|\alpha - \alpha_n\| \leq p^{-n}$ and $\alpha - \alpha_n \in \widehat{K}_{\mathrm{cyc},\infty}^{\hat{G}_m\text{-la}}$. We may suppose $\{r(n)\}_{n \geq 0}$ is an increasing sequence.

Definition 7.7 ([BC16, §4.4], [Gao22, Def. 3.2.6]). Let $(H, \|\cdot\|)$ be a \mathbb{Q}_p -Banach algebra such that $\|\cdot\|$ is sub-multiplicative and let $W \subset H$ is a \mathbb{Q}_p -subalgebra. For any $n \geq 0$, let $W\{\{T\}\}_n$ denote the vector space consisting of $\sum_{k \geq 0} a_k T^k$ with $a_k \in W$ and $p^{nk} a_k \rightarrow 0$ when $k \rightarrow +\infty$.

Proposition 7.8. (1) $R_{\widehat{K}_{\mathrm{cyc},\infty}}^{\hat{G}\text{-la}} = \bigcup_n R_{K_{r(n)}}\{\{\alpha - \alpha_n\}\}_n$.

- (2) $R_{\hat{K}_{\text{cyc},\infty}}^{\hat{G}\text{-la}, \nabla_\gamma=0} = R \otimes_K K_{\text{cyc},\infty} = R_{K_{\text{cyc},\infty}}.$
 (3) $R_{\hat{K}_{\text{cyc},\infty}}^{\tau\text{-la}, \gamma^{p^m}=1} = R \otimes_K K_\infty(\zeta_{p^m}) = R_{K_\infty(\zeta_{p^m})}.$

Proof. (1) The proof is essentially the same as [BC16, Prop. 4.12]. We now recall the proof therein. Suppose $x \in (R_{\hat{K}_{\text{cyc},\infty}})^{G_n\text{-an}}$. For $i \geq 0$, we let

$$y_i = \sum_{k \geq 0} (-1)^k (\alpha - \alpha_m)^k \frac{\nabla_\tau^{k+i}(x)}{(k+i)!} \binom{k+i}{k},$$

where $\nabla_\tau = \frac{\log(\tau^{p^k})}{p^k}$ for $k \gg 0$. Then by similar arguments in the proof of [BC16, Thm. 4.2], there exists $m \geq n$ such that $y_i \in R_{\hat{K}_{\text{cyc},\infty}}^{G_m\text{-an}}$ for all i and $x = \sum_{i \geq 0} y_i (\alpha - \alpha_m)^i$ in $(R_{\hat{K}_{\text{cyc},\infty}})^{G_m\text{-an}}$. By Construction 7.6, we see that $\nabla_\tau(\alpha - \alpha_m) = 1$. From this, we deduce that $\nabla_\tau(y_i) = 0$ and hence that $\tau^{p^m}(y_i) = y_i$. So $y_i \in (R_{\widehat{K_m(\zeta_{p^\infty})}})^{\text{Gal}(K_m(\zeta_{p^\infty})/K_m)\text{-an}} = R_{K_m}$. We are done.

(2) We obtain from Construction 7.6 that $\nabla_\gamma(\alpha - \alpha_n) = -1$. Therefore, for any $n \geq 0$, we have $(R_{K_{r(n)}}\{\{\alpha - \alpha_n\}\})^{\nabla_\gamma=0} = R_{K_{r(n)}}$. Then Item (2) follows from Item (1) immediately.

(3) This follows from Item (2) by taking γ^{p^m} -invariants. \square

We are now ready to study the Sen theory in the Kummer case.

Theorem 7.9. *Let $W \in \text{Rep}_{R_{\hat{K}_{\text{cyc},\infty}}}^{\text{free}}(\hat{G})$ of rank d and let*

$$D_{\text{Sen}, K_\infty}(W) := W^{\tau\text{-la}, \gamma=1}.$$

Then $D_{\text{Sen}, K_\infty}(W)$ is a finite projective R_{K_∞} -module of rank d . Moreover, there are identifications

$$D_{\text{Sen}, K_\infty}(W) \otimes_{R_{K_\infty}} R_{\hat{K}_{\text{cyc},\infty}}^{\hat{G}\text{-la}} = W^{\hat{G}\text{-la}} = D_{\text{Sen}, K_{\text{cyc}}}(W) \otimes_{R_{K_{\text{cyc}}}} R_{\hat{K}_{\text{cyc},\infty}}^{\hat{G}\text{-la}}.$$

Proof. By [GMW22, Lem. 8.5], we see that $W^{\hat{G}\text{-la}}$ is a finite free $(R_{\hat{K}_{\text{cyc},\infty}})^{\hat{G}\text{-la}}$ -module with

$$W^{\hat{G}\text{-la}} \otimes_{R_{\hat{K}_{\text{cyc},\infty}}^{\hat{G}\text{-la}}} R_{\hat{K}_{\text{cyc},\infty}} \cong W.$$

Furthermore, by [GMW22, Prop. 8.9], we see that $W^{\hat{G}\text{-la}, \nabla_\gamma=0}$ is a finite free $R_{K_{\text{cyc},\infty}}$ -module with

$$W^{\hat{G}\text{-la}, \nabla_\gamma=0} \otimes_{R_{K_{\text{cyc},\infty}}} R_{\hat{K}_{\text{cyc},\infty}}^{\hat{G}\text{-la}} \cong W^{\hat{G}\text{-la}}.$$

Note that $W^{\hat{G}\text{-la}, \nabla_\gamma=0}$ admits a $\text{Gal}(K_{\text{cyc},\infty}/K_\infty)$ -action. As $\text{Gal}(K_{\text{cyc},\infty}/K_\infty)$ is topologically finitely generated, we can find a finite free R_{K_m} -module W_m of rank d with a $\text{Gal}(K_m/K(\pi^{\frac{1}{p^m}}))$ -action for some $m \gg 0$, such that $W_m \otimes_{R_{K_m}} R_{K_{\text{cyc},\infty}} \cong W^{\hat{G}\text{-la}, \nabla_\gamma=0}$ is a $\text{Gal}(K_{\text{cyc},\infty}/K_\infty)$ -equivariant isomorphism. Now by Galois descent, we get a finite projective $R_{K(\pi^{\frac{1}{p^m}})}$ -module $W_m^{\gamma=1}$ such that $W_m^{\gamma=1} \otimes_{R_{K(\pi^{\frac{1}{p^m}})}} R_{K_m} \cong W_m$.

We claim that the finite projective R_{K_∞} -module $W_m^{\gamma=1} \otimes_{R_{K(\pi^{\frac{1}{p^m}})}} R_{K_\infty}$ is exactly $D_{\text{Sen}, K_\infty}(W)$. To see this, we first show that $W^{\tau\text{-la}} = W_m^{\gamma=1} \otimes_{R_{K(\pi^{\frac{1}{p^m}})}} R_{\hat{K}_{\text{cyc},\infty}}^{\tau\text{-la}}$. Note that $W^{\hat{G}\text{-la}, \nabla_\gamma=0}$ is contained in $W^{\tau\text{-la}}$. Then by [BC16, Prop. 2.3], we have

$$W^{\tau\text{-la}} = W^{\hat{G}\text{-la}, \nabla_\gamma=0} \otimes_{R_{K_{\text{cyc},\infty}}} R_{\hat{K}_{\text{cyc},\infty}}^{\tau\text{-la}} = W_m^{\gamma=1} \otimes_{R_{K(\pi^{\frac{1}{p^m}})}} R_{\hat{K}_{\text{cyc},\infty}}^{\tau\text{-la}}.$$

Now, the claim follows from Proposition 7.8 (3) after taking γ -invariants. We win. \square

Similar to the cyclotomic case, there is also a linear operator in the Kummer case. Before we move on, we recall some constructions in [Gao22].

Construction 7.10. (1) For any \hat{G} -locally analytic representation W , there are two Lie algebra differential operators:

- (a) $\nabla_\gamma := \frac{\log(g)}{\log(\chi(g))}$ for $g \in \text{Gal}(K_{\text{cyc},\infty}/K_\infty)$ close enough to 1. (See also Theorem 7.4.)

- (b) $\nabla_\tau := \frac{\log(\tau^{p^n})}{p^n}$ for $n \gg 0$. (See also the proof of Proposition 7.8 (1).)
- (2) Let $\tilde{\lambda} := \prod_{n \geq 0} \varphi^n(\frac{E([\pi^b])}{E(0)}) \in B_{\text{cris}}^+$.⁸ Then we can define $\mathfrak{t} := \frac{t}{p\lambda}$ which turns out to be in A_{inf} .
- (3) Define $N_\nabla : R_{\hat{K}_{\text{cyc},\infty}}^{\hat{G}\text{-la}} \rightarrow R_{\hat{K}_{\text{cyc},\infty}}^{\hat{G}\text{-la}}$ by setting $N_\nabla := \frac{1}{p\theta(\mathfrak{t})}\nabla_\tau$. Recall we always have $K_\infty \cap K_{\text{cyc}} = K$ (cf. Convention 1.19). By [Gao22, Lem. 3.2.1], $\theta(\mathfrak{t}) \in \hat{K}_{\text{cyc},\infty}^{\hat{G}\text{-la}}$ is non-zero and hence the operator N_∇ is well-defined.

Let $W \in \text{Rep}_{R_{\hat{K}_{\text{cyc},\infty}}}^{\text{free}}(\hat{G})$. Then $D_{\text{Sen},K_\infty}(W)$ consists of locally analytic vectors of W and we have the following operator

$$N_\nabla : D_{\text{Sen},K_\infty}(W) \rightarrow W^{\hat{G}\text{-la}}.$$

Theorem 7.11. *Let $W \in \text{Rep}_{R_{\hat{L}}}^{\text{free}}(\hat{G})$. Then there is an R_{K_∞} -linear operator*

$$\frac{1}{\theta(u\tilde{\lambda}')} N_\nabla : D_{\text{Sen},K_\infty}(W) \rightarrow D_{\text{Sen},K_\infty}(W)$$

where $\tilde{\lambda}'$ is the u -derivative of $\tilde{\lambda}$. After extending $R_{\hat{K}_{\text{cyc},\infty}}$ -linearly, the operator $\frac{1}{\theta(u\tilde{\lambda}')} N_\nabla : W \rightarrow W$ is exactly the Sen operator in Theorem 7.4.

Proof. Note that for any $g \in \text{Gal}(K_{\text{cyc},\infty}/K_\infty)$, $g(\mathfrak{t}) = \chi(g)\mathfrak{t}$ and $g\tau g^{-1} = \tau^{\chi(g)}$. We see that N_∇ commutes with the action of $\text{Gal}(K_{\text{cyc},\infty}/K_\infty)$. So $N_\nabla : D_{\text{Sen},K_\infty}(W) \rightarrow D_{\text{Sen},K_\infty}(W)$ is well-defined.

By [Gao22, Cor. 3.2.4] and Proposition 7.8, we see that the operator $\nabla_{\text{Sen}} := \frac{1}{\theta(u\tilde{\lambda}')} N_\nabla + \nabla_\gamma$ acts $R_{\hat{K}_{\text{cyc},\infty}}^{\hat{G}\text{-la}}$ -linearly on $W^{\hat{G}\text{-la}}$. Since ∇_γ kills $D_{\text{Sen},K_\infty}(W)$, the operator ∇_{Sen} is equal to $\frac{1}{\theta(u\tilde{\lambda}')} N_\nabla$ on $W^{\hat{G}\text{-la}} = D_{\text{Sen},K_\infty}(W) \otimes_{R_{K_\infty}} R_{\hat{K}_{\text{cyc},\infty}}^{\hat{G}\text{-la}}$. Similarly, as N_∇ kills $D_{\text{Sen},K_{\text{cyc}}}(W)$, the operator ∇_{Sen} is equal to ∇_γ on $W^{\hat{G}\text{-la}} = D_{\text{Sen},K_{\text{cyc}}}(W) \otimes_{R_{K_{\text{cyc}}}} R_{\hat{K}_{\text{cyc},\infty}}^{\hat{G}\text{-la}}$. So we are done. \square

Now we come back to the prismatic theory and show how it is related to the Sen operator. Let $\mathfrak{X} = \text{Spf}(R^+)$ be a small smooth p -adic formal scheme over \mathcal{O}_K with the adic generic fiber $X = \text{Spa}(R, R^+)$. We will apply the above results to $R_{\hat{K}_{\text{cyc},\infty}}$ -representations of \hat{G} coming from enhanced Higgs modules in $\text{HIG}_*^{\text{nil}}(R)$ in the sense of Construction 6.8 and Remark 6.10.

Theorem 7.12 (Hui Gao). *Let $F : \text{HIG}_*^{\text{nil}}(R) \rightarrow \text{HIG}_{\hat{G}}(R_{\hat{K}_{\text{cyc},\infty}})$ be the functor defined in Construction 6.8 (together with Remark 6.10). For any $(H, \theta_H, \phi_H) \in \text{HIG}_*^{\text{nil}}(R)$, let $W := F(H, \theta_H, \phi_H)$ be the underlying $R_{\hat{K}_{\text{cyc},\infty}}$ -representations of \hat{G} . Assume furthermore that H is finite free over R . Then*

- (1) $D_{\text{Sen},K_\infty}(W) = H \otimes_R R_{K_\infty}$.
- (2) The operator $\frac{-\phi_H}{E'(\pi)}$ is exactly the operator $\frac{1}{\theta(u\tilde{\lambda}')} N_\nabla$ on $D_{\text{Sen},K_\infty}(W)$ after extending linearly to R_{K_∞} . Moreover, $\frac{-\phi_H}{E'(\pi)}$ is the Sen operator of W .

Proof. (1) By the Galois action described in Lemma 6.9, we see that $H \subset D_{\text{Sen},K_\infty}(W)$. By [BC16, Prop. 2.3], we see that $W^{\tau\text{-la}} = H \otimes_R R_{\hat{K}_{\text{cyc},\infty}}^{\tau\text{-la}}$. Since $W^{\gamma=1} = H \otimes_R R_{\hat{K}_{\text{cyc},\infty}}^{\gamma=1}$, we get that

$$D_{\text{Sen},K_\infty}(W) = (H \otimes_R R_{\hat{K}_{\text{cyc},\infty}}^{\tau\text{-la}}) \cap (H \otimes_R R_{\hat{K}_{\text{cyc},\infty}}^{\gamma=1}) = H \otimes_R (R_{\hat{K}_{\text{cyc},\infty}}^{\tau\text{-la},\gamma=1}) = H \otimes_R R_{K_\infty},$$

where the last equality follows from Proposition 7.8 (3).

(2) By Lemma 6.9, we see that ∇_τ acts on $D_{\text{Sen},K_\infty}(W)$ via

$$\begin{aligned} \lim_{n \rightarrow +\infty} \frac{\tau^{p^n} - 1}{p^n} &= \lim_{n \rightarrow +\infty} \frac{1}{p^n} ((1 + \pi E'(\pi)(\zeta_p - 1)\lambda p^n)^{-\frac{\phi_H}{E'(\pi)}} - 1) \\ &= \lim_{n \rightarrow +\infty} \sum_{k \geq 1} \prod_{i=0}^{k-1} (\phi_H + iE'(\pi)) \frac{(-\pi(\zeta_p - 1)\lambda)^k p^{(k-1)n}}{k!} \\ &= -\pi(\zeta_p - 1)\lambda \phi_H. \end{aligned}$$

⁸Note that $\tilde{\lambda}$ is denoted by λ in [Gao22, Def. 2.2.2]. To distinguish it from the $\lambda := \theta(\frac{\xi}{E([\pi^b])})$ defined in this paper, we denote it by $\tilde{\lambda}$.

Therefore, we deduce that

$$\frac{1}{\theta(u\tilde{\lambda}')}N_{\nabla} = \frac{1}{p\theta(u\tilde{\lambda}'t)}\nabla_{\tau} = \frac{\pi E'(\pi)(\zeta_p - 1)\lambda}{p\theta(u\tilde{\lambda}'t)}\left(-\frac{\phi_H}{E'(\pi)}\right).$$

To conclude Item (2), we are reduced to showing that

$$\frac{\pi E'(\pi)(\zeta_p - 1)\lambda}{p\theta(u\tilde{\lambda}'t)} = 1,$$

which follows exactly from the calculations in the proof of [Gao22, Thm. 4.3.3 (2)]. \square

Remark 7.13. As we have mentioned in the beginning of this section, when $R = K$, Theorem 7.12 is conjectured in [MW21b] and proved by Hui Gao in [Gao22]. The method to prove Theorem 7.12 is a generalisation of Gao's work [Gao22].

When $R = W(k)[\frac{1}{p}]$ i.e. in the unramified case, Bhatt and Lurie has proved a similar result in [BL22a]. More precisely, by using the prism $(W(k)[[\tilde{p}]], (\tilde{p}))$ (corresponding to the cyclotomic case), which is the \mathbb{F}_p^\times -invariants of the q -de Rham prism, instead of the Breuil–Kisin prism $(W(k)[[u]], (u - p))$ (corresponding to the Kummer case), they proved that the linear operator associated with a Hodge–Tate crystal is exactly the classical Sen operator in the cyclotomic case. Unlike the q -de Rham prism, its \mathbb{F}_p^\times -invariants behaves quite similarly to the Breuil–Kisin prism. It will be very interesting to figure out whether such a prism exists in the ramified and geometric case.

7.3. Arithmetic Higgs bundles. In this subsection, let's assume \mathfrak{X} is a smooth quasi-compact p -adic formal scheme over \mathcal{O}_K with adic generic fiber X . Let \mathcal{X} be the ringed space introduced in Notation 7.1.

Definition 7.14. By an *arithmetic Higgs bundle* of rank l on X , we mean a triple $(\mathcal{E}, \theta_{\mathcal{E}}, \phi_{\mathcal{E}})$ satisfying

- (1) $(\mathcal{E}, \theta_{\mathcal{E}})$ is a Higgs bundle on \mathcal{X} (see Notation 7.1); in other words, \mathcal{E} is a vector bundle on \mathcal{X} and $\theta_{\mathcal{E}} : \mathcal{E} \rightarrow \mathcal{E} \otimes_{\mathcal{O}_X} \Omega_{X/K}^1(-1)$ is a Higgs field on \mathcal{E} (i.e. $\theta_{\mathcal{E}}$ is $\mathcal{O}_{\mathcal{X}}$ -linear such that $\theta_{\mathcal{E}} \wedge \theta_{\mathcal{E}} = 0$);
- (2) $\phi_{\mathcal{E}}$ is an $\mathcal{O}_{\mathcal{X}}$ -linear endomorphism of \mathcal{E} satisfying $[\phi_{\mathcal{E}}, \theta_{\mathcal{E}}] = -\theta_{\mathcal{E}}$; that is, the following diagram

$$\begin{array}{ccccccc} \mathcal{E} & \xrightarrow{\theta_{\mathcal{E}}} & \mathcal{E} \otimes_{\mathcal{O}_X} \Omega_{X/K}^1(-1) & \rightarrow \cdots \rightarrow & \mathcal{E} \otimes_{\mathcal{O}_X} \Omega_{X/K}^d(-d) \\ \phi_{\mathcal{E}} \downarrow & & \phi_{\mathcal{E}} - \text{id}_{\mathcal{E}} \downarrow & & \phi_{\mathcal{E}} - d \text{id}_{\mathcal{E}} \downarrow \\ \mathcal{E} & \xrightarrow{\theta_{\mathcal{E}}} & \mathcal{E} \otimes_{\mathcal{O}_X} \Omega_{X/K}^1(-1) & \rightarrow \cdots \rightarrow & \mathcal{E} \otimes_{\mathcal{O}_X} \Omega_{X/K}^d(-d) \end{array}$$

is commutative. We denote its total complex by $\text{HIG}(\mathcal{E}, \theta_{\mathcal{E}}, \phi_{\mathcal{E}})$.

Let $\text{HIG}^{\text{arith}}(X, \mathcal{O}_{\mathcal{X}})$ be the category of arithmetic Higgs bundles on X .

Similarly, we have the following definition.

Definition 7.15. Let L/K be any Galois extension of K in \overline{K} containing K_{cyc} . By an *arithmetic Higgs bundle* of rank l on $X_{\widehat{L}}$, we mean a triple $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$ satisfying

- (1) $(\mathcal{H}, \theta_{\mathcal{H}})$ is a Higgs bundle on $X_{\widehat{L}}$;
- (2) $\phi_{\mathcal{H}}$ is an $\mathcal{O}_{X_{\widehat{L}}}$ -linear endomorphism of \mathcal{H} satisfying $[\phi_{\mathcal{H}}, \theta_{\mathcal{H}}] = -\theta_{\mathcal{H}}$; that is, the following diagram

$$\begin{array}{ccccccc} \mathcal{H} & \xrightarrow{\theta_{\mathcal{H}}} & \mathcal{H} \otimes_{\mathcal{O}_X} \Omega_{X/K}^1(-1) & \rightarrow \cdots \rightarrow & \mathcal{H} \otimes_{\mathcal{O}_X} \Omega_{X/K}^d(-d) \\ \phi_{\mathcal{H}} \downarrow & & \phi_{\mathcal{H}} - \text{id}_{\mathcal{H}} \downarrow & & \phi_{\mathcal{H}} - d \text{id}_{\mathcal{H}} \downarrow \\ \mathcal{H} & \xrightarrow{\theta_{\mathcal{H}}} & \mathcal{H} \otimes_{\mathcal{O}_X} \Omega_{X/K}^1(-1) & \rightarrow \cdots \rightarrow & \mathcal{H} \otimes_{\mathcal{O}_X} \Omega_{X/K}^d(-d) \end{array}$$

is commutative. We denote its total complex by $\text{HIG}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$.

Let $\text{HIG}^{\text{arith}}(X_{\widehat{L}})$ be the category of arithmetic Higgs bundles on $X_{\widehat{L}}$.

The following result follows from Proposition 7.2 directly.

Corollary 7.16. *Assume X is a quasi-compact smooth rigid analytic space over K .*

(1) *There exists an equivalence of categories*

$$\mathrm{HIG}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(X, \mathcal{O}_X) \simeq \mathrm{HIG}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(X_{\widehat{K}_{\mathrm{cyc}}}),$$

which preserves ranks, tensor products and dualities. Here, $\mathrm{HIG}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(X, \mathcal{O}_X)$ denotes the category of Higgs bundles on X equipped with continuous $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ -actions which are compatible with Higgs fields.

(2) *For any extension L of K containing K_{cyc} , there is a faithful functor*

$$\mathrm{HIG}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(X, \mathcal{O}_X) \rightarrow \mathrm{HIG}^{\mathrm{arith}}(X, \mathcal{O}_X).$$

(3) *There is faithful functors*

$$\mathrm{HIG}^{\mathrm{arith}}(X, \mathcal{O}_X) \rightarrow \mathrm{HIG}^{\mathrm{arith}}(X_{\widehat{L}}).$$

Proof. (1) This follows from Proposition 7.2 directly.

(2) For any $(\mathcal{H}, \theta_{\mathcal{H}}) \in \mathrm{HIG}_{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}(X, \mathcal{O}_X)$, let $(\mathcal{E}, \theta_{\mathcal{E}}, \phi_{\mathcal{E}}) \in \mathrm{HIG}^{\mathrm{arith}}(X, \mathcal{O}_X)$ such that $(\mathcal{E}, \theta_{\mathcal{E}}) = (\mathcal{H}, \theta_{\mathcal{H}})$ and that $\phi_{\mathcal{E}}$ is the unique operator determined by (7.2). Then the functor sending $(\mathcal{H}, \theta_{\mathcal{H}})$ to $(\mathcal{E}, \theta_{\mathcal{E}}, \phi_{\mathcal{E}})$ is well-defined. We claim it is faithful. For this purpose, we may assume $X = \mathrm{Spa}(R, R^+)$ is affinoid and are reduced to showing that

$$(E^{\theta_E=0})^{\mathrm{Gal}(K_{\mathrm{cyc}}/K)} \subset E^{\theta_E=0, \phi_E=0},$$

where (E, θ_E, ϕ_E) denotes the global section of $(\mathcal{E}, \theta_{\mathcal{E}}, \phi_{\mathcal{E}})$ on X , which also inherits a $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$ -action from $(\mathcal{H}, \theta_{\mathcal{H}})$. However, for any $x \in (E^{\theta_E=0})^{\mathrm{Gal}(K_{\mathrm{cyc}}/K)}$, by (7.2), we have

$$\phi_E(x) = \lim_{\gamma \rightarrow 1} \frac{\gamma(x) - x}{\log \chi(\gamma)} = 0,$$

which is exactly what we want.

(3) We do have the desired functors by considering corresponding scalar extensions. To see the faithfulness, we may assume $X = \mathrm{Spa}(R, R^+)$ admits a toric chart such that (H, θ_H, ϕ_H) , the global section of $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) \in \mathrm{HIG}^{\mathrm{arith}}(X, \mathcal{O}_X)$ on X , has H finite free over $R \otimes_K K_{\mathrm{cyc}}$. So we need to show that

$$H^{\theta_H=0, \phi_H=0} \subset (H \otimes_R R_{\widehat{L}})^{\theta_H=0, \phi_H=0}.$$

This follows from the faithful flatness of $R \rightarrow R_{\widehat{L}}$. □

Now, we give a relative version of classical Sen theory.

Corollary 7.17. *There exists a faithful functor*

$$\mathrm{D} : \mathrm{Vect}(X_{\mathrm{pro\acute{e}t}}, \widehat{\mathcal{O}}_X) \rightarrow \mathrm{HIG}^{\mathrm{arith}}(X_C)$$

from the category of generalised representations on $X_{\mathrm{pro\acute{e}t}}$ to the category of arithmetic Higgs bundles on X_C , which preserves ranks, tensor products and dualities.

Proof. Just combine Theorem 6.4 (for replacing C and G_K by $\widehat{K}_{\mathrm{cyc}}$ and $\mathrm{Gal}(K_{\mathrm{cyc}}/K)$) with Corollary 7.16. □

Remark 7.18. When $X = \mathrm{Spa}(K, \mathcal{O}_K)$, we see that $\mathrm{Vect}(X_{\mathrm{pro\acute{e}t}}, \widehat{\mathcal{O}}_X) = \mathrm{Rep}_{G_K}(C)$ is the category of C -representations of G_K and that $\mathrm{HIG}^{\mathrm{arith}}(X_C) = \mathrm{Mod}_*(C)$ is the category of pairs (V, ϕ_V) consisting of finite dimensional C -spaces together with endomorphisms ϕ_V of V . Then the functor in Corollary 7.17 coincides with the classical functor

$$\mathrm{D} : \mathrm{Rep}_{G_K}(C) \rightarrow \mathrm{Mod}_*(C)$$

introduced in [Sen80]. Note that even in this case, the above functor is not fully faithful.

Recall that for any $V \in \mathrm{Rep}_{G_K}(C)$, the underlying C -space of $\mathrm{D}(V)$ is V and the induced Sen operator ϕ_V is indeed defined over K (cf. [Sen80, Thm. 5]) by using matrix theory and hence D upgrades to a functor from $\mathrm{Rep}_{G_K}(C)$ to $\mathrm{Mod}_*(K)$, the category of pairs (V, ϕ_V) consisting of finite dimensional K -spaces together with endomorphisms ϕ_V of V . So the following question appears naturally.

Question 7.19. For a generalised representation $\mathcal{L} \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ with the induced arithmetic Higgs bundle $D(\mathcal{L})$ over X_C , is $D(\mathcal{L})$ always defined over X itself?

On the other hand, not any pairs (V, ϕ_V) in $\text{Mod}_*(C)$ with ϕ_V defined over K comes from a C -representation V of G_K (cf. [Sen80, Thm. 7]). So one may also ask

Question 7.20. Is any arithmetic Higgs bundle $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$ over X_C of the form $D(\mathcal{L})$ for some generalised representation $\mathcal{L} \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$?

Remark 7.21. When $\mathfrak{X} = \text{Spf}(\mathcal{O}_K)$, Question 7.20 was solved by Fontaine [Fon04] after classifying all C -representations of G_K , using the representation theory of certain algebraic group over K . His method looks difficult to generalised to the relative case. For example, even for small affine \mathfrak{X} , we do not have a classification of generalised representations on $X_{\text{proét}}$. We even do not know whether the Hodge–Tate weights of a generalised representation at each classical point is constant, unless the generalised representations comes from \mathbb{Q}_p -local systems (cf. [Shi18]).

We are going to give partial answers of the above two questions in sequels.

Construction 7.22. For any enhanced Higgs bundle $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) \in \text{HIG}_*^{\text{nil}}(X)$, define

- (1) the Kummer case: $F_{\infty}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) = (\mathcal{H} \otimes_{\mathcal{O}_{\mathfrak{X}}} \mathcal{O}_{X_C}, -(\zeta_p - 1)\lambda\theta_{\mathcal{H}}, -\frac{\phi_{\mathcal{H}}}{E'(\pi)}) \in \text{HIG}^{\text{arith}}(X_C)$, and
- (2) the cyclotomic case: $F_{\text{cyc}}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) \in \text{HIG}^{\text{arith}}(X_C)$ as the image under the composition of

$$\text{HIG}_*^{\text{nil}}(X) \xrightarrow{F_S} \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X) \xrightarrow{D} \text{HIG}^{\text{arith}}(X_C),$$

where F_S is the inverse Simpson functor and D is defined in Corollary 7.17.

Then we get two functors

$$F_{\infty}, F_{\text{cyc}} : \text{HIG}_*^{\text{nil}}(X) \rightarrow \text{HIG}^{\text{arith}}(X_C).$$

Let F be as in Construction 6.13. Then we see that F_{cyc} coincides with the composition

$$\text{HIG}_*^{\text{nil}}(X) \xrightarrow{F} \text{HIG}_{G_K}(X_C) \xrightarrow{\text{Corollary 7.16}} \text{HIG}^{\text{arith}}(X_C).$$

Theorem 7.23. *There is an equivalence of functors $F_{\infty} \simeq F_{\text{cyc}}$.*

Proof. Fix an $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) \in \text{HIG}_*^{\text{nil}}(X)$. By Construction 6.13, we see that the underlying Higgs bundles of $F_{\infty}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) \in \text{HIG}_*^{\text{nil}}(X)$ and $F_{\Delta}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) \in \text{HIG}_*^{\text{nil}}(X)$ are same. So we are reduced to comparing the corresponding Sen operators. In other words, we have to show the Sen operator of the underlying G_K -vector bundle of $F(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) \in \text{HIG}_*^{\text{nil}}(X)$ is exactly $-\frac{\phi_{\mathcal{H}}}{E'(\pi)}$. By the uniqueness criterion of Sen operator in Proposition 7.2, we may assume $\mathfrak{X} = \text{Spf}(R^+)$ is small affine such that $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$ is induced by an enhanced Higgs module (H, θ_H, ϕ_H) over R with H finite free. Then the result follows from Theorem 7.12. \square

Corollary 7.24. *Let \mathfrak{X} be a quasi-compact formal scheme over \mathcal{O}_K with rigid generic fiber X .*

- (1) *Let \mathcal{L} be a generalised representation with the corresponding arithmetic Higgs bundle $D(\mathcal{L}) \in \text{HIG}^{\text{arith}}(X_C)$. If \mathcal{L} belongs to the essential image of the functor*

$$\text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \xrightarrow{\text{Res}} \text{Vect}((\mathfrak{X})_{\Delta}^{\text{perf}}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}]) \xrightarrow{\simeq} \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X),$$

then $D(\mathcal{L})$ is defined over X .

- (2) *Let $(\mathcal{E}, \theta_{\mathcal{E}}, \phi_{\mathcal{E}})$ be an arithmetic Higgs bundle over X_C . If it is enhanced in the sense that it lies in the essential image of F_{∞} , then there is a generalised representation $\mathcal{L} \in \text{Vect}(X_{\text{proét}}, \widehat{\mathcal{O}}_X)$ such that $D(\mathcal{L}) \cong (\mathcal{E}, \theta_{\mathcal{E}}, \phi_{\mathcal{E}})$.*

Proof. (1) Assume \mathcal{L} is associated to $\text{Res}(\mathbb{M})$ for some $\mathbb{M} \in \text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$ via the equivalence in Theorem 5.1. By Theorem 7.23, we see that $D(\mathcal{L}) \cong F_{\text{cyc}}(\rho(\mathbb{M}))$ is defined over X , where ρ is the equivalence defined in Theorem 6.4.

(2) Assume $(\mathcal{E}, \theta_{\mathcal{E}}, \phi_{\mathcal{E}}) = F_{\infty}(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$ for some enhance Higgs bundle $(\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}}) \in \text{HIG}_*^{\text{nil}}(X)$. Let $\mathbb{M} \in \text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$ such that $\rho(\mathbb{M}) = (\mathcal{H}, \theta_{\mathcal{H}}, \phi_{\mathcal{H}})$. Let \mathcal{L} be the generalised representation on $X_{\text{proét}}$ corresponding to $\text{Res}(\mathbb{M})$ via the equivalence in Theorem 5.1. Then by Theorem 7.23, we see that $(\mathcal{E}, \theta_{\mathcal{E}}, \phi_{\mathcal{E}}) \cong D(\mathcal{L})$ as desired. \square

Note that Corollary 7.24 answers Question 7.19 and Question 7.20 partially.

Remark 7.25. Let \mathfrak{X} be a quasi-compact formal scheme over \mathcal{O}_K with rigid generic fiber X as before.

Let \mathcal{L} be a generalised representation on $X_{\text{proét}}$. Note that for any classical point $x \in X$, \mathcal{L}_x is a $\widehat{k(x)}$ -representation of $\text{Gal}(\widehat{k(x)}/k(x))$, where $k(x)$ denotes the residue field of x . Let ϕ_x be the Sen operator of \mathcal{L}_x .

Assume \mathcal{L} is induced from some rational Hodge–Tate crystal $\mathbb{M} \in \text{Vect}((\mathfrak{X})_{\Delta}, \overline{\mathcal{O}}_{\Delta}[\frac{1}{p}])$. Then by Theorem 7.23, we see from Definition 6.2 (2) that ϕ_x is topologically nilpotent in the sense that

$$\lim_{n \rightarrow +\infty} E'(\pi)^n \prod_{i=0}^{n-1} (\phi_V - i) = 0.$$

Therefore, \mathcal{L}_x is a *nearly Hodge–Tate representation* of $\text{Gal}(\widehat{k(x)}/k(x))$ over $\widehat{k(x)}$ in the sense of [Gao22].

However, assume that for any classical point $x \in X$, \mathcal{L}_x is nearly Hodge–Tate (or even Hodge–Tate). We do not know whether \mathcal{L} is induced from a rational Hodge–Tate crystal on $(\mathfrak{X})_{\Delta}$.

REFERENCES

- [AGT16] A. Abbes, M. Gros, T. Tsuji: The p -adic Simpson Correspondence, *Annals of Mathematics Studies*. 193 (2016).
- [ALB19] J. Anschütz and A.-C. Le Bras: *Prismatic Dieudonné theory*, arxiv:1907.10525v3 (2019).
- [ALB20] J. Anschütz and A.-C. Le Bras: *The p -completed cyclotomic trace in degree 2*, *Ann. K-Th.* 5, pp. 539–580, (2020).
- [AHLB22] J. Anschütz, B. Heuer and A.-C. Le Bras: *v -vector bundles on p -adic fields and Sen theory via the Hodge-Tate stack*, arxiv:2211.08470, (2022).
- [BC08] L. Berger and P. Colmez: *Familles de représentations de de Rham et monodromie p -adique*, *Astérisque*, tome 319 (2008), pp. 303–337.
- [BC16] L. Berger and P. Colmez: *Théorie de Sen et vecteurs localement analytiques*, *Ann. Sci. École Norm. Sup.* 49 (2016), no. 4, 947–970, (2016).
- [BdJ11] B. Bhatt and A. J. de Jong: *Crystalline cohomology and de Rham cohomology*, arXiv:1110.5001, (2011).
- [Bha16] B. Bhatt: *Specializing varieties and their cohomology from characteristic 0 to characteristic p* , arxiv:1606.01463v2, (2012).
- [Bha] B. Bhatt: *Lecture 4: Perfect prism and perfectoid rings*, available at <http://www-personal.umich.edu/bhattb/teaching/prismatic-columbia/lecture4-perfect-prisms-and-perfectoids.pdf>.
- [BL22a] B. Bhatt and J. Lurie: *Absolute prismatic cohomology*, arXiv:2201.06120, (2022).
- [BL22b] B. Bhatt and J. Lurie: *The prismaticization of p -adic formal schemes*, arXiv:2201.06124, (2022).
- [BMS18] B. Bhatt, M. Morrow and P. Scholze: *Integral p -adic Hodge theory*, *Publications mathématiques de l’IHÉS* volume 128, pages 219–397 (2018).
- [BS22] B. Bhatt and P. Scholze: *Prisms and prismatic cohomology*, *Ann. of Math. (2)* 196(3): 1135–1275, (2022).
- [BS21] B. Bhatt and P. Scholze: *Prismatic F -crystals and crystalline Galois representations*, arxiv:2106.14735v1, (2021).
- [CS19] K. Česnavičius and P. Scholze: *Purity for flat cohomology*, arXiv:1912.10932v2, (2019).
- [DL21] H. Du and T. Liu: *A prismatic approach to (φ, \hat{G}) -modules and F -crystals*, arXiv:2107.12240v1, (2021).
- [DLLZ18] H. Diao, K.-W. Lan, R. Liu, X. Zhu: *Logarithmic Riemann-Hilbert correspondences for rigid varieties*, arXiv:1803.05786v2 (2018).
- [DLMS22] H. Du, T. Liu, Y.-S. Moon, K. Shimizu: *Completed prismatic F -crystals and crystalline \mathbb{Z}_p -local systems*, arxiv:2203.03444, (2022).
- [Dr21] V. Drinfeld: *A 1-dimensional formal group over the prismaticization of $\mathrm{Spf}\mathbb{Z}_p$* , arxiv:2107.11466, (2021).
- [DW05] C. Deninger, A. Werner: *Vector bundles on p -adic curves and parallel transport*, *Ann. Sci. Ecole Norm. Sup.* 38, pp. 553–597 (2005).
- [DW20] C. Deninger, A. Werner: *Parallel transport for vector bundles on p -adic varieties*, *J. Algebraic Geom.* 29, no. 1, pp. 1–52, (2020).
- [EG19] M. Emerton and T. Gee: *Moduli stack of étale (φ, Γ) -modules and the existence of crystalline lifts*, arXiv:1908.07185v2, (2020).
- [Fal05] G. Faltings: *A p -adic Simpson correspondence*, *Advances in Mathematics*. 198 pp. 847–862 (2005).
- [Fon04] J.-M. Fontaine: *Arithmétique des représentations galoisiennes p -adiques*, *Astérisque* 295, pages xi, 1–115. *Cohomologies p -adiques et applications arithmétiques. III.* (2004).
- [FvdP04] J. Frensel, M. van der Put: *Rigid analytic geometry and its applications*, *Progress in mathematics* (Boston, Mass.), v. 218, (2004).
- [Gao22] H. Gao: *Hodge-Tate prismatic crystals and Sen theory*, arXiv:2201.10136, (2022).
- [GM02] L. Grunewald and M. Mastnak: *Cohomology of abelian matched pairs and the Kac sequence*, *Journal of Algebra* 276, 706–736 (2002).
- [GLSQ20] M. Gros, B. Le Stum and A. Quirós: *Twisted differential operators and q -crystals*, arxiv:2004.14320.
- [GLSQ22] M. Gros, B. Le Stum and A. Quirós: *Cartier transform and prismatic crystals*, arxiv:2203.09897.
- [GMW22] H. Gao, Y. Min and Y. Wang: *de Rham crystals on the prismatic site of \mathcal{O}_K* , arXiv:2206.10276, (2022).
- [GR22] H. Guo, E. Reinecke: *Prismatic F -crystals and crystalline local systems*, arxiv:2203.09490, (2022).
- [Gro57] A. Grothendieck: *Sur quelques points d’algèbre homologique*, *Tohoku Math. J. (2)* 9, 119–221, (1957).
- [Heu22a] B. Heuer: *Line bundles on rigid spaces in the v -topology*, *Forum of Mathematics, Sigma*, 10, E82., pp. 1–36 (2022).
- [Heu22b] B. Heuer: *Moduli spaces in p -adic non-abelian Hodge theory*, arXiv:2207.13819, (2022).
- [HMW22] Ben Heuer, Lucas Mann, Annette Werner: *The p -adic Corlette-Simpson correspondence for abeloids*, *Math. Ann.* (2022).
- [Hum72] J.E. Humphreys: *Introduction to Lie algebras and representation theory*, *Graduate texts in mathematics*. Vol 9, Springer-Verlag, New York, 1972.
- [Hy89] O. Hyodo: *On variation of Hodge-Tate structures*, *Math. Ann.* 284, pp. 7–22 (1989).
- [Ked16] K.S. Kedlaya: *The Hochschild-Serre property for some p -adic analytic group actions*, *Ann. Math. Québec* (2016) 40:149–157.
- [KL15] K.S. Kedlaya and R. Liu: *Relative p -adic Hodge theory: Foundations*, *Astérisque* 371, (2015).
- [KL16] K.S. Kedlaya and R. Liu: *Relative p -adic Hodge theory, II: Imperfect period rings*, 1602.06899v3 (2016).

- [Laz65] M. Lazard: *Groupes analytiques p -adiques*, Publications mathématiques de l’IHÉS., tome 26, (1965), pp. 5-219.
- [Liu08] T. Liu: *On lattices in semi-stable representations: a proof of a conjecture of Breuil*, Compositio Mathematica, Vol. 144, Issue 1, pp. 61-88, (2008).
- [LZ17] R. Liu, X. Zhu: *Rigidity and a Riemann-Hilbert correspondence for p -adic local systems*, Inv. Math. 207, pp. 291-343 (2017).
- [MT20] M. Morrow and T. Tsuji: *Generalised representations as q -connections in integral p -adic Hodge theory*, arXiv:2010.04059v2, (2020).
- [MW21a] Y. Min and Y. Wang: *Relative (φ, Γ) -modules and prismatic F -crystals*, arXiv:2110.06076, (2021).
- [MW21b] Y. Min and Y. Wang: *On the Hodge–Tate prismatic crystals over \mathcal{O}_K* , arXiv:2112.10140. (2021).
- [Og22] A. Ogus: *Crystalline prisms: Reflection on the present and the past*, arxiv: 2204.06621, (2022).
- [Pan20] , L. Pan: *On locally analytic vectors of the completed cohomology of modular curves*, Forum of Mathematics, Pi , Volume 10, (2022), e7.
- [Pet20] A. Petrov : *Geometrically irreducible p -adic local systems are de Rham up to a twist*, arxiv:2012.23372v3, (2020). to appear in Duke Math. J..
- [Por22] G. Porat : *Locally analytic vector bundles on the Fargues-Fontaine curve*, arxiv.2204.11354v2, (2022).
- [Sch12] P. Scholze: *Perfectoid spaces*, Publ.math.IHES 116, 245–313 (2012).
- [Sch13] P. Scholze: *p -adic Hodge theory for rigid-analytic varieties*, Forum of Mathematics, Pi, 1, e1, 77 pages, (2013).
- [Sch17] P. Scholze: *Étale cohomology of diamonds*, arXiv:1709.07343v2, (2017).
- [Sen80] S. Sen : *Continuous cohomology and p -adic Galois representations* Inventiones math. 62, 89-116 (1980).
- [Shi18] K. Shimizu : *Constancy of generalized Hodge-Tate weights of a local system* Compositio Math. 154, 2606-2642 (2018).
- [Sim92] C.T. Simpson: *Higgs bundles and local systems*, Publ. math. de l’IHÉS 75, pp. 5-95 (1992).
- [SW20] P. Scholze and J. Weinstein: *Berkeley lectures on p -adic geometry*, Princeton University Press, Princeton, NJ, (2020).
- [Tam15] G. Tamme: *On an analytic version of Lazard’s isomorphism*, Algebra Number Theory 9 (4), 937-956, (2015).
- [Tang22] L. Tang: *Syntomic cycle classes and prismatic Poincaré duality*, arxiv:2210.14279.
- [Tian21] Y. Tian: *Finiteness and duality for the cohomology of prismatic crystals*, arXiv:2109.00801v1, (2021).
- [Tsu18] T. Tsuji: *Notes on the local p -adic Simpson correspondence*, Math. Ann. 371, 795–881, (2018).
- [Wang17] X. Wang: *Weight elimination in two dimensions when $p = 2$* , arXiv:1711.09035, (2017).
- [Wang21] Y. Wang: *A p -adic Simpson correspondence for rigid analytic varieties*, arxiv:2101.00263v3, (2021).
- [Wei04] C.A. Weibel: *An introduction to homological algebra*, Camb. Stu. in. Adv. Math. (2004).
- [Wu21] Z. Wu: *Galois representations, (φ, Γ) -modules and prismatic F -crystals*, arXiv:2104.12105v3, (2021).
- [Xu17] D. Xu: *Parallel transport and the p -adic Simpson correspondence*, Forum of Mathematics, Sigma, Vol. 5 (2017), e13, 127 pages.
- [Xu22] D. Xu: *Parallel transport for Higgs bundles over p -adic curves*, arXiv:2201.06697, (2022).

Y. MIN, DEPARTMENT OF MATHEMATICS, IMPERIAL COLLEGE LONDON, LONDON SW7 2RH, UK.
Email address: y.min@imperial.ac.uk

Y. WANG, MORNINGSIDE CENTER OF MATHEMATICS NO.505, CHINESE ACADEMY OF SCIENCES, ZHONGGUAN-CUN EAST ROAD 55, BEIJING, 100190, CHINA.
Email address: wangyupeng@amss.ac.cn