A note on composition operators between weighted spaces of smooth functions

Andreas Debrouwere and Lenny Neyt

Abstract For certain weighted locally convex spaces *X* and *Y* of one real variable smooth functions, we characterize the smooth functions $\varphi : \mathbb{R} \to \mathbb{R}$ for which the composition operator $C_{\varphi} : X \to Y$, $f \mapsto f \circ \varphi$ is well-defined and continuous. This problem has been recently considered for X = Y being the space \mathscr{S} of rapidly decreasing smooth functions [1] and the space \mathscr{O}_M of slowly increasing smooth functions [2]. In particular, we recover both these results as well as obtain a characterization for X = Y being the space \mathscr{O}_C of very slowly increasing smooth functions.

1 Introduction

One of the most fundamental questions in the study of composition operators is to characterize when such an operator is well-defined and continuous in terms of its symbol. The goal of this article is to consider this question for weighted locally convex spaces of one real variable smooth functions.

Let $\varphi : \mathbb{R} \to \mathbb{R}$ be smooth. In [1] Galbis and Jordá showed that the composition operator $C_{\varphi} : \mathscr{S} \to \mathscr{S}, f \mapsto f \circ \varphi$, with \mathscr{S} the space of rapidly decreasing smooth functions [3], is well-defined (continuous) if and only if

$$\exists N \in \mathbb{Z}_+ : \sup_{x \in \mathbb{R}} \frac{1 + |x|}{(1 + |\varphi(x)|)^N} < \infty$$

Andreas Debrouwere

Department of Mathematics and Data Science, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium, e-mail: andreas.debrouwere@vub.be

Lenny Neyt

Department of Mathematics: Analysis, Logic and Discrete Mathematics, Ghent University, Krijgslaan 281, 9000 Ghent, Belgium, e-mail: lenny.neyt@UGent.be

Andreas Debrouwere and Lenny Neyt

and

$$\forall p \in \mathbb{Z}_+ \; \exists N \in \mathbb{N} \; : \; \sup_{x \in \mathbb{R}} \frac{|\varphi^{(p)}(x)|}{(1+|\varphi(x)|)^N} < \infty.$$

Albanese et. al [2] proved that the composition operator $C_{\varphi} : \mathcal{O}_M \to \mathcal{O}_M$, with \mathcal{O}_M the space of slowly increasing smooth functions [3], is well-defined (continuous) if and only if $\varphi \in \mathcal{O}_M$. In [2, Remark 2.6] they also pointed out that the corresponding result for the space \mathcal{O}_C of very slowly increasing smooth functions [3] is false, namely, they showed that $\sin(x^2) \notin \mathcal{O}_C$, while, obviously, $\sin x, x^2 \in \mathcal{O}_C$.

Inspired by these results, we study in this article the following general question: Given two weighted locally convex spaces X and Y of smooth functions, when is the composition operator $C_{\varphi} : X \to Y$ well-defined (continuous)? We shall consider this problem for X and Y both being Fréchet spaces, (LF)-spaces, or (PLB)-spaces.

We now state a particular instance of our main result that covers many well-known spaces. We need some preparation. Given a positive continuous function v on \mathbb{R} , we write \mathscr{B}_v^n , $n \in \mathbb{N}$, for the Banach space consisting of all $f \in C^n(\mathbb{R})$ such that

$$||f||_{v,n} = \max_{p \le n} \sup_{x \in \mathbb{R}} \frac{|f^{(p)}(x)|}{v(x)} < \infty.$$

For $v \ge 1$ we consider the following three weighted spaces of smooth functions

$$\mathcal{K}_{v} = \lim_{\substack{N \in \mathbb{N} \\ N \in \mathbb{N}}} \mathcal{B}_{1/v^{N}}^{N},$$
$$\mathcal{O}_{C,v} = \lim_{\substack{N \in \mathbb{N} \\ N \in \mathbb{N}}} \lim_{n \in \mathbb{N}} \mathcal{B}_{v^{N}}^{n},$$
$$\mathcal{O}_{M,v} = \lim_{\substack{n \in \mathbb{N} \\ n \in \mathbb{N}}} \lim_{N \in \mathbb{N}} \mathcal{B}_{v^{N}}^{n}.$$

Theorem 2 below implies the following result:

Theorem 1 Let $v, w : \mathbb{R} \to [1, \infty)$ be continuous functions such that

$$\sup_{x,t\in\mathbb{R}, |t|\leq 1} \frac{v(x+t)}{v^{\lambda}(x)} < \infty \quad and \quad \sup_{x,t\in\mathbb{R}, |t|\leq 1} \frac{w(x+t)}{w^{\mu}(x)} < \infty,$$

for some $\lambda, \mu > 0$. Let $\varphi : \mathbb{R} \to \mathbb{R}$ be smooth. Then,

(I) The following statements are equivalent:

(i) $C_{\varphi}(\mathscr{K}_{v}) \subseteq \mathscr{K}_{w}.$ (ii) $C_{\varphi}: \mathscr{K}_{v} \to \mathscr{K}_{w}$ is continuous. (iii) φ satisfies the following two properties (a) $\exists \lambda > 0$: $\sup_{x \in \mathbb{R}} \frac{w(x)}{v^{\lambda}(\varphi(x))} < \infty.$

(b)
$$\forall p \in \mathbb{Z}_+ \exists \lambda > 0$$
 : $\sup_{x \in \mathbb{R}} \frac{|\varphi^{(p)}(x)|}{v^{\lambda}(\varphi(x))} < \infty.$

(II) The following statements are equivalent:

A note on composition operators between weighted spaces of smooth functions

(i)
$$C_{\varphi}(\mathcal{O}_{C,v}) \subseteq \mathcal{O}_{C,w}$$
.
(ii) $C_{\varphi}: \mathcal{O}_{C,v} \to \mathcal{O}_{C,w}$ is continuous.
(iii) φ satisfies the following two properties
(a) $\exists \mu > 0$: $\sup_{x \in \mathbb{R}} \frac{v(\varphi(x))}{w^{\mu}(x)} < \infty$.
(b) $\forall p, k \in \mathbb{Z}_{+}$: $\sup_{x \in \mathbb{R}} \frac{|\varphi^{(p)}(x)|}{w^{1/k}(x)} < \infty$.

(III) The following statements are equivalent:

$$\begin{array}{l} (i) \ C_{\varphi}(\mathcal{O}_{M,v}) \subseteq \mathcal{O}_{M,w}.\\ (ii) \ C_{\varphi}: \mathcal{O}_{M,v} \to \mathcal{O}_{M,w} \ is \ continuous.\\ (iii) \ \varphi \ satisfies \ the \ following \ two \ properties\\ (a) \ \exists \mu > 0 \ : \ \sup_{x \in \mathbb{R}} \frac{v(\varphi(x))}{w^{\mu}(x)} < \infty.\\ (b) \ \forall p \in \mathbb{Z}_{+} \ \exists \mu > 0 \ : \ \sup_{x \in \mathbb{R}} \frac{|\varphi^{(p)}(x)|}{w^{\mu}(x)} < \infty. \end{array}$$

By setting v(x) = w(x) = 1 + |x| in Theorem 1 we recover the above results about \mathscr{S} and \mathscr{O}_M from [1, 2] as well as the following characterization for the space \mathscr{O}_C of very slowly increasing smooth functions: $C_{\varphi} : \mathscr{O}_C \to \mathscr{O}_C$ is well defined (continuous) if and only if

$$\exists N \in \mathbb{N} : \sup_{x \in \mathbb{R}} \frac{|\varphi(x)|}{(1+|x|)^N} < \infty \quad \text{and} \quad \forall p, k \in \mathbb{Z}_+ : \sup_{x \in \mathbb{R}} \frac{|\varphi^{(p)}(x)|}{(1+|x|)^{1/k}} < \infty.$$

For v = w = 1, Theorem 1 gives the following result for the Fréchet space \mathscr{B} of smooth functions that are bounded together will all their derivatives [3]: $C_{\varphi} : \mathscr{B} \to \mathscr{B}$ is well defined (continuous) if and only if $\varphi' \in \mathscr{B}$. Another interesting choice is $v(x) = w(x) = e^{|x|}$, for which Theorem 2 characterizes composition operators on spaces of exponentially decreasing/increasing smooth functions [4, 5]. We leave it to the reader to explicitly formulate this and other examples.

2 Statement of the main result

A pointwise non-decreasing sequence $V = (v_N)_{N \in \mathbb{N}}$ of positive continuous functions on \mathbb{R} is called a *weight system* if $v_0 \ge 1$ and

$$\forall N \; \exists M \ge N \; : \; \sup_{x,t \in \mathbb{R}, |t| \le 1} \frac{v_N(x+t)}{v_M(x)} < \infty.$$

We shall also make use of the following condition on a weight system $V = (v_N)_{N \in \mathbb{N}}$:

$$\forall N, M \exists K \ge N, M : \sup_{x \in \mathbb{R}} \frac{v_N(x)v_M(x)}{v_K(x)} < \infty.$$
(1)

Example 1 Let $v : \mathbb{R} \to [1, \infty)$ be a continuous function satisfying

$$\sup_{x,t\in\mathbb{R},|t|\leq 1}\frac{v(x+t)}{v^N(x)}<\infty.$$

for some $N \in \mathbb{N}$ (cf. Theorem 1). Then,

$$V_v = (v^N)_{N \in \mathbb{N}}$$

is a weight system satisfying (1).

Recall that for a positive continuous function v on \mathbb{R} and $n \in \mathbb{N}$, we write \mathscr{B}_v^n for the Banach space consisting of all $f \in C^n(\mathbb{R})$ such that

$$||f||_{v,n} = \max_{p \le n} \sup_{x \in \mathbb{R}} \frac{|f^{(p)}(x)|}{v(x)} < \infty.$$

Let $V = (v_N)_{N \in \mathbb{N}}$ be a weight system. We shall be concerned with the following weighted spaces of smooth functions

$$\mathcal{K}_{V} = \lim_{\substack{N \in \mathbb{N} \\ N \in \mathbb{N}}} \mathcal{B}_{1/v_{N}}^{N},$$
$$\mathcal{O}_{C,V} = \lim_{\substack{N \in \mathbb{N} \\ N \in \mathbb{N}}} \lim_{n \in \mathbb{N}} \mathcal{B}_{v_{N}}^{n},$$
$$\mathcal{O}_{M,V} = \lim_{\substack{n \in \mathbb{N} \\ n \in \mathbb{N}}} \lim_{N \in \mathbb{N}} \mathcal{B}_{v_{N}}^{n}.$$

Note that \mathscr{K}_V is a Fréchet space, $\mathscr{O}_{C,V}$ is an (LF)-space, and $\mathscr{O}_{M,V}$ is a (PLB)-space. Furthermore, we have the following continuous inclusions

$$\mathscr{D}(\mathbb{R}) \subset \mathscr{K}_V \subset \mathscr{O}_{C,V} \subset \mathscr{O}_{M,V} \subset C^{\infty}(\mathbb{R}),$$

where $\mathscr{D}(\mathbb{R})$ denotes the space of compactly supported smooth functions. The spaces \mathscr{K}_V were introduced and studied by Gelfand and Shilov [6], while we refer to [7] for more information on the spaces $\mathscr{O}_{C,V}$. For $N, n \in \mathbb{N}$ fixed we will also need the following spaces

$$\mathscr{B}_{v_N} = \underset{n \in \mathbb{N}}{\lim} \mathscr{B}_{v_N}^n, \qquad \mathscr{O}_{M,V}^n = \underset{N \in \mathbb{N}}{\lim} \mathscr{B}_{v_N}^n.$$

The goal of this article is to show the following result.

Theorem 2 Let $V = (v_N)_{N \in \mathbb{N}}$ and $W = (w_M)_{M \in \mathbb{N}}$ be two weight systems and let $\varphi : \mathbb{R} \to \mathbb{R}$ be smooth.

- (I) Suppose that V satisfies (1). The following statements are equivalent:
 - (*i*) $C_{\varphi}(\mathscr{K}_V) \subseteq \mathscr{K}_W$.

A note on composition operators between weighted spaces of smooth functions

(ii)
$$C_{\varphi} : \mathscr{K}_{V} \to \mathscr{K}_{W}$$
 is continuous.
(iii) φ satisfies the following two properties
(a) $\forall M \exists N : \sup_{x \in \mathbb{R}} \frac{w_{M}(x)}{v_{N}(\varphi(x))} < \infty.$
(b) $\forall p \in \mathbb{Z}_{+} \exists N : \sup_{x \in \mathbb{R}} \frac{|\varphi^{(p)}(x)|}{v_{N}(\varphi(x))} < \infty.$

(II) Suppose that W satisfies (1). The following statements are equivalent:

 $\begin{array}{l} (i) \ C_{\varphi}(\mathscr{O}_{C,V}) \subseteq \mathscr{O}_{C,W}. \\ (ii) \ C_{\varphi}: \mathscr{O}_{C,V} \to \mathscr{O}_{C,W} \ is \ continuous. \\ (iii) \ \forall N \ \exists M \ such \ that \ C_{\varphi}: \mathscr{B}_{v_N} \to \mathscr{B}_{w_M} \ is \ continuous. \\ (iv) \ \varphi \ satisfies \ the \ following \ two \ properties \\ (a) \ \forall N \ \exists M \ : \ \sup_{x \in \mathbb{R}} \frac{v_N(\varphi(x))}{w_M(x)} < \infty. \\ (b) \ \exists M \ \forall p, k \in \mathbb{Z}_+ \ : \ \sup_{x \in \mathbb{R}} \frac{|\varphi^{(p)}(x)|}{w_M^{1/k}(x)} < \infty. \end{array}$

(III) Suppose that W satisfies (1). The following statements are equivalent:

$$\begin{array}{l} (i) \ C_{\varphi}(\mathscr{O}_{M,V}) \subseteq \mathscr{O}_{M,W}.\\ (ii) \ C_{\varphi}: \mathscr{O}_{M,V} \to \mathscr{O}_{M,W} \ is \ continuous.\\ (iii) \ C_{\varphi}: \mathscr{O}_{M,V}^{n} \to \mathscr{O}_{M,W}^{n} \ is \ continuous \ for \ all \ n \in \mathbb{N}.\\ (iv) \ \varphi \ satisfies \ the \ following \ two \ properties\\ (a) \ \forall N \ \exists M \ : \ \sup_{x \in \mathbb{R}} \frac{v_{N}(\varphi(x))}{w_{M}(x)} < \infty.\\ (b) \ \forall p \in \mathbb{Z}_{+} \ \exists M \ : \ \sup_{x \in \mathbb{R}} \frac{|\varphi^{(p)}(x)|}{w_{M}(x)} < \infty. \end{array}$$

The proof of Theorem 2 will be given in the next section. The spaces \mathscr{K}_v , $\mathscr{O}_{C,v}$ and $\mathscr{O}_{M,v}$ from the introduction can be written as

$$\mathscr{K}_v = \mathscr{K}_{V_v}, \qquad \mathscr{O}_{C,v} = \mathscr{O}_{C,V_v}, \qquad \mathscr{O}_{M,v} = \mathscr{O}_{M,V_v},$$

where $V_v = (v^N)_{N \in \mathbb{N}}$ is the weight system from Example 1. Hence, Theorem 1 is a direct consequence of Theorem 2 with $V = V_v$ and $W = V_w$.

3 Proof of the main result

Throughout this section we fix a smooth symbol $\varphi : \mathbb{R} \to \mathbb{R}$. We need two lemmas in preparation for the proof of Theorem 2. For $n \in \mathbb{N}$ we set

$$||f||_n = ||f||_{1,n} = \max_{p \le n} \sup_{x \in \mathbb{R}} |f^{(p)}(x)|.$$

Lemma 1 Let v, \tilde{v}, w be three positive continuous functions on \mathbb{R} such that

$$C_0 = \sup_{x,t \in \mathbb{R}, |t| \le 1} \frac{v(x+t)}{\widetilde{v}(x)} < \infty.$$

Let $p, n \in \mathbb{N}$ be such that

$$\|C_{\varphi}(f)\|_{w,p} \le C_1 \|f\|_{\widetilde{v},n}, \qquad \forall f \in \mathscr{D}(\mathbb{R}),$$
(2)

for some $C_1 > 0$. Then,

$$\sup_{x \in \mathbb{R}} \frac{v(\varphi(x))}{w(x)} < \infty, \tag{3}$$

and, if $p \ge 1$, also

$$\sup_{x \in \mathbb{R}} \frac{v(\varphi(x))|\varphi'(x)|^p}{w(x)} < \infty,$$
(4)

and

$$\sup_{x \in \mathbb{R}} \frac{v(\varphi(x))|\varphi^{(p)}(x)|}{w(x)} < \infty.$$
(5)

Proof Given $f \in \mathscr{D}(\mathbb{R})$ with supp $f \subseteq [-1, 1]$, we set $f_x = f(\cdot - \varphi(x))$ for $x \in \mathbb{R}$. Note that

$$\|f_x\|_{\widetilde{v},n} \le \frac{C_0 \|f\|_n}{v(\varphi(x))}, \qquad x \in \mathbb{R}.$$
(6)

We first show (3). Choose $f \in \mathscr{D}(\mathbb{R})$ with supp $f \subseteq [-1, 1]$ such that f(0) = 1. For all $x \in \mathbb{R}$ it holds that

$$\|C_{\varphi}(f_x)\|_{w,p} \geq \frac{C_{\varphi}(f_x)(x)}{w(x)} = \frac{1}{w(x)}.$$

Hence, by (2) and (6), we obtain that

$$\frac{v(\varphi(x))}{w(x)} \le C_0 C_1 ||f||_n, \qquad \forall x \in \mathbb{R}.$$

Now assume that $p \ge 1$. We prove (4). Choose $f \in \mathscr{D}(\mathbb{R})$ with supp $f \subseteq [-1, 1]$ such that $f^{(j)}(0) = 0$ for j = 1, ..., p - 1 and $f^{(p)}(0) = 1$. Faà di Bruno's formula implies that for all $x \in \mathbb{R}$

$$\|C_{\varphi}(f_x)\|_{w,p} \ge \frac{|C_{\varphi}(f_x)^{(p)}(x)|}{w(x)} = \frac{|\varphi'(x)|^p}{w(x)}.$$

Similarly as in the proof of (3), the result now follows from (2) and (6). Finally, we show (5). Choose $f \in \mathcal{D}(\mathbb{R})$ with supp $f \subseteq [-1, 1]$ such that f'(0) = 1 and $f^{(j)}(0) = 0$ for j = 2, ..., p. Faà di Bruno's formula implies that for all $x \in \mathbb{R}$

$$\|C_{\varphi}(f_x)\|_{w,p} \ge \frac{|C_{\varphi}(f_x)^{(p)}(x)|}{w(x)} = \frac{|\varphi^{(p)}(x)|}{w(x)}.$$

As before, the result is now a consequence of (2) and (6).

Lemma 2 Let v and w be positive continuous functions on \mathbb{R} . Then,

(*i*) *If*

$$\sup_{x \in \mathbb{R}} \frac{v(\varphi(x))}{w(x)} < \infty$$

then $C_{\varphi} : \mathscr{B}_{v}^{0} \to \mathscr{B}_{w}^{0}$ is well-defined and continuous. (ii) Let $n \in \mathbb{Z}_{+}$. If

$$\sup_{x \in \mathbb{R}} \frac{v(\varphi(x))}{w(x)} \prod_{p=1}^{n} |\varphi^{(p)}(x)|^{k_p} < \infty$$

for all $(k_1, \ldots, k_n) \in \mathbb{N}^n$ with $\sum_{j=1}^p jk_j \leq p$ for all $p = 1, \ldots, n$, then C_{φ} : $\mathscr{B}_{v}^{n} \to \mathscr{B}_{w}^{n}$ is well-defined and continuous.

Proof (i) Obvious.

(*ii*) This is a direct consequence of (*i*) and Faà di Bruno's formula.

Proof (of Theorem 2) (I) (i) \Rightarrow (ii): Since $C_{\varphi} : C^{\infty}(\mathbb{R}) \to C^{\infty}(\mathbb{R})$ is continuous, this follows from the closed graph theorem for Fréchet spaces. $(ii) \Rightarrow (iii)$: For all $p, M \in \mathbb{N}$ there are $n, L \in \mathbb{N}$ such that

$$\|C_{\varphi}(f)\|_{p,1/w_M} \le C \|f\|_{n,1/v_L}, \qquad \forall f \in \mathscr{K}_V.$$

Choose $N \ge L$ such that

$$\sup_{x,t \in \mathbb{R}, |t| \le 1} \frac{v_L(x+t)}{v_N(x)} = \sup_{x,t \in \mathbb{R}, |t| \le 1} \frac{1/v_N(x+t)}{1/v_L(x)} < \infty.$$

Lemma 1 with $w = 1/w_M$, $v = 1/v_N$ and $\tilde{v} = 1/v_L$ yields that

$$\sup_{x \in \mathbb{R}} \frac{w_M(x)}{v_N(\varphi(x))} < \infty$$

and (recall that $w_M \ge 1$)

$$\sup_{x \in \mathbb{R}} \frac{|\varphi^{(p)}(x)|}{v_N(\varphi(x))} < \infty.$$

 $(iii) \Rightarrow (i)$: As V satisfies (1), this follows from Lemma 2.

(11) (i) \Rightarrow (ii): Since C_{φ} : $C^{\infty}(\mathbb{R}) \rightarrow C^{\infty}(\mathbb{R})$ is continuous, this follows from De Wilde's closed graph theorem.

 $(ii) \Rightarrow (iii)$: This is a consequence of Grothendieck's factorization theorem. $(iii) \Rightarrow (iv)$: Fix an arbitrary $N \in \mathbb{N}$. Choose $L \ge N$ such that

$$\sup_{x,t\in\mathbb{R},|t|\leq 1}\frac{v_N(x+t)}{v_L(x)}<\infty.$$

Choose $K \in \mathbb{N}$ such that $C_{\varphi} : \mathscr{B}_{v_L} \to \mathscr{B}_{w_K}$ is continuous. For all $m \in \mathbb{Z}_+$ there are $n \in \mathbb{Z}_+$ and C > 0 such that

$$\|C_{\varphi}(f)\|_{m,w_{K}} \leq C \|f\|_{n,v_{L}}, \qquad \forall f \in \mathscr{B}_{v_{L}}.$$

Lemma 1 with $w = w_K$, $v = v_N$ and $\tilde{v} = v_L$ yields that

$$\sup_{x \in \mathbb{R}} \frac{v_N(\varphi(x))}{w_K(x)} < \infty \tag{7}$$

and (recall that $v_N \ge 1$)

$$\sup_{x \in \mathbb{R}} \frac{|\varphi'(x)|}{w_K^{1/m}(x)} < \infty \quad \text{and} \quad \sup_{x \in \mathbb{R}} \frac{|\varphi^{(m)}(x)|}{w_K(x)} < \infty.$$
(8)

Equation (7) shows (*a*). We now prove (*b*). To this end, we will make use of the following Landau-Kolmogorov type inequality due to Gorny [8]: For all $j \le m \in \mathbb{Z}_+$ there is C > 0 such that

$$\|g^{(j)}\| \le C \|g\|^{1-j/m} \left(\max\{\|g\|, \|g^{(m)}\|\} \right)^{j/m}, \qquad \forall g \in C^{\infty}([-1, 1]), \quad (9)$$

where $\|\cdot\|$ denotes the sup-norm on [-1, 1]. Choose $M \ge K$ such that

$$\sup_{x,t\in\mathbb{R},|t|\leq 1}\frac{w_K(x+t)}{w_M(x)}<\infty.$$

Let $p, k \in \mathbb{Z}_+$ and $x \in \mathbb{R}$ be arbitrary. Equation (8) yields that for all $m \in \mathbb{Z}_+$ there is C > 0 such that

$$\|\varphi'(x+\cdot)\| \le Cw_M^{1/m}(x)$$
 and $\|\varphi^{(m)}(x+\cdot)\| \le Cw_M(x).$

By applying (9) to $g = \varphi'(x + \cdot)$ and $m \ge p$ such that

$$\left(1 - \frac{p-1}{m}\right)\frac{1}{m} + \frac{p-1}{m} \le \frac{1}{k}$$

we find that (recall that $w_M \ge 1$)

$$\begin{split} |\varphi^{(p)}(x)| &\leq \|\varphi^{(p)}(x+\cdot)\| \\ &\leq C \|\varphi'(x+\cdot)\|^{1-(p-1)/m} \left(\max\{\|\varphi'\|, \|\varphi^{(m+1)}\|\} \right)^{(p-1)/m} \\ &\leq C' w_M^{1/k}(x). \end{split}$$

 $(iv) \Rightarrow (i)$: As W satisfies (1), this follows from Lemma 2.

(*III*) (*iii*) \Rightarrow (*ii*) \Rightarrow (*ii*): Obvious. (*i*) \Rightarrow (*iv*): Fix arbitrary $p \in \mathbb{Z}_+$ and $N \in \mathbb{N}$. Choose $L \ge N$ such that A note on composition operators between weighted spaces of smooth functions

$$\sup_{x,t\in\mathbb{R},|t|\leq 1}\frac{v_N(x+t)}{v_L(x)}<\infty.$$

Since $\mathscr{B}_{v_L} \subset \mathscr{O}_{M,V}$ and $\mathscr{O}_{M,W} \subset \mathscr{O}_{M,W}^p$, we obtain that $C_{\varphi}(\mathscr{B}_{v_L}) \subset \mathscr{O}_{M,W}^p$. As $C_{\varphi} : C^{\infty}(\mathbb{R}) \to C^p(\mathbb{R})$ is continuous, De Wilde's closed graph theorem implies that $C_{\varphi} : \mathscr{B}_{v_L} \to \mathscr{O}_{M,W}^p$ is continuous. Grothendieck's factorization theorem yields that there is $M \in \mathbb{N}$ such that $C_{\varphi} : \mathscr{B}_{v_L} \to \mathscr{B}_{w_M}^p$ is well-defined and continuous, and thus that

$$\|C_{\varphi}(f)\|_{p,w_{M}} \le C \|f\|_{n,v_{L}}, \qquad \forall f \in \mathscr{B}_{v_{L}},$$

for some $n \in \mathbb{N}$ and C > 0. Lemma 1 with $w = w_M$, $v = v_N$ and $\tilde{v} = v_L$ yields that

$$\sup_{x \in \mathbb{R}} \frac{v_N(\varphi(x))}{w_M(x)} < \infty$$

and (recall that $v_N \ge 1$)

$$\sup_{x\in\mathbb{R}}\frac{|\varphi^{(p)}(x)|}{w_M(x)}<\infty.$$

 $(iv) \Rightarrow (iii)$: As W satisfies (1), this follows from Lemma 2.

Acknowledgements L. Neyt gratefully acknowledges support by FWO-Vlaanderen through the postdoctoral grant 12ZG921N.

References

- Galbis, A., Jordá, E.: Composition operators on the Schwartz space. Rev. Mat. Iberoam. 34, 397–412 (2018).
- 2. Albanese, A.A., Jordá, E. Mele, C.: Dynamics of composition operators on function spaces defined by local and global properties. Preprint. Available on arXiv:2112.08699.
- 3. Schwartz, L.: Théorie des distributions. Hermann, Paris (1966).
- Hasumi, M.: Note on the *n*-dimensional tempered ultra-distributions. Tôhoku Math. J. 13, 94–104 (1961).
- 5. Zieleźny, Z.: On the space of convolution operators in \mathcal{K}'_1 . Studia Math. **31**, 111–124 (1968).
- Gel'fand, I.M., Shilov, G.E.: Generalized functions. Vol. 2: Spaces of fundamental and generalized functions. Academic Press, New York-London (1968).
- 7. Debrouwere, A., Vindas, J.: Topological properties of convolutor spaces via the short-time Fourier transform. Trans. Amer. Math. Soc. **374**, 829–861 (2021).
- Gorny, A.: Contribution à l'étude des fonctions dérivables d'une variable réelle. Acta Math. 71, 317–358 (1939).