

One-Dimensional Quantum Systems with Ground-State of Jastrow Form Are Integrable

Jing Yang^{1,*} and Adolfo del Campo^{1,2,†}

¹*Department of Physics and Materials Science, University of Luxembourg, L-1511 Luxembourg, Luxembourg*

²*Donostia International Physics Center, E-20018 San Sebastián, Spain*

(Dated: January 21, 2022)

The exchange operator formalism (EOF) describes many-body integrable systems using phase-space variables in terms of which the Hamiltonian takes a decoupled form. We establish an equivalence between models described by EOF and the complete infinite family of parent Hamiltonians (PHJ) describing quantum many-body models with ground-states of Jastrow form. This makes it possible to identify the invariants of motion for any model in the PHJ family and establish its integrability, even in the presence of an external potential. Using this construction we establish the integrability of the long-range Lieb-Liniger model, describing bosons in a harmonic trap and subject to contact and Coulomb interactions in one dimension.

Integrability in both classical and quantum many-body systems is associated with the existence of conserved quantities. At the quantum level, the latter correspond to operators that commute with the system Hamiltonian and govern the nonequilibrium dynamics and thermalization of a system in isolation [1, 2]. Over the last decades, an increasing number of integrable models has been realized in the laboratory, prompting their use as a test-bed for quantum many-body physics, statistical mechanics, and nonequilibrium phenomena [3, 4].

In a specific system, integrability may be proven by explicitly finding the set of conserved quantities. In one spatial dimension, this is possible in systems that are exactly solved using Bethe ansatz, which posits that any quantum state can be described by a combination of plane waves. The integrals of motion are the corresponding quasimomenta, known as Bethe roots or rapidities, that serve as “good” quantum numbers [5, 6]. An alternative framework is the exchange operator formalism [7, 8], in which the Hamiltonian of the quantum system admits a decoupled form in terms of generalized momenta, which readily allow for the identification of integrals of motion. An encompassing notion of quantum integrability relies on scattering without diffraction, encoded in the Yang-Baxter equation [9–11], when collisions between particles can be described exclusively as a sequence of two-body scattering events. The system is then solvable by algebraic Bethe ansatz, i.e., using the quantum inverse scattering method. Integrals of motion may then be identified taking derivative of a transfer matrix [11] or invoking the asymptotic Bethe ansatz for systems in the continuum [12, 13]. While a definite notion of quantum integrability remains under debate, many of these approaches are closely interrelated [6, 14]. In particular, the exchange operator formalism is related to the Yang-Baxter equation and asymptotic Bethe ansatz [15, 16].

An important class of quantum systems is characterized by a ground-state of (Bijl-Dingle-) Jastrow form, in which the wave function is simply the pairwise product of a pair function [17–19]. This facilitates the computation of correlation functions in these systems [12]. The family of parent Hamiltonians with Jastrow wave functions (PHJ, for short) can be determined by solving an inverse problem: by acting with the kinetic energy operator in the ground-state wave function, one can recast the resulting terms in the form of a many-body

Schrödinger equation, thus identifying the parent Hamiltonian. This approach has its roots in the early works by Calogero and Sutherland [20–22]. It has been extended in a number of ways [23, 24] and by now, for identical particles without internal degrees of freedom, the complete family of PHJ is known both in one and higher spatial dimensions, provided that the ground-state wave function includes at most the product of one-particle and two-particle functions [25, 26]. The corresponding Hamiltonians generally contain two-body and three-body interactions. It was shown by Kane et al [27] that the three-body contribution does not affect the low-energy physics. Further, the conditions for the three-body term to vanish or reduce to a constant have been long-established in the homogeneous case, in the absence of an external potential [12, 28, 29].

Paradigmatic instances of PHJ are integrable. Hard-core bosons in the Tonks-Girardeau regime realized in the laboratory with ultracold gases [30, 31], have ground state of Jastrow form [32–34] and are integrable, being related to noninteracting fermions via the Bose-Fermi duality [3, 32, 35]. The Calogero-Sutherland model with a Jastrow ground state has a harmonic spectrum, it can be mapped to a set of independent harmonic oscillators [36–38] and satisfies the asymptotic Bethe ansatz [12, 13]. Similarly, the attractive Lieb-Liniger (LL) model of bosons subject to contact interactions, used to describe ultracold gases in tight waveguides [39, 40], has a bright quantum soliton as Jastrow ground-state [41]. This system is solvable by coordinate Bethe ansatz, which yields the Bethe roots as integrals of motion [5, 6, 42, 43].

One may thus wonder the extent to which the ground-state correlations can determine the complete integrability of the system, and what are the required conditions for this to be the case. In this Letter, we show that the complete family of one-dimensional many-body quantum models with ground-state of Jastrow form is integrable. To this end, we first establish the equivalence between this family and models described by the exchange operator formalism. In doing so, we identify explicitly the integrals of motion. Our construction holds in the presence of an external potential, which allows us to show the integrability of the long-range Lieb-Liniger model, describing bosons confined in a harmonic trap and subject to both contact and Coulomb interactions in one spatial dimension [25, 44].

Exchange operator formalism (EOF).— Consider the family of one-dimensional systems of identical particles without internal degrees of freedom. It will prove useful to consider those models subject to pair-wise interactions that are possibly supplemented with three-body interactions. In this context, EOF is a powerful framework due to Polychronakos that explicitly exhibits the integrability of a many-body quantum system in one spatial dimension [7, 8]. Its application has been particularly fruitful in the family of Calogero-Sutherland-Moser systems in which one-dimensional identical particles are subject to two-body inverse-square interactions and their generalizations [12, 20, 21, 23, 45, 46], as discussed in [7, 8]. The pairwise potential has in this case the same scaling dimension as the kinetic energy operator.

Let M_{ij} denote the exchange operator, which exchanges the positions of two particles labelled by i and j , respectively. This operator is Hermitian, idempotent $M_{ij}^2 = \mathbb{I}$ and symmetric with respect to the indices, i.e., $M_{ij} = M_{ji}$. For any 1-body operator $A_j \equiv A(x_j)$, it obeys the relations $M_{ij}A_j = A_iM_{ij}$ and $M_{ij}A_k = A_kM_{ij}$ for distinct i, j, k [7, 8, 47]. Note that when for spinless identical particles, M_{ij} can be identified with the permutation of two particles. In terms of the canonical position and momentum coordinates, x_i and $p_j = -i\hbar\partial/\partial x_j$, one can introduce the generalized momenta

$$\pi_i = p_i + i \sum_{j \neq i} V_{ij} M_{ij}, \quad (1)$$

for particles $j = 1, \dots, N$. The so-called prepotential function $V_{ij} = V(x_i - x_j)$ should be antisymmetric (i.e., $V_{ij} = -V_{ji}$) to guarantee the Hermiticity of the generalized momenta. Using the latter, one can construct a permutation-invariant quantities $I_n \equiv \sum_i \pi_i^n$. In particular, I_2 is quadratic in p_i 's, which resembles the Hamiltonian of many-body systems. To describe states of N particles, consider the tensor product of the single-particle Hilbert space \mathcal{H} , i.e., $\mathcal{H}^{\otimes N}$. For indistinguishable particles, states are restricted to the bosonic or fermionic subspaces of $\mathcal{H}^{\otimes N}$, denoted as \mathcal{H}_ζ with $\zeta = +1$ for spinless bosons and $\zeta = -1$ for spinless fermions. We define the projector \mathcal{P}_ζ onto \mathcal{H}_ζ as [48] $\mathcal{P}_\zeta\psi(x_1, x_2, \dots, x_N) = \frac{1}{N!} \sum_\sigma \zeta^\sigma \psi(x_{\sigma_1}, x_{\sigma_2}, \dots, x_{\sigma_N})$, where σ denotes a permutation of the tuple $(1, 2, \dots, N)$. Projecting I_2 onto the subspace \mathcal{H}_ζ and using

$$M_{ij}\mathcal{P}_\zeta = \mathcal{P}_\zeta M_{ij} = \zeta\mathcal{P}_\zeta, \quad (2)$$

we obtain $\mathcal{P}_\zeta I_2 \mathcal{P}_\zeta / (2m) = \mathcal{P}_\zeta H_0 \mathcal{P}_\zeta$, where H_0 is the translation-invariant quantum many-body Hamiltonian defined as follows

$$H_0 = \sum_i \frac{p_i^2}{2m} + \frac{1}{m} \left[\sum_{i < j} (\zeta \hbar V'_{ij} + V_{ij}^2) - \sum_{i < j < k} V_{ijk} \right], \quad (3)$$

where $V_{ijk} = V_{ij}V_{jk} + V_{jk}V_{ki} + V_{ki}V_{ij}$ is fully symmetric and a prime denotes the spatial derivative. It will play an important role in proving the integrability of the family of Hamiltonians generated by EOF and PHJ. Specific choices of the prepotential function $V(x)$ gives rise to well-known models. For

$V(x) = \lambda/x$, V_{ijk} vanishes by permutation symmetry and one recovers the Hamiltonian of identical particles with inverse-square interactions [12, 20]. For $V(x) = \lambda \cot(ax)$, V_{ijk} is constant and H_0 involves the inverse sine square potentials. The case $V(x) = c \operatorname{sgn}(x)$, corresponding V_{ijk} being a negative constant, gives rise to the celebrated Lieb-Liniger (LL) model [42, 43] describing ultracold gases in tight-waveguides [3, 39]. For all these cases where V_{ijk} vanishes or is constant, I_n commute with each other. As the system Hamiltonian coincides with I_2 on the bosonic or fermionic sector, the set of I_n can be identified as invariants of motion, i.e., $[I_n, I_m] = 0$. We note that all the models that have been shown to be integrable by the EOF in Ref. [7] happen to have a ground-state wave function of Jastrow form, which we discuss next.

Parent Hamiltonians with Jastrow ground-state.— Consider a homogenous one-dimensional many-body quantum system described by a ground state of Jastrow form [17–19],

$$\Phi_0(x_1, \dots, x_N) = \prod_{i < j} f_{ij}, \quad (4)$$

this is, the pairwise product of the correlation function $f_{ij} = f(x_i - x_j)$ [12]. In the “beautiful models” that concern us here, quantum statistics is encoded in the symmetry of $f(x)$ which is an even function for bosons and odd for fermions, i.e., without resorting to the use of permanents or determinants. The case of one dimensional anyons can similarly be taken into account by including a phase factor θ , i.e., $f(x) = e^{-i\theta} f(-x)$ [49–51]. The complete family of PHJ of ground-state of the Jastrow form (4) has been identified in one spatial dimension [25] and includes paradigmatic models such as the LL gas with contact interactions [42, 43] and the rational Calogero-Sutherland model with inverse-square interactions [20, 21], as well as the recently introduced long-range LL model [44], among other examples. For a given choice of f , the parent Hamiltonian H_0 takes the form

$$H_0 = \sum_i \frac{p_i^2}{2m} + \frac{\hbar^2}{m} \left[\sum_{i < j} \frac{f''_{ij}}{f_{ij}} + \sum_{i < j < k} \left(\frac{f'_{ij} f'_{ik}}{f_{ij} f_{ik}} - \frac{f'_{ij} f'_{jk}}{f_{ij} f_{jk}} + \frac{f'_{ik} f'_{jk}}{f_{ik} f_{jk}} \right) \right]. \quad (5)$$

Here, f' and f'' denote the first and second spatial derivatives of f , respectively. The explicit expressions for this Hamiltonian directly follow from evaluating the Laplacian on the Jastrow wave function (4) and recasting all resulting terms in the form of a Schrödinger equation.

Equivalence of EOF and PHJ for spinless indistinguishable particles.— We now establish the correspondence between EOF and PHJ for spinless identical particles. Comparing EOF and PHJ Hamiltonians in Eqs. (3) and (5), we find that for the two-body terms to be equal it is required that $\hbar^2 f''(x_{ij})/f(x_{ij}) = \zeta \hbar V'_{ij} + V_{ij}^2$. Thus, the prepotential reads

$$V_{ij} = \zeta \hbar \frac{d}{dx_{ij}} \log(f_{ij}) = \zeta \hbar \frac{f'_{ij}}{f_{ij}}. \quad (6)$$

Independently of whether the pair function is symmetric or antisymmetric, its logarithmic derivative is guaranteed to be odd

$f'_{ij}/f_{ij} = -f'_{ji}/f_{ji}$. Thus, this property holds for spinless bosons and fermions. The antisymmetry of the prepotential in Eq. (6) guarantees the Hermiticity condition of the associated generalized momenta in the EOF,

$$\pi_i = p_i + i\zeta\hbar \sum_{j \neq i} \frac{f'_{ij}}{f_{ij}} M_{ij}. \quad (7)$$

The prepotential in Eq. (6) further ensures the equivalence of the three-body interaction in the EOF and the PHJ. Thus, any spinless system described by EOF, as in Eq. (3), has a ground-state of Jastrow form with a pair function $f_{ij} = \exp[\int^{x_{ij}} dy V(y)/(\zeta\hbar)]$. Conversely, the complete infinite family of PHJ can be recast in the EOF provided (6) is satisfied. This makes it possible to identify the class of PHJ that is integrable as we shall see later.

Embedding in an external potential.— The embedding of a system described by EOF in an external potential is known in the case of a harmonic trap [7]. Can we extend the embedding of a homogeneous system to the case of an arbitrary anharmonic trapping potential? To this end, we draw inspirations from supersymmetric quantum mechanics [52] and introduce the one-body superpotential $W_i \equiv W(x_i)$ in terms of which the external trapping potential U_i will be identified. We thus consider a many-particle generalization of supersymmetric quantum mechanics and define the operators

$$a_i = \frac{\pi_i}{\sqrt{2m}} - iW_i, \quad a_i^\dagger = \frac{\pi_i}{\sqrt{2m}} + iW_i, \quad (8)$$

and the permutation-invariant quantities $\tilde{I}_n \equiv \sum_i h_i^n$, where $h_i \equiv a_i^\dagger a_i$. Projecting \tilde{I}_1 onto \mathcal{H}_ζ , we find $\mathcal{P}_\zeta \tilde{I}_1 \mathcal{P}_\zeta = \mathcal{P}_\zeta H \mathcal{P}_\zeta$, where H is the Hamiltonian of the system in the presence of the trap, i.e.,

$$H = H_0 + \sum_i U_i - \zeta \sqrt{\frac{2}{m}} \sum_{i < j} V_{ij} (W_i - W_j), \quad (9)$$

with the external potential U_i being determined by the Riccati equation

$$U_i = W_i^2 - \frac{\hbar}{\sqrt{2m}} W_i'. \quad (10)$$

As a familiar example, when H_0 is the homogeneous Calogero model with inverse-square interactions [20] and $W_i = \sqrt{m/2}\omega x_i$, Eq. (9) reduces to the Calogero-Sutherland model [21, 45] embedded in a harmonic trap.

In PHJ, the ground-state wave functions is not limited to homogeneous form (4), but also includes more general ground-states

$$\Psi_0 = \prod_{i < j} f_{ij} \prod_i \exp(v_i) = \Phi_0 \prod_i \exp(v_i), \quad (11)$$

where the one-body function $v_i = v(x_i)$ accounts for the role of the external potential $U_i = U(x_i)$ that breaks translational

invariance [25]. Specifically, if H_0 is the parent hamiltonian of Φ_0 in Eq. (4), then Ψ_0 has the parent Hamiltonian

$$H = H_0 + \sum_i U_i + \frac{\hbar^2}{m} \sum_{i < j} (v_i' - v_j') \frac{f'_{ij}}{f_{ij}}, \quad (12)$$

with the one-body local external potential U_i given in terms of the function v_i by

$$U_i = \frac{\hbar^2}{m} [(v_i')^2 + v_i'']. \quad (13)$$

As in the homogenous case, the explicit forms of H and U_i are readily obtained by computing the action of the canonical kinetic energy operator $\sum_i p_i^2/(2m)$ on the Jastrow wave function Ψ_0 and recasting the resulting terms as a Schrödinger equation. As a result, the Hamiltonian H includes the external potential U_i and an additional pairwise (two-body) potential which is generally of long-range character.

The requirement of the equivalence between EOF and PHJ imposes two constraints that the one-body and additional long-range potentials in Eqs. (9, 12) must coincide with each other, respectively. Remarkably, these two constraints are compatible with each other. Comparing Eq. (10) and Eq. (13), the superpotential W_i and the function v_i entering the one-body function of the Jastrow form are related by

$$W_i = -\frac{\hbar}{\sqrt{2m}} v_i'. \quad (14)$$

Upon substituting Eq. (14) into Eqs. (9, 12), we find that the additional long-range potentials coincide, given the correspondence Eq. (6) is identified. The ground state of the Hamiltonian with the external potential in terms of the prepotential and the superpotential is

$$\Psi_0 = \exp\left[-\frac{\sqrt{2m}}{\hbar} \sum_i \int^{x_i} W(y) dy\right] \prod_{i < j} \exp\left[\frac{\int^{x_{ij}} dy V(y)}{\zeta\hbar}\right]. \quad (15)$$

This establishes the equivalence between EOF and PHJ in the presence of external potential.

Integrability via projection formalism.— As emphasized, a common definition of quantum integrability relies on scattering without diffraction, that is encoded the Yang-Baxter equations [12]. For quantum systems with classical analog, which is the case here, one can define quantum integrability by promoting the Poisson bracket into commutators in the definition of classical integrability. Polychronakos [7] pursued along this line and showed that $I_n \equiv \sum_i \pi_i^n$ become integrals of motion, i.e., $[I_n, I_m] = 0, \forall n, m$, in the restricted case in which V_{ijk} vanishes or is constant. The latter applies to the Calogero models [7], in which EOF can be further related to Yang-Baxter integrability [15]. Having shown that any spinless model described by EOF is a PHJ with a Jastrow ground state, we next establish the integrability of the complete family of PHJ models, i.e., without restrictions on the three-body potential V_{ijk} or the external potential U_i .

Note that any physical observable O for spinless indistinguishable particles must be permutation invariant, i.e., $O(x_1, x_2, \dots, x_N) = O(x_{\sigma_1}, x_{\sigma_2}, \dots, x_{\sigma_N}), \forall$ permutation σ . As a consequence,

$$[\mathcal{P}_\zeta, O] = 0, \quad (16)$$

which can be easily checked that by acting on any wave function in $\mathcal{H}^{\otimes N}$ [47]. Eq. (16) implies the a permutation-invariant observable is block diagonal on \mathcal{H}_ζ and its orthogonal complement. We define an observable is local if it only involves derivatives with respect to the coordinates up to a finite order. Then permutation invariance and locality implies that if a permutation-invariant and local observable O vanishes on \mathcal{H}_ζ , then it also vanishes on the full Hilbert space $\mathcal{H}^{\otimes N}$. That is [53],

$$\mathcal{P}_\zeta O \mathcal{P}_\zeta = \mathcal{P}_\zeta O = O \mathcal{P}_\zeta = 0 \iff O = 0, \quad (17)$$

for a permutation-invariant and local observable O .

Eqs. (16, 17) lead to the following theorem regarding the commutators of two permutation-invariant observables, which is extremely useful in proving integrability.

Theorem 1. *For two permutation-invariant and local observable O_n and O_m , the following three conditions are equivalent to each other (i) $[O_n, O_m] = 0$ (ii) $\mathcal{P}_\zeta [O_n, O_m] \mathcal{P}_\zeta = 0$ (iii) $[\mathcal{P}_\zeta O_n \mathcal{P}_\zeta, \mathcal{P}_\zeta O_m \mathcal{P}_\zeta] = 0$.*

The equivalence between (i) and (ii) is a consequence of Eq. (17). The equivalence between (ii) and (iii) follows from

$$\begin{aligned} \mathcal{P}_\zeta [O_n, O_m] \mathcal{P}_\zeta &= \mathcal{P}_\zeta O_n O_m \mathcal{P}_\zeta^2 - \mathcal{P}_\zeta O_m O_n \mathcal{P}_\zeta^2 \\ &= \mathcal{P}_\zeta O_n \mathcal{P}_\zeta O_m \mathcal{P}_\zeta - \mathcal{P}_\zeta O_m \mathcal{P}_\zeta O_n \mathcal{P}_\zeta \\ &= [\mathcal{P}_\zeta O_n \mathcal{P}_\zeta, \mathcal{P}_\zeta O_m \mathcal{P}_\zeta], \end{aligned}$$

where we have used Eq. (16).

Theorem 2. *Both the quantum mechanical homogenous model (3) and the inhomogeneous model (9) generated in EOF are integrable, with the integral of motion being I_n for the homogenous model and \tilde{I}_n for the inhomogeneous model.*

To prove Theorem 2, let us first observe a very interesting property due to the projection \mathcal{P}_ζ and the exchange operator M_{ij} . Although the generalized momentum π_i involves N degrees of freedom due to the prepotential term, when it is multiplied by M_{ij} from the left, it still satisfies the exchange rule for 1-body operators [47]. As a consequence,

$$\mathcal{P}_\zeta \pi_i^n \mathcal{P}_\zeta = \mathcal{P}_\zeta M_{ij}^2 \pi_i^n \mathcal{P}_\zeta = \mathcal{P}_\zeta M_{ij} \pi_i^n M_{ij} \mathcal{P}_\zeta = \mathcal{P}_\zeta \pi_j^n \mathcal{P}_\zeta. \quad (18)$$

A similar equation also holds for h_i . For a more general version of this identity, see Lemma 1 in [47].

On the other hand, since the integrals of motions I_n and \tilde{I}_n are permutation invariant and local, one can reduce their commutativity to condition (iii) in Theorem 1. Using Eq. (18) or the analogous equation for h_i , it follows that the condition (iii)

in Theorem 1 is satisfied, with $O_n = I_n$ or $O_n = \tilde{I}_n$. This concludes the proof of the integrability of the Hamiltonians (3) and (9).

A few comments are in order. First, since we have proved the equivalence between EOF and PHJ, the Hamiltonians (5) and (12) are therefore also integrable. Second, it is possible to build integrals of motions for families of classical models generated by the EOF and PHJ according to the quantum-classical correspondence. One can expand the powers in I_n and \tilde{I}_n and compute $\mathcal{P}_\zeta I_n \mathcal{P}_\zeta$ and $\mathcal{P}_\zeta \tilde{I}_n \mathcal{P}_\zeta$ explicitly with Eq. (2). Then one is left with the expressions $\mathcal{P}_\zeta K_n \mathcal{P}_\zeta$ and $\mathcal{P}_\zeta \tilde{K}_n \mathcal{P}_\zeta$, where K_n and \tilde{K}_n contain only the phase space variables but no exchange operators. In particular, we note that $K_2 = H_0$ and $\tilde{K}_1 = H$. An example of this procedure is described in Sec. III in the Supplemental Material [47], where one obtains H_0 by projecting I_2 onto \mathcal{H}_ζ . According to Theorem 1, K_n 's and \tilde{K}_n 's must commute on the whole Hilbert space $\mathcal{H}^{\otimes N}$, respectively. Transitioning to the classical model, where the commutator is demoted to Poisson brackets, the Poisson brackets of K_n 's and \tilde{K}_n 's must vanish, respectively. Thus, we see that K_n 's and \tilde{K}_n 's are also the integrals of motions for the classical model with Hamiltonians (3, 5) and (9, 12), respectively.

Discussion.— It is worth noting that the Jastrow wave functions Φ_0, Ψ_0 may not be the true eigenstate of the generated Hamiltonian if they cannot be properly normalized. If interaction potential in Eqs. (3, 5) becomes asymptotically constant as particles become far apart then one would expect as $|x_{ij}| \rightarrow \infty$ either $|\Phi_0|, |\Psi_0| \rightarrow 0$ (bound states) or $|\Phi_0|, |\Psi_0| \rightarrow \text{constant}$ (scattering states). If interaction potential or Eqs. (9, 12) tends to infinity as particles become far apart, one would expect $|\Phi_0|, |\Psi_0| \rightarrow 0$ as $|x_{ij}| \rightarrow \infty$ [54]. Nevertheless, it should be noted that the family of models generated by EOF and PHJ is always integrable, regardless of the normalization of the Jastrow wave function.

For example, if $f_{ij} = \exp(g|x_{ij}|)$, H_0 becomes the well-known LL model [25]. However, the Jastrow wave function is normalizable only when $g < 0$, which corresponds to the McGuire bright soliton [41]. Therefore Φ_0 is no longer the ground state wave function of the repulsive LL model. However, as we have discussed previously, the integrability of the Hamiltonian is not affected, so our result reproduces the integrability of the LL model with the integral of motion being I_n or K_n . More interestingly, upon introducing the external harmonic potential, according to Eq. (11), Ψ_0 becomes normalizable even if Φ_0 is not and Eq. (5) corresponds to the Lieb-Liniger-Coulomb model introduced in Refs. [44], i.e.,

$$H = \sum_i \left[\frac{p_i^2}{2m} + \frac{1}{2} m \omega^2 x_i^2 \right] + g \sum_{i < j} \left[\frac{2\hbar^2}{m} \delta(x_{ij}) - \frac{m\omega}{\hbar} |x_{ij}| \right], \quad (19)$$

with ground state $E_0 = \frac{N\hbar\omega}{2} - \frac{g^2\hbar^2}{m} \frac{N(N^2-1)}{6}$. This system describes harmonically confined bosons subject to contact and Coulomb interactions or gravitational attraction in one spatial dimension. Ref. [44] characterized its EOF representation and ground state properties. Using Theorem 2, we conclude that this system is integrable, with the integrals of motion being

$$\tilde{I}_n \equiv \sum_i h_i^n.$$

Conclusion.— We have established the equivalence between the families of one-dimensional many-body quantum systems generated by the exchange operator formalism and parent Hamiltonians with a ground-state wavefunction of Jastrow form, describing indistinguishable particles with no internal degrees of freedom. We have further generalized this equivalence when embedding the systems in an external potential. Making use of the projection operator onto the spinless bosonic or fermionic subspace, we have proved the integrability of all these systems by constructing explicitly the corresponding integrals of motion. Embedding these translation-invariant models in an external potential preserves the integrability, in the presence of long-range interactions, as we have shown in the long-range Lieb-Liniger model.

The findings advance the study of many-body physics by uncovering the implications of ground-state correlations on integrability and should lead to manifold applications in nonlinear physics, quantum quenches, quantum simulation, Wigner crystals, quantum gases, among others. Our results bear also implications on numerical methods for strongly-correlated systems such as variational methods and quantum Monte Carlo algorithms, in which the ubiquitous use of Jastrow wavefunction ansatz may impose integrability on systems lacking it. It may be possible to extend our results to higher spatial dimensions [26], the inclusion of spin degrees of freedom [8], mixtures of different species [34], and distinguishable particles [55, 56].

Acknowledgement.— It is a pleasure to acknowledge discussions with Pablo Martinez Azcona and Aritra Kundu.

* jing.yang@uni.lu

† adolfo.delcampo@uni.lu

- [1] J. Dziarmaga, *Advances in Physics* **59**, 1063 (2010).
- [2] T. Mori, T. N. Ikeda, E. Kaminishi, and M. Ueda, *Journal of Physics B: Atomic, Molecular and Optical Physics* **51**, 112001 (2018).
- [3] M. A. Cazalilla, R. Citro, T. Giamarchi, E. Orignac, and M. Rigol, *Rev. Mod. Phys.* **83**, 1405 (2011).
- [4] X.-W. Guan, M. T. Batchelor, and C. Lee, *Rev. Mod. Phys.* **85**, 1633 (2013).
- [5] M. Takahashi, *Thermodynamics of One-Dimensional Solvable Models* (Cambridge, Cambridge, 1999).
- [6] M. Gaudin, *The Bethe Wavefunction*, edited by J.-S. Caux (Cambridge University Press, 2014).
- [7] A. P. Polychronakos, *Phys. Rev. Lett.* **69**, 703 (1992).
- [8] A. P. Polychronakos, *Journal of Physics A: Mathematical and General* **39**, 12793 (2006).
- [9] C. N. Yang, *Physical Review Letters* **19**, 1312 (1967).
- [10] R. J. Baxter, *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences* **289**, 315 (1978).
- [11] V. E. Korepin, N. M. Bogoliubov, and A. G. Izergin, *Quantum Inverse Scattering Method and Correlation Functions* (Cambridge, Cambridge, 1997).
- [12] B. Sutherland, *Beautiful Models* (World Scientific, 2004).
- [13] B. Sutherland, *Phys. Rev. Lett.* **75**, 1248 (1995).
- [14] J.-S. Caux and J. Mossel, *Journal of Statistical Mechanics: Theory and Experiment* **2011**, P02023 (2011).
- [15] H. Ujino and M. Wadati, *Journal of the Physical Society of Japan* **64**, 4121 (1995).
- [16] A. P. Polychronakos, *Nuclear Physics B* **961**, 115243 (2020).
- [17] A. Bijl, *Physica* **7**, 869 (1940).
- [18] R. B. Dingle, *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science* **40**, 573 (1949).
- [19] R. Jastrow, *Phys. Rev.* **98**, 1479 (1955).
- [20] F. Calogero, *Journal of Mathematical Physics* **12**, 419 (1971).
- [21] B. Sutherland, *Journal of Mathematical Physics* **12**, 246 (1971).
- [22] B. Sutherland, *Phys. Rev. A* **4**, 2019 (1971).
- [23] F. Calogero and C. Marchioro, *Journal of Mathematical Physics* **14**, 182 (1973).
- [24] P. J. Gambardella, *Journal of Mathematical Physics* **16**, 1172 (1975).
- [25] A. del Campo, *Phys. Rev. Research* **2**, 043114 (2020).
- [26] M. Beau and A. del Campo, *SciPost Phys. Core* **4**, 30 (2021).
- [27] C. L. Kane, S. Kivelson, D. H. Lee, and S. C. Zhang, *Phys. Rev. B* **43**, 3255 (1991).
- [28] F. Calogero, *Lett. Nuovo Cim.* **3**, 411 (1975).
- [29] L. Šamaj and Z. Bajnok, *Introduction to the Statistical Physics of Integrable Many-body Systems* (Cambridge University Press, 2013).
- [30] T. Kinoshita, T. Wenger, and D. S. Weiss, *Science* **305**, 1125 (2004).
- [31] B. Paredes, A. Widera, V. Murg, O. Mandel, S. Fölling, I. Cirac, G. V. Shlyapnikov, T. W. Hänsch, and I. Bloch, *Nature* **429**, 277 (2004).
- [32] M. Girardeau, *Journal of Mathematical Physics* **1**, 516 (1960).
- [33] M. D. Girardeau, E. M. Wright, and J. M. Triscari, *Phys. Rev. A* **63**, 033601 (2001).
- [34] M. D. Girardeau and A. Minguzzi, *Phys. Rev. Lett.* **99**, 230402 (2007).
- [35] M. Girardeau, H. Nguyen, and M. Olshanii, *Optics Communications* **243**, 3 (2004).
- [36] N. Kawakami, *Journal of the Physical Society of Japan* **62**, 4163 (1993).
- [37] K. Vacek, A. Okiji, and N. Kawakami, *Journal of Physics A: Mathematical and General* **27**, L201 (1994).
- [38] N. Gurappa and P. K. Panigrahi, *Phys. Rev. B* **59**, R2490 (1999).
- [39] M. Olshanii, *Phys. Rev. Lett.* **81**, 938 (1998).
- [40] E. H. Lieb, R. Seiringer, and J. Yngvason, *Phys. Rev. Lett.* **91**, 150401 (2003).
- [41] J. B. McGuire, *Journal of Mathematical Physics* **5**, 622 (1964).
- [42] E. H. Lieb and W. Liniger, *Phys. Rev.* **130**, 1605 (1963).
- [43] E. H. Lieb, *Phys. Rev.* **130**, 1616 (1963).
- [44] M. Beau, S. M. Pittman, G. E. Astrakharchik, and A. del Campo, *Phys. Rev. Lett.* **125**, 220602 (2020).
- [45] B. Sutherland, *Journal of Mathematical Physics* **12**, 251 (1971).
- [46] J. Moser, *Advances in Mathematics* **16**, 197 (1975).
- [47] J. Yang and A. del Campo, *Supplemental Material* (2022).
- [48] J. W. Negele and H. Orland, *Quantum Many-particle Systems* (Perseus Books, Reading, MA, 1998).
- [49] A. Kundu, *Phys. Rev. Lett.* **83**, 1275 (1999).
- [50] M. D. Girardeau, *Phys. Rev. Lett.* **97**, 100402 (2006).
- [51] M. T. Batchelor, X.-W. Guan, and N. Oelkers, *Phys. Rev. Lett.* **96**, 210402 (2006).
- [52] F. Cooper, A. Khare, and U. Sukhatme, *Physics Reports* **251**, 267 (1995).
- [53] This observation was put forward first by Polychakros [7], using a heuristic locality argument. However, we emphasize the argument misses an important ingredient, the permutation in-

variance. In [47], we give a formal proof with inspirations from the coordinate Bethe ansatz [6, 29].

- [54] In this case, only bound states are possible. No scattering states are allowed since particles can not tunnel through an infinite potential barrier.
- [55] S. R. Jain and A. Khare, *Physics Letters A* **262**, 35 (1999).
- [56] S. M. Pittman, M. Beau, M. Olshanii, and A. del Campo, *Phys. Rev. B* **95**, 205135 (2017).

Supplemental Material for “One-Dimensional Quantum Systems with Ground-State of Jastrow Form Are Integrable”

CONTENTS

- I. The exchange operator
 - A. Properties of the two-particle position exchange operator M_{ij}
 - B. Properties of the three-particle position exchange operators M_{ijk}
- II. Projection onto the bosonic and fermionic subspaces
- III. Derivation of Eq. (3)
- IV. Results related to integrability
 - A. Proof of Eq. (16)
 - B. Proof of the block-diagonal structure of permutation-invariant observables.
 - C. Proof of Eq. (17)
 - D. Effective 1-body operator

I. THE EXCHANGE OPERATOR

In this section, we give a rigorous definition of the exchange operator used in the main text and derive its properties from first principles.

A. Properties of the two-particle position exchange operator M_{ij}

Consider the N -fold tensor-product of a single-particle Hilbert space \mathcal{H} , i.e., $\mathcal{H}^{\otimes N}$, where we do not assume a particular exchange statistics for the particles. This means that the particles may be distinguishable or not. The position exchange operator M_{ij} acting on the spatial coordinates x_i and x_j on a many-particle state $\psi_s \in \mathcal{H}^{\otimes N}$ is defined as

$$M_{ij}\psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N) = \psi_s(x_1, \dots, x_j, \dots, x_i, \dots, x_N), \quad i \neq j, \quad (\text{S1})$$

where $\psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N)$ is the wave function on $\mathcal{H}^{\otimes N}$ and s denotes the internal degrees of freedom, which can account for spin in identical particles. By definition, we find

$$M_{ji} = M_{ij}. \quad (\text{S2})$$

Furthermore, according to this definition, it follows that

$$M_{ij}^2\psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N) = \psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N), \quad (\text{S3})$$

which implies that M_{ij} is idempotent, i.e.,

$$M_{ij}^2 = \mathbb{I}. \quad (\text{S4})$$

The inner product on $\mathcal{H}^{\otimes N}$ is

$$(\phi, \psi) \equiv \sum_s \int \prod_{k=1}^N dx_k \phi_s^*(x_1, \dots, x_i, \dots, x_j, \dots, x_N) \psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N). \quad (\text{S5})$$

The adjoint of M_{ij} is defined as $(M_{ij}^\dagger \phi, \psi) \equiv (\phi, M_{ij} \psi)$. Explicitly expanding the inner product according to Eq (S5), we arrive at

$$\begin{aligned}
& \sum_s \int \prod_{k=1}^N dx_k [M_{ij}^\dagger \phi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N)]^* \psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N) \\
&= \sum_s \int \prod_{k=1}^N dx_k \phi_s^*(x_1, \dots, x_i, \dots, x_j, \dots, x_N) \psi_s(x_1, \dots, x_j, \dots, x_i, \dots, x_N) \\
&= \sum_s \int \prod_{k=1}^N dx_k \phi_s^*(x_1, \dots, x_j, \dots, x_i, \dots, x_N) M_{ij} \psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N),
\end{aligned} \tag{S6}$$

which indicates that

$$M_{ij}^\dagger \phi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N) = \phi_s(x_1, \dots, x_j, \dots, x_i, \dots, x_N). \tag{S7}$$

Comparing above equation with Eq. (S1), we conclude that M_{ij} is a Hermitian operator,

$$M_{ij} = M_{ij}^\dagger. \tag{S8}$$

We define operators or observables

$$A_i \equiv A(x_i), \tag{S9}$$

$$A_{ij} \equiv A(x_i, x_j), \tag{S10}$$

$$A_{ijk} \equiv A(x_i, x_j, x_k), \tag{S11}$$

⋮

where the dependence on the internal degrees of freedom of O is suppressed. Then one can easily find that

$$(\phi, M_{ij} A_k \psi) = (M_{ij}^\dagger \phi, A_k \psi) = \sum_s \int \prod_{l=1}^N dx_l \phi_s^*(x_1, \dots, x_j, \dots, x_i, \dots, x_N) O(x_k) \psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N) \tag{S12}$$

For $k \neq i, j$,

$$\begin{aligned}
& \sum_s \int \prod_{l=1}^N dx_l \phi_s^*(x_1, \dots, x_j, \dots, x_i, \dots, x_N) O(x_k) \psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N) \\
&= \sum_s \int \prod_{l=1}^N dx_l \phi_s^*(x_1, \dots, x_j, \dots, x_i, \dots, x_N) O(x_k) \psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N) \\
&= \sum_s \int \prod_{l=1}^N dx_l \phi_s^*(x_1, \dots, x_j, \dots, x_i, \dots, x_N) O(x_k) M_{ji} \psi_s(x_1, \dots, x_j, \dots, x_i, \dots, x_N) \\
&= (\phi, O_k M_{ji})
\end{aligned} \tag{S13}$$

We conclude that

$$M_{ij} A_k = A_k M_{ji} = A_k M_{ij}, \quad \text{for } i, j, k, \text{ distinct.} \tag{S14}$$

If $k = i$, then we find

$$\begin{aligned}
(\phi, M_{ij} A_i \psi) &= \sum_s \int \prod_{l=1}^N dx_l \phi_s^*(x_1, \dots, x_j, \dots, x_i, \dots, x_N) O(x_i) \psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N) \\
&= \sum_s \int \prod_{l=1}^N dx_l \phi_s^*(x_1, \dots, x_j, \dots, x_i, \dots, x_N) A(x_i) M_{ji} \psi_s(x_1, \dots, x_j, \dots, x_i, \dots, x_N) \\
&= \sum_s \int \prod_{l=1}^N dx_l \phi_s^*(x_1, \dots, x_i, \dots, x_j, \dots, x_N) A(x_j) M_{ij} \psi_s(x_1, \dots, x_i, \dots, x_j, \dots, x_N) \\
&= (\phi, A_j M_{ij} \psi),
\end{aligned} \tag{S15}$$

where we have used change of the dummy indices $x_i \rightarrow x_j$ and $x_j \rightarrow x_i$ in the second last equation. We find

$$M_{ij}A_i = A_jM_{ij} \quad (\text{S16})$$

and similarly

$$M_{ij}A_j = A_iM_{ij}. \quad (\text{S17})$$

One can extend the above arguments to many-body operators $A_{ijk\dots}$ with little effort. For example, for two-body operators A_{kl} , one can find

$$M_{ij}A_{kl} = A_{kl}, \quad \text{for } i, j, k, l, \text{ distinct} \quad (\text{S18})$$

$$M_{ij}A_{jk} = A_{ik}M_{ij}, \quad \text{for } i, j, k, \text{ distinct} \quad (\text{S19})$$

$$M_{ij}A_{kj} = A_{ki}M_{ij}, \quad \text{for } i, j, k, \text{ distinct} \quad (\text{S20})$$

$$M_{ij}A_{ij} = A_{ji}M_{ij}, \quad (\text{S21})$$

$$M_{ij}A_{ji} = A_{ij}M_{ij}. \quad (\text{S22})$$

B. Properties of the three-particle position exchange operators M_{ijk}

The particle exchange operator with three indices is defined as

$$M_{ijk} = M_{ij}M_{jk}, \quad \text{for } i, j, k, \text{ distinct}. \quad (\text{S23})$$

With the properties of the two-particle exchange operator, one can easily show that M_{ijk} is invariant under cyclic permutation, i.e.,

$$M_{ijk} = M_{jki} = M_{kij} \quad (\text{S24})$$

However, it is *not* fully symmetric in its indices. In particular, $M_{ijk} \neq M_{jik}$. Furthermore, for all distinct i, j, k, l ,

$$M_{ijk}A_l = A_lM_{ijk} \quad (\text{S25})$$

since A_l commute with M_{ij} and M_{jk} . Furthermore, one can explicit check that

$$M_{ijk}A_i = M_{ij}A_iM_{jk} = A_jM_{ijk}, \quad (\text{S26})$$

$$M_{ijk}A_j = M_{ij}A_kM_{jk} = A_kM_{ijk}, \quad (\text{S27})$$

$$M_{ijk}A_k = M_{ij}A_jM_{jk} = A_iM_{ijk}. \quad (\text{S28})$$

II. PROJECTION ONTO THE BOSONIC AND FERMIONIC SUBSPACES

In EOF, one may restrict the construction to the bosonic or fermionic subspace to simplify the calculation or motivated on physical grounds. For example, it was shown by Polychronakos [7, 8] that if the prepotential is $V_{ij} = l/x_{ij}$, then

$$\frac{1}{2}I_2 = \frac{1}{2m} \sum_i \pi_i^2 = \frac{1}{2m} \sum_i p_i^2 + \frac{\hbar^2}{m} \sum_{i>j} \frac{l(l - M_{ij})}{(x_i - x_j)^2}. \quad (\text{S29})$$

This Hamiltonian is still defined on $\mathcal{H}^{\otimes N}$. Recall that the Hilbert space \mathcal{H}_ζ of identical particles is a subspace of $\mathcal{H}^{\otimes N}$. We define the projection operator on $\mathcal{H}^{\otimes N}$ to \mathcal{H}_ζ as \mathcal{P}_ζ . It can be readily checked that for any $\psi(x_1, x_2, \dots, x_N) \in \mathcal{H}^{\otimes N}$,

$$\begin{aligned} M_{ij}\mathcal{P}_\zeta\psi(x_1, x_2, \dots, x_N) &= \frac{1}{N!} \sum_{\sigma} (\pm 1)^\sigma M_{ij}\psi(x_{\sigma_1}, \dots, x_{\sigma_i}, \dots, x_{\sigma_j}, \dots, x_{\sigma_N}) \\ &= \frac{1}{N!} \sum_{\sigma} (\pm 1)^{\sigma+1} \psi(x_{\sigma_1}, \dots, x_{\sigma_i}, \dots, x_{\sigma_j}, \dots, x_{\sigma_N}) \\ &= \pm \mathcal{P}_\zeta\psi(x_1, x_2, \dots, x_N), \end{aligned} \quad (\text{S30})$$

$$\begin{aligned}
\mathcal{P}_\zeta M_{ij} \psi(x_1, x_2, \dots, x_N) &= \mathcal{P}_\zeta \psi(x_1, \dots, x_j, \dots, x_i, \dots, x_N) \\
&= (\pm 1) \mathcal{P}_\zeta \psi(x_1, \dots, x_i, \dots, x_j, \dots, x_N) \\
&= \frac{1}{N!} \sum_{\sigma} (\pm 1)^{\sigma+1} \psi(x_{\sigma_1}, \dots, x_{\sigma_i}, \dots, x_{\sigma_j}, \dots, x_{\sigma_N}) \\
&= \pm \mathcal{P}_\zeta \psi(x_1, x_2, \dots, x_N).
\end{aligned} \tag{S31}$$

Thus, we conclude

$$[\mathcal{P}_\zeta, M_{ij}] = 0, \tag{S32}$$

$$\mathcal{P}_\zeta M_{ij} = M_{ij} \mathcal{P}_\zeta = \zeta \mathcal{P}_\zeta. \tag{S33}$$

Therefore, when projecting onto the bosonic or fermionic subspace, Eq. (S29) becomes

$$\frac{1}{2} \mathcal{P}_\zeta I_2 \mathcal{P}_\zeta = \frac{1}{2} \sum_i \pi_i^2 = \mathcal{P}_\zeta H_{\text{CS}} \mathcal{P}_\zeta, \tag{S34}$$

where

$$H_{\text{CS}} = \frac{1}{2m} \sum_i p_i^2 + \frac{\hbar^2}{m} \sum_{i>j} \frac{l(l \mp \zeta)}{(x_i - x_j)^2} \tag{S35}$$

is the rational Calogero model. The procedure in Eq. (S34) is effectively equivalent to replacing $M_{ij} = \zeta \mathbb{1}$.

However, we warn the audience that when focusing on spinless bosons or fermions, setting $M_{ij} = \zeta \mathbb{1}$ requires some caution. For example, when calculating $[p_i, \sum_{k \neq j} V_{jk} M_{jk}]$ with $i \neq j$, had one setting $M_{jk} = \zeta \mathbb{1}$ before actually calculating the commutator, one would obtain

$$[p_i, \sum_{k \neq j} V_{jk} M_{jk} \mathcal{P}_\zeta] \stackrel{?}{=} \zeta i \hbar V'_{ij}. \tag{S36}$$

On the other hand, a rigorous calculation following Eqs. (S14-S22) shows that

$$\begin{aligned}
[p_i, \sum_{k \neq j} V_{jk} M_{jk}] &= [p_j, V_{ji} M_{ji}] = [p_i, V_{ji}] M_{ji} + V_{ji} [p_i, M_{ji}] \\
&= i \hbar V'_{ij} M_{ij} - V_{ij} (p_i - p_j) M_{ij}, \quad (i \neq j)
\end{aligned} \tag{S37}$$

Upon projecting this relation onto \mathcal{H}_α by setting $M_{jk} = \mathbb{1}$, which yields

$$[p_i, \sum_{k \neq j} V_{jk} M_{jk}] \mathcal{P}_\zeta = \zeta i \hbar V'_{ij} - \zeta V_{ij} (p_i - p_j). \tag{S38}$$

This example clearly shows that $[\mathcal{A}, \mathcal{B}] \mathcal{P}_\zeta \neq [\mathcal{A}, \mathcal{B} \mathcal{P}_\zeta]$ and in particular

$$[\mathcal{A}, \mathcal{B} \mathcal{P}_\zeta] = [\mathcal{A}, \mathcal{B}] \mathcal{P}_\zeta + \mathcal{B} [\mathcal{A}, \mathcal{P}_\zeta]. \tag{S39}$$

The caveat we would like to give to the audience is that whenever one would like to set $M_{ij} = \pm \mathbb{1}$, say, on physical grounds, one should bear in mind that at the formal level a projection has been introduced, which has to be taken into account to work out the correct algebra.

Finally, we mention the following interesting lemma for spinless indistinguishable particles, thanks to the EOF and the projection operator:

Lemma 1. *For any k -body operator $A_{i_1 \dots i_k}$ acting on states of N -spinless indistinguishable particles, $\mathcal{P}_\zeta A_{i_1 \dots i_k} \mathcal{P}_\zeta$ is “super-symmetric”, i.e.,*

$$\mathcal{P}_\zeta A_{i_1 \dots i_k} \mathcal{P}_\zeta = \mathcal{P}_\zeta A_{j_1 \dots j_k} \mathcal{P}_\zeta, \tag{S40}$$

where (j_1, j_2, \dots, j_k) is arbitrary k -tuple, not necessarily a permutation of (i_1, \dots, i_k)

Proof. The proof is to take advantage of Eq. (S4) and Eq. (2). The intuition for the proof can be easily seen from the case where indices (j_1, j_2, \dots, j_k) are distinct:

$$\begin{aligned}
\mathcal{P}_\zeta A_{i_1 \dots i_k} \mathcal{P}_\zeta &= \mathcal{P}_\zeta M_{i_1 j_1} \dots M_{i_k j_k} M_{i_k j_k} \dots M_{i_1 j_1} A_{i_1 \dots i_k} \mathcal{P}_\zeta \\
&= \mathcal{P}_\zeta M_{i_1 j_1} \dots M_{i_k j_k} A_{j_1 \dots j_k} M_{i_k j_k} M_{i_k j_k} M_{i_1 j_1} \mathcal{P}_\zeta \\
&= \mathcal{P}_\zeta \zeta^k A_{j_1 \dots j_k} \zeta^k \mathcal{P}_\zeta \\
&= \mathcal{P}_\zeta A_{j_1 \dots j_k} \mathcal{P}_\zeta.
\end{aligned} \tag{S41}$$

With the above intuition, one can easily prove the general case without any difficulty. \square

III. DERIVATION OF EQ. (3)

Given the generalized momenta

$$\pi_i = p_i + i \sum_{j \neq i} V_{ij} M_{ij},$$

the EOF Hamiltonian is defined as

$$\begin{aligned}
\frac{1}{2m} I_2 &= \frac{1}{2m} \sum_i \pi_i^2 \\
&= \frac{1}{2m} \sum_i \left(p_i + i \sum_{j \neq i} V_{ij} M_{ij} \right)^2 \\
&= \frac{1}{2m} \sum_i \left[p_i^2 + i \sum_{j \neq i} (p_i V_{ij} M_{ij} + V_{ij} M_{ij} p_i) - \sum_{j, k \neq i} V_{ij} M_{ij} V_{ik} M_{ik} \right],
\end{aligned} \tag{S42}$$

which is the sum of a purely kinetic one-body term, a two-body term, and a three-body term. We next use the fact the two-body term can be rewritten making use of $[p_i, V_{ij}] = -i\hbar$, $\partial_i V_{ij} = -i\hbar V'_{ij}$ together with $V_{ij} M_{ij} p_i = V_{ij} p_j M_{ij}$.

In addition, we note that the third term on r.h.s. of Eq. (S42) with $j = k$ becomes a two-body term $-V_{ij} M_{ij} V_{ij} M_{ij} = -V_{ij} V_{ji} M_{ij}^2 = +V_{ij}^2$, using the fact that V_{ij} is antisymmetric in its indices. Thus

$$\begin{aligned}
\frac{1}{2m} I_2 &= \frac{1}{2m} \sum_i \left[p_i^2 + \sum_{j \neq i} (\hbar V'_{ij} M_{ij} + i V_{ij} (p_i + p_j) M_{ij} + V_{ij}^2) - \sum_{i \neq j \neq k \neq i} V_{ij} M_{ij} V_{ik} M_{ik} \right] \\
&= \frac{1}{2m} \sum_i \left[p_i^2 + \sum_{j \neq i} (\hbar V'_{ij} M_{ij} + V_{ij}^2) - \sum_{i \neq j \neq k \neq i} V_{ij} V_{jk} M_{ij} M_{ik} \right] \\
&= \frac{1}{2m} \sum_i \left[p_i^2 + \sum_{j \neq i} (\hbar V'_{ij} M_{ij} + V_{ij}^2) - \sum_{i \neq j \neq k \neq i} V_{ij} V_{ki} M_{ijk} \right],
\end{aligned} \tag{S43}$$

where in the last step we have swapped the indices i and j and used the fact that V_{ij} is antisymmetric in its indices, and that M_{ijk} is the three-particle exchange operator defined in Eq. (S23). Taking advantage of the invariance of M_{ijk} under the cyclic permutation of i, j, k , we can further symmetrize the last term on the r.h.s. of Eq. (S43) as

$$\sum_{i \neq j \neq k \neq i} V_{ij} V_{ki} M_{ijk} = \frac{1}{3} \sum_{j \neq k \neq i} (V_{ij} V_{jk} + V_{jk} V_{ki} + V_{ki} V_{ij}) M_{ijk} = \frac{1}{3} \sum_{j \neq k \neq i} V_{ijk} M_{ijk}, \tag{S44}$$

where $V_{ijk} = V_{ij} V_{jk} + V_{jk} V_{ki} + V_{ki} V_{ij}$. Upon projecting to the bosonic or fermionic subspace, we find,

$$\frac{1}{2m} \mathcal{P}_\zeta I_2 \mathcal{P}_\zeta = \mathcal{P}_\zeta \left(\frac{1}{2m} \sum_i p_i^2 + \frac{1}{2m} \sum_{i \neq j} (\zeta \hbar V'_{ij} + V_{ij}^2) - \frac{1}{6m} \sum_{j \neq k \neq i} V_{ijk} \right) \mathcal{P}_\zeta \tag{S45}$$

Since V'_{ij} and V_{ij}^2 are symmetric in the indices i and j and V_{ijk} is fully symmetric in permutation of any pair of the indices, the sums in above equation can be further rewritten as

$$\frac{1}{2m}\mathcal{P}_\zeta I_2 \mathcal{P}_\zeta = \mathcal{P}_\zeta \left(\frac{1}{2m} \sum_i p_i^2 + \frac{1}{m} \sum_{i<j} (\zeta \hbar V'_{ij} + V_{ij}^2) - \frac{1}{m} \sum_{i<j<k} V_{ijk} \right) \mathcal{P}_\zeta. \quad (\text{S46})$$

Therefore, we find Eq. (3).

IV. RESULTS RELATED TO INTEGRABILITY

In this section, we discuss some results that are used in the proof of integrability discussed in the main text.

A. Proof of Eq. (16)

For any permutation-invariant observable $O(x_1, x_2, \dots, x_N)$,

$$\begin{aligned} \mathcal{P}_\zeta O(x_1, x_2, \dots, x_N) \psi(x_1, x_2, \dots, x_N) &= \frac{1}{N!} \sum_{\sigma} O(x_{\sigma_1}, x_{\sigma_2}, \dots, x_{\sigma_N}) \psi(x_{\sigma_1}, x_{\sigma_2}, \dots, x_{\sigma_N}) \\ &= O(x_1, x_2, \dots, x_N) \mathcal{P}_\zeta \psi(x_1, x_2, \dots, x_N), \end{aligned} \quad (\text{S47})$$

for any $\psi(x_1, x_2, \dots, x_N) \in \mathcal{H}^{\otimes N}$. Therefore, we conclude that

$$[\mathcal{P}_\zeta, O] = 0. \quad (\text{S48})$$

B. Proof of the block-diagonal structure of permutation-invariant observables.

Proposition. A permutation-invariant observable is block diagonal on \mathcal{H}_ζ and \mathcal{H}_ζ^\perp , where \mathcal{H}_ζ^\perp is the orthogonal complement of \mathcal{H}_ζ on $\mathcal{H}^{\otimes N}$.

Proof. We denote the projector onto \mathcal{H}_ζ^\perp as \mathcal{P}_ζ^\perp . Then, for a permutation-invariant observable O , according to Eq. (16), it can be easily seen

$$\mathcal{P}_\zeta O \mathcal{P}_\zeta^\perp = O \mathcal{P}_\zeta \mathcal{P}_\zeta^\perp = 0, \quad (\text{S49})$$

$$\mathcal{P}_\zeta^\perp O \mathcal{P}_\zeta = \mathcal{P}_\zeta^\perp \mathcal{P}_\zeta^\perp O = 0, \quad (\text{S50})$$

whence it follows that

$$O = \mathcal{P}_\zeta O \mathcal{P}_\zeta \oplus \mathcal{P}_\zeta^\perp O \mathcal{P}_\zeta^\perp, \quad (\text{S51})$$

which concludes the proof. \square

C. Proof of Eq. (17)

Eq. (17) is equivalent to the following proposition:

Proposition. If a permutation-invariant and local observable O vanishes on \mathcal{H}_ζ , then it must vanish on the full Hilbert space $\mathcal{H}^{\otimes N}$.

Proof. Assuming $\exists \psi(x_1, x_2, \dots, x_N) \in \mathcal{H}^{\otimes N}$, which is not necessarily symmetric or anti-symmetric. We borrow the inspirations from coordinate Bethe ansatz [6, 29] and construct another symmetric or anti-symmetric wave function $\phi(x_1, x_2, \dots, x_N) \in \mathcal{H}_\zeta$

$$\phi(x_1, x_2, \dots, x_N) \equiv \sum_{\sigma} \zeta^{\sigma} \psi(x_{\sigma_1}, x_{\sigma_2}, \dots, x_{\sigma_N}) \theta_{\text{H}}(x_{\sigma_N} - x_{\sigma_{N-1}}) \cdots \theta_{\text{H}}(x_{\sigma_3} - x_{\sigma_2}) \theta_{\text{H}}(x_{\sigma_2} - x_{\sigma_1}) \quad (\text{S52})$$

where σ is a permutation of the tuple $(1, 2, \dots, N)$ and $\theta_H(x)$ is the Heaviside function. It is worth to note that ϕ is not $\mathcal{P}_\zeta\psi$. For example, if $\psi \in \mathcal{H}_\zeta$, then $\mathcal{P}_\zeta\psi = 0$ while $\phi \neq 0$ as long as $\psi \neq 0$. Since \mathcal{O} vanishes on \mathcal{H}_ζ therefore

$$\mathcal{O}(x_1, x_2, \dots, x_N)\phi(x_1, x_2, \dots, x_N) = 0 \quad (\text{S53})$$

which implies

$$\mathcal{O}(x_1, x_2, \dots, x_N)\psi(x_{\sigma_1}, x_{\sigma_2}, \dots, x_{\sigma_N}) = 0 \quad (\text{S54})$$

for $x_{\sigma_1} < x_{\sigma_2} < \dots < x_{\sigma_N}$. Eq. (S54) implies that

$$\mathcal{O}(x_1, x_2, \dots, x_N)\psi(x_1, x_2, \dots, x_N) = 0 \quad (\text{S55})$$

for $x_1 < x_2 < \dots < x_N$. However, our goal is to show it holds in all the regions. To reach this goal, let us take advantage of permutation invariance. Note that r.h.s. of Eq. (S54) implies the x_i 's on l.h.s. are dummy indices. Therefore, Eq. (S54) is equivalence as

$$\mathcal{O}(x_{\sigma_1^{-1}}, x_{\sigma_2^{-1}}, \dots, x_{\sigma_N^{-1}})\psi(x_1, x_2, \dots, x_N) = \mathcal{O}(x_1, x_2, \dots, x_N)\psi(x_1, x_2, \dots, x_N) = 0 \quad (\text{S56})$$

where σ^{-1} is the inverse permutation of σ . So far, we have shown that Eq. (S55) holds on regions where (x_1, x_2, \dots, x_N) . On the boundary where at least two of the coordinates coincide, locality of \mathcal{O} implies that the value of $\mathcal{O}\psi$ on the boundary is full determined by the its value on the neighborhood outside the boundary, where $\mathcal{O}\psi$ vanishes. Thus, $\mathcal{O}\psi$ must also vanishes on the boundary. \square

It can be shown that [7]

$$[\pi_i, \pi_j] = \sum_{k \neq i, j} V_{ijk}(M_{ijk} - M_{jik}). \quad (\text{S57})$$

Thus $(M_{ijk} - M_{jik})\mathcal{P}_\zeta = 0$ and we obtain

$$[\pi_i, \pi_j]\mathcal{P}_\zeta = 0. \quad (\text{S58})$$

Furthermore $[\pi_i, \pi_j]$ is local since they only involve at most second derivative of the coordinate. However, it does not vanish on the whole Hilbert space $\mathcal{H}^{\otimes N}$ because $[\pi_i, \pi_j]$ is not permutation-invariant. So here we can see for a local, but not permutation-invariant operator, even it vanishes on \mathcal{H}_ζ , it may not vanish on $\mathcal{H}^{\otimes N}$.

D. Effective 1-body operator

The operator

$$W_i = i \sum_{k \neq i} V_{ik} M_{ik} \quad (\text{S59})$$

acts an effectively 1-body potential, in the sense that it satisfies the exchange rule for 1-body operators, even if it involves N degrees of freedom! Indeed,

$$\begin{aligned} M_{ij}W_j &= i \sum_{k \neq j} M_{ij}V_{jk}M_{jk} = i \sum_{k \neq i, j} M_{ij}V_{jk}M_{jk} + iM_{ij}V_{ji}M_{ji} \\ &= i \sum_{k \neq i, j} V_{ik}M_{ik}M_{ij} + iV_{ij}M_{ij}M_{ij} \\ &= i \sum_{k \neq i} V_{ik}M_{ik}M_{ij} = W_iM_{ij}. \end{aligned} \quad (\text{S60})$$

Further,

$$\begin{aligned} M_{ij}W_k &= i \sum_{l \neq k} M_{ij}V_{kl}M_{kl} = i \sum_{l \neq k, i, j} M_{ij}V_{kl}M_{kl} + iM_{ij}V_{ki}M_{ki} + iM_{ij}V_{kj}M_{kj} \\ &= i \sum_{l \neq k, i, j} V_{kl}M_{kl}M_{ij} + iV_{ki}M_{ki}M_{ij} + iV_{kj}M_{kj}M_{ij} \\ &= i \sum_{l \neq k} V_{kl}M_{kl}M_{ij} = W_kM_{ij}. \end{aligned} \quad (\text{S61})$$

Therefore, π_i , a_i , a_i^\dagger and h_i act all as effectively 1-body operators, i.e., they satisfy the exchange rules of a 1-body operator.