

Original Paper

Comment on Causal Networks and Freedom of Choice in Bell's **Theorem**

Marian Kupczynski

Département d'informatique et d'ingénierie, Université du Québec en Outaouais, Gatineau, QC, Canada

E-Mail: marian.kupczynski@uqo.ca

Received: 22 March 2022 / Accepted: 30 March 2022 / Published: 31 March 2022

Abstract: Bell inequalities may only be derived, if hidden variables do not depend on the experimental settings. The stochastic independence of hidden and setting variables is called: freedom of choice, free will, measurement independence (MI) or no conspiracy. By imbedding the Bell causal structure in a larger causal network the authors correctly prove, that one can explain and quantify possible violation of MI without evoking super-determinism. They assume the independence of the variables that causally determine the settings and investigate how they might become correlated with hidden variables (e.g., when the cosmic photons enter the laboratory). Using their extended causal networks they derive a contextual probabilistic model on which their further correct results are based. The authors seem to ignore that contextual probabilistic model may be derived directly using only probabilistic concepts and incorporating correctly setting dependent variables describing measuring instruments. In these contextual probabilistic models experimenters' freedom of choice is not compromised and the results of Bell Tests including an apparent violation of Einsteinian nonsignaling may be explained in a locally causal way. Talking about freedom of choice is misleading and is rooted in incorrect understanding of Bayes Theorem. We explain why MI should be called noncontextuality and why its violation in Bell Tests confirms only the contextual character of quantum observables. Therefore, contextuality and not experimenters' freedom of choice are important resources in quantum information.

Keywords: Bell inequalities; measurement independence; contextuality; local causality

In Bell scenario we have 4 random experiments performed using 4 incompatible pairs of settings [1, 2]. A source S is sending two correlated photonic signals to Alice and Bob in distant laboratories. Before 'photons' arrive to corresponding measuring instruments Alice and Bob choose independently their experimental settings (x, y). In quantum mechanics (QM) and in Bell model, x=1 or 2 and y=1 or 2 are labels indicating which pair of experimental setting are chosen. In Bell tests (x, y) are values of two binary random variables (X, Y) describing outcomes of supplementary random experiments (e.g. flipping two fair coins, generation of pseudo-random numbers) or coded information from some random phenomena (e.g. arrivals of cosmic photons [3-4], unpredictable bits created by smartphones and other internet-connected devices [5]. After passing by distant polarization beam splitters the 'photons' produce two time-series of clicks on distant detectors coded by corresponding random variables: $A_x = \pm 1$ and $B_y = \pm 1$.

In Bell 64 [1,2] local realistic hidden variable model (LRHVM), a source is producing a mixed statistical ensemble Λ of particle pairs carrying strictly correlated properties λ and outcomes ± 1 are predetermined by these properties for all the settings. The randomness of the outcomes is created at the source and not during the measurement:

$$E(A_x B_y) = \sum_{\lambda \in \Lambda} A_x(\lambda) B_y(\lambda) p(\lambda)$$
 (1)

where $A_x(\lambda) = \pm 1$ and $B_y(\lambda) = \pm 1$. According to the model (1), there exists a counterfactual joint probability distribution (JP) of random variables describing the outcomes obtained in all different experimental setting. In particular:

$$E(A_x A_{x'} B_y B_{y'}) = \sum_{\lambda \in \Lambda} A_x(\lambda) A_{x'}(\lambda) B_y(\lambda) B_{y'}(\lambda) p(\lambda)$$
 (2)

Using (1) and (2) one derives easily Clauser-Horne-Shimony –Holt (CHSH) inequalities [6]:

$$|E(A_{r}B_{v}) + E(A_{r}B_{v}) + E(A_{r}B_{v}) - E(A_{r}B_{v})| \le 2.$$
 (3)

As Fine demonstrated [7, 8], CHSH are necessary and sufficient conditions for the existence of a counterfactual JP of A_x , A_x , B_y and B_y .

In LRHVM [1], λ describe only entangled 'particle pairs', thus it is reasonable to assume that setting variables and hidden variables in (1) are stochastically independent:

$$p(x, y, \lambda) = p(x, y) p(\lambda), p(x y | \lambda) = p(x, y), p(\lambda | x, y) = p(\lambda).$$
 (4)

The stochastic independence (4) is called: no-conspiracy, freedom of choice, free will or measurement independence (MI) [9]. Since the inequalities (3) are violated for some settings by quantum predictions and by experimental data, it has been debated which assumptions in the model (1) might be relaxed. One might relax locality (e.g. $A_x=A_x(\lambda, y)$ and $B_y(\lambda, x)$) or relax MI [10, 11, 12] and assume that: $p(\lambda|x, y)\neq p(\lambda)$.

However, the relaxation of locality would allow for spooky influences between distant measuring instruments and $p(x y | \lambda) \neq p(x, y)$ is often believed to imply the *super-determinism* [13]. Such beliefs are unjustified. As we explained in [14,15], conditional probabilities in Bayes Theorem do not admit, in general, any causal interpretation and the violation of MI does not constraint *experimenters'* freedom of choice, which is in fact the prerequisite of science.

We also explained, that MI should be called *noncontextuality*, because it allows to describe random experiments performed in incompatible experimental settings using a counterfactual joint probability distribution (JP) on a unique probability space Λ . The violation of Bell inequalities proves only that pairwise expectations in the inequality (3) are not marginal expectations derived from a non-existing JP [16, 17]. CHSH are simply *noncontextuality inequalities* for 4-cyclic scenarios [18].

By embedding the Bell causal structure in a larger causal network ,Rafael Chavez et al. [12] prove , that it is not necessary to evoke *super-determinism* in order to justify the violation of MI . They assume the independence of the variables that causally determine the settings and investigate how they might become correlated with hidden variables (e.g., when the cosmic photons enter the laboratory [3, 4]). Using their extended causal networks they derive a contextual probabilistic hidden variable model on which their further correct results are based:

$$p(a,b \mid x,y) = \sum_{\lambda} p(a \mid x,\lambda) p(b \mid y,\lambda) p(\lambda \mid x,y)$$
 (5)

However, they do not realize that calling the statistical independence (4) *freedom of choice* is misleading. They also seem to ignore, that that the simplest way to justify the contextual probabilistic model (5) is the inclusion of local setting dependent hidden variables describing the measuring instruments at the moment of the their interaction with the correlated photonic signals arriving from the source.

Such models were proposed several years ago and have been recently updated and reviewed extensively in [15,17]. They provide a locally causal explanation not only of imperfect correlations between distant clicks, but also the explanation of an apparent violation of Einsteinian non-signalling discovered in the data of several Bell Tests [19-22].

Let us consider a spin polarisation correlation experiment (SPCE) using a protocol similar to that of Weihs et al. [23]. Alice and Bob have each two detectors and register time series of clicks. From raw data one has to extract final samples in order to estimate correlations and perform a Bell test. A detailed discussion, how it is done may be found in [19, 20, 23, 24]. As we explained in [14]:

- 1. Raw time-tagged data are two samples: $S_A(x, y) = \{(a_k, t_k) \mid k=1...n_x\}$ and $S_B(x, y) = \{(b_m, t_m^2) \mid j=1...n_y\}$ with $a_k = \pm 1$ and $b_m = \pm 1$.
- 2. Using fixed synchronized time-windows of width W and keeping only time -windows in which there is no click at all or a click on one of Alice's or/and Bob's detectors two new samples are created: $S_A(x, y, W) = \{a_s \mid s=1,...N_x\}, S_B(x, y, W) = \{b_t \mid t=1...N_y\}$ with $a_s=0$, ± 1 and $b_t=0$, ± 1
- 3. Now by keeping only synchronized time-windows, in which both Alice and Bob observed a click on one of their detectors, new sample of paired outcomes is obtained: $S'_{AB}(x, y, W) = \{(a_r, b_r) | r=1,...N_{xy}\}$ with $a_r=\pm 1$ and $b_r=\pm 1$.

Samples, obtained at the step 2, may be described by the following non-signalling contextual hidden variable model

$$E(A_x B_y) = \sum_{\lambda \in \Lambda_{xy}} A_x(\lambda_1, \lambda_x) B_y(\lambda_2, \lambda_y) p_x(\lambda_x) p_y(\lambda_y) p(\lambda_1, \lambda_2)$$
(6)

where A_x (λ_1 , λ_x)=0,±1, B_y (λ_2 , λ_y)=0,±1. We have 4 random variables (L_1 , L_2 , L_x , L_y) taking values λ = (λ_1 , λ_2 , λ_x , λ_y), where (λ_1 , λ_2) describe photonic signals and (λ_x , λ_y) the measuring instruments, how they are 'perceived' by the signals at the moment of interaction. Four incompatible experiments are described by probability distributions defined on disjoint hidden variable spaces:

$$\Lambda_{xy} = \Lambda_{12} \times \Lambda_x \times \Lambda_y; \Lambda_{x'y} = \Lambda_{12} \times \Lambda_{x'} \times \Lambda_y; \Lambda_{xy'} = \Lambda_{12} \times \Lambda_x \times \Lambda_{y'}; \Lambda_{x'y'} = \Lambda_{12} \times \Lambda_{x'} \times \Lambda_{y'}$$
(7)

where $\Lambda_x \cap \Lambda_{x'} = \Lambda_y \cap \Lambda_{y'} = \emptyset$. Probability distributions of hidden variables depend on the settings: $p(\lambda|x, y) = p_x(\lambda_x) p_y(\lambda_y)$ and:

$$p(a,b \mid x,y) = \sum_{\lambda \in \Lambda_{ab}} p_x(\lambda_x) p_y(\lambda_y) p(\lambda_1,\lambda_2)$$
(8)

where $\Lambda_{ab} = \{\lambda \in \Lambda_{xy} | A_x(\lambda_1, \lambda_x) = a \text{ and } B_x(\lambda_2, \lambda_y) = b\}$. Using the model (6-8) one may not derive CHSH.

In order to perform Bell Test (3) the samples obtained in the step 3 have to be used. From (6-8): we easily derive required expectations:

$$E(A_x B_y \mid A_x B_y \neq 0) = \sum_{\lambda \in \Lambda'_{xy}} A_x(\lambda_1, \lambda_x) B_y(\lambda_2, \lambda_y) p_{xy}(\lambda)$$
(9)

$$E(A_x \mid A_x B_y \neq 0) = \sum_{\lambda \in \Lambda'_{xy}} A_x(\lambda_1, \lambda_x) p_{xy}(\lambda)$$
 (10)

$$E(\mathbf{B}_{y} \mid A_{x}B_{y} \neq 0) = \sum_{\lambda \in \Lambda'_{xy}} B_{y}(\lambda_{1}, \lambda_{y}) p_{xy}(\lambda)$$
(11)

where $p_{xy}(\lambda) = p_x(\lambda_x) p_y(\lambda_y) p(\lambda_1, \lambda_2)$, $\Lambda_{xy} = \Lambda_1 \times \Lambda_2 \times \Lambda_x \times \Lambda_y$ and

$$\Lambda_{xy}^{2} = \left\{ \lambda \stackrel{.}{\circ} \Lambda_{xy} \mid A_{x} \left(\lambda_{1}, \lambda_{x} \right) \neq 0, B_{y} \left(\lambda_{2}, \lambda_{y} \right) \neq 0 \right\}. \tag{12}$$

The model (9-12) allows explaining the reported violation of Einsteinian non-signalling:

$$E(A_x \mid A_x B_y \neq 0) \neq E(A_x \mid A_x B_y \neq 0); E(B_y \mid A_x B_y \neq 0) \neq E(B_y \mid A_x B_y \neq 0)$$
 (13)

This apparent violation of non-signalling does not prove the nonlocality of Nature, it is the result of the setting dependent post -selection (PSL) of events, such that both Alice and Bob register clicks on one of their detectors in corresponding synchronized time-windows.

The model (6-13) contains enough free parameters to fit experimental data [17] and to reproduce any violation of the inequality (3). The violation of CHSH inequality is not surprising, because it may not be derived for the pairwise expectations defined by (9).

The inequalities (13) prove that random variables describing the data obtained in the step 3 are inconsistently connected, thus a rigorous description of Bell Test should be done using the *Contextuality –by- Default* (CbD) approach of Dzhafarov and Kujala [25-28]. In CbD, random variables measuring the same content in different contexts are *stochastically unrelated* and they are labelled by contexts in which they are measured.

Therefore in Bell Tests we do not have a 4- cyclic system $X = \{A_x, A_{x'}, B_x, B_{x'}\}$ of 4 random variables but a system $X' = \{A_{xy}, A_{xy'}, B_{xy}, B_{x'y}, A_{x'y}, A_{x'y'}, B_{xy'}, B_{xy'}\}$ of 8 binary inconsistently connected random variables : $E(A_{xy}) \neq E(A_{xy'}), \dots E(B_{xy'}) \neq E(B_{x'y'})$ [15]. The only inequality which may be derived for these random variables without additional assumptions is:

$$S' = E(A_{xy}B_{yy}) + E(A_{yy}B_{yy'}) + E(A_{yy}B_{yy'}) + E(A_{yy}B_{yy}) - E(A_{yy}B_{yy}) - E(A_{yy}B_{yy}) \le 4$$
 (14)

where $E(A_{xy}B_{xy}) = E(A_xB_y \mid A_xB_y \neq 0)$ etc. By imposing maximal couplings between observables measuring the same content in different contexts one derives more restrictive inequalities, which may be compared with the experimental data [15].

There is an intimate relation between probabilistic models and experimental protocols [16, 29]. Let us analyze this relationship in the contextual model (6-8). For each setting (x, y), we have 4 random variables (X,Y,A_x,B_y) describing observed outcomes; (x, y, a_x, b_y) and 4 hidden random variables:

 $(L_1,L_2,\,L_x,\,L_y)$ taking values $(\lambda_1,\,\lambda_2,\,\lambda_x\,,\lambda_y)$, The experimental protocol may be subdivided into the following "random experiments":

- 1. Generate variables $(\lambda_1, \lambda_2) \in \Lambda_{12}$ describing and an entangled pair.
- 1. Generate labels (x, y).
- 2. Generate variables $\lambda_r \in \Lambda_r$ and $\lambda_v \in \Lambda_v$ describing the instruments.
- 3. Evaluate: $a_x = A_x(\lambda_1, \lambda_x)$ and $b_y = B_y(\lambda_2, \lambda_y)$.
- 4. Output: x, y, a_x, b_y .

The setting variables X, Y are causally independent from **all** other random variables. They also do not stochastically depend on L_1 and L_2 . The variables L_1 and L_2 are stochastically dependent. The variables A $_x$, B $_y$ are causally independent, but are stochastically dependent. The setting variables (X, Y) and instrument variables (L_x, L_y) are also stochastically dependent:

$$p(\lambda) = p_x(\lambda_x) p_y(\lambda_y) p(x, y) p(\lambda_1, \lambda_2) = p(x, y, \lambda) \Rightarrow p(x, y \mid \lambda) = \frac{p(x, y, \lambda)}{p(\lambda)} = 1$$
 (15)

It does not mean that the experimenters' freedom of choice is compromised. It means only: if a "hidden event' $\{\lambda_1, \lambda_2, \lambda_x, \lambda_y\}$ "happened", thus the settings (x, y) were chosen.

Therefore, the stochastic independence of setting and hidden variables (4) should be called *noncontextuality* and not *freedom of choice*. A much more detailed discussion of MI, Bayes Theorem and its implications may be found in [14].

Misleading terminology is a source of speculations about *super-determinism*, *retro-causation*, and *quantum nonlocality* not only on the social media but also in serious scientific journals. It is high time to understand that adding the instrument variables is the simplest explanation of statistical dependence of setting and hidden variables.

Some incorrectly believe that Bell rejected this solution in 1970, because after averaging over instrument variables in (6), CHSH inequalities may still be derived [30]. However he did not realized, that after such averaging the resulting probabilistic model describes different random experiments with experimental protocols which are impossible to implement [14, 17, 29].

Long time ago, several authors explained, that the violation of Bell inequalities proved only, that incompatible spin polarization correlation experiments do not allow for a *noncontextual* description in terms of a joint probability on a unique probability space e.g. [7, 8, 16. 31-37]. We cited here only few papers more references may be found in [17] and in the papers cited therein.

Bell inequalities were proven assuming predetermination of experimental results, called also local realism or counterfactual definiteness (CFD). The inequalities are violated, thus as Peres [38] resumed: unperformed experiments have no results.

The results do not exist because in quantum mechanics measuring instruments play an active role. Nieuwenhuizen [39.40] pointed out that LRHVM suffered from incurable *contextuality loophole*, because this model did not include correctly variables describing measuring instruments.

Some authors claim that inequalities are violated because of *quantum nonlocality*. Khrennikov [41] clearly explained that *quantum nonlocality* is a misleading notion, which should be abandoned.

Therefore the important resource in quantum information is not the violation of *freedom of choice*, as it was called in [11,12], but *contextuality*. It is consistent with the contextual character of quantum observables, Bohr's complementarity [44] and *KS-contextuality* [45].

Experimenters' freedom of choice is the prerequisite of science and the violation of inequalities in various Bell Tests does not allow doubting in its validity.

References

- 1. J. S. Bell, "On the Einstein Podolsky Rosen paradox," Physics Physique Fizika 1, 195 (1964).
- 2. J.S. Bell, *Speakable and unspeakable in quantum mechanics*. Cambridge, UK: Cambridge University Press (2004).
- 3. J. Handsteiner, A. S. Friedman, D. Rauch, J. Gallicchio, B. Liu, H. Hosp, J. Kofler, D. Bricher, M. Fink, C. Leung, et al., "Cosmic Bell test measurement settings from Milky Way stars," Phys. Rev. Lett. **118**, 060401 (2017).
- 4. D. Rauch, J. Handsteiner, A. Hochrainer, J. Gallicchio, A. S. Friedman, C. Leung, B. Liu, L. Bulla, S. Ecker, F. Steinlechner, et al., "Cosmic Bell test using random measurement settings from high-redshift quasars," Phys. Rev. Lett. **121**, 080403 (2018).
- 5. BIG Bell Test Collaboration and others, "Challenging local realism with human choices," Nature **557**, 212 (2018).
- 6. J. F. Clauser, M. A. Horne, A. Shimony, and R. A. Holt, "Proposed experiment to test local hidden-variable theories," Phys. Rev. Lett. 23, 880 (1969).
- 7. A. Fine, "Hidden variables, joint probability and the Bell inequalities," Phys. Rev. Lett. **48**, 291-295(1982).
- 8. A. Fine, "Joint distributions, quantum correlations, and commuting observables," J. Math. Phys. **23**, 1306-1310(1982).
- 9. W. Myrvold, M. Genovese and A. Shimony, "Bell's Theorem", The Stanford Encyclopedia of Philosophy(Fall 2020 Edition), Edward N. Zalta (ed.), URL = https://plato.stanford.edu/archives/fall2020/entries/bell-theorem/.
- 10. M.J.W. Hall, "Local deterministic model of singlet state correlations based on relaxing measurement independence", Phys.Rev.Lett.105, 250404. doi:10.1103/PhysRevLett.105.250404).
- 11. P. Blasiak, E. M. Pothos, J. M. Yearsley, C. Gallus, and E. Borsuk, "Violations of locality and free choice are equivalent resources in Bell experiments," Proc. Nat. Ac. Sci. **118**, e2020569118 (2021).
- R. Chaves, G. Moreno, E. Polino, D. Poderini, I. Agresti, A. Suprano, M. R. Barros, G. Carvacho, E. Wolfe, A. Canabarro, R. W. Spekkens, and F. Sciarrino, "Causal Networks and Freedom of Choice in Bell's Theorem," PRX Quantum (2021)2040323, DOI: 10.1103/PRXQuantum.2.040323.
- 13. S. Hossenfelder and T. Palmer, "Rethinking superdeterminism," Front. Phys. 8, 139 (2020).
- 14. M. Kupczynski, "Can we close the Bohr-Einstein quantum debate?," Phil.Trans.R.Soc.A., 2017, 20160392., DOI: 0.1098/rsta.2016,0392.
- 15. M. Kupczynski, "Contextuality-by-Default Description of Bell Tests: Contextuality as the Rule and Not as an Exception," Entropy 2021, 23(9), 1104; https://doi.org/10.3390/e23091104.
- 16. M. Kupczynski, "Bertrand's paradox and Bell's inequalities," Phys.Lett. A121, 205-207(1987).
- 17. M. Kupczynski, "Is the Moon there when nobody looks: Bell inequalities and physical reality," Front. Phys., 23 September 2020 |.https://doi.org/10.3389/fphy.2020.00273 (2020).
- 18. M. Araujo, M.T. Quintino., C. Budroni, M.T. Cunha., and A. Cabello, "All noncontextuality inequalities for the n-cycle scenario," Phys. Rev. A. **88**, 022118 (2013).
- 19. G. Adenier and A. Khrennikov., "Is the fair sampling assumption supported by EPR experiments,? J. Phys. B: Atom. Mol. Opt. Phys. **40**, 131-141(2007).
- 20. H. De Raedt , F. Jin , and K. Michielsen , "Data analysis of Einstein-Podolsky-Rosen-Bohm laboratory experiments," Proc. of SPIE 8832, 88321N1-11(2013).

- 21. G. Adenier and A. Khrennikov, "Test of the no-signaling principle in the Hensen loophole-free CHSH experiment," Fortschritte der Physik (Prog. Phys.) 2016, 65, 1600096. (2017).
- 22. A. Bednorz, "Analysis of assumptions of recent tests of local realism,", Phys. Rev. A **95**, 042118 (2017).
- 23. G. Weihs; T. Jennewein, C. Simon, H. Weinfurther and A. Zeilinger, "Violation of Bell's inequality under strict Einstein locality conditions," Phys. Rev. Lett. **81**, 5039-5043 (1998).
- 24. M. Kupczynski, "Is Einsteinian no-signalling violated in Bell tests?" Open Physics **15**, 739-753, DOI: https://doi.org/10.1515/phys-2017-0087(2017).
- 25. E.N. Dzhafarov and J.V. Kujala, "Contextuality is about identity of random variables," Physica Scripta T163:014009 (2014).
- 26. E.N. Dzhafarov. J.V. Kujala, and J.-Å. Larsson, "Contextuality in three types of quantum-mechanical systems, 'Foundations of Physics 7, 762-782.(2015).
- 27. J.V. Kujala, E.N. Dzhafarov and J-Å Larsson," Necessary and sufficient conditions for extended noncontextuality in a broad class of quantum mechanical systems," Physical Review Letters 115,150401(2015).
- 28. E.N. Dzhafarov, "Assumption-Free Derivation of the Bell-Type Criteria of contextuality/ Nonlocality," *Entropy* 2021, 23(11),1543; https://doi.org/10.3390/e2311154343 (2021).
- 29. M. Kupczynski, "Bell Inequalities, Experimental Protocols and Contextuality," Found. Phys., **45**, 735-753 (2015).
- 30. J.S.Bell ,"Introduction to the hidden variable question." In 49th Int. School of Physics 'Enrico Fermi': Foundations of quantum mechanics, Varenna, Italy, 29 June, pp. 171–181. Report CERNTH-1220 (1970).
- 31. L. Accardi, "Some loopholes to save quantum nonlocality," AIP.Conf. Proc.750, 1-19 (2005).
- 32. K. Hess and W. Philipp, "Bell's theorem: critique of proofs with and without inequalities," AIP Conf. Proc. **750**, 150-157 (2005).
- 33. K.Hess, K. Michielsen and H.De Raedt, "Possible experience: from Boole to Bell". *Europhys. Lett.* **87**, 60007. (doi:10.1209/0295-5075/87/60007) (2009).
- 34. A. Yu. Khrennikov," Bell's inequality: nonlocality, "death of reality", or incompatibility of random variables, "AIP Conf. Proc. **962**, 121-131(2007).
- 35. A. Khrennikov, "Contextual Approach to Quantum Formalism," Springer: Dordrecht, The Netherlands, 2009.
- 36. V. M. De Muynck, W. De Baere and H. Martens," Interpretations of quantum mechanics, joint measurement of incompatible observables and counterfactual definiteness," Found. Phys. 24, 1589-1664 (1994).
- 37. I. Pitowsky, "George Boole's 'conditions of possible experience' and the quantum puzzle," *Br. J. Phil. Sci.* **45**, 95–125. (doi:10.1093/bjps/45.1.95) (1994).
- 38. A. Peres, "Unperformed experiments have no results," *Am J Phys.***46**:745–7. doi: 10.1119/1.11393 (1978).
- 39. T.M. Nieuwenhuizen," Where Bell went wrong," AIP Conf. Proc. 1101, 127 (2009).
- 40. T.M. Nieuwenhuizen and M.Kupczynski, 'The contextuality loophole is fatal for derivation of Bell inequalities: Reply to a Comment by I. Schmelzer, Found. Phys. **47**, 316-319, DOI: 10.1007/s10701-017-0062-y (2017).
- 41. A. Khrennikov, Two Faced Janus of Quantum Nonlocality ,"Entropy **2020**, 22(3), 303; https://doi.org/10.3390/e22030303.
- 42. R. Raussendorf, "Contextuality in measurement-based quantum computation," Physical Review A **88**, 022322.
- 43. M.Frembs, S. Roberts and S. D. Bartlett, "Contextuality as a resource for measurement-based quantum computation beyond qubits," *New J. Phys.* **20** 103011 (2018).
- 44. N.Bohr, The Philosophical Writings of Niels Bohr. Ox Bow Press: Woodbridge, UK, 1987.
- 45. S. Kochen and E.P. Specker, "The problem of hidden variables in quantum mechanics," Journal of Mathematics and Mechanics **17**, 59–87 (1967).

International Journal of Quantum Foundations 8 (2022)

Copyright © 2022 by Marian Kupczynski. This article is an Open Access article distributed under the terms and conditions of the Creative Commons Attribution license (http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction, provided the original work is properly cited.