

High-Dimensional Inference over Networks: Linear Convergence and Statistical Guarantees

Ying Sun^{1*}, Marie Maros², Gesualdo Scutari², Guang Cheng^{3*}

¹The Pennsylvania State University, ysun@psu.edu

²Purdue University, mmaros@purdue.edu, gscutari@purdue.edu

³The University of California, Los Angeles, guangcheng@ucla.edu

Abstract

We study sparse linear regression over a network of agents, modeled as an undirected graph and no server node. The estimation of the s -sparse parameter is formulated as a constrained LASSO problem wherein each agent owns a subset of the N total observations. We analyze the convergence rate and statistical guarantees of a distributed projected gradient tracking-based algorithm under high-dimensional scaling, allowing the ambient dimension d to grow with (and possibly exceed) the sample size N . Our theory shows that, under standard notions of restricted strong convexity and smoothness of the loss functions, suitable conditions on the network connectivity and algorithm tuning, the distributed algorithm converges globally at a *linear* rate to an estimate that is within the centralized *statistical precision* of the model, $O(s \log d/N)$. When $s \log d/N = o(1)$, a condition necessary for statistical consistency, an ε -optimal solution is attained after $O(\kappa \log(1/\varepsilon))$ gradient computations and $O(\kappa/(1-\rho) \log(1/\varepsilon))$ communication rounds, where κ is the restricted condition number of the loss function and ρ measures the network connectivity. The computation cost matches that of the centralized projected gradient algorithm despite having data distributed; whereas the communication rounds reduce as the network connectivity improves. Overall, our study reveals interesting connections between statistical efficiency, network connectivity & topology, and convergence rate in high dimensions.

Keywords— High-dimensional inference, distributed convex optimization, linear convergence

1 Introduction

Datasets with massive sample size and high-dimensionality are ubiquitous in modern science and engineering; examples include genomics and biomedical data, social media, financial time-series, and e-commerce data, just to name a few. The archetypal scenario is the one wherein data are generated, acquired, or stored at multiple sources at different points in time and space; the sheer volume and spatial/temporal disparity of data render centralized processing and storage on standalone machines prohibitory or infeasible. This has motivated in recent years a surge of interest in developing distributed methods that enable analytics and computations across multiple machines (hereafter referred to as agents), connected by a communication network. Broadly speaking, two architectures have captured most of the interest of research communities, namely: (i) the *master-worker* topology and (ii) a general, connected, multiagent architecture (a.k.a. *mesh network*); they are depicted in Fig. 1 and briefly commented next.

*This work was conducted when Sun and Cheng were at Purdue University.




Figure 1: Two network architectures: master-worker and mesh networks.

In a *master-worker* architecture there exists one (or multiple) node(s)—generally called master node(s)—that is connected to all the other nodes, the so-called workers. The workers store part of the data, run (in parallel) intermediate computations, and communicate (iteratively) suitable outcomes to the master node, which maintains and updates the authoritative copy of the optimization variables, producing eventually the final output. These infrastructures have been adopted in several applications to parallelize and decompose a variety of learning and optimization tasks (e.g., [23, 26, 5]). The statistical community is best acquainted with *divide-and-conquer* (D&C) methods whereby the master node combines the estimators produced by the workers using their local data set. The idea has been widely applied in statistical estimation and inference such as M-estimation [51, 10, 39], sparse high dimensional models [3, 22], PCA [14], matrix factorization [25], quantile regression [44, 9] and non- and semi-parametric methods including [32, 38, 52, 50, 21], just to name a few.

There are many applications, however, where these infrastructures may be impractical or unattractive. For instance, in large scale, the master node may become the bottleneck of the system: (i) its failure will cause the entire network to fail, (ii) its communication resources may become insufficient as the number of nodes grows (e.g., due to limited bandwidth), and (iii) the implementation of some algorithms may require prohibitive computation efforts on standalone machines [15] (e.g., when high dimensional and large data sets are processed). Also, there are scenarios where establishing a star topology (or a spanning tree) is simply not feasible or too expensive; this is the case, e.g., of wireless networks with low-power devices, which can communicate only with nodes in their physical proximity. To address some of these issues, a natural approach is to eliminate master nodes, leading to peer-to-peer systems (*mesh* networks) wherein each node is connected only to a subset of the other nodes (see Fig. 1). This however makes algorithms designed for master-worker networks not directly implementable, due to the lack of centralized nodes. This motivated a surge of studies producing a variety of distributed schemes suitable for mesh networks; see [27, 31, 2, 36] for some tutorials on the topic. The common approach in such designs is to interleave (iteratively) local computation steps at agents' sides with communication among neighboring nodes, the latter aiming at aligning agents' estimates and updating local information. Despite the vast literature on distributed optimization (see Sec. 1.2 for an overview of relevant works), existing results on mesh networks are of pure optimization type; they lack any statistical guarantee, even for simple high-dimensional learning tasks. This paper along with the companion work [19] represents a first step towards an understanding of statistical and computational guarantees of widely adopted distributed first order methods solving sparse regression over mesh networks.

1.1 Sparse linear regression over networks

We study sparse linear regression over a mesh network of m agents, modeled as a general connected, undirected graph. Each agent i takes n linear measures of an s -sparse signal $\theta^* \in \mathbb{R}^d$:

$$\mathbf{y}_i = \mathbf{X}_i \theta^* + \mathbf{n}_i, \quad (1)$$

where $\mathbf{y}_i \in \mathbb{R}^n$ is the vector of measurements, $\mathbf{X}_i \in \mathbb{R}^{n \times d}$ is the design matrix, and $\mathbf{n}_i \in \mathbb{R}^n$ is the observation noise. For simplicity we assume that all agents take the same number n of measurements, for a total over the network of $N = m \cdot n$. Our interest is in the high-dimensional setting: the ambient dimension d exceeds the sample size N .

A standard approach to estimate $\boldsymbol{\theta}^*$ from $\{(\mathbf{y}_i, \mathbf{X}_i)\}_{i=1}^m$ is solving the LASSO problem, whose constrained form reads

$$\hat{\boldsymbol{\theta}} \in \operatorname{argmin}_{\|\boldsymbol{\theta}\|_1 \leq r} \left\{ \mathcal{L}(\boldsymbol{\theta}) \triangleq \frac{1}{m} \sum_{i=1}^m \mathcal{L}_i(\boldsymbol{\theta}) \right\}, \quad (2)$$

where $\mathcal{L}_i$ is the least squares loss of agent i , defined as

$$\mathcal{L}_i(\boldsymbol{\theta}) \triangleq \frac{1}{2n} \|\mathbf{y}_i - \mathbf{X}_i \boldsymbol{\theta}\|^2, \quad (3)$$

and the ℓ_1 -norm constraint aims at promoting sparsity on the solution $\hat{\boldsymbol{\theta}}$.

To solve (2), we focus on distributed *first-order* methods. Specifically, we aim at decentralizing the projected gradient descent (PGD) algorithm. In a centralized setting (e.g., on master-worker architecture), the iterates generated by the PGD read: given $\boldsymbol{\theta}^0 \in \mathbb{R}^d$,

$$\boldsymbol{\theta}^{t+1} = \prod_{\|\boldsymbol{\theta}\|_1 \leq r} (\boldsymbol{\theta}^t - \gamma \nabla \mathcal{L}(\boldsymbol{\theta}^t)), \quad t = 0, 1, \dots, \quad (4)$$

where $\prod_{\|\boldsymbol{\theta}\|_1 \leq r}(\bullet)$ denotes the orthogonal projection of its argument onto the ℓ_1 -norm ball of radius $r > 0$, and $\nabla \mathcal{L}(\boldsymbol{\theta}^t)$ is the gradient of the loss $\mathcal{L}$ at $\boldsymbol{\theta}^t$. Classical optimization-based analyses (see, e.g., [33]), certifying linear convergence of the PGD under strong convexity and smoothness of $\mathcal{L}$, break down in high-dimensions: (i) $\mathcal{L}$ is convex but not strongly convex on the entire space, as the Hessian matrix $\mathbf{X}^\top \mathbf{X}$ is rank deficient; and (ii) for commonly used design matrices $\mathbf{X}$, the Lipschitz constant of $\nabla \mathcal{L}$ (the largest eigenvalue of $\mathbf{X}^\top \mathbf{X}$) grows indefinitely under the high-dimensional scaling $d/N \rightarrow \infty$; as a consequence, the step-size γ reduces to zero. A more thoughtful analysis exploiting the statistical setting (see [1]) showed that suitably restricted notions of strong convexity and smoothness are sufficient for the global linear convergence of PGD. In particular, these restricted strong convexity (RSC) and smoothness (RSM) do hold with high probability (w.h.p.) for the loss $\mathcal{L}$ along the iterates (4). This leads to the following desirable statistical and computational guarantees for PGD (which hold w.h.p.). **(P.1)** The optimization error $\|\boldsymbol{\theta}^t - \hat{\boldsymbol{\theta}}\|$ decays *linearly* up to a nonzero tolerance. Under $s \log d/N = o(1)$, a condition necessary for statistical consistency, this tolerance is made of a smaller order than the intrinsic statistical error of the model:

$$\|\boldsymbol{\theta}^t - \hat{\boldsymbol{\theta}}\|^2 \leq \lambda_0^t \|\boldsymbol{\theta}^0 - \hat{\boldsymbol{\theta}}\|^2 + o(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|^2), \quad (5)$$

for $\lambda_0 \in (0, 1)$. Therefore, every limit point $\boldsymbol{\theta}^\infty$ of $\{\boldsymbol{\theta}^t\}$ solves the linear regression problem up to the statistical precision, that is

$$\|\boldsymbol{\theta}^\infty - \boldsymbol{\theta}^*\| = \|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\| + o(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|).$$

(P.2) Under the high-dimensional scaling $s, d/N \rightarrow \infty$ and $s \log d/N = o(1)$, the rate λ_0 approaches $1 - \kappa^{-1}$, remaining thus invariant, where $\kappa \geq 1$ is the restricted condition number of the centralized loss $\mathcal{L}$.

Open questions: The PGD (4) is not implementable on mesh networks: agents cannot compute the full gradient $\nabla \mathcal{L}$, as they do not have access to the gradients $\nabla \mathcal{L}_j$ of all the others. This poses the following natural questions: **(i)** Given (2) with data distributed over a mesh network, can one decentralize the PGD while retaining the desirable properties **P.1** (linear rate up to *centralized* statistical precision) and **P.2** (rate invariance under high dimensional scaling)? Of particular interest is the unexplored regime wherein the local sample size n is not sufficient for statistical consistency while the total one N is so. **(ii)** What is the role/impact of the network (and its topology) towards **P.1** and **P.2**? **(iii)** How are sample and convergence rates affected by (e.g., scale with) the network parameters (connectivity, number of agents)?

To the best of our knowledge, current literature on distributed optimization does not provide a satisfactory answer to the above questions (see Sec. 1.2); new techniques and analyses are needed, revealing connections between statistical and computational efficiency of distributed algorithms in high-dimensional estimation, which is indeed the main contribution of this work (see Sec. 1.3).

1.2 Related works and challenges

The last decade has witnessed a flurry of works studying statistical procedures over master-worker architectures. As already mentioned, the majority of these works focused on divide and conquer methods, e.g., [51, 10, 39, 3, 22, 14, 25, 44, 9, 32, 38, 52, 50, 21]. These algorithms only need one round of communication from the workers to the server, but statistical guarantees are achieved under a constraint on the number of workers, which limits the range of applicability of these methods to small-size networks. To cope with this issue, iterative procedures have been proposed, minimizing a centralized risk function over all the data; examples include [20, 13, 45]. At the cost of multiple communication rounds, these methods do not impose any constraint on the number of worker nodes. However, none of the above algorithms is implementable on a mesh network, because of the lack of a server node.

Solving the LASSO problem (2) over mesh networks falls under the umbrella of distributed optimization. Despite the vast literature, given the focus of the paper, we comment next only relevant first-order methods solving (strongly) convex optimization problems over mesh networks modeled as undirected graphs.

Early attempts to decentralize the (sub)gradient algorithm have been proposed in [29, 30]; a prototypical form of such Distributed Gradient Algorithms (DGD) applied to (2) reads

$$\boldsymbol{\theta}_i^{t+1} = \prod_{\|\boldsymbol{\theta}\|_1 \leq r} \left(\sum_{j \in \mathcal{N}_i}^m w_{ij} \boldsymbol{\theta}_j^t - \gamma \nabla \mathcal{L}_i(\boldsymbol{\theta}_i^t) \right), \quad t = 0, 1, \dots, \quad (6)$$

where $\boldsymbol{\theta}_i^t$ is an estimate of the optimization variable $\boldsymbol{\theta}$ at agent i 's side after t iterations; $\mathcal{N}_i$ is the set of neighbors of agent i ; and w_{ij} 's are suitably chosen nonnegative weights. After receiving $\boldsymbol{\theta}_j^t$ from its neighbors $j \in \mathcal{N}_i$, in (6), each agent i first averages the $\boldsymbol{\theta}_j^t$'s by the weights w_{ij} 's, and then corrects the resulting iterate by the local gradient $\nabla \mathcal{L}_i(\boldsymbol{\theta}_i^t)$ (recall that agents do not have access to the entire gradient $\nabla \mathcal{L}(\boldsymbol{\theta}_i^t)$)—with $\gamma > 0$ being a common step-size—followed by the projection onto the ℓ_1 ball. The goal of the averaging step is to enforce consensus among the local estimates $\boldsymbol{\theta}_i$'s.

Subsequent works studied alternative decentralizations of the gradient algorithm [7, 8, 48, 49]; roughly speaking, the difference among such DGD-like schemes is in what information gets exchanged in the communication step and in the precise way this information and agents' gradients are combined to form the update, not necessarily using (6), as well as in the assumptions made about agents' losses and network topology. Although different in the updates, convergence guarantees of such DGD-like algorithms, including (6), can be roughly summarized as follows: (i) when the agents' loss functions $\mathcal{L}_i$ are smooth, strongly convex (or the average-loss $\mathcal{L}$ is smooth and satisfies the Kurdyka-Lojasiewicz property [49, 11]) and there are no constraints, DGD-like schemes equipped with a constant step-size, converge at linear rate but only to a neighborhood of the minimizer [48, 49]. The size of the neighborhood is monotonic in the step-size. Convergence to the exact minimizer is achieved only using diminishing step-size rules, at the price of slower sublinear rate [49, 18].

These results are unsatisfactory when applied to the LASSO problem. First, they show a gap with the convergence properties of the centralized PGD (see Sec. 1.1); for instance, they would predict sublinear convergence rate (recall $d > N$), while PGD enjoys linear convergence up to statistical precision (property **P.1**). Second, these studies are of pure optimization type: no statistical analysis of the limit points is provided, and they are not suitable for the high-dimensional regime when d grows faster than N . In fact, they all require *global smoothness* of the loss function which, as discussed in Sec. 1.1, cannot hold for $\mathcal{L}$ under the high-dimensional scaling $d/N \rightarrow \infty$. Third, these theoretical predictions are confuted by experiments. In Fig. 2 (left panel), we plot the estimation error generated by DGD (6) versus the iterations, for different step-size values γ ; the estimation error is defined as $(1/m) \cdot \sum_{i=1}^m \|\boldsymbol{\theta}_i^t - \boldsymbol{\theta}^*\|^2$. As benchmark, we also plot the estimation error achieved by the PGD (4). The curves show that DGD converges linearly, and not sublinearly as theory predicts; furthermore, it slows down as smaller statistical errors are required. When $d \rightarrow \infty$, with $N = O(s \log d)$ and fixed network topology, the right panel of the figure shows that the rate of DGD *deteriorates*, thus violating **P.2**. These experiments suggest that DGD does not inherit the desirable properties **P.1** and **P.2** of PGD (4).




Figure 2: DGD algorithm (6) in high-dimension. **Left panel:** Estimation error vs. iterations for different step-size values: statistical accuracy is achieved at the price of slow convergence. **Right panel:** Increasing (d, N) , with $N = O(s \log d)$: DGD slows down! This contrasts with properties **P.1** and **P.2** of PGD (4).

A first statistical analysis of a DGD-like algorithm for the Lagrangian formulation of the LASSO problem has been developed in our companion paper [19]. For commonly used random design matrices $\mathbf{X}$ and under certain regularity conditions, the iterates generated by the algorithm enter an ε -neighborhood of a statistically optimal estimate of θ^* after

$$\mathcal{O}\left(\frac{\kappa}{1-\rho} \cdot d \cdot m^2 \log m \cdot \log \frac{1}{\varepsilon}\right) \quad (7)$$

number of iterations, where $\rho \in [0, 1)$ is a measure of the connectivity of the network (the smaller ρ , the more connected the graph); and κ is the restricted condition number of $\mathcal{L}$. This shows that this instance of DGD achieves centralized statistical accuracy at a linear rate, which however scales undesirably as $\mathcal{O}(d)$, proving the ‘speed-accuracy dilemma’, as suffered also from the DGD (6) (see Fig. 2).

The above discussion seems to support, theoretically and numerically, the fact that known DGD-like algorithms are not suitable decentralization of the PGD in high-dimensions, if one wants to retain properties **P.1** and **P.2**. To cope with the speed-accuracy dilemma of DGD (already known in the literature when $d < N$), a variety of other distributed algorithms have been proposed. While different in the execution, they share a common idea, that is, employing some form of correction of the local gradient direction in the agents’ updates, so that a constant stepsize can be used, still preserving exact convergence to the minimizer of the average-loss. Examples include: gradient tracking methods [34, 28, 47, 12, 43] and primal-dual schemes [17, 41, 16, 40, 42, 24], just to name a few. Linear convergence is certified under global smoothness of agents’ losses and strong convexity of the average-loss (or local losses). These algorithms converge sublinearly when the problem is merely convex. This means that, if applied to the LASSO problem (2) (with $d > N$), the theory predicts sublinear convergence, which contrasts with linear convergence of PGD (4) [1]. Furthermore, these studies lack of any statistical analysis and, as those for DGD-like methods, they break down under high-dimension scaling $d/N \rightarrow \infty$.

In conclusion, current literature on distributed optimization leads to the apparent dilemma that even though the LASSO estimator $\hat{\theta}$ defined in (2) can consistently estimate θ^* when $d > N$, it is not computable in a distributed way. New analysis is needed to understand the behavior of distributed algorithms in high-dimensions. This paper takes a first step toward this direction, showing that existing optimization theory on distributively solving the LASSO problem can be strengthened by injecting the statistical properties into the analysis. Our major contributions are summarized next.

1.3 Major contributions

- (i) **Distributed PGD via gradient tracking:** We study a decentralization of the PGD for (2) that leverages the idea of gradient tracking [47, 12, 43]: in contrast with DGD-like approaches, agents employ a correction of their gradient direction $\nabla \mathcal{L}_i(\theta_i^t)$ forming a local estimate of the centralized

gradient $\nabla \mathcal{L}(\theta_i^t)$. This is achieved via a suitably designed dynamic consensus mechanism (a.k.a. gradient tracking), which only requires neighboring information. We name the algorithm *Network LASSO* (NetLASSO).

- (ii) **Statistical-computational guarantees:** Under the same notions of RSC & RSM of $\mathcal{L}$ as in the centralized setting [1] (and RSM of $\mathcal{L}_i$'s), and suitable conditions on the RSC/RSM parameters (implied by $s \log d/N = o(1)$), network connectivity, and algorithm tuning, the sequence generated by NetLASSO provably converges at a *linear rate* to an estimate whose distance to $\hat{\theta}$ is *smaller than the centralized statistical precision*—mimicking thus the behavior of PDG (property **P.1**). Specifically, to enter an ε -neighborhood of such a statistically optimal estimate, it takes

$$O(\kappa \log(1/\varepsilon)) \quad (8)$$

iterations (gradient evaluations), as long as $\rho \leq \text{poly}(m, \kappa)^{-1}$, that is, the network is sufficiently connected. Recall that κ is the restricted condition number of $\mathcal{L}$ and ρ is the network connectivity. The condition on ρ , when not met by the given graph, can be enforced via multiple rounds of communication per iteration, resulting in

$$O\left(\frac{\kappa}{1-\rho} \log(m\kappa) \log(1/\varepsilon)\right) \quad (9)$$

overall communication rounds.

This closes the gap between statistics and optimization theory in distributed settings, showing that statistically optimal estimators are in fact computable (at linear rate), under *centralized* statistical consistency, $s \log d/N = o(1)$. Furthermore, the iteration complexity (8) matches that of the PGD (in order).

- (iii) **Sample/network scaling:** The communication complexity (9) is proved to be invariant, under high-dimensional scaling $d, N \rightarrow \infty$, with $s \log d/N = O(1)$ and for a fixed network—this matches property **P.2** of PGD. We also study the scalability of (9) with respect to the network size m , revealing tradeoffs between communication rounds and communication costs for different graph topologies, as summarized in Table 1. This sheds light on which network architectures are more favorable for high-dimensional estimation. For instance, among the commonly used network architectures listed in Fig. 3, NetLASSO running on the Erdős-Rényi graph with edge connecting probability $p = \log m/m$ achieves the lowest communication cost at the busiest node, and ties with the implementation on the star topology in the amount of total channel use.

1.4 Paper organization

The rest of the paper is organized as follows. Sec. 2 introduces the setup and necessary background. The NetLASSO algorithm is given in Sec. 3. Statistical-computational guarantees of the algorithm are stated and commented in Sec. 4; proofs are reported in Sec. 5, with some complementary and intermediated results provided in the supporting material. Numerical simulations corroborating our theory are discussed in Sec. 6. Finally, Sec. 7 draws some conclusions.

2 Setup and Background

In this section we introduce the main assumptions on the data model and optimization problem (cf. Sec. 2.1) as well as the network setting (cf. Sec. 2.2) underlying our analysis.

2.1 Problem setting

We recall here the key properties of the LASSO problem (2), which enable fast convergence of the PGD, namely: the geometry of the constraint set and the landscape of the loss $\mathcal{L}$. They will serve as foundations for the design and analysis of the proposed decentralized LASSO algorithm.

The geometry of the constraint set. The ℓ_1 ball-constraint plays a crucial role in shaping the sparsity of feasible points. Specifically, for any $\boldsymbol{\theta} \in \mathbb{R}^d$ satisfying $\|\boldsymbol{\theta}\|_1 \leq r$, $\boldsymbol{\theta} - \hat{\boldsymbol{\theta}}$ is approximately s -sparse, as quantified next.

Lemma 1 ([1]). *Let $\hat{\boldsymbol{\theta}}$ be any optimal solution of Problem (2) such that $\|\hat{\boldsymbol{\theta}}\|_1 = r$ (the constraint is active). Then for any $\|\boldsymbol{\theta}\|_1 \leq r$, there holds*

$$\|\boldsymbol{\theta} - \hat{\boldsymbol{\theta}}\|_1 \leq 2\sqrt{s}\|\boldsymbol{\theta} - \hat{\boldsymbol{\theta}}\| + \underbrace{2\|\Delta^*\|_1 + 2\sqrt{s}\|\Delta^*\|}_{\nu}, \quad (10)$$

where $\Delta^* \triangleq \hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*$ is the statistical error.

The landscape of the loss function: As already commented, fast convergence of the PGD does not require strong convexity and smoothness of $\mathcal{L}$ globally, but only on a restricted set of directions. Given the sparsity of the optimization error $\boldsymbol{\theta}^t - \hat{\boldsymbol{\theta}}$ (Lemma 10), it is sufficient to enforce such a benign landscape of $\mathcal{L}$ along sparse directions, as captured by the following widely-used notions of RSC and RSM [1].

Assumption 2 (Global RSC/RSS). *Given $\mathbf{X} = [\mathbf{X}_1^\top, \dots, \mathbf{X}_m^\top]^\top \in \mathbb{R}^{N \times d}$, with $N = n \cdot m$, the following conditions hold:*

$$\begin{aligned} \text{(Global RSC/RSM)} \quad & \frac{\|\mathbf{X}\mathbf{u}\|^2}{N} \geq \mu_\Sigma \|\mathbf{u}\|^2 - \tau_\mu \|\mathbf{u}\|_1^2, \\ & \frac{\|\mathbf{X}\mathbf{u}\|^2}{N} \leq L_\Sigma \|\mathbf{u}\|^2 + \tau_g \|\mathbf{u}\|_1^2, \end{aligned} \quad \forall \mathbf{u} \in \mathbb{R}^d, \quad (11)$$

where (μ_Σ, τ_μ) and (L_Σ, τ_g) are positive RSC/RSM parameters.

The RSC/RSM conditions above can be certified with high probability by a variety of random design matrices $\mathbf{X}$. Here we consider the following.

Proposition 3 ([35]). *Given m i.i.d. distributed random matrices $\mathbf{X}_i \in \mathbb{R}^{n \times d}$ drawn from the Σ -Gaussian ensemble. Let $\mathbf{X} = [\mathbf{X}_1^\top, \dots, \mathbf{X}_m^\top]^\top \in \mathbb{R}^{N \times d}$, with $N = n \cdot m$. Then, there exist universal constants $c_0, c_1 > 0$ such that*

$$\begin{aligned} \text{(Global RSC/RSM)} \quad & \frac{\|\mathbf{X}\mathbf{u}\|^2}{N} \geq \frac{1}{2} \|\Sigma^{1/2} \mathbf{u}\|^2 - c_1 \zeta \frac{\log d}{N} \|\mathbf{u}\|_1^2, \\ & \frac{\|\mathbf{X}\mathbf{u}\|^2}{N} \leq 2 \|\Sigma^{1/2} \mathbf{u}\|^2 + c_1 \zeta \frac{\log d}{N} \|\mathbf{u}\|_1^2, \end{aligned} \quad \forall \mathbf{u} \in \mathbb{R}^d; \quad (12)$$

with probability greater than $1 - \exp(-c_0 N)$, where $\zeta = \max_j \Sigma_{jj}$.

We conclude this section introducing a local RSM condition for $\mathbf{X}_i$, instrumental to prove convergence of the proposed distributed algorithm.

Assumption 4 (Local RSM). *For each local design matrix $\mathbf{X}_i$, the following holds:*

$$\frac{\|\mathbf{X}_i \mathbf{u}\|^2}{n} \leq \ell_\Sigma \|\mathbf{u}\|^2 + \tau_\ell \|\mathbf{u}\|_1^2, \quad \forall \mathbf{u} \in \mathbb{R}^d, \quad (13)$$

where (ℓ_Σ, τ_ℓ) are positive RSM parameters.

Assumption 4 holds with high probability for Gaussian design matrices.

Proposition 5. *Given a random matrix $\mathbf{X}_i \in \mathbb{R}^{n \times d}$ drawn from the Σ -Gaussian ensemble, there exists universal constants $c_0, c_1 > 0$ such that*

$$\frac{\|\mathbf{X}_i \mathbf{u}\|^2}{n} \leq 16m \|\Sigma^{1/2} \mathbf{u}\|^2 + c_1 \zeta \frac{m \log d}{n} \|\mathbf{u}\|_1^2, \quad \forall \mathbf{u} \in \mathbb{R}^d, i \in [m] \quad (14)$$

with probability greater than $1 - \exp(-c_0 N)$, where $\zeta = \max_j \Sigma_{jj}$.

Proof. The proof is a slight modification of that in [35, Theorem 1], and is reported in Appendix I (supplementary material). $\square$

2.2 Network setting

The network of agents is modeled as an undirected graph $\mathcal{G} \triangleq (\mathcal{V}, \mathcal{E})$; where $\mathcal{V} = \{1, \dots, m\}$ is the set of m nodes and $\mathcal{E} \subseteq \mathcal{V} \times \mathcal{V}$ is the set of edges, corresponding to the communication links: $(i, j) \in \mathcal{E}$ iff there exists a communication link between agent i and j . We denote by $\mathcal{N}_i = \{j \in \mathcal{V} : (i, j) \in \mathcal{E}\} \cup \{i\}$ the set of neighbors of agent i (including the agent itself). We make the following standard assumption on the graph connectivity, which is necessary to achieve consensus over the network.

Assumption 6 (Connected network). *The graph $\mathcal{G}$ is connected.*

3 Decentralized projected gradient method with tracking

We aim at decentralizing the PGD (4) over mesh networks while preserving properties **P.1** and **P.2**. The challenge is to acquire, at each agent's side, faithful estimates of the global gradient $\nabla \mathcal{L}$ using only neighboring information and still guaranteeing convergence at fast rate. Our idea is to estimate locally the gradient $\nabla \mathcal{L}$ via gradient-tracking [47, 12, 43]. The resulting algorithm, termed Network Lasso (NetLASSO), is formally introduced in Algorithm 1, and commented next.

Each agent i maintains and updates iteratively a local copy $\theta_i \in \mathbb{R}^d$ of the global optimization variable θ along with the auxiliary variable $\mathbf{g}_i \in \mathbb{R}^d$ that estimates the centralized gradient $\nabla \mathcal{L}$; θ_i^t (resp. $\mathbf{g}_i^t$) denotes the value of θ_i (resp. $\mathbf{g}_i$) after t iterations. The algorithm alternates optimization steps [cf. (S.2)] and communications rounds among neighboring agents [cf. (S.1)]. In (S.2), given θ_i^t and $\mathbf{g}_i^t$, each agent solves the optimization problem (18), where $\mathbf{g}_i^t$ plays the role of the centralized gradient $\nabla \mathcal{L}(\theta^t)$ in (4). Therefore, (18) can be interpreted as the *local* counterpart of the PGD update (4). In (S.1), based upon exchange of the two vectors $\theta_i^{t-\frac{1}{2}}$ and $\mathbf{g}_i^{t-1} + \nabla \mathcal{L}_i(\theta_i^t) - \nabla \mathcal{L}_i(\theta_i^{t-1})$ with their immediate neighbors, each agent updates first the estimate $\theta_i^{t-\frac{1}{2}} \rightarrow \theta_i^t$ via the consensus step (16) and $\mathbf{g}_i^{t-1} \rightarrow \mathbf{g}_i^t$ via the perturbed consensus (17). The gradient-tracking update (17) can be informally explained as follows: At each iteration t , first, each agent i subtracts the outdated gradient $\nabla \mathcal{L}_i(\theta_i^{t-1})$ at the previous iteration and adds the new one $\nabla \mathcal{L}_i(\theta_i^t)$; this refreshes the “memory” of $\mathbf{g}_i$ and ensures that the sum of the $\mathbf{g}_i^t$'s is equal to the sum gradient $\sum_{i=1}^m \nabla \mathcal{L}_i(\theta_i^t)$. Then, agents perform a local averaging of the same form of (16), forcing thus consensus among the $\mathbf{g}_i$'s. If both θ_i 's and $\mathbf{g}_i$'s are asymptotically consensual (a fact that will be proved in Sec.5), then

$$\mathbf{g}_i^t \rightarrow \frac{1}{m} \sum_{j=1}^m \mathbf{g}_j^t = \frac{1}{m} \sum_{j=1}^m \nabla \mathcal{L}_j(\theta_j^t) \rightarrow \frac{1}{m} \sum_{j=1}^m \nabla \mathcal{L}_j(\theta^t) \quad \text{as } t \rightarrow \infty. \quad (15)$$

That is, $\mathbf{g}_i^t$ converges to $\nabla \mathcal{L}(\theta^t)$, as desired.

The weights $(w_{ij})_{ij}$ in (16)-(17) are free design quantities and subject to the following conditions, where $\mathcal{P}_K$ denotes the set of polynomials with degree no larger than K .

Assumption 7. *[On the weight matrix $\mathbf{W}$] The matrix $\mathbf{W} = (w_{ij})_{i,j=1}^m$ satisfies the following:*

(a) $\mathbf{W} = P^K(\overline{\mathbf{W}})$, where $P_K \in \mathcal{P}_K$ with $P_K(1) = 1$, and $\overline{\mathbf{W}} \triangleq (\bar{w}_{ij})_{i,j=1}^m$ has a sparsity pattern compliant with $\mathcal{G}$, that is

i) $\bar{w}_{ii} > 0$, for all $i = 1, \dots, m$;

ii) $\bar{w}_{ij} > 0$, if $(i, j) \in \mathcal{E}$; and $\bar{w}_{ij} = 0$ otherwise.

Furthermore, $\overline{\mathbf{W}}$ is doubly stochastic, that is, $\mathbf{1}^\top \overline{\mathbf{W}} = \mathbf{1}^\top$ and $\overline{\mathbf{W}} \mathbf{1} = \mathbf{1}$.

(b) Let $\mathbf{J} \triangleq \frac{1}{m} \mathbf{1} \mathbf{1}^\top$ be the projection onto the consensus space and ρ be the operator norm of $\mathbf{W} - \mathbf{J}$ induced by the ℓ_2 norm, i.e., $\rho \triangleq \|\mathbf{W} - \mathbf{J}\|_2$. The operator norm of $(\mathbf{W} - \mathbf{J})^t$, $t = 1, 2, \dots$, induced by the ℓ_∞ norm satisfies $\|(\mathbf{W} - \mathbf{J})^t\|_\infty \leq c_m \rho^t$, for some $c_m > 0$.

Remark 8. *Assumption 6 and Assumption 7(a) imply $\rho < 1$ and $\|(\mathbf{W} - \mathbf{J})^t\|_2 = \rho^t$. Using the norm bound $\|(\mathbf{W} - \mathbf{J})^t\|_\infty \leq \sqrt{m} \|(\mathbf{W} - \mathbf{J})^t\|_2$, we infer that Assumption 7(b) holds with $c_m \leq \sqrt{m}$.*

Algorithm 1: Network LASSO (NetLASSO)

Data: $\boldsymbol{\theta}_i^{\frac{1}{2}} = \boldsymbol{\theta}_i^0 = \mathbf{0}$, $\mathbf{g}_i^0 = \nabla \mathcal{L}_i(\boldsymbol{\theta}_i^0)$, for all $i \in [m]$;

Iterate: $t = 1, 2, \dots$

[S.1] **Communication:** Each agent i computes

$$\boldsymbol{\theta}_i^t = \sum_{j \in \mathcal{N}_i} w_{ij} \boldsymbol{\theta}_j^{t-\frac{1}{2}}, \quad (16)$$

$$\mathbf{g}_i^t = \sum_{j \in \mathcal{N}_i} w_{ij} \left(\mathbf{g}_j^{t-1} + \nabla \mathcal{L}_j(\boldsymbol{\theta}_j^t) - \nabla \mathcal{L}_j(\boldsymbol{\theta}_j^{t-1}) \right); \quad (17)$$

[S.2] **Local optimization:** each agent i solves

$$\boldsymbol{\theta}_i^{t+\frac{1}{2}} = \underset{\|\boldsymbol{\theta}_i\|_1 \leq r}{\operatorname{argmin}} (\mathbf{g}_i^t)^\top (\boldsymbol{\theta}_i - \boldsymbol{\theta}_i^t) + \frac{\gamma}{2} \|\boldsymbol{\theta}_i - \boldsymbol{\theta}_i^t\|^2. \quad (18)$$

Algorithm 2: PGD over star-networks

Data: $\boldsymbol{\theta}_i^0 = \mathbf{0}$, for all $i \in [m]$;

Iterate: $t = 1, 2, \dots$

[S.1] **Communications:**

(i) The master node broadcasts $\boldsymbol{\theta}^t$;

(ii) Each worker $j = 2, \dots, m$ evaluates $\nabla \mathcal{L}_j(\boldsymbol{\theta}^t)$ and sends it to the master;

[S.2] **Local optimization:** The master evaluates $\nabla \mathcal{L}(\boldsymbol{\theta}^t) = \frac{1}{m} \sum_{j=1}^m \nabla \mathcal{L}_j(\boldsymbol{\theta}^t)$ and solves

$$\boldsymbol{\theta}^{t+1} = \underset{\|\boldsymbol{\theta}\|_1 \leq r}{\operatorname{argmin}} \left\{ \nabla \mathcal{L}(\boldsymbol{\theta}^t)^\top (\boldsymbol{\theta} - \boldsymbol{\theta}^t) + \frac{\gamma}{2} \|\boldsymbol{\theta} - \boldsymbol{\theta}^t\|^2 \right\}. \quad (19)$$

Several rules of choosing $\overline{\mathbf{W}}$ have been proposed in the literature satisfying Assumption 7, such as the Laplacian, the Metropolis-Hasting, and the maximum-degree weights rules; see, e.g., [27] and references therein. When $K > 1$, K rounds of communications per iteration t are employed. For instance, this can be performed using the same given reference matrix $\overline{\mathbf{W}}$ (satisfying Assumption 7) in each communication exchange, resulting in $\bar{\rho} = \|\overline{\mathbf{W}} - \mathbf{J}\|_2$ and $\mathbf{W} = \overline{\mathbf{W}}^K$. Such a $\mathbf{W}$ satisfies Assumption 7, with $\rho = \bar{\rho}^K$. Faster information mixing can be obtained using suitably designed polynomials $P_K(\overline{\mathbf{W}})$, such as Chebyshev [46, 37] or orthogonal (a.k.a. Jacobi) [4] polynomials.

It is worth noting that the NetLASSO algorithm contains as special instance, the PGD when the graph $\mathcal{G}$ is a star-network. In this setting, by electing the first node as the master node (star center), a natural implementation of the PGD is described in Algorithm 2. Clearly, the iterates $\{\boldsymbol{\theta}^t\}$ generated by Algorithm 2 coincide with the iterates $\{\boldsymbol{\theta}_i^t\}$ of Algorithm 1, $i = 1, \dots, m$, using the mixing matrix $\mathbf{W} = \mathbf{W}'\mathbf{W}'^\top$, with $\mathbf{W}' = 1/\sqrt{m}[\mathbf{1}_m, \mathbf{0}_{m \times m-1}]$. Note that the resulting $\mathbf{W}$ is the same as that of a complete graph, but the implementation as in Algorithm 2 is more communication efficient.

Algorithm 1 can be interpreted as an instance of the framework recently introduced in [43] for strongly

convex optimization problems. However, as discussed in Sec. 1.2, the algorithmic tuning and convergence results in [43] are not suitable for the high-dimensional setting, which calls for a new convergence analysis.

4 Summary of Results

This section summarizes the main convergence properties of NetLASSO and their consequences. A formal statements of these results along with their proofs can be found in Sec. 5.

Our statements consists of two parts. First, we establish conditions under which the network average optimization error (in the ℓ_2 norm) $\frac{1}{m} \sum_{i=1}^m \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|^2$ shrinks linearly up to a tolerance $o(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|^2)$, and provide an explicit expression of the rate (cf. Sec. 4.1). Then, we show that for the statistical model subsumed in (1), this result holds with high probability (cf. Sec. 4.2). We also discuss the impact of the network topology on the algorithm's performance.

4.1 Convergence under RSC/RSM

Consider the LASSO problem (2), with data set $\{\mathbf{X}_i, \mathbf{y}_i\}_{i=1}^m$ generated according to the linear model (1) and ground truth $\boldsymbol{\theta}^*$ s -sparse. Assume that the global loss function $\mathcal{L}$ is RSC and RSM with parameters (μ_Σ, τ_μ) and (L_Σ, τ_g) , respectively. Further, each local loss function $\mathcal{L}_i$ is RSM with parameter (ℓ_Σ, τ_ℓ) . Define

$$\Delta_{\text{stat}} = \frac{\rho}{2L_\Sigma} \left(\frac{\ell_\Sigma}{\mu_\Sigma} \frac{5C_2^2 c_m^2}{(1-\rho)^2} \right) \tau_\ell \nu^2 + \frac{C_1(\tau_\mu + \tau_g)}{L_\Sigma} \nu^2, \quad \text{with} \quad \nu = 2\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_1 + 2\sqrt{s}\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|,$$

where $C_1, C_2 > 0$ are universal constants. Observe Δ_{stat} consists of a network dependent and a network independent term. Finally, the NetLASSO algorithm runs on a connected communication graph $\mathcal{G}$, with weight matrix $\mathbf{W}$ satisfying Assumption 7. Convergence is summarized next.

Theorem 18 (informal). *Let $\{(\boldsymbol{\theta}_i^t)_{i=1}^m\}$ be the sequence generated by the NetLASSO algorithm in the setting above; further assume that τ_μ and τ_g are sufficiently small and*

$$\rho \leq \text{poly} \left(m, \frac{s\tau_\ell}{\mu_\Sigma}, \frac{\ell_\Sigma}{\mu_\Sigma} \right)^{-1}. \quad (20)$$

Then for Problem (2) with optimal solution $\hat{\boldsymbol{\theta}}$ satisfying $\|\hat{\boldsymbol{\theta}}\|_1 = r$, it holds

$$\frac{1}{m} \sum_{i=1}^m \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|^2 \leq \left(\frac{1 - (2\kappa)^{-1} + C_1 s(\tau_\mu + \tau_g)/L_\Sigma}{1 - 2C_1 s\tau_g/L_\Sigma} \right)^t \cdot B + O(\Delta_{\text{stat}}), \quad (21)$$

where $\kappa = L_\Sigma/\mu_\Sigma$ is the restricted condition number of $\mathcal{L}$, and $B > 0$.

The residual error Δ_{stat} depends on the parameters of the RSC/RSM conditions, the network connectivity ρ , and the statistical error $\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|^2$. The next corollary shows that these parameters can be chosen such that $\Delta_{\text{stat}} = o(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|^2)$.

Corollary 19. *Instate Theorem 18; and further suppose $r \leq \|\boldsymbol{\theta}^*\|_1$ and*

$$s(\tau_\mu + \tau_g) = o(1) \quad \text{and} \quad s \cdot \rho c_m^2 \frac{\ell_\Sigma}{\mu_\Sigma} \tau_\ell = o(1). \quad (22)$$

Then,

$$\frac{1}{m} \sum_{i=1}^m \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|^2 \leq \left(\frac{1 - (2\kappa)^{-1} + C_1 s(\tau_\mu + \tau_g)/L_\Sigma}{1 - 2C_1 s\tau_g/L_\Sigma} \right)^t \cdot B + o(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|^2). \quad (23)$$

Corollary 19 guarantees linear convergence up to a tolerance that is smaller than statistical precision. In Sec. 4.2 we show that, when considering the statistical model underlying (1), conditions in (22) are satisfied for near optimal sample complexity and sufficiently small ρ .

• **Iteration complexity.** The contraction factor determining the linear decay of the optimization error depends, as expected, on the restricted condition number κ and the RSC/RSM parameters, the latter due to the lack of strong convexity and smoothness in a global sense. We anticipate that a closer look at the proof of Theorem 18 shows that this degradation on the rate due to the RSC/RSM is limited, resulting in a worst-case rate of $1 - (4\kappa)^{-1}$. This matches the convergence rate of the centralized PGD up to a constant factor and improves on existing analyses of distributed algorithms whose convergence to a solution of (2) is certified only at *sublinear* rate, due to the lack of strong convexity in the global sense (see Sec. 1.2).

When special settings are considered, the convergence rate reduces to well-known expressions. In particular, when strong convexity and smoothness hold globally on $\mathbb{R}^d$, i.e., $\tau_\mu = \tau_g = \tau_\ell = 0$, we have $\Delta_{\text{stat}} = 0$, implying that NetLASSO converges to an ε -solution of (1) in $O(\kappa \log(1/\varepsilon))$ iterations. This recovers the result in [43]. When the network is fully connected, the first term in Δ_{stat} vanishes (as $\rho = 0$), implying that NetLASSO converges linearly to a neighborhood of $\hat{\theta}$ of size $O((\tau_\nu + \tau_g)\nu^2/L_\Sigma)$. This rate matches that of the centralized PGD under RSC/RSM (up to constant factors) established in [1].

• **Interplay between optimization and communication.** Convergence of NetLASSO is established, in particular, under condition (20). This reveals an interesting interplay between communication cost and hardness of the optimization problem. The quantity ℓ_Σ/μ_Σ can be viewed as the restricted condition number of the local losses $\mathcal{L}_i$ (cf. Assumption 4); larger ℓ_Σ/μ_Σ and $s\tau_\ell/\mu_\Sigma$ correspond to more ill conditioned $\mathcal{L}_i$. Therefore, the more ill conditioned $\mathcal{L}_i$, the more connected the network must be to achieve convergence the rate of centralized PGD. When the network topology is given, the associated ρ might not satisfy (20). If so, one can still enforce the desired threshold on ρ by employing multiple rounds of communications. This will be discussed in detail in the next section.

4.2 Guarantees for sparse vector regression

We present some consequences of Theorem 18 and Corollary 19 for the statistical model (1). We assume next that $\{\mathbf{X}_i\}_{i=1}^m$ are i.i.d. random matrices drawn from the Σ -Gaussian ensemble, and the elements of $\{\mathbf{n}_i\}_{i=1}^m$ are i.i.d. σ^2 -sub-Gaussian random variables and independent of the $\mathbf{X}_i$'s. In contrast with the deterministic arguments underlying Theorem 18 and Corollary 19, convergence results in this section are of probabilistic nature, and build upon Proposition 3 and Proposition 5 to establish that the RSC/RSM properties hold with high-probability.

Theorem 22 (informal). *Instate the above statistical model, and further assume that*

$$\|\hat{\theta}\|_1 = r \leq \|\theta^*\|_1, \quad s \log d/N = O(1), \quad \text{and} \quad \rho \leq \text{poly}(m, \kappa)^{-1}. \quad (24)$$

Then, the iterates $\{(\theta_i^t)_{i=1}^m\}$ generated by NetLASSO satisfy

$$\frac{1}{m} \sum_{i=1}^m \|\theta_i^t - \hat{\theta}\|^2 \leq \left(\frac{1 - (2\kappa)^{-1} + C' s \log d/N}{1 - C' s \log d/N} \right)^t \cdot B + O\left(\frac{s \log d}{N} \cdot \|\hat{\theta} - \theta^*\|^2 \right), \quad (25)$$

with high probability, where $C' > 0$ is some universal constant.

Theorem 22 reveals several interesting properties of NetLASSO, as discussed next.

• **Scalability with respect to the problem dimension.** For fixed network whose connectivity ρ satisfies (24), the dependency of the convergence rate on the ambient dimension d , the total sample size N , and sparsity level s is only through the ratio $s \log d/N$. This implies that the convergence rate in (25) is invariant under the high-dimensional scaling $s, N, d \rightarrow \infty$ and $s \log d/N = O(1)$ —resulting in the global sample scaling $N = O(s \log d)$.




Figure 3: Some commonly used graphs $\mathcal{G}$ for the communication network. Dashed line in the Erdős-Rényi graph stands for the existence of an edge with some probability.

• **Near optimal sample complexity.** When $s \log d/N = o(1)$, the residual error in (25) is of smaller order than the statistical precision $\|\hat{\theta} - \theta^*\|^2$. Therefore, NetLASSO takes $O(\kappa \log(1/\varepsilon))$ iterations to reach an ε -neighborhood of a statistically optimal solution; this rate is of the same order of the centralized PGD under RSC/RSM [1]. Note also that the condition $s \log d/N = o(1)$ (almost) matches the optimal sample complexity $N = \Omega(s \log(d/s))$ for model (1), which is necessary for any centralized method to consistently estimate the d -dimensional, s -sparse θ^* from N samples.

Since $N = n \cdot m$, the condition $s \log d/N = o(1)$ reveals an interesting interplay between statistical error, convergence rate, and network connectivity/communication cost, which is peculiar of the distributed setting: when agents do not have enough local samples n for statistical consistency but the overall sample size N across the network suffices—a situation that happens, e.g., when m is large (large-scale network), NetLASSO still achieves centralized statistical accuracy at linear rate provided that the network is “sufficiently” connected, i.e., $\rho \leq \text{poly}(m, \kappa)^{-1}$. Therefore, the insufficiency of the local sample size is compensated by higher communication costs. The impact of network parameters and topology on the convergence of NetLASSO is further elaborated next.

	path	2-d grid	star	complete	push-pull (star)	Erdős-Rényi	Erdős-Rényi
$(1 - \rho)^{-1}$	$O(m^2)$	$O(m \log m)$	$O(m^2)$	1	1	$O(1)$ [$p = \log m/m$]	$O(1)$ [$p = O(1)$]
max degree	2	4	m	m	m	$O(\log m)$	$O(m)$
channel use/comm.	$O(m)$	$O(m)$	$O(m)$	$O(m^2)$	$O(m)$	$O(m \log m)$	$O(m^2)$
max channel use	$\tilde{O}(m^2)$	$\tilde{O}(m)$	$\tilde{O}(m^3)$	$\tilde{O}(m)$	$\tilde{O}(m)$	$\tilde{O}(1)$	$\tilde{O}(m)$
total channel use	$\tilde{O}(m^3)$	$\tilde{O}(m^2)$	$\tilde{O}(m^3)$	$\tilde{O}(m^2)$	$\tilde{O}(m)$	$\tilde{O}(m)$	$\tilde{O}(m^2)$

Table 1: Scalability of the communication complexity of NetLASSO with the network size m , under different network topology. Push-pull(star): NetLASSO over star-networks as in Algorithm 2 (it reduces to the PGD). **Max degree**: the maximum degree of the nodes; **channel use/comm.**: total channel use per communication round; **max channel use**: channel use of the busiest node to reach ε optimization precision; **total channel use**: total channel uses to reach ε optimization precision. $\tilde{O}$ hides the logarithmic factors in communication complexity.

• **Logarithmic scalability of the communication cost with the network size.** When the graph $\mathcal{G}$ is not part of the design but given a-priori along with $\overline{\mathbf{W}}$ (satisfying Assumption 7), the condition $\rho \leq \text{poly}(m, \kappa)^{-1}$ can be otherwise satisfied by running multiple rounds of consensus steps. More precisely, since $\bar{\rho} = \|\overline{\mathbf{W}} - \mathbf{J}\| < 1$ (Assumption 7), one can construct $\mathbf{W} = \overline{\mathbf{W}}^K$ such that $\rho = \bar{\rho}^K \leq \text{poly}(m, \kappa)^{-1}$. This corresponds to running K consensus steps over the graph $\mathcal{G}$ using each time the matrix $\overline{\mathbf{W}}$ in Step [S.1]. Note that for fixed m (graph), such K is always finite. In fact, one can see that taking $K = O(\log(m\kappa) \cdot (1 - \rho)^{-1})$ fulfills

the requirement. If Chebyshev acceleration is used to employ multiple rounds of communications, coupled with a symmetric matrix $\overline{\mathbf{W}}$ (satisfying (Assumption 7)), one can show that the number of communication steps K reduces to $O(\log(m\kappa) \cdot (1-\rho)^{-1/2})$. Combining this fact with the iteration complexity (25), one can conclude that, in the setting of Theorem 22, NetLASSO based on $\mathbf{W} = \overline{\mathbf{W}}^K$ drives $\frac{1}{m} \sum_{i=1}^m \|\theta_i^t - \hat{\theta}\|^2$ within an ε -neighborhood of a statistical optimal solution in the following number of communication steps

$$O\left(\frac{\kappa}{1-\rho} \log(m\kappa) \log(1/\varepsilon)\right). \quad (26)$$

Notice that this improves on the communication complexity of DGD-like algorithms, such as [19]: contrary to (7), (26) does not depend on the ambient dimension d ; hence, NetLASSO does not suffer from the speed-accuracy dilemma as DGD algorithms. This is the first time that a distributed algorithm exhibits such a desirable property in high-dimensions.

• **Impact of network topology.** The network topology affects the convergence rate of NetLASSO through parameters m and ρ . A larger m or ρ increases communication complexity, as indicated by (26). In addition, for networks of the same size m , the efficiency of information propagation depends on the graph's topology, which in (26) is captured by the dependency of ρ on m [27]. Table 1 provides, for some commonly used network topologies (see Fig. 3 for some examples), the dependency of ρ on m , when the lazy Metropolis rule is adopted; for other rules, see [6]. The “push-pull” (star) in Table 1 refers specifically to the implementation of the NetLASSO as in Algorithm 2 over a star network (which coincides with the PGD). While $(1-\rho)^{-1}$ affects the total number of communications, it does not capture the overall cost of communications, which also depends on how dense the graph is. Hence, counting as one channel use per communication, with each edge shared between two nodes, Table 1 reports also the following quantities capturing different communications costs: total channel use per communication round, channel use of the busiest node and total channel uses to reach an ε neighborhood of a statistically optimal solution.

The following comments are in order. From the first row of the table and (26), we infer that NetLASSO running over a graph with stronger connectivity (smaller ρ) requires less communication rounds to converge. On the other hand, as tradeoff, better connectivity generally results from denser graphs, and thus yields higher communications costs (heavier channel uses), as reported in the table. We deduce that NetLASSO running on the Erdős-Rényi graph with edge connecting probability $p = \log m/m$ achieves the lowest communication cost at the busiest node, and ties with the implementation on the star (“push-pull”) in the amount of total channel use¹. For the path graph and 2-d grid graph, even though the channel use per communication round is low, NetLASSO takes more iterations to converge, and overall a higher total communication cost. Comparing the two instances of the Erdős-Rényi graphs (the last two columns in the table), one may prefer a more sparse graph at the price of more communication rounds to a dense graph and less communication rounds. Finally, even though the Erdős-Rényi is relatively efficient for NetLASSO, in practice one may not be able to construct such a topology, e.g., due to geographic constraints. In such scenarios, a path/grid graph is still a valuable choice.

5 Convergence Analysis

5.1 Preliminaries

We begin rewriting the iterates (16)-(18) in matrix/vector form and introducing some notation, which is convenient for our developments. Given $(\theta_i^t)_{i=1}^m$, $(\theta_i^{t+\frac{1}{2}})_{i=1}^m$, $(\mathbf{g}_i^t)_{i=1}^m$, $(\Delta\theta_i^t)_{i=1}^m$ and $(\nabla\mathcal{L}(\theta_i^t))_{i=1}^m$, we introduce the following associated matrices:

$$\begin{aligned} \Theta^t &\triangleq [\theta_1^t, \dots, \theta_m^t]^\top, & \Theta^{t+\frac{1}{2}} &\triangleq [\theta_1^{t+\frac{1}{2}}, \dots, \theta_m^{t+\frac{1}{2}}]^\top, & \Delta\Theta^t &\triangleq [\Delta\theta_1^t, \dots, \Delta\theta_m^t]^\top = \Theta^{t+\frac{1}{2}} - \Theta^t, \\ \mathbf{G}^t &\triangleq [\mathbf{g}_1^t, \dots, \mathbf{g}_m^t]^\top, & \text{and} & & \nabla\mathcal{L}(\Theta^t) &\triangleq [\nabla\mathcal{L}_1(\theta_1^t), \dots, \nabla\mathcal{L}_m(\theta_m^t)]^\top. \end{aligned} \quad (27)$$

¹We note that the performance on “push-pull” (star) are better than those on the Erdős-Rényi when the log factor in $\tilde{O}$ is explicitly accounted.

Using (27), we can rewrite (16)-(18) in a compact form as

$$\Theta^t = \mathbf{W} (\Theta^{t-1} + \Delta \Theta^{t-1}), \quad (28a)$$

$$\mathbf{G}^t = \mathbf{W} (\mathbf{G}^{t-1} + \nabla \mathcal{L}(\Theta^t) - \nabla \mathcal{L}(\Theta^{t-1})). \quad (28b)$$

Introducing the average quantities

$$\bar{\theta}^t \triangleq \frac{1}{m} \sum_{i=1}^m \theta_i^t \quad \text{and} \quad \bar{\mathbf{g}}^t \triangleq \frac{1}{m} \sum_{i=1}^m \mathbf{g}_i^t, \quad (29)$$

we define the consensus errors on the θ , $\mathbf{g}$ -vectors as

$$\theta_{i,\perp}^t \triangleq \theta_i^t - \bar{\theta}^t \quad \text{and} \quad \mathbf{g}_{i,\perp}^t \triangleq \mathbf{g}_i^t - \bar{\mathbf{g}}^t, \quad i = 1, \dots, m. \quad (30)$$

Note that in our setting, $\theta_{i,\perp}^0 = 0$, for all $i = 1, \dots, m$. The matrix counterparts of (30) are

$$\Theta_{\perp}^t \triangleq [\theta_{1,\perp}^t, \dots, \theta_{m,\perp}^t]^\top = (\mathbf{I} - \mathbf{J})\Theta^t \quad \text{and} \quad \mathbf{G}_{\perp}^t \triangleq [\mathbf{g}_{1,\perp}^t, \dots, \mathbf{g}_{m,\perp}^t]^\top = (\mathbf{I} - \mathbf{J})\mathbf{G}^t, \quad (31)$$

where $\mathbf{J} \triangleq \mathbf{1}\mathbf{1}^\top/m$.

Using (28), the dynamics of (31) read

$$\Theta_{\perp}^t = (\mathbf{W} - \mathbf{J})(\Theta_{\perp}^{t-1} + \Delta \Theta^{t-1}) \quad (32a)$$

$$\mathbf{G}_{\perp}^t = (\mathbf{W} - \mathbf{J})(\mathbf{G}_{\perp}^{t-1} + \nabla \mathcal{L}(\Theta^t) - \nabla \mathcal{L}(\Theta^{t-1})). \quad (32b)$$

where we used $(\mathbf{I} - \mathbf{J})\mathbf{W} = \mathbf{W} - \mathbf{J}$.

5.2 Inexact descent

We begin with the analysis of the optimization step (18). Since each $\mathbf{g}_i$ aims to estimate the local gradient $\nabla \mathcal{L}(\theta_i)$, step (18) can be regarded as an inexact (“noisy”) version of the proximal gradient update:

$$\theta_i^{t+\frac{1}{2}} = \underset{\|\theta\|_1 \leq r}{\operatorname{argmin}} \left\{ \nabla \mathcal{L}(\theta_i^t)^\top (\theta_i - \theta_i^t) + \frac{\gamma}{2} \|\theta_i - \theta_i^t\|^2 \right\}. \quad (33)$$

Therefore, we base our proof on the analysis of (33) while taking into account the gradient estimation error $\nabla \mathcal{L}(\theta_i) - \mathbf{g}_i$, due to the use in (33) of the gradient tracking vector $\mathbf{g}_i$ rather than $\nabla \mathcal{L}(\theta_i)$.

Proposition 9. *Consider the optimization problem (2) over the network $\mathcal{G}$, under Assumptions 6 and 2; and $\hat{\theta}$ such that $\|\hat{\theta}\| = r$. Let $\{\theta_i^t\}$ be the sequence generated by Algorithm 1 under Assumption 7. Then, the following holds:*

$$\begin{aligned} \left(1 - \frac{C_1 s \tau_g}{\gamma}\right) \|\Theta^{t+\frac{1}{2}} - \hat{\Theta}\|^2 &\leq \left(1 - \frac{\mu_\Sigma}{\gamma} + \frac{C_1 s (\tau_\mu + \tau_g)}{\gamma}\right) \|\Theta^t - \hat{\Theta}\|^2 \\ &\quad - \left(1 - \frac{L_\Sigma}{\gamma}\right) \|\Delta \Theta^t\|^2 + \underbrace{\frac{2}{\gamma} \sum_{i=1}^m (\nabla \mathcal{L}(\theta_i^t) - \mathbf{g}_i^t)^\top (\theta_i^{t+\frac{1}{2}} - \hat{\theta})}_{\triangleq \delta^t} + \frac{C_1 (\tau_\mu + \tau_g)}{\gamma} m \nu^2, \end{aligned} \quad (34)$$

for some absolute constant $C_1 > 0$.

Proof. See Appendix A (supplementary material). $\square$

Note that when the network is fully connected we have $\nabla \mathcal{L}(\theta_i^t) - \mathbf{g}_i^t = \mathbf{0}$ and $\theta_i^t = \theta_j^t$ for all $i, j = 1, \dots, m$, and $t = 1, \dots$. Letting $\gamma = L_\Sigma$, Eq. (34) recovers the result provided in [1, Thm. 1] for the centralized PGD algorithm. Therefore, the next step is to analyze the dynamics of the gradient tracking error δ^t in (34), which is the subject of the next section.

5.3 Bounding the tracking error

To understand the evolution of the tracking error

$$\delta^t = \sum_{i=1}^m (\nabla \mathcal{L}(\boldsymbol{\theta}_i^t) - \mathbf{g}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}),$$

we need to resort to the dynamics of $\mathbf{g}_i^t$. Recalling the definition of $\bar{\mathbf{g}}^t$ [cf. (29)] and the tracking dynamics (17), we have

$$\bar{\mathbf{g}}^t = \bar{\mathbf{g}}^{t-1} + \frac{1}{m} \sum_{i=1}^m (\nabla \mathcal{L}_i(\boldsymbol{\theta}_i^t) - \nabla \mathcal{L}_i(\boldsymbol{\theta}_i^{t-1})).$$

Hence,

$$\bar{\mathbf{g}}^t = \frac{1}{m} \sum_{i=1}^m \nabla \mathcal{L}_i(\boldsymbol{\theta}_i^t), \quad (35)$$

where we used the initialization $\bar{\mathbf{g}}^0 = \frac{1}{m} \sum_{i=1}^m (\nabla \mathcal{L}_i(\boldsymbol{\theta}_i^0))$.

This average preserving property of the sum-gradient naturally suggests the following decomposition of the tracking error δ^t :

$$\begin{aligned} \delta^t &= \sum_{i=1}^m \left(\nabla \mathcal{L}(\boldsymbol{\theta}_i^t) - \frac{1}{m} \sum_{j=1}^m \nabla \mathcal{L}_j(\boldsymbol{\theta}_j^t) + \bar{\mathbf{g}}^t - \mathbf{g}_i^t \right)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}) \\ &= \sum_{i=1}^m \left(\frac{1}{m} \sum_{j=1}^m \nabla \mathcal{L}_j(\boldsymbol{\theta}_j^t) - \frac{1}{m} \sum_{j=1}^m \nabla \mathcal{L}_j(\boldsymbol{\theta}_j^t) \right)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}) + \sum_{i=1}^m (\bar{\mathbf{g}}^t - \mathbf{g}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}). \end{aligned} \quad (36)$$

Clearly, δ^t vanishes if $\boldsymbol{\theta}_i^{t+\frac{1}{2}}$ converges to $\widehat{\boldsymbol{\theta}}$, or $\{\boldsymbol{\theta}_i^t\}_i$ and $\{\mathbf{g}_i^t\}_i$ are asymptotically consensual. Furthermore, we may intuitively expect a more connected network supports faster information propagation, which potentially yield a faster vanishing rate of the tracking error.

We are now ready to established the bound of δ^t .

Proposition 10. *Under the setting of Prop. 9 and the local restricted smoothness Assumption 4, the gradient tracking error δ^t ($t = 1, \dots$) can be bounded as*

$$\begin{aligned} \delta^t &\leq \frac{C_2}{2} \left\{ \frac{\rho \cdot c_m \epsilon}{1 - \rho} (\ell_\Sigma + s\tau_\ell) \|\boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 \right. \\ &\quad + \frac{c_m \ell_\Sigma}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta \boldsymbol{\Theta}^s\|_F^2 + \frac{c_m \ell_\Sigma}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}_\perp^s\|_F^2 \\ &\quad + c_m s \tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^{s+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 + c_m s \tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^{s-\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 \\ &\quad \left. + m c_m \rho^t (A_3 \cdot \epsilon^{-1} + s c_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu) + \frac{\rho \cdot m c_m}{1 - \rho} \tau_\ell \nu^2 (\epsilon + \epsilon^{-1}) \right\} \end{aligned} \quad (37)$$

for some $C_2 > 0$; where $\epsilon > 0$, $\boldsymbol{\Theta}^{-\frac{1}{2}} \triangleq \boldsymbol{\Theta}^0$ and

$$\begin{aligned} A_3 &\triangleq \ell_\Sigma \|\boldsymbol{\theta}^*\|^2 + \tau_\ell \|\boldsymbol{\theta}^*\|_1^2, \\ c_g &\triangleq \|\nabla \mathcal{L}_j(\boldsymbol{\theta}^*)\|_\infty + \|\nabla \mathcal{L}(\boldsymbol{\theta}^*)\|_\infty. \end{aligned} \quad (38)$$

Proof. See Appendix B (supplementary material). $\square$

Proposition 10 reveals that the gradient tracking error δ^t can be bounded by the discounted cumulative sum of historical optimization errors $\{\|\Theta^{s-1/2} - \hat{\Theta}\|\}_{s=0}^{t+1}$, step-length history $\{\|\Delta\Theta^s\|\}_{s=0}^{t-1}$, and consensus errors $\{\|\Theta_\perp^s\|\}_{s=0}^{t-1}$. Furthermore, it is not difficult to check that δ^t decreases if ρ , m , or the RSS tolerance parameter τ_ℓ decrease, which is a consequence of a smaller, more connected network, and larger local sample size n , respectively.

We can now combine the statements obtained in Propositions 9 and 10, and write

$$\begin{aligned}
& \left(1 - \frac{1}{\gamma} \left(C_1 s \tau_g + \frac{C_2 \rho \cdot c_m \epsilon}{1 - \rho} (\ell_\Sigma + s \tau_\ell)\right)\right) \|\Theta^{t+\frac{1}{2}} - \hat{\Theta}\|^2 \\
& \leq \left(1 - \frac{\mu_\Sigma}{\gamma} + \frac{C_1 s (\tau_\mu + \tau_g)}{\gamma}\right) \|\Theta^t - \hat{\Theta}\|^2 - \left(1 - \frac{L_\Sigma}{\gamma}\right) \|\Delta\Theta^t\|^2 + \frac{C_1 (\tau_\mu + \tau_g)}{\gamma} m \nu^2 \\
& \quad + \frac{C_2}{\gamma} \frac{c_m \ell_\Sigma}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta\Theta^s\|_F^2 + \frac{C_2}{\gamma} \frac{c_m \ell_\Sigma}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta_\perp^s\|_F^2 \\
& \quad + \frac{C_2}{\gamma} c_m s \tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+\frac{1}{2}} - \hat{\Theta}\|_F^2 + \frac{C_2}{\gamma} c_m s \tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s-\frac{1}{2}} - \hat{\Theta}\|_F^2 \\
& \quad + \frac{C_2}{\gamma} m c_m \rho^t (A_3 \cdot \epsilon^{-1} + s c_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu) + \frac{C_2}{\gamma} \frac{\rho \cdot m c_m}{1 - \rho} \tau_\ell \nu^2 (\epsilon + \epsilon^{-1}).
\end{aligned} \tag{39}$$

Our last step bounds the consensus error $\|\Theta_\perp^t\|$ using $\{\|\Delta\Theta^s\|\}_{s=0}^{t-1}$, as given next.

Proposition 11. *The consensus error $\Theta_\perp^t$ can be bounded as*

$$\|\Theta_\perp^t\|_F^2 \leq 2\rho^{2t} \underbrace{\|\Theta_\perp^0\|_F^2}_{=0} + \frac{2\rho}{1 - \rho} \cdot \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta\Theta^s\|_F^2. \tag{40}$$

Proof. See Appendix D (supplementary material). $\square$

5.4 Linear convergence up to a finite ball under RSC/RSM

To establish linear convergence of NetLASSO from (39)-(40), we will employ the z -transform for finite length sequences. We begin briefly reviewing the relevant properties of the z -transform (cf. Sec. 5.4.1) and then apply it to the system (39)-(40) (cf. Sec. 5.4.2).

5.4.1 The z -transform and properties

Definition 12. *Let $\{a(t)\}$ be a sequence of nonnegative numbers. The K -length z -transform of $\{a(t)\}$ is defined as*

$$A^K(z) \triangleq \sum_{t=1}^K a(t) z^{-t}, \quad z \in \mathbb{R}_{++}. \tag{41}$$

Some properties of the z -transform instrumental to our developments are summarized next.

Lemma 13. *Let $A^K(z)$ be the length K z -transform of a nonnegative sequence $\{a(t)\}$, and $\rho \in (0, 1)$. Then, the following hold:*

- (i) $\sum_{t=1}^K a(t+1) z^{-t} \geq z A^K(z) - a(1), \quad \forall z \in \mathbb{R}_{++};$
- (ii) $\sum_{t=1}^K \left(\sum_{s=0}^{t-1} \rho^{t-s} a(s)\right) z^{-t} \leq \frac{\rho}{z - \rho} \cdot (A^K(z) + a(0)), \quad \forall z \in (\rho, 1);$

$$(iii) \sum_{t=1}^K \left(\sum_{s=0}^{t-1} \rho^{t-s} a(s+1) \right) z^{-t} \leq z \cdot \frac{\rho}{z-\rho} A^K(z), \quad \forall z \in (\rho, 1).$$

Proof. (i): For the time-shifted sequence $\{a(t)\}_{t=2}^{K+1}$, it holds

$$\begin{aligned} \sum_{t=1}^K a(t+1)z^{-t} &= z \sum_{t=1}^K a(t+1)z^{-t-1} \\ &= z \left(\sum_{t=1}^K a(t)z^{-t} + a(K+1)z^{-K-1} - a(1)z^{-1} \right) \\ &= zA^K(z) - a(1) + a(K+1)z^{-K} \geq zA^K(z) - a(1). \end{aligned}$$

(ii): For the sequence $\{\sum_{s=0}^{t-1} \rho^{t-s} a(s)\}_{t=1}^K$, we have

$$\sum_{t=1}^K \sum_{s=0}^{t-1} z^{s-t} \rho^{t-s} a(s) z^{-s} = \sum_{s=0}^{K-1} a(s) z^{-s} \sum_{t=s+1}^K z^{s-t} \rho^{t-s} \leq \frac{\rho}{z-\rho} \cdot (A^K(z) + a(0)), \quad \forall z \in (\rho, 1).$$

(iii): Similarly, the sequence $\{\sum_{s=0}^{t-1} \rho^{t-s} a(s+1)\}_{t=1}^K$, we have

$$\sum_{t=1}^K \sum_{s=0}^{t-1} \rho^{t-s} a(s+1) z^{-t} = \sum_{s=0}^{K-1} a(s+1) z^{-s-1} \sum_{t=s+1}^K \rho^{t-s} z^{-t+s+1} \leq z \cdot \frac{\rho}{z-\rho} A^K(z).$$

□

Lemma 14. Suppose a nonnegative sequence $\{a(t)\}$ satisfies

$$\sum_{t=1}^K a(t) z^{-t} \leq B + c \cdot \sum_{t=1}^K z^{-t}, \quad \text{for all } K \geq 1 \quad \text{and } z \in (\bar{z}, 1), \quad (42)$$

and some $B, c > 0$. Then

$$a(t) \leq B \cdot \bar{z}^t + \frac{c}{1-\bar{z}}. \quad (43)$$

When $B < +\infty$, $\{a(t)\}$ converges linearly at rate $\bar{z}$ up to a constant.

Proof. Since $a(t) \geq 0$, (42) implies that for all $K \geq 1$ and $z \in (\bar{z}, 1)$,

$$a(K) z^{-K} \leq B + c \cdot \sum_{t=1}^K z^{-t}. \quad (44)$$

Therefore, for all $z \in (\bar{z}, 1)$

$$a(K) \leq B \cdot z^K + c \cdot \sum_{t=1}^K z^{K-t} \leq B \cdot z^K + \frac{c}{1-z}. \quad (45)$$

Letting $z \rightarrow \bar{z}^+$ concludes the proof. □

5.4.2 The z-transform system

Introduce the following quantities:

$$D_{\Theta}^K(z) = \sum_{t=1}^K z^{-t} \|\Theta^{t-\frac{1}{2}} - \hat{\Theta}\|^2, \quad \Delta_{\Theta}^K(z) = \sum_{t=1}^K z^{-t} \|\Delta\Theta^t\|^2, \quad \Theta_{\perp}^K(z) = \sum_{t=1}^K z^{-t} \|\Theta_{\perp}^t\|^2. \quad (46)$$

To apply (46) to (39) we first bound the term $\|\Theta^t - \hat{\Theta}\|^2$ on the RHS there as $\|\Theta^t - \hat{\Theta}\|^2 \leq \|\Theta^{t-\frac{1}{2}} - \hat{\Theta}\|^2$, yielding

$$\begin{aligned} & \left(1 - \frac{1}{\gamma} \left(C_1 s \tau_g + \frac{C_2 \rho \cdot c_m \epsilon}{1 - \rho} (\ell_{\Sigma} + s \tau_{\ell})\right)\right) \|\Theta^{t+\frac{1}{2}} - \hat{\Theta}\|^2 \\ & \leq \left(1 - \frac{\mu_{\Sigma}}{\gamma} + \frac{C_1 s (\tau_{\mu} + \tau_g)}{\gamma}\right) \|\Theta^{t-\frac{1}{2}} - \hat{\Theta}\|^2 - \left(1 - \frac{L_{\Sigma}}{\gamma}\right) \|\Delta\Theta^t\|^2 + \frac{C_1 (\tau_{\mu} + \tau_g)}{\gamma} m \nu^2 \\ & \quad + \frac{C_2}{\gamma} \frac{c_m \ell_{\Sigma}}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta\Theta^s\|_F^2 + \frac{C_2}{\gamma} \frac{c_m \ell_{\Sigma}}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta_{\perp}^s\|_F^2 \\ & \quad + \frac{C_2}{\gamma} c_m s \tau_{\ell} \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+\frac{1}{2}} - \hat{\Theta}\|_F^2 + \frac{C_2}{\gamma} c_m s \tau_{\ell} \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s-\frac{1}{2}} - \hat{\Theta}\|_F^2 \\ & \quad + \frac{C_2}{\gamma} m c_m \rho^t (A_3 \cdot \epsilon^{-1} + s c_g^2 \ell_{\Sigma}^{-1} \epsilon^{-1} + c_g \nu) + \frac{C_2}{\gamma} \frac{\rho \cdot m c_m}{1 - \rho} \tau_{\ell} \nu^2 (\epsilon + \epsilon^{-1}) \end{aligned} \quad (47)$$

as long as

$$\gamma \geq \mu_{\Sigma} - C_1 s (\tau_{\mu} + \tau_g). \quad (48)$$

Multiplying by z^{-t} , $z > \rho$, both sides of (47) and (40) while summing up from $t = 1$ to $t = K$ and applying Lemma 13 (recall that $\Delta\Theta^0 = \mathbf{0}$), we obtain

$$\begin{aligned} & \left(1 - \frac{1}{\gamma} \left\{C_1 s \tau_g + \frac{C_2 \rho \cdot c_m \epsilon}{1 - \rho} (\ell_{\Sigma} + s \tau_{\ell})\right\}\right) \left(z D_{\Theta}^K(z) - \|\Theta^{1/2} - \hat{\Theta}\|^2\right) \\ & \leq \left(1 - \frac{\mu_{\Sigma}}{\gamma} + \frac{C_1 s (\tau_{\mu} + \tau_g)}{\gamma}\right) D_{\Theta}^K(z) - \left(1 - \frac{L_{\Sigma}}{\gamma}\right) \Delta_{\Theta}^K(z) \\ & \quad + \frac{C_2}{\gamma} \frac{c_m \ell_{\Sigma}}{\epsilon} \frac{\rho}{z - \rho} \Delta_{\Theta}^K(z) + \frac{C_2}{\gamma} \frac{c_m \ell_{\Sigma}}{\epsilon} \frac{\rho}{z - \rho} \Theta_{\perp}^K(z) \\ & \quad + \frac{C_2}{\gamma} c_m s \tau_{\ell} \epsilon^{-1} \frac{z \rho}{z - \rho} D_{\Theta}^K(z) + \frac{C_2}{\gamma} c_m s \tau_{\ell} \epsilon^{-1} \frac{\rho}{z - \rho} \left(D_{\Theta}^K(z) + \|\Theta^0 - \hat{\Theta}\|^2\right) \\ & \quad + \frac{C_2}{\gamma} m c_m (A_3 \cdot \epsilon^{-1} + s c_g^2 \ell_{\Sigma}^{-1} \epsilon^{-1} + c_g \nu) \frac{\rho}{z - \rho} \\ & \quad + \left\{ \frac{C_2}{\gamma} \frac{\rho \cdot m c_m}{1 - \rho} \tau_{\ell} \nu^2 (\epsilon + \epsilon^{-1}) \frac{C_1 (\tau_{\mu} + \tau_g)}{\gamma} m \nu^2 \right\} \sum_{t=1}^K z^{-t} \end{aligned} \quad (49)$$

and

$$\Theta_{\perp}^K(z) \leq \frac{2\rho}{1 - \rho} \cdot \frac{\rho}{z - \rho} \Delta_{\Theta}^K(z), \quad (50)$$

respectively, under the condition

$$\gamma \geq C_1 s \tau_g + \frac{C_2 \rho \cdot c_m \epsilon}{1 - \rho} (\ell_{\Sigma} + s \tau_{\ell}). \quad (51)$$

Substituting (50) in (49) yields

$$\begin{aligned}
& \left(1 - \frac{1}{\gamma} \left\{ C_1 s \tau_g + \frac{C_2 \rho \cdot c_m \epsilon}{1 - \rho} (\ell_\Sigma + s \tau_\ell) + \frac{C_2 \rho \cdot c_m \epsilon^{-1} s \tau_\ell}{z - \rho} \right\} \right) z D_\Theta^K(z) \\
& - \left(1 - \frac{\mu_\Sigma}{\gamma} + \frac{C_1 s (\tau_\mu + \tau_g)}{\gamma} + \frac{C_2}{\gamma} c_m s \tau_\ell \epsilon^{-1} \frac{\rho}{z - \rho} \right) D_\Theta^K(z) \\
& \leq - \left(1 - \frac{L_\Sigma}{\gamma} - \frac{C_2}{\gamma} \frac{c_m \ell_\Sigma}{\epsilon} \frac{\rho}{z - \rho} - \frac{C_2}{\gamma} \frac{c_m \ell_\Sigma}{\epsilon} \frac{2\rho}{1 - \rho} \cdot \left(\frac{\rho}{z - \rho} \right)^2 \right) \Delta_\Theta^K(z) \\
& + \left\{ \frac{C_2}{\gamma} \frac{\rho \cdot m c_m}{1 - \rho} \tau_\ell \nu^2 (\epsilon + \epsilon^{-1}) + \frac{C_1 (\tau_\mu + \tau_g)}{\gamma} m \nu^2 \right\} \sum_{t=1}^K z^{-t} \\
& + \frac{C_2}{\gamma} c_m s \tau_\ell \epsilon^{-1} \frac{\rho}{z - \rho} \|\Theta^0 - \widehat{\Theta}\|^2 + \frac{C_2}{\gamma} m c_m (A_3 \cdot \epsilon^{-1} + s c_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu) \frac{\rho}{z - \rho} \\
& + \left(1 - \frac{1}{\gamma} \left\{ C_1 s \tau_g + \frac{C_2 \rho \cdot c_m \epsilon}{1 - \rho} (\ell_\Sigma + s \tau_\ell) \right\} \right) \|\Theta^{1/2} - \widehat{\Theta}\|^2.
\end{aligned} \tag{52}$$

Accordingly, we define the following quantities

$$\begin{aligned}
B(z) & \triangleq \frac{C_2}{\gamma} c_m s \tau_\ell \epsilon^{-1} \frac{\rho}{z - \rho} \|\Theta^0 - \widehat{\Theta}\|^2 + \frac{C_2}{\gamma} m c_m (A_3 \cdot \epsilon^{-1} + s c_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu) \frac{\rho}{z - \rho} \\
& + \left(1 - \frac{1}{\gamma} \left\{ C_1 s \tau_g + \frac{C_2 \rho \cdot c_m \epsilon}{1 - \rho} (\ell_\Sigma + s \tau_\ell) \right\} \right) \|\Theta^{1/2} - \widehat{\Theta}\|^2
\end{aligned} \tag{53}$$

and

$$\overline{\Delta}_{\text{stat}} = \frac{C_2}{\gamma} \frac{\rho \cdot c_m}{1 - \rho} \tau_\ell \nu^2 (\epsilon + \epsilon^{-1}) + \frac{C_1 (\tau_\mu + \tau_g)}{\gamma} \nu^2. \tag{54}$$

The following conditions are sufficient to invoke Lemma 14 on the sequence $\{(1/m) \|\Theta^{t+\frac{1}{2}} - \widehat{\Theta}\|_F^2\}$: i) (48) and (51) hold; and ii) there exists some $z \in (\rho, 1)$ such that

$$\begin{aligned}
& \left[1 - \frac{1}{\gamma} \left(C_1 s \tau_g + \frac{C_2 \rho \cdot c_m \epsilon}{1 - \rho} (\ell_\Sigma + s \tau_\ell) + \frac{C_2 \rho \cdot c_m \epsilon^{-1} s \tau_\ell}{z - \rho} \right) \right] z \\
& > 1 - \frac{\mu_\Sigma}{\gamma} + \frac{C_1 s (\tau_\mu + \tau_g)}{\gamma} + \frac{C_2}{\gamma} c_m s \tau_\ell \epsilon^{-1} \frac{\rho}{z - \rho};
\end{aligned} \tag{55a}$$

$$1 - \frac{L_\Sigma}{\gamma} - \frac{C_2}{\gamma} \frac{c_m \ell_\Sigma}{\epsilon} \frac{\rho}{z - \rho} - \frac{C_2}{\gamma} \frac{c_m \ell_\Sigma}{\epsilon} \frac{2\rho}{1 - \rho} \cdot \left(\frac{\rho}{z - \rho} \right)^2 > 0. \tag{55b}$$

5.4.3 Path to linear convergence (Theorem 18 & Corollary 19)

We derive next conditions for the existence of a $z \in (\rho, 1)$ satisfying (55). We organize the proof in the following steps:

- **Step 1:** We provide choices of ϵ and γ for the existence of some $z > \rho$ that satisfies (55a)-(55b);
- **Step 2:** We use the findings in Step 1 to bound the expression of $B(z)$ [cf. (53)] and $\overline{\Delta}_{\text{stat}}$ [cf. (54)];
- **Step 3:** By combining Step 1 and Step 2, we establish conditions for the linear convergence of $\{(1/m) \|\Theta^{t+\frac{1}{2}} - \widehat{\Theta}\|_F^2\}_t$ up to a residual error smaller than the statistical precision $\|\widehat{\theta} - \theta^*\|^2$.

• **Step 1: Expression of the rate from (55a)-(55b)**

Lemma 15 and 16 below provide conditions and the range of $z > \rho$ for (55b) and (55a) to hold, respectively. Throughout this section we tacitly assume that γ satisfies (48) and (51) (strictly when needed). Furthermore, we set

$$\epsilon = \frac{\mu_\Sigma}{\ell_\Sigma} \frac{1-\rho}{2C_2c_m}, \quad (56)$$

which we found a convenient choice for our purposes.

Lemma 15. *Given $\gamma > L_\Sigma$, condition (55b) is satisfied for all $z > z_\rho$, with*

$$z_\rho \triangleq \rho + \frac{4\rho^2}{1-\rho} \left(\sqrt{1 + (\gamma - L_\Sigma) \left(\frac{\mu_\Sigma}{\ell_\Sigma^2} \frac{2\rho}{C_2^2 c_m^2} \right) - 1} \right)^{-1}. \quad (57)$$

Proof. See Appendix E (supplementary material). $\square$

We focus now on condition (55a). Denote for simplicity

$$\begin{aligned} A &\triangleq \frac{1 - \frac{\mu_\Sigma}{\gamma} + \frac{C_1 s(\tau_\mu + \tau_g)}{\gamma}}{\frac{C_2}{\gamma} c_m s \tau_\ell \epsilon^{-1}} = \frac{\gamma - \mu_\Sigma + C_1 s(\tau_\mu + \tau_g)}{\frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_2^2 c_m^2}{1-\rho} s \tau_\ell}, \\ D &\triangleq \frac{1 - \frac{1}{\gamma} \left\{ C_1 s \tau_g + \frac{C_2 \rho \cdot c_m \epsilon}{1-\rho} (\ell_\Sigma + s \tau_\ell) \right\}}{\frac{C_2}{\gamma} c_m s \tau_\ell \epsilon^{-1}} = \frac{\gamma - \left\{ C_1 s \tau_g + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma} (\ell_\Sigma + s \tau_\ell) \right\}}{\frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_2^2 c_m^2}{1-\rho} s \tau_\ell}, \end{aligned} \quad (58)$$

and

$$\sigma \triangleq \frac{A}{D} = \frac{\gamma - \mu_\Sigma + C_1 s(\tau_\mu + \tau_g)}{\gamma - \left\{ C_1 s \tau_g + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma} (\ell_\Sigma + s \tau_\ell) \right\}}. \quad (59)$$

Note that, under (48) and (51) strictly satisfied, $A, B > 0$. Then (55a) can be rewritten succinctly as

$$z \geq \frac{A + G_{\text{net}}(z)}{D - G_{\text{net}}(z)}, \quad G_{\text{net}}(z) \triangleq \frac{\rho}{z - \rho}. \quad (60)$$

Lemma 16. *Condition (55a) is satisfied for all $z > z_\sigma$, with*

$$z_\sigma \triangleq \frac{A}{D} + \rho \cdot \left(1 + \frac{1}{D} \right), \quad (61)$$

and A and B defined in (58).

Proof. See Appendix F (supplementary material). $\square$

Combining Lemma 15 and 16 we conclude that, for any given $\gamma > L_\Sigma$, (55a)-(55b) are satisfied by

$$z > \bar{z}, \quad \text{with} \quad \bar{z} \triangleq \max\{z_\rho, z_\sigma\}, \quad (62)$$

where z_ρ and z_σ are defined in (57) and (61), respectively.

In the remaining of this section, we will simplify the expression of $\bar{z}$ by making a specific choice of γ . We do not aim to find the optimal value of γ but a convenient feasible choice yielding an insightful expression of the rate. We set

$$\gamma = L_\Sigma + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4}, \quad (63)$$

where $C_3 > 0$ is some absolute constant (to be determined). Recall that γ must also satisfy (48) and (51), which will impose conditions on the parameters on the right hand side of (63). With this choice we have

$$\begin{aligned}
z_\rho &= \rho + \sqrt{\rho}(1-\rho)^3 \frac{C_2^2}{C_3} \left(\sqrt{1 + \frac{\rho\sqrt{\rho}}{(1-\rho)^4} \left(\frac{2C_3}{C_2^2} \right)} + 1 \right) \\
&\leq \rho + \sqrt{\rho}(1-\rho)^3 \frac{C_2^2}{C_3} \left(\frac{\sqrt{\rho\sqrt{\rho}}}{(1-\rho)^2} \cdot \frac{\sqrt{2C_3}}{C_2} + 2 \right) \\
&\leq \rho + \frac{\sqrt{2}C_2}{\sqrt{C_3}} \rho(1-\rho) + \frac{2C_2^2}{C_3} \sqrt{\rho}(1-\rho)^3 \triangleq \bar{z}_\rho.
\end{aligned} \tag{64}$$

We focus now on z_σ . In view of Lemma 16, we bound next the quantities A/B and $\rho + \rho/B$. Define

$$\sigma_0 \triangleq \frac{1 - \kappa^{-1} + C_1 s(\tau_\mu + \tau_g)/L_\Sigma}{1 - C_1 s\tau_g/L_\Sigma}, \quad \kappa \triangleq \frac{L_\Sigma}{\mu_\Sigma}, \tag{65}$$

where we implicitly assume $1 - C_1 s\tau_g/L_\Sigma > 0$; we have the following.

Lemma 17. *Consider A and D defined in (58); and suppose that the following conditions hold*

$$\mu_\Sigma > 4C_1 s(\tau_\mu + \tau_g) + \frac{\ell_\Sigma}{\mu_\Sigma} \frac{8C_2^2 c_m^2 \rho}{(1-\rho)^2} s\tau_\ell + \rho \cdot \frac{\mu_\Sigma}{\ell_\Sigma} s\tau_\ell, \tag{66}$$

and $\rho \leq 1/2$. Then, A/D and $\rho + \rho/D$ can be bounded as

$$\frac{A}{D} \leq \sigma_0 + \sqrt{\rho} \cdot \frac{16C_3 c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} + \frac{1}{2} \left(\kappa^{-1} + \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell \right)}{1 - C_1 s\tau_g/L_\Sigma}, \tag{67}$$

$$\rho + \frac{\rho}{D} \leq \rho + \rho \cdot \frac{\frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \cdot 4C_2^2 c_m^2 s\tau_\ell + \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell}{1 - 2C_1 s\tau_g/L_\Sigma}. \tag{68}$$

Proof. See Appendix G (supplementary material). □

Using Lemma 16 and Lemma 17, z_σ can be readily bounded as

$$\begin{aligned}
z_\sigma &= \frac{A}{D} + \rho + \frac{\rho}{D} \\
&\leq \sigma_0 + \sqrt{\rho} \cdot \frac{16C_3 c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} + \frac{1}{2} \left(\kappa^{-1} + \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell \right)}{1 - C_1 s\tau_g/L_\Sigma} + \rho + \rho \cdot \frac{\frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \cdot 4C_2^2 c_m^2 s\tau_\ell + \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell}{1 - 2C_1 s\tau_g/L_\Sigma} \\
&\leq \sigma_0 + \sqrt{\rho} \cdot \frac{18C_3 c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} + \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \cdot 4C_2^2 c_m^2 s\tau_\ell + \frac{2\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell}{1 - 2C_1 s\tau_g/L_\Sigma},
\end{aligned} \tag{69}$$

where in the last inequality we have used the fact that $C_3 c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \geq 1$, $\rho \leq 1$ and $\kappa^{-1} \leq 1$. To make the bound not vacuous, we need to assume

$$1 - 2C_1 s\tau_g/L_\Sigma > 0. \tag{70}$$

Using (64) and (69), we we can then bound $\bar{z}$ in (62) as

$$\bar{z} \leq \max \left\{ \rho + \frac{\sqrt{2}C_2}{\sqrt{C_3}} \rho(1-\rho) + \frac{2C_2^2}{C_3} \sqrt{\rho}(1-\rho)^3, \right.$$

$$\sigma_0 + \sqrt{\rho} \cdot \frac{18C_3c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} + \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \cdot 4C_2^2 c_m^2 s\tau_\ell + \frac{2\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell}{1 - 2C_1 s\tau_g/L_\Sigma} \Bigg\}. \quad (71)$$

We further simplify (71), as follows. Recall the definition of σ_0 as in (65). We enforce the second argument in the max-expression in (71) to be no larger than

$$\frac{1 - (2\kappa)^{-1} + C_1 s(\tau_\mu + \tau_g)/L_\Sigma}{1 - 2C_1 s\tau_g/L_\Sigma},$$

by letting

$$\sqrt{\rho} \leq \left\{ 2\kappa \left(18C_3c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} + \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \cdot 4C_2^2 c_m^2 s\tau_\ell + \frac{2\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell \right) \right\}^{-1}.$$

Therefore, choosing $C_3 = 4$, we can upper bound the RHS of (71) as

$$\bar{z} \leq \bar{z}_{\text{up}} \triangleq \max \left\{ \rho + C_2\rho + \frac{C_2^2}{2}\sqrt{\rho}, \frac{1 - (2\kappa)^{-1} + C_1 s(\tau_\mu + \tau_g)/L_\Sigma}{1 - 2C_1 s\tau_g/L_\Sigma} \right\}, \quad (72)$$

where ρ satisfies

$$\begin{aligned} \rho &\leq \left\{ 2\kappa \left(72c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} + \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \cdot 4C_2^2 c_m^2 s\tau_\ell + \frac{2\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell \right) \right\}^{-2} \\ &= \left\{ 2 \left(72c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma^2} + \frac{\ell_\Sigma}{\mu_\Sigma^2} \cdot 4C_2^2 c_m^2 s\tau_\ell + \frac{2}{\ell_\Sigma} s\tau_\ell \right) \right\}^{-2}. \end{aligned} \quad (73)$$

Therefore, we can conclude that (55a)-(55b) are satisfied by any z and γ such that $z > \bar{z}_{\text{up}}$, with $\bar{z}_{\text{up}}$ defined in (72), and

$$\gamma = L_\Sigma + 4 \frac{\ell_\Sigma^2}{\mu_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4}, \quad (74)$$

as long as the following conditions hold: (51), with ϵ chosen according to (56); (66); and (73) [note that conditions (48) and (70) are already implied by (74) and (66), respectively].

Next, we further simplify these conditions.

First, (51), with ϵ given by (56), is a consequence of (66) and (74). In fact, substituting the expression of γ and ϵ into (51), (51) reads

$$\begin{aligned} L_\Sigma + 4 \frac{\ell_\Sigma^2}{\mu_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4} &> C_1 s\tau_g + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma} (\ell_\Sigma + s\tau_\ell) = C_1 s\tau_g + \frac{\rho}{2} \mu_\Sigma + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma} (s\tau_\ell) \\ \Leftrightarrow \quad 2L_\Sigma - \rho\mu_\Sigma + 8 \frac{\ell_\Sigma^2}{\mu_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4} &> 2C_1 s\tau_g + \rho \frac{\mu_\Sigma}{\ell_\Sigma} (s\tau_\ell). \end{aligned}$$

Since $\rho \leq 1$ and $\mu_\Sigma \leq L_\Sigma$, clearly (66) implies (51).

Second, we claim that (66) is implied by (73) and the following condition on the RSC/RSM parameters

$$\mu_\Sigma > 36C_1 s(\tau_\mu + \tau_g). \quad (75)$$

Using the facts that $c_m \geq 1$ and $\ell_\Sigma/\mu_\Sigma \geq 1$, (73) implies the following: $\rho \leq 1/4$ and

$$2\sqrt{\rho} \left(\frac{\ell_\Sigma}{\mu_\Sigma^2} \cdot 4C_2^2 c_m^2 s\tau_\ell + \frac{2}{\ell_\Sigma} s\tau_\ell \right) \leq 1. \quad (76)$$

Consequently, we can upperbound the second and third summands on the RHS of (66) as

$$\frac{\ell_\Sigma}{\mu_\Sigma} \frac{8C_2^2 c_m^2 \rho}{(1-\rho)^2} s\tau_\ell + \rho \cdot \frac{\mu_\Sigma}{\ell_\Sigma} s\tau_\ell$$

$$\begin{aligned}
&\leq \frac{\ell_\Sigma}{\mu_\Sigma} \frac{8C_2^2 c_m^2 \rho}{9/16} s\tau_\ell + \rho \cdot \frac{\mu_\Sigma}{\ell_\Sigma} s\tau_\ell \\
&\leq \frac{16}{9} \cdot \rho \left(\frac{\ell_\Sigma}{\mu_\Sigma} 8C_2^2 c_m^2 s\tau_\ell + \frac{4\mu_\Sigma}{\ell_\Sigma} s\tau_\ell \right) \leq \frac{16}{9} \sqrt{\rho} \mu_\Sigma \leq \frac{8}{9} \mu_\Sigma.
\end{aligned}$$

This together with (75) implies (66).

In summary, we proved that (55a)-(55b) are satisfied by any $z > \bar{z}_{\text{up}}$, with $\bar{z}_{\text{up}}$ defined in (72), and γ given by (74), if conditions (73) and (75) hold.

• **Step 2: Bounding $B(z)$ [cf. (53)] and Δ_{stat} [cf. (54)]**

We begin providing the bound of $B(z)$. According to Lemma 15 and (64), z is lower bounded by

$$\begin{aligned}
z &\geq \rho + \sqrt{\rho}(1-\rho)^3 \frac{C_2^2}{C_3} \left(\sqrt{1 + \frac{\rho\sqrt{\rho}}{(1-\rho)^4} \left(\frac{4C_3}{C_2^2} \right) + 1} \right) \\
&\geq \rho + \sqrt{\rho}(1-\rho)^3 \frac{C_2^2}{C_3}.
\end{aligned} \tag{77}$$

Substituting the expression of γ [cf. (74)] and ϵ [cf. (56)] into (53):

$$\begin{aligned}
B(z) &= \frac{C_2 s\tau_\ell}{L_\Sigma + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4}} \frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_2 c_m^2}{1-\rho} \frac{\rho}{z-\rho} \|\Theta^0 - \hat{\Theta}\|^2 \\
&\quad + \frac{C_2}{L_\Sigma + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4}} m c_m \left(A_3 \cdot \frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_2 c_m}{1-\rho} + s c_g^2 \frac{1}{\mu_\Sigma} \frac{2C_2 c_m}{1-\rho} + c_g \nu \right) \frac{\rho}{z-\rho} \\
&\quad + \left(1 - \frac{1}{L_\Sigma + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4}} \left\{ C_1 s\tau_g + \frac{\rho}{2} (\ell_\Sigma + s\tau_\ell) \frac{\mu_\Sigma}{\ell_\Sigma} \right\} \right) \|\Theta^{1/2} - \hat{\Theta}\|^2 \\
&\leq \frac{C_2 s\tau_\ell}{L_\Sigma} \frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_2 c_m^2}{1-\rho} \frac{\rho}{z-\rho} \|\Theta^0 - \hat{\Theta}\|^2 \\
&\quad + \frac{C_2}{L_\Sigma} m c_m \left(A_3 \cdot \frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_2 c_m}{1-\rho} + s c_g^2 \frac{1}{\mu_\Sigma} \frac{2C_2 c_m}{1-\rho} + c_g \nu \right) \frac{\rho}{z-\rho} + \|\Theta^{1/2} - \hat{\Theta}\|^2 \\
&\stackrel{(77)}{\leq} \frac{s\tau_\ell}{L_\Sigma} \frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_3 c_m^2 \sqrt{\rho}}{(1-\rho)^4} \|\Theta^0 - \hat{\Theta}\|^2 \\
&\quad + \frac{1}{L_\Sigma} m c_m \sqrt{\rho} \left(A_3 \cdot \frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_2 c_m}{1-\rho} + s c_g^2 \frac{1}{\mu_\Sigma} \frac{2C_2 c_m}{1-\rho} + c_g \nu \right) \frac{C_3}{(1-\rho)^3 C_2} + \|\Theta^{1/2} - \hat{\Theta}\|^2.
\end{aligned} \tag{78}$$

Substituting the expression of A_3 [cf. Prop. 10] in the above bound and rearranging terms yield

$$\begin{aligned}
B(z) &\leq \sqrt{\rho} \frac{s\tau_\ell}{L_\Sigma} \frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_3 c_m^2}{(1-\rho)^4} \|\Theta^0 - \widehat{\Theta}\|^2 \\
&\quad + m\sqrt{\rho} \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \frac{2C_3 c_m^2}{(1-\rho)^4} (\ell_\Sigma \|\Theta^*\|^2 + \tau_\ell \|\Theta^*\|_1^2) \\
&\quad + \frac{1}{L_\Sigma} m c_m \sqrt{\rho} \left(s c_g^2 \frac{1}{\mu_\Sigma} \frac{2C_2 c_m}{1-\rho} + c_g \nu \right) \frac{C_3}{(1-\rho)^3 C_2} + \|\Theta^{1/2} - \widehat{\Theta}\|^2 \\
&\leq \sqrt{\rho} \frac{s\tau_\ell}{L_\Sigma} \frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_3 c_m^2}{(1-\rho)^4} \|\Theta^0 - \widehat{\Theta}\|^2 \\
&\quad + m\sqrt{\rho} \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \frac{2C_3 c_m^2}{(1-\rho)^4} (\ell_\Sigma \|\Theta^*\|^2 + s\tau_\ell \|\Theta^*\|^2) \quad (\Theta^* \text{ is } s\text{-sparse}) \\
&\quad + \frac{1}{L_\Sigma} m c_m \sqrt{\rho} \left(s c_g^2 \frac{1}{\mu_\Sigma} \frac{2C_2 c_m}{1-\rho} + c_g \nu \right) \frac{C_3}{(1-\rho)^3 C_2} + \|\Theta^{1/2} - \widehat{\Theta}\|^2.
\end{aligned} \tag{79}$$

Since (73) is equivalent to

$$2\sqrt{\rho}\kappa \left(18C_3 c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} + \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \cdot 4C_2^2 c_m^2 s\tau_\ell + \frac{2\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell \right) \leq 1, \tag{80}$$

together with the fact that $\kappa \geq 1$, we have

$$\sqrt{\rho} \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} C_3 c_m^2 \leq \frac{1}{36} \quad \text{and} \quad \sqrt{\rho} \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \cdot C_2^2 c_m^2 s\tau_\ell \leq \frac{1}{8}. \tag{81}$$

In addition, from (73) we get $\rho \leq 1/4$, and thus we conclude that there exists a constant $C_4 > 0$ such that

$$\frac{B(z)}{m} \leq \frac{C_4}{m} \left(\|\Theta^0 - \widehat{\Theta}\|^2 + \|\Theta^*\|^2 + \|\Theta^{1/2} - \widehat{\Theta}\|^2 + \frac{m c_m \sqrt{\rho}}{L_\Sigma} \left(\frac{c_m s c_g^2}{\mu_\Sigma} + c_g \nu \right) \right) \triangleq B. \tag{82}$$

We bound now $\overline{\Delta}_{\text{stat}}$. Using the expression of γ [cf. (74)] and ϵ [cf. (56)], we have

$$\begin{aligned}
\overline{\Delta}_{\text{stat}} &= \frac{C_2}{\gamma} \frac{\rho \cdot c_m}{1-\rho} \tau_\ell \nu^2 (\epsilon + \epsilon^{-1}) + \frac{C_1(\tau_\mu + \tau_g)}{\gamma} \nu^2 \\
&\leq \frac{C_2}{L_\Sigma} \frac{\rho \cdot c_m}{1-\rho} \tau_\ell \nu^2 \left(\frac{\mu_\Sigma}{\ell_\Sigma} \frac{1-\rho}{2C_2 c_m} + \frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_2 c_m}{1-\rho} \right) + \frac{C_1(\tau_\mu + \tau_g)}{L_\Sigma} \nu^2 \\
&= \left(\frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} + \rho \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \frac{2C_2^2 c_m^2}{(1-\rho)^2} \right) \tau_\ell \nu^2 + \frac{C_1(\tau_\mu + \tau_g)}{L_\Sigma} \nu^2 \\
&\leq \frac{\rho}{2L_\Sigma} \left(1 + \frac{\ell_\Sigma}{\mu_\Sigma} \frac{4C_2^2 c_m^2}{(1-\rho)^2} \right) \tau_\ell \nu^2 + \frac{C_1(\tau_\mu + \tau_g)}{L_\Sigma} \nu^2 \\
&\leq \underbrace{\frac{\rho}{2L_\Sigma} \left(\frac{\ell_\Sigma}{\mu_\Sigma} \frac{5C_2^2 c_m^2}{(1-\rho)^2} \right) \tau_\ell \nu^2 + \frac{C_1(\tau_\mu + \tau_g)}{L_\Sigma} \nu^2}_{\Delta_{\text{stat}}} \quad (\text{given } C_2 \geq 1).
\end{aligned} \tag{83}$$

• **Step 3: Linear convergence up to $o(\|\widehat{\Theta} - \Theta^*\|^2)$**

We can now state our main convergence result building on Steps 1 & 2 and Lemma 14.

Recalling the definitions of B and Δ_{stat} , namely:

$$B = \frac{C_4}{m} \left(\|\Theta^0 - \widehat{\Theta}\|^2 + \|\Theta^*\|^2 + \|\Theta^{1/2} - \widehat{\Theta}\|^2 + \frac{m c_m \sqrt{\rho}}{L_\Sigma} \left(\frac{c_m s c_g^2}{\mu_\Sigma} + c_g \nu \right) \right), \tag{84}$$

with

$$c_g = \max_{j \in [m]} \|\nabla \mathcal{L}_j(\boldsymbol{\theta}^*)\|_\infty + \|\nabla \mathcal{L}(\boldsymbol{\theta}^*)\|_\infty \quad (85)$$

$$\nu = 2\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_1 + 2\sqrt{s}\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|; \quad (86)$$

and

$$\Delta_{\text{stat}} = \frac{\rho}{2L_\Sigma} \left(\frac{\ell_\Sigma}{\mu_\Sigma} \frac{5C_2^2 c_m^2}{(1-\rho)^2} \right) \tau_\ell \nu^2 + \frac{C_1(\tau_\mu + \tau_g)}{L_\Sigma} \nu^2, \quad (87)$$

we have the following.

Theorem 18. *Given the linear model (1), with $\boldsymbol{\theta}^*$ s -sparse, consider the LASSO problem (2) over a network $\mathcal{G}$ satisfying Assumption 6. Assume that the global $\mathcal{L}$ and local $\mathcal{L}_i$ loss functions satisfy Assumption 2 and Assumption 4, respectively. Let $\{(\boldsymbol{\theta}_i^t)_{i=1}^m\}$ be the sequence generated by the NetLASSO algorithm with step size γ chosen according to (74) and the weight matrix $\mathbf{W}$ satisfying Assumption 7. Further, assume that the following two conditions hold:*

$$\mu_\Sigma > 36C_1 s(\tau_\mu + \tau_g) \quad (88)$$

and

$$\rho \leq \left\{ 2 \left(75c_m^2 C_2^2 \frac{\ell_\Sigma^2}{\mu_\Sigma^2} + \frac{\ell_\Sigma}{\mu_\Sigma^2} \cdot 6C_2^2 c_m^2 s \tau_\ell \right) \right\}^{-2}, \quad (89)$$

for some constants $C_1 > 0$ and $C_2 \geq 1$.

Then, for any optimal solution $\hat{\boldsymbol{\theta}}$ of the LASSO problem satisfying $\|\hat{\boldsymbol{\theta}}\|_1 = r$, we have

$$\frac{1}{m} \sum_{i=1}^m \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|^2 \leq B \cdot \lambda^t + \frac{\Delta_{\text{stat}}}{1-\lambda}, \quad \forall t = 1, 2, \dots, \quad (90)$$

where

$$\lambda = \frac{1 - (2\kappa)^{-1} + C_1 s(\tau_\mu + \tau_g)/L_\Sigma}{1 - 2C_1 s \tau_g/L_\Sigma} < 1, \quad (91)$$

$\kappa = L_\Sigma/\mu_\Sigma$, and B and Δ_{stat} are defined in (84) and (87), respectively.

Proof. First of all, note that (89) implies (73), due to the following chain of inequalities

$$\begin{aligned} & 2 \left(72c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma^2} + \frac{\ell_\Sigma}{\mu_\Sigma^2} \cdot 4C_2^2 c_m^2 s \tau_\ell + \frac{2}{\ell_\Sigma} s \tau_\ell \right) \\ & \leq 2 \left(72c_m^2 C_2^2 \frac{\ell_\Sigma^2}{\mu_\Sigma^2} + 3C_2^2 c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma^2} + \frac{\ell_\Sigma}{\mu_\Sigma^2} \cdot 4C_2^2 c_m^2 s \tau_\ell + \frac{2}{\ell_\Sigma} s \tau_\ell \right) \\ & \leq 2 \left(75c_m^2 C_2^2 \frac{\ell_\Sigma^2}{\mu_\Sigma^2} + \frac{\ell_\Sigma}{\mu_\Sigma^2} \cdot 4C_2^2 c_m^2 s \tau_\ell + \frac{2}{\ell_\Sigma} s \tau_\ell \right) \\ & \leq 2 \left(75c_m^2 C_2^2 \frac{\ell_\Sigma^2}{\mu_\Sigma^2} + \frac{\ell_\Sigma}{\mu_\Sigma^2} \cdot 4C_2^2 c_m^2 s \tau_\ell + \frac{2}{\ell_\Sigma} \frac{\ell_\Sigma^2}{\mu_\Sigma^2} C_2^2 c_m^2 s \tau_\ell \right) \\ & \leq 2 \left(75c_m^2 C_2^2 \frac{\ell_\Sigma^2}{\mu_\Sigma^2} + \frac{\ell_\Sigma}{\mu_\Sigma^2} \cdot 6C_2^2 c_m^2 s \tau_\ell \right). \end{aligned}$$

Furthermore, $\rho < (3C_2^2)^{-2}$.

Therefore (55a)-(55b) are satisfied by any $z > \bar{z}_{\text{up}}$ [cf. Steps 1 & 2], with $\bar{z}_{\text{up}}$ defined in (72). It remains to show that i) $\lambda < 1$; and ii) $\bar{z}_{\text{up}} \leq \lambda$.

i) It is not difficult to check that $\mu_\Sigma > 36C_1 s(\tau_\mu + \tau_g)$ implies $z < 1 - (4\kappa)^{-1}$.

ii) We prove

$$\rho + C_2\rho + \frac{C_2^2}{2}\sqrt{\rho} \leq \lambda,$$

and thus $\bar{z}_{\text{up}} = \lambda$. It is sufficient to check that the following chain of inequalities hold:

$$\begin{aligned} & \left(\rho + C_2\rho + \frac{C_2^2}{2}\sqrt{\rho} \right) (1 - 2C_1s\tau_g/L_\Sigma) \\ & \leq 3C_2^2\sqrt{\rho} \stackrel{(89)}{\leq} \frac{1}{50} \cdot \frac{\mu_\Sigma^2}{c_m^2\ell_\Sigma^2} \leq \frac{1}{2\kappa} \leq 1 - \frac{1}{2\kappa} + C_1s(\tau_\mu + \tau_g)/L_\Sigma, \end{aligned} \quad (92)$$

where we used $\rho < 1$, $c_m, C_2 \geq 1$, $\ell_\Sigma \geq L_\Sigma$ and $\kappa \geq 1$.

Given the setting above, the convergence as stated in (90) follows from Lemma 14. $\square$

The residual error Δ_{stat} depends, in particular, on the parameters of the RSC/RSM and the statistical error $\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*$. The next corollary provides sufficient conditions for $\Delta_{\text{stat}} = o(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|^2)$, implying thus linear convergence up to a residual error smaller than the statistical precision.

Corollary 19. *Instate assumptions of Theorem 18; and suppose $r \leq \|\boldsymbol{\theta}^*\|_1$ and*

$$s(\tau_\mu + \tau_g) = o(1) \quad \text{and} \quad s \cdot \rho c_m^2 \frac{\ell_\Sigma}{\mu_\Sigma} \tau_\ell = o(1). \quad (93)$$

Then, we have

$$\frac{1}{m} \sum_{i=1}^m \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|^2 \leq \lambda^t \cdot B + o\left(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|^2\right). \quad (94)$$

Proof. Using [1, Lemma 5] yields $\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_1 \leq 2\sqrt{s}\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|$. Using $\rho \leq 1/4$ and conditions in (93) it is not difficult to verify that Δ_{stat} in (87) is $o(\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|^2)$. $\square$

5.5 Linear convergence for sparse vector regression (Theorem 22)

We customize now Theorem 18 under the parameter settings $(\mu_\Sigma, L_\Sigma, \ell_\Sigma)$, $(\tau_\mu, \tau_g, \tau_\ell)$, and c_g guaranteed by Propositions 3 and 5 for the sparse vector regression model (1). This will lead to the formal statement and proof of Theorem 22, informally discussed in Sec. 4.

We begin deriving a high probability bound for $c_g = \max_{j \in [m]} \|\nabla \mathcal{L}_j(\boldsymbol{\theta}^*)\|_\infty + \|\nabla \mathcal{L}(\boldsymbol{\theta}^*)\|_\infty$.

Lemma 20. *Consider the linear model (1). Suppose $\{\mathbf{X}_i\}_{i=1}^m$ are i.i.d. random matrices drawn from the Σ -Gaussian ensemble; and the elements of $\{\mathbf{n}_i\}_{i=1}^m$ are i.i.d. σ^2 -sub-Gaussian random variables, independent of the $\mathbf{X}_i$'s. Then*

$$\mathbb{P} \left(\max_{j \in [m]} \left\| \frac{\mathbf{X}_j^\top \mathbf{n}_j}{n} \right\|_\infty \geq \sqrt{\zeta} \sigma \cdot \max \left\{ \sqrt{\frac{2 \log md}{c_3 n}}, \frac{2 \log md}{c_3 n} \right\} \right) \leq 2 \exp(-\log d) \quad (95)$$

and

$$\mathbb{P} \left(\left\| \frac{\mathbf{X}^\top \mathbf{n}}{N} \right\|_\infty \geq \sqrt{\zeta} \sigma \cdot \max \left\{ \sqrt{\frac{2 \log d}{c_3 N}}, \frac{2 \log d}{c_3 N} \right\} \right) \leq 2 \exp(-\log d) \quad (96)$$

for some $c_3 > 0$.

Proof. See Appendix H (supplementary material). $\square$

According to (1) we have $\|\nabla \mathcal{L}(\boldsymbol{\theta}^*)\|_\infty = \|\mathbf{X}^\top \mathbf{n}/N\|_\infty$ and $\|\nabla \mathcal{L}_j(\boldsymbol{\theta}^*)\|_\infty = \|\mathbf{X}_j^\top \mathbf{n}_j/n\|_\infty$. Using Lemma 20 we can readily obtain the following bound for c_g^2 .

Corollary 21. *Reintate the conditions of Lemma 20; there holds:*

$$\begin{aligned} c_g^2 &\leq 4\zeta\sigma^2 \left(\frac{2 \log md}{c_3 n} + \left(\frac{2 \log md}{c_3 n} \right)^2 + \frac{2 \log d}{c_3 N} + \left(\frac{2 \log d}{c_3 N} \right)^2 \right) \\ &\leq 4\zeta\sigma^2 \left(\frac{4 m \log md}{c_3 N} + \left(\frac{4 m \log md}{c_3 N} \right)^2 \right), \end{aligned} \quad (97)$$

with probability greater than $1 - 4 \exp(-c_4 \log d)$, for some $c_4 > 0$.

We can now leverage Theorem 18 with the parameter settings $(\mu_\Sigma, L_\Sigma, \ell_\Sigma)$, $(\tau_\mu, \tau_g, \tau_\ell)$, and c_g guaranteed by Proposition 3, Proposition 5, and Corollary 21; we have the following.

Theorem 22. *Given the linear model (1), suppose $\{\mathbf{X}_i\}_{i=1}^m$ are i.i.d. random matrices drawn from the Σ -Gaussian ensemble; and the elements of $\{\mathbf{n}_i\}_{i=1}^m$ are i.i.d. σ^2 -sub-Gaussian random variables, independent of the $\mathbf{X}_i$'s. Consider the LASSO problem (2) over a network $\mathcal{G}$ satisfying Assumption 6; further assume for (2): $\|\hat{\boldsymbol{\theta}}\|_1 = r \leq \|\boldsymbol{\theta}^*\|_1$, $\frac{\zeta}{\mu_\Sigma} \frac{s \log d}{N} < c_5$ and $c_6 \rho m^8 \kappa^4 < 1$, for some constants $c_5, c_6 > 0$. Let $\{(\boldsymbol{\theta}_i^t)_{i=1}^m\}$ be the sequence generated by the NetLASSO algorithm with step size γ chosen according to (74) and the weight matrix $\mathbf{W}$ satisfying Assumption 7. Then, the following holds:*

$$\frac{1}{m} \sum_{i=1}^m \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|^2 \leq B \cdot \left(\frac{1 - (2\kappa)^{-1} + C' s \log d/N}{1 - C' s \log d/N} \right)^t + c_7 \left(\frac{\zeta}{L_\Sigma} \frac{s \log d}{N} \right) \|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|^2, \quad (98)$$

with probability greater than $1 - c_8 \exp(-c_9 \log d)$, for some constants $C', c_7, c_8, c_9 > 0$. The expression of $B > 0$ is given in (104).

Proof. The proof consists in applying Theorem 18 under the setting guaranteed by Proposition 3, Proposition 5, and Corollary 21. Specifically, let

$$\begin{aligned} \mu_\Sigma &= \frac{1}{2} \sigma_{\min}(\Sigma) \quad \text{and} \quad \tau_\mu = c_1 \zeta \frac{\log d}{N} \quad (\text{for the RSC}), \\ L_\Sigma &= 2\sigma_{\max}(\Sigma) \quad \text{and} \quad \tau_g = c_1 \zeta \frac{\log d}{N} \quad (\text{for the RSM}), \end{aligned} \quad (99)$$

and

$$\ell_\Sigma = 16m\sigma_{\max}(\Sigma) \quad \text{and} \quad \tau_\ell = c_1 \zeta m^2 \frac{\log d}{N} \quad (\text{for the local RSM}); \quad (100)$$

Propositions 3 & 5 and Corollary 21 guarantee that, with the above choices for $(\mu_\Sigma, L_\Sigma, \ell_\Sigma)$ and $(\tau_\mu, \tau_g, \tau_\ell)$, the global RSC and RSM conditions (cf. Assumption 2), the local RSM property (cf. Assumption 4), and the bound (97) on c_g all hold with probability at least $1 - c_8 \exp(-c_9 \log d)$, where we invoked the union bound and used $\log d \leq s \log d \leq c_5 \frac{\mu_\Sigma}{\zeta} N \leq c_5 N$, due to $\frac{\zeta}{\mu_\Sigma} \frac{s \log d}{N} < c_5$. So are conditions (88)-(89) in Theorem 18, as shown next.

Since $\frac{\zeta}{\mu_\Sigma} \frac{s \log d}{N} < c_5$, (88) is satisfied for sufficiently small c_5 . A sufficient condition for (89) is

$$\rho \leq C_6 \left(m^3 \kappa^2 + m^4 \kappa^2 \frac{\zeta}{L_\Sigma} \frac{s \log d}{N} \right)^{-2}, \quad (101)$$

for some $C_6 > 0$; (101) is implied by $c_6 \rho m^8 \kappa^4 < 1$ and $\frac{\zeta}{\mu_\Sigma} \frac{s \log d}{N} < c_5$, for sufficiently large $c_6 > 0$.

Finally, we use (99)-(100) to further simplify the expressions of the statistical error (83), the rate (91), and B in (84).

The statistical error can be bounded as

$$\begin{aligned}
\Delta_{\text{stat}} &= \frac{\rho}{2L_\Sigma} \left(\frac{\ell_\Sigma}{\mu_\Sigma} \frac{5C_2^2 c_m^2}{(1-\rho)^2} \right) \tau_\ell \nu^2 + \frac{C_1(\tau_\mu + \tau_g)}{L_\Sigma} \nu^2 \\
&= \rho \left(\frac{4}{\mu_\Sigma} \frac{5C_2^2 m^2}{(1-\rho)^2} \right) c_1 \zeta \frac{m^2 \log d}{N} \nu^2 + \frac{C_1}{L_\Sigma} \left(2c_1 \zeta \frac{\log d}{N} \right) \nu^2 \\
&\leq C_5 \left(\frac{m^4 \rho \kappa}{(1-\rho)^2} \frac{\zeta}{L_\Sigma} \frac{\log d}{N} + \frac{\zeta}{L_\Sigma} \frac{\log d}{N} \right) \nu^2 \\
&\leq c_7 \left(\frac{\zeta}{L_\Sigma} \frac{s \log d}{N} \right) \|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|^2,
\end{aligned} \tag{102}$$

for some $C_5, c_7 > 0$, where in the last inequality we used $c_6 \rho m^8 \kappa^4 < 1$, $\nu^2 = \mathcal{O}(s \|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|^2)$ (due to $\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_1 \leq 2\sqrt{s} \|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|$ [1, Lemma 5]), and $\rho < (3C_2^2)^{-2}$ [see, e.g., (89)].

The rate λ in (91) can be bounded as

$$\lambda = \left(1 - (2\kappa)^{-1} + 2C_1 c_1 \frac{\zeta}{L_\Sigma} \frac{s \log d}{N} \right) \left(1 - 2C_1 c_1 \frac{\zeta}{L_\Sigma} \frac{s \log d}{N} \right)^{-1} < 1, \tag{103}$$

where the inequality is due to (88) [cf. Theorem 18].

Finally, for B in (84), we have

$$\begin{aligned}
B &= \frac{C_4}{m} \left(\|\boldsymbol{\Theta}^0 - \hat{\boldsymbol{\Theta}}\|^2 + \|\boldsymbol{\Theta}^*\|^2 + \|\boldsymbol{\Theta}^{1/2} - \hat{\boldsymbol{\Theta}}\|^2 + \frac{m c_m \sqrt{\rho}}{L_\Sigma} \left(\frac{c_m s c_g^2}{\mu_\Sigma} + c_g \nu \right) \right) \\
&\leq \frac{C_4}{m} \left(\|\boldsymbol{\Theta}^0 - \hat{\boldsymbol{\Theta}}\|^2 + \|\boldsymbol{\Theta}^*\|^2 + \|\boldsymbol{\Theta}^{1/2} - \hat{\boldsymbol{\Theta}}\|^2 \right) \\
&\quad + C_4 \frac{4\zeta \sigma^2}{\mu_\Sigma L_\Sigma} m s \sqrt{\rho} \left(\frac{4}{c_3} \frac{m \log md}{N} + \left(\frac{4}{c_3} \frac{m \log md}{N} \right)^2 \right) \\
&\quad + C_4 \frac{12\sqrt{\zeta} \sigma}{L_\Sigma} \sqrt{m \rho s} \left(\sqrt{\frac{4}{c_3} \frac{m \log md}{N}} + \frac{4}{c_3} \frac{m \log md}{N} \right) \|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|,
\end{aligned} \tag{104}$$

where we used again $\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_1 \leq 2\sqrt{s} \|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|$.

The proof is completed invoking Theorem 18. $\square$

6 Numerical Results

In this section we provide some numerical results that validate the accuracy of our theoretical findings, in particular Theorem 22 and its implications.

We consider the following problem setup. Given the distributed linear regression model (1), the unknown s -sparse vector $\boldsymbol{\theta}^*$ is generated with the first s elements being i.i.d. $\mathcal{N}(0, 1)$; the remaining coordinates are set to zero. Each row in $\mathbf{X}_i \in \mathbb{R}^{n \times d}$ is generated according to $\mathcal{N}(\mathbf{0}, \mathbf{I})$; each element of $\mathbf{n}_i$ follows $\mathcal{N}(0, 0.25)$; and r is set $r = \|\boldsymbol{\theta}^*\|_1$. Unless otherwise specified, we generate the communication network among the agents using a random Erdős-Rényi graph $G(m, p)$, where m is the number of nodes/agents and each edge is included in the graph independently with probability p . The specific values of problem parameters (N, d, s) and network parameters (m, p) are specified in each simulation.

Aiming at validating Theorem 22, we report next experiments showing the following: (i) the scalability of the algorithm with respect to the total sample size N and problem dimension d , for a fixed network; and (ii) the impact of network topology on the algorithm.




Figure 4: Plot of the log-normalized estimation error $\log\left(\frac{1}{m} \sum_{i=1}^m \|\theta_i^t - \theta^*\|^2 / \|\theta^*\|^2\right)$ for $\alpha \approx 0.2$. In these problems $d \in \{5000, 10000, 20000\}$, $s = \lceil \sqrt{d} \rceil$, $m = 50$, $n \in \{61, 94, 142\}$. Dashed lines indicate estimation error if using $\hat{\theta}$.

6.1 Scalability with respect to problem parameters for a fixed network

Theorem 22 states that, for a fixed, sufficiently connected network, the average optimization error of NetLASSO among all agents decreases at a linear, up to a tolerance depending on the statistical precision $\|\hat{\theta} - \theta^*\|$. Moreover, the expression of both the convergence rate and the tolerance term depend exclusively on the ratio $\alpha = \frac{s \log d}{N}$. To support these theoretical findings, we conduct the following two simulations over an Erdős-Rényi graph with $m = 50$ nodes and edge activation probability $p = 0.5$. The reported results are the average of 100 trials with the same graph and independently randomly generated data sets.

(i) Fixed ratio α and increasing dimension d : We set $\alpha \approx 0.2$ and consider three sets of problem parameters, namely: $d = 5000, 10000, 20000$ and $s = \lceil \sqrt{d} \rceil$. Consequently, we set the number of total samples N and local sample size n to $N = 3050, 4700, 7100$ and $n = 61, 94, 142$, respectively. Observe that $N = m \cdot n$ with $m = 50$. Since the expression of the proximal parameter γ may be conservative in practice, we tune it manually as described next. We first generate one instance of the LASSO problem with $d = 5000$ and run NetLASSO using the γ that yields the fastest convergence rate. This value is selected via grid search. The same γ is used for all problem instances. Fig. 4 plots the average normalized estimation error $(1/m) \sum_{i=1}^m \|\theta_i^t - \theta^*\|^2 / \|\theta^*\|^2$ versus the number of iterations, for $d = 5000, 10000, 20000$. The dashed-line curves represent the normalized statistical precision $(\|\hat{\theta} - \theta^*\|^2 / \|\theta^*\|^2)^{-2}$ for the three sets of problems. The following comments are in order. First, in all cases the estimation error decays at a linear rate before reaching the statistical precision. Second, the rate is invariant with respect to the change of dimension d as long as the ratio α is fixed. This is revealed by the fact that the slope of the linear part of the three curves is the same for different choices of d , s and N , as long as α remains constant. This is exactly what Theorem 22 predicts.

(ii) Increasing α : This experiment explores the relation between the convergence rate of NetLASSO and α . We set $d = 20000$, $s = \lceil (\log d)/2 \rceil$ and consider three local sample sizes, $n = 1, 5$, and 25 . Correspondingly, we have $\alpha \approx \{1, 0.2, 0.04\}$. For each value of n , we choose the proximal parameter γ via grid search. The one that yields fastest convergence rate is kept fixed for all independent trials. Fig. 5 shows that the algorithm's performance degrades as α increases—both the convergence rate of NetLASSO and the statistical precision get worse. Note that this behavior is in agreement with Theorem 22.




Figure 5: Plots of the log-normalized estimation error $\log \left(\frac{1}{m} \sum_{i=1}^m \|\theta_i^t - \theta^*\|^2 / \|\theta^*\|^2 \right)$ for different values of α . The problem parameters are set to be $d = 20000$, $s = \lceil \log(d)/2 \rceil$, $m = 50$ and $\alpha = \frac{s \log d}{N}$ with $\alpha \in \{0.04, 0.2, 1\}$. Dashed lines indicate normalized estimation error if using $\hat{\theta}$. The network is composed of $m = 50$ agents with connectivity $\rho = 0.0638$.

6.2 Impact of network scale and connectivity


This set of simulations investigates the scalability of NetLASSO with respect to network parameters (m, ρ) , for fixed problem parameters (N, d, s) . In particular, Theorem 22 established the convergence of NetLASSO requiring $\rho \lesssim m^{-8} \kappa^{-4}$. To validate this condition, we conduct the following simulations.

(i) Fixed network connectivity ρ and increasing m : The condition $\rho \lesssim m^{-8} \kappa^{-4}$ requires a better connectivity for a network involving more agents. To test the necessity of this condition, we run NetLASSO on the networks with increasing size $m = 50, 625, 1250, 2500$. Correspondingly we varied the edge connectivity probability $p = 0.87, 0.4, 0.23, 0.15$ in order to keep ρ roughly constant, equal to ≈ 0.18 . The problem parameters are set to be $N = 2500$, $d = 5000$, $s = \lceil \sqrt{d} \rceil$. The step size is chosen via grid search choosing the one yielding fastest convergence. All curves reported for this simulation in one trial are conducted on the same problem instance for different network configurations. The result is averaged over 100 trials and is reported in Fig 6a. Observe that keeping the connectivity of the graph constant while increasing the number of agents leads to a degradation of the achievable rate and estimation error.


Additionally, we empirically demonstrate that for Erdős-Rényi graphs with connectivity $\rho = \mathcal{O}(1/\sqrt{m})$ —achieved by fixing the network connectivity probability p —NetLASSO converges under a growing network size. The result is reported in Fig. 6b. The simulation is conducted under the same setting as Fig. 6a except p is set to be 0.87. This suggests that the condition $\rho \lesssim m^{-8} \kappa^{-4}$ might be more restrictive than what needed in practice, which is an artifact of our analysis.

(ii) Effect of multiple consensus rounds: Our last simulation shows that for any given connected network, running multiple rounds of consensus per computation improves the convergence rate of NetLASSO by reducing the effective ρ exponentially. To this end, we test three settings, with the LASSO problem generated in the same way as Sec. 6.1 (i). The network size is $m = 50$ and the edge connectivity p is chosen so that $\rho = 0.06, 0.4$, and 0.78 . Fig. 7 shows that, for networks with $\rho = 0.4$ and $\rho = 0.78$, one can improve their connectivity by running 3 and 9 rounds of consensus per computation, respectively. By doing so, the effective communication networks across the three settings have approximately the same (effective) connectivity. Consequently, the iteration complexity of NetLASSO remains invariant.

In Table 8 and Figure 9 we provide the value of the communication costs incurred on some of the graphs, when using Metropolis weights. In particular, Table 8 reports the smallest number of communication rounds k required to meet the condition $\rho^k \leq m^{-8}$ over a Erdős-Rényi graph with $p = 0.87$, and a line graph for $m \in \{50, 625, 1250, 2500\}$. Figure 9 shows the convergence of NetLASSO, in the setting of Table 8. The



(a) Erdős-Rényi graphs with $\rho \approx 0.18$.



(b) Erdős-Rényi graphs with link probability 0.87.

Figure 6: Plot of the log-normalized estimation error for $m \in \{50, 625, 1250, 2500\}$. In this problem $d = 5000$, $s = \lceil \sqrt{d} \rceil$, $n \in \{50, 4, 2, 1\}$ and $N = 2500$. All curves in each panel are obtained using the same data points. Black horizontal lines indicate the normalized estimation error if using $\hat{\theta}$.




Figure 7: Plots of the log-normalized estimation error $\log\left(\frac{1}{m} \sum_{i=1}^m \|\theta_i^t - \theta^*\|^2 / \|\theta^*\|^2\right)$ for different values of α and network graphs: $d = 20000$, $s = \lceil \log(d)/2 \rceil$, $m = 50$ and $\alpha = \frac{s \log d}{N}$ with $\alpha \in \{0.04, 0.2, 1\}$. Dashed-line curves indicate normalized estimation error using $\hat{\theta}$. The left panel shows convergence over a network with $\rho = 0.0638$, using a single round of communication per computation step. The mid panel shows convergence over a network with $\rho = 0.4038$, using 3 rounds of communications per computation step, resulting in $\rho^3 = 0.0658$. The right panel shows convergence for a network with $\rho = 0.781$, using 9 rounds of communications per computation step, thus $\rho^9 = 0.058$. In the three sets of simulations, the iteration complexity is the same while the communication cost increases by 3 times, due to the multi-rounds consensus procedure.

	Erdős-Rényi ($p = 0.87$)	Line graph
$m = 50$	18	23775
$m = 625$	18	6115118
$m = 1250$	18	27094153
$m = 2500$	19	118911225

Figure 8: Number of communication rounds required for each network to satisfy $\rho^k \leq m^{-8}$.

problem parameters are $d = 5000$, $s = \lceil \sqrt{d} \rceil$, $m \in \{50, 625, 1250, 2500\}$, and $n \in \{50, 4, 2, 1\}$, thus $N = 2500$. The results are consistent with what predicted in Table 8. For the Erdős-Rényi graph, a target estimation error is achieved within the same number of communication rounds, irrespective of the network size (for the range of values of m simulated). This is not the case for line graphs, which instead call for more rounds of communications, as the network size increases.




Figure 9: Plots of the log-normalized estimation error for Erdős-Rényi and line graphs, with $m \in \{50, 625, 1250, 2500\}$. In this problem, $d = 5000$, $s = \lceil \sqrt{d} \rceil$, $n \in \{50, 4, 2, 1\}$ and $N = 2500$.

7 Discussion

We studied sparse linear regression over mesh networks. We established statistical and computational guarantees in high-dimensions for NetLASSO, a decentralization of the PGD based on gradient tracking. We proved that (near) optimal sample complexity $\mathcal{O}(s \log d/N)$ for the distributed estimator is achievable over networks, even when local sample size is not sufficient for statistical consistency. Convergence at linear rate was proved, of the same order of that of PGD. This improves on DGD-like methods [19], which instead suffer from speed-accuracy dilemma.

Our work, together with the companion paper [19], represents the first attempt of understanding the behavior of some distributed algorithms for high-dimensional estimation over mesh networks. A future investigation will be the study of other distributed methods; of particular interest is understanding the role of other forms of gradient corrections towards statistical, computation, and communication tradeoffs.

8 Acknowledgment

The work of Sun, Maros and Scutari has been partially supported by the Office of Naval Research, under the Grant #N00014-21-1-2673, and the Army Research Office, under the grant W911NF1810238. The work of Cheng has been partially supported by NSF– SCALE MoDL (2134209).

References

- [1] Alekh Agarwal, Sahand Negahban, Martin J Wainwright, et al. Fast global convergence of gradient methods for high-dimensional statistical recovery. *The Annals of Statistics*, 40(5):2452–2482, 2012.
- [2] Waheed U. Bajwa, Volkan Cevher, Dimitris Papailiopoulos, and Anna Scaglione. Machine learning from distributed, streaming data [from the guest editors]. *IEEE Signal Processing Magazine*, 37(3):11–13, 2020.
- [3] Heather Battey, Jianqing Fan, Han Liu, Junwei Lu, and Ziwei Zhu. Distributed testing and estimation under sparse high dimensional models. *Annals of Statistics*, 46(3):1352, 2018.
- [4] Raphael Berthier, Francis Bach, and Pierre Gaillard. Accelerated gossip in networks of given dimension using jacobi polynomial iterations. *SIAM J. on Mathematics of Data Science*, 1:24–47, 2020.
- [5] Stephen Boyd, Neal Parikh, and Eric Chu. *Distributed optimization and statistical learning via the alternating direction method of multipliers*. Now Publishers Inc, 2011.
- [6] Bernadette Charron-Bost. Geometric bounds for convergence rates of averaging algorithms. *arXiv preprint arXiv:2007.04837*, 2020.
- [7] A. I. Chen and A. Ozdaglar. A fast distributed proximal-gradient method. In *2012 50th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pages 601–608, October 2012.
- [8] J. Chen and A. H. Sayed. Diffusion adaptation strategies for distributed optimization and learning over networks. *IEEE Transactions on Signal Processing*, 60(8):4289–4305, August 2012.
- [9] Xi Chen, Weidong Liu, and Yichen Zhang. Quantile regression under memory constraint. *The Annals of Statistics*, 47(6):3244–3273, 2019.
- [10] Xueying Chen and Min-ge Xie. A split-and-conquer approach for analysis of extraordinarily large data. *Statistica Sinica*, pages 1655–1684, 2014.
- [11] A. Daneshmand, G. Scutari, and V. Kungurtsev. Second-order guarantees of distributed gradient algorithms. *SIAM Journal on Optimization*, 30(4):3029–3068, January 2020.
- [12] P. Di Lorenzo and G. Scutari. NEXT: In-network nonconvex optimization. *IEEE Trans. Signal Inf. Process. Netw.*, 2(2):120–136, June 2016.
- [13] Jianqing Fan, Yongyi Guo, and Kaizheng Wang. Communication-efficient accurate statistical estimation. *Journal of the American Statistical Association*, 0(0):1–11, 2021.
- [14] Jianqing Fan, Dong Wang, Kaizheng Wang, and Ziwei Zhu. Distributed estimation of principal eigenspaces. *Annals of statistics*, 47(6):3009, 2019.
- [15] Hadrien Hendrikx, Lin Xiao, Sebastien Bubeck, Francis Bach, and Laurent Massoulié. Statistically preconditioned accelerated gradient method for distributed optimization. In *International Conference on Machine Learning*, pages 4203–4227. PMLR, 2020.

- [16] D. Jakovetić. A unification and generalization of exact distributed first-order methods. *IEEE Transactions on Signal and Information Processing over Networks*, 5:31–46, September 2019.
- [17] D. Jakovetić, J. Xavier, and JMF. Moura. Cooperative convex optimization in networked systems: augmented lagrangian algorithms with directed gossip communication. *IEEE Transactions on Signal Processing*, 59(8):3889–3902, July 2011.
- [18] D. Jakovetić, J. Xavier, and JMF. Moura. Fast distributed gradient methods. *IEEE Transactions on Automatic Control*, 59:1131–1146, December 2014.
- [19] Y. Ji, G. Scutari, Y. Sun, and H. Honnappa. Distributed sparse regression via penalization. *arXiv preprint, arXiv:2111.06530*, Nov. 2021.
- [20] Michael I Jordan, Jason D Lee, and Yun Yang. Communication-efficient distributed statistical inference. *Journal of the American Statistical Association*, 2018.
- [21] Ariel Kleiner, Ameet Talwalkar, Purnamrita Sarkar, and Michael I Jordan. A scalable bootstrap for massive data. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 76(4):795–816, 2014.
- [22] Jason D Lee, Qiang Liu, Yuekai Sun, and Jonathan E Taylor. Communication-efficient sparse regression. *The Journal of Machine Learning Research*, 18(1):115–144, 2017.
- [23] Tian Li, Anit Kumar Sahu, Ameet Talwalkar, and Virginia Smith. Federated learning: challenges, methods, and future directions. *IEEE Signal Processing Magazine*, 37(3):50–60, 2020.
- [24] Zhi Li, Wei Shi, and Ming Yan. A decentralized proximal-gradient method with network independent step-sizes and separated convergence rates. *IEEE Transactions on Signal Processing*, 67(17):4494–4506, 2019.
- [25] Lester Mackey, Michael Jordan, and Ameet Talwalkar. Divide-and-conquer matrix factorization. In J. Shawe-Taylor, R. Zemel, P. Bartlett, F. Pereira, and K. Q. Weinberger, editors, *Advances in Neural Information Processing Systems*, volume 24. Curran Associates, Inc., 2011.
- [26] Brendan McMahan, Eider Moore, Daniel Ramage, Seth Hampson, and Blaise Aguera y Arcas. Communication-efficient learning of deep networks from decentralized data. In Aarti Singh and Jerry Zhu, editors, *Proceedings of the 20th International Conference on Artificial Intelligence and Statistics*, volume 54 of *Proceedings of Machine Learning Research*, pages 1273–1282. PMLR, 20–22 Apr 2017.
- [27] A. Nedić, A. Olshevsky, and M. G. Rabbat. Network topology and communication-computation tradeoffs in decentralized optimization. *Proceedings of the IEEE*, 106:953–976, 2018.
- [28] A. Nedić, A. Olshevsky, and W. Shi. Achieving geometric convergence for distributed optimization over time-varying graphs. *SIAM Journal on Optimization*, 27:2597–2633, July 2016.
- [29] A. Nedić and A. Ozdaglar. Distributed subgradient methods for multi-agent optimization. *IEEE Transactions on Automatic Control*, 54(1):48–61, January 2009.
- [30] A. Nedić, A. Ozdaglar, and P. A. Parrilo. Constrained consensus and optimization in multi-agent networks. *IEEE Transactions on Automatic Control*, 55(4):922–938, April 2010.
- [31] Angelia Nedic. Distributed gradient methods for convex machine learning problems in networks. *IEEE Signal Processing Magazine*, 10:92–101, 2020.
- [32] Willie Neiswanger, Chong Wang, and Eric Xing. Asymptotically exact, embarrassingly parallel mcmc, 2014.

- [33] Yurii Nesterov. *Lectures on convex optimization*, volume 137. Springer, 2018.
- [34] G. Qu and N. Li. Harnessing smoothness to accelerate distributed optimization. *IEEE Transactions on Control of Network Systems*, 5:1245–1260, April 2017.
- [35] Garvesh Raskutti, Martin J Wainwright, and Bin Yu. Restricted eigenvalue properties for correlated gaussian designs. *The Journal of Machine Learning Research*, 11:2241–2259, 2010.
- [36] A. H. Sayed. Adaptation, learning, and optimization over networks. *Foundations and Trends in Machine Learning*, 7:311–801, January 2014.
- [37] Kevin Scaman, Francis Bach, Sébastien Bubeck, Yin Tat Lee, and Laurent Massoulié. Optimal algorithms for smooth and strongly convex distributed optimization in networks. In *Proceedings of the 34th International Conference on Machine Learning*, volume 70, pages 3027–3036, 2017.
- [38] Zuofeng Shang and Guang Cheng. Computational limits of a distributed algorithm for smoothing spline. *Journal of Machine Learning Research*, 18(108):1–37, 2017.
- [39] Chengchun Shi, Wenbin Lu, and Rui Song. A massive data framework for M-estimators with cubic-rate. *Journal of the American Statistical Association*, 113(524):1698–1709, 2018.
- [40] W. Shi, Q. Ling, G. Wu, and W. Yin. EXTRA: An exact first-order algorithm for decentralized consensus optimization. *SIAM Journal on Optimization*, 25(2):944–966, November 2015.
- [41] W. Shi, Q. Ling, K. Yuan, G. Wu, and W. Yin. On the linear convergence of the admm in decentralized consensus optimization. *IEEE Transactions on Signal Processing*, 62:1750–1761, July 2014.
- [42] Wei Shi, Qing Ling, Gang Wu, and Wotao Yin. A proximal gradient algorithm for decentralized composite optimization. *IEEE Transactions on Signal Processing*, 63(22):6013–6023, 2015.
- [43] Y Sun, A Daneshmand, and G Scutari. Distributed optimization based on gradient-tracking revisited: Enhancing convergence rate via surrogation. *arXiv preprint arXiv:1905.02637*, 2019.
- [44] Stanislav Volgushev, Shih-Kang Chao, and Guang Cheng. Distributed inference for quantile regression processes. *The Annals of Statistics*, 47(3):1634–1662, 2019.
- [45] Jialei Wang, Mladen Kolar, Nathan Srebro, and Tong Zhang. Efficient distributed learning with sparsity. In *International Conference on Machine Learning*, pages 3636–3645. PMLR, 2017.
- [46] Auzinger Wien. *Iterative solution of large linear systems*. Lecture Notes, TU Wien, 2011.
- [47] J. Xu, S. Zhu, Y. C. Soh, and L. Xie. Convergence of asynchronous distributed gradient methods over stochastic networks. *IEEE Transactions on Automatic Control*, 63(2):434–448, July 2018.
- [48] K. Yuan, Q. Ling, and W. Yin. On the convergence of decentralized gradient descent. *SIAM Journal on Optimization*, 26(3):1835–1854, January 2016.
- [49] J. Zeng and W. Yin. On nonconvex decentralized gradient descent. *IEEE Transactions on Signal Processing*, 66(11):2834–2848, June 2018.
- [50] Yuchen Zhang, John Duchi, and Martin Wainwright. Divide and conquer kernel ridge regression. In *Conference on Learning Theory*, pages 592–617. PMLR, 2013.
- [51] Yuchen Zhang, John C Duchi, and Martin J Wainwright. Communication-efficient algorithms for statistical optimization. *The Journal of Machine Learning Research*, 14(1):3321–3363, 2013.
- [52] Tianqi Zhao, Guang Cheng, and Han Liu. A partially linear framework for massive heterogeneous data. *Annals of Statistics*, 44(4):1400, 2016.

Appendix

A Proof of Proposition 9

First note that the optimality of $\boldsymbol{\theta}_i^{t+\frac{1}{2}}$ implies:

$$\left(\boldsymbol{\theta} - \boldsymbol{\theta}_i^{t+\frac{1}{2}}\right)^\top (\mathbf{g}_i^t + \gamma \Delta \boldsymbol{\theta}_i^t) \geq 0, \quad \forall \|\boldsymbol{\theta}\|_1 \leq r, \quad (105)$$

where we recall $\Delta \boldsymbol{\theta}_i^t \triangleq \boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t$.

We use now the global RSC/RSS property of $\mathcal{L}$ (cf. Assumption 2) to compute the reduction of optimality gap $\mathcal{L}(\boldsymbol{\theta}) - \mathcal{L}(\hat{\boldsymbol{\theta}})$ by executing step (18).

$$\begin{aligned} & \mathcal{L}(\boldsymbol{\theta}_i^{t+\frac{1}{2}}) \\ &= \mathcal{L}(\boldsymbol{\theta}_i^t) + \nabla \mathcal{L}(\boldsymbol{\theta}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t) + \frac{1}{2} (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t)^\top \left(\frac{\mathbf{X}^\top \mathbf{X}}{N} \right) (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t) \\ &\stackrel{(RSS)}{\leq} \mathcal{L}(\boldsymbol{\theta}_i^t) + \nabla \mathcal{L}(\boldsymbol{\theta}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t) + \frac{L_\Sigma}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|^2 + \frac{\tau_g}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|_1^2 \\ &= \mathcal{L}(\hat{\boldsymbol{\theta}}) + \nabla \mathcal{L}(\boldsymbol{\theta}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}}) - \frac{1}{2} (\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}})^\top \left(\frac{\mathbf{X}^\top \mathbf{X}}{N} \right) (\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}) \\ &\quad + \frac{L_\Sigma}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|^2 + \frac{\tau_g}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|_1^2 \\ &\stackrel{(RSC)}{\leq} \mathcal{L}(\hat{\boldsymbol{\theta}}) + \nabla \mathcal{L}(\boldsymbol{\theta}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}}) - \frac{\mu_\Sigma}{2} \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|^2 + \frac{L_\Sigma}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|^2 \\ &\quad + \frac{\tau_\mu}{2} \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|_1^2 + \frac{\tau_g}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|_1^2. \end{aligned} \quad (106)$$

Letting $\boldsymbol{\theta} = \hat{\boldsymbol{\theta}}$ in (105) and adding to the above inequality we have

$$\begin{aligned} & \mathcal{L}(\boldsymbol{\theta}_i^{t+\frac{1}{2}}) \\ &\leq \mathcal{L}(\hat{\boldsymbol{\theta}}) - \gamma (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}})^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t) + (\nabla \mathcal{L}(\boldsymbol{\theta}_i^t) - \mathbf{g}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}}) \\ &\quad - \frac{\mu_\Sigma}{2} \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|^2 + \frac{L_\Sigma}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|^2 + \frac{\tau_\mu}{2} \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|_1^2 + \frac{\tau_g}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|_1^2 \\ &= \mathcal{L}(\hat{\boldsymbol{\theta}}) - \frac{\gamma}{2} \left(\|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}}\|^2 - \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|^2 + \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|^2 \right) + (\nabla \mathcal{L}(\boldsymbol{\theta}_i^t) - \mathbf{g}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}}) \\ &\quad - \frac{\mu_\Sigma}{2} \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|^2 + \frac{L_\Sigma}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|^2 + \frac{\tau_\mu}{2} \|\boldsymbol{\theta}_i^t - \hat{\boldsymbol{\theta}}\|_1^2 + \frac{\tau_g}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|_1^2. \end{aligned}$$

Now applying the triangle inequality to the last term and Lemma 1 yields:

$$\begin{aligned}
& \mathcal{L}(\boldsymbol{\theta}_i^{t+\frac{1}{2}}) \\
& \leq \mathcal{L}(\widehat{\boldsymbol{\theta}}) - \frac{\gamma}{2} \left(\|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 - \|\boldsymbol{\theta}_i^t - \widehat{\boldsymbol{\theta}}\|^2 + \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|^2 \right) + (\nabla \mathcal{L}(\boldsymbol{\theta}_i^t) - \mathbf{g}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}) \\
& \quad - \frac{\mu_\Sigma}{2} \|\boldsymbol{\theta}_i^t - \widehat{\boldsymbol{\theta}}\|^2 + \frac{L_\Sigma}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|^2 + \frac{\tau_\mu}{2} (8s \|\boldsymbol{\theta}_i^t - \widehat{\boldsymbol{\theta}}\|^2 + 2\nu^2) \\
& \quad + \frac{\tau_g}{2} (16s \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 16s \|\boldsymbol{\theta}_i^t - \widehat{\boldsymbol{\theta}}\|^2 + 8\nu^2) \\
& \leq \mathcal{L}(\widehat{\boldsymbol{\theta}}) - \frac{\gamma}{2} \left(\|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 - \|\boldsymbol{\theta}_i^t - \widehat{\boldsymbol{\theta}}\|^2 + \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|^2 \right) + (\nabla \mathcal{L}(\boldsymbol{\theta}_i^t) - \mathbf{g}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}) \\
& \quad - \frac{\mu_\Sigma}{2} \|\boldsymbol{\theta}_i^t - \widehat{\boldsymbol{\theta}}\|^2 + \frac{L_\Sigma}{2} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|^2 + 8s\tau_g \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + (4s\tau_\mu + 8s\tau_g) \|\boldsymbol{\theta}_i^t - \widehat{\boldsymbol{\theta}}\|^2 \\
& \quad + (\tau_\mu + 4\tau_g)\nu^2.
\end{aligned}$$

Rearranging terms and using the fact that $\mathcal{L}(\boldsymbol{\theta}_i^{t+\frac{1}{2}}) \geq \mathcal{L}(\widehat{\boldsymbol{\theta}})$, yields

$$\begin{aligned}
\left(1 - \frac{C_1 s \tau_g}{\gamma}\right) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 & \leq \left(1 - \frac{\mu_\Sigma}{\gamma} + \frac{C_1 s (\tau_\mu + \tau_g)}{\gamma}\right) \|\boldsymbol{\theta}_i^t - \widehat{\boldsymbol{\theta}}\|^2 \\
& \quad - \left(1 - \frac{L_\Sigma}{\gamma}\right) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \boldsymbol{\theta}_i^t\|^2 + \frac{2}{\gamma} \cdot (\nabla \mathcal{L}(\boldsymbol{\theta}_i^t) - \mathbf{g}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}) + \frac{C_1 (\tau_\mu + \tau_g)}{\gamma} \nu^2. \quad (107)
\end{aligned}$$

Summing over $i = 1, \dots, m$, completes the proof. $\square$

B Proof of Proposition 10

The difference $\frac{1}{m} \sum_{j=1}^m \nabla \mathcal{L}_j(\boldsymbol{\theta}_i^t) - \frac{1}{m} \sum_{j=1}^m \nabla \mathcal{L}_j(\boldsymbol{\theta}_j^t)$ can be expressed as

$$\begin{aligned}
& \frac{1}{m} \sum_{j=1}^m \nabla \mathcal{L}_j(\boldsymbol{\theta}_i^t) - \frac{1}{m} \sum_{j=1}^m \nabla \mathcal{L}_j(\boldsymbol{\theta}_j^t) \\
& = \frac{1}{m} \sum_{j=1}^m \frac{1}{n} \mathbf{X}_j^\top \mathbf{X}_j (\boldsymbol{\theta}_i^t - \boldsymbol{\theta}_j^t) \\
& = \frac{1}{N} \mathbf{X}^\top \mathbf{X} (\boldsymbol{\theta}_i^t - \bar{\boldsymbol{\theta}}^t) + \frac{1}{m} \sum_{j=1}^m \frac{1}{n} \mathbf{X}_j^\top \mathbf{X}_j (\bar{\boldsymbol{\theta}}^t - \boldsymbol{\theta}_j^t).
\end{aligned} \quad (108)$$

Substituting into the expression of δ^t we can split it into three error terms:

$$\begin{aligned}
\delta^t & = \underbrace{\sum_{i=1}^m (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}})^\top \left(\frac{1}{N} \mathbf{X}^\top \mathbf{X} \right) (\boldsymbol{\theta}_i^t - \bar{\boldsymbol{\theta}}^t)}_{T_1} + \underbrace{\frac{1}{m} \sum_{i=1}^m \sum_{j=1}^m (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}})^\top \left(\frac{1}{n} \mathbf{X}_j^\top \mathbf{X}_j \right) (\bar{\boldsymbol{\theta}}^t - \boldsymbol{\theta}_j^t)}_{T_2} \\
& \quad + \underbrace{\sum_{i=1}^m (\bar{\mathbf{g}}^t - \mathbf{g}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}})}_{T_3}.
\end{aligned} \quad (109)$$

We are now ready to bound δ^t . For this, we substitute the expressions of $\boldsymbol{\Theta}_\perp^t$ and $\mathbf{G}_\perp^t$ [cf. (32a)-(32b)] into (109). We then use the RSS (cf. Assumption 2) to upper bound δ^t by quantities that depend exclusively on the sequences $\|\boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|$ and $\|\Delta \boldsymbol{\Theta}^t\|$.

1) **Bounding T_1 :** Iterating (32a) telescopically to $t = 0$, yields

$$\boldsymbol{\Theta}_\perp^t = (\mathbf{W} - \mathbf{J})\boldsymbol{\Theta}_\perp^0 + \sum_{s=0}^{t-1} (\mathbf{W} - \mathbf{J})^{t-s} \Delta \boldsymbol{\Theta}^s, \quad t = 1, 2, \dots, \quad (110)$$

This shows that $\boldsymbol{\Theta}_\perp^t$ lies in the span of $\{\boldsymbol{\Theta}_\perp^0, \Delta \boldsymbol{\Theta}^0, \dots, \Delta \boldsymbol{\Theta}^{t-1}\}$. Therefore, $\boldsymbol{\theta}_i^t - \bar{\boldsymbol{\theta}}^t$ can be expressed as

$$\boldsymbol{\theta}_i^t - \bar{\boldsymbol{\theta}}^t = \sum_{j=1}^m \ell_{ij}^{(t)} (\boldsymbol{\theta}_{j,\perp}^0) + \sum_{j=1}^m \sum_{s=0}^{t-1} \ell_{ij}^{(t-s)} \Delta \boldsymbol{\theta}_j^s, \quad t = 1, 2, \dots, \quad (111)$$

where $\ell_{ij}^{(t)}$ denotes the ij -th element of the matrix $(\mathbf{W} - \mathbf{J})^t$.

Substituting (111) in the expression of T_1 , we can bound it as

$$\begin{aligned} T_1 &= \sum_{i=1}^m (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}})^\top \left(\frac{1}{N} \mathbf{X}^\top \mathbf{X} \right) (\boldsymbol{\theta}_i^t - \bar{\boldsymbol{\theta}}^t) \\ &= \sum_{i=1}^m (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}})^\top \left(\frac{1}{N} \mathbf{X}^\top \mathbf{X} \right) \left(\sum_{j=1}^m \ell_{ij}^{(t)} \boldsymbol{\theta}_{j,\perp}^0 + \sum_{j=1}^m \sum_{s=0}^{t-1} \ell_{ij}^{(t-s)} \Delta \boldsymbol{\theta}_j^s \right) \\ &\leq \sum_{i=1}^m \sum_{j=1}^m |\ell_{ij}^{(t)}| \left\| \frac{1}{\sqrt{N}} \mathbf{X} (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}}) \right\| \left\| \frac{1}{\sqrt{N}} \mathbf{X} \boldsymbol{\theta}_{j,\perp}^0 \right\| \\ &\quad + \sum_{i=1}^m \sum_{j=1}^m \sum_{s=0}^{t-1} |\ell_{ij}^{(t-s)}| \left\| \frac{1}{\sqrt{N}} \mathbf{X} (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}}) \right\| \left\| \frac{1}{\sqrt{N}} \mathbf{X} \Delta \boldsymbol{\theta}_j^s \right\| \\ &\leq \sum_{i=1}^m \sum_{j=1}^m |\ell_{ij}^{(t)}| \left(\frac{\epsilon}{2} \left\| \frac{1}{\sqrt{N}} \mathbf{X} (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}}) \right\|^2 + \frac{1}{2\epsilon} \left\| \frac{1}{\sqrt{N}} \mathbf{X} \boldsymbol{\theta}_{j,\perp}^0 \right\|^2 \right) \\ &\quad + \sum_{i=1}^m \sum_{j=1}^m \sum_{s=0}^{t-1} |\ell_{ij}^{(t-s)}| \left(\frac{\epsilon}{2} \left\| \frac{1}{\sqrt{N}} \mathbf{X} (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}}) \right\|^2 + \frac{1}{2\epsilon} \left\| \frac{1}{\sqrt{N}} \mathbf{X} \Delta \boldsymbol{\theta}_j^s \right\|^2 \right), \end{aligned} \quad (112)$$

where we have applied the Cauchy-Schwartz and Young's inequality with $\epsilon > 0$. Invoking the RSS (Assumption 2) along with Assumption 7 and Lemma 1, we can bound the RHS of (112) as (cf. Appendix C.1): for all $t \geq 1$,

$$\begin{aligned} T_1 &\leq \frac{\rho \cdot c_m \epsilon}{1 - \rho} (L_\Sigma + 8s\tau_g) \left\| \boldsymbol{\Theta}^{t+\frac{1}{2}} - \hat{\boldsymbol{\Theta}} \right\|_F^2 + \frac{c_m L_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta \boldsymbol{\Theta}^s\|_F^2 \\ &\quad + c_m \cdot 8s\tau_g \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^{s+\frac{1}{2}} - \hat{\boldsymbol{\Theta}}\|_F^2 + c_m \cdot 8s\tau_g \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^s - \hat{\boldsymbol{\Theta}}\|_F^2 \\ &\quad + \underbrace{\frac{\rho \cdot mc_m}{1 - \rho} \cdot 4\tau_g \nu^2 (\epsilon + \epsilon^{-1}) + \frac{1}{2\epsilon} c_m \rho^t \left(L_\Sigma \|\boldsymbol{\Theta}_\perp^0\|_F^2 + \tau_g \sum_{j=1}^m \|\boldsymbol{\theta}_j^0 - \bar{\boldsymbol{\theta}}^0\|_1^2 \right)}_{=0 \text{ by initialization}}. \end{aligned} \quad (113)$$

2) **Bounding T_2 :** Recall that

$$T_2 = \frac{1}{m} \sum_{i=1}^m \sum_{j=1}^m \underbrace{(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \hat{\boldsymbol{\theta}})^\top \left(\frac{1}{n} \mathbf{X}_j^\top \mathbf{X}_j \right) (\bar{\boldsymbol{\theta}}^t - \boldsymbol{\theta}_j^t)}_{\triangleq T_2^{ij}}. \quad (114)$$

Focusing on the ij -th summand and using (111) we can write

$$T_2^{ij} = (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}})^\top \left(\frac{1}{n} \mathbf{X}_j^\top \mathbf{X}_j \right) \left(\sum_{k=1}^m (-\ell_{jk}^{(t)}) \boldsymbol{\theta}_{k,\perp}^0 + \sum_{k=1}^m \sum_{s=0}^{t-1} (-\ell_{jk}^{(t-s)}) \Delta \boldsymbol{\theta}_k^s \right). \quad (115)$$

Note that T_2^{ij} has the same form as T_1 and thus following similar steps as in (113) while using the local RSS (cf. Assumption 4), we can bound T_2 as (cf. Appendix C.2): for all $t \geq 1$,

$$\begin{aligned} T_2 &\leq \frac{\rho \cdot c_m \epsilon}{1 - \rho} (\ell_\Sigma + 8s\tau_\ell) \left\| \boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}} \right\|_F^2 + \frac{c_m \ell_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta \boldsymbol{\Theta}^s\|_F^2 \\ &\quad + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^{s+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^s - \widehat{\boldsymbol{\Theta}}\|_F^2 \\ &\quad + \underbrace{\frac{\rho \cdot mc_m}{1 - \rho} 4\tau_\ell \nu^2 \cdot (\epsilon + \epsilon^{-1}) + \frac{1}{2\epsilon} c_m \rho^t \left(\ell_\Sigma \|\boldsymbol{\Theta}_\perp^0\|_F^2 + \tau_\ell \sum_{k=1}^m \|\boldsymbol{\theta}_k^0 - \bar{\boldsymbol{\theta}}^0\|_1^2 \right)}_{=0}. \end{aligned} \quad (116)$$

3) Bounding T_3 : Using (32b) we can express $(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}})^\top (\bar{\mathbf{g}}^t - \mathbf{g}_i^t)$ in T_3 as

$$\begin{aligned} &(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}})^\top (\bar{\mathbf{g}}^t - \mathbf{g}_i^t) \\ &= (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}})^\top \left(\sum_{j=1}^m (-\ell_{ij}^{(t)}) \mathbf{g}_{j,\perp}^0 + \sum_{s=0}^{t-1} \sum_{j=1}^m (-\ell_{ij}^{(t-s)}) \left(\frac{1}{n} \mathbf{X}_j^\top \mathbf{X}_j \right) (\boldsymbol{\theta}_j^{s+1} - \boldsymbol{\theta}_j^s) \right) \\ &\leq \sum_{j=1}^m |\ell_{ij}^{(t)}| \underbrace{\left| \left(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}} \right)^\top \mathbf{g}_{j,\perp}^0 \right|}_{T_{3,1}} + \sum_{s=0}^{t-1} \sum_{j=1}^m |\ell_{ij}^{(t-s)}| \underbrace{\left| \left(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}} \right)^\top \left(\frac{1}{n} \mathbf{X}_j^\top \mathbf{X}_j \right) (\boldsymbol{\theta}_j^{s+1} - \boldsymbol{\theta}_j^s) \right|}_{T_{3,2}}. \end{aligned} \quad (117)$$

Following again similar steps as in the bounds of T_1 and T_2 , we can obtain the following bound for T_3 (cf. Appendix C.3):

$$\begin{aligned} T_3 &\leq \frac{3}{2} \cdot \frac{\rho \cdot c_m \epsilon}{1 - \rho} (2\ell_\Sigma + 8s\tau_\ell) \left\| \boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}} \right\|_F^2 + \frac{c_m \ell_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^{s+1} - \boldsymbol{\Theta}^s\|_F^2 \\ &\quad + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^{s+1} - \widehat{\boldsymbol{\Theta}}\|_F^2 + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^s - \widehat{\boldsymbol{\Theta}}\|_F^2 \\ &\quad + mc_m \rho^t (A_3 \epsilon^{-1} + sc_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu) + \frac{\rho \cdot mc_m}{1 - \rho} 4\tau_\ell \nu^2 (\epsilon + \epsilon^{-1}), \end{aligned} \quad (118)$$

where

$$\begin{aligned} A_3 &\triangleq \frac{1}{2} (\ell_\Sigma \|\boldsymbol{\theta}^*\|^2 + \tau_\ell \|\boldsymbol{\theta}^*\|_1^2 + \frac{1}{m} \ell_\Sigma \|\boldsymbol{\Theta}^*\|_F^2 + \tau_\ell \|\boldsymbol{\theta}^*\|_1^2), \\ c_g &\triangleq \|\nabla \mathcal{L}_j(\boldsymbol{\theta}^*)\|_\infty + \|\nabla \mathcal{L}(\boldsymbol{\theta}^*)\|_\infty. \end{aligned} \quad (119)$$

4) Combining the bounds of T_1 , T_2 , and T_3 : We can finally bound δ^t as (see Appendix C.4 for more details):

$$\begin{aligned} \delta^t &= \sum_{i=1}^m (\nabla \mathcal{L}(\boldsymbol{\theta}_i^t) - \mathbf{y}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}) \\ &\leq T_1 + T_2 + T_3 \end{aligned}$$

$$\begin{aligned}
&\stackrel{(a)}{\leq} \frac{7}{2} \cdot \frac{\rho \cdot c_m \epsilon}{1 - \rho} (2\ell_\Sigma + 8s\tau_\ell) \|\Theta^{t+\frac{1}{2}} - \widehat{\Theta}\|_F^2 \\
&\quad + \frac{c_m \ell_\Sigma}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta \Theta^s\|_F^2 + \frac{c_m \ell_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+1} - \Theta^s\|_F^2 \\
&\quad + c_m 24\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+\frac{1}{2}} - \widehat{\Theta}\|_F^2 + c_m 24\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^s - \widehat{\Theta}\|_F^2 \\
&\quad + mc_m \rho^t (A_3 \cdot \epsilon^{-1} + sc_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu) + \frac{\rho \cdot mc_m}{1 - \rho} 12\tau_\ell \nu^2 (\epsilon + \epsilon^{-1}) \\
&\stackrel{(b)}{\leq} \frac{7}{2} \cdot \frac{\rho \cdot c_m \epsilon}{1 - \rho} (2\ell_\Sigma + 8s\tau_\ell) \|\Theta^{t+\frac{1}{2}} - \widehat{\Theta}\|_F^2 \\
&\quad + \frac{2c_m \ell_\Sigma}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta \Theta^s\|_F^2 + \frac{2c_m \ell_\Sigma}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta_\perp^s\|_F^2 \\
&\quad + c_m 24s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+\frac{1}{2}} - \widehat{\Theta}\|_F^2 + c_m 24s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s-\frac{1}{2}} - \widehat{\Theta}\|_F^2 \\
&\quad + mc_m \rho^t (A_3 \cdot \epsilon^{-1} + sc_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu) + \frac{\rho \cdot mc_m}{1 - \rho} 12\tau_\ell \nu^2 (\epsilon + \epsilon^{-1}),
\end{aligned}$$

where in (a) we used (113), (116), and (118); $L_\Sigma \leq \ell_\Sigma$ and $\tau_g \leq \tau_\ell$; and

$$\|\Theta^{t+1} - \widehat{\Theta}\|_F^2 = \|\mathbf{W}(\Theta^{t+\frac{1}{2}} - \widehat{\Theta})\|_F^2 \leq \|\Theta^{t+\frac{1}{2}} - \widehat{\Theta}\|_F^2; \quad (120)$$

while in (b) we used again (120) and

$$\|\Theta^{s+1} - \Theta^s\|_F^2 \stackrel{(28a)}{=} \|\mathbf{W}(\Theta^s + \Delta \Theta^s) - \Theta^s\|_F^2 \leq 4\|\Theta_\perp^s\|_F^2 + 2\|\Delta \Theta^s\|_F^2. \quad (121)$$

□

C Supplement to Appendix B

We bound next the tracking errors T_1 , T_2 , and T_3 in (109).

C.1 Bounding T_1 in (109)

Invoking the RSC property (Assumption 2) and Lemma 1, we have

$$\begin{aligned}
\left\| \frac{1}{\sqrt{N}} \mathbf{X}(\theta_i^{t+\frac{1}{2}} - \widehat{\theta}) \right\|^2 &\leq L_\Sigma \left\| \theta_i^{t+\frac{1}{2}} - \widehat{\theta} \right\|^2 + \tau_g \left\| \theta_i^{t+\frac{1}{2}} - \widehat{\theta} \right\|_1^2 \\
&\leq (L_\Sigma + 8s\tau_g) \left\| \theta_i^{t+\frac{1}{2}} - \widehat{\theta} \right\|^2 + 2\tau_g \nu^2.
\end{aligned}$$

Similarly, the other two terms in (112) can be bounded respectively as

$$\left\| \frac{1}{\sqrt{N}} \mathbf{X} \theta_{j,\perp}^0 \right\|^2 \leq L_\Sigma \|\theta_j^0 - \bar{\theta}^0\|^2 + \tau_g \|\theta_j^0 - \bar{\theta}^0\|_1^2$$

and

$$\left\| \frac{1}{\sqrt{N}} \mathbf{X} \Delta \theta_j^s \right\|^2 \leq L_\Sigma \|\Delta \theta_j^s\|^2 + \tau_g \|\theta_j^{s+\frac{1}{2}} - \theta_j^s\|_1^2$$

$$\begin{aligned}
&\leq L_\Sigma \|\Delta \boldsymbol{\theta}_j^s\|^2 + 2\tau_g \|\boldsymbol{\theta}_j^{s+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|_1^2 + 2\tau_g \|\boldsymbol{\theta}_j^s - \widehat{\boldsymbol{\theta}}\|_1^2 \\
&\leq L_\Sigma \|\Delta \boldsymbol{\theta}_j^s\|^2 + 16s\tau_g \|\boldsymbol{\theta}_j^{s+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 16s\tau_g \|\boldsymbol{\theta}_j^s - \widehat{\boldsymbol{\theta}}\|^2 + 8\tau_g \nu^2.
\end{aligned}$$

Summing up the terms and using $\|(\mathbf{W} - \mathbf{J})^t\|_\infty \leq c_m \rho^t$ (due to Assumption 7), we arrive to the following bound for T_1 :

$$\begin{aligned}
T_1 &\leq \sum_{i=1}^m \sum_{j=1}^m |\ell_{ij}^{(t)}| \frac{\epsilon}{2} \left((L_\Sigma + 8s\tau_g) \left\| \boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}} \right\|^2 + 2\tau_g \nu^2 \right) \\
&\quad + \sum_{i=1}^m \sum_{j=1}^m |\ell_{ij}^{(t)}| \frac{1}{2\epsilon} \left(L_\Sigma \|\boldsymbol{\theta}_j^0 - \bar{\boldsymbol{\theta}}^0\|^2 + \tau_g \|\boldsymbol{\theta}_j^0 - \bar{\boldsymbol{\theta}}^0\|_1^2 \right) \\
&\quad + \sum_{i=1}^m \sum_{j=1}^m \sum_{s=0}^{t-1} |\ell_{ij}^{(t-s)}| \frac{\epsilon}{2} \left((L_\Sigma + 8s\tau_g) \left\| \boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}} \right\|^2 + 2\tau_g \nu^2 \right) \\
&\quad + \sum_{i=1}^m \sum_{j=1}^m \sum_{s=0}^{t-1} |\ell_{ij}^{(t-s)}| \frac{1}{2\epsilon} \left(L_\Sigma \|\Delta \boldsymbol{\theta}_j^s\|^2 + 16s\tau_g \|\boldsymbol{\theta}_j^{s+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 16s\tau_g \|\boldsymbol{\theta}_j^s - \widehat{\boldsymbol{\theta}}\|^2 + 8\tau_g \nu^2 \right) \\
&\leq \frac{\epsilon}{2} c_m \rho^t (L_\Sigma + 8s\tau_g) \left\| \boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}} \right\|_F^2 + \frac{\epsilon}{2} \cdot m c_m \rho^t \cdot 2\tau_g \nu^2 \\
&\quad + \frac{1}{2\epsilon} c_m \rho^t \left(L_\Sigma \|\boldsymbol{\Theta}_\perp^0\|_F^2 + \tau_g \sum_{j=1}^m \|\boldsymbol{\theta}_j^0 - \bar{\boldsymbol{\theta}}^0\|_1^2 \right) \\
&\quad + \sum_{s=0}^{t-1} c_m \rho^{t-s} \frac{\epsilon}{2} (L_\Sigma + 8s\tau_g) \left\| \boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}} \right\|_F^2 + \sum_{s=0}^{t-1} m c_m \rho^{t-s} \cdot \frac{\epsilon}{2} \cdot 2\tau_g \nu^2 \\
&\quad + \sum_{s=0}^{t-1} c_m \rho^{t-s} \frac{1}{2\epsilon} L_\Sigma \|\Delta \boldsymbol{\Theta}^s\|_F^2 + \sum_{s=0}^{t-1} c_m \rho^{t-s} \cdot \frac{1}{2\epsilon} \cdot 16s\tau_g \|\boldsymbol{\Theta}^{s+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 \\
&\quad + \sum_{s=0}^{t-1} c_m \rho^{t-s} \frac{1}{2\epsilon} \cdot 16s\tau_g \|\boldsymbol{\Theta}^s - \widehat{\boldsymbol{\Theta}}\|_F^2 + \sum_{s=0}^{t-1} m c_m \rho^{t-s} \frac{1}{2\epsilon} 8\tau_g \nu^2 \\
&\leq \frac{\rho \cdot c_m \epsilon}{1 - \rho} (L_\Sigma + 8s\tau_g) \left\| \boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}} \right\|_F^2 + \frac{c_m L_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta \boldsymbol{\Theta}^s\|_F^2 \\
&\quad + c_m \cdot 8s\tau_g \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^{s+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 + c_m \cdot 8s\tau_g \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^s - \widehat{\boldsymbol{\Theta}}\|_F^2 \\
&\quad + \frac{\rho \cdot m c_m}{1 - \rho} \cdot 4\tau_g \nu^2 (\epsilon + \epsilon^{-1}) + \frac{1}{2\epsilon} c_m \rho^t \left(L_\Sigma \|\boldsymbol{\Theta}_\perp^0\|_F^2 + \tau_g \sum_{j=1}^m \|\boldsymbol{\theta}_j^0 - \bar{\boldsymbol{\theta}}^0\|_1^2 \right). \tag{122}
\end{aligned}$$

C.2 Bounding T_2 in (109)

Using the local RSS (cf. Assumption 4) and Lemma 1, we can bound the summands in T_2^{ij} [cf. (115)] as

$$\begin{aligned}
&(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}})^\top \left(\frac{1}{n} \mathbf{X}_j^\top \mathbf{X}_j \right) \ell_{jk}^{(t)} \boldsymbol{\theta}_{k,\perp}^0 \\
&\leq |\ell_{jk}^{(t)}| \left\| \frac{1}{\sqrt{n}} \mathbf{X}_j (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}) \right\| \cdot \left\| \frac{1}{\sqrt{n}} \mathbf{X}_j (\boldsymbol{\theta}_k^0 - \bar{\boldsymbol{\theta}}^0) \right\| \\
&\leq |\ell_{jk}^{(t)}| \frac{\epsilon}{2} \left((L_\Sigma + 8s\tau_\ell) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 2\tau_\ell \nu^2 \right) + |\ell_{jk}^{(t)}| \frac{1}{2\epsilon} \left(L_\Sigma \|\boldsymbol{\theta}_k^0 - \bar{\boldsymbol{\theta}}^0\|^2 + \tau_\ell \|\boldsymbol{\theta}_k^0 - \bar{\boldsymbol{\theta}}^0\|_1^2 \right)
\end{aligned}$$

and

$$\begin{aligned}
& (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}})^\top \left(\frac{1}{n} \mathbf{X}_j^\top \mathbf{X}_j \right) \ell_{jk}^{(t-s)} \Delta \boldsymbol{\theta}_k^s \\
& \leq |\ell_{jk}^{(t-s)}| \left\| \frac{1}{\sqrt{n}} \mathbf{X}_j (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}) \right\| \left\| \frac{1}{\sqrt{n}} \mathbf{X}_j \Delta \boldsymbol{\theta}_k^s \right\| \\
& \leq |\ell_{jk}^{(t-s)}| \frac{\epsilon}{2} \left((\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 2\tau_\ell \nu^2 \right) \\
& \quad + |\ell_{jk}^{(t-s)}| \frac{1}{2\epsilon} \left(\ell_\Sigma \|\Delta \boldsymbol{\theta}_k^s\|^2 + 16s\tau_\ell \|\boldsymbol{\theta}_k^{s+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 16s\tau_\ell \|\boldsymbol{\theta}_k^s - \widehat{\boldsymbol{\theta}}\|^2 + 8\tau_\ell \nu^2 \right),
\end{aligned}$$

respectively.

Summing over i, j, k and using $\|(\mathbf{W} - \mathbf{J})^t\|_\infty \leq c_m \rho^t$, we can bound T_2 as:

$$\begin{aligned}
& m \cdot T_2 \\
& \leq \sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^m |\ell_{jk}^{(t)}| \frac{\epsilon}{2} \left((\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 2\tau_\ell \nu^2 \right) \\
& \quad + \sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^m |\ell_{jk}^{(t)}| \frac{1}{2\epsilon} \left(\ell_\Sigma \|\boldsymbol{\theta}_k^0 - \bar{\boldsymbol{\theta}}^0\|^2 + \tau_\ell \|\boldsymbol{\theta}_k^0 - \bar{\boldsymbol{\theta}}^0\|_1^2 \right) \\
& \quad + \sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^m \sum_{s=0}^{t-1} |\ell_{jk}^{(t-s)}| \frac{\epsilon}{2} \left((\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 2\tau_\ell \nu^2 \right) \\
& \quad + \sum_{i=1}^m \sum_{j=1}^m \sum_{k=1}^m \sum_{s=0}^{t-1} |\ell_{jk}^{(t-s)}| \frac{1}{2\epsilon} \left(\ell_\Sigma \|\Delta \boldsymbol{\theta}_k^s\|^2 + 16s\tau_\ell \|\boldsymbol{\theta}_k^{s+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 16s\tau_\ell \|\boldsymbol{\theta}_k^s - \widehat{\boldsymbol{\theta}}\|^2 + 8\tau_\ell \nu^2 \right) \\
& \leq mc_m \rho^t \cdot \frac{\epsilon}{2} (\ell_\Sigma + 8s\tau_\ell) \left\| \boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}} \right\|_F^2 + m^2 c_m \rho^t \cdot \tau_\ell \nu^2 \cdot \epsilon \\
& \quad + mc_m \rho^t \cdot \frac{1}{2\epsilon} \ell_\Sigma \|\boldsymbol{\Theta}_\perp^0\|_F^2 + mc_m \rho^t \cdot \frac{1}{2\epsilon} \tau_\ell \sum_{k=1}^m \|\boldsymbol{\theta}_k^0 - \bar{\boldsymbol{\theta}}^0\|_1^2 \\
& \quad + \sum_{s=0}^{t-1} mc_m \rho^{t-s} \frac{\epsilon}{2} (\ell_\Sigma + 8s\tau_\ell) \left\| \boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}} \right\|_F^2 + \sum_{s=0}^{t-1} m^2 c_m \rho^{t-s} \tau_\ell \nu^2 \cdot \epsilon \\
& \quad + \sum_{s=0}^{t-1} mc_m \rho^{t-s} \frac{1}{2\epsilon} \ell_\Sigma \|\Delta \boldsymbol{\Theta}^s\|_F^2 + \sum_{s=0}^{t-1} mc_m \rho^{t-s} \frac{1}{\epsilon} 8s\tau_\ell \|\boldsymbol{\Theta}^{s+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 \\
& \quad + \sum_{s=0}^{t-1} mc_m \rho^{t-s} \frac{1}{\epsilon} 8s\tau_\ell \|\boldsymbol{\Theta}^s - \widehat{\boldsymbol{\Theta}}\|_F^2 + \sum_{s=0}^{t-1} m^2 c_m \rho^{t-s} \frac{1}{\epsilon} 4\tau_\ell \nu^2 \\
& \leq \frac{\rho \cdot mc_m \epsilon}{1 - \rho} (\ell_\Sigma + 8s\tau_\ell) \left\| \boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}} \right\|_F^2 + \frac{mc_m \ell_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta \boldsymbol{\Theta}^s\|_F^2 \\
& \quad + mc_m \frac{1}{\epsilon} 8s\tau_\ell \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^{s+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 + mc_m \frac{1}{\epsilon} 8s\tau_\ell \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^s - \widehat{\boldsymbol{\Theta}}\|_F^2 \\
& \quad + \frac{\rho \cdot m^2 c_m}{1 - \rho} 4\tau_\ell \nu^2 \cdot (\epsilon + \epsilon^{-1}) + \frac{1}{2\epsilon} mc_m \rho^t \left(\ell_\Sigma \|\boldsymbol{\Theta}_\perp^0\|_F^2 + \tau_\ell \sum_{k=1}^m \|\boldsymbol{\theta}_k^0 - \bar{\boldsymbol{\theta}}^0\|_1^2 \right). \tag{123}
\end{aligned}$$

C.3 Bounding T_3 in (109)

We bound first $T_{3,1}$ and $T_{3,2}$.

Invoking the definition of $\mathbf{g}_{j,\perp}^0$ we have

$$\begin{aligned}
T_{3,1} &= \left| \left(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}} \right)^\top \left(\nabla \mathcal{L}_j(\boldsymbol{\theta}_j^0) - \frac{1}{m} \sum_{k=1}^m \nabla \mathcal{L}_k(\boldsymbol{\theta}_k^0) \right) \right| \\
&= \left| \left(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}} \right)^\top \left(\frac{1}{n} \mathbf{X}_j^\top (\mathbf{X}_j \boldsymbol{\theta}_j^0 - \mathbf{y}_j) - \frac{1}{mn} \sum_{k=1}^m \mathbf{X}_k^\top (\mathbf{X}_k \boldsymbol{\theta}_k^0 - \mathbf{y}_k) \right) \right| \\
&= \left| \left(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}} \right)^\top \left(\frac{1}{n} \mathbf{X}_j^\top (\mathbf{X}_j \boldsymbol{\theta}_j^0 - \mathbf{X}_j \boldsymbol{\theta}^* + \mathbf{X}_j \boldsymbol{\theta}^* - \mathbf{y}_j) - \frac{1}{mn} \sum_{k=1}^m \mathbf{X}_k^\top (\mathbf{X}_k \boldsymbol{\theta}_k^0 - \mathbf{X} \boldsymbol{\theta}^* + \mathbf{X} \boldsymbol{\theta}^* - \mathbf{y}_k) \right) \right| \\
&\leq \left| \left(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}} \right)^\top \left(\frac{1}{n} \mathbf{X}_j^\top \mathbf{X}_j (\boldsymbol{\theta}_j^0 - \boldsymbol{\theta}^*) - \frac{1}{mn} \sum_{k=1}^m \mathbf{X}_k^\top \mathbf{X}_k (\boldsymbol{\theta}_k^0 - \boldsymbol{\theta}^*) \right) \right| \\
&\quad + \left| \left(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}} \right)^\top \nabla \mathcal{L}_j(\boldsymbol{\theta}^*) \right| + \left| \left(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}} \right)^\top \nabla \mathcal{L}(\boldsymbol{\theta}^*) \right|.
\end{aligned}$$

Therefore,

$$\begin{aligned}
T_{3,1} &\leq \frac{\epsilon}{2} \left(\ell_\Sigma \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + \tau_\ell \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|_1^2 \right) + \frac{1}{2\epsilon} \left(\ell_\Sigma \|\boldsymbol{\theta}_j^0 - \boldsymbol{\theta}^*\|^2 + \tau_\ell \|\boldsymbol{\theta}_j^0 - \boldsymbol{\theta}^*\|_1^2 \right) \\
&\quad + \frac{1}{m} \sum_{k=1}^m \left(\frac{\epsilon}{2} \left(\ell_\Sigma \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + \tau_\ell \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|_1^2 \right) + \frac{1}{2\epsilon} \left(\ell_\Sigma \|\boldsymbol{\theta}_k^0 - \boldsymbol{\theta}^*\|^2 + \tau_\ell \|\boldsymbol{\theta}_k^0 - \boldsymbol{\theta}^*\|_1^2 \right) \right) \\
&\quad + \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|_1 \underbrace{(\|\nabla \mathcal{L}_j(\boldsymbol{\theta}^*)\|_\infty + \|\nabla \mathcal{L}(\boldsymbol{\theta}^*)\|_\infty)}_{c_g} \\
&\stackrel{(a)}{\leq} \frac{\epsilon}{2} \left((\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 2\tau_\ell \nu^2 \right) + \frac{1}{2\epsilon} \left(\ell_\Sigma \|\boldsymbol{\theta}_j^0 - \boldsymbol{\theta}^*\|^2 + \tau_\ell \|\boldsymbol{\theta}_j^0 - \boldsymbol{\theta}^*\|_1^2 \right) \\
&\quad + \frac{1}{m} \sum_{k=1}^m \left(\frac{\epsilon}{2} \left((\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 2\tau_\ell \nu^2 \right) \right) + \frac{1}{2\epsilon} \frac{1}{m} \left(\ell_\Sigma \|\boldsymbol{\Theta}^0 - \boldsymbol{\Theta}^*\|_F^2 + \tau_\ell \sum_{k=1}^m \|\boldsymbol{\theta}_k^0 - \boldsymbol{\theta}^*\|_1^2 \right) \\
&\quad + c_g \left(2\sqrt{s} \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\| + \nu \right) \\
&\stackrel{(b)}{\leq} \epsilon \left((\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 2\tau_\ell \nu^2 \right) \\
&\quad + \frac{1}{2\epsilon} \left(\ell_\Sigma \|\boldsymbol{\theta}_j^0 - \boldsymbol{\theta}^*\|^2 + \tau_\ell \|\boldsymbol{\theta}_j^0 - \boldsymbol{\theta}^*\|_1^2 + \frac{1}{m} \ell_\Sigma \|\boldsymbol{\Theta}^0 - \boldsymbol{\Theta}^*\|_F^2 + \tau_\ell \frac{1}{m} \sum_{k=1}^m \|\boldsymbol{\theta}_k^0 - \boldsymbol{\theta}^*\|_1^2 \right) \\
&\quad + \epsilon \ell_\Sigma \cdot \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + sc_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu \\
&\leq \epsilon \left((2\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + 2\tau_\ell \nu^2 \right) \\
&\quad + \frac{1}{\epsilon} \frac{1}{2} \underbrace{\left(\ell_\Sigma \|\boldsymbol{\theta}_j^0 - \boldsymbol{\theta}^*\|^2 + \tau_\ell \|\boldsymbol{\theta}_j^0 - \boldsymbol{\theta}^*\|_1^2 + \frac{1}{m} \ell_\Sigma \|\boldsymbol{\Theta}^0 - \boldsymbol{\Theta}^*\|_F^2 + \tau_\ell \frac{1}{m} \sum_{k=1}^m \|\boldsymbol{\theta}_k^0 - \boldsymbol{\theta}^*\|_1^2 \right)}_{A_3} \\
&\quad + sc_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu,
\end{aligned} \tag{124}$$

where (a) follows from Lemma 1 while in (b) used the bound $2 \cdot (c_g \sqrt{s}) \cdot \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\| \leq \epsilon \ell_\Sigma \cdot \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + sc_g^2 \ell_\Sigma^{-1} \epsilon^{-1}$.

Similarly, we can bound $T_{3,2}$ as

$$\begin{aligned}
T_{3,2} &= \left| \left(\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}} \right)^\top \left(\frac{1}{n} \mathbf{X}_j^\top \mathbf{X}_j \right) (\boldsymbol{\theta}_j^{s+1} - \boldsymbol{\theta}_j^s) \right| \\
&\leq \frac{\epsilon}{2} \left(\ell_\Sigma \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 + \tau_\ell \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|_1^2 \right) + \frac{1}{2\epsilon} \left(\ell_\Sigma \|\boldsymbol{\theta}_j^{s+1} - \boldsymbol{\theta}_j^s\|^2 + \tau_\ell \|\boldsymbol{\theta}_j^{s+1} - \boldsymbol{\theta}_j^s\|_1^2 \right) \\
&\leq \frac{\epsilon}{2} (\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 \\
&\quad + \frac{1}{2\epsilon} \left(\ell_\Sigma \|\boldsymbol{\theta}_j^{s+1} - \boldsymbol{\theta}_j^s\|^2 + 16s\tau_\ell \|\boldsymbol{\theta}_j^{s+1} - \widehat{\boldsymbol{\theta}}\|^2 + 16s\tau_\ell \|\boldsymbol{\theta}_j^s - \widehat{\boldsymbol{\theta}}\|^2 \right) + \tau_\ell \nu^2 (\epsilon + 4\epsilon^{-1})
\end{aligned} \tag{125}$$

Using (124) and (125) along with $\|(\mathbf{W} - \mathbf{J})^t\|_\infty \leq c_m \rho^t$, we can bound T_3 as follows:

$$\begin{aligned}
T_3 &= \sum_{i=1}^m (\bar{\mathbf{g}}^t - \mathbf{g}_i^t)^\top (\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}) \\
&\leq \sum_{i=1}^m \sum_{j=1}^m |\ell_{ij}^{(t)}| \cdot T_{3,1} + \sum_{s=0}^{t-1} \sum_{i=1}^m \sum_{j=1}^m |\ell_{ij}^{(t-s)}| \cdot T_{3,2} \\
&\leq \sum_{i=1}^m \sum_{j=1}^m |\ell_{ij}^{(t)}| \left(\epsilon (2\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 \right) + \sum_{i=1}^m \sum_{j=1}^m |\ell_{ij}^{(t)}| \left(\epsilon \cdot 2\tau_\ell \nu^2 + \epsilon^{-1} A_3 + s c_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu \right) \\
&\quad + \sum_{s=0}^{t-1} \sum_{i=1}^m \sum_{j=1}^m |\ell_{ij}^{(t-s)}| \left(\frac{\epsilon}{2} (\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\theta}_i^{t+\frac{1}{2}} - \widehat{\boldsymbol{\theta}}\|^2 \right) \\
&\quad + \sum_{s=0}^{t-1} \sum_{i=1}^m \sum_{j=1}^m |\ell_{ij}^{(t-s)}| \frac{1}{2\epsilon} \left(\ell_\Sigma \|\boldsymbol{\theta}_j^{s+1} - \boldsymbol{\theta}_j^s\|^2 + 16s\tau_\ell \|\boldsymbol{\theta}_j^{s+1} - \widehat{\boldsymbol{\theta}}\|^2 + 16s\tau_\ell \|\boldsymbol{\theta}_j^s - \widehat{\boldsymbol{\theta}}\|^2 \right) \\
&\quad + \sum_{s=0}^{t-1} \sum_{i=1}^m \sum_{j=1}^m |\ell_{ij}^{(t-s)}| \tau_\ell \nu^2 (\epsilon + 4\epsilon^{-1}) \\
&\leq c_m \rho^t \epsilon (2\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 + m c_m \rho^t \left(\epsilon \cdot 2\tau_\ell \nu^2 + \epsilon^{-1} A_3 + s c_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu \right) \\
&\quad + \sum_{s=0}^{t-1} c_m \rho^{t-s} \frac{\epsilon}{2} (\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 \\
&\quad + \sum_{s=0}^{t-1} c_m \rho^{t-s} \frac{1}{2\epsilon} \ell_\Sigma \|\boldsymbol{\Theta}^{s+1} - \boldsymbol{\Theta}^s\|_F^2 + \sum_{s=0}^{t-1} c_m \rho^{t-s} \frac{1}{\epsilon} 8s\tau_\ell \|\boldsymbol{\Theta}^{s+1} - \widehat{\boldsymbol{\Theta}}\|^2 + \sum_{s=0}^{t-1} c_m \rho^{t-s} \frac{1}{\epsilon} 8s\tau_\ell \|\boldsymbol{\Theta}^s - \widehat{\boldsymbol{\Theta}}\|^2 \\
&\quad + \sum_{s=0}^{t-1} m c_m \rho^{t-s} \tau_\ell \nu^2 (\epsilon + 4\epsilon^{-1}) \\
&\leq \frac{3}{2} \cdot \frac{\rho \cdot c_m \epsilon}{1 - \rho} (2\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 + \frac{c_m \ell_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^{s+1} - \boldsymbol{\Theta}^s\|_F^2 \\
&\quad + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^{s+1} - \widehat{\boldsymbol{\Theta}}\|_F^2 + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^s - \widehat{\boldsymbol{\Theta}}\|_F^2 \\
&\quad + m c_m \rho^t (A_3 \epsilon^{-1} + s c_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu) + \frac{\rho \cdot m c_m}{1 - \rho} 4\tau_\ell \nu^2 (\epsilon + \epsilon^{-1}) \\
&\leq \frac{3}{2} \cdot \frac{\rho \cdot c_m \epsilon}{1 - \rho} (2\ell_\Sigma + 8s\tau_\ell) \|\boldsymbol{\Theta}^{t+\frac{1}{2}} - \widehat{\boldsymbol{\Theta}}\|_F^2 + \frac{c_m \ell_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\boldsymbol{\Theta}^{s+1} - \boldsymbol{\Theta}^s\|_F^2
\end{aligned}$$

$$\begin{aligned}
& + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+1} - \widehat{\Theta}\|_F^2 + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^s - \widehat{\Theta}\|_F^2 \\
& + mc_m \rho^t (A_3 \epsilon^{-1} + sc_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + c_g \nu) + \frac{\rho \cdot mc_m}{1-\rho} 4\tau_\ell \nu^2 (\epsilon + \epsilon^{-1}).
\end{aligned}$$

C.4 Bounding δ^t in (109)

Using the bounds on T_1 , T_2 , and T_3 , we can finally bound δ^t :

$$\begin{aligned}
\delta^t &= \sum_{i=1}^m (\nabla \mathcal{L}(\theta_i^t) - \mathbf{y}_i^t)^\top (\theta_i^{t+\frac{1}{2}} - \widehat{\theta}) \\
&\leq T_1 + T_2 + T_3 \\
&\leq \frac{\rho \cdot c_m \epsilon}{1-\rho} (L_\Sigma + 8s\tau_g) \left\| \Theta^{t+\frac{1}{2}} - \widehat{\Theta} \right\|_F^2 + \frac{c_m L_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta \Theta^s\|_F^2 \\
&\quad + c_m \cdot 8s\tau_g \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+\frac{1}{2}} - \widehat{\Theta}\|_F^2 + c_m \cdot 8s\tau_g \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^s - \widehat{\Theta}\|_F^2 \\
&\quad + \frac{\rho \cdot mc_m}{1-\rho} 4\tau_g \nu^2 (\epsilon + \epsilon^{-1}) \\
&\quad + \frac{\rho \cdot c_m \epsilon}{1-\rho} (\ell_\Sigma + 8s\tau_\ell) \left\| \Theta^{t+\frac{1}{2}} - \widehat{\Theta} \right\|_F^2 + \frac{c_m \ell_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta \Theta^s\|_F^2 \\
&\quad + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+\frac{1}{2}} - \widehat{\Theta}\|_F^2 + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^s - \widehat{\Theta}\|_F^2 \\
&\quad + \frac{\rho \cdot mc_m}{1-\rho} 4\tau_\ell \nu^2 \cdot (\epsilon + \epsilon^{-1}) \\
&\quad + \frac{3}{2} \cdot \frac{\rho \cdot c_m \epsilon}{1-\rho} (2\ell_\Sigma + 8s\tau_\ell) \left\| \Theta^{t+\frac{1}{2}} - \widehat{\Theta} \right\|_F^2 + \frac{c_m \ell_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+1} - \Theta^s\|_F^2 \\
&\quad + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+1} - \widehat{\Theta}\|_F^2 + c_m 8s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^s - \widehat{\Theta}\|_F^2 \\
&\quad + mc_m \rho^t (A_3 \epsilon^{-1} + sc_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + 1) + \frac{\rho \cdot mc_m}{1-\rho} 4\tau_\ell \nu^2 (\epsilon + \epsilon^{-1}) + \frac{\rho \cdot mc_m}{1-\rho} c_g^2 \nu^2 \\
&\stackrel{(a)}{\leq} \frac{7}{2} \cdot \frac{\rho \cdot c_m \epsilon}{1-\rho} (2\ell_\Sigma + 8s\tau_\ell) \left\| \Theta^{t+\frac{1}{2}} - \widehat{\Theta} \right\|_F^2 \\
&\quad + \frac{c_m \ell_\Sigma}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta \Theta^s\|_F^2 + \frac{c_m \ell_\Sigma}{2\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+1} - \Theta^s\|_F^2 \\
&\quad + c_m 24\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+\frac{1}{2}} - \widehat{\Theta}\|_F^2 + c_m 24\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^s - \widehat{\Theta}\|_F^2 \\
&\quad + mc_m \rho^t (A_3 \cdot \epsilon^{-1} + sc_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + 1) + \frac{\rho \cdot mc_m}{1-\rho} 12\tau_\ell \nu^2 (\epsilon + \epsilon^{-1}) + \frac{\rho \cdot mc_m}{1-\rho} c_g^2 \nu^2 \\
&\stackrel{(b)}{\leq} \frac{7}{2} \cdot \frac{\rho \cdot c_m \epsilon}{1-\rho} (2\ell_\Sigma + 8s\tau_\ell) \left\| \Theta^{t+\frac{1}{2}} - \widehat{\Theta} \right\|_F^2 \\
&\quad + \frac{2c_m \ell_\Sigma}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta \Theta^s\|_F^2 + \frac{2c_m \ell_\Sigma}{\epsilon} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta_\perp^s\|_F^2
\end{aligned}$$

$$\begin{aligned}
& + c_m 24s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s+\frac{1}{2}} - \widehat{\Theta}\|_F^2 + c_m 24s\tau_\ell \epsilon^{-1} \sum_{s=0}^{t-1} \rho^{t-s} \|\Theta^{s-\frac{1}{2}} - \widehat{\Theta}\|_F^2 \\
& + mc_m \rho^t (A_3 \cdot \epsilon^{-1} + sc_g^2 \ell_\Sigma^{-1} \epsilon^{-1} + 1) + \frac{\rho \cdot mc_m}{1-\rho} 12\tau_\ell \nu^2 (\epsilon + \epsilon^{-1}) + \frac{\rho \cdot mc_m}{1-\rho} c_g^2 \nu^2,
\end{aligned}$$

where in (a) we used $L_\Sigma \leq \ell_\Sigma$, $\tau_g \leq \tau_\ell$, and (120); and in (b) we used again (120) and (121). $\square$

D Proof of Proposition 11

Using Assumptions 6 and 7 on (110) we have

$$\|\Theta_\perp^t\| \leq \rho^t \|\Theta_\perp^0\| + \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta\Theta^s\|. \quad (126)$$

Squaring both sides gives

$$\begin{aligned}
\|\Theta_\perp^t\|^2 & \leq \left(\rho^t \|\Theta_\perp^0\| + \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta\Theta^s\| \right)^2 \\
& \leq 2\rho^{2t} \|\Theta_\perp^0\|^2 + 2 \left(\sum_{s=0}^{t-1} \rho^{t-s} \|\Delta\Theta^s\| \right)^2 \\
& \leq 2\rho^{2t} \|\Theta_\perp^0\|^2 + \frac{2\rho}{1-\rho} \cdot \sum_{s=0}^{t-1} \rho^{t-s} \|\Delta\Theta^s\|^2,
\end{aligned} \quad (127)$$

where the last inequality follows from Cauchy-Schwartz inequality.

E Proof of Lemma 15

Recall $G_{\text{net}}(z) = \frac{\rho}{z-\rho}$. Since (55b) is quadratic in $G_{\text{net}}(z)$, we obtain the following feasible range of $G_{\text{net}}(z)$:

$$\begin{aligned}
G_{\text{net}}(z) & \leq \frac{\frac{C_2}{\gamma} \frac{c_m \ell_\Sigma}{\epsilon} - \sqrt{\left(\frac{C_2}{\gamma} \frac{c_m \ell_\Sigma}{\epsilon}\right)^2 + \frac{4C_2}{\gamma} \frac{c_m \ell_\Sigma}{\epsilon} \frac{2\rho}{1-\rho} \left(1 - \frac{L_\Sigma}{\gamma}\right)}}{-\frac{2C_2}{\gamma} \frac{c_m \ell_\Sigma}{\epsilon} \frac{2\rho}{1-\rho}} \\
& = \frac{1-\rho}{4\rho} \left(\sqrt{1 + \frac{8\rho}{1-\rho} \left(1 - \frac{L_\Sigma}{\gamma}\right) \left(\frac{\gamma}{C_2} \frac{\epsilon}{c_m \ell_\Sigma}\right)} - 1 \right) \\
& \stackrel{(56)}{=} \frac{1-\rho}{4\rho} \left(\sqrt{1 + (\gamma - L_\Sigma) \left(\frac{\mu_\Sigma}{\ell_\Sigma^2} \frac{2\rho}{C_2^2 c_m^2}\right)} - 1 \right).
\end{aligned} \quad (128)$$

Therefore, condition (55b) is fulfilled for all z in the form

$$z = \rho + \frac{\rho}{G_{\text{net}}(z)} \geq \rho + \frac{4\rho^2}{1-\rho} \left(\sqrt{1 + (\gamma - L_\Sigma) \left(\frac{\mu_\Sigma}{\ell_\Sigma^2} \frac{2\rho}{C_2^2 c_m^2}\right)} - 1 \right)^{-1}. \quad (129)$$

$\square$

F Proof of Lemma 16

We substitute $G_{\text{net}}(z) = \frac{\rho}{z-\rho}$ into (60) and solve the equation

$$z = \frac{A + \frac{\rho}{z-\rho}}{D - \frac{\rho}{z-\rho}}. \quad (130)$$

Let Δ denote the discriminant of the second order equation in z (60). Since

$$\begin{aligned} \Delta &= (D\rho + \rho + A)^2 - 4D(A-1)\rho \\ &= (D\rho + \rho)^2 + A^2 + 2(D\rho + \rho)A - 4D(A-1)\rho \\ &= (D\rho + \rho)^2 + A^2 - 2(D\rho + \rho)A - 4D(A-1)\rho + 4(D\rho + \rho)A \\ &= (D\rho + \rho - A)^2 + 4(A+D)\rho \geq 0, \end{aligned} \quad (131)$$

Eq. (130) has two real roots ($z_1 \leq z_2$)

$$z_{1,2} = \frac{(D+1)\rho + A \pm \sqrt{(D\rho + \rho + A)^2 - 4D(A-1)\rho}}{2D}. \quad (132)$$

By inspection, we have that $z_1 \geq 0$. Hence, (60) is satisfied for all $z \in (0, z_1] \cup [z_2, 1)$.

Next we show $z_1 \leq \rho$ and $z_2 \geq \rho$. As $z > \rho$ must be satisfied, $z \in [z_2, 1)$. Suppose on the contrary that $z_1 > \rho$, which implies that

$$\begin{aligned} \rho + A - D\rho &> \sqrt{(D\rho + \rho + A)^2 - 4D(A-1)\rho} \\ \Leftrightarrow (\rho + A - D\rho)^2 &> (D\rho + \rho + A)^2 - 4D(A-1)\rho \quad \text{and} \quad \rho + A - D\rho > 0. \end{aligned} \quad (133)$$

After simplification, the first inequality becomes

$$4D\rho^2 + 4D\rho < 0, \quad (134)$$

which is impossible. Therefore we have proved that $z_1 \leq \rho$.

As both z_1 and z_2 are nonnegative, we can simply upperbound z_2 as

$$z_2 \leq z_1 + z_2 = \frac{(D+1)\rho + A}{D}. \quad (135)$$

This completes the proof. $\square$

G Proof of Lemma 17

Substituting γ into the expression of A and D we get

$$\begin{aligned} \frac{A}{D} &= \frac{\gamma - \mu_\Sigma + C_1 s(\tau_\mu + \tau_g)}{\gamma - \left\{ C_1 s\tau_g + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma} (\ell_\Sigma + s\tau_\ell) \right\}} \\ &= \frac{L_\Sigma - \mu_\Sigma + C_1 s(\tau_\mu + \tau_g) + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4}}{L_\Sigma - C_1 s\tau_g - \frac{\rho}{2} \mu_\Sigma - \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma} s\tau_\ell + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4}} \\ &= \frac{1 - \kappa^{-1} + C_1 s(\tau_\mu + \tau_g)/L_\Sigma + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4}}{1 - C_1 s\tau_g/L_\Sigma - \frac{\rho}{2} \kappa^{-1} - \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4}}. \end{aligned} \quad (136)$$

Dividing both the numerator and denominator by $1 - C_1 s\tau_g/L_\Sigma$ and using the definition of σ_0 we have

$$\begin{aligned}
\frac{A}{D} &= \frac{\sigma_0 + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4} (1 - C_1 s\tau_g/L_\Sigma)^{-1}}{1 + \left(-\frac{\rho}{2} \kappa^{-1} - \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4} \right) (1 - C_1 s\tau_g/L_\Sigma)^{-1}} \\
&\stackrel{A \leq B}{\leq} \frac{\sigma_0 + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4} (1 - C_1 s\tau_g/L_\Sigma)^{-1} + \left(\frac{\rho}{2} \kappa^{-1} + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell \right) (1 - C_1 s\tau_g/L_\Sigma)^{-1}}{1 + \left(C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4} \right) (1 - C_1 s\tau_g/L_\Sigma)^{-1}} \\
&\leq \sigma_0 + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4} (1 - C_1 s\tau_g/L_\Sigma)^{-1} + \frac{\rho}{2} \left(\kappa^{-1} + \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell \right) (1 - C_1 s\tau_g/L_\Sigma)^{-1} \\
&= \sigma_0 + \sqrt{\rho} \cdot \frac{C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4} + \frac{1}{2} \left(\kappa^{-1} + \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell \right)}{1 - C_1 s\tau_g/L_\Sigma} \\
&\stackrel{\rho \leq 1/2}{\leq} \sigma_0 + \sqrt{\rho} \cdot \frac{16C_3 c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} + \frac{1}{2} \left(\kappa^{-1} + \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell \right)}{1 - C_1 s\tau_g/L_\Sigma},
\end{aligned} \tag{137}$$

where for $A \leq D$ to hold, we require

$$\left(1 - \frac{\rho}{2} \right) \mu_\Sigma \geq C_1 s\tau_g + C_1 s(\tau_\mu + \tau_g) + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma} s\tau_\ell. \tag{138}$$

Similarly, we can bound $\rho + \rho/D$ as

$$\begin{aligned}
\rho + \frac{\rho}{D} &= \rho + \frac{\rho \frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_2^2 c_m^2}{1-\rho} s\tau_\ell}{\gamma - \left\{ C_1 s\tau_g + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma} (\ell_\Sigma + s\tau_\ell) \right\}} \\
&= \rho + \frac{\rho \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \frac{2C_2^2 c_m^2}{1-\rho} s\tau_\ell}{1 - C_1 s\tau_g/L_\Sigma + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4} - \left\{ \frac{\rho}{2} \kappa^{-1} + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell \right\}} \\
&\stackrel{\rho \leq B}{\leq} \rho + \frac{\rho \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \frac{2C_2^2 c_m^2}{1-\rho} s\tau_\ell + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell}{1 - C_1 s\tau_g/L_\Sigma - \frac{\rho}{2} \kappa^{-1} + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4}} \\
&\stackrel{\rho \kappa^{-1} \leq 1}{\leq} \rho + \frac{\rho \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \frac{2C_2^2 c_m^2}{1-\rho} s\tau_\ell + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell}{\frac{1}{2} - C_1 s\tau_g/L_\Sigma + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4}},
\end{aligned} \tag{139}$$

where $\rho \leq D$ requires

$$L_\Sigma + C_3 \frac{\ell_\Sigma^2}{\mu_\Sigma} \frac{c_m^2 \sqrt{\rho}}{(1-\rho)^4} \geq \rho \frac{\ell_\Sigma}{\mu_\Sigma} \frac{2C_2^2 c_m^2}{1-\rho} s\tau_\ell + C_1 s\tau_g + \frac{\rho}{2} \mu_\Sigma + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma} s\tau_\ell. \tag{140}$$

Using the fact that $\rho \leq 1$ and $L_\Sigma \geq \mu_\Sigma$, it suffices

$$\mu_\Sigma \geq \rho \frac{\ell_\Sigma}{\mu_\Sigma} \frac{4C_2^2 c_m^2}{1-\rho} s\tau_\ell + 2C_1 s\tau_g + \rho \frac{\mu_\Sigma}{\ell_\Sigma} s\tau_\ell, \tag{141}$$

Note that (66) implies both (138) and (141).

Finally, we further simplify the bound in (139) as follows. Multiplying both the numerator and denominator by $(1-\rho)^4$ we get

$$\rho + \frac{\rho}{D} \leq \rho + \frac{\rho \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \cdot 2C_2^2 c_m^2 s\tau_\ell (1-\rho)^3 + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell (1-\rho)^4}{\left(\frac{1}{2} - C_1 s\tau_g/L_\Sigma \right) (1-\rho)^4 + C_3 c_m^2 \sqrt{\rho} \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma}}. \tag{142}$$

Define function

$$f(\rho) = \left(\frac{1}{2} - C_1 s\tau_g / L_\Sigma \right) (1 - \rho)^4 + C_3 c_m^2 \sqrt{\rho} \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma}. \quad (143)$$

Differentiating f with respect to ρ yields

$$f'(\rho) = -4 \left(\frac{1}{2} - C_1 s\tau_g / L_\Sigma \right) (1 - \rho)^3 + C_3 c_m^2 \frac{\ell_\Sigma^2}{\mu_\Sigma L_\Sigma} \frac{1}{2\sqrt{\rho}} > 0, \quad (144)$$

for $C_3 \geq 4$, due to $c_m \geq 1$, $\ell_\Sigma \geq \mu_\Sigma$, and $\ell_\Sigma \geq L_\Sigma$. Therefore, $f(\rho)$ is monotonically increasing in $(0, 1)$ and it holds

$$\begin{aligned} \rho + \frac{\rho}{D} &\leq \rho + \frac{\rho \frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \cdot 2C_2^2 c_m^2 s\tau_\ell + \frac{\rho}{2} \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell}{\frac{1}{2} - C_1 s\tau_g / L_\Sigma} \\ &= \rho + \rho \cdot \frac{\frac{\ell_\Sigma}{\mu_\Sigma L_\Sigma} \cdot 4C_2^2 c_m^2 s\tau_\ell + \frac{\mu_\Sigma}{\ell_\Sigma L_\Sigma} s\tau_\ell}{1 - 2C_1 s\tau_g / L_\Sigma}. \end{aligned} \quad (145)$$

□

H Proof of Lemma 20

We first bound $\max_{j \in [m]} \|\mathbf{X}_j^\top \mathbf{n}_j / n\|_\infty$. Each column of $\mathbf{X}_j$ is an n -dimensional Gaussian random vector with independent entries. The maximum variance of these entries is bounded by $\zeta = \max_{j \in [d]} \Sigma_{jj}$. Since $\mathbf{X}_j$ is independent of $\mathbf{n}_j$ and the elements of $\mathbf{n}_j$ are i.i.d. σ^2 -sub-Gaussian, each element of $\mathbf{X}_j^\top \mathbf{n}_j$ is the sum of n independent sub-exponential random variables with sub-exponential norm no larger than $\sqrt{\zeta}\sigma$. Applying the Bernstein's inequality and the union bound we get

$$\mathbb{P} \left(\max_{j \in [m]} \left\| \frac{\mathbf{X}_j^\top \mathbf{n}_j}{n} \right\|_\infty \geq t \right) \leq 2 \exp \left(-c_3 \min \left\{ \frac{t^2}{\zeta \sigma^2}, \frac{t}{\sqrt{\zeta} \sigma} \right\} n + \log md \right) \quad (146)$$

for some $c_3 > 0$. When $\log md \geq (c_3/2) \cdot n$, we choose $t = (2/c_3) \sqrt{\zeta} \sigma \cdot \log md / n \geq \sqrt{\zeta} \sigma$, which leads to

$$\mathbb{P} \left(\max_{j \in [m]} \left\| \frac{\mathbf{X}_j^\top \mathbf{n}_j}{n} \right\|_\infty \geq \sqrt{\zeta} \sigma \cdot \frac{2 \log md}{c_3 n} \right) \leq 2 \exp(-\log md). \quad (147)$$

On the other hand, when $\log md < (c_3/2) \cdot n$, we choose $t = \sqrt{\zeta} \sigma \cdot \sqrt{(2/c_3) \log md / n} < \sqrt{\zeta} \sigma$, which leads to

$$\mathbb{P} \left(\max_{j \in [m]} \left\| \frac{\mathbf{X}_j^\top \mathbf{n}_j}{n} \right\|_\infty \geq \sqrt{\zeta} \sigma \cdot \sqrt{\frac{2 \log md}{c_3 n}} \right) \leq 2 \exp(-\log md). \quad (148)$$

Combining the two cases we obtain (95).

Applying the same procedure to $\|\nabla \mathcal{L}(\boldsymbol{\theta}^*)\|_\infty$, proves (96). □

I Proof of Proposition 5

The proof is a modification of [35]. Next, we only highlight the key differences, for completeness. Define the set

$$V(r) = \{r \in \mathbb{R}^d \mid \|\Sigma^{\frac{1}{2}} v\|_2 = 1, \|v\|_1 \leq r\},$$

for fixed radius $r > 0$. Accordingly, define the random variable

$$M(r, \mathbf{X}_i) = \sup_{v \in V(r)} \frac{\|\mathbf{X}_i v\|}{\sqrt{n}}.$$

Following similar steps as in [35, Lemma 1], and using the Sudakov-Fernique inequality for Gaussian processes, we can bound the expectation of $M(r, \mathbf{X}_i)$ as

$$\mathbb{E}M(r, \mathbf{X}_i) \leq 1 + 3\rho(\Sigma)\sqrt{\frac{\log d}{n}}r \triangleq t(r), \quad (149)$$

where $\rho^2(\Sigma) = \max_j \Sigma_{jj}$. Then applying the concentration inequality in the same way as in [35, Lemma 2] we obtain

$$\mathbb{P}(M(r, \mathbf{X}_i) > t(r) + \sqrt{mt}(r)) \leq 2\exp(-Nt(r)^2/2) \leq 2\exp(-n((\sqrt{m}+1)t(r))^2/8). \quad (150)$$

Finally, define the event

$$\mathcal{T}_i = \{\exists v \in \mathbb{R}^d \text{ s.t. } \|\Sigma^{1/2}v\|_2 = 1 \text{ and } \|\mathbf{X}_i v\|_2/\sqrt{n} > 2(\sqrt{m}+1)t(\|v\|_1)\}.$$

Applying [35, Lemma 3] with

$$f(v, \mathbf{X}_i) = \|\mathbf{X}_i v\|_2/\sqrt{n}, \quad h(v) = \|v\|_1, \quad g(r) = (\sqrt{m}+1)t(r), \quad a_n = n/8 \quad \text{and} \quad \mu = \sqrt{m},$$

we have

$$\mathbb{P}(\mathcal{T}_i^c) \geq 1 - c'_2 \exp(-c'_0 N),$$

for some $c'_0 \geq 1/2$.

Let us use now the union bound:

$$\mathbb{P}(\cup_i \mathcal{T}_i) \leq \sum_{i=1}^m \mathbb{P}(\mathcal{T}_i) \leq c'_2 m \exp(-c'_0 N) = c'_2 \exp(-c'_0 N + \log m). \quad (151)$$

Therefore

$$\frac{\|\mathbf{X}_i v\|}{\sqrt{n}} \leq 2(\sqrt{m}+1)\|\Sigma^{1/2}v\| + 6(\sqrt{m}+1)\rho(\Sigma)\sqrt{\frac{\log d}{n}}\|v\|_1, \quad \forall v \in \mathbb{R}^d, \quad \forall i \in [m], \quad (152)$$

with probability larger than $1 - c'_2 \exp(-c'_0 N + \log m)$. Since $N/2 \geq m/2 > \log m$, it holds $1 - \exp(-c'_0 N + \log m) \geq 1 - \exp(-c''_0 N)$, for some $c''_0 > 0$. $\square$

This figure "DocFig1to3.png" is available in "png" format from:

<http://arxiv.org/ps/2201.08507v1>

This figure "DocFig4.png" is available in "png" format from:

<http://arxiv.org/ps/2201.08507v1>

This figure "DocFig5.png" is available in "png" format from:

<http://arxiv.org/ps/2201.08507v1>

This figure "DocFig6.png" is available in "png" format from:

<http://arxiv.org/ps/2201.08507v1>

This figure "DocFig7.png" is available in "png" format from:

<http://arxiv.org/ps/2201.08507v1>

This figure "DocFig8.png" is available in "png" format from:

<http://arxiv.org/ps/2201.08507v1>

This figure "preview.jpeg" is available in "jpeg" format from:

<http://arxiv.org/ps/2201.08507v1>