BAHADUR EFFICIENCY FOR CERTAIN GOODNESS-OF-FIT TESTS BASED ON THE EMPIRICAL CHARACTERISTIC FUNCTION

Simos G. Meintanis^{*,b}, Bojana Milošević[‡], and Marko Obradović[‡]

^aDepartment of Economics, National and Kapodistrian University of Athens, Athens, Greece ^bPure and Applied Analytics, North–West University, Potchefstroom, South Africa ^cFaculty of Mathematics, University of Belgrade, Studenski trg 16, Belgrade, Serbia

ABSTRACT

We study the Bahadur efficiency of several weighted L2–type goodness–of–fit tests based on the empirical characteristic function. The methods considered are for normality and exponentiality testing, and for testing goodness–of–fit to the logistic distribution. Our results are helpful in deciding which specific test a potential practitioner should apply. For the celebrated BHEP and energy tests for normality we obtain novel efficiency results, with some of them in the multivariate case, while in the case of the logistic distribution this is the first time that efficiencies are computed for any composite goodness–of–fit test.

Keywords Goodness–of–fit test · Bahadur efficiency · Empirical characteristic function · Normality test · Exponentiality test

1 Introduction

Let $X_1, ..., X_n$, denote independent copies of an arbitrary random variable $X \in \mathbb{R}^p$, and consider the problem of testing the composite goodness–of–fit (GOF) null hypothesis,

$$\mathcal{H}_0: \text{The law of } X \in \mathcal{F}_{\vartheta}, \text{ for some } \vartheta, \tag{1.1}$$

where $\mathcal{F}_{\vartheta} := \{F_{\vartheta}, \vartheta \in \Theta\}$ denotes a class of distributions indexed by the parameter $\vartheta \in \Theta \subseteq \mathbb{R}^q$, $q \ge 1$.

For certain popular distributions, such as the normal and the exponential distribution, there exist many GOF tests, while for others such as the logistic distribution the range of methods available is not so extended. In either case though a potential practitioner would like to have some quality measure on the basis of which one could choose amongst the existing GOF methods. In this connection, one of the most popular methods for test comparison is the so–called Bahadur efficiency that allows to compare the efficiency of any given GOF test vis–á–vis its optimal counterpart, which is the likelihood ratio (LR) test for the distribution under test against a specified alternative.

^{*}simosmei@econ.uoa.gr

[†]bojana@matf.bg.ac.rs

[‡]marcone@matf.bg.ac.rs

In this work we consider GOF tests that are based on the weighted L2-type test statistic

$$T_{n,w} = n \int_{\mathbb{R}^p} \left| \varphi_n(t) - \varphi_{\widehat{\vartheta}_n}(t) \right|^2 w(t) \mathrm{d}t,$$
(1.2)

where $\varphi_{\vartheta}(\cdot)$ denotes the characteristic function corresponding to F_{ϑ} ,

$$\varphi_n(t) = \frac{1}{n} \sum_{j=1}^n e^{it^\top X_j},\tag{1.3}$$

is the empirical characteristic function (ECF) and $\hat{\vartheta}_n := \hat{\vartheta}_n(X_1, ..., X_n)$ is an estimator of ϑ obtained on the basis of $(X_j, j = 1, ..., n)$.

The test statistic $T_{n,w}$, besides the family \mathcal{F}_{ϑ} being tested and the corresponding estimator $\widehat{\vartheta}_n$ employed, also depends on the the weight function $w \ge 0$ figuring in (1.2). For certain choices of $w(\cdot)$, we obtain here efficiency results for the BHEP GOF test for normality of Epps and Pulley (1983) and Henze and Wagner (1997) as well as for the generalized energy (GE) test for normality first put forward by Székely and Rizzo (2005). In doing so we provide extension of the efficiency results of Ebner and Henze (2021a) and Tenreiro (2009) for the BHEP test as well as extension of the results of Móri et al. (2021) from the original energy statistic to its generalized counterpart suggested by Székely and Rizzo (2013). Furthermore we obtain for the first time analogous results for the GOF test for the logistic distribution of Meintanis (2004) and for the exponentiality test suggested by Henze and Meintanis (2005), including efficiency comparisons with alternative tests. As already mentioned these efficiency results will facilitate the choice of the specific weight function $w(\cdot)$ that figures in the test statistics defined by (1.2), and thus provide some guidance on which method to apply amongst the many tests available for the normal, the exponential distribution and the logistic distribution.

The rest of this work unfolds as follows. In Section 2 we review the basic theory of Bahadur efficiency and the theory of the limit null distribution of the test statistics, while in Section 3 the necessary efficiency computations are discussed. In Section 4 we provide efficiency results and discussion for the BHEP and energy statistics, as well as for the two aforementioned ECF–based tests, one for the logistic and the other for the exponential distribution. Analogous efficiency results for the normality tests in the bivariate case are reported in Section 5, and we conclude in Section 6 with discussion.

2 Limit null distribution and Bahadur slopes

The limit in distribution of the L2-type test statistics figuring in (1.2) is given by

$$T_{n,w} \xrightarrow{\mathcal{D}} \sum_{j=1}^{\infty} \lambda_j N_j^2,$$
 (2.1)

where N_j , $j \ge 1$, are independent copies of a standard normally distributed random variable, and $\lambda_1 \ge \lambda_2 \ge ...$, are the eigenvalues of the integral equation

$$\lambda f(s) = \int K(s,t)f(t)w(t)dt; \qquad (2.2)$$

where K(s,t) is a covariance kernel associated, besides the weight function $w(\cdot)$, with the given test statistic and the estimator employed in estimating the distributional parameter ϑ ; see for instance Meintanis and Swanepoel (2007).

In this section, we briefly review the essence of the Bahadur theory, and associate that theory with the limit law of $T_{n,w}$ figuring in (2.1)–(2.2); for more details we refer to Bahadur (1971) and Nikitin (1995), and to the more recent article

by Grané and Tchirina (2013). To this end let $\mathcal{G}_{\theta} = \{G_{\theta}, \theta > 0\}$ be a family of alternative distribution functions, such that G_0 is the null family of distributions for some typical value $\vartheta \in \Theta$. Assume that the regularity conditions for V–statistics with weakly degenerate kernels in Nikitin and Peaucelle (2004), are satisfied.

Also recall that LR tests are optimal tests in the Bahadur sense. Hence, for close alternatives from \mathcal{G}_{θ} , the absolute local approximate Bahadur efficiency for any sequence of test statistics, say $\{T_n\}$, is defined as the ratio of the Bahadur approximate slope of the considered test statistic to the corresponding slope of the LR test, i.e.

$$\operatorname{eff}(T_n) = \lim_{\theta \to 0} \frac{c_T(\theta)}{2K(\theta)},\tag{2.3}$$

where

$$c_T(\theta) = \frac{b_T(\theta)}{\lambda_1} = \frac{b_T''(0)}{2\lambda_1}\theta^2 + o(\theta^2), \qquad (2.4)$$

(the last equation obtained by a Taylor expansion), with λ_1 being the largest eigenvalue figuring in (2.1) and

$$\frac{T_n}{n} \xrightarrow{\mathbb{P}} b_T(\theta), \tag{2.5}$$

where $K(\theta)$ is equal to minimal Kullback–Leibler (KL) distance from the given alternative to the class of distributions within the null hypothesis.

Before closing this section we wish to point out that the BHEP and GE tests are well known to be location/scale invariant, i.e. for both tests it holds,

$$T_{n,w}(a+bX_1,...,a+bX_n) = T_{n,w}(X_1,...,X_n), \text{ for each } a \in \mathbb{R}, b > 0,$$
(2.6)

and therefore their respective limit null distributions are independent of the true values of the mean and variance of the underlying Gaussian law. Note that this property extends to the multivariate version of these tests. In this connection, the other tests studied herein are also invariant, whereby invariance is understood within the context of the specific family \mathcal{F}_{ϑ} being tested. For instance if this family is confined to the positive real axis (such is the exponential family of distributions), invariance is understood only with respect to scale, i.e. (2.6) holds for a = 0, and b > 0.

3 KL distance and eigenvalue approximation

It can be seen from (2.3)–(2.5) that the basic components of an approximate Bahadur slope are the computation of the KL distance and of the largest eigenvalue. In this section we consider these components with emphasis on location–scale families of distributions.

3.1 KL distance for location-scale families

Assume that the family \mathcal{F}_{ϑ} is a location–scale family, i.e. a family generated by the density function f_{ϑ} corresponding to F_{ϑ} , where ϑ contains the location and scale parameters, μ and σ , respectively. We set $f_0(\cdot)$ and $F_0(\cdot)$ for the density and distribution function, respectively, of the standardized variable $Z := \sigma^{-1}(X - \mu)$. Recall also that for the family \mathcal{G}_{θ} with density $g_{\theta}(\cdot)$, $\theta > 0$, we have that $g_0 \equiv f_0$, and assume that certain regularity conditions are satisfied. Then the next theorem gives the local behavior of the KL distance $K(\theta; \mu, \sigma)$ between $g_{\theta} \in \mathcal{G}_{\theta}$ and $f_0 \in \mathcal{F}_{\vartheta}$,

$$K(\theta;\mu,\sigma) = \int \log(g_{\theta}(x))g_{\theta}(x)dx - \int \log(f_0(x))g_{\theta}(x)dx.$$
(3.1)

The proof of Theorem 3.1 is postponed to the Appendix.

Theorem 3.1 The KL distance from the alternative $g_{\theta}(\cdot)$ in \mathcal{G}_{θ} to the closest null distribution

$$K(\theta) = \inf_{\mu,\sigma} K(\theta;\mu,\sigma)$$

admits the representation

$$2K(\theta) = \left(\int \frac{h^2(x)}{f_0(x)} dx - (\sigma'(0))^2 + \int \frac{1}{f_0(x)} (f'_0(x))^2 (\mu'(0) + x\sigma'(0))^2 dx + 2\int \frac{1}{f_0(x)} f'_0(x) (\mu'(0) + x\sigma'(0)) h(x) dx \right) \theta^2 + o(\theta^2), \quad \theta \to 0,$$
(3.2)

where $(\mu(\theta), \sigma(\theta)) = \operatorname{arginf} K(\theta; \mu, \sigma)$ and $h(x) = \frac{\partial}{\partial \theta} g_{\theta}(x)|_{\theta=0}$.

In the case of some specific distribution the expression above can be simplified. Since the maximum likelihood (ML) estimators minimize the KL distance, the functions $\mu(\theta)$ and $\sigma(\theta)$ are equal to population counterparts, i.e. probability limits of the ML estimators of the location and scale parameter, respectively.

For example, in the case that \mathcal{F}_{ϑ} is the normal location–scale family, we have

$$\mu(\theta) = \int_{-\infty}^{\infty} x g_{\theta}(x) \mathrm{d}x; \ \sigma(\theta) = \left(\int_{-\infty}^{\infty} x^2 g_{\theta}(x) \mathrm{d}x - \mu^2(\theta)\right)^{1/2}$$

and thus the KL distance reduces to (see also Milošević et al. (2021))

$$2K(\theta) = \left(\int_{-\infty}^{\infty} \sqrt{2\pi}h^2(x)e^{\frac{x^2}{2}}dx - \left(\int_{-\infty}^{\infty} xh(x)dx\right)^2 - \frac{1}{2}\left(\int_{-\infty}^{\infty} x^2h(x)dx\right)^2\right)\theta^2 + o(\theta^2).$$

For the exponential scale family, we have

$$\mu(\theta) = 0; \ \sigma(\theta) = \int_0^\infty x g_\theta(x) \mathrm{d}x,$$

and thus the KL distance reduces to (see also Nikitin and Tchirina (1996))

$$2K(\theta) = \left(\int_0^\infty h^2(x)e^x dx - \left(\int_0^\infty xh(x)dx\right)^2\right)\theta^2 + o(\theta^2).$$

However, in the case that \mathcal{F}_{ϑ} is the logistic location–scale, the parameters $(\mu(\theta), \sigma(\theta))$ that minimize the KL distance figuring in (3.1) admit no closed–form expression. Nevertheless, by applying the implicit function theorem we obtain

$$\mu'(0) = 6 \int_{-\infty}^{\infty} \frac{h(x)}{1 + e^{-x}} dx; \ \sigma'(0) = \frac{9}{\pi^2 + 3} \int_{-\infty}^{\infty} \frac{(1 - e^{-x})xh(x)}{1 + e^{-x}} dx,$$

and thus by plugging these expressions in (3.2) we can obtain the corresponding KL distance.

3.2 Eigenvalue approximation

In this section we briefly review a method for eigenvalue approximation proposed in Božin et al. (2020). Recall that these eigenvalues are solutions of an integral equation involving a specific operator. At first we replace the original operator by a symmetric operator that has the same eigenvalues, and then (i) consider a truncated version of the symmetrized operator, and (ii) consider a discretized version of the truncated operator. In these two steps the truncated operator is chosen so that as the amount of truncation diminishes this operator converges to the symmetrized operator,

and likewise, the discretized operator approaches the truncated operator as the grid of discretization becomes more fine and extended.

Specifically we replace the original operator

$$Af(s) = \int_{-\infty}^{\infty} K(s,t)f(t)w(t)dt$$
$$\overline{A}f(s) = \int_{-\infty}^{\infty} K(s,t)f(t)\sqrt{w(t)w(s)}dt,$$
(3.3)

with the operator

that has the same spectrum as A, but is symmetric. Then in the first step we define the truncated operator
$$\overline{A_B}$$
 acting
on the set of real functions with support $[-B, B]$, $B > 0$, defined by

$$\overline{A_B}f(s) = \int_{-\infty}^{\infty} K(s,t)f(t)\sqrt{w(t)w(s)} \ 1(|t| \le B) \mathrm{d}t,$$

which clearly and for sufficiently large B, is close to \overline{A} .

In the second step of approximation we employ a sequence of symmetric linear operators $M^{(m)}$, which converges in norm to $\overline{A_B}$, as $m \to \infty$. This discretized sequence can be defined by $(2m + 1) \times (2m + 1)$ matrices $M^{(m)} = ||m_{i,j}^{(m)}||$, $-m \le i \le m, -m \le j \le m$, with elements

$$m_{i,j}^{(m)} = \frac{2B}{(2m+1)} K\left(\frac{iB}{m}, \frac{jB}{m}\right) \sqrt{w\left(\frac{iB}{m}\right) w\left(\frac{jB}{m}\right)}.$$
(3.4)

Using the perturbation theory–see (Kato, 2013, Theorem 4.10, page 291)–we have that the spectra of these two operators are at a distance that tends to zero. Hence within the degree of approximation, the sequence $\lambda_1^{(m)}$ of the largest eigenvalues of $M^{(m)}$ will converge to the largest eigenvalue $\lambda_1(B)$ of $\overline{A_B}$, which in turn approaches λ_1 . Consequently, the eigenvalues $\lambda_1^{(m)}$ and λ_1 will coincide up to any desired accuracy, provided that the pair of approximation parameters (m, B) is large enough.

4 Tests for the normal and the logistic distribution

When testing for GOF to a symmetric distribution such as the normal or the logistic, it is customary to use general purpose nonparametric alternatives parametrized by θ , a parameter that usually controls skewness. A discussion on the construction and applications of these alternatives is available in Jones (2015) and Ley (2015). Specifically we consider the following alternatives:

· Lehmann alternatives

$$g_{\theta}^{(1)}(x) = (1+\theta)F_0^{\theta}(x)f_0(x)$$

• first Ley-Paindaveine alternatives

$$g_{\theta}^{(2)}(x) = f_0(x)e^{-\theta(1-F_0(x))}(1+\theta F_0(x))$$

· second Ley-Paindaveine alternatives

$$g_{\theta}^{(3)}(x) = f_0(x)(1 - \theta\pi\cos(\pi F_0(x)))$$

• contamination alternatives

$$g^{(4)}_{\theta}(x;\mu,\sigma^2) = (1-\theta)f_0(x) + \theta f(x;\mu,\sigma^2)$$

Note that in the above alternatives, $f_0(\cdot)$ and $F_0(\cdot)$ are the null density and distribution function, respectively, so that for $\theta = 0$ each of these alternatives reduces to the null distribution under test.

4.1 Tests for normality

4.1.1 The GE test

The GE test may be formulated as

$$T_{n,w} = n \int_{-\infty}^{\infty} \left| \phi_n(t) - e^{-\frac{t^2}{2}} \right|^2 w(t) \mathrm{d}t,$$
(4.1)

with the ECF $\phi_n(t)$ obtained as in (1.3) with X_j replaced by

$$Z_j = \frac{X_j - \bar{X}_n}{S_n}, \ j = 1, ..., n,$$
(4.2)

where \bar{X}_n denotes the sample mean and S_n^2 the sample variance of $(X_j, j = 1, ..., n)$. As weight function Móri et al. (2021) use $w(t) = |t|^{-2}$. Here we consider the GE test of Székely and Rizzo (2013) whereby $w(t) = |t|^{-1-\gamma}$, $0 < \gamma < 2$, is adopted as weight function, thus rendering the convenient form

$$T_{n,w} = 2\sum_{j=1}^{n} \mathbb{E}|Z_j - X_1|^{\gamma} - n\mathbb{E}|X_1 - X_2|^{\gamma} - \frac{1}{n}\sum_{j,k=1}^{n} |Z_j - Z_k|^{\gamma}.$$
(4.3)

⁴ Note that the expectations in (4.3) are taken with respect to the standard Gaussian distribution, and consequently the GE test statistic may be explicitly expressed by using the equations

$$\mathbb{E}|X_1 - X_2|^{\gamma} = \frac{2^{\gamma}}{\sqrt{\pi}} \Gamma\left(\frac{1+\gamma}{2}\right),$$

and

$$\mathbb{E}|x - X_{1}|^{\gamma} = \frac{2^{\frac{\gamma}{2}}}{\sqrt{\pi}} e^{-\frac{x^{2}}{2}} \Gamma\left(\frac{1+\gamma}{2}\right) {}_{1}F_{1}\left(\frac{1}{2} + \frac{\gamma}{2}, \frac{1}{2}, \frac{x^{2}}{2}\right)$$
$$= \frac{2^{\frac{\gamma}{2}}}{\sqrt{\pi}} \Gamma\left(\frac{1+\gamma}{2}\right) {}_{1}F_{1}\left(-\frac{\gamma}{2}, \frac{1}{2}, -\frac{x^{2}}{2}\right),$$

where $_1F_1(a, b, c)$ stands for the Kummer confluent hypergeometric function; see Gradshteyn and Ryzhik (1994). For the covariance kernel K(s, t) corresponding to the GE statistic $T_{n,w}$ we refer to Móri et al. (2021), while calculation of the eigenvalues λ_j , $j \ge 1$, figuring in (2.1)–(2.2) is carried out using method described in §3.2.

Letting $(\mu(\theta), \sigma^2(\theta))$ be the probability limit of the estimator $(\bar{X_n}, \bar{S}_n^2)$, i.e.,

$$\mu(\theta) = \int_{-\infty}^{\infty} x g_{\theta}(x) dx; \quad \sigma^2(\theta) = \int_{-\infty}^{\infty} (x - \mu(\theta))^2 g_{\theta}(x) dx,$$

we obtain the probability limit figuring in (2.5) for the GE test statistic as

$$\frac{T_{n,w}}{n} \xrightarrow{\mathbb{P}} b(\theta) = \frac{2^{1+\frac{\gamma}{2}}}{\sqrt{\pi}} \Gamma\left(\frac{1+\gamma}{2}\right) \mathbb{E}_{\theta} \left[{}_{1}F_{1}\left(-\frac{\gamma}{2}, \frac{1}{2}, -\frac{(X_{1}-\mu(\theta))^{2}}{2\sigma^{2}(\theta)}\right) \right] - \frac{2^{\gamma}}{\sqrt{\pi}} \Gamma\left(\frac{1+\gamma}{2}\right) - \frac{1}{\sigma(\theta)^{\gamma}} \mathbb{E}_{\theta} |X_{1}-X_{2}|^{\gamma}.$$
(4.4)

In order to calculate the quantity b''(0) figuring in (2.4) we use numerical integration in Wolfram Mathematica facilitated by differentiation under the integral sign, and thereby obtain the expressions

$$\mu'(0) = \int_{-\infty}^{\infty} xh(x)dx; \quad \mu''(0) = \int_{-\infty}^{\infty} xu(x)dx$$

$$\sigma'(0) = \frac{1}{2} \int_{-\infty}^{\infty} x^2h(x)dx$$

$$\sigma''(0) = \frac{1}{2} \int_{-\infty}^{\infty} x^2u(x)dx - \left(\int_{-\infty}^{\infty} xh(x)dx\right)^2 - \frac{1}{4} \left(\int_{-\infty}^{\infty} x^2h(x)dx\right)^2,$$

⁴The original weight function also includes the constant $2\sqrt{\pi}\Gamma(1-(\gamma/2))/(\gamma 2^{\gamma}\Gamma((1+\gamma)/2))$

γ	$g^{(1)}$	$g^{(2)}$	$g^{(3)}$	$g^{(4)}(1,1)$	$g^{(4)}(0.5,1)$	$g^{(4)}(0, 0.5)$
0.1	0.501	0.714	0.843	0.323	0.431	0.630
0.2	0.520	0.734	0.861	0.336	0.447	0.636
0.3	0.538	0.754	0.877	0.349	0.464	0.640
0.4	0.556	0.772	0.892	0.362	0.480	0.643
0.5	0.573	0.790	0.906	0.374	0.496	0.645
0.6	0.590	0.806	0.918	0.387	0.512	0.645
0.7	0.608	0.821	0.930	0.399	0.527	0.645
0.8	0.623	0.837	0.940	0.411	0.542	0.643
0.9	0.639	0.851	0.950	0.423	0.558	0.640
1.0	0.655	0.865	0.958	0.434	0.572	0.636
1.1	0.670	0.877	0.966	0.446	0.586	0.632
1.2	0.685	0.889	0.973	0.457	0.600	0.626
1.3	0.699	0.900	0.978	0.468	0.614	0.620
1.4	0.713	0.911	0.984	0.479	0.628	0.613
1.5	0.727	0.921	0.988	0.490	0.641	0.605
1.6	0.740	0.930	0.991	0.501	0.654	0.596
1.7	0.753	0.939	0.994	0.511	0.666	0.587
1.8	0.765	0.947	0.997	0.521	0.679	0.577
1.9	0.777	0.954	0.998	0.531	0.691	0.567

Table 1: LABE of the energy test for normality

(where $h(x) = \frac{\partial}{\partial \theta} g_{\theta}(x)|_{\theta=0}$ and $u(x) = \frac{\partial^2}{\partial \theta^2} g_{\theta}(x)|_{\theta=0}$), and compute the local approximate Bahadur relative efficiencies (LABEs) of the GE test with respect to the LR test. These efficiences are reported in Table 1.

4.1.2 The BHEP test

The BHEP test may be formulated as in (4.1) with weight function $w(t) = e^{-\gamma t^2}$, $\gamma > 0$, which leads to the convenient expression

$$T_{n,w} = \sqrt{\frac{\pi}{\gamma}} \frac{1}{n} \sum_{j,k=1}^{n} e^{-\frac{(Z_j - Z_k)^2}{4\gamma}} + n\sqrt{\frac{\pi}{1+\gamma}} - 2\sqrt{\frac{2\pi}{1+2\gamma}} \sum_{j=1}^{n} e^{-\frac{Z_j^2}{2+4\gamma}}.$$
(4.5)

The asymptotic null distribution of the BHEP test along with the expression for the covariance kernel K(s, t) may be found in Henze and Wagner (1997), and corresponding eigenvalues have been recently computed by Ebner and Henze (2021b).

The probability limit of the BHEP test is obtained analogously to (4.4) as

$$b(\theta) = \sqrt{\frac{\pi}{1+\gamma}} + \sqrt{\frac{\pi}{\gamma}} \mathbb{E}_{\theta} \left[\exp\left(-\frac{(X_1 - X_2)^2}{4\gamma\sigma^2(\theta)}\right) \right] - 2\sqrt{\frac{2\pi}{1+\gamma}} \mathbb{E}_{\theta} \left[\exp\left(-\frac{(X_1 - \mu(\theta))^2}{(2+4\gamma)\sigma^2(\theta)}\right) \right],$$

while the calculation of efficiencies is also carried out in the same way as in the previous subsection by means of the approximation outlined in §3.2. These efficiency results (LABEs) for the BHEP test are reported in Table 2.

γ	$g^{(1)}$	$g^{(2)}$	$g^{(3)}$	$g^{(4)}(1,1)$	$g^{(4)}(0.5,1)$	$g^{(4)}(0, 0.5)$
0.1	0.477	0.701	0.840	0.302	0.406	0.654
0.2	0.582	0.814	0.929	0.376	0.501	0.676
0.3	0.655	0.879	0.968	0.429	0.568	0.658
0.4	0.710	0.921	0.986	0.471	0.620	0.628
0.5	0.752	0.948	0.992	0.505	0.661	0.593
0.6	0.785	0.967	0.993	0.532	0.695	0.559
0.7	0.812	0.979	0.990	0.555	0.722	0.527
0.8	0.834	0.987	0.986	0.574	0.745	0.497
0.9	0.853	0.993	0.980	0.591	0.764	0.469
1	0.868	0.997	0.974	0.605	0.780	0.443
2	0.941	0.992	0.917	0.681	0.865	0.281
3	0.963	0.975	0.879	0.711	0.896	0.202
4	0.972	0.961	0.855	0.726	0.910	0.158
5	0.977	0.951	0.839	0.735	0.918	0.129
6	0.979	0.942	0.826	0.741	0.923	0.109
7	0.980	0.936	0.817	0.745	0.926	0.094
8	0.981	0.931	0.810	0.746	0.929	0.083
9	0.981	0.926	0.804	0.750	0.930	0.074
10	0.981	0.923	0.799	0.751	0.932	0.067

Table 2: LABE of the BHEP test for normality

4.1.3 Discussion

From Tables 1 and 2 we can see that there is a significant influence of the tuning parameter on the efficiency of both tests. Specifically, in the case of the GE test, the efficiencies generally grow with γ , hence a high value of the tuning parameter (close to the boundary value $\gamma = 2$) can be recommended for this test. On the other hand, the corresponding LABEs of the BHEP test exhibit no such consistent pattern, with the impact on the tuning parameter depending very much on the specific alternative. Specifically for Lehmann and location/scale contamination alternatives higher values of the tuning parameter should be used, while for all other alternatives an "in-between" value in the neighborhood of $\gamma = 1$ yields better efficiency for the corresponding BHEP test. Moreover the aforementioned value seems to be a good compromise, and if one must choose a single test between the GE and BHEP tests, then the latter test with $\gamma = 1$ appears to yield a good overall efficiency. We can also compare the efficiency of the ECF-based tests to the efficiencies of corresponding tests based on the empirical distribution function (EDF) provided by Milošević et al. (2021). In this connection, a close inspection of the corresponding efficiency figures shows a significant advantage of the ECF-based tests over their EDF counterparts.

4.2 Tests for the Logistic distribution

In complete analogy to the GE and BHEP tests formulated as in (1.2) and (4.1), Meintanis (2004) defines a GOF test statistic for the logistic distribution as

$$T_{n,w} = n \int_{-\infty}^{\infty} \left| \phi_n(t) - \frac{\pi t}{\sinh(\pi t)} \right|^2 w(t) \mathrm{d}t, \tag{4.6}$$

with the ECF $\phi_n(t)$ obtained as in (1.3) with X_j replaced by $Z_j = (X_j - \hat{\mu}_n)/\hat{\sigma}_n$, where $(\hat{\mu}_n, \hat{\sigma}_n)$ denote the moment estimators or the ML estimators of the parameters (μ, σ) of the logistic distribution. An explicit test statistic formula corresponding to the weight function $w(t) = e^{-\gamma |t|}, \gamma > 0$, is given by

$$T_{n,w} = \frac{2\gamma}{n} \sum_{j,k=1}^{n} \frac{1}{\gamma^2 + (Z_j - Z_k)^2} - \frac{2}{\pi} \sum_{j=1}^{n} \left[S_{\gamma}^{(1)}(Z_j) - \frac{Z_j^2}{2\pi^2} S_{\gamma}^{(2)}(Z_j) \right] + \frac{n}{\pi} \left[2\zeta_2(1 + \frac{\gamma}{2\pi}) - \frac{\gamma}{\pi} \zeta_3(1 + \frac{\gamma}{2\pi}) \right],$$
(4.7)

where $\zeta_{\gamma}(x) = \sum_{k=0}^{\infty} (k+x)^{-\gamma}$, and

$$S_{\gamma}^{(m)}(x) = \sum_{k=0}^{\infty} \left[\left(\frac{x}{2\pi}\right)^2 + \left(\frac{\gamma + \pi}{2\pi} + k\right)^2 \right]^{-m}.$$

The expression for the covariance kernel K(s,t) may also be found in Meintanis (2004), along with a Monte Carlo study of the power of the test based on $T_{n,w}$ against the classical tests based on the EDF. These results, nicely complemented by results from Gulati and Shapiro (2009), suggest that the test based on (4.7) is an overall competitive test.

Turning to efficiency calculations we obtain the corresponding probability limit (recall (2.5)) as

$$b(\theta) = \mathbb{E}_{\theta} \left[\frac{2\gamma}{\gamma^2 + \left(\frac{X_1 - X_2}{\sigma(\theta)}\right)^2} \right] - \frac{2}{\pi} \mathbb{E}_{\theta} \left[S_{\gamma}^{(1)} \left(\frac{X - \mu(\theta)}{\sigma(\theta)}\right) \right] \\ - \frac{1}{\pi^3} \mathbb{E}_{\theta} \left[\left(\frac{X - \mu(\theta)}{\sigma(\theta)}\right)^2 S_{\gamma}^{(2)} \left(\frac{X - \mu(\theta)}{\sigma(\theta)}\right) \right] + \frac{1}{\pi} \left(2\zeta_2 (1 + \frac{\gamma}{2\pi}) - \frac{\gamma}{\pi} \zeta_3 (1 + \frac{\gamma}{2\pi}) \right),$$

where $\mu(\theta)$ and $\sigma(\theta)$ denote the probability limits of $\hat{\mu}_n$ and $\hat{\sigma}_n$, respectively.

In order to calculate b''(0), we use as before numerical integration in Wolfram Mathematica facilitated by the formulae for the first and second derivatives at zero of $\mu(\theta)$ and $\sigma(\theta)$. In this connection and since for the ML estimators there exist no closed expressions for $\mu(\theta)$ and $\sigma(\theta)$, the necessary derivatives are obtained via the implicit function theorem as

$$\begin{split} \mu'(0) &= 6 \int_{-\infty}^{\infty} \frac{h(x)}{1 + e^{-x}} dx; \ \ \sigma'(0) = \frac{9}{\pi^2 + 3} \int_{-\infty}^{\infty} \frac{(1 - e^{-x})xh(x)}{1 + e^{-x}} dx, \\ \mu''(0) &= 6 \Big(\int \frac{u(x)}{1 + e^{-x}} dx + \frac{1}{6} \mu'(0)\sigma'(0) - 2 \int \frac{(\mu'(0) + x\sigma'(0))e^{-x}h(x)}{(1 + e^{-x})^2} dx \Big), \\ \sigma''(0) &= \frac{9}{3 + \pi^2} \Big(\int xu(x)dx + \frac{1}{2} \mu'(0)^2 - 4\mu'(0)\sigma'(0) + \frac{1}{18}(\pi^2 - 6)\sigma'(0)^2 \\ &- 4 \int \frac{(e^x(x - 1) - 1)\mu'(0) + e^x x^2 \sigma'(0)}{(1 + e^x)^2} h(x)dx - 2 \int \frac{xe^{-x}u(x)}{(1 + e^{-x})} \Big). \end{split}$$

γ	$g^{(1)}$	$g^{(2)}$	$g^{(3)}$	$g^{(4)}(1,1)$	$g^{(4)}(0.5,1)$	$g^{(4)}(0, 0.5)$
0.1	0.274	0.456	0.641	0.468	0.463	0.702
0.2	0.314	0.508	0.710	0.525	0.516	0.759
0.3	0.347	0.548	0.762	0.570	0.558	0.795
0.4	0.378	0.581	0.804	0.607	0.593	0.820
0.5	0.406	0.608	0.839	0.641	0.622	0.837
0.6	0.432	0.632	0.868	0.670	0.648	0.847
0.7	0.457	0.652	0.893	0.695	0.669	0.852
0.8	0.481	0.668	0.913	0.718	0.688	0.854
0.9	0.503	0.683	0.931	0.738	0.704	0.853
1	0.524	0.695	0.945	0.756	0.718	0.850
2	0.693	0.728	0.989	0.844	0.768	0.745
3	0.802	0.673	0.945	0.837	0.728	0.605
4	0.870	0.581	0.876	0.787	0.650	0.473
5	0.908	0.479	0.812	0.722	0.559	0.362
6	0.928	0.381	0.767	0.663	0.473	0.274
7	0.938	0.299	0.745	0.616	0.402	0.210
8	0.943	0.237	0.739	0.584	0.351	0.166
9	0.946	0.194	0.741	0.565	0.315	0.136
10	0.948	0.163	0.744	0.552	0.230	0.116

Table 3: LABE of the test for the logistic distribution (ML estimation)

On the other hand, when $\hat{\mu}_n$ and $\hat{\sigma}_n$ are obtained by the method of moments, and by using the corresponding probability limits

$$\mu(\theta) = \int_{-\infty}^{\infty} xg_{\theta}(x)dx,$$

$$\sigma(\theta) = \frac{\sqrt{3}}{\pi} \left(\int_{-\infty}^{\infty} x^2 g_{\theta}(x)dx - \mu^2(\theta) \right)^{\frac{1}{2}},$$

we obtain the necessary derivatives as

$$\mu'(0) = \int_{-\infty}^{\infty} xh(x)dx; \quad \mu''(0) = \int_{-\infty}^{\infty} xu(x)dx;$$

$$\sigma'(0) = \frac{3}{2\pi^2} \int_{-\infty}^{\infty} x^2h(x)dx;$$

$$\sigma''(0) = \frac{3}{2\pi^2} \int_{-\infty}^{\infty} x^2u(x)dx - \frac{3}{\pi^2} \Big(\int_{-\infty}^{\infty} xh(x)dx\Big)^2 - \frac{3}{4\pi^2} \int_{-\infty}^{\infty} x^2h(x)dx.$$

(recall $h(x) = \frac{\partial}{\partial \theta} g_{\theta}(x)|_{\theta=0}$ and $u(x) = \frac{\partial^2}{\partial \theta^2} g_{\theta}(x)|_{\theta=0}$), and thereby calculate the efficiencies of the test for the logistic distribution based on (4.7). These efficiencies (LABEs) are reported in Table 3 (ML estimation) and Table 4 (moment estimation).

For comparison purposes we also provide in Table 5 corresponding efficiencies for three EDF-based tests, namely the Kolmogorov-Smirnov (KS), the Cramér-von Mises (CM), and the Anderson-Darling (AD) test. The calculations were

γ	$g^{(1)}$	$g^{(2)}$	$g^{(3)}$	$g^{(4)}(1,1)$	$g^{(4)}(0.5,1)$	$g^{(4)}(0, 0.5)$
0.1	0.485	0.695	0.811	0.680	0.693	0.878
0.2	0.539	0.754	0.874	0.741	0.752	0.924
0.3	0.582	0.794	0.915	0.786	0.794	0.947
0.4	0.619	0.824	0.944	0.820	0.825	0.957
0.5	0.651	0.846	0.965	0.846	0.849	0.958
0.6	0.670	0.863	0.980	0.868	0.867	0.954
0.7	0.706	0.875	0.989	0.884	0.880	0.945
0.8	0.729	0.883	0.995	0.890	0.890	0.933
0.9	0.751	0.889	0.998	0.908	0.896	0.919
1	0.770	0.891	0.998	0.916	0.901	0.904
2	0.898	0.841	0.931	0.910	0.863	0.724
3	0.941	0.729	0.816	0.828	0.760	0.559
4	0.773	0.504	0.580	0.601	0.532	0.355
5	0.581	0.323	0.387	0.405	0.348	0.213
6	0.422	0.202	0.254	0.266	0.222	0.126
7	0.304	0.127	0.169	0.176	0.142	0.076
8	0.221	0.081	0.114	0.118	0.092	0.047
9	0.161	0.052	0.079	0.080	0.061	0.029
10	0.120	0.034	0.055	0.056	0.041	0.019
	(1)	(2)	(2)	(4) ((1) (2, 1, 1)	(4) (5 5 11)
Test	$g^{(1)}$				$g^{(4)}(0.5,1)$	
KS	0.191	0.312	0.455	0.329	0.318	0.435
СМ	0.475	0.677	0.920	0.715	0.693	0.866
AD	0.754	0.640	0.999	0.808	0.697	0.718

Table 4: LABE of the test for the logistic distribution (moment estimation)

Table 5: LABE of EDF tests for the logistic distribution

carried out using the same method used by Milošević et al. (2021) for normality tests, while the covariance function of the corresponding empirical process is available from Stephens (1979).

4.2.1 Discussion

We note that this is the first time that efficiency results for a test for the composite hypothesis of GOF to the logistic distribution are obtained. In particular, there exist no results for the Bahadur efficiencies for classical GOF tests in the case of the logistic distribution, or in fact for any GOF test for this distribution in the case of the "fully" composite null hypothesis, i.e. when both the location and scale parameters are unknown.

When ECF based tests are compared among themselves, from Tables 3 and 4 it may be inferred that the efficiencies vary considerably between estimation methods, and that the variant of the test which uses the method of moments is somewhat more efficient than its counterpart based on ML estimation, a finding that is in line with the Monte Carlo

results of Meintanis (2004). This phenomenon of (generally) less efficient estimators, such as the moment estimators, producing more powerful tests has already been pointed out by Gürtler and Henze (2000) under similar circumstances, and dates back at least to Drost et al. (1990). Turning to the tuning parameter we observe that its impact on test performance is noticeable, and that, by taking into account all considered alternatives, we can recommend $\gamma = 3$ in the case of ML estimation and $\gamma = 1$ in the case of moment estimation as overall good choices.

As far as classical tests are concern, we notice from Table 5 that the AD test is more efficient than the other EDF-based tests and competitive with the ECF-based tests. Nevertheless the moment-based ECF test appears to be preferable overall against the AD in the neighborhood of the suggested value $\gamma = 1$. The other EDF-based tests are also less efficient than the ECF tests for the recommended value of the tuning parameter γ .

4.3 Tests for the exponential distribution

In extensive Monte Carlo studies (see for instance Henze and Meintanis (2005) and Grané and Fortiana (2011)), certain tests based on the ECF are found to be competitive for testing GOF to the exponential distribution. One such test is also a weighted L2–type test, but the corresponding test statistic admits a slightly different formulation from that in (1.2). Specifically we have

$$T_{n,w} = n \, \int_{-\infty}^{\infty} \left(|\phi_n(t)|^2 - C_n(t) \right)^2 w(t) \mathrm{d}t, \tag{4.8}$$

with $|\phi_n(t)|$ denoting the modulus and $C_n(t)$ the real part of the ECF obtained from (1.3) by replacing X_j by

$$Z_j = X_j / \bar{X}_n, \ j = 1, ..., n.$$
(4.9)

As weight functions Henze and Meintanis (2005) suggest $w_{\beta}(t) = e^{-\gamma |t|^{\beta}}$, $\gamma > 0$, $\beta = 1, 2$, and provide explicit expressions for the resulting test statistics. The corresponding kernel K(s, t) figuring in (2.2) may be computed as

$$K(s,t) = \frac{s^2 t^2 (1+s^2+t^2)}{(1+s^2)(1+t^2)(1+(s-t)^2)(1+(s+t)^2)}.$$

For the weight function $w_1(t) = e^{-\gamma |t|}$, the probability limit figuring in (2.5) is given by

$$b_{1}(\theta) = \gamma \mathbb{E}_{\theta} \left[\frac{1}{\gamma^{2} + (\frac{X_{1} - X_{2}}{\sigma(\theta)})^{2}} + \frac{1}{\gamma^{2} + (\frac{X_{1} + X_{2}}{\sigma(\theta)})^{2}} \right] - 2\gamma \mathbb{E}_{\theta} \left[\frac{1}{\gamma^{2} + (\frac{X_{1} - X_{2} - X_{3}}{\sigma(\theta)})^{2}} + \frac{1}{\gamma^{2} + (\frac{X_{1} - X_{2} + X_{3}}{\sigma(\theta)})^{2}} \right] + \gamma \mathbb{E}_{\theta} \left[\frac{1}{\gamma^{2} + (\frac{X_{1} - X_{2} - X_{3} + X_{4}}{\sigma(\theta)})^{2}} + \frac{1}{\gamma^{2} + (\frac{X_{1} - X_{2} + X_{3} - X_{4}}{\sigma(\theta)})^{2}} \right]$$

while for $w_2(t) = e^{-\gamma t^2}$, the same probability limit is given by

$$b_{2}(\theta) = \frac{1}{2}\sqrt{\frac{\pi}{\gamma}} \mathbb{E}_{\theta} \left[\exp\left(\frac{(X_{1} - X_{2})^{2}}{4\gamma\sigma^{2}(\theta)}\right) + \left(\exp\left(\frac{(X_{1} + X_{2})^{2}}{4\gamma\sigma^{2}(\theta)}\right)\right] \\ - \sqrt{\frac{\pi}{\gamma}} \mathbb{E}_{\theta} \left[\exp\left(\frac{(X_{1} - X_{2} - X_{3})^{2}}{4\gamma\sigma^{2}(\theta)}\right) + \left(\exp\left(\frac{(X_{1} - X_{2} + X_{3})^{2}}{4\gamma\sigma^{2}(\theta)}\right)\right] \\ + \frac{1}{2}\sqrt{\frac{\pi}{\gamma}} \mathbb{E}_{\theta} \left[\exp\left(\frac{(X_{1} - X_{2} - X_{3} + X_{4})^{2}}{4\gamma\sigma^{2}(\theta)}\right) + \left(\exp\left(\frac{(X_{1} - X_{2} + X_{3} - X_{4})^{2}}{4\gamma\sigma^{2}(\theta)}\right)\right],$$

where, recall, $\sigma(\theta) = \int_{-\infty}^{\infty} x g_{\theta}(x) dx$.

As far as alternatives are concerned, the exponential distribution has some common close alternatives. Specifically we consider the following alternatives:

γ	Weibull	Gamma	Makeham	LFR	ME(3)	ME(6)
0.1	0.502	0.619	0.404	0.125	0.585	0.831
0.2	0.552	0.625	0.470	0.155	0.669	0.841
0.3	0.584	0.625	0.523	0.183	0.724	0.834
0.4	0.610	0.632	0.580	0.203	0.763	0.819
0.5	0.625	0.627	0.621	0.231	0.792	0.802
0.6	0.637	0.626	0.657	0.252	0.813	0.784
0.7	0.649	0.626	0.688	0.275	0.829	0.767
0.8	0.659	0.624	0.713	0.295	0.842	0.750
0.9	0.671	0.622	0.737	0.315	0.852	0.735
1	0.676	0.620	0.757	0.334	0.859	0.720
2	0.722	0.596	0.881	0.494	0.874	0.616
3	0.738	0.573	0.925	0.611	0.850	0.553
4	0.741	0.553	0.938	0.699	0.821	0.510
5	0.737	0.534	0.936	0.765	0.792	0.478
6	0.730	0.518	0.928	0.815	0.766	0.453
7	0.721	0.504	0.917	0.853	0.743	0.434
8	0.713	0.487	0.905	0.882	0.723	0.418
9	0.704	0.478	0.893	0.905	0.705	0.404
10	0.696	0.461	0.881	0.923	0.690	0.393

Table 6: LABE of the test for exponentiality with weight function w_1

• the Weibull distribution with density

$$g_{\theta}(x) = e^{-x^{1+\theta}} (1+\theta) x^{\theta}, \theta > 0, x \ge 0;$$

• the gamma distribution with density

$$g_{\theta}(x) = \frac{x^{\theta} e^{-x}}{\Gamma(\theta+1)}, \theta > 0, x \ge 0;$$

• the Makeham distribution with density

$$g_{\theta}(x) = e^{-x - \theta e^x} (1 + \theta e^x), \theta > 0, x \ge 0;$$

• the linear failure rate (LFR) distribution with density

$$g_{\theta}(x) = e^{-x - \theta \frac{x^2}{2}} (1 + \theta x), \theta > 0, x \ge 0;$$

• the mixture of exponential distributions with negative weights (ME(β)) with density

$$g_{\theta}(x;\beta) = (1+\theta)e^{-x} - \theta\beta e^{-\beta x}, \theta \in \left(0, \frac{1}{\beta - 1}\right], x \ge 0;$$

These alternatives are also used by e.g. Cuparić et al. (2019); Milošević (2016); Milošević and Obradović (2016); Nikitin and Volkova (2016). The resulting efficiencies are calculated by following analogous steps as in the previous sections and are reported in Table 6 and Table 7, for the test based on $w_1(t) = e^{-\gamma |t|}$ and $w_2(t) = e^{-\gamma |t|^2}$, respectively.

γ	Weibull	Gamma	Makeham	LFR	ME(3)	ME(6)
0.1	0.581	0.585	0.580	0.198	0.756	0.749
0.2	0.619	0.584	0.676	0.261	0.807	0.698
0.3	0.639	0.581	0.735	0.312	0.828	0.665
0.4	0.657	0.579	0.773	0.350	0.838	0.640
0.5	0.670	0.576	0.802	0.384	0.843	0.621
0.6	0.679	0.573	0.824	0.413	0.845	0.605
0.7	0.687	0.571	0.842	0.439	0.845	0.592
0.8	0.693	0.568	0.856	0.463	0.844	0.580
0.9	0.698	0.566	0.867	0.485	0.842	0.570
1	0.703	0.564	0.877	0.504	0.840	0.562
2	0.722	0.544	0.918	0.643	0.811	0.505
3	0.725	0.529	0.925	0.724	0.785	0.474
4	0.723	0.516	0.921	0.778	0.763	0.452
5	0.719	0.505	0.915	0.816	0.746	0.436
6	0.714	0.497	0.908	0.844	0.731	0.424
7	0.710	0.489	0.900	0.866	0.718	0.414
8	0.705	0.481	0.892	0.892	0.708	0.406
9	0.700	9.476	0.885	0.885	0.698	0.399
10	0.697	0.467	0.879	0.910	0.690	0.393

Table 7: LABE of the test for exponentiality with weight function w_2

4.3.1 Discussion

From Tables 6 and 7 one can notice that the type of monotonicity of the efficiencies with respect to the tuning parameter γ varies amongst alternatives. However, for any fixed alternative, both tests exhibit the same behaviour, and if we compare maximal reached efficiencies, the test with weight function $w_1(\cdot)$ appears to have a slight edge. Next we compare the efficiencies obtained here with the efficiencies reported by Cuparić et al. (2022) for a wide variety of exponentiality tests and for four of the alternatives considered herein (e.g., Weibull, Gamma, LFR and ME(3)). Specifically the tests based on $T_{n,w}$ seem to be less efficient than some of the best tests considered by Cuparić et al. (2022). When restricting comparison to the classical tests based on the EDF, we see that the new tests are more efficient than the KS test, but overall less efficient than the CM and AD tests, with the exception of the LFR alternative where, with proper choice of γ , our tests compete well with these two tests.

5 Tests for the bivariate normal distribution

We consider the problem of testing the null hypothesis \mathcal{H}_0 that the sample comes from a bivariate normal distribution, using the BHEP and GE tests presented in Section 4.1. Efficiencies are calculated for two cases: The simple hypothesis of known mean/covariance, that the law of X is $\mathcal{N}_2(0, I)$, and the hybrid case of unknown mean/known covariance, i.e. that the law of X is $\mathcal{N}_2(\mu, I)$, where $\mathcal{N}_p(\mu, \Sigma)$ denotes the multivariate normal distribution with mean vector μ and covariance matrix equal to Σ , and I stands for the identity matrix in the corresponding dimension. The covariance kernels of the test statistics are available from Móri et al. (2021) (GE test) and Henze and Wagner (1997) (BHEP test), while for the simple hypothesis case with $\gamma = 1/2$, eigenvalues of the BHEP test in arbitrary dimension may be obtained from Baringhaus (1996).⁵

The method of calculation of efficiencies is the same as for univariate tests. Specifically, consider an alternative with density $g(x, y; \theta)$, and for small θ , compute the double KL distance of this alternative to the null hypothesis, as

• simple hypothesis case:

$$2K(\theta) = \left(\iint_{\mathbb{R}^2} \frac{h^2(x,y)}{f_0(x,y)} \mathrm{d}x \mathrm{d}y\right) \theta^2 + o(\theta^2),$$

• estimated mean case:

$$2K(\theta) = \left(\iint_{\mathbb{R}^2} \frac{\left(h(x,y) - f_0(x,y)(x\mu'_X(0) + y\mu'_Y(0))\right)^2}{f_0(x,y)} \mathrm{d}x\mathrm{d}y\right) \theta^2 + o(\theta^2),$$

where

$$\mu_X(\theta) = \iint_{\mathbb{R}^2} xg(x, y; \theta) \mathrm{d}x \mathrm{d}y; \ \mu_Y(\theta) = \iint_{\mathbb{R}^2} yg(x, y; \theta) \mathrm{d}x \mathrm{d}y$$

and $f_0(\cdot, \cdot)$ denotes the density of the standard normal distribution. The proof is analogous to that of Theorem 3.1, so we omit it here.

The multidimensional version of the eigenvalue approximation procedure described in Section 3 is computationally very complex, hence here we obtained the largest eigenvalues using the Monte Carlo procedure described in Móri et al. (2021). In this connection we have noticed that the variance of the Monte Carlo estimator of the eigenvalue is small in the case of BHEP test, which makes the obtained values reliable. On the other hand, for the GE test, the variance is significant, making the approximation not so accurate. In addition, this problem drastically increases with the increase of the tuning parameter γ , therefore we excluded the cases when $\gamma > 1$ from the study.

In order to introduce our alternatives write $f(\mu_1, \mu_2, \sigma_1, \sigma_2, \rho)$ for the bivariate normal density with mean vector $\mu = (\mu_1, \mu_2)^{\top}$ and covariance matrix Σ with elements $\sigma_{11} = \sigma_1^2$, $\sigma_{22} = \sigma_2^2$, and $\sigma_{12} = \sigma_{21} = \rho$. With this notation we consider the following alternatives for the case of the simple hypothesis tests:

- location alternative $g_l(\theta) = f(\theta, 0, 1, 1, 0)$
- correlation alternative $g_c(\theta) = f(0, 0, 1, 1, \theta)$
- single scale alternative $g_{s1}(\theta) = f(0, 0, 1 \theta, 1, 0)$
- double scale alternative $g_{s2}(\theta) = f(0, 0, 1 \theta, 1 \theta, 0)$
- contamination alternative

$$g_{cn}(\mu_1, \mu_2, \sigma_1, \sigma_2, \rho, \theta) = (1 - \theta)f(0, 0, 1, 1, 0) + \theta f(\mu_1, \mu_2, \sigma_1, \sigma_2, \rho),$$

for the specific choices of parameters given in Table 8.

⁵As a numerical confirmation, we have compared the efficiencies obtained by using Baringhaus' eigenvalues with those using the Monte Carlo method and found them to be quite close in all cases

	μ_1	μ_2	σ_1	σ_2	ρ
$g_{cn}^{(1)}$	0.1	0	1	1	0
$g_{cn}^{(2)}$	0.5	0	1	1	0
$g_{cn}^{(3)}$	0.9	0	1	1	0
$g_{cn}^{(4)}$	1.5	0	1	1	0
$g_{cn}^{(5)}$	0	0	1	1	0.1
$g_{cn}^{(6)}$	0	0	1	1	0.5
$g_{cn}^{(7)}$	0	0	1	1	0.9
$g_{cn}^{(8)}$	0	0	0.5	1	0
$g_{cn}^{(9)}$	0	0	0.7	1	0
$g_{cn}^{(10)}$	0	0	0.9	1	0
$g_{cn}^{(11)}$	0	0	1.1	1	0

Table 8: Parameters of the contaminating distribution

5.1 Discussion

Table 9 (simple hypothesis case) and Table 10 (estimated mean case)⁶ report efficiency results for the GE and BHEP tests for bivariate normality. To facilitate comparison, for each alternative we report in italics the best efficiency of the test under discussion against γ , while a bold entry indicates the best efficiency value across both tests.

For the GE test the monotonicity is clear, since better efficiencies are obtained as γ increases, in both the simple and the hybrid case test, a behaviour that mimics the univariate tests. On the other hand, also analogously to the univariate case, no such pattern is visible for the BHEP test in the simple hypothesis case, as efficiencies are better at the one or the other end of the tuning parameter interval, or even in-between values of γ , depending on the type of alternative. Things are somewhat more clear though for the BHEP test with estimated mean, and specifically in this case better efficiencies are observed for larger values of γ , nearly uniformly over alternatives.

When comparing the GE and BHEP tests with each other, one cannot suggest a single test as being uniformly more efficient over all alternatives. Nevertheless in the simple hypothesis case, the GE test with $\gamma = 1$ appears to perform better against location contamination alternatives, while the BHEP test with $\gamma = 0.25, 0.50$ shows higher efficiency in nearly all other cases of alternatives. With a few exceptions, the BHEP test with $\gamma = 2$ seems to be also preferable over the GE test in the case of testing for bivariate normality with unknown mean.

6 Conclusion

We consider test optimality in the Bahadur sense for certain ECF-based GOF tests for the normal, the logistic and the exponential distribution. In the case of testing for normality we compare the efficiencies of the celebrated generalized energy and BHEP tests, and we offer suggestions as to which tests should be used on the basis of our efficiency comparisons, in the univariate as well as the bivariate case, with or without estimated parameters. In the case of testing for the logistic distribution efficiency comparisons of an ECF-based test against classical tests based on the EDF are

⁶For the hybrid case we only present results for contamination alternatives as location alternative are clearly excluded and efficiencies for the correlation and scale alternatives coincide with those of the simple hypothesis case

Test		GE			BHEP			
γ	0.5	0.7	1	0.25	0.5	1	2	
g_l	0.285	0.564	0.961	0.631	0.762	0.860	0.927	
g_c	0.054	0.092	0.119	0.252	0.254	0.215	0.154	
g_{s1}	0.080	0.138	0.180	0.379	0.381	0.323	0.232	
g_{s2}	0.106	0.184	0.240	0.505	0.508	0.430	0.309	
$g_{cn}^{(1)}$	0.284	0.562	0.956	0.627	0.735	0.853	0.918	
$g_{cn}^{(2)}$	0.248	0.492	0.840	0.542	0.637	0.743	0.808	
$g_{cn}^{(3)}$	0.178	0.355	0.609	0.378	0.449	0.531	0.582	
$g_{cn}^{(4)}$	0.068	0.137	0.238	0.134	0.163	0.200	0.225	
$g_{cn}^{(5)}$	0.053	0.092	0.119	0.253	0.245	0.213	0.150	
$g_{cn}^{(6)}$	0.046	0.078	0.101	0.237	0.215	0.172	0.117	
$g_{cn}^{(7)}$	0.019	0.031	0.038	0.110	0.082	0.055	0.033	
$g_{cn}^{(8)}$	0.098	0.159	0.193	0.588	0.454	0.310	0.180	
$g_{cn}^{(9)}$	0.102	0.170	0.213	0.563	0.486	0.369	0.233	
$g_{cn}^{(10)}$	0.090	0.156	0.199	0.451	0.425	0.355	0.245	
$g_{cn}^{(11)}$	0.067	0.116	0.154	0.303	0.303	0.275	0.206	

Table 9: LABE for the GE and BHEP tests-simple hypothesis

Table 10: LABE for the GE and BHEP tests - estimated mean

Test		GE		BHEP			
γ	0.5	0.7	1	0.25	0.5	1	2
$g_{cn}^{(1)}$	0.218	0.366	0.641	0.455	0.553	0.637	0.700
$g_{cn}^{(2)}$	0.200	0.330	0.563	0.409	0.499	0.577	0.636
$g_{cn}^{(3)}$	0.155	0.257	0.440	0.313	0.386	0.453	0.505
$g_{cn}^{(4)}$	0.075	0.126	0.217	0.146	0.184	0.222	0.258
$g_{cn}^{(5)}$	0.147	0.243	0.412	0.305	0.369	0.423	0.465
$g_{cn}^{(6)}$	0.128	0.208	0.348	0.285	0.323	0.348	0.364
$g_{cn}^{(7)}$	0.054	0.083	0.130	0.137	0.124	0.111	0.103
$g_{cn}^{(8)}$	0.269	0.422	0.667	0.710	0.683	0.614	0.546
$g_{cn}^{(9)}$	0.281	0.451	0.736	0.680	0.731	0.731	0.704
$g_{cn}^{(10)}$	0.250	0.409	0.687	0.545	0.639	0.704	0.740
$g_{cn}^{(11)}$	0.185	0.309	0.531	0.363	0.455	0.544	0.624

reported for moment as well as maximum likelihood estimation of parameters, whereas for testing exponentiality we compare our efficiency findings with those of Cuparić et al. (2022) for a wide range of alternative tests, including classical ones. Overall ECF tests appear to compare well and often outperform competitors. Moreover as ECF–based tests involve a tuning parameter, the results reported herein may be used in determining which value of this parameter should be employed by the user, a problem that in various forms occupies researchers to this date; see for instance the contributions of Ebner and Henze (2021a), Ebner and Henze (2021b), Tenreiro (2019), and Allison and Santana (2015).

Appendix

Proof of Theorem 3.1. Let $\mu = \mu(\theta)$ and $\sigma = \sigma(\theta)$ be the values that minimize (3.1). Then, differentiating (3.1) with respect to θ we get

$$\begin{split} K'(\theta) &= \int \log g_{\theta}(x) g'_{\theta}(x) dx - \int \frac{1}{f(\frac{x-\mu(\theta)}{\sigma(\theta)})} f'\Big(\frac{x-\mu(\theta)}{\sigma(\theta)}\Big)\Big(\frac{x-\mu(\theta)}{\sigma(\theta)}\Big)' g_{\theta}(x) \\ &+ \frac{\sigma'(\theta)}{\sigma(\theta)} - \int \log f\Big(\frac{x-\mu(\theta)}{\sigma(\theta)}\Big) g'_{\theta}(x) dx. \end{split}$$

It is easy to show that K'(0) = 0. Differentiating (3.1) once more we obtain at $\theta = 0$,

$$\begin{split} K''(0) &= \int \frac{h^2(x)}{f(x)} dx + \int \log f(x)u(x)dx + \sigma''(0) - (\sigma'(0))^2 \\ &+ \int \frac{1}{f(x)} (f'(x))^2 (\mu'(0) + x\sigma'(0))^2 dx - \int f''(x)(\mu'(0) + x\sigma'(0))^2 dx \\ &- \int f'(x)(-\mu''(0) + 2\mu'(0)\sigma'(0) + x \left(2\sigma'(0)^2 - \sigma''(0)\right)) dx \\ &+ \int \frac{1}{f(x)} f'(x)(\mu'(0) + x\sigma'(0))h(x)dx \\ &+ \int \frac{1}{f(x)} f'(x)(\mu'(0) + x\sigma'(0))h(x)dx - \int \log f(x)u(x)dx, \end{split}$$

where $h(x) = \frac{\partial}{\partial \theta} g_{\theta}(x)|_{\theta=0}$ and $u(x) = \frac{\partial^2}{\partial \theta^2} g_{\theta}(x)|_{\theta=0}$. Rearranging terms in the above expression, and expanding $K(\theta)$ into a Maclaurin expansion completes the proof.

References

- Allison, J. and L. Santana (2015). On a data-dependent choice of the tuning parameter appearing in certain goodnessof-fit tests. *Journal of Statistical Computation and Simulation* 85(16), 3276–3288.
- Bahadur, R. R. (1971). Some limit theorems in statistics. SIAM.
- Baringhaus, L. (1996). Fibonacci numbers, Lucas numbers and integrals of certain Gaussian processes. *Proceeding* of the American Mathematical Society 124(12), 3875–3884.
- Božin, V., B. Milošević, Ya. Yu. Nikitin, and M. Obradović (2020). New characterization-based symmetry tests. Bulletin of the Malaysian Mathematical Sciences Society 43(1), 297–320.
- Cuparić, M., B. Milošević, and M. Obradović (2019). New L²-type exponentiality tests. SORT-Statistics and Operations Research Transactions 43(1), 25–50.

- Cuparić, M., B. Milošević, and M. Obradović (2022). New consistent exponentiality tests based on V-empirical Laplace transforms with comparison of efficiencies. *Revista de la Real Academia de Ciencias Exactas, Físicas y Naturales. Serie A. Matemáticas 116*(42), 1–26.
- Drost, F., W. Kallenberg, and J. Oosterhoff (1990). The power of EDF tests of fit under non-robust estimation of nuisance parameters. *Statistics & Decisions* 8, 167–182.
- Ebner, B. and N. Henze (2021a). Bahadur efficiencies of the Epps–Pulley test for normality. *Zapiski Nauchnykh Seminarov POMI 501*, 302–314.
- Ebner, B. and N. Henze (2021b). On the eigenvalues associated with the limit null distribution of the Epps–Puley test for normality. *arXiv* 2109.04897v1.
- Epps, T. and L. Pulley (1983). A test for normality based on the empirical characteristic function. *Biometrika* 70(3), 723–726.
- Gradshteyn, I. and I. Ryzhik (1994). Tables of Integrals, Series, and Products. Academic Press, New York.
- Grané, A. and J. Fortiana (2011). A directional test of exponentiality based on maximum correlations. *Metrika* 73(2), 255–274.
- Grané, A. and A. Tchirina (2013). Asymptotic properties of goodness–of–fit test based on maximum correlations. *Statistics* 47(1), 202–205.
- Gulati, S. and S. Shapiro (2009). A new goodness of fit test for the logistic distribution. *Journal of Statistical Theory and Practice* 3(3), 567–576.
- Gürtler, N. and N. Henze (2000). Goodness-of-fit tests for the Cauchy distribution based on the empirical characteristic function. *Annals of the Institute of Statistical Mathematics* 52(2), 267–286.
- Henze, N. and S. G. Meintanis (2005). Recent and classical tests for exponentiality: a partial review with comparisons. *Metrika* 61(1), 29–45.
- Henze, N. and T. Wagner (1997). A new approach to the BHEP tests for multivariate normality. *Journal of Multivariate Analysis* 62(1), 1–23.
- Jones, M. C. (2015). On families of distributions with shape parameters. *International Statistical Review* 83(2), 175–192.
- Kato, T. (2013). Perturbation Theory for Linear Operators, Volume 132. Springer Science & Business Media.
- Ley, C. (2015). Flexible modelling in statistics: past, present and future. *Journal de la Société Française de Statistique 156*(1), 76–96.
- Meintanis, S. G. (2004). Goodness-of-fit tests for the logistic distribution based on empirical transforms. *Sankhyā: The Indian Journal of Statistics* 66(2), 306–326.
- Meintanis, S. G. and J. Swanepoel (2007). Bootstrap goodness-of-fit tests with estimated parameters based on empirical transforms. *Statistics & Probability Letters* 77(10), 1004–1013.
- Milošević, B. (2016). Asymptotic efficiency of new exponentiality tests based on a characterization. *Metrika* 79(2), 221–236.
- Milošević, B. and M. Obradović (2016). Some characterization based exponentiality tests and their Bahadur efficiencies. *Publications de L'Institut Mathematique 100*(114), 107–117.

- Milošević, B., Ya. Yu. Nikitin, and M. Obradović (2021). Bahadur efficiency of EDF based normality tests when parameters are estimated. *Zapiski Nauchnykh Seminarov POMI 501*, 203–217.
- Móri, T. F., G. J. Székely, and M. L. Rizzo (2021). On energy tests of normality. *Journal of Statistical Planning and Inference 213*, 1–15.
- Nikitin, Ya. Yu. (1995). Asymptotic Efficiency of Nonparametric Tests. Cambridge University Press, New York.
- Nikitin, Ya. Yu. and I. Peaucelle (2004). Efficiency and local optimality of nonparametric tests based on U- and V-statistics. *Metron* 62(2), 185–200.
- Nikitin, Ya. Yu. and A. V. Tchirina (1996). Bahadur efficiency and local optimality of a test for the exponential distribution based on the Gini statistic. *Journal of the Italian Statistical Society* 5(1), 163–175.
- Nikitin, Ya. Yu. and K. Yu. Volkova (2016). Efficiency of exponentiality tests based on a special property of exponential distribution. *Mathematical Methods of Statistics* 25(1), 54–66.
- Stephens, M. A. (1979). Tests of fit for the logistic distribution based on the empirical distribution function. *Biometrika* 66(3), 591–595.
- Székely, G. J. and M. L. Rizzo (2005). A new test for multivariate normality. *Journal of Multivariate Analysis 93*(1), 58–80.
- Székely, G. J. and M. L. Rizzo (2013). Energy statistics: A class of statistics based on distances. *Journal of Statistical Planning and Inference 143*(8), 1249–1272.
- Tenreiro, C. (2009). On the choice of the smoothing parameter for the BHEP goodness-of-fit test. *Computational Statistics & Data Analysis 53*(4), 1038–1053.
- Tenreiro, C. (2019). On the automatic selection of the tuning parameter appearing in certain families of goodness–of– fit tests. *Journal of Statistical Computation and Simulation 89*(10), 1780–1797.