

Signature of the gluon orbital angular momentum

Shohini Bhattacharya,^{1,*} Renaud Boussarie,^{2,†} and Yoshitaka Hatta^{1,3,‡}

¹*Physics Department, Brookhaven National Laboratory, Upton, NY 11973, USA*

²*CPHT, CNRS, Ecole Polytechnique, Institut Polytechnique de Paris, 91128 Palaiseau, France*

³*RIKEN BNL Research Center, Brookhaven National Laboratory, Upton, NY 11973, USA*

We propose a novel observable for the experimental detection of the gluon orbital angular momentum (OAM) that constitutes the proton spin sum rule. We consider longitudinal double spin asymmetry in exclusive dijet production in electron-proton scattering and demonstrate that the $\cos\phi$ azimuthal angle correlation between the scattered electron and proton is a sensitive probe of the gluon OAM at small- x and its interplay with the gluon helicity. We also present a numerical estimate of the cross section for the kinematics of the Electron-Ion Collider.

1. Introduction—After more than 20 years of operation, the Relativistic Heavy Ion Collider (RHIC) spin program at Brookhaven National Laboratory has revealed that the gluon helicity contribution ΔG to the proton spin sum rule

$$\frac{1}{2} = \frac{1}{2}\Delta\Sigma + \Delta G + L_q + L_g, \quad (1)$$

is nonvanishing and likely sizable [1–5]. Together with the known quark helicity contribution $\Delta\Sigma \sim 0.3$, the result indicates that parton helicities account for a significant fraction of the proton spin. Yet, there still remain huge uncertainties about the small- x contribution to $\Delta G = \int_0^1 dx \Delta G(x)$ defined as the first moment of the polarized gluon distribution function. Resolving this issue is one of the major goals of the future Electron-Ion Collider (EIC) [6].

Another obvious goal of the EIC is to measure the orbital angular momentum (OAM) of quarks and gluons $L_{q,g}$. However, progress in this direction is relatively slow although there have been continuing theory efforts (for recent works, see, e.g., [7–10]). Currently, there does not seem to be a consensus in the community about which observables need to be measured at the EIC in order to constrain $L_{q,g}$. This is so even after the first wave of proposals for experimental observables appeared several years ago [10–14] (see also an earlier attempt [15]). These works exploit the known connection [16–18] between parton OAMs and the Wigner distributions [19], or equivalently, the generalized transverse momentum dependent distributions (GTMDs). However, the required processes typically involve very exclusive final states which are challenging to measure. Besides, observables are often related to GTMDs via complicated multi-dimensional convolutions even at the leading order. Clearly, more theoretical efforts are needed to increase the accuracy of predictions or devise new observables better suited for the purpose.

With this in mind, we take a fresh look at the process considered in Refs. [11, 12]. There, the authors have

proposed to measure longitudinal single spin asymmetry (SSA) in exclusive dijet production in electron-proton collisions $ep \rightarrow \gamma^* p \rightarrow jjp'$ where the incoming proton is longitudinally polarized. It has been shown that the following angular-dependent part of the cross section

$$d\sigma^{h_p} \sim h_p \sin(\phi_{q_\perp} - \phi_{\Delta_\perp})(z - \bar{z}) \Im(A_2 A_3^*), \quad (2)$$

is an experimental probe of the gluon OAM. $h_p = \pm 1$ is the proton helicity and z ($\bar{z} = 1 - z$) is the momentum fraction of the virtual photon carried by the quark (antiquark) jet. ϕ_{q_\perp} is the azimuthal angle of the relative transverse momentum of the two jets $q_\perp = q_{1\perp} - q_{2\perp}$ in a frame in which the proton and virtual photon are collinear, and ϕ_{Δ_\perp} is that of the scattered proton. A_2 and A_3 are certain twist-two and twist-three amplitudes, respectively, and the latter is sensitive to the gluon OAM. As is familiar in the context of transverse single spin asymmetry, one takes the imaginary part of their interference terms.

In this paper, we propose a new observable for the gluon OAM by implementing two major changes in the above proposal. First, we consider *double* spin asymmetry (DSA) in dijet production $ep \rightarrow e' jjp'$ where both the electron and incoming proton are longitudinally polarized. The outgoing lepton must be tagged and its azimuthal angle ϕ_{l_\perp} measured. Electron polarization brings in an extra factor of i to the cross section, so this time one takes the real part of the interference terms. The formula we shall arrive at is

$$d\sigma^{h_p h_l} \sim h_p h_l \cos(\phi_{l_\perp} - \phi_{\Delta_\perp}) \Re(A'_2 A'_3{}^*), \quad (3)$$

where $h_l = \pm 1$ is the electron helicity. Experimentally, the term (3) can be isolated by forming the linear combination $d\sigma^{++} - d\sigma^{+-} - d\sigma^{-+} + d\sigma^{--}$. Unlike in (2), the prefactor does not vanish for symmetric jet configurations ($z = 1/2$), a fact which will turn out to be important. We shall argue that DSA (3) is more advantageous than SSA (2) from both theoretical and practical points of view. Second, we point out that there is another contribution to the asymmetry coming from the gluon helicity generalized parton distribution (GPD). Such a contribution was overlooked in [11, 12], but is parameterically as important as that from the OAM. We shall perform the leading order calculation

* sbhattach@bnl.gov

† renaud.boussarie@polytechnique.edu

‡ yhatta@bnl.gov

of both contributions and numerically evaluate them. The result demonstrates that DSA in dijet production is a unique observable which allows us to directly probe into the gluon OAM L_g and its interplay with the gluon helicity ΔG .

2. Orbital angular momentum and GTMDs—Let us first quickly recapitulate the connection between GTMDs and parton OAMs. Following [20, 21], we parameterize the leading-twist gluon GTMDs as

$$\begin{aligned} & x f_g(x, \xi, \tilde{k}_\perp, \tilde{\Delta}_\perp) \\ &= \int \frac{d^3 z}{(2\pi)^3 P^+} e^{i x P^+ z^- - i \tilde{k}_\perp \cdot z_\perp} \langle p' | F_a^{+i}(-z/2) F_a^{+i}(z/2) | p \rangle \\ &= \frac{1}{2M} \bar{u}(p') \left[F_{1,1} + i \frac{\sigma^{j+}}{P^+} (\tilde{k}_\perp^j F_{1,2} + \tilde{\Delta}_\perp^j F_{1,3}) \right. \\ &\quad \left. + i \frac{\sigma^{ij} \tilde{k}_\perp^i \tilde{\Delta}_\perp^j}{M^2} F_{1,4} \right] u(p), \end{aligned} \quad (4)$$

where $P^\mu = \frac{p+p'}{2}$, $\Delta^\mu = p'^\mu - p^\mu$ and $\xi = (p^+ - p'^+)/2P^+$. $i, j = 1, 2$ are two-dimensional vector indices. All the GTMDs are a function of $x, \xi, \tilde{k}_\perp^2, \tilde{\Delta}_\perp^2$ and $\tilde{k}_\perp \cdot \tilde{\Delta}_\perp$. The usual GPDs are obtained by integrating over \tilde{k}_\perp ,

$$\int d^2 k_\perp x f_g = \frac{1}{2P^+} \bar{u}(p') \left(H_g \gamma^+ + E_g \frac{i \sigma^{\nu+} \Delta_\nu}{2M} \right) u(p) \quad (5)$$

normalized as $H_g(x) = xG(x)$ in the forward limit. In the following we shall encounter the integrals

$$x L_g(x, \xi) \equiv - \int d^2 \tilde{k}_\perp \frac{\tilde{k}_\perp^2}{M^2} F_{1,4}(x, \xi, \tilde{\Delta}_\perp = 0), \quad (6)$$

$$O(x, \xi) \equiv \int d^2 \tilde{k}_\perp \frac{\tilde{k}_\perp^2}{M^2} F_{1,2}(x, \xi, \tilde{\Delta}_\perp = 0). \quad (7)$$

In the limit $\xi \rightarrow 0$, $L_g(x)$ is the parton distribution function of the gluon OAM [22] normalized as $\int_0^1 dx L_g(x) = L_g$. The imaginary part of $F_{1,2}$ is called the spin-dependent Odderon [23] and its k_\perp -moment is related to the three-gluon correlator relevant to transverse single spin asymmetry. The real part of $F_{1,2}$ is proportional to ξ , but otherwise unconstrained. In (4), the GTMDs are defined in the ‘symmetric’ frame where $P_\perp = 0$ so that $\tilde{p}'_\perp = \tilde{\Delta}_\perp/2 = -\tilde{p}_\perp$. The advantage of this frame is that one can exploit PT (parity & time-reversal) symmetry to constrain the dependence of GTMDs on variables. However, this frame is inconvenient and practically not used when describing actual experimental processes. We shall instead work in the so-called hadron frame where the incoming virtual photon and proton are collinear along the x^3 direction, namely, $p_\perp = 0$. The two frames are related by the so-called transverse boost, a Lorentz transformation that leaves invariant the plus component of a four-vector $V^+ = \tilde{V}^+$, $V_\perp = \tilde{V}_\perp + C_\perp \tilde{V}^+$, $V^- = \tilde{V}^- + C_\perp \cdot \tilde{V}_\perp + \frac{C_\perp^2}{2} \tilde{V}^+$. Applying this transformation to the matrix element (4) with $C_\perp = \tilde{\Delta}_\perp/(2p^+)$, we see that if one considers a scattering process in the




FIG. 1. Exclusive dijet production in electron-proton scattering.

hadron frame where $p_\perp = 0$, transverse momentum transfer $p'_\perp = \Delta_\perp$ and t -channel gluons with transverse momentum k_\perp , the GTMDs $F_{1,n}(x, \xi, \tilde{k}_\perp, \tilde{\Delta}_\perp)$ should be evaluated at

$$\tilde{k}_\perp = k_\perp - \frac{x}{2} \Delta_\perp, \quad \tilde{\Delta}_\perp = (1 + \xi) \Delta_\perp. \quad (8)$$

Many of the previous phenomenological applications of GTMDs have adopted the small- x kinematics $x \ll 1$ which also implies $\xi \ll 1$. In such cases, the difference (8) is negligible to first approximation.

3 Double spin asymmetry in diffractive dijet production—We consider exclusive dijet production in electron-proton scattering depicted in Fig. 1. This process has attracted a lot of attention in the literature in different contexts [11, 12, 24–32]. However, longitudinal double spin asymmetry has not been studied to our knowledge. In the hadron frame, the longitudinally polarized proton moves fast in the $+x^3$ direction and the virtual photon with virtuality $q^2 = -Q^2$ in the $-x^3$ direction. Two jets in the final state have longitudinal momentum fractions (of the photon) $z = p \cdot q_1/p \cdot q$ and $\bar{z} = 1 - z$, and transverse momenta $q_{1\perp} - z\Delta_\perp$ and $-q_{1\perp} - \bar{z}\Delta_\perp$, respectively, where Δ_\perp is the transverse momentum of the recoiling proton. q_\perp is related to skewness ξ and the γ^*p center-of-mass energy $W^2 = (p + q)^2$ as

$$\xi = \frac{q_\perp^2 + z\bar{z}Q^2}{-q_\perp^2 + z\bar{z}(Q^2 + 2W^2)}. \quad (9)$$

The momenta of the incoming lepton is parameterized as $l^\mu = (l^+, l^-, l_\perp) = (\frac{Q(1-y)}{\sqrt{2y}}, \frac{Q}{\sqrt{2y}}, \frac{Q\sqrt{1-y}}{y} n_\perp)$ where $y = p \cdot q/p \cdot l$ as usual and $n_\perp = (\cos \phi_{l_\perp}, \sin \phi_{l_\perp})$ is a unit vector in the transverse plane. The spin-dependent part of the lepton tensor is $L^{\mu\nu} \sim -2i\epsilon^{\mu\nu\rho\sigma} s_\rho q_\sigma$. For a longitudinally polarized lepton, $s_\rho = h_l l_\rho$ where $h_l = \pm 1$ is the helicity. In order to be sensitive to the azimuthal angle of the lepton plane ϕ_{l_\perp} , the index ρ has to be transverse. Since $\sigma = \pm$ is longitudinal, one of μ, ν must be

longitudinal and the other transverse. Namely, we should look at the interference effect between the longitudinal A_L and transverse A_T^λ ($\lambda = 1, 2$) virtual photon amplitudes

$$A_L = A_L^2 + A_L^3, \quad A_T^\lambda = \epsilon^{\lambda i} (A_T^{2i} + A_T^{3i}). \quad (10)$$

The twist-2 part $A_{L/T}^2$ is proportional to gluon GPDs and has been calculated in [25] in the two-gluon exchange approximation (see Fig. 1). The twist-3 part $A_{L/T}^3$ involves GTMDs and retains one factor of t -channel gluon transverse momentum k_\perp . It has been calculated in [11] and here we reproduce the result

$$\begin{aligned} A_T^3 &= -\frac{ig_s^2 e_{em} e_q}{N_c} \frac{2(\bar{z} - z)}{(q_\perp^2 + \mu^2)^2} \bar{u}(q_1) \epsilon_\perp \cdot \gamma_\perp v(q_2) \\ &\times \int dx \frac{x}{(x^2 - \xi^2 + i\xi\epsilon)^2} \left(2\xi + \frac{(2\xi)^3(1 - 2\beta)}{(x^2 - \xi^2 + i\xi\epsilon)} \right) \\ &\times \int d^2 k_\perp q_\perp \cdot k_\perp x f_g(x, \xi, k_\perp, \Delta_\perp) \\ &- \frac{ig_s^2 e_{em} e_q}{N_c} \frac{2(2\xi)^2 z \bar{z} W}{(q_\perp^2 + \mu^2)^2} \bar{u}(q_1) \gamma^- v(q_2) \\ &\times \int dx \frac{x}{(x^2 - \xi^2 + i\xi\epsilon)^2} \\ &\times \int d^2 k_\perp \epsilon_\perp \cdot k_\perp x f_g(x, \xi, k_\perp, \Delta_\perp), \quad (11) \end{aligned}$$

$$\begin{aligned} A_L^3 &= \frac{ig_s^2 e_{em} e_q}{N_c} \frac{16\xi^2(\bar{z} - z)z\bar{z}QW}{(q_\perp^2 + \mu^2)^3} \bar{u}(q_1) \gamma^- v(q_2) \\ &\times \int dx \frac{x}{(x^2 - \xi^2 + i\xi\epsilon)^2} \left(1 + \frac{8\xi^2(1 - \beta)}{(x^2 - \xi^2 + i\xi\epsilon)} \right) \\ &\times \int d^2 k_\perp q_\perp \cdot k_\perp x f_g(x, \xi, k_\perp, \Delta_\perp), \quad (12) \end{aligned}$$

where $\mu^2 = z\bar{z}Q^2$ and $\beta = \frac{\mu^2}{q_\perp^2 + \mu^2}$. The k_\perp -weighted integrals of f_g lead to the moments (6) and (7). Importantly, in both the longitudinal and transverse amplitudes, the x -integral contains a third pole at $x = \pm\xi$. Such poles often imply the breakdown of collinear factorization due to diverging x -integrals [33]. (Gluon GPDs may contain terms proportional to $\theta(\xi - x)(x^2 - \xi^2)^2$ which are not integrable if there is a third pole.) Fortunately, these potentially dangerous terms can be dropped by setting $z = 1/2$, after which $A_L^3 = 0$ and only a second pole remains in A_T^3 . Note that, if one considers SSA [11], one cannot set $z = 1/2$ because the asymmetry (2) vanishes at this point. After integrating over the jet azimuthal angle ϕ_{q_\perp} , we obtain the following contribution to DSA at $z = 1/2$

$$\begin{aligned} &\frac{d\sigma}{dydQ^2 d\phi_\perp dzdq_\perp^2 d^2\Delta_\perp} \\ &= \frac{\alpha_{em} y}{2^{11} \pi^7 Q^4} \frac{\int d\phi_{q_\perp} L^{\mu\nu} A_\mu^* A_\nu}{(W^2 + Q^2)(W^2 - M_J^2)z\bar{z}}, \quad (13) \end{aligned}$$

where $M_J^2 = q_\perp^2/(z\bar{z}) = 4q_\perp^2$ is the invariant mass of dijet and

$$\begin{aligned} &\int d\phi_{q_\perp} L^{\mu\nu} A_\mu^* A_\nu \\ &= -\frac{2^{10} \pi^4}{N_c} h_l h_p \alpha_s^2 \alpha_{em} e_q^2 \frac{(1 + \xi)\xi Q^2}{(q_\perp^2 + \mu^2)^2} |l_\perp| |\Delta_\perp| \\ &\times \Re \left[\left\{ \mathcal{H}_g^{(1)*} - \frac{\xi^2}{1 - \xi^2} \mathcal{E}_g^{(1)*} \right. \right. \\ &\quad \left. \left. + \frac{4q_\perp^2}{q_\perp^2 + \mu^2} \left(\mathcal{H}_g^{(2)*} - \frac{\xi^2}{1 - \xi^2} \mathcal{E}_g^{(2)*} \right) \right\} \mathcal{L}_g \right. \\ &\quad \left. + \left(\mathcal{E}_g^{(1)*} + \frac{4q_\perp^2}{q_\perp^2 + \mu^2} \mathcal{E}_g^{(2)*} \right) \frac{\mathcal{O}}{2} \right] \cos(\phi_{l_\perp} - \phi_{\Delta_\perp}). \quad (14) \end{aligned}$$

The details of the calculation, including the case $z \neq 1/2$, will be presented elsewhere [34]. The various ‘Compton form factors’ are defined as

$$\mathcal{H}_g^{(1)}(\xi) = \int_{-1}^1 dx \frac{H_g(x, \xi)}{(x - \xi + i\epsilon)(x + \xi - i\epsilon)}, \quad (15)$$

$$\mathcal{H}_g^{(2)}(\xi) = \int_{-1}^1 dx \frac{\xi^2 H_g(x, \xi)}{(x - \xi + i\epsilon)^2 (x + \xi - i\epsilon)^2}, \quad (16)$$

$$\mathcal{L}_g(\xi) = \int_{-1}^1 dx \frac{x^2 L_g(x, \xi)}{(x - \xi + i\epsilon)^2 (x + \xi - i\epsilon)^2}, \quad (17)$$

$$\mathcal{O}(\xi) = \int_{-1}^1 dx \frac{x \mathcal{O}(x, \xi)}{(x - \xi + i\epsilon)^2 (x + \xi - i\epsilon)^2}, \quad (18)$$

and $\mathcal{E}_g^{(1,2)}(\xi)$ is defined from $E_g(x, \xi)$ similarly to $\mathcal{H}_g^{(1,2)}(\xi)$. Assuming $|H_g| \gg |E_g|$, we see that the cross section is directly proportional to the Compton form factor of the gluon OAM \mathcal{L}_g . The characteristic correlation $k_\perp \times \Delta_\perp \sim \sin(\phi_{k_\perp} - \phi_{\Delta_\perp})$ of OAM manifests itself as a cosine correlation between the outgoing electron and proton angles. A similar transfer of angular correlations has been noticed in [35] for the $\cos 2(\phi_{k_\perp} - \phi_{\Delta_\perp})$ dependence of the elliptic gluon GTMD [27, 36]. Away from the point $z = 1/2$, there are corrections proportional to $(z - 1/2)^2$, but collinear factorization is suspect for them as already mentioned. Instead, one should use the k_\perp -factorization approach to calculate the corrections, although their connection to the OAM is less clear.

4. *DSA from the gluon helicity*—Next we discuss another source of DSA from the gluon helicity GPDs

$$\begin{aligned} &\epsilon_{ij} \int \frac{dz^-}{\pi} e^{ixP^+ z^-} \langle p' | F_a^{+i}(-z/2) F_a^{+j}(z/2) | p \rangle \\ &= \bar{u}(p') \left(i\tilde{H}_g \gamma_5 \gamma^+ - i\tilde{E}_g \frac{\gamma_5 \Delta^+}{2M} \right) u(p), \quad (19) \end{aligned}$$

where $\tilde{H}_g(x) = x\Delta G(x)$ in the forward limit. This originates from the interference between the unpolarized and

polarized gluon GPDs in the amplitude and the complex-conjugate amplitude.¹ An entirely analogous contribution should be added to the result for SSA in [11, 12]. The $\cos(\phi_{l_\perp} - \phi_{\Delta_\perp})$ asymmetry due to this mechanism is actually known in the context of Deeply Virtual Compton Scattering (DVCS) [37]. Unlike in DVCS, in dijet production there is no contamination from the Bethe-Heitler process. The asymmetry can be calculated purely within the GPD framework by setting $k_\perp = 0$ but keeping one factor of Δ_\perp in the hard part. In general, the cross section contains integrals with a third pole such as

$$\int dx \frac{H_g(x, \xi)}{(x^2 - \xi^2 + i\xi\epsilon)^3}, \quad \int dx \frac{x\tilde{H}_g(x, \xi)}{(x^2 - \xi^2 + i\xi\epsilon)^3}. \quad (20)$$

Remarkably, however, these factorization-breaking terms all vanish at $z = 1/2$ and we find [34]

$$\begin{aligned} \int d\phi_{q_\perp} L^{\mu\nu} A_\mu A_\nu &= \frac{2^{10}\pi^4}{N_c} h_l h_p \alpha_s^2 \alpha_{em} e_q^2 \frac{(1 - \xi^2)\xi Q^2}{(q_\perp^2 + \mu^2)^2} \\ &\times \Re \left[\left(\mathcal{H}_g^{(1)*} - \frac{\xi^2}{1 - \xi^2} \mathcal{E}_g^{(1)*} \right) \left(\tilde{\mathcal{H}}_g^{(2)} - \frac{\xi^2}{1 - \xi^2} \tilde{\mathcal{E}}_g^{(2)} \right) \right] \\ &\times |l_\perp| |\Delta_\perp| \cos(\phi_{l_\perp} - \phi_{\Delta_\perp}), \end{aligned} \quad (21)$$

with

$$\begin{aligned} \tilde{\mathcal{H}}_g^{(2)}(\xi) &= \int dx \frac{x\tilde{H}_g(x, \xi)}{(x^2 - \xi^2 + i\xi\epsilon)^2}, \\ \tilde{\mathcal{E}}_g^{(2)}(\xi) &= \int dx \frac{x\tilde{E}_g(x, \xi)}{(x^2 - \xi^2 + i\xi\epsilon)^2}. \end{aligned} \quad (22)$$

In the following, we shall be mainly interested in the small- ξ region $\xi \lesssim 10^{-3}$. In this region $\mathcal{H}_g^{(1,2)}(\xi)$ are dominated by the imaginary part, and one can show that $\Im \mathcal{H}_g^{(1)} \approx -2\Im \mathcal{H}_g^{(2)}$.² Combining (21) with (14) and neglecting \mathcal{E}_g and $\tilde{\mathcal{E}}_g$, we find that the asymmetry is roughly proportional to the combination

$$\mathcal{H}_g^{(1)*} \left(\tilde{\mathcal{H}}_g^{(2)} + \frac{q_\perp^2 - \mu^2}{q_\perp^2 + \mu^2} \mathcal{L}_g \right). \quad (23)$$

Depending on the sign of $q_\perp^2 - \mu^2 = q_\perp^2 - Q^2/4$, the helicity and OAM contributions interfere positively or negatively. Note that $\tilde{\mathcal{H}}_g^{(2)} \pm \mathcal{L}_g \sim \Delta G(x) \pm L_g(x)$, and

$\Delta G(x) \approx -L_g(x)$ at small- x [7, 12, 38, 39] (see however, [8]). Thus, the two contributions have the same sign when $q_\perp^2 < Q^2/4$ but tend to cancel each other when $q_\perp^2 > Q^2/4$. By varying Q^2 , we should be able to see this very interesting interplay between the helicity and the OAM.

5. Numerical results—We now present a numerical estimate of the cross section. We neglect E_g, \tilde{E}_g altogether. $H_g(x, \xi), \tilde{H}_g(x, \xi)$ and $xL_g(x, \xi)$ are reconstructed from their PDF counterparts $xG(x), x\Delta G(x)$ and $xL_g(x)$, respectively, using the method of double distributions [40, 41]. We use the JAM [4, 42] gluon PDFs $xG(x)$ and $x\Delta G(x)$ as inputs. As for $xL_g(x)$, we employ the Wandzura-Wilczek (WW) approximation [22]

$$L_g(x) \approx x \int_x^1 \frac{dx'}{x'^2} x' G(x') - 2x \int_x^1 \frac{dx'}{x'^2} \Delta G(x'), \quad (24)$$

although we are eventually interested in constraining the genuine twist-three part neglected in this approximation. We integrate over Δ_\perp assuming a Gaussian form factor $e^{-b\Delta_\perp^2}$ with $b = 5 \text{ GeV}^{-2}$ [25] and change variables $q_\perp \rightarrow \xi$ according to (9). The other parameters are fixed as $h_p h_l = 1$, $\sqrt{s_{ep}} = 120 \text{ GeV}$ and $y = 0.7$. The resulting cross section (only the DSA part)

$$\frac{d\sigma}{dy dQ^2 dz d\xi d\delta\phi}, \quad (25)$$

is shown in Fig. 2 at $\delta\phi = \phi_{l_\perp} - \phi_{\Delta_\perp} = 0$ for three different values of Q^2 (2.7 GeV², 4.8 GeV² and 10 GeV²). The plots correspond to $1 < q_\perp < 3 \text{ GeV}$ ($1 < q_\perp < 2.35 \text{ GeV}$ in the $Q^2 = 2.7 \text{ GeV}^2$ case). Typical jet rapidities in the laboratory frame are $-2 < \eta < -1$ at the top EIC energy $E_p \sim 250 \text{ GeV}$. We see that the OAM (14) and helicity (21) contributions are comparable in magnitude, though the latter tends to be larger because of the cancellation between $\mathcal{H}_g^{(1)}$ and $\mathcal{H}_g^{(2)}$. As a result of this cancellation, we observe a clear sign change of the OAM contribution with increasing Q^2 , see (23). It should be mentioned that there are large uncertainties in our prediction even in the helicity part because currently $\Delta G(x)$ is poorly constrained, including even the sign, especially in the small- x region but also in the large- x region. (See a recent discussion [43] on this point.) Besides, nothing is known about $L_g(x)$ experimentally at the moment, and our model for \mathcal{L}_g involves key assumptions (the WW approximation and the use of the double distribution technique) whose validity needs to be investigated. The above result should thus be regarded as an exploratory study to be significantly improved in future. Nonetheless, our calculation adequately demonstrates the feasibility of accessing the OAM from DSA. Ultimately, \mathcal{L}_g can be extracted from future experimental data, and for this purpose an accurate determination of $xG(x)$ and $x\Delta G(x)$ down to $x \sim 10^{-3}$ is crucial.

¹ We mention in passing that if one starts out with the GTMD version of Eq. (19), then there will be another contribution to the asymmetry proportional to certain gluon helicity GTMDs and the gluon helicity GPD. However, we expect this contribution to be orders of magnitude smaller than the one we discuss here, because the helicity GPD would be much smaller than the unpolarized GPD in the kinematics we consider. We plan to explain these subtleties in a follow-up work [34].

² More precisely, one can show that

$$\Im \mathcal{H}_g^{(2)}(\xi) = -\frac{\pi}{2} \frac{d}{dx} H_g(x, \xi) \Big|_{x=\xi} + \frac{\pi}{2\xi} H_g(\xi, \xi).$$

In the limit $\xi \ll 1$, the second term dominates.




FIG. 2. The DSA part of the differential cross section (25) at $Q^2 = 2.7 \text{ GeV}^2$ (left), $Q^2 = 4.8 \text{ GeV}^2$ (middle) and $Q^2 = 10 \text{ GeV}^2$ (right). The labels ‘OAM’ and ‘Helicity’ refer to the two contributions (14) and (21), respectively.

6. Conclusions—We have proposed DSA in exclusive dijet production as a novel observable for the gluon OAM L_g that can be measured at the EIC. Compared to SSA (2) previously suggested in [11, 12], it has a number of advantages. Most importantly, the third poles at $x = \pm\xi$ in (11), (12) and (20) which are potentially dangerous for QCD factorization can be eliminated by setting $z = 1/2$, but this is not possible in SSA. In practice, measurements are done in some window in z . We expect that the cross section $d\sigma/dz$ varies mildly around $z \sim 1/2$, but this needs to be substantiated in future investigations. Second, unlike the jet angle $\phi_{q\perp}$, the electron angle $\phi_{l\perp}$ is not affected by final state QCD radiations. The former is integrated over in DSA, and this greatly simplifies the cross section formula without losing sensitivity to the OAM. We thus expect that DSA is more robust against higher order QCD corrections to this process [28]. Furthermore, in the limit $x \approx \xi \rightarrow 0$, $\mathcal{H}_g^{(1,2)}$ are dominantly imaginary, and the extraction of the imaginary part in (2) turned out to be a delicate problem within the effective theory of high energy QCD [12]. For DSA, such a concern is simply absent. (We however note that in the present GPD-like approach, the real and imaginary parts

of \mathcal{L}_g are comparable in magnitude.)

The present calculation can be straightforwardly extended to the quark exchange channel important in the low-energy (low- W , high- ξ) region. We expect an additional contribution proportional to the product of the quark GPD and the quark OAM $\sim \mathcal{H}_q^* \mathcal{L}_q$. This will be a nice addition to the finding in [13] which is so far the only observable known to be sensitive to L_q .

Acknowledgements—We thank Feng Yuan and Yong Zhao for explaining to us the results in [11] and for discussion. S. B. and Y. H. were supported by the U.S. Department of Energy under Contract No. DE-SC0012704, and also by Laboratory Directed Research and Development (LDRD) funds from Brookhaven Science Associates. S. B. has also been supported by the U.S. Department of Energy, Office of Science, Office of Nuclear Physics and Office of Advanced Scientific Computing Research within the framework of Scientific Discovery through Advance Computing (SciDAC) award Computing the Properties of Matter with Leadership Computing Resources.

-
- [1] L. Adamczyk *et al.* [STAR], Phys. Rev. Lett. **115** (2015), 092002 [arXiv:1405.5134 [hep-ex]].
- [2] D. de Florian, R. Sassot, M. Stratmann and W. Vogelsang, Phys. Rev. Lett. **113** (2014), 012001 [arXiv:1404.4293 [hep-ph]].
- [3] E. R. Nocera *et al.* [NNPDF], Nucl. Phys. B **887** (2014), 276-308 [arXiv:1406.5539 [hep-ph]].
- [4] J. J. Ethier, N. Sato and W. Melnitchouk, Phys. Rev. Lett. **119** (2017), 132001 [arXiv:1705.05889 [hep-ph]].
- [5] M. S. Abdallah *et al.* [STAR], [arXiv:2110.11020 [hep-ex]].
- [6] R. Abdul Khalek, A. Accardi, J. Adam, D. Adamiak, W. Akers, M. Albaladejo, A. Al-bataineh, M. G. Alexeev, F. Ameli and P. Antonioli, *et al.* [arXiv:2103.05419 [physics.ins-det]].
- [7] R. Boussarie, Y. Hatta and F. Yuan, Phys. Lett. B **797** (2019), 134817 [arXiv:1904.02693 [hep-ph]].
- [8] Y. V. Kovchegov, JHEP **03** (2019), 174 [arXiv:1901.07453 [hep-ph]].
- [9] M. Engelhardt, J. R. Green, N. Hasan, S. Krieg, S. Meinel, J. Negele, A. Pochinsky and S. Syritsyn, Phys. Rev. D **102** (2020), 074505 [arXiv:2008.03660 [hep-lat]].
- [10] Y. Guo, X. Ji and K. Shiells, Nucl. Phys. B **969** (2021), 115440 [arXiv:2101.05243 [hep-ph]].
- [11] X. Ji, F. Yuan and Y. Zhao, Phys. Rev. Lett. **118** (2017), 192004 [arXiv:1612.02438 [hep-ph]].
- [12] Y. Hatta, Y. Nakagawa, F. Yuan, Y. Zhao and B. Xiao, Phys. Rev. D **95** (2017), 114032 [arXiv:1612.02445 [hep-ph]].
- [13] S. Bhattacharya, A. Metz and J. Zhou, Phys. Lett. B **771** (2017), 396-400 [erratum: Phys. Lett. B **810** (2020), 135866] [arXiv:1702.04387 [hep-ph]].
- [14] S. Bhattacharya, A. Metz, V. K. Ojha, J. Y. Tsai and J. Zhou, [arXiv:1802.10550 [hep-ph]].
- [15] A. Courtoy, G. R. Goldstein, J. O. Gonzalez Hernandez, S. Liuti and A. Rajan, Phys. Lett. B **731** (2014), 141-147

- [arXiv:1310.5157 [hep-ph]].
- [16] C. Lorce and B. Pasquini, Phys. Rev. D **84** (2011), 014015 [arXiv:1106.0139 [hep-ph]].
- [17] Y. Hatta, Phys. Lett. B **708** (2012), 186-190 [arXiv:1111.3547 [hep-ph]].
- [18] C. Lorce, B. Pasquini, X. Xiong and F. Yuan, Phys. Rev. D **85** (2012), 114006 [arXiv:1111.4827 [hep-ph]].
- [19] A. V. Belitsky, X. d. Ji and F. Yuan, Phys. Rev. D **69** (2004), 074014 [arXiv:hep-ph/0307383 [hep-ph]].
- [20] S. Meissner, A. Metz and M. Schlegel, JHEP **08** (2009), 056 [arXiv:0906.5323 [hep-ph]].
- [21] C. Lorcé and B. Pasquini, JHEP **09** (2013), 138 [arXiv:1307.4497 [hep-ph]].
- [22] Y. Hatta and S. Yoshida, JHEP **10** (2012), 080 [arXiv:1207.5332 [hep-ph]].
- [23] J. Zhou, Phys. Rev. D **89** (2014), 074050 [arXiv:1308.5912 [hep-ph]].
- [24] J. Bartels, H. Lotter and M. Wüsthoff, Phys. Lett. B **379** (1996), 239-248 [erratum: Phys. Lett. B **382** (1996), 449-449] [arXiv:hep-ph/9602363 [hep-ph]].
- [25] V. M. Braun and D. Y. Ivanov, Phys. Rev. D **72** (2005), 034016 [arXiv:hep-ph/0505263 [hep-ph]].
- [26] T. Altinoluk, N. Armesto, G. Beuf and A. H. Rezaeian, Phys. Lett. B **758** (2016), 373-383 [arXiv:1511.07452 [hep-ph]].
- [27] Y. Hatta, B. W. Xiao and F. Yuan, Phys. Rev. Lett. **116** (2016), 202301 [arXiv:1601.01585 [hep-ph]].
- [28] R. Boussarie, A. V. Grabovsky, L. Szymanowski and S. Wallon, JHEP **11** (2016), 149 [arXiv:1606.00419 [hep-ph]].
- [29] Y. Hagiwara, Y. Hatta, R. Pasechnik, M. Tasevsky and O. Teryaev, Phys. Rev. D **96** (2017), 034009 [arXiv:1706.01765 [hep-ph]].
- [30] H. Mäntysaari, N. Mueller and B. Schenke, Phys. Rev. D **99** (2019), 074004 [arXiv:1902.05087 [hep-ph]].
- [31] F. Salazar and B. Schenke, Phys. Rev. D **100** (2019), 034007 [arXiv:1905.03763 [hep-ph]].
- [32] D. Boer and C. Setyadi, Phys. Rev. D **104** (2021), 074006 [arXiv:2106.15148 [hep-ph]].
- [33] Z. L. Cui, M. C. Hu and J. P. Ma, Eur. Phys. J. C **79** (2019), 812 [arXiv:1804.05293 [hep-ph]].
- [34] S. Bhattacharya, R. Boussarie and Y. Hatta, work in progress.
- [35] J. Zhou, Phys. Rev. D **94** (2016), 114017 [arXiv:1611.02397 [hep-ph]].
- [36] Y. Hagiwara, C. Zhang, J. Zhou and Y. j. Zhou, Phys. Rev. D **104** (2021), 094021.
- [37] A. V. Belitsky, D. Mueller, L. Niedermeier and A. Schafer, Nucl. Phys. B **593** (2001), 289-310 [arXiv:hep-ph/0004059 [hep-ph]].
- [38] Y. Hatta and D. J. Yang, Phys. Lett. B **781** (2018), 213-219 [arXiv:1802.02716 [hep-ph]].
- [39] J. More, A. Mukherjee and S. Nair, Eur. Phys. J. C **78** (2018), 389 [arXiv:1709.00943 [hep-ph]].
- [40] A. V. Radyushkin, Phys. Rev. D **59** (1999), 014030 [arXiv:hep-ph/9805342 [hep-ph]].
- [41] A. V. Radyushkin, [arXiv:hep-ph/0101225 [hep-ph]].
- [42] N. Sato *et al.* [JAM], Phys. Rev. D **101** (2020), 074020 [arXiv:1905.03788 [hep-ph]].
- [43] Y. Zhou, N. Sato and W. Melnitchouk, [arXiv:2201.02075 [hep-ph]].