




















**Detection of Near-Infrared Water Ice at the Surface of the (pre)Transitional Disk of AB Aur:
Informing Icy Grain Abundance, Composition, and Size**S. K. BETTI ¹, K. FOLLETTE ², S. JORQUERA ³, G. DUCHÊNE ^{4,5}, J. MAZOYER ⁶, M. BONNEFOY ⁵,
G. CHAUVIN ⁵, L. M. PÉREZ ^{3,7}, A. BOCCALETTI ⁶, C. PINTÉ ^{8,5}, A. J. WEINBERGER ⁹, C. GRADY ¹⁰,
L. M. CLOSE ¹¹, D. DEFRÈRE ¹², E. C. DOWNEY ¹¹, P. M. HINZ ^{11,13}, F. MÉNARD ⁵, G. SCHNEIDER ¹¹,
A. J. SKEMER ¹⁴ AND A. VAZ¹¹¹*Department of Astronomy, University of Massachusetts, Amherst, MA 01003, USA*²*Department of Physics and Astronomy, Amherst College, Amherst, MA 01003, USA*³*Departamento de Astronomía, Universidad de Chile, Camino El Observatorio 1515, Las Condes, Santiago, Chile*⁴*Department of Astronomy, University of California, Berkeley, CA 94720, USA*⁵*Université Grenoble Alpes, CNRS, IPAG, F-38000 Grenoble, France*⁶*LESIA, Observatoire de Paris, Université PSL, CNRS, Sorbonne Université, Université de Paris, 92195 Meudon, France*⁷*Núcleo Milenio de Formación Planetaria (NPF), Chile*⁸*School of Physics & Astronomy, Monash University, Clayton VIC 3800, Australia*⁹*Department of Terrestrial Magnetism, Carnegie Institution for Science, Washington, DC 20015, USA*¹⁰*Eureka Scientific, Oakland, CA 96002, USA*¹¹*Steward Observatory, Department of Astronomy, University of Arizona, Tucson, AZ 85721, USA*¹²*Institute of Astronomy, KU Leuven, Celestijnenlaan 200D, 3001, Leuven, Belgium*¹³*Center for Adaptive Optics, UC Santa Cruz, 1156 High St., Santa Cruz, CA 95064, USA*¹⁴*Department of Astronomy and Astrophysics, University of California, Santa Cruz, Santa Cruz, CA 95064, USA***ABSTRACT**

We present near-infrared Large Binocular Telescope Interferometer LMIRCam imagery of the disk around the Herbig Ae/Be star AB Aurigae. A comparison of surface brightness at K_s ($2.16\ \mu\text{m}$), H_2O narrowband ($3.08\ \mu\text{m}$), and L' ($3.7\ \mu\text{m}$) allows us to probe the presence of icy grains in this (pre)transitional disk environment. By applying Reference Differential Imaging PSF subtraction, we detect the disk at high signal to noise in all three bands. We find strong morphological differences between bands, including asymmetries consistent with observed spiral arms within 100 AU in L' . An apparent deficit of scattered light at $3.08\ \mu\text{m}$ relative to bracketing wavelengths (K_s and L') is evocative of ice absorption at the disk surface layer. However, the $\Delta(K_s - \text{H}_2\text{O})$ color is consistent with grains with little to no ice (0 – 5% by mass). The $\Delta(\text{H}_2\text{O} - L')$ color, conversely, suggests grains with a much higher ice mass fraction (~ 0.68), and the two colors cannot be reconciled under a single grain population model. Additionally, we find the extremely red $\Delta(K_s - L')$ disk color cannot be reproduced under conventional scattered light modeling with any combination of grain parameters or reasonable local extinction values. We hypothesize that the scattering surfaces at the three wavelengths are not co-located, and optical depth effects result in each wavelength probing the grain population at different disk surface depths. The morphological similarity between K_s and H_2O suggests their scattering surfaces are near one another, lending credence to the $\Delta(K_s - \text{H}_2\text{O})$ disk color constraint of $< 5\%$ ice mass fraction for the outermost scattering disk layer.

1. INTRODUCTION

The structure of the circumstellar environment of a star affects its evolution and the formation of any bound companions. The formation of substellar objects within

disks is highly dependent on location, stellar and circumstellar activity, and the distribution and state of the materials necessary for formation (Terada et al. 2007). One key to understanding planet/substellar formation within a disk is understanding the role water ice plays in the formation of both terrestrial and giant planets.

Water ice in circumstellar disks has been detected in various forms, including as vapor with *Spitzer Space Telescope* (Carr & Najita 2008; Salyk et al. 2008; Blevins

et al. 2016), and as crystalline water ice at 44 and 62 μm (Malfait et al. 1999; Meeus et al. 2001; McClure et al. 2012, 2015). Near-infrared (NIR) absorption from water ice has also been detected (Pontoppidan et al. 2005; Terada et al. 2007; Aikawa et al. 2012) using spectroscopy of edge-on disks, but these methods lack the angular resolution to resolve the spatial distribution of ice. The presence or absence of ice in circumstellar disks, and in particular the location of the snow line, is important to inform the reservoir of volatiles available to build giant planet cores (Drakowska & Alibert 2017).

In 2008, Inoue et al. proposed that the presence of icy grains in the surface layers of protoplanetary disks might be inferred from a deficit of NIR scattered light at 3.09 μm . Icy grains should absorb rather than scatter light at this wavelength preferentially, such that disks with an excess of icy grains will be fainter at 3.09 μm than the surrounding wavelengths. This method of constraining the ice distribution and spatial properties has been successfully implemented with detections of icy grains on the surface of the transitional disks of HD 142527 (Honda et al. 2009) and HD 100546 (Honda et al. 2016).

However, the recent results of Tazaki et al. (2021a) suggest that a more sophisticated approach to modeling the 3.09 μm ice absorption feature results in disks that require a substantially lower abundance of ice-rich grains in order to match observations. This was due to a relaxation of the assumptions of simple disk geometry and isotropic scattering present in the initial modeling (Inoue et al. 2008). Tazaki et al. (2021a) reanalyzed the results of Honda et al. (2009) using anisotropic scattering, and their results suggest a much lower ice abundance, as well as the presence of large micron-sized grains on the surface of HD 142527.

If we want to understand the reservoir of material available for formation of substellar objects beyond the snow line, we first need to be able to accurately measure its location. The disk surface ice line as probed by scattered light, and the midplane ice line as probed by mm emission (Notsu et al. 2017, 2018) may not necessarily align, however, the reservoir of icy grains available both radially and vertically throughout the disk is an important constraint for constraining planet formation theories, mapping grain growth, and informing disk evolutionary processes such as vertical transport. The so-called “grain filtration” theories are also of particular relevance, as these models predict that small grains, particularly at the disk surface, can easily flow inward past a forming planet while larger grains are impeded by a positive pressure gradient at the edge of the gap formed by a planet and remain there (Zhu et al. 2012).

AB Aurigae (AB Aur) is one of the most widely-studied Herbig Ae/Be stars. This A0 spectral type star is located approximately 162.9 ± 1.5 pc away (Gaia Collaboration et al. 2018) with an estimated mass of $2.4 \pm 0.2 M_{\odot}$, and an age of 4 ± 1 Myr (van den Ancker et al. 1997; DeWarf et al. 2003). It hosts a large, flared, inclined ($i \sim 23.2^{\circ}$) circumstellar disk with a range of substructures. The extended ($r \sim 450$ AU) and massive ($\sim 10 M_{\text{J}}$) disk around AB Aur was originally inferred from millimeter observations (Henning et al. 1998). Both spiral structure in the outer disk ($r \gtrsim 200$ AU, Grady et al. 1999; Fukagawa et al. 2004; Perrin et al. 2009; Tang et al. 2012) and a ring-like gap between 40 and 100 AU (Hashimoto et al. 2011) have been detected in scattered light. More recent work using resolved sub-mm and NIR imaging of AB Aur has led to the discovery of inner spiral arms (Hashimoto et al. 2011; Tang et al. 2017, Joquera et al. in prep), including a twist in the spiral at 30 AU that could be a result of a planet (Boccaletti et al. 2020).

Furthermore, millimeter continuum and CO gas emission reveal a radial cavity in the disk at 70 AU (Piétu et al. 2005), as well as midplane CO spirals in the inner disk, which could be a result of two or more planets at 30 and 60–80 AU from the star (Tang et al. 2017). AB Aur is also accreting at a high rate of $\sim 10^{-7} M_{\odot} \text{ yr}^{-1}$ (Salyk et al. 2013), indicating ongoing radial transport in the disk. Due to its intrinsic merit as a possible site of ongoing planet formation and its relatively face-on inclination, we targeted AB Aur to search for icy grains within the disk as they relate to potentially on-going planet formation.

In Section 2, we present new LBTI/LMIRCam observations and discuss our data reduction, while in Section 3 we detail the methods to obtain accurate disk surface brightnesses. We present the detection of absorption by icy grains in the disk of AB Aur in Section 4. We then discuss how this absorption informs our understanding of ice abundance, grain properties, and optical depth effects in Section 5. We conclude in Section 6 by summarizing the main results.

2. OBSERVATIONS AND DATA REDUCTION

2.1. Observations

AB Aur was observed with the L/M-band InfraRed Camera (LMIRCam) using one of the two 8.4 m mirrors of the Large Binocular Telescope Interferometer (LBTI) on 2015 January 04 (Skrutskie et al. 2010) in pupil-stabilized mode. The target was observed in two narrow band H₂O ice filters (Ice1: $\lambda_c = 3.05 \mu\text{m}$, $\Delta\lambda = 0.16 \mu\text{m}$ and Ice2: $\lambda_c = 3.08 \mu\text{m}$, $\Delta\lambda = 0.14 \mu\text{m}$), and the broadband K_s filter ($\lambda_c = 2.16 \mu\text{m}$, $\Delta\lambda = 0.32 \mu\text{m}$). Due to a

significant number of ghosts ($n = 5$) in the Ice1 band, a decision was made to obtain additional observations in the ghost-free Ice2 band filter. To allow for the removal and characterization of thermal emission from the background sky and telescope, the star was nodded up/down by $6''.3$ on the detector every 120 frames. Each frame had an exposure time of 2 s, resulting in saturation for the very bright ($K = 4.23$, $L = 3.24$) central star within $0''.15$.

The total exposure times for each filter (after removing 43 (1%) images unusable due to AO system effects such as an elongated point spread function (PSF)) were 31.7 min, 64 min, and 32 min in the Ice1, Ice2, and K_s bands, respectively, resulting in 152° of on-sky rotation. The full width at half maximum (FWHM) averaged around $0''.19$ for all wavelengths.

Two PSF reference stars were imaged in the same Ice and K_s bands with the same exposure time as AB Aur the same night shortly after the AB Aur observations. The two reference stars, HIP 22138 (G8III, $K = 4.627$, $L = 4.582$) and HIP 24447 (K0, $K = 4.219$, $L = 4.136$) (nearby and at a similar brightness and color as AB Aur), were observed in each band with 2 s exposures. HIP 22138 was observed for a total of 24 min, 16 min, and 16 min in the Ice1, Ice2, and K_s bands, respectively. HIP 24447 was observed for 16 mins in each of the Ice2 and K_s band and was not observed in Ice1. For flux calibration/photometry, unsaturated images of AB Aur ($t_{K_s} = 0.015$ s, $t_{Ice2} = 0.058$ s) and HIP 24447 ($t_{K_s} = 0.015$ s, $t_{Ice2} = 0.058$ s) were collected in Ice2 and K_s . See Table 1 for a summary of the observations.

We also obtained L' band ($\lambda_c = 3.70 \mu\text{m}$, $\Delta\lambda = 0.58 \mu\text{m}$) observations of AB Aur from Jorquera et al. (2022, in press). These were collected on 2014 February 12 in non-interferometric pupil-stabilized mode at the LBT. The star was nodded up/down on the LMIRCam detector every 50 co-adds. Each co-added consisted of 5 frames, each with exposure times of 0.175 s, resulting in 0.875 s per co-add. This resulted in saturation of the central star within $0''.15$. The total exposure time in L' was 133.5 min. During the observations, AB Aur transited within 2° of the zenith, resulting in an unstable AO loop due to rapid field rotation, resulting in the loss of a number ($n = 62$) of left (SX) image frames. The FWHM averaged around $0''.21$. One PSF reference star, HD 39925 (K5, $K = 3.899$, $L = 3.52$), was imaged at the same exposure time as AB Aur the same night immediately following the AB Aur observation for a total of 64 mins of integration. For flux calibration and photometry, unsaturated images of AB Aur ($t = 1.455$ s) were collected using a neutral density filter with a 0.9% transmission (no unsaturated HD 39925 were observed).

2.2. Data Reduction

The Ice and K_s band data were reduced using a custom reduction pipeline following the procedure given in Crepp et al. (2018). The L' band data was re-reduced using the same pipeline for self-consistency between the three bands. After removing unusable images, bad pixels ($> 8\sigma$) were corrected by replacing their value with the median of the surrounding 8 good pixels. Due to offsets between detector channels caused by bias drifts (64 columns), we removed striping from the images by subtracting the 3σ -clipped median of each channel from its columns; row offsets were removed similarly. Sky background was removed by subtracting the median of the 120 (50 for L') images taken closest in time at the opposite nod position. Distortion correction was also applied, as LMIRCam images contain significant optical distortion from camera flexure. The distortion is expected to be constant in time (Maire et al. 2015); therefore, we use an established distortion solution for the DX and SX aperture from 8 November 2017 provided by the LBTO team¹. Finally, LMIRCam is designed to be used with both apertures; since AB Aur was imaged on one side in K_s and Ice2, the images are oversampled, and we therefore bin each image into 2×2 pixels, giving a final plate scale of $0''.022$. We treat each eye taken with L' independently in order to match K_s and Ice. Therefore, we also bin each image into 2×2 pixels.

Images are registered across all three bands using cross correlation, and the unsaturated images are median combined. To fit for and remove the effects of the central star before derotating and stacking the frames, we use reference differential imaging (RDI) with Karhunen-Loève Image Projection (KLIP; Soummer et al. 2012) using the pyKLIP python package (Wang et al. 2015). KLIP works by creating a model of the stellar PSF for each frame from a library of reference images, which can then be subtracted to remove the stellar PSF. In the case of RDI, these reference images are images of one or several ‘reference’ stars, selected to be close in brightness and type, and without disks or planet detected around them.

We select the 100 most correlated reference star images to form the PSF library for each frame, and then use KLIP principal component analysis (PCA) to construct the model PSF and subtract it from the target image. The number of eigenvectors (KL modes basis vectors) used controls the complexity of the constructed PSF. We apply a simple (3 KL mode) model in or-

¹ <https://sites.google.com/a/lbto.org/lbti/data-retrieval-reduction/distortion-correction-and-astrometric-solution>

Table 1. Log of the LMIRCam AB Aur observations

Date	Time (UT)	Target	Band	Seeing (")	Air mass	θ (°)	t_{exp} (s)	N_{exp}	N_{nods}	Comment
2014-02-13	02:18–03:39	AB Aur	L'	0.66-1.05	1.01/1.02	-66/76	0.875	50	23.5	
2014-02-13	04:04–04:36	HD 39925	L'	0.73-1.42	1.00/1.01	62/75	0.875	50	13.5	PSF reference
2014-02-13	04:46–06:53	AB Aur	L'	0.68-1.41	1.02/1.63	77/70	0.875	50	27	
2014-02-13	07:06–07:31	HD 39925	L'	1.02-1.83	1.35/1.48	73/72	0.875	50	11	PSF reference
2015-01-04	02:33–03:20	AB Aur	H ₂ O-3.05 μ m	1.20-1.00	1.23/1.13	-75/-76	2	120	4	
2015-01-04	03:21–03:48	HIP 22138	H ₂ O-3.05 μ m	–	1.09/1.05	-73/-72	2	120	3	PSF reference
2015-01-04	03:48–04:07	HIP 22138	H ₂ O-3.08 μ m	–	1.05/1.03	-72/-69	2	120	2	PSF reference
2015-01-04	04:04–04:26	HIP 22138	K_s	0.74-1.03	1.03/1.01	-70/-65	2	120	2.5	PSF reference
2015-01-04	04:42–04:59	AB Aur	K_s	0.76-1.19	1.01/1.00	-74/-65	2	120	2	
2015-01-04	05:01–05:22	AB Aur	H ₂ O-3.08 μ m	unk-1.19	1.00/1.00	-63/-20	2	120	2	
2015-01-04	05:23–05:40	AB Aur	K_s	unk-3.18	1.00/1.00	2/60	2	120	2	
2015-01-04	05:42–06:35	AB Aur	H ₂ O-3.08 μ m	0.87-0.96	1.00/1.04	61/77	2	120	6	
2015-01-04	06:54–07:11	HIP 24447	H ₂ O-3.08 μ m	0.70-0.92	1.03/1.06	80/80	2	120	2	PSF reference
2015-01-04	07:13–07:30	HIP 24447	K_s	0.68-0.76	1.06/1.09	79/90	2	120	2	PSF reference
Unsaturated										
2014-02-13	03:39–04:04	AB Aur	L'	0.89-0.95	1.02/1.04	76/77	1.455	50	2.5	ND 0.9%
2014-02-13	06:53–07:05	AB Aur	L'	1.28-1.02	1.64/1.67	70/69	1.455	50	1	ND 0.9%
2015-01-04	06:36–06:38	AB Aur	H ₂ O-3.08 μ m	0.99	1.04/1.04	77/77	0.058	100	1	
2015-01-04	06:39–06:46	AB Aur	K_s	0.70	1.04/1.05	77/77	0.015	100	1.5	
2015-01-04	06:47–06:48	HIP 24447	K_s	–	1.03/1.03	80/80	0.015	100	1	PSF reference
2015-01-04	06:49–06:51	HIP 24447	H ₂ O-3.08 μ m	0.85-0.86	1.03/1.03	80/80	0.058	100	1	PSF reference

NOTE— θ refers to the parallactic angle in degrees, t_{exp} refers to the exposure time per image, N_{exp} is the number of images per nod, N_{nods} is the number of up/down nod sequences.

der to avoid, to the extent possible, oversubtraction of the nearly symmetrical and nearly face-on disk of AB Aur. Finally, KLIP derotates and stacks all the AB Aur frames to create a final image (Figure 1).

In order to demonstrate its effectiveness and estimate uncertainty, we also applied KLIP-RDI to the reference images, using one PSF reference star as a reference library for the other. If the disk structure around AB Aur is robust, similar structures should not appear on the PSF-PSF KLIP-RDI images. As the PSF reference stars are at the same brightness as AB Aur for their observed relevant bands (see Table 2), the resulting images give an estimate of the resulting noise due to PSF mismatch (bottom panel of Figure 1). Following Chen et al. (2020), we determine the uncertainty on recovered flux at each radial location by calculating the standard deviation in concentric annuli around the star with a width equal to the FWHM of the PSF (~ 4 pixels).

3. DISK SURFACE BRIGHTNESS

In order to quantify flux loss as a result of KLIP-RDI over-subtraction and PSF mismatch, which must be corrected for in order to report accurate photometry, we forward model synthetic MCFOST disks to determine the “throughput” of the algorithm at each band. These forward models were selected to provide a good match to the the surface brightness of the disk, as this is what we aim to quantify. We note that we were unable to find

Table 2. Properties of Observed Targets

	AB Aur	HIP22138	HD 24447	HD 39925
R.A. (J2000)*	04:55:45.84	04:45:50.07	05:14:39.43	05:57:00.23
Decl. (J2000)*	+30:33:04.29	+28:39:38.47	+31:24:06.93	+30:36:34.72
Observed band	all	K_s /Ice	K_s /Ice	L'
K (mag) [×]	4.23 \pm 0.02	4.62 \pm 0.02	4.22 \pm 0.02	3.89 \pm 0.23
L (mag) [^]	3.24 \pm 0.13	4.58 \pm 0.07	4.13 \pm 0.10	3.52 \pm 0.11
$K - L$ (mag)	0.99 \pm 0.14	0.04 \pm 0.08	0.09 \pm 0.10	0.37 \pm 0.26

References—*Gaia Collaboration et al. (2018), [×]Cutri et al. (2003), [^]Cutri & et al. (2012)

MCFOST models that both (a) converged on physically-realistic values of geometric and grain properties and (b) provided a good match to the surface brightness of the disk. Therefore, we use the models only for correction of surface brightness flux loss from KLIP-RDI, and do not make inferences from them about the physical or grain properties of AB Aur. See Appendix A for full details of our modeling attempts.

3.1. Throughput Correction

After computing the best fit models for each wavelength independently, we compute the throughput (TP) of the KLIP-RDI algorithm for these models. The TP is computed as the ratio of the initial, convolved model




Figure 1. **top)** KLIP-RDI reduced images of the disk around AB Aur at K_s (left), $3.08 \mu\text{m}$ Ice (center), and L' (right). **middle)** Same at top panel but multiplied by r^2 to enhance the outer substructure. **bottom)** Uncertainty noise maps from the KLIP-RDI reduction of the PSF reference stars. The field of view is $3'' \times 3''$ with north-up and east-left. A software mask (black) of $0''.15$ in radius is centered on each image. A black software mask also covers a ghost in the H_2O $3.08 \mu\text{m}$ noise map NW of the central mask.

disk to the final KLIP-RDI forward modeled image, with a value of 1 indicating no flux loss, and 0 indicating all disk flux was removed (oversubtracted) by RDI.

In Figure 2, the left column show the throughput maps computed for the best-fitting MCFOST model. As the substructure seen in the TP is mainly a result of broadly symmetric speckle noise or post processing artifacts, we azimuthally average² the TP using annuli of width $\sim \text{FWHM}$ (as shown in the right column of Figure 2), which smooths over these spurious variations. We find a consistent degree of flux loss ($\text{TP} \sim 15 - 25\%$) in $3.08 \mu\text{m}$ ice and L' bands, and almost 45% flux loss in

K_s band. At the outer boundary of the disk region, the model flux drops below the speckle noise in the forward model. The final reduced KLIP-RDI AB Aur images are divided by the final azimuthally averaged TPs to correct for algorithmic flux loss before conducting photometry on the disk. In Appendix B, we show that varying the surface brightness and geometry of the initial model has a negligible effect on the measured throughput ($\lesssim 2.5\%$).

3.2. Photometric Calibration

To determine the surface brightness of the throughput corrected disk, we photometrically calibrate the disk following the procedure of Quanz et al. (2011). The SED of AB Aur (Figure 3) shows strong IR excess from $1-4 \mu\text{m}$ in both photometry and spectroscopy indicating that the innermost disk component (within 0.3 AU ; Tannirkulam

² Boxcar smoothing the TP map to remove the variations gives similar final results/conclusions. We have chosen to azimuthally average the TP to remove any non-axisymmetric variations.




Figure 2. Map of throughput resulting from the KLIP RDI reduction for K_s (top), $3.08 \mu\text{m}$ Ice (center), and L' (bottom) filters. The left column shows the true throughput and the right column is the azimuthally averaged throughput. A value of 0 means all flux is removed by RDI, while a value of 1 means no flux is lost. The “point source” in the west of the L' TP map is an artifact in the post-processing.

et al. 2008; Natta et al. 2001) is completely unresolved. Therefore, we can assume that our unsaturated images of AB Aur contain both star and excess inner disk thermal radiation. We use the unsaturated images of AB Aur to determine both the instrumental zeropoint and the photometric system zeropoint since the known photometry and spectroscopy of AB Aur also contain this excess IR emission. To quantify the instrumental zeropoint, we use aperture photometry to measure the total intensity of the unsaturated AB Aur images in each band and translate this to an instrumental magnitude. We find 1.9×10^6 counts/s, 1.0×10^6 counts/s, and 9.4×10^5 counts/s in the K_s , $3.08 \mu\text{m}$ and L' filters, respectively. For K_s band, the transmission bandpass is centered at 2MASS K_s band; therefore, we can use

the known 2MASS AB Aur K_s magnitude ($K_s = 4.23 \pm 0.016$; Cutri et al. 2003) to calculate the zeropoint.

We calculate the $3.08 \mu\text{m}$ and L' zeropoints by convolving the LMIRCam transmission profiles with the published ISO/SWS spectrum of AB Aur ($2.4 - 45 \mu\text{m}$ van den Ancker et al. 2000). The transmission curves for the filters were obtained from the SVO Filter Profile Service (Rodrigo et al. 2012; Rodrigo & Solano 2020). The $2.6 - 40 \mu\text{m}$ flux calibrated spectrum (Figure 3 left panel) covers the Ice2 and L' LMIRCam bands, allowing us to synthesize the photometric flux in each band (see right panel Figure 3). Using the LMIRCam known photometric system zeropoints ($3.08 \mu\text{m}$: 359.73 Jy, L' : 257.10 Jy; Rodrigo et al. 2012; Rodrigo & Solano 2020), we convert flux to magnitudes. We then calculate and apply the instrumental zeropoints to find the surface brightness of the disk using the photometric system zeropoints (K_s : 666.7 Jy, $3.08 \mu\text{m}$ and L' as listed above).

4. RESULTS

We detect the disk of AB Aur with a signal to noise ratio (SNR) of ~ 2.5 from ~ 25 to ~ 175 AU in all three bands, as shown in Figure 1. Overall, as shown by Figure 4a, the K_s disk structure and morphology closely match the morphology of the H -band polarized intensity images of Hashimoto et al. (2011) and Boccaletti et al. (2020). As PI imagery is generally considered extremely robust, and is virtually unaffected by stellar emission, this match lends credence to the integrity of our KLIP-RDI total intensity disk recoveries in general, and K_s band in particular. The SE rim of the disk is markedly brighter than the NW in K_s and H_2O $3.08 \mu\text{m}$ band, similar to the H band polarized images of Hashimoto et al. (2011), suggesting strong forward-scattering at K_s . The disk appears more elliptical in L' (see Figure 4b), with a shape and spirals features similar to those reported in Jorquera et al. (2022, in press), Boccaletti et al. (2020), and Oppenheimer et al. (2008). Figure 5 show the locations of known spiral arms and substructure (Boccaletti et al. 2020) overlaid on the L' RDI image. The location of the spiral arms aligns with visible substructure in L' , visually confirming that we are seeing the spirals.

4.1. Surface Brightness Profiles

Figure 6 shows disk surface brightness profiles along the major (panels a, c, e) and minor (panels b, d, f) axes of the disk in contrast units (panels a and b), flux units (panels c and d) and $K_s - H_2O$ and $K_s - L'$ disk colors (panels e and f) space. Locations of apparent negative flux are indicated by dashed lines. We are able to detect the disk down to 10^{-5} contrast. The substructure




Figure 3. *left*) optical to FIR SED of AB Aur. A reddening extinction correction is applied using $A_V = 0.431$ mag (Vioque et al. 2018) assuming an Cardelli et al. (1989) extinction law. The blue circles indicate known observed photometry from literature (photometry downloaded from VizieR photometry viewer). The dark red spectra is from 1 – 2.46 μm taken from ARC 3.5m TripleSpec at APO. The red spectra is from ISO SWS (van den Ancker et al. 2000). The gold line shows the best fit Kurucz stellar atmosphere model in the optical (T= 9772 K, log g = 4.0). The thick solid black line is the best fit MCFOST model. *right*) AB Aur spectra from 2.4 – 5 μm overlaid with the literature photometry (blue), and the LMIRCam H2O-Ice2 (thin black line) and L' filters (magenta line). Synthetic photometry for each filter is shown by the corresponding colored star.

in L' band is clearly seen in panels c) and d) as peaks in the radial profile, while the structure in K_s and ice bands matches that of Hashimoto et al. (2011). From these profiles, we also see that the Ice band is fainter than K_s and L' at all distances, lending support to absorption by icy grains. Finally, in panels e) and f), we measure the color excess of the scattered light from the disk (see Section 4.3 for more details). We find that the disk is markedly red in $\Delta(K_s - L')$ at all separations, indicating a positive spectral slope. Conversely the disk is blue in $\Delta(K_s - \text{H}_2\text{O})$ at most separations, suggesting the presence of water ice in the disk surface layer. Red $K_s - L'$ disk colors have been associated in the literature with large grains that strongly forward scatter at shorter wavelengths; the grain scattering efficiency at these short wavelengths is lowered causing the longer wavelength light (L') to appear brighter (Mulders et al. 2013; Tazaki et al. 2019). We discuss the strong red disk color in Section 5.3.

4.2. Icy Grains Across the Disk

The model predictions of Honda et al. (2016) and Tazaki et al. (2021a) suggest that the depth of water ice absorption across the disk can vary even if the ice abundance remains fairly consistent. However, we find that the observed spectral dip in Figure 7 remains relatively constant across the disk, though there is a decrease in

depth along the SE direction. This can be seen qualitatively in the images, as the surface brightness of the 3.08 μm Ice band in Figure 1 is dimmer along the major axis (NE-SW). The radial profiles along the NE-SW direction also show dimmer surface brightness in the 3.08 μm ice line from 50 – 150 AU compared to the opposite direction. The SE direction shows the shallowest absorption feature. Since the depth of the absorption feature depends on the scattering angle (Tazaki et al. 2021a) for anisotropic scattering, this could be a result of the inclination of the disk, since the near-side of the disk is the SE.

The scattered light absorption feature at 3.08 μm is an effect of the wavelength dependence of the albedo for optically thick disks (Inoue et al. 2008, see their Fig 2). Dust albedo is dependent on both grain size and composition (Mulders & Dominik 2012). The overall disk scattered light surface brightness, in turn, depends on both the dust albedo and the geometry of the disk, especially the flaring index, since it is the amount of starlight scattered off the disk as a function of the radius (Mulders et al. 2013). We find the absorption feature to be strongest in the outer regions along the major axis (NE-SW), where the flaring is the most pronounced, which could imply more icy grains at the disk surface.




Figure 4. **a)** *Main panel:* K_s band total intensity imagery from this work overlain with H -band polarized intensity contours from Hashimoto et al. (2011) (cyan) and Boccaletti et al. (2020) (magenta). *Right hand inset:* The three individual K_s band total intensity morphology closely matches that of H -band PI imagery, for which PSF subtraction is unnecessary, and suggests that KLIP-RDI is able to accurately recover the disk surface brightness morphology. **b)** *Main panel:* K_s band total intensity imagery overlain with contours of the H_2O $3.08 \mu m$ band (cyan), and L' band (magenta). *Right hand inset:* The three individual images are on equivalent scales (top to bottom: K_s , $3.08 \mu m$, and L'). These reveal that the observed H_2O $3.08 \mu m$ morphology most closely resembles K_s band. This suggests that the scattering surfaces at these two wavelengths are likely at similar depths in the disk surface layers. The poor qualitative and quantitative match of the L' morphology to the other two wavelengths suggests that the L' scattering surface is likely located deeper in the disk surface layer.

4.3. Spectro-Photometry

We construct three band scattered light photometric spectra along the major and minor axes in the NE (position angle $PA = 54^\circ$), SW ($PA = 234^\circ$), and SE ($PA = 144^\circ$) directions. The images are deprojected by




Figure 5. L' KLIP-RDI reduced image of the disk of AB Aur overlaid with the locations of the spiral arms from Boccaletti et al. (2020) in green. The spiral arms align with areas of brighter flux in the L' image indicating they are substructure in the disk and not artifacts from data reduction.

the assumed inclination ($i = 23.2^\circ$; Tang et al. 2017) in order to cleanly extract profiles without inclination biases. The NW direction was excluded due to apparent negative flux (a result of disk oversubtraction) in the L' band reduction in that direction.

As shown in Figure 7, spectra are extracted at 4 radial separations along each axis in $0''.132 \times 0''.066$ wide rectangles. These rectangles are centered at $0''.528$, $0''.660$ and $0''.792$. The inset images in each panel of Figure 7 show the location of each extracted region overlain on the $3.08 \mu m$ disk image. We see a dip in flux at $3.08 \mu m$ relative to K_s and L' at all distances from AB Aur. This deficit is not present in the spectrum of the central region (Figure 3), suggesting that the presence of icy grains leading to decreased scattering efficiency at this wavelength. As evidenced by Figure 6e and f, the slope from K_s to L' is consistently positive in the NE, SW, and in the SE and grows steeper at larger radial distances in the NE. In the southern portion of the disk the $\Delta(K_s - L')$ color averages around 1.08 ± 0.40 mag from 50-150 AU, while in the NE it increases from an average color of 1.61 ± 0.44 mag to a maximum color of 4.04 ± 0.02 mag by 150 AU away from the central star. This strong red color is seen in the steep slope by $0''.792$ in the NE spectra in Figure 7. Compared to the stellar $K - L$ color of 0.99 ± 0.13 mag, the disk is consistently redder indicating we are seeing more than just stellar radiation influencing the disk color.

Following Honda et al. (2009), Honda et al. (2016) and Tazaki et al. (2021a), we determine the extent to which the observed $3.08 \mu m$ absorption is consistent with the presence of water ice grains, as well as the physical properties of these icy grains, using a color-difference dia-

gram (Inoue et al. 2008). The original model predictions derived in Inoue et al. (2008) are calculated for the K , L' , and H_2O ice ($3.02 - 3.16 \mu\text{m}$ rectangular) filters on Subaru/CIAO instead of LBT/LMIRCam, and as they assume isotropic scattering (Tazaki et al. 2021a), may overpredict ice abundance as well. We therefore compare our observations to MCFOST anisotropic scattered light disk model spectra. The method to compute MCFOST models is described in Section 4.4. To derive K_s , H_2O ice, and L' magnitudes from the MCFOST models (in order to calculate color), we convolve the model spectra with LBTI/LMIRCam filter profiles (as discussed in Section 3.2) for each filter. As we only want to see differential scattering imposed by the grains themselves, and not the intrinsic colors of the source of scattering photons (AB Aur), we remove its contribution following Honda et al. (2009) as:

$$\Delta(K_s - \text{H}_2\text{O}) = (K_s - \text{H}_2\text{O})_{sca} - (K_s - \text{H}_2\text{O})_{source} \quad (1)$$

$$\Delta(K_s - L') = (K_s - L')_{sca} - (K_s - L')_{source}, \quad (2)$$

where *sca* represents the color of scattered light and *source* is the intrinsic color of AB Aur. We show model predictions, observed colors for each region, as well as the combined average of all regions in Figure 8. We have corrected for extinction ($A_V = 0.431$; Vioque et al. 2018) using the extinction law of Cardelli et al. (1989).

While the observed $\Delta(K_s - L')$ colors at all computed locations in the disk lie redward of the predicted models, they are closest to models of an optically thick disk with a 0 – 15% ice mass fraction for grain sizes $> 3 \mu\text{m}$. This prediction is in agreement with grains sizes predicted by Perrin et al. (2009) based on their detailed scattered light modeling and within the large range ($0.1 - 40 \mu\text{m}$) predicted by Bouwman et al. (2000) for a cold 50 – 400 K disk from 28 – 175 AU. Most of the grains are bluer in $\Delta(K_s - L')$ in the SE direction, and redder in the NE. We find that the colors along the major and minor axes vary in $\Delta(K_s - L')$ color; the NE direction is redder compared to the southern part of the disk.

The original models by Inoue et al. (2008) did not utilize a grain size distribution, assuming instead a “typical” particle size. Tazaki et al. (2021a) found that the grain-size distribution plays an important role in explaining both the depth of absorption and red $K_s - L'$ scattered light colors of previous observations. When they rederived the Inoue et al. (2008) results using anisotropic scattering, they found that a) a log-normal size distribution can better explain redder scattered light, and b) the wavelength dependence of the phase function in turn helps increase the depth of the ice ab-

sorption feature for larger ($> 1 \mu\text{m}$) grains. This is due to the decreased amount of the angular distribution of the light intensity at $3.08 \mu\text{m}$ compared K and L' for scattering angles $> 30^\circ$. However, though the models utilize Mie scattering, our results still lie redward of the model extremes in $\Delta(K_s - L')$.

4.4. Ice Abundance and Grain Size

We compare the observed scattered light photometric spectra to model spectra for various size distributions, grain sizes, and ice abundances. We use the fixed model parameters from Table 4), and an inclination of 23.2° (Tang et al. 2017), PA of 328° (Tang et al. 2017), and scale height of 14 AU (Perrin et al. 2009). We consider three different maximum grain sizes: 1.0, 3.0, and $10.0 \mu\text{m}$ and four different grain size distribution power law indices (q ; $dN/da \propto a^q$): -2.0 , -2.5 , -3.0 , -3.5 . Changing the grain size distribution, should, in effect, change the amount of large grains and therefore the spectral slope. We use the same astronomical silicates and ice refractive indices, mass abundances, and material densities as in Inoue et al. (2008). The refractive indices for the astronomical silicates are from Draine & Lee (1984) while the ices are from Irvine & Pollack (1968). The components are mixed using an effective mixing theory following the Bruggeman mixing rule to obtain the effective optical index of the new “mixed” grain. The optical properties are then obtained via Mie theory. We assume a porosity of 60% (Perrin et al. 2009) and an inner radius of 0.3 AU. To change the relative mass abundance of ices to silicates, we adjust the mass fraction within each grain so that there are either 0%, 5%, 15% or 68% water ice by mass, and the rest composed of astrosilicates. In order to confirm the validity and global energy budget of the models, we validate the modeled SEDs against the observed SED of AB Aur (see Figure 3). We do not resolve the innermost disk emission (within 0.3 AU, < 1 pixel) in our LMIRCam data. Therefore, we have added a 1900 K blackbody inner disk component (Tannirkulam et al. 2008) to the stellar spectrum to include this additional excess NIR flux seen in the observed spectrum.

We synthesize scattered light images of the disk using MCFOST for this small grid (q , a_{max} , and ice mass fraction) from $2 - 4 \mu\text{m}$ with a resolution of $0.1 \mu\text{m}$. Each image is convolved with a Gaussian PSF with FWHM equal to the FWHM of the nearest LMIRCam data (since the observed reference PSF stars are only in 3 distinct wavelength bands). We then extract the scattered light spectra at $0''.792$ from these model images. Finally, we interpolated over these spectra in order to




Figure 6. Radial profiles along the major (left) and minor (right) axis of the disk of AB Aur. K_s band is shown in blue, $3.08 \mu\text{m}$ Ice is shown in black, and L' is shown in pink. **top)** Disk to star contrast of the observations as a function of distance from the central star. **middle)** Surface brightness of the disk as a function of distance from the central star. **bottom)** Scattered light color excess as a function of distance from of the central star. $\Delta(K_s - \text{H}_2\text{O})$ is shown in green, while $\Delta(K_s - L')$ is shown in purple.

synthesize a continuous spectral model; the results are shown in Figure 9.

For grains composed of 0–68% water ice by mass, the mass abundance does not greatly affect the $K_s - L'$ color; however, the deficit at $3.08 \mu\text{m}$ increases by almost an order of magnitude as the icy grain abundance increases from 0 to 15% icy grains. We also find that an increase in maximum grain size has the opposite effect, causing the absorption at $3.08 \mu\text{m}$ to decrease. Decreasing the size distribution power law index does redden the spectral slope; however, its effect is small relative to the change in maximum grain size, which reddens the slope by almost 50% at L' from $1 \mu\text{m}$ to $10 \mu\text{m}$.

From Figure 7, we find that the $\Delta(K_s - \text{H}_2\text{O})$ slopes are relatively consistent. Therefore, to compare the observed spectra to the above models, we show the spectra at $0''.792$ for each direction. We first normalize the spectra to K_s (left panel of Figure 9) and find

that the observed surface brightness from K_s to H_2O in the SW (square) direction can be reproduced with a model with 5% icy grains by mass with a grain size $> 3 \mu\text{m}$ and $q = [-2.0, -3.5]$ ($r^2 = 0.98$). In the SE (diamond) and NE (circle) direction, a model with 5% icy grains by mass with a grain size $> 10.0 \mu\text{m}$ and $q = [-2.0, -3.0]$ can reproduce the observed surface brightness ($r^2 \sim 0.91$). However, models with no icy grains can also reproduce observed ice surface brightness. This is easily seen in the top panel of Figure 10, where the $\Delta(K_s - \text{H}_2\text{O})$ color is most consistent with models with grains composed of 0–5% ice by mass and grain sizes $\gtrsim 3 \mu\text{m}$. We discuss the implications of having a grain population composed of mostly large grains below and in Section 5.2.

These results could indicate that the observed disk surface is ice-poor and primarily composed of larger grains, perhaps due to intense photoevaporation or tur-




Figure 7. Surface brightness spectra along with NE (right), SW (center), and SE (left) direction of the disk of AB Aur. Each spectra is extracted within a $0''.132 \times 0''.066$ wide rectangle. The spectra are centered at $0''.528$, $0''.660$ and $0''.792$ along each axis. The inset $3.08 \mu\text{m}$ Ice band deprojected images shows the location of each box along each direction. In all cases, a dip at $3.08 \mu\text{m}$ is seen.




Figure 8. Color-color difference diagram of the MCFOST scattered light spectra from Figure 7. The black lines are models for maximum grain sizes of 1.0 , 3.0 , and $10.0 \mu\text{m}$ (left to right), while the dashed lines show the ice mass fraction (circle - 0% , diamond - 5% , square - 15% , triangle 68% .) The individual colored markers show the color difference for each extracted spectra (SE: orange diamond, NE: olive circle, SW: blue square) from Figure 7, with the size increasing further with distance from the star. The black square indicate the average of all spectra. The black arrow is the interstellar reddening vector.

bulence. None of the models we constructed were able to reproduce the observed red $\Delta(K_s - L')$ color while normalized to K_s band, indicating the power law size distribution may not be sufficient to describe this slope. The same conclusion was made by Tazaki et al. (2021a), who found that a log-normal distribution reproduces red scattered light profiles. This is because the power law size distribution is dominated by $1 \mu\text{m}$ grains, causing the scattered light spectra to become almost gray (the albedo and scattering cross section are wavelength independent for $\gtrsim 1 \mu\text{m}$ grains Tazaki et al. 2021a). If we instead assume a log-normal distribution will match the observed slope, then Tazaki et al. (2021a) claims that a top-heavy power law (e.g. smaller power law index) distribution should give the closest match to the log-normal distribution as there will be more large grains. However, as shown in Figure 9, even decreasing the power law index does not redden the slope enough, indicating the redward slope might have another cause (see Section 5.3).

Therefore, we also normalize the spectra to L' , as shown in the right panel of Figure 9, to quantify the fraction of icy grains necessary to match the $\Delta(\text{H}_2\text{O} - L')$ slope. We find that only grains that are $\gg 5\%$ water ice by mass can reproduce the observed spectral slope. This is highlighted in the bottom panel of Figure 10, where the color is consistent with models with grains $\sim 68\%$ ice by mass and grain sizes $> 1 \mu\text{m}$. As we discuss in Section 5, this composition is unlikely within 200 AU , lending credence to our assertion that the L' scattering

surface is likely at a different location in the disk surface layer.

5. DISCUSSION

We have found a shallow observed dip at the $3.08\ \mu\text{m}$ ice line relative to K_s , and a deeper observed dip relative to L' , suggestive, though unlikely, of more icy grains by mass. When compared to models, the shallow dip relative to K_s suggests that the surface is very ice-poor with a ice mass fraction of $\sim 0 - 5\%$ if the grains are at least $3\ \mu\text{m}$ in size.

5.1. Comparison to HD 142527 and HD 100546

As only two other disks, HD 142527 and HD 100546, have been studied with the water ice deficit technique (Honda et al. 2009, 2016; Tazaki et al. 2021a), we discuss the similarities among the three disks below, with their relevant properties listed in Table 3. Like AB Aur, HD 142527 and HD 100546 are both Herbig Ae/Be stars with strong absorption at $3.09\ \mu\text{m}$, as well as a red $\Delta(K - L')$ slope.

HD 100546 shows changes in inferred feature depth as a function of distance from the star. Honda et al. (2016) found that the optical depth increases from ~ 0.5 to ~ 1.0 further from the star, potentially due to an increase in ice grains outward in the disk. They also find strong asymmetry in the ice absorption optical depth along the minor axis of the disk, which they attribute to the disk geometry; larger optical depth for the near side of the disk. However, Tazaki et al. (2021a) found the opposite to be true, concluding that this asymmetry must be due to ice abundance variation. AB Aur does not have a strong asymmetry; there is an increased optical depth along the major axis which decreases along the minor axis. The optical depth also decreases closer to the central star, which could be due to a decrease in icy grains. These disks show remarkably similar ice abundances and ice line optical depths that could merely be due to their similar evolution and stellar properties.

Honda et al. (2009) found a ice to silicate mass ratio of > 2.2 along the SW of the disk of HD 142527, though Tazaki et al. (2021a) attributes this high ratio to the assumption of isotropic scattering in the Inoue et al. (2008) models used to interpret the data. With the inclusion of anisotropic scattering, Tazaki et al. (2021a) find a mass ratio of $0.06 - 0.2$, similar to the ratio of 0.05 found in AB Aur. They also find a reddish scattered spectral similar to AB Aur, through the disk of AB Aur is far redder at L' band compared to HD 142527 (see Table 3). The predicted surface snowline of HD 142527 is located between $100 - 300\ \text{AU}$ according to models of photodesorption from far-ultraviolet photons from Oka

Table 3. Comparison of Disk Properties

	AB Aur	HD 142527	HD 100546	Refs ^a
inclination (deg)	23.2	28 ± 0.5	32-42	(1)/(2)/(3)
age (Myr)	4 ± 1	~ 1	> 10	(4)/(5)/(6)
avg $\Delta(K - \text{H}_2\text{O})$	-1.11 ± 0.22	-1.01 ± 0.41	–	(7)/(8)/–
avg $\Delta(K - L')$	1.27 ± 0.61	0.44 ± 0.28	–	(7)/(8)/–
ice/silicate mass ratio	0-0.05	0.06-0.2	0.06-0.2	(7)/(9)/(9)

^aReference for AB Aur/HD 142527/HD 100546

References—(1) Tang et al. (2017); (2) Perez et al. (2015); (3) Pineda et al. (2019); (4) Rodriguez et al. (2014); (5) Fujiwara et al. (2006); (6) Grady et al. (2001); (7) this work; (8) Honda et al. (2009); (9) Tazaki et al. (2021a)

et al. (2012). Tazaki et al. (2021a) attributes the low ice abundance to being near the surface water snowline. Overall, these disks are quite similar in ice abundance, composition, and color. More work needs to be done to understand if these are typical or unique in their grain properties among Herbig Ae/Be stars.

5.2. Red Scattered Light Spectra

Though the red $\Delta(K_s - L')$ scattered light color that we see in AB Aur has also been seen in the disks of HD 142527 and HD 100546, blue NIR to L' disk colors have also been detected (e.g. GG Tau; Duchêne et al. 2004). This extreme color variation among disks at similar evolutionary stages highlights the complex nature of scattering and disk surface brightness. Differences in grain properties (grain size, porosity, vertical dust settling, polarization fraction), as well as wavelength dependencies of the albedo and scattering phase function can all affect the NIR colors of the disk (Duchêne et al. 2004; Mulders et al. 2013; Tazaki et al. 2019). Therefore, we review a range of factors that could explain the red scattered light color of AB Aur below.

We first quantify how much local extinction, beyond the $0.431\ \text{mag}$ observed interstellar extinction for AB Aur (Vioque et al. 2018), would be necessary to produce the observed $\Delta(K_s - L')$ slope. As shown in Figure 11, if we deredden the observed spectra by various values of A_V , we are able to match the modeled extinction free slope. However, we find that this requires $A_V \sim 40 - 50\ \text{mags}$, which is nonphysical. Therefore, other explanations beyond local extinction are required to reproduce the red $\Delta(K_s - L')$ scattered light color.

Tazaki et al. (2021a) attributes the red scattered light color of AB Aur to the spiral structure in the disk (Hunziker et al. 2021). One possibility for the formation of these spirals arms is that they result from gravitational instabilities (GI) in the disk. GI can help bring large grains to the surface of the disk, which then forward




Figure 9. Scattered light spectra at a radius of $0''.792$ normalized to K_s band (lefthand column) or L' band (righthand column) overlain with modeled spectra for various ice abundances. The points indicate the relative photometry in the SE (diamond), NE (circle), and SW (square) directions. Models with maximum grain sizes of 10 (green), 3.0 (gold), and $1.0\ \mu\text{m}$ (purple), are shown overplotted for a range of power law grain size distribution indices, namely: -2.0 (dotted), -2.5 (solid), -3.0 (dash-dotted), -3.5 (dashed). Vertically, the panels range from high to low ice mass composition fractions, with values of 68%, 15%, 5%, and 0% ice. Grains that are 0% or 5% ice best match the K_s - L' slope (lower left panels), while the $3.08\ \mu\text{m}$ - L' slope is best matched by grains with very high ice fractions ($\sim 68\%$, upper right panel).

scatter light at short wavelength thereby reducing the scattering efficiency of grains and causing the red slope (Tazaki et al. 2019, 2021a; Mulders et al. 2013). Recent studies (Dullemond et al. 2019; Kuffmeier et al. 2020) claim that the spiral arms around AB Aur are formed from late infall of material on the disk, and therefore, there should be blue-scattering only in the NIR (Tazaki et al. 2021a).

However, there are two caveats that can help explain our redward slope. First, previous observations of the disk of AB Aur have not gone as red as L' band. Therefore, while the disk of AB Aur shows a blueward slope in the bluer end of the NIR supporting the claim of blue-scattering, the spiral arms in L' are quite prominent and could help influence the slope at the redder end of the NIR. Second, the most prominent spiral arms in AB Aur extend outward from 200 AU to 400 AU (Fukagawa




Figure 10. $\Delta(K_s - H_2O)$ color (top) and $\Delta(H_2O - L')$ color (bottom) of scattered light spectra as a function of grain size. The dashed lines show the ice mass fraction for MCFOST models with grain size distribution power law index $q = -3.5$. The solid horizontal lines are the colors of the AB Aur disk with LMIRCam (SE: orange, NE: olive, SW: blue) in Figure 9. The best-fitting ice mass fraction differs substantially between the two colors from 0 – 15% in $\Delta(K_s - H_2O)$ to $\sim 68\%$ in $\Delta(H_2O - L')$.

et al. 2004; Boccaletti et al. 2020) in the outer disk. The spirals we see in L' are within 200 AU, in the innermost part of the disk and could therefore not be part of the later infalling material, and instead be formed from gravitational instability (Cadman et al. 2021). Therefore, the reddish scattered light in the inner disk of AB Aur could be explained by the spiral arms effectively diffusing large grains to the disk surface, similar to HD 142527 and HD 100546.

(Hunziker et al. 2021) found that for HD 142527, the red scattered light spectrum of the disk could be explained by a simple scattering model with strong forward scattering induced by large, highly porous aggregate dust grains. They also found that this dust model was able to explain the high polarization fraction seen in HD 142527. Due to the strong similarities between HD 142527 and AB Aur (disk inclination, age,




Figure 11. Effect of dereddening on the AB Aur spectra in order to match the modeled scattered light spectra. The darker colors indicate a larger dereddening correction (A_V). The models are as in Figure 9 for an ice mass composition fraction of 5% ice. L' must have an extra ~ 45 –50 mags of extinction to get its observed spectral slope.

spectral type), we postulate that a population of large porous aggregate grains could also help to explain the red $\Delta(K_s - L')$ color and the high polarization fraction ($\sim 40\%$; Perrin et al. 2009) seen in the disk of AB Aur. On the other hand, Tazaki et al. (2021b) found from scattered light disk simulations that while porous grains will have higher polarization fractions (Kirchschlager & Wolf 2014; Min et al. 2012), only large compact dust grains (with lower polarization fractions) can produce red scattered light. Detailed dust grain modeling of AB Aur is needed to disentangle the relationship between porosity, disk color, and polarization fraction.

5.3. Optical Depth Effects

We investigate the varying ice abundances necessary to explain the $\Delta(K_s - H_2O)$ and $\Delta(H_2O - L')$ colors. The differences in ice mass fraction between $\Delta(K_s - H_2O)$ and $\Delta(H_2O - L')$ colors along with the strong morphological differences could be a result of these wavelengths probing different depths of the disk atmosphere. This is supported by the fact that the K_s and L' images show marked differences in shape and morphology (notably the presence of spiral arm features, though their apparent absence in K_s could be a result of its reduced brightness contrast to the surrounding disk at shorter wavelength along with its more limited sensitivity). If the opacity of the disk is reduced at L' , then scattered light at this wavelength will probe deeper into the disk, allowing us to image the spirals. If this is the case, then the dust populations probed at K_s and L' are likely different in size and/or composition, as these properties should have radial and vertical gra-

dients within the disk. This is our favored explanation for the ensemble of observations, as it explains both the apparent morphological differences and the fact that the three wavelength photometry is not well-matched to any theoretical predictions for icy grains.

Visually, we find that the K_s and the $3.08\ \mu\text{m}$ ice images are morphologically similar (Figure 4b), indicating they likely probe nearby components of the disk surface, though the ice component is most likely at a somewhat lower depth given its longer wavelength. The L' morphology is markedly different from both K_s and $3.08\ \mu\text{m}$, suggesting that it may be scattered from significantly deeper in the disk. The discrepancy in the icy grain fractions required to match the $\Delta(K_s - \text{H}_2\text{O})$ (0–5%) and $\Delta(\text{H}_2\text{O} - L')$ (68%) colors lend additional support to this conclusion. Given that an ice to silicate mass ratio of 68% is likely nonphysical on the disk surface (especially if there is strong photodesorption; see Section 5.4), our hypothesis that the K_s and $3.08\ \mu\text{m}$ scattering surfaces are roughly co-located is further supported. However, care must be taken when interpreting both the $K_s - L'$ and $\text{H}_2\text{O} - L'$ slopes, and more work is needed to disentangle optical depth effects from K_s to L' .

5.4. Temperature Dependence on the Survival of Surface Icy Grains

Rivière Marichalar et al. (2020) derived the surface temperature of the disk of AB Aur using $^{12}\text{CO}(1-0)$ gas kinematics. They estimate a temperature of 70 K near the central star, decreasing to 10 K by 650 AU. The average dust temperature across the disk is around 25–30 K (Piétu et al. 2005; Woitke et al. 2019); though warmer than other similar disks (e.g. HD 142527), it is still below the temperature necessary for water ice desorption ($\sim 120 - 180$ K). Due to this, the detection of ice absorption could be possible. For these large icy grains to exist on the disk surface, turbulent diffusion or strong disk winds (via photoevaporation) must be evoked to explain their presence, as evidenced by Tazaki et al. (2021a).

For Herbig Ae/Be stars, Agúndez et al. (2018) found that the photodissociation rate of water ice at 100 AU is $1 \times 10^{-4} - 1 \times 10^{-5}\ \text{s}^{-1}$ ($\sim 310 - 3100\ \text{yr}^{-1}$). This is driven by stellar radiation entering the disk at a low grazing angle in flared disks, causing it to be the dominant source of radiation on their surfaces. By 4 Myr (approximate age of AB Aur), water at the surface of the disk is highly dissociated, and the vertical extent of icy grains is controlled by photodesorption, shifting the snowline down and inward (or pushing the surface snow line out). Models by Oka et al. (2012) found pho-

todesorption to be quite efficient for high temperature ($T \sim 10000$ K) and luminosity ($L \sim 40\ L_\odot$) Herbig Ae/Be stars (see their Figure 9).

At these high temperatures, they conclude that the surface snow line should be significantly further away from the central star, > 1000 AU, due to the high rate of photodesorption. Icy grains should, in effect, not exist on the surface of disks around hot stars (where $T_* > T_c$, the critical temperature between sublimation/condensation that controls the radius of the snowline and photodesorption/condensation mechanisms), as the ice-condensation front (where condensation of icy grains is equal to the rate of photodesorption) forms at the surface of the disk. AB Aur, with a temperature ~ 9700 K and luminosity of $\sim 40\ L_\odot$, has a critical temperature of $T_c \sim 4500$ K assuming an average dust temperature of 36 K (Woitke et al. 2019), and should therefore have exceedingly strong photodesorption. As shown by Figure 9, the $\Delta(K_s - \text{H}_2\text{O})$ color aligns with models with 0–5% icy grains by mass (and is the more likely ice abundance) indicating surface water ice could be affected by photodesorption. If we assume that the K_s and $3.08\ \mu\text{m}$ ice components are at approximately the same location on the disk surface, then the 0% icy grains by mass within 150 AU is consistent with Oka et al. (2012). However, though small, models with 5% icy grains by mass do match the observed $\Delta(K_s - \text{H}_2\text{O})$ color, indicating that if the ice surface is at a slightly lower disk surface depth compared to K_s , the icy grains at ~ 100 AU could both exist and be shielded from some photodesorption on the surface.

5.5. H_2O Snowline in Relation to Planet Formation in AB Aur

Within the disk of AB Aur, two indirectly inferred planet candidates have been reported at 30 and 140 AU (Boccaletti et al. 2020). The inner planet candidate mass is tentatively estimated at $4 - 13\ M_{\text{Jup}}$ (Boccaletti et al. 2020; Piétu et al. 2005; Tang et al. 2017), while the outer is tentatively estimated at $3\ M_{\text{Jup}}$ (Boccaletti et al. 2020); the interpretation is indirect, based on a substructure that can be interpreted as a planet-induced perturbation (but other explanations are plausible; Jorquera et al. 2022, in press). The estimated masses and locations of the planets are consistent with formation through gravitational instability Cadman et al. (2021).

Our detection of reduced ice absorption at ~ 100 AU would place the potential outer companion within the region where we see little to no ice absorption. Far-infrared spectroscopic observations of AB Aur (Fedele et al. 2013) did not detect H_2O emission, indicating that water molecules in the outer disk beyond the snow line

have likely been photodissociated by stellar radiation. Fedele et al. (2013) conclude that this would result in a lack of water reservoir in the outer disk necessary for giant icy planet core formation. Since the location of the midplane snowline has not been measured, it is not certain if there are icy volatiles at this separation in the disk midplane. Therefore, whether or not there are enough icy volatiles for icy core formation for an outer planet is unknown, and would require further observations and detections of volatile snow lines.

6. CONCLUSION

In this paper, we observe the (pre)transitional disk around AB Aur in order to explore the presence of water ice in the surface layers of the scattered light disk. Using LBTI LMIRCam at K_s , the $3.08\ \mu\text{m}$ water ice line, and L' , we look for a deficient of NIR scattered light at the ice line, as icy grains absorb preferentially at this wavelength. We use the $\Delta(K_s - \text{H}_2\text{O})$ and $\Delta(K_s - L')$ color differences in order to probe the grain size distribution and ice abundance within this disk. We find the following.

1. We detect the disk of AB Aur at high SNR in all three LBTI LMIRCam bands, K_s , $3.08\ \mu\text{m}$ ice, and L' . After removing the starlight using KLIP-RDI, the disk is visible out to 175 AU ($1''.1$). We find strong morphological differences between K_s and L' band along with prominent L' inner spiral arms. The K_s image closely resembles the morphology found in H polarized intensity imaging. The $3.08\ \mu\text{m}$ ice band image has a closer morphology to K_s than L' .
2. After extracting scattered light photometry along the major and minor axes for each band, we detect absorption at $3.08\ \mu\text{m}$ from 90 – 120 AU. Using a color-color difference diagram (Inoue et al. 2008), we argue that our observations are consistent with the presence of icy grains on the surface of the disk of AB Aur with sizes $3\ \mu\text{m}$.
3. By comparing to MCFOST models of varying ice abundances and maximum grain sizes, we find the $\Delta(K_s - \text{H}_2\text{O})$ slope most closely matches a model with a ice/silicate mass ratio ~ 0.05 (5% icy grains by mass) with a maximum grain size of 3 – 10 μm . The reddish K_s and L' continuum slope can be explained by the presence of these large grains (though the MCFOST models cannot predict this slope). However, these large grains can only exist on the disk surface due to either disk turbulence or winds (via photoevaporation) preventing grain settling.
4. We argue that the strong $\Delta(K_s - L')$ slope seen in the scattered light spectra is most likely to be a result of probing different scattering surfaces at different depths within the disk with the L' disk component probed at a deeper disk surface. Further investigation is necessary to update model predictions of disk colors under the assumption of differential optical depths between NIR wavelengths.

These results suggest that the surface of the disk of AB Aur is highly ice-poor resulting from strong photodesorption of large grains. At the higher stellar temperature and luminosity of AB Aur, the surface snowline lies beyond 100 AU as the stellar radiation field pushes the ice condensation front outward along the surface, potentially as far as 1000 AU (far beyond our detection limits). The small (0 – 5%) presence of icy grains at the disk surface is suggestive of vertical mixing in the disk or that the grains are shielded from photodesorption (due to optical depth effects) in order to survive. Therefore, if we want to strongly detect the surface snow line, sensitive observations of the outer disk, as well as millimeter observations of the midplane snow line, are necessary.

ACKNOWLEDGMENTS

We thank Esther Buenzli, Vanessa Bailey, Timothy Rodigas, and Jared Males for contributing to or obtaining the original imagery. We thank the anonymous referee for their careful review. We thank Kellen Lawson for his helpful suggestion to look into the differential evolution algorithm as a way to fit the forward modeled disks. SKB and KBF acknowledge support from NSF AST-2009816. S.J. acknowledges support from the National Agency for Research and Development (ANID), Scholarship Program, Doctorado Becas Nacionales/2020 - 21212356. L.P. gratefully acknowledges support by the ANID BASAL project FB210003, and by ANID, – Millennium Science Initiative Program – NCN19_171. We acknowledge the use of the Large Binocular Telescope Interferometer (LBTI) and the support from the LBTI team. MCFOST is funded from the Australian Research Council under contracts FT170100040 and DP180104235, from Agence Nationale pour la Recherche (ANR) of France under contract ANR-16-CE31-0013. This research has made use of the SVO Filter Profile Service (<http://svo2.cab.inta-csic.es/theory/fps/>) supported from the Spanish MINECO through grant AYA2017-84089. This publication makes use of data products from the Wide-field Infrared Survey Explorer, which is a joint project of the University of California, Los Angeles, and the Jet Propulsion Laboratory/California Institute of Technology, funded by the National Aeronautics and Space Administration.

Facilities: LBT(LMIRCam)

Software: pyKLIP (Wang et al. 2015), diskFM (Mazoyer et al. 2020), MCFOST (Pinte et al. 2006, 2009), Astropy (Astropy Collaboration et al. 2013, 2018), numpy,

scipy, matplotlib (Hunter 2007), Astrolib PySynphot (STScI Development Team 2013), Dewarp (Spalding & Stone 2019), DebrisDiskFM (Ren et al. 2019), pymcfost (<https://github.com/cpinte/pymcfost>), mcfost-python (<https://github.com/swolff9/mcfost-python>)

REFERENCES

- Agúndez, M., Roueff, E., Le Petit, F., & Le Bourlot, J. 2018, *A&A*, 616, A19, doi: [10.1051/0004-6361/201732518](https://doi.org/10.1051/0004-6361/201732518)
- Aikawa, Y., Kamuro, D., Sakon, I., et al. 2012, *A&A*, 538, A57, doi: [10.1051/0004-6361/201015999](https://doi.org/10.1051/0004-6361/201015999)
- Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, *A&A*, 558, A33, doi: [10.1051/0004-6361/201322068](https://doi.org/10.1051/0004-6361/201322068)
- Astropy Collaboration, Price-Whelan, A. M., Sipőcz, B. M., et al. 2018, *AJ*, 156, 123, doi: [10.3847/1538-3881/aabc4f](https://doi.org/10.3847/1538-3881/aabc4f)
- Blevins, S. M., Pontoppidan, K. M., Banzatti, A., et al. 2016, *ApJ*, 818, 22, doi: [10.3847/0004-637X/818/1/22](https://doi.org/10.3847/0004-637X/818/1/22)
- Boccaletti, A., Di Folco, E., Pantin, E., et al. 2020, *A&A*, 637, L5, doi: [10.1051/0004-6361/202038008](https://doi.org/10.1051/0004-6361/202038008)
- Bouwman, J., de Koter, A., van den Ancker, M. E., & Waters, L. B. F. M. 2000, *A&A*, 360, 213
- Cadman, J., Rice, K., & Hall, C. 2021, *MNRAS*, 504, 2877, doi: [10.1093/mnras/stab905](https://doi.org/10.1093/mnras/stab905)
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245, doi: [10.1086/167900](https://doi.org/10.1086/167900)
- Carr, J. S., & Najita, J. R. 2008, *Science*, 319, 1504, doi: [10.1126/science.1153807](https://doi.org/10.1126/science.1153807)
- Chen, C., Mazoyer, J., Poteet, C. A., et al. 2020, *ApJ*, 898, 55, doi: [10.3847/1538-4357/ab9aba](https://doi.org/10.3847/1538-4357/ab9aba)
- Crepp, J. R., Gonzales, E. J., Bowler, B. P., et al. 2018, *ApJ*, 864, 42, doi: [10.3847/1538-4357/aad381](https://doi.org/10.3847/1538-4357/aad381)
- Cutri, R. M., & et al. 2012, *VizieR Online Data Catalog*, II/311
- Cutri, R. M., Skrutskie, M. F., van Dyk, S., et al. 2003, *2MASS All Sky Catalog of point sources*.
- DeWarf, L. E., Sepinsky, J. F., Guinan, E. F., Ribas, I., & Nadalin, I. 2003, *ApJ*, 590, 357, doi: [10.1086/374979](https://doi.org/10.1086/374979)
- di Folco, E., Dutrey, A., Chesneau, O., et al. 2009, *A&A*, 500, 1065, doi: [10.1051/0004-6361/200809902](https://doi.org/10.1051/0004-6361/200809902)
- Draine, B. T., & Lee, H. M. 1984, *ApJ*, 285, 89, doi: [10.1086/162480](https://doi.org/10.1086/162480)
- Drakowska, J., & Alibert, Y. 2017, *A&A*, 608, A92, doi: [10.1051/0004-6361/201731491](https://doi.org/10.1051/0004-6361/201731491)
- Duchêne, G., McCabe, C., Ghez, A. M., & Macintosh, B. A. 2004, *ApJ*, 606, 969, doi: [10.1086/383126](https://doi.org/10.1086/383126)
- Dullemond, C. P., Küffmeier, M., Goicovic, F., et al. 2019, *A&A*, 628, A20, doi: [10.1051/0004-6361/201832632](https://doi.org/10.1051/0004-6361/201832632)
- Fedele, D., Bruderer, S., van Dishoeck, E. F., et al. 2013, *A&A*, 559, A77, doi: [10.1051/0004-6361/201321118](https://doi.org/10.1051/0004-6361/201321118)
- Fujiwara, H., Honda, M., Kataza, H., et al. 2006, *ApJL*, 644, L133, doi: [10.1086/505597](https://doi.org/10.1086/505597)
- Fukagawa, M., Hayashi, M., Tamura, M., et al. 2004, *ApJL*, 605, L53, doi: [10.1086/420699](https://doi.org/10.1086/420699)
- Gaia Collaboration, Brown, A. G. A., Vallenari, A., et al. 2018, *A&A*, 616, A1, doi: [10.1051/0004-6361/201833051](https://doi.org/10.1051/0004-6361/201833051)
- Grady, C. A., Woodgate, B., Bruhweiler, F. C., et al. 1999, *ApJL*, 523, L151, doi: [10.1086/312270](https://doi.org/10.1086/312270)
- Grady, C. A., Polomski, E. F., Henning, T., et al. 2001, *AJ*, 122, 3396, doi: [10.1086/324447](https://doi.org/10.1086/324447)
- Hashimoto, J., Tamura, M., Muto, T., et al. 2011, *ApJL*, 729, L17, doi: [10.1088/2041-8205/729/2/L17](https://doi.org/10.1088/2041-8205/729/2/L17)
- Henning, T., Burkert, A., Launhardt, R., Leinert, C., & Stecklum, B. 1998, *A&A*, 336, 565
- Honda, M., Inoue, A. K., Fukagawa, M., et al. 2009, *ApJL*, 690, L110, doi: [10.1088/0004-637X/690/2/L110](https://doi.org/10.1088/0004-637X/690/2/L110)
- Honda, M., Kudo, T., Takatsuki, S., et al. 2016, *ApJ*, 821, 2, doi: [10.3847/0004-637X/821/1/2](https://doi.org/10.3847/0004-637X/821/1/2)
- Hunter, J. D. 2007, *Computing in Science & Engineering*, 9, 90, doi: [10.1109/MCSE.2007.55](https://doi.org/10.1109/MCSE.2007.55)
- Hunziker, S., Schmid, H. M., Ma, J., et al. 2021, *A&A*, 648, A110, doi: [10.1051/0004-6361/202040166](https://doi.org/10.1051/0004-6361/202040166)
- Inoue, A. K., Honda, M., Nakamoto, T., & Oka, A. 2008, *PASJ*, 60, 557, doi: [10.1093/pasj/60.3.557](https://doi.org/10.1093/pasj/60.3.557)
- Irvine, W. M., & Pollack, J. B. 1968, *Icarus*, 8, 324, doi: [10.1016/0019-1035\(68\)90083-3](https://doi.org/10.1016/0019-1035(68)90083-3)
- Kirchschrager, F., & Wolf, S. 2014, *A&A*, 568, A103, doi: [10.1051/0004-6361/201323176](https://doi.org/10.1051/0004-6361/201323176)
- Kuffmeier, M., Goicovic, F. G., & Dullemond, C. P. 2020, *A&A*, 633, A3, doi: [10.1051/0004-6361/201936820](https://doi.org/10.1051/0004-6361/201936820)
- Lawson, K., Currie, T., Wisniewski, J. P., et al. 2020, *AJ*, 160, 163, doi: [10.3847/1538-3881/ababa6](https://doi.org/10.3847/1538-3881/ababa6)
- Maire, A. L., Skemer, A. J., Hinz, P. M., et al. 2015, *A&A*, 576, A133, doi: [10.1051/0004-6361/201425185](https://doi.org/10.1051/0004-6361/201425185)
- Malfait, K., Waelkens, C., Bouwman, J., de Koter, A., & Waters, L. B. F. M. 1999, *A&A*, 345, 181
- Mazoyer, J., Arriaga, P., Hom, J., et al. 2020, in *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series*, Vol. 11447, Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series, 1144759, doi: [10.1117/12.2560091](https://doi.org/10.1117/12.2560091)

- McClure, M. K., Manoj, P., Calvet, N., et al. 2012, *ApJL*, 759, L10, doi: [10.1088/2041-8205/759/1/L10](https://doi.org/10.1088/2041-8205/759/1/L10)
- McClure, M. K., Espaillat, C., Calvet, N., et al. 2015, *ApJ*, 799, 162, doi: [10.1088/0004-637X/799/2/162](https://doi.org/10.1088/0004-637X/799/2/162)
- Meeus, G., Waters, L. B. F. M., Bouwman, J., et al. 2001, *A&A*, 365, 476, doi: [10.1051/0004-6361:20000144](https://doi.org/10.1051/0004-6361:20000144)
- Min, M., Canovas, H., Mulders, G. D., & Keller, C. U. 2012, *A&A*, 537, A75, doi: [10.1051/0004-6361/201117333](https://doi.org/10.1051/0004-6361/201117333)
- Mulders, G. D., & Dominik, C. 2012, *A&A*, 539, A9, doi: [10.1051/0004-6361/201118127](https://doi.org/10.1051/0004-6361/201118127)
- Mulders, G. D., Min, M., Dominik, C., Debes, J. H., & Schneider, G. 2013, *A&A*, 549, A112, doi: [10.1051/0004-6361/201219522](https://doi.org/10.1051/0004-6361/201219522)
- Natta, A., Prusti, T., Neri, R., et al. 2001, *A&A*, 371, 186, doi: [10.1051/0004-6361:20010334](https://doi.org/10.1051/0004-6361:20010334)
- Notsu, S., Nomura, H., Ishimoto, D., et al. 2017, *ApJ*, 836, 118, doi: [10.3847/1538-4357/836/1/118](https://doi.org/10.3847/1538-4357/836/1/118)
- Notsu, S., Nomura, H., Walsh, C., et al. 2018, *ApJ*, 855, 62, doi: [10.3847/1538-4357/aaaa72](https://doi.org/10.3847/1538-4357/aaaa72)
- Oka, A., Inoue, A. K., Nakamoto, T., & Honda, M. 2012, *ApJ*, 747, 138, doi: [10.1088/0004-637X/747/2/138](https://doi.org/10.1088/0004-637X/747/2/138)
- Oppenheimer, B. R., Brenner, D., Hinkley, S., et al. 2008, *ApJ*, 679, 1574, doi: [10.1086/587778](https://doi.org/10.1086/587778)
- Perez, S., Casassus, S., Ménard, F., et al. 2015, *ApJ*, 798, 85, doi: [10.1088/0004-637X/798/2/85](https://doi.org/10.1088/0004-637X/798/2/85)
- Perrin, M. D., Schneider, G., Duchene, G., et al. 2009, *ApJL*, 707, L132, doi: [10.1088/0004-637X/707/2/L132](https://doi.org/10.1088/0004-637X/707/2/L132)
- Piétu, V., Guilloteau, S., & Dutrey, A. 2005, *A&A*, 443, 945, doi: [10.1051/0004-6361:20042050](https://doi.org/10.1051/0004-6361:20042050)
- Pineda, J. E., Szulágyi, J., Quanz, S. P., et al. 2019, *The Astrophysical Journal*, 871, 48, doi: [10.3847/1538-4357/aaf389](https://doi.org/10.3847/1538-4357/aaf389)
- Pinte, C., Harries, T. J., Min, M., et al. 2009, *A&A*, 498, 967, doi: [10.1051/0004-6361/200811555](https://doi.org/10.1051/0004-6361/200811555)
- Pinte, C., Ménard, F., Duchêne, G., & Bastien, P. 2006, *A&A*, 459, 797, doi: [10.1051/0004-6361:20053275](https://doi.org/10.1051/0004-6361:20053275)
- Pontoppidan, K. M., Dullemond, C. P., van Dishoeck, E. F., et al. 2005, *ApJ*, 622, 463, doi: [10.1086/427688](https://doi.org/10.1086/427688)
- Quanz, S. P., Schmid, H. M., Geissler, K., et al. 2011, *ApJ*, 738, 23, doi: [10.1088/0004-637X/738/1/23](https://doi.org/10.1088/0004-637X/738/1/23)
- Ren, B., & Perrin, M. 2018, *DebrisDiskFM*, v1.0, Zenodo, doi: [10.5281/zenodo.2398963](https://doi.org/10.5281/zenodo.2398963), 1.0, Zenodo, doi: [10.5281/zenodo.2398963](https://doi.org/10.5281/zenodo.2398963)
- Ren, B., Choquet, É., Perrin, M. D., et al. 2019, *ApJ*, 882, 64, doi: [10.3847/1538-4357/ab3403](https://doi.org/10.3847/1538-4357/ab3403)
- . 2021, *ApJ*, 914, 95, doi: [10.3847/1538-4357/ac03b9](https://doi.org/10.3847/1538-4357/ac03b9)
- Rivière Marichalar, P., Fuente, A., Le Gal, R., et al. 2020, *arXiv e-prints*, [arXiv:2008.00751](https://arxiv.org/abs/2008.00751), <https://arxiv.org/abs/2008.00751>
- Rodrigo, C., & Solano, E. 2020, in *Contributions to the XIV.0 Scientific Meeting (virtual) of the Spanish Astronomical Society*, 182
- Rodrigo, C., Solano, E., & Bayo, A. 2012, *SVO Filter Profile Service Version 1.0*, IVOA Working Draft 15 October 2012, doi: [10.5479/ADS/bib/2012ivoa.rept.1015R](https://doi.org/10.5479/ADS/bib/2012ivoa.rept.1015R)
- Rodríguez, L. F., Zapata, L. A., Dzib, S. A., et al. 2014, *ApJL*, 793, L21, doi: [10.1088/2041-8205/793/1/L21](https://doi.org/10.1088/2041-8205/793/1/L21)
- Salyk, C., Herczeg, G. J., Brown, J. M., et al. 2013, *ApJ*, 769, 21, doi: [10.1088/0004-637X/769/1/21](https://doi.org/10.1088/0004-637X/769/1/21)
- Salyk, C., Pontoppidan, K. M., Blake, G. A., et al. 2008, *ApJL*, 676, L49, doi: [10.1086/586894](https://doi.org/10.1086/586894)
- Skrutskie, M. F., Jones, T., Hinz, P., et al. 2010, in *Ground-based and Airborne Instrumentation for Astronomy III*, ed. I. S. McLean, S. K. Ramsay, & H. Takami, Vol. 7735, International Society for Optics and Photonics (SPIE), 1368 – 1378, doi: [10.1117/12.857724](https://doi.org/10.1117/12.857724)
- Soummer, R., Pueyo, L., & Larkin, J. 2012, *ApJL*, 755, L28, doi: [10.1088/2041-8205/755/2/L28](https://doi.org/10.1088/2041-8205/755/2/L28)
- Spalding, E., & Stone, J. 2019, *Dewarp*, Astrophysics Source Code Library. <http://ascl.net/1907.008>
- Storm, R., & Kenneth, P. 1997, *Journal of Global Optimization*, 11, 341, doi: [10.1023/A:1008202821328](https://doi.org/10.1023/A:1008202821328)
- STScI Development Team. 2013, *pysynphot*: Synthetic photometry software package. <http://ascl.net/1303.023>
- Tang, Y. W., Guilloteau, S., Piétu, V., et al. 2012, *A&A*, 547, A84, doi: [10.1051/0004-6361/201219414](https://doi.org/10.1051/0004-6361/201219414)
- Tang, Y.-W., Guilloteau, S., Dutrey, A., et al. 2017, *ApJ*, 840, 32, doi: [10.3847/1538-4357/aa6af7](https://doi.org/10.3847/1538-4357/aa6af7)
- Tannirkulam, A., Monnier, J. D., Harries, T. J., et al. 2008, *ApJ*, 689, 513, doi: [10.1086/592346](https://doi.org/10.1086/592346)
- Tazaki, R., Murakawa, K., Muto, T., Honda, M., & Inoue, A. K. 2021a, *arXiv e-prints*, [arXiv:2108.08637](https://arxiv.org/abs/2108.08637), <https://arxiv.org/abs/2108.08637>
- . 2021b, *ApJ*, 910, 26, doi: [10.3847/1538-4357/abdd3d](https://doi.org/10.3847/1538-4357/abdd3d)
- Tazaki, R., Tanaka, H., Muto, T., Kataoka, A., & Okuzumi, S. 2019, *MNRAS*, 485, 4951, doi: [10.1093/mnras/stz662](https://doi.org/10.1093/mnras/stz662)
- Terada, H., Tokunaga, A. T., Kobayashi, N., et al. 2007, *ApJ*, 667, 303, doi: [10.1086/520951](https://doi.org/10.1086/520951)
- van den Ancker, M. E., Bouwman, J., Wesselius, P. R., et al. 2000, *A&A*, 357, 325, <https://arxiv.org/abs/astro-ph/0002440>
- van den Ancker, M. E., The, P. S., Tjin A Djie, H. R. E., et al. 1997, *A&A*, 324, L33
- Vioque, M., Oudmaijer, R. D., Baines, D., Mendigutía, I., & Pérez-Martínez, R. 2018, *A&A*, 620, A128, doi: [10.1051/0004-6361/201832870](https://doi.org/10.1051/0004-6361/201832870)

- Wang, J. J., Ruffio, J.-B., De Rosa, R. J., et al. 2015, pyKLIP: PSF Subtraction for Exoplanets and Disks. <http://ascl.net/1506.001>
- Weingartner, J. C., & Draine, B. T. 2001, ApJ, 548, 296, doi: [10.1086/318651](https://doi.org/10.1086/318651)
- Woitke, P., Kamp, I., Antonellini, S., et al. 2019, PASP, 131, 064301, doi: [10.1088/1538-3873/aaf4e5](https://doi.org/10.1088/1538-3873/aaf4e5)
- Zhu, Z., Nelson, R. P., Dong, R., Espaillat, C., & Hartmann, L. 2012, ApJ, 755, 6, doi: [10.1088/0004-637X/755/1/6](https://doi.org/10.1088/0004-637X/755/1/6)

APPENDIX

A. GEOMETRIC DISK MODELING

We use previously measured AB Aur observations and model parameters to fix the majority of the non-geometric parameters (e.g. stellar temperature and luminosity, dust mass and composition, grain porosity and radius, grain size power law etc.; Perrin et al. 2009; di Folco et al. 2009; Tang et al. 2017). Following Perrin et al. (2009) and di Folco et al. (2009), we use a mixture of astrosilicate grains (Draine & Lee 1984; Weingartner & Draine 2001) with 60% porosity from $0.005 - 1 \mu\text{m}$ and with grain sizes described by the power law $dN/da \propto a^q$, where the power law index (q) is -3.5 . We set the inner radius to 0.2 AU (Perrin et al. 2009), and the outer radius to the outer edge of each band’s KLIP-RDI image ($K_s = 189 \text{ AU}$, $3.08 \mu\text{m} = 168 \text{ AU}$, $L' = 175 \text{ AU}$). See Table 4. As the purpose of modeling the disk of AB Aur at each wavelength is to determine the throughput and reproduce the observed surface brightness, our primary concern is finding a model that reproduces the brightness and morphology of the disk, and derived parameters may be unphysical.

We investigate four different geometric parameters to achieve a best fit: inclination, (i), position angle of the disk along the minor axis (PA), scale height at 100 AU (h_{100}), and flaring index (β ; $h(r) = h_0(r/r_0)^\beta$ where $r_0 = 100 \text{ AU}$ and $h_0 = h_{100}$). Though all of these parameters have been previously fit for AB Aur, our initial tests (fitting the bands both simultaneously and separately) using best fit values from other work ($i = 23.2^\circ$ (ALMA; Tang et al. 2017), PA = 328° (ALMA; Tang et al. 2017; Perrin et al. 2009), $h_{100} = 14 \text{ AU}$ (Perrin et al. 2009) or $h_{100} = 8 \text{ AU}$ (di Folco et al. 2009), $\beta = 1.3$ (Perrin et al. 2009; di Folco et al. 2009)) provided ill-fitting models. This led us to model the disk at K_s , $3.08 \mu\text{m}$, and L' separately, and without limiting ourselves to physically-motivated parameters.

To forward model, we create $2.16 \mu\text{m}$, $3.08 \mu\text{m}$, and $3.70 \mu\text{m}$ scattered light models of an AB Aur analog disk using the MCFOST Monte Carlo radiative transfer code (Pinte et al. 2006, 2009), which utilizes Mie theory. The MCFOST model is then convolved by the unsaturated PSF of AB Aur and multiplied by an additional fit parameter, a scaling parameter (A). This scaled and convolved image is then forward modeled using the pyKLIP DiskFM package (Mazoyer et al. 2020) to produce a simulated KLIP-RDI reduced disk model under the same KLIP parameters as the observations.

To find the best fit model, we tested a variety of optimization methods, including a Markov-Chain Monte-Carlo (MCMC) and a coarse grid search. The near face-on nature of the disk, together with the distinct and complex substructure between the three bands, indicates parameter degeneracies. Local minima, as well as differences in geometric parameters between bands, is both possible and expected.

We first attempted to fit the three wavelengths separately using MCMC by combining the pyKLIP diskFM implementation from Mazoyer et al. (2020) and Chen et al. (2020) with the MCMC implementation (Ren & Perrin 2018) from Ren et al. (2019) and Ren et al. (2021). However, due to the degeneracy between several variables and a broad parameter space, the MCMC walkers could not find global minima that were qualitatively good fits to the data.

We next attempted a coarse grid search across the four geometric free parameters (i , PA, β , h_{100}) and the scaling amplitude (A). The range in parameter space for each free parameter was determined either from previous literature estimates (β , h_{100} , PA) or from initial model testing (i , A). We initially required the inclination to be fixed at 23.2° , the inclination found from ALMA dust continuum kinematics (Tang et al. 2017); however, this inclination produced a significantly poor fit to the disk at all three wavelengths, while more face-on inclinations ($i < 15^\circ$) produced better fits. Though our coarse grid (5 parameters, 3–20 values for each parameter) found best-fit models, these models were dependent on the grid structure. To expand to a finely sampled grid would be computationally time consuming and would expand to an unfeasible number of models.

Therefore, following Lawson et al. (2020), we turned to a combination of these approaches via a differential evolution optimization (DE; Storm & Kenneth 1997). For a detailed description of the DE algorithm and its application to modeling scattered light disks, see Lawson et al. (2020). In short, DE is a global optimization and search algorithm that can quickly and efficiently explore multi-dimensional, correlated, non-differential and/or nonlinear parameter spaces to find a global minimum. By only exploring within specified parameter boundaries, DE starts by initializing a population of models randomly sampling the parameter space. These models then iteratively evolve (or mutate) towards solutions that best fit the data. This evolution allows the walkers to explore multiple local minima and converge on the global minimum relatively quickly with a smaller (compared to a grid search or MCMC) number of

Table 4. MCFOST Fixed Parameters

Parameter	Value	References
T_{eff} (K)	9772	(1)
R (R_{\odot})	2.5	(2)
M (M_{\odot})	2.4	(2)
R_{inner} (AU)	0.2	(1)
R_{outer} (AU)	175	
M_{dust} (M_{\odot})	1×10^{-4}	(1)
a_{max} (μm)	1.0	(1)
q	-3.5	(1)
Dust Composition	Astrosilicates	(1, 3, 4)

References—(1) Perrin et al. (2009, and references therein); (2) DeWarf et al. (2003); (3) Draine & Lee (1984); (4) Weingartner & Draine (2001)

samples. Following (Lawson et al. 2020), we initialize $N_{\text{pop}} = 50$ models, $10\times$ the number of free parameters ($M=5$). These models are then forward modeled using `pyKLIP` diskFM, as discussed above, and the forwarded models are compared to the reduced observations by computing

$$\chi_r^2 = \frac{\chi^2}{\nu} = \frac{1}{\nu} \sum_S \frac{(\text{Data} - \text{Forward Model})^2}{\text{Uncertainty}^2} \quad (\text{A1})$$

following Chen et al. (2020), where ν is the number of degrees of freedom given by N-M (N is the number of pixels within S), and χ^2 is computed over S , a circular annulus. The uncertainty is measured as discussed in Section 2.2. Then, the population mutates following the "best/1/bin" strategy of Storm & Kenneth (1997) with the mutation constant in the range [0.5, 1.0]. Each mutation replaces its "parent" if a combined version of the two models (determined by changing select individual parameters of the parent with the same parameters of the mutant if a random probability $P \geq 0.7$ is found for that parameter) yields a better fit (e.g. smaller χ_r^2). This mutation, recombination, and replacement is continued until the population converges to a single solution with the global minimum χ_r^2 . Since the population is initialized across the whole grid, the algorithm can robustly explore across local minima without getting trapped.

Since the goal of this model fitting is to match the surface brightness of the disk and not to infer the real, physical disk properties, we are less concerned about finding physically-motivated or matching values between the three bands. For each band, the DE algorithm converged on a best fit solution which minimized χ_r^2 . The boundaries, best fit values, and acceptable ranges for each parameter are shown in Table 5. Following Lawson et al. (2020), models whose solutions are consistent with our observations are those with $\chi_r^2 \leq \chi_{r,\text{min}}^2 + \sqrt{2/\nu}$. We show the minimum χ_r^2 , the maximum χ_r^2 for acceptable solutions, and the number of models to reach convergence in Table 6. Figures 12 - 14 shows the parameter space explored by the DE algorithm with the best fit model shown by the red cross. The models are binned with each color showing the minimum χ_r^2 found in that bin. Figures 15 - 17 shows the best fit model compared to the original data. The residuals are only shown for the region within which χ_r^2 is calculated. Of the three bands, L' had the worst convergence, most likely due to the complicated substructure in L' band, namely the known spiral arms (see Jorquera et al. (2022, in press) and Figure 5). As we assume a simple, radially-symmetric disk model without taking into account the spiral features, the larger L' band residuals are perhaps unsurprising.

B. THROUGHPUT VARIABILITY

The surface brightness and disk colors are dependent on the amount of flux loss from KLIP-RDI. Therefore, we test the variability in throughput on the brightness and geometry of the best fit model disks found in Appendix A in order to see how the throughput will change with varying surface brightness and inclination. This will test the validity of using the best fit models even if the scaling factor or inclination are not perfect matches to the observed data. Using the best fit DE solutions for each band, we vary the amplitude by a factor of 2 and the inclination within $\pm 6 - 14^\circ$. The model is then forward modeled using the same process outlined in Appendix A. We then calculate the azimuthally averaged radial throughput. Figure 18 shows the azimuthally averaged radial profile of the throughput for varying amplitudes and inclinations. The black throughput curve is the best fit model throughput. At 100 AU, where there is

Table 5. Differential Evolution Modeling and Solutions

Parameter	Parameter Boundaries			Best Fit			Acceptable Range		
	K_s	3.08 μm	L'	K_s	3.08 μm	L'	K_s	3.08 μm	L'
i (deg)	5–30	15–25	5–40	11.05	22.46	34.98	10.68–12.04	19.82–24.49	34.62–34.98
PA (deg)	326–330	326–330	326–330	329.29	328.02	329.12	326.66–329.79	326.95–329.95	328.58–329.9
β	1.1–1.95	1.1–1.4	1.1–1.95	1.88	1.35	1.93	1.86–1.95	1.33–1.36	1.93–1.94
h_{100} (AU)	6–35	8–14	4–20	29.99	8.25	6.04	29.57–29.99	8.01–9.06	6.01–6.05
A	0.1–5	0.5–5	6–30	0.63	4.33	10.81	0.58–0.66	3.90–4.70	10.81–11.22

NOTE—PA is along the semi-minor axis, measured east of north. “Acceptable Range” indicates parameter ranges that give $\chi_r^2 \leq \chi_{r,min}^2 + \sqrt{2/\nu}$.

Table 6. Differential Evolution χ^2 values

Band	Minimum χ_r^2	Acceptable χ_r^2	# Total Models
K_s	2.31	2.33	1519
3.08 μm	1.67	1.69	773
L'	10.62	10.64	1593

NOTE—Acceptable χ_r^2 indicates parameter ranges that give $\chi_r^2 \leq \chi_{r,min}^2 + \sqrt{2/\nu}$.

the large deviation in TP, it is still quite small, ranging from 0.001 in L' to 0.15 in K_s . At most, the TP varies by 40% (at 100 AU in K_s , at the AO control radius). However, on average, the TP only varies by 2.5%. Overall, varying the amplitude and inclination affects the TP by a factor of 1.02. Therefore, differences between the models and observed images will not cause large variations in the throughput and therefore disk color.


LMIRCam K_s band

Figure 12. DE optimization of the best fit model for K_s band KLIP-RDI reduction of AB Aur. The off diagonal plots show solutions for two parameters binned 6×6 . The color of each bin represents the χ_r^2 of the best-fitting model within that bin. Lighter colors indicate smaller values of χ_r^2 . The diagonal plots show a one dimensional view of each parameter of the minimum χ_r^2 as a function of the parameter range, and colored by χ_r^2 . The best fit solutions are shown by the red square/line, while the solutions within the “acceptable range” ($\chi_r^2 \leq \chi_{r,min}^2 + \sqrt{2/\nu}$) are those whose χ_r^2 fall below the dashed black line in the diagonal plots.


LMIRCam 3.08 μm band

Figure 13. As Figure 12, but for 3.08 μm Ice band DE optimization.

LMIRCam L' band




Figure 14. As Figure 12, but for L' band DE optimization.




Figure 15. Best fit model and residuals from the DE optimization for K_s . **top left)** Best fit model image **top center)** Best fit convolved and forward modeled image **top right)** Residuals between the model convolved + FM model and the KLIP reduced data for just the area of interest (i.e. the disk of AB Aur). **bottom left)** Best fit model image after convolved with an observed LMIRCam PSF (shown in the offset on the bottom left of the image) **bottom center)** Observed K_s band image of AB Aur after pyKLIP RDI reduction. **bottom right)** χ^2 residuals between the model convolved + FM model and the KLIP reduced data for the area of interest.




Figure 16. As Figure 15, but for $3.08 \mu\text{m}$ Ice band best fit model and residuals.




Figure 17. As Figure 15, but for L' band best fit model and residuals.




Figure 18. Azimuthally averaged throughput radial profiles for varying amplitude and inclination. All other parameters are the fixed and best fit solutions in Tables 4 and 5. The black curve is the best fit profile.