

ON EXPLICIT BIRATIONAL GEOMETRY FOR MINIMAL n -FOLDS OF CANONICAL DIMENSION $n - 1$

MENG CHEN, LOUIS ESSER, AND CHENGXI WANG

ABSTRACT. Let $n \geq 2$ be any integer. We study the optimal lower bound $v_{n,n-i}$ of the canonical volume and the optimal upper bound $r_{n,n-i}$ of the canonical stability index for minimal projective n -folds of general type, which are canonically fibered by i -folds ($i = 0, 1$). The results for $i = 0$, $v_{n,n} = 2$ and $r_{n,n} = n + 2$, are known to experts. In this article, we show that $v_{n,n-1} = \frac{6}{2n+(n \bmod 3)}$ and $r_{n,n-1} = \frac{1}{3}(5n + 3 + (n \bmod 3))$. The machinery is applicable to all canonical dimensions $n - i$.

1. INTRODUCTION

In birational geometry, minimal varieties of general type form one of the basic building blocks of the minimal model program. A crucial step toward classifying these varieties is to study the distribution of their birational invariants. Since every smooth variety of general type has a minimal model, this will give the distribution of invariants in the smooth case as well. Throughout, we will work over any algebraically closed field of characteristic zero.

We begin by recalling a few examples of birational invariants. Let $n \geq 2$ be any integer and X be a minimal projective n -fold of general type, so that X has at worst \mathbb{Q} -factorial terminal singularities and the canonical divisor K_X is nef. The m th *plurigenus* $P_m(X)$ of X is the dimension of the space of sections $H^0(X, mK_X)$ for $m \geq 1$. The *volume* of X is a measure of the asymptotic growth of the plurigenera, defined as

$$\mathrm{Vol}(X) := \limsup_{m \rightarrow \infty} \frac{n!P_m(X)}{m^n},$$

which is a positive rational number. Since X is minimal, so that K_X is nef, the Riemann-Roch formula shows that $\mathrm{Vol}(X) = K_X^n$, the top intersection number of K_X . We define the *canonical stability index* of X by

$$r_s(X) := \min\{p \in \mathbb{Z}_{>0} \mid \varphi_{m,X} \text{ is birational for all } m \geq p\},$$

where $\varphi_{m,X}$ denotes the m -canonical map of X . For any dimension n , the n th *canonical stability index* is defined as:

$$r_n := \max\{r_s(X) \mid X \text{ is any minimal projective } n\text{-fold of general type}\}.$$

Date: January 25, 2022.

M. Chen was supported by NSFC for Innovative Research Groups (#12121001), National Key Research and Development Program of China (#2020YFA0713200) and NSFC (#12071078, #11731004). L. Esser was partially supported by NSF grant DMS-2054553.

Similarly, we define

$$v_n := \inf\{\text{Vol}(X) \mid X \text{ is any minimal projective } n\text{-fold of general type}\}.$$

A remarkable theorem of Hacon-McKernan [16], Takayama [21] and Tsuji [23] states that the canonical stability index is uniformly bounded in each dimension, so that $r_n < +\infty$. As a consequence, we have $v_n > 0$ (more precisely, $v_n \geq \frac{1}{(r_n)^n}$). Once boundedness is established, the natural next step in classification is to find explicit values for these bounds (see, for instance, Hacon-McKernan [16, Problem 1.5, Question 1.6]).

We list some known results concerning r_n and v_n :

- By Bombieri [3], $r_2 = 5$ and $v_2 = 1$.
- By Iano-Fletcher [18], $r_3 \geq 27$ and $v_3 \leq \frac{1}{420}$; by Chen-Chen [4–6] and Chen [12], $r_3 \leq 57$ and $v_3 \geq \frac{1}{1680}$.
- By Esser-Totaro-Wang [22], for $n \geq 3$, $r_n > 2^{2^{(n-2)/2}}$ and $v_n < \frac{1}{2^{2^{n/2}}}$.

For a given minimal variety X of general type, the *canonical dimension* of X is defined as $d_1 := \dim \varphi_1(X)$. For $1 \leq i < n$, we define $r_{n,n-i}$ and $v_{n,n-i}$ to be the maximal canonical stability index and the infimum of volumes, respectively, among all minimal n -folds of general type with canonical dimension $n-i$. Similarly, $r_{n,0}$ and $v_{n,0}$ (resp. $r_{n,-\infty}$ and $v_{n,-\infty}$) are the corresponding values for minimal n -folds with geometric genus $p_g = 1$ (resp. $p_g = 0$). With these definitions,

$$\begin{aligned} r_n &= \max\{r_{n,j} \mid -\infty \leq j \leq n\} \text{ and} \\ v_n &= \min\{v_{n,j} \mid -\infty \leq j \leq n\}. \end{aligned}$$

When X has canonical dimension $n-i$ for small i , we can analyze it using its canonical fibration by i -folds. In particular, the values $r_{n,n-1}$ have long been studied with applications to the classification of algebraic varieties, beginning with surfaces. In this paper, we study values of $r_{n,j}$ and $v_{n,j}$ for $j \geq n-1$ and prove the following theorem:

Theorem 1.1. *Let $n \geq 2$ be any integer. The following statements hold:*

- (1) $v_{n,n} = 2$ and $r_{n,n} = n+2$;
- (2) $v_{n,n-1} = \frac{6}{2n+(n \bmod 3)}$ and $r_{n,n-1} = \frac{1}{3}(5n+3+(n \bmod 3))$;

where “ $n \bmod 3$ ” is the minimal non-negative residue of n modulo 3.

In each dimension, we will present optimal examples achieving the bounds in (1) and (2), which will be weighted projective hypersurfaces of general type. For an introduction to weighted projective hypersurfaces, see [22, Section 2].

Remark 1.2. Some special values of $v_{n,j}$ and $r_{n,j}$ are already known:

- The surface case is due to Bombieri [3];
- The threefold case is proved by one of the authors (Chen) [9, 10];
- The value of $v_{n,n}$ is originally due to Kobayashi [19];
- The values of $v_{4,3}$, $v_{4,2}$, $v_{5,4}$ and $v_{5,3}$ are already obtained in Chen-Jiang-Li [13].

Throughout, we make use of the following notation:

- ^o For two \mathbb{Q} -divisors D_1 and D_2 , $D_1 \sim_{\mathbb{Q}} D_2$ (resp., $D_1 \equiv D_2$) means that D_1 is \mathbb{Q} -linearly equivalent (resp., numerically equivalent) to D_2 . Similarly, $D_1 \geq_{\text{num}} D_2$ (resp., $D_1 \geq_{\mathbb{Q}} D_2$) means that $D_1 - D_2$ is numerically (resp., \mathbb{Q} -linearly) equivalent to an effective \mathbb{Q} -divisor.
- ^o For two linear systems $|A|$ and $|B|$ on a variety, we write $|A| \succsim |B|$ (or symmetrically, $|B| \precsim |A|$) if there exists an effective divisor F such that

$$|A| \supseteq |B| + F.$$

2. A KEY TECHNICAL THEOREM

Theorem 2.1. *Let X be a minimal projective n -fold of general type. Let X' be a nonsingular projective variety and $\pi : X' \rightarrow X$ be a birational morphism. Assume that there exists a chain of smooth subvarieties of general type*

$$Z_1 \subset Z_2 \subset \cdots \subset Z_{n-1} \subset Z_n = X'$$

with $\dim Z_j = j$ for $j = 1, 2, \dots, n-1$. Suppose that the following conditions hold:

- (i) $\pi^*(K_X)|_{Z_i}$ is big for each $i = 2, \dots, n-1$;
- (ii) $\pi^*(K_X)|_{Z_i} \equiv \beta_i Z_{i-1} + \Delta_{i-1}$ where β_i is a positive rational number, Δ_{i-1} is an effective \mathbb{Q} -divisor on Z_i and

$$Z_{i-1} \notin \text{Supp}(\Delta_{i-1} + \sum_{j \geq i} \Delta_j|_{Z_i})$$

for each $i = 2, \dots, n$;

- (iii) the number $\xi := (\pi^*(K_X) \cdot Z_1) > 0$.

Then each of the following statements holds:

- (1) *For any integer m with $\alpha_m := (m-1 - \sum_{i=2}^n \frac{1}{\beta_i})\xi > 1$, the inequality*

$$m\xi \geq (2g(Z_1) - 2) + \lceil \alpha_m \rceil \quad (2.1)$$

holds. In particular, under this situation, one has

$$\xi \geq \frac{2g(Z_1) - 2}{1 + \sum_{i=2}^n \frac{1}{\beta_i}}. \quad (2.2)$$

- (2) *The canonical volume of X has the lower bound:*

$$K_X^n \geq \beta_2 \beta_3 \cdots \beta_n \xi. \quad (2.3)$$

Proof. First of all, if we replace X' by any nonsingular birational model (which dominates X') for which no Z_i is contained in the image of the exceptional locus, then we may replace each Z_i by its strict transform. After performing this replacement, the assumption still holds for the same values of β_i and the number ξ remains unchanged by the projection formula. Hence, fixing an effective divisor $K_1 \sim K_X$ at the beginning, we may and do assume that, on X' , the union of the divisor $\pi^*(K_1)|_{Z_i}$, Δ_{i-1} (for all $i = 2, \dots, n$) and exceptional divisors has simple normal crossing supports.

Suppose $m > 1 + \sum_{i=2}^n \frac{1}{\beta_i}$. Since the \mathbb{Q} -divisor

$$(m-1)\pi^*(K_X)|_{Z_n} - \frac{1}{\beta_n}\Delta_{n-1} - Z_{n-1} \equiv \left(m-1 - \frac{1}{\beta_n}\right)\pi^*(K_X)|_{Z_n}$$

is nef and big, and the fractional part has snc supports, the Kawamata-Viehweg vanishing theorem [20, 24] implies:

$$\begin{aligned} |mK_{X'}|_{Z_{n-1}} &\succcurlyeq |K_{X'} + \lceil (m-1)\pi^*(K_X) - \frac{1}{\beta_n}\Delta_{n-1} \rceil|_{Z_{n-1}} \\ &\succcurlyeq |K_{Z_{n-1}} + \lceil ((m-1)\pi^*(K_X) - Z_{n-1} - \frac{1}{\beta_n}\Delta_{n-1})|_{Z_{n-1}} \rceil|. \end{aligned} \quad (2.4)$$

By induction, one clearly gets, for $i = 2, \dots, n-1$, that

$$\begin{aligned} |mK_{X'}|_{Z_{i-1}} &\succcurlyeq |K_{Z_i} + \lceil (m-1)\pi^*(K_X)|_{Z_i} - \sum_{l=i+1}^n (Z_{l-1} + \frac{1}{\beta_l}\Delta_{l-1})|_{Z_i} - \frac{1}{\beta_i}\Delta_{i-1} \rceil|_{Z_{i-1}} \\ &\succcurlyeq |K_{Z_{i-1}} + \lceil (m-1)\pi^*(K_X)|_{Z_{i-1}} - \sum_{l=i}^n (Z_{l-1} + \frac{1}{\beta_l}\Delta_{l-1})|_{Z_{i-1}} \rceil|. \end{aligned} \quad (2.5)$$

By (2.4) and (2.5) while repeatedly using [8, Lemma 2.7], we have

$$M_m|_{Z_1} \geq \text{Mov}|K_{Z_1} + \lceil Q_m \rceil|, \quad (2.6)$$

where $Q_m := (m-1)\pi^*(K_X)|_{Z_1} - \sum_{i=2}^n (Z_{i-1} + \frac{1}{\beta_i}\Delta_{i-1})|_{Z_1}$ and $M_m := \text{Mov}|mK_{X'}|$. Note that, by Condition (ii), the divisor $Z_{i-1}|_{Z_{i-1}}$ never appears in fractional part of all above \mathbb{Q} -divisors.

Whenever $\alpha_m = \deg(Q_m) > 1$, $Q_m \equiv (m-1 - \sum_{i=2}^n \frac{1}{\beta_i})\pi^*(K_X)|_{Z_1}$ is nef and big, and $\text{Mov}|K_{Z_1} + \lceil Q_m \rceil| = K_{Z_1} + \lceil Q_m \rceil$, because divisors of degree at least $2g(Z_1)$ are base point free. Since $m\pi^*(K_X) \geq M_m$ and $\deg(\lceil Q_m \rceil) \geq \deg(Q_m)$, taking degrees on both sides of (2.6) proves (2.1) in (1).

Taking a sufficiently large integer m' so that $\alpha_{m'} > 1$, (2.2) is a consequence of (2.1). Statement (2) then directly follows from the assumptions and the fact that $\pi^*(K_X)$ is nef. \square

Definition 2.2. [11, Definition 2.3] A *generic irreducible element* S of a movable linear system $|N|$ on a variety Z is a generic irreducible component in a general member of $|N|$. Thus, it is a general member of $|N|$ whenever $\dim \varphi_{|N|}(V) \geq 2$. If $|N|$ is composed with a pencil, i.e., $\dim \varphi_{|N|}(V) = 1$, one has $N \equiv tS$ for some integer $t \geq 1$.

Definition 2.3. [6, Definition 2.6] Let $|N|$ be a movable linear system on a variety Z . Pick two different generic irreducible elements S', S'' in $|N|$. We say that a linear system $|M|$ (resp. a rational map φ corresponding to a linear system) *distinguishes* S' and S'' if $\varphi_{|M|}(S') \neq \varphi_{|M|}(S'')$ (resp. if $\varphi(S') \neq \varphi(S'')$).

In proving birationality of the rational map φ_Λ , where $\Lambda \subset |L|$ and L a divisor on a projective variety Z , we tacitly use the following rule:

Let $|M|$ be a base point free linear system on Z and denote by S a generic irreducible element of $|M|$. If φ_Λ distinguishes different generic irreducible elements of $|M|$ and $\varphi_\Lambda|_S$ is birational, then φ_Λ is birational. Conversely, if φ_Λ is birational, it is clear that, for any base point free linear system $|M|$, φ_Λ distinguishes different generic irreducible elements of $|M|$ and that $\varphi_\Lambda|_S$ is birational.

We'll also make use of the following [5, Section 2.7]:

Proposition 2.4 (Birationality principle). *Let D and M be two divisors on a smooth projective variety Z . Assume that $|M|$ is base point free. Take the Stein factorization of $\varphi_{|M|} : Z \xrightarrow{f} W \rightarrow \mathbb{P}^{h^0(Z,M)-1}$, where f is a fibration onto a normal variety W . The rational map $\varphi_{|D|}$ is birational onto its image if one of the following conditions is satisfied:*

- $\dim \varphi_{|M|}(Z) \geq 2$, $|D - M| \neq \emptyset$ and $\varphi_{|D|}|_S$ is birational for a general member S of $|M|$
- $\dim \varphi_{|M|}(Z) = 1$, $\varphi_{|D|}$ distinguishes general fibers of f and $\varphi_{|D|}|_S$ is birational for a general fiber S of f .

In practice, in order to prove the birationality of $\varphi_{|D|}$, it is sufficient to find two sublinear systems $\Lambda_i \subset |D|$ ($i = 1, 2$) such that φ_{Λ_1} distinguishes different generic irreducible elements of $|M|$ while $\varphi_{\Lambda_2}|_S$ is birational for a generic irreducible element S of $|M|$.

Corollary 2.5. *Keep the same assumptions and notation as in Theorem 2.1. Let m be a positive integer. Assume that the following conditions are satisfied:*

For each $i = 2, \dots, n$,

- (1) *there exists a base point free linear system $|N_i|$ on Z_i such that Z_{i-1} is the generic irreducible element of $|N_i|$;*
- (2) *the linear system $|mK_{X'}|_{Z_i}$ distinguishes different generic irreducible elements of $|N_i|$;*
- (3) $\alpha_m > 2$.

Then $\varphi_{m,X}$ is birational onto its image.

Proof. We refer to the proof of Theorem 2.1. Applying the birationality principle repeatedly, by Assumptions (1) - (2), Relation (2.4) and Relation (2.5), $\varphi_{m,X}$ is birational if and only if so is $\varphi_{m,X'}|_{Z_{n-1}}$, by inductions, \dots , if and only if so is $\varphi_{m,X'}|_{Z_2}$, if and only if so is $\varphi_{m,X'}|_{Z_1}$. Now we know that $\varphi_{|K_{Z_1} + [Q_m]|}$ is birational as $\alpha_m > 2$. From the proof of Theorem 2.1, we have seen that $|K_{Z_1} + [Q_m]| \preccurlyeq |mK_{X'}|_{Z_1}$. Hence $\varphi_{m,X'}|_{Z_1}$ is birational. We are done. \square

3. THE CANONICAL MAP OF VARIETIES OF GENERAL TYPE

3.1. Set up for $\varphi_{1,X}$.

Let X be a minimal projective n -fold ($n \geq 2$) of general type with $p_g(X) \geq 2$ and canonical dimension d_1 . Clearly we have $1 \leq d_1 \leq n$. Let $\pi : X' \rightarrow X$ be a succession of blow-ups along nonsingular centers such that $|M_1|$, the linear system of $\text{Mov}|K_{X'}|$, is base point free and that the union of the fixed part of $|K_{X'}|$ and exceptional divisors of π has simple normal crossing supports. Take $g_1 := \varphi_{1,X} \circ \pi$, which is a projective morphism. Denote by $f : X' \rightarrow \Gamma$ be the induced fibration after taking the Stein factorization of g_1 . So $\dim \Gamma = d_1$.

Set $|N_n| := |M_1|$ and $Z_n := X'$. For any integer k with $n - d_1 + 2 \leq k \leq n$, inductively, take Z_{k-1} to be the generic irreducible element of $|N_k|$ and set $|N_{k-1}| := |M_1|_{Z_{k-1}}$. By Bertini's theorem, $|N_{k-1}|$ is base point free and is not composed with a pencil for each k . We have defined the following chain

of nonsingular subvarieties:

$$Z_{n-d_1+1} \subset \cdots \subset Z_{n-1} \subset Z_n = X'.$$

Let us review some established inequalities in Chen-Jiang-Li [13, Step 0, Proof of Theorem 5.1]. Remodify π (if necessary) such that, for $1 \leq j \leq d_1 - 1$, Z_{n-j} dominates a minimal model $Z_{n-j,0}$ and, in particular, there is a birational morphism $\pi_{Z_{n-j,0}} : Z_{n-j} \rightarrow Z_{n-j,0}$. By induction, for $1 \leq j \leq d_1 - 1$, one has

$$h^0(X', M_1|_{Z_{n-j}}) \geq p_g(X') - j \quad (3.1)$$

Similar to Chen-Jiang-Li [13, Inequality (5.4)], we have

$$\pi^*(K_X)|_{Z_{n-j}} \geq \frac{1}{j+1} \pi_{Z_{n-j}}^*(K_{Z_{n-j,0}}) \quad (3.2)$$

for $1 \leq j \leq d_1 - 1$. In particular, we have

$$\pi^*(K_X)|_{Z_{n-d_1+1}} \geq \frac{1}{d_1} \pi_{Z_{n-d_1+1}}^*(K_{Z_{n-d_1+1,0}}). \quad (3.3)$$

3.2. Proof of Theorem 1.1: the case $d_1 = n$.

Theorem 2.1 and Corollary 2.5 naturally apply to this very special case. We have already defined the chain of subvarieties:

$$Z_1 \subset Z_2 \subset \cdots \subset Z_{n-1} \subset Z_n = X'.$$

By construction, we have $\beta_i \geq 1$ for each i with $2 \leq i \leq n$. Since $|M_1|_{Z_2}|$ is base point free and is not composed with a pencil, we have

$$\xi \geq (M_1 \cdot Z_1)_{X'} \geq (Z_1|_{Z_2})^2 \geq 2.$$

Hence, by (2.3), $v_{n,n} \geq 2$.

Take any integer $m > n + 1$. Since $p_g(X) > 0$, the proof of Theorem 2.1 simply implies $|mK_{X'}|_{Z_i} \not\supseteq |N_i|$ for $2 \leq i \leq n$. Because each linear system $|N_i|$ is not composed with a pencil, the linear system $|mK_{X'}|_{Z_i}$ naturally distinguishes generic irreducible elements of $|N_i|$. Conditions Corollary 2.5(1) and (2) are therefore automatically satisfied. Since $\alpha_m > \xi \geq 2$, we have proved that $r_s(X) \leq n + 2$ by Corollary 2.5. Hence $r_{n,n} \leq n + 2$.

Example 3.1. For any integer $n \geq 2$, the general hypersurface

$$X = V_{2(n+2)} \subseteq \mathbb{P}(1^{(n+1)}, n+2)$$

(of degree $2(n+2)$) is a smooth canonical n -fold with $\omega_X \cong \mathcal{O}_X(1)$, $\text{Vol}(X) = 2$ and $r_s(X) = n + 2$. Clearly, X has canonical dimension $d_1 = n$.

So we conclude that Theorem 2.1 (1) is true, i.e. $v_{n,n} = 2$ and $r_{n,n} = n + 2$. Conversely, if X has canonical dimension n and $\text{Vol}(X) = v_{n,n} = 2$, X is a double cover of \mathbb{P}^n [19, Proposition 2.5].

3.3. Further setting for the case $d_1 < n$.

We take Z_{n-d_1} to be the generic irreducible element of $|N_{n-d_1+1}| := |M_1|_{Z_{n-d_1+1}}|$. Since $|N_{n-d_1+1}|$ is composed with a pencil and by (3.1), we have

$$N_{n-d_1+1} \equiv \beta_{n-d_1+1} Z_{n-d_1}$$

where $\beta_{n-d_1+1} \geq h^0(M_1|_{Z_{n-d_1+1}}) - 1 \geq p_g(X) - d_1$.

In addition, by our definitions, we have $\beta_i \geq 1$ for each i with $n - d_1 + 2 \leq i \leq n$.

4. PROOF OF THEOREM 1.1: THE CASE $d_1 = n - 1$

We have defined the chain $\{Z_i | i = 1, \dots, n\}$ according to Subsections 3.1 and 3.3. Note that $|N_2|$ is composed with a pencil of curves on the smooth surface Z_2 . By our definition, $C := Z_1$ is the generic irreducible element of $|N_2|$. Furthermore, we have $\beta_i \geq 1$ for $3 \leq i \leq n$ and $\beta_2 \geq p_g(X) - n + 1 \geq 1$. Since Z_1 moves in an algebraic family and $\pi^*(K_X)|_{Z_2}$ is nef and big, we have $\xi > 0$ by the Hodge index theorem.

4.1. Volume estimation.

Noting that $g(Z_1) \geq 2$, by (2.2), we have $\xi \geq \frac{2}{n}$. We optimize the estimation by distinguishing three exclusive cases:

Case 4.1.1. $n = 3k + 2$, $k \geq 0$.

Take $m_1 = \lfloor \frac{9}{2}k + 4 \rfloor$. Then, since $\alpha_{m_1} > 1$, we have

$$\xi \geq \frac{8}{9k + 8} = \frac{2 \cdot 4}{(2 \cdot 4 + 1)k + 2 \cdot 4}$$

by (2.1). We will repeatedly use (2.1) to optimize the estimation of ξ . In fact, suppose that we already know

$$\xi \geq \frac{2 \cdot 4^l}{(2 \cdot 4^l + 1)k + 2 \cdot 4^l}$$

for an integer $l > 0$. Take

$$m_l = \lfloor \frac{(2 \cdot 4^l + 1)k}{2 \cdot 4^l} \rfloor + 3k + 4.$$

Then we see $\alpha_{m_l} > 1$ and (2.1) implies

$$\xi \geq \frac{2 \cdot 4^{l+1}}{(2 \cdot 4^{l+1} + 1)k + 2 \cdot 4^{l+1}}.$$

Taking the limit, as $l \mapsto +\infty$, we have

$$\xi \geq \frac{1}{k + 1} = \frac{3}{n + 1}. \quad (4.1)$$

Case 4.1.2. $n = 3k + 1$, $k \geq 1$.

Take $m_1 = \lfloor \frac{9k+1}{2} + 2 \rfloor$. Then, since $\alpha_{m_1} > 1$, we have

$$\xi \geq \frac{8}{9k + 5} = \frac{2 \cdot 4}{(2 \cdot 4 + 1)k + 4 + 1}$$

by (2.1). Assume that we have shown

$$\xi \geq \frac{2 \cdot 4^l}{(2 \cdot 4^l + 1)k + (4^l + 4^{l-1} + \dots + 1)}$$

for some integer $l > 0$. Take $m_l = \lfloor \frac{(2 \cdot 4^l + 1)k + (4^l + 4^{l-1} + \dots + 1)}{2 \cdot 4^l} \rfloor + 3k + 2$. Then, since $\alpha_{m_l} > 1$, we get

$$\xi \geq \frac{2 \cdot 4^{l+1}}{(2 \cdot 4^{l+1} + 1)k + (4^{l+1} + 4^l + \dots + 1)}.$$

Taking the limit, as $l \mapsto +\infty$, we have $\xi \geq \frac{1}{k+2/3}$. Finally, taking $m = 4k+2$, then we get

$$\xi \geq \frac{2}{2k+1} = \frac{6}{2n+1}. \quad (4.2)$$

Case 4.1.3. $n = 3k$, $k \geq 1$.

Take $m_1 = \lfloor \frac{9k}{2} + 1 \rfloor$. Then, since $\alpha_{m_1} > 1$, we get

$$\xi \geq \frac{8}{9k+2} > \frac{2 \cdot 4}{(2 \cdot 4 + 1)k + 4}$$

by (2.1). Suppose that, for some $l > 0$, we have already shown

$$\xi \geq \frac{2 \cdot 4^l}{(2 \cdot 4^l + 1)k + (4^l + \dots + 4)}.$$

Take

$$m_l = \lfloor \frac{(2 \cdot 4^l + 1)k + (4^l + \dots + 4)}{2 \cdot 4^l} \rfloor + 3k + 2.$$

Then, since $\alpha_{m_l} > 1$, we have

$$\xi \geq \frac{2 \cdot 4^{l+1}}{(2 \cdot 4^{l+1} + 1)k + (4^{l+1} + \dots + 4)}$$

by (2.1). Taking the limit, as $l \mapsto +\infty$, we have $\xi \geq \frac{1}{k+2/3}$. Take $m = 4k+1$.

Then $\xi \geq \frac{4}{4k+1} = \frac{1}{k+1/4}$.

We continue the optimization for the lower bound of ξ . Assume that, for some integer $t \geq 4$, we already know $\xi \geq \frac{1}{k+1/t}$. Take $m_t = (t+1)k+1$. Then, since $\alpha_{m_t} > t-2 \geq 2$, we have

$$\xi \geq \frac{t+1}{(t+1)k+1} = \frac{1}{k+1/(t+1)}$$

by (2.1). Taking the limit, as $t \mapsto +\infty$, then we have

$$\xi \geq \frac{1}{k} = \frac{3}{n}. \quad (4.3)$$

By (4.1), (4.2) and (4.3), we have proved the following:

Corollary 4.1. *For any integer $n \geq 2$, $v_{n,n-1} \geq \frac{6}{2n+(n \bmod 3)}$.*

4.2. The canonical stability index. As we have seen, for $3 \leq i \leq n$, $|N_i|$ is not composed with a pencil. Applying the Kawamata-Viehweg vanishing theorem in the similar way to that of (2.4) and (2.5), for any $m > n-2$ and any j with $3 \leq j \leq n-1$, we have

$$\begin{aligned} |mK_{X'}||_{Z_j} &\succcurlyeq |K_{X'} + \lceil (m-n+2)\pi^*(K_X) \rceil + (n-3)M_1||_{Z_j} \\ &\succcurlyeq |M_1|_{Z_j}. \end{aligned} \quad (4.4)$$

Hence $|mK_{X'}||_{Z_j}$ distinguishes different generic irreducible elements of $|N_j|$ whenever $m > n-2$ and $3 \leq j \leq n$.

We study the situation on $|N_2|$, which is composed with a pencil of curves. If $|N_2|$ is composed with a rational pencil of curves, as long as $m > n-1$, $|mK_{X'}||_{Z_2}$ distinguishes different generic irreducible elements of $|N_2|$ since $|mK_{X'}||_{Z_2} \succcurlyeq |N_2|$. If $|N_2|$ is composed with a non-rational pencil

of curves, picking two different generic irreducible elements C_1, C_2 of $|N_2|$, then $(M_1|_{Z_2} - C_1 - C_2)$ is nef and

$$(M_1|_{Z_2} - C_i)|_{C_i} \sim 0$$

for $i = 1, 2$. Hence, by the vanishing theorem and for any $m > n$, one has the relation:

$$\begin{aligned} |mK_{X'}|_{Z_2} &\supseteq |K_{X'} + [(m - n)\pi^*(K_X)] + (n - 1)M_1| \\ &\supseteq |K_{Z_2} + [(m - n)\pi^*(K_X)|_{Z_2}] + M_1|_{Z_2}| \end{aligned}$$

and the surjective map:

$$\begin{aligned} &H^0(Z_2, K_{Z_2} + [(m - n)\pi^*(K_X)|_{Z_2}] + M_1|_{Z_2}) \\ \longrightarrow &H^0(C_1, K_{C_1} + D_1) \oplus H^0(C_2, K_{C_2} + D_2) \end{aligned}$$

where

$$D_i := ([(m - n)\pi^*(K_X)|_{Z_2}] + M_1|_{Z_2} - C_i)|_{C_i} = [(m - n)\pi^*(K_X)|_{Z_2}]|_{C_i}$$

for $i = 1, 2$. Note that $H^0(C_i, K_{C_i} + D_i) \neq 0$ since $\deg(D_i) > 0$. Therefore $|mK_{X'}|$ can distinguish different generic irreducible elements of $|N_2|$ in this case. In summary, for any $m > n$, Conditions (1) and (2) of Corollary 2.5 are satisfied.

We have proved that $\xi \geq \frac{6}{2n+(n \bmod 3)}$. For any $m \geq \frac{5n+3+(n \bmod 3)}{3}$, we have $\alpha_m > 2$. By Corollary 2.5, whenever $m \geq \frac{5n+3+(n \bmod 3)}{3}$, $\varphi_{m,X}$ is birational onto its image. We have proved the following:

Corollary 4.2. *For any integer $n \geq 2$, $r_{n,n-1} \leq \frac{5n+3+(n \bmod 3)}{3}$.*

4.3. Examples.

Example 4.3. (cf. [13, Example 6.6(1)]) Let $n = 3k + 2$. The general hypersurface $X = V_{10k+10} \subset \mathbb{P}(1^{(3k+2)}, 2(k+1), 5(k+1))$ of degree $10(k+1)$ has at worst canonical singularities and is a minimal n -fold with $\omega_X = \mathcal{O}_X(1)$, canonical dimension $n - 1$, and volume $\text{Vol}(X) = \frac{1}{k+1} = \frac{3}{n+1} = \frac{6}{2n+(n \bmod 3)}$. Furthermore, it is clear that $r_s(X) = 5k + 5 = \frac{5n+3+(n \bmod 3)}{3}$.

Example 4.4. (cf. [13, Example 6.6(2)]) Let $n = 3k + 1$. The general hypersurface $X = V_{10k+6} \subset \mathbb{P}(1^{(3k+1)}, 2k+1, 5k+3)$ of degree $10k+6$ has at worst canonical singularities and is a minimal n -fold with $\omega_X = \mathcal{O}_X(1)$, canonical dimension $n - 1$, and volume $\text{Vol}(X) = \frac{6}{2n+1} = \frac{6}{2n+(n \bmod 3)}$. Furthermore, it is clear that $r_s(X) = 5k + 3 = \frac{5n+3+(n \bmod 3)}{3}$.

Example 4.5. Let $n = 3k$. The general hypersurface $X = V_{20k+2} \subset \mathbb{P}(1, 2^{(3k-1)}, 4k, 10k+1)$ of degree $20k+2$ has at worst canonical singularities and is a minimal n -fold with $\omega_X = \mathcal{O}_X(2)$, canonical dimension $n - 1$, and volume $\text{Vol}(X) = \frac{3}{n} = \frac{6n}{2n+(n \bmod 3)}$. Furthermore, it is clear that $r_s(X) = 5k + 1 = \frac{5n+3+(n \bmod 3)}{3}$.

Theorem 1.1(2) follows directly from Corollary 4.1, Corollary 4.2, Example 4.3, Example 4.4 and Example 4.5.

5. GEOMETRY OF OPTIMAL EXAMPLES

In this section, we summarize an alternate approach proving the volume bound in Theorem 1.1(2). This method also shows that any X realizing the minimal value of $v_{n,n-1}$ must strongly resemble one of the weighted projective hypersurfaces in Section 4.3. We omit some of the proof details and continue to use the notation of Section 3.1.

Claim 5.1. *Let X be a minimal variety of general type with dimension $n \geq 3$ and canonical dimension $n - 1$ such that*

$$\mathrm{Vol}(X) = \frac{6}{2n + (n \bmod 3)}. \quad (5.1)$$

Then the canonical map of X is a dominant rational map to \mathbb{P}^{n-1} with general fiber a genus 2 curve. Furthermore, the general surface Z_2 has minimal volume in the sense of Proposition 5.2 below. Its minimal model $Z_{2,0}$ contains a tree of rational curves contracted by the morphism from X' to its canonical model, described in Cases 5.1, 5.2, and 5.3.

Proof. Let X be a minimal variety realizing (5.1) as above. We immediately see that $g(Z_1) = 2$, or else $\xi \geq \frac{4}{n}$ by (2.2). Further, $p_g(X) = n$, or else $\beta_2 \geq 2$, so that $K_X^n \geq 2 \cdot \frac{2}{n} = \frac{4}{n}$ by (2.3). The image $\overline{\varphi_1(X)}$ of the canonical map is then a subvariety of \mathbb{P}^{n-1} of dimension $n - 1$, so $\overline{\varphi_1(X)} = \mathbb{P}^{n-1}$. Therefore, the canonical map is a dominant rational map to \mathbb{P}^{n-1} whose general fiber is a smooth genus 2 curve.

We can also say a great deal about Z_2 and the properties of the birational contraction $\pi : X' \rightarrow X_{\mathrm{can}}$, restricted to Z_2 . Here $X_{\mathrm{can}} = \mathrm{Proj} \bigoplus_{m=1}^{\infty} H^0(X, mK_X)$ is the canonical model of the variety X (equivalently, of X') and $K_{X_{\mathrm{can}}}^n = K_X^n$.

The key idea is to find a commutative diagram

$$\begin{array}{ccccc} X' & \xrightarrow{\pi_{\mathrm{can}}} & X_{\mathrm{can}} & & \\ \uparrow & & \uparrow \psi & & \\ Z_2 & \xrightarrow{g} & Z_{2,0} & \xrightarrow{f} & S, \end{array}$$

where $g : Z_2 \rightarrow Z_{2,0}$ is the minimal model, S is a projective normal surface with K_S \mathbb{Q} -Cartier, and $f : Z_{2,0} \rightarrow S$ is a birational morphism whose exceptional divisors have nonpositive discrepancies.

By inequality (3.3), $(n-1)\pi|_{Z_2}^*(K_X) \geq g^*(K_{Z_{2,0}})$. Taking the pushforward to S gives $(n-1)\psi^*(K_{X_{\mathrm{can}}}) \geq K_S$ by the commutativity of the diagram and because $K_{Z_{2,0}}$ and $f^*(K_S)$ differ by exceptional divisors contracted by f . Both $\psi^*(K_{X_{\mathrm{can}}})$ and K_S are nef divisors, the latter by the assumption on discrepancies. Therefore,

$$((n-1)\psi^*(K_{X_{\mathrm{can}}}))^2 \geq (n-1)\psi^*(K_{X_{\mathrm{can}}}) \cdot K_S \geq K_S^2. \quad (5.2)$$

Combining the inequality (5.2) with $K_X^n \geq (\pi_{\mathrm{can}}^*(K_{X_{\mathrm{can}}})|_{Z_2})^2$, we get

$$(n-1)^2 K_X^n \geq ((n-1)\pi_{\mathrm{can}}^*(K_{X_{\mathrm{can}}})|_{Z_2})^2 = ((n-1)\psi^*(K_{X_{\mathrm{can}}}))^2 \geq K_S^2.$$

Moreover, the intersection number K_S^2 is determined by the volume of the minimal surface $Z_{2,0}$ and the divisors are contracted by f . We may obtain sharper bounds on K_X^n via this method than from only considering smooth models of Z_2 because often, $K_S^2 > \text{Vol}(Z_2)$. Therefore, we seek to factor $\pi_{\text{can}}|_{Z_2} : Z_2 \rightarrow X_{\text{can}}$ through contractions to certain singular surfaces S .

The minimal model $Z_{2,0}$ of Z_2 is a fibration of genus 2 curves over \mathbb{P}^1 . The general fiber is a genus 2 curve Z_1 . We know by the adjunction formula that Z_2 satisfies $K_{Z_2} \geq (n-1)Z_1$. Supposing $n \geq 3$, $p_g(Z_2) \geq h^0(Z_2, (n-1)Z_1) \geq n \geq 3$. Under these assumptions, we have the following properties (cf. [7, Proposition 2.9]):

Proposition 5.2.

$$\text{Vol}(Z_2) = (K_{Z_{2,0}})^2 \geq \begin{cases} \frac{8n-16}{3}, & n \equiv 2 \pmod{3} \\ \frac{8n-14}{3}, & n \equiv 1 \pmod{3} \\ \frac{8n-12}{3}, & n \equiv 0 \pmod{3} . \end{cases} \quad (5.3)$$

The fibration $Z_2 \rightarrow \mathbb{P}^1$ factors through $Z_{2,0}$ and $K_{Z_{2,0}} = (n-1)Z_1 + G$, where G is an effective divisor.

If we have equality in (5.3), then $G = V + 2E$, where V is a vertical divisor, $E \cong \mathbb{P}^1$ is a section of the fibration, and $E^2 = -\frac{1}{3}(n+1+V \cdot E)$. The divisor V satisfies $K_{Z_{2,0}} \cdot V = 0$ so that all irreducible components of V are (-2) -curves contained in fibers. Finally, $V \cdot E = 0, 1, 2$, in the cases $n = 3k+2$, $n = 3k+1$, and $n = 3k$, respectively.

One can show that if K_X^n is small enough to compete with the examples in Section 4.3, then $\text{Vol}(Z_2)$ also must be equal to the minimum in (5.3). Further, $\pi_{\text{can}}|_{Z_2} : Z_2 \rightarrow X_{\text{can}}$ factors through the minimal model $g : Z_2 \rightarrow Z_{2,0}$. Using Proposition 5.2, we may identify some additional curves on $Z_{2,0}$ which are contracted by π_{can} . In each case, we will contract a connected curve of arithmetic genus 0 with negative-definite self-intersection matrix. This results in a projective surface with rational singularities (see [1, Theorem 2.3] and [2, Proposition 1]), which is therefore \mathbb{Q} -Gorenstein.

Case 5.1. $n = 3k+2$, $k \geq 1$.

This is the easiest case, because we can define the appropriate $f : Z_{2,0} \rightarrow S$ by only contracting the curve E of Proposition 5.2 on $Z_{2,0}$. This gives $K_{Z_{2,0}}^2 = f^*K_S + aE$ with $a = \frac{-k+1}{k+1} \leq 0$. Note that the $n = 2$ case must be treated separately. Assuming that $K_{Z_{2,0}}^2$ is its minimum of $\frac{8n-16}{3}$, we may compute

$$(n-1)^2 K_X^n \geq K_S^2 = \frac{8n-16}{3} - (aE)^2 = \frac{3(n-1)^2}{n+1}.$$

Therefore, $K_X^n \geq \frac{3}{n+1}$ and we recover the same volume bound.

We can see this curve E explicitly in Example 4.3: the general weighted projective hypersurface

$$X_{10(k+1)} \subset \mathbb{P}^{n+1}(1^{(n)}, 2(k+1), 5(k+1))$$

has canonical dimension $n-1$, with the canonical map a rational map $X \dashrightarrow \mathbb{P}^{n-1}$ having generic fiber a curve of genus 2.

Now, consider the intersection of $n - 2$ general sections of the reflexive sheaf K_X . This is a general surface $S = S_{10(k+1)} \subset \mathbb{P}^3(1, 1, 2(k+1), 5(k+1))$ with one singular point p , which is a quotient singularity of type $\frac{1}{k+1}(1, 1)$. This is a cone over the rational normal curve of degree $k + 1$, so we can explicitly resolve it with a single blowup at p . Denote this blowup as $f : Z_{2,0} \rightarrow S$. This has the same numerical properties as in the general case above. From the description as a weighted projective hypersurface, we know $K_S^2 = \frac{(3k+1)^2}{k+1}$. Therefore,

$$K_{Z_{2,0}}^2 = \frac{8n - 16}{3},$$

so that the smooth model has volume matching the minimum in Proposition 5.2. As the notation suggests, this is the same $Z_{2,0}$ obtained by desingularizing X with a blowup and applying the machinery of Section 3.1.

Case 5.2. $n = 3k + 1$, $k \geq 1$. If $K_{Z_{2,0}}^2$ achieves the minimum of $\frac{8n-14}{3}$, then by Proposition 5.2, $V \cdot E = 1$. In general, V could have several (-2) -curves as irreducible components, but a unique one, say V_1 , intersects E in a single point. There is a birational morphism $f : Z_{2,0} \rightarrow S$ contracting exactly the connected curve $V_1 \cup E$.

The morphism $\pi_{\text{can}} : Z_2 \rightarrow X_{\text{can}}$ contracts both these curves on the minimal model, so it factors through $f : Z_{2,0} \rightarrow S$ and $K_{Z_{2,0}} = f^*K_S + aE + bV_1$, where $a = \frac{1-k}{2k+1}$ and $b = \frac{2-2k}{2k+1}$ are both nonpositive. One can calculate

$$(n-1)^2 K_X^n \geq K_S^2 = \frac{8n-14}{3} - (aE + bV_1)^2 = \frac{6(n-1)^2}{2n+1},$$

so that $K_X^n \geq \frac{6}{2n+1}$, in agreement with Corollary 4.1.

Once again, these exceptional curves appear in Example 4.4. The general weighted projective hypersurface

$$X_{10k+6} \subset \mathbb{P}^{n+1}(1^{(n)}, 2k+1, 5k+3)$$

has canonical dimension $n - 1$ with the canonical map a rational map $X \dashrightarrow \mathbb{P}^{n-1}$ having generic fiber a curve of genus 2. As above, we consider the intersection of $n - 2$ general sections of K_X to obtain $S = S_{10k+6} \subset \mathbb{P}^3(1, 1, 2k+1, 5k+3)$. Then $K_S^2 = \frac{6(n-1)^2}{2n+1}$ and S has only one singular point, at $[0 : 1 : 0 : 0]$, which is a quotient singularity of type $\frac{1}{2k+1}(1, 5k+3) = \frac{1}{2k+1}(1, k+1)$. The resolution of any surface quotient singularity of the form $\frac{1}{P}(1, Q)$ is related to the *Hirzebruch-Jung continued fraction* of $\frac{P}{Q}$ (see, for example, [15, Section 2]). In this case,

$$\frac{2k+1}{k+1} = 2 - \frac{1}{k+1},$$

so $\frac{P}{Q} = [2, k+1]$ is the continued fraction representation. This means that the exceptional locus has two rational curves as irreducible components, with self-intersections -2 and $-(k+1) = \frac{1}{3}(n+2)$, corresponding to V_1 and E above, respectively. It's straightforward to verify (e.g. using [15, Section

2]) that the discrepancies also agree with the above, so that

$$K_{Z_{2,0}}^2 = \frac{8n - 14}{3},$$

as expected. For $n = 4$, the singularity $\frac{1}{3}(1, 2)$ is canonical (in particular, an A_2 singularity) and the resolution is crepant. For $n \geq 7$, S has worse than canonical singularities so $K_{Z_{2,0}}^2 < K_S^2$.

Case 5.3. $n = 3k$, $k \geq 1$. If $K_{Z_{2,0}}^2$ achieves the minimum of $\frac{8n-12}{3}$, then by Proposition 5.2, $V \cdot E = 2$. Unlike in the previous two cases, these intersection numbers do not pin down a unique component graph for the connected component of $V_{\text{red}} \cup E$ containing E . The simplest configuration is that there are two irreducible components V_1 and V_2 of V intersecting E , each in a single point, and no other components of V intersecting these. Fortunately, one can verify that the other possible configurations yield the same volume bound; suppose for clarity that we have the simple configuration. Once again, there is a birational morphism $f : Z_{2,0} \rightarrow S$ contracting exactly $E \cup V_1 \cup V_2$ and π_{can} factors through this morphism. $K_{Z_{2,0}} = f^*K_X + aE + bV_1 + cV_2$ with $a = c = \frac{1-k}{2k}$ and $b = \frac{1-k}{k}$ (both nonpositive). We have

$$(n-1)^2 K_X^n \geq K_S^2 = \frac{8n-12}{3} - (aE + bV_1 + cV_2)^2 = \frac{3(n-1)^2}{n}.$$

This gives $K_X^n \geq \frac{3}{n}$, as desired.

Example 4.5 again displays the same behavior. For any positive integer $n = 3k$, $k \geq 1$, the general weighted projective hypersurface

$$X_{20k+2} \subset \mathbb{P}^{n+1}(1, 2^{(n-1)}, 4k, 10k+1)$$

has canonical dimension $n - 1$. It has $K_X = \mathcal{O}_X(2)$ and volume $\frac{1}{k} = \frac{3}{n}$. The rational map $|K_X| = |\mathcal{O}_X(2)|$ has image $\mathbb{P}^{n-1}(1, 2^{(n-1)}) \cong \mathbb{P}^{n-1}$. The intersection of $n - 2$ general sections of K_X can be identified with a general $S_{20k+2} \subset \mathbb{P}^3(1, 2, 4k, 10k+1)$. To see this, let x_1 have degree 1 and x_2, \dots, x_n have degree 2, so that a section of $\mathcal{O}_X(2)$ is a polynomial of the form $a_1 x_1^2 + a_2 x_2 + \dots + a_n x_n$. After a suitable change of coordinates within x_2, \dots, x_n , we may assume that the common zero set of $n - 2$ general sections is given by equations $x_1^2 = x_3, x_1^2 = x_4, \dots, x_1^2 = x_n$, which yields the result.

This surface S satisfies $K_S^2 = \frac{3(n-1)^2}{n}$ and is smooth away from the weighted \mathbb{P}^1 with weights 2 and $4k$. The general S only intersects this stratum at the coordinate point of $4k$. At this base point, there is a singularity of type $\frac{1}{4k}(1, 2k+1)$. The corresponding Hirzebruch-Jung continued fraction is

$$\frac{4k}{2k+1} = 2 - \frac{1}{(k+1) - \frac{1}{2}},$$

also represented as $\frac{4k}{2k+1} = [2, k+1, 2]$. Therefore, the minimal resolution of this singularity is a chain of three exceptional rational curves with self-intersections -2 , $-(k+1) = \frac{1}{3}(n+3)$, and -2 . These play the role of

Remark 5.4. In the explicit examples in Section 4.3 and Example 5.3, the smooth models $Z_{2,0}$ of surfaces that appear are *Horikawa surfaces*, that is, they are on the Noether line $K_{Z_{2,0}}^2 = 2p_g(Z_{2,0}) - 4$. For such varieties, $|K_{Z_{2,0}}|$ gives a ramified double cover to a ruled surface and the types of ruled surfaces and branch loci that appear are completely classified (see [17]). This suggests the following question:

Question 5.5. If X is a minimal variety of general type of dimension n , canonical dimension $n - 1$, and smallest possible volume, is the minimal model of the associated general surface Z_2 a Horikawa surface?

Acknowledgments

M. Chen is a member of the Key Laboratory of Mathematics for Nonlinear Sciences, Fudan University. We would like to thank Chen Jiang and Burt Totaro for useful discussions and suggestions.

REFERENCES

- [1] M. Artin. *Some numerical criteria for contractibility of curves on algebraic surfaces*. Am. J. Math. **84** (1962), 485–496.
- [2] M. Artin. *On isolated rational singularities of surfaces*. Am. J. Math. **88** (1966), no. 1, 129–136.
- [3] E. Bombieri. *Canonical models of surfaces of general type*. Inst. Hautes études Sci. Publ. Math. **42** (1973), 171–219.
- [4] J. A. Chen, M. Chen, *Explicit birational geometry of threefolds of general type, I*, Ann. Sci. Éc. Norm. Supér. (4) **43** (2010), no.3, 365–394.
- [5] J. A. Chen, M. Chen, *Explicit birational geometry of 3-folds of general type, II*, J. Differential Geom. **86** (2010), 237–271.
- [6] J. A. Chen, M. Chen, *Explicit birational geometry for 3-folds and 4-folds of general type, III*, Compos. Math. **151** (2015), no. 6, 1041–1082.
- [7] J. A. Chen, M. Chen, and C. Jiang. *The Noether inequality for algebraic threefolds (with an appendix by J. Kollár)*. Duke Math. J. **169** (2020) 1603–1645. Addendum, Duke Math. J. **169** (2020) 2199–2204.
- [8] M. Chen, *Canonical stability in terms of singularity index for algebraic threefolds*. Math. Proc. Cambridge Philos. Soc. **131** (2001), no. 2, 241–264.
- [9] M. Chen, *Canonical stability of 3-folds of general type with $p_g \geq 3$* . Internat. J. Math. **14** (2003), no. 5, 515–528.
- [10] M. Chen, *A sharp lower bound for the canonical volume of 3-folds of general type*. Math. Ann. **337** (2007), no. 4, 887–908.
- [11] M. Chen, D.-Q. Zhang. *Characterization of the 4-canonical birationality of algebraic threefolds*. Math. Z. **258** (2008), no. 3, 565–585.
- [12] M. Chen, *On minimal 3-folds of general type with maximal pluricanonical section index*. Asian J. Math. **22** (2018), no. 2, 257–268.
- [13] M. Chen, C. Jiang, B. Li. *On minimal varieties growing from quasismooth weighted hypersurfaces*, arXiv: 2005.09828.
- [14] I. Dolgachev, *Weighted projective varieties*, In: *Group Actions and Vector Fields*, pp. 34–71. Lecture Notes in Math. 956, Springer-Verlag, Berlin, 1982.
- [15] P. Hacking, J. Tevelev, and G. Urzúa. *Flipping surfaces*. J. Algebraic Geom. **26**(2) (2017), 279–345.
- [16] C.D. Hacon, J. McKernan, *Boundedness of pluricanonical maps of varieties of general type*, Invent. Math. **166** (2006), no.1, 1–25.
- [17] E. Horikawa. *Algebraic surfaces of general type with small c_1^2 , I*. Ann. Math. **104** (1976), 357–387.

- [18] A.R. Iano-Fletcher, *Working with weighted complete intersections*, In: Explicit birational geometry of 3-folds, pp. 101–173, London Math. Soc. Lecture Note Ser., 281, Cambridge Univ. Press, Cambridge, 2000.
- [19] M. Kobayashi. *On Noether's inequality for threefolds*. J. Math. Soc. Japan **1** (1992), 145–156.
- [20] Y. Kawamata. *A generalization of Kodaira-Ramanujam's vanishing theorem*. Math. Ann. **261** (1982), no. 1, 43–46.
- [21] S. Takayama. *Pluricanonical systems on algebraic varieties of general type*. Invent. Math. **165** (2006), no. 3, 551–587.
- [22] L. Esser, B. Totaro, and C. Wang. *Varieties of general type with doubly exponential asymptotics*. arXiv:2109.13383.
- [23] H. Tsuji. *Pluricanonical systems of projective varieties of general type. I*. Osaka J. Math. **43** (2006), no. 4, 967–995.
- [24] E. Viehweg. *Vanishing theorems*. J. Reine Angew. Math. **335** (1982), 1–8.

School of Mathematical Sciences, Fudan University, Shanghai 200433, China
Email address: `mchen@fudan.edu.cn`

UCLA MATHEMATICS DEPARTMENT, BOX 951555, LOS ANGELES, CA 90095-1555
Email address: `esserl@math.ucla.edu`

UCLA MATHEMATICS DEPARTMENT, BOX 951555, LOS ANGELES, CA 90095-1555
Email address: `chwang@math.ucla.edu`