
Faster Algorithms for Sparse ILP and Hypergraph Multi-Packing/Multi-Cover Problems

D. V. Griбанov · D. S. Malyshev ·
N. Yu. Zolotykh

Received: date / Accepted: date

Abstract In our paper, we consider the following general problems: check feasibility, count the number of feasible solutions, find an optimal solution, and count the number of optimal solutions in $\mathcal{P} \cap \mathbb{Z}^n$, assuming that \mathcal{P} is a polyhedron, defined by systems $Ax \leq b$ or $Ax = b, x \geq \mathbf{0}$ with a sparse matrix A . We develop algorithms for these problems that outperform state of the art ILP and counting algorithms on sparse instances with bounded elements.

We use known and new methods to develop new exponential algorithms for *Edge/Vertex Multi-Packing/Multi-Cover Problems* on graphs and hypergraphs. This framework consists of many different problems, such as the *Stable Multi-set*, *Vertex Multi-cover*, *Dominating Multi-set*, *Set Multi-cover*, *Multi-set Multi-cover*, and *Hypergraph Multi-matching* problems, which are natural generalizations of the standard *Stable Set*, *Vertex Cover*, *Dominating Set*, *Set Cover*, and *Maximal Matching* problems.

Keywords Integer Linear Programming · Counting Problem · Parameterized Complexity · Multipacking · Multicover · Stable Set · Vertex Cover · Dominating Set · Multiset Multicover · Hypergraph Matching · Sparse Matrix

D. V. Griбанov

Laboratory of Algorithms and Technologies for Network Analysis, National Research University Higher School of Economics, 25/12 Bolshaja Pecherskaja Ulitsa, Nizhny Novgorod, 603155, Russian Federation
E-mail: dimitry.griбанov@gmail.com

D. S. Malyshev

Laboratory of Algorithms and Technologies for Network Analysis, National Research University Higher School of Economics, 25/12 Bolshaja Pecherskaja Ulitsa, Nizhny Novgorod, 603155, Russian Federation
E-mail: dsmalyshev@rambler.ru

N. Yu. Zolotykh

Mathematics of Future Technologies Center, Lobachevsky State University of Nizhni Novgorod, 23 Gagarin ave, Nizhni Novgorod, 603950, Russian Federation
E-mail: nikolai.zolotykh@itmm.unn.ru

1 Introduction

Let a polytope \mathcal{P} of dimension $d := \dim(\mathcal{P})$ be defined by one of the following ways:

(i) **System in the canonical form:**

$$\mathcal{P} = \{x \in \mathbb{R}^n : Ax \leq b\}, \quad \text{where } A \in \mathbb{Z}^{m \times n} \text{ and } b \in \mathbb{Q}^m; \quad (\text{Canon-Form})$$

(ii) **System in the standard form:**

$$\mathcal{P} = \{x \in \mathbb{R}_+^n : Ax = b\}, \quad \text{where } A \in \mathbb{Z}^{k \times n}, b \in \mathbb{Q}^k. \quad (\text{Standard-Form})$$

If \mathcal{P} is defined by a system in the form Standard-Form with an additional constraint $x \leq u$, for given $u \in \mathbb{Z}_{>0}^n$, we call such a system as **the system in the standard form with multiplicities**.

We consider the following problems:

Problem 1 (Feasibility)

Find a point x inside $\mathcal{P} \cap \mathbb{Z}^n$ or declare that $\mathcal{P} \cap \mathbb{Z}^n = \emptyset$. (Feasibility-IP)

Problem 2 (Counting)

Compute the value of $|\mathcal{P} \cap \mathbb{Z}^n|$ or declare that $|\mathcal{P} \cap \mathbb{Z}^n| = +\infty$.
(Count-IP)

Problem 3 (Optimization) Given $c \in \mathbb{Z}^n$, compute some $x^* \in \mathcal{P} \cap \mathbb{Z}^n$, such that

$$c^\top x^* = \max\{c^\top x : x \in \mathcal{P} \cap \mathbb{Z}^n\}. \quad (\text{Opt-IP})$$

Declare, if $\mathcal{P} \cap \mathbb{Z}^n = \emptyset$ or if the maximization problem is unbounded.

Problem 4 (Optimization and Counting) Given $c \in \mathbb{Z}^n$, compute the number of x^* , such that

$$c^\top x^* = \max\{c^\top x : x \in \mathcal{P} \cap \mathbb{Z}^n\}, \quad (\text{Opt-And-Count-IP})$$

and find an example of x^* , if such exists. Declare, if $\mathcal{P} \cap \mathbb{Z}^n = \emptyset$ or if the maximization problem is unbounded.

In our work, we analyze these problems in the assumption that the matrix A is sparse. To estimate the sparsity of A , it is convenient to use the maximal number of non-zero elements in rows and columns of A :

$$\text{rs}(A) := \max_i \|A_{i*}\|_0 \quad \text{and} \quad \text{cs}(A) := \max_j \|A_{*j}\|_0.$$

Here, $\|x\|_0 = |\{i : x_i \neq 0\}|$ denotes the number of non-zeros in a vector x and A_{i*} , A_{*j} denote the i -th row and the j -th column of A , respectively.

Additionally, we define the *total sparsity of A* as the minimum between above parameters:

$$\text{ts}(A) = \min\{\text{rs}(A), \text{cs}(A)\}.$$

For our purposes, we sometimes need to use a slightly weaker parameters that estimate the number of non-zero elements in non-degenerate square sub-matrices. The reason is that the matrix A can have multiple rows and columns in some problem definitions. We want to avoid these multiplicities, estimating the sparsity of the matrices. For arbitrary $A \in \mathbb{Z}^{m \times n}$, we define

$$\begin{aligned}\overline{\text{rs}}(A) &:= \max\{\text{rs}(B) : B \text{ is non-degenerate sub-matrix of } A\}, \\ \overline{\text{cs}}(A) &:= \max\{\text{cs}(B) : B \text{ is non-degenerate sub-matrix of } A\} \quad \text{and} \\ \overline{\text{ts}}(A) &= \min\{\overline{\text{rs}}(A), \overline{\text{cs}}(A)\}.\end{aligned}$$

Clearly, the new sparsity parameters are weaker than the standard $\text{rs}(A)$ and $\text{cs}(A)$:

$$\overline{\text{ts}}(A) \leq \text{ts}(A).$$

Another useful parameters to estimate the sparsity of A and to formulate our results are the matrix norms. We recall the definitions. The maximum absolute value of entries of a matrix A (also known as *the matrix max-norm*) is denoted by $\|A\|_{\max} = \max_{i,j} |A_{ij}|$. For a matrix A , by $\|A\|_p$ we denote the matrix norm, induced by the l_p vector norm. It is known that

$$\begin{aligned}\|A\|_1 &= \max_i \|A_{i*}\|_1 = \max_i \sum_j |A_{ij}| \quad \text{and} \\ \|A\|_{\infty} &= \max_j \|A_{*j}\|_1 = \max_j \sum_i |A_{ij}|.\end{aligned}$$

Again, we need a similar definition of a norm with respect to non-degenerate sub-matrices B of A :

$$\gamma_p(A) = \max\{\|B\|_p : B \text{ is a non-degenerate sub-matrix of } A\}.$$

The last important matrix parameters that are related to the sparsity are the values of matrix sub-determinants.

Definition 1 For a matrix $A \in \mathbb{Z}^{m \times n}$, by

$$\Delta_k(A) = \max\left\{|\det(A_{\mathcal{I}\mathcal{J}})| : \mathcal{I} \subseteq \{1, \dots, m\}, \mathcal{J} \subseteq \{1, \dots, n\}, |\mathcal{I}| = |\mathcal{J}| = k\right\},$$

we denote the maximum absolute value of determinants of all the $k \times k$ sub-matrices of A . Here, the symbol $A_{\mathcal{I}\mathcal{J}}$ denotes the sub-matrix of A , which is generated by all the rows with indices in \mathcal{I} and all the columns with indices in \mathcal{J} .

Note that $\Delta_1(A) = \|A\|_{\max}$. By $\Delta_{\text{gcd}}(A, k)$, we denote the greatest common divisor of determinants of all the $k \times k$ sub-matrices of A . Additionally, let $\Delta(A) = \Delta_{\text{rank}(A)}(A)$ and $\Delta_{\text{gcd}}(A) = \Delta_{\text{gcd}}(A, \text{rank}(A))$. The matrix A with $\Delta(A) \leq \Delta$, for some $\Delta > 0$, is called Δ -modular.

Due to the Hadamard's inequality and since $\det(B) = \det(B^\top)$ and $\|x\|_2 \leq \|x\|_1$, for any $B \in \mathbb{R}^{n \times n}$ and $x \in \mathbb{R}^n$, the following inequalities connect $\Delta_k(A)$, the matrix norms, and $\text{ts}(A)$:

$$\Delta_k(A) \leq \min\{\gamma_1(A), \gamma_\infty(A)\}^k \leq \min\{\|A\|_1, \|A\|_\infty\}^k, \quad (1)$$

$$\Delta_k(A) \leq (\|A\|_{\max})^k \cdot \overline{\text{ts}}(A)^{k/2} \leq (\|A\|_{\max})^k \cdot \text{ts}(A)^{k/2}. \quad (2)$$

Denoting $\gamma_{1,\infty}(A) = \min\{\gamma_1(A), \gamma_\infty(A)\}$, the inequality (1) becomes

$$\Delta_k(A) \leq \gamma_{1,\infty}(A)^k. \quad (3)$$

2 Results on Sparse ILP Problems and the Related Work

The state of the art algorithm, due to Dadush, Peikert & Vempala [23, 25] (see also [19, 38, 77] for more general formulations), for the problems Feasibility-IP and Opt-IP has the complexity bound $O(d)^d \cdot \text{poly}(\phi)$, where $\phi = \text{size}(A, b, c)$ is the input bit-encoding length. We will use the notation ϕ for it in the whole article. It is a long-standing open problem to break the $O(d)^d$ -dependence in the complexity of ILP algorithms. Useful discussion and new ideas on this problem can be found, for example, in [11, 23, 24].

The asymptotically fastest algorithm for the problem Count-IP in a fixed dimension can be obtained, using the approach of A. Barvinok [9] with modifications, due to Dyer & Kannan [27] and Barvinok & Pommersheim [8]. A complete exposition of the Barvinok's approach can be found in [7, 8, 10, 13, 26], additional discussion can be found in the book [60], due to J. Lasserre. An important notion of the *half-open sign decomposition* and other variant of the Barvinok's algorithm that is more efficient in practise is given by Köppe & Verdoolaeghe in [54]. The paper [10] of Barvinok & Woods gives important generalizations of the original techniques and adapts them to a wider range of problems to handle projections of polytopes. Using the fastest deterministic Shortest Lattice Vector Problem (SVP) solver by Micciancio & Voulgaris [67], the Barvinok's algorithm computational complexity can be estimated by

$$\nu \cdot 2^{O(d)} \cdot (\log_2(\Delta))^{d \log d}, \quad (4)$$

where $\Delta := \Delta(A)$ and $\nu := \nu(A)$ is the maximal number of vertices in the class of polytopes, defined by systems in the canonical or the standard forms with a fixed A and varying b . Since any polytope can be transformed to an integer-equivalent simple polytope, using a slight perturbation of the r.h.s. vector b , the parameterization by ν is correct (see, for example, [40, Theorem 3]). Let us assume that \mathcal{P} is defined by the form Canon-Form. Due to the seminal work of P. McMullen [66], the value of ν attains its maximum at the class of polytopes, which is dual to the class of cyclic polytopes. Consequently, due to [45, Section 4.7], $\nu = O(m/n)^{n/2}$. Due to [72, Chapter 3.2, Theorem 3.2], we have $\Delta \leq 2^\phi$. In notation with ϕ , the bound (5) becomes

$$O\left(\frac{m}{n}\right)^{n/2} \cdot (\log_2(\Delta))^{n \log n} = O\left(\frac{m}{n}\right)^{n/2} \cdot \phi^{n \log n}, \quad (5)$$

which gives a polynomial-time algorithm in a fixed dimension for the problem Count-IP.

The papers [39, 40, 41, 42] try to deal with the parameter Δ to give pseudo-polynomial algorithms, which will be more effective in a varying dimension. The next theorem, due to Griбанov & Malyshev [40], gives a polynomial-time complexity bound with respect to ν , d , and Δ .

Theorem 1 (Griбанov & Malyshev[40]) *Let \mathcal{P} be a polytope, given by a system in the standard or the canonical forms. Then, the problem Count-IP can be solved by an algorithm with the arithmetic complexity bound*

$$O(\nu \cdot d^3 \cdot \Delta^4 \cdot \log_2(\Delta)),$$

where $\nu := \nu(A)$ is the maximal possible number of vertices in \mathcal{P} , for fixed A and varying b .

We improve the last result in our paper, and it will be our main tool for sparse problems.

Theorem 2 *In assumptions of Theorem 1, the problem Count-IP can be solved by an algorithm with the arithmetic complexity bound:*

$$O(\nu \cdot d^3 \cdot \Delta^3).$$

The theorem proof is given in Subsection 2.2.1.

Using Theorem 1 and different ways to estimate ν , the paper [40] gives new interesting arithmetic complexity bounds for the Feasibility-IP and Count-IP problems. Let us present them, taking into account the improvement made in the Theorem 2:

- The bound

$$O\left(\frac{m}{n}\right)^{\frac{n}{2}} \cdot n^3 \cdot \Delta^3$$

for systems in the form Canon-Form that is polynomial on m and Δ , for any fixed n . In comparison with the bound (5), this bound has a much better dependence on n , considering Δ as a parameter. For example, taking $m = O(n)$ and $\Delta = 2^{O(n)}$, the above bound becomes $2^{O(n)}$, which is even faster, than the state of the art algorithm for the problem Feasibility-IP, due to Dadush, Peikert & Vempala [23, 25], with the complexity bound $O(n)^n \cdot \text{poly}(\phi)$;

- The general bound, for systems in the canonical or the standard forms,

$$O(d)^{3+\frac{d}{2}} \cdot \Delta^{3+d}$$

that is polynomial on Δ , for any fixed d . Taking $\Delta = d^{o(d)}$, the last bound becomes

$$O(d)^{\frac{d}{2}+o(d)},$$

which again gives a faster algorithm for the ILP feasibility problem, than the state of the art $O(n)^n \cdot \text{poly}(\phi)$ -complexity algorithm, due to [23, 25].

– The bound

$$O(n/k)^{k+3} \cdot \Delta^3 \quad (6)$$

for systems in the form Standard-Form, which is also valid for systems in the form Canon-Form with $k = m - n$, that is polynomial on n and Δ , for $k = O(1)$. Taking $k = 1$, it gives an $O(n^4 \cdot \Delta^3)$ -algorithm to compute the number of integer points in a simplex. The last result can be used to count solutions of the Unbounded Subset-Sum problem, which is formulated as follows. Given numbers w_1, \dots, w_n and W , we need to count the number of ways to exchange the value W by an unlimited number of coins with costs w_1, \dots, w_n . It can be done by algorithms with the arithmetic complexity bound

$$O(n^4 \cdot w_{\max}^3).$$

Moreover, this result can be used to handle the k -dimensional variant of the Unbounded Subset-Sum problem, when the costs w_i and W are represented by k -dimensional vectors. Using the Hadamard's bound, it gives the following arithmetic complexity bound:

$$O(n^{k+3} \cdot k^{k/2} \cdot w_{\max}^{3k}),$$

where $w_{\max} = \max_i \|w_i\|_{\infty}$.

Note that the earlier paper of Lasserre & Zeron [59] also gives a counting FPT-algorithm for the Unbounded Subset-Sum problem, parameterized by w_{\max} , but an exact complexity bound was not given.

In the current work, we try to estimate the value of ν in a different way, to handle ILP problems with sparse matrices. Additionally, we generalize Theorem 2 to work with the problem Opt-And-Count-IP. The resulting theorem is following:

Theorem 3 *Let \mathcal{P} be a polyhedron, defined by the system in Canon-Form. Then, the problems Feasibility-IP and Count-IP can be solved by an algorithm with the complexity bounds*

$$\begin{aligned} & \gamma_{1,\infty}(A)^{4n} \cdot 2^n \cdot \text{poly}(\phi), \\ & (\|A\|_{\max})^{4n} \cdot \overline{\text{ts}}(A)^{2.5n} \cdot 2^n \cdot \text{poly}(\phi), \end{aligned}$$

respectively.

The problem Opt-And-Count-IP can be solved by an algorithm with the complexity bounds

$$\begin{aligned} & \gamma_{1,\infty}(A)^{5n} \cdot (\|c\|_{\infty})^3 \cdot 4^n \cdot \text{poly}(\phi), \\ & (\|A\|_{\max})^{5n} \cdot (\|c\|_{\infty})^3 \cdot \overline{\text{ts}}(A)^{3.5n} \cdot 4^n \cdot \text{poly}(\phi), \end{aligned}$$

respectively.

Table 1: The complexity bounds for the problems Feasibility-IP, Count-IP, Opt-IP, and Opt-And-Count-IP in the form Canon-Form

Problems:	Time: ¹	Reference:
Feasibility-IP and Opt-IP	$O(n)^n$	Dadush, Peikert, Vempala [23, 25]
Count-IP	$O(m/n)^{n/2} \cdot \phi^{n \log n}$	Barvinok et al. [8, 9, 27]
Feasibility-IP and Count-IP	$\gamma_{1,\infty}(A)^{4n} \cdot 2^n$ $(\ A\ _{\max})^{4n} \cdot \overline{\text{ts}}(A)^{2.5n} \cdot 2^n$	this work
Opt-And-Count-IP	$\gamma_{1,\infty}(A)^{5n} \cdot (\ c\ _{\infty})^3 \cdot 4^n$ $(\ A\ _{\max})^{5n} \cdot (\ c\ _{\infty})^3 \cdot \overline{\text{ts}}(A)^{3.5n} \cdot 4^n$	this work

¹ The multiplicative factor $\text{poly}(\phi)$ is skipped.

The theorem proof is given in Subsection 2.2.2.

This new complexity bounds, applied to the problems in the form Canon-Form, are emphasised in Table 1. As the reader can see, in the assumptions that $\|A\|_{\max} = o(n)$ and $\|c\|_{\infty} = 2^{n \cdot o(\log n)}$ our complexity bounds, applied to sparse instances, are better than the general state of the art bound $O(n)^n \cdot \text{poly}(\phi)$.

The following corollary, which is a straightforward consequence of Theorem 3, shows that, in some assumptions, the Count-IP and Opt-And-Count-IP problems can be solved by a faster algorithm than the complexity bound (5) gives.

Corollary 1 *In the notation of Theorem 3, assuming that $\|A\|_{\max} = n^{O(1)}$ and $\|c\|_{\infty} = n^{O(n)}$, the problems Count-IP and Opt-And-Count-IP can be solved by algorithms with the complexity bound*

$$n^{O(n)} \cdot \text{poly}(\phi).$$

Due to Kratsch [58], the sparse ILP problems attain a polynomial kernelization with respect to the parameter $n + u$, where u is the maximal variable range. More precisely, it was shown that any ILP can be reduced to an equivalent ILP with $O(u^r \cdot n^r)$ variables and constraints with the coefficients bit-encoding length $O(\log(nu))$, where $r := \text{rs}(A)$. On the contrary, it was shown that if the range u is unbounded, then r -row-sparse ILP problems do not admit a polynomial kernelization unless $NP \subseteq \text{coNP}/\text{poly}$.

There are many other interesting works about the ILP's complexity with respect to the parameter Δ . Since a good survey is given in the work [41], we mention only the most remarkable results. The first paper that discovers fundamental properties of the bimodular ILP problem ($\Delta = 2$) is [76], due to Veselov & Chirkov. Using results of [76], a strong polynomial-time solvability of the bimodular ILP problem was proved by Artmann, Weismantel & Zenklusen in [3]. Very recently, this result was generalised by Glanzer, Stallknecht & Weismantel in [34]. More precisely, it was shown that if the set of all the minors of A consists of only at most 3 non-zero different absolute values, then the corresponding ILP can be solved by a polynomial-time algorithm.

Unfortunately, not much is known for $\Delta(A) \geq 3$. Very recently, it was shown by Fiorini, Joret, Weltge & Yuditsky in [29] that the ILP problem is polynomial-time solvable, for any fixed Δ , if the matrix A has at most 2 non-zeros per row or per column. Previously, a weaker result, based on the same reduction, was known, due to [1]. It states that any ILP with a $\{0, 1\}$ -matrix A , which has at most two non-zeros per row and a fixed value of $\Delta\left(\begin{smallmatrix} 1 \\ A \end{smallmatrix}\right)$, can be solved by a linear-time algorithm.

Additionally, we note that, due to Bonifas, Di Summa et al. [15], there are no polynomial-time algorithms for the ILP problems with $\Delta(A) = \Omega(n^\varepsilon)$, for any $\varepsilon > 0$, unless the ETH (the Exponential Time Hypothesis) is false. Consequently, with the same assumption, there are no algorithms for the Δ -modular ILP problem with the complexity bound $\Delta^{O(1)} \cdot \text{poly}(\phi)$, despite the fact that algorithms with complexities $\Delta^{f(\Delta)} \cdot \text{poly}(\phi)$ or $\phi^{f(\Delta)}$ may still exist.

The last fact is the reason why we need to use both parameters ν and Δ . As it was noted, the complexity bound $\Delta^{O(1)} \cdot \text{poly}(\phi)$ is unlikely to exist, while the bound $\nu \cdot \Delta^3 \cdot \text{poly}(\phi)$ is presented in this paper, which gives the opportunity to give efficient algorithms for sparse problems.

2.1 ILP Problems with a Bounded Co-dimension

In this subsection, we consider ILP problems in the form Standard-Form. Since in our definition $k = \text{rank}(A)$, it is essential to call the parameter k as the *co-dimension of \mathcal{P}* .

We are interested in the complexity bounds for bounded values of k . Let us survey some remarkable results. The following result, due to Griбанov et al. [41, see Theorem 8 and Corollary 9], gives a parameterization by k and $\Delta := \Delta(A)$.

Theorem 4 (Griбанov et al. [41]) *Assume that some $k \times k$ non-degenerate sub-matrix B of A is given and $\eta = \Delta/|\det(B)|$. Then, the problem *Opt-IP* can be solved by an algorithm with the arithmetic complexity bound*

$$O(k)^{k+1} \cdot \eta^{2k} \cdot \Delta^2 \cdot \log(\Delta_{\text{gcd}}) \cdot \log(k \cdot \Delta),$$

where $\Delta_{\text{gcd}} := \Delta_{\text{gcd}}(A)$.

Due to [41], we can assume that $\eta = O(\log(k))^k$, and the previous complexity bound becomes

$$O(\log(k))^{2k^2} \cdot k^{k+1} \cdot \Delta^2 \cdot \log(\Delta_{\text{gcd}}) \cdot \log(k \cdot \Delta).$$

As an alternative variant, we can directly compute a non-degenerate $n \times n$ sub-matrix B of A with $|\det(B)| = \Delta$. After such a preprocessing, we can assume that $\eta = 1$, which gives the following arithmetic complexity bound:

$$T_{\det}(m, n) + O(k)^{k+1} \cdot \Delta^2 \cdot \log(\Delta_{\text{gcd}}) \cdot \log(k \cdot \Delta),$$

where $T_{\det}(m, n)$ is the arithmetic complexity to find such B .

For the case, when A has non-negative elements, the basic dynamic-programming scheme from [14] can be used to derive an algorithm, parameterized by $\|b\|_\infty$ and k . Using fast $(\min, +)$ -convolution algorithms (see, for example, [37] or [70]), the same complexity bound can be used for systems in the Standard-Form form with multiplicities. We emphasize it to the following statement:

Proposition 1 *The problem Opt-IP in the form Standard-Form with multiplicities can be solved by an algorithm with the arithmetic complexity bound*

$$O\left(n \cdot (\|b\|_\infty + 1)^k\right).$$

Due to the works [22] and [30] of Cunningham & Geelen and Fomin et al., the parameter k in the term $(\|b\|_\infty + 1)^k$ can be replaced by stronger parameters 2ω or $\rho + 1$, where ω is the branch-width and ρ is the path-width of the column matroid of A .

The approach, which is most important for us, is based on the notion of the *hereditary discrepancy* of A .

Definition 2 For a matrix $A \in \mathbb{R}^{k \times n}$, its *discrepancy* and its *hereditary discrepancy* is defined by the formulas

$$\begin{aligned} \text{disc}(A) &= \min_{z \in \{-1/2, 1/2\}^n} \|Az\|_\infty, \\ \text{herdisc}(A) &= \max_{\mathcal{I} \subseteq \{1, \dots, n\}} \text{disc}(A_* \mathcal{I}). \end{aligned}$$

The paper [49], due to Jansen and Rohwedder, gives a powerful ILP algorithm, parameterized by $\text{herdisc}(A)$ and k , which will be our second main tool.

Theorem 5 (Jansen & Rohwedder [49]) *Let $H = \text{herdisc}(A)$ and assume that there exists an optimal solution x^* of the problem Opt-IP with $\|x^*\|_1 \leq K$. Then, the problem Opt-IP can be solved by an algorithm with the complexity bound*

$$O(H)^{2k} \cdot \log(K).$$

Different bounds on $\text{herdisc}(A)$ can be used to develop different complexity bounds for ILP problems. Due to the works [64] and [74] of Lovász, Spencer, & Vesztergombi, and Spencer, it is known that

$$\text{herdisc}(A) \leq 2 \text{disc}(A) \leq \eta_k \cdot \|A\|_{\max}, \quad \text{where } \eta_k \leq 12 \cdot \sqrt{k}. \quad (7)$$

Due to Beck and Fiala [12], the value of $\text{herdisc}(A)$ is bounded by the l_1 -norm of columns. More precisely,

$$\text{herdisc}(A) < \|A\|_\infty. \quad (8)$$

Additionally, Beck and Fiala conjectured that $\text{herdisc}(A) = O(\sqrt{\|A\|_\infty})$ and settling this has been an elusive open problem. The best known result in this direction is due to Banaszczyk [6]:

$$\text{herdisc}(A) = O\left(\sqrt{\|A\|_\infty \cdot \log(n)}\right). \quad (9)$$

The important matrix characteristic that is closely related to $\text{herdisc}(A)$ is $\text{detlb}(A)$. Due to Lovász, Spencer, & Vesztergombi [64], it can be defined as follows:

$$\text{detlb}(A) = \max_{t \in \{1, \dots, k\}} \sqrt[t]{\Delta_t(A)},$$

and it was shown in [64] that

$$\text{herdisc}(A) \geq 1/2 \cdot \text{detlb}(A).$$

Matoušek in [65] showed that $\text{detlb}(A)$ can be used to produce tight upper bounds on $\text{herdisc}(A)$:

$$\text{herdisc}(A) = O\left(\text{detlb}(A) \cdot \log(kn) \cdot \sqrt{\log(n)}\right).$$

The previous bound of Matoušek was improved by Jiang & Reis in [51]:

$$\text{herdisc}(A) = O\left(\text{detlb}(A) \cdot \sqrt{\log(k) \cdot \log(n)}\right). \quad (10)$$

Next, let us consider the problems Count-IP and Opt-And-Count-IP. Clearly, the number of vertices in a polyhedron, defined by a system in the Canon-Form, can be estimated by $\binom{n}{k} = O(n/k)^k$. The last fact in combination with Theorem 2 results to the following corollary, which gives a parameterization by $\Delta := \Delta(A)$ and k .

Corollary 2 *Assume that \mathcal{P} is bounded, then the problem Count-IP can be solved by an algorithm with the arithmetic complexity bound*

$$O(n/k)^k \cdot n^3 \cdot \Delta^3.$$

Remark 1 Note that if we already know an optimal solution x^* of the problem Opt-IP, we can solve the problem Opt-And-Count-IP, using Corollary 2 just by adding the equality $c^\top x = c^\top x^*$ to the problem's definition. Clearly, the resulting arithmetic complexity bound is

$$O(n/k)^{k+1} \cdot n^3 \cdot \|c\|_\infty^3 \cdot \Delta^3. \quad (11)$$

The next theorem considers the ILP problems in the standard form with sparse A . In this theorem, we just summarize the combinations of Theorem 5 with the different bounds on $\text{herdisc}(A)$, described in Subsection 2.1, to solve the problems Feasibility-IP and Opt-IP. Additionally, here we use Corollary 2 to solve the problems Count-IP and Opt-And-Count-IP.

Note that the 5-th complexity bound of the next theorem has already been proven in [49], we put it here for the sake of completeness.

Theorem 6 *Let \mathcal{P} be a polyhedron, defined by the form Standard-Form. The problems Feasibility-IP and Opt-IP can be solved by algorithms with the following complexity bounds:*

1. $O(\gamma_\infty(A))^{2k} = O(\|A\|_{\max})^{2k} \cdot \overline{\text{CS}}(A)^{2k}$,

2. $O(\gamma_\infty(A))^k \cdot 2^{k \cdot \log \log(n)} = O(\|A\|_{\max})^k \cdot \overline{\text{cs}}(A)^k \cdot 2^{k \cdot \log \log(n)}$,
3. $O(\gamma_1(A))^{2k} \cdot 2^{k \cdot \log(\log(k) \cdot \log(n))}$,
4. $O(\|A\|_{\max})^{2k} \cdot \overline{\text{rs}}(A)^k \cdot 2^{k \cdot \log(\log(k) \cdot \log(n))}$,
5. $O(\|A\|_{\max})^{2k} \cdot k^k$.

The problem Count-IP can be solved by algorithms with the following complexity bounds:

6. $O(n/k)^k \cdot \gamma_{1,\infty}(A)^{3k}$,
7. $O(n/k)^k \cdot (\|A\|_{\max})^{3k} \cdot \overline{\text{ts}}(A)^{1.5k}$.

The problem Opt-And-Count-IP can be solved by the same algorithm with the cost of an additional multiplicative term $(\|c\|_\infty)^3$ in the complexity bound.

Everywhere in the complexity bounds, we skip the $\text{poly}(\phi)$ multiplicative term.

Proof Due to Theorem 5, the problems Feasibility-IP and Opt-IP can be solved by algorithms with the arithmetic complexity bound $O(H)^{2k} \cdot \log(K)$, where $H = \text{herdisc}(A)$ and $K = \|x^*\|_1$, for any optimal solution x^* . It is known that the problem has an optimal solution x^* with $\text{size}(x^*) = \text{poly}(\phi)$, so $\log(K) = \text{poly}(\phi)$.

Now, the first bound follows from the inequality (8). The second bound follows from the inequality (9). To establish the third and the fourth bounds, we use the equality (10): $\text{herdisc}(A) = O(\det\text{lb}(A) \cdot \sqrt{\log(k) \cdot \log(n)})$. Due to the inequalities (2) and (3), we clearly have

$$\begin{aligned} \det\text{lb}(A) &\leq \|A\|_{\max} \cdot \sqrt{\overline{\text{rs}}(A)}, \\ \det\text{lb}(A) &\leq \|A\|_1. \end{aligned}$$

Putting these bounds to (10), it gives the third and the fourth complexity bounds. The fifth complexity bound directly follows from the inequality 7.

Now, let us consider the problems Count-IP and Opt-And-Count-IP. The 6-th and 7-th complexity bounds straightforwardly follow from the inequalities (2) and (3) on Δ and Corollary 2. To satisfy its prerequisites, \mathcal{P} needs to be bounded. If \mathcal{P} is unbounded, then we can check that $|\mathcal{P} \cap \mathbb{Z}^n| = 0$, using the algorithm for the problem Feasibility-IP. As it was already mentioned, its complexity can be estimated by $O(\|A\|_{\max})^{2k} \cdot k^k$, which has no effect on the desired bound. In the opposite case, we have $|\mathcal{P} \cap \mathbb{Z}^n| = +\infty$. So, we can assume that \mathcal{P} is bounded, and the result is true. Note additionally that, if $\mathcal{P} \cap \mathbb{Z}^n \neq \emptyset$, then we can use the same algorithm for the problem Feasibility-IP to find some $x \in \mathcal{P} \cap \mathbb{Z}^n$.

Finally, using the same reasoning, the complexity bounds for the problem Opt-And-Count-IP just follows from Corollary 2 and its Remark 1. \square

2.1.1 ILP problems in the form Standard-Form with multiplicities

Finally, before we will finish Subsection 2.1, let us consider ILP problems in the form Standard-Form with multiplicities.

Using the basic dynamic programming scheme from [28], combined with a linear-time algorithm for the (min, +)-convolution (see, for example, [41, Theorem 7], [37] or [70]), it is easy to prove the following proposition.

Proposition 2 *The problem Opt-IP in the form Standard-Form with multiplicities can be solved by an algorithm with the arithmetic complexity bound*

$$O(\chi + k)^k \cdot (\|A\|_{\max})^k,$$

where $\Delta := \Delta(A)$ and χ is a value of the l_1 -proximity bound. That is

$$\chi = \max_{x^*} \min_{z^*} \|x^* - z^*\|_1,$$

where x^* and z^* are optimal solutions of the LP relaxation and of the original ILP, respectively.

Different bounds on χ give different algorithms, based on Proposition 2. The paper [28] of Eisenbrand & Weismantel gives

$$\chi \leq k \cdot (2k \cdot \|A\|_{\max} + 1)^k.$$

The paper [62], due to Lee, Paat et al., gives

$$\chi \leq (2k + 1)^k \cdot \Delta.$$

Recent result of Lee, Paat et al. [63] states that

$$\chi \leq k \cdot (k + 1)^2 \cdot \Delta^3 + (k + 1) \cdot \Delta = O(k^3 \cdot \Delta^3).$$

The dependence on Δ in the last bound can be reduced by Averkov & Schymura [4]

$$\chi = O(k^5 \cdot \Delta^2). \quad (12)$$

Using Proposition 2 with the bound (12), we see that the ILP in the form Standard-Form with multiplicities can be solved by an algorithm with the arithmetic complexity bound

$$(\|A\|_{\max})^k \cdot O(\Delta)^{2k} \cdot k^{5k}.$$

Using the inequalities (3) and (2), the last bound transforms to the bounds

$$\begin{aligned} & O(\gamma_{1,\infty}(A))^{2k^2 + O(k \log k)}, \\ & O(\|A\|_{\max})^{2k^2 + O(k)} \cdot \overline{\text{fs}}(A)^{k^2 + O(k \log k)}. \end{aligned} \quad (13)$$

In Table 2, we summarize all the facts, mentioned in the current subsection. The complexity bounds for the problems Feasibility-IP, Opt-IP, Count-IP, Opt-And-Count-IP without multiplicities are taken from Theorem 6 and Remark 1. To handle the problems with multiplicities, we just take the complexity bound (13). We also mention that the existence of algorithms for the problems Count-IP and Opt-And-Count-IP in the form Standard-Form with multiplicities, parameterized by k and polynomial by n , is open, and it is a good direction for further research.

Table 2: New complexity bounds for ILP problems in the form Standard-Form

Problems:	Time : ¹
Opt-IP without mult.	$O(\gamma_\infty(A))^{2k} = O(\ A\ _{\max})^{2k} \cdot \overline{\text{cs}}(A)^{2k}$ $O(\gamma_\infty(A))^k \cdot 2^{k \cdot \log \log(n)} = O(\ A\ _{\max})^k \cdot \overline{\text{cs}}(A)^k \cdot 2^{k \cdot \log \log(n)}$ $O(\gamma_1(A))^{2k} \cdot 2^{k \cdot \log(\log(k) \cdot \log(n))}$ $O(\ A\ _{\max})^{2k} \cdot \overline{\text{fs}}(A)^k \cdot 2^{k \cdot \log(\log(k) \cdot \log(n))}$ $O(\ A\ _{\max})^k \cdot k^k$, due to Jansen & Rohwedder [50]
Count-IP without mult. ²	$O(n)^k \cdot \gamma_{1,\infty}(A)^{3k}$ $O(n)^k \cdot O(\ A\ _{\max})^{3k} \cdot \overline{\text{fs}}(A)^{1.5k}$
Opt-IP with mult.	$O(\gamma_{1,\infty}(A))^{2k^2 + O(k \log k)}$ $O(\ A\ _{\max})^{2k^2 + O(k)} \cdot \overline{\text{fs}}(A)^{k^2 + O(k \log k)}$
Count-IP with mult.	open problem

¹ The multiplicative factor $\text{poly}(\phi)$ is skipped.² To solve the problem Opt-And-Count-IP, we need to pay an additional multiplicative factor $(\|c\|_\infty)^3$.

2.2 Proofs of the main Theorems 2 and 3

We need some additional definitions to prove Theorem 2.

Definition 3 For a polyhedron \mathcal{P} and a vector $c \in \mathbb{Z}^n$, we denote

$$f(\mathcal{P}, c; \tau) = \sum_{z \in \mathcal{P} \cap \mathbb{Z}^n} e^{\langle c, z \rangle \tau}.$$

Definition 4 Let \mathcal{V} be a vector space and $\mathcal{A} \subseteq \mathcal{V}$ be a set. The *indicator* $[\mathcal{A}]$ of \mathcal{A} is the function $[\mathcal{A}]: \mathcal{V} \rightarrow \mathbb{R}$ defined by

$$[\mathcal{A}](x) = \begin{cases} 1, & \text{if } x \in \mathcal{A} \\ 0, & \text{if } x \notin \mathcal{A}. \end{cases}$$

Definition 5 Let us fix the Euclidean space \mathcal{V} and a lattice $\Lambda \subseteq \mathcal{V}$.

The polyhedron $\mathcal{P} \subseteq \mathcal{V}$ is called *rational*, if it can be defined by a system of finitely many inequalities

$$\langle a_i, x \rangle \leq b_i,$$

where $a_i \in \Lambda^\circ$ and $b_i \in \mathbb{Z}$.

The *algebra of rational polyhedra* $\mathcal{P}(\mathbb{Q}\mathcal{V})$ is the vector space, defined as the span of the indicator functions of all the rational polyhedra $\mathcal{P} \subset \mathcal{V}$.

We recall the following restatement of the theorem, proved by Khovanski & Pukhlikov [71], and, independently, by Lawrence [61], declared as Theorem 13.8b in [7, Section 13].

Theorem 7 (Khovanski & Pukhlikov [71], Lawrence [61]) *Let us fix the Euclidean space \mathcal{V} with $\dim(\mathcal{V}) = n$ and a lattice $\Lambda \subseteq \mathcal{V}$.*

Let $\mathcal{R}(\mathcal{V})$ be the space of functions in \mathcal{V} , spanned by functions of the type

$$\frac{e^{\langle c, v \rangle}}{(1 - e^{\langle c, u_1 \rangle}) \cdots (1 - e^{\langle c, u_n \rangle})},$$

where $v \in \Lambda$ and $u_i \in \Lambda \setminus \{\mathbf{0}\}$, for $i \in \{1, \dots, n\}$. Then, there exists a linear transformation

$$\mathcal{F}: \mathcal{P}(\mathbb{Q}\mathcal{V}) \rightarrow \mathcal{R}(\mathcal{V}),$$

such that the following properties hold:

- (1) Let $\mathcal{P} \subseteq \mathcal{V}$ be a non-empty rational polyhedron without lines and let $\mathcal{R} := \mathcal{R}_{\mathcal{P}} \subseteq \mathcal{V}$ be its recession cone. Then, for all $c \in \text{int}(K^\circ)$, the series

$$\sum_{z \in \mathcal{P} \cap \Lambda} e^{\langle c, z \rangle}$$

converges absolutely to a function $\mathcal{F}([\mathcal{P}])$.

- (2) If \mathcal{P} contains a line, then $\mathcal{F}([\mathcal{P}]) = 0$.

Note that hereafter we will use this Theorem 7 just with $\mathcal{V} = \mathbb{R}^n$ and $\Lambda = \mathbb{Z}^n$. The following lemma represents a core of Theorem 2 and contains a main improvement with respect to the counting algorithm from [40].

Lemma 1 Let $A \in \mathbb{Z}^{n \times n}$, $b \in \mathbb{Z}^n$, $\Delta = |\det(A)| > 0$. Let us consider the polyhedron $\mathcal{P} = \{x \in \mathbb{R}^n : Ax \leq b\}$.

Assume that $c \in \mathbb{Z}^n$ is given, such that $\langle c, h_i \rangle > 0$, where h_i are the columns of $A^* = \Delta \cdot A^{-1}$, for $i \in \{1, \dots, n\}$. Denote $\chi = \max_{i \in \{1, \dots, n\}} \{|\langle c, h_i \rangle|\}$.

Let, additionally, $S = PAQ$ be the SNF of A , where $P, Q \in \mathbb{Z}^{n \times n}$ are unimodular, and denote $\sigma = S_{nn}$.

Then, for any $\tau > 0$, the series $\mathfrak{f}(\mathcal{P}, c; \tau)$ converges absolutely to a function of the type

$$\frac{\sum_{i=-n \cdot \sigma \cdot \chi}^{n \cdot \sigma \cdot \chi} \epsilon_i \cdot e^{\alpha_i \cdot \tau}}{(1 - e^{\beta_1 \cdot \tau})(1 - e^{\beta_2 \cdot \tau}) \cdots (1 - e^{\beta_n \cdot \tau})},$$

where $\epsilon_i \in \mathbb{Z}_{\geq 0}$, $\beta_i \in \mathbb{Z}_{< 0}$, and $\alpha_i \in \mathbb{Z}$.

This representation can be found with an algorithm, having the arithmetic complexity bound

$$O(T_{\text{SNF}}(n) + \Delta \cdot n^2 \cdot \sigma \cdot \chi),$$

where $T_{\text{SNF}}(n)$ is the arithmetic complexity of computing the SNF for $n \times n$ integer matrices.

Proof After the unimodular map $x = Qx'$ and introducing slack variables y , the system $\{x \in \mathbb{Z}^n : Ax \leq b\}$ becomes

$$\begin{cases} Sx + Py = Pb \\ x \in \mathbb{Z}^n \\ y \in \mathbb{Z}_{\geq 0}^n. \end{cases}$$

Since P is unimodular, the last system is equivalent to the system

$$\begin{cases} Py = Pb \pmod{S\mathbb{Z}^n} \\ y \in \mathbb{Z}_{\geq 0}^n. \end{cases} \quad (14)$$

Denoting $\mathcal{G} = \mathbb{Z}^n / S\mathbb{Z}^n$, $g_0 = Pb \pmod{S\mathbb{Z}^n}$, $g_i = P_{*i} \pmod{S\mathbb{Z}^n}$, the last system (14) can be rewritten:

$$\begin{cases} \sum_{i=1}^n y_i g_i = g_0 \\ y \in \mathbb{Z}_{\geq 0}^n. \end{cases} \quad (15)$$

Note that points $x \in \mathcal{P} \cap \mathbb{Z}^n$ and the solutions y of the system (15) are connected by the bijective map $x = A^{-1}(b - y)$.

Let $r_i = |\langle g_i \rangle|$, for $i \in \{1, \dots, n\}$, and $r_{\max} := \max_{i \in \{1, \dots, n\}} \{r_i\}$. Clearly, $|\mathcal{G}| = |\det(S)| = \Delta$ and $r_{\max} \leq \sigma$.

For $k \in \{1, \dots, n\}$ and $g' \in \mathcal{G}$, let $\mathcal{M}_k(g')$ be the solutions set of the auxiliary system

$$\begin{cases} \sum_{i=1}^k y_i g_i = g' \\ y \in \mathbb{Z}_{\geq 0}^k, \end{cases}$$

and define

$$\mathfrak{g}_k(g'; \tau) = \sum_{y \in \mathcal{M}_k(g')} e^{-\langle c, \sum_{i=1}^k h_i y_i \rangle \tau}$$

Note that

$$\begin{aligned} f(\mathcal{P}, c; \tau) &= \sum_{z \in \mathcal{P} \cap \mathbb{Z}^n} e^{\langle c, z \rangle \tau} = \sum_{y \in \mathcal{M}_n(g_0)} e^{\langle c, A^{-1}(b-y) \rangle \tau} = \\ &= e^{\langle c, A^{-1}b \rangle \tau} \cdot \sum_{y \in \mathcal{M}_n(g_0)} e^{-\frac{1}{\Delta} \langle c, A^* y \rangle \tau} = e^{\langle c, A^{-1}b \rangle \tau} \cdot \mathfrak{g}_n(g_0; \frac{\tau}{\Delta}). \end{aligned} \quad (16)$$

The following formulas for $\mathfrak{g}_k(g'; \tau)$ were proven formally in [40]. Note that, since the cone $\{x \in \mathbb{R}^n : Ax \leq \mathbf{0}\}$ is pointed, the ring of such series is correctly defined.

$$\mathfrak{g}_1(g'; \tau) = \frac{e^{-\langle c, sh_1 \rangle \tau}}{1 - e^{-\langle c, r_1 h_1 \rangle \tau}}, \quad \text{where } s = \min\{y_1 \in \mathbb{Z}_{\geq 0} : y_1 \cdot g_1 = g'\}, \quad (17)$$

$$\mathfrak{g}_k(g'; \tau) = \frac{1}{1 - e^{-\langle c, r_k h_k \rangle \tau}} \cdot \sum_{i=0}^{r_k-1} e^{-\langle c, i h_k \rangle \tau} \cdot \mathfrak{g}_{k-1}(g' - i \cdot g_k; \tau), \quad (18)$$

$$\mathfrak{g}_k(g'; \tau) = \frac{\sum_{i=-k \cdot \sigma \cdot \chi}^{k \cdot \sigma \cdot \chi} \epsilon_i(k, g') \cdot e^{-i\tau}}{(1 - e^{-\langle c, r_1 h_1 \rangle \tau})(1 - e^{-\langle c, r_2 h_2 \rangle \tau}) \dots (1 - e^{-\langle c, r_k h_k \rangle \tau})}. \quad (19)$$

where $\epsilon_i(k, g') \in \mathbb{Z}_{\geq 0}$ are some coefficients, depending on k and g' . If the equation $\{y_1 \in \mathbb{Z}_{\geq 0} : y_1 g_1 = g'\}$ has no solutions, it is set $\mathfrak{g}_1(g'; \tau) := 0$.

It directly follows from the definitions of the vector c , from the formulas (17), (18) and the induction principle that the power series $\mathfrak{g}_k(g'; \tau)$ absolutely converges to the corresponding functions, for any $\tau > 0$.

Let us estimate the complexity to compute the representation (19) of $\mathfrak{g}_k(g'; \tau)$, for all $k \in \{1, \dots, n\}$ and $g' \in \mathcal{G}$, using the recurrence (18). In comparison to the paper [40], we will use a bit more sophisticated and efficient algorithm to do that.

Consider a quotient group $\mathcal{Q}_k = \mathcal{G} / \langle g_k \rangle$ and fix $\mathcal{Q} \in \mathcal{Q}_k$. Clearly, $\mathcal{Q} = q + \langle g_k \rangle$, where $q \in \mathcal{G}$ is a member of \mathcal{Q} , and $r_k = |\mathcal{Q}|$. For $j \in \{0, \dots, r_k - 1\}$, define

$$\mathfrak{h}_k(j; \tau) = (1 - e^{-\langle c, r_1 h_1 \rangle \tau}) \cdot \dots \cdot (1 - e^{-\langle c, r_k h_k \rangle \tau}) \cdot \mathfrak{g}_k(q + j \cdot g_k; \tau). \quad (20)$$

The formulas (17), (18) and (19) become

$$\mathfrak{h}_1(j; \tau) = e^{-\langle c, s h_1 \rangle \tau}, \quad \text{where } s = \min\{y_1 \in \mathbb{Z}_{\geq 0} : y_1 g_1 = q + j \cdot g_1\}, \quad (21)$$

$$\mathfrak{h}_k(j; \tau) = \sum_{i=0}^{r_k-1} e^{-\langle c, i h_k \rangle \tau} \cdot \mathfrak{h}_{k-1}(j - i; \tau), \quad (22)$$

$$\mathfrak{h}_k(j; \tau) = \sum_{i=-k \cdot \sigma \cdot \chi}^{k \cdot \sigma \cdot \chi} \epsilon_i(k, q + j \cdot g_k) \cdot e^{-i\tau}. \quad (23)$$

Assume first that $k = 1$. Then, clearly, all the values

$$\mathfrak{h}_1(0; \tau), \mathfrak{h}_1(1; \tau), \dots, \mathfrak{h}_1(r_1 - 1; \tau)$$

can be computed with $O(r_1)$ operations.

Assume now that $k \geq 2$ and that $(k-1)$ -th level has already been computed. By the k -th level, we mean all the functions $\mathfrak{h}_k(j; \tau)$, for $j \in \{1, \dots, r_k - 1\}$.

Due to the formula (23), $\mathfrak{h}_k(j; \tau)$ contains $O(k \cdot \sigma \cdot \chi)$ monomials. Hence, the function $\mathfrak{h}_k(0; \tau)$ can be computed directly using the formula (22) with $O(r_k \cdot k \cdot \sigma \cdot \chi)$ operations. For $j \geq 1$, we have

$$\begin{aligned} \mathfrak{h}_k(j; \tau) &= \sum_{i=0}^{r_k-1} e^{-\langle c, i h_k \rangle \tau} \cdot \mathfrak{h}_{k-1}(j - i; \tau) = \\ &= \sum_{i=-1}^{r_k-2} e^{-\langle c, (i+1) h_k \rangle \tau} \cdot \mathfrak{h}_{k-1}(j - 1 - i; \tau) = \\ &= e^{-\langle c, h_k \rangle \tau} \cdot \mathfrak{h}_k(j - 1; \tau) + \\ &= \mathfrak{h}_{k-1}(j; \tau) + e^{-\langle c, (r_k-1) h_k \rangle \tau} \cdot \mathfrak{h}_{k-1}(j - (r_k - 1); \tau). \end{aligned} \quad (24)$$

Consequently, in the assumption that the $(k-1)$ -th level has already been computed and that $\mathfrak{h}_k(0; \tau)$ is known, all the functions $\mathfrak{h}_k(1; \tau), \dots, \mathfrak{h}_k(r_k -$

$1; \tau)$ can be computed with $O(r_k \cdot k \cdot \sigma \cdot \chi)$ operations, using the last formula (24).

When the functions $\mathfrak{h}_k(j; \tau)$, for $j \in \{0, \dots, r_k - 1\}$, are already computed, we can return to the functions $\mathfrak{g}_k(g'; \tau)$, for $g' = q + j \cdot g_k$, using the formula (20). It will consume additional $O(r_k)$ group operations to compute $g' = q + j \cdot g_k$. By the definition of \mathcal{G} , the arithmetic cost of a one group operation can be estimated by the number of elements on the diagonal of S that are not equal to 1. Clearly, this number is bounded by $\min\{n, \log_2(\Delta)\}$. Consequently, the arithmetic cost of the last step is $O(r_k \cdot n)$, which is negligible in comparison with the $\mathfrak{h}_k(j; \tau)$ computational cost.

Summarizing, we need $O(r_k \cdot k \cdot \sigma \cdot \chi)$ operations to compute $\mathfrak{g}_k(g'; \tau)$, for $g' = q + j \cdot g_k$ and $j \in \{0, \dots, r_k\}$. Therefore, since $|\mathcal{Q}| = \Delta/r_k$, the arithmetic computational cost to compute k -th level of $\mathfrak{g}_k(\cdot)$ is

$$O(\Delta \cdot k \cdot \sigma \cdot \chi),$$

and the total arithmetic cost to compute all the levels is

$$O(\Delta \cdot n^2 \cdot \sigma \cdot \chi).$$

Finally, using the formula (16), we construct the function

$$\begin{aligned} \mathfrak{f}(\mathcal{P}, c; \tau) &= e^{\langle c, A^{-1}b \rangle \tau} \cdot \mathfrak{g}_n(g_0; \frac{\tau}{\Delta}) = \\ &= \frac{\sum_{i=-k \cdot \sigma \cdot \chi}^{k \cdot \sigma \cdot \chi} \epsilon_i \cdot e^{\frac{1}{\Delta} (\langle c, A^*b \rangle - i) \tau}}{(1 - e^{-\langle c, \frac{r_1}{\Delta} h_1 \rangle \tau}) (1 - e^{-\langle c, \frac{r_2}{\Delta} h_2 \rangle \tau}) \dots (1 - e^{-\langle c, \frac{r_n}{\Delta} h_n \rangle \tau})}, \end{aligned}$$

where $\epsilon_i := \epsilon_i(n, g_0)$, which gives the desired representation of $\mathfrak{f}(\mathcal{P}, c; \tau)$. Since $\mathfrak{g}_n(g_0; \tau)$ converges absolutely, for all $\tau > 0$, the same is true for $\mathfrak{f}(\mathcal{P}, c; \tau)$. Clearly, the arithmetic cost of the last transformation is proportional to the nominator length of $\mathfrak{g}_n(g_0; \tau)$, which is $O(n \cdot \sigma \cdot \chi)$. \square

2.2.1 The proof of Theorem 2

Proof Since any system in the standard form can be straightforwardly transformed to a system in the canonical form without changing the solutions set, assume that the polytope \mathcal{P} is defined by a system $Ax \leq b$, where $A \in \mathbb{Z}^{m \times n}$, $b \in \mathbb{Q}^m$, and $\Delta(A) = \Delta$.

Since \mathcal{P} is bounded, it follows that $\text{rank}(A) = n$. Since b is a rational vector, we can assume that $\text{gcd}(A_j) = 1$, for all $j \in \{1, \dots, m\}$.

Now, let us assume that $\dim(\mathcal{P}) < n$. Clearly, it is equivalent to the existence of an index $j \in \{1, \dots, m\}$, such that $A_j x = b_j$, for all $x \in \mathcal{P}$. Note that such j could be found by a polynomial-time algorithm. W.l.o.g., assume that $j = 1$. Let $A_1 = HQ$, where $H \in \mathbb{Z}^{1 \times n}$ is the *Hermite Normal Form (HNF)* of A_1 and $Q \in \mathbb{Z}^{n \times n}$ is unimodular. Since $\text{gcd}(A_1) = 1$, we have $H = (1 \ \mathbf{0}_{n-1})$.

After the unimodular map $x' = Qx$, the system $Ax \leq b$ transforms to the integrally equivalent system

$$\begin{pmatrix} 1 & \mathbf{0}_{n-1} \\ h & B \end{pmatrix} x \leq b,$$

where $h \in \mathbb{Z}^{m-1}$ and $B \in \mathbb{Z}^{(m-1) \times (n-1)}$. Note that $\Delta(B) = \Delta(A) = \Delta$. Since the first inequality always holds as an equality on the solutions set, we can just substitute $x_1 = b_1$. As the result we achieve a new integrally equivalent system with $n - 1$ variables $Bx \leq b'$, where $b' = b_{\{2, \dots, m\}} - b_1 \cdot h$.

Due to the proposed reasoning, we can assume that $\dim(\mathcal{P}) = n$. Let us do some more assumptions on \mathcal{P} . Due to [40, Section 2.3], using $O(n)$ operations, we can produce a new r.h.s. vector b' , such that a new polytope, defined by $Ax \leq b'$, will be simple and integrally equivalent to \mathcal{P} . Consequently, we can assume that \mathcal{P} is simple.

Let $v \in \text{vert}(\mathcal{P})$, denote

$$\begin{aligned} \mathcal{J}(v) &= \{j : A_j v = b_j\}, \text{ and} \\ \mathcal{P}_v &= \{x \in \mathcal{V} : A_{\mathcal{J}(v)} x \leq b_{\mathcal{J}(v)}\}. \end{aligned}$$

Since \mathcal{P} is simple, it follows that $A_{\mathcal{J}(v)} \in \mathbb{Z}^{n \times n}$ and $0 < \det(A_{\mathcal{J}(v)}) \leq \Delta$.

Due to the seminal work [5], all vertices of the simple polyhedron \mathcal{P} can be enumerated with $O(m \cdot n \cdot |\text{vert}(\mathcal{P})|)$ arithmetic operations. Due to [63], we can assume that $m = O(n^2 \cdot \Delta^2)$. Hence, all the polyhedrons \mathcal{P}_v can be constructed with $O(\nu \cdot n^3 \cdot \Delta^2)$ operations.

Let us set $c := \sum_{i \in \{1, \dots, n\}} B_{*i}$, where B^\top is some non-degenerate $n \times n$ submatrix of A . As it was shown in [40], such the choice of c satisfies requirements of Lemma 1, applied to any \mathcal{P}_v , for $v \in \text{vert}(\mathcal{P})$. Additionally, it was shown that $\chi \leq n\Delta$. We use Lemma 1 to all \mathcal{P}_v with the proposed choice of c , and construct the corresponding functions $f_v(\tau)$. Since $\sigma \leq \Delta$, $\chi \leq n\Delta$ and, due to Storjohann [75], $T_{SNF}(n) = O(n^3)$, the arithmetic complexity of the last operation can be estimated by

$$O(\nu \cdot n^3 \cdot \Delta^3).$$

Denote, additionally, $f_{\mathcal{P}}(\tau) = \sum_{v \in \text{vert}(\mathcal{P})} f_v(\tau)$.

Due to Brion's theorem [18] (see also [7, Chapter 6]), we have

$$[\mathcal{P}] = \sum_{v \in \text{vert}(\mathcal{P})} [\mathcal{P}_v] \quad \text{modulo polyhedra with lines.} \quad (25)$$

Consequently, it follows from Theorem 7 and the last formula (25) that, for any $\tau \in \mathbb{R}$, the series $\mathfrak{f}(\mathcal{P}, c; \tau)$ absolutely converges to the function $f_{\mathcal{P}}(\tau)$. Therefore, to calculate $|\mathcal{P} \cap \mathbb{Z}^n|$, we need to compute $\lim_{\tau \rightarrow 0} f_{\mathcal{P}}(\tau)$.

We follow to [7, Chapter 14], to compute $|\mathcal{P} \cap \mathbb{Z}^n| = \lim_{\tau \rightarrow 0} f_{\mathcal{P}}(\tau)$ as a constant term in the Taylor decomposition of $f_{\mathcal{P}}(\tau)$. Clearly, the constant term

of $f_{\mathcal{P}}(\tau)$ is just the sum of constant terms of $f_v(\tau)$, for $v \in \text{vert}(\mathcal{P})$. By this reason, let us fix some v and consider

$$f_v(\tau) = \frac{\sum_{i=-n \cdot \sigma \cdot \chi}^{n \cdot \sigma \cdot \chi} \epsilon_i \cdot e^{\alpha_i \cdot \tau}}{(1 - e^{\beta_1 \cdot \tau})(1 - e^{\beta_2 \cdot \tau}) \dots (1 - e^{\beta_n \cdot \tau})},$$

where $\epsilon_i \in \mathbb{Z}_{\geq 0}$, $\beta_i \in \mathbb{Z}_{< 0}$, and $\alpha_i \in \mathbb{Z}$.

Due to [7, Chapter 14], we can see that the constant term in the Taylor decomposition for $f_v(\mathcal{P})$ is exactly

$$\sum_{i=-n \cdot \sigma \cdot \chi}^{n \cdot \sigma \cdot \chi} \frac{\epsilon_i}{\beta_1 \dots \beta_n} \sum_{j=0}^n \frac{\alpha_i^j}{j!} \cdot \text{td}_{n-j}(\beta_1, \dots, \beta_n), \quad (26)$$

where $\text{td}_j(\beta_1, \dots, \beta_n)$ is a homogeneous polynomial of degree j , called the j -th Todd polynomial on β_1, \dots, β_n . Due to [26, Theorem 7.2.8, p. 137], the values of $\text{td}_j(\beta_1, \dots, \beta_n)$, for $j \in \{1, \dots, n\}$, can be computed with an algorithm that is polynomial in n and the bit-encoding length of β_1, \dots, β_n . Moreover, it follows from the theorem's proof that the arithmetic complexity can be bounded by $O(n^3)$. Therefore, it is not hard to see that we need $O(n^3)$ operations to compute the value of (26), and the total arithmetic cost to find the constant term in the Taylor's decomposition of the whole function $f_{\mathcal{P}}(\tau)$ is $O(\nu \cdot n^3)$, which is negligible with respect to $O(\nu \cdot n^3 \cdot \Delta^3)$ operations to compute $f_{\mathcal{P}}(\tau)$. So, the desired complexity bound is achieved. \square

The following lemmas help to estimate the number of vertices in a polyhedron, defined by a sparse system. We will use this bound to prove Theorem 3.

Lemma 2 *Let $A \in \mathbb{Z}^{n \times n}$, $\det(A) \neq 0$, and $\|\cdot\|: \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$ be any vector norm, which is symmetric with respect to any coordinate, i.e. $\|x\| = \|x - 2x_i \cdot e_i\|$, for any $x \in \mathbb{R}^n$ and $i \in \{1, \dots, n\}$.*

Let us consider a sector $\mathcal{U} = \mathbb{B}_{\|\cdot\|} \cap \text{cone}(A)$, where $\mathbb{B}_{\|\cdot\|} = \{x \in \mathbb{R}^n: \|x\| \leq 1\}$ is the unit ball with respect to the $\|\cdot\|$ -norm. Then,

$$\text{vol}(\mathcal{U}) \geq \frac{|\det(A)|}{2^n} \cdot \text{vol}(r \cdot \mathbb{B}_{\|\cdot\|}), \quad (27)$$

where $r \cdot \mathbb{B}_{\|\cdot\|}$ is the $\|\cdot\|$ -ball of the maximum radius r , inscribed into the set $\{x \in \mathbb{R}^n: \|Ax\| \leq 1\}$.

Consequently, let $\mathcal{U}_1 = \mathbb{B}_1 \cap \text{cone}(A)$ and $\mathcal{U}_\infty = \mathbb{B}_\infty \cap \text{cone}(A)$. Then,

$$\text{vol}(\mathcal{U}_1) \geq \frac{|\det(A)|}{(2\|A\|_\infty)^n} \cdot \text{vol}(\mathbb{B}_1) \geq \frac{|\det(A)|}{(2\|A\|_{\max} \cdot \text{cs}(A))^n} \cdot \text{vol}(\mathbb{B}_1); \quad (28)$$

$$\text{vol}(\mathcal{U}_\infty) \geq \frac{|\det(A)|}{(2\|A\|_1)^n} \cdot \text{vol}(\mathbb{B}_\infty) \geq \frac{|\det(A)|}{(2\|A\|_{\max} \cdot \text{rs}(A))^n} \cdot \text{vol}(\mathbb{B}_\infty). \quad (29)$$

Proof Let us prove the inequality (27). Clearly,

$$\text{vol}(\mathcal{U}) = |\det(A)| \cdot \text{vol}(\mathcal{K} \cap \text{cone}(I_{n \times n})),$$

where $\mathcal{K} = \{x \in \mathbb{R}^n : \|Ax\| \leq 1\}$.

By definition of r , we have $\mathcal{K} \supseteq r \cdot \mathbb{B}_{\|\cdot\|}$. Consequently,

$$\text{vol}(\mathcal{U}) \geq |\det(A)| \cdot \text{vol}(r \cdot \mathbb{B}_{\|\cdot\|} \cap \text{cone}(I_{n \times n})) \geq \frac{|\det(A)|}{2^n} \cdot \text{vol}(r \cdot \mathbb{B}_{\|\cdot\|}).$$

Now, let us prove the inequality (28). To this end, we just need to prove the inequality $r \geq \frac{1}{\|A\|_\infty}$ with respect to the l_1 -norm. Let us consider the set \mathcal{K} . It can be represented as the set of solutions of the following inequality:

$$\sum_{i=1}^n |A_{i*}x| \leq 1. \quad (30)$$

Let us consider the $2n$ points $\pm p_i = \pm \frac{1}{\|A\|_\infty} \cdot e_i$, for $i \in \{1, \dots, n\}$. Substituting $\pm p_j$ to the inequality (30), we have

$$\sum_{i=1}^n |A_{i*}p_j| = \frac{1}{\|A\|_\infty} \cdot \sum_{i=1}^n |A_{i*}e_j| = \frac{1}{\|A\|_\infty} \cdot \sum_{i=1}^n |A_{ij}| \leq 1.$$

Hence, all the points $\pm p_i$, for $i \in \{1, \dots, n\}$, satisfy the inequality (30). Since \mathcal{K} is convex, we have $\frac{1}{\|A\|_\infty} \cdot \mathbb{B}_1 \subseteq \mathcal{K}$, and, consequently, $r \geq \frac{1}{\|A\|_\infty}$.

Finally, let us prove the inequality (29). Again, we need to show that $r \geq \frac{1}{\|A\|_1}$ with respect to the l_1 -norm. In the current case, the set \mathcal{K} can be represented as the set of solutions of the following system:

$$\forall i \in \{1, \dots, n\}, \quad |A_{i*}x| \leq 1. \quad (31)$$

Let us consider the set $\mathcal{M} = \{\frac{1}{\|A\|_1} \cdot (\pm 1, \pm 1, \dots, \pm 1)^\top\}$ of 2^n points. Substituting any point $p \in \mathcal{M}$ to the j -th inequality of the system (31), we have

$$|A_{j*}p| \leq \sum_{i=1}^n |A_{ji}| |p_i| = \frac{1}{\|A\|_1} \cdot \sum_{i=1}^n |A_{ji}| \leq 1.$$

Hence, all the points $p \in \mathcal{M}$ satisfy the inequality (31). Since \mathcal{K} is convex, we have $\frac{1}{\|A\|_1} \cdot \mathbb{B}_\infty \subseteq \mathcal{K}$, and, consequently, $r \geq \frac{1}{\|A\|_1}$. \square

Lemma 3 *Let $A \in \mathbb{Z}^{m \times n}$, $b \in \mathbb{Q}^m$, and $\text{rank}(A) = n$. Let \mathcal{P} be a polyhedron, defined by a system $Ax \leq b$. Then,*

$$|\text{vert}(\mathcal{P})| \leq 2^n \cdot \gamma_{1,\infty}(A)^n \leq (2\|A\|_{\max})^n \cdot \overline{\text{ts}}(A)^n.$$

Proof Let $\mathcal{N}(v) = \text{cone}(A_{\mathcal{J}(v)}^\top)$ be the normal cone of a vertex $v \in \text{vert}(\mathcal{P})$, where $\mathcal{J}(v) = \{j \in \{1, \dots, m\} : A_{j*}v = b_j\}$. Since $\text{rank}(A) = n$, we have $\dim(\mathcal{N}(v)) = n$, for any $v \in \text{vert}(\mathcal{P})$. It is a known fact that $\dim(\mathcal{N}(v_1) \cap \mathcal{N}(v_2)) < n$, for different $v_1, v_2 \in \text{vert}(\mathcal{P})$. Next, we will use the following trivial inclusion

$$\bigcup_{v \in \text{vert}(\mathcal{P})} \mathcal{N}(v) \cap \mathbb{B} \subseteq \mathbb{B}, \quad (32)$$

where \mathbb{B} is the unit ball with respect to any vector norm $\|\cdot\|: \mathbb{R}^n \rightarrow \mathbb{R}_{\geq 0}$.

Again, since $\text{rank}(A) = n$, each matrix $A_{\mathcal{J}(v)}^\top$ contains a non-degenerate $n \times n$ sub-matrix. Taking $\mathbb{B} := \mathbb{B}_1$ or $\mathbb{B} := \mathbb{B}_\infty$, by Lemma 2, we have $\text{vol}(\mathcal{N}(v) \cap \mathbb{B}) \geq \frac{\text{vol}(\mathbb{B})}{(2\gamma)^n}$, where $\gamma := \gamma_{1,\infty}(A)$. Finally, due to (32), we have

$$\frac{\text{vol}(\mathbb{B})}{(2\gamma)^n} \cdot |\text{vert}(\mathcal{P})| \leq \text{vol}(\mathbb{B}). \quad \square$$

2.2.2 The proof of Theorem 3

Proof Consider first the case, when \mathcal{P} is unbounded. We need to distinguish between two possibilities: $|\mathcal{P} \cap \mathbb{Z}^n| = 0$ and $|\mathcal{P} \cap \mathbb{Z}^n| = +\infty$. Due to [72, Theorem 17.1], if $|\mathcal{P} \cap \mathbb{Z}^n| \neq 0$, then there exists $v \in \mathcal{P} \cap \mathbb{Z}^n$ such that $\|v\|_\infty \leq (n+1) \cdot \Delta_{ext}$, where $\Delta_{ext} = \Delta(A_{ext})$ and $A_{ext} = (Ab)$ is the extended matrix of the system $Ax \leq b$. Consequently, to transform the unbounded case to the bounded one, we just need to add the inequalities $|x_i| \leq n^{n/2} \cdot (\|A_{ext}\|_{\max})^n$, for $i \in \{1, \dots, n\}$, to the original system $Ax \leq b$.

Now, we can assume that \mathcal{P} is bounded, and consequently $\text{rank}(A) = n$. Due to Theorem 2, the counting problem can be solved by an algorithm with the arithmetic complexity bound

$$O(\nu \cdot n^3 \cdot \Delta^3), \quad (33)$$

where ν is the maximal number of vertices in polyhedra with fixed A and varying b . In our case, the value of ν can be estimated by Lemma 3. To estimate the value of Δ , we use the inequalities (2) and (3).

The inequalities for ν and $\Delta(A)$, together with the bound (33), give the desired complexity bound for the problem Count-IP.

Let us show how to find some point z inside of $\mathcal{P} \cap \mathbb{Z}^n$ in the case $|\mathcal{P} \cap \mathbb{Z}^n| > 0$, to handle the problem Feasibility-IP. For $\alpha, \beta \in \mathbb{Z}$, let us consider the polyhedron $\mathcal{P}'(\alpha, \beta)$, defined by the system $Ax \leq b$ with the additional inequality $\alpha \leq x_1 \leq \beta$. Let v be some vertex of \mathcal{P} , which can be found by a polynomial-time algorithm. Due to [21], if $\mathcal{P} \cap \mathbb{Z}^n \neq \emptyset$, then there exists a point $z \in \mathcal{P} \cap \mathbb{Z}^n$ such that $\|v - z\|_\infty \leq n \cdot \Delta$. So, the value of z_1 can be found, using the binary search with questions to the $\mathcal{P}'(\alpha, \beta) \cap \mathbb{Z}^n$ -feasibility oracle. Clearly, we need $O(\log(n\Delta))$ calls to the oracle. After the moment, when we already know the value of z_1 , we just add the equality $x_1 = z_1$ to the system $Ax \leq b$ and start a similar search procedure for the value of z_2 . The total number of calls to the binary search oracle to compute all the components of z is $O(n \cdot \log(n\Delta))$.

Finally, let us explain how to deal with the problem Opt-And-Count-IP. Let $\alpha, \beta \in \mathbb{Z}$, consider the polyhedron $\mathcal{P}'(\alpha, \beta)$, defined by the system $Ax \leq b$ with the additional inequality $\alpha \leq c^\top x \leq \beta$.

Let $A' \in \mathbb{Z}^{(m+2) \times n}$ be the matrix that defines $\mathcal{P}'(\alpha, \beta)$, i.e. $A' = \begin{pmatrix} c^\top \\ -c^\top \\ A \end{pmatrix}$.

Expanding sub-determinants of A' by the c^\top -row, we have $\Delta(A') \leq \|c\|_1 \cdot \Delta_{n-1}(A)$. The value of $\Delta_{n-1}(A)$ can be estimated in the same way, as it was done for $\Delta(A)$.

Let us estimate the number of vertices in $\mathcal{P}'(\alpha, \beta)$. The polyhedron $\mathcal{P}'(\alpha, \beta)$ is the intersection of the polyhedron \mathcal{P} with the slab $\{x \in \mathbb{R}^n : \alpha \leq c^\top x \leq \beta\}$. Clearly, the new vertices may appear only on edges of \mathcal{P} , by at most 2 new vertices per edge. The number of edges in \mathcal{P} is bounded by $|\text{vert}(\mathcal{P})|^2/4$. In turn, the value of $|\text{vert}(\mathcal{P})|^2/4$ can be estimated, using Lemma 3.

Due to Theorem 2, the value $|\mathcal{P}'(\alpha, \beta) \cap \mathbb{Z}^n|$ can be computed by an algorithm with the desired complexity bounds. To complete the proof, we note that, using the binary search method, the original optimization problem can be reduced to a polynomial number of feasibility questions in the set $\mathcal{P}'(\alpha, \beta) \cap \mathbb{Z}^n$ for different α, β . \square

3 Applications: The Vertex/Edge Multi-Packing and Multi-Cover Problems on Graphs and Hypergraphs.

To define a hypergraph, we will often use the notation $\mathcal{H} = (\mathcal{V}, \mathcal{E})$, where \mathcal{V} is the set of vertices, represented by an arbitrary finite set, and $\mathcal{E} \subseteq 2^\mathcal{V}$ is a set of hyperedges. To denote a single vertex and a single hyperedge of \mathcal{H} , we will use the symbols $v \in \mathcal{V}$ and $\mathcal{E} \in \mathcal{E}$. Additionally, we denote $\mathbf{n} = |\mathcal{V}|$, $\mathbf{m} = |\mathcal{E}|$, $\mathbf{d} = \max_{v \in \mathcal{V}} \deg(v)$, and $\mathbf{r} = \max_{\mathcal{E} \in \mathcal{E}} |\mathcal{E}|$. In other words, the symbols \mathbf{n} , \mathbf{m} , \mathbf{d} , and \mathbf{r} denote the number of vertices, the number of hyperedges, the maximal vertex degree, and the maximal edge cardinality, respectively. We use this notation to avoid ambiguity with the notation n , m , and d from the subsections, considering ILP problems.

In some problem formulations, we need to deal with hypergraphs $\mathcal{H} = (\mathcal{V}, \mathcal{E})$ having multiple hyperedges. That is, \mathcal{E} is a multi-set of sets $\mathcal{E} \in 2^\mathcal{V}$. In this case, by $\deg(v)$ we denote the number of **unique** hyperedges that are incident to v , and \mathbf{d} denotes the maximal vertex degree with respect to **unique** hyperedges.

In our work, we consider two types of combinatorial multi-packing/multi-cover problems: *vertex-based* problems and *edge-based* problems. In vertex-based problems, we need to pack vertices into hyperedges or to cover the hyperedges by vertices. In edge-based problems, we need to pack hyperedges or to cover vertices by hyperedges. The word "multi" means that we can choose a multi-set of vertices or edges to satisfy cover constraints or to not violate packing constraints. Before we give formal definitions, we present a few examples. The *Stable Multi-set* problem, which was introduced by Koster and

Zymolka in [55] as a natural generalization of the standard *Stable Set* problem, is an example of a vertex-based multi-packing problem. Similarly, the *Vertex Multi-cover* problem, which is a natural generalization of the standard *Vertex Cover* problem, is an example of vertex-based multi-cover problem. Some properties of the Stable Multi-set problem polyhedron were investigated in [56, 57], which had given a way to construct effective branch & bound algorithms for this problem. We cannot find a reference to the paper that introduces the Vertex Multi-cover problem, but this problem can be interpreted as a blocking problem for the Stable Multi-set problem (introduction to the theory of blocking and anti-blocking can be found in [31, 32], see also [44, p. 225]).

The examples of edge-based problems are the *Set Multi-cover*, *Multi-set Multi-cover*, and *Hypergraph Multi-matching* problems. The Set Multi-cover problem is a natural generalization of the classic *Set Cover* problem, where we need to choose a multi-set of hyperedges to cover the vertices by a given number of times. In the Multi-Set Multi-Cover problem, the input hypergraph \mathcal{H} can have multiple hyperedges. This problem has received quite a lot of attention in the recent papers [16, 17, 36, 47, 48, 53]. An exact $O((c_{\max} + 1)^n \cdot m)$ arithmetic complexity algorithm for the Multi-Set Multi-Cover problem, parameterized by n and the maximum coverage constraint number c_{\max} , is given by Hua, Wang, Yu & Lau in [47, 48]. A double exponential $2^{2^{O(n \log n)}}$ poly(ϕ)-complexity FPT-algorithm, parameterized by n , is given in Bredereck et al. [16]. The last algorithm was improved to a $n^{O(n^2)} \cdot \text{poly}(\phi)$ -complexity algorithm by Knop, Koutecký & Mních in [53]. A polynomial-time approximate algorithm can be found in Gorgi et al. [36]. The *Hypergraph Multimatching* problem is a very natural generalization of the *Hypergraph Matching* problem (see, for example, [2, 52]), which in turn is a generalization of the standard *Maximal Matching* problem in simple graphs. We cannot find a reference to the paper that formally introduces the Hypergraph Multi-matching problem, but, again, this problem can be interpreted as a blocking problem for the Multi-set Multi-cover problem.

The papers [17, 33] give good surveys and contain new ideas to use the ILP theory in combinatorial optimization setting.

Now, let us give some formal definitions. The vertex-based multi-packing/multi-cover problems can easily be modeled, using the following template problem:

Problem 5 (Hypergraph Vertex-Based Multi-Packing/Multi-Cover)

Let $\mathcal{H} = (\mathcal{V}, \mathcal{E})$ be a hypergraph. Given numbers $c_{\mathcal{E}}, p_{\mathcal{E}} \in \mathbb{Z}_{\geq 0}$, for $\mathcal{E} \in \mathcal{E}$, compute a multi-subset of \mathcal{V} , represented by natural numbers x_v , for $v \in \mathcal{V}$, such that

- (i) $c_{\mathcal{E}} \leq x(\mathcal{E}) \leq p_{\mathcal{E}}$, for any $\mathcal{E} \in \mathcal{E}$;
- (ii) $x(\mathcal{V}) \rightarrow \max / \min$.

Here, $x(\mathcal{M}) = \sum_{v \in \mathcal{M}} x_v$, for any $\mathcal{M} \subseteq \mathcal{V}$.

In other words, we need to solve the following ILP:

$$\begin{aligned} & \mathbf{1}^\top x \rightarrow \max / \min \\ & \begin{cases} c \leq A(\mathcal{H})^\top x \leq p \\ x \in \mathbb{Z}_{\geq 0}^{\mathcal{V}} \end{cases} \end{aligned} \quad (\text{Vertex-Based-ILP})$$

where $A(\mathcal{H})$ denotes the vertex-hyperedge incidence matrix of \mathcal{H} , and the vectors c and p are composed of the values $p_{\mathcal{E}}$ and $c_{\mathcal{E}}$, respectively.

It is natural to think that \mathcal{H} does not contain multiple hyperedges, because the multiple edge-constraints can easily be replaced by a stronger one.

If $c_{\mathcal{E}} = -\infty$, for all $\mathcal{E} \in \mathcal{E}$, and $x(\mathcal{V})$ is maximized, it can be considered as the *Stable Multi-set Problem on Hypergraphs*, when we need to find a multi-set of vertices of the maximum size, such that each hyperedge $\mathcal{E} \in \mathcal{E}$ is triggered at most $p_{\mathcal{E}}$ times.

Similarly, if $p_{\mathcal{E}} = +\infty$, for all $\mathcal{E} \in \mathcal{E}$, and $x(\mathcal{V})$ is minimized, it can be considered as the *Vertex Multi-cover Problem on Hypergraphs*, when we need to find a multi-set of vertices of the minimum size, such that each hyperedge $\mathcal{E} \in \mathcal{E}$ is triggered at least $c_{\mathcal{E}}$ times.

For the case, when \mathcal{H} is a simple graph, these problems can be considered as very natural generalizations of the classical Stable Set and Vertex Cover problems. Following to [55], the first one is called the *Stable Multi-set Problem*. As it was previously discussed, it is natural to call the second problem as the *Vertex Multi-cover Problem*.

Remark 2 Given numbers $u_v \in \mathbb{Z}_{\geq 0}$, for $v \in \mathcal{V}$, we can add an additional constraints $x_v \leq u_v$ to any of the problems above. We call such a problem as a *problem with multiplicities*.

Similarly, given $w_v \in \mathbb{Z}$, for $v \in \mathcal{V}$, we can consider the objective function $\sum_{v \in \mathcal{V}} w_v x_v$ instead of $x(\mathcal{V}) = \sum_{v \in \mathcal{V}} x_v$. We call such a problem as a *weighted problem*. The maximal weight is denoted by $w_{\max} = \max_{v \in \mathcal{V}} |w_v|$.

Similarly, the edge-based multi-packing/multi-cover problems can be modeled using the following template problem:

Problem 6 (Hypergraph Edge-Based Multi-packing/Multi-cover) Let $\mathcal{H} = (\mathcal{V}, \mathcal{E})$ be a hypergraph. Given numbers $c_v, p_v \in \mathbb{Z}_{\geq 0}$, for $v \in \mathcal{V}$, compute a multi-subset of \mathcal{E} , represented by the natural numbers $x_{\mathcal{E}}$, for $\mathcal{E} \in \mathcal{E}$, such that

- (i) $c_v \leq x(\delta(v)) \leq p_v$, for any $v \in \mathcal{V}$;
- (ii) $x(\mathcal{E}) \rightarrow \max / \min$.

Here, $x(\mathcal{M}) = \sum_{\mathcal{E} \in \mathcal{M}} x_{\mathcal{E}}$, for any $\mathcal{M} \subseteq \mathcal{E}$, and $\delta(v) = \{\mathcal{E} \in \mathcal{E} : v \in \mathcal{E}\}$ denotes the set of hyperedges that are incident to the vertex v .

The problem can be represented by the following ILP:

$$\begin{aligned} & \mathbf{1}^\top x \rightarrow \max / \min \\ & \begin{cases} c \leq A(\mathcal{H})x \leq p \\ x \in \mathbb{Z}_{\geq 0}^{\mathcal{E}} \end{cases} \end{aligned} \quad (\text{Edge-Based-ILP})$$

where the vectors c, p are composed of the values c_v and p_v .

Again, it is natural to think that \mathcal{H} does not contain multiple hyperedges, because the multiple edge-variables can be easily glued to one variable.

If $c_v = -\infty$, for all $v \in \mathcal{V}$, and $x(\mathcal{E})$ is maximized, it can be considered as the *Hypergraph Multi-matching* problem, when we need to find a multi-set of hyperedges of the maximum size, such that each vertex $v \in \mathcal{V}$ is triggered at most p_v times.

Similarly, if $p_v = +\infty$, for all $v \in \mathcal{V}$, and $x(\mathcal{E})$ is minimized, it can be considered as the *Set Multi-cover* problem, when we need to find a multi-set of hyper-edges of the minimum size, such that each vertex $v \in \mathcal{V}$ is triggered at least c_v times.

For the case, when \mathcal{H} is a simple graph, these problems can be considered as very natural generalizations of the classical Matching and Edge Cover problems. It seems natural to call these problems as the *Maximal Multi-matching* and *Edge Multi-cover* problems. The definition of the *Edge Multi-cover* problem can be found, for example, in the work [20], due to Cohen and Nutov. For the *Maximal Multi-matching* problem, we did not find a correct reference.

In a similar way, we can introduce the *Dominating Multi-set Problem* on simple graphs, which is a natural generalization of the classical *Dominating Set* problem. In this problem, we need to find a multi-set of vertices of the minimal size, such that all the vertices of a given graph will be covered given number of times by neighbors of the constructed vertex multi-set. The Dominating Multi-set Problem can be straightforwardly reduced to the Set Multi-cover Problem. To do that, we just need to construct the set system $\mathcal{H} = (\mathcal{V}, \mathcal{E})$, where \mathcal{V} coincides with the set of vertices of a given graph, and \mathcal{E} is constituted by neighbors of its vertices.

Remark 3 By analogy with Remark 2, we can introduce the *weighted* variants and variants *with multiplicities* for the all edge-based multi-packing/multi-cover problems discussed above.

Note that for the weighted Set Multi-cover and Hypergraph Multi-matching problems with multiplicities, it is essential to think that the hypergraph \mathcal{H} contains multiple hyperedges, since two equivalent hyperedges can correspond to different weights.

Problem 7 The weighted Set Multi-cover with multiplicities is known in literature as the *Weighted Multi-Set multi-Cover* problem, see, for example, [47, 48, 53].

Let us explain our motivation with respect to the specified combinatorial problems. The classical Stable Set and Vertex Cover Problems on graphs and hypergraphs admit trivial $2^{O(n)} \cdot \text{poly}(\phi)$ -complexity algorithms. However, the Stable Multi-set and Vertex Multi-cover Problems does not admit such a trivial algorithm. But, both problems can be modeled as the ILP problem (Vertex-Based-ILP) with n variables. Consequently, both problems can be solved by the previously mentioned $O(n)^n \cdot \text{poly}(\phi)$ -complexity general ILP algorithm. Here $\phi = \text{size}(c, p, w, u)$.

Is it possible to give a faster algorithm? Is it possible to give a positive answer to this question, considering a more complex variant with multiplicities? We show that these problems on hypergraphs can be solved by a $\min\{\mathfrak{d}, \mathfrak{r}\}^{O(n)} \cdot \text{poly}(\phi)$ -complexity algorithm. Consequently, the Stable Multi-set and Vertex Multi-cover Problems on simple graphs can be solved by $2^{O(n)} \cdot \text{poly}(\phi)$ -complexity algorithms. Our complexity results for these problems, together with the Multi-set Multi-cover, Hypergraph Multi-matching, and Dominating Multi-set problems, are gathered in Theorem 8.

Theorem 8 *Let us consider the Opt-And-Count-IP-variants of the problems Stable Multi-set, Vertex Multi-cover, Set Multi-cover, Hypergraph Multi-matching, and Dominating Multi-set with multiplicities. Note that the Set Multi-cover problem with multiplicities is also known as the Multi-set Multi-cover problem.*

The following complexity bounds hold:

<i>Problems:</i>	<i>Time :</i>
<i>Stable Multi-set and Vertex Multi-cover on hypergraps</i>	$\min\{\mathfrak{d}, \mathfrak{r}\}^{3.5n} \cdot 4^n$
<i>Stable Multi-set and Vertex Multi-cover on simple graphs</i>	2^{5n}
<i>Dominating Multi-set</i>	$\mathfrak{d}^{3.5n} \cdot 4^n$
<i>Set Multi-cover and Hypergraph Multi-matching</i>	$\min\{\mathfrak{d}, \mathfrak{r}\}^{3.5m} \cdot 4^m$

The complexity bounds for the weighted variants of the considered problems contain additional multiplicative term $(\|c\|_\infty)^3$. Everywhere in the complexity bounds, we skip the $\text{poly}(\phi)$ multiplicative term.

Proof To prove the theorem, we use Theorem 3 for the problems' definitions 5 and 6, equipped by Remarks 2 and 3. This approach gives us the desired complexity bounds for all the problems, except for the Stable Multiset and Vertex Multicover problems on simple graphs. For these exceptions, we will give a more accurate analysis.

We follow to the proof of Theorem 3, using a more accurate bound for $\Delta_{n-1}(M)$. Let $A := A(\mathcal{G})$ be the incidence matrix of the corresponding simple graph \mathcal{G} . Due to [43], it is known that $\Delta_i(A) \leq 2^{\tau_0}$, where $\tau_0 = \tau_0(\mathcal{G})$ is the odd tulgeity of \mathcal{G} , which is defined as the maximum number of vertex-disjoint odd cycles of \mathcal{G} . Clearly, $\tau_0 \leq n/3$, so, $\Delta_{n-1}(M) \leq 2^{n/3}$. Using this bound for $\Delta_{n-1}(A)$ in the proof of Theorem 3, it gives the desired complexity bounds for the Stable Multiset and Vertex Multicover Problems. \square

3.1 The Multi-set Multi-cover and Hypergraph Multi-matching Problems Parameterized by the number of vertices n .

In Theorem 8, we have presented $\min\{\mathfrak{d}, \mathfrak{r}\}^{3.5m} \cdot 4^m \cdot \text{poly}(\phi)$ -complexity algorithms for the Opt-And-Count-IP-variant of the Set Multi-cover and Hypergraph Multi-matching problems with multiplicities. Due to Knop, Koutecký & Mnich [53], the weighted Opt-IP-variants of these problems admit an $n^{O(n^2)}$ $\cdot \text{poly}(\phi)$ -complexity algorithm, which is faster than our algorithm for $m =$

$\Omega(n^{2+\varepsilon})$ and any $\varepsilon > 0$. In other words, our last complexity bound is good only for sufficiently sparse hypergraphs.

So, the motivation of this subsection is to present faster algorithms for the Opt-And-Count-IP- and Opt-IP-variants of the weighted Multi-set Multi-cover and Hypergraph Multi-matching problems with and without multiplicities, parameterized by n instead of m . Our results for these problems are gathered in Table 3.

Table 3: New complexity bounds for the Set Multi-cover and Hypergraph Multi-matching problems

Version: ²	Time : ¹
Opt-IP, without multipl.	$O(\tau)^{2n}$ $O(\tau)^n \cdot 2^n \cdot \log(\tau \log(n))$ $O(\mathfrak{d})^n \cdot 2^n \cdot \log(\log(\mathfrak{d} n) \log(n))$ $O(n)^n$
Opt-And-Count-IP, without multipl.	$\min\{\tau, \mathfrak{d}\}^{1.5n} \cdot O(m)^n \cdot w_{\max}^3$ $\tau^{1.5n} \cdot O(n)^{(\tau+1)n} + O(\tau) \cdot w_{\max}^3$ $\mathfrak{d}^{2.5n} \cdot O(n)^n \cdot w_{\max}^3$ $2^{n^2+o(n^2)} \cdot w_{\max}^3$
Opt-IP, with multipl.	$O(\min\{\mathfrak{d}, \tau\})^{n^2} + O(n \log n)$
Opt-And-Count-IP, with multipl.	open problem

¹ The multiplicative factor $\text{poly}(\phi)$ is skipped.

² Only the variants without multiplicities are considered.

Remark 4 Let us have a little discussion about the complexity bounds, presented in Table 3. Firstly, let us consider the problems without multiplicities. As the reader can see, for fixed τ , the weighted Opt-IP-variant of the considered problems can be solved by $2^{O(n)}$ -complexity algorithms (the $\text{poly}(\phi)$ -term is ignored). For $\tau = \log(n)^{O(1)}$, the best complexity bound is $2^{O(n \cdot \log \log(n))}$.

Another interesting case is $\mathfrak{d} = o(n)$, which gives the $o(n)^n \cdot 2^{O(n \cdot \log \log(n))}$ -complexity bound. For other values of parameters, the general $O(n)^n$ -complexity bound holds.

For the unweighted Opt-And-Count-IP-variant of the considered problems, if τ is fixed, then the $n^{O(n)}$ -complexity algorithm exists. The same is true if $\mathfrak{d} = n^{O(1)}$ or $m = n^{O(1)}$. For the general values of τ , \mathfrak{d} , and m , it is better to use the complexity bound $\min\{\mathfrak{d}, \tau\}^{1.5n} \cdot O(m)^n$. Since $m \leq 2^n$, it straightforwardly gives the general $2^{n^2+o(n^2)}$ -complexity bound.

Note that the considered complexity bounds for the problems without multiplicities are sufficiently outperform the best complexity bound that we know $n^{O(n^2)}$, due to Knop, Kouřtecký, and Mních [53].

Now, let us consider the weighted problems with multiplicities. Note again that the weighted Set Multi-cover problem with multiplicities is also known as the weighted Multi-set Multi-cover problem. In comparison with the state of the art complexity bound $n^{O(n^2)}$, our bound $O(\min\{\mathfrak{d}, \tau\})^{n^2} + O(n \log n)$ has a

lower exponent base and it gives a constant-estimate in the exponent power. Unfortunately, we are not able to present a complexity bound, parameterized by \mathbf{n} , for the Opt-And-Count-IP-variant, and it seems to be an interesting open problem.

We omit proofs of the results, presented in Table 3, because they straightforwardly follow from the complexity bounds, described in Theorem 6 and Table 2. Indeed, the weighted Multi-set Multi-cover and Hypergraph Multi-matching problems with or without multiplicities can be represented by the following ILP's in the standard form:

$$\begin{array}{ll} w^\top x \rightarrow \max & w^\top x \rightarrow \min \\ \left\{ \begin{array}{l} (A(\mathcal{H}) I_{\mathbf{n} \times \mathbf{n}})x = p \\ \mathbf{0} \leq x \leq u \\ x \in \mathbb{Z}^{\mathbf{n} + \mathbf{m}} \end{array} \right. & \left\{ \begin{array}{l} (-A(\mathcal{H}) I_{\mathbf{n} \times \mathbf{n}})x = -c \\ \mathbf{0} \leq x \leq u \\ x \in \mathbb{Z}^{\mathbf{n} + \mathbf{m}}, \end{array} \right. \end{array}$$

where the constraint $x \leq u$ need to be omitted for the variants without multiplicities. The co-dimension of these formulations is \mathbf{n} . Using simple quantities $\mathbf{m} \leq 2^{\mathbf{n}}$, $\mathbf{m} \leq \mathfrak{d} \mathbf{n}$, and $\mathbf{m} = O(\mathbf{n})^{\mathfrak{r}+1}$ that are valid for the problems without multiplicities, the desired complexity bounds of Table 3 can be easily obtained. Note that the equality $\mathbf{m} = O(\mathbf{n})^{\mathfrak{r}+1}$ directly follows from the inequality $\mathbf{m} \leq \sum_{i=1}^{\mathfrak{r}} \binom{\mathbf{n}}{i}$.

4 Additional Notes: Expected ILP Complexity

It was shown by Oertel, Paat & Weismantel in [69] that, for almost all r.h.s. $b \in \mathbb{Z}^m$, the original ILP problem in the form Canon-Form is equivalent to the problem $\max\{c^\top x: A_{\mathcal{B}}x \leq b_{\mathcal{B}}, x \in \mathbb{Z}^n\}$, where $A_{\mathcal{B}}$ is a non-degenerate $n \times n$ sub-matrix of A , induced by some optimal LP base \mathcal{B} . It was noted by Shevchenko in [73, Paragraph 3.3., p. 42–43] (see also [41, Chapter 5.2]) that such a square ILP problem is equivalent to the group minimization problem, described by R. Gomory in the seminal work [35] (see also [46, Chapter 19]). Consequently, due to [35] and [46, Chapter 19], such an ILP can be solved by an algorithm with the arithmetic complexity bound

$$O(\min\{n, \Delta\} \cdot \Delta \cdot \log(\Delta)), \quad (34)$$

where $\Delta := \Delta(A)$. The result of Oertel, Paat & Weismantel [69] was refined by Griбанov et al. in [41, Chapter 5.5.1], where a stronger probability argument was given.

A similar result for the problems in the form Standard-Form is given in the paper of Oertel, Paat & Weismantel [68]. Another way is to reduce the problem in the form Standard-Form to the problem in the form Canon-Form, using [41, Lemma 5]. It follows from [35] and [68] (or from [41, Lemma 5]) and result for the form Canon-Form) that, for almost all r.h.s. $b \in \mathbb{Z}^k$, the ILP

problem in the form Standard-Form of co-dimension k can be solved by an algorithm with the arithmetic complexity bound

$$O((n - k) \cdot \Delta \cdot \log(\Delta)). \quad (35)$$

It is also easy to see that this fact also holds for problems with multiplicities, the simplest way is to reduce the problem into the form Canon-Form.

The bounds (34) and (35), together with the inequalities (3) and (2), give the following complexity bounds, described in Table 4, for the sparse problem Opt-IP in the canonical and the standard forms, respectively.

Table 4: Expected complexity bounds for almost all b for the problem Opt-IP in the standard and the canonical forms

Problems: ¹	Time : ¹
The form Canon-Form, for almost all $b \in \mathbb{Z}^m$	$\gamma_{1,\infty}(A)^n$ $(\ A\ _{\max})^n \cdot \overline{\text{ts}}(A)^{n/2}$
The form Standard-Form, for almost all $b \in \mathbb{Z}^k$	$\gamma_{1,\infty}(A)^k$ $(\ A\ _{\max})^k \cdot \overline{\text{ts}}(A)^{k/2}$

¹ The multiplicative factor $\text{poly}(\phi)$ is skipped.

These bounds can be used to give expected-case complexity bounds for the combinatorial problems, described in Table 5.

Table 5: Expected complexity bounds for combinatorial packing/cover problems with multiplicities, for almost all r.h.s. p or c

Problems: ¹	Time : ²
Stable Multi-set on hypergraps and Hypergraph Multi-matching, for almost all r.h.s. p	$\min\{\mathfrak{d}, \mathfrak{r}\}^{n/2}$
Vertex Multi-cover on hypergraps and Multi-set Multi-cover, for almost all r.h.s. c	$\min\{\mathfrak{d}, \mathfrak{r}\}^{n/2}$
Dominating Multi-set, for almost all r.h.s. c	$\mathfrak{d}^{n/2}$
Stable Multi-set and Vertex Multi-cover on simple graphs, for almost all r.h.s. p, c resp. ³	$2^{n/3}$

¹ All the considered problems are weighted problems with multiplicities.

² The multiplicative factor $\text{poly}(\phi)$ is skipped.

³ The bound $2^{n/2}$ is trivial, to achieve the bound $2^{n/3}$, see the proof of Theorem 8.

5 Conclusion

Here we give a summary of results, notes, and implications of our work.

- We give new algorithms for the problems Feasibility-IP, Count-IP, Opt-IP, and Opt-And-Count-IP. Our algorithms outperform the general state of the art $O(n)^n \cdot \text{poly}(\phi)$ -complexity algorithm, due to Dadush, Peikert & Vempala [23, 25] for sparse instances with bounded elements. Details can be found in Table 1 and Theorem 3.
- We show that in the assumptions $\|A\|_{\max} = n^{O(1)}$ and $\|c\|_{\infty} = n^{O(n)}$, the problems Count-IP and Opt-And-Count-IP can be solved by algorithms with the complexity bound $n^{O(n)} \cdot \text{poly}(\phi)$, which outperforms the state of the art bound (5). For details, see Corollary 1.
- We give new algorithms for the Opt-And-Count-IP-variant of the Stable Multi-set, Vertex Multi-cover, Set Multi-cover, Multi-matching, and Dominating Multi-set problems with respect to simple graphs and hypergraphs, see the definitions 5 and 6. The weighted variants and the variants with the multiplicities of the above problems are handled, see Remarks 2 and 3. Note that the weighted Set Multi-cover problem with multiplicities is also known as the weighted Multi-set Multi-cover problem. Our algorithms outperform the general state of the art $O(n)^n \cdot \text{poly}(\phi)$ -complexity algorithm, applied to these problems. Details can be found in Theorem 8.
- We summarise known results and new methods to give new algorithms for the Feasibility-IP-, Count-IP-, Opt-IP-, Opt-And-Count-IP-variants of ILP problems in the standard form with and without multiplicities, parameterised by $\|A\|_{\max}$ and the co-dimension of $Ax = b$. The new complexity bounds outperform general-case bounds on sparse instances. Details can be found in Subsection 2.1, Table 2, and Theorem 6.
- Using our notes for sparse problems in the standard form, we give new algorithms for the Opt-IP- and Opt-And-Count-IP-variants of the Set Multi-cover and Hypergraph Multi-matching problems with and without multiplicities, parameterized by the number of vertices n . The weighted variants are handled. Tighter complexity bounds with respect to the parameters n , m , r , and d are considered. The most general complexity bounds for the problems without multiplicities are: $O(n)^n \cdot \text{poly}(\phi)$, for the Opt-IP-variant, and $2^{n^2 + o(n^2)} \cdot \text{poly}(\phi)$, for the Opt-And-Count-IP-variant. For the Opt-IP-variant with multiplicities, we give the $O(\min\{d, r\})^{n^2 + O(n \log n)} \cdot \text{poly}(\phi)$ -complexity bound. Unfortunately, we are not able to present a complexity bound, parameterized by n , for the Opt-And-Count-IP-variant with multiplicities, it seems to be an interesting open problem. Our complexity bounds for the considered problems outperform the state of the art $n^{O(n^2)} \cdot \text{poly}(\phi)$ -complexity bound, due to Knop, Kouřtecký, and Mních [53]. Details can be found in Subsection 3.1 and Table 3. Discussion can be found in Remark 4.
- We note that expected versions of the considered sparse problems admit much more efficient algorithms. Saying "expected", we mean that the related complexity bounds are valid for almost all r.h.s. For the details, see Subsection 4 and Tables 4, 5.

Open problems:

- As it was noted before, we are not able to present an algorithm for the problem Count-IP in the form Standard-Form with multiplicities, which will be polynomial on n , Δ or $\|A\|_{\max}$, for any fixed co-dimension k . More precisely, given $A \in \mathbb{Z}^{k \times n}$, $b \in \mathbb{Q}^k$, and $u \in \mathbb{Z}^n$, let us consider the polytope \mathcal{P} , defined by the following system

$$\begin{cases} Ax = b \\ 0 \leq x \leq u. \end{cases}$$

The problem is to develop an algorithm to compute $|\mathcal{P} \cap \mathbb{Z}^n|$, whose complexity will be polynomial on n , $\Delta(A)$ or $\|A\|_{\max}$, for any fixed k . Despite considerable effort, we are not able to present such an algorithm. The main difficulty is that our methods work well only in the scenarios, when the value of $|\text{vert}(\mathcal{P})|$ is sufficiently small. But, in the current case, the value of $|\text{vert}(\mathcal{P})|$ can be equal to 2^n . Note that the positive solution for this problem can grant new more efficient algorithms for the Multi-set Multi-cover problem and its weighted variant.

- Our general complexity bounds (see Theorems 3 and 6) for sparse variants of the problem Count-IP contain a term of the type $(\|A\|_{\max})^{O(n)}$ or of the type $(\|A\|_{\max})^{O(k)}$. Could we develop an algorithm, which will be polynomial on $\|A\|_{\max}$ and more efficient for sparse problems with respect to the general state of the art algorithms? Could we do this for the simpler problem Feasibility-IP?
- Our complexity bounds for sparse problems depend mainly on the total number of variables n , which can be significantly bigger than an actual dimension $d = \dim(\mathcal{P})$ of a polyhedron. The known state of the art algorithms can be easily adapted to work with the parameter d instead of n . For example, the state of the art algorithm, due to Dadush, Peikert & Vempala [23, 25], gives the $O(d)^d \cdot \text{poly}(\phi)$ complexity bound. Unfortunately, at the current moment, we cannot adapt our methods for sparse problems to work with the parameter d . The difficulty is concentrated in Lemma 3, which estimates the number of vertices of a polyhedron. The proof of such a lemma, based on a parameter d , is an interesting open question, which will guaranty the existence of an algorithm for sparse problems, parameterized by d instead of n .

Acknowledgement

Results of the Section 2 were prepared within the framework of the Basic Research Program at the National Research University Higher School of Economics (HSE). Results of the Subsection 2.1 and Sections 3, 4 were prepared under financial support of Russian Science Foundation grant No 21-11-00194.

References

1. Alekseev, V., E., Zakharova, D., V.: Independent sets in the graphs with bounded minors of the extended incidence matrix. *Journal of Applied and Industrial Mathematics* **5**(1), 14–18. doi:10.1134/S1990478911010029
2. Alon, N., Yuster, R.: On a hypergraph matching problem. *Graphs and Combinatorics* **21**(4), 377–384 (2005)
3. Artmann, S., Weismantel, R., Zenklusen, R.: A strongly polynomial algorithm for bi-modular integer linear programming. In: *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*. pp. 1206–1219. STOC 2017, Association for Computing Machinery, New York, NY, USA (2017). doi:10.1145/3055399.3055473
4. Averkov, G., Schymura, M.: On the maximal number of columns of δ -modular matrix. In: *International Conference on Integer Programming and Combinatorial Optimization*. pp. 29–42. Springer (2022)
5. Avis, D., Fukuda, K.: A pivoting algorithm for convex hulls and vertex enumeration of arrangements and polyhedra. *Discrete & Computational Geometry* **8**(3), 295–313 (1992). doi:10.1007/BF02293050, <https://doi.org/10.1007/BF02293050>
6. Banaszczyk, W.: Balancing vectors and gaussian measures of n-dimensional convex bodies. *Random Structures & Algorithms* **12**(4), 351–360 (1998)
7. Barvinok, A.: *Integer Points in Polyhedra*. European Mathematical Society, ETH-Zentrum, Zürich, Switzerland (2008)
8. Barvinok, A., Pommersheim, J.: An algorithmic theory of lattice points in polyhedra. *New Perspect. Algebraic Combin.* **38** (1999)
9. Barvinok, A.: A polynomial time algorithm for counting integral points in polyhedra when the dimension is fixed. In: *Proceedings of 1993 IEEE 34th Annual Foundations of Computer Science*. pp. 566–572 (1993). doi:10.1109/SFCS.1993.366830
10. Barvinok, A., Woods, K.: Short rational generating functions for lattice point problems. *Journal of the American Mathematical Society* **16**(4), 957–979 (2003), <http://www.jstor.org/stable/30041461>
11. Basu, A., Oertel, T.: Centerpoints: a link between optimization and convex geometry. *SIAM Journal on Optimization* **27**(2), 866–889 (2017). doi:10.1137/16M1092908
12. Beck, J., Fiala, T.: “integer-making” theorems. *Discrete Applied Mathematics* **3**(1), 1–8 (1981)
13. Beck, M., Robins, S.: *Computing the continuous discretely*. Springer, New York (2015)
14. Bellman, R.: Dynamic programming. *Science* **153**(3731), 34–37 (1966)
15. Bonifas, N., Di Summa, M., Eisenbrand, F., Hähnle, N., Niemeier, M.: On sub-determinants and the diameter of polyhedra. *Discrete & Computational Geometry* **52**, 102–115 (2014). doi:10.1007/s00454-014-9601-x, <https://doi.org/10.1007/s00454-014-9601-x>
16. Bredereck, R., Faliszewski, P., Niedermeier, R., Skowron, P., Talmon, N.: Elections with few candidates: Prices, weights, and covering problems. In: *International Conference on Algorithmic Decision Theory*. pp. 414–431. Springer (2015). doi:10.1007/978-3-319-23114-3_25
17. Bredereck, R., Faliszewski, P., Niedermeier, R., Skowron, P., Talmon, N.: Mixed integer programming with convex/concave constraints: Fixed-parameter tractability and applications to multicovering and voting. *Theoretical Computer Science* **814**, 86–105 (2020). doi:10.1016/j.tcs.2020.01.017
18. Brion, M.: Points entiers dans les polyèdres convexes. *Annales scientifiques de l’École Normale Supérieure 4e série*, **21**(4), 653–663 (1988). doi:10.24033/asens.1572, <http://www.numdam.org/articles/10.24033/asens.1572/>
19. Chirkov, A., Y., Griбанov, D., V., Malyshev, D., S., Pardalos, P., M., Veselov, S., I., Zolotykh, N., Y.: On the complexity of quasiconvex integer minimization problem. *Journal of Global Optimization* **73**(4), 761–788 (2019). doi:10.1007/s10898-018-0729-8
20. Cohen, N., Nutov, Z.: Approximating minimum power edge-multi-covers. *Journal of Combinatorial Optimization* **30**(3), 563–578 (2015)
21. Cook, W., Gerards, A.M.H., Schrijver, A., Tardos, E.: Sensitivity theorems in integer linear programming. *Mathematical Programming* **34**(3), 251–261 (1986). doi:10.1007/BF01582230

22. Cunningham, W.H., Geelen, J.: On integer programming and the branch-width of the constraint matrix. In: International Conference on Integer Programming and Combinatorial Optimization. pp. 158–166. Springer (2007)
23. Dadush, D.: Integer programming, lattice algorithms, and deterministic volume estimation. Georgia Institute of Technology, ProQuest Dissertations Publishing, Ann Arbor (2012)
24. Dadush, D.: On approximating the covering radius and finding dense lattice subspaces. In: Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing. pp. 1021–1026 (2019). doi:10.1145/3313276.3316397
25. Dadush, D., Peikert, C., Vempala, S.: Enumerative lattice algorithms in any norm via m-ellipsoid coverings. In: 2011 IEEE 52nd Annual Symposium on Foundations of Computer Science. pp. 580–589 (2011). doi:10.1109/FOCS.2011.31
26. De Loera, Jesús, A., Hemmecke, R., Köppe, M.: Algebraic And geometric ideas in the theory of discrete optimization. Society for Industrial and Applied Mathematics, Philadelphia, USA (2013)
27. Dyer, M., Kannan, R.: On barvinok’s algorithm for counting lattice points in fixed dimension. *Mathematics of Operations Research* **22**(3), 545–549 (1997). doi:10.1287/moor.22.3.545
28. Eisenbrand, F., Weismantel, R.: Proximity Results and Faster Algorithms for Integer Programming Using the Steinitz Lemma. *ACM Transactions on Algorithms* **16**(1) (Nov 2019). doi:10.1145/3340322, <https://doi.org/10.1145/3340322>
29. Fiorini, S., Joret, G., Weltge, S., Yuditsky, Y.: Integer programs with bounded subdeterminants and two nonzeros per row (2021), <https://arxiv.org/abs/2106.05947>
30. Fomin, F.V., Panolan, F., Ramanujan, M., Saurabh, S.: On the optimality of pseudo-polynomial algorithms for integer programming. *Mathematical Programming* pp. 1–33 (2022)
31. Fulkerson, D.R.: Blocking and anti-blocking pairs of polyhedra. *Mathematical programming* **1**(1), 168–194 (1971). doi:10.1007/BF01584085
32. Fulkerson, D.R.: Anti-blocking polyhedra. *Journal of Combinatorial Theory, Series B* **12**(1), 50–71 (1972). doi:10.1016/0095-8956(72)90032-9
33. Gavenčiak, T., Koutecký, M., Knop, D.: Integer programming in parameterized complexity: Five miniatures. *Discrete Optimization* (2020). doi:10.1016/j.disopt.2020.100596
34. Glanzer, C., Stallknecht, I., Weismantel, R.: Notes on $\{a, b, c\}$ -modular matrices (2021), <https://arxiv.org/abs/2106.14980>
35. Gomory, R.E.: On the relation between integer and noninteger solutions to linear programs. *Proceedings of the National Academy of Sciences* **53**(2), 260–265 (1965). doi:10.1073/pnas.53.2.260, <https://www.pnas.org/content/53/2/260>
36. Gorgi, A., El Ouali, M., Srivastav, A., Hachimi, M.: Approximation algorithm for the multicovering problem. *Journal of Combinatorial Optimization* **41**(2), 433–450 (2021). doi:10.1007/s10878-020-00688-9
37. Griбанov, D., Shumilov, I., Malyshev, D.: Structured (min, +)-convolution and its applications for the shortest vector, closest vector, and separable nonlinear knapsack problems (2022)
38. Griбанov, D.V., Malyshev, D.S.: Integer conic function minimization based on the comparison oracle. In: Khachay, M., Kochetov, Y., Pardalos, P. (eds.) *Mathematical Optimization Theory and Operations Research*. pp. 218–231. Springer International Publishing, Cham (2019). doi:10.1007/978-3-030-22629-9_16
39. Griбанov, D., V.: An FPTAS for the Δ -Modular Multidimensional Knapsack Problem (2021). doi:https://doi.org/10.1007/978-3-030-77876-7_6
40. Griбанov, D., V., Malyshev, D., S.: Faster algorithm for counting of the integer points number in δ -modular polyhedra (2021), <https://arxiv.org/abs/2110.01732?context=cs>
41. Griбанov, D., V., Shumilov, I., A., Malyshev, D., S., Pardalos, P., M.: On δ -modular integer linear problems in the canonical form and equivalent problems (2021), <https://arxiv.org/abs/2002.01307v5>
42. Griбанov, Dmitry, V., Zolotykh, N., Y.: On lattice point counting in δ -modular polyhedra. *Optimization Letters* pp. 1–28 (2021). doi:10.1007/s11590-021-01744-x

43. Grossman, J.W., Kulkarni, D.M., Schochetman, I.E.: On the minors of an incidence matrix and its smith normal form. *Linear Algebra and its Applications* **218**, 213–224 (1995). doi:10.1016/0024-3795(93)00173-W
44. Grötschel, M., Lovász, L., Schrijver, A.: *Geometric algorithms and combinatorial optimization*. Springer Science & Business Media (2012)
45. Grünbaum, B.: *Convex Polytopes*. Graduate Texts in Mathematics, Springer-Verlag, New York (2011)
46. Hu, T., C.: *Integer programming and network flows*. Addison-Wesley Publishing Company, London (1970)
47. Hua, Q.S., Wang, Y., Yu, D., Lau, F.C.: Dynamic programming based algorithms for set multicover and multiset multicover problems. *Theoretical Computer Science* **411**(26–28), 2467–2474 (2010). doi:10.1016/j.tcs.2010.02.016
48. Hua, Q.S., Yu, D., Lau, F.C., Wang, Y.: Exact algorithms for set multicover and multiset multicover problems. In: *International Symposium on Algorithms and Computation*. pp. 34–44. Springer (2009). doi:10.1007/978-3-642-10631-6_6
49. Jansen, K., Rohwedder, L.: On integer programming, discrepancy, and convolution (2018), <https://arxiv.org/abs/1803.04744>
50. Jansen, K., Rohwedder, L.: On integer programming and convolution. In: *10th Innovations in Theoretical Computer Science Conference (ITCS 2019)*. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik (2018)
51. Jiang, H., Reis, V.: A tighter relation between hereditary discrepancy and determinant lower bound. In: *Symposium on Simplicity in Algorithms (SOSA)*. pp. 308–313. SIAM (2022)
52. Keevash, P., Mycroft, R.: A geometric theory for hypergraph matching. *American Mathematical Soc.* (2014)
53. Knop, D., Koutecký, M., Mnich, M.: Combinatorial n-fold integer programming and applications. *Mathematical Programming* **184**(1), 1–34 (2020). doi:10.1007/s10107-019-01402-2
54. Köppe, M., Verdoolaege, S.: Computing parametric rational generating functions with a primal barvinok algorithm. *The electronic journal of combinatorics* **15** (2008). doi:10.37236/740, <https://doi.org/10.37236/740>
55. Koster, A.M., Zymolka, A.: Stable multi-sets. *Mathematical Methods of Operations Research* **56**(1), 45–65 (2002). doi:10.1007/s001860200199
56. Koster, A.M., Zymolka, A.: Polyhedral investigations on stable multi-sets. *Tech. Rep. 03–10*, ZIB, Takustr. 7, 14195 Berlin (2003)
57. Koster, A.M., Zymolka, A.: On cycles and the stable multi-set polytope. *Discrete Optimization* **2**(3), 241–255 (2005). doi:10.1016/j.disopt.2005.06.004
58. Kratsch, S.: On polynomial kernels for sparse integer linear programs. *Journal of Computer and System Sciences* **82**(5), 758–766 (2016)
59. Lasserre, Jean, B., Zeron, Eduardo, S.: Solving the knapsack problem via z-transform. *Operations Research Letters* **30**(6), 394–400 (2002)
60. Lasserre, J.B.: *Linear and integer programming vs linear integration and counting: a duality viewpoint*. Springer Science & Business Media, New York (2009)
61. Lawrence, J.: *Rational-function-valued valuations on polyhedra*. *Discrete and computational geometry* (New Brunswick, NJ, 1989/1990) **6**, 199–208 (1991)
62. Lee, J., Paat, J., Stallknecht, I., Xu, L.: Improving proximity bounds using sparsity. In: Baïou, M., Gendron, B., Günlük, O., Mahjoub, A.R. (eds.) *Combinatorial Optimization*. pp. 115–127. Springer International Publishing, Cham (2020). doi:https://doi.org/10.1007/978-3-030-53262-8_10
63. Lee, J., Paat, J., Stallknecht, I., Xu, L.: Polynomial upper bounds on the number of differing columns of an integer program. *arXiv preprint arXiv:2105.08160v2 [math.OC]* (2021), <https://arxiv.org/abs/2105.08160>
64. Lovász, L., Spencer, J., Vesztergombi, K.: Discrepancy of set-systems and matrices. *European Journal of Combinatorics* **7**(2), 151–160 (1986). doi:[https://doi.org/10.1016/S0195-6698\(86\)80041-5](https://doi.org/10.1016/S0195-6698(86)80041-5), <https://www.sciencedirect.com/science/article/pii/S0195669886800415>
65. Matoušek, J.: The determinant bound for discrepancy is almost tight. *Proceedings of the American Mathematical Society* **141**(2), 451–460 (2013)

66. McMullen, P.: The maximum numbers of faces of a convex polytope. *Mathematika* **17**(2), 179–184 (1970). doi:10.1112/S0025579300002850
67. Micciancio, D., Voulgaris, P.: A deterministic single exponential time algorithm for most lattice problems based on voronoi cell computations. *SIAM Journal on Computing* **42**(3), 1364–1391 (2013)
68. Oertel, T., Paat, J., Weismantel, R.: The distributions of functions related to parametric integer optimization. *SIAM Journal on Applied Algebra and Geometry* **4**(3), 422–440 (2020). doi:10.1137/19M1275954, <https://doi.org/10.1137/19M1275954>
69. Paat, J., Schlöter, M., Weismantel, R.: The integrality number of an integer program. *Mathematical Programming* pp. 1–21 (2021). doi:10.1007/s10107-021-01651-0, <https://doi.org/10.1007/s10107-021-01651-0>
70. Polak, A., Rohwedder, L., Węgrzycki, K.: Knapsack and subset sum with small items. arXiv:2105.04035v1 [cs.DS] (2021), <https://arxiv.org/abs/2105.04035>
71. Pukhlikov, A.V., Khovanskii, A.G.: The riemann–roch theorem for integrals and sums of quasipolynomials on virtual polytopes (russian). *Algebra i analiz* **4**(4), 188–216 (1992)
72. Schrijver, A.: *Theory of linear and integer programming*. John Wiley & Sons, Chichester (1998)
73. Shevchenko, V.N.: *Qualitative topics in integer linear programming*. American Mathematical Soc., Providence, Rhode Island (1996)
74. Spencer, J.: Six standard deviations suffice. *Transactions of the American mathematical society* **289**(2), 679–706 (1985). doi:10.1090/S0002-9947-1985-0784009-0, <https://doi.org/10.1090/S0002-9947-1985-0784009-0>
75. Storjohann, A.: Near optimal algorithms for computing Smith normal forms of integer matrices. In: *Proceedings of the 1996 International Symposium on Symbolic and Algebraic Computation*. pp. 267–274. ISSAC '96, Association for Computing Machinery, New York, NY, USA (1996). doi:10.1145/236869.237084
76. Veselov, S., Chirkov, A.: Integer program with bimodular matrix. *Discrete Optimization* **6**(2), 220–222 (2009). doi:<https://doi.org/10.1016/j.disopt.2008.12.002>, <https://www.sciencedirect.com/science/article/pii/S1572528608000881>
77. Veselov, S., Gribanov, D., Zolotykh, N., Chirkov, A.: A polynomial algorithm for minimizing discrete convex functions in fixed dimension. *Discrete Applied Mathematics* **283**, 11–19 (2020). doi:<https://doi.org/10.1016/j.dam.2019.10.006>, <https://www.sciencedirect.com/science/article/pii/S0166218X19304561>