arXiv:2201.09301v3 [math.CA] 2 Feb 2023

ON AN EXPANSION FORMULA OF LIU

BING HE

Abstract. In this paper, we extend an expansion formula of Liu to multiple basic hypergeometric series over the root system A_n. The usefulness of Liu's expansion formula in special functions and number theory has been shown by Liu and many others. We first establish a very general multiple expansion formula over the root system A_n and then deduce several A_n extensions of Liu's expansion formula. From these multiple formulas, we derive two groups of multiple expansion formulas for infinite products. As applications, we deduce an A_n Rogers' ${}_6\phi_5$ summation, an A_n extension of Sylvester's identity, some multiple expansion formulas for $(q)_\infty^m, \pi_q$ and $1/\pi_q$, two A_n extensions of the Rogers-Fine identity, an A_n extension of Liu's extension of Rogers' nonterminating ${}_6\phi_5$ summation, an A_n extension of a generalization of Fang's identity and an A_n extension of Andrews' expansion formula.

1. Introduction

In a battery of papers, Liu [15, 18, 19] extended many summation formulas and transformation formulas in the theory of basic hypergeometric series and proved the following well-known expansion formula [15, Theorem 1.1][18, Theorem 1.2]:

Theorem 1.1. *Suppose that $f(x)$ is an analytic function near $x = 0$, then, under suitable convergence conditions,*

$$\begin{aligned}&\frac{(\alpha q, \alpha ab/q)_\infty}{(\alpha a, \alpha b)_\infty} f(\alpha a)\\ &= \sum_{n=0}^{\infty} \frac{(1-\alpha q^{2n})(\alpha, q/a)_n (a/q)^n}{(1-\alpha)(q, \alpha a)_n} \sum_{k=0}^{n} \frac{(q^{-n}, \alpha q^n)_k q^k}{(q, \alpha b)_k} f(\alpha q^{k+1}),\end{aligned} \tag{1.1}$$

where the q-shifted factorial $(z)_\infty$ is defined by

$$(z)_\infty := (z;q)_\infty = \prod_{k=0}^{\infty} (1 - zq^k)$$

and

$$(z)_n = \frac{(z)_\infty}{(zq^n)_\infty}.$$

Here and in what follows, we always assume that $0 < |q| < 1$ and we also use the compact notation:

$$(a_1, a_2, \cdots, a_n)_k = (a_1)_k (a_2)_k \cdots (a_n)_k, \ k \text{ is a nonnegative integer and } \infty.$$

2000 *Mathematics Subject Classification.* 33D67; 33D15.

Key words and phrases. expansion formula; transformation formula; root system A_n; A_n extension; basic hypergeometric series; infinite product; Rogers' ${}_6\phi_5$ summation; Sylvester's identity; $(q)_\infty^m$; π_q; Rogers-Fine identity; Liu's extension of Rogers' ${}_6\phi_5$ summation; a generalization of Fang's identity; Andrews' expansion formula.

The expansion formula (1.1) is very useful and was used by Liu in [15] to extend Rogers' non-terminating ${}_6\phi_5$ summation formula [12, eq.(2.7.1)], to give a new proof of the orthogonality relation [12, eq.(7.5.15)] for the Askey-Wilson polynomials and to derive an interesting double summation formula which is equivalent to an identity of Andrews [2, Theorem 5].

Our work belongs to the context of multiple basic hypergeometric series over the root system A_n. Series of this type have been investigated systematically over the last many years and many important results in basic hypergeometric series have been extended. See Schlosser [26] for a comprehensive survey of this type of series. It should be mentioned that S.C. Milne made important contributions to multiple basic hypergeometric series over the root system A_n. See [5], [6], [9] and [27] for certain generalizations of Milne's A_n q-summation formulas. Recently, G. Bhatnager and S. Rai [4] extended some expansion formulas to the context of multiple series over root systems. A_n Extensions of Liu's results in [16, 18] have not been fully studied yet, even though his resuls are very useful in special functions and number theory [7, 15, 18, 19]. Inspired by their work, we will extend (1.1) to multiple basic hypergeometric series over the root system A_n.

Before stating our results, we need to mention some notations on multiple basic hypergeometric series over the root system A_n. The series we will discuss are of the form:

$$\sum_{\substack{k_r\geq 0\\ r=1,\cdots,n}} S(\boldsymbol{k}),$$

where $\boldsymbol{k}=(k_1,\cdots,k_n)$ and $k_1,\cdots,k_n$ are nonnegative integers. Similarly, $\boldsymbol{j}$ and $\boldsymbol{m}$ are multi-indices of nonnegative integers. This type of series reduces to classical basic hypergeometric series when $n=1$. We also use the notation $|\boldsymbol{k}|:=k_1+\cdots+k_n$. Series of this type are called A_n basic hypergeometric series if they only have

$$\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}$$

as a factor of $S(\boldsymbol{k})$.

The main result of this paper is to extend (1.1) to the following multiple transformation formula over the root system A_n.

Theorem 1.2. *Let $K(y)$ and $\beta(j)$ be chosen so that the series on both sides are analytic in $y_1,y_2,\cdots,y_n$ and converge absolutely for each y_r in a disk around the origin; or, $y_r=q^{N_r}$, for $r=1,2,\cdots,n$, and so the series on both sides terminate. Then*

$$\begin{aligned}
&K(\boldsymbol{y})\sum_{\substack{j_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}(x_r/x_sy_s)_{j_r}\\
&\quad\times\frac{1-bq^{2|\boldsymbol{j}|}}{(1-b)(bqy_1y_2\cdots y_n)_{|\boldsymbol{j}|}}(y_1y_2\cdots y_n)^{|\boldsymbol{j}|}q^{\sum_{r=1}^n(r-1)j_r}\beta(\boldsymbol{j})\\
&=\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(x_r/x_sy_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\quad\times\frac{(1-aq^{2|\boldsymbol{k}|})(a)_{|\boldsymbol{k}|}}{(1-a)(aqy_1y_2\cdots y_n)_{|\boldsymbol{k}|}}(y_1y_2\cdots y_n)^{|\boldsymbol{k}|}q^{\sum_{r=1}^n(r-1)k_r}
\end{aligned}\tag{1.2}$$

$$\times \sum_{\substack{0\le j_r\le k_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} (q^{-k_s}x_r/x_s)_{j_r}$$
$$\times \frac{(aq^{|\boldsymbol{k}|},bq)_{|\boldsymbol{j}|}}{(b)_{2|\boldsymbol{j}|}}(-1)^{|\boldsymbol{j}|}q^{\sum_{r=1}^n rj_r+\binom{|\boldsymbol{j}|}{2}}\beta(\boldsymbol{j})$$
$$\times \sum_{\substack{0\le m_r\le k_r-j_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{m_r-m_s+j_r-j_s}x_r/x_s}{1-q^{j_r-j_s}x_r/x_s} q^{\sum_{r=1}^n rm_r}K(q^{\boldsymbol{m}+\boldsymbol{j}})$$
$$\times \prod_{r,s=1}^{n} \frac{(q^{-k_s+j_r}x_r/x_s)_{m_r}}{(q^{1+j_r-j_s}x_r/x_s)_{m_r}} \frac{(aq^{|\boldsymbol{k}|+|\boldsymbol{j}|},bq^{1+|\boldsymbol{j}|})_{|\boldsymbol{m}|}}{(aq)_{|\boldsymbol{m}|+|\boldsymbol{j}|}(bq^{2|\boldsymbol{j}|+1})_{|\boldsymbol{m}|}}.$$

The identity (1.2) in Theorem 1.2 is an A_n extension of the following expansion formula:

$$\begin{aligned} &K(y)\sum_{j=0}^{\infty}\frac{(1-bq^{2j})(1/y)_j}{(1-b)(bqy)_j}y^j\beta_j\\ &=\sum_{k=0}^{\infty}\frac{(1-aq^{2k})(1/y,a)_k}{(1-a)(q,aqy)_k}y^k\sum_{j=0}^{k}\frac{(q^{-k},aq^k,bq)_j}{(b)_{2j}}(-1)^jq^{\binom{j+1}{2}}\beta_j\\ &\times\sum_{m=0}^{k-j}\frac{(q^{-k+j},aq^{j+k},bq^{j+1})_m}{(q,bq^{1+2j})_m(aq)_{j+m}}q^mK(q^{j+m}). \end{aligned} \tag{1.3}$$

Take $y_1\mapsto y$ in the $n=1$ case of (1.2) in Theorem 1.2 to obtain (1.3) easily. Another A_n extension of (1.3) can be found in [4, Theorem 2.2].

Theorem 1.2 plays an important role in multiple q-series over the root system A_n, from which many important results can be derived. In this paper, we will use it to deduce several summation formulas and transformation formulas for q-series over the root system A_n.

We first apply the identity in Theorem 1.2 to obtain four A_n extensions of (1.1).

Theorem 1.3. (First A_n extension of (1.1)) *Let $f(y)$ be such that the series on both sides are analytic in $a_1,a_2,\cdots,a_n$ and converge absolutely for each a_r in a disk around the origin; or, $a_r=q^{N_r}$, for $r=1,2,\cdots,n$, and so the series terminate. Then*

$$\begin{aligned} &f(\alpha a_1\cdots a_nq^{1-n})\frac{(\alpha q,\alpha a_1\cdots a_nb/q^n)_\infty}{(\alpha a_1\cdots a_nq^{1-n},\alpha b)_\infty}\\ &=\sum_{\substack{k_r\ge 0\\ r=1,2,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(qx_r/x_sa_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\ &\times\frac{(1-\alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)(\alpha a_1\cdots a_nq^{1-n})_{|\boldsymbol{k}|}}(a_1\cdots a_n)^{|\boldsymbol{k}|}q^{\sum_{r=1}^n(r-1)k_r-n|\boldsymbol{k}|}\\ &\times\sum_{\substack{0\le m_r\le k_r\\ r=1,2,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\\ &\times\frac{(\alpha q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}}{(\alpha b)_{|\boldsymbol{m}|}}q^{\sum_{r=1}^n rm_r}f(\alpha q^{1+|\boldsymbol{m}|}). \end{aligned} \tag{1.4}$$

Theorem 1.4. (Second A_n extension of (1.1)) *Let $f(y_1, y_2, \cdots, y_n)$ be such that the series on both sides are analytic in $a_1, a_2, \cdots, a_n$ and converge absolutely for each a_r in a disk around the origin; or, $a_r = q^{N_r}$, for $r = 1, 2, \cdots, n$, and so the series terminate. Then*

$$\begin{aligned}
&f(\alpha a_1, \cdots, \alpha a_n)\frac{(\alpha q, \alpha a_1 \cdots a_n b/q^n)_\infty}{(\alpha b, \alpha a_1 \cdots a_n q^{1-n})_\infty} \\
&= \sum_{\substack{k_r \geq 0 \\ r=1,2,\cdots,n}} \prod_{1 \leq r < s \leq n} \frac{1 - q^{k_r - k_s} x_r/x_s}{1 - x_r/x_s} \prod_{r,s=1}^{n} \frac{(qx_r/x_s a_s)_{k_r}}{(qx_r/x_s)_{k_r}} \\
&\times \frac{(1 - \alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)(\alpha a_1 \cdots a_n q^{1-n})_{|\boldsymbol{k}|}} (a_1 \cdots a_n)^{|\boldsymbol{k}|} q^{\sum_{r=1}^n (r-1)k_r - n|\boldsymbol{k}|} \qquad (1.5) \\
&\times \sum_{\substack{0 \leq m_r \leq k_r \\ r=1,2,\cdots,n}} \prod_{1 \leq r < s \leq n} \frac{1 - q^{m_r - m_s} x_r/x_s}{1 - x_r/x_s} \prod_{r,s=1}^{n} \frac{(q^{-k_s} x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}} \\
&\times \frac{(\alpha q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}}{(\alpha b)_{|\boldsymbol{m}|}} q^{\sum_{r=1}^n r m_r} f(\alpha q^{1+m_1}, \cdots, \alpha q^{1+m_n}).
\end{aligned}$$

Theorem 1.5. (Third A_n extension of (1.1)) *Let $f(y)$ be such that the series on both sides are analytic in $a_1, a_2, \cdots, a_n$ and converge absolutely for each a_r in a disk around the origin; or, $a_r = q^{N_r}$, for $r = 1, 2, \cdots, n$, and so the series terminate. Then*

$$\begin{aligned}
&f(\alpha a_1 \cdots a_n q^{1-n})\frac{(\alpha q)_\infty}{(\alpha a_1 \cdots a_n q^{1-n})_\infty} \prod_{r=1}^{n} \frac{(\alpha a_r b x_r/q)_\infty}{(\alpha b x_r)_\infty} \\
&= \sum_{\substack{k_r \geq 0 \\ r=1,2,\cdots,n}} \prod_{1 \leq r < s \leq n} \frac{1 - q^{k_r - k_s} x_r/x_s}{1 - x_r/x_s} \prod_{r,s=1}^{n} \frac{(qx_r/x_s a_s)_{k_r}}{(qx_r/x_s)_{k_r}} \\
&\times \frac{(1 - \alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)(\alpha a_1 \cdots a_n q^{1-n})_{|\boldsymbol{k}|}} (a_1 \cdots a_n)^{|\boldsymbol{k}|} q^{\sum_{r=1}^n (r-1)k_r - n|\boldsymbol{k}|} \qquad (1.6) \\
&\times \sum_{\substack{0 \leq m_r \leq k_r \\ r=1,2,\cdots,n}} \prod_{1 \leq r < s \leq n} \frac{1 - q^{m_r - m_s} x_r/x_s}{1 - x_r/x_s} \prod_{r,s=1}^{n} \frac{(q^{-k_s} x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}} \\
&\times \frac{(\alpha q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}}{\prod_{r=1}^n (\alpha b x_r)_{m_r}} q^{\sum_{r=1}^n r m_r} f(\alpha q^{1+|\boldsymbol{m}|}).
\end{aligned}$$

Theorem 1.6. (Fourth A_n extension of (1.1)) *Let $f(y_1, y_2, \cdots, y_n)$ be such that the series on both sides are analytic in $a_1, a_2, \cdots, a_n$ and converge absolutely for each a_r in a disk around the origin; or, $a_r = q^{N_r}$, for $r = 1, 2, \cdots, n$, and so the series terminate. Then*

$$\begin{aligned}
&f(\alpha a_1, \cdots, \alpha a_n)\frac{(\alpha q)_\infty}{(\alpha a_1 a_2 \cdots a_n q^{1-n})_\infty} \prod_{r=1}^{n} \frac{(\alpha a_r b x_r/q)_\infty}{(\alpha b x_r)_\infty} \\
&= \sum_{\substack{k_r \geq 0 \\ r=1,2,\cdots,n}} \prod_{1 \leq r < s \leq n} \frac{1 - q^{k_r - k_s} x_r/x_s}{1 - x_r/x_s} \prod_{r,s=1}^{n} \frac{(qx_r/x_s a_s)_{k_r}}{(qx_r/x_s)_{k_r}} \qquad (1.7) \\
&\times \frac{(1 - \alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)(\alpha a_1 a_2 \cdots a_n q^{1-n})_{|\boldsymbol{k}|}} (a_1 a_2 \cdots a_n)^{|\boldsymbol{k}|} q^{\sum_{r=1}^n (r-1)k_r - n|\boldsymbol{k}|}
\end{aligned}$$

$$\times \sum_{\substack{0\le m_r\le k_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}$$

$$\times \frac{(\alpha q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}}{\prod_{r=1}^{n}(\alpha b x_r)_{m_r}} q^{\sum_{r=1}^{n} r m_r} f(\alpha q^{1+m_1},\cdots,\alpha q^{1+m_n}).$$

Two other A_n extensions of (1.1) are given in the following theorems.

Theorem 1.7. (Fifth A_n extension of (1.1)) *Let $f(y)$ be such that the series on both sides are analytic in $a_1, a_2, \cdots, a_n$ and converge absolutely for each a_r in a disk around the origin; or, $a_r = q^{N_r}$, for $r = 1, 2, \cdots, n$, and so the series terminate. Then*

$$\begin{aligned}
&f(\alpha a_1\cdots a_n q^{1-n}) \prod_{r=1}^{n} \frac{(\alpha q x_r, \alpha a_r b x_r/q)_\infty}{(\alpha b x_r, \alpha a_r x_r)_\infty}\\
&= \sum_{\substack{k_r\ge 0\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{(qx_r/x_s a_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times \prod_{r=1}^{n} \frac{1-\alpha x_r q^{k_r+|\boldsymbol{k}|}}{1-\alpha x_r} \prod_{r=1}^{n} \frac{(\alpha x_r)_{|\boldsymbol{k}|}}{(\alpha x_r a_r)_{|\boldsymbol{k}|}} (a_1\cdots a_n)^{|\boldsymbol{k}|}\, q^{\sum_{r=1}^{n}(r-1)k_r - n|\boldsymbol{k}|} \qquad (1.8)\\
&\times \sum_{\substack{0\le m_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_r/x_s\right)_{m_r}}{(qx_r/x_s)_{m_r}}\\
&\times \prod_{r=1}^{n} \frac{\left(\alpha x_r q^{|\boldsymbol{k}|}\right)_{m_r}}{(\alpha b x_r)_{m_r}} q^{\sum_{r=1}^{n} r m_r} f(\alpha q^{1+|\boldsymbol{m}|}).
\end{aligned}$$

Theorem 1.8. (Sixth A_n extension of (1.1)) *Let $f(y_1, y_2, \cdots, y_n)$ be a function of n variables such that the series on both sides are analytic in $a_1, a_2, \cdots, a_n$ and converge absolutely for each a_r in a disk around the origin; or, $a_r = q^{N_r}$, for $r = 1, 2, \cdots, n$, and so the series terminate. Then*

$$\begin{aligned}
&f(\alpha a_1, \cdots, \alpha a_n) \prod_{r=1}^{n} \frac{(\alpha q x_r, \alpha a_r b x_r/q)_\infty}{(\alpha b x_r, \alpha x_r a_r)_\infty}\\
&= \sum_{\substack{k_r\ge 0\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{(qx_r/x_s a_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times \prod_{r=1}^{n} \frac{1-\alpha x_r q^{k_r+|\boldsymbol{k}|}}{1-\alpha x_r} \prod_{r=1}^{n} \frac{(\alpha x_r)_{|\boldsymbol{k}|}}{(\alpha x_r a_r)_{|\boldsymbol{k}|}} (a_1\cdots a_n/q^n)^{|\boldsymbol{k}|}\, q^{\sum_{r=1}^{n}(r-1)k_r} \qquad (1.9)\\
&\times \sum_{\substack{0\le m_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_r/x_s\right)_{m_r}}{(qx_r/x_s)_{m_r}}\\
&\times \prod_{r=1}^{n} \frac{\left(\alpha x_r q^{|\boldsymbol{k}|}\right)_{m_r}}{(\alpha b x_r)_{m_r}} q^{\sum_{r=1}^{n} r m_r} f(\alpha q^{1+m_1},\cdots,\alpha q^{1+m_n}).
\end{aligned}$$

The identities in Theorems 1.3–1.8 are all A_n extensions of (1.1); we set $x_r \mapsto x_r/x_n, a_1 \mapsto a$ in the $n = 1$ case of the identities in Theorems 1.3–1.8 to obtain (1.1) readily.

Two groups of multiple expansion formulas for infinite products can be derived from the identities in Theorems 1.3–1.8. We set

$$f(x)=\frac{(cx/q,dx/q)_\infty}{(\beta x/q,\gamma x/q)_\infty},$$

in (1.4), (1.6) and (1.8), and then simplify to obtain the following three multiple expansion formulas for infinite products.

Theorem 1.9. *Let $\alpha,\beta,\gamma,a_1,\cdots,a_n,b,c,d$ be such that $\max\{|\alpha a_1\cdots a_nq^{1-n}|,|\alpha b|,|\alpha c|,|\alpha d|,|\alpha\beta a_1\cdots a_n/q^n|,|\alpha\gamma a_1\cdots a_n/q^n|\}<1$; or, $a_r=q^{N_r}$, for $r=1,2,\cdots,n$, and so the series terminate. Suppose that none of the denominators in the following identity vanishes, then*

$$\begin{aligned}
&\frac{(\alpha q,\alpha a_1\cdots a_nb/q^n,\alpha a_1\cdots a_nc/q^n,\alpha a_1\cdots a_nd/q^n,\alpha\beta,\alpha\gamma)_\infty}{(\alpha a_1\cdots a_nq^{1-n},\alpha b,\alpha c,\alpha d,\alpha\beta a_1\cdots a_n/q^n,\alpha\gamma a_1\cdots a_n/q^n)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(qx_r/x_sa_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times\frac{(1-\alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)(\alpha a_1\cdots a_nq^{1-n})_{|\boldsymbol{k}|}}(a_1\cdots a_n)^{|\boldsymbol{k}|}q^{\sum_{r=1}^n(r-1)k_r-n|\boldsymbol{k}|}\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\\
&\quad\times\frac{(\alpha q^{|\boldsymbol{k}|},\alpha\beta,\alpha\gamma)_{|\boldsymbol{m}|}}{(\alpha b,\alpha c,\alpha d)_{|\boldsymbol{m}|}}q^{\sum_{r=1}^n rm_r}.
\end{aligned}$$

Theorem 1.10. *Let $\alpha,\beta,\gamma,a_1,\cdots,a_n,b,c,d,x_1,\cdots,x_n$ be such that $\max\{|\alpha c|,|\alpha d|,|\alpha a_1\cdots a_nq^{1-n}|,|\alpha\beta a_1\cdots a_n/q^n|,|\alpha\gamma a_1\cdots a_n/q^n|\}<1$ and $\max\{|\alpha bx_r|:1\leq r\leq n\}<1$; or, $a_r=q^{N_r}$, for $r=1,2,\cdots,n$, and so the series terminate. Suppose that none of the denominators in the following identity vanishes, then*

$$\begin{aligned}
&\frac{(\alpha q,\alpha a_1\cdots a_nc/q^n,\alpha a_1\cdots a_nd/q^n,\alpha\beta,\alpha\gamma)_\infty}{(\alpha a_1\cdots a_nq^{1-n},\alpha c,\alpha d,\alpha a_1\cdots a_n\beta/q^n,\alpha a_1\cdots a_n\gamma/q^n)_\infty}\prod_{r=1}^{n}\frac{(\alpha a_rbx_r/q)_\infty}{(\alpha bx_r)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(qx_r/x_sa_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times\frac{(1-\alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)(\alpha a_1\cdots a_nq^{1-n})_{|\boldsymbol{k}|}}(a_1\cdots a_n)^{|\boldsymbol{k}|}q^{\sum_{r=1}^n(r-1)k_r-n|\boldsymbol{k}|}\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\\
&\quad\times\frac{(\alpha q^{|\boldsymbol{k}|},\alpha\beta,\alpha\gamma)_{|\boldsymbol{m}|}}{(\alpha c,\alpha d)_{|\boldsymbol{m}|}\prod_{r=1}^n(\alpha bx_r)_{m_r}}q^{\sum_{r=1}^n rm_r}.
\end{aligned}$$

Theorem 1.11. *Let $\alpha,\beta,\gamma,a_1,\cdots,a_n,b,c,d,x_1,\cdots,x_n$ be such that $\max\{|\alpha c|,|\alpha d|,|\alpha\beta a_1\cdots a_n/q^n|,|\alpha\gamma a_1\cdots a_n/q^n|\}<1$ and $\max\{|\alpha ax_r|,|\alpha bx_r|:1\leq r\leq n\}<1$; or, $a_r=q^{N_r}$, for $r=1,2,\cdots,n$, and so the series terminate. Suppose that none*

of the denominators in the following identity vanishes, then

$$
\begin{aligned}
&\frac{(\alpha\beta,\alpha\gamma,\alpha a_1\cdots a_n cq^{-n},\alpha a_1\cdots a_n dq^{-n})_\infty}{(\alpha c,\alpha d,\alpha\beta a_1\cdots a_n q^{-n},\alpha\gamma a_1\cdots a_n q^{-n})_\infty}\prod_{r=1}^{n}\frac{(\alpha q x_r,\alpha a_r b x_r/q)_\infty}{(\alpha b x_r,\alpha a_r x_r)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(qx_r/x_sa_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times\prod_{r=1}^{n}\frac{1-\alpha x_r q^{k_r+|\boldsymbol{k}|}}{1-\alpha x_r}\prod_{r=1}^{n}\frac{(\alpha x_r)_{|\boldsymbol{k}|}}{(\alpha x_r a_r)_{|\boldsymbol{k}|}}(a_1\cdots a_n)^{|\boldsymbol{k}|}\,q^{\sum_{r=1}^{n}(r-1)k_r-n|\boldsymbol{k}|}\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{\left(q^{-k_s}x_r/x_s\right)_{m_r}}{(qx_r/x_s)_{m_r}}\\
&\times\prod_{r=1}^{n}\frac{\left(\alpha x_r q^{|\boldsymbol{k}|}\right)_{m_r}}{(\alpha b x_r)_{m_r}}\frac{(\alpha\beta,\alpha\gamma)_{|\boldsymbol{m}|}}{(\alpha c,\alpha d)_{|\boldsymbol{m}|}}q^{\sum_{r=1}^{n}rm_r}.
\end{aligned}
\tag{1.10}
$$

The second group can be obtained as follows. We now take

$$f(y_1,y_2,\cdots,y_n)=\prod_{r=1}^{n}\frac{(cx_ry_r/q,dx_ry_r/q)_\infty}{(\beta x_ry_r/q,\gamma x_ry_r/q)_\infty}$$

in (1.5), (1.7) and (1.9), and then simplify to give three other multiple expansion formulas for infinite products.

Theorem 1.12. *Let $\alpha,\beta,\gamma,a_1,a_2,\cdots,a_n,b,c,d,x_1,\cdots,x_n$ be such that $\max\{|\alpha b|,|\alpha a_1\cdots a_nq^{1-n}|\}<1$ and $\max\{|\alpha cx_r|,|\alpha dx_r|,|\alpha\beta a_rx_r/q|,|\alpha\gamma a_rx_r/q|:1\leq r\leq n\}<1$; or, $a_r=q^{N_r}$, for $r=1,2,\cdots,n$, and so the series terminate. Suppose that none of the denominators in the following identity vanishes, then*

$$
\begin{aligned}
&\frac{(\alpha q,\alpha a_1\cdots a_nb/q^n)_\infty}{(\alpha b,\alpha a_1\cdots a_nq^{1-n})_\infty}\prod_{r=1}^{n}\frac{(\alpha a_rcx_r/q,\alpha a_rdx_r/q,\alpha\beta x_r,\alpha\gamma x_r)_\infty}{(\alpha cx_r,\alpha dx_r,\alpha\beta a_rx_r/q,\alpha\gamma a_rx_r/q)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(qx_r/x_sa_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times\frac{(1-\alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)(\alpha a_1\cdots a_nq^{1-n})_{|\boldsymbol{k}|}}(a_1\cdots a_n)^{|\boldsymbol{k}|}q^{\sum_{r=1}^{n}(r-1)k_r-n|\boldsymbol{k}|}\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\\
&\times\frac{(\alpha q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}}{(\alpha b)_{|\boldsymbol{m}|}}\prod_{r=1}^{n}\frac{(\alpha\beta x_r,\alpha\gamma x_r)_{m_r}}{(\alpha cx_r,\alpha dx_r)_{m_r}}q^{\sum_{r=1}^{n}rm_r}.
\end{aligned}
$$

Theorem 1.13. *Let $\alpha,\beta,\gamma,a_1,\cdots,a_n,b,c,d,x_1,\cdots,x_n$ be such that $\max\{|\alpha a_1\cdots a_nq^{1-n}|\}<1$ and $\max\{|\alpha bx_r|,|\alpha cx_r|,|\alpha dx_r|,|\alpha\beta a_rx_r/q|,|\alpha\gamma a_rx_r/q|:1\leq r\leq n\}<1$; or, $a_r=q^{N_r}$, for $r=1,2,\cdots,n$, and so the series terminate. Suppose that none of the denominators in the following identity vanishes, then*

$$\frac{(\alpha q)_\infty}{(\alpha a_1 a_2\cdots a_n q^{1-n})_\infty}\prod_{r=1}^{n}\frac{(\alpha a_r bx_r/q,\alpha a_r cx_r/q,\alpha a_r dx_r/q,\alpha\beta x_r,\alpha\gamma x_r)_\infty}{(\alpha bx_r,\alpha cx_r,\alpha dx_r,\alpha\beta a_r x_r/q,\alpha\gamma a_r x_r/q)_\infty}$$
$$=\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(qx_r/x_sa_s)_{k_r}}{(qx_r/x_s)_{k_r}}$$
$$\times\frac{(1-\alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)(\alpha a_1a_2\cdots a_nq^{1-n})_{|\boldsymbol{k}|}}(a_1a_2\cdots a_n)^{|\boldsymbol{k}|}q^{\sum_{r=1}^n(r-1)k_r-n|\boldsymbol{k}|}$$
$$\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}$$
$$\times(\alpha q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}\prod_{r=1}^{n}\frac{(\alpha\beta x_r,\alpha\gamma x_r)_{m_r}}{(\alpha bx_r,\alpha cx_r,\alpha dx_r)_{m_r}}q^{\sum_{r=1}^n rm_r}.$$

Theorem 1.14. *Let $\alpha,\beta,\gamma,a_1,\cdots,a_n,b,c,d,x_1,\cdots,x_n$ be such that $\max\{|\alpha ax_r|,|\alpha bx_r|,|\alpha cx_r|,|\alpha dx_r|,|\alpha\beta a_rx_r/q|,|\alpha\gamma a_rx_r/q|:1\leq r\leq n\}<1$; or, $a_r=q^{N_r}$, for $r=1,2,\cdots,n$, and so the series terminate. Suppose that none of the denominators in the following identity vanishes, then*

$$\begin{aligned}
&\prod_{r=1}^{n}\frac{(\alpha qx_r,\alpha a_rbx_r/q,\alpha a_rcx_r/q,\alpha a_rdx_r/q,\alpha\beta x_r,\alpha\gamma x_r)_\infty}{(\alpha a_rx_r,\alpha bx_r,\alpha cx_r,\alpha dx_r,\alpha\beta a_rx_r/q,\alpha\gamma a_rx_r/q)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(qx_r/x_sa_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times\prod_{r=1}^{n}\frac{1-\alpha x_rq^{k_r+|\boldsymbol{k}|}}{1-\alpha x_r}\prod_{r=1}^{n}\frac{(\alpha x_r)_{|\boldsymbol{k}|}}{(\alpha x_ra_r)_{|\boldsymbol{k}|}}(a_1\cdots a_n/q^n)^{|\boldsymbol{k}|}\,q^{\sum_{r=1}^n(r-1)k_r}\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{\left(q^{-k_s}x_r/x_s\right)_{m_r}}{(qx_r/x_s)_{m_r}}\\
&\times\prod_{r=1}^{n}\frac{\left(\alpha x_rq^{|\boldsymbol{k}|},\alpha\beta x_r,\alpha\gamma x_r\right)_{m_r}}{(\alpha bx_r,\alpha cx_r,\alpha dx_r)_{m_r}}q^{\sum_{r=1}^n rm_r}.
\end{aligned}\tag{1.11}$$

Take $x_r\mapsto x_r/x_n, a_1\mapsto a$ in the $n=1$ case of the identities in Theorems 1.9–1.14. We can obtain the following expansion formula [15, Theorem 1.2]: for $\max\{|\alpha a|,|\alpha b|,|\alpha c|,|\alpha d|,|\alpha\beta a/q|,|\alpha\gamma a/q|\}<1$, we have

$$\begin{aligned}
&\frac{(\alpha q,\alpha ab/q,\alpha ac/q,\alpha ad/q,\alpha\beta,\alpha\gamma)_\infty}{(\alpha a,\alpha b,\alpha c,\alpha d,\alpha\beta a/q,\alpha\gamma a/q)_\infty}\\
&=\sum_{n=0}^{\infty}\frac{(1-\alpha q^{2n})(\alpha,q/a)_n(a/q)^n}{(1-\alpha)(q,\alpha a)_n}\sum_{k=0}^{n}\frac{(q^{-n},\alpha q^n,\alpha\beta,\alpha\gamma)_kq^k}{(q,\alpha b,\alpha c,\alpha d)_k}.
\end{aligned}\tag{1.12}$$

Therefore, the identities in Theorems 1.9–1.14 are all A_n extensions of (1.12). The identity (1.12) is very useful in [15]. It was used by Liu [15] to give a new proof of the orthogonality relation for the Askey-Wilson polynomials and to derive a useful

identity which is equivalent to the well-known identity of Andrews in [2, Theorem 5].

Several multiple expansion formulas for $(q)_\infty^3$ and $(q)_\infty^{-2}$ can be easily derived from the identities in Theorems 1.9–1.14. For example, we set $\alpha = q, \beta = \gamma = 1, a_1 = \cdots = a_n = b = c = d = 0$ or $\alpha = 1, \beta = \gamma = q, a_1 = \cdots = a_n = b = c = d = 0$ in the identity of Theorem 1.9 to get two multiple expansion formulas for $(q)_\infty^3$:

$$\begin{aligned}(q)_\infty^3 = &\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}} \prod_{1\leq r<s\leq n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-q^{2|\boldsymbol{k}|+1})(q)_{|\boldsymbol{k}|}}{\prod_{r,s=1}^n (qx_r/x_s)_{k_r}}\\ &\times(-1)^{n|\boldsymbol{k}|}q^{\sum_{r=1}^n (r-1)k_r+n\sum_{r=1}^n\binom{k_r}{2}}\prod_{r=1}^n x_r^{nk_r-|\boldsymbol{k}|}\\ &\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\\ &\times\prod_{r,s=1}^n\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\cdot(q^{|\boldsymbol{k}|+1},q,q)_{|\boldsymbol{m}|}q^{\sum_{r=1}^n rm_r}\end{aligned}\tag{1.13}$$

and

$$\begin{aligned}(q)_\infty^3 = 1+&\sum_{\substack{k_1\geq 0,\cdots,k_n\geq 0\\ |\boldsymbol{k}|\geq 1}} \prod_{1\leq r<s\leq n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-q^{2|\boldsymbol{k}|})(q)_{|\boldsymbol{k}|-1}}{\prod_{r,s=1}^n (qx_r/x_s)_{k_r}}\\ &\times(-1)^{n|\boldsymbol{k}|}q^{\sum_{r=1}^n (r-1)k_r+n\sum_{r=1}^n\binom{k_r}{2}}\prod_{r=1}^n x_r^{nk_r-|\boldsymbol{k}|}\\ &\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\\ &\times\prod_{r,s=1}^n\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\cdot(q^{|\boldsymbol{k}|},q,q)_{|\boldsymbol{m}|}q^{\sum_{r=1}^n rm_r}.\end{aligned}\tag{1.14}$$

Taking $\alpha = q, \beta = \gamma = a_1 = \cdots = a_n = 0, b = q, c = d = 1$ or $\alpha = 1, \beta = \gamma = a_1 = \cdots = a_n = 0, b = c = d = q$ in the identity of Theorem 1.9 we obtain two multiple expansion formulas for $(q)_\infty^{-2}$:

$$\begin{aligned}\frac{1}{(q)_\infty^2} = &\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}} \prod_{1\leq r<s\leq n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-q^{2|\boldsymbol{k}|+1})(q)_{|\boldsymbol{k}|}}{(1-q)\prod_{r,s=1}^n (qx_r/x_s)_{k_r}}\\ &\times(-1)^{n|\boldsymbol{k}|}q^{\sum_{r=1}^n (r-1)k_r+n\sum_{r=1}^n\binom{k_r}{2}}\prod_{r=1}^n x_r^{nk_r-|\boldsymbol{k}|}\\ &\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\\ &\times\prod_{r,s=1}^n\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\cdot\frac{(q^{|\boldsymbol{k}|+1})_{|\boldsymbol{m}|}}{(q^2,q,q)_{|\boldsymbol{m}|}}q^{\sum_{r=1}^n rm_r}\end{aligned}\tag{1.15}$$

and

$$\begin{aligned}\frac{1}{(q)_\infty^2} = 1 + &\sum_{\substack{k_1\geq 0,\cdots,k_n\geq 0\\ |\boldsymbol{k}|\geq 1}} \prod_{1\leq r<s\leq n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-q^{2|\boldsymbol{k}|})(q)_{|\boldsymbol{k}|-1}}{\prod_{r,s=1}^n (qx_r/x_s)_{k_r}}\\ &\times(-1)^{n|\boldsymbol{k}|}q^{\sum_{r=1}^n (r-1)k_r+n\sum_{r=1}^n\binom{k_r}{2}}\prod_{r=1}^n x_r^{nk_r-|\boldsymbol{k}|}\\ &\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\\ &\times\prod_{r,s=1}^n\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\cdot\frac{(q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}}{(q,q,q)_{|\boldsymbol{m}|}}q^{\sum_{r=1}^n rm_r}.\end{aligned}\tag{1.16}$$

For any interger m, some general multiple expansion formulas for $(q)_\infty^m$ will be showed in Section 5.

The rest of this paper is organized as follows. In the next section, we first construct two inverse matrices equivalent to the matrices in [21, Theorem 8.26] and then employ a inverse relation satisfied by these two matrices to prove Theorem 1.2. The identity in Theorem 1.2 is very useful. In Section 3, we will apply the identity (1.2) in Theorem 1.2 and a multiple series expansion formula in [4, Theorem 2.2] to show the identities in Theorems 1.3–1.8. As applications, in Section 4, we will apply the identity in Theorem 1.11 to deduce an A_n Rogers' ${}_6\phi_5$ summation formula which is equivalent to the identity in [23, Theorem A.3]. From this formula, we will derive an A_n extension of Sylvester's identity. This A_n extension is equivalent to the identity in [20, Theorem 5.19]; in Section 5, we will employ the identities in Theorems 1.3 and 1.9 to deduce some multiple expansion formulas for $(q)_\infty^m, \pi_q$ and $1/\pi_q$; in Section 6, we will apply the identity (1.2) of Theorem 1.2 to present two A_n extensions of the Rogers-Fine identity [11, eq.(14.1)]; in Section 7, we will employ the identity (1.8) in Theorem 1.8 to give an A_n extension of Liu's extension [17, Theorem 3] of Rogers' non-terminating ${}_6\phi_5$ summation [12, eq.(II.20)]; in Section 8, we will use the identity (1.2) of Theorem 1.2 to prove an A_n extension of a generalization of Fang's identity [8, eq.(9)]; and in the last section, we will apply the identity in Theorem 1.9 to deduce an A_n extension of Andrews' expansion formula [2, Theorem 5].

2. Proof of Theorem 1.2

We first construct two inverse, infinite matrices, which are equivalent to the matrices in [21, Theorem 8.26]. Let

$$H_{\boldsymbol{km}}(a) = \prod_{r,s=1}^n\left(q^{1+m_r-m_s}\frac{x_r}{x_s}\right)_{k_r-m_r}^{-1}(aq)_{|\boldsymbol{k}|+|\boldsymbol{m}|}^{-1}$$

and

$$\begin{aligned}I_{\boldsymbol{km}}(a) = &(1-aq^{2|\boldsymbol{k}|})(aq)_{|\boldsymbol{k}|+|\boldsymbol{m}|-1}\prod_{r,s=1}^n\left(q^{1+m_r-m_s}\frac{x_r}{x_s}\right)_{k_r-m_r}^{-1}\\ &\times(-1)^{|\boldsymbol{k}|+|\boldsymbol{m}|}q^{\binom{|\boldsymbol{k}|-|\boldsymbol{m}|}{2}}\end{aligned}$$

with rows and columns ordered lexicographically. Then $\boldsymbol{H} = (H_{\boldsymbol{km}}(a))$ and $\boldsymbol{I} = (I_{\boldsymbol{km}}(a))$ are inverse, infinite matrices.

Recall the following important result [21, Lemma 4.3]:

$$\begin{aligned}&\prod_{r,s=1}^{n}\left(q^{1+m_r-m_s}\frac{x_r}{x_s}\right)_{k_r-m_r}^{-1}\\ &=(-1)^{|\boldsymbol{m}|}q^{|\boldsymbol{k}||\boldsymbol{m}|-\binom{|\boldsymbol{m}|}{2}+\sum_{r=1}^{n}(r-1)m_r}\\ &\quad\times\prod_{1\le r<s\le n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{k_r}}.\end{aligned}\tag{2.1}$$

Let $\boldsymbol{D}$ and $\boldsymbol{E}$ be two diagonal matrices with diagonal entries given by

$$D_{kk}=(aq)_{|\boldsymbol{k}|}\prod_{r,s=1}^{n}(qx_r/x_s)_{k_r}$$

and

$$E_{mm}=(-1)^{|\boldsymbol{m}|}q^{\binom{|\boldsymbol{m}|}{2}}(1-aq^{2|\boldsymbol{m}|}).$$

Set $\boldsymbol{F}=\boldsymbol{DHE}$ and $\boldsymbol{G}=\boldsymbol{E}^{-1}\boldsymbol{ID}^{-1}$. Then, by (2.1),

$$\begin{aligned}F_{\boldsymbol{km}}(a)=&\prod_{1\le r<s\le n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}q^{|\boldsymbol{k}||\boldsymbol{m}|+\sum_{r=1}^{n}(r-1)m_r}\\ &\times\frac{1-aq^{2|\boldsymbol{m}|}}{(aq^{|\boldsymbol{k}|+1})_{|\boldsymbol{m}|}}\prod_{r,s=1}^{n}(q^{-k_s}x_r/x_s)_{m_r}\end{aligned}\tag{2.2}$$

and

$$\begin{aligned}G_{\boldsymbol{km}}(a)=&\prod_{1\le r<s\le n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}q^{\sum_{r=1}^{n}rm_r}\\ &\times\frac{(aq^{|\boldsymbol{m}|})_{|\boldsymbol{k}|}}{1-aq^{|\boldsymbol{m}|}}\prod_{r,s=1}^{n}\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{k_r}(qx_r/x_s)_{m_r}}.\end{aligned}\tag{2.3}$$

Since $\boldsymbol{H}$ and $\boldsymbol{I}$ are inverses, the matrices $\boldsymbol{F}$ and $\boldsymbol{G}$ are also inverses. Note that $F_{\boldsymbol{km}}(a)=G_{\boldsymbol{km}}(a)=0$ unless $m_r\le k_r$ for $r=1,2,\cdots,n$. We have

$$\sum_{\substack{m_r\le k_r\le N_r\\ r=1,2,\cdots,n}}F_{\boldsymbol{Nk}}(a)G_{\boldsymbol{km}}(a)=\prod_{r=1}^{n}\delta_{N_rm_r},\tag{2.4}$$

where

$$\delta_{ij}=\begin{cases}1, & \text{if } i=j,\\ 0, & \text{otherwise.}\end{cases}$$

We now prove (1.2). Let $f_{\boldsymbol{k}}(\boldsymbol{y};a)$ be a function of $\boldsymbol{y}=(y_1,y_2,\cdots,y_n)$ given by

$$\begin{aligned}f_{\boldsymbol{k}}(\boldsymbol{y};a)=&\prod_{1\le r<s\le n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}q^{\sum_{r=1}^{n}(r-1)k_r}\\ &\times\frac{1-aq^{2|\boldsymbol{k}|}}{(aqy_1y_2\cdots y_n)_{|\boldsymbol{k}|}}(y_1y_2\cdots y_n)^{|\boldsymbol{k}|}\prod_{r,s=1}^{n}(x_r/x_sy_s)_{k_r}.\end{aligned}$$

Then $f_{\boldsymbol{k}}(\boldsymbol{y};a)$ is analytic in $\boldsymbol{y}$ in a disk around the origin and

$$f_{\boldsymbol{k}}(q^{\boldsymbol{m}};a)=F_{\boldsymbol{mk}}(a). \tag{2.5}$$

Let

$$g(\boldsymbol{y})=K(\boldsymbol{y})\sum_{\substack{j_r\geq 0\\ r=1,2,\cdots,n}} f_{\boldsymbol{j}}(\boldsymbol{y};b)\beta(\boldsymbol{j}). \tag{2.6}$$

It follows from (2.4) that

$$\begin{aligned} g(q^{\boldsymbol{N}}) &= \sum_{\substack{0\leq m_r\leq N_r\\ r=1,2,\cdots,n}}\ \sum_{\substack{m_r\leq k_r\leq N_r\\ r=1,2,\cdots,n}} F_{\boldsymbol{Nk}}(a)G_{\boldsymbol{km}}(a)g(q^{\boldsymbol{m}})\\ &= \sum_{\substack{0\leq k_r\leq N_r\\ r=1,2,\cdots,n}} F_{\boldsymbol{Nk}}(a)\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}} G_{\boldsymbol{km}}(a)g(q^{\boldsymbol{m}}). \end{aligned} \tag{2.7}$$

Substituting (2.6) into both sides of (2.7) we have

$$\begin{aligned} &K(q^{\boldsymbol{N}})\sum_{\substack{j_r\geq 0\\ r=1,2,\cdots,n}} f_{\boldsymbol{j}}(q^{\boldsymbol{N}};b)\beta(\boldsymbol{j})\\ &= \sum_{\substack{0\leq k_r\leq N_r\\ r=1,2,\cdots,n}} F_{\boldsymbol{Nk}}(a)\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}} G_{\boldsymbol{km}}(a)K(q^{\boldsymbol{m}})\sum_{\substack{0\leq j_r\leq m_r\\ r=1,2,\cdots,n}} f_{\boldsymbol{j}}(q^{\boldsymbol{m}};b)\beta(\boldsymbol{j})\\ &= \sum_{\substack{0\leq k_r\leq N_r\\ r=1,2,\cdots,n}} F_{\boldsymbol{Nk}}(a)\sum_{\substack{0\leq j_r\leq k_r\\ r=1,2,\cdots,n}}\beta(\boldsymbol{j})\sum_{\substack{0\leq m_r\leq k_r-j_r\\ r=1,2,\cdots,n}} G_{\boldsymbol{k},\boldsymbol{m}+\boldsymbol{j}}(a)K(q^{\boldsymbol{m}+\boldsymbol{j}})f_{\boldsymbol{j}}(q^{\boldsymbol{m}+\boldsymbol{j}};b), \end{aligned} \tag{2.8}$$

where we interchanged the inner sums using the identity

$$\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\ \sum_{\substack{0\leq j_r\leq m_r\\ r=1,2,\cdots,n}} S(\boldsymbol{m},\boldsymbol{j})=\sum_{\substack{0\leq j_r\leq k_r\\ r=1,2,\cdots,n}}\ \sum_{\substack{0\leq m_r\leq k_r-j_r\\ r=1,2,\cdots,n}} S(\boldsymbol{m}+\boldsymbol{j},\boldsymbol{j}).$$

Replacing N_r by m_r+j_r, y_r by j_r in the identity of [21, Lemma 4.3] we get

$$\begin{aligned} &\prod_{r,s=1}^{n}\frac{(q^{-m_s-j_s}x_r/x_s)_{j_r}}{(qx_r/x_s)_{m_r+j_r}}\\ &=\prod_{r,s=1}^{n}(q^{1+j_r-j_s}x_r/x_s)_{m_r}^{-1}\prod_{1\leq r<s\leq n}\frac{1-x_r/x_s}{1-q^{j_r-j_s}x_r/x_s}\\ &\quad\times(-1)^{|\boldsymbol{j}|}q^{\binom{|\boldsymbol{j}|}{2}-\sum_{r=1}^{n}(r-1)j_r-(|\boldsymbol{m}|+|\boldsymbol{j}|)|\boldsymbol{j}|}. \end{aligned} \tag{2.9}$$

Dividing both sides of (2.8) by $1-b$, then substituting (2.2), (2.3), (2.5) and (2.9) into (2.8) and finally simplifying using the identities

$$\begin{aligned} \frac{(aq^{m+j})_k}{1-aq^{m+j}}&=\frac{(a)_k(aq^k)_j(aq^{k+j})_m}{(1-a)(aq)_{m+j}},\\ \frac{1-bq^{2j}}{(1-b)(bq^{m+j+1})_j}&=\frac{(bq)_j(bq^{1+j})_m}{(b)_{2j}(bq^{2j+1})_m} \end{aligned}$$

we obtain

$$
\begin{aligned}
&K(q^{\boldsymbol{N}}) \sum_{\substack{0\le j_r\le N_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} (q^{-N_s}x_r/x_s)_{j_r}\\
&\quad\times \frac{1-bq^{2|\boldsymbol{j}|}}{(1-b)(bq^{|\boldsymbol{N}|+1})_{|\boldsymbol{j}|}} q^{|\boldsymbol{N}||\boldsymbol{j}|+\sum_{r=1}^{n}(r-1)j_r}\beta(\boldsymbol{j})\\
&=\sum_{\substack{0\le k_r\le N_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{(q^{-N_s}x_r/x_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times \frac{(1-aq^{2|\boldsymbol{k}|})(a)_{|\boldsymbol{k}|}}{(1-a)(aq^{|\boldsymbol{N}|+1})_{|\boldsymbol{k}|}} q^{|\boldsymbol{N}||\boldsymbol{k}|+\sum_{r=1}^{n}(r-1)k_r}\\
&\times \sum_{\substack{0\le j_r\le k_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} (q^{-k_s}x_r/x_s)_{j_r}\\
&\quad\times \frac{(aq^{|\boldsymbol{k}|},bq)_{|\boldsymbol{j}|}}{(b)_{2|\boldsymbol{j}|}}(-1)^{|\boldsymbol{j}|}q^{\sum_{r=1}^{n} rj_r+\binom{|\boldsymbol{j}|}{2}}\beta(\boldsymbol{j})\\
&\quad\times \sum_{\substack{0\le m_r\le k_r-j_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{m_r-m_s+j_r-j_s}x_r/x_s}{1-q^{j_r-j_s}x_r/x_s} q^{\sum_{r=1}^{n} rm_r}K(q^{\boldsymbol{m}+\boldsymbol{j}})\\
&\qquad\times \prod_{r,s=1}^{n} \frac{(q^{-k_s+j_r}x_r/x_s)_{m_r}}{(q^{1+j_r-j_s}x_r/x_s)_{m_r}} \frac{(aq^{|\boldsymbol{k}|+|\boldsymbol{j}|},bq^{1+|\boldsymbol{j}|})_{|\boldsymbol{m}|}}{(aq)_{|\boldsymbol{m}|+|\boldsymbol{j}|}(bq^{2|\boldsymbol{j}|+1})_{|\boldsymbol{m}|}}.
\end{aligned}
$$

This indicates that the identity (1.2) holds when $y_r = q^{N_r}$ for $r = 1, 2, \cdots, n$. Assume that both sides of (1.2) are analytic in $y_1, y_2, ..., y_n$ in a disk around the origin. Note that $N_r = 0, 1, \cdots$ for $r = 1, 2, \cdots, n$. Thus the identity (1.2) is also true in the same disk around the origin by analytic continuation. This completes the proof of Theorem 1.2. □

3. Proofs of Theorems 1.3–1.8

Proofs of Theorems 1.3–1.6. Taking $\beta(\boldsymbol{j}) = \prod_{r=1}^{n} \delta_{j_r,0}$ in (1.2) we have

$$
\begin{aligned}
K(\boldsymbol{y}) = &\sum_{\substack{k_r\ge 0\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{(x_r/x_sy_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times \frac{(1-aq^{2|\boldsymbol{k}|})(a)_{|\boldsymbol{k}|}}{(1-a)(aqy_1y_2\cdots y_n)_{|\boldsymbol{k}|}}(y_1y_2\cdots y_n)^{|\boldsymbol{k}|}q^{\sum_{r=1}^{n}(r-1)k_r}\\
&\times \sum_{\substack{0\le m_r\le k_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s} q^{\sum_{r=1}^{n} rm_r}K(q^{\boldsymbol{m}})\\
&\times \prod_{r,s=1}^{n} \frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}} \frac{(aq^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}}{(aq)_{|\boldsymbol{m}|}}.
\end{aligned}
\tag{3.1}
$$

Then (1.4) follows readily by setting

$$K(\boldsymbol{y}) = f(\alpha qy_1y_2\cdots y_n)\frac{(\alpha q,\alpha by_1y_2\cdots y_n)_\infty}{(\alpha b,\alpha qy_1y_2\cdots y_n)_\infty}$$

in (3.1), and then replacing a by α, y_r by a_r/q in the resulting identity.

Putting

$$K(\boldsymbol{y}) = f(\alpha q y_1, \cdots, \alpha q y_n) \frac{(\alpha q, \alpha b y_1 y_2 \cdots y_n)_\infty}{(\alpha b, \alpha q y_1 y_2 \cdots y_n)_\infty}$$

in (3.1), and then replacing a by α, y_r by a_r/q we easily obtain (1.5).

Take

$$K(\boldsymbol{y}) = f(\alpha q y_1 y_2 \cdots y_n) \frac{(\alpha q)_\infty}{(\alpha q y_1 y_2 \cdots y_n)_\infty} \prod_{r=1}^{n} \frac{(\alpha b x_r y_r)_\infty}{(\alpha b x_r)_\infty}$$

in (3.1), and then replace a by α, y_r by a_r/q we can get (1.6).

The identity (1.7) follows easily by substituting

$$K(\boldsymbol{y}) = f(\alpha q y_1, \cdots, \alpha q y_n) \frac{(\alpha q)_\infty}{(\alpha q y_1 y_2 \cdots y_n)_\infty} \prod_{r=1}^{n} \frac{(\alpha b x_r y_r)_\infty}{(\alpha b x_r)_\infty}$$

into (3.1), and then replacing a by α, y_r by a_r/q. □

Proofs of Theorems 1.7–1.8. Recall a multiple series expansion formula in [4, Theorem 2.2]:

$$\begin{aligned}
&K(\boldsymbol{y}) \sum_{\substack{j_r \geq 0 \\ r=1,\ldots,n}} \prod_{1 \leq r < s \leq n} \frac{1 - q^{j_r - j_s} x_r/x_s}{1 - x_r/x_s} \prod_{r,s=1}^{n} (x_r/x_s y_s)_{j_r} \\
&\times \prod_{r=1}^{n} \frac{1 - b x_r q^{j_r + |\boldsymbol{j}|}}{(1 - b x_r)(b q x_r y_r)_{|\boldsymbol{j}|}} (y_1 \cdots y_n)^{|\boldsymbol{j}|} \, q^{\sum_{r=1}^{n} (r-1) j_r} \beta(\boldsymbol{j}) \\
&= \sum_{\substack{k_r \geq 0 \\ r=1,\cdots,n}} \prod_{1 \leq r < s \leq n} \frac{1 - q^{k_r - k_s} x_r/x_s}{1 - x_r/x_s} \prod_{r,s=1}^{n} \frac{(x_r/x_s y_s)_{k_r}}{(q x_r/x_s)_{k_r}} \\
&\times \prod_{r=1}^{n} \frac{1 - a x_r q^{k_r + |\boldsymbol{k}|}}{1 - a x_r} \prod_{r=1}^{n} \frac{(a x_r)_{|\boldsymbol{k}|}}{(a q x_r y_r)_{|\boldsymbol{k}|}} (y_1 \cdots y_n)^{|\boldsymbol{k}|} \, q^{\sum_{r=1}^{n} (r-1) k_r} \\
&\times \sum_{\substack{0 \leq j_r \leq k_r \\ r=1,\cdots,n}} \prod_{1 \leq r < s \leq n} \frac{1 - q^{j_r - j_s} x_r/x_s}{1 - x_r/x_s} \prod_{r,s=1}^{n} \left(q^{-k_s} x_r/x_s\right)_{j_r} \\
&\times \prod_{r=1}^{n} \frac{\left(a x_r q^{|\boldsymbol{k}|}, b q x_r\right)_{j_r}}{(b x_r)_{j_r + |\boldsymbol{j}|}} (-1)^{|\boldsymbol{j}|} q^{\sum_{r=1}^{n} r j_r + \binom{|\boldsymbol{j}|}{2}} \beta(\boldsymbol{j}) \\
&\times \sum_{\substack{0 \leq m_r \leq k_r - j_r \\ r=1,\cdots,n}} \prod_{1 \leq r < s \leq n} \frac{1 - q^{j_r - j_s + m_r - m_s} x_r/x_s}{1 - q^{j_r - j_s} x_r/x_s} q^{\sum_{r=1}^{n} r m_r} K\left(q^{\boldsymbol{j} + \boldsymbol{m}}\right) \\
&\times \prod_{r,s=1}^{n} \frac{\left(q^{j_r - k_s} x_r/x_s\right)_{m_r}}{\left(q^{1 + j_r - j_s} x_r/x_s\right)_{m_r}} \prod_{r=1}^{n} \frac{\left(a x_r q^{j_r + |\boldsymbol{k}|}, b x_r q^{1 + j_r}\right)_{m_r}}{(a q x_r)_{j_r + m_r} \left(b x_r q^{1 + j_r + |\boldsymbol{j}|}\right)_{m_r}}.
\end{aligned} \tag{3.2}$$

Then the identity (1.8) follows readily by taking

$$K(\boldsymbol{y}) = f(\alpha q y_1 y_2 \cdots y_n) \prod_{r=1}^{n} \frac{(\alpha q x_r, \alpha b x_r y_r)_\infty}{(\alpha b x_r, \alpha q x_r y_r)_\infty},$$

and $\beta(\boldsymbol{j}) = \prod_{r=1}^{n} \delta_{j_r, 0}$ in (3.2) and then replacing a by α, y_r by a_r/q in the resulting formula.

Setting

$$K(\boldsymbol{y}) = f(\alpha q y_1, \cdots, \alpha q y_n) \prod_{r=1}^{n} \frac{(\alpha q x_r, \alpha b x_r y_r)_\infty}{(\alpha b x_r, \alpha q x_r y_r)_\infty},$$

and $\beta(\boldsymbol{j}) = \prod_{r=1}^{n} \delta_{j_r,0}$ in (3.2) and then substituting $a \mapsto \alpha, y_r \mapsto a_r/q$ into the resulting identity we easily obtain (1.9). □

4. An A_n Rogers' ${}_6\phi_5$ summation and an A_n extension of Sylvester's identity

In this section, we apply the identity (1.10) in Theorem 1.11 deduce an A_n Rogers' ${}_6\phi_5$ summation formula which is equivalent to the identity in [23, Theorem A.3]. From this summation formula, we derive an A_n extension of Sylvester's identity. This A_n extension is equivalent to the identity in [20, Theorem 5.19].

4.1. **An A_n Rogers' ${}_6\phi_5$ summation formula.** Milne proved the following well-known identity [21, Theorem 4.1]:

$$\begin{aligned}
&\frac{(c/a)_{|\boldsymbol{N}|}}{(c/ab)_{|\boldsymbol{N}|}} \prod_{i=1}^{n} \frac{(cx_i/b)_{N_i}}{(cx_i)_{N_i}} \\
&= \sum_{\substack{0\le y_i\le N_i \\ i=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{y_r-y_s}x_r/x_s}{1-x_r/x_s} \\
&\times \prod_{r,s=1}^{n} \frac{\left(q^{-N_s}x_r/x_s\right)_{y_r}}{\left(qx_r/x_s\right)_{y_r}} \prod_{i=1}^{n} \frac{(ax_i)_{y_i}}{(cx_i)_{y_i}} \\
&\times \frac{(b)_{|\boldsymbol{y}|}}{(abq^{1-|\boldsymbol{N}|}/c)_{|\boldsymbol{y}|}} q^{\sum_{r=1}^{n} r y_r},
\end{aligned} \tag{4.1}$$

where $N_1, N_2, \cdots, N_n$ are nonnegative integers. The $N_s \mapsto k_s, a \mapsto \alpha q^{|\boldsymbol{k}|}, b \mapsto \alpha bc/q, c \mapsto \alpha b$ case of (4.1) is

$$\begin{aligned}
&\sum_{\substack{0\le y_i\le k_i \\ i=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{y_r-y_s}x_r/x_s}{1-x_r/x_s} \\
&\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_r/x_s\right)_{y_r}}{\left(qx_r/x_s\right)_{y_r}} \prod_{i=1}^{n} \frac{(\alpha q^{|\boldsymbol{k}|}x_i)_{y_i}}{(\alpha b x_i)_{y_i}} \\
&\times \frac{(\alpha bc/q)_{|\boldsymbol{y}|}}{(\alpha\alpha b)_{|\boldsymbol{y}|}} q^{\sum_{r=1}^{n} r y_r} \\
&= \frac{(bq^{-|\boldsymbol{k}|})_{|\boldsymbol{k}|}}{(q^{1-|\boldsymbol{k}|}/\alpha c)_{|\boldsymbol{k}|}} \prod_{i=1}^{n} \frac{(qx_i/c)_{k_i}}{(\alpha b x_i)_{k_i}} \\
&= \frac{(q/b)_{|\boldsymbol{k}|}}{(\alpha c)_{|\boldsymbol{k}|}} \left(\frac{\alpha bc}{q}\right)^{|\boldsymbol{k}|} \prod_{i=1}^{n} \frac{(qx_i/c)_{k_i}}{(\alpha b x_i)_{k_i}},
\end{aligned}$$

where, in the last equality we have used [12, eq.(I.8)]. Letting $\beta = bc/q, \gamma = d$ in (1.10) of Theorem 1.11 and then applying the above identity in the inner sum on the right side of the resulting identity we get

Theorem 4.1. *Let $\alpha, a_1, \cdots, a_n, b, c$ be such that $|\alpha a_1 \cdots a_n bc/q^{1+n}| < 1$; or, $a_r = q^{N_r}$, for $r = 1, 2, \cdots, n$, and so the series terminate. Suppose that none of the denominators in the following identity vanishes, then*

$$\begin{aligned}
&\frac{(\alpha bc/q, \alpha a_1 \cdots a_n c q^{-n})_\infty}{(\alpha c, \alpha a_1 \cdots a_n b c q^{-(1+n)})_\infty} \prod_{r=1}^{n} \frac{(\alpha q x_r, \alpha a_r b x_r/q)_\infty}{(\alpha b x_r, \alpha a_r x_r)_\infty} \\
&= \sum_{\substack{k_r \geq 0 \\ r=1,\cdots,n}} \prod_{1 \leq r < s \leq n} \frac{1 - q^{k_r - k_s} x_r/x_s}{1 - x_r/x_s} \prod_{r,s=1}^{n} \frac{(q x_r/x_s a_s)_{k_r}}{(q x_r/x_s)_{k_r}} \\
&\times \prod_{r=1}^{n} \frac{1 - \alpha x_r q^{k_r + |\boldsymbol{k}|}}{1 - \alpha x_r} \prod_{r=1}^{n} \frac{(\alpha x_r)_{|\boldsymbol{k}|}}{(\alpha x_r a_r)_{|\boldsymbol{k}|}} q^{\sum_{r=1}^{n} (r-1) k_r} \\
&\times \frac{(q/b)_{|\boldsymbol{k}|}}{(\alpha c)_{|\boldsymbol{k}|}} \prod_{r=1}^{n} \frac{(q x_r/c)_{k_r}}{(\alpha b x_r)_{k_r}} \cdot \left(\frac{\alpha a_1 \cdots a_n bc}{q^{1+n}} \right)^{|\boldsymbol{k}|}.
\end{aligned} \tag{4.2}$$

This identity is an A_n extension of Rogers' non-terminating ${}_6\phi_5$ summation formula [12, eq.(II.20)]:

$$\begin{aligned}
&{}_6\phi_5 \left(\begin{matrix} \alpha, q\sqrt{\alpha}, -q\sqrt{\alpha}, q/a, q/b, q/c \\ \sqrt{\alpha}, -\sqrt{\alpha}, \alpha a, \alpha b, \alpha c \end{matrix} ; q, \frac{\alpha abc}{q^2} \right) \\
&= \frac{(\alpha q, \alpha ab/q, \alpha ac/q, \alpha bc/q)_\infty}{(\alpha a, \alpha b, \alpha c, \alpha abc/q^2)_\infty},
\end{aligned}$$

where $|\alpha abc/q^2| < 1$ and

$${}_{r+1}\phi_r \left(\begin{matrix} a_1, a_2, \cdots, a_{r+1} \\ b_1, b_2, \cdots, b_r \end{matrix} ; q, z \right) := \sum_{n=0}^{\infty} \frac{(a_1, a_2, \cdots, a_{r+1})_n}{(q, b_1, b_2, \cdots, b_r)_n} z^n.$$

Replacing q/a_r by a_r $(r = 1, 2, \cdots, n)$, q/b by b, q/c by c in the identity (4.2) of Theorem 4.1, we have

$$\begin{aligned}
&\frac{(\alpha q/bc, \alpha q/a_1 \cdots a_n c)_\infty}{(\alpha q/c, \alpha q/a_1 \cdots a_n bc)_\infty} \prod_{r=1}^{n} \frac{(\alpha q x_r, \alpha q x_r/a_r b)_\infty}{(\alpha q x_r/b, \alpha q x_r/a_r)_\infty} \\
&= \sum_{\substack{k_r \geq 0 \\ r=1,\cdots,n}} \prod_{1 \leq r < s \leq n} \frac{1 - q^{k_r - k_s} x_r/x_s}{1 - x_r/x_s} \prod_{r,s=1}^{n} \frac{(a_s x_r/x_s)_{k_r}}{(q x_r/x_s)_{k_r}} \\
&\times \prod_{r=1}^{n} \frac{1 - \alpha x_r q^{k_r + |\boldsymbol{k}|}}{1 - \alpha x_r} \prod_{r=1}^{n} \frac{(\alpha x_r)_{|\boldsymbol{k}|}}{(\alpha q x_r/a_r)_{|\boldsymbol{k}|}} q^{\sum_{r=1}^{n} (r-1) k_r} \\
&\times \frac{(b)_{|\boldsymbol{k}|}}{(\alpha q/c)_{|\boldsymbol{k}|}} \prod_{r=1}^{n} \frac{(c x_r)_{k_r}}{(\alpha q x_r/b)_{k_r}} \cdot \left(\frac{\alpha q}{a_1 \cdots a_n bc} \right)^{|\boldsymbol{k}|}.
\end{aligned}$$

This identity is equivalent to the A_n nonterminating ${}_6\phi_5$ summation formula in [23, Theorem A.3].

4.2. **An A_n extension of Sylvester's identity.** The Euler Pentagonal Number Theorem may be one of the most interesting formulas in basic hypergeometric series, which states that [12, eq.(8.10.10)]

$$(q)_\infty = 1 + \sum_{n=1}^{\infty} (-1)^n (q^{n(3n-1)/2} + q^{n(3n+1)/2}). \tag{4.3}$$

Sylvester gave the following beautiful refinement [3, eq.(9.2.3)] of Euler's Pentagonal Number Theorem.

Theorem 4.2. (Sylvester's identity) *We have*

$$\frac{1}{(\alpha)_\infty}\sum_{k=0}^{\infty}(-\alpha)^k q^{k^2+\binom{k}{2}}(1-\alpha q^{2k})\frac{(\alpha)_k}{(q)_k}=1.$$

In this subsection we will derive the following A_n extension of Sylvester's identity.

Theorem 4.3. *Suppose that none of the denominators in the following identity vanishes, then*

$$\begin{aligned}1=&\frac{1}{\prod_{r=1}^n(\alpha x_r)_\infty}\sum_{\substack{k_r\geq 0\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n(qx_r/x_s)_{k_r}^{-1}\\ &\times\prod_{r=1}^n\{(1-\alpha x_r q^{k_r+|\boldsymbol{k}|})(\alpha x_r)_{|\boldsymbol{k}|}\}\prod_{r=1}^n x_r^{(n+1)k_r-|\boldsymbol{k}|}\\ &\times(-1)^{n|\boldsymbol{k}|}\alpha^{|\boldsymbol{k}|}q^{(n+1)\sum_{r=1}^n\binom{k_r}{2}+\binom{|\boldsymbol{k}|}{2}+\sum_{r=1}^n rk_r}.\end{aligned}$$

Proof. The result follows readily by multiplying both sides of the identity in Theorem 4.1 by $\prod_{r=1}^n(\alpha q x_r)_\infty^{-1}$, setting $a_r\to 0\,(r=1,2,\cdots,n), b\to 0, c\to 0$ in the resulting identity and then using the limiting relation $\lim_{a\to 0}(u/a)_n a^n=(-u)^n q^{\binom{n}{2}}$. □

Actually, this A_n extension is equivalent to the identity in [20, Theorem 5.19]. Replacing a by α, z_i/z_n by x_i on both sides of [20, Theorem 5.19] and then multiplying both sides of the resulting identity by $\prod_{i=1}^n(\alpha q x_i)_\infty^{-1}$ we can easily obtain the identity in Theorem 4.3.

Take $\alpha=1$ in the identity of Theorem 4.3 and then multiply both sides of the resulting identity by $\prod_{r=1}^n(qx_r)_\infty$. We get

$$\begin{aligned}\prod_{r=1}^n(qx_r)_\infty=&\frac{1}{\prod_{r=1}^n(1-x_r)}\sum_{\substack{k_r\geq 0\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n(qx_r/x_s)_{k_r}^{-1}\\ &\times\prod_{r=1}^n\{(1-x_r q^{k_r+|\boldsymbol{k}|})(x_r)_{|\boldsymbol{k}|}\}\prod_{r=1}^n x_r^{(n+1)k_r-|\boldsymbol{k}|}\\ &\times(-1)^{n|\boldsymbol{k}|}q^{(n+1)\sum_{r=1}^n\binom{k_r}{2}+\binom{|\boldsymbol{k}|}{2}+\sum_{r=1}^n rk_r}.\end{aligned}$$

This identity can be regarded as an A_n extension of the Euler Pentagonal Number Theorem (4.3).

5. Multiple expansion formulas for $(q)_\infty^m, \pi_q$ and $1/\pi_q$

In this section, we first employ the identity (1.4) in Theorem 1.3 to deduce some multiple expansion formulas for $(q)_\infty^m$ and then apply the identity in Theorem 1.9 to derive several multiple expansion formulas for π_q and $1/\pi_q$.

5.1. **Multiple expansion formulas for $(q)_\infty^m$.** For any integer m, we will give some multiple expansion formulas for $(q)_\infty^m$. These formulas can be divided into two groups: the first one involves $(q)_\infty^m$ with positive integer m while the second one involves $(q)_\infty^m$ with non-positive integer m. All of these expansion formulas are A_n extensions of some well-known identities of Liu [18].

The first group of multiple expansion formulas for $(q)_\infty^m$ is as follows.

Theorem 5.1. *Let m be a nonnegative integer. Then*

$$\begin{aligned}(q)_\infty^{m+1} = &\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}} \prod_{1\leq r<s\leq n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-q^{2|\boldsymbol{k}|+1})(q)_{|\boldsymbol{k}|}}{\prod_{r,s=1}^n (qx_r/x_s)_{k_r}}\\ &\times(-1)^{n|\boldsymbol{k}|}q^{\sum_{r=1}^n (r-1)k_r+n\sum_{i=1}^n\binom{k_i}{2}}\prod_{r=1}^n x_r^{nk_r-|\boldsymbol{k}|}\\ &\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\\ &\times\prod_{r,s=1}^n\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\cdot(q^{|\boldsymbol{k}|+1})_{|\boldsymbol{m}|}(q)_{|\boldsymbol{m}|}^m q^{\sum_{r=1}^n rm_r}\end{aligned}\tag{5.1}$$

and

$$\begin{aligned}(q)_\infty^{m+1} = 1+&\sum_{\substack{k_1,k_2,\cdots,k_n\geq 0\\ |\boldsymbol{k}|\geq 1}} \prod_{1\leq r<s\leq n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-q^{2|\boldsymbol{k}|})(q)_{|\boldsymbol{k}|-1}}{\prod_{r,s=1}^n (qx_r/x_s)_{k_r}}\\ &\times(-1)^{n|\boldsymbol{k}|}q^{\sum_{r=1}^n (r-1)k_r+n\sum_{i=1}^n\binom{k_i}{2}}\prod_{r=1}^n x_r^{nk_r-|\boldsymbol{k}|}\\ &\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\\ &\times\prod_{r,s=1}^n\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\cdot(q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}(q)_{|\boldsymbol{m}|}^m\cdot q^{\sum_{r=1}^n rm_r}.\end{aligned}\tag{5.2}$$

Proof. If

$$f(y)=\prod_{j=1}^m\frac{(b_jy/q)_\infty}{(c_jy/q)_\infty},$$

then

$$f(\alpha a_1\cdots a_nq^{1-n})=\prod_{j=1}^m\frac{(\alpha a_1\cdots a_nb_j/q^n)_\infty}{(\alpha a_1\cdots a_nc_j/q^n)_\infty},\tag{5.3}$$

and

$$f(\alpha q^{1+|\boldsymbol{m}|})=\prod_{j=1}^m\frac{(\alpha b_jq^{|\boldsymbol{m}|})_\infty}{(\alpha c_jq^{|\boldsymbol{m}|})_\infty}=\prod_{j=1}^m\frac{(\alpha b_j)_\infty}{(\alpha c_j)_\infty}\prod_{j=1}^m\frac{(\alpha c_j)_{|\boldsymbol{m}|}}{(\alpha b_j)_{|\boldsymbol{m}|}}.\tag{5.4}$$

Setting

$$f(y)=\prod_{j=1}^m\frac{(b_jy/q)_\infty}{(c_jy/q)_\infty}$$

in the identity (1.4) of Theorem 1.3, substituting (5.3) and (5.4) into the resulting identity and then taking $a_1 = a_2 = \cdots = a_n = 0$ we get

$$\begin{aligned}
&\frac{(\alpha q)_\infty}{(\alpha b)_\infty}\prod_{j=1}^{m}\frac{(\alpha c_j)_\infty}{(\alpha b_j)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-\alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)\prod_{r,s=1}^{n}(qx_r/x_s)_{k_r}}\\
&\times(-1)^{n|\boldsymbol{k}|}q^{\sum_{r=1}^{n}(r-1)k_r+n\sum_{i=1}^{n}\binom{k_i}{2}}\prod_{r=1}^{n}x_r^{nk_r-|\boldsymbol{k}|} \qquad (5.5)\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\\
&\times\frac{(\alpha q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}}{(\alpha b)_{|\boldsymbol{m}|}}\prod_{j=1}^{m}\frac{(\alpha c_j)_{|\boldsymbol{m}|}}{(\alpha b_j)_{|\boldsymbol{m}|}}q^{\sum_{r=1}^{n}rm_r}.
\end{aligned}$$

The $b = b_1 = \cdots = b_m = 0$ case of (5.5) is

$$\begin{aligned}
&(\alpha q)_\infty\prod_{j=1}^{m}(\alpha c_j)_\infty\\
&=\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-\alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)\prod_{r,s=1}^{n}(qx_r/x_s)_{k_r}}\\
&\times(-1)^{n|\boldsymbol{k}|}q^{\sum_{r=1}^{n}(r-1)k_r+n\sum_{i=1}^{n}\binom{k_i}{2}}\prod_{r=1}^{n}x_r^{nk_r-|\boldsymbol{k}|} \qquad (5.6)\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\\
&\times(\alpha q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}\prod_{j=1}^{m}(\alpha c_j)_{|\boldsymbol{m}|}\cdot q^{\sum_{r=1}^{n}rm_r}.
\end{aligned}$$

Then the identities (5.1) and (5.2) follow by setting $\alpha = q, c_1 = c_2 = \cdots = c_m = 1$ and $\alpha \to 1, c_1 = c_2 = \cdots = c_m = q$ in (5.6) respectively and then simplifying. □

The identity (5.1) in Theorem 5.1 is an A_n extension of the well-known identity in [18, Proposition 3.1] while (5.2) can be regarded as an A_n extension of the identity in [18, Proposition 3.2]. The identities (1.13) and (1.14) are the special $m = 2$ case of (5.1) and (5.2) respectively.

The second group of multiple expansion formulas for $(q)_\infty^m$ is showed in the following theorem.

Theorem 5.2. *Let m be a nonnegative integer. Then*

$$\begin{aligned}\frac{1}{(q)_\infty^m} &= \sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}} \prod_{1\leq r<s\leq n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-q^{2|\boldsymbol{k}|+1})(q)_{|\boldsymbol{k}|}}{\prod_{r,s=1}^n (qx_r/x_s)_{k_r}}\\ &\quad\times(-1)^{n|\boldsymbol{k}|}q^{\sum_{r=1}^n (r-1)k_r+n\sum_{i=1}^n\binom{k_i}{2}}\prod_{r=1}^n x_r^{nk_r-|\boldsymbol{k}|}\\ &\quad\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\\ &\quad\times\prod_{r,s=1}^n\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\cdot\frac{(q^{|\boldsymbol{k}|+1})_{|\boldsymbol{m}|}}{(q)_{|\boldsymbol{m}|}^{m+1}}q^{\sum_{r=1}^n rm_r},\end{aligned}\tag{5.7}$$

and

$$\begin{aligned}\frac{1}{(q)_\infty^m} &= 1+\sum_{\substack{k_1,k_2,\cdots,k_n\geq 0\\ |\boldsymbol{k}|\geq 1}} \prod_{1\leq r<s\leq n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-q^{2|\boldsymbol{k}|})(q)_{|\boldsymbol{k}|-1}}{\prod_{r,s=1}^n (qx_r/x_s)_{k_r}}\\ &\quad\times(-1)^{n|\boldsymbol{k}|}q^{\sum_{r=1}^n (r-1)k_r+n\sum_{i=1}^n\binom{k_i}{2}}\prod_{r=1}^n x_r^{nk_r-|\boldsymbol{k}|}\\ &\quad\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\\ &\quad\times\prod_{r,s=1}^n\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\cdot\frac{(q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}}{(q)_{|\boldsymbol{m}|}^{m+1}}q^{\sum_{r=1}^n rm_r}.\end{aligned}\tag{5.8}$$

Proof. Putting $c_1=\cdots=c_m=0$ in (5.5) we arrive at

$$\begin{aligned}&\frac{(\alpha q)_\infty}{(\alpha b)_\infty\prod_{j=1}^m(\alpha b_j)_\infty}\\ &=\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}} \prod_{1\leq r<s\leq n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-\alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)\prod_{r,s=1}^n (qx_r/x_s)_{k_r}}\\ &\quad\times(-1)^{n|\boldsymbol{k}|}q^{\sum_{r=1}^n (r-1)k_r+n\sum_{i=1}^n\binom{k_i}{2}}\prod_{r=1}^n x_r^{nk_r-|\boldsymbol{k}|}\\ &\quad\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}}\\ &\quad\times\frac{(\alpha q^{|\boldsymbol{k}|})_{|\boldsymbol{m}|}}{(\alpha b)_{|\boldsymbol{m}|}\prod_{j=1}^m(\alpha b_j)_{|\boldsymbol{m}|}}q^{\sum_{r=1}^n rm_r}.\end{aligned}\tag{5.9}$$

We take $\alpha=q, b=b_1=\cdots=b_m=1$ and $\alpha\to 1, b=b_1=\cdots=b_m=q$ in (5.9) to obtain (5.7) and (5.8) respectively. □

The identity (5.7) is an A_n extension of the identity in [18, Proposition 3.3] while (5.8) can be considered as an A_n extension of the identity in [18, Proposition 3.4]. The identity (1.16) is the special $m=2$ case of (5.8).

5.2. **Multiple expansion formulas for π_q and $1/\pi_q$.** In [12], W. Gosper introduced a q-analogue of the irrational number π :

$$\pi_q := (1-q^2)q^{1/4}\frac{(q^2;q^2)_\infty^2}{(q;q^2)_\infty^2}.$$

It satisfies the following limiting relation:

$$\lim_{q\to 1}\pi_q = \pi.$$

In this subsection, we give some multiple expansion formulas for π_q and $1/\pi_q$.

Theorem 5.3. *We have*

$$\begin{aligned}\pi_q = (1-q^2)&\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{2k_r-2k_s}x_r/x_s}{1-x_r/x_s}\prod_{r=1}^n x_r^{nk_r-|\boldsymbol{k}|}\\
&\times\frac{(1-q^{4|\boldsymbol{k}|+2})(q^2;q^2)_{|\boldsymbol{k}|}}{\prod_{r,s=1}^n(q^2x_r/x_s;q^2)_{k_r}}\cdot(-1)^{n|\boldsymbol{k}|}q^{2\sum_{r=1}^n(r-1)k_r+2n\sum_{r=1}^n\binom{k_r}{2}+\frac{1}{4}}\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{2m_r-2m_s}x_r/x_s}{1-x_r/x_s}\\
&\times\prod_{r,s=1}^n\frac{(q^{-2k_s}x_r/x_s;q^2)_{m_r}}{(q^2x_r/x_s;q^2)_{m_r}}\cdot\frac{(q^{2|\boldsymbol{k}|+2},q^2;q^2)_{|\boldsymbol{m}|}}{(q,q;q^2)_{|\boldsymbol{m}|}}q^{2\sum_{r=1}^n rm_r},\end{aligned}$$

and

$$\begin{aligned}\pi_q = (1-q^2)&q^{1/4}\\
&+(1-q^2)\sum_{\substack{k_r\geq 0\\ |\boldsymbol{k}|\geq 1}}\prod_{1\leq r<s\leq n}\frac{1-q^{2k_r-2k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-q^{4|\boldsymbol{k}|})(q^2;q^2)_{|\boldsymbol{k}|-1}}{\prod_{r,s=1}^n(q^2x_r/x_s;q^2)_{k_r}}\\
&\times(-1)^{n|\boldsymbol{k}|}q^{2\sum_{r=1}^n(r-1)k_r+2n\sum_{r=1}^n\binom{k_r}{2}+\frac{1}{4}}\prod_{r=1}^n x_r^{nk_r-|\boldsymbol{k}|}\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{2m_r-2m_s}x_r/x_s}{1-x_r/x_s}\\
&\times\prod_{r,s=1}^n\frac{(q^{-2k_s}x_r/x_s;q^2)_{m_r}}{(q^2x_r/x_s;q^2)_{m_r}}\cdot\frac{(q^{2|\boldsymbol{k}|},q^2;q^2)_{|\boldsymbol{m}|}}{(q,q;q^2)_{|\boldsymbol{m}|}}q^{2\sum_{r=1}^n rm_r}.\end{aligned}$$

Proof. Replacing q by q^2 in the identity of Theorem 1.9 and then taking $a_1=\cdots=a_n=0$ and $\gamma=d$ we have

$$\begin{aligned}&\frac{(\alpha q^2,\alpha\beta;q^2)_\infty}{(\alpha b,\alpha c;q^2)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{2k_r-2k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-\alpha q^{4|\boldsymbol{k}|})(\alpha;q^2)_{|\boldsymbol{k}|}}{(1-\alpha)\prod_{r,s=1}^n(q^2x_r/x_s;q^2)_{k_r}}\end{aligned}\tag{5.10}$$

$$\times (-1)^{n|\boldsymbol{k}|}q^{2\sum_{r=1}^{n}(r-1)k_r+2n\sum_{r=1}^{n}\binom{k_r}{2}}\prod_{r=1}^{n}x_r^{nk_r-|\boldsymbol{k}|}$$
$$\times \sum_{\substack{0\le m_r\le k_r\\ r=1,2,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{2m_r-2m_s}x_r/x_s}{1-x_r/x_s}$$
$$\times \prod_{r,s=1}^{n}\frac{(q^{-2k_s}x_r/x_s;q^2)_{m_r}}{(q^2x_r/x_s;q^2)_{m_r}}\cdot\frac{(\alpha q^{2|\boldsymbol{k}|},\alpha\beta;q^2)_{|\boldsymbol{m}|}}{(\alpha b,\alpha c;q^2)_{|\boldsymbol{m}|}}q^{2\sum_{r=1}^{n}rm_r}.$$

Then the first identity follows easily by setting $\alpha\mapsto q^2,\beta\mapsto 1,b\mapsto q^{-1},c\mapsto q^{-1}$ in (5.10) and then multiplying both sides of the resulting identity by $(1-q^2)q^{1/4}$.

We put $\alpha\mapsto 1,\beta\mapsto q^2,b\mapsto q,c\mapsto q$ in (5.10) and then multiply both sides of the resulting identity by $(1-q^2)q^{1/4}$ to get the second identity. □

Theorem 5.4. *We have*

$$1/\pi_q=\frac{1}{1-q^2}\sum_{\substack{k_r\ge 0\\ r=1,2,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{2k_r-2k_s}x_r/x_s}{1-x_r/x_s}\cdot\frac{(1-q^{4|\boldsymbol{k}|+1})(q;q^2)_{|\boldsymbol{k}|}}{\prod_{r,s=1}^{n}(q^2x_r/x_s;q^2)_{k_r}}$$
$$\times (-1)^{n|\boldsymbol{k}|}q^{2\sum_{r=1}^{n}(r-1)k_r+2n\sum_{r=1}^{n}\binom{k_r}{2}-\frac{1}{4}}\prod_{r=1}^{n}x_r^{nk_r-|\boldsymbol{k}|}$$
$$\times \sum_{\substack{0\le m_r\le k_r\\ r=1,2,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{2m_r-2m_s}x_r/x_s}{1-x_r/x_s}$$
$$\times \prod_{r,s=1}^{n}\frac{(q^{-2k_s}x_r/x_s;q^2)_{m_r}}{(q^2x_r/x_s;q^2)_{m_r}}\cdot\frac{(q^{2|\boldsymbol{k}|+1},q;q^2)_{|\boldsymbol{m}|}}{(q^2,q^2;q^2)_{|\boldsymbol{m}|}}q^{2\sum_{r=1}^{n}rm_r},$$

Proof. This identity follows readily by taking $\alpha\mapsto q,\beta\mapsto 1,b\mapsto q,c\mapsto q$ in (5.10) and then multiplying both sides of the resulting identity by $(1-q^2)^{-1}q^{-1/4}$. □

6. Two A_n extensions of the Rogers-Fine identity

The Rogers-Fine identity [11, eq.(14.1)] may be one of the fundamental formulas in basic hypergeometric series. It can be stated in the following form:

$$\sum_{n=0}^{\infty}\frac{(a)_n}{(b)_n}z^n=\sum_{n=0}^{\infty}\frac{(1-azq^{2n})(a,azq/b)_n}{(b)_n(z)_{n+1}}(bz)^nq^{n^2-n},$$

where $|z|<1$. This identity was first proved by L.J. Rogers [25] and also discovered by N.J. Fine [11] independently. In [11], Fine used it to deduce many important results in partitions, modular equations and mock theta functions. In [16], Liu gave an alternate proof of this result by using an important expansion formula [16, Theorem 2] for q-series.

In this section, we will establish two A_n extensions of the Rogers-Fine identity by applying the identity (1.2) in Theorem 1.2.

Theorem 6.1. (First A_n extension of the Rogers-Fine identity) *Let a,z be such that $|z/a^{n-1}|<1$. Suppose that none of the denominators in the following identity*

vanishes, then

$$\begin{aligned}
&\sum_{\substack{j_r\geq 0\\ r=1,2,\cdots,n}} \prod_{1\leq r<s\leq n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{(ax_r/x_s)_{j_r}}{(qx_r/x_s)_{j_r}}\\
&\times \frac{(q)_{|\boldsymbol{j}|}}{(b)_{|\boldsymbol{j}|}}(z/a^{n-1})^{|\boldsymbol{j}|}q^{\sum_{r=1}^{n}(r-1)j_r}\\
&= \sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}} \prod_{1\leq r<s\leq n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{(ax_r/x_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times \frac{(1-azq^{2|\boldsymbol{k}|})(q,azq/b)_{|\boldsymbol{k}|}}{(b)_{|\boldsymbol{k}|}(z/a^{n-1})_{|\boldsymbol{k}|+1}}(bz/a^{n-1})^{|\boldsymbol{k}|}q^{\sum_{r=1}^{n}(r-1)k_r+2\binom{|\boldsymbol{k}|}{2}}.
\end{aligned}$$

Proof. We recall a $U(n+1)$ q-Gauss summation in [21, Theorem 7.6]:

$$\begin{aligned}
&\sum_{\substack{y_k\geq 0\\ k=1,2,\cdots,n}} \prod_{1\leq r<s\leq n} \frac{1-\frac{x_r}{x_s}q^{y_r-y_s}}{1-\frac{x_r}{x_s}} \prod_{r,s=1}^{n} \frac{\left(\frac{x_r}{x_s}a_s\right)_{y_r}}{\left(q\frac{x_r}{x_s}\right)_{y_r}}\\
&\quad\times \frac{(b)_{|\boldsymbol{y}|}}{(c)_{|\boldsymbol{y}|}}\left(\frac{c}{a_1\cdots a_n b}\right)^{|\boldsymbol{y}|} q^{y_2+2y_3+\cdots+(n-1)y_n}\\
&= \frac{(c/b, c/a_1\cdots a_n)_\infty}{(c, c/a_1\cdots a_n b)_\infty},
\end{aligned}$$

where $|c|<|a_1\cdots a_n b|$. If $a_i=q^{-N_i}$ for $i=1,2,\cdots,n$, then the above identity becomes

$$\begin{aligned}
&\sum_{\substack{0\leq y_k\leq N_k\\ k=1,2,\cdots,n}} \prod_{1\leq r<s\leq n} \frac{1-\frac{x_r}{x_s}q^{y_r-y_s}}{1-\frac{x_r}{x_s}} \prod_{r,s=1}^{n} \frac{\left(\frac{x_r}{x_s}q^{-N_s}\right)_{y_r}}{\left(q\frac{x_r}{x_s}\right)_{y_r}}\\
&\quad\times \frac{(b)_{|\boldsymbol{y}|}}{(c)_{|\boldsymbol{y}|}}\left(\frac{cq^{|\boldsymbol{N}|}}{b}\right)^{|\boldsymbol{y}|} q^{y_2+2y_3+\cdots+(n-1)y_n}\\
&= \frac{(c/b)_{|\boldsymbol{N}|}}{(c)_{|\boldsymbol{N}|}}.
\end{aligned}$$

Applying

$$(A;q)_n=(A^{-1};q^{-1})_n(-A)^n q^{\binom{n}{2}}$$

in this formula, replacing each parameter, including q and $x_1,\cdots,x_n$, by its reciprocal and then simplifying we get

$$\begin{aligned}
&\sum_{\substack{0\leq y_r\leq N_r\\ r=1,2,\cdots,n}} \prod_{1\leq r<s\leq n} \frac{1-\frac{x_r}{x_s}q^{y_r-y_s}}{1-\frac{x_r}{x_s}}\\
&\quad\times \prod_{r,s=1}^{n} \frac{\left(\frac{x_r}{x_s}q^{-N_s}\right)_{y_r}}{\left(q\frac{x_r}{x_s}\right)_{y_r}} \frac{(b)_{|\boldsymbol{y}|}}{(c)_{|\boldsymbol{y}|}} q^{y_1+2y_2+\cdots+ny_n}\\
&= \frac{(c/b)_{|\boldsymbol{N}|}}{(c)_{|\boldsymbol{N}|}} b^{|\boldsymbol{N}|}.
\end{aligned} \tag{6.1}$$

Taking $N_r \mapsto k_r - j_r, x_r \mapsto q^{j_r}x_r, b \mapsto azq^{|\boldsymbol{k}|+|\boldsymbol{j}|}, c \mapsto azq^{|\boldsymbol{j}|}$ in (6.1) we have

$$\begin{aligned}
&\sum_{\substack{0\le m_r\le k_r-j_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{m_r-m_s+j_r-j_s}x_r/x_s}{1-q^{j_r-j_s}x_r/x_s} q^{\sum_{r=1}^n rm_r}\\
&\times \prod_{r,s=1}^{n} \frac{(q^{-k_s+j_r}x_r/x_s)_{m_r}}{(q^{1+j_r-j_s}x_r/x_s)_{m_r}} \frac{(azq^{|\boldsymbol{k}|+|\boldsymbol{j}|})_{|\boldsymbol{m}|}}{(azq^{|\boldsymbol{j}|})_{|\boldsymbol{m}|}}\\
&= \frac{(q^{-|\boldsymbol{k}|})_{|\boldsymbol{k}|-|\boldsymbol{j}|}}{(azq^{|\boldsymbol{j}|})_{|\boldsymbol{k}|-|\boldsymbol{j}|}} (azq^{|\boldsymbol{k}|+|\boldsymbol{j}|})^{|\boldsymbol{k}|-|\boldsymbol{j}|}\\
&= \frac{(q)_{|\boldsymbol{k}|}(-az)^{|\boldsymbol{k}|-|\boldsymbol{j}|}}{(azq^{|\boldsymbol{j}|})_{|\boldsymbol{k}|-|\boldsymbol{j}|}(q)_{|\boldsymbol{j}|}} q^{\binom{|\boldsymbol{k}|}{2}-\binom{|\boldsymbol{j}|}{2}},
\end{aligned} \tag{6.2}$$

where, in the last equality we have used the identity [12, eq.(I.14)].

Setting $N_r \mapsto k_r, b \mapsto azq^{|\boldsymbol{k}|}, c \mapsto b$ in (6.1) we see that

$$\begin{aligned}
&\sum_{\substack{0\le j_r\le k_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s}\\
&\times \prod_{r,s=1}^{n} \frac{(q^{-k_s}x_r/x_s)_{j_r}}{(qx_r/x_s)_{j_r}} \frac{(azq^{|\boldsymbol{k}|})_{|\boldsymbol{j}|}}{(b)_{|\boldsymbol{j}|}} q^{\sum_{r=1}^n rj_r}\\
&= \frac{(bq^{-|\boldsymbol{k}|}/az)_{|\boldsymbol{k}|}}{(b)_{|\boldsymbol{k}|}} (azq^{|\boldsymbol{k}|})^{|\boldsymbol{k}|}\\
&= \frac{(azq/b)_{|\boldsymbol{k}|}}{(b)_{|\boldsymbol{k}|}} (-b)^{|\boldsymbol{k}|} q^{\binom{|\boldsymbol{k}|}{2}},
\end{aligned} \tag{6.3}$$

where, in the last equality we applied the formula [12, eq.(I.8)].

Replacing a by az in the $b = 0$ case of (1.2) in Theorem 1.2 and then taking

$$K(y) = \frac{1-azy_1\cdots y_n}{1-az}$$

and

$$\beta(\boldsymbol{j}) = \frac{(q)_{|\boldsymbol{j}|}(az)^{|\boldsymbol{j}|}}{(b)_{|\boldsymbol{j}|}\prod_{r,s=1}^n (qx_r/x_s)_{j_r}}$$

we obtain

$$\begin{aligned}
&\frac{1-azy_1\cdots y_n}{1-az} \sum_{\substack{j_r\ge 0\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s}\\
&\quad\times \prod_{r,s=1}^{n} \frac{(x_r/x_sy_s)_{j_r}}{(qx_r/x_s)_{j_r}} \frac{(q)_{|\boldsymbol{j}|}}{(b)_{|\boldsymbol{j}|}} (azy_1y_2\cdots y_n)^{|\boldsymbol{j}|} q^{\sum_{r=1}^n (r-1)j_r}\\
&= \sum_{\substack{k_r\ge 0\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{(x_r/x_sy_s)_{k_r}}{(qx_r/x_s)_{k_r}}
\end{aligned}$$

$$\times \frac{(1-azq^{2|\boldsymbol{k}|})(az)_{|\boldsymbol{k}|}}{(1-az)(azqy_1y_2\cdots y_n)_{|\boldsymbol{k}|}}(y_1y_2\cdots y_n)^{|\boldsymbol{k}|}q^{\sum_{r=1}^n(r-1)k_r}$$
$$\times \sum_{\substack{0\le j_r\le k_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n}\frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{(q^{-k_s}x_r/x_s)_{j_r}}{(qx_r/x_s)_{j_r}}$$
$$\times \frac{(azq^{|\boldsymbol{k}|},q)_{|\boldsymbol{j}|}}{(b,az)_{|\boldsymbol{j}|}}(-az)^{|\boldsymbol{j}|}q^{\sum_{r=1}^n rj_r+\binom{|\boldsymbol{j}|}{2}}$$
$$\times \sum_{\substack{0\le m_r\le k_r-j_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n}\frac{1-q^{m_r-m_s+j_r-j_s}x_r/x_s}{1-q^{j_r-j_s}x_r/x_s}q^{\sum_{r=1}^n rm_r}$$
$$\times \prod_{r,s=1}^n\frac{(q^{-k_s+j_r}x_r/x_s)_{m_r}}{(q^{1+j_r-j_s}x_r/x_s)_{m_r}}\frac{(azq^{|\boldsymbol{k}|+|\boldsymbol{j}|})_{|\boldsymbol{m}|}}{(azq^{|\boldsymbol{j}|})_{|\boldsymbol{m}|}}.$$

Substituting (6.2) into the inner-most sum $\sum_{\substack{0\le m_r\le k_r-j_r\\ r=1,2,\cdots,n}}$ on the right side of the above identity and then applying (6.3) in the second sum $\sum_{\substack{0\le j_r\le k_r\\ r=1,2,\cdots,n}}$ on the right side of the resulting identity yields

$$\frac{1-azy_1\cdots y_n}{1-az}\sum_{\substack{j_r\ge 0\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n}\frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{(x_r/x_sy_s)_{j_r}}{(qx_r/x_s)_{j_r}}$$
$$\times \frac{(q)_{|\boldsymbol{j}|}}{(b)_{|\boldsymbol{j}|}}(azy_1y_2\cdots y_n)^{|\boldsymbol{j}|}q^{\sum_{r=1}^n(r-1)j_r}$$
$$= \sum_{\substack{k_r\ge 0\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{(x_r/x_sy_s)_{k_r}}{(qx_r/x_s)_{k_r}}$$
$$\times \frac{(1-azq^{2|\boldsymbol{k}|})(q,azq/b)_{|\boldsymbol{k}|}}{(1-az)(b,azqy_1y_2\cdots y_n)_{|\boldsymbol{k}|}}(abzy_1y_2\cdots y_n)^{|\boldsymbol{k}|}q^{\sum_{r=1}^n(r-1)k_r+2\binom{|\boldsymbol{k}|}{2}}.$$

Then the result follows by multiplying both sides of this identity by $\frac{1-az}{1-azy_1\cdots y_n}$ and then setting $y_1=y_2=\cdots=y_n=1/a$. □

A second A_n extension of the Rogers-Fine identity is as follows.

Theorem 6.2. (Second A_n extension of the Rogers-Fine identity) *Let a, z be such that $|z/a^{n-1}|<1$. Suppose that none of the denominators in the following identity vanishes, then*

$$\sum_{\substack{j_r\ge 0\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n}\frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{(ax_r/x_s)_{j_r}}{(qx_r/x_s)_{j_r}}$$
$$\times \frac{(q)_{|\boldsymbol{j}|}}{\prod_{i=1}^n(bx_i)_{j_i}}(z/a^{n-1})^{|\boldsymbol{j}|}q^{\sum_{r=1}^n(r-1)j_r}$$
$$= \sum_{\substack{k_r\ge 0\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{(ax_r/x_s)_{k_r}}{(qx_r/x_s)_{k_r}}$$

$$\times \frac{(1-azq^{2|\boldsymbol{k}|})(q)_{|\boldsymbol{k}|}}{(z/a^{n-1})_{|\boldsymbol{k}|+1}} \prod_{i=1}^{n} \frac{(azq^{1+|\boldsymbol{k}|-k_i}/bx_i)_{k_i}}{(bx_i)_{k_i}}$$
$$\times (bz/a^{n-1})^{|\boldsymbol{k}|} q^{\sum_{r=1}^{n}(r-1)k_r+\binom{|\boldsymbol{k}|}{2}+\sum_{i=1}^{n}\binom{k_i}{2}} \prod_{i=1}^{n} x_i^{k_i}.$$

Proof. We first take $a \mapsto az, b \mapsto 0, K(y) \mapsto \frac{1-azy_1\cdots y_n}{1-az}$ and

$$\beta(\boldsymbol{j}) \mapsto \frac{(q)_{|\boldsymbol{j}|}(az)^{|\boldsymbol{j}|}}{\prod_{i=1}^{n}(bx_i)_{j_i}\prod_{r,s=1}^{n}(qx_r/x_s)_{j_r}}$$

in (1.2) of Theorem 1.2, then apply (6.2) and the identity in [21, Theorem 5.10] and finally set $y_i \mapsto 1/a$ for $i = 1, 2, \cdots, n$ to obtain this A_n extension of the Rogers-Fine identity. □

7. An A_n extension of Liu's extension of Rogers' non-terminating ${}_6\phi_5$ summation

Rogers' non-terminating ${}_6\phi_5$ summation [12, eq.(II.20)] may be one of the most important formulas in q-series. It can be stated in the following equivalent form.

Theorem 7.1. *For $|\alpha abc/q^2| < 1$, we have*

$${}_6\phi_5\left(\begin{matrix}\alpha, q\sqrt{\alpha}, -q\sqrt{\alpha}, q/a, q/b, q/c \\ \sqrt{\alpha}, -\sqrt{\alpha}, \alpha a, \alpha b, \alpha c\end{matrix}; q, \frac{\alpha abc}{q^2}\right)$$
$$= \frac{(\alpha q, \alpha ab/q, \alpha ac/q, \alpha bc/q)_\infty}{(\alpha a, \alpha b, \alpha c, \alpha abc/q^2)_\infty}.$$

Using the operator method, Liu proved the following extension [17, Theorem 3] of the Rogers non-terminating ${}_6\phi_5$ summation formula.

Theorem 7.2. *For $\max\{|\alpha\beta abc/q^2|, |\alpha\gamma abc/q^2|\} < 1$, we have*

$$\sum_{n=0}^{\infty} \frac{(1-\alpha q^{2n})(\alpha, q/a, q/b, q/c)_n}{(q, \alpha a, \alpha b, \alpha c)_n}\left(\frac{\alpha abc}{q^2}\right)^n {}_4\phi_3\left(\begin{matrix}q^{-n}, \alpha q^n, \beta, \gamma \\ q/a, q/b, \alpha\beta\gamma abc/q\end{matrix}; q, q\right)$$
$$= \frac{(\alpha, \alpha ac/q, \alpha bc/q, \alpha\beta ab/q, \alpha\gamma ab/q, \alpha\beta\gamma abc/q^2)_\infty}{(\alpha a, \alpha b, \alpha c, \alpha\beta abc/q^2, \alpha\gamma abc/q^2, \alpha\beta\gamma ab/q)_\infty}.$$

Applying an expansion theorem [19, Proposition 1.6] for the analytic functions, Liu provided a simple proof of the above summation formula.

In this section, we will establish an A_n extension of Liu's extension by using the identity (1.8) in Theorem 1.8.

Theorem 7.3. *Let $\alpha, \beta, \gamma, a, b, c_1, \cdots, c_n$ be such that $\max\{|\alpha\beta abc_1\cdots c_n/q^{n+1}|, |\alpha\gamma abc_1\cdots c_n/q^{n+1}|\} < 1$; or, $c_r = q^{N_r}$, for $r = 1, 2, \cdots, n$, and so the series terminate. Suppose that none of the denominators in the following identity vanishes,*

then

$$\begin{aligned}
&\frac{(\alpha\beta ab/q,\alpha\gamma ab/q,\alpha ac_1\cdots c_n/q^n,\alpha bc_1\cdots c_n/q^n)_\infty}{(\alpha a,\alpha b,\alpha\beta abc_1\cdots c_n/q^{n+1},\alpha\gamma abc_1\cdots c_n/q^{n+1})_\infty}\\
&\times\prod_{i=1}^n\frac{(\alpha x_i,\alpha\beta\gamma abc_ix_i/q^2)_\infty}{(\alpha\beta\gamma abx_i/q,\alpha c_ix_i)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{(qx_r/x_sc_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times\prod_{r=1}^n\frac{(1-\alpha x_rq^{k_r+|\boldsymbol{k}|})\,(\alpha x_r)_{|\boldsymbol{k}|}}{(\alpha c_rx_r)_{|\boldsymbol{k}|}}q^{\sum_{r=1}^n(r-1)k_r}\\
&\times(\alpha a,\alpha b)_{|\boldsymbol{k}|}^{-1}\prod_{i=1}^n(q/ax_i,qx_i/b)_{k_i}\\
&\times\left(\frac{\alpha abc_1\cdots c_n}{q^{n+1}}\right)^{|\boldsymbol{k}|}q^{e_2(k_1,k_2\cdots,k_n)}\prod_{i=1}^nx_i^{k_i}\\
&\times\sum_{\substack{0\leq m_i\leq k_i\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}q^{\sum_{r=1}^n rm_r}\\
&\times\prod_{r,s=1}^n\frac{\left(q^{-k_s}x_r/x_s\right)_{m_r}}{(qx_r/x_s)_{m_r}}\prod_{i=1}^n\frac{(\alpha q^{|\boldsymbol{k}|}x_i,\beta x_i,\gamma x_i)_{m_i}}{(\alpha\beta\gamma abx_i/q,qx_i/a,qx_i/b)_{m_i}},
\end{aligned}$$

where $e_2(k_1,\cdots,k_n)$ is the second elementary symmetric function of $\{k_1,\cdots,k_n\}$.

Proof. Replacing a_r by qy_r for $r=1,2,\cdots,n$ in the identity (1.8) of Theorem 1.8 we have

$$\begin{aligned}
&f(\alpha qy_1,\cdots,\alpha qy_n)\prod_{i=1}^n\frac{(\alpha qx_i,\alpha bx_iy_i)_\infty}{(\alpha bx_i,\alpha qx_iy_i)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{(x_r/x_sy_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times\prod_{r=1}^n\frac{1-\alpha x_rq^{k_r+|\boldsymbol{k}|}}{1-\alpha x_r}\prod_{r=1}^n\frac{(\alpha x_r)_{|\boldsymbol{k}|}}{(\alpha qx_ry_r)_{|\boldsymbol{k}|}}\,(y_1\cdots y_n)^{|\boldsymbol{k}|}\,q^{\sum_{r=1}^n(r-1)k_r}\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{\left(q^{-k_s}x_r/x_s\right)_{m_r}}{(qx_r/x_s)_{m_r}}\\
&\times\prod_{r=1}^n\frac{\left(\alpha x_rq^{|\boldsymbol{k}|}\right)_{m_r}}{(\alpha bx_r)_{m_r}}q^{\sum_{r=1}^n rm_r}f(\alpha q^{1+m_1},\cdots,\alpha q^{1+m_n}).
\end{aligned}\tag{7.1}$$

Now we choose

$$\begin{aligned}
f(y_1,\cdots,y_n)=&\frac{(A_2,A_3,B_2y_1\cdots y_n/\alpha^nq^n,B_3y_1\cdots y_n/\alpha^nq^n)_\infty}{(B_2,B_3,A_2y_1\cdots y_n/\alpha^nq^n,A_3y_1\cdots y_n/\alpha^nq^n)_\infty}\\
&\cdot\prod_{i=1}^n\frac{(A_1x_i,B_1x_iy_i/\alpha q,\alpha bx_i,x_iy_i)_\infty}{(B_1x_i,A_1x_iy_i/\alpha q,\alpha qx_i,bx_iy_i/q)_\infty}.
\end{aligned}$$

Then

$$f(\alpha q y_1,\cdots,\alpha q y_n)=\frac{(A_2,A_3,B_2y_1\cdots y_n,B_3y_1\cdots y_n)_\infty}{(B_2,B_3,A_2y_1\cdots y_n,A_3y_1\cdots y_n)_\infty}\cdot\prod_{i=1}^n\frac{(A_1x_i,B_1x_iy_i,\alpha bx_i,\alpha qx_iy_i)_\infty}{(B_1x_i,A_1x_iy_i,\alpha qx_i,\alpha bx_iy_i)_\infty} \tag{7.2}$$

and

$$f(\alpha q^{1+m_1},\cdots,\alpha q^{1+m_n})=\frac{(A_2,A_3)_{|\boldsymbol{m}|}}{(B_2,B_3)_{|\boldsymbol{m}|}}\prod_{i=1}^n\frac{(A_1x_i,\alpha bx_i)_{m_i}}{(B_1x_i,\alpha qx_i)_{m_i}}. \tag{7.3}$$

Substituting (7.2) and (7.3) into (7.1) and then simplifying we get

$$\begin{aligned}
&\frac{(A_2,A_3,B_2y_1\cdots y_n,B_3y_1\cdots y_n)_\infty}{(B_2,B_3,A_2y_1\cdots y_n,A_3y_1\cdots y_n)_\infty}\prod_{i=1}^n\frac{(A_1x_i,B_1x_iy_i)_\infty}{(B_1x_i,A_1x_iy_i)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{(x_r/x_sy_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times\prod_{r=1}^n\frac{1-\alpha x_rq^{k_r+|\boldsymbol{k}|}}{1-\alpha x_r}\prod_{r=1}^n\frac{(\alpha x_r)_{|\boldsymbol{k}|}}{(\alpha qx_ry_r)_{|\boldsymbol{k}|}}(y_1\cdots y_n)^{|\boldsymbol{k}|}\,q^{\sum_{r=1}^n(r-1)k_r}\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{\left(q^{-k_s}x_r/x_s\right)_{m_r}}{(qx_r/x_s)_{m_r}}q^{\sum_{r=1}^n rm_r}\\
&\times\frac{(A_2,A_3)_{|\boldsymbol{m}|}}{(B_2,B_3)_{|\boldsymbol{m}|}}\prod_{r=1}^n\frac{\left(\alpha x_rq^{|\boldsymbol{k}|},A_1x_r\right)_{m_r}}{(B_1x_r,\alpha qx_r)_{m_r}}.
\end{aligned}$$

Taking $A_1\mapsto\alpha q, A_2\mapsto\alpha\beta ab/q, A_3\mapsto\alpha\gamma ab/q, B_1\mapsto\alpha\beta\gamma ab/q, B_2\mapsto\alpha a, B_3\mapsto\alpha b$ in the above identity and then multiplying both sides of the resulting identity by $\prod_{i=1}^n(1-\alpha x_i)$ we arrive at

$$\begin{aligned}
&\frac{(\alpha\beta ab/q,\alpha\gamma ab/q,\alpha ay_1\cdots y_n,\alpha by_1\cdots y_n)_\infty}{(\alpha a,\alpha b,\alpha\beta aby_1\cdots y_n/q,\alpha\gamma aby_1\cdots y_n/q)_\infty}\prod_{i=1}^n\frac{(\alpha x_i,\alpha\beta\gamma abx_iy_i/q)_\infty}{(\alpha\beta\gamma abx_i/q,\alpha qx_iy_i)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{(x_r/x_sy_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times\prod_{r=1}^n\frac{(1-\alpha x_rq^{k_r+|\boldsymbol{k}|})(\alpha x_r)_{|\boldsymbol{k}|}}{(\alpha qx_ry_r)_{|\boldsymbol{k}|}}(y_1\cdots y_n)^{|\boldsymbol{k}|}\,q^{\sum_{r=1}^n(r-1)k_r}\\
&\times\sum_{\substack{0\leq m_r\leq k_r\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{\left(q^{-k_s}x_r/x_s\right)_{m_r}}{(qx_r/x_s)_{m_r}}q^{\sum_{r=1}^n rm_r}\\
&\times\frac{(\alpha\beta ab/q,\alpha\gamma ab/q)_{|\boldsymbol{m}|}}{(\alpha a,\alpha b)_{|\boldsymbol{m}|}}\prod_{r=1}^n\frac{\left(\alpha x_rq^{|\boldsymbol{k}|}\right)_{m_r}}{(\alpha\beta\gamma abx_r/q)_{m_r}}.
\end{aligned} \tag{7.4}$$

It follows from [12, eq.(I.8)] that

$$(ax_iq^{-k_i},bq^{-k_i}/x_i)_{k_i}=(q/ax_i,qx_i/b)_{k_i}(ab)^{k_i}q^{-k_i-k_i^2}.$$

Then

$$\prod_{i=1}^{n}(ax_iq^{-k_i},bq^{-k_i}/x_i)_{k_i}=(ab)^{|\boldsymbol{k}|}q^{-|k|-\sum_{i=1}^n k_i^2}\prod_{i=1}^{n}(q/ax_i,qx_i/b)_{k_i}. \tag{7.5}$$

Recall a $U(n+1)$ Sears ${}_4\phi_3$ transformation [24, Theorem 5.2]:

$$\begin{aligned}
&\sum_{\substack{0\le y_r\le N_r\\ r=1,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{y_r-y_s}x_r/x_s}{1-x_r/x_s}q^{\sum_{r=1}^n rj_r}\\
&\times\prod_{r,s=1}^{n}\frac{\left(q^{-N_s}x_r/x_s\right)_{y_r}}{(qx_r/x_s)_{y_r}}\prod_{r=1}^{n}\frac{(ax_r)_{y_r}}{(dx_r)_{y_r}}\cdot\frac{(b,c)_{|\boldsymbol{y}|}}{(e,f)_{|\boldsymbol{y}|}}\\
&=(e,f)_{|\boldsymbol{N}|}^{-1}\prod_{i=1}^{n}(ex_iq^{|\boldsymbol{N}|-N_i}/a,fq^{|\boldsymbol{N}|-N_i}/ax_i)_{N_i}\\
&\times a^{|\boldsymbol{N}|}q^{-e_2(N_1,N_2\cdots,N_n)}\prod_{i=1}^{n}x_i^{N_i}\\
&\times\sum_{\substack{0\le y_i\le N_i\\ r=1,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{y_r-y_s}x_r/x_s}{1-x_r/x_s}q^{\sum_{r=1}^n ry_r}\\
&\times\prod_{r,s=1}^{n}\frac{\left(q^{-N_s}x_r/x_s\right)_{y_r}}{(qx_r/x_s)_{y_r}}\prod_{i=1}^{n}\frac{(ax_i,\ dx_i/b,dx_i/c)_{y_i}}{(dx_i,ax_iq^{1-|\boldsymbol{N}|}/e,ax_iq^{1-|\boldsymbol{N}|}/f)_{y_i}},
\end{aligned}$$

where $abc=defq^{|\boldsymbol{N}|-1}$ and $e_2(N_1,N_2,\cdots,N_n)$ is the second elementary symmetric function of $\{N_1,N_2,\cdots,N_n\}$. Applying the $U(n+1)$ Sears ${}_4\phi_3$ transformation with $N_r\mapsto k_r, a\mapsto\alpha q^{|\boldsymbol{k}|}, b\mapsto\alpha\beta ab/q, c\mapsto\alpha\gamma ab/q, d\mapsto\alpha\beta\gamma ab/q, e\mapsto\alpha a, f\mapsto\alpha b$ and then substituting (7.5) into the resulting identity we get

$$\begin{aligned}
&\sum_{\substack{0\le m_r\le k_r\\ r=1,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{\left(q^{-k_s}x_r/x_s\right)_{m_r}}{(qx_r/x_s)_{m_r}}q^{\sum_{r=1}^n rm_r}\\
&\quad\times\frac{(\alpha\beta ab/q,\alpha\gamma ab/q)_{|\boldsymbol{m}|}}{(\alpha a,\alpha b)_{|\boldsymbol{m}|}}\prod_{r=1}^{n}\frac{\left(\alpha x_rq^{|\boldsymbol{k}|}\right)_{m_r}}{(\alpha\beta\gamma abx_r/q)_{m_r}}\\
&=(\alpha a,\alpha b)_{|\boldsymbol{k}|}^{-1}\prod_{i=1}^{n}(q/ax_i,qx_i/b)_{k_i}\\
&\times(\alpha ab/q)^{|\boldsymbol{k}|}q^{e_2(k_1,k_2\cdots,k_n)}\prod_{i=1}^{n}x_i^{k_i}\\
&\times\sum_{\substack{0\le m_i\le k_i\\ r=1,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}q^{\sum_{r=1}^n rm_r}\\
&\times\prod_{r,s=1}^{n}\frac{\left(q^{-k_s}x_r/x_s\right)_{m_r}}{(qx_r/x_s)_{m_r}}\prod_{i=1}^{n}\frac{(\alpha q^{|\boldsymbol{k}|}x_i,\beta x_i,\gamma x_i)_{m_i}}{(\alpha\beta\gamma abx_i/q,qx_i/a,qx_i/b)_{m_i}}.
\end{aligned}\tag{7.6}$$

Then the result follows by substituting (7.6) into (7.4) and then replacing y_i by c_i/q for $i=1,2,\cdots,n$ in the resulting identity. □

The summation formula in Theorem 7.3 is an A_n extension of Liu's extension of the Rogers non-terminating ${}_6\phi_5$ summation formula; replace x_i by x_i/x_n in the $n=1$ case of the identity in Theorem 7.3.

If $\gamma=0$, then the identity in Theorem 7.3 becomes the following summation formula.

Theorem 7.4. *Let $\alpha,\beta,a,b,c_1,\cdots,c_n$ be such that $|\alpha\beta abc_1\cdots c_n/q^{n+1}|<1$; or, $c_r=q^{N_r}$, for $r=1,2,\cdots,n$, and so the series terminate. Suppose that none of the denominators in the following identity vanishes, then*

$$\begin{aligned}
&\frac{(\alpha\beta ab/q,\alpha ac_1\cdots c_n/q^n,\alpha bc_1\cdots c_n/q^n)_\infty}{(\alpha a,\alpha b,\alpha\beta abc_1\cdots c_n/q^{n+1})_\infty}\prod_{i=1}^n\frac{(\alpha x_i)_\infty}{(\alpha c_ix_i)_\infty}\\
&=\sum_{\substack{k_r\geq 0\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^n\frac{(qx_r/x_sc_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times\prod_{r=1}^n\frac{\left(1-\alpha x_rq^{k_r+|\boldsymbol{k}|}\right)(\alpha x_r)_{|\boldsymbol{k}|}}{(\alpha c_rx_r)_{|\boldsymbol{k}|}}q^{\sum_{r=1}^n(r-1)k_r}\\
&\times(\alpha a,\alpha b)_{|\boldsymbol{k}|}^{-1}\prod_{i=1}^n(q/ax_i,qx_i/b)_{k_i}\\
&\times\left(\frac{\alpha abc_1\cdots c_n}{q^{n+1}}\right)^{|\boldsymbol{k}|}q^{e_2(k_1,k_2\cdots,k_n)}\prod_{i=1}^nx_i^{k_i}\\
&\times\sum_{\substack{0\leq m_i\leq k_i\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s}q^{\sum_{r=1}^n rm_r}\\
&\times\prod_{r,s=1}^n\frac{\left(q^{-k_s}x_r/x_s\right)_{m_r}}{(qx_r/x_s)_{m_r}}\prod_{i=1}^n\frac{(\alpha q^{|\boldsymbol{k}|}x_i,\beta x_i)_{m_i}}{(qx_i/a,qx_i/b)_{m_i}},
\end{aligned}$$

where $e_2(k_1,\cdots,k_n)$ is the second elementary symmetric function of $\{k_1,\cdots,k_n\}$.

The identity in Theorem 7.4 is an A_n extension of the following summation formula:

$$\begin{aligned}
&\sum_{n=0}^\infty\frac{(1-\alpha q^{2n})(\alpha,q/a,q/b,q/c)_n}{(q,\alpha a,\alpha b,\alpha c)_n}\left(\frac{\alpha abc}{q^2}\right)^n{}_3\phi_2\left(\begin{matrix}q^{-n},\alpha q^n,\beta\\ q/a,q/b\end{matrix};q,q\right)\\
&=\frac{(\alpha,\alpha ac/q,\alpha bc/q,\alpha\beta ab/q)_\infty}{(\alpha a,\alpha b,\alpha c,\alpha\beta abc/q^2)_\infty},
\end{aligned}$$

where $|\alpha\beta abc/q^2|<1$. This formula is equivalent to a summation theorem due to Ismail, Rahman and Suslov [14, Theorem 5.1].

8. An A_n extension of a generalization of Fang's identity

Fang [8] applied a q-exponential operator constructed in [10] to the Rogers-Fine identity to establish the following interesting identity [8, eq.(9)]:

$$\sum_{n=0}^{M}\frac{(q^{-M},c)_n}{(\beta,bc)_n}\tau^n=\sum_{n=0}^{M}\frac{(q^{-M},q^{1-M}\tau/\beta,c)_n(1-\tau q^{2n-M})}{(\beta,bc)_n(\tau)_{n+1}}(\beta\tau)^n q^{n^2-n}\times{}_3\phi_2\left(\begin{matrix}q^{-n},b,q^{1-n}/\beta\\ q^{1-M}\tau/\beta,q^{1-n}/c\end{matrix};q,q\right),\tag{8.1}$$

where M is a nonnegative integer. Applying this q-exponential operator repeatedly, Fang obtained a very general identity, from which an extension of the terminating very-well-poised ${}_6\phi_5$ summation formula was derived. The formula (8.1) is important in Fang's paper [8]. It can be extended to the following non-terminating identity:

$$\sum_{n=0}^{\infty}\frac{(1/y,c)_n}{(\beta,bc)_n}(y\tau)^n=\sum_{n=0}^{\infty}\frac{(1/y,q\tau/\beta,c)_n(1-\tau q^{2n})}{(\beta,bc)_n(y\tau)_{n+1}}(y\beta\tau)^n q^{n^2-n}\times{}_3\phi_2\left(\begin{matrix}q^{-n},b,q^{1-n}/\beta\\ q\tau/\beta,q^{1-n}/c\end{matrix};q,q\right),\tag{8.2}$$

where $|y\tau|<1$. Applying the terminating ${}_3\phi_2$ transformation formula in [12, eq.(III.13)] to the inner sum ${}_3\phi_2$ on the right side of (8.2), we obtain an equivalent identity:

$$\sum_{n=0}^{\infty}\frac{(1/y,c)_n}{(\beta,bc)_n}(y\tau)^n=\sum_{n=0}^{\infty}\frac{(1/y,q\tau/\beta)_n(1-\tau q^{2n})}{(\beta)_n(y\tau)_{n+1}}(y\beta\tau)^n q^{n^2-n}\times{}_3\phi_2\left(\begin{matrix}q^{-n},\tau q^n,b\\ q\tau/\beta,bc\end{matrix};q,cq/\beta\right),\tag{8.3}$$

where $|y\tau|<1$. Let $y\mapsto q^M,\tau\mapsto\tau/q^M$. Then the formula (8.2) reduces to (8.1). Thus, (8.2) and (8.3) are both generalizations of Fang's identity (8.1).

In this section, we deduce an A_n extension of the identity (8.3) using the identity (1.2) in Theorem 1.2.

Theorem 8.1. *Let $y_1,y_2,\cdots,y_n,\tau$ be such that $|y_1y_2\cdots y_n\tau|<1$. Suppose that none of the denominators in the following identity vanishes, then*

$$\begin{aligned}&\sum_{\substack{j_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(x_r/x_sy_s)_{j_r}}{(qx_r/x_s)_{j_r}}\\ &\quad\times(y_1y_2\cdots y_n\tau)^{|\boldsymbol{j}|}q^{\sum_{r=1}^n(r-1)j_r}\frac{(q,c)_{|\boldsymbol{j}|}}{(bc)_{|\boldsymbol{j}|}}\prod_{r=1}^{n}(\beta x_r)_{j_r}^{-1}\\ =&\sum_{\substack{k_r\geq 0\\ r=1,2,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(x_r/x_sy_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\ &\times\frac{(1-\tau q^{2|\boldsymbol{k}|})(q)_{|\boldsymbol{k}|}}{(y_1y_2\cdots y_n\tau)_{|\boldsymbol{k}|+1}}(y_1y_2\cdots y_n\beta\tau)^{|\boldsymbol{k}|}q^{\sum_{r=1}^n(r-1)k_r+\binom{|\boldsymbol{k}|}{2}+\sum_{r=1}^n\binom{k_r}{2}}\\ &\times\prod_{r=1}^{n}\frac{(q^{1+|\boldsymbol{k}|-k_r}\tau/\beta x_r)_{k_r}}{(\beta x_r)_{k_r}}\prod_{r=1}^{n}x_r^{k_r}\sum_{\substack{0\leq j_r\leq k_r\\ r=1,\cdots,n}}\prod_{1\leq r<s\leq n}\frac{1-q^{k_s-k_r+j_r-j_s}x_s/x_r}{1-q^{k_s-k_r}x_s/x_r}\end{aligned}$$

$$\times q^{|\boldsymbol{k}||\boldsymbol{j}|-\sum_{r=1}^n k_r j_r - e_2(j_1,\cdots,j_n)+\sum_{r=1}^n r j_r}$$
$$\times \prod_{r,s=1}^n \frac{\left(q^{-k_r}x_s/x_r\right)_{j_r}}{\left(q^{1+k_s-k_r}x_s/x_r\right)_{j_r}} \prod_{r=1}^n \left\{ x_r^{-j_r} \left(\tau q^{1-k_r+|\boldsymbol{k}|}/\beta x_r\right)_{j_r}^{-1} \right\}$$
$$\times \left(\frac{c}{\beta}\right)^{|\boldsymbol{j}|} \frac{(\tau q^{|\boldsymbol{k}|})_{|\boldsymbol{j}|}(b)_{|\boldsymbol{j}|}}{(bc)_{|\boldsymbol{j}|}},$$

where $e_2(j_1,\cdots,j_n)$ is the second elementary symmetric function of $\{j_1,\cdots,j_n\}$.

Proof. Recall a relation between (6.5a) and (6.5c) in [22, Theorem 6.5]:

$$\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^n \frac{\left(q^{-k_s}x_r/x_s\right)_{j_r}}{\left(qx_r/x_s\right)_{j_r}}$$
$$\times \prod_{r=1}^n \frac{(cx_r)_{j_r}}{(ex_r)_{j_r}} \frac{(a)_{|\boldsymbol{j}|}(b)_{|\boldsymbol{j}|}}{(d)_{|\boldsymbol{j}|}(f)_{|\boldsymbol{j}|}} q^{\sum_{r=1}^n r j_r}$$
$$= a^{|\boldsymbol{k}|} \frac{(f/a)_{|\boldsymbol{k}|}}{(f)_{|\boldsymbol{k}|}} \prod_{r=1}^n \frac{(ex_r/a)_{k_r}}{(ex_r)_{k_r}}$$
$$\times \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_s-k_r+j_r-j_s}x_s/x_r}{1-q^{k_s-k_r}x_s/x_r} q^{\sum_{r=1}^n r j_r}$$
$$\times \prod_{r,s=1}^n \frac{\left(q^{-k_r}x_s/x_r\right)_{j_r}}{\left(q^{1+k_s-k_r}x_s/x_r\right)_{j_r}} \prod_{r=1}^n \frac{\left(dq^{|\boldsymbol{k}|-k_r}/cx_r\right)_{j_r}}{\left(aq^{1-k_r}/ex_r\right)_{j_r}}$$
$$\times \frac{(a)_{|\boldsymbol{j}|}(d/b)_{|\boldsymbol{j}|}}{(d)_{|\boldsymbol{j}|}(aq^{1-|\boldsymbol{k}|}/f)_{|\boldsymbol{j}|}},$$

where $def = abcq^{1-|\boldsymbol{N}|}$. This is actually an A_n Sears transformation formula. Substituting $f = abcq^{1-|\boldsymbol{N}|}/de$ into the above identity and then taking $c \to 0$ we obtain an A_n terminating ${}_3\phi_2$ transformation formula:

$$\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^n \frac{\left(q^{-k_s}x_r/x_s\right)_{j_r}}{\left(qx_r/x_s\right)_{j_r}}$$
$$\times \prod_{r=1}^n (ex_r)_{j_r}^{-1} \frac{(a)_{|\boldsymbol{j}|}(b)_{|\boldsymbol{j}|}}{(d)_{|\boldsymbol{j}|}} q^{\sum_{r=1}^n r j_r}$$
$$= a^{|\boldsymbol{k}|} \prod_{r=1}^n \frac{(ex_r/a)_{k_r}}{(ex_r)_{k_r}} \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_s-k_r+j_r-j_s}x_s/x_r}{1-q^{k_s-k_r}x_s/x_r}$$
$$\times q^{|\boldsymbol{k}||\boldsymbol{j}|-\sum_{r=1}^n k_r j_r - e_2(j_1,\cdots,j_n)+\sum_{r=1}^n r j_r}$$
$$\times \prod_{r,s=1}^n \frac{\left(q^{-k_r}x_s/x_r\right)_{j_r}}{\left(q^{1+k_s-k_r}x_s/x_r\right)_{j_r}} \prod_{r=1}^n \left\{ x_r^{-j_r} \left(aq^{1-k_r}/ex_r\right)_{j_r}^{-1} \right\}$$
$$\times \left(\frac{b}{e}\right)^{|\boldsymbol{j}|} \frac{(a)_{|\boldsymbol{j}|}(d/b)_{|\boldsymbol{j}|}}{(d)_{|\boldsymbol{j}|}}.$$

The $a \mapsto \tau q^{|\boldsymbol{k}|}, b \mapsto c, d \mapsto bc, e \mapsto \beta$ case of this identity is

$$
\begin{aligned}
&\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_r/x_s\right)_{j_r}}{\left(qx_r/x_s\right)_{j_r}} \\
&\times \prod_{r=1}^{n} (\beta x_r)_{j_r}^{-1} \frac{(\tau q^{|\boldsymbol{k}|})_{|\boldsymbol{j}|}(c)_{|\boldsymbol{j}|}}{(bc)_{|\boldsymbol{j}|}} q^{\sum_{r=1}^n r j_r} \\
&= (\tau q^{|\boldsymbol{k}|})^{|\boldsymbol{k}|} \prod_{r=1}^{n} \frac{(\beta x_r q^{-|\boldsymbol{k}|}/\tau)_{k_r}}{(\beta x_r)_{k_r}} \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_s-k_r+j_r-j_s}x_s/x_r}{1-q^{k_s-k_r}x_s/x_r} \\
&\times q^{|\boldsymbol{k}||\boldsymbol{j}|-\sum_{r=1}^n k_r j_r - e_2(j_1,\cdots,j_n)+\sum_{r=1}^n r j_r} \\
&\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_r}x_s/x_r\right)_{j_r}}{\left(q^{1+k_s-k_r}x_s/x_r\right)_{j_r}} \prod_{r=1}^{n} \left\{ x_r^{-j_r} \left(\tau q^{1-k_r+|\boldsymbol{k}|}/\beta x_r\right)_{j_r}^{-1} \right\} \\
&\times \left(\frac{c}{\beta}\right)^{|\boldsymbol{j}|} \frac{(\tau q^{|\boldsymbol{k}|})_{|\boldsymbol{j}|}(b)_{|\boldsymbol{j}|}}{(bc)_{|\boldsymbol{j}|}}.
\end{aligned} \tag{8.4}
$$

It follows from [12, eq.(I.8)] that

$$
(\beta x_r q^{-|\boldsymbol{k}|}/\tau)_{k_r} = \left(\frac{q^{1+|\boldsymbol{k}|-k_r}\tau}{\beta x_r}\right)_{k_r} \left(-\frac{\beta x_r q^{k_r-|\boldsymbol{k}|-1}}{\tau}\right)^{k_r} q^{-\binom{k_r}{2}}.
$$

Then

$$
\prod_{r=1}^{n} (\beta x_r q^{-|\boldsymbol{k}|}/\tau)_{k_r} = \left(-\frac{\beta}{\tau}\right)^{|\boldsymbol{k}|} q^{\sum_{r=1}^n \binom{k_r}{2} - |\boldsymbol{k}|^2} \prod_{r=1}^{n} \left(\frac{q^{1+|\boldsymbol{k}|-k_r}\tau}{\beta x_r}\right)_{k_r} \prod_{r=1}^{n} x_r^{k_r}.
$$

Substituting this identity into (8.4) gives

$$
\begin{aligned}
&\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_r/x_s\right)_{j_r}}{\left(qx_r/x_s\right)_{j_r}} \\
&\times \prod_{r=1}^{n} (\beta x_r)_{j_r}^{-1} \frac{(\tau q^{|\boldsymbol{k}|})_{|\boldsymbol{j}|}(c)_{|\boldsymbol{j}|}}{(bc)_{|\boldsymbol{j}|}} q^{\sum_{r=1}^n r j_r} \\
&= \prod_{r=1}^{n} \frac{(q^{1+|\boldsymbol{k}|-k_r}\tau/\beta x_r)_{k_r}}{(\beta x_r)_{k_r}} (-\beta)^{|\boldsymbol{k}|} q^{\sum_{r=1}^n \binom{k_r}{2}} \prod_{r=1}^{n} x_r^{k_r} \\
&\times \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_s-k_r+j_r-j_s}x_s/x_r}{1-q^{k_s-k_r}x_s/x_r} \prod_{r,s=1}^{n} \frac{\left(q^{-k_r}x_s/x_r\right)_{j_r}}{\left(q^{1+k_s-k_r}x_s/x_r\right)_{j_r}} \\
&\times \prod_{r=1}^{n} \left\{ x_r^{-j_r} \left(\tau q^{1-k_r+|\boldsymbol{k}|}/\beta x_r\right)_{j_r}^{-1} \right\} \cdot q^{|\boldsymbol{k}||\boldsymbol{j}|-\sum_{r=1}^n k_r j_r - e_2(j_1,\cdots,j_n)+\sum_{r=1}^n r j_r} \\
&\times \left(\frac{c}{\beta}\right)^{|\boldsymbol{j}|} \frac{(\tau q^{|\boldsymbol{k}|})_{|\boldsymbol{j}|}(b)_{|\boldsymbol{j}|}}{(bc)_{|\boldsymbol{j}|}}.
\end{aligned} \tag{8.5}
$$

Taking $N_r \mapsto k_r - j_r, x_r \mapsto q^{j_r}x_r, b \mapsto \tau q^{|\boldsymbol{k}|+|\boldsymbol{j}|}, c \mapsto \tau q^{|\boldsymbol{j}|}$ in (6.1) we have

$$\begin{aligned}
&\sum_{\substack{0\le m_r\le k_r-j_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{m_r-m_s+j_r-j_s}x_r/x_s}{1-q^{j_r-j_s}x_r/x_s} q^{\sum_{r=1}^n rm_r}\\
&\times \prod_{r,s=1}^n \frac{(q^{-k_s+j_r}x_r/x_s)_{m_r}}{(q^{1+j_r-j_s}x_r/x_s)_{m_r}} \frac{(\tau q^{|\boldsymbol{k}|+|\boldsymbol{j}|})_{|\boldsymbol{m}|}}{(\tau q^{|\boldsymbol{j}|})_{|\boldsymbol{m}|}}\\
&= \frac{(q^{-|\boldsymbol{k}|})_{|\boldsymbol{k}|-|\boldsymbol{j}|}}{(\tau q^{|\boldsymbol{j}|})_{|\boldsymbol{k}|-|\boldsymbol{j}|}} (\tau q^{|\boldsymbol{k}|+|\boldsymbol{j}|})^{|\boldsymbol{k}|-|\boldsymbol{j}|}\\
&= \frac{(q)_{|\boldsymbol{k}|}(-\tau)^{|\boldsymbol{k}|-|\boldsymbol{j}|}}{(\tau q^{|\boldsymbol{j}|})_{|\boldsymbol{k}|-|\boldsymbol{j}|}(q)_{|\boldsymbol{j}|}} q^{\binom{|\boldsymbol{k}|}{2}-\binom{|\boldsymbol{j}|}{2}},
\end{aligned} \tag{8.6}$$

where we have used the identity [12, eq.(I.14)] in the last equality.

We first set $b \mapsto 0, a \mapsto \tau$, and then take

$$K(y) \mapsto 1 - y_1y_2\cdots y_n\tau$$

and

$$\beta(\boldsymbol{j}) \mapsto \frac{(q,c)_{|\boldsymbol{j}|}}{(bc)_{|\boldsymbol{j}|}} \tau^{|\boldsymbol{j}|} \prod_{r=1}^n (\beta x_r)_{j_r}^{-1} \prod_{r,s=1}^n (qx_r/x_s)_{j_r}^{-1}$$

in the identity (1.2) of Theorem 1.2 to get

$$\begin{aligned}
&(1-y_1y_2\cdots y_n\tau) \sum_{\substack{j_r\ge 0\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^n \frac{(x_r/x_sy_s)_{j_r}}{(qx_r/x_s)_{j_r}}\\
&\quad\times (y_1y_2\cdots y_n\tau)^{|\boldsymbol{j}|} q^{\sum_{r=1}^n (r-1)j_r} \frac{(q,c)_{|\boldsymbol{j}|}}{(bc)_{|\boldsymbol{j}|}} \prod_{r=1}^n (\beta x_r)_{j_r}^{-1}\\
&= \sum_{\substack{k_r\ge 0\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^n \frac{(x_r/x_sy_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times \frac{(1-\tau q^{2|\boldsymbol{k}|})(\tau)_{|\boldsymbol{k}|}}{(\tau q y_1y_2\cdots y_n)_{|\boldsymbol{k}|}} (y_1y_2\cdots y_n)^{|\boldsymbol{k}|} q^{\sum_{r=1}^n (r-1)k_r}\\
&\times \sum_{\substack{0\le j_r\le k_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^n \frac{(q^{-k_s}x_r/x_s)_{j_r}}{(qx_r/x_s)_{j_r}}\\
&\times \frac{(\tau q^{|\boldsymbol{k}|},q,c)_{|\boldsymbol{j}|}}{(\tau,bc)_{|\boldsymbol{j}|}} (-\tau)^{|\boldsymbol{j}|} q^{\sum_{r=1}^n rj_r+\binom{|\boldsymbol{j}|}{2}} \prod_{r=1}^n (\beta x_r)_{j_r}^{-1}\\
&\times \sum_{\substack{0\le m_r\le k_r-j_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{m_r-m_s+j_r-j_s}x_r/x_s}{1-q^{j_r-j_s}x_r/x_s} q^{\sum_{r=1}^n rm_r}\\
&\quad\times \prod_{r,s=1}^n \frac{(q^{-k_s+j_r}x_r/x_s)_{m_r}}{(q^{1+j_r-j_s}x_r/x_s)_{m_r}} \frac{(\tau q^{|\boldsymbol{k}|+|\boldsymbol{j}|})_{|\boldsymbol{m}|}}{(\tau q^{|\boldsymbol{j}|})_{|\boldsymbol{m}|}}.
\end{aligned}$$

Then the result follows by substituting (8.6) into the inner-most sum $\sum_{\substack{0\le m_r\le k_r-j_r\\ r=1,2,\cdots,n}}$ on the right side of the above identity, then applying (8.5) to the second sum $\sum_{\substack{0\le j_r\le k_r\\ r=1,2,\cdots,n}}$ and finally multiplying both sides by $\frac{1}{1-y_1y_2\cdots y_n\tau}$. □

9. An A_n extension of Andrews' expansion formula

Applying a Bailey pair in [2, Theorem 3], Andrews [2, Theorem 5] proved the following celebrated expansion formula[1] for an infinite product:

$$\frac{(q,bq/y,aq)_\infty}{(bq,q/y,aq/y)_\infty}=\sum_{n=0}^{\infty}\frac{(1-aq^{2n})(a,y)_n(a/y)^nq^{n^2}}{(1-a)(aq/y,bq)_n}\sum_{j=0}^{n}\frac{(b)_j}{(q)_j}(1/a)^jq^{j(1-n)}. \tag{9.1}$$

The formula (9.1) is very useful and was used by Andrews to deduce Hecke modular form identities and generating functions for the number of representations of an integer as a sum of three squares and the number of representations of an integer as a sum of three triangular numbers.

In this section, we apply the identity in Theorem 1.9 to give an A_n extension of the identity (9.1).

Theorem 9.1. *Suppose that none of the denominators in the following identity vanishes, then*

$$\begin{aligned}
&\frac{(q,aq,bq/y_1\cdots y_n)_\infty}{(aq/y_1\cdots y_n,bq,q/y_1\cdots y_n)_\infty}\\
&=\sum_{\substack{k_r\ge 0\\ r=1,2,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s}\prod_{r,s=1}^{n}\frac{(y_sx_r/x_s)_{k_r}}{(qx_r/x_s)_{k_r}}\\
&\times\frac{(1-aq^{2|\boldsymbol{k}|})(a,q)_{|\boldsymbol{k}|}(a/y_1\cdots y_n)^{|\boldsymbol{k}|}}{(1-a)(aq/y_1\cdots y_n,bq)_{|\boldsymbol{k}|}}q^{\sum_{r=1}^n(r-1)k_r+|\boldsymbol{k}|^2}\\
&\times\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s}\cdot q^{\sum_{r=1}^n rj_r-|\boldsymbol{k}||\boldsymbol{j}|}\\
&\times\prod_{r,s=1}^{n}\frac{\left(q^{-k_s}x_r/x_s\right)_{j_r}}{\left(qx_r/x_s\right)_{j_r}}\frac{(b)_{|\boldsymbol{j}|}(1/a)^{|\boldsymbol{j}|}}{(q^{-|\boldsymbol{k}|})_{|\boldsymbol{j}|}}.
\end{aligned}\tag{9.2}$$

Proof. We first prove the following identity: if $k_1,k_2,\cdots,k_n$ are non-negative integers, then

$$\begin{aligned}
&\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s}\\
&\times\prod_{r,s=1}^{n}\frac{\left(q^{-k_s}x_r/x_s\right)_{j_r}}{\left(qx_r/x_s\right)_{j_r}}\frac{(aq^{|\boldsymbol{k}|},q)_{|\boldsymbol{j}|}}{(cq)_{|\boldsymbol{j}|}}q^{\sum_{r=1}^n rj_r}\\
&=a^{|\boldsymbol{k}|}q^{|\boldsymbol{k}|^2}\frac{(q)_{|\boldsymbol{k}|}}{(cq)_{|\boldsymbol{k}|}}\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}}\prod_{1\le r<s\le n}\frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s}
\end{aligned}\tag{9.3}$$

[1]Note that the representations for A_n in [2, (4.2) and (4.5)] are equal. Therefore, for brevity, we choose the former representation for A_n in [2, Theorem 5].

$$\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_r/x_s\right)_{j_r}}{\left(qx_r/x_s\right)_{j_r}} \frac{(c)_{|\boldsymbol{j}|}(1/a)^{|\boldsymbol{j}|}}{(q^{-|\boldsymbol{k}|})_{|\boldsymbol{j}|}} q^{\sum_{r=1}^{n} rj_r - |\boldsymbol{k}||\boldsymbol{j}|}.$$

Recall an A_n Sears transformation formula between (6.5a) and (6.5c) in [22, Theorem 6.5]: Suppose that $def = abcq^{1-|\boldsymbol{k}|}$, then

$$\begin{aligned}
&\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_r/x_s\right)_{j_r}}{\left(qx_r/x_s\right)_{j_r}}\\
&\times \prod_{r=1}^{n} \frac{(cx_r)_{j_r}}{(ex_r)_{j_r}} \frac{(a)_{|\boldsymbol{j}|}(b)_{|\boldsymbol{j}|}}{(d)_{|\boldsymbol{j}|}(f)_{|\boldsymbol{j}|}} \cdot q^{\sum_{r=1}^{n} rj_r}\\
&= a^{|\boldsymbol{k}|} \frac{(f/a)_{|\boldsymbol{k}|}}{(f)_{|\boldsymbol{k}|}} \prod_{r=1}^{n} \frac{(ex_r/a)_{k_r}}{(ex_r)_{k_r}}\\
&\times \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_s-k_r+j_r-j_s}x_s/x_r}{1-q^{k_s-k_r}x_s/x_r} q^{\sum_{r=1}^{n} rj_r}\\
&\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_r}x_s/x_r\right)_{j_r}}{\left(q^{1+k_s-k_r}x_s/x_r\right)_{j_r}} \prod_{r=1}^{n} \frac{\left(dq^{|\boldsymbol{k}|-k_r}/cx_r\right)_{j_r}}{\left(aq^{1-k_r}/ex_r\right)_{j_r}}\\
&\times \frac{(a)_{|\boldsymbol{j}|}(d/b)_{|\boldsymbol{j}|}}{(d)_{|\boldsymbol{j}|}(aq^{1-|\boldsymbol{k}|}/f)_{|\boldsymbol{j}|}}.
\end{aligned} \tag{9.4}$$

Substituting $e = abcq^{1-|\boldsymbol{k}|}/df$ into (9.4), setting $c \to 0$, and then replacing f by e, and a by c, we obtain

$$\begin{aligned}
&\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s}\\
&\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_r/x_s\right)_{j_r}}{\left(qx_r/x_s\right)_{j_r}} \frac{(b)_{|\boldsymbol{j}|}(c)_{|\boldsymbol{j}|}}{(d)_{|\boldsymbol{j}|}(e)_{|\boldsymbol{j}|}} q^{\sum_{r=1}^{n} rj_r}\\
&= c^{|\boldsymbol{k}|} \frac{(e/c)_{|\boldsymbol{k}|}}{(e)_{|\boldsymbol{k}|}} \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_s-k_r+j_r-j_s}x_s/x_r}{1-q^{k_s-k_r}x_s/x_r}\\
&\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_r}x_s/x_r\right)_{j_r}}{\left(q^{1+k_s-k_r}x_s/x_r\right)_{j_r}} \cdot \left(\frac{bq}{e}\right)^{|\boldsymbol{j}|} q^{\sum_{r=1}^{n}(r-1)j_r}\\
&\times \frac{(c)_{|\boldsymbol{j}|}(d/b)_{|\boldsymbol{j}|}}{(d)_{|\boldsymbol{j}|}(cq^{1-|\boldsymbol{k}|}/e)_{|\boldsymbol{j}|}}.
\end{aligned} \tag{9.5}$$

Replace b by d/b, e by $cq^{1-|\boldsymbol{k}|}/e, x_r$ by $x_r q^{-k_r}$ in (9.5) to arrive at

$$\begin{aligned}
&\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_s/x_r}{1-x_s/x_r} \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_s/x_r\right)_{j_r}}{\left(qx_s/x_r\right)_{j_r}}\\
&\times \frac{(b)_{|\boldsymbol{j}|}(c)_{|\boldsymbol{j}|}}{(d)_{|\boldsymbol{j}|}(e)_{|\boldsymbol{j}|}} \cdot \left(\frac{deq^{|\boldsymbol{k}|}}{bc}\right)^{|\boldsymbol{j}|} q^{\sum_{r=1}^{n}(r-1)j_r}
\end{aligned} \tag{9.6}$$

$$= \frac{(e/c)_{|\boldsymbol{k}|}}{(e)_{|\boldsymbol{k}|}} \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s-k_r+k_s}x_r/x_s}{1-q^{-k_r+k_s}x_r/x_s}$$
$$\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_r}x_r/x_s\right)_{j_r}}{\left(q^{1-k_r+k_s}x_r/x_s\right)_{j_r}} \frac{(c)_{|\boldsymbol{j}|}(d/b)_{|\boldsymbol{j}|}}{(d)_{|\boldsymbol{j}|}(cq^{1-|\boldsymbol{k}|}/e)_{|\boldsymbol{j}|}} q^{\sum_{r=1}^n rj_r}.$$

Let $b\mapsto aq^{|\boldsymbol{k}|}, d\mapsto 1/t, e\mapsto cq$ in (9.6) we have

$$\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_s/x_r}{1-x_s/x_r} \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_s/x_r\right)_{j_r}}{\left(qx_s/x_r\right)_{j_r}}$$
$$\times \frac{(c)_{|\boldsymbol{j}|}(aq^{|\boldsymbol{k}|})_{|\boldsymbol{j}|}}{(1/t)_{|\boldsymbol{j}|}(cq)_{|\boldsymbol{j}|}} \cdot \left(\frac{q}{at}\right)^{|\boldsymbol{j}|} q^{\sum_{r=1}^n (r-1)j_r}$$
$$= \frac{(q)_{|\boldsymbol{k}|}}{(cq)_{|\boldsymbol{k}|}} \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s-k_r+k_s}x_r/x_s}{1-q^{-k_r+k_s}x_r/x_s}$$
$$\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_r}x_r/x_s\right)_{j_r}}{\left(q^{1-k_r+k_s}x_r/x_s\right)_{j_r}} \frac{(c)_{|\boldsymbol{j}|}(1/atq^{|k|})_{|\boldsymbol{j}|}}{(1/t)_{|\boldsymbol{j}|}(q^{-|\boldsymbol{k}|})_{|\boldsymbol{j}|}} q^{\sum_{r=1}^n rj_r}$$

Taking the limit $t\to 0$ on both sides of this identity gives

$$\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_s/x_r}{1-x_s/x_r} \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_s/x_r\right)_{j_r}}{\left(qx_s/x_r\right)_{j_r}}$$
$$\times \frac{(aq^{|\boldsymbol{k}|},c)_{|\boldsymbol{j}|}}{(cq)_{|\boldsymbol{j}|}} \left(-\frac{1}{a}\right)^{|\boldsymbol{j}|} q^{\sum_{r=1}^n rj_r-\binom{|\boldsymbol{j}|}{2}}$$
$$= \frac{(q)_{|\boldsymbol{k}|}}{(cq)_{|\boldsymbol{k}|}} \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s-k_r+k_s}x_r/x_s}{1-q^{-k_r+k_s}x_r/x_s}$$
$$\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_r}x_r/x_s\right)_{j_r}}{\left(q^{1-k_r+k_s}x_r/x_s\right)_{j_r}} \frac{(c)_{|\boldsymbol{j}|}(1/a)^{|\boldsymbol{j}|}}{(q^{-|\boldsymbol{k}|})_{|\boldsymbol{j}|}} q^{\sum_{r=1}^n rj_r-|\boldsymbol{k}||\boldsymbol{j}|} \tag{9.7}$$

We set $c\mapsto aq^{|\boldsymbol{k}|}, b\mapsto c, d\mapsto cq, e\mapsto 1/t$ in (9.6) to get

$$\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_s/x_r}{1-x_s/x_r} \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_s/x_r\right)_{j_r}}{\left(qx_s/x_r\right)_{j_r}}$$
$$\times \frac{(aq^{|\boldsymbol{k}|})_{|\boldsymbol{j}|}(c)_{|\boldsymbol{j}|}}{(cq)_{|\boldsymbol{j}|}(1/t)_{|\boldsymbol{j}|}} \cdot \left(\frac{q}{at}\right)^{|\boldsymbol{j}|} q^{\sum_{r=1}^n (r-1)j_r}$$
$$= \frac{(1/atq^{|\boldsymbol{k}|})_{|\boldsymbol{k}|}}{(1/t)_{|\boldsymbol{k}|}} \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s-k_r+k_s}x_r/x_s}{1-q^{-k_r+k_s}x_r/x_s}$$
$$\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_r}x_r/x_s\right)_{j_r}}{\left(q^{1-k_r+k_s}x_r/x_s\right)_{j_r}} \frac{(aq^{|\boldsymbol{k}|})_{|\boldsymbol{j}|}(q)_{|\boldsymbol{j}|}}{(cq)_{|\boldsymbol{j}|}(aqt)_{|\boldsymbol{j}|}} q^{\sum_{r=1}^n rj_r}.$$

Taking the limit $t \to 0$ on both sides of this identity yields

$$
\begin{aligned}
&\sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_s/x_r}{1-x_s/x_r} \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_s/x_r\right)_{j_r}}{(qx_s/x_r)_{j_r}} \\
&\times \frac{(aq^{|\boldsymbol{k}|},c)_{|\boldsymbol{j}|}}{(cq)_{|\boldsymbol{j}|}} \left(-\frac{1}{a}\right)^{|\boldsymbol{j}|} q^{\sum_{r=1}^n rj_r-\binom{|\boldsymbol{j}|}{2}} \\
&= (1/a)^{|\boldsymbol{k}|}q^{-|\boldsymbol{k}|^2} \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s-k_r+k_s}x_r/x_s}{1-q^{-k_r+k_s}x_r/x_s} \\
&\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_r}x_r/x_s\right)_{j_r}}{(q^{1-k_r+k_s}x_r/x_s)_{j_r}} \frac{(aq^{|\boldsymbol{k}|})_{|\boldsymbol{j}|}(q)_{|\boldsymbol{j}|}}{(cq)_{|\boldsymbol{j}|}} q^{\sum_{r=1}^n rj_r}
\end{aligned} \tag{9.8}
$$

Then the identity (9.3) follows easily by combining (9.7) and (9.8) and then replacing x_r by $x_rq^{k_r}$ in the resulting identity.

We now show (9.2). Let $c=d=\gamma=0$ in the identity of Theorem 1.9 and then replace β by q/α in the resulting identity. We get

$$
\begin{aligned}
&\frac{(q,\alpha q,\alpha a_1\cdots a_nb/q^n)_\infty}{(\alpha a_1\cdots a_nq^{1-n},\alpha b,a_1\cdots a_nq^{1-n})_\infty} \\
&= \sum_{\substack{k_r\ge 0\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{(qx_r/x_sa_s)_{k_r}}{(qx_r/x_s)_{k_r}} \\
&\times \frac{(1-\alpha q^{2|\boldsymbol{k}|})(\alpha)_{|\boldsymbol{k}|}}{(1-\alpha)(\alpha a_1\cdots a_nq^{1-n})_{|\boldsymbol{k}|}} (a_1\cdots a_n)^{|\boldsymbol{k}|}q^{\sum_{r=1}^n(r-1)k_r-n|\boldsymbol{k}|} \\
&\times \sum_{\substack{0\le m_r\le k_r\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{m_r-m_s}x_r/x_s}{1-x_r/x_s} \\
&\quad\times \prod_{r,s=1}^{n} \frac{(q^{-k_s}x_r/x_s)_{m_r}}{(qx_r/x_s)_{m_r}} \frac{(\alpha q^{|\boldsymbol{k}|},q)_{|\boldsymbol{m}|}}{(\alpha b)_{|\boldsymbol{m}|}} q^{\sum_{r=1}^n rm_r}.
\end{aligned}
$$

Replacing a by α, c by $\alpha b/q$ in the identity (9.3) and then substituting the resulting identity into the above identity we have

$$
\begin{aligned}
&\frac{(q,\alpha q,\alpha a_1\cdots a_nb/q^n)_\infty}{(\alpha a_1\cdots a_nq^{1-n},\alpha b,a_1\cdots a_nq^{1-n})_\infty} \\
&= \sum_{\substack{k_r\ge 0\\ r=1,2,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{k_r-k_s}x_r/x_s}{1-x_r/x_s} \prod_{r,s=1}^{n} \frac{(qx_r/x_sa_s)_{k_r}}{(qx_r/x_s)_{k_r}} \\
&\times \frac{(1-\alpha q^{2|\boldsymbol{k}|})(\alpha,q)_{|\boldsymbol{k}|}(\alpha a_1\cdots a_n)^{|\boldsymbol{k}|}}{(1-\alpha)(\alpha a_1\cdots a_nq^{1-n},\alpha b)_{|\boldsymbol{k}|}} q^{\sum_{r=1}^n(r-1)k_r-n|\boldsymbol{k}|+|\boldsymbol{k}|^2} \\
&\times \sum_{\substack{0\le j_r\le k_r\\ r=1,\cdots,n}} \prod_{1\le r<s\le n} \frac{1-q^{j_r-j_s}x_r/x_s}{1-x_r/x_s} \cdot q^{\sum_{r=1}^n rj_r-|\boldsymbol{k}||\boldsymbol{j}|} \\
&\times \prod_{r,s=1}^{n} \frac{\left(q^{-k_s}x_r/x_s\right)_{j_r}}{(qx_r/x_s)_{j_r}} \frac{(\alpha b/q)_{|\boldsymbol{j}|}(1/\alpha)^{|\boldsymbol{j}|}}{(q^{-|\boldsymbol{k}|})_{|\boldsymbol{j}|}}.
\end{aligned}
$$

Then the identity (9.2) follows readily by replacing a_i by q/y_i for $i = 1, 2, \cdots, n$, b by bq/a, and α by a in this identity. □

References

[1] G.E. Andrews, Parity in partition identities, Ramanujan J. 23(1–3)(2010), 45–90.

[2] G.E. Andrews, The fifth and seventh order mock theta functions. Trans. Am. Math. Soc. 293(1)(1986), 113–134

[3] G.E. Andrews, The theory of partitions, in Encyclopedia of Mathematics and its Applications (G.-C. Rota, ed.), Vol. 2, Addison-Wesley, Reading, MA, 1976.

[4] G. Bhatnagar and S. Rai, Expansion formulas for multiple basic hypergeometric series over root systems, Adv. in Appl. Math. 137(2022), Paper No. 102329, 29 pp.

[5] J. Cao, Homogeneous q-difference equations and generating functions for q-hypergeometric polynomials. Ramanujan J. 40(1)(2016), 177–192.

[6] J. Cao, Homogeneous q-partial difference equations and some applications. Adv. in Appl. Math. 84(2017), 47–72.

[7] H.H. Chan and Z.-G. Liu, On certain series of Hecke-type. New Zealand J. Math. 48(2018), 1–10.

[8] J.-P. Fang, A note on the Rogers-Fine identity. Electron. J. Combin. 14(1)(2007), Note 17, 5 pp.

[9] J.-P. Fang, Generalizations of Milne's $U(n+1)$ q-Chu-Vandermonde summation. Czechoslovak Math. J. 66(141)(2016), 395–407.

[10] J.-P. Fang, q-Differential operator identities and applications, J. Math. Anal. Appl., 332(2007), 1393–1407.

[11] N.J. Fine, Basic Hypergeometric Series and Applications, With a foreword by George E. Andrews. Mathematical Surveys and Monographs, 27. American Mathematical Society, Providence, RI, 1988.

[12] G. Gasper and M. Rahman, Basic Hypergeometric Series, Volume 96 of Encyclopedia of Mathematics and its Applications, Cambridge University Press, Cambridge, second edition, 2004.

[13] R.W. Gosper, Experiments and discoveries in q-trigonometry, in: F.G. Garvan, M.E.H. Ismail (Eds.), Symbolic Computation, Number Theory, Special Functions, Physics and Combinatorics, Kluwer, Dordrecht, Netherlands, 2001, pp.79–105.

[14] M. Ismail, M. Rahman, K. Suslov, Some summation theorems and transformations for q-series, Can. J. Math. 49(3)(1997), 543–567.

[15] Z.-G. Liu, A q-series expansion formula and the Askey-Wilson polynomials. Ramanujan J., 30(2)(2013), 193–210.

[16] Z.-G. Liu, An expansion formula for q-series and applications. Ramanujan J. 6(4)(2002), 429–447.

[17] Z.-G. Liu, An extension of the non-terminating ${}_6\phi_5$ summation and the Askey-Wilson polynomials, J. Difference Equ. Appl. 17(10)(2011), 1401–1411.

[18] Z.-G. Liu, On the q-derivative and q-series expansions, Int. J. Number Theory, 9(8)(2013), 2069–2089.

[19] Z.-G. Liu, On the q-partial differential equations and q-series. In The legacy of Srinivasa Ramanujan, Volume 20 of Ramanujan Math. Soc. Lect. Notes Ser., pages 213–250. Ramanujan Math. Soc., Mysore, 2013.

[20] S.C. Milne, A q-analog of a Whipple's transformation for hypergeometric series in $U(n)$. Adv. Math., 108(1)(1994), 1–76.

[21] S.C. Milne, Balanced ${}_3\phi_2$ summation theorems for $U(n)$ basic hypergeometric series, Adv. Math. 131(1)(1997), 93–187.

[22] S.C. Milne and G.M. Lilly, Consequences of the A_l and C_l Bailey transform and Bailey lemma. Formal power series and algebraic combinatorics (Montreal, PQ, 1992). Discrete Math. 139(1-3)(1995), 319–346.

[23] S.C. Milne and J.W. Newcomb. Nonterminating q-Whipple transformations for basic hypergeometric series in $U(n)$. In Partitions, q-series, and modular forms, volume 23 of Dev. Math., pages 181–224. Springer, New York, 2012.

[24] S.C. Milne and J.W. Newcomb, $U(n)$ very-well-poised ${}_{10}\phi_9$ transformations. J. Comput. Appl. Math. 68(1–2)(1996), 239–285.

[25] L.J. Rogers, On two theorems of combinatory analysis and some allied identities, Proc. London Math. Soc. 16(1917), 315–336.
[26] M. J. Schlosser, Hypergeometric and basic hypergeometric series and integrals associated with root systems. In Encyclopedia of Special Functions: The Askey-Bateman Project, Volume II, Multivariable Special Functions, T. H. Koornwinder and J. V. Stokman (eds.), pages 122–158. Cambridge University Press, 2020.
[27] M.J. Wang, Generalizations of Milne's $U(n+1)$ q-binomial theorems. Comput. Math. Appl. 58(1)(2009), 80–87.
[28] G.N. Watson, A new proof of the Rogers-Ramanujan identities, J. London Math. Soc. 4(1)(1929), 4–9.

School of Mathematics and Statistics, HNP-LAMA, Central South University, Changsha 410083, Hunan, People's Republic of China
Email address: yuhelingyun@foxmail.com; yuhe001@foxmail.com