

COVID-19 mortality prediction: A case study for İstanbul

Erkan Yilmaz^a, Özgür Ökcü^b, Ekrem Aydiner^{b,*}

^a*Institute of Graduate Studies in Science, Istanbul University, 34134, İstanbul, Turkey*

^b*Department of Physics, Faculty of Science, Istanbul University, İstanbul, 34134, Turkey*

Abstract

In this paper, we use SEIR equations to make predictions for the number of mortality due to COVID-19 in İstanbul. Using excess mortality method, we find the number of mortality for the previous three waves in 2020 and 2021. We show that the predictions of our model is consistent with number of mortalities for each wave. Furthermore, we predict the number of mortality for the second wave of 2021. We also extend our analysis for Germany, Italy and Turkey to compare the basic reproduction number R_0 for Istanbul. Finally, we calculate the number of infected people in Istanbul for herd immunity.

Keywords: Sars Cov-2, Covid-19, Excess Mortality, SIR and SEIR Model, Pandemic, Mortality Prediction, Basic Reproduction Number

1. Introduction

The world has struggled to control the deadly coronavirus disease 2019 (COVID-19) which is caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). After the initial outbreak in China, the novel COVID-19 spread across the world in a short time. COVID-19 was declared a pandemic on 11th March 2020 by World Health Organisation (WHO) [1]. In the same day, the first case in Turkey was reported by the Ministry of Health [2].

In order to struggle with diseases more effectively and detect their characteristics such as symptoms, spread dynamics, transmission, etc., the humans firstly began to record the diseases. Those records paved the way to fight against the diseases. The scientists have modelled the pandemics to reveal the pandemics characteristics. Modelling a pandemic is very important since it may provides how to control the disease and how to reduce the mortality. For example, London mortality statistics were published weekly in Bills of Mortality from the 17th century to 1830s [3]. In 1662, Graunt studied mortality rates and causes in his book Natural and Political Observations Made Upon the Bills of Mortality [4].

The first mathematical model on the characteristic behaviours of contagious diseases was proposed by Bernoulli in 1766 [5]. He investigated the smallpox disease and variolation method which aimed to immunize people against smallpox with the material taken from patients. Bernoulli showed that variolation was necessary in his mathematical model [5]. In 1906, Hamer analysed the measles epidemic and proposed a discrete time model [6]. This model is important since it shows that the new cases of an epidemic depend on the numbers of patients and individuals susceptible to disease. In 1911, Ross proposed a differential equation model related to the number of cases and control of malaria [7]. In 1927, Kermack and McKendrick published their paper entitled A Contribution to the Mathematical Theory of Epidemics [8]. In their paper, they considered an interactive model for an isolated society. They proposed SIR model. In Refs.[9, 10], they made the model more useful by adding population dynamics and developed SIR model.

Building a mathematical model provides predictions for pandemics, effective control and political strategies [11, 12, 13, 14, 15]. It also provides accurate future predictions and effective struggle against the diseases [16, 17, 18]. Differential equations can be used to predict the time evolution of the pandemic. Although the mathematical models are useful to predict the characteristics of a disease, they have some difficulties. It is not easy to determine the model parameters. Besides, the factors such as mutations, health infrastructure, disasters, etc. affect the evolution of a disease. Therefore, the predictions from differential equations may become harder.

*Corresponding author

Email addresses: erkany@ogr.iu.edu.tr (Erkan Yilmaz), ozgur.okcu@ogr.iu.edu.tr (Özgür Ökcü), ekrem.aydiner@istanbul.edu.tr (Ekrem Aydiner)

In order to determine the COVID-19 characteristics, the COVID-19 has been widely studied in the literature for two years [16, 19, 20, 21, 22, 23, 24]. In order to predict the trend of COVID-19 in Italy, Wangping et. al. used the extended SIR model [16]. They also estimated the basic reproductive number R_0 from Markov Chain Monte Carlo methods. Danon et. al. adapted an existing national-scale metapopulation model to get the spread of CoVID-19 in England and Wales [19]. They found that the size and spread rate of epidemic highly depend on the seasonal transmission. Fanalli and Piazza analyzed the temporal dynamics of the COVID-19 outbreak in China, Italy and France [20]. Roosa et. al. [21] reported short-term estimation of the cumulative number of reported cases of the COVID-19 epidemic in Hubei province and other provinces in China as of February 9, 2020. Using the data until February 9, 2020, they also estimated the number of reported cases between 37,415 and 38,028 in Hubei Province and 11,588–13,499 in other provinces by February 24, 2020. In Ref. [22], Li et. al. studied the transmission process of the COVID-19 via based on the official data. Based on Ref. [23], Mangoni and Pistilli [24] developed a generalised SEIR model to estimate the dynamics of COVID-19.

Up to now, Covid-19 has the best statistics. However, recording exact and reliable data include serious difficulties. For example, PCR tests expected success rate is very low. In addition to medical issues, political and economic issues affect the reliability of these data. On the other hand, mortality records of a country are more reliable, and excess mortality data are independent from political, economic and medical issues.

As before mentioned, the differential models provides to determine the behaviours of an epidemic despite its disadvantages. SEIR is the one of these models [11]. In thse paper, we use the SEIR model to predict the COVID-19 waves for Istanbul. To the best of our knowledge, we calculate the parameters of SEIR model from excess mortality methods for the first time. Therefore, we believe our results are reliable since they are based on excess mortality.

The paper is organised as follows: In Section 2, we review the SIR and SEIR models. We present the details on determining the parameters β , η and ϵ of SEIR model, excess mortality methods, and time series for COVID-19. In Section 3, we determine the time series from excess mortality methods. We also find the SEIR model parameters. We calculate the basic reproduction number R_0 for each wave. We make predictions for the second wave of 2021. In order to compare to value of R_0 for Istanbul, we also computed R_0 for Germany, Italy and Turkey. In Section 4, we finally discuss our results and estimate the number of infected people for herd immunity.

2. Model

2.1. Mathematical models of the pandemic

In this study, we use the SEIR model [11] which is the modification of SIR model. Just like SIR model, SEIR is also a set of coupled differential equations. It is possible to construct many disease spread models from SIR and modified SIR models[8, 9, 10, 15, 16, 23, 24]. Recently, much attention has been paid to these models due to COVID-19. Because these models provide to estimate numbers of dead, recovered and infected individuals for local and global regions during a pandemic. These models completely depend on the parameters and give the reliable results. However, contagious diseases may regard as catastrophic events. Because stochastic factors such as vaccine, mutations, curfew or quarantine precautions may make the process more chaotic. Therefore, it is difficult to make the accurate predictions. On the other hand, it is not impossible to make accurate predictions for a short term since all stochastic effects can be expressed in macroscopic forms in the limit cases.

SIR equations are given by [8]

$$\begin{aligned}\frac{dS}{dt} &= -\beta SI, \\ \frac{dI}{dt} &= \beta SI - \eta I, \\ \frac{dR}{dt} &= \eta I,\end{aligned}\tag{1}$$

where S , I and R correspond to susceptible, infected and recovered individuals, respectively. Total probability is $S + I + R = 1$. Let us explain S , I and R , respectively. S represents the individuals that may have tendency to get a disease. I represents the individuals that are infected and spread a disease. R represents the individuals that have not got any possibilities to get a disease. Individuals which is recovered or died from a disease can be considered in R . The parameter β corresponds to disease spread speed while the parameter η is related with illness duration. We define the parameter basic reproduction number as $R_0 = \frac{\beta}{\eta}$. It is important for the disease spread dynamics.

There are some disadvantages of SIR model. Getting a disease and contagiousness are considered as mono-type. In fact, these factors vary from individual to individual. The spread rate of an epidemic may depend on seasons. A disease may have different effects from human to human since humans have different genetics.

A more realistic model is the modification of SIR model, i.e., SEIR model [11]. In SEIR model, there is an extra parameter E which represents the number of individuals exposed to COVID-19. When people exposed to coronavirus, they do not immediately become ill. There is a time period between getting the virus and transmitting to humans. This period is called incubation period. The incubation period changes the date and value of peak. SEIR equations are given by [11]

$$\begin{aligned}
 \frac{dS}{dt} &= -\beta SI \\
 \frac{dE}{dt} &= \beta SI - \epsilon E \\
 \frac{dI}{dt} &= \epsilon E - \eta I \\
 \frac{dR}{dt} &= \eta I
 \end{aligned}
 \tag{2}$$

where the parameter ϵ is related with the incubation period of epidemic.

2.2. Determining the model parameters

Excess mortality determines the number of deaths from all reported number of deaths during a crisis and expected deaths for the same time in the absence of the crisis [25, 26]. Excess mortality method was used to determine the mortalities of the Great Plague of London in 1665 [27], influenza epidemic in London in 1875 [28, 29], influenza pandemics of 1918, 1957, 1968, 2009 [30, 31], as well as seasonal influenza epidemics [32]. Excess mortality is given by [25]

$$\text{Excess Mortality} = \text{Reported Deaths} - \text{Expected Deaths}.
 \tag{3}$$




Figure 1: Numbers of reported death for 2015-2019 (yellow), 2020 (blue) and 2021 (red).

3. Istanbul Covid-19 Waves

3.1. The first wave of 2020

This method is more reliable to detect the number of deaths from COVID-19 [33]. Using excess mortality method, WHO has obtained the number of deaths from seasonal influenza, pandemics and other public health threats since 2009 [34].




Figure 2: Using excess mortality method, numbers of death due to the COVID-19 in 2020 (blue) and 2021 (red).

We take numbers of daily deaths in İstanbul from 2015 to 2019 [35] to predict the expected number of death in 2020-2021. In the calculation of expected deaths, 40% deaths in 2019, 30% deaths in 2018, 20% deaths in 2017, 5% deaths in 2016 and 5% deaths in 2015 were taken. In order to make our time series more reliable, we consider the following average over seven days

$$d = \frac{d_i + d_{i-1} + d_{i-2} + d_{i-3} + d_{i-4} + d_{i-5} + d_{i-6}}{7} \quad (4)$$

In Fig. (1), we present the reported numbers of deaths in 2015-2019, 2020, 2021. Yellow line represents the numbers of average mortality between 2015 and 2019. Note that value of yellow line is between 200 and 250. Furthermore, numbers of reported death for 2020 (blue line) and 2021 (red line) reach bigger values than yellow line's. It is clearly obvious that numbers of mortality dramatically increased during COVID-19 pandemic. In Fig. (2), using excess mortality method for the data between 2015-2019, we show numbers of death in 2020 and 2021 due to COVID-19. In this figure, it is clear that there are two COVID-19 waves for each year.

The first case in Turkey was reported on March 2020 and COVID-19 quickly spread across the country. The first wave in Istanbul occurred in the spring of 2020. Pandemic started on 15th March 2020 and 1653 people died from COVID-19 until pandemic's first peak 10th April 2020. Then it started to decrease. During the decrease of the first wave, 2798 people died from COVID-19.

The first wave in Istanbul reached the peak in 26 days, then it decreased in 43 days. The total time of the first wave is 69 days. 4451 people died from COVID-19 during the first wave. In Fig. (3), we show daily and total mortalities. It can be seen that positive slope of graphics is bigger than negative slope. This implies that the value of R_0 is high.

In order to detect the parameters β , η and ϵ of the first wave, we must obtain the numerical solutions. First of all, we define the SEIR model equations and data in matlab program. We obtain the first wave parameter within 1 or 2 percentage of error. The parameters and their percentages of error can be found in Fig.(4). Comparing SEIR model with data gives more than 10,000 results within 5 percentage of error. Best ten results are listed in Table 1. The results in Table 1 are obtained less than 1 percentage of error. We use the fourth result in Table 1 to produce Fig. (3) since it is close to average value.

3.2. The Second Wave of 2020

The second wave started on 22nd October 2020. It peaked on 26th November 2020, and ended on 14th January 2021. Mortality increased for 35 days, and decreased for 49 days. The second wave lasted 84 days. 4754 people




Figure 3: Mortality numbers of the first wave in 2020 (a)Daily (b)Total




Figure 4: Percentages of error for the first wave in 2020 (a) The parameter β (b) The parameter η for the first wave.

Table 1: The First Wave Parameters of 2020

$R_0 = \frac{\beta}{\eta}$	β	η	ϵ	Error %
3.19	0.232846715	0.072992701	3	0.867405
3.11	0.232089552	0.074626866	3	0.868163
3.22	0.23	0.071428571	4	0.868588
3.16	0.232352941	0.073529412	3	0.868712
3.22	0.233333333	0.072463768	3	0.868797
3.14	0.232592593	0.074074074	3	0.86956
3.14	0.22919708	0.072992701	4	0.869781
3.25	0.230496454	0.070921986	4	0.869876
3.08	0.231578947	0.07518797	3	0.870172
3.17	0.229710145	0.072463768	4	0.870274

died from COVID-19 until the peak. During the decrease of the second wave, 6433 people died. 11187 people died from COVID-19 during the second wave.

Comparing the first and second waves, we can easily see differences between them. The second wave is more symmetrical than the first wave. It means that the value of R_0 is more smaller than the first wave's. The number of mortality for the second wave is more higher than the first wave's. The mutation of the virus or seasonal effects may have caused more deaths.

The numbers of mortality for the second wave are given in Fig. (5). We calculate the parameters for the second wave. We found the percentages of error for the parameters β and η in Fig. (6). In Fig. (6.a), one can see that the parameter β has a low percentage of error between 0.215 and 0.235. Its percentage of error has the lowest value near 0.23. In Fig. (6.b), the parameter η has a low percentage of error between 0.13 and 0.15. It has the lowest percentage of error near 0.14. Ten results which give the lowest percentage of error are given in Table (2). We use the first result in Table (2) to produce the Fig. 5.




Figure 5: Mortality numbers of the second wave in 2020 (a)Daily (b)Total




Figure 6: Percentages of error for the second wave in 2020 (a) The parameter β (b) The parameter η for the second wave.

Table 2: The Second Wave Parameter of 2020

$R_0 = \frac{\beta}{\eta}$	β	η	ϵ	Error %
1.62	0.231098431	0.142653352	3	2.655150333
1.62	0.230769231	0.142450142	3	2.656684896
1.62	0.231428571	0.142857143	3	2.661003256
1.62	0.229787234	0.141843972	3	2.661754709
1.63	0.230878187	0.141643059	3	2.662501288
1.62	0.230113636	0.142045455	3	2.662901328
1.63	0.229901269	0.141043724	3	2.663813534
1.63	0.231205674	0.141843972	3	2.664873865
1.62	0.230440967	0.142247511	3	2.665545267
1.63	0.229577465	0.14084507	3	2.666175529

3.3. The first wave of 2021

The first wave of 2021 started on 11th March 2021. It peaked on 20th April 2021, and ended on 8th June 2021. Mortality increased for 44 days, and decreased for 45 days. 3886 people died until the peak. During the decrease of the wave, 4422 people died. 8308 people died during the first wave of 2021.

The numbers of mortality are more higher than the numbers of mortality in the first wave of 2020. The first wave of 2021 is more symmetrical than the first wave of 2020.

The numbers of mortality are given in Fig. (7). The percentages of error for parameters β and η are given in Fig. (8). In Fig. (8.a), it can be seen that the parameter β has the low percentage of error between 0.21 and 0.24. Its percentage of error is the lowest near 0.23. In Fig (8.b), one can see that the parameter η has a low percentage of error between 0.13 and 0.15. In Table 3, its percentage of error is the lowest near 0.14. We list the parameters which have the lowest error of percentage. We use the second result of Table (3) to produce the Fig. (7).

3.4. The predictions on the second wave of 2021

Up to now, we have modelled the COVID-19 by using SEIR and found the model parameters β , η and ϵ . These parameters determine the behaviours of the pandemic. The parameter ϵ does not effect the trend of epidemic but




Figure 7: Mortality numbers of the first wave in 2021 (a)Daily (b)Total




Figure 8: Percentages of error for the first wave in 2021 (a) The parameter β (b) The parameter η for the first wave.

Table 3: The First Wave Parameters of 2021

$R_0 = \frac{\beta}{\eta}$	β	η	ϵ	Error %
1.61	0.23	0.142857143	3.56	1.324750255
1.61	0.229344729	0.142450142	3.75	1.328061544
1.6	0.228571429	0.142857143	4.19	1.330648971
1.61	0.228693182	0.142045455	3.96	1.331516693
1.6	0.228571429	0.142857143	4.12	1.331578669
1.61	0.228045326	0.141643059	4.19	1.335107124
1.6	0.227920228	0.142450142	4.44	1.335252725
1.62	0.229461756	0.141643059	3.53	1.338365318
1.61	0.22740113	0.141242938	4.53	1.340113226
1.62	0.228813559	0.141242938	3.73	1.340357901

changes the peak date of epidemic. Now, using the Tables 1, 2 and 3, we will predict the numbers of deaths for the second wave of 2021. In Table 4, we give the results of each wave for lowest error of percentage.

Table 4: The averages of ten results which have the lowest percentages of error.

Wave	$R_0 = \frac{\beta}{\eta}$	β	η
First wave of 2020	3.168	0.231419776	0.073068182
Second wave of 2020	1.624	0.230520067	0.14194734
First wave of 2021	1.609	0.228682277	0.142128916

As it can be seen in Table 4, the values of R_0 are 3.168 for the first wave of 2020, 1.624 for the second wave of 2020 and 1.609 for the first wave of 2021. The second and third values of R_0 are lower than the first value. It means that the spread speed decreased. Because taking precautions against COVID-19 is very effective. We assume that the parameters value of the second wave of 2021 is nearly close to last two waves. These parameters may decrease due to vaccination or may increase due to mutation. Therefore, we estimate lower and upper bounds for the new

wave. In Fig. (9), we present our predictions for the second wave of 2021. In the figure, we show the numbers of mortality for 2020 and 2021. It can easily be seen that our model (black line) is fit with 2020 (blue line) and 2021 (red line) data. We present our predictions with black dashed line.




Figure 9: Numbers of mortality for last three COVID-19 waves. Dashed black lines correspond to predictions for the current COVID-19 wave.

3.5. Calculation of R_0 for Turkey, Germany and Italy

Up to now, we have considered R_0 for Istanbul. Taking precautions cause to decrease the number of cases. Therefore, the parameter R_0 has tendency to decrease. In this section, we want to study the behaviour of R_0 for Germany, Italy and Turkey. The value of basic production number R_0 for the first wave of 2020 is more bigger than the value of R_0 for the other waves. In order to test it, we found the values of R_0 for the numbers of case for Germany, Italy and Turkey.

We take the data from the web site of Republic of Turkey Ministry of Health [2]. The first wave of 2020 and the first wave of 2021 are estimated for the numbers of case. However, we take into account the numbers of mortality for the second wave of 2020 due to the change of the COVID-19 new case announcement system. In Table 5, one can see the values of R_0 decreasing for Turkey.

Table 5: R_0 for the COVID-19 Waves in Turkey

Year	Wave	$R_0 = \frac{\beta}{\eta}$
2020	First	4.2 - 4.7
2020	Second	1.3 - 1.47
2021	First	1.75 - 2.1

In Table 6, using the data in Ref. ([36]), we give R_0 for Germany. It can be seen that R_0 gets the values between 5 and 9 for the first wave of 2020. If such huge values were permanent, all the people in the world would nearly have COVID-19. For the second wave of 2020, R_0 has the values between 1.3 and 2.1. Finally, for the first wave of 2021, R_0 has the values between 1.4 and 1.8.

For Italy, using the data in Ref. ([36]), one can see the values of R_0 in Table 7. The values of R_0 are between 4 and 5 for the first wave of 2020. The values of R_0 are between 3.5 and 5 for the second wave of 2020. For the first wave of 2021, R_0 has the values between 1.3 and 1.9.

Our analysis show that R_0 may have the tendency to decrease in the presence of precautions.

Table 6: R_0 for the COVID-19 Waves in Germany

Year	Wave	$R_0 = \frac{\beta}{\eta}$
2020	First	5.0 - 9.0
2020	Second	1.3 - 2.1
2021	First	1.4 - 1.8

Table 7: R_0 for the COVID-19 Waves in Italy

Year	Wave	$R_0 = \frac{\beta}{\eta}$
2020	First	4.0 - 5.0
2020	Second	3.5 - 5.0
2021	First	1.3 - 1.9




Figure 10: R_0 for Istanbul (a) The first wave of 2020 (b) The second wave of 2020 (c) The first wave of 2021

4. Conclusions and discussions

In this paper, using the reported numbers of mortality for 2014-2021, we have determined the numbers of COVID-19 mortality in Istanbul via excess mortality method. Our analysis shows that there are two deadly pandemic waves for a year. We have modelled each wave via SEIR equations. We have calculated the parameters β , η , ϵ and R_0 . Our model is compatible with the previous three waves. Then, we have estimated the second wave of 2021. Interestingly, we have found that R_0 may have tendency to decrease for precautions. In order to confirm the behaviour of R_0 , we have extended our analysis for Germany, Italy and Turkey. It seems R_0 may decrease for precautions.¹

Before finishing the paper, we give some comments on the number of infected people to gain herd immunity in Istanbul. Solving Eq. (1), one can determine the how many individuals get infected with COVID-19 until the end of the pandemic.² Using

$$S + I + R = 1 \quad (5)$$

¹Mutation may change this case. Omicron is one of the recent mutation, which is very contagious. In this paper, we do not take into account any mutations. We will left this case for future study.

²In fact, we can use SEIR equations, but the result does not change. The extra parameter ϵ only changes the peak date of an epidemic.

Table 8: R_0 and R_f values for different waves in the last two years

Wave	R_0	R_f
First wave of 2020	2.5	89.3%
	4.5	98.9%
Second wave of 2020	1.6	64.2%
	1.65	66.8%
First wave of 2021	1.6	64.2%
	1.63	65.8%

with the initial conditions $S(0) = 1$ and $R(0) = 0$, one finds

$$S(t) = e^{-R_0 R(t)}. \quad (6)$$

At the end of COVID-19, the value of I is zero. Therefore, employing Eq. (6) in Eq. (5), we obtain

$$R_f + e^{-R_0 R_f} = 1, \quad (7)$$

where R_f is the number of recovered people at the end of COVID-19. We show numerical solutions of this equation in Fig. (11). In Fig. (11), one can see that all people nearly get infected for $R_0 = 5$. For example, varying the




Figure 11: The behavior of R_f with respect to R_0 .

value of R_0 from 7 to 5 does not effect the number of infected people. However, varying the value of R_0 from 5 to 3 effects the number of infected people.

In Fig. (10), we show the R_0 for the previous three waves in Istanbul. Using the data in Fig. (10), we give the percentage of infected people for herd immunity in Table (8). One can easily see that R_f depends on R_0 . For the first wave of 2020, 89.3%-98.9% of people in Istanbul should get infected to gain herd immunity. The rates for two next waves decrease. Therefore, number of infected people for herd immunity decreases.

We see that the first and second main waves of 2020 and 2021 show seasonal dependence. It can be concluded that the first main mortality wave appears in the spring period which correspond to the increasing of the human

metabolic activity. Similarly, the second main mortality wave appears in between December and February. This period corresponds to the decreasing of the human metabolic activity and increasing air pollution.

The results obtained in this type of study may contribute to the development strategies of the control, quarantine, physical distance, vaccination and determination of health load. On the other hand, the presented model can be used to make predictions not only local region but also country scale.

Acknowledgments

Authors would like to thank Professor Ayşe H. Bilge (Kadir Has University) for useful discussions.

References

- [1] World Health Organization, Coronavirus disease 2019 (COVID-19): Situation report, 52 (2020). Available online at: <https://apps.who.int/iris/handle/10665/331476>
- [2] Republic of Turkey Ministry of Health, (2020). Available online at: <https://covid19.saglik.gov.tr/TR-66494/pandemi.html>
- [3] K. Heitman, Authority, autonomy and the first London Bills of Mortality, *Centaurus*, **62**, (2020) 275–284.
- [4] D. V. Glass, John Graunt and his natural and political observations, *Proc. R. Soc. Lond. B.*, **159**, (1963) 2-37.
- [5] D. Bernoulli, S. Blower, An attempt at a new analysis of the mortality caused by smallpox and of the advantages of inoculation to prevent it, *Rev. Med. Virol*, **14**, (2004) 275-288.
- [6] W. H. Hamer, *The Milroy Lectures*, Bedford Press (1906).
- [7] N. Bacaer, *Ross and Malaria (1911)*, A short history of mathematical population dynamics, Springer (2011).
- [8] W. O. Kermack, A. G. McKendrick, (1927). Contributions to the mathematical theory of epidemics, *Proc. R. Soc. Lond. A*, **115**, (1927) 700-721.
- [9] W. O. Kermack, A. G. Kendrick, A Contribution to the mathematical theory of epidemics part II. The Problem of Endemicity, *Proc. R. Soc. Lond. A*, **138**, (1932) 55-83.
- [10] W. O. Kermack, A. G. McKendrick, Contributions to the mathematical theory of epidemics. III.—Further studies of the problem of endemicity, *Proc. R. Soc. A*, **141**, (1933) 94-122.
- [11] R. M. Anderson, R. M. May, B. Anderson, *Infectious diseases of humans: Dynamics and control*, Oxford Science Publications (1991).
- [12] S. V. Scarpino, G. Petri, On the predictability of infectious disease outbreaks, *Nat. Commun.*, **10**, (2019) 898.
- [13] M. Chinazzi, J. T. Davis, M. Ajelli, C. Gioannini, M. Litvinova, S. Merler, A. P. Y. Piontti, K. Mu, L. Rossi, K. Sun, C. Viboud, X. Xiong, H. Yu, M. E. Halloran, I. M. Longini Jr, A. Vespignani, The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak, *Science*, **368**, (2020) 395–400.
- [14] S. Rüdiger, A. Plietzsch, F. Sagués, Epidemics with mutating infectivity on small-world networks, *Sci. Rep.*, **10**, (2020) 1–11.
- [15] A. H. Bilge, Ö. Pekcan, M. V. Gürol, Application of epidemic models to phase transitions, *Phase Transitions*, **85**, (2012) 1009-1017.
- [16] J. Wangping, H. Ke, S. Yang, C. Wenzhe, W. Shengshu, Y. Shanshan, W. Jianwei, K. Fuyin, T. Penggang, L. Jing, L. Miao, H. Yao, Extended SIR prediction of the epidemics trend of COVID-19 in Italy and compared with Hunan, China, *Front. Med.*, **7**, (2020) 169.

- [17] A. J. Kucharski, T. W. Russell, C. Diamond, Y. Liu, J. Edmunds, S. Funk, R. M. Eggo, Early dynamics of transmission and control of COVID-19: A mathematical modelling study, *Lancet Infect Dis.*, **20**, (2020) 553-558.
- [18] Q. Li, B. Tang, N. L. Bragazzi, Y. Xiao, J. Wu, Modeling the impact of mass influenza vaccination and public health interventions on COVID-19 epidemics with limited detection capability, *Math. Biosci.*, **325**, (2020) 108378.
- [19] L. Danon, E. Brooks-Pollock, M. Bailey, M. Keeling, A spatial model of CoVID-19 transmission in England and Wales: Early spread and peak timing, *medRxiv 2020.02.12.20022566*, (2020).
- [20] D. Fanelli, F. Piazza, Analysis and forecast of COVID-19 spreading in China, Italy and France, *Chaos, Solitons and Fractals*, (2020) **134**, (2020) 109761.
- [21] K. Roosa, Y. Lee, R. Luo, A. Kirpich, R. Rothenberg, J. M. Hyman, P. Yan, G. Chowell, Real-time forecasts of the COVID-19 epidemic in China from February 5th to February 24th, 2020, *Infect. Dis. Model.*, **5**, (2020) 256–263.
- [22] L. Li, Z. Yang, Z. Dang, C. Meng, J. Huang, H. Meng, D. Wang, G. Chen, J. Zhang, H. Peng, Y. Shao, Propagation analysis and prediction of the COVID-19, *Infect. Dis. Model.*, **5**, (2020) 282–292.
- [23] L. Peng, W. Yang, D. Zhang, C. Zhuge, L. Hong, Epidemic analysis of COVID-19 in China by dynamical modeling, *medRxiv 2020.02.16.20023465*, (2020).
- [24] L. Mangoni, M. Pistilli, Epidemic analysis of COVID-19 in Italy by dynamical modelling, *SSRN Electronic Journal*, (2020). Available online at: <https://ssrn.com/abstract=3567770>
- [25] H. Ritchie, E. Mathieu, L. Rodés-Guirao, C. Appel, C. Giattino, E. Ortiz-Ospina, J. Hasell, B. Macdonald, D. Beltekian, M. Roser, *Coronavirus Pandemic (COVID-19)*, (2020). Available online at: <https://ourworldindata.org/coronavirus>
- [26] F. Checchi, L. Roberts, Interpreting and using mortality data in humanitarian emergencies, *Humanitarian Practice Network*, (2005).
- [27] D. M. Boka, H. Wainer, How can we estimate the death toll from COVID-19?, *Chance*, **33**, (2020) 67–72.
- [28] W. Farr 1885. *Vital statistics: A memorial volume of selections from the reports and writings of William Farr.* (edited for the Sanitary Institute of Great Britain by Noel A. Humphreys) London: The Sanitary Institute, (1885).
- [29] A. D. Langmuir, William Farr: Founder of modern concepts of surveillance, *International Journal of Epidemiology*, **5**, (1976) 13–18.
- [30] C. J. Murray, A. D. Lopez, B. Chin, D. Feehan, K. H. Hill 2006. Estimation of potential global pandemic influenza mortality on the basis of vital registry data from the 1918-20 pandemic: a quantitative analysis, *The Lancet*, **368**, (2006) 2211–2218.
- [31] C. Viboud, R. F. Grais, B. A. Lafont, M. A. Miller, L. Simonsen, Multinational influenza seasonal mortality study group. Multinational impact of the 1968 Hong Kong influenza pandemic: evidence for a smoldering pandemic, *J Infect Dis.*, **192**, (2005) 233–248.
- [32] J. Housworth, A. D. Langmuir, Excess mortality from epidemic influenza, 1957-1966. *Am. J. Epidemiol.*, **100**, (1974) 40–48.
- [33] H. Krelle, C. Barclay, C. Tallack, Understanding excess mortality: What is the fairest way to compare Covid-19 deaths internationally? (2020) Available at online: <https://www.health.org.uk/news-and-comment/charts-and-infographics/understanding-excess-mortality-the-fairest-way-to-make-international-comparisons>
- [34] WHO, ECDC. EuroMOMO. <http://www.euromomo.eu/>. Published 2020. Accessed 15.05.2020
- [35] Republic of Turkey, e-Government Portal, Available online at: <https://giris.turkiye.gov.tr/Giris/>
- [36] <https://ourworldindata.org/>