

CLASSICAL MDS ON METRIC MEASURE SPACES

SUNHYUK LIM AND FACUNDO MÉMOLI

ABSTRACT. We study a generalization of the classical Multidimensional Scaling procedure (cMDS) to the setting of general metric measure spaces. We identify certain crucial spectral properties of the generalized cMDS operator thus providing a natural and rigorous mathematical formulation of cMDS. Furthermore, we characterize the cMDS output of several continuous exemplar metric measures spaces such as high dimensional spheres and tori (both with their geodesic distance). In particular, the case of spheres (with geodesic distance) requires that we establish that its cMDS operator is trace class, a condition which is natural in the context when the cMDS operator has infinite rank. Finally, we establish the stability of the generalized cMDS method with respect to the Gromov-Wasserstein distance.

CONTENTS

1. Introduction	2
Contributions	3
Organization	4
Acknowledgements	4
2. Preliminaries	4
2.1. Metric measure spaces and the Gromov-Wasserstein distance	4
2.2. Functional analysis and spectral theory of operators	5
2.3. Spherical harmonics, Legendre Polynomials, and Funk-Hecke Theorem	10
3. cMDS: The case of finite metric spaces	13
3.1. Classical MDS of finite metric spaces	13
4. Generalized cMDS on metric measure spaces	20
4.1. The formulation of cMDS for metric measure spaces	24
4.2. A metric measure space with non-traceable cMDS operator	28
4.3. Distortion incurred by the cMDS operator	29
4.4. Optimality of the generalized cMDS	31
5. Thickness and spectrum of \mathfrak{K}_χ for euclidean data	34
6. Traceability of cMDS on exemplar metric measures	36
6.1. Generalized cMDS for product spaces	36
6.2. The case of \mathbb{S}^1	38
6.3. The case of $\mathbb{T}^N = (\mathbb{S}^1)^N$	41
6.4. The case of \mathbb{S}^{d-1}	41
6.5. Metric graphs and fractal spaces	55
7. Stability of Generalized cMDS	56
7.1. Consistency	60

MAX PLANCK INSTITUTE FOR MATHEMATICS IN THE SCIENCES.

THE OHIO STATE UNIVERSITY.

E-mail addresses: sulim@mis.mpg.de, facundo.memoli@gmail.com.

1. INTRODUCTION

The direct manipulation of high dimensional data is often difficult if not intractable. Nevertheless, many areas of science (e.g., neuroscience, climate science, network science, etc) have increasingly been confronted with large volumes of high dimensional data whose acquisition or generation has been facilitated by rapid technological advances.

Dimension reduction techniques have become central to data science in the past several decades due to their ability to obtain low-dimensional data representations which retain meaningful intrinsic properties of the original data.

When a dataset is modeled as a collection of points in some euclidean space, principal component analysis (PCA) provides a natural change of basis which can be used for ulterior processing such as dimension reduction [Pea01]. However, PCA cannot be directly applied to “non-euclidean” datasets such as those given in the form of dissimilarity matrices.

Classical Multidimensional Scaling (cMDS, from now on) is a *nonlinear* dimension reduction procedure which is applicable to data given as a dissimilarity or distance matrix. For a given dataset, modeled as finite metric space (X, d_X) , the goal of cMDS is to find an *optimal* embedding of X into Euclidean space. This optimality condition requires minimizing a certain notion of distortion (cf. Theorem 16).

From a mathematical perspective, the following result proved by Schoenberg in the early 1930s [Sch35] (cf. Theorem 15) can be regarded as a cornerstone of the theory supporting cMDS:

A finite metric space $\mathcal{X} = (X, d_X)$ can be embedded into a Euclidean space if and only if the matrix $B_{\mathcal{X}} := ((B_X)_{ij})$, where

$$(B_X)_{ij} := -\frac{1}{2} \left(d_X^2(x_i, x_j) - \frac{1}{n} \sum_{s=1}^n d_X^2(x_i, x_s) - \frac{1}{n} \sum_{r=1}^n d_X^2(x_r, x_j) + \frac{1}{n^2} \sum_{r,s=1}^n d_X^2(x_r, x_s) \right),$$

is positive semi-definite.

Schoenberg’s theorem not only provides a direct method for ascertaining the embedability of a given finite metric space into some euclidean space, but also it reveals that the obstacle to achieving this dwells in the presence of negative eigenvalues of the matrix $B_{\mathcal{X}}$. In this way it leads to one idea about how to produce an embedding of \mathcal{X} to Euclidean space by abandoning the negative eigenvalues of $B_{\mathcal{X}}$, which is exactly what cMDS does (cf. Algorithm 1).

Schoenberg’s theorem was independently rediscovered by scientists working in psychophysics [Ric38, YH38] whose original motivation came from the area of *sensory analysis*. More precisely, their goal was to estimate the dimensionality of the stimulus space, as well as the positions of the stimuli within this space [Mea92]. A few years later, Torgerson and Gower gave the formulation of cMDS that we know today and provided a supporting mathematical theory [Tor52, Tor58, Gow66].

The list of dimension reduction techniques has been constantly growing. Some notable examples are isomap [TSL00], locally linear embeddings [RS00], Laplacian eigenmaps [BN03], diffusion maps [CL06], Hessian eigenmaps [DG03], and more recently t-SNE [VdMH08] and

UMAP [MHSG18]. While their differences mainly come from how the distance (or, proximity) of data points is computed, these methods all attempt to preserve the local structure of data.

Contributions. The cMDS method has been mostly considered through its applicability to data analysis tasks, which means that its understanding has been mostly restricted to the case of finite datasets. Properties such as its stability and consistency (i.e. the behavior of the result of applying cMDS to a sample of a space as the cardinality of the sample increases) have not yet been clarified. In order to tackle these and other questions, in this paper we study a generalization of Schoenberg’s result to the *metric measure space* setting [Gro07, Mém11] which will permit (1) modeling both these ideal limiting objects and their finite approximations within a flexible framework and (2) reasoning about such questions in an effective manner.

A *metric measure space* is a triple (X, d_X, μ_X) where $\mathcal{X} = (X, d_X)$ is a compact metric space and μ_X is a full-support Borel probability measure on X . In this setting, we aim to quantify how much distortion will be incurred when trying to embed \mathcal{X} into Euclidean space of a given dimension via a certain generalized formulation of the cMDS procedure. This extension of cMDS to metric measure spaces was first considered in [Kas19, ABK20] where the authors were able to compute the spectrum of the cMDS operator for the case of the circle \mathbb{S}^1 with geodesic distance and normalized length measure. They also formulated a number of questions, such as extending the analysis to the case of spheres with arbitrary dimension (with geodesic distance and normalized volume measure) and establishing the stability of the cMDS procedure. The results in the present paper fully answer those and other questions about cMDS on continuous spaces.

When studying the spectral properties for the generalized cMDS operator in the setting of possible continuous metric measure spaces (such as spheres with their geodesic distance), one must pay careful attention to the possibility that the rank of the generalized cMDS operator can be infinite. This leads to considering on the collection of metric measure spaces with *traceable* cMDS operators.

Ascertaining whether the cMDS operator associated to a given metric measure space is traceable (i.e. in trace class) or not is far from trivial. In this paper we do this for some concrete examples¹, including spheres of arbitrary dimension with geodesic distances, tori and metric graphs (with their geodesic distances). In particular, we compute all eigenvalues and eigenfunctions of the cMDS operator for spheres \mathbb{S}^{d-1} with its geodesic distance $d_{\mathbb{S}^{d-1}}$ and normalized volume measure (i.e. regarded as a metric measure space), and establish that the cMDS operator is indeed trace class. Furthermore, we fully characterize the metric induced on \mathbb{S}^{d-1} by the cMDS operator: we prove that this metric precisely coincides with $\sqrt{\pi \cdot d_{\mathbb{S}^{d-1}}}$ (cf. Theorem 23). In order to highlight the difficulty in establishing that a given metric measure space induces a traceable cMDS operator, we describe the construction of one such space and formulate a number of questions suggested by our analysis.

Finally, we prove results which clarify the stability properties of the cMDS procedure with respect to the Gromov-Wasserstein distance.

This paper is part of the first author’s PhD dissertation [Lim21, Chapter 5].

Kroshnin, Stepanov and Trevisan have independently developed related results – we refer the reader to their preprint [KST22].

¹Note that all finite metric measure spaces induce traceable cMDS operators.

Organization. In §2 we give an account of Gromov-Wasserstein distances, spectral theory, and spherical harmonics which is then used in subsequent sections.

In §3, we provide some necessary definitions and results on the classical multidimensional scaling on finite metric spaces.

In §4, we consider a generalized version of cMDS for metric measure spaces and prove some basic properties. We define a notion of distortion involving the negative eigenvalues of the cMDS operator, and show that it somehow measures non-flatness of a metric measure space. Also, we prove that this generalized cMDS operator is the optimal choice in a certain rigorous sense.

In §5, we prove that a certain notion of thickness upper bounds the smallest positive eigenvalue of the cMDS operator for a subset of Euclidean space.

In §6, we provide a wide variety of nontrivial examples for the generalized cMDS. These include spheres \mathbb{S}^{d-1} , products such as the n -torus, and also metric graphs. In particular, we compute the spectrum of the generalized cMDS for the sphere \mathbb{S}^{d-1} endowed with its geodesic metric and prove the operator is in trace class.

In §7, we prove that, in a certain sense, the generalized cMDS process is stable with respect to Gromov-Wasserstein distances and briefly discuss an application to proving consistency of cMDS.

Acknowledgements. We are grateful to Henry Adams and Boris Mityagin for helpful conversations related to this work. We also thank Eugene Stepanov for pointing out to us a mistake in the proof of Proposition 4.12. This work was supported by NSF grants DMS 1547357, CCF-1526513, IIS-1422400, and CCF-1740761.

2. PRELIMINARIES

In this section we recall concepts that will be used in later sections to formulate and analyze the generalized cMDS procedure. Readers familiar with the Gromov-Wasserstein distance, spectral theory, and spherical harmonics may want to directly approach §3.

2.1. Metric measure spaces and the Gromov-Wasserstein distance. In this section we introduce machinery that permits modeling and comparing metric datasets.

In [Mém07, Mém11], the author defines the Gromov-Wasserstein distances combining ideas related to the Wasserstein distance and the Gromov-Hausdorff distance. Let (X, d_X) be a compact metric space and $\mathcal{P}(X)$ be the set of all Borel probability measures on X . Before proceeding, let us recall the definition of the coupling measure and support of a measure.

Definition 2.1. Suppose X is a topological space and μ_X is a measure on its Borel σ -algebra Σ_X . Then the support of μ_X is defined as the smallest closed set C satisfying $\mu_X(X \setminus C) = 0$ and denoted by $\text{supp}[\mu_X]$. In particular, if X is discrete,

$$\text{supp}[\mu_X] = \{x \in X : \mu_X(x) > 0\}.$$

Definition 2.2. Let (X, Σ_X, μ_X) and (Y, Σ_Y, μ_Y) be two measure spaces. Then, we call a measure μ on $X \times Y$ a coupling measure between μ_X and μ_Y if $\mu(A \times Y) = \mu_X(A)$ and $\mu(X \times B) = \mu_Y(B)$ whenever A and B are any measurable subsets of X and Y , respectively. Also, we denote with $\Gamma(\mu_X, \mu_Y)$ the set of all coupling measures.

Remark 2.3. Note that $\Gamma(\mu_X, \mu_Y)$ is nonempty since this set always contains the product measure $\mu_X \otimes \mu_Y$.

The Wasserstein distance on $\mathcal{P}(X) \times \mathcal{P}(X)$ is defined as follows.

Definition 2.4 (Wasserstein distances [Vil03]). Assume (X, d_X) is a compact metric space. For each $p \in [1, \infty]$ and $\alpha, \beta \in \mathcal{P}(X)$,

$$d_{W,p}^X(\alpha, \beta) = \inf_{\mu \in \Gamma(\alpha, \beta)} \left(\int_X d_X(x, x')^p d\mu(x, x') \right)^{\frac{1}{p}} = \inf_{\mu \in \Gamma(\alpha, \beta)} \|d_X\|_{L^p(\mu)}$$

for $p \in [1, \infty)$, and

$$d_{W,\infty}^X(\alpha, \beta) = \inf_{\mu \in \Gamma(\alpha, \beta)} \sup_{(x, x') \in \text{supp}[\mu]} d_X(x, x') = \inf_{\mu \in \Gamma(\alpha, \beta)} \|d_X\|_{L^\infty(\text{supp}[\mu])}.$$

Now, consider the collection of compact metric measure spaces, that is triples $\mathcal{X} = (X, d_X, \mu_X)$ where (X, d_X) is a compact metric space and μ_X is a fully supported Borel probability measure on X . This collection will be denoted by \mathcal{M}_w .

Definition 2.5 (Gromov-Wasserstein distances [Mém11]). For two compact metric measure spaces $\mathcal{X} = (X, d_X, \mu_X)$ and $\mathcal{Y} = (Y, d_Y, \mu_Y)$ and each $p \in [1, \infty]$,

$$\begin{aligned} d_{\text{GW},p}(\mathcal{X}, \mathcal{Y}) &= \frac{1}{2} \inf_{\mu \in \Gamma(\mu_X, \mu_Y)} \left(\int_{X \times Y \times X \times Y} |d_X(x, x') - d_Y(y, y')|^p d(\mu \otimes \mu)(x, y, x', y') \right)^{\frac{1}{p}} \\ &= \frac{1}{2} \inf_{\mu \in \Gamma(\mu_X, \mu_Y)} \|d_X - d_Y\|_{L^p(\mu \otimes \mu)} \end{aligned}$$

for $p \in [1, \infty)$, and

$$\begin{aligned} d_{\text{GW},\infty}(\mathcal{X}, \mathcal{Y}) &= \frac{1}{2} \inf_{\mu \in \Gamma(\mu_X, \mu_Y)} \sup_{(x, y), (x', y') \in \text{supp}[\mu]} |d_X(x, x') - d_Y(y, y')| \\ &= \frac{1}{2} \inf_{\mu \in \Gamma(\mu_X, \mu_Y)} \|d_X - d_Y\|_{L^\infty(\text{supp}[\mu] \times \text{supp}[\mu])}. \end{aligned}$$

The Gromov-Wasserstein distance $d_{\text{GW},p}$ defines a pseudometric on \mathcal{M}_w and it can be extended to a proper metric on the collection of isomorphism classes of compact metric measure spaces. We say two compact metric measure spaces $\mathcal{X} = (X, d_X, \mu_X)$ and $\mathcal{Y} = (Y, d_Y, \mu_Y)$ are isomorphic if there is a bijective map $f: X \rightarrow Y$ satisfying $d_X(x, x') = d_Y(f(x), f(x'))$ for all $x, x' \in X$ and $\mu_Y(B) = \mu_X(f^{-1}(B))$ for any borel set $B \subseteq Y$. i.e., $\mu_Y = f_*\mu_X$. We call this μ_Y the pushforward measure induced by f .

For each $p \in [1, \infty]$, let

$$(1) \quad \text{diam}_p(\mathcal{X}) := \|d_X\|_{L^p(\mu_X \otimes \mu_X)}.$$

Then one sees that

$$d_{\text{GW},p}(\mathcal{X}, \star) = \frac{1}{2} \text{diam}_p(\mathcal{X}).$$

2.2. Functional analysis and spectral theory of operators. We review the basics of spectral theory here. In particular, compact and self-adjoint operators on Hilbert spaces will be important for later sections. For the details, see [Ree12] and [RR06, Chs. 6 and 8].

Definition 2.6 (bounded operator). Let $(\mathcal{B}, \|\cdot\|_{\mathcal{B}})$ and $(\mathcal{C}, \|\cdot\|_{\mathcal{C}})$ be Banach spaces, and $\mathfrak{A} : \mathcal{B} \rightarrow \mathcal{C}$ be an operator from \mathcal{B} to \mathcal{C} . Then, \mathfrak{A} is said to be **bounded** if

$$\|\mathfrak{A}\| := \sup_{b \in \mathcal{B} \setminus \{0\}} \frac{\|\mathfrak{A}b\|_{\mathcal{C}}}{\|b\|_{\mathcal{B}}} < \infty.$$

From now on, $\mathcal{L}(\mathcal{B}, \mathcal{C})$ denotes the set of bounded operators from \mathcal{B} to \mathcal{C} . Also, for the simplicity, we use $\mathcal{L}(\mathcal{B})$ instead of $\mathcal{L}(\mathcal{B}, \mathcal{B})$.

Theorem 1 (adjoint [Ree12, Section VI.2]). *Let \mathcal{H} be a Hilbert space. Then, for any $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$, there is the unique bounded operator $\mathfrak{A}^* : \mathcal{H} \rightarrow \mathcal{H}$ satisfying:*

$$\langle u, \mathfrak{A}v \rangle = \langle \mathfrak{A}^*u, v \rangle$$

for any $u, v \in \mathcal{H}$.

From now on, the operator \mathfrak{A}^* will be called the **adjoint** of \mathfrak{A} .

Definition 2.7 (self-adjoint operator). Let \mathcal{H} be a Hilbert space. Then, $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$ is said to be **self-adjoint** if $\mathfrak{A} = \mathfrak{A}^*$. Equivalently, if \mathfrak{A} satisfies the following:

$$\langle \mathfrak{A}u, v \rangle = \langle u, \mathfrak{A}v \rangle$$

for any $u, v \in \mathcal{H}$.

Definition 2.8. Let \mathcal{H} be a Hilbert space and $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$. Then, a complex number λ is said to be in the **resolvent set** $\rho(\mathfrak{A})$ of \mathfrak{A} if $\lambda I - \mathfrak{A}$ is a bijection with a bounded inverse. If $\lambda \notin \rho(\mathfrak{A})$, then λ is said to be in the **spectrum** $\sigma(\mathfrak{A})$ of \mathfrak{A} .

Definition 2.9. Let \mathcal{H} be a Hilbert space and $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$.

- (1) An $u \neq 0$ which satisfies $\mathfrak{A}u = \lambda u$ for some $\lambda \in \mathbb{C}$ is called an **eigenvector** of \mathfrak{A} ; λ is called the corresponding eigenvalue. If λ is an eigenvalue, then $\lambda I - \mathfrak{A}$ is not injective so $\lambda \in \sigma(\mathfrak{A})$. The set of all eigenvalues is called the **point spectrum** of \mathfrak{A} .
- (2) If λ is not an eigenvalue and if the range of $\lambda I - \mathfrak{A}$ is not dense, then λ is said to be in the **residual spectrum**.

Theorem 2 ([Ree12, Theorem VI.6, Theorem VI.8]). *Let \mathfrak{A} be a bounded and self-adjoint operator on a Hilbert space \mathcal{H} . Then,*

- (1) \mathfrak{A} has no residual spectrum.
- (2) The spectrum $\sigma(\mathfrak{A})$ of \mathfrak{A} is a subset of \mathbb{R} . In particular, $\sup_{\lambda \in \sigma(\mathfrak{A})} |\lambda| = \|\mathfrak{A}\|$.
- (3) Eigenvectors corresponding to distinct eigenvalues of \mathfrak{A} are orthogonal.

Definition 2.10 (positive semi-definite operator). Let \mathcal{H} be a Hilbert space and $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$. \mathfrak{A} is called **positive semi-definite** if $\langle u, \mathfrak{A}u \rangle \geq 0$ for any $u \in \mathcal{H}$. We write $\mathfrak{A} \succeq 0$ if \mathfrak{A} is positive semi-definite.

Remark 2.11. *If \mathfrak{A} is a bounded operator on a Hilbert space \mathcal{H} . Then, $\mathfrak{A}^*\mathfrak{A}$ is always positive semi-definite.*

Theorem 3 ([Ree12, Theorem VI.9]). *Let \mathfrak{A} be a positive semi-definite and bounded operator on a Hilbert space \mathcal{H} . Then, there is the unique positive semi-definite and bounded operator \mathfrak{B} on \mathcal{H} satisfying $\mathfrak{B}^2 = \mathfrak{A}$. Furthermore, \mathfrak{B} commutes with every bounded operator which commutes with \mathfrak{A} .*

By Theorem 3 and Remark 2.11, we define $|\mathfrak{A}|$ in the following way.

Definition 2.12. Let \mathfrak{A} be a bounded operator on a Hilbert space \mathcal{H} . Then $|\mathfrak{A}| := \sqrt{\mathfrak{A}^*\mathfrak{A}}$, $\mathfrak{A}^+ := \frac{1}{2}(|\mathfrak{A}| + \mathfrak{A})$, and $\mathfrak{A}^- := \frac{1}{2}(|\mathfrak{A}| - \mathfrak{A})$. Note that $\mathfrak{A} = \mathfrak{A}^+ - \mathfrak{A}^-$.

Definition 2.13 (compact operator). Let $(\mathcal{B}, \|\cdot\|_{\mathcal{B}})$ and $(\mathcal{C}, \|\cdot\|_{\mathcal{C}})$ be Banach spaces. An operator $\mathfrak{A} \in \mathcal{L}(\mathcal{B}, \mathcal{C})$ is called **compact** if \mathfrak{A} takes bounded sets in \mathcal{B} into precompact sets in \mathcal{C} .

Theorem 4 (Riesz-Schauder theorem, [Ree12, Theorem VI.15]). *Let \mathcal{H} be a separable Hilbert space and $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$ be a compact operator. Then, $\sigma(\mathfrak{A})$ is a discrete set having no limit points except perhaps $\lambda = 0$. Further, any nonzero $\lambda \in \sigma(\mathfrak{A})$ is an eigenvalue of finite multiplicity (i.e. the corresponding space of eigenvectors is finite dimensional).*

Theorem 5 (Hilbert-Schmidt Theorem, [Ree12, Theorem VI.16] and [RR06, Theorem 8.94]). *Let \mathcal{H} be a separable Hilbert space and $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$ be compact and self-adjoint. Then, there is a complete orthonormal basis $\{\phi_i\}_{i=1}^N$ for \mathcal{H} where $N = \dim(\mathcal{H})$ so that $\mathfrak{A}\phi_i = \lambda_i\phi_i$, $|\lambda_i|$ is monotone nonincreasing, and if $N = \infty$, $\lim_{i \rightarrow \infty} \lambda_i = 0$. Furthermore, \mathfrak{A} can be represented by*

$$\mathfrak{A}(\cdot) = \sum_{i=1}^N \lambda_i \langle \phi_i, \cdot \rangle \phi_i.$$

Definition 2.14 (singular values). Let \mathcal{H} be a separable Hilbert space and $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$ be a compact operator. Then, the non-negative square roots of eigenvalues of the self-adjoint operator $\mathfrak{A}^*\mathfrak{A}$ are called **singular values** of \mathfrak{A} .

For a given metric measure space $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$, the following theorem gives conditions under which the Hilbert space $L^2(\mu_X)$ is separable.

Proposition 2.15 ([Coh13, Proposition 3.4.5]). *Let (X, Σ_X, μ_X) be a measure space, and let p satisfy $1 \leq p < \infty$. If μ_X is σ -finite and Σ_X is countably generated, then $L^p(X, \Sigma_X, \mu_X)$ is separable.*

In particular, if (X, d_X) is complete and separable metric space (every compact metric space satisfies these conditions) and μ_X is Borel probability measure, then $L^2(\mu_X)$ is separable.

Next, we introduce particularly important kinds of compact operators: Trace class and Hilbert-Schmidt operators.

Theorem 6 ([Ree12, Theorem VI.18]). *Let \mathcal{H} be a separable Hilbert space, and $\{\phi_i\}_{i=1}^N$ be an orthonormal basis of \mathcal{H} . Then for any positive semi-definite operator $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$, the value*

$$\sum_{i=1}^N \langle \phi_i, \mathfrak{A}\phi_i \rangle$$

does not depend on the choice of basis.

Definition 2.16 (trace of a positive semi-definite operator). Let \mathcal{H} be a separable Hilbert space and $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$ be a positive semi-definite operator. Then,

$$\text{tr}(\mathfrak{A}) := \sum_{n=1}^N \langle \phi_n, \mathfrak{A}\phi_n \rangle$$

where $\{\phi_n\}_{n=1}^N$ is an orthonormal basis of \mathcal{H} .

Definition 2.17 (trace class). Let \mathcal{H} be a separable Hilbert space. Then, for any $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$, we define the **trace norm** of \mathfrak{A} in the following way:

$$\|\mathfrak{A}\|_1 := \text{tr}(|\mathfrak{A}|).$$

Moreover, \mathfrak{A} is called **trace class** if $\|\mathfrak{A}\|_1 < \infty$. The set of all traceable operators in $\mathcal{L}(\mathcal{H})$ is denoted by $\mathcal{L}_1(\mathcal{H})$.

Theorem 7 ([Ree12, Theorem VI.19, Theorem VI.20]). *Let \mathcal{H} be a separable Hilbert space. Then,*

- (1) $\mathcal{L}_1(\mathcal{H})$ is a Banach space with the trace norm $\|\cdot\|_1$.
- (2) If $\mathfrak{A} \in \mathcal{L}_1(\mathcal{H})$ and $\mathfrak{B} \in \mathcal{L}(\mathcal{H})$, then $\mathfrak{A}\mathfrak{B} \in \mathcal{L}_1(\mathcal{H})$ and $\mathfrak{B}\mathfrak{A} \in \mathcal{L}_1(\mathcal{H})$.
- (3) If $\mathfrak{A} \in \mathcal{L}_1(\mathcal{H})$, then $\mathfrak{A}^* \in \mathcal{L}_1(\mathcal{H})$.

Theorem 8 ([Ree12, Theorem VI.21]). *If \mathfrak{A} is trace class, then \mathfrak{A} is compact. Moreover, a compact operator \mathfrak{A} is trace class if and only if $\sum_{i=1}^N \lambda_i < \infty$ where $\{\lambda_i\}_{i=1}^N$ are the singular values of \mathfrak{A} .*

Moreover, the notion of trace can be extended to trace class operators via the following theorem.

Theorem 9 ([Ree12, Theorem VI.24]). *Let \mathcal{H} be a separable Hilbert space, and $\{\phi_i\}_{i=1}^N$ be an orthonormal basis of \mathcal{H} . Then for any $\mathfrak{A} \in \mathcal{L}_1(\mathcal{H})$, the value*

$$\sum_{i=1}^N \langle \phi_i, \mathfrak{A}\phi_i \rangle$$

converges absolutely and the limit does not depend on the choice of basis.

Definition 2.18 (trace of a trace class operator). Let \mathcal{H} be a separable Hilbert space and $\mathfrak{A} \in \mathcal{L}_1(\mathcal{H})$. Then,

$$\text{Tr}(\mathfrak{A}) := \sum_{n=1}^N \langle \phi_n, \mathfrak{A}\phi_n \rangle$$

where $\{\phi_n\}_{n=1}^N$ is an orthonormal basis of \mathcal{H} . Observe that $\text{Tr}(\mathfrak{A}) = \text{tr}(\mathfrak{A})$ if \mathfrak{A} is positive semi-definite.

Theorem 10 ([Ree12, Theorem VI.25]). (1) $\text{Tr}(c\mathfrak{A} + \mathfrak{B}) = c\text{Tr}(\mathfrak{A}) + \text{Tr}(\mathfrak{B})$ for any $\mathfrak{A}, \mathfrak{B} \in \mathcal{L}_1(\mathcal{H})$ and $c \in \mathbb{C}$.
 (2) $\text{Tr}(\mathfrak{A}) = \overline{\text{Tr}(\mathfrak{A}^*)}$ for any $\mathfrak{A} \in \mathcal{L}_1(\mathcal{H})$.
 (3) $\text{Tr}(\mathfrak{A}\mathfrak{B}) = \text{Tr}(\mathfrak{B}\mathfrak{A})$ if $\mathfrak{A} \in \mathcal{L}_1(\mathcal{H})$ and $\mathfrak{B} \in \mathcal{L}(\mathcal{H})$.

Next, we introduce the notion of Hilbert-Schmidt operators.

Definition 2.19 (Hilbert-Schmidt operator). Let \mathcal{H} be a separable Hilbert space. Then, for any $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$, we define the **Hilbert-Schmidt norm** of \mathfrak{A} in the following way:

$$\|\mathfrak{A}\|_2 := (\text{Tr}(\mathfrak{A}^*\mathfrak{A}))^{\frac{1}{2}}.$$

Moreover, \mathfrak{A} is called **Hilbert-Schmidt operator** if $\|\mathfrak{A}\|_2 < \infty$. The set of all Hilbert-Schmidt operators in $\mathcal{L}(\mathcal{H})$ is denoted by $\mathcal{L}_2(\mathcal{H})$.

Let us collect some useful properties of trace, trace class and Hilbert-Schmidt operators.

Lemma 2.20 ([Ree12, Theorem VI.19, Theorem VI.21, Theorem VI.22, Theorem VI.25]). *Let \mathcal{H} be a separable Hilbert space and $\mathfrak{A}, \mathfrak{B}, \mathfrak{C} \in \mathcal{L}(\mathcal{H})$. Then,*

- (1) $\|\mathfrak{A}\| \leq \|\mathfrak{A}\|_2 \leq \|\mathfrak{A}\|_1$.
- (2) *If \mathfrak{A} is Hilbert-Schmidt operator, then \mathfrak{A} is compact. Moreover, a compact operator \mathfrak{A} is Hilbert-Schmidt if and only if $\sum_{i=1}^N \lambda_i^2 < \infty$ where $\{\lambda_i\}_{i=1}^N$ are the singular values of \mathfrak{A} .*
- (3) \mathfrak{A} *is trace class if and only if $\mathfrak{A} = \mathfrak{B}\mathfrak{C}$ for some Hilbert-Schmidt operators $\mathfrak{B}, \mathfrak{C}$.*
- (4) $\text{Tr}(\mathfrak{A}\mathfrak{B}) = \text{Tr}(\mathfrak{B}\mathfrak{A})$ *if both of \mathfrak{A} and \mathfrak{B} are Hilbert-Schmidt operators.*

Proof. To prove item (4), which is the only one having no proof in [Ree12], we use item (3) of this lemma and follow the proof of [Ree12, Theorem VI.25]. \square

The following theorem connects Hilbert-Schmidt operator and integral kernel.

Theorem 11 ([Ree12, Theorem VI.23]). *Let (X, Σ_X, μ_X) be a measure space with the separable Hilbert space $L^2(\mu_X)$. Then, $\mathfrak{K} \in \mathcal{L}(L^2(\mu_X))$ is Hilbert-Schmidt if and only if there is a function*

$$K \in L^2(\mu_X \otimes \mu_X)$$

with

$$\mathfrak{K}\phi(\cdot) = \int_X K(\cdot, x') \phi(x') d\mu_X(x').$$

Moreover,

$$\|\mathfrak{K}\|_2^2 = \|K\|_{L^2(\mu_X \otimes \mu_X)}^2.$$

Furthermore, the integral kernel associated to a given Hilbert-Schmidt operator on $L^2(\mu_X)$ is uniquely determined (save for a set with zero measure) as one can see from the following lemma and corollary.

Lemma 2.21 ([Wof18]). *Suppose a measure space (X, Σ_X, μ_X) and the separable Hilbert space $L^2(\mu_X)$ are given. Let $K \in L^2(\mu_X \otimes \mu_X)$ and let $\mathfrak{K} \in \mathcal{L}(L^2(\mu_X))$ be the Hilbert-Schmidt operator associated to K . If $\mathfrak{K} = 0$, then $K = 0$ ($\mu_X \otimes \mu_X$)-almost everywhere.*

Proof. For each $x \in X$, let $K_x := K(x, \cdot)$. Then, since $K \in L^2(\mu_X \otimes \mu_X)$, we have $K_x \in L^2(\mu_X)$ for μ_X -almost every x . Observe that $\mathfrak{K}\psi(x) = \langle \psi, K_x \rangle$ for any $\psi \in L^2(\mu_X)$. Now, choose a complete orthonormal basis $\{\phi_i\}_{i=1}^N$ of $L^2(\mu_X)$. For each i , $\mathfrak{K}\phi_i = 0$ implying that K_x is orthogonal to ϕ_i μ_X -almost every x . Since a countable union of null sets is null, K_x is orthogonal to ϕ_i for all i and for μ_X -almost every x . Hence, $K_x = 0$ almost everywhere for almost every $x \in X$. Therefore, $K = 0$ ($\mu_X \otimes \mu_X$)-almost everywhere. \square

Corollary 2.22. *Suppose a measure space (X, Σ_X, μ_X) and the separable Hilbert space $L^2(\mu_X)$ are given. Let $K_1, K_2 \in L^2(\mu_X \otimes \mu_X)$ and $\mathfrak{K}_1, \mathfrak{K}_2 \in \mathcal{L}(L^2(\mu_X))$ be the Hilbert-Schmidt operators associated to K_1 and K_2 respectively. If $\mathfrak{K}_1 = \mathfrak{K}_2$, then $K_1 = K_2$ ($\mu_X \otimes \mu_X$)-almost everywhere.*

Also, when (X, Σ_X, μ_X) is a measure space with the separable Hilbert space $L^2(\mu_X)$, the set of kernels $K \in L^2(\mu_X \otimes \mu_X)$ whose associated integral operator $\mathfrak{K} \in \mathcal{L}(L^2(\mu_X))$ is positive semi-definite is denoted by $L^2_{\geq 0}(\mu_X \otimes \mu_X)$.

Definition 2.23 (symmetric kernel). (X, Σ_X, μ_X) is a measure space. A kernel $K : X \times X \rightarrow \mathbb{R}$ is said to be **symmetric** if $K(x, x') = K(x', x)$ $\mu_X \otimes \mu_X$ -a.e.

Theorem 12 ([Bri91, Corollary 3.2]). *Let μ_X be a σ -finite Borel measure on a second countable space X , and let K be the integral kernel of a trace class operator \mathfrak{K} on $L^2(\mu_X)$. If K is continuous at (x, x) for μ_X -almost every x then*

$$\mathrm{Tr}(\mathfrak{K}) = \int_X K(x, x) d\mu_X(x).$$

2.3. Spherical harmonics, Legendre Polynomials, and Funk-Hecke Theorem. For any $d \geq 2$, we consider the standard unit $(d - 1)$ -dimensional sphere

$$\mathbb{S}^{d-1} := \{(x_1, \dots, x_d) \in \mathbb{R}^d : \sum_{i=1}^d x_i^2 = 1\}$$

with the canonical Riemannian metric. Let $d_{\mathbb{S}^{d-1}}$ be the geodesic (round) distance of \mathbb{S}^{d-1} and $\mathrm{vol}_{\mathbb{S}^{d-1}}$ be the volume measure of \mathbb{S}^{d-1} induced by the canonical Riemannian metric. Since $\mathrm{vol}_{\mathbb{S}^{d-1}}(\mathbb{S}^{d-1})$, the volume of \mathbb{S}^{d-1} , will be used multiple times later, let us introduce quick notation $|\mathbb{S}^{d-1}| := \mathrm{vol}_{\mathbb{S}^{d-1}}(\mathbb{S}^{d-1})$. The following equality is well-known ([AH12, (1.19)]).

$$(2) \quad |\mathbb{S}^{d-1}| = \frac{2\pi^{\frac{d}{2}}}{\Gamma(\frac{d}{2})}.$$

Note that $(\mathbb{S}^{d-1}, d_{\mathbb{S}^{d-1}}, \mathrm{nvol}_{\mathbb{S}^{d-1}} := \frac{1}{|\mathbb{S}^{d-1}|} \mathrm{vol}_{\mathbb{S}^{d-1}})$ is a metric measure space in \mathcal{M}_w .

Now we review basic results on spherical harmonics. Our main reference is [AH12].

For any $d \geq 2$ and $n \geq 0$, let \mathbb{H}_n^d be the space of all homogeneous polynomials of degree n in d dimensions.

Example 2.24. *Here are some examples of \mathbb{H}_n^d .*

- $\mathbb{H}_2^2 = \{a_1x_1^2 + a_2x_1x_2 + a_3x_2^2 : a_1, a_2, a_3 \in \mathbb{R}\}$.
- $\mathbb{H}_3^2 = \{a_1x_1^3 + a_2x_1x_2^2 + a_3x_1^2x_2 + a_4x_2^3 : a_1, a_2, a_3, a_4 \in \mathbb{R}\}$.
- $\mathbb{H}_2^3 = \{a_1x_1^2 + a_2x_2^2 + a_3x_3^2 + a_4x_1x_2 + a_5x_2x_3 + a_6x_3x_1 : a_1, \dots, a_6 \in \mathbb{R}\}$.

Recall that a function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is called **harmonic** if $\Delta f = 0$. The space of homogeneous harmonics of degree n in d dimension is denoted by $\mathbb{Y}_n(\mathbb{R}^d)$. More precisely,

$$\mathbb{Y}_n(\mathbb{R}^d) := \{f \in \mathbb{H}_n^d : \Delta f = 0\}.$$

Let $N_{n,d} := \dim(\mathbb{Y}_n(\mathbb{R}^d))$. The following equality is well-known ([AH12, (2.10)]).

$$(3) \quad N_{n,d} = \frac{(2n + d - 2)(n + d - 3)!}{n!(d - 2)!}.$$

Moreover, $N_{n,d} = O(n^{d-2})$ for n sufficiently large ([AH12, (2.12)]).

Finally, we are ready to introduce spherical harmonics.

Definition 2.25 (spherical harmonics). $\mathbb{Y}_n^d := \mathbb{Y}_n(\mathbb{R}^d)|_{\mathbb{S}^{d-1}}$ is called the spherical harmonic space of size n in d dimensions. Any function in \mathbb{Y}_n^d is called a spherical harmonic of order n in d dimensions.

Remark 2.26. *It is known that $\dim(\mathbb{Y}_n^d) = \dim(\mathbb{Y}_n(\mathbb{R}^d)) = N_{n,d}$.*

Now, let us introduce Legendre polynomials, which will be useful later. A more detailed explanation on Legendre polynomials can be found in [AH12, Section 2].

Definition 2.27 (Legendre polynomial). The map

$$P_{n,d} : [-1, 1] \longrightarrow \mathbb{R}$$

$$t \longmapsto n! \Gamma\left(\frac{d-1}{2}\right) \sum_{k=0}^{\lfloor n/2 \rfloor} (-1)^k \frac{(1-t^2)^k t^{n-2k}}{4^k k! (n-2k)! \Gamma\left(k + \frac{d-1}{2}\right)}$$

is said to be the *Legendre polynomial of degree n in d dimensions*.

The following theorem connects spherical harmonics and Legendre polynomials.

Theorem 13 (addition Theorem [AH12, Theorem 2.9]). *Let $\{Y_{n,j}^d : 1 \leq j \leq N_{n,d}\}$ be an orthonormal basis of \mathbb{Y}_n^d , i.e.,*

$$\int_{\mathbb{S}^{d-1}} Y_{n,j}^d(u) Y_{n,k}^d(u) d\text{vol}_{\mathbb{S}^{d-1}}(u) = \delta_{jk}.$$

Then

$$\sum_{j=1}^{N_{n,d}} Y_{n,j}^d(u) Y_{n,j}^d(v) = \frac{N_{n,d}}{|\mathbb{S}^{d-1}|} P_{n,d}(\langle u, v \rangle)$$

for any $u, v \in \mathbb{S}^{d-1}$.

Lemma 2.28 ([AH12, Corollary 2.15]). *For $m \neq n$, $\mathbb{Y}_m^d \perp \mathbb{Y}_n^d$ in $L^2(\text{nvoll}_{\mathbb{S}^{d-1}})$.*

Corollary 2.29. *For any $n > 0$ and $Y_n^d \in \mathbb{Y}_n^d$, $\int_{\mathbb{S}^{d-1}} Y_n^d(u) d\text{nvoll}_{\mathbb{S}^{d-1}}(u) = 0$.*

Proof. Apply Lemma 2.28 and the fact $Y_0^d \equiv 1 \in \mathbb{Y}_0^d$. \square

Let us collect some useful properties of Legendre polynomials which will be used in the sequel.

Lemma 2.30. *For any $d \geq 3$, $m, n \geq 0$, and $t \in [-1, 1]$,*

- (1) $|P_{n,d}(t)| \leq 1 = P_{n,d}(1)$ (see [AH12, (2.39)]).
- (2) $\int_{\mathbb{S}^{d-1}} |P_{n,d}(\langle u, v \rangle)|^2 d\text{vol}_{\mathbb{S}^{d-1}}(v) = \frac{|\mathbb{S}^{d-1}|}{N_{n,d}}$ for any $u \in \mathbb{S}^{d-1}$ (see [AH12, (2.40)]).
- (3) (Rodrigues representation formula) $P_{n,d}(t) = (-1)^n R_{n,d} (1-t^2)^{\frac{3-d}{2}} \left(\frac{d}{dt}\right)^n (1-t^2)^{n+\frac{d-3}{2}}$
where $R_{n,d} = \frac{\Gamma(\frac{d-1}{2})}{2^n \Gamma(n+\frac{d-1}{2})}$ (see [AH12, Theorem 2.23]).
- (4) $P_{n,d}(-t) = (-1)^n P_{n,d}(t)$ (see [AH12, (2.73)]).
- (5) If $f \in C^n([-1, 1])$, then

$$\int_{-1}^1 f(t) P_{n,d}(t) (1-t^2)^{\frac{d-3}{2}} dt = R_{n,d} \int_{-1}^1 f^{(n)}(t) (1-t^2)^{n+\frac{d-3}{2}} dt$$

$$R_{n,d} = \frac{\Gamma(\frac{d-1}{2})}{2^n \Gamma(n+\frac{d-1}{2})} \text{ (see [AH12, Proposition 2.26])}.$$

- (6) $\int_{-1}^1 P_{m,d}(t) P_{n,d}(t) (1-t^2)^{\frac{d-3}{2}} dt = \frac{|\mathbb{S}^{d-1}|}{N_{n,d} |\mathbb{S}^{d-2}|} \delta_{mn}$ (see [AH12, (2.79)]).
- (7) (Poisson identity) For any $r \in (-1, 1)$ and $t \in [-1, 1]$,

$$\sum_{n=0}^{\infty} N_{n,d} P_{n,d}(t) r^n = \frac{1-r^2}{(1+r^2-2rt)^{\frac{d}{2}}}.$$

Moreover, for each fixed $r \in (-1, 1)$, the convergence is uniform as a function of t (see [AH12, Proposition 2.28]).

(8) (*Green-Beltrami identity*) For any $f \in C^2(\mathbb{S}^{d-1})$ and any $g \in C^1(\mathbb{S}^{d-1})$, we have

$$\int_{\mathbb{S}^{d-1}} g \Delta f \, d\text{vol}_{\mathbb{S}^{d-1}} = - \int_{\mathbb{S}^{d-1}} \langle \nabla g, \nabla f \rangle \, d\text{vol}_{\mathbb{S}^{d-1}}$$

(see [AH12, Proposition 3.3]).

(9) For any $f \in C^1(\mathbb{S}^{d-1})$ and $u \in \mathbb{S}^{d-1}$, we have

$$\int_{\mathbb{S}^{d-1}} \nabla f(v) \cdot \nabla_v P_{n,d}(\langle u, v \rangle) \, d\text{vol}_{\mathbb{S}^{d-1}}(v) = n(n+d-2) \int_{\mathbb{S}^{d-1}} f(v) P_{n,d}(\langle u, v \rangle) \, d\text{vol}_{\mathbb{S}^{d-1}}(v)$$

(see [AH12, (3.21)]).

(10) For any $u \in \mathbb{S}^{d-1}$, we have

$$\int_{\mathbb{S}^{d-1}} \|\nabla_v P_{n,d}(\langle u, v \rangle)\|^2 \, d\text{vol}_{\mathbb{S}^{d-1}}(v) = n(n+d-2) \frac{|\mathbb{S}^{d-1}|}{N_{n,d}}$$

(see [AH12, Proposition 3.6]).

Proof. For item (7) (Poisson identity), the uniform convergence holds because of the following argument: Observe that, for any positive integer $k > 0$,

$$\left| \sum_{n=k}^{\infty} N_{n,d} P_{n,d}(t) r^n \right| \leq \sum_{n=k}^{\infty} N_{n,d} |P_{n,d}(t)| |r|^n = \sum_{n=k}^{\infty} N_{n,d} |r|^n \quad (\because \text{item (1)}).$$

But, since

$$\sum_{n=0}^{\infty} N_{n,d} |r|^n = \sum_{n=0}^{\infty} N_{n,d} P_{n,d}(1) |r|^n = \frac{1+|r|}{(1-|r|)^d} < \infty,$$

$\sum_{n=k}^{\infty} N_{n,d} |r|^n$ becomes arbitrarily small for large enough k .

For the proof of other items, see [AH12]. \square

Let us introduce a few more notation. For any real-valued measurable map f on $[-1, 1]$,

$$\|f\|_{L^1_{\frac{d-3}{2}}([-1,1])} := \int_{-1}^1 |f(t)| (1-t^2)^{\frac{d-3}{2}} \, dt,$$

and

$$L^1_{\frac{d-3}{2}}([-1, 1]) := \left\{ f \text{ measurable on } [-1, 1] : \|f\|_{L^1_{\frac{d-3}{2}}([-1,1])} < \infty \right\}.$$

Now we are ready to state our main tool for studying the output of the cMDS method when applied to \mathbb{S}^{d-1} .

Theorem 14 (Funk-Hecke theorem [AH12, Theorem 2.22]). *Let $f \in L^1_{\frac{d-3}{2}}([-1, 1])$, $u \in \mathbb{S}^{d-1}$ and $Y_n^d \in \mathbb{Y}_n^d$. Then the Funk-Hecke formula*

$$\int_{\mathbb{S}^{d-1}} f(\langle u, v \rangle) Y_n^d(v) \, d\text{vol}_{\mathbb{S}^{d-1}}(v) = \lambda_n Y_n^d(u)$$

holds with

$$\lambda_n = \int_{\mathbb{S}^{d-1}} P_{n,d}(\langle u, v \rangle) f(\langle u, v \rangle) \, d\text{vol}_{\mathbb{S}^{d-1}}(v) = |\mathbb{S}^{d-2}| \int_{-1}^1 P_{n,d}(t) f(t) (1-t^2)^{\frac{d-3}{2}} \, dt.$$

3. cMDS: THE CASE OF FINITE METRIC SPACES

In this paper, \mathbb{R}^k stands for the canonical k -dimensional Euclidean space, and the set of square-summable sequences $\{(x_1, x_2, \dots) : x_i \in \mathbb{R} \text{ for each } i \in \mathbb{Z}_{>0} \text{ and } \sum_{i=1}^{\infty} x_i^2 < \infty\}$ with the Euclidean inner product structure will be denoted by ℓ^2 . As an abuse of notation, $\|\cdot\|$ will also denote the metric induced by the Euclidean norm. Finally, m_k will denote the Lebesgue measure on \mathbb{R}^k .

3.1. Classical MDS of finite metric spaces. The main reference for this subsection is [MKB79, Chapter 14]. For each $n \in \mathbb{Z}_{>0}$, let \mathcal{M}_n be the collection of all finite metric spaces $\mathcal{X} = (X, d_X)$ with $\text{card}(X) = n$. Schoenberg's Theorem (Theorem 15) provides a criterion for checking whether a finite metric space $\mathcal{X} = (X, d_X) \in \mathcal{M}_n$ admits an isometric embedding into Euclidean space \mathbb{R}^k . We need some preparation before stating this theorem.

For a finite metric space $\mathcal{X} = (X = \{x_1, \dots, x_n\}, d_X) \in \mathcal{M}_n$, consider an n by n matrix $A_{\mathcal{X}}$ where $(A_{\mathcal{X}})_{ij} := d_X^2(x_i, x_j)$ for each $i, j = 1, \dots, n$. Next, construct another n by n matrix

$$B_{\mathcal{X}} := -\frac{1}{2}H_n A_{\mathcal{X}} H_n$$

where $H_n := I_n - \frac{1}{n}e_n e_n^T$, I_n is the n by n identity matrix, and e_n is the n by 1 column vector with every entry is 1.

Let us collect some of simple properties of $A_{\mathcal{X}}$ and $B_{\mathcal{X}}$.

Lemma 3.1. $\mathcal{X} = (X, d_X) \in \mathcal{M}_n$ is given. $A_{\mathcal{X}}$ and $B_{\mathcal{X}}$ are as before. Then,

- (1) $A_{\mathcal{X}}$ and $B_{\mathcal{X}}$ are symmetric matrices.
- (2) $(B_{\mathcal{X}})_{ij} = -\frac{1}{2} \left((A_{\mathcal{X}})_{ij} - \frac{1}{n} \sum_{s=1}^n (A_{\mathcal{X}})_{is} - \frac{1}{n} \sum_{r=1}^n (A_{\mathcal{X}})_{rj} + \frac{1}{n^2} \sum_{r,s=1}^n (A_{\mathcal{X}})_{rs} \right)$ for each $i, j = 1, \dots, n$.
- (3) $(A_{\mathcal{X}})_{ij} = d_X^2(x_i, x_j) = (B_{\mathcal{X}})_{ii} + (B_{\mathcal{X}})_{jj} - 2(B_{\mathcal{X}})_{ij}$ for each $i, j = 1, \dots, n$.
- (4) e_n is an eigenvector of $B_{\mathcal{X}}$ with zero eigenvalue. In particular, $\dim(\ker B_{\mathcal{X}}) \geq 1$.
- (5) If $v \in \mathbb{R}^n$ is an eigenvector for a nonzero eigenvalue of $B_{\mathcal{X}}$, then $\sum_{i=1}^n v(i) = 0$.

Proof. (1) $A_{\mathcal{X}}$ is obviously symmetric because of the properties of metric. Also, $B_{\mathcal{X}}$ is symmetric since both of $A_{\mathcal{X}}$ and H_n are symmetric.

(2) From the definition,

$$B_{\mathcal{X}} = -\frac{1}{2} \left(A_{\mathcal{X}} - \frac{1}{n} e_n e_n^T A_{\mathcal{X}} - \frac{1}{n} A_{\mathcal{X}} e_n e_n^T + \frac{1}{n^2} e_n e_n^T A_{\mathcal{X}} e_n e_n^T \right).$$

After direct computation, one obtains (2).

(3) Use item (2).

(4) For each $i = 1, \dots, n$,

$$\begin{aligned} (B_{\mathcal{X}} e_n)(i) &= \sum_{j=1}^n (B_{\mathcal{X}})_{ij} \\ &= -\frac{1}{2} \left(\sum_{j=1}^n (A_{\mathcal{X}})_{ij} - \sum_{s=1}^n (A_{\mathcal{X}})_{is} - \frac{1}{n} \sum_{r,j=1}^n (A_{\mathcal{X}})_{rj} + \frac{1}{n} \sum_{r,s=1}^n (A_{\mathcal{X}})_{rs} \right) \\ &= 0 \end{aligned}$$

for each $i = 1, \dots, n$ where the second equality comes from (2). Hence, $B_{\mathcal{X}} e_n = 0$ as we required.

(5) Let v be an eigenvector of $B_{\mathcal{X}}$ with nonzero eigenvalue λ . Then,

$$\begin{aligned} \lambda \sum_{i=1}^n v(i) &= \sum_{i=1}^n (B_{\mathcal{X}}v)(i) \\ &= \sum_{i=1}^n \sum_{j=1}^n (B_{\mathcal{X}})_{ij} v(j) \\ &= \sum_{j=1}^n v(j) \sum_{i=1}^n (B_{\mathcal{X}})_{ij} = 0 \end{aligned}$$

since $\sum_{i=1}^n (B_{\mathcal{X}})_{ij} = 0$ for each $i = 1, \dots, n$ by item (1) and item (4). Hence, $\sum_{i=1}^n v(i) = 0$. □

Remark 3.2. Observe that, by item (2) of Lemma 3.1,

$$\text{Tr}(B_{\mathcal{X}}) = \frac{1}{2n} \sum_{r,s=1}^n d_X^2(x_r, x_s).$$

Hence, for any non-singleton metric space $\mathcal{X} = (X, d_X)$, $B_{\mathcal{X}}$ must have at least one strictly positive eigenvalue.

Theorem 15 (Schoenberg's Theorem [Sch35, YH38, MKB79]). *Let $(X = \{x_1, \dots, x_n\}, d_X) \in \mathcal{M}_n$ and define $B_{\mathcal{X}}$ as before. Then, X is isometric to some $\{z_1, \dots, z_n\} \subset \mathbb{R}^k$ for some positive integer k if and only if $B_{\mathcal{X}}$ is positive semi-definite.*

Proof. (\Rightarrow) We prove the *only if* direction first. Because of the assumption, there exist $\{z_1, \dots, z_n\} \subset \mathbb{R}^k$ such that $d_X(x_i, x_j) = \|z_i - z_j\|$ for each $i, j = 1, \dots, n$. Therefore,

$$(A_{\mathcal{X}})_{ij} = d_X^2(x_i, x_j) = \langle z_i - z_j, z_i - z_j \rangle = \langle z_i, z_i \rangle + \langle z_j, z_j \rangle - 2\langle z_i, z_j \rangle$$

and,

$$\begin{aligned} \frac{1}{n} \sum_{s=1}^n (A_{\mathcal{X}})_{is} &= \frac{1}{n} \sum_{s=1}^n \langle z_s, z_s \rangle + \langle z_i, z_i \rangle - 2\langle z_i, \bar{z} \rangle, \\ \frac{1}{n} \sum_{r=1}^n (A_{\mathcal{X}})_{rj} &= \frac{1}{n} \sum_{r=1}^n \langle z_r, z_r \rangle + \langle z_j, z_j \rangle - 2\langle \bar{z}, z_j \rangle, \\ \frac{1}{n^2} \sum_{r,s=1}^n (A_{\mathcal{X}})_{rs} &= \frac{1}{n} \sum_{s=1}^n \langle z_s, z_s \rangle + \frac{1}{n} \sum_{r=1}^n \langle z_r, z_r \rangle - 2\langle \bar{z}, \bar{z} \rangle \end{aligned}$$

for each $i, j = 1, \dots, n$ where \bar{z} is the mean point of z_1, \dots, z_n . Also, by Lemma 3.1 item (2),

$$(B_{\mathcal{X}})_{ij} = -\frac{1}{2} \left((A_{\mathcal{X}})_{ij} - \frac{1}{n} \sum_{s=1}^n (A_{\mathcal{X}})_{is} - \frac{1}{n} \sum_{r=1}^n (A_{\mathcal{X}})_{rj} + \frac{1}{n^2} \sum_{r,s=1}^n (A_{\mathcal{X}})_{rs} \right)$$

for each $i, j = 1, \dots, n$. Hence,

$$(B_{\mathcal{X}})_{ij} = \langle z_i - \bar{z}, z_j - \bar{z} \rangle$$

for each $i, j = 1, \dots, n$. This implies that $B_{\mathcal{X}} = (ZH_n)^T(ZH_n)$ where Z is the k by n matrix whose i -th column is the z_i so that $ZH_n = Z - \frac{1}{n}Ze_n e_n^T$ is the n by k matrix whose i -th column is the $z_i - \bar{z}$. Therefore, for any nonzero vector $x \in \ell^2(n)$,

$$x^T B_{\mathcal{X}} x = (ZH_n x)^T (ZH_n x) \geq 0.$$

Therefore, $B_{\mathcal{X}}$ is positive semi-definite.

(\Leftarrow) Now, let's prove the *if* direction. Since $B_{\mathcal{X}}$ is symmetric and positive semi-definite, we have eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n \geq 0$ and corresponding orthonormal eigenvectors v_1, \dots, v_n . Moreover,

$$B_{\mathcal{X}} = V\Lambda V^T$$

where $V = [v_1 \ v_2 \ \dots \ v_n]$ and Λ is the diagonal matrix with $\lambda_1, \dots, \lambda_n$. Now, let $k = \text{rank}(B_{\mathcal{X}})$. It means that $\lambda_1 \geq \dots \geq \lambda_k > 0 = \lambda_{k+1} = \dots = \lambda_n$. Then,

$$B_{\mathcal{X}} = \widehat{V}\widehat{\Lambda}\widehat{V}^T$$

where $\widehat{V} = [v_1 \ \dots \ v_k]$ and $\widehat{\Lambda}$ is the k by k diagonal matrix with $\lambda_1, \dots, \lambda_k$. Let $Z := \widehat{\Lambda}^{\frac{1}{2}}\widehat{V}^T$ and let $z_1, \dots, z_n \in \mathbb{R}^k$ be its column vectors. Then we have $(B_{\mathcal{X}})_{ij} = \langle z_i, z_j \rangle$ since $B_{\mathcal{X}} = Z^T Z$. Hence,

$$\|z_i - z_j\|^2 = \langle z_i - z_j, z_i - z_j \rangle = (B_{\mathcal{X}})_{ii} + (B_{\mathcal{X}})_{jj} - 2(B_{\mathcal{X}})_{ij} = d_X^2(x_i, x_j)$$

by Lemma 3.1 item (3). Hence $\{z_1, \dots, z_n\} \subset \mathbb{R}^k$ is isometric to X . \square

Remark 3.3. *As one sees in the proof, if X is isometric to $\{z_1, \dots, z_n\} \subset \mathbb{R}^k$, then the rank of $B_{\mathcal{X}}$ is less than or equal to k since $B_{\mathcal{X}} = (ZH_n)^T(ZH_n)$ where ZH_n is a k by n matrix.*

Definition 3.4. For each square symmetric matrix A of size n ,

$$\text{pr}(A) := \text{the number of positive eigenvalues of } A.$$

Of course, not every finite metric space can be isometrically embedded into Euclidean space. However, Schoenberg's result shows that the negative eigenvalues of $B_{\mathcal{X}}$ are the obstruction to achieving this. Indeed, by eliminating those negative eigenvalues, the classical cMDS algorithm [MKB79] gives a precise way of building a cMDS embedding into \mathbb{R}^k $\Phi_{\mathcal{X},k} : X \rightarrow \mathbb{R}^k$ together with its induced finite metric space $\widehat{\mathcal{X}}_k := (\Phi_{\mathcal{X},k}(X) \subset \mathbb{R}^k, \|\cdot\|)$:

Algorithm 1.

- (1) **Input:** $\mathcal{X} = (X, d_X) \in \mathcal{M}_n$.
- (2) Construct $A_{\mathcal{X}}$ and $B_{\mathcal{X}}$ as before.
- (3) Compute the eigenvalues $\lambda_1 \geq \dots \geq \lambda_{\text{pr}(B_{\mathcal{X}})} > 0 = \lambda_{\text{pr}(B_{\mathcal{X}})+1} \geq \dots \geq \lambda_n$ and corresponding orthonormal eigenvectors v_1, \dots, v_n of $B_{\mathcal{X}}$.
- (4) Choose $k \leq \text{pr}(B_{\mathcal{X}})$.
- (5) Construct $\widehat{\Lambda}_k^{\frac{1}{2}}\widehat{V}_k^T$ where $\widehat{\Lambda}_k := \text{diag}(\lambda_1, \lambda_2, \dots, \lambda_k)$ and $\widehat{V}_k := [v_1 \ \dots \ v_k]$. Let $\Phi_{\mathcal{X},k}(x_i) \in \mathbb{R}^k$ be the i -th column vector of $\widehat{\Lambda}_k^{\frac{1}{2}}\widehat{V}_k^T$ for each $i = 1, \dots, n$.
- (6) **Output:** $\Phi_{\mathcal{X},k} : X \rightarrow \mathbb{R}^k$, and $\widehat{\mathcal{X}}_k := (\Phi_{\mathcal{X},k}(X), \|\cdot\|)$.

Remark 3.5. *Here are some simple facts from the above cMDS procedure.*

- (1) $\Phi_{\mathcal{X},k}(x_i) = (\lambda_1^{\frac{1}{2}}v_1(i), \dots, \lambda_k^{\frac{1}{2}}v_k(i))^T$ for each $i = 1, \dots, n$.
- (2) *The mean point of $\Phi_{\mathcal{X},k}(x_1), \dots, \Phi_{\mathcal{X},k}(x_n)$ is 0 by Lemma 3.1 item (5).*

- (3) $\lambda_1 \geq \dots \geq \lambda_k$ are all positive eigenvalues of $B_{\hat{\mathcal{X}}_k}$ and all the other eigenvalues are zero.




FIGURE 1. Description of the cMDS procedure into \mathbb{R}^2 from Example 3.6. Blue lines indicate length 1, red lines indicate length 2, green lines indicate length $\frac{2}{\sqrt{3}}$, and purple lines indicate length 2.

Example 3.6. Consider $(X = \{x_0, x_1, x_2, x_3\}, d_X) \in \mathcal{M}_4$ such that $d_X(x_0, x_i) = 1$ for each $i = 1, 2, 3$ and $d_X(x_i, x_j) = 2$ for each $i \neq j \in \{1, 2, 3\}$. Then,

$$A_X = \begin{pmatrix} 0 & 1 & 1 & 1 \\ 1 & 0 & 4 & 4 \\ 1 & 4 & 0 & 4 \\ 1 & 4 & 4 & 0 \end{pmatrix}$$

and,

$$B_X = \frac{1}{16} \cdot \begin{pmatrix} -3 & 1 & 1 & 1 \\ 1 & 21 & -11 & -11 \\ 1 & -11 & 21 & -11 \\ 1 & -11 & -11 & 21 \end{pmatrix}$$

The eigenvalues of B_X are $2, 2, 0, -1/4$. Their corresponding mutually orthogonal eigenvectors are $(0, 1, 0, -1)$, $(0, -\frac{1}{2}, 1, -\frac{1}{2})$, $(1, 1, 1, 1)$, $(-3, 1, 1, 1)$.

Therefore, the result of cMDS into \mathbb{R}^2 is the following: $\Phi_{X,2}(x_0) = (0, 0)$, $\Phi_{X,2}(x_1) = (1, -\frac{1}{\sqrt{3}})$, $\Phi_{X,2}(x_2) = (0, \frac{2}{\sqrt{3}})$, and $\Phi_{X,2}(x_3) = (-1, -\frac{1}{\sqrt{3}})$.

Moreover, this $\hat{\mathcal{X}}_k$ is the optimal one in the following sense.

Theorem 16 ([MKB79, Mar78]). Suppose $\mathcal{X} = (X, d_X) \in \mathcal{M}_n$ and $k \leq \text{pr}(B_X)$ are given. Then, for any positive semi-definite matrix A of size n with $\text{rank}(A) \leq k$, we have

$$\|B_X - A\|_F \geq \|B_X - B_{\hat{\mathcal{X}}_k}\|_F$$

where $\|\cdot\|_F$ is the Frobenius norm.

For pedagogical purposes, we provide a full proof² of Theorem 16 which will be suitably generalised in Theorem 19.

²We follow the general lines from [MKB79, CH14].

Theorem 17 (rearrangement inequality [HLP67, Theorem 368]). *Suppose we have two sequences of real numbers $x_1 \leq \dots \leq x_n$ and $y_1 \leq \dots \leq y_n$ for some positive integer n . Then,*

$$x_n y_1 + x_{n-1} y_2 + \dots + x_1 y_n \leq x_{\sigma(1)} y_1 + x_{\sigma(2)} y_2 + \dots + x_{\sigma(n)} y_n \leq x_1 y_1 + x_2 y_2 + \dots + x_n y_n$$

for any permutation σ on $[n] = \{1, \dots, n\}$.

Definition 3.7 (doubly stochastic matrix). A square matrix $A = (a_{ij})_{i,j=1}^n$ of size n is called **doubly stochastic** if it satisfies:

- (1) $a_{ij} \geq 0$ for any $i, j \in \{1, \dots, n\}$.
- (2) $\sum_{j=1}^n a_{ij} = 1$ for any $i \in \{1, \dots, n\}$.
- (3) $\sum_{i=1}^n a_{ij} = 1$ for any $j \in \{1, \dots, n\}$.

Theorem 18 (Birkhoff–von Neumann theorem [Bru06, Theorem 1.7.1]). *Any doubly stochastic matrix is a convex combination of permutation matrices.*

Lemma 3.8 ([MKB79, Exercise 14.4.2]). *Let $M = \text{diag}(\lambda_1, \dots, \lambda_n)$ where $\lambda_1 \geq \dots \geq \lambda_n$, and $N = \text{diag}(\xi_1, \dots, \xi_n)$ where $\xi_1 \geq \dots \geq \xi_n$. Then,*

$$\text{Tr}(M - UNU^T)^2 \geq \text{Tr}(M - N)^2$$

for any orthogonal matrix $U = (u_{ij})_{i,j=1}^n$.

Proof. Since $\text{Tr}(M - UNU^T)^2 = \text{Tr} M^2 + \text{Tr} N^2 - 2\text{Tr}(MUNU^T)$ and $\text{Tr}(M - N)^2 = \text{Tr} M^2 + \text{Tr} N^2 - 2\text{Tr}(MN)$, it is enough to show $\text{Tr}(MUNU^T) \leq \text{Tr}(MN)$. First, observe that

$$\text{Tr}(MUNU^T) = \sum_{i=1}^n \lambda_i \left(\sum_{j=1}^n \xi_j u_{ij}^2 \right) = \sum_{i=1}^n \lambda_i h_i$$

where $h_i = \sum_{j=1}^n \xi_j u_{ij}^2$ for each $i = 1, \dots, n$. Since U is orthogonal, $\sum_{j=1}^n u_{ij}^2 = 1$ for each $i = 1, \dots, n$, and $\sum_{i=1}^n u_{ij}^2 = 1$ for each $j = 1, \dots, n$. In other words, $(u_{ij}^2)_{i,j=1}^n$ is a doubly stochastic matrix. Hence, by Theorem 18, $(u_{ij}^2)_{i,j=1}^n$ is a convex combination of permutation matrices. Hence, there are non-negative numbers c_1, \dots, c_k such that $c_1 + \dots + c_k = 1$ and permutations $\sigma_1, \dots, \sigma_n$ on $\{1, \dots, n\}$ such that

$$(h_1, \dots, h_n) = c_1(\xi_{\sigma_1(1)}, \dots, \xi_{\sigma_1(n)}) + \dots + c_k(\xi_{\sigma_k(1)}, \dots, \xi_{\sigma_k(n)}).$$

Therefore, by Theorem 17, we have

$$\text{Tr}(MUNU^T) = \sum_{i=1}^n \lambda_i h_i \leq \sum_{i=1}^n \lambda_i \xi_i = \text{Tr}(MN)$$

as we wanted. □

Proof of Theorem 16. Let $\lambda_1 \geq \dots \geq \lambda_{\text{pr}(B_{\mathcal{X}})} > 0 = \lambda_{\text{pr}(B_{\mathcal{X}})+1} \geq \dots \geq \lambda_n$ be the eigenvalues of $B_{\mathcal{X}}$. Then, we have $B_{\mathcal{X}} = VMV^T$ for some orthogonal matrix V and $M = \text{diag}(\lambda_1, \dots, \lambda_n)$. Similarly, let $\xi_1 \geq \dots \geq \xi_k \geq 0 = \xi_{k+1} = \dots = \xi_n$ be the eigenvalues of A . Then, $A = WNW^T$ for some orthogonal matrix W and $N = \text{diag}(\xi_1, \dots, \xi_n)$. Therefore,

$$V^T A V = V^T W N W^T V = UNU^T$$

where $U = V^T W$ is another orthogonal matrix. Hence,

$$\begin{aligned}
\|B_{\mathcal{X}} - A\|_{\text{F}}^2 &= \text{Tr}(B_{\mathcal{X}} - A)(B_{\mathcal{X}} - A)^T \\
&= \text{Tr}(B_{\mathcal{X}} - A)^2 \\
&= \text{Tr}(V^T(B_{\mathcal{X}} - A)V V^T(B_{\mathcal{X}} - A)V) \\
&= \text{Tr}(M - U N U^T)(M - U N U^T) \\
&= \text{Tr}(M - U N U^T)^2 \\
&\geq \text{Tr}(M - N)^2 \quad (\text{by Lemma 3.8}) \\
&= \sum_{i=1}^n (\lambda_i - \xi_i)^2 \\
&= \sum_{i=1}^k (\lambda_i - \xi_i)^2 + \sum_{i=k+1}^n \lambda_i^2 \\
&\geq \sum_{i=k+1}^n \lambda_i^2 = \|B_{\mathcal{X}} - B_{\hat{\mathcal{X}}_k}\|_{\text{F}}^2.
\end{aligned}$$

□

Now that we know that the cMDS methods gives an optimal embedding into \mathbb{R}^k (with respect to the Frobenius norm), the next natural goal is to estimate the metric distortion incurred during this process. To understand this, we need the following definition.

Definition 3.9. For each square symmetric matrix A of size n ,

$$\text{Tr}_{\text{neg}}(A) := \sum_{i=1}^n (\lambda_i)^-$$

where $\lambda_1, \dots, \lambda_n$ are eigenvalues of A . Equivalently, $\text{Tr}_{\text{neg}}(A)$ is the sum of absolute values of all negative eigenvalues of A .

The following proposition shows the the metric distortion incurred during the cMDS method is controlled by $\text{Tr}_{\text{neg}}(B_{\mathcal{X}})$.

Proposition 3.10 (cMDS bounds). *Let $\mathcal{X} = (X = \{x_1, \dots, x_n\}, d_X) \in \mathcal{M}_n$. Then the following bounds hold:*

- (1) For any $i, j = 1, \dots, n$ and $k \leq \text{pr}(B_{\mathcal{X}})$,

$$d_X(x_i, x_j) \leq \|\Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_i) - \Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_j)\|$$

and

$$\|\Phi_{\mathcal{X}, k}(x_i) - \Phi_{\mathcal{X}, k}(x_j)\| \leq \|\Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_i) - \Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_j)\|.$$

- (2) (L^∞ bound) For any $i, j = 1, \dots, n$ and $k \leq \text{pr}(B_{\mathcal{X}})$,

$$\|\Phi_{\mathcal{X}, k}(x_i) - \Phi_{\mathcal{X}, k}(x_j)\| \leq d_X(x_i, x_j) + \sqrt{2 \text{Tr}_{\text{neg}}(B_{\mathcal{X}})}.$$

- (3) (L^2 bound)

$$\left(\frac{1}{n^2} \sum_{i, j=1}^n (\|\Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_i) - \Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_j)\| - d_X(x_i, x_j))^2 \right)^{\frac{1}{2}} \leq 2 \sqrt{\frac{\text{Tr}_{\text{neg}}(B_{\mathcal{X}})}{n}}.$$

This result will be generalized to arbitrary (trace class) metric measure spaces in Proposition 4.22.

Proof of Proposition 3.10. (1) Let $\lambda_1 \geq \dots \geq \lambda_{\text{pr}(B_{\mathcal{X}})} > 0 = \lambda_{\text{pr}(B_{\mathcal{X}})+1} \geq \dots \geq \lambda_n$ be the eigenvalues of $B_{\mathcal{X}}$ and $\{v_m\}_{m=1}^n$ be corresponding orthonormal eigenvectors. Then, by item (3) of Lemma 3.1 and item (1) of Remark 3.5,

$$\begin{aligned} d_X^2(x_i, x_j) &= \sum_{m=1}^n \lambda_m (v_m(i) - v_m(j))^2, \\ \|\Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_i) - \Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_j)\|^2 &= \sum_{m=1}^{\text{pr}(B_{\mathcal{X}})} \lambda_m (v_m(i) - v_m(j))^2, \text{ and} \\ \|\Phi_{\mathcal{X}, k}(x_i) - \Phi_{\mathcal{X}, k}(x_j)\|^2 &= \sum_{m=1}^k \lambda_m (v_m(i) - v_m(j))^2 \end{aligned}$$

for any $i, j = 1, \dots, n$. Then, because of the sign of eigenvalues,

$$\|\Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_i) - \Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_j)\|^2 - d_X^2(x_i, x_j) = \sum_{m=\text{pr}(B_{\mathcal{X}})+1}^n -\lambda_m (v_m(i) - v_m(j))^2 \geq 0, \text{ and}$$

$$\|\Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_i) - \Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_j)\|^2 - \|\Phi_{\mathcal{X}, k}(x_i) - \Phi_{\mathcal{X}, k}(x_j)\|^2 = \sum_{m=k+1}^{\text{pr}(B_{\mathcal{X}})} \lambda_m (v_m(i) - v_m(j))^2 \geq 0$$

for any $i, j = 1, \dots, n$. The required results follow.

(2) Observe that, for any $i, j = 1, \dots, n$,

$$\begin{aligned} \|\Phi_{\mathcal{X}, k}(x_i) - \Phi_{\mathcal{X}, k}(x_j)\|^2 - d_X^2(x_i, x_j) &= \sum_{m=k+1}^n -\lambda_m (v_m(i) - v_m(j))^2 \\ &\leq \sum_{m=\text{pr}(B_{\mathcal{X}})+1}^n -\lambda_m (v_m(i) - v_m(j))^2 \\ &\leq \sum_{m=\text{pr}(B_{\mathcal{X}})+1}^n -2\lambda_m (v_m(i)^2 + v_m(j)^2) \\ &\leq 2 \text{Tr}_{\text{neg}}(B_{\mathcal{X}}). \end{aligned}$$

where the last inequality is true because $(v_m(i)^2 + v_m(j)^2) \leq 1$ for any $m = 1, \dots, n$. Hence,

$$\begin{aligned} \|\Phi_{\mathcal{X}, k}(x_i) - \Phi_{\mathcal{X}, k}(x_j)\| &\leq \sqrt{d_X^2(x_i, x_j) + 2 \text{Tr}_{\text{neg}}(B_{\mathcal{X}})} \\ &\leq d_X(x_i, x_j) + \sqrt{2 \text{Tr}_{\text{neg}}(B_{\mathcal{X}})}. \end{aligned}$$

(3) We now obtain an L^2 bound. Since we already know that

$$d_X(x_i, x_j) \leq \|\Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_i) - \Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_j)\|$$

for any $i, j = 1, \dots, n$,

$$\begin{aligned}
(\|\Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_i) - \Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_j)\| - d_X(x_i, x_j))^2 &\leq \|\Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_i) - \Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_j)\|^2 - d_X^2(x_i, x_j) \\
&= \sum_{m=\text{pr}(B_{\mathcal{X}})+1}^n -\lambda_m (v_m(i) - v_m(j))^2 \\
&\leq \sum_{m=\text{pr}(B_{\mathcal{X}})+1}^n -2\lambda_m (v_m(i)^2 + v_m(j)^2)
\end{aligned}$$

for any $i, j = 1, \dots, n$. Therefore,

$$\begin{aligned}
\sum_{i,j=1}^n (\|\Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_i) - \Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_j)\| - d_X(x_i, x_j))^2 &\leq 2 \sum_{m=\text{pr}(B_{\mathcal{X}})+1}^n - \left(\sum_{i,j=1}^n (v_m(i)^2 + v_m(j)^2) \right) \lambda_m \\
&= 4n \text{Tr}_{\text{neg}}(B_{\mathcal{X}}).
\end{aligned}$$

It follows that

$$\left(\frac{1}{n^2} \sum_{i,j=1}^n (\|\Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_i) - \Phi_{\mathcal{X}, \text{pr}(B_{\mathcal{X}})}(x_j)\| - d_X(x_i, x_j))^2 \right)^{\frac{1}{2}} \leq 2\sqrt{\frac{\text{Tr}_{\text{neg}}(B_{\mathcal{X}})}{n}}.$$

□

4. GENERALIZED cMDS ON METRIC MEASURE SPACES

Assume a metric measure space $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$ is given. Partially motivated by [ZZ03] and cMDS for finite metric spaces, and following [Kas19, ABK20], we now define a generalized cMDS procedure for the metric measure space \mathcal{X} in the following way.

First, define the integral kernel $K_{\mathcal{X}} : X \times X \rightarrow \mathbb{R}$ by

$$\begin{aligned}
&K_{\mathcal{X}}(x, x') \\
&:= -\frac{1}{2} \left(d_X^2(x, x') - \int_X d_X^2(x, v) d\mu_X(v) - \int_X d_X^2(u, x') d\mu_X(u) + \iint_{X \times X} d_X^2(u, v) d\mu_X(u) d\mu_X(v) \right)
\end{aligned}$$

for any $x, x' \in X$.

Remark 4.1. Observe that the last term in the above definition is $(\text{diam}_2(\mathcal{X}))^2$, c.f. equation (1).

Note that, since X is compact, $\text{diam}(X) < \infty$ and

$$\sup_{x, x' \in X} |K_{\mathcal{X}}(x, x')| \leq 2 (\text{diam}(X))^2.$$

Next, define the integral operator $\mathfrak{K}_{\mathcal{X}} : L^2(\mu_X) \rightarrow L^2(\mu_X)$ given by

$$\phi \mapsto \mathfrak{K}_{\mathcal{X}}(\phi) := \int_X K_{\mathcal{X}}(x, x') \phi(x') d\mu_X(x').$$

Let us collect some of simple properties of $K_{\mathcal{X}}$ and $\mathfrak{K}_{\mathcal{X}}$.

Lemma 4.2. Let $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$ be given and let $K_{\mathcal{X}}$ and $\mathfrak{K}_{\mathcal{X}}$ be as above. Then,

- (1) $K_{\mathcal{X}}(x, x') = K_{\mathcal{X}}(x', x)$ for any $x, x' \in X$. In particular, $\mathfrak{K}_{\mathcal{X}}$ is a self-adjoint operator.
- (2) $\mathfrak{K}_{\mathcal{X}}$ is a Hilbert-Schmidt operator. In particular, it is a compact operator.

- (3) $d_X^2(x, x') = K_{\mathcal{X}}(x, x) + K_{\mathcal{X}}(x', x') - 2K_{\mathcal{X}}(x, x')$ for any $x, x' \in X$.
 (4) $\phi \equiv 1$ is an eigenfunction of $\mathfrak{K}_{\mathcal{X}}$ with zero eigenvalue. In particular,

$$\dim(\ker(\mathfrak{K}_{\mathcal{X}})) \geq 1.$$

- (5) If $\phi \in L^2(\mu_X)$ is an eigenfunction for a nonzero eigenvalue of $\mathfrak{K}_{\mathcal{X}}$, then

$$\int_X \phi(x) d\mu_X(x) = 0.$$

- (6) If $\phi \in L^2(\mu_X)$ is an eigenfunction for a nonzero eigenvalue of $\mathfrak{K}_{\mathcal{X}}$, then there is a uniformly continuous function ϕ' such that $\phi = \phi'$ μ_X -almost everywhere.

Proof. (1) Obvious from the definition since d_X is metric.

- (2) Observe that $\int_X \int_X (K_{\mathcal{X}}(x, x'))^2 d\mu_X(x') d\mu_X(x) \leq 4 (\text{diam}(X))^4 < \infty$. This implies that $\mathfrak{K}_{\mathcal{X}}$ is a Hilbert-Schmidt operator. In particular, this is a compact operator.

- (3) Easy to show by using the definition and direct computation.

- (4) Let $\phi \equiv 1$. Then,

$$\begin{aligned} \mathfrak{K}_{\mathcal{X}}(\phi)(x) &= \int_X K_{\mathcal{X}}(x, x') \phi(x') d\mu_X(x') \\ &= \int_X K_{\mathcal{X}}(x, x') d\mu_X(x') \\ &= -\frac{1}{2} \left(\int_X d_X^2(x, x') d\mu_X(x') - \int_X d_X^2(x, v) d\mu_X(v) \right. \\ &\quad \left. - \iint_{X \times X} d_X^2(u, x') d\mu_X(u) d\mu_X(x') + \iint_{X \times X} d_X^2(u, v) d\mu_X(u) d\mu_X(v) \right) \\ &= 0 \end{aligned}$$

for any $x \in X$.

- (5) Obvious by item (3) of Theorem 2 and item (4) of this lemma.

- (6) Let ϕ be an eigenfunction of $\mathfrak{K}_{\mathcal{X}}$ with nonzero eigenvalue λ . This implies $\phi(x) = \frac{1}{\lambda} \mathfrak{K}_{\mathcal{X}} \phi(x)$ for μ_X -a.e. $x \in X$. Hence, it is enough to show that $\phi' := \frac{1}{\lambda} \mathfrak{K}_{\mathcal{X}} \phi$ is a uniformly continuous function on X . Since $K_{\mathcal{X}}$ is continuous on a compact space $X \times X$, it is uniformly continuous. Therefore, for arbitrary $\varepsilon > 0$, there exists $\delta > 0$ such that $|K_{\mathcal{X}}(x, x'') - K_{\mathcal{X}}(x', x'')| < \varepsilon$ for any $x, x', x'' \in X$ whenever $d_X(x, x') < \delta$. Moreover, observe that $\int_X |\phi| d\mu_X = C_{\phi} > 0$ since otherwise $\phi = 0$ μ_X -almost everywhere so it contradicts to the fact that $\lambda \neq 0$. Also, C_{ϕ} cannot be infinity because $L^2(\mu_X) \subset L^1(\mu_X)$ (see [Fol13, Proposition 6.12]). Now, observe that

$$\begin{aligned} \left| \frac{1}{\lambda} \mathfrak{K}_{\mathcal{X}} \phi(x) - \frac{1}{\lambda} \mathfrak{K}_{\mathcal{X}} \phi(x') \right| &= \frac{1}{\lambda} \left| \int_X (K_{\mathcal{X}}(x, x'') - K_{\mathcal{X}}(x', x'')) \phi(x'') d\mu_X(x'') \right| \\ &\leq \frac{1}{\lambda} \int_X |K_{\mathcal{X}}(x, x'') - K_{\mathcal{X}}(x', x'')| |\phi(x'')| d\mu_X(x'') \\ &\leq \frac{\varepsilon}{\lambda} \int_X |\phi(x'')| d\mu_X(x'') = \frac{C_{\phi} \varepsilon}{\lambda} \end{aligned}$$

for any $x, x' \in X$ with $d_X(x, x') < \delta$. Since $\varepsilon > 0$ is arbitrary, this concludes the proof. \square

Even though the definition of $K_{\mathcal{X}}$ is rather complicated, finding the spectrum of $\mathfrak{K}_{\mathcal{X}}$ can be reduced to a simpler eigenproblem as one can see in the following results.

Lemma 4.3. *Suppose $\mathcal{X} \in \mathcal{M}_w$ is given. If $\lambda \in \mathbb{R}$ and $0 \neq \phi \in L^2(\mu_X)$ satisfy the following two conditions:*

- (1) $-\frac{1}{2} \int_X d_X^2(x, x') \phi(x') d\mu_X(x') = \lambda \phi(x)$ for μ_X -a.e $x \in X$, and
- (2) $\int_X \phi(x) d\mu_X(x) = 0$,

then λ is an eigenvalue of $\mathfrak{K}_{\mathcal{X}}$ and ϕ is a associated eigenfunction.

Proof. Assume ϕ and λ satisfy the two conditions above. Observe that, for any $x \in X$, we have

$$\begin{aligned} & \int_X K_{\mathcal{X}}(x, x') \phi(x') d\mu_X(x') \\ &= -\frac{1}{2} \left(\int_X d_X^2(x, x') \phi(x') d\mu_X(x') - \int_X d_X^2(x, v) d\mu_X(v) \cdot \int_X \phi(x') d\mu_X(x') \right. \\ & \quad \left. - \iint_{X \times X} d_X^2(u, x') \phi(x') d\mu_X(u) d\mu_X(x') + \iint_{X \times X} d_X^2(u, v) d\mu_X(u) d\mu_X(v) \cdot \int_X \phi(x') d\mu_X(x') \right) \\ &= \lambda \phi(x) - \lambda \int_X \phi(u) d\mu_X(u) \\ &= \lambda \phi(x). \end{aligned}$$

This completes the proof. □

Moreover, the reverse implication of Lemma 4.3 holds for nonzero eigenvalues in the setting of **two-point homogeneous spaces** (cf. Corollary 4.6).

Definition 4.4 (two-point homogeneous space). A metric measure space $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$ is said to be **two-point homogeneous** if it satisfies the following condition:

For arbitrary $x, x' \in X$ there is an isometry $f : X \rightarrow X$ such that $f_*\mu_X = \mu_X$ and $f(x) = x'$.

Proposition 4.5. *Suppose $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$ is two-point homogeneous. Then,*

$$K_{\mathcal{X}}(x, x') = -\frac{1}{2} \left(d_X^2(x, x') - (\text{diam}_2(\mathcal{X}))^2 \right)$$

for any $x, x' \in X$.

Proof. Fix arbitrary $x, x' \in X$. Then, since \mathcal{X} is two-point homogeneous, there exist an isometry f such that $f(x) = x'$ and $f_*\mu_X = \mu_X$. Hence,

$$\begin{aligned} \int_X d_X^2(x, u) d\mu_X(u) &= \int_X d_X^2(f(x), f(u)) d\mu_X(u) = \int_X d_X^2(x', f(u)) d\mu_X(u) \\ &= \int_X d_X^2(x', u') df_*\mu_X(u') = \int_X d_X^2(x', u') d\mu_X(u'). \end{aligned}$$

This implies that the map $x \mapsto \int_X d_X^2(x, u) d\mu_X(u)$ is constant, and therefore the constant is equal to its average $(\text{diam}_2(\mathcal{X}))^2$. Hence,

$$\begin{aligned} K_{\mathcal{X}}(x, x') &= -\frac{1}{2} \left(d_X^2(x, x') - \int_X d_X^2(x, v) d\mu_X(v) - \int_X d_X^2(u, x') d\mu_X(u) + \iint_{X \times X} d_X^2(u, v) d\mu_X(u) d\mu_X(v) \right) \\ &= -\frac{1}{2} (d_X^2(x, x') - \text{diam}_2(\mathcal{X})^2) \end{aligned}$$

for any $x, x' \in X$ as we required. \square

Corollary 4.6. *Suppose $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$ is two-point homogeneous. Then, if λ is a nonzero eigenvalue of $\mathfrak{K}_{\mathcal{X}}$ with an associated eigenfunction $\phi \in L^2(\mu_X)$, then the following two equalities hold:*

- (1) $-\frac{1}{2} \int_X d_X^2(x, x') \phi(x') d\mu_X(x') = \lambda \phi(x)$ for μ_X -a.e $x \in X$, and
- (2) $\int_X \phi(x) d\mu_X(x) = 0$.

Proof. Apply Proposition 4.5 and item (5) of Lemma 4.2. \square

The above corollary will be useful when studying the spectrum of $\mathfrak{K}_{\mathbb{S}^{d-1}}$ (see §6.2 and §6.4).

Remark 4.7. *For a compact subset X of \mathbb{R}^k with $m_k(X) > 0$, we consider*

$$\mathcal{X} = (X, \|\cdot\|, \overline{m}_k) \in \mathcal{M}_w$$

where $\|\cdot\|$ indicates the metric on X induced by the Euclidean norm, and $\overline{m}_k := \frac{1}{m_k(X)} m_k$ is the normalized k -dimensional Lebesgue measure on X . Then, one can easily show that

$$K_{\mathcal{X}}(x, x') = \langle x - \text{cm}(\mathcal{X}), x' - \text{cm}(\mathcal{X}) \rangle$$

for any $x, x' \in X$ where $\text{cm}(\mathcal{X}) := \int_X x d\overline{m}_k(x) \in \mathbb{R}^k$ is the center of mass of \mathcal{X} . This means that $\mathfrak{K}_{\mathcal{X}}$ is positive semi-definite, since

$$\langle \mathfrak{K}_{\mathcal{X}} \phi, \phi \rangle = \left\| \int_X (x - \text{cm}(\mathcal{X})) \phi(x) d\overline{m}_k(x) \right\|^2 \geq 0$$

for any $\phi \in L^2(\overline{m}_k)$. Moreover, observe that $\mathfrak{K}_{\mathcal{X}} = \mathfrak{K}_2 \circ \mathfrak{K}_1$ where

$$\begin{aligned} \mathfrak{K}_1 : L^2(\overline{m}_k) &\longrightarrow \mathbb{R}^k \\ \phi &\longmapsto \int_X (x - \text{cm}(\mathcal{X})) \phi(x) d\overline{m}_k(x), \end{aligned}$$

and

$$\begin{aligned} \mathfrak{K}_2 : \mathbb{R}^k &\longrightarrow L^2(\overline{m}_k) \\ x &\longmapsto \langle \cdot - \text{cm}(\mathcal{X}), x \rangle. \end{aligned}$$

In particular, this implies that the rank of $\mathfrak{K}_{\mathcal{X}}$ is at most k since the domain of \mathfrak{K}_2 is \mathbb{R}^k .

4.1. **The formulation of cMDS for metric measure spaces.** Observe that, by item (1) and (2) of Lemma 4.2, Theorem 4, and Theorem 5, \mathfrak{K}_X has discrete spectrum and one can apply spectral theory.

Definition 4.8. Let \mathfrak{A} be a bounded operator on a Hilbert space \mathcal{H} . Then,

$$\begin{aligned} \text{pr}(\mathfrak{A}) &:= \text{the rank of } \mathfrak{A}^+ \\ \text{nr}(\mathfrak{A}) &:= \text{the rank of } \mathfrak{A}^-. \end{aligned}$$

Finally, cMDS is generalized in the following way:

Algorithm 2.

- (1) **Input:** $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$.
- (2) Construct K_X and \mathfrak{K}_X as before.
- (3) Compute the positive eigenvalues $\lambda_1 \geq \dots \geq \lambda_{\text{pr}(\mathfrak{K}_X)} > 0$ ($\text{pr}(\mathfrak{K}_X)$ can be infinite) and corresponding orthonormal eigenfunctions $\{\phi_i\}_{i=1}^{\text{pr}(\mathfrak{K}_X)}$ of \mathfrak{K}_X .
- (4) Choose any positive integer $k \leq \text{pr}(\mathfrak{K}_X)$.
- (5) $\Phi_{\mathcal{X},k}(x) := (\lambda_1^{\frac{1}{2}}\phi_1(x), \lambda_2^{\frac{1}{2}}\phi_2(x), \dots, \lambda_k^{\frac{1}{2}}\phi_k(x)) \in \mathbb{R}^k$ for each $x \in X$.
- (6) **Output:** $\Phi_{\mathcal{X},k} : X \rightarrow \mathbb{R}^k$, and $\widehat{\mathcal{X}}_k := (\Phi_{\mathcal{X},k}(X), \|\cdot\|, (\Phi_{\mathcal{X},k})_*\mu_X) \in \mathcal{M}_w$.

Remark 4.9. By item (6) of Lemma 4.2, one can assume $\Phi_{\mathcal{X},k}$ is continuous.

Definition 4.10 (generalized cMDS into \mathbb{R}^k). Given a metric measure space $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$ and orthonormal eigenfunctions $a = \{\phi_i\}_{i=1}^{\text{pr}(\mathfrak{K}_X)}$ of \mathfrak{K}_X for the corresponding positive eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots > 0$, one defines the following map:

$$\begin{aligned} \Phi_{\mathcal{X},k}^a : X &\longrightarrow \mathbb{R}^k \\ x &\longmapsto (\lambda_1^{\frac{1}{2}}\phi_1(x), \lambda_2^{\frac{1}{2}}\phi_2(x), \dots, \lambda_k^{\frac{1}{2}}\phi_k(x)). \end{aligned}$$

This map $\Phi_{\mathcal{X},k}^a$ is called **generalized cMDS embedding into \mathbb{R}^k** of \mathcal{X} , and the resulting metric measure space is denoted by

$$\widehat{\mathcal{X}}_k := (\Phi_{\mathcal{X},k}^a(X), \|\cdot\|, (\Phi_{\mathcal{X},k}^a)_*\mu_X) \in \mathcal{M}_w.$$

In practice, the superscript a will be omitted in many cases for the simplicity, unless we have to specify the set of orthonormal eigenfunctions.

One critical difference between the cMDS method for finite metric spaces (see §3) and the generalized cMDS method is the possibility that there can exist infinitely many positive eigenvalues of \mathfrak{K}_X . \mathbb{S}^{d-1} ($d \geq 2$) and \mathbb{T}^N ($N \geq 2$) are actual examples with such property, and they are studied in §6.2, §6.3, and §6.4. To encompass such cases where $\text{pr}(\mathfrak{K}_X) = \infty$, we consider a variant of the cMDS method where the target space is infinite dimensional, namely ℓ^2 . As a result of applying the generalized cMDS method to a given metric measure space X into ℓ^2 , we formally expect to be able to build the following embedding:

$$\begin{aligned} \Phi_X^a : X &\longrightarrow \ell^2 \\ x &\longmapsto \left(\lambda_1^{\frac{1}{2}}\phi_1(x), \lambda_2^{\frac{1}{2}}\phi_2(x), \dots \right) \end{aligned}$$

for a fixed family of orthonormal eigenfunctions $a = \{\phi_i\}_{i=1}^{\text{pr}(\mathfrak{K}_X)}$.

However, in order to guarantee that $\Phi_{\mathcal{X}}^a(x)$ is a point in ℓ^2 , square summability of the sequence $\left(\lambda_1^{\frac{1}{2}}\phi_1(x), \lambda_2^{\frac{1}{2}}\phi_2(x), \dots\right)$ is required. By assuming that the cMDS operator $\mathfrak{K}_{\mathcal{X}}$ is trace class (cf. Definition 2.17), one can guarantee that the sequence $\left(\lambda_1^{\frac{1}{2}}\phi_1(x), \lambda_2^{\frac{1}{2}}\phi_2(x), \dots\right)$ is indeed square summable μ_X -a.e (see Proposition 4.12). Therefore, we will define this version of cMDS which maps into ℓ^2 under the assumption that the operator $\mathfrak{K}_{\mathcal{X}}$ is trace class (see Algorithm 3 and Definition 4.13).

Definition 4.11. The collection of all metric measure spaces $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$ such that $\mathfrak{K}_{\mathcal{X}} : L^2(\mu_X) \rightarrow L^2(\mu_X)$ is trace class is said to be the collection of all **traceable compact metric measure spaces** and is denoted by $\mathcal{M}_w^{\text{tr}}$.

Naturally, establishing canonical examples of metric measure spaces which belong to $\mathcal{M}_w^{\text{tr}}$ will be an important topic of the later part of this paper. However, there exists metric measure spaces which *do not* induce traceable cMDS operators. A construction of one such a metric measure space will be explained in §4.2.

Before we formally introduce a generalized cMDS formulation for metric measure in $\mathcal{M}_w^{\text{tr}}$ which produces embeddings into ℓ^2 , it is worth mentioning that there is an alternative, *coordinate-free* interpretation of generalized cMDS. Even though in this scenario we lose precise coordinate representations, this alternative formulation is well defined even in the case when the cMDS operator is not trace class.

4.1.1. *Coordinate-free interpretation of generalized cMDS.* Fix an arbitrary $\mathcal{X} \in \mathcal{M}_w$. Let $L_{\text{sym}}^2(\mu_X \otimes \mu_X)$ denote the subset of symmetric kernels in $L^2(\mu_X \otimes \mu_X)$. Observe that $K_{\mathcal{X}} \in L_{\text{sym}}^2(\mu_X \otimes \mu_X)$. Also, let $L_{\geq 0, \text{sym}}^2(\mu_X \otimes \mu_X)$ denote the subset of symmetric kernels in $L_{\geq 0}^2(\mu_X \otimes \mu_X)$. Note that $L_{\geq 0, \text{sym}}^2(\mu_X \otimes \mu_X)$ is a convex subset of $L_{\text{sym}}^2(\mu_X \otimes \mu_X)$.

Finally, let P_{μ_X} be the nearest point projection (with respect to the $L^2(\mu_X \otimes \mu_X)$ -norm) from $L_{\text{sym}}^2(\mu_X \otimes \mu_X)$ to the closure of $L_{\geq 0, \text{sym}}^2(\mu_X \otimes \mu_X)$. Let

$$K_{\mathcal{X}}^+ := P_{\mu_X}(K_{\mathcal{X}})$$

be the nearest point projection of the cMDS operator $\mathfrak{K}_{\mathcal{X}}$. Then, the following map

$$K_{\mathcal{X}} \longmapsto K_{\mathcal{X}}^+$$

can be regarded as an alternative, coordinate free, interpretation of cMDS into ℓ^2 .

4.1.2. *Coordinate-dependent interpretation of generalized cMDS into ℓ^2 .* Now we introduce the coordinate-dependent version of cMDS mapping into ℓ^2 that we promised. The following proposition shows that the traceability of $\mathfrak{K}_{\mathcal{X}}$ guarantees that the μ_X -almost everywhere square summability of the sequence $\left(\lambda_1^{\frac{1}{2}}\phi_1(x), \lambda_2^{\frac{1}{2}}\phi_2(x), \dots\right)$.

Proposition 4.12. *Let $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w^{\text{tr}}$. Then,*

$$\sum_{i=1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})} \lambda_i(\phi_i(x))^2 < \infty$$

for μ_X -almost every $x \in X$ where $\lambda_1 \geq \dots \geq \lambda_{\text{pr}(\mathfrak{K}_{\mathcal{X}})} > 0$ are all positive eigenvalues and $\{\phi_i\}_{i=1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})}$ are corresponding orthonormal eigenfunctions of $\mathfrak{K}_{\mathcal{X}}$.

Proof. If $\text{pr}(\mathfrak{K}_X) < \infty$, the claim is obvious. Consider the $\text{pr}(\mathfrak{K}_X) = \infty$ case. We define the following map

$$\begin{aligned} f : X &\longrightarrow [0, \infty] \\ x &\longmapsto \sum_{i=1}^{\infty} \lambda_i (\phi_i(x))^2. \end{aligned}$$

Note that f is measurable since it is the pointwise limit of a sequence of continuous functions (cf. item (6) of Lemma 4.2).

Therefore, by the monotone convergence theorem,

$$\int_X f d\mu_X(x) = \sum_{i=1}^{\infty} \lambda_i \int_X (\phi_i(x))^2 d\mu_X(x) = \sum_{i=1}^{\infty} \lambda_i \leq \text{Tr}(|\mathfrak{K}_X|) < \infty.$$

Hence, $\sum_{i=1}^{\infty} \lambda_i (\phi_i(x))^2 < \infty$ for μ_X -almost every $x \in X$. \square

Hence, for any $\mathcal{X} \in \mathcal{M}_w^{\text{tr}}$, we consider the following type of cMDS embedding where the target space ℓ^2 is infinite dimensional:

Algorithm 3.

- (1) **Input:** $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w^{\text{tr}}$.
- (2) Construct K_X and \mathfrak{K}_X as before.
- (3) Compute the positive eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots > 0$ and corresponding orthonormal eigenfunctions $\{\phi_i\}_{i=1}^{\text{pr}(\mathfrak{K}_X)}$ of \mathfrak{K}_X .
- (4) $\Phi_X(x) := \begin{cases} (\lambda_1^{\frac{1}{2}} \phi_1(x), \lambda_2^{\frac{1}{2}} \phi_2(x), \dots, \lambda_k^{\frac{1}{2}} \phi_{\text{pr}(\mathfrak{K}_X)}(x), 0, 0, \dots) & \text{if } \text{pr}(\mathfrak{K}_X) < \infty \\ (\lambda_1^{\frac{1}{2}} \phi_1(x), \lambda_2^{\frac{1}{2}} \phi_2(x), \dots) & \text{if } \text{pr}(\mathfrak{K}_X) = \infty \end{cases}$
for each $x \in X$. Note that $\Phi_X(x) \in \ell^2$ μ_X -a.e. by Proposition 4.12.
- (5) **Output:** $\Phi_X : X \rightarrow \ell^2$, and $\widehat{\mathcal{X}} := (\Phi_X(X), \|\cdot\|, (\Phi_X)_* \mu_X)$.

Definition 4.13 (generalized cMDS into ℓ^2). Given a metric measure space $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w^{\text{tr}}$ and orthonormal eigenfunctions $a = \{\phi_i\}_{i=1}^{\text{pr}(\mathfrak{K}_X)}$ of \mathfrak{K}_X for the corresponding positive eigenvalues $\lambda_1 \geq \lambda_2 \geq \dots > 0$, one defines the following map:

$$\begin{aligned} \Phi_X^a : X &\longrightarrow \ell^2 \\ x &\longmapsto \begin{cases} (\lambda_1^{\frac{1}{2}} \phi_1(x), \lambda_2^{\frac{1}{2}} \phi_2(x), \dots, \lambda_k^{\frac{1}{2}} \phi_{\text{pr}(\mathfrak{K}_X)}(x), 0, 0, \dots) & \text{if } \text{pr}(\mathfrak{K}_X) < \infty \\ (\lambda_1^{\frac{1}{2}} \phi_1(x), \lambda_2^{\frac{1}{2}} \phi_2(x), \dots) & \text{if } \text{pr}(\mathfrak{K}_X) = \infty \end{cases} \end{aligned}$$

This map Φ_X^a is called **generalized cMDS embedding into ℓ^2** of \mathcal{X} , and the resulting metric measure space is denoted by

$$\widehat{\mathcal{X}} := (\Phi_X^a(X), \|\cdot\|, (\Phi_X^a)_* \mu_X).$$

Unless we have to specify the set of orthonormal eigenfunctions, the superscript a will be omitted, for simplicity.

The following proposition shows that the coordinate-free interpretation agrees with the coordinate-dependent interpretation when the traceability of the cMDS operator is assumed.

Proposition 4.14. *Let $\mathcal{X} \in \mathcal{M}_w^{\text{tr}}$. Then, $K_X^+ = K_{\widehat{\mathcal{X}}}$.*

Proof. Theorem 19 below establishes that $K_{\widehat{\mathcal{X}}}$ is the nearest projection of $K_{\mathcal{X}}$ to $L_{\geq 0, \text{sym}}^2(\mu_X \otimes \mu_X)$. Since there is the unique nearest projection point to a closed convex subset, we can conclude that $K_{\mathcal{X}}^+ = K_{\widehat{\mathcal{X}}}$ in this case. \square

Remark 4.15. Suppose $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$ and positive integer $k \leq \text{pr}(\mathfrak{K}_{\mathcal{X}})$ are given. Then, observe that

$$\|\phi_{\mathcal{X},k}(x) - \phi_{\mathcal{X},k}(x')\|^2 = \sum_{i=1}^k \lambda_i (\phi_i(x) - \phi_i(x'))^2$$

for any $x, x' \in X$ where $\lambda_1 \geq \lambda_2 \geq \dots > 0$ are positive eigenvalues of $\mathfrak{K}_{\mathcal{X}}$, and $\{\phi_i\}_{i=1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})}$ are corresponding orthonormal eigenfunctions. Then, by the definition, it is easy to check

$$K_{\widehat{\mathcal{X}}_k}(x, x') = \sum_{i=1}^k \lambda_i \phi_i(x) \phi_i(x')$$

for any $x, x' \in X$. Similarly, if $\mathcal{X} \in \mathcal{M}_w^{\text{tr}}$, we have

$$\|\phi_{\mathcal{X}}(x) - \phi_{\mathcal{X}}(x')\|^2 = \sum_{i=1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})} \lambda_i (\phi_i(x) - \phi_i(x'))^2,$$

and

$$K_{\widehat{\mathcal{X}}}(x, x') = \sum_{i=1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})} \lambda_i \phi_i(x) \phi_i(x').$$

Remark 4.16. If $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w^{\text{tr}}$, by Theorem 12 and the definition of $K_{\mathcal{X}}$, we have

$$\text{Tr}(\mathfrak{K}_{\mathcal{X}}) = \int_X K_{\mathcal{X}}(x, x) d\mu_X(x) = \frac{1}{2} \iint_{X \times X} d_X^2(u, v) d\mu_X(u) d\mu_X(v) = \frac{1}{2} (\text{diam}_2(\mathcal{X}))^2.$$

This implies that, unless $\text{diam}_2(\mathcal{X}) = 0$, there must be at least one positive eigenvalue of $\mathfrak{K}_{\mathcal{X}}$.

Now, let us collect some simple examples of spaces which belong to $\mathcal{M}_w^{\text{tr}}$. More advanced examples of metric measure space in $\mathcal{M}_w^{\text{tr}}$ can be found in §6.2, §6.3, and §6.4.

Remark 4.17. If $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$ is a finite metric measure space, then \mathcal{X} belongs to $\mathcal{M}_w^{\text{tr}}$ since there are only finitely many eigenvalues.

Remark 4.18. For a compact subset X of \mathbb{R}^k with $m_k(X) > 0$, we consider

$$\mathcal{X} = (X, \|\cdot\|, \overline{m}_k) \in \mathcal{M}_w$$

as in Remark 4.7. Then, there are only finitely many eigenvalues of $\mathfrak{K}_{\mathcal{X}}$ since $\text{rank}(\mathfrak{K}_{\mathcal{X}}) \leq k$. Hence, $\mathcal{X} \in \mathcal{M}_w^{\text{tr}}$.

Example 4.19. Consider the interval $\mathcal{X} = (X, \|\cdot\|, \overline{m}_1) \in \mathcal{M}_w$ where $X := [-1, 1]$ and $\overline{m}_1 := \frac{1}{2}m_1$. Then, by Remark 4.7, $K_{\mathcal{X}}(x, x') = xx'$ for any $x, x' \in X$ since the center of mass of \mathcal{X} is the origin. Then, observe that

$$\begin{aligned} \mathfrak{K}_{\mathcal{X}} \text{Id}_{[-1,1]}(x) &= \int_{[-1,1]} x(x')^2 d\overline{m}_1(x') = \frac{1}{2} \int_{[-1,1]} x(x')^2 dm_1(x') \\ &= \frac{1}{3}x = \frac{1}{3} \text{Id}_{[-1,1]}(x) \end{aligned}$$

for any $x \in X$. Therefore, $\frac{1}{3}$ is the unique nonzero eigenvalue of \mathfrak{K}_X and the identity map is its eigenfunction.

4.2. A metric measure space with non-traceable cMDS operator. We adapt a construction due to Lyons [L⁺18, Lyo13] which arose in the context of the notion of distance covariance between metric spaces.

The construction. For each metric measure space $\mathcal{X} = (X, d_X, \mu_X)$, we define the following operator.

$$\begin{aligned} \mathfrak{T}_{\mathcal{X}} : L^2(\mu_X) &\longrightarrow L^2(\mu_X) \\ \phi &\longmapsto \int_X d_X(\cdot, x') \phi(x') d\mu_X(x'). \end{aligned}$$

Also, for each prime q that is congruent to 1 modulo 4, let $\mathcal{G}_q := (G_q, d_q, \mu_q)$ be the metric measure such that (G_q, d_q) is the **Paley graph** consisting of q vertices with the shortest path metric and μ_q is the uniform probability measure. It is well-known that the Paley graph G_q is strongly regular with parameters $(q, (q-1)/2, (q-5)/4, (q-1)/4)$ meaning that the number of vertices is q , the degree equals $(q-1)/2$ for all vertices, each pair of adjacent vertices has $(q-5)/4$ common neighbors, and each pair of nonadjacent neighbors has $(q-1)/4$ common neighbors [BH11, §9.1.2]. In particular, all graph distances are 0, 1, or 2.

Another well-known fact about G_q is that the adjacency matrix of G_q has eigenvalues $\frac{q-1}{2}$ (with multiplicity 1) and $\frac{-1 \pm \sqrt{q}}{2}$ (with multiplicity $\frac{q-1}{2}$ each) [BH11, Proposition 9.1.1]. Moreover, the eigenvectors of its adjacency matrix are either constant functions or are orthogonal to constant functions. The constant eigenvectors correspond to the eigenvalue $\frac{q-1}{2}$, whereas eigenvectors orthogonal to constant functions correspond to the eigenvalue pair $\frac{-1 \pm \sqrt{q}}{2}$. With these facts and the strong regularity of G_q , one can easily check that (1) the eigenvalues of $\mathfrak{T}_{\mathcal{G}_q}$ are $A_q := \frac{3}{2} \left(1 - \frac{1}{q}\right)$ (with multiplicity 1) and $B_q^\pm := \frac{-3 \pm \sqrt{q}}{2q}$ (each with multiplicity $\frac{q-1}{2}$) and (2) eigenvectors of $\mathfrak{T}_{\mathcal{G}_q}$ are either constant functions or orthogonal to constant functions: Constant eigenvectors correspond to the eigenvalue A_q , and the eigenvectors orthogonal to constant functions correspond to the eigenvalue pair B_q^\pm . At any rate, note that the trace norm (cf. Definition 2.17) $\|\mathfrak{T}_{\mathcal{G}_q}\|_1 = \text{Tr}(|\mathfrak{T}_{\mathcal{G}_q}|)$ is approximately $\frac{\sqrt{q}}{2}$ as $q \rightarrow \infty$.

Now, choose a sequence $(q_n)_n$ of primes $q_n \equiv 1 \pmod{4}$ so that $\sum_n \frac{1}{\sqrt{q_n}} =: \frac{1}{c} < \infty$. We let $\mathcal{X} := (X, d_X, \mu_X)$ be the metric measure space defined as follows:

- $X := \sqcup_n G_{q_n}$,
- $d_X|_{G_{q_n} \times G_{q_n}} = d_{q_n}$ for each n , whereas, for $n \neq m$, $d_X|_{G_{q_n} \times G_{q_m}} = 1$.
- μ_X is the probability measure on X which assigns mass $c q_n^{-3/2}$ to each vertex of \mathcal{G}_{q_n} .

We will prove that $\mathfrak{T}_{\mathcal{X}}$ is not trace class.

A calculation shows that for any eigenvector $v_n \perp \mathbf{1}$ of $\mathfrak{T}_{\mathcal{G}_{q_n}}$ (necessarily) with eigenvalue $\lambda = B_{q_n}^\pm$, the function $f_n : X \rightarrow \mathbb{R}$ defined by

$$f_n(x) := \begin{cases} v_n(x) & \text{if } x \in G_{q_n} \\ 0 & \text{otherwise} \end{cases}$$

is an eigenfunction of $\mathfrak{T}_{\mathcal{X}}$ for the eigenvalue $\frac{\lambda \cdot c}{\sqrt{q_n}}$. Therefore, for each n , an eigenvalue of $\mathfrak{T}_{\mathcal{G}_{q_n}}$ other than A_{q_n} is also an eigenvalue of $\mathfrak{T}_{\mathcal{X}}$ with multiplicity at least $\frac{q_n-1}{2}$. This implies that the trace norm of $\mathfrak{T}_{\mathcal{X}}$ satisfies

$$\|\mathfrak{T}_{\mathcal{X}}\|_1 \geq \sum_n \left(\|\mathfrak{T}_{\mathcal{G}_{q_n}}\|_1 - A_{q_n} \right) \cdot \frac{c}{\sqrt{q_n}}.$$

Hence, $\|\mathfrak{T}_{\mathcal{X}}\|_1$ cannot be finite since

$$\lim_{n \rightarrow \infty} \left(\|\mathfrak{T}_{\mathcal{G}_{q_n}}\|_1 - A_{q_n} \right) \cdot \frac{c}{\sqrt{q_n}} = \lim_{n \rightarrow \infty} \frac{\sqrt{q_n}}{2} \cdot \frac{c}{\sqrt{q_n}} = \frac{c}{2} > 0.$$

Therefore, $\mathfrak{T}_{\mathcal{X}}$ is *not* trace class.

Finally, by Lemma 4.3, if we consider the metric measure space $\mathcal{X}' := (X, d'_X, \mu_X)$ where $d'_X(x, x') := \sqrt{d_X(x, x')}$ for $x, x' \in X$, then $\mathfrak{K}_{\mathcal{X}'}$ also fails to be trace class.

We point out that the space \mathcal{X}' constructed above is not compact. This motivates:

Question 1. *Is it true that any compact metric measure space induces a cMDS operator which is trace class?*

4.3. Distortion incurred by the cMDS operator. We define the following notions in order to measure the metric distortion incurred by the generalized cMDS method.

Definition 4.20. Let \mathcal{H} be a Hilbert space. For each self-adjoint and compact operator $\mathfrak{A} \in L(\mathcal{H})$,

$$\mathrm{Tr}_{\mathrm{neg}}(\mathfrak{A}) := \sum_{i=1}^{\mathrm{nr}(\mathfrak{A})} |\zeta_i|$$

where $\zeta_1 \leq \dots \leq \zeta_{\mathrm{nr}(\mathfrak{A})} < 0$ are the negative eigenvalues of \mathfrak{A} .

Definition 4.21. Let $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w^{\mathrm{tr}}$. For each $k \leq \mathrm{pr}(\mathfrak{K}_{\mathcal{X}})$, we define:

$$\mathrm{dis}_k(\mathcal{X}) := \left(\int_X \int_X \left(\|\Phi_{\mathcal{X},k}(x) - \Phi_{\mathcal{X},k}(x')\| - d_X(x, x') \right)^2 d\mu_X(x') d\mu_X(x) \right)^{\frac{1}{2}},$$

and

$$\mathrm{dis}(\mathcal{X}) := \left(\int_X \int_X \left(\|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\| - d_X(x, x') \right)^2 d\mu_X(x') d\mu_X(x) \right)^{\frac{1}{2}}.$$

Proposition 4.22 (Generalized cMDS bounds). *Let $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w^{\mathrm{tr}}$. Then the following bounds hold:*

(1) (**cMDS is non-contracting**) For $\mu_X \otimes \mu_X$ -almost every $(x, x') \in X \times X$,

$$d_X(x, x') \leq \|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\|$$

and for any integer $k \leq \mathrm{pr}(\mathfrak{K}_{\mathcal{X}})$ and $\mu_X \otimes \mu_X$ -almost every $(x, x') \in X \times X$,

$$\|\Phi_{\mathcal{X},k}(x) - \Phi_{\mathcal{X},k}(x')\| \leq \|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\|.$$

(2) (**L^2 bound**)

$$\mathrm{dis}(\mathcal{X}) \leq 2\sqrt{\mathrm{Tr}_{\mathrm{neg}}(\mathfrak{K}_{\mathcal{X}})}.$$

Remark 4.23. Note that item (1) of Proposition 4.22 implies that

$$d_X(x, x') \leq \|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\| \geq \|\Phi_{\mathcal{X},k}(x) - \Phi_{\mathcal{X},k}(x')\|$$

for any integer $k \leq \text{pr}(\mathfrak{K}_{\mathcal{X}})$ and $\mu_X \otimes \mu_X$ -almost every $(x, x') \in X \times X$. Also, in general it is not true that $d_X(x, x') \leq \|\Phi_{\mathcal{X},k}(x) - \Phi_{\mathcal{X},k}(x')\|$ for $\mu_X \otimes \mu_X$ -almost every $(x, x') \in X \times X$. Indeed, one such example is given by the case when $\mathcal{X} = \mathbb{S}^1$ and $k = 1$ (see §6.2). By Proposition 6.7, we can choose a cMDS embedding of \mathbb{S}^1 into \mathbb{R}^1

$$\begin{aligned} \Phi_{\mathcal{X},1} : \mathbb{S}^1 &\longrightarrow \mathbb{R} \\ \theta &\longmapsto \sqrt{2} \cos \theta. \end{aligned}$$

Then, $d_{\mathbb{S}^1}(0, \pi) = \pi > 2\sqrt{2} = |\Phi_{\mathcal{X},1}(0) - \Phi_{\mathcal{X},1}(\pi)|$.

Proof of Proposition 4.22. (1) Let $\lambda_1 \geq \dots \geq \lambda_{\text{pr}(\mathfrak{K}_{\mathcal{X}})} > 0$ be all positive eigenvalues and $\{\phi_i\}_i$ be the corresponding orthonormal eigenfunctions of $\mathfrak{K}_{\mathcal{X}}$. Also, let $\zeta_1 \leq \dots \leq \zeta_{\text{nr}(\mathfrak{K}_{\mathcal{X}})} < 0$ be all negative eigenvalues and $\{\psi_i\}_i$ be the corresponding orthonormal eigenfunctions of $\mathfrak{K}_{\mathcal{X}}$.

Since $\mathfrak{K}_{\mathcal{X}}$ is trace class, we have that

$$K_{\widehat{\mathcal{X}}}(x, x') = \sum_{i=1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})} \lambda_i \phi_i(x) \phi_i(x'),$$

and

$$K_{\mathcal{X}}^-(x, x') := \sum_{i=1}^{\text{nr}(\mathfrak{K}_{\mathcal{X}})} (-\zeta_i) \psi_i(x) \psi_i(x')$$

are well-defined for $\mu_X \otimes \mu_X$ -almost every $(x, x') \in X \times X$. Moreover, because both kernels $K_{\widehat{\mathcal{X}}} - K_{\mathcal{X}}$ and $K_{\mathcal{X}}^-$ generate the same operator, by Corollary 2.22 we have

$$K_{\widehat{\mathcal{X}}}(x, x') - K_{\mathcal{X}}(x, x') = K_{\mathcal{X}}^-(x, x') = \sum_{i=1}^{\text{nr}(\mathfrak{K}_{\mathcal{X}})} (-\zeta_i) \psi_i(x) \psi_i(x')$$

for $\mu_X \otimes \mu_X$ -almost every (x, x') . Therefore,

$$\begin{aligned} &\|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\|^2 - d_X^2(x, x') \\ &= (K_{\widehat{\mathcal{X}}}(x, x) - K_{\mathcal{X}}(x, x)) + (K_{\widehat{\mathcal{X}}}(x', x') - K_{\mathcal{X}}(x', x')) - 2(K_{\widehat{\mathcal{X}}}(x, x') - K_{\mathcal{X}}(x, x')) \\ &= \sum_{i=1}^{\text{nr}(\mathfrak{K}_{\mathcal{X}})} (-\zeta_i) (\psi_i(x) - \psi_i(x'))^2 \geq 0 \end{aligned}$$

Therefore, we have $\|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\| \geq d_X(x, x')$ for $\mu_X \otimes \mu_X$ -almost every (x, x') . Also,

$$\|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\|^2 - \|\Phi_{\mathcal{X},k}(x) - \Phi_{\mathcal{X},k}(x')\|^2 = \sum_{i=k+1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})} \lambda_i ((\phi_i(x) - \phi_i(x'))^2) \geq 0$$

for $\mu_X \otimes \mu_X$ -almost every (x, x') . Hence, we have $\|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\| \geq \|\Phi_{\mathcal{X},k}(x) - \Phi_{\mathcal{X},k}(x')\|$ for $\mu_X \otimes \mu_X$ -almost every (x, x') .

- (2) Since we already know $\|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\| \geq d_X(x, x')$ for $\mu_X \otimes \mu_X$ -almost every (x, x') , it is easy to observe the following:

$$(\|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\| - d_X(x, x'))^2 \leq \|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\|^2 - d_X^2(x, x')$$

for $\mu_X \otimes \mu_X$ -almost every (x, x') . Hence,

$$\begin{aligned} & \int_X \int_X (\|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\| - d_X(x, x'))^2 d\mu_X(x') d\mu_X(x) \\ & \leq \int_X \int_X \|\Phi_{\mathcal{X}}(x) - \Phi_{\mathcal{X}}(x')\|^2 - d_X^2(x, x') d\mu_X(x') d\mu_X(x) \\ & = \int_X \int_X \sum_{i=1}^{\text{nr}(\mathfrak{K}_{\mathcal{X}})} (-\zeta_i) ((\psi_i(x) - \psi_i(x'))^2) d\mu_X(x') d\mu_X(x) \\ & \leq 2 \int_X \int_X \sum_{i=1}^{\text{nr}(\mathfrak{K}_{\mathcal{X}})} (-\zeta_i) (\psi_i(x)^2 + \psi_i(x')^2) d\mu_X(x') d\mu_X(x) \\ & = 4 \sum_{i=1}^{\text{nr}(\mathfrak{K}_{\mathcal{X}})} (-\zeta_i) = 4 \text{Tr}_{\text{neg}}(\mathfrak{K}_{\mathcal{X}}). \end{aligned}$$

By applying the square root, we have the required upper bound of $\text{dis}(\mathcal{X})$. \square

4.4. Optimality of the generalized cMDS. We are able to prove the following optimality result for generalized cMDS in a manner is analogous to Theorem 16 (which is only applicable to finite metric spaces).

Theorem 19. *Suppose $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w^{\text{tr}}$. Then,*

- (1) $\|K_{\mathcal{X}} - K\|_{L^2(\mu_X \otimes \mu_X)} \geq \|K_{\mathcal{X}} - K_{\widehat{\mathfrak{K}}_k}\|_{L^2(\mu_X \otimes \mu_X)}$ for any positive integer $k \leq \text{pr}(\mathfrak{K}_{\mathcal{X}})$, and $K \in L_{\geq 0, \text{sym}}^2(\mu_X \otimes \mu_X)$ with $\text{rank}(\mathfrak{K}) \leq k$ where \mathfrak{K} is the positive semi-definite operator induced by K .
- (2) $\|K_{\mathcal{X}} - K\|_{L^2(\mu_X \otimes \mu_X)} \geq \|K_{\mathcal{X}} - K_{\widehat{\mathfrak{K}}}\|_{L^2(\mu_X \otimes \mu_X)}$ for any $K \in L_{\geq 0, \text{sym}}^2(\mu_X \otimes \mu_X)$.

Definition 4.24 (doubly substochastic matrix). A square matrix $A = (a_{ij})_{i,j=1}^n$ of size n is called **doubly substochastic** if it satisfies:

- (1) $a_{ij} \geq 0$ for any $i, j \in \{1, \dots, n\}$.
- (2) $\sum_{j=1}^n a_{ij} \leq 1$ for any $i \in \{1, \dots, n\}$.
- (3) $\sum_{i=1}^n a_{ij} \leq 1$ for any $j \in \{1, \dots, n\}$.

Theorem 20 ([HHJ94, Theorem 3.2.6]). *A square matrix $A = (a_{ij})_{i,j=1}^n$ of size n is doubly substochastic matrix if and only if there is a doubly stochastic matrix $B = (b_{ij})_{i,j=1}^n$ of size n such that $0 \leq a_{ij} \leq b_{ij}$ for any $i, j = 1, \dots, n$.*

Lemma 4.25. *Let \mathcal{H} be a separable Hilbert space, $\mathfrak{A} \in \mathcal{L}(\mathcal{H})$ be Hilbert-Schmidt and self-adjoint, and $\mathfrak{B} \in \mathcal{L}(\mathcal{H})$ be Hilbert-Schmidt, self-adjoint, and positive semi-definite. Then, we have*

$$\text{Tr}(\mathfrak{A}\mathfrak{B}) \leq \sum_{i=1}^M \lambda_i \lambda'_i$$

where $M := \min\{\text{rank}(\mathfrak{B}), \text{pr}(\mathfrak{A})\}$, $\lambda_1 \geq \lambda_2 \geq \dots > 0$ are the positive eigenvalues of \mathfrak{A} , and $\lambda'_1 \geq \lambda'_2 \geq \dots > 0$ are the positive eigenvalues of \mathfrak{B} .

Proof. Let $\{\lambda_i\}_{i=1}^{\text{pr}(\mathfrak{A})}$ be the positive eigenvalues with corresponding orthonormal eigenfunctions $\{\phi_i\}_{i=1}^{\text{pr}(\mathfrak{A})}$ of \mathfrak{A} , $\{\zeta_i\}_{i=1}^{\text{nr}(\mathfrak{A})}$ be the negative eigenvalues with corresponding orthonormal eigenfunctions $\{\psi_i\}_{i=1}^{\text{nr}(\mathfrak{A})}$ of \mathfrak{A} , and $\{\lambda'_i\}_{i=1}^{\text{rank}(\mathfrak{B})}$ be the positive eigenvalues with corresponding orthonormal eigenfunctions $\{\phi'_i\}_{i=1}^{\text{rank}(\mathfrak{B})}$ of \mathfrak{B} . Then,

$$\begin{aligned} \text{Tr}(\mathfrak{A}\mathfrak{B}) &= \sum_{j=1}^{\text{rank}(\mathfrak{B})} \langle \phi'_j, \mathfrak{A}\mathfrak{B}\phi'_j \rangle = \sum_{j=1}^{\text{rank}(\mathfrak{B})} \lambda'_j \langle \phi'_j, \mathfrak{A}\phi'_j \rangle \\ &= \sum_{j=1}^{\text{rank}(\mathfrak{B})} \lambda'_j \left\langle \phi'_j, \sum_{i=1}^{\text{pr}(\mathfrak{A})} \lambda_i \langle \phi_i, \phi'_j \rangle \phi_i + \sum_{i=1}^{\text{nr}(\mathfrak{A})} \zeta_i \langle \psi_i, \phi'_j \rangle \psi_i \right\rangle \\ &= \sum_{j=1}^{\text{rank}(\mathfrak{B})} \sum_{i=1}^{\text{pr}(\mathfrak{A})} \lambda'_j \lambda_i \langle \phi'_j, \phi_i \rangle^2 + \sum_{j=1}^{\text{rank}(\mathfrak{B})} \sum_{i=1}^{\text{nr}(\mathfrak{A})} \lambda'_j \zeta_i \langle \phi'_j, \psi_i \rangle^2 \\ &\leq \sum_{j=1}^{\text{rank}(\mathfrak{B})} \sum_{i=1}^{\text{pr}(\mathfrak{A})} \lambda'_j \lambda_i \langle \phi'_j, \phi_i \rangle^2 \end{aligned}$$

First, let's consider $M = \text{rank}(\mathfrak{B})$ case. Observe that $\sum_{j=1}^{\text{rank}(\mathfrak{B})} \langle \phi'_j, \phi_i \rangle^2 \leq 1$ for each $i = 1, \dots, \text{pr}(\mathfrak{A})$, and $\sum_{i=1}^{\text{pr}(\mathfrak{A})} \langle \phi'_j, \phi_i \rangle^2 \leq 1$ for each $j = 1, \dots, \text{rank}(\mathfrak{B})$ by the orthonormal property. Now, let's do case-by-case analysis.

(1) If $\text{pr}(\mathfrak{A}) < \infty$: Observe that, if we define $A = (a_{ji})_{j,i=1}^{\text{pr}(\mathfrak{A})}$ where

$$a_{ji} = \begin{cases} \langle \phi'_j, \phi_i \rangle^2 & \text{if } j = 1, \dots, \text{rank}(\mathfrak{B}) \\ 0 & \text{if } j = \text{rank}(\mathfrak{B}) + 1, \dots, \text{pr}(\mathfrak{A}), \end{cases}$$

then A is substochastic matrix. Then, by Theorem 20, A is (entrywisely) upper bounded by some doubly stochastic matrix $B = (b_{ji})_{j,i=1}^{\text{pr}(\mathfrak{A})}$. Hence,

$$\begin{aligned} \sum_{j=1}^{\text{rank}(\mathfrak{B})} \sum_{i=1}^{\text{pr}(\mathfrak{A})} \lambda'_j \lambda_i \langle \phi'_j, \phi_i \rangle^2 &= \sum_{j=1}^{\text{rank}(\mathfrak{B})} \lambda_j \left(\sum_{i=1}^{\text{pr}(\mathfrak{A})} a_{ji} \lambda_i \right) \\ &\leq \sum_{j=1}^{\text{rank}(\mathfrak{B})} \lambda_j \left(\sum_{i=1}^{\text{pr}(\mathfrak{A})} b_{ji} \lambda_i \right) \\ &\leq \sum_{i=1}^{\text{rank}(\mathfrak{B})} \lambda_i \lambda'_i \end{aligned}$$

as we wanted, where the last inequality holds by Theorem 17 and Theorem 18.

(2) If $\text{pr}(\mathfrak{A}) = \infty$:

- (a) If $\text{rank}(\mathfrak{B}) < \infty$: Choose arbitrary positive integer $l \geq k$. Then, by the similar argument of the previous case (1), we have

$$\sum_{j=1}^{\text{rank}(\mathfrak{B})} \sum_{i=1}^l \lambda'_j \lambda_i \langle \phi'_j, \phi_i \rangle^2 \leq \sum_{i=1}^{\text{rank}(\mathfrak{B})} \lambda_i \lambda'_i.$$

Since l is arbitrary,

$$\sum_{j=1}^{\text{rank}(\mathfrak{B})} \sum_{i=1}^{\infty} \lambda'_j \lambda_i \langle \phi'_j, \phi_i \rangle^2 \leq \sum_{i=1}^{\text{rank}(\mathfrak{B})} \lambda_i \lambda'_i$$

as we wanted.

- (b) If $\text{rank}(\mathfrak{B}) = \infty$: Choose arbitrary positive integers $l \leq l'$. Then, by the similar argument of the previous case (1), we have

$$\sum_{j=1}^l \sum_{i=1}^{l'} \lambda'_j \lambda_i \langle \phi'_j, \phi_i \rangle^2 \leq \sum_{i=1}^l \lambda_i \lambda'_i.$$

Since l, l' are arbitrary, we have

$$\sum_{j=1}^{\infty} \sum_{i=1}^{\infty} \lambda'_j \lambda_i \langle \phi'_j, \phi_i \rangle^2 \leq \sum_{i=1}^{\infty} \lambda_i \lambda'_i$$

as we wanted.

The case when $M = \text{pr}(\mathfrak{A})$ is proved in a similar manner, which we omit. \square

Proof of Theorem 19. Let's prove the first claim.

Let $\{\lambda_i\}_{i=1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})}$ be the positive eigenvalues with corresponding orthonormal eigenfunctions $\{\phi_i\}_{i=1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})}$ of $\mathfrak{K}_{\mathcal{X}}$, $\{\zeta_i\}_{i=1}^{\text{nr}(\mathfrak{K}_{\mathcal{X}})}$ be the negative eigenvalues with corresponding orthonormal eigenfunctions $\{\psi_i\}_{i=1}^{\text{nr}(\mathfrak{K}_{\mathcal{X}})}$ of $\mathfrak{K}_{\mathcal{X}}$, and $\{\lambda'_i\}_{i=1}^{\text{rank}(\mathfrak{K})}$ be the positive eigenvalues with corresponding orthonormal eigenfunctions $\{\phi'_i\}_{i=1}^{\text{rank}(\mathfrak{K})}$ of \mathfrak{K} . Note that $\text{rank}(\mathfrak{K}) \leq k \leq \text{pr}(\mathfrak{K}_{\mathcal{X}})$. Then,

$$\begin{aligned} \|K_{\mathcal{X}} - K\|_{L^2(\mu_{\mathcal{X}} \otimes \mu_{\mathcal{X}})}^2 &= \|\mathfrak{K}_{\mathcal{X}} - \mathfrak{K}\|_2^2 = \text{Tr}((\mathfrak{K}_{\mathcal{X}} - \mathfrak{K})^*(\mathfrak{K}_{\mathcal{X}} - \mathfrak{K})) \\ &= \text{Tr}((\mathfrak{K}_{\mathcal{X}} - \mathfrak{K})^2) = \text{Tr}(\mathfrak{K}_{\mathcal{X}}^2) + \text{Tr}(\mathfrak{K}^2) - 2 \text{Tr}(\mathfrak{K}_{\mathcal{X}} \mathfrak{K}) \quad (\because \text{item (4) of Lemma 2.20}) \\ &\geq \sum_{i=1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})} \lambda_i^2 + \sum_{i=1}^{\text{nr}(\mathfrak{K}_{\mathcal{X}})} \zeta_i^2 + \sum_{j=1}^{\text{rank}(\mathfrak{K})} (\lambda'_j)^2 - 2 \sum_{i=1}^{\text{rank}(\mathfrak{K})} \lambda_i \lambda'_i \quad (\because \text{Lemma 4.25}) \\ &= \sum_{i=1}^{\text{rank}(\mathfrak{K})} (\lambda_i - \lambda'_i)^2 + \sum_{i=\text{rank}(\mathfrak{K})+1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})} \lambda_i^2 + \sum_{i=1}^{\text{nr}(\mathfrak{K}_{\mathcal{X}})} \zeta_i^2 \\ &\geq \sum_{i=\text{rank}(\mathfrak{K})+1}^{\text{pr}(\mathfrak{K}_{\mathcal{X}})} \lambda_i^2 + \sum_{i=1}^{\text{nr}(\mathfrak{K}_{\mathcal{X}})} \zeta_i^2 \\ &= \|K_{\mathcal{X}} - K_{\hat{\mathcal{X}}_k}\|_{L^2(\mu_{\mathcal{X}} \otimes \mu_{\mathcal{X}})}^2. \end{aligned}$$

So, this concludes the first claim. The second claim is proved by the similar argument. \square

5. THICKNESS AND SPECTRUM OF $\mathfrak{K}_{\mathcal{X}}$ FOR EUCLIDEAN DATA

In this section, we prove that if \mathcal{X} is a compact subset of Euclidean space, then the smallest positive eigenvalue of $\mathfrak{K}_{\mathcal{X}}$ is upper bounded by a certain notion of *thickness* of \mathcal{X} (Theorem 21). This result exemplifies the phenomenon that the spectrum of $\mathfrak{K}_{\mathcal{X}}$ is controlled by the underlying geometry of \mathcal{X} .

We begin with an example.

Example 5.1. For $a \geq b > 0$ consider the following ellipse in \mathbb{R}^2 : $\mathcal{X} := (X, \|\cdot\|, \overline{m}_2)$ where $\overline{m}_2 = \frac{1}{ab\pi}m_2$ is the normalized Lebesgue measure and

$$X := \left\{ x = (x_1, x_2) \in \mathbb{R}^2 : \left(\frac{x_1}{a}\right)^2 + \left(\frac{x_2}{b}\right)^2 \leq 1 \right\}.$$

Then, by Remark 4.7, $K_{\mathcal{X}}(x, x') = \langle x, x' \rangle$ for any $x, x' \in X$, $\mathfrak{K}_{\mathcal{X}}$ is positive semi-definite, and the rank of $\mathfrak{K}_{\mathcal{X}}$ is at most 2. Let $\lambda_1 \geq \lambda_2 \geq 0$ be the two largest eigenvalues of $\mathfrak{K}_{\mathcal{X}}$. We will show that $\pi_1 := \langle \cdot, e_1 \rangle$ and $\pi_2 := \langle \cdot, e_2 \rangle$ are their corresponding eigenfunctions. First, consider π_1 :

$$\begin{aligned} \int_X (\pi_1(x))^2 d\overline{m}_2(x) &= \frac{1}{ab\pi} \int_X x_1^2 dm_2(x) \\ &= \frac{1}{ab\pi} \int_0^{2\pi} \int_0^1 (ar \cos \theta)^2 ab r dr d\theta \quad (\because \text{use polar coordinates}) \\ &= \frac{a^2}{4}. \end{aligned}$$

Hence, $\|\pi_1\| = \frac{a}{2}$. Also,

$$\begin{aligned} \mathfrak{K}_{\mathcal{X}}\pi_1(x) &= \int_X \langle x, x' \rangle \pi_1(x') d\overline{m}_2(x') \\ &= \frac{1}{ab\pi} \int_X x_1(x_1')^2 dm_2(x') + \frac{1}{ab\pi} \int_X x_2 x_1' x_2' dm_2(x') \\ &= \frac{x_1}{ab\pi} \int_0^{2\pi} \int_0^1 (ar \cos \theta)^2 ab r dr d\theta + \frac{x_2}{ab\pi} \int_0^{2\pi} \int_0^1 a^2 b^2 r^3 \cos \theta \sin \theta dr d\theta \\ &= \frac{a^2 x_1}{4} = \frac{a^2}{4} \pi_1(x) \end{aligned}$$

for any $x \in X$. In a similar way, we are able to show that the norm of $\pi_2 = \langle \cdot, e_2 \rangle$ is $\frac{b}{2}$ and $\mathfrak{K}_{\mathcal{X}}\pi_2 = \frac{b^2}{4}\pi_2$. Therefore $\frac{a^2}{4} \geq \frac{b^2}{4} > 0$ are all positive eigenvalues, and $\frac{2\pi_1}{a}, \frac{2\pi_2}{b}$ are the corresponding orthonormal eigenfunctions.

With the aid of the above calculations, it is easy to check $\Phi_{\mathcal{X},1}(x) = \frac{a}{2} \cdot \frac{2}{a}\pi_1(x) = x_1$ for any $x = (x_1, x_2) \in X$. Hence, if we only retain the 1st coordinate produced by cMDS, we

have the distortion:

$$\begin{aligned}
 \text{dis}_1(\mathcal{X}) &= \left(\int_X \int_X (\|\Phi_{\mathcal{X},1}(x) - \Phi_{\mathcal{X},1}(x')\| - \|x - x'\|)^2 d\bar{m}_2(x') d\bar{m}_2(x) \right)^{\frac{1}{2}} \\
 &= \left(\int_X \int_X (|x_1 - x'_1| - \sqrt{(x_1 - x'_1)^2 + (x_2 - x'_2)^2})^2 d\bar{m}_2(x') d\bar{m}_2(x) \right)^{\frac{1}{2}} \\
 &\leq \left(\int_X \int_X ((x_1 - x'_1)^2 + (x_2 - x'_2)^2 - (x_1 - x'_1)^2) d\bar{m}_2(x') d\bar{m}_2(x) \right)^{\frac{1}{2}} \\
 &= \left(\int_X \int_X (x_2 - x'_2)^2 d\bar{m}_2(x') d\bar{m}_2(x) \right)^{\frac{1}{2}} \\
 &= \left(2 \int_X x_2^2 d\bar{m}_2(x) - 2 \left(\int_X x_2 d\bar{m}_2(x) \right)^2 \right)^{\frac{1}{2}} \\
 &= \sqrt{\frac{b^2}{2}} = \frac{b}{\sqrt{2}}.
 \end{aligned}$$

The bound $\text{dis}_1(\mathcal{X}) \leq \frac{b}{\sqrt{2}}$ is consistent with the fact that, for thin ellipses ($a \gg b$), the cMDS method with 1 coordinate causes small distortion.

We now introduce a notion of thickness which is applicable in general.

Definition 5.2. For a given metric measure space $\mathcal{X} = (X, \|\cdot\|, \bar{m}_k) \in \mathcal{M}_w$ where X is a compact subset of \mathbb{R}^k with $m_k(X) > 0$ and $\bar{m}_k := \frac{1}{m_k(X)} m_k$, the *thickness* of \mathcal{X} is the number

$$\text{Th}(\mathcal{X}) := \inf_{u \in \mathbb{S}^{k-1}} \left(\int_X (\pi_u(x - \text{cm}(\mathcal{X})))^2 d\bar{m}_k(x) \right)^{\frac{1}{2}}.$$

Example 5.3. For the ellipse considered above, its thickness is $\frac{b}{2}$.

Theorem 21. Suppose $\mathcal{X} = (X, \|\cdot\|, \bar{m}_k) \in \mathcal{M}_w$ is given, where X is a compact subset of \mathbb{R}^k with $m_k(X) > 0$ and $\bar{m}_k := \frac{1}{m_k(X)} m_k$. Then, the k -th eigenvalue λ_k of $K_{\mathcal{X}}$ satisfies the following inequality:

$$\lambda_k \leq (\text{Th}(\mathcal{X}))^2.$$

Proof. Fix any $u \in \mathbb{S}^{k-1}$. Without loss of generality, we may assume $\text{cm}(\mathcal{X}) = 0$. Choose an orthonormal basis $v_1, \dots, v_{k-1}, v_k = u$ of \mathbb{R}^k . Then, consider the following map

$$\begin{aligned}
 \pi_{u^\perp} : \mathbb{R}^k &\longrightarrow \mathbb{R}^{k-1} \\
 x &\longmapsto (\pi_{v_1}(x), \dots, \pi_{v_{k-1}}(x)),
 \end{aligned}$$

and

$$\begin{aligned}
 K_{u^\perp} : X \times X &\longrightarrow \mathbb{R} \\
 (x, x') &\longmapsto \langle \pi_{u^\perp}(x), \pi_{u^\perp}(x') \rangle.
 \end{aligned}$$

Then, by Theorem 19, we have the following inequality:

$$\|K_{\mathcal{X}} - K_{u^\perp}\|_{L^2(\bar{m}_k \otimes \bar{m}_k)} \geq \|K_{\mathcal{X}} - K_{\hat{\mathcal{X}}_{k-1}}\|_{L^2(\bar{m}_k \otimes \bar{m}_k)}.$$

Observe that it is easy to show $\|K_{\mathcal{X}} - K_{\hat{\mathcal{X}}_{k-1}}\|_{L^2(\bar{m}_k \otimes \bar{m}_k)} = \lambda_k$. Also, by Remark 4.7, we have $K_{\mathcal{X}}(x, x') = \langle x, x' \rangle$ for any $x, x' \in X$. Therefore,

$$\begin{aligned} K_{\mathcal{X}}(x, x') - K_{u^\perp}(x, x') &= \sum_{i=1}^k \pi_{v_i}(x) \pi_{v_i}(x') - \sum_{i=1}^{k-1} \pi_{v_i}(x) \pi_{v_i}(x') \\ &= \pi_{v_k}(x) \pi_{v_k}(x') = \pi_u(x) \pi_u(x') \end{aligned}$$

for any $x, x' \in X$. Hence,

$$\begin{aligned} \|K_{\mathcal{X}} - K_{u^\perp}\|_{L^2(\bar{m}_k \otimes \bar{m}_k)}^2 &= \int_X \int_X (\pi_u(x) \pi_u(x'))^2 d\bar{m}_k(x') d\bar{m}_k(x) \\ &= \left(\int_X (\pi_u(x))^2 d\bar{m}_k(x) \right)^2. \end{aligned}$$

Hence, we have $\|K_{\mathcal{X}} - K_{u^\perp}\|_{L^2(\bar{m}_k \otimes \bar{m}_k)} = \int_X (\pi_u(x))^2 d\bar{m}_k(x)$. Therefore,

$$\lambda_k \leq \int_X (\pi_u(x))^2 d\bar{m}_k(x).$$

The conclusion follows. \square

6. TRACEABILITY OF cMDS ON EXEMPLAR METRIC MEASURES

In this section, we study the traceability of the generalized cMDS procedure when applied to exemplar metric measures such as spheres, tori, and metric graphs.

6.1. Generalized cMDS for product spaces. Let $\mathcal{X} = (X, d_X, \mu_X)$ and $\mathcal{Y} = (Y, d_Y, \mu_Y)$ be two metric measure spaces in \mathcal{M}_w . Now, we consider the following product metric measure space:

$$\mathcal{X} \times \mathcal{Y} := \left(X \times Y, d_{X \times Y} = \sqrt{d_X^2 + d_Y^2}, \mu_X \otimes \mu_Y \right) \in \mathcal{M}_w.$$

Lemma 6.1. *Two metric measure spaces $\mathcal{X} = (X, d_X, \mu_X)$ and $\mathcal{Y} = (Y, d_Y, \mu_Y)$ are given. Then, $K_{\mathcal{X} \times \mathcal{Y}}((x, y), (x', y')) = K_{\mathcal{X}}(x, x') + K_{\mathcal{Y}}(y, y')$ for any $x, x' \in X$ and $y, y' \in Y$.*

Proof. Fix arbitrary $x, x' \in X$ and $y, y' \in Y$. Then,

$$\begin{aligned} &K_{\mathcal{X} \times \mathcal{Y}}((x, y), (x', y')) \\ &= -\frac{1}{2} d_{X \times Y}^2((x, y), (x', y')) + \frac{1}{2} \int_{X \times Y} d_{X \times Y}^2((x, y), (u, v)) d\mu_X(u) d\mu_Y(v) \\ &\quad + \frac{1}{2} \int_X d_{X \times Y}^2((s, t), (x', y')) d\mu_X(s) d\mu_Y(t) \\ &\quad - \frac{1}{2} \int_{X \times Y} \int_{X \times Y} d_{X \times Y}^2((s, t), (u, v)) d\mu_X(s) d\mu_Y(t) d\mu_X(u) d\mu_X(v) \\ &= -\frac{1}{2} \left(d_X^2(x, x') - \int_X d_X^2(x, u) d\mu_X(u) - \int_X d_X^2(s, x') d\mu_X(s) + \iint_{X \times X} d_X^2(s, u) d\mu_X(s) d\mu_X(u) \right) \\ &\quad - \frac{1}{2} \left(d_Y^2(y, y') - \int_Y d_Y^2(y, v) d\mu_Y(v) - \int_Y d_Y^2(t, y') d\mu_Y(t) + \iint_{Y \times Y} d_Y^2(t, v) d\mu_Y(t) d\mu_Y(v) \right) \\ &= K_{\mathcal{X}}(x, x') + K_{\mathcal{Y}}(y, y'). \end{aligned}$$

This concludes the proof. \square

Remark 6.2. Let $\mathcal{X}_i = (X_i, d_i, \mu_i) \in \mathcal{M}_w$ for each $i = 1, \dots, N$, and $\mathcal{X} = \mathcal{X}_1 \times \dots \times \mathcal{X}_N$ be the product metric measure space. Then, by using the induction on N and Lemma 6.1, we have the following:

$$K_{\mathcal{X}}((x_1, \dots, x_n), (x'_1, \dots, x'_n)) = \sum_{i=1}^N K_{\mathcal{X}_i}(x_i, x'_i)$$

for any $(x_1, \dots, x_n), (x'_1, \dots, x'_n) \in \mathcal{X}$.

The following proposition clarifies what the spectrum of $\mathfrak{K}_{\mathcal{X} \times \mathcal{Y}}$ is for given $\mathcal{X}, \mathcal{Y} \in \mathcal{M}_w$.

Proposition 6.3. Two metric measure spaces $\mathcal{X} = (X, d_X, \mu_X), \mathcal{Y} = (Y, d_Y, \mu_Y) \in \mathcal{M}_w$ are given. Then,

- (1) Let ϕ be an eigenfunction of $\mathfrak{K}_{\mathcal{X}}$ for some nonzero eigenvalue, and ψ be an eigenfunction of $\mathfrak{K}_{\mathcal{Y}}$ for some nonzero eigenvalue. Then, $\phi \times \psi$ becomes an eigenfunction of $\mathfrak{K}_{\mathcal{X} \times \mathcal{Y}}$ for zero eigenvalue.
- (2) Let ϕ be an eigenfunction of $\mathfrak{K}_{\mathcal{X}}$ for some nonzero eigenvalue λ , and ψ be an eigenfunction of $\mathfrak{K}_{\mathcal{Y}}$ for zero eigenvalue. Then, $\phi \times \psi$ becomes an eigenfunction of $\mathfrak{K}_{\mathcal{X} \times \mathcal{Y}}$ for the eigenvalue $\lambda \cdot \int_Y \psi d\mu_Y$.
- (3) Let ϕ be an eigenfunction of $\mathfrak{K}_{\mathcal{X}}$ for zero eigenvalue, and ψ be an eigenfunction of $\mathfrak{K}_{\mathcal{Y}}$ for zero eigenvalue. Then, $\phi \times \psi$ becomes an eigenfunction of $\mathfrak{K}_{\mathcal{X} \times \mathcal{Y}}$ for zero eigenvalue.

Proof. By Lemma 6.1,

$$\begin{aligned} & \int_{X \times Y} K_{\mathcal{X} \times \mathcal{Y}}((x, y), (x', y')) \phi(x') \psi(y') d\mu_X(x') d\mu_Y(y') \\ &= \int_Y \int_X K_{\mathcal{X}}(x, x') \phi(x') \psi(y') d\mu_X(x') d\mu_Y(y') + \int_Y \int_X K_{\mathcal{Y}}(y, y') \phi(x') \psi(y') d\mu_X(x') d\mu_Y(y') \\ &= \int_X K_{\mathcal{X}}(x, x') \phi(x') d\mu_X(x') \cdot \int_Y \psi(y') d\mu_Y(y') + \int_Y K_{\mathcal{Y}}(y, y') \psi(y') d\mu_Y(y') \cdot \int_X \phi(x') d\mu_X(x') \end{aligned}$$

for any $(x, y) \in X \times Y$. Hence,

- (1) If ϕ be an eigenfunction of $\mathfrak{K}_{\mathcal{X}}$ for some nonzero eigenvalue, and ψ be an eigenfunction of $\mathfrak{K}_{\mathcal{Y}}$ for some nonzero eigenvalue: Then,

$$\int_{X \times Y} K_{\mathcal{X} \times \mathcal{Y}}((x, y), (x', y')) \phi(x') \psi(y') d\mu_X(x') d\mu_Y(y') = 0$$

since $\int_Y \psi_j(y') d\mu_Y(y') = \int_X \phi_i(x') d\mu_X(x') = 0$ by item (5) of Lemma 4.2. Hence, $\phi \times \psi$ is an eigenfunction for zero eigenvalue.

- (2) If ϕ be an eigenfunction of $\mathfrak{K}_{\mathcal{X}}$ for some nonzero eigenvalue λ , and ψ be an eigenfunction of $\mathfrak{K}_{\mathcal{Y}}$ for zero eigenvalue: Then,

$$\int_{X \times Y} K_{\mathcal{X} \times \mathcal{Y}}((x, y), (x', y')) \phi(x') \psi(y') d\mu_X(x') d\mu_Y(y') = \left(\lambda \cdot \int_Y \psi d\mu_Y \right) \phi(x)$$

since $\int_X \phi_i(x') d\mu_X(x') = 0$ by item (5) of Lemma 4.2. Hence, $\phi \times \psi$ is an eigenfunction for $\lambda \cdot \int_Y \psi d\mu_Y$.

- (3) If ϕ be an eigenfunction of $\mathfrak{K}_{\mathcal{X}}$ for zero eigenvalue, and ψ be an eigenfunction of $\mathfrak{K}_{\mathcal{Y}}$ for zero eigenvalue: Then,

$$\int_{X \times Y} K_{\mathcal{X} \times \mathcal{Y}}((x, y), (x', y')) \phi(x') \psi(y') d\mu_X(x') d\mu_Y(y') = 0$$

by the assumption on ϕ and ψ . Hence, $\phi \times \psi$ is an eigenfunction for zero eigenvalue. \square

Proposition 6.4. *Suppose $\mathcal{X}_i = (X_i, d_i, \mu_i) \in \mathcal{M}_w$ for $i = 1, \dots, N$ are given. Let $\mathcal{X} = \mathcal{X}_1 \times \dots \times \mathcal{X}_N$ be the product metric measure space. Then, if λ_i is an eigenvalue of $\mathfrak{K}_{\mathcal{X}_i}$ with an eigenfunction ϕ_i for each $i = 1, \dots, N$, the $\phi_1 \times \dots \times \phi_N$ is an eigenfunction of $\mathfrak{K}_{\mathcal{X}}$ with eigenvalue:*

$$\begin{cases} 0 & \text{if there are at least two of nonzero } \lambda_i \text{'s.} \\ \lambda_i \cdot \left(\prod_{j \neq i} \int_{X_j} \phi_j d\mu_j \right) & \text{if } \lambda_i \neq 0 \text{ and } \lambda_j = 0 \text{ for any } j \neq i. \\ 0 & \text{if } \lambda_i = 0 \text{ for any } i = 1, \dots, N. \end{cases}$$

Proof. Follow the proof of Proposition 6.3 and use the observation of Remark 6.2, \square

Corollary 6.5. *Suppose $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w^{\text{tr}}$ satisfies $\dim(\ker \mathfrak{K}_{\mathcal{X}}) = 1$. Let $\mathcal{X}^{\otimes N} = \mathcal{X} \times \dots \times \mathcal{X}$ (the product of N \mathcal{X} 's). Then,*

- (1) $\text{Tr}(|\mathfrak{K}_{\mathcal{X}^{\otimes N}}|) = N \cdot \text{Tr}(|\mathfrak{K}_{\mathcal{X}}|)$. In particular, this implies $\mathcal{X}^{\otimes N} \in \mathcal{M}_w^{\text{tr}}$.
- (2) $\text{Tr}(\mathfrak{K}_{\mathcal{X}^{\otimes N}}) = N \cdot \text{Tr}(\mathfrak{K}_{\mathcal{X}})$.
- (3) $\text{Tr}_{\text{neg}}(\mathfrak{K}_{\mathcal{X}^{\otimes N}}) = N \cdot \text{Tr}_{\text{neg}}(\mathfrak{K}_{\mathcal{X}})$.
- (4) For any $k \leq \text{pr}(\mathfrak{K}_{\mathcal{X}})$ and $(x_1, \dots, x_N), (x'_1, \dots, x'_N) \in \mathcal{X}^{\otimes N}$, we have

$$\begin{aligned} \|\Phi_{\mathcal{X}^{\otimes N}, kN}(x_1, \dots, x_N) - \Phi_{\mathcal{X}^{\otimes N}, kN}(x'_1, \dots, x'_N)\|^2 &= \sum_{i=1}^N \|\Phi_{\mathcal{X}, k}(x_i) - \Phi_{\mathcal{X}, k}(x'_i)\|^2, \text{ and} \\ \|\Phi_{\mathcal{X}^{\otimes N}}(x_1, \dots, x_N) - \Phi_{\mathcal{X}^{\otimes N}}(x'_1, \dots, x'_N)\|^2 &= \sum_{i=1}^N \|\Phi_{\mathcal{X}}(x_i) - \Phi_{\mathcal{X}}(x'_i)\|^2. \end{aligned}$$

Proof. Observe that the assumption $\dim(\ker \mathfrak{K}_{\mathcal{X}}) = 1$ implies that constant functions are the only eigenfunctions for zero eigenvalue. Then, this corollary is a direct result of Proposition 6.4. \square

6.2. The case of \mathbb{S}^1 . We will compute eigenvalues and corresponding eigenfunctions of the cMDS operator on \mathbb{S}^1 (with geodesic metric) and apply generalized cMDS. In this subsection, we will view \mathbb{S}^1 as $\mathbb{R}/2\pi$.

Lemma 6.6. *For any $n \in \mathbb{Z} \setminus \{0\}$,*

- (1) $\int_{-\pi}^{\pi} u^2 \cos(nu) du = \frac{(-1)^n 4\pi}{n^2}$, and
- (2) $\int_{-\pi}^{\pi} u^2 \sin(nu) du = 0$.

Proof. Fix arbitrary $n \in \mathbb{Z} \setminus \{0\}$. Then,

(1) For the first claim,

$$\begin{aligned}
 \int_{-\pi}^{\pi} u^2 \cos(nu) du &= \left[\frac{1}{n} u^2 \sin(nu) \right]_{u=-\pi}^{u=\pi} - \frac{2}{n} \int_{-\pi}^{\pi} u \sin(nu) du = -\frac{2}{n} \int_{-\pi}^{\pi} u \sin(nu) du \\
 &= -\frac{2}{n} \left[-\frac{1}{n} u \cos(nu) \right]_{u=-\pi}^{u=\pi} + \frac{2}{n^2} \int_{-\pi}^{\pi} \cos(nu) du \\
 &= \frac{4\pi}{n^2} \cos(n\pi) = \frac{(-1)^n 4\pi}{n^2}.
 \end{aligned}$$

(2) For the second claim,

$$\begin{aligned}
 \int_{-\pi}^{\pi} u^2 \sin(nu) du &= \left[-\frac{1}{n} u^2 \cos(nu) \right]_{u=-\pi}^{u=\pi} + \frac{2}{n} \int_{-\pi}^{\pi} u \cos(nu) du = \frac{2}{n} \int_{-\pi}^{\pi} u \cos(nu) du \\
 &= \frac{2}{n} \left[\frac{1}{n} u \sin(nu) \right]_{u=-\pi}^{u=\pi} - \frac{2}{n^2} \int_{-\pi}^{\pi} \sin(nu) du = 0.
 \end{aligned}$$

□

The following result has been described in [ABK20, Proposition 5.1]. However, we provide a proof here in order to give a standalone treatment which will be contrasted with the much harder case of \mathbb{S}^{d-1} for $d \geq 3$ (which will be dealt with in §6.4).

Proposition 6.7. *For each nonzero integer $n \in \mathbb{Z} \setminus \{0\}$, $\theta \mapsto \cos(n\theta)$ and $\theta \mapsto \sin(n\theta)$ are eigenfunctions of $\mathfrak{K}_{\mathbb{S}^1}$ for the eigenvalue $\frac{(-1)^{n+1}}{n^2}$.*

Proof. Fix arbitrary $n \in \mathbb{Z} \setminus \{0\}$. Then,

$$\begin{aligned}
 -\frac{1}{2} \int_{\mathbb{S}^1} d_{\mathbb{S}^1}^2(\theta, \theta') \cos(n\theta') d\theta' &= -\frac{1}{2} \cdot \frac{1}{2\pi} \int_{\theta-\pi}^{\theta+\pi} (\theta' - \theta)^2 \cos(n\theta') d\theta' \\
 &= -\frac{1}{4\pi} \int_{-\pi}^{\pi} u^2 \cos(nu + n\theta) du \quad (\text{by substituting } u = \theta' - \theta) \\
 &= -\cos(n\theta) \cdot \left(\frac{1}{4\pi} \int_{-\pi}^{\pi} u^2 \cos(nu) du \right) + \sin(n\theta) \cdot \left(\frac{1}{4\pi} \int_{-\pi}^{\pi} u^2 \sin(nu) du \right) \\
 &= \frac{(-1)^{n+1}}{n^2} \cos(n\theta) \quad (\because \text{Lemma 6.6}).
 \end{aligned}$$

for any $\theta \in \mathbb{S}^1$. Also,

$$\begin{aligned}
 -\frac{1}{2} \int_{\mathbb{S}^1} d_{\mathbb{S}^1}^2(\theta, \theta') \sin(n\theta') d\theta' &= -\frac{1}{2} \cdot \frac{1}{2\pi} \int_{\theta-\pi}^{\theta+\pi} (\theta' - \theta)^2 \sin(n\theta') d\theta' \\
 &= -\frac{1}{4\pi} \int_{-\pi}^{\pi} u^2 \sin(nu + n\theta) du \quad (\text{by substituting } u = \theta' - \theta) \\
 &= -\cos(n\theta) \cdot \left(\frac{1}{4\pi} \int_{-\pi}^{\pi} u^2 \sin(nu) du \right) - \sin(n\theta) \cdot \left(\frac{1}{4\pi} \int_{-\pi}^{\pi} u^2 \cos(nu) du \right) \\
 &= \frac{(-1)^{n+1}}{n^2} \sin(n\theta) \quad (\because \text{Lemma 6.6}).
 \end{aligned}$$

for any $\theta \in \mathbb{S}^1$. Moreover, since $\int_{-\pi}^{\pi} \cos(n\theta) d\theta = \int_{-\pi}^{\pi} \sin(n\theta) d\theta = 0$, the proof is completed by invoking Lemma 4.3. □

Proposition 6.7 implies that $\{\sqrt{2}\cos(n\theta), \sqrt{2}\sin(n\theta)\}_{n=1}^{\infty} \subset L^2(\text{nvols}_{\mathbb{S}^1})$ are orthonormal eigenfunctions of $\mathfrak{K}_{\mathbb{S}^1}$ corresponding to nonzero eigenvalues $\left\{\frac{(-1)^{n+1}}{n^2}\right\}_{n=1}^{\infty}$.

- Corollary 6.8.** (1) $\text{pr}(\mathfrak{K}_{\mathbb{S}^1}) = \infty$ and $\text{nr}(\mathfrak{K}_{\mathbb{S}^1}) = \infty$.
(2) $\text{Tr}(|\mathfrak{K}_{\mathbb{S}^1}|) = 2 \cdot \sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{3}$. In particular, $\mathbb{S}^1 \in \mathcal{M}_w^{\text{tr}}$.
(3) $\text{Tr}(\mathfrak{K}_{\mathbb{S}^1}) = 2 \cdot \left(\sum_{\text{odd } n \geq 1} \frac{1}{n^2} - \sum_{\text{even } n \geq 1} \frac{1}{n^2}\right) = \frac{\pi^2}{6}$.
(4) $\text{Tr}_{\text{neg}}(\mathfrak{K}_{\mathbb{S}^1}) = 2 \cdot \sum_{\text{even } n \geq 1} \frac{1}{n^2} = \frac{1}{2} \sum_{m=1}^{\infty} \frac{1}{m^2} = \frac{\pi^2}{12}$.
(5) For all $\theta, \theta' \in \mathbb{S}^1$,

$$\|\Phi_{\mathbb{S}^1}(\theta) - \Phi_{\mathbb{S}^1}(\theta')\| = \sqrt{\pi}(d_{\mathbb{S}^1}(\theta, \theta'))^{\frac{1}{2}}.$$

Note that item (5) provides a conclusive answer to the discussion at the end of Section 5 [ABK20].

Proof of Corollary 6.8. Items (1),(2),(3), and (4) are achieved from the well-known fact $\sum_{n=1}^{\infty} \frac{1}{n^2} = \frac{\pi^2}{6}$ and Proposition 6.7. Let's prove item (5).

Without loss of generality, we may assume that $\theta \geq \theta'$ and $2\pi \geq \theta - \theta' \geq 0$. Also,

$$\begin{aligned} \|\Phi_{\mathbb{S}^1}(\theta) - \Phi_{\mathbb{S}^1}(\theta')\|^2 &= \sum_{\text{odd } n \geq 1} \frac{1}{n^2} (2(\cos(n\theta) - \cos(n\theta'))^2 + 2(\sin(n\theta) - \sin(n\theta'))^2) \\ &= \sum_{\text{odd } n \geq 1} \frac{4}{n^2} (1 - (\cos(n\theta)\cos(n\theta') + \sin(n\theta)\sin(n\theta'))) \\ &= \sum_{\text{odd } n \geq 1} \frac{4}{n^2} (1 - \cos(n(\theta' - \theta))) \\ &= 4 \sum_{\text{odd } n \geq 1} \frac{1}{n^2} - 4 \sum_{\text{odd } n \geq 1} \frac{\cos(n(\theta' - \theta))}{n^2} \end{aligned}$$

Here, by [BS13, Section 21.6], we have

$$\sum_{\text{odd } n \geq 1} \frac{\cos(n(\theta' - \theta))}{n^2} = \begin{cases} \frac{\pi}{4}(\frac{\pi}{2} - (\theta' - \theta)) & \text{if } 0 \leq \theta' - \theta \leq \pi \\ \frac{\pi}{4}((\theta' - \theta) - \frac{3\pi}{2}) & \text{if } \pi < \theta' - \theta \leq 2\pi. \end{cases}$$

Hence, combining the above with the fact $\sum_{\text{odd } n \geq 1} \frac{1}{n^2} = \frac{\pi^2}{8}$, one concludes

$$\begin{aligned} \|\Phi_{\mathbb{S}^1}(\theta) - \Phi_{\mathbb{S}^1}(\theta')\|^2 &= 4 \left(\frac{\pi^2}{8} - \left(\frac{\pi^2}{8} - \frac{\pi}{4} d_{\mathbb{S}^1}(\theta, \theta') \right) \right) \\ &= \pi d_{\mathbb{S}^1}(\theta, \theta'). \end{aligned}$$

This completes the proof. □

Item (5) of Corollary 6.8 motivates the following conjecture which we arrived at via numerical computations.

Conjecture 1. For any integer $n \geq 1$

$$\|\Phi_{\mathbb{S}^n}(\theta) - \Phi_{\mathbb{S}^n}(\theta')\| = \sqrt{\pi}(d_{\mathbb{S}^n}(\theta, \theta'))^{\frac{1}{2}}.$$

It turns out that, the above conjecture is true. See Theorem 23.

Remark 6.9. We now compute the LHS and RHS of the inequality from item (2) of Proposition 4.22 for the case of \mathbb{S}^1 . First,

$$\begin{aligned}
 (\text{dis}(\mathbb{S}^1))^2 &= \frac{1}{(2\pi)^2} \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} (\|\Phi_{\mathbb{S}^1}(\theta) - \Phi_{\mathbb{S}^1}(\theta')\| - d_{\mathbb{S}^1}(\theta, \theta'))^2 d\theta d\theta' \\
 &= \frac{1}{(2\pi)^2} \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} (\sqrt{\pi}(d_{\mathbb{S}^1}(\theta, \theta'))^{\frac{1}{2}} - d_{\mathbb{S}^1}(\theta, \theta'))^2 d\theta d\theta' \\
 &= \frac{1}{(2\pi)^2} \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} \pi d_{\mathbb{S}^1}(\theta, \theta') d\theta d\theta' - \frac{1}{(2\pi)^2} \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} 2\sqrt{\pi}(d_{\mathbb{S}^1}(\theta, \theta'))^{\frac{3}{2}} d\theta d\theta' \\
 &\quad + \frac{1}{(2\pi)^2} \int_{\mathbb{S}^1} \int_{\mathbb{S}^1} (d_{\mathbb{S}^1}(\theta, \theta'))^2 d\theta d\theta' \\
 &= \int_0^\pi \theta d\theta - \frac{2}{\sqrt{\pi}} \int_0^\pi \theta^{\frac{3}{2}} d\theta + \frac{1}{\pi} \int_0^\pi \theta^2 d\theta \\
 &= \frac{\pi^2}{2} - \frac{4\pi^2}{5} + \frac{\pi^2}{3} = \frac{\pi^2}{30}.
 \end{aligned}$$

Hence, $\text{dis}(\mathbb{S}^1) = \frac{\pi}{\sqrt{30}}$. Also, $2\sqrt{\text{Tr}_{\text{neg}}(\mathbb{S}^1)} = \frac{\pi}{\sqrt{3}}$ by item (3) of Corollary 6.8. Indeed, they satisfy the inequality in item (2) of Proposition 4.22. However, we still do not know whether there is a constant $C > 0$ such that $\text{RHS} \leq C \cdot \text{LHS}$.

Lemma 6.10. The dimension of $\ker(\mathfrak{K}_{\mathbb{S}^1})$ is exactly 1. In other words, there are no eigenfunctions for zero eigenvalue other than constant functions (see item (4) of Lemma 4.2).

Proof. Observe that $\phi_0 \equiv 1$, $\phi_n(\theta) = \sqrt{2} \cos(n\theta)$, and $\psi_n(\theta) = \sqrt{2} \sin(n\theta)$ for $n \geq 1$ form a complete orthonormal basis of $L^2(\text{nvols}_{\mathbb{S}^1})$. Also, ϕ_n and ψ_n are the eigenfunctions for a nonzero eigenvalue $\frac{(-1)^{n+1}}{n^2}$ for each $n \geq 1$. This completes the proof. \square

6.3. The case of $\mathbb{T}^N = (\mathbb{S}^1)^N$. Applying the results about product spaces (see §6.1) we have the following result.

Corollary 6.11. Let $\mathbb{T}^N = (\mathbb{S}^1)^N$ be the N -torus. Then,

- (1) $\text{Tr}(|\mathfrak{K}_{\mathbb{T}^N}|) = \frac{N\pi^2}{3}$. In particular, $\mathbb{T}^N \in \mathcal{M}_w^{\text{tr}}$.
- (2) $\text{Tr}(\mathfrak{K}_{\mathbb{T}^N}) = \frac{N\pi^2}{6}$.
- (3) $\text{Tr}_{\text{neg}}(\mathbb{T}^N) = \frac{N\pi^2}{12}$.
- (4) For any positive integer k and $(\alpha_1, \dots, \alpha_N), (\beta_1, \dots, \beta_N) \in \mathbb{T}^N$, we have

$$\begin{aligned}
 \|\Phi_{\mathbb{T}^N, kN}(\alpha_1, \dots, \alpha_N) - \Phi_{\mathbb{T}^N, kN}(\beta_1, \dots, \beta_N)\|^2 &= \sum_{i=1}^N \|\Phi_{\mathbb{S}^1, k}(\alpha_i) - \Phi_{\mathbb{S}^1, k}(\beta_i)\|^2, \text{ and} \\
 \|\Phi_{\mathbb{T}^N}(\alpha_1, \dots, \alpha_N) - \Phi_{\mathbb{T}^N}(\beta_1, \dots, \beta_N)\|^2 &= \sum_{i=1}^N \|\Phi_{\mathbb{S}^1}(\alpha_i) - \Phi_{\mathbb{S}^1}(\beta_i)\|^2 = \pi \sum_{i=1}^N d_{\mathbb{S}^1}(\alpha_i, \beta_i).
 \end{aligned}$$

Proof. Apply Corollary 6.5, Corollary 6.8, and Lemma 6.10. \square

6.4. The case of \mathbb{S}^{d-1} . We will solve the eigenvalue problem of $\mathfrak{K}_{\mathbb{S}^{d-1}}$ for $d \geq 3$. Since \mathbb{S}^{d-1} is two-point homogeneous, it is equivalent to solving the following integral equations by Lemma 4.3 and Corollary 4.6:

- (1) $-\frac{1}{2} \int_{\mathbb{S}^{d-1}} d_{\mathbb{S}^{d-1}}^2(u, v) \phi(v) \, d\text{vol}_{\mathbb{S}^{d-1}}(v) = -\frac{1}{2|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} d_{\mathbb{S}^{d-1}}^2(u, v) \phi(v) \, d\text{vol}_{\mathbb{S}^{d-1}}(v) = \lambda \phi(u)$
for any $u \in \mathbb{S}^{d-1}$, and
- (2) $\int_{\mathbb{S}^{d-1}} \phi(u) \, d\text{vol}(u) = \frac{1}{|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} \phi(u) \, d\text{vol}(u) = 0.$

Theorem 22. *Let $d \geq 3$ and $n > 0$ be given. Then, any $Y_n^d \in \mathbb{Y}_n^d$ is an eigenfunction of $\mathfrak{K}_{\mathbb{S}^{d-1}}$ with the eigenvalue*

$$\lambda_{n,d} = -\frac{|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \int_{-1}^1 P_{n,d}(t) \arccos^2(t) (1-t^2)^{\frac{d-3}{2}} dt$$

Proof. Recall that $d_{\mathbb{S}^{d-1}}(u, v) = \arccos(\langle u, v \rangle)$ for any $u, v \in \mathbb{S}^{d-1}$. Then, by Funke-Hecke theorem (Theorem 14),

$$\begin{aligned} -\frac{1}{2|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} d_{\mathbb{S}^{d-1}}^2(u, v) Y_n^d(v) \, d\text{vol}_{\mathbb{S}^{d-1}}(v) &= -\frac{1}{2|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} \arccos^2(\langle u, v \rangle) Y_n^d(v) \, d\text{vol}_{\mathbb{S}^{d-1}}(v) \\ &= \lambda_{n,d} Y_n^d(u) \end{aligned}$$

for any $u \in \mathbb{S}^{d-1}$ where

$$\lambda_{n,d} = -\frac{|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \int_{-1}^1 P_{n,d}(t) \arccos^2(t) (1-t^2)^{\frac{d-3}{2}} dt.$$

Also, $\int_{\mathbb{S}^{d-1}} Y_n^d(u) \, d\text{vol}_{\mathbb{S}^{d-1}}(u) = 0$ by Corollary 2.29. This concludes the proof by Lemma 4.3. \square

Via Theorem 22 we now explicitly compute $\lambda_{n,3}$ for small n in the following examples.

Example 6.12. *When $d = 3$, by Theorem 22, the eigenvalues $\lambda_{n,3}$ of $\mathfrak{K}_{\mathbb{S}^2}$ have the following expression:*

$$\lambda_{n,3} = -\frac{1}{4} \int_{-1}^1 P_{n,d}(t) \arccos^2(t) dt.$$

Hence,

(1)

$$\begin{aligned} \lambda_{1,3} &= -\frac{1}{4} \int_{-1}^1 P_{1,3}(t) \arccos^2(t) dt \\ &= -\frac{1}{4} \int_{-1}^1 t \arccos^2(t) dt = \frac{\pi^2}{16}. \end{aligned}$$

(2)

$$\begin{aligned} \lambda_{2,3} &= -\frac{1}{4} \int_{-1}^1 P_{2,3}(t) \arccos^2(t) dt \\ &= -\frac{1}{4} \int_{-1}^1 \left(\frac{3}{2}t^2 - \frac{1}{2} \right) \arccos^2(t) dt = -\frac{1}{9}. \end{aligned}$$

(3)

$$\begin{aligned} \lambda_{3,3} &= -\frac{1}{4} \int_{-1}^1 P_{3,3}(t) \arccos^2(t) dt \\ &= -\frac{1}{4} \int_{-1}^1 \left(\frac{5}{2}t^3 - \frac{3}{2}t \right) \arccos^2(t) dt = \frac{\pi^2}{256}. \end{aligned}$$

(4)

$$\begin{aligned}\lambda_{4,3} &= -\frac{1}{4} \int_{-1}^1 P_{4,3}(t) \arccos^2(t) dt \\ &= -\frac{1}{4} \int_{-1}^1 \left(\frac{35}{8}t^4 - \frac{15}{4}t^2 + \frac{3}{8} \right) \arccos^2(t) dt = -\frac{4}{225}.\end{aligned}$$

6.4.1. *Explicit formula for the eigenvalues of $\mathfrak{K}_{\mathbb{S}^{d-1}}$.* Let's use the Taylor expansion of $\arccos(t)$.

Lemma 6.13. *For any $t \in [-1, 1]$,*

$$\arccos(t) = \frac{\pi}{2} - \sum_{n=0}^{\infty} a_n x^{2n+1}$$

where $a_n := \frac{(2n)!}{2^{2n}(n!)^2(2n+1)}$. This implies,

$$\arccos^2(t) = \frac{\pi^2}{4} - \pi \sum_{n=0}^{\infty} a_n x^{2n+1} + \sum_{n,m=0}^{\infty} a_n a_m x^{2(n+m+1)}.$$

Lemma 6.14. *For any $n, m > 0$, $k \geq 0$, and $d \geq 3$,*

$$\int_{-1}^1 P_{n,d}(t) t^m (1-t^2)^{\frac{d-3}{2}} dt = \begin{cases} 0 & \text{if the parities of } n, m \text{ are different} \\ 0 & \text{if the parities of } n, m \text{ are the same, and } n > m \\ \frac{(n+2k)! \Gamma(k+\frac{1}{2}) \Gamma(\frac{d-1}{2})}{2^n (2k)! \Gamma(k+n+\frac{d}{2})} & \text{if the parity of } n, m \text{ are the same, and } m = n + 2k \end{cases}$$

Proof. Let's do case-by-case analysis.

- (1) If the parities of n, m are different: Then, by item (4) of Lemma 2.30 and the parity assumption, $P_{n,d}(t)t^m(1-t^2)^{\frac{d-3}{2}}$ is odd function. So, $\int_{-1}^1 P_{n,d}(t)t^m(1-t^2)^{\frac{d-3}{2}} dt = 0$.
- (2) If the parities of n, m are the same, and $n > m$: It is easy to show that,

$$t^m = \sum_{k=0}^m b_k P_{k,d}(t)$$

since each $P_{k,d}(t)$ is a k -dimensional polynomial of t . So, by the orthogonality (see item (6) of Lemma 2.30) and $n > m$, we have

$$\int_{-1}^1 P_{n,3}(t)t^m(1-t^2)^{\frac{d-3}{2}} dt = 0.$$

- (3) If the parities of n, m are the same, and $m = n + 2k$: Recall the Rodrigues representation formula (see item (3) of Lemma 2.30):

$$P_{n,d}(t) = (-1)^n R_{n,d} (1-t^2)^{\frac{3-d}{2}} \left(\frac{d}{dt} \right)^n (1-t^2)^{n+\frac{d-3}{2}}$$

$$\text{where } R_{n,d} = \frac{\Gamma(\frac{d-1}{2})}{2^n \Gamma(n+\frac{d-1}{2})}.$$

Therefore,

$$\begin{aligned}
\int_{-1}^1 P_{n,d}(t)t^{n+2k}(1-t^2)^{\frac{d-3}{2}} dt &= (-1)^n R_{n,d} \int_{-1}^1 \left(\frac{d}{dt}\right)^n (1-t^2)^{n+\frac{d-3}{2}} t^{n+2k} dt \\
&= R_{n,d} \frac{(n+2k)!}{(2k)!} \int_{-1}^1 (1-t^2)^{n+\frac{d-3}{2}} t^{2k} dt \quad (\because \text{integration by parts } n \text{ times}) \\
&= R_{n,d} \frac{(n+2k)!}{(2k)!} \int_0^1 (1-s)^{n+\frac{d-3}{2}} s^{k-\frac{1}{2}} ds \\
&= R_{n,d} \frac{(n+2k)!}{(2k)!} B\left(k+\frac{1}{2}, n+\frac{d-1}{2}\right)
\end{aligned}$$

where $B(x, y)$ represents Beta function. It is well-known that $B(x, y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}$ (see [AH12, 1.14]). Hence,

$$\begin{aligned}
\int_{-1}^1 P_{n,d}(t)t^{n+2k}(1-t^2)^{\frac{d-3}{2}} dt &= R_{n,d} \frac{(n+2k)!}{(2k)!} \frac{\Gamma(k+\frac{1}{2})\Gamma(n+\frac{d-1}{2})}{\Gamma(k+n+\frac{d}{2})} \\
&= \frac{(n+2k)!}{2^n(2k)!} \frac{\Gamma(k+\frac{1}{2})\Gamma(\frac{d-1}{2})}{\Gamma(k+n+\frac{d}{2})}.
\end{aligned}$$

□

Proposition 6.15. *For any $n > 0$ and $d \geq 3$,*

$$\lambda_{n,d} = \begin{cases} \frac{\pi^{|\mathbb{S}^{d-2}|}}{2^{|\mathbb{S}^{d-1}|}} \sum_{k=0}^{\infty} a_{\frac{n+2k-1}{2}} \cdot \frac{(n+2k)!}{2^n(2k)!} \frac{\Gamma(k+\frac{1}{2})\Gamma(\frac{d-1}{2})}{\Gamma(k+n+\frac{d}{2})} & \text{if } n \text{ is odd} \\ -\frac{|\mathbb{S}^{d-2}|}{2^{|\mathbb{S}^{d-1}|}} \sum_{k=0}^{\infty} \left(\sum_{i,j:2(i+j+1)=n+2k} a_i a_j \right) \frac{(n+2k)!}{2^n(2k)!} \frac{\Gamma(k+\frac{1}{2})\Gamma(\frac{d-1}{2})}{\Gamma(k+n+\frac{d}{2})} & \text{if } n \text{ is even} \end{cases}$$

In particular, $\lambda_{n,d} > 0$ if n is odd and $\lambda_{n,d} < 0$ if n is even.

Proof. By Theorem 22 and Lemma 6.13,

$$\begin{aligned}
\lambda_{n,d} &= -\frac{|\mathbb{S}^{d-2}|}{2^{|\mathbb{S}^{d-1}|}} \int_{-1}^1 P_{n,d}(t) \arccos^2(t) (1-t^2)^{\frac{d-3}{2}} dt \\
&= -\frac{|\mathbb{S}^{d-2}|}{2^{|\mathbb{S}^{d-1}|}} \left(\frac{\pi^2}{4} \int_{-1}^1 P_{n,d}(t) (1-t^2)^{\frac{d-3}{2}} dt - \pi \sum_{i=0}^{\infty} a_i \int_{-1}^1 P_{n,d}(t) t^{2i+1} (1-t^2)^{\frac{d-3}{2}} dt \right. \\
&\quad \left. + \sum_{i,j=0}^{\infty} a_i a_j \int_{-1}^1 P_{n,d}(t) t^{2(i+j+1)} (1-t^2)^{\frac{d-3}{2}} dt \right).
\end{aligned}$$

Also, $\int_{-1}^1 P_{n,d}(t) (1-t^2)^{\frac{d-3}{2}} dt = 0$ for any $n > 0$ by item (5) of Lemma 2.30.

If n is odd, $\int_{-1}^1 P_{n,d}(t) t^{2(i+j+1)} (1-t^2)^{\frac{d-3}{2}} dt = 0$ and $\int_{-1}^1 P_{n,d}(t) t^{2i+1} (1-t^2)^{\frac{d-3}{2}} dt = 0$ if $2i+1 < n$ by Lemma 6.14. Also, for any $k \geq 0$,

$$\int_{-1}^1 P_{n,d}(t) t^{n+2k} (1-t^2)^{\frac{d-3}{2}} dt = \frac{(n+2k)!}{2^n(2k)!} \frac{\Gamma(k+\frac{1}{2})\Gamma(\frac{d-1}{2})}{\Gamma(k+n+\frac{d}{2})}$$

by Lemma 6.14. Therefore,

$$\lambda_{n,d} = \frac{\pi|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \sum_{k=0}^{\infty} a_{n+2k-1} \cdot \frac{(n+2k)! \Gamma(k + \frac{1}{2}) \Gamma(\frac{d-1}{2})}{2^n (2k)! \Gamma(k + n + \frac{d}{2})} > 0$$

If n is even, $\int_{-1}^1 P_{n,d}(t) t^{2i+1} (1-t^2)^{\frac{d-3}{2}} dt = 0$ and $\int_{-1}^1 P_{n,d}(t) t^{2(i+j+1)} (1-t^2)^{\frac{d-3}{2}} dt = 0$ if $2(i+j+1) < n$ by Lemma 6.14. Also, for any $k \geq 0$,

$$\int_{-1}^1 P_{n,d}(t) t^{n+2k} (1-t^2)^{\frac{d-3}{2}} dt = \frac{(n+2k)! \Gamma(k + \frac{1}{2}) \Gamma(\frac{d-1}{2})}{2^n (2k)! \Gamma(k + n + \frac{d}{2})}$$

by Lemma 6.14. Therefore,

$$\lambda_{n,d} = -\frac{|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \sum_{k=0}^{\infty} \left(\sum_{i,j:2(i+j+1)=n+2k} a_i a_j \right) \frac{(n+2k)! \Gamma(k + \frac{1}{2}) \Gamma(\frac{d-1}{2})}{2^n (2k)! \Gamma(k + n + \frac{d}{2})} < 0.$$

□

There is another way to compute $\lambda_{n,d}$ when n is odd, and it is based on the following result.

Lemma 6.16 ([BBS07, Section 6]). *For any $d \geq 3$, $n \geq 1$, $u \in \mathbb{S}^{d-1}$, and $Y_n^d \in \mathbb{Y}_n^d$,*

$$\frac{1}{|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} d_{\mathbb{S}^{d-1}}(u, v) Y_n^d(v) d\text{vol}_{\mathbb{S}^{d-1}}(v) = \eta_{n,d} Y_n^d(u)$$

where

$$\eta_{n,d} = \begin{cases} -\frac{2^{\frac{n-1}{2}} \pi^{\frac{d+1}{2}} n \Gamma(\frac{d}{2}) \Gamma(\frac{n}{2})}{n! (\Gamma(\frac{n+d}{2}))^2 |\mathbb{S}^{d-1}|} & \text{if } n \text{ is odd} \\ 0 & \text{if } n \text{ is even} \end{cases}$$

Our proof is basically the same as [BBS07, Section 6]. But, we write it for the completeness of the paper.

Proof. By Theorem 14, for any $u \in \mathbb{S}^{d-1}$ and $Y_n^d \in \mathbb{Y}_n^d$,

$$\frac{1}{|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} d_{\mathbb{S}^{d-1}}(u, v) Y_n^d(v) d\text{vol}_{\mathbb{S}^{d-1}}(v) = \eta_{n,d} Y_n^d(u)$$

holds with

$$\eta_{n,d} = \frac{|\mathbb{S}^{d-2}|}{|\mathbb{S}^{d-1}|} \int_{-1}^1 P_{n,d}(t) \arccos(t) (1-t^2)^{\frac{d-3}{2}} dt.$$

Moreover, by Erdelyi [Bat53, pg.248], it is shown that

$$\eta_{n,d} = \frac{i^n (2\pi)^{\frac{d}{2}}}{|\mathbb{S}^{d-1}|} \int_{-\infty}^{\infty} t^{1-\frac{d}{2}} J_{n+\frac{d}{2}-1}(t) \hat{f}(t) dt$$

where $J_{n+\frac{d}{2}-1}(t)$ denotes Bessel function [Bat53, Chapter VII], and

$$\hat{f}(t) = \frac{1}{2\pi} \int_{-1}^1 e^{-ixt} \arccos(x) dx.$$

Moreover, by substituting $x = \cos \theta$,

$$\hat{f}(t) = \frac{1}{2\pi} \int_{-1}^1 e^{-ixt} \arccos(x) dx = \frac{1}{2\pi} \int_0^\pi \theta \sin \theta e^{-it \cos \theta} d\theta = \frac{1}{2it} (e^{it} - J_0(t))$$

Corresponding to the two terms in $\hat{f}(t)$ there are two terms in $\eta_{n,d}$. The integral including e^{it} is zero for all $n \neq 0$ and the integral with $J_0(t)$ is zero for even n . For odd n ,

$$\eta_{n,d} = -\frac{i^{n-1}(2\pi)^{\frac{d}{2}}}{|\mathbb{S}^{d-1}|} \int_0^\infty t^{-\frac{d}{2}} J_{n+\frac{d}{2}-1}(t) J_0(t) dt$$

The last integral can be computed using the following integral [Bat53, 7.7.4.30]

$$\int_0^\infty t^{-\rho} J_\mu(t) J_\nu(t) dt = \frac{\Gamma(\rho) \Gamma(\frac{1}{2}(\nu + \mu + 1 - \rho))}{2^\rho \Gamma(\frac{1}{2}(1 + \nu - \mu + \rho)) \Gamma(\frac{1}{2}(1 + \nu + \mu + \rho)) \Gamma(\frac{1}{2}(1 + \mu - \nu + \rho))}.$$

Hence, the final result is

$$\begin{aligned} \eta_{n,d} &= -\frac{1}{|\mathbb{S}^{d-1}|} (-1)^{\frac{n-1}{2}} \pi^{\frac{d}{2}} \frac{\Gamma(\frac{d}{2}) \Gamma(\frac{n}{2})}{\Gamma(1 - \frac{n}{2}) (\Gamma(\frac{n+d}{2}))^2} \\ &= -\frac{\pi^{\frac{d-1}{2}} n!! \Gamma(\frac{d}{2}) \Gamma(\frac{n}{2})}{n 2^{\frac{n-1}{2}} (\Gamma(\frac{n+d}{2}))^2 |\mathbb{S}^{d-1}|} \end{aligned}$$

as we wanted. □

Proposition 6.17. *For any $d \geq 3$ and odd $n \geq 1$,*

$$\lambda_{n,d} = -\frac{\pi}{2} \eta_{n,d}.$$

Proof. Recall that

$$\lambda_{n,d} = -\frac{|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \int_{-1}^1 P_{n,d}(t) \arccos^2(t) (1-t^2)^{\frac{d-3}{2}} dt,$$

and

$$\eta_{n,d} = \frac{|\mathbb{S}^{d-2}|}{|\mathbb{S}^{d-1}|} \int_{-1}^1 P_{n,d}(t) \arccos(t) (1-t^2)^{\frac{d-3}{2}} dt.$$

Moreover, since n is odd, $P_{n,d}(-t) = P_{n,d}(t)$ for any $t \in [-1, 1]$. Hence,

$$\begin{aligned} & \int_{-1}^1 P_{n,d}(t) \arccos^2(t) (1-t^2)^{\frac{d-3}{2}} dt \\ &= \left(\int_0^1 P_{n,d}(t) \arccos^2(t) (1-t^2)^{\frac{d-3}{2}} dt - \int_0^1 P_{n,d}(t) \arccos^2(-t) (1-t^2)^{\frac{d-3}{2}} dt \right) \\ &= \left(\int_0^1 P_{n,d}(t) \arccos^2(t) (1-t^2)^{\frac{d-3}{2}} dt - \int_0^1 P_{n,d}(t) (\pi - \arccos(t))^2 (1-t^2)^{\frac{d-3}{2}} dt \right) \\ &= -\pi^2 \int_0^1 P_{n,d}(t) (1-t^2)^{\frac{d-3}{2}} dt + 2\pi \int_0^1 P_{n,d}(t) \arccos(t) (1-t^2)^{\frac{d-3}{2}} dt. \end{aligned}$$

Also,

$$\begin{aligned}
 & \int_{-1}^1 P_{n,d}(t) \arccos(t)(1-t^2)^{\frac{d-3}{2}} dt \\
 &= \int_0^1 P_{n,d}(t) \arccos(t)(1-t^2)^{\frac{d-3}{2}} dt - \int_0^1 P_{n,d}(t) \arccos(-t)(1-t^2)^{\frac{d-3}{2}} dt \\
 &= \int_0^1 P_{n,d}(t) \arccos(t)(1-t^2)^{\frac{d-3}{2}} dt - \int_0^1 P_{n,d}(t)(\pi - \arccos(-t))(1-t^2)^{\frac{d-3}{2}} dt \\
 &= 2 \int_0^1 P_{n,d}(t) \arccos(t)(1-t^2)^{\frac{d-3}{2}} dt - \pi \int_0^1 P_{n,d}(t)(1-t^2)^{\frac{d-3}{2}} dt.
 \end{aligned}$$

Hence,

$$\int_0^1 P_{n,d}(t) \arccos(t)(1-t^2)^{\frac{d-3}{2}} dt = \frac{1}{2} \left(\int_{-1}^1 P_{n,d}(t) \arccos(t)(1-t^2)^{\frac{d-3}{2}} dt + \pi \int_0^1 P_{n,d}(t)(1-t^2)^{\frac{d-3}{2}} dt \right).$$

Therefore,

$$\int_{-1}^1 P_{n,d}(t) \arccos^2(t)(1-t^2)^{\frac{d-3}{2}} dt = \pi \int_{-1}^1 P_{n,d}(t) \arccos(t)(1-t^2)^{\frac{d-3}{2}} dt.$$

This implies, $\lambda_{n,d} = -\frac{\pi}{2}\eta_{n,d}$ as we required. \square

Moreover, Proposition 6.17 help in settling Conjecture 1.

Theorem 23. For any $d \geq 3$, and $u, v \in \mathbb{S}^{d-1}$,

$$\|\phi_{\mathbb{S}^{d-1}}(u) - \phi_{\mathbb{S}^{d-1}}(v)\|^2 = \pi \cdot d_{\mathbb{S}^{d-1}}(u, v).$$

Proof. Observe that, by Lemma 6.16,

$$(4) \quad d_{\mathbb{S}^{d-1}}(u, v) = \eta_{0,d} + \sum_{\text{odd } n \geq 1} \eta_{n,d} \left(\sum_{j=1}^{N_{n,d}} Y_{n,j}^d(u) Y_{n,j}^d(v) \right)$$

where $\eta_{0,d} = \frac{1}{|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} d_{\mathbb{S}^{d-1}}(x, y) d\text{vol}_{\mathbb{S}^{d-1}}(y) = \text{diam}_1(\mathbb{S}^{d-1})$, and $\{Y_{n,j}^d\}_{j=1}^{N_{n,d}}$ is an orthonormal basis of \mathbb{Y}_n^d for any $u, v \in \mathbb{S}^{d-1}$. In particular, if we choose $u = v$, one obtains

$$0 = \eta_{0,d} + \sum_{\text{odd } n \geq 1} \eta_{n,d} \left(\sum_{j=1}^{N_{n,d}} (Y_{n,j}^d(u))^2 \right) = \eta_{0,d} + \sum_{\text{odd } n \geq 1} \eta_{n,d} N_{n,d}$$

since $\left(\sum_{j=1}^{N_{n,d}} (Y_{n,j}^d(u))^2 \right) = N_{n,d}$ by the addition theorem (Theorem 13).

Also, recall that

$$\|\phi_{\mathbb{S}^{d-1}}(u) - \phi_{\mathbb{S}^{d-1}}(v)\|^2 = K_{\widehat{\mathbb{S}^{d-1}}}(u, u) + K_{\widehat{\mathbb{S}^{d-1}}}(v, v) - 2K_{\widehat{\mathbb{S}^{d-1}}}(u, v)$$

for any $u, v \in \mathbb{S}^{d-1}$. Moreover, since $\lambda_{n,d} > 0$ only for odd n by Proposition 6.15,

$$\begin{aligned} K_{\widehat{\mathbb{S}}^{d-1}}(u, u) &= \sum_{\text{odd } n \geq 1} \lambda_{n,d} \left(\sum_{j=1}^{N_{n,d}} (Y_{n,j}^d(u))^2 \right) = \sum_{\text{odd } n \geq 1} \lambda_{n,d} N_{n,d} \\ &= \sum_{\text{odd } n \geq 1} \left(-\frac{\pi}{2} \eta_{n,d} \right) N_{n,d} = -\frac{\pi}{2} \sum_{\text{odd } n \geq 1} \eta_{n,d} N_{n,d}. \end{aligned}$$

where the the third equality holds by Proposition 6.17.

Similarly, $K_{\widehat{\mathbb{S}}^{d-1}}(v, v) = -\frac{\pi}{2} \sum_{\text{odd } n \geq 1} \eta_{n,d} N_{n,d}$. Hence,

$$K_{\widehat{\mathbb{S}}^{d-1}}(u, u) + K_{\widehat{\mathbb{S}}^{d-1}}(v, v) = \pi \left(- \sum_{n \geq 1, \text{odd}} \eta_{n,d} N_{n,d} \right) = \pi \cdot \eta_{0,d}.$$

Also,

$$\begin{aligned} K_{\widehat{\mathbb{S}}^{d-1}}(u, v) &= \sum_{\text{odd } n \geq 1} \lambda_{n,d} \left(\sum_{j=1}^{N_{n,d}} Y_{n,j}^d(u) Y_{n,j}^d(v) \right) \\ &= -\frac{\pi}{2} \sum_{\text{odd } n \geq 1} \eta_{n,d} \left(\sum_{j=1}^{N_{n,d}} Y_{n,j}^d(u) Y_{n,j}^d(v) \right) \end{aligned}$$

Finally, one concludes that

$$\|\phi_{\mathbb{S}^{d-1}}(u) - \phi_{\mathbb{S}^{d-1}}(v)\|^2 = \pi \left(\eta_{0,d} + \sum_{n \geq 1, \text{odd}} \eta_{n,d} \left(\sum_{j=1}^{N_{n,d}} Y_{n,j}^d(u) Y_{n,j}^d(v) \right) \right) = \pi d_{\mathbb{S}^{d-1}}(u, v)$$

where the last equality holds because of equation (4). \square

Remark 6.18. Note that Theorem 23 implies that

$$\delta(d-1) := \text{dis}(\mathbb{S}^{d-1}) = \left((\text{diam}_2(\mathbb{S}^{d-1}))^2 + \pi \text{diam}_1(\mathbb{S}^{d-1}) - 2\sqrt{\pi} (\text{diam}_{\frac{3}{2}}(\mathbb{S}^{d-1}))^{\frac{3}{2}} \right)^{\frac{1}{2}}.$$

This suggests, via concentration of measure considerations, that $\delta(\infty) = \frac{\pi}{2} (3 - 2\sqrt{2})^{1/2}$.

Question 2. In analogy with Remark 6.9: What is the precise value of $\text{Tr}_{\text{neg}}(\mathbb{S}^{d-1})$ for arbitrary $d \geq 3$?

6.4.2. *Traceability of $\mathfrak{K}_{\mathbb{S}^{d-1}}$.* We already know that $\mathfrak{K}_{\mathbb{S}^1}$ is in trace class. Furthermore, by employing the Funk-Hecke theorem, we prove the traceability of $\mathfrak{K}_{\mathbb{S}^{d-1}}$ for any $d \geq 3$.

Lemma 6.19. For any $d \geq 3$,

$$\Delta d_{\mathbb{S}^{d-1}}(e_d, \cdot)(u) = -(d-2) \cdot \cot d_{\mathbb{S}^{d-1}}(e_d, u)$$

for any $u = (u_1, \dots, u_d) \in \mathbb{S}^{d-1} \setminus \{e_d, -e_d\}$ where $e_d = (0, \dots, 0, 1) \in \mathbb{S}^{d-1}$ is the north pole of \mathbb{S}^{d-1} .

Proof. Observe that $d_{\mathbb{S}^{d-1}}(e_d, u) = \arccos(\langle e_d, u \rangle) = \arccos(u_d)$. We use the following expression of the Laplace-Beltrami operator Δ of \mathbb{S}^{d-1} (see [AH12, (3.18)]):

$$\Delta = - \sum_{1 \leq j < i \leq d} \left(u_i \frac{\partial}{\partial u_j} - u_j \frac{\partial}{\partial u_i} \right)^2.$$

Then,

$$\begin{aligned} \Delta d_{\mathbb{S}^{d-1}}(e_d, \cdot)(u) &= - \sum_{1 \leq j < i \leq d} \left(u_i \frac{\partial}{\partial u_j} - u_j \frac{\partial}{\partial u_i} \right)^2 \arccos(u_d) \\ &= - \sum_{j=1}^{d-1} \left(u_d \frac{\partial}{\partial u_j} - u_j \frac{\partial}{\partial u_d} \right) \frac{u_j}{\sqrt{1-u_d^2}} \\ &= - \sum_{j=1}^{d-1} \left(\frac{u_d}{\sqrt{1-u_d^2}} - \frac{u_j^2 u_d}{(1-u_d^2)^{\frac{3}{2}}} \right) \\ &= -(d-1) \cdot \frac{u_d}{\sqrt{1-u_d^2}} + \frac{u_d}{(1-u_d^2)^{\frac{3}{2}}} \cdot \sum_{j=1}^{d-1} u_j^2 \\ &= -(d-1) \cdot \frac{u_d}{\sqrt{1-u_d^2}} + \frac{u_d}{\sqrt{1-u_d^2}} \left(\because \sum_{j=1}^{d-1} u_j^2 = 1 - u_d^2 \right) \\ &= -(d-2) \cdot \frac{u_d}{\sqrt{1-u_d^2}} = -(d-2) \cdot \cot d_{\mathbb{S}^{d-1}}(e_d, u). \end{aligned}$$

□

Lemma 6.20. *For any $d \geq 3$, $n \geq 1$, and $\delta \in (0, 1)$,*

$$|\lambda_{n,d}| \leq \frac{C''_{\delta,d}}{n(n+d-2)} N_{n,d}^{-\frac{1}{1+\delta}}$$

for some constant $C''_{\delta,d} > 0$.

Proof. Let

$$\begin{aligned} F : \mathbb{S}^{d-1} &\longrightarrow \mathbb{R} \\ u &\longmapsto \arccos^2(\langle e_d, u \rangle). \end{aligned}$$

Then, by Theorem 22,

$$\lambda_{n,d} = - \frac{|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \int_{-1}^1 P_{n,d}(t) \arccos^2(t) (1-t^2)^{\frac{d-3}{2}} dt = - \frac{1}{2|\mathbb{S}^{d-1}|} \int_{\mathbb{S}^{d-1}} P_{n,d}(\langle e_d, u \rangle) F(u) d \text{vol}_{\mathbb{S}^{d-1}}(u).$$

Observe that, for any $\delta \in (0, 1)$,

$$\begin{aligned}
|\lambda_{n,d}| &= \frac{1}{2|\mathbb{S}^{d-1}|} \left| \int_{\mathbb{S}^{d-1}} P_{n,d}(e_d \cdot u) F(u) d \text{vol}_{\mathbb{S}^{d-1}}(u) \right| \\
&= \frac{1}{2n(n+d-2)|\mathbb{S}^{d-1}|} \left| \int_{\mathbb{S}^{d-1}} \langle \nabla_u P_{n,d}(e_d \cdot u), \nabla F(u) \rangle d \text{vol}_{\mathbb{S}^{d-1}}(u) \right| \quad (\because \text{item (9) of Lemma 2.30}) \\
&= \frac{1}{2n(n+d-2)|\mathbb{S}^{d-1}|} \left| \int_{\mathbb{S}^{d-1}} P_{n,d}(e_d \cdot u) \Delta F(u) d \text{vol}_{\mathbb{S}^{d-1}}(u) \right| \quad (\because \text{item (8) of Lemma 2.30}) \\
&\leq \frac{1}{2n(n+d-2)|\mathbb{S}^{d-1}|} \|P_{n,d}(e_d \cdot u) \Delta F(u)\|_{L^1(\text{vol}_{\mathbb{S}^{d-1}})} \\
&\leq \frac{1}{2n(n+d-2)|\mathbb{S}^{d-1}|} \|P_{n,d}(e_d \cdot u)\|_{L^{1+\frac{1}{\delta}}(\text{vol}_{\mathbb{S}^{d-1}})} \cdot \|\Delta F(u)\|_{L^{1+\delta}(\text{vol}_{\mathbb{S}^{d-1}})} \quad (\because \text{Holder's inequality})
\end{aligned}$$

where the second and third equality make sense because ∇f and Δf are well-defined almost everywhere on \mathbb{S}^{d-1} . Now, let's prove that $\|\Delta F(u)\|_{L^{1+\delta}(\text{vol}_{\mathbb{S}^{d-1}})}$ is bounded by a constant $C_{\delta,d} > 0$: First, note that

$$\begin{aligned}
\Delta F &= \Delta d_{\mathbb{S}^{d-1}}^2(e_d, \cdot) = \nabla \cdot \nabla d_{\mathbb{S}^{d-1}}^2(e_d, \cdot) = 2\nabla \cdot (d_{\mathbb{S}^{d-1}}(e_d, \cdot) \nabla d_{\mathbb{S}^{d-1}}(e_d, \cdot)) \\
&= 2(\|\nabla d_{\mathbb{S}^{d-1}}(e_d, \cdot)\|^2 + d_{\mathbb{S}^{d-1}}(e_d, \cdot) \Delta d_{\mathbb{S}^{d-1}}(e_d, \cdot)).
\end{aligned}$$

Also, since $d_{\mathbb{S}^{d-1}}(e_d, \cdot)$ is 1-Lipschitz, $\|\nabla d_{\mathbb{S}^{d-1}}(e_d, \cdot)\| \leq 1$. Moreover,

$$\begin{aligned}
&\int_{\mathbb{S}^{d-1}} |d_{\mathbb{S}^{d-1}}(e_d, u) \Delta d_{\mathbb{S}^{d-1}}(e_d, u)|^{(1+\delta)} d \text{vol}_{\mathbb{S}^{d-1}}(u) \\
&= (d-2) \int_{\mathbb{S}^{d-1}} |d_{\mathbb{S}^{d-1}}(e_d, u) \cot d_{\mathbb{S}^{d-1}}(e_d, u)|^{(1+\delta)} d \text{vol}_{\mathbb{S}^{d-1}}(u) \quad (\because \text{Lemma 6.19}) \\
&= (d-2) |\mathbb{S}^{d-2}| \int_0^\pi |\theta \cot \theta|^{1+\delta} (\sin \theta)^{d-2} d\theta \\
&\leq (d-2) |\mathbb{S}^{d-2}| \int_0^\pi |\theta \cot \theta|^{1+\delta} \sin \theta d\theta \quad (\because \sin \theta \in [0, 1] \text{ for } \theta \in [0, \pi]) \\
&= (d-2) |\mathbb{S}^{d-2}| \int_0^\pi \frac{\theta^{1+\delta} |\cos \theta|^{1+\delta}}{(\sin \theta)^\delta} d\theta.
\end{aligned}$$

Let's bound $\int_0^\pi \frac{\theta^{1+\delta} |\cos \theta|^{1+\delta}}{(\sin \theta)^\delta} d\theta$. Observe that,

$$\begin{aligned}
\int_0^{\frac{\pi}{2}} \frac{\theta^{1+\delta} |\cos \theta|^{1+\delta}}{(\sin \theta)^\delta} d\theta &\leq \int_0^{\frac{\pi}{2}} \frac{\theta^{1+\delta}}{(\sin \theta)^\delta} d\theta \quad (\because \cos \theta \in [-1, 1]) \\
&\leq \int_0^{\frac{\pi}{2}} \theta \left(\frac{\pi}{2}\right)^\delta d\theta \quad \left(\because \frac{\theta}{\sin \theta} \leq \frac{\pi}{2}\right) < \infty,
\end{aligned}$$

and

$$\begin{aligned}
 \int_{\frac{\pi}{2}}^{\pi} \frac{\theta^{1+\delta} |\cos \theta|^{1+\delta}}{(\sin \theta)^\delta} d\theta &\leq \pi^{1+\delta} \int_{\frac{\pi}{2}}^{\pi} \frac{|\cos \theta|}{(\sin \theta)^\delta} d\theta \quad (\because \cos \theta \in [-1, 1]) \\
 &= -\pi^{1+\delta} \int_{\frac{\pi}{2}}^{\pi} \frac{\cos \theta}{(\sin \theta)^\delta} d\theta \\
 &= \pi^{1+\delta} \int_0^1 \frac{1}{t^\delta} dt = \frac{\pi^{1+\delta}}{1-\delta} < \infty.
 \end{aligned}$$

So, we conclude that $\|\Delta F(u)\|_{L^{1+\delta}(\text{vol}_{\mathbb{S}^{d-1}})} \leq C_{\delta,d}$ for some positive integer $C_{\delta,d} > 0$.

Next,

$$\begin{aligned}
 \int_{\mathbb{S}^{d-1}} |P_{n,d}(\langle e_d, u \rangle)|^{1+\frac{1}{\delta}} d \text{vol}_{\mathbb{S}^{d-1}}(u) &= \int_{\mathbb{S}^{d-1}} |P_{n,d}(\langle e_d, u \rangle)|^2 \cdot |P_{n,d}(\langle e_d, u \rangle)|^{\frac{1}{\delta}-1} d \text{vol}_{\mathbb{S}^{d-1}}(u) \\
 &\leq \int_{\mathbb{S}^{d-1}} |P_{n,d}(\langle e_d, u \rangle)|^2 d \text{vol}_{\mathbb{S}^{d-1}}(u) \quad (\because \text{item (1) of Lemma 2.30}) \\
 &= \frac{|\mathbb{S}^{d-1}|}{N_{n,d}} \quad (\because \text{item (2) of Lemma 2.30}).
 \end{aligned}$$

This implies $\|P_{n,d}(e_d \cdot u)\|_{L^{1+\frac{1}{\delta}}(\mathbb{S}^{d-1})} \leq C'_{\delta,d} N_{n,d}^{-\frac{1}{1+\frac{1}{\delta}}}$ for some positive constant $C'_{\delta,d} > 0$. Finally, we conclude that

$$|\lambda_{n,d}| \leq \frac{C''_{\delta,d}}{n(n+d-2)} N_{n,d}^{-\frac{1}{1+\frac{1}{\delta}}}$$

for some positive constant $C''_{\delta,d} > 0$. □

Theorem 24. $\mathfrak{K}_{\mathbb{S}^2}$ is trace class.

Proof. Fix an arbitrary $\delta \in (0, 1)$. Then, by Lemma 6.20, we know $|\lambda_{n,3}| \leq \frac{C''_{\delta,3}}{n(n+1)} N_{n,3}^{-\frac{1}{1+\frac{1}{\delta}}}$. Therefore,

$$\sum_{n=1}^{\infty} N_{n,3} |\lambda_{n,3}| \leq \sum_{n=1}^{\infty} \frac{C''_{\delta,3}}{n(n+1)} N_{n,3}^{1-\frac{1}{1+\frac{1}{\delta}}}$$

But, since $N_{n,3} = 2n + 1$,

$$\frac{N_{n,3}^{1-\frac{1}{1+\frac{1}{\delta}}}}{n(n+1)} = O(n^{\frac{1}{1+\delta}-2}).$$

Since $\frac{1}{1+\delta} - 2 < -1$, we conclude $\sum_{n=1}^{\infty} N_{n,3} |\lambda_{n,3}|$ converges so that $\mathfrak{K}_{\mathbb{S}^2}$ is trace class. □

Unfortunately, the technique used in the proof of Theorem 24 cannot be generalized to higher dimensional spheres. In Section 6.4.3, this technique will nevertheless yield interesting conclusions regarding *metric transformations* (such as those arising from applying the snowflake transform) to 2-dimensional spheres.

For general $d \geq 3$, the following strategy exploiting the Poisson identity yields a successful approach.

Lemma 6.21. *Suppose a positive integer $d \geq 3$ is given. Then,*

$$\int_{-1}^1 \frac{\arccos^2(t)(1-t^2)^{\frac{d-3}{2}}}{(1+r^2+2rt)^{\frac{d}{2}}} dt \in \left[0, \frac{2\pi^2}{1-r^2}\right]$$

for each $r \in (0, 1)$.

Proof. Observe that $\arccos^2(t), (1-t^2), (1+r^2+2rt) \geq 0$ for any $r \in (0, 1)$ and $t \in [-1, 1]$.

So, obviously $\int_{-1}^1 \frac{\arccos^2(t)(1-t^2)^{\frac{d-3}{2}}}{(1+r^2+2rt)^{\frac{d}{2}}} dt \geq 0$. Moreover,

$$\begin{aligned} \int_{-1}^1 \frac{\arccos^2(t)(1-t^2)^{\frac{d-3}{2}}}{(1+r^2+2rt)^{\frac{d}{2}}} dt &\leq \pi^2 \int_{-1}^1 \frac{(1-t^2)^{\frac{d-3}{2}}}{(1+r^2+2rt)^{\frac{d}{2}}} dt \quad (\because \arccos(t) \in [0, \pi]) \\ &\leq \pi^2 \int_{-1}^1 \frac{(1+r^2+2rt)^{\frac{d-3}{2}}}{(1+r^2+2rt)^{\frac{d}{2}}} dt \quad (\because 1-t^2 \leq 1+r^2+2rt) \\ &= \pi^2 \int_{-1}^1 \frac{1}{(1+r^2+2rt)^{\frac{3}{2}}} dt \\ &= \pi^2 \left[-\frac{1}{r}(1+r^2+2rt)^{-\frac{1}{2}} \right]_{t=-1}^{t=1} \\ &= \frac{\pi^2}{r} \left(\frac{1}{1-r} - \frac{1}{1+r} \right) = \frac{2\pi^2}{1-r^2} \end{aligned}$$

as we wanted. □

Theorem 25. *For any $d \geq 3$, $\mathfrak{K}_{\mathbb{S}^{d-1}}$ is trace class.*

Proof. To prove $\mathfrak{K}_{\mathbb{S}^{d-1}}$ is trace class, it is enough to show that $\sum_{n=1}^{\infty} |\lambda_{n,d}| N_{n,d} < \infty$ (see Theorem 8). Now, fix arbitrary $r \in (0, 1)$. Then,

$$\begin{aligned} &\sum_{n=1}^{\infty} |\lambda_{n,d}| N_{n,d} r^n \\ &= - \sum_{n=1}^{\infty} \lambda_{n,d} (-r)^n N_{n,d} \quad (\because \lambda_{n,d} > 0 \text{ for odd } n \text{ and } \lambda_{n,d} < 0 \text{ for even } n \text{ by Proposition 6.15}) \\ &= \frac{|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \sum_{n=1}^{\infty} \int_{-1}^1 P_{n,d}(t) \arccos^2(t)(1-t^2)^{\frac{d-3}{2}} (-r)^n N_{n,d} dt \\ &= \frac{|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \left(\sum_{n=0}^{\infty} \int_{-1}^1 P_{n,d}(t) \arccos^2(t)(1-t^2)^{\frac{d-3}{2}} (-r)^n N_{n,d} dt - \int_{-1}^1 \arccos^2(t)(1-t^2)^{\frac{d-3}{2}} dt \right) \\ &= \frac{|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \sum_{n=0}^{\infty} \int_{-1}^1 P_{n,d}(t) \arccos^2(t)(1-t^2)^{\frac{d-3}{2}} (-r)^n N_{n,d} dt - \frac{1}{2} (\text{diam}_2(\mathbb{S}^{d-1}))^2 \\ &= \frac{|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \int_{-1}^1 \sum_{n=0}^{\infty} P_{n,d}(t) \arccos^2(t)(1-t^2)^{\frac{d-3}{2}} (-r)^n N_{n,d} dt - \frac{1}{2} (\text{diam}_2(\mathbb{S}^{d-1}))^2. \end{aligned}$$

Here, note that the last equality holds by the dominated convergence theorem, since

$$\sum_{n=0}^k P_{n,d}(t) \arccos^2(t)(1-t^2)^{\frac{d-3}{2}}(-r)^n N_{n,d}$$

uniformly converges, as $k \rightarrow \infty$, to the following non-negative and integrable function

$$\frac{(1-r^2) \arccos^2(t)(1-t^2)^{\frac{d-3}{2}}}{(1+r^2+2rt)^{\frac{d}{2}}}$$

by item (7) of Lemma 2.30 and Lemma 6.21. Then,

$$\begin{aligned} & \frac{|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \int_{-1}^1 \sum_{n=0}^{\infty} P_{n,d}(t) \arccos^2(t)(1-t^2)^{\frac{d-3}{2}}(-r)^n N_{n,d} dt - \frac{1}{2}(\text{diam}_2(\mathbb{S}^{d-1}))^2 \\ &= \frac{|\mathbb{S}^{d-2}|}{2|\mathbb{S}^{d-1}|} \int_{-1}^1 \frac{(1-r^2) \arccos^2(t)(1-t^2)^{\frac{d-3}{2}}}{(1+r^2+2rt)^{\frac{d}{2}}} dt - \frac{1}{2}(\text{diam}_2(\mathbb{S}^{d-1}))^2 \\ &\leq \frac{|\mathbb{S}^{d-2}|\pi^2}{|\mathbb{S}^{d-1}|} - \frac{1}{2}(\text{diam}_2(\mathbb{S}^{d-1}))^2 \quad (\because \text{Lemma 6.21}). \end{aligned}$$

Let $C_d := \frac{|\mathbb{S}^{d-2}|\pi^2}{|\mathbb{S}^{d-1}|} - \frac{1}{2}(\text{diam}_2(\mathbb{S}^{d-1}))^2 > 0$. We have just proved that

$$\sum_{n=1}^{\infty} |\lambda_{n,d}| N_{n,d} r^n < C_d \quad \text{for any } r \in (0, 1).$$

Now, to obtain a contradiction, suppose $\sum_{n=1}^{\infty} |\lambda_{n,d}| N_{n,d} = \infty$. Then, we choose a positive integer $m > 0$ such that

$$\sum_{n=1}^m |\lambda_{n,d}| N_{n,d} > 2C_d.$$

Also, choose large enough $r \in (0, 1)$ so that $r > 2^{-\frac{1}{m}}$. Then,

$$\sum_{n=1}^{\infty} |\lambda_{n,d}| N_{n,d} r^n \geq \sum_{n=1}^m |\lambda_{n,d}| N_{n,d} r^n \geq \left(\sum_{n=1}^m |\lambda_{n,d}| N_{n,d} \right) r^m > 2C_d r^m > C_d,$$

which is a contradiction. Hence,

$$\sum_{n=1}^{\infty} |\lambda_{n,d}| N_{n,d} < \infty$$

as we required. □

6.4.3. *Metric transforms of \mathbb{S}^2 .* Suppose $f : [-1, 1] \rightarrow \mathbb{R}$ is a continuous function such that

$$\begin{aligned} d_f : \mathbb{S}^2 \times \mathbb{S}^2 &\longrightarrow \mathbb{R} \\ (u, v) &\longmapsto f(\langle u, v \rangle) \end{aligned}$$

becomes a metric on \mathbb{S}^2 compatible with the canonical topology of \mathbb{S}^2 . Then, this leads to considering the metric measure space $\mathbb{S}_f^2 := (\mathbb{S}^2, d_f, \text{nvols}_{\mathbb{S}^2})$ where the metric on the sphere is now given by d_f instead of the usual euclidean or geodesic metrics. Some examples of such f are snowflake transforms of the euclidean or geodesic distances on \mathbb{S}^2 :

- $f(t) = (\arccos(t))^{\frac{1}{p}}$ for $p \geq 1$ (if $p = 1$, d_f is the geodesic metric on \mathbb{S}^2).
- $f(t) = (\sqrt{2-2t})^{\frac{1}{p}}$ for $p \geq 1$ (if $p = 1$, d_f is the Euclidean metric on \mathbb{S}^2 inherited from \mathbb{R}^d for any $d \geq 2$).

By emulating the argument used in the proof of Theorem 24, we are able to prove the following proposition.³

Proposition 6.22. *Suppose $\mathbb{S}_f^2 \in \mathcal{M}_w$ is given for a continuous map $f : [-1, 1] \rightarrow \mathbb{R}$. Let*

$$\begin{aligned} F : \mathbb{S}^2 &\longrightarrow \mathbb{R} \\ u &\longmapsto d_f^2(e_3, u). \end{aligned}$$

If $\|\Delta F\|_{L^{1+\delta}(\text{vol}_{\mathbb{S}^2})} < \infty$ for some $\delta \in (0, 1)$, then $\mathbb{S}_f^2 \in \mathcal{M}_w^{\text{tr}}$.

Example 6.23. *Let $f(t) = (2-2t)^{\frac{1}{4}}$ and consider \mathbb{S}_f^2 (i.e., d_f is the square root of the Euclidean metric). Also, let*

$$\begin{aligned} F : \mathbb{S}^2 &\longrightarrow \mathbb{R} \\ u &\longmapsto d_f^2(e_3, u). \end{aligned}$$

For this case, we verify that $\|\Delta F\|_{L^{1+\delta}(\text{vol}_{\mathbb{S}^2})} < \infty$ when $\delta = \frac{1}{2}$.

Indeed, it is easy to check that in the coordinate system where $\mathbb{S}^2 \ni u = (\sqrt{1-t^2} \cos \theta, \sqrt{1-t^2} \sin \theta, t)$, where $t \in [-1, 1]$ and $\theta \in [0, 2\pi]$ we have:

$$\|\nabla d_f(e_3, \cdot)(\sqrt{1-t^2} \cos \theta, \sqrt{1-t^2} \sin \theta, t)\| = \frac{\sqrt{1+t}}{2^{\frac{3}{2}}(2-2t)^{\frac{1}{4}}},$$

and

$$\Delta d_f(e_3, \cdot)(\sqrt{1-t^2} \cos \theta, \sqrt{1-t^2} \sin \theta, t) = -\frac{(2-2t)^{\frac{1}{4}}}{4} + \frac{(1+t)}{8(2-2t)^{\frac{3}{4}}}.$$

Therefore,

$$\begin{aligned} \Delta F(\sqrt{1-t^2} \cos \theta, \sqrt{1-t^2} \sin \theta, t) &= 2(\|\nabla d_f(e_3, \cdot)\|^2 + d_f(e_3, \cdot)\Delta d_f(e_3, \cdot)) \\ &= \frac{(1+t)}{4\sqrt{2-2t}} - \frac{\sqrt{2-2t}}{4} \end{aligned}$$

for any $\theta \in \mathbb{R}$ and $t \in (-1, 1)$. The $L^{\frac{3}{2}}(\text{vol}_{\mathbb{S}^2})$ -norm of the second term is obviously finite since $|\frac{\sqrt{2-2t}}{4}| \leq \frac{1}{2}$. For the first term, observe that:

$$\begin{aligned} \int_{\mathbb{S}^2} \left| \frac{(1+t)}{4\sqrt{2-2t}} \right|^{\frac{3}{2}} d\text{vol}_{\mathbb{S}^2}(\sqrt{1-t^2} \cos \theta, \sqrt{1-t^2} \sin \theta, t) &= 2\pi \int_{-1}^1 \left(\frac{(1+t)}{4\sqrt{2-2t}} \right)^{\frac{3}{2}} dt \\ &\leq \frac{\pi}{\sqrt{2}} \int_{-1}^1 \frac{1}{(2-2t)^{\frac{3}{4}}} dt = 2\pi < \infty. \end{aligned}$$

Therefore $\|\Delta F\|_{L^{\frac{3}{2}}(\text{vol}_{\mathbb{S}^2})} < \infty$ as we required. Hence, $\mathbb{S}_f^2 \in \mathcal{M}_w^{\text{tr}}$ by Proposition 6.22.

³Since the proof is essentially the same as the one of Theorem 24, we omit it.

Note that the definitions of d_f and \mathbb{S}_f^{d-1} can both be extended to the case of an arbitrary $d \geq 3$. Unfortunately, the argument via the Poisson identity used in Theorem 25 does not seem applicable to the study \mathbb{S}_f^{d-1} with an arbitrary f . The general classification of f such that $\mathbb{S}_f^{d-1} \in \mathcal{M}_w^{\text{tr}}$ is currently not known. Related questions can be traced back to some classical papers by Von Neumann and Schoenberg such as [VNS41].

6.5. Metric graphs and fractal spaces. In this section we consider a fairly large class of one-dimensional metric spaces called *metric graphs* and prove that they all induce traceable cMDS operators.

The literature on metric graphs in the computational geometry and computational topology is keep growing. In particular, metric graphs are one of the important tools for the shape comparison and the topological data analysis [HSKK01, SK91, BLM21] and in the study of quantum graphs [Kuc08]. Furthermore, the class of metric graphs also includes metric trees, which feature prominently in phylogenetics.

Definition 6.24 ([BBI01, Definition 3.2.11]). A metric graph is the result of gluing of a disjoint collection of segments $\{E_i\}$ and points $\{v_j\}$ (regarded with the length metric of disjoint union) along an equivalence relation R defined on the union of the set $\{v_j\}$ and the set of the endpoints of the segments.

Alternatively, a metric graph can be regarded as a geodesic metric space which is homeomorphic to a topological graph. Note that any compact geodesic metric space can be approximated by finite graphs with respect to the Gromov-Hausdorff distance [BBI01, Theorem 7.5.5].

Recall that a subset Y of a metric space (X, d_X) is said to be ε -net if $X = \cup_{y \in Y} B_\varepsilon(y, X)$.

Definition 6.25 (metric entropy). Let (X, d_X) be a totally bounded metric space. Then, given $\varepsilon > 0$, the **metric entropy** $N_M(\varepsilon, X)$ is the smallest n such that there exists a ε -net $\{x_1, \dots, x_n\}$ for (X, d_X) .

Then, we have the following result.

Theorem 26. *If $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$ satisfies $N_M(\varepsilon, X) \leq C \varepsilon^{-d}$ for some $d \in (0, 2)$, $C > 0$, and for all $\varepsilon \in (0, 1)$, then $\mathfrak{K}_{\mathcal{X}}$ is trace class. Hence, $\mathcal{X} \in \mathcal{M}_w^{\text{tr}}$.*

We now apply this result to metric graphs. Similar consequences follow for certain fractal metric measure spaces such as those considered in [LLL21].

Corollary 6.26. *If $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w$ is a metric graph with uniform probability measure, then \mathcal{X} belongs to $\mathcal{M}_w^{\text{tr}}$.*

Proof. Since \mathcal{X} is a metric graph with the uniform measure, $N_M(\varepsilon, X) \leq C \varepsilon^{-1}$ for some $c > 0$. Hence, if we apply Theorem 26, we are able to completes the proof. \square

In order to prove the above theorem, we need the following theorem from [GBD93] relating traceability and metric entropy.

Theorem 27 ([GBD93, Theorem 1]). *Let $\mathcal{X} = (X, d_X, \mu_X), \mathcal{Y} = (Y, d_Y, \mu_Y) \in \mathcal{M}_w$ be two metric measure spaces, $K \in L^2(X \times Y, \mu_X \otimes \mu_Y)$ where*

$$\mathfrak{K} : L^2(\mu_Y) \rightarrow L^2(\mu_X)$$

is the integral operator generated by K , and $K_Y := \{K(\cdot, y) : y \in Y\} \subseteq L^2(\mu_X)$. Suppose K is such that for some $M < \infty$ and $r < 2$, we have $N_M(\varepsilon, K_Y) \leq M\varepsilon^{-r}$ for any $0 < \varepsilon < 1$. Then \mathfrak{K} is trace class.

Proof of Theorem 26. First, observe that

$$\begin{aligned} & K_{\mathcal{X}}(x, x') - K_{\mathcal{X}}(x, x'') \\ &= -\frac{1}{2} \left(d_X^2(x, x') - d_X^2(x, x'') - \int_X d_X^2(u, x') d\mu_X(u) + \int_X d_X^2(u, x'') d\mu_X(u) \right). \end{aligned}$$

for any $x, x', x'' \in X$. Hence, because of the triangle inequality and the fact that $a^2 - b^2 = (a + b)(a - b)$,

$$|K_{\mathcal{X}}(x, x') - K_{\mathcal{X}}(x, x'')| \leq 2 \operatorname{diam}(X) d_X(x', x'')$$

for any $x, x', x'' \in X$. Hence, $K_{\mathcal{X}}$ is a $(2 \operatorname{diam}(X))$ -Lipschitz function with respect to the second variable. Now, fix arbitrary $\varepsilon \in (0, 1)$. Then, from the assumption, we are able to choose $\{x_1, \dots, x_N\} \subseteq X$ which is an ε -net of X and $N \leq C\varepsilon^{-d}$.

Now, choose arbitrary $x' \in X$. Then, there exist x_i such that $d_X(x', x_i) \leq \varepsilon$. Therefore,

$$\begin{aligned} \|K_{\mathcal{X}}(\cdot, x') - K_{\mathcal{X}}(\cdot, x_i)\|_{L^2(\mu_X)}^2 &= \int_X |K_{\mathcal{X}}(x, x') - K_{\mathcal{X}}(x, x_i)|^2 d\mu_X(x) \\ &\leq 4 \operatorname{diam}(X)^2 (d_X(x', x_i))^2 \end{aligned}$$

Hence,

$$\|K_{\mathcal{X}}(\cdot, x') - K_{\mathcal{X}}(\cdot, x_i)\|_{L^2(\mu_X)} \leq 2 \operatorname{diam}(X) d_X(x', x_i) \leq 2 \operatorname{diam}(X) \varepsilon.$$

This implies that

$$N_M(2 \operatorname{diam}(X) \varepsilon, K_{\mathcal{X}}(\cdot, X)) \leq N_M(\varepsilon, X) \leq C \cdot \varepsilon^{-d}.$$

Hence, finally,

$$N_M(\varepsilon, K_{\mathcal{X}}(\cdot, X)) \leq N_M\left(\frac{\varepsilon}{2 \operatorname{diam}(X)}, X\right) \leq C \cdot (2 \operatorname{diam}(X))^d \cdot \varepsilon^{-d}.$$

So, by [GBD93, Theorem 1], $\mathfrak{K}_{\mathcal{X}}$ is trace class. \square

7. STABILITY OF GENERALIZED cMDS

It is reasonable to expect that, the cMDS process is stable with respect to perturbations of metric measure spaces. The following results confirms that the expectation is true.

Theorem 28 (stability of generalized cMDS). *For any $\mathcal{X} = (X, d_X, \mu_X)$, $\mathcal{Y} = (Y, d_Y, \mu_Y) \in \mathcal{M}_w^{\operatorname{tr}}$,*

$$\inf_{\mu \in \Gamma(\mu_X, \mu_Y)} \|K_{\hat{\mathcal{X}}} - K_{\hat{\mathcal{Y}}}\|_{L^2(\mu \otimes \mu)} \leq 7 \max\{\operatorname{diam}(X), \operatorname{diam}(Y)\} \cdot d_{\operatorname{GW}, 2}(\mathcal{X}, \mathcal{Y}).$$

We need a few more observations to prove the stability theorem.

Theorem 29 ([GR84, Theorem 3.6]). *The (unique) nearest point projection onto a closed convex subset of a Hilbert space is nonexpansive.*

Lemma 7.1. *Let $\mathcal{X}_1 = (X, d_1, \mu_X)$, $\mathcal{X}_2 = (X, d_2, \mu_X) \in \mathcal{M}_w^{\operatorname{tr}}$ be two metric measure spaces with the same underlying set and measure, but different metric. Then*

$$\|K_{\hat{\mathcal{X}}_1} - K_{\hat{\mathcal{X}}_2}\|_{L^2(\mu_X \otimes \mu_X)} \leq \|K_{\mathcal{X}_1} - K_{\mathcal{X}_2}\|_{L^2(\mu_X \otimes \mu_X)}.$$

Proof. Note that $K_{\widehat{\mathcal{X}}_1}$ is the nearest point projection of $K_{\mathcal{X}_1}$ to the closure of $L^2_{\geq 0, \text{sym}}(\mu_X \otimes \mu_X)$ by Theorem 19. Similarly, $K_{\widehat{\mathcal{X}}_2}$ is the nearest point projection of $K_{\mathcal{X}_2}$ to the closure of $L^2_{\geq 0, \text{sym}}(\mu_X \otimes \mu_X)$. Since $L^2_{\geq 0, \text{sym}}(\mu_X \otimes \mu_X)$ is convex, the proof is complete by Theorem 29. \square

Lemma 7.2. *Suppose $\mathcal{X} = (X, d_X, \mu_X), \mathcal{Y} = (Y, d_Y, \mu_Y) \in \mathcal{M}_w$ and a coupling measure $\mu \in \Gamma(\mu_X, \mu_Y)$ are given. Then,*

$$\begin{aligned} & \|K_{\mathcal{X}} - K_{\mathcal{Y}}\|_{L^2(\mu \otimes \mu)} \\ & \leq 3 \max\{\text{diam}(X), \text{diam}(Y)\} \cdot \|d_X - d_Y\|_{L^2(\mu \otimes \mu)} + \max\{\text{diam}(X), \text{diam}(Y)\} \cdot d_{\text{GW},2}(\mathcal{X}, \mathcal{Y}). \end{aligned}$$

Proof. By the definition of $K_{\mathcal{X}}$ and $K_{\mathcal{Y}}$, it is easy to check

$$\begin{aligned} & 2|K_{\mathcal{X}}(x, x') - K_{\mathcal{Y}}(y, y')| \\ & \leq |d_X^2(x, x') - d_Y^2(y, y')| + \left| \int_X d_X^2(x, u) d\mu_X(u) - \int_Y d_Y^2(y, v) d\mu_Y(v) \right| \\ & \quad + \left| \int_X d_X^2(x', u) d\mu_X(u) - \int_Y d_Y^2(y', v) d\mu_Y(v) \right| + |(\text{diam}_2(\mathcal{X}))^2 - (\text{diam}_2(\mathcal{Y}))^2| \end{aligned}$$

for any $x, x' \in X$ and $y, y' \in Y$. Also,

$$\begin{aligned} |d_X^2(x, x') - d_Y^2(y, y')| & = |d_X(x, x') + d_Y(y, y')| \cdot |d_X(x, x') - d_Y(y, y')| \\ & \leq 2 \max\{\text{diam}(X), \text{diam}(Y)\} \cdot |d_X(x, x') - d_Y(y, y')|, \end{aligned}$$

and

$$\begin{aligned} & \left| \int_X d_X^2(x, u) d\mu_X(u) - \int_Y d_Y^2(y, v) d\mu_Y(v) \right| \\ & = \left| \int_{X \times Y} (d_X^2(x, u) - d_Y^2(y, v)) d\mu(u, v) \right| \\ & \leq \int_{X \times Y} |d_X^2(x, u) - d_Y^2(y, v)| d\mu(u, v) \\ & \leq 2 \max\{\text{diam}(X), \text{diam}(Y)\} \cdot \int_{X \times Y} |d_X(x, u) - d_Y(y, v)| d\mu(u, v) \\ & \leq 2 \max\{\text{diam}(X), \text{diam}(Y)\} \cdot \left(\int_{X \times Y} |d_X(x, u) - d_Y(y, v)|^2 d\mu(u, v) \right)^{\frac{1}{2}}, \end{aligned}$$

Similarly

$$\begin{aligned} & \left| \int_X d_X^2(x', u) d\mu_X(u) - \int_Y d_Y^2(y', v) d\mu_Y(v) \right| \\ & \leq 2 \max\{\text{diam}(X), \text{diam}(Y)\} \cdot \left(\int_{X \times Y} |d_X(x', u) - d_Y(y', v)|^2 d\mu(u, v) \right)^{\frac{1}{2}} \end{aligned}$$

for any $x, x' \in X$ and $y, y' \in Y$. Furthermore,

$$\begin{aligned}
|\text{diam}_2^2(\mathcal{X}) - \text{diam}_2^2(\mathcal{Y})| &= 2 \max\{\text{diam}_2(X), \text{diam}_2(Y)\} \cdot |\text{diam}_2(\mathcal{X}) - \text{diam}_2(\mathcal{Y})| \\
&\leq 2 \max\{\text{diam}_2(X), \text{diam}_2(Y)\} \cdot d_{\text{GW},2}(\mathcal{X}, \mathcal{Y}) \\
&\leq 2 \max\{\text{diam}(X), \text{diam}(Y)\} \cdot d_{\text{GW},2}(\mathcal{X}, \mathcal{Y}).
\end{aligned}$$

So, by taking $L^2(\mu \otimes \mu)$ -norm,

$$\begin{aligned}
\|K_{\mathcal{X}} - K_{\mathcal{Y}}\|_{L^2(\mu \otimes \mu)} \\
\leq 3 \max\{\text{diam}(X), \text{diam}(Y)\} \cdot \|d_X - d_Y\|_{L^2(\mu \otimes \mu)} + \max\{\text{diam}(X), \text{diam}(Y)\} \cdot d_{\text{GW},2}(\mathcal{X}, \mathcal{Y}).
\end{aligned}$$

□

Lemma 7.3. *Suppose $\mathcal{X} = (X, d_X, \mu_X), \mathcal{Y} = (Y, d_Y, \mu_Y) \in \mathcal{M}_w$, a coupling measure $\mu \in \Gamma(\mu_X, \mu_Y)$, and the canonical projection*

$$\begin{aligned}
\pi_X : X \times Y &\longrightarrow X \\
(x, y) &\longmapsto x
\end{aligned}$$

are given. Also, let $\mathcal{X}^* := (X \times Y, \pi_X^* d_X, \mu) \in \mathcal{M}_w$. Then,

- (1) $\langle \phi, \psi \rangle_{L^2(\mu_X)} = \langle \pi_X^*(\phi), \pi_X^*(\psi) \rangle_{L^2(\mu)}$ for any $\phi, \psi \in L^2(\mu_X)$,
- (2) $K_{\mathcal{X}^*}((x, y), (x', y')) = K_{\mathcal{X}}(x, x')$ for any $(x, y), (x', y') \in X \times Y$.
- (3) $\phi \in L^2(\mu_X)$ is an eigenfunction of $\mathfrak{K}_{\mathcal{X}}$ with an eigenvalue λ if and only if $\pi_X^*(\phi)$ is an eigenfunction of $\mathfrak{K}_{\mathcal{X}^*}$ with the eigenvalue λ .
- (4) If $\psi \in L^2(\mu) \setminus \pi_X^*(L^2(\mu_X))$ is an eigenfunction of $\mathfrak{K}_{\mathcal{X}^*}$ with an eigenvalue λ , then $\lambda = 0$.
- (5) $K_{\widehat{\mathcal{X}^*}}((x, y), (x', y')) = K_{\widehat{\mathcal{X}}}(x, x')$ for any $(x, y), (x', y') \in X \times Y$.

Proof. (1) Note that $(\pi_X)_*\mu = \mu_X$ because μ is a coupling measure. Now, fix arbitrary $\phi, \psi \in L^2(\mu_X)$. Then,

$$\langle \pi_X^*(\phi), \pi_X^*(\psi) \rangle_{L^2(\mu)} = \int_{X \times Y} \phi(\pi_X(x, y)) \psi(\pi_X(x, y)) d\mu(x, y) = \int_X \phi(x) \psi(x) d\mu_X(x) = \langle \phi, \psi \rangle_{L^2(\mu_X)}.$$

- (2) Obvious, since $\pi_X^* d_X((x, y), (x', y')) = d_X(x, x')$ for any $(x, y), (x', y') \in X \times Y$ and $(\pi_X)_*\mu = \mu_X$.

- (3) By item (2), we have

$$\int_{X \times Y} K_{\mathcal{X}}(\pi(x, y), \pi(x', y')) \phi(\pi(x', y')) d\mu(x', y') = \int_X K_{\mathcal{X}}(x, x') \phi(x') d\mu_X(x')$$

for any $(x, y) \in X \times Y$. Hence, the claim is proved.

- (4) Suppose not. Then, by item (6) of Lemma 4.2, we may assume ψ is continuous. Now, I claim that there are $(x, y), (x, y') \in X \times Y$ such that $\psi(x, y) \neq \psi(x, y')$. Since otherwise, there is a map $\phi : X \longrightarrow \mathbb{R}$ such that $\phi \circ \pi_X = \psi$ (one can define $\phi(x) := \psi(x, y) \forall (x, y) \in X \times Y$). Moreover, $\psi^{-1}(C) = \pi_X^{-1}(\phi^{-1}(C))$ is closed for arbitrary closed set $C \subseteq \mathbb{R}$ since ψ is continuous. Hence, $\phi^{-1}(C) = \pi_X(\psi^{-1}(C))$ is also closed for arbitrary closed set $C \subseteq \mathbb{R}$. Therefore, ϕ is continuous, and $\phi \in L^2(\mu_X)$. This contradicts to the assumption that $\psi \in L^2(\mu) \setminus \pi_X^*(L^2(\mu_X))$. So, we can guarantee that there are $(x, y), (x, y') \in X \times Y$ such that $\psi(x, y) \neq \psi(x, y')$. Then,

$$\begin{aligned}\lambda\psi(x, y) &= \int_{X \times Y} K_{\mathcal{X}}(\pi_X(x, y), \pi_X(x'', y''))\psi(x'', y''') d\mu(x'', y''') \\ &= \int_{X \times Y} K_{\mathcal{X}}(x, x'')\psi(x'', y''') d\mu(x'', y''')\end{aligned}$$

and

$$\begin{aligned}\lambda\psi(x, y') &= \int_{X \times Y} K_{\mathcal{X}}(\pi_X(x, y'), \pi_X(x'', y'''))\psi(x'', y''') d\mu(x'', y''') \\ &= \int_{X \times Y} K_{\mathcal{X}}(x, x'')\psi(x'', y''') d\mu(x'', y''')\end{aligned}$$

Hence, by subtracting the above two equalities, we have

$$\lambda(\psi(x, y) - \psi(x, y')) = 0.$$

Since $\psi(x, y) - \psi(x, y') \neq 0$, this implies $\lambda = 0$. This completes the proof by contradiction.

- (5) By item (3) and (4), every positive eigenvalue and its corresponding eigenfunction of $K_{\mathcal{X}^*}$ comes from $K_{\mathcal{X}}$. So, by the spectral expansion, the proof is complete. \square

Finally, we are ready to prove the stability theorem.

Proof of Theorem 28. Fix arbitrary $\varepsilon > d_{\text{GW},2}(\mathcal{X}, \mathcal{Y})$. Then, there is a coupling measure $\mu \in \Gamma(\mu_X, \mu_Y)$ such that

$$\|d_X - d_Y\|_{L^2(\mu \otimes \mu)} < 2\varepsilon.$$

Let $\pi_X : X \times Y \rightarrow X$ and $\pi_Y : X \times Y \rightarrow Y$ be the canonical projections.

$$\begin{array}{ccc} & X \times Y & \\ \pi_X \swarrow & & \searrow \pi_Y \\ X & & Y \end{array}$$

Let $\mathcal{X}^* := (X \times Y, (\pi_X)^*d_X, \mu)$ and $\mathcal{Y}^* := (X \times Y, (\pi_X)^*d_Y, \mu)$ be two metric measure spaces with the same underlying set and measure, but different metric. Then,

$$\begin{aligned}\|K_{\hat{\mathcal{X}}} - K_{\hat{\mathcal{Y}}}\|_{L^2(\mu \otimes \mu)} &= \|K_{\hat{\mathcal{X}}^*} - K_{\hat{\mathcal{Y}}^*}\|_{L^2(\mu \otimes \mu)} \quad (\because \text{item (5) of Lemma 7.3}) \\ &\leq \|K_{\mathcal{X}^*} - K_{\mathcal{Y}^*}\|_{L^2(\mu \otimes \mu)} \quad (\because \text{Lemma 7.1}) \\ &\leq \|K_{\mathcal{X}} - K_{\mathcal{Y}}\|_{L^2(\mu \otimes \mu)} \quad (\because \text{item (2) of Lemma 7.3}) \\ &\leq 3 \max\{\text{diam}(X), \text{diam}(Y)\} \cdot \|d_X - d_Y\|_{L^2(\mu \otimes \mu)} + \max\{\text{diam}(X), \text{diam}(Y)\} \cdot d_{\text{GW},2}(\mathcal{X}, \mathcal{Y}) \\ &\quad (\because \text{Lemma 7.2}) \\ &< 6 \max\{\text{diam}(X), \text{diam}(Y)\}\varepsilon + \max\{\text{diam}(X), \text{diam}(Y)\} \cdot d_{\text{GW},2}(\mathcal{X}, \mathcal{Y}).\end{aligned}$$

This implies,

$$\inf_{\mu \in \Gamma(\mu_X, \mu_Y)} \|K_{\hat{\mathcal{X}}} - K_{\hat{\mathcal{Y}}}\|_{L^2(\mu \otimes \mu)} < 6 \max\{\text{diam}(X), \text{diam}(Y)\}\varepsilon + \max\{\text{diam}(X), \text{diam}(Y)\} \cdot d_{\text{GW},2}(\mathcal{X}, \mathcal{Y}).$$

Finally, since ε is arbitrary, we have

$$\inf_{\mu \in \Gamma(\mu_X, \mu_Y)} \|K_{\hat{\mathcal{X}}} - K_{\hat{\mathcal{Y}}}\|_{L^2(\mu \otimes \mu)} \leq 7 \max\{\text{diam}(X), \text{diam}(Y)\} \cdot d_{\text{GW},2}(\mathcal{X}, \mathcal{Y}).$$

□

7.1. Consistency. A basic consequence of our stability bounds is that cMDS is consistent for traceable metric measure spaces.

It is useful to define the following notion of convergence of 4-tuples $\mathcal{X} := (X, \Sigma_X, P_X, \mu_X)$ where (X, Σ_X, μ_X) is a measure space endowed with a σ -algebra Σ_X , a probability measure μ_X , and $P_X \in L^2_{\geq 0}(\mu_X \otimes \mu_X)$ is a kernel on X whose associated integral operator is positive semi-definite (see §2.2). We say that a sequence $(\mathcal{X}_n = (X_n, \Sigma_n, P_{X_n}, \mu_{X_n}))_n$ of such 4-tuples *converges weakly* to $\mathcal{X} = (X, \Sigma_X, P_X, \mu_X)$ if for each n there exist a couplings $\mu_n \in \Gamma(\mu_X, \mu_{X_n})$ such that $\|P_X - P_{X_n}\|_{L^2(\mu_n \otimes \mu_n)} \rightarrow 0$ as $n \rightarrow \infty$. We denote this by $P_{\mathcal{X}_n} \xrightarrow{n} P_{\mathcal{X}}$.

Together with Theorem 28, Theorem 5.1 item (e) of [Mém11] imply:

Corollary 7.4. *Let $\mathcal{X} = (X, d_X, \mu_X) \in \mathcal{M}_w^{\text{tr}}$ and let $X_n = X_n(\omega) = \{\mathbf{x}_i(\omega), i = 1, \dots, n\}$ be any collection of i.i.d. random variables chosen from X w.r.t. the measure μ_X for $\omega \in \Omega$, for some probability space Ω . Let $\mathcal{X}_n = \mathcal{X}_n(\omega) = (X_n, d_X|_{X_n \times X_n}, \mu_{X_n})$ be the random metric measure where μ_{X_n} is the uniform measure on X_n . Then, for μ_X -almost every $\omega \in \Omega$:*

$$K_{\hat{\mathcal{X}}_n(\omega)} \xrightarrow{n} K_{\hat{\mathcal{X}}}.$$

This Corollary provides a partial answer to [ABK20, Conjecture 6.1].

REFERENCES

- [ABK20] Henry Adams, Mark Blumstein, and Lara Kassab. Multidimensional scaling on metric measure spaces. *Rocky Mountain Journal of Mathematics*, 50(2):397–413, 2020.
- [AH12] Kendall Atkinson and Weimin Han. *Spherical harmonics and approximations on the unit sphere: an introduction*, volume 2044. Springer Science & Business Media, 2012.
- [Bat53] Harry Bateman. *Higher transcendental functions volume ii*, volume 1. McGraw-Hill Book Company, 1953.
- [BBI01] Dmitri Burago, Yuri Burago, and Sergei Ivanov. *A course in metric geometry*, volume 33. American Mathematical Society Providence, 2001.
- [BBS07] E Bogomolny, O Bohigas, and C Schmit. Distance matrices and isometric embeddings. *arXiv preprint arXiv:0710.2063*, 2007.
- [BH11] Andries E Brouwer and Willem H Haemers. *Spectra of graphs*. Springer Science & Business Media, 2011.
- [BLM21] Ulrich Bauer, Claudia Landi, and Facundo Mémoli. The Reeb graph edit distance is universal. *Foundations of Computational Mathematics*, 21(5):1441–1464, 2021.
- [BN03] Mikhail Belkin and Partha Niyogi. Laplacian eigenmaps for dimensionality reduction and data representation. *Neural computation*, 15(6):1373–1396, 2003.
- [Bri91] Christopher Brislawn. Traceable integral kernels on countably generated measure spaces. *Pacific Journal of Mathematics*, 150(2):229–240, 1991.
- [Bru06] Richard A Brualdi. *Combinatorial matrix classes*, volume 13. Cambridge University Press, 2006.
- [BS13] Ilia Nikolaevich Bronshtein and Konstantin A Semendyayev. *Handbook of mathematics*. Springer Science & Business Media, 2013.
- [CL06] Ronald R Coifman and Stéphane Lafon. Diffusion maps. *Applied and computational harmonic analysis*, 21(1):5–30, 2006.
- [Coh13] Donald L Cohn. *Measure theory*. Springer, 2013.
- [DG03] David L Donoho and Carrie Grimes. Hessian eigenmaps: Locally linear embedding techniques for high-dimensional data. *Proceedings of the National Academy of Sciences*, 100(10):5591–5596, 2003.

- [Fol13] Gerald B Folland. *Real analysis: modern techniques and their applications*. John Wiley & Sons, 2013.
- [GBD93] José M González-Barrios and Richard M Dudley. Metric entropy conditions for an operator to be of trace class. *Proceedings of the American Mathematical Society*, 118(1):175–180, 1993.
- [Gow66] John C Gower. Some distance properties of latent root and vector methods used in multivariate analysis. *Biometrika*, 53(3-4):325–338, 1966.
- [GR84] Kazimierz Goebel and Simeon Reich. *Uniform Convexity, Hyperbolic Geometry, and Nonexpansive Mappings*. CRC Press, 1984.
- [Gro07] Mikhail Gromov. *Metric structures for Riemannian and non-Riemannian spaces*. Springer Science & Business Media, 2007.
- [HHJ94] Roger A Horn, Roger A Horn, and Charles R Johnson. *Topics in matrix analysis*. Cambridge university press, 1994.
- [HLP67] GH Hardy, JE Littlewood, and G Pólya. *Inequalities. Cambridge Mathematical Library Series*. Cambridge University Press, 1967.
- [HSKK01] Masaki Hilaga, Yoshihisa Shinagawa, Taku Kohmura, and Tosiyasu L Kunii. Topology matching for fully automatic similarity estimation of 3d shapes. In *Proceedings of the 28th annual conference on Computer graphics and interactive techniques*, pages 203–212, 2001.
- [Kas19] Lara Kassab. Multidimensional scaling: infinite metric measure spaces. Master’s thesis, Colorado State University, 2019.
- [KST22] Alexey Kroshnin, Eugene Stepanov, and Dario Trevisan. Infinite multidimensional scaling for metric measure spaces. *arXiv preprint arXiv:2201.05885*, 2022.
- [Kuc08] Peter Kuchment. Quantum graphs: an introduction and a brief survey. *arXiv preprint arXiv:0802.3442*, 2008.
- [L⁺18] Russell Lyons et al. Errata to “Distance covariance in metric spaces”. *Annals of Probability*, 46(4):2400–2405, 2018.
- [Lim21] Sunhyuk Lim. *Injective Metric Spaces, through Gromov-type Distances, to Generalized MDS*. PhD thesis, The Ohio State University, 2021.
- [LLL21] Therese-Marie Landry, Michel L Lapidus, and Frederic Latremoliere. Metric approximations of spectral triples on the Sierpiński gasket and other fractal curves. *Advances in Mathematics*, 385:107771, 2021.
- [Lyo13] Russell Lyons. Distance covariance in metric spaces. *The Annals of Probability*, 41(5):3284–3305, 2013.
- [Mar78] Kanti V Mardia. Some properties of classical multi-dimensional scaling. *Communications in Statistics-Theory and Methods*, 7(13):1233–1241, 1978.
- [Mea92] Al Mead. Review of the development of multidimensional scaling methods. *Journal of the Royal Statistical Society: Series D (The Statistician)*, 41(1):27–39, 1992.
- [Mém07] Facundo Mémoli. On the use of Gromov-Hausdorff distances for shape comparison. 2007.
- [Mém11] Facundo Mémoli. Gromov-Wasserstein distances and the metric approach to object matching. *Foundations of computational mathematics*, 11(4):417–487, 2011.
- [MHSG18] Leland McInnes, John Healy, Nathaniel Saul, and Lukas Großberger. Umap: Uniform manifold approximation and projection. *Journal of Open Source Software*, 3(29):861, 2018.
- [MKB79] KV Mardia, JT Kent, and JM Bibby. *Multivariate analysis*. Academic Press, 1979.
- [Pea01] Karl Pearson. Liii. on lines and planes of closest fit to systems of points in space. *The London, Edinburgh, and Dublin philosophical magazine and journal of science*, 2(11):559–572, 1901.
- [Ree12] Michael Reed. *Methods of modern mathematical physics: Functional analysis*. Elsevier, 2012.
- [Ric38] Marion Webster Richardson. Multidimensional psychophysics. *Psychological Bulletin*, 35:659–660, 1938.
- [RR06] Michael Renardy and Robert C Rogers. *An introduction to partial differential equations*, volume 13. Springer Science & Business Media, 2006.
- [RS00] Sam T Roweis and Lawrence K Saul. Nonlinear dimensionality reduction by locally linear embedding. *science*, 290(5500):2323–2326, 2000.
- [Sch35] Isaac J Schoenberg. Remarks to Maurice Frechet’s article “sur la definition axiomatique d’une classe d’espace distances vectoriellement applicable sur l’espace de hilbert. *Annals of Mathematics*, pages 724–732, 1935.

- [SK91] Yoshihisa Shinagawa and Toshiyasu L Kunii. Constructing a Reeb graph automatically from cross sections. *IEEE Computer Graphics and Applications*, 11(06):44–51, 1991.
- [Tor52] Warren S Torgerson. Multidimensional scaling: I. theory and method. *Psychometrika*, 17(4):401–419, 1952.
- [Tor58] Warren S Torgerson. Theory and methods of scaling. 1958.
- [TSL00] Joshua B Tenenbaum, Vin de Silva, and John C Langford. A global geometric framework for nonlinear dimensionality reduction. *science*, 290(5500):2319–2323, 2000.
- [VdMH08] Laurens Van der Maaten and Geoffrey Hinton. Visualizing data using t-SNE. *Journal of machine learning research*, 9(11), 2008.
- [Vil03] Cédric Villani. *Topics in optimal transportation*. Number 58. American Mathematical Soc., 2003.
- [VNS41] John Von Neumann and Isaac Jacob Schoenberg. Fourier integrals and metric geometry. *Transactions of the American Mathematical Society*, 50(2):226–251, 1941.
- [Wof18] Eric Wofsey. Math stackexchange discussion <https://math.stackexchange.com/questions/2914719/question> 2018.
- [YH38] Gale Young and Aiston S Householder. Discussion of a set of points in terms of their mutual distances. *Psychometrika*, 3(1):19–22, 1938.
- [ZZ03] Hongyuan Zha and Zhenyue Zhang. Isometric embedding and continuum isomap. In *Proceedings of the 20th International Conference on Machine Learning (ICML-03)*, pages 864–871, 2003.