

# A INEQUALITY FOR NON-MICROSTATES FREE ENTROPY DIMENSION FOR CROSSED PRODUCTS BY FINITE ABELIAN GROUPS

DIMITRI SHLYAKHTENKO

*To the memory of V.F.R. Jones.*

**ABSTRACT.** For certain generating sets of the subfactor pair  $M \subset M \rtimes G$  where  $G$  is a finite abelian group we prove an approximate inequality between their non-microstates free entropy dimension, resembling the Shreier formula for ranks of finite index subgroups of free groups. As an application, we give bounds on free entropy dimension of generating sets of crossed products of the form  $M \rtimes (\mathbb{Z}/2\mathbb{Z})^{\oplus \infty}$  for a large class of algebras  $M$ .

## 1. INTRODUCTION.

The famous Jones index [Jon83] is the von Neumann algebra extension of the group-subgroup index and is defined for any inclusion  $M_0 \subset M_1$  of  $\text{II}_1$  factors. An open question in von Neumann algebra theory is to find an analog of the Shreier's formula for the number of generators of finite-index subgroups of non-abelian free groups  $\mathbb{F}_n$ . For example, one expects that the “number of generators” of a finite-index subfactor  $M_0 \subset M_1 = L(\mathbb{F}_n)$  should be  $1 + (n - 1)[M : N]$ . Indeed, specific subfactors of  $L(\mathbb{F}_r)$  constructed via amalgamated free products [Pop93, Răd94, PS03, GJS10, GJS11] have sets of generators for which such a formula holds. However, there is little that is known in general, even for index 2.

Returning to group theory, let  $H \subset G$  be a finite-index inclusion of groups. Denoting by  $\beta_j^2(G)$  the  $L^2$ -Betti numbers of  $G$  one has the following generalization of Shreier's formula (see e.g. [Lüc02]):

$$\beta_j^{(2)}(G) = [G : H]^{-1} \beta_j^{(2)}(H).$$

(Schreier's formula corresponds to the case  $j = 1$  and involves the equality  $\beta_1^{(2)}(\mathbb{F}_r) = r - 1$ ). A similar formula is true for finite-index inclusions of tracial algebras [Tho09].

Voiculescu's free entropy dimension  $\delta_0$  [Voi94, Voi96] takes the value  $r$  on a set of generators of  $L(\mathbb{F}_r)$ ; more generally, its value is related to  $L^2$  Betti numbers [CS05, MS05, Shl21]. Therefore one expects a statement of the following kind: given a finite index inclusion  $M_0 \subset M_1$ , for any generating set  $S_0$  for  $M_0$  there exists a generating set for  $M_1$  (and conversely, for any generating set  $S_1$  for  $M_1$  there exists a generating set  $S_0$  for  $M_0$ ) so that

$$(1.1) \quad \delta_0(S_0) - 1 = [M_1 : M_0]^{-1} (\delta_0(S_1) - 1).$$

However, at present only inequalities of the form  $\delta_0(S_1) \leq \delta_0(S_0)$  are available [Jun04].

In this paper we show that for very specific examples of subfactors, namely subfactors of the form  $M_0 \subset M_1 = M_0 \rtimes G$  with  $G$  a finite abelian group, we can find generators  $M_0$  and  $M_1$  for which the non-microstates free entropy dimension [Voi98a] analog of (1.1) holds with an arbitrary small error. More precisely, we prove that given  $\epsilon > 0$  there exist generating sets  $S_0$  for  $M_0$  and  $S_1$  for  $M_1$  for which

$$\delta^*(S_0) - 1 = [M_1 : M_0]^{-1} (\delta^*(S_1) - 1) + \epsilon.$$

Our result is interesting in connection with the following question. Let  $H$  be a finitely generated group, and let  $\alpha$  be an action of some infinite group  $G$  on  $H$ . Then it is known [Gab02, Theorem 6.8] that  $\beta_1^{(2)}(H \rtimes_{\alpha} G) = 0$ . In case that  $G = \mathbb{Z}$  and  $H$  is finitely presented sofic, this implies that the von Neumann algebra  $M_1 = L(H) \rtimes \mathbb{Z}$  is strongly one-bounded [Jun07, Hay15, Shl21]; in particular for any generating  $S$  set of  $M_1$ ,  $\delta_0(S) = 1$ . This leads us to the following conjecture:

**Conjecture 1.** *Let  $M_0$  be a finitely-generated von Neumann algebra, and let  $G$  be an action of a infinite group on  $M_0$ . Then  $M_1 = M_0 \rtimes_{\alpha} G$  is strongly 1-bounded.*

If true, the conjecture has a somewhat surprising consequence: it would imply non-isomorphism of free group factors. Indeed, let  $G$  be any infinite group so that  $L(G)$  is  $R^\omega$ -embeddable (e.g.  $G = \mathbb{Z}$  or  $G$  amenable), and regard  $\mathbb{F}_\infty$  as the infinite free product of copies of  $\mathbb{Z}$  indexed by  $G$ . Then  $G$  acts on this index set by permutations and thus on  $\mathbb{F}_\infty$ ; call this action  $\alpha$ . One can easily see that the resulting semi-direct product is  $\mathbb{Z} * G$ , corresponding to the extension  $e \rightarrow \mathbb{F}_{|G|} \rightarrow \mathbb{Z} * G \xrightarrow{e*id} G \rightarrow e$ . Thus

$$M_1 = M_0 \rtimes_\alpha G = L(\mathbb{F}_\infty \rtimes_\alpha G) = L(\mathbb{Z} * G) \cong L(\mathbb{Z}) * L(G).$$

The latter von Neumann algebra is known to be not strongly 1-bounded; in fact it has a generating set whose free entropy dimension is strictly above 1. On the other hand, if we were to assume that  $L(\mathbb{F}_\infty) \cong L(\mathbb{F}_2)$  (or even finitely-generated), our conjecture (for the specific group  $G$ ) would imply that  $M_1$  is strongly 1-bounded and thus all of its generating sets have free entropy dimension 1, which would be a contradiction.

While we are unable to even come close to proving the conjecture, we are able to show that if  $M_0$  is finitely-generated and in addition  $M_0 \cong M_{2 \times 2}(M_0)$  and  $G = (\mathbb{Z}/2\mathbb{Z})^{\oplus \infty}$  then for any  $\epsilon > 0$ ,  $M_1$  has generating sets with free entropy dimension bounded by  $1 + \epsilon$ . The proof is reminiscent of Gaboriau's proof [Gab00] and it is this connection that inspired us to study the behavior of free entropy dimension under crossed products.

**Acknowledgements.** The author is grateful to B. Hayes for showing him the reference [Gab00] and for useful discussions. This research was supported by NSF grant DMS-2054450.

## 2. ESTIMATES ON NON-MICROSTATES FREE ENTROPY DIMENSION.

**2.1. Special generators for crossed product subfactors.** Let  $M$  be a  $\text{II}_1$  factor and let  $\alpha$  be a properly outer action of a finite abelian group  $G$  on  $M$ . Consider the inclusion of factors

$$M^G \subset M \subset M \rtimes_\alpha G$$

where  $M^G$  is the fixed point algebra for the action  $\alpha$ . It follows from Takesaki duality that  $M \rtimes_\alpha G \cong M_{|G| \times |G|}(M^G)$ , and moreover that the inclusion  $M^G \subset M$  is isomorphic to an inclusion of the form  $M^G \subset M^G \rtimes_{\alpha'} \hat{G}$ , where  $\hat{G}$  is the group dual of  $G$  and  $\alpha'$  is a certain action related to the dual action of  $\hat{G}$  on  $M \rtimes_\alpha G$ .

Assume now that  $M$  is finitely generated; therefore also  $M \rtimes_\alpha G$  is finitely generated. Since  $M_{|G| \times |G|}(M^G) \cong M \rtimes_\alpha G$  we also know that  $M^G$  is finitely generated. Let  $X = (X_1, \dots, X_d)$  be a set of generators for  $M^G$ . Denote by  $\hat{u}_g \in M^G \rtimes \hat{G}$ ,  $g \in \hat{G}$ , the unitaries implementing  $\alpha'$ . Using the isomorphism  $(M^G \subset M) \subset (M^G \subset M^G \rtimes_{\alpha'} \hat{G})$ , we may view these unitaries as elements of  $M$ . Then the set  $X \cup (\hat{u}_g)_{g \in \hat{G}}$  generates  $M$ . Furthermore, if we denote by  $u_g \in M \rtimes_\alpha G$ ,  $g \in G$ , the unitaries implementing  $\alpha$ , then  $X \cup (\hat{u}_g)_{g \in \hat{G}} \cup (u_g)_{g \in G}$  form a generating set of  $M \rtimes_\alpha G$ . Note that we have the following relations:

$$\begin{aligned} u_g X_j u_g^* &= X_j, & g \in G, j = 1, \dots, d \\ u_g v_h u_g^* &= \langle g, h \rangle v_h, & g \in G, h \in \hat{G} \end{aligned}$$

where we use  $\langle \cdot, \cdot \rangle$  to denote the pairing between the elements of  $G$  and its dual  $\hat{G}$ .

**2.2. Estimates on non-microstates free entropy dimension  $\delta^*$ .** In this paper it will be convenient to work with a non-selfadjoint version of free entropy dimension. Given a non-commutative random variables  $Y$  in a tracial von Neumann algebra  $M$  and a sub-algebra  $B \subset M$ , let  $A = *\text{-alg}(Y, B)$  and let  $C$  be a circular element free from  $A$ ; we normalize  $C$  so that  $\tau(C^*C) = 2$ . Consider the derivation  $\partial_Y : A \rightarrow \text{span}\{ACA + AC^*A\}$  determined by the Leibniz rule and by  $\partial_Y(Y) = C$ ,  $\partial_Y(Y^*) = C^*$ ,  $\partial_Y(b) = 0$  for all  $b \in B$ . The vector  $J(Y : B) \in L^2(A, \tau)$ , if it exists, is called the conjugate variable to  $Y$  and is uniquely determined by

$$\langle J(Y : B), P \rangle_{L^2(A)} = \langle C^*, P \rangle_{L^2(A * W^*(C))}, \quad \forall P \in A.$$

Given  $Y_i : i \in I$  the free entropy dimension is then determined by

$$\delta^*(Y_i : i \in I) = 2|I| - \limsup_{t \rightarrow 0} t \|J(Y_i : (Y_j : j \in I \setminus \{i\}))\|_2^2.$$

It is not hard to see that our definition is equivalent to the usual definition of  $\delta^*$  for self-adjoint variables, in that  $\delta^*(Y_i : i \in I) = \delta_{s.a.}^*(\text{Re}Y_i, \text{Im}Y_i : i \in I)$ .

**Lemma 2.** Suppose that  $\alpha$  is an action of a finite abelian group on a tracial von Neumann algebra  $M$ , and suppose that  $Y_j \in M$ ,  $j \in I$  are (not necessarily self-adjoint) generators of  $M$ . Let  $C_j^{(g)}$ ,  $g \in G$  be circular elements  $*$ -free from  $M$ , and extend  $\alpha$  to  $W^*(Y_i : i \in I) * W^*(C_i^{(g)} : i \in I, g \in G)$  by setting  $\alpha_g(C_i^{(g')}) = C_i^{(gg')}$ . Let finally  $Y_j^t = Y_j + \sqrt{t}C_j^{(e)}$ , where  $e \in G$  is the neutral element. Let

$$\xi_j^t = J(Y_j^t : (Y_i^t : i \in I \setminus \{j\}))$$

be the free conjugate variables.

For each  $h \in \hat{G}$ , denote by  $(\bar{C}_j^{(h)})^*$  the projection of  $(C_j^{(e)})^*$  on to the linear subspace of  $\text{span}\{\alpha_g((C_j^{(e)})^*) : g \in G\}$  consisting of vectors  $x$  satisfying  $\alpha_g(x) = \langle g, h \rangle x$ .

Then

$$(2.1) \quad \xi_j^t = |G|^{1/2} t^{-1/2} E_{W^*(Y_i^t : i \in I)}((\bar{C}_j^{(h)})^*), \quad \forall h \in \hat{G}$$

and also

$$\langle \xi_j^t, (\bar{C}_j^{(h)})^* \rangle = \frac{1}{|G|^{1/2}} \langle \xi_j^t, (C_j^{(e)})^* \rangle = \frac{t^{1/2}}{|G|^{1/2}} \|\xi_j^t\|_2^2, \quad \forall j \in I, h \in \hat{G}.$$

In particular, if we denote by  $\bar{\xi}_j^{t,(h)}$  the projection of  $\xi_j^t$  onto the subspace of  $\text{span}\{\alpha_g(\xi_j^t) : g \in G\}$  consisting of vectors  $x$  satisfying  $\alpha_g(x) = \langle g, h \rangle x$ , then

$$\begin{aligned} \|\bar{\xi}_j^{t,(h)}\|_2^2 &= \frac{1}{|G|} \|\xi_j^t\|^2, \quad j \in I, h \in \hat{G} \\ \langle \bar{\xi}_j^{t,(h)}, (C_j^{(e)})^* \rangle &= \frac{t^{1/2}}{|G|^{1/2}} \|\xi_j^t\|_2^2. \end{aligned}$$

so that all of the orthogonal components  $\bar{\xi}_j^{t,(h)}$  in the decomposition  $\xi_j^t = \sum_{h \in \hat{G}} \bar{\xi}_j^{t,(h)}$  have the same length and the same inner product with  $(C_j^{(e)})^*$ . (It is worth noting that  $\bar{\xi}_j^{t,(h)} \notin W^*(Y_j^t : j \in I)$  since that algebra is not invariant under the action  $\alpha$ .)

*Proof.* By [Shl98] we may assume that there exists a family of free creation operators  $\ell_j^{(g)}$ ,  $\hat{\ell}_j^{(g)}$  satisfying, for all  $g, g' \in G$ ,  $j, j' \in I$ ,  $y \in M$ :

$$\begin{aligned} (\ell_j^{(g)})^* y \ell_{j'}^{(g')} &= (\hat{\ell}_j^{(g)})^* y \hat{\ell}_{j'}^{(g')} = \delta_{g=g'} \delta_{j=j'} \tau(y) \\ (\hat{\ell}_j^{(g)})^* y \ell_{j'}^{(g')} &= (\ell_j^{(g)})^* y \hat{\ell}_{j'}^{(g')} = 0 \end{aligned}$$

and so that  $C_j^{(g)} = \ell_j^{(g)} + (\hat{\ell}_j^{(g)})^*$ . The action  $\alpha$  can be extended by putting  $\alpha_g(\ell_j^{(g')}) = \ell_j^{(gg')}$  and  $\alpha_g(\hat{\ell}_j^{(g')}) = \hat{\ell}_j^{(g^{-1}g')}$ . Denote by  $\bar{\ell}_j^{(h)}$  (resp.,  $\bar{\hat{\ell}}_j^{(h)}$ ) the projection of  $\ell_j^{(h)}$  (resp.,  $\hat{\ell}_j^{(h)}$ ) onto the linear subspace of  $\text{span}\{\hat{\ell}_j^{(g)} : g \in G\}$  (resp.,  $\text{span}\{\ell_j^{(g)} : g \in G\}$ ) consisting of vectors  $x$  satisfying  $\alpha_g(x) = \langle g, h \rangle x$ ; in this way we get that  $\bar{C}_j^{(h)} = \bar{\ell}_j^{(h)} + (\bar{\hat{\ell}}_j^{(h)})^*$ . We can now verify the following equations:

$$\begin{aligned} (\bar{\ell}_j^{(h)})^* y \bar{\ell}_{j'}^{(h')} &= (\bar{\hat{\ell}}_j^{(h)})^* y \bar{\hat{\ell}}_{j'}^{(h')} = |G|^{-1} \delta_{h=h'} \delta_{j=j'} \tau(y) \\ (\bar{\hat{\ell}}_j^{(h)})^* y \bar{\ell}_{j'}^{(h')} &= (\bar{\ell}_j^{(h)})^* y \bar{\hat{\ell}}_{j'}^{(h')} = 0 \end{aligned}$$

From this it follows that  $(\bar{C}_j^{(h)} : j \in I, h \in \hat{G})$  is a family of free circular operators (of variance  $|G|^{-1}$ ) which are free from  $M$ . Since for each  $j \in I$ ,  $C_j^{(e)} = \sum_{h \in \hat{G}} \bar{C}_j^{(h)}$ , it follows from [Voi98a] and the equalities

$J(C_j) = C_j^*$ ,  $J(\bar{C}_j^{(h)}) = |G|^{-1/2}(\bar{C}_j^{(h)})^*$  that

$$\begin{aligned}\xi_j^t &= J(Y_j + \sqrt{t}C_j^{(e)} : (Y_i^t : i \in I \setminus \{j\})) \\ &= J((Y_j + \sqrt{t} \sum_{h' \neq h} \bar{C}_j^{(h')}) + \sqrt{t}\bar{C}_j^{(h)} : (Y_i^t : i \in I \setminus \{j\})) \\ &= E_{W^*(Y_j^t : j \in I)} J(\sqrt{t}\bar{C}_j^{(h)} : (Y_j + \sqrt{t} \sum_{h' \neq h} \bar{C}_j^{(h')}) \cup (Y_i^t : i \in I \setminus \{j\})) \\ &= |G|^{1/2} t^{-1/2} E_{W^*(Y_j^t : j \in I)} ((\bar{C}_j^{(h)})^*),\end{aligned}$$

the last equality by freeness. This gives (2.1). On the other hand,

$$\xi_j^t = t^{-1/2} E_{W^*(Y_j^t : j \in I)} (C_j^{(e)}).$$

It follows that

$$\begin{aligned}\langle \xi_j^t, (C_j^{(e)})^* \rangle &= \langle \xi_j^t, E_{W^*(Y_j^t : j \in I)} ((C_j^{(e)})^*) \rangle \\ &= t^{1/2} \langle \xi_j^t, \xi_j^t \rangle \\ &= |G|^{-1/2} \langle \xi_j^t, E_{W^*(Y_j^t : j \in I)} ((\bar{C}_j^{(h)})^*) \rangle \\ &= |G|^{-1/2} \langle \xi_j^t, (\bar{C}_j^{(h)})^* \rangle,\end{aligned}$$

which readily implies the remaining statements of the Lemma.  $\square$

**Theorem 3.** *Let  $X$  be an arbitrary generating set for  $M^G$ , and let  $X \cup (\hat{u}_g)_{g \in \hat{G}}$  be the generating set for  $M$  and  $X \cup (\hat{u}_g)_{g \in \hat{G}} \cup (u_g)_{g \in G}$  be the generating set for  $M \rtimes_\alpha G$  as constructed in §2.1. Then for any  $\epsilon > 0$  there exists a  $\lambda > 0$  so that*

$$(2.2) \quad \delta^*((\lambda X_i : i \in I) \cup (\lambda \hat{u}_h : h \in \hat{G}) \cup (u_g : g \in G) - 1) \leq |G|^{-1}(\delta^*(Y_i : i \in I) - 1) + \epsilon.$$

*Proof.* Let  $I = \{1, \dots, d\} \sqcup \hat{G}$ ; for  $i \in I$  set  $Y_i = X_i$  if  $i \in \{1, \dots, d\}$  and  $Y_i = u_h$  if  $i = h \in \hat{G}$ . Let also  $\omega_i : G \rightarrow \mathbb{C}$  be given by  $\omega_i(g) = 1$  if  $i \in \{1, \dots, d\}$  and  $\omega_i(g) = \langle g, h \rangle$  if  $i = h \in \hat{G}$ . We then have the following relations:

$$u_g Y_i u_g^* = \omega_i(g) Y_i, \quad i \in I, \quad g \in G.$$

For  $i \in I$  let  $C_i$  be a circular system, free from  $W^*(Y_i : i \in I, u_g : g \in G)$ , and for  $g \in G$  let  $C'_g$  be another circular system free from  $W^*(Y_i, C_i : i \in I, u_g : g \in G)$ . We then have:

$$\begin{aligned}\delta^*(X \cup (\hat{u}_h)_{h \in \hat{G}} \cup (u_g)_{g \in G}) &= 2(d + |\hat{G}| + |G|) \\ &\quad - \limsup_{t \rightarrow 0} t \left[ \sum_{j \in I} \|J(Y_j + \sqrt{t}C_j) : (Y_i : i \in I \setminus \{j\}) \cup (u_g + \sqrt{t}C'_g : g \in G)\|_2^2 \right. \\ &\quad \left. + \sum_{g \in G} \|J(u_g + \sqrt{t}C'_g : (Y_i : i \in I \setminus \{j\}) \cup (u_{g'} + \sqrt{t}C_{g'} : g' \in G \setminus \{g\}))\|_2^2 \right].\end{aligned}$$

Denote by  $M_t$  the von Neumann algebra  $W^*(Y_j + \sqrt{t}C_j : j \in I)$  and by  $\hat{M}_t$  the von Neumann algebra  $W^*(M_t, u_g : g \in G)$ .

Using [Voi98a] we note that

$$\begin{aligned}\|J(u_g + \sqrt{t}C'_g : (Y_i : i \in I \setminus \{j\}) \cup (u_{g'} + \sqrt{t}C_{g'} : g' \in G \setminus \{g\}))\|_2^2 \\ \geq \|J(u_g + \sqrt{t}C'_g : (u'_g + \sqrt{t}C_{g'} : g' \in G \setminus \{g\}))\|_2^2\end{aligned}$$

and by [MS05] we get that

$$\begin{aligned}2|G| - \limsup_{t \rightarrow 0} t \|J(u_g + \sqrt{t}C'_g : (u'_g + \sqrt{t}C_{g'} : g' \in G \setminus \{g\}))\|_2^2 \\ = \delta^*(u_g : g \in G) = \beta_1^{(2)}(G) - \beta_0^{(2)}(G) + 1 = 1 - |G|^{-1}.\end{aligned}$$

Let  $\zeta_j^t = J(Y_j + \sqrt{t}C_j : (Y_i : i \in I \setminus \{j\}) \cup (u_g + \sqrt{t}C'_g : g \in G))$  and set  $\xi_j^t = J(Y_j + \sqrt{t}C_j : (Y_i : i \in I \setminus \{j\}))$ . For  $h \in \hat{G}$ , denote by  $\bar{C}_j^{(h)}$  the projection of  $C_j$  onto the subspace of  $\text{span}\{u_g C_j u_g^* : g \in G\}$  consisting of vectors  $x$  so that  $\{u_g(x)u_g^* = \langle g, h \rangle x\}$ . Let also

$$\bar{\zeta}_j^{(h)} = t^{-1/2} E_{\hat{M}_t}((\bar{C}_j^{(h)})^*).$$

Then by applying Lemma 2 with  $C_j^{(e)} = C_j$ ,  $C_j^{(g)} = u_g C_j u_g^*$ , we have

$$\xi_j^t = |G|^{1/2} t^{-1/2} E_{M_t}((\bar{C}_j^{(h)})^*)$$

It follows that

$$\begin{aligned} \langle t^{-1/2} \zeta_j^t, (\bar{C}_j^{(h)})^* \rangle &= \langle E_{\hat{M}_t}(C_j^*), (\bar{C}_j^{(h)})^* \rangle \\ &= \langle C_j^*, E_{\hat{M}_t}(\bar{C}_j^{(h)})^* \rangle \\ &= \langle C_j^*, E_{\hat{M}_t}(|G|^{-1} \sum_{g \in G} \langle h, g \rangle u_g (\bar{C}_j^{(h)})^* u_g^*) \rangle \\ &= \langle C_j^*, E_{\hat{M}_t}(|G|^{-1} \sum_{g \in G} \langle h, g \rangle (u_g + \sqrt{t}C'_g) (\bar{C}_j^{(h)})^* (u_g^* + \sqrt{t}C'_g)) \rangle + O(t^{1/2}) \\ &= \langle C_j^*, |G|^{-1} \sum_{g \in G} \langle h, g \rangle (u_g + \sqrt{t}C'_g) E_{\hat{M}_t}((\bar{C}_j^{(h)})^*) (u_g^* + \sqrt{t}C'_g) \rangle + O(t^{1/2}) \\ &= \langle C_j^*, |G|^{-1} \sum_{g \in G} \langle h, g \rangle (u_g) E_{\hat{M}_t}((\bar{C}_j^{(h)})^*) (u_g^*) \rangle + O(t^{1/2}) \\ &= \langle |G|^{-1} \sum_{g \in G} \langle g, h \rangle u_{g^{-1}} C_j^* u_g, E_{\hat{M}_t}((\bar{C}_j^{(h)})^*) \rangle + O(t^{1/2}) \\ &= \langle (\bar{C}_j^{(h)})^*, E_{\hat{M}_t}((\bar{C}_j^{(h)})^*) \rangle + O(t^{1/2}) \\ &= \|E_{\hat{M}_t}((\bar{C}_j^{(h)})^*)\|_2^2 + O(t^{1/2}) \\ &\geq \|E_{M_t}((\bar{C}_j^{(h)})^*)\|_2^2 + O(t^{1/2}) \\ &= t|G|^{-1} \|\xi_j^t\|_2^2 + O(t^{1/2}). \end{aligned}$$

We now claim that for all by one  $h \in \hat{G}$ ,  $\langle t^{1/2} \zeta_j^t, (\bar{C}_j^{(h)})^* \rangle$  is almost  $2|G|^{-1}$ .

Let us use the notation  $x(t) \in \hat{M}_t + O(t^\gamma)$  to signify that there exists an element  $y(t) \in \hat{M}_t$  so that  $\|x(t) - y(t)\|_2 = O(t^\gamma)$ . Then, since  $(u_g + \sqrt{t}C'_g)(Y_j + \sqrt{t}C_j)(u_g + \sqrt{t}C'_g)^* \in \hat{M}_t$  and

$$\begin{aligned} (u_g + \sqrt{t}C'_g)(Y_j + \sqrt{t}C_j)(u_g + \sqrt{t}C'_g)^* &= u_g Y_j u_g^* + \sqrt{t}(C'_g Y_j u_g^* + u_g Y_j (C'_g)^* + u_g C_j u_g^*) + O(t) \\ &= \omega_j(g) Y_j + \sqrt{t}(C'_g Y_j u_g^* + u_g Y_j (C'_g)^* + u_g C_j u_g^*) + O(t), \end{aligned}$$

we deduce that

$$(u_g + \sqrt{t}C'_g)(Y_j + \sqrt{t}C_j)(u_g + \sqrt{t}C'_g)^* - \omega_j(g)(Y_j + \sqrt{t}C_j) \in \hat{M}_t + O(t).$$

Thus also

$$C'_g Y_j u_g^* + u_g Y_j (C'_g)^* - \omega_j(g) C_j + u_g C_j u_g^* \in \hat{M}_t + O(t).$$

Hence

$$C'_g Y_j u_g^* + u_g Y_j (C'_g)^* - \omega_j(g) C_j + u_g C_j u_g^* \in \hat{M}_t + O(t^{1/2}).$$

Since  $u_g^* u_g = 1$ , we similarly deduce

$$C'_g u_g^* + u_g (C'_g)^* \in \hat{M}_t + O(t^{1/2}),$$

so that, noting that  $u_g \in \hat{M}_t + O(t^{1/2})$

$$(C'_g)^* + u_g^* C'_g u_g^* \in \hat{M}_t + O(t^{1/2}).$$

This gives

$$C'_g Y_j u_g^* - u_g Y_j u_g^* C'_g u_g^* - \omega_j(g) C_j + u_g C_j u_g^* \in \hat{M}_t + O(t^{1/2}).$$

Projecting onto eigenspaces for the  $G$  action (noting again that  $u_g \in \hat{M}_t + O(t^{1/2})$ ) gives us

$$(\bar{C}')_g^{(h \cdot \omega_j)} Y_j u_g^* - u_g Y_j u_g^* \bar{C}'_g^{(h \cdot \omega_j)} u_g^* - \omega_j(g) \bar{C}_j^{(h)} + u_g \bar{C}_j^{(h)} u_g^* \in M_t + O(t^{1/2}).$$

We note also that  $(\bar{C}'_g)^{(h \cdot \omega_j)} u_g^* = \langle h, g \rangle \omega_j(g) \cdot u_g^* (\bar{C}'_g)^{(h)}$  and  $u_g C_j^{(h)} u_g^* = \langle h, g \rangle C_j^{(h)}$  whence

$$(\bar{C}')_g^{(h \cdot \omega_j)} Y_j u_g^* - \langle h, g \rangle \omega_j(g) u_g Y_j u_g^* u_g^* \bar{C}'_g^{(h \cdot \omega_j)} + (\langle h, g \rangle - \omega_j(g)) \bar{C}_j^{(h)} \in M_t + O(t^{1/2}).$$

It follows that

$$\begin{aligned} \langle t^{1/2} \zeta_j^t, (\bar{C}_j^{(h)})^* \rangle &= \langle E_{\hat{M}_t}(C_j), \bar{C}_j^{(h)} \rangle \\ &= \langle C_j, E_{\hat{M}_t}(\bar{C}_j^{(h)}) \rangle \\ &= \sum_{h' \in \hat{G}} \langle \bar{C}_j^{(h')}, E_{\hat{M}_t}(\bar{C}_j^{(h)}) \rangle \\ &= \langle \bar{C}_j^{(h)}, E_{\hat{M}_t}(\bar{C}_j^{(h)}) \rangle + O(t^{1/2}) \\ &= \|E_{\hat{M}_t}(\bar{C}_j^{(h)})\|_2^2 + O(t^{1/2}) \\ &\geq \max_g \left| \left\langle \bar{C}_j^{(h)}, \frac{(\bar{C}')_g^{(h \cdot \omega_j)} Y_j u_g^* - \langle h, g \rangle \omega_j(g) u_g Y_j u_g^* u_g^* \bar{C}'_g^{(h \cdot \omega_j)} + (\langle h, g \rangle - \omega_j(g)) \bar{C}_j^{(h)}}{\|(\bar{C}')_g^{(h \cdot \omega_j)} Y_j u_g^* - \langle h, g \rangle \omega_j(g) u_g Y_j u_g^* u_g^* \bar{C}'_g^{(h \cdot \omega_j)} + (\langle h, g \rangle - \omega_j(g)) \bar{C}_j^{(h)}\|} \right\rangle \right|^2 \\ &\quad + O(t^{1/2}) \\ &= 2|G|^{-1} \max_g \frac{|\omega_j(g) - \langle h, g \rangle|^2}{|\omega_j(g) - \langle h, g \rangle|^2 + 4\|Y_j\|^2} + O(t^{1/2}). \end{aligned}$$

Thus if  $h \neq \omega_j$ , we have

$$\langle t^{1/2} \zeta_j^t, (\bar{C}_j^{(h)})^* \rangle \geq \frac{2|G|^{-1}}{1 + \kappa} + O(t^2),$$

where

$$\kappa = \sup_{j, h} \min_g \frac{4\|Y_j\|_2^2}{|\omega_j(g) - \langle h, g \rangle|^2} = C \sup_j \|Y_j\|_2^2,$$

where  $C$  is some constant that only depends on  $G$ .

We can now compute, using that  $\zeta_j^t = t^{-1/2} E_{\hat{M}_t}(C_j^*)$ ,

$$\begin{aligned} \limsup_{t \rightarrow 0} t \sum_{j \in I} \|J(Y_j + \sqrt{t} C_j) : (Y_i : i \in I \setminus \{j\}) \cup (u_g + \sqrt{t} C'_g : g \in G)\|_2^2 \\ &= \limsup_{t \rightarrow \infty} \langle t^{1/2} \zeta_j^t, t^{1/2} \zeta_j^t \rangle = \limsup_{t \rightarrow \infty} \langle t^{1/2} \zeta_j^t, C_j^* \rangle \\ &= \limsup_{t \rightarrow 0} \sum_{h \in \hat{G}} \sum_{j \in I} \langle t^{1/2} \zeta_j^t, (\bar{C}_j^{(h)})^* \rangle + O(t^{1/2}) \\ &\geq 2|I|(|G| - 1)|G|^{-1}(1 + \kappa)^{-1} + \limsup_{t \rightarrow 0} |G|^{-1} t \|\xi_j^t\|_2^2 + O(t^{1/2}) \\ &\geq 2|I|(|G| - 1)|G|^{-1}(1 + \kappa)^{-1} + 2|I||G|^{-1} - |G|^{-1} \delta^*(Y_i : i \in I) \\ &= 2|I|(1 + \kappa)^{-1} - 2|I||G|^{-1}(1 + \kappa)^{-1} + 2|I||G|^{-1} - |G|^{-1} \delta^*(Y_i : i \in I) \\ &= 2|I|(1 + \kappa)^{-1} - |G|^{-1} \delta^*(Y_i : i \in I) + 2|I||G|^{-1}(1 - (1 + \kappa)^{-1}). \end{aligned}$$

Putting all this together gives us

$$\begin{aligned}
\delta^*((Y_i : i \in I) \cup (u_g : g \in G)) &= 2|I| + 2|G| \\
&- \limsup_{t \rightarrow 0} \left[ \sum_{j \in I} \|J(Y_j + \sqrt{t}C_j) : (Y_i : i \in I \setminus \{j\}) \cup (u_g + \sqrt{t}C'_g : g \in G)\|_2^2 \right. \\
&\quad \left. + \sum_{g \in G} \|J(u_g + \sqrt{t}C'_g) : (Y_i : i \in I \setminus \{j\}) \cup (u_{g'} + \sqrt{t}C_{g'} : g' \in G \setminus \{g\})\|_2^2 \right] \\
&\leq 2|I| + 2|G| - 2|I|(1 + \kappa)^{-1} + |G|^{-1}\delta^*(Y_i : i \in I) - 2|G|^{-1}(1 - (1 + \kappa)^{-1}) - 2|G| + 1 - |G|^{-1} \\
&= |G|^{-1}\delta^*(Y_i : i \in I) + 1 - |G|^{-1} + 2|I|(1 - |G|^{-1})(1 - (1 + \kappa)^{-1}).
\end{aligned}$$

Thus

$$\begin{aligned}
\delta^*((Y_i : i \in I) \cup (u_g : g \in G)) - 1 &\leq |G|^{-1}\delta^*(Y_i : i \in I) - |G|^{-1} + 2|I|(1 - |G|^{-1})(1 - (1 + \kappa)^{-1}) \\
&= |G|^{-1}(\delta^*(Y_i : i \in I) - 1) + 2|I|(1 - |G|^{-1})(1 - (1 + \kappa)^{-1}).
\end{aligned}$$

Suppose now we rescale  $Y_i$  by replacing  $Y_i$  with  $\lambda Y_i$ . Then

$$\delta^*(Y_i : i \in I) = \delta^*(\lambda Y_i : i \in I)$$

so that we get

$$\begin{aligned}
\delta^*(\lambda Y_i : i \in I) \cup (u_g : g \in G) - 1 \\
\leq |G|^{-1}(\delta^*(Y_i : i \in I) - 1) + 2|I|(1 - |G|^{-1})(1 - (1 + \lambda^2 \kappa)^{-1}).
\end{aligned}$$

Thus choosing  $\lambda$  small enough we can ensure that

$$\delta^*(\lambda Y_i : i \in I) \cup (u_g : g \in G) - 1 \leq |G|^{-1}(\delta^*(Y_i : i \in I) - 1) + \epsilon$$

as claimed.  $\square$

Let  $Z_1, \dots, Z_n$  be generators of a tracial von Neumann algebra, it would be natural to expect that  $\delta^*(Z_1, \dots, Z_n)$  is an algebraic invariant: if  $Z'_1, \dots, Z'_{n'}$  is another set of generators for the (non-closed) algebra  $*\text{-alg}(Z_1, \dots, Z_n)$ , then  $\delta^*(Z_1, \dots, Z_n) = \delta^*(Z'_1, \dots, Z'_{n'})$ . In particular, one expects that for any nonzero numbers  $\lambda_1, \dots, \lambda_n$ ,

$$(2.3) \quad \delta^*(Z_1, \dots, Z_n) = \delta^*(\lambda_1 Z_1, \dots, \lambda_n Z_n).$$

If this were true, then we could combine the inequality in Theorem 3 with the equality  $\delta^*((\lambda Y_i : i \in I) \cup (u_g : g \in G)) = \delta^*((Y_i : i \in I) \cup (u_g : g \in G))$  to deduce that  $\delta^*((Y_i : i \in I) \cup (u_g : g \in G)) - 1 \leq |G|^{-1}(\delta^*(Y_i : i \in I) - 1) + \epsilon$  for all  $\epsilon > 0$ , and conclude that  $\delta^*((Y_i : i \in I) \cup (u_g : g \in G)) - 1 \leq |G|^{-1}(\delta^*(Y_i : i \in I) - 1)$ . However, to our embarrassment, we could not find a proof of (2.3). Note, however, that when  $*\text{-alg}(Z_1, \dots, Z_n)$  is isomorphic to a group algebra, then algebraic invariance holds [MS05].

The difficulty in the proof of Theorem 3 arises from the complicated form that the relation  $u_g^* Y_i u_g = \omega_i(g) Y_i$  takes when we substitute  $u_g + \sqrt{t}C'_g$  for  $u_g$  and  $Y_i + \sqrt{t}C_i$  for  $Y_i$ . If we instead redefine  $\hat{M}_t$  as  $W^*(M_t, u_g : g \in G)$  and set  $\zeta_j^t = t^{-1/2} E_{\hat{M}_t}(C_j)$ , then it is easy to show that  $\|\bar{\zeta}_j^{t, (h)}\|_2^2 = 2t^{-1}$  if  $h \neq \omega_i$ . Indeed, since now  $u_g \in \hat{M}_t$  we see that

$$\hat{M}_t \ni u_g(Y_i + \sqrt{t}C_g)u_g^* - \omega_i(g)(Y_i + \sqrt{t}C_g) = \sqrt{t}(u_g C_g u_g^* - \omega_i(g)C_g),$$

so that  $u_g C_g u_g^* - \omega_i(g)C_g \in \hat{M}_t$ . Decomposing into orthogonal components according to  $h \in \hat{G}$  then gives that  $\langle h, g \rangle - \omega_j(g) \bar{C}_j^{(h)} \in \hat{M}_t$  so that  $\bar{C}_j^{(h)} \in \hat{M}_t$  whenever  $h \neq \omega_j$ ; thus  $\bar{\zeta}_j^{t, h} = t^{-1/2} E_{\hat{M}_t}(\bar{C}_j^{(h)}) = t^{-1/2} \bar{C}_j^{(h)}$  and the claimed equality on the norm follows. Using this we can prove the following inequality:

$$\delta^*((Y_i : i \in I) \mid (u_g : g \in G)) + (1 - |G|^{-1}) \leq |G|^{-1}(\delta^*(Y_i : i \in I) - 1)$$

where we define a kind of “relative non-microstates free entropy dimension”  $\delta^*((Y_i : i \in I) \mid (u_g : g \in G)) = 2|I| - \limsup_t t \sum_i \|J(Y_k : (Y_j : j \in I \setminus \{i\}) \cup (u_g : g \in G))\|_2^2$ . We summarize this as:

*Remark 4.* Let  $X$  be an arbitrary generating set for  $M^G$ , and let  $X \cup (\hat{u}_g)_{g \in \hat{G}}$  be the generating set for  $M$  and  $X \cup (\hat{u}_g)_{g \in \hat{G}} \cup (u_g)_{g \in G}$  be the generating set for  $M \rtimes_\alpha G$  as constructed in §2.1. Then

$$\delta^*((Y_i : i \in I) \mid (u_g : g \in G)) + (1 - |G|^{-1}) \leq |G|^{-1}(\delta^*(Y_i : i \in I) - 1).$$

### 3. SOME APPLICATIONS.

**Corollary 5.** *Suppose that  $G$  is a finite abelian group acting properly outer on a factor  $M$ . Suppose that  $M^G$  is generated by  $d$  elements. Then  $M \rtimes_\alpha G$  has a generating set  $S$  satisfying  $\delta_0(S) \leq (2d + 2)|G|^{-1} + 1$ .*

*Proof.* If  $|G| = 1$ , there is nothing to prove, so let us assume that  $|G| \geq 2$ . By Theorem 3 we have, for every  $\epsilon > 0$  the existence of  $\lambda > 0$  so that

$$\delta^*(\lambda X \cup (\lambda \hat{u}_g)_{g \in \hat{G}} \cup (u_g)_{g \in G}) - 1 \leq |G|^{-1} \cdot [\delta^*(X \cup (\hat{u}_h)_{h \in \hat{G}}) - 1] + \epsilon.$$

where  $X$  is any generating set for  $M^G$ . We can thus assume that  $|X| \leq d$ .

$$\begin{aligned} \delta^*(X \cup (\hat{u}_h)_{h \in \hat{G}}) &\leq \delta^*(X) + \delta^*((\hat{u}_h)_{h \in \hat{G}}) \\ &\leq 2d + 1 - |\hat{G}|^{-1}. \end{aligned}$$

since  $\hat{G}$  is abelian [MS05].

Let  $\epsilon = |\hat{G}|^{-1}$ . Then by Theorem 3 there exists some  $\lambda > 0$  so that if we set  $S = X \cup (\hat{u}_g)_{g \in \hat{G}} \cup (u_g)_{g \in G}$  so that (2.2) holds. Combining this with the remarkable inequality between microstates and non-microstates free entropy [BCG03] and invariance of  $\delta_0$  under algebraic changes of variables [Voi98b] we obtain:

$$\begin{aligned} \delta_0(S) &= \delta_0(\lambda X \cup (\lambda \hat{u}_g)_{g \in \hat{G}} \cup (u_g)_{g \in G}) \\ &\leq \delta^*(\lambda X \cup (\lambda \hat{u}_g)_{g \in \hat{G}} \cup (u_g)_{g \in G}) \\ &\leq |G|^{-1}(2d + 1 - |\hat{G}|^{-1}) + 1 + \epsilon = 2d|G|^{-1} + |G|^{-1} - |G|^{-2} + 1 + |G|^{-1} \\ &\leq (2d + 2)|G|^{-1} + 1 = (2d + 2)|G|^{-1} + 1. \end{aligned}$$

as claimed.  $\square$

**Theorem 6.** *Let  $M$  be a finitely generated factor, and assume that  $M \cong M_{2 \times 2}(M)$  and that  $\alpha$  is a properly outer action of  $G = (\mathbb{Z}/2\mathbb{Z})^{\oplus \infty}$  on  $M$ . Then for every  $\epsilon > 0$  there exists a finite generating set  $S$  for  $M \rtimes_\alpha G$  so that  $\delta_0(S) \leq 1 + \epsilon$ .*

*Proof.* Suppose that  $M$  is generated by  $d$  elements. Choose  $m$  so that  $2^{-m}(2d + 2m + 2) < \epsilon$ .

Denote by  $G_m$  the subgroup of  $G$  generated by first  $m$  copies of  $\mathbb{Z}/2\mathbb{Z}$ . Then  $M^{G_m} \cong N \rtimes G_m$  where  $N$  is a  $\text{II}_1$  factor so that  $M_{2^m \times 2^m}(N) \cong M$ . Thus by assumption  $N \cong M$ , so  $N$  can be generated by  $d$  elements. Thus  $M^{G_m}$  is generated by a set  $S'$  of at most  $d + m$  elements. Applying now Corollary 5 we deduce that there exists a generating set  $S_1$  for  $M \rtimes G_m$  so that  $\delta_0(S_1) \leq 2^{-m}(2d + 2m + 2) + 1$ ; moreover,  $S_1$  includes the set  $S_0$  consisting of generators of  $G_m$ .

$G$  is an infinite abelian group whose topological dual is isomorphic to the Cantor set. Thus the von Neumann algebra of  $G$  is generated by a single unitary  $w$ . Let  $S_2$  be that unitary. Then  $S = S_0 \cup S_1 \cup S_2$  is a generating set for  $M \rtimes_\alpha G$ .

By the hyperfinite inequality for  $\delta_0$  [Jun06], we have

$$\begin{aligned} \delta_0(S) &= \delta_0(S_1 \cup S_2 \cup S_0) \leq \delta_0(S_1 \cup S_0) + \delta_0(S_2 \cup S_0) - \delta_0(S_0) \\ &\leq 2^{-m}(2d + 2m + 2) + 1 + 1 - \delta_0(S_0). \end{aligned}$$

Since  $S_0$  generates  $G_m$  which is a finite abelian group,  $\delta_0(S_0) = 1 - |G_m|^{-1} = 1 - 2^{-m}$  [Jun06, Voi96]. Substituting this into the inequality above gives

$$\delta_0(S) \leq 2^{-m}(2d + 2m + 2) + 1 - (1 - 2^{-m}) = 2^{-m}(2d + 2m + 2) + 1 < 1 + \epsilon$$

as claimed.  $\square$



## REFERENCES

- [BCG03] P. Biane, M. Capitaine, and A. Guionnet. Large deviation bounds for matrix Brownian motion. *Invent. Math.*, 152(2):433–459, 2003.
- [CS05] A. Connes and D. Shlyakhtenko.  $L^2$ -homology for von Neumann algebras. *J. Reine Angew. Math.*, 586:125–168, 2005.
- [Gab00] D. Gaboriau. Cout des relations d’équivalence et des groupes. *Invent. Math.*, 139(1):41–98, 2000.
- [Gab02] D. Gaboriau. Invariants  $\ell^2$  de relations d’équivalence et de groupes. *Publ. Math. Inst. Hautes Études Sci.*, 95:93–150, 2002.
- [GJS10] A. Guionnet, V.F.R. Jones, and D. Shlyakhtenko. Random matrices, free probability, planar algebras and subfactors. In *Quanta of maths*, Clay Math. Proc., 11, pages 201–239. Amer. Math. Soc., Providence, RI, 2010.
- [GJS11] A. Guionnet, V.F.R. Jones, and D. Shlyakhtenko. A semi-finite algebra associated to a planar algebra. *J. Func. Anal.*, 261:1345–1360S, 2011.
- [Hay15] B. Hayes. 1-bounded entropy and regularity problems in von Neumann algebras. Preprint arXiv:1505.06682, 2015.
- [Jon83] V.F.R. Jones. Index for subfactors. *Invent. Math.*, 72:1–25, 1983.
- [Jun04] K. Jung. Some free entropy dimension inequalities for subfactors. Preprint arXiv:math/0410594, 2004.
- [Jun06] K. Jung. A hyperfinite inequality for free entropy dimension. *Proc. Amer. Math. Soc.*, 134(7):2099–2108 (electronic), 2006.
- [Jun07] K. Jung. Strongly 1-bounded von Neumann algebras. *Geom. Funct. Anal.*, 17(4):1180–1200, 2007.
- [Lüc02] W. Lück.  $L^2$ -invariants: theory and applications to geometry and  $K$ -theory, volume 44 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 2002.
- [MS05] I. Mineyev and D. Shlyakhtenko. Non-microstates free entropy dimension for groups. *Geom. Func. Anal.*, 15:476–490, 2005.
- [Pop93] S. Popa. Markov traces on universal Jones algebras and subfactors of finite index. *Invent. Math.*, 111:375–405, 1993.
- [PS03] S. Popa and D. Shlyakhtenko. Universal properties of  $L(\mathbf{F}_\infty)$  in subfactor theory. *Acta Math.*, 191(2):225–257, 2003.
- [Răd94] F. Rădulescu. Random matrices, amalgamated free products and subfactors of the von Neumann algebra of a free group, of noninteger index. *Invent. math.*, 115:347–389, 1994.
- [Shl98] D. Shlyakhtenko. Some applications of freeness with amalgamation. *J. reine angew. Math.*, 500:191–212, 1998.
- [Shl21] D. Shlyakhtenko. Von neumann algebras of sofic groups with  $\beta_1^{(2)} = 0$  are strongly 1-bounded. *J. Operator Theory*, 85:217–228, 2021.
- [Tho09] Andreas Thom.  $L^2$ -Betti numbers for subfactors. *J. Operator Theory*, 61(2):295–299, 2009.
- [Voi94] D.-V. Voiculescu. The analogues of entropy and of Fisher’s information measure in free probability theory II. *Invent. Math.*, 118:411–440, 1994.
- [Voi96] D.-V. Voiculescu. The analogues of entropy and of Fisher’s information measure in free probability theory, III. *Geometric and Functional Analysis*, 6:172–199, 1996.
- [Voi98a] D.-V. Voiculescu. The analogues of entropy and of Fisher’s information measure in free probability, V. *Invent. Math.*, 132:189–227, 1998.
- [Voi98b] D.-V. Voiculescu. A strengthened asymptotic freeness result for random matrices with applications to free entropy. *IMRN*, 1:41 – 64, 1998.

UCLA DEPARTMENT OF MATHEMATICS, UCLA, LOS ANGELES, CA 90095  
 Email address: shlyakht@math.ucla.edu