

# Asymptotic theory in a class of directed networks with a differentially private bi-degree sequence

Lu Pan

Jianwei Hu

Peiyan Li

Central China Normal University \*

## Abstract

Although the asymptotic properties of the parameter estimator have been derived in the  $p_0$  model for directed graphs with the differentially private bi-degree sequence, asymptotic theory in general models is still lacking. In this paper, we release the bi-degree sequence of directed graphs via the discrete Laplace mechanism, which satisfies differential privacy. We use the moment method to estimate the unknown parameter. We establish a unified asymptotic result, in which consistency and asymptotic normality of the differentially private estimator hold. We apply the unified theoretical result to the Probit model. Simulations and the real data demonstrate our theoretical findings.

**Key words:** Asymptotic normality; Consistency; Differential privacy; Directed networks; Moment estimation.

## 1 Introduction

As more and more network data are being made public, data privacy has received wide attention because data may contain sensitive information about individuals and their relationships (e.g., sexual relationships, e-mail exchanges). Using anonymous or non-anonymous nodes to publish these sensitive data may cause serious privacy problems and even lead to legal proceedings. For example, [Jernigan and Mistree \(2009\)](#) successfully

---

\*Department of Statistics, Central China Normal University, Wuhan, 430079, China. **Emails:** pan-lulu@mails.ccn.u.edu.cn. jwhu@mail.ccn.u.edu.cn. lipeiyan@mail.ccn.u.edu.cn.

predicted the sexual orientation of Facebook users by using their friendships' public information. It has been proven that this anonymous method can expose privacy through re-identification technology [e.g., [Hay et al. \(2008\)](#); [Narayanan and Shmatikov \(2009\)](#)].

[Dwork et al. \(2006\)](#) developed a rigorous privacy standard, called *differential privacy*, to achieve privacy protection. It guarantees that changes to one person's data will not significantly affect the output distribution. The differential privacy is a good framework for privacy protection, which reduces the leakage of privacy risks and ensures the utility of the data, and has been widely used as a privacy standard when releasing network data [e.g., [Hay et al. \(2009\)](#); [Lu and Miklau \(2014\)](#); [Karwa and Slavković \(2016\)](#)].

In many cases, the degree sequence is the only information available and many other important properties are constrained by it. However, the degree may carry confidential and sensitive information, such as the sexually transmitted disease [[Helleringer and Kohler \(2007\)](#)]. To solve it, we can add noises into degrees. For example, [Hay et al. \(2009\)](#) used the Laplace mechanism to release the degree partition, and proposed an efficient algorithm to find the solution. [Karwa and Slavković \(2016\)](#) used a discrete Laplace mechanism to release the degree sequence, derived differentially private estimators of parameters in the  $\beta$ -model, and proved that they are consistent and asymptotically normally distributed. These works are concerned with undirected graphs. In directed graphs, [Yan \(2021\)](#) proved that differentially private estimators of parameters in the  $p_0$  model are consistent and asymptotically normally distributed.

In this paper, we aim to establish the unified asymptotic theory in a class of directed random graph models with the differentially private bi-degree sequence. We release bi-degree sequences by adding discrete Laplacian noises and using the moment estimation to estimate the unknown parameters. This is inspired by [Yan \(2021\)](#). In a general class of directed random graph models, we establish the consistency and asymptotic normal distribution of the differentially private estimator as the number of nodes goes to infinity. In the case of nondifferential private, [Fan \(2023\)](#) established the unified theoretical framework for directed graphs with bi-degree sequence. Our work is about differential private, which is different from [Fan \(2023\)](#).

The rest of this article is organized as follows. Section 2 introduces the necessary background of differential privacy and presents the estimation of noisy bi-degree sequence based on the moment equation, and obtains unified asymptotic properties for the differentially private estimation as the number of nodes goes to infinity. Section 3 illustrates our theoretical result by the Probit model. Section 4 carries out the simulations and real data under the Probit model. Some further discussion is given in Section 5. All proofs are deferred to the appendix section.

## 2 Main results

Consider a simple directed graph  $G_n$  with no multiple edges and no self-loops on  $n$  nodes labeled by “1, ...,  $n$ ”. Let  $A = (a_{i,j})_{n \times n}$  be the adjacency matrix of  $G_n$ , where  $a_{i,j}$  is a variable of the directed edge from node  $i$  to node  $j$ . Define  $d_i^+ = \sum_{j \neq i} a_{i,j}$  as the out-degree of node  $i$  and  $\mathbf{d}^+ = (d_1^+, \dots, d_n^+)^\top$  as the out-degree sequence of the graph  $G_n$ . Similarly, define  $d_j^- = \sum_{i \neq j} a_{i,j}$  as the in-degree of node  $j$  and  $\mathbf{d}^- = (d_1^-, \dots, d_n^-)^\top$  as the in-degree sequence. The pair  $(\mathbf{d}^+, \mathbf{d}^-)$  or  $\{(d_1^+, d_1^-), \dots, (d_n^+, d_n^-)\}$  is called the bi-degree sequence.

Assume that the probability mass function of the edge weights  $a_{i,j}$  ( $i \neq j$ ) have the following general form [Fan (2023)]:

$$a_{i,j} = a | \alpha_i, \beta_j \sim f((\alpha_i + \beta_j)a), \quad (2.1)$$

where  $f(\cdot)$  is a probability mass function,  $\alpha_i$  denotes the strength parameter of node  $i$  from an outgoing edge and  $\beta_j$  denotes the strength parameter of node  $j$  from an incoming edge. We note that the value of  $f(\cdot)$  in Equation (2.1) is invariant under the transforms  $(\boldsymbol{\alpha}, \boldsymbol{\beta})$  to  $(\boldsymbol{\alpha} - c, \boldsymbol{\beta} + c)$  for a constant  $c$ . For the model identification, we set  $\beta_n = 0$  hereafter.

### 2.1 Differential privacy

Consider two databases  $D$  and  $D'$ , which differ only in a single element. In addition, we consider a randomized mechanism  $Q$  that takes  $D$  as input and outputs a sanitized database  $S = (S_1, \dots, S_k)$  for public use. In general, the size of  $S$  may not be the same as  $D$ . The mechanism  $Q(\cdot|D)$  defines a conditional probability distribution of the output  $S$  given  $D$ . Let  $\epsilon$  be a positive real number and  $\mathcal{S}$  denote the sample space of  $Q$ . This mechanism is differentially private if the results of  $Q(\cdot|D)$  and  $Q(\cdot|D')$  are almost indistinguishable for every choice of  $D$  and  $D'$ .

More formally, the data releasing mechanism  $Q$  is  $\epsilon$ -differentially private if for all subsets of output  $B \subset \mathcal{S}$  and databases  $D$  and  $D'$  [Dwork et al. (2006)],

$$Q(S \in B|D) \leq e^\epsilon \times Q(S \in B|D'),$$

where  $\epsilon$  is the privacy level that denotes the level of privacy guarantees such that a smaller value means more privacy protection.

Differential privacy requires that the distribution of the output is almost the same whether or not an individual's record appears in database. Within network data, depending on the definition of the graph neighbor, *differential privacy* is divided into *node*

*differential privacy* [Kasiviswanathan et al. (2013)] and *edge differential privacy* [Nissim et al. (2007)]. *Difference privacy* is *edge difference privacy* if two graphs are called neighbors when they differ in exactly one edge. Analogously, we can define *node differential privacy* by letting graphs be neighbors if one can be obtained from the other by removing a node and its adjacent edges. In this paper, we focus on edge differential privacy [Hay et al. (2009)]. Let  $\delta(G, G')$  denote the number of different edges of graphs  $G$  and  $G'$ . The formal definition of edge differential privacy is as follows.

**Definition 1.** (Edge Differential Privacy). *Let  $\epsilon > 0$  be a privacy parameter. A randomized mechanism  $Q(\cdot|G)$  is  $\epsilon$ -edge differentially private if*

$$\sup_{G, G' \in \mathcal{G}, \delta(G, G')=1} \sup_{S \in \mathcal{S}} \frac{Q(S|G)}{Q(S|G')} \leq e^\epsilon,$$

where  $\mathcal{G}$  is the set of all directed graphs of interest on  $n$  nodes and  $\mathcal{S}$  is the set of all possible outputs.

Let  $f : \mathcal{G} \rightarrow \mathbb{R}^k$  be a function,  $\Delta f$  represents the global sensitivity of the function  $f$ . The magnitude of the noise added in the differentially private algorithm  $Q$  mainly depends on the global sensitivity. If the outputs are the network statistics, then a simple algorithm to guarantee edge difference privacy is the Laplace mechanism [Dwork et al. (2006)], which adds the Laplace noise proportional to the global sensitivity of  $f$ . When  $f(G)$  is integer, we can use a discrete Laplace random variables as the noise as in Karwa and Slavković (2016).

**Lemma 1.** (Discrete Laplace Mechanism). *Let  $f : \mathcal{G} \rightarrow \mathbb{R}^k$ . The discrete Laplace mechanism works by adding noise to  $f$ :*

$$Q(\cdot|G) = f(G) + (e_1, \dots, e_k),$$

where  $e_1, \dots, e_k$  be independent and identically distributed (i.i.d.) discrete Laplace random variables with probability mass function defined as follows:

$$\mathbb{P}(X = x) = \frac{1 - \lambda}{1 + \lambda} \lambda^{|x|}, \quad x \in \{0, \pm 1, \dots\}, \lambda \in (0, 1).$$

The discrete Laplace mechanism is  $\epsilon$ -edge differentially private, where  $\epsilon = -\Delta(f) \log \lambda$ , and  $\Delta(f)$  is the  $L_1$  sensitivity of  $f$  and is defined as:

$$\Delta(f) = \max_{\delta(G, G')=1} \|f(G) - f(G')\|_1,$$

where  $\|\cdot\|_1$  is the  $L_1$ -norm.

**Post-Processing:** Differentially private mechanisms are immune to post-processing. This means that any function with a differentially private mechanism will remain differentially private. More formally, if a mechanism  $f$  is  $\epsilon$ -differentially private and  $g$  is any function. Then  $g(f(G))$  is also  $\epsilon$ -differentially private. This result indicates that any post-processing done on the noisy bi-degree sequences obtained as an output of a differentially private mechanism is also differentially private.

## 2.2 Differential privacy in directed random graph models

We use the discrete Laplace mechanism in Lemma 1 to release the bi-degree sequence  $\mathbf{d} = (\mathbf{d}^+, \mathbf{d}^-)$  to guarantee edge differential privacy. Since adding or removing a directed edge will increase or decrease the degrees of two corresponding nodes by one each. Therefore, the global sensitivity of the bi-degree sequence is two.

---

**Algorithm 1** *Input:* A graph  $G_n$  and privacy parameter  $\epsilon$ . *Output:* Differentially private answer to the sequence of  $G_n$ .

---

```

1: Let  $(\mathbf{d}^+, \mathbf{d}^-)$  be the bi-degree sequence of  $G_n$ 
2: for  $i = 1 \rightarrow n$  do
3:   for  $j = 1 \rightarrow n$  do
4:     Generate two independent  $e_i^+$  and  $e_j^-$  from discrete Laplace with  $\lambda = \exp(-\epsilon/2)$ 
5:     Let  $z_i^+ = d_i^+ + e_i^+$  and  $z_j^- = d_j^- + e_j^-$ 
6:   end for
7: end for

```

---

The random variables  $\{e_i^+\}_{i=1}^n$  and  $\{e_j^-\}_{j=1}^n$  are independently generated from symmetric discrete Laplace distribution with the same parameter  $\lambda$ . So  $\mathbb{E}(e_i^+) = \mathbb{E}(e_j^-) = 0$ . Let  $\mu(\cdot)$  denote the expectation of  $f(\cdot)$  and  $\boldsymbol{\theta} = (\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_{n-1})^\top$ . Define the following system of functions:

$$\begin{aligned}
F_i(\boldsymbol{\theta}) &= z_i^+ - \sum_{k=1; k \neq i}^n \mu(\alpha_i + \beta_k), \quad i = 1, \dots, n, \\
F_{n+j}(\boldsymbol{\theta}) &= z_j^- - \sum_{k=1; k \neq j}^n \mu(\alpha_k + \beta_j), \quad j = 1, \dots, n, \\
F(\boldsymbol{\theta}) &= (F_1(\boldsymbol{\theta}), \dots, F_{2n-1}(\boldsymbol{\theta}))^\top.
\end{aligned} \tag{2.2}$$

Obviously, the solution to  $F(\boldsymbol{\theta}) = 0$  is the differentially private estimator of  $\boldsymbol{\theta}$ , which is

induced by the following moment equations:

$$\begin{aligned} z_i^+ &= \sum_{k=1; k \neq i}^n \mu(\hat{\alpha}_i + \hat{\beta}_k), i = 1, \dots, n, \\ z_j^- &= \sum_{k=1; k \neq j}^n \mu(\hat{\alpha}_k + \hat{\beta}_j), j = 1, \dots, n-1, \end{aligned} \quad (2.3)$$

where  $\hat{\boldsymbol{\theta}} = (\hat{\alpha}_1, \dots, \hat{\alpha}_n, \hat{\beta}_1, \dots, \hat{\beta}_{n-1})^\top$  and  $\hat{\beta}_n = 0$ . Since  $\mathbf{z}^+$  and  $\mathbf{z}^-$  satisfy edge differentially privacy,  $\hat{\boldsymbol{\theta}}$  is edge differentially private estimator of  $\boldsymbol{\theta}$  according to post-processing. In the following, we will conduct a rigorous analysis of the asymptotic properties of  $\hat{\boldsymbol{\theta}}$ .

## 2.3 Asymptotic properties of the differentially private estimator

We first state the parameter space and some technical conditions. Let  $\boldsymbol{\theta}^*$  be the true parameter vector satisfying  $-Q_n \leq \alpha_i^* + \beta_j^* \leq Q_n$  ( $1 \leq i \neq j \leq n$ ) for variable  $Q_n$ . We consider the parameter space

$$\Theta = \{\boldsymbol{\theta} : -Q_n - 2r \leq \alpha_i + \beta_j \leq Q_n + 2r, \text{ for all } 1 \leq i \neq j \leq n\}, \quad (2.4)$$

where  $r = \|[F'(\boldsymbol{\theta}^*)]^{-1}F(\boldsymbol{\theta}^*)\|_\infty$ . Let  $\mathbf{g} = (d_1^+, \dots, d_n^+, d_1^-, \dots, d_{n-1}^-)^\top$ ,  $g_{2n} = d_n^-$  and  $\tilde{\mathbf{g}} = (z_1^+, \dots, z_n^+, z_1^-, \dots, z_{n-1}^-)^\top$ ,  $\tilde{g}_{2n} = z_n^-$ . By Fan (2023), the following condition is given. When  $\boldsymbol{\theta} \in \Theta$ , for  $i = 1, \dots, n, j = 1, \dots, n-1, j \neq i$ , the derivatives of  $\mu(\cdot)$  satisfy

$$m \leq \left| \frac{\partial \mu(\alpha_i + \beta_j)}{\partial \alpha_i} \right| = \left| \frac{\partial \mu(\alpha_i + \beta_j)}{\partial \beta_j} \right| \leq M, \quad (2.5)$$

$$\max_{i,j} \left\{ \left| \frac{\partial^2 \mu(\alpha_i + \beta_j)}{\partial \alpha_i^2} \right|, \left| \frac{\partial^2 \mu(\alpha_i + \beta_j)}{\partial \beta_j \partial \alpha_i} \right|, \left| \frac{\partial^2 \mu(\alpha_i + \beta_j)}{\partial \beta_j^2} \right| \right\} \leq \eta_1, \quad (2.6)$$

where  $M$ ,  $m$  and  $\eta_1$  are functions on variable  $Q_n$ . Moreover, we give the following lemma, whose proof is given in the Appendix.

**Lemma 2.** *With probability approaching one,*

$$\max \left\{ \max_{i=1, \dots, n} |z_i^+ - E(z_i^+)|, \max_{j=1, \dots, n} |z_j^- - E(z_j^-)| \right\} \leq \sqrt{n \log n} + \kappa \sqrt{\log n}, \quad (2.7)$$

where  $\kappa = 2(-\log \lambda)^{-1} = 4/\epsilon$ .

Now, we present the existence and consistency of  $\hat{\boldsymbol{\theta}}$  under some mild conditions. This is proved by constructing a Newton iterative sequence:  $\boldsymbol{\theta}^{(k+1)} = \boldsymbol{\theta}^{(k)} - [F'(\boldsymbol{\theta}^{(k)})]^{-1}F(\boldsymbol{\theta}^{(k)})$ . If the true value  $\boldsymbol{\theta}^*$  is chosen as the initial value, we derive the error between  $\boldsymbol{\theta}^*$  and  $\hat{\boldsymbol{\theta}}$ . We state the consistency of  $\hat{\boldsymbol{\theta}}$ , whose proof is given in the Appendix.

**Theorem 1.** *If inequalities (2.5) and (2.6) hold, and*

$$\frac{M^2}{nm^3}(\sqrt{n \log n} + \kappa \sqrt{\log n}) = o(1), \quad (2.8)$$

$$\frac{M^4 \eta_1}{nm^6}(\sqrt{n \log n} + \kappa \sqrt{\log n}) = o(1), \quad (2.9)$$

*then with probability approaching one as  $n$  goes to infinity, the differentially private estimator  $\hat{\boldsymbol{\theta}}$  exists and satisfies*

$$\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_\infty = O_p\left(\frac{M^2}{nm^3}(\sqrt{n \log n} + \kappa \sqrt{\log n})\right) = o_p(1). \quad (2.10)$$

Next we state the asymptotic normality of  $\hat{\boldsymbol{\theta}}$ . Before introducing our main theorem, we need to provide some preliminary results on the covariance matrix of bi-degree sequences. Let  $U = u_{i,j} = \text{Cov}\{\mathbf{g} - E(\mathbf{g})\}$ . It follows that  $\text{Var}(a_{i,j}) = u_{i,j}$  for  $i = 1, \dots, n$  and  $j = 1, \dots, n-1$ . Obviously, if

$$m_u \leq \min_{i,j} \text{Var}(a_{i,j}) \leq \max_{i,j} \text{Var}(a_{i,j}) \leq M_u, \quad (2.11)$$

then  $U \in \mathcal{L}_n(m_u, M_u)$ ,  $u_{2n,2n} = \text{Var}(d_n^-)$  and  $(n-1)m_u \leq u_{i,i} \leq (n-1)M_u$ . If  $M_u/m_u = o(n)$ , then  $nm_u \rightarrow \infty$ . So we have  $u_{i,i} \rightarrow \infty$  as  $n \rightarrow \infty$ . Then we give the following lemma.

**Lemma 3.** *Let  $\kappa = 2(-\log \lambda)^{-1}$ , where  $\lambda = \exp(-\epsilon/2)$ . (i) If  $\frac{\kappa(\log n)^{1/2}}{m_u} = o(1)$  and  $M_u/m_u = o(n)$ , then for any fixed  $k \geq 1$ , as  $n \rightarrow \infty$ , the vector consisting of the first  $k$  elements of  $S(\tilde{\mathbf{g}} - \mathbb{E}\mathbf{g})$  is asymptotically multivariate normal with mean zero and covariance matrix  $Z$ , where  $Z$  is given as follows:*

$$\text{diag}\left(\frac{u_{1,1}}{v_{1,1}^2}, \dots, \frac{u_{k,k}}{v_{k,k}^2}\right) + \left(\frac{u_{2n,2n}}{v_{2n,2n}^2}\right) \mathbf{1}_k \mathbf{1}_k^\top, \quad (2.12)$$

*where  $\mathbf{1}_k$  is a  $k$ -dimensional column vector with all entries 1.*

*(ii) Let*

$$s_n^2 = \text{Var}\left(\sum_{i=1}^n e_i^+ - \sum_{j=1}^{n-1} e_j^-\right) = (2n-1) \frac{2\lambda}{(1-\lambda)^2}.$$

*Assume that  $s_n/v_{2n,2n}^{1/2} \rightarrow c$  for some constant  $c$ . For any fixed  $k \geq 1$ , the vector consisting of the first  $k$  elements of  $S(\tilde{\mathbf{g}} - \mathbb{E}\mathbf{g})$  is asymptotically  $k$ -dimensional multivariate normal distribution with mean zero and covariance matrix  $B = (b_{i,j})_{k \times k}$ , where  $B$  is given as follows:*

$$\text{diag}\left(\frac{u_{1,1}}{v_{1,1}^2}, \dots, \frac{u_{k,k}}{v_{k,k}^2}\right) + \left(\frac{u_{2n,2n}}{v_{2n,2n}^2} + \frac{s_n^2}{v_{2n,2n}^2}\right) \mathbf{1}_k \mathbf{1}_k^\top, \quad (2.13)$$

where  $\mathbf{1}_k$  is a  $k$ -dimensional column vector with all entries 1.

We prove Lemma 3 in the Appendix. Finally, we state the asymptotic normality of the differentially private estimator  $\widehat{\boldsymbol{\theta}}$ , as shown below.

**Theorem 2.** *If inequalities (2.6), (2.10) hold and*

$$\frac{(n + 2\kappa n^{1/2} + \kappa^2)M^6\eta_1 \log n}{m^9 n^{3/2}} = o(1), \quad (2.14)$$

(i) if  $\frac{\kappa(\log n)^{1/2}}{m_u} = o(1)$ , then for any fixed  $k \geq 1$ , as  $n \rightarrow \infty$ , the vector consisting of the first  $k$  elements of  $(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*)$  is asymptotically multivariate normal with mean zero and covariance matrix  $Z$ , defined in (2.12).

(ii) then for any fixed  $k \geq 1$ , the vector consisting of the first  $k$  elements of  $(\widehat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*)$  is asymptotically  $k$ -dimensional multivariate normal distribution with mean zero and covariance matrix  $B$ , defined in (2.13).

**Remark 1.** It is meaningful to compare the above theorem with Theorem 2.2 in Fan (2023). The key differences in that the asymptotic variance of  $\widehat{\theta}_i$  has an additional variance factor  $s_n^2/v_{2n,2n}^2$ . This is due to the fact that they only consider nondifferential private case. The asymptotic expression of  $\widehat{\theta}_i$  contains a term  $\sum_{i=1}^n e_i^+ - \sum_{j=1}^{n-1} e_j^-$ . Its variance is in the magnitude of  $ne^{-\epsilon/2}$ . When  $\epsilon$  becomes small, the variance increases quickly, such that its impact on  $\widehat{\theta}_i$  cannot be ignored when it increases to a certain level. This leads to the appearance of the additional variance factor.

### 3 Application

In this section, we provide application of the unified theoretical result to the Probit model satisfying (2.1). The Probit model can be formulated using an array of mutually independent Bernoulli random variables  $a_{i,j}$  ( $1 \leq i \neq j \leq n$ ), with probability mass function:

$$P(a_{i,j} = 1) = \int_{-\infty}^{\alpha_i + \beta_j} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx = \Phi(\alpha_i + \beta_j), \quad (3.1)$$

where  $\Phi(\cdot)$  is the cumulative distribution function of the standard normal distribution. In this case, we get

$$\mu(\alpha_i + \beta_j) = \int_{-\infty}^{\alpha_i + \beta_j} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}} dx.$$

By direct calculations, we have

$$\frac{\partial \mu(\alpha_i + \beta_j)}{\partial \alpha_i} = \frac{\partial \mu(\alpha_i + \beta_j)}{\partial \beta_j} = \frac{1}{\sqrt{2\pi}} e^{-\frac{(\alpha_i + \beta_j)^2}{2}}.$$



The function  $h(x) = \exp(-x^2/2)$  is symmetric on  $x$  and is a decreasing function when  $x \geq 0$ . So we have the following inequalities,

$$\frac{1}{\sqrt{2\pi}}e^{-\frac{(Q_n+2r)^2}{2}} \leq \left| \frac{\partial \mu(\alpha_i + \beta_j)}{\partial \alpha_i} \right| \leq \frac{1}{\sqrt{2\pi}}, \frac{1}{\sqrt{2\pi}}e^{-\frac{Q_n^2}{2}} \leq \left| \frac{\partial \mu(\alpha_i + \beta_j)}{\partial \alpha_i^*} \right| \leq \frac{1}{\sqrt{2\pi}}.$$

Thus  $F'(\boldsymbol{\theta}^*) \in \mathcal{L}_n(m^*, M^*)$ , where  $M^* = \frac{1}{\sqrt{2\pi}}, m^* = \frac{1}{\sqrt{2\pi}}e^{-\frac{Q_n^2}{2}}$ . Following [Yan \(2021\)](#) and [Wang et al. \(2020\)](#), we gain the following inequality holds,

$$\max\{\max_i |z_i^+ - \mathbb{E}(z_i^+)|, \max_j |z_j^- - \mathbb{E}(z_j^-)|\} \leq \sqrt{n \log n} + \kappa \sqrt{\log n}. \quad (3.2)$$

Again, by (3.2) and Lemma 5 in the Appendix, if  $\kappa = o_p(n^{1/2})$ , we obtain

$$r = O\left(e^{3Q_n^2/2} \sqrt{\frac{\log n}{n}}\right).$$

If  $e^{3Q_n^2/2} = o((n/\log n)^{1/2})$ , then  $r \rightarrow 0$  as  $n$  goes to infinity. Thus  $r$  can be small enough to ignore, for any  $\boldsymbol{\theta} \in \Omega(\boldsymbol{\theta}^*, 2r)$ , we obtain  $F'(\boldsymbol{\theta}) \in \mathcal{L}_n(m, M)$ , where

$$M = \frac{1}{\sqrt{2\pi}}, m = \frac{1}{\sqrt{2\pi}}e^{-\frac{Q_n^2}{2}}. \quad (3.3)$$

On the other hand, for any  $i = 1, \dots, n, j = 1, \dots, n-1, j \neq i$ ,

$$\frac{\partial^2 \mu(\alpha_i + \beta_j)}{\partial \alpha_i^2} = \frac{\partial^2 \mu(\alpha_i + \beta_j)}{\partial \beta_j^2} = \frac{\partial^2 \mu(\alpha_i + \beta_j)}{\partial \alpha_i \partial \beta_j} = \frac{\partial^2 \mu(\alpha_i + \beta_j)}{\partial \beta_j \partial \alpha_i} = -\frac{(\alpha_i + \beta_j)e^{-\frac{(\alpha_i + \beta_j)^2}{2}}}{\sqrt{2\pi}}.$$

The function  $g(x) = -x \exp(-x^2/2)$  is decreasing function when  $-1 < x < 1$ ; otherwise, it is the increasing function. Moreover, the function value is negative when  $x > 0$  and the function value is positive when  $x < 0$ . So, the function value reaches the maximum  $e^{-1/2}$  when  $x = -1$ . Then, we have

$$\left| -\frac{\alpha_i + \beta_j}{\sqrt{2\pi}}e^{-\frac{(\alpha_i + \beta_j)^2}{2}} \right| \leq \frac{1}{\sqrt{2\pi}e}.$$

Thus,  $\eta_1 = \frac{1}{\sqrt{2\pi}e}$ . If  $e^{Q_n^2} = o((n/\log n)^{1/6})$  and  $\kappa = o_p(n^{1/2})$ ,

$$\frac{M^4 \eta_1}{nm^6}(\sqrt{n \log n} + \kappa \sqrt{\log n}) = O\left(e^{3Q_n^2} \sqrt{\frac{\log n}{n}}\right) = o(1),$$

where  $M$  and  $m$  are given in (3.3), then (2.10) is satisfied. By Theorem 1, the consistency of  $\hat{\boldsymbol{\theta}}$  is as follows.

**Corollary 1.** *If  $e^{Q_n^2} = o((n/\log n)^{1/6})$  and  $\kappa = o_p(n^{1/2})$ , then with probability approaching one as  $n$  goes to infinity, the differentially private estimator  $\hat{\boldsymbol{\theta}}$  exists and satisfies*

$$\|\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*\|_\infty = O_p\left(e^{3Q_n^2/2} \sqrt{\frac{\log n}{n}}\right) = o_p(1). \quad (3.4)$$

Again, note that both  $d_i^+ = \sum_{j \neq i} a_{i,j}$  and  $d_j^- = \sum_{j \neq i} a_{i,j}$  are sums of  $n-1$  independent random variables, respectively. It can be shown that  $U = \text{Cov}\{\mathbf{g} - \mathbb{E}(\mathbf{g})\} \in \mathcal{L}_n(m_u, M_u)$ , where

$$m_u = \Phi(-Q_n)(1 - \Phi(-Q_n)), M_u = 1/4.$$

Since  $\Phi(x)(1 - \Phi(x))$  is an increasing function on  $x$  when  $x \leq \Phi^{-1}(1/2)$  and a decreasing function when  $x \geq \Phi^{-1}(1/2)$ , we have

$$(n-1)\Phi(-Q_n)(1 - \Phi(-Q_n)) \leq u_{i,i} \leq \frac{n-1}{4}, \quad i = 1, \dots, 2n.$$

If  $e^{Q_n^2/2} = o(n^{1/18}/(\log n)^{1/9})$  and  $\kappa = o_p(n^{1/2})$ , then

$$\frac{M^6 \eta_1 \log n (n + 2\kappa n^{1/2} + \kappa^2)}{m^9 n^{3/2}} = O\left(\frac{e^{9Q_n^2/2} \log n}{n^{1/2}}\right) = o(1).$$

By Theorem 2 and Lemma 3, the asymptotic normality of  $\hat{\boldsymbol{\theta}}$  is as below.

**Corollary 2.** *Assume that  $e^{Q_n^2/2} = o(n^{1/18}/(\log n)^{1/9})$  and  $\kappa = o_p(n^{1/2})$ .*

(i) *If  $\frac{\kappa(\log n)^{1/2}}{m_u} = o(1)$ , then for any fixed  $k \geq 1$ , as  $n \rightarrow \infty$ , the vector consisting of the first  $k$  elements of  $(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*)$  is asymptotically multivariate normal with mean zero and covariance matrix  $Z$ .*

(ii) *Let*

$$s_n^2 = \text{Var}\left(\sum_{i=1}^n e_i^+ - \sum_{j=1}^{n-1} e_j^-\right) = (2n-1) \frac{2\lambda}{(1-\lambda)^2}.$$

*Assume that  $s_n/v_{2n,2n}^{1/2} \rightarrow c$  for some constant  $c$ . For any fixed  $k \geq 1$ , the vector consisting of the first  $k$  elements of  $(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*)$  is asymptotically  $k$ -dimensional multivariate normal distribution with mean  $\mathbf{0}$  and covariance matrix  $B$ .*

## 4 Simulations

In this section, we evaluate the asymptotic results for Probit model (3.1) of directed graph model (2.1) through numerical simulations. Similar to Yan et al. (2016), we determine the parameter values using a linear form. In particular, we set  $\alpha_{i+1}^* = (n-1-i)L/(n-1)$  for  $i = 0, \dots, n-1$ ; let  $\beta_i^* = \alpha_i^*$  for  $i = 1, \dots, n-1$  for simplicity and  $\beta_n^* = 0$  by default. We consider three different values for  $L$ ,  $L = 0$ ,  $\log(\log n)$  and  $(\log n)^{1/2}$ , respectively. We simulate two different values for  $\epsilon$ ,  $\epsilon = 2$  and  $\log(n)/n^{1/4}$ . We carry out simulations under two different sizes of networks:  $n = 100$  and  $n = 200$ . Under each simulation setting, 10,000 datasets are generated.

By Corollary 2,

$$\begin{aligned}\hat{\xi}_{i,j} &= [\hat{\alpha}_i - \hat{\alpha}_j - (\alpha_i^* - \alpha_j^*)]/(\hat{z}_{i,i} + \hat{z}_{j,j})^{1/2}, \\ \hat{\zeta}_{i,j} &= [\hat{\alpha}_i + \hat{\beta}_j - (\alpha_i^* + \beta_j^*)]/(\hat{z}_{i,i} + \hat{z}_{n+j,n+j})^{1/2}, \\ \hat{\eta}_{i,j} &= [\hat{\beta}_i - \hat{\beta}_j - (\beta_i^* - \beta_j^*)]/(\hat{z}_{n+i,n+i} + \hat{z}_{n+j,n+j})^{1/2}\end{aligned}$$

are all asymptotically distributed as standard normal random variables, where  $\hat{z}_{i,i}$  is the estimate of  $z_{i,i}$  by replacing  $\theta_i^*$  with  $\hat{\theta}_i$ . Three special pairs  $(1, 2)$ ,  $(n/2, n/2 + 1)$  and  $(n-1, n)$  for  $(i, j)$  are discussed.

We apply the quantile-quantile (QQ) plot to demonstrate the asymptotic normality of  $\hat{\xi}_{i,j}$ ,  $\hat{\zeta}_{i,j}$ ,  $\hat{\eta}_{i,j}$  and the results are similar, thus we here only show the QQ-plot for  $\hat{\xi}_{i,j}$  under the case of  $n = 100$  and  $\epsilon = 2$  in Figure 1 to save space. From this Figure, we see that the empirical quantiles coincide well with the ones of the standard normality for estimate.




Figure 1: The QQ plot of  $\hat{\xi}_{i,j}$ . The horizontal and vertical axes are the theoretical and empirical quantiles, respectively, and the red lines correspond to the reference lines  $y = x$ .

Table 1 reports the coverage frequencies of the 95% confidence interval for  $\alpha_i - \alpha_j$ , the length of the confidence interval, and the frequency that the estimate does not exist. As expected, the length of the confidence interval increases as  $L$  increases and decreases as  $n$  increases.

Table 1: Estimated coverage probabilities ( $\times 100\%$ ) of  $\alpha_i - \alpha_j$  for pair  $(i, j)$  as well as the length of confidence intervals (in square brackets), and the probabilities ( $\times 100\%$ ) that the estimates do not exist (in parentheses).

n	$(i, j)$	$L = 0$	$L = \log(\log n)$	$L = (\log n)^{1/2}$
$\epsilon = 2$				
100	(1, 2)	93.80[0.36](0)	93.61[0.41](0.06)	92.63[0.46](1.13)
	(50, 51)	93.49[0.36](0)	92.78[0.48](0.06)	91.28[0.57](1.13)
	(99, 100)	93.96[0.36](0)	92.73[0.54](0.06)	91.14[0.64](1.13)
200	(1, 2)	94.32[0.25](0)	94.00[0.29](0)	94.29[0.33](0)
	(100, 101)	94.64[0.25](0)	93.61[0.35](0)	92.97[0.41](0)
	(199, 200)	94.66[0.25](0)	93.94[0.39](0)	93.02[0.46](0)
$\epsilon = \log n / n^{1/4}$				
100	(1, 2)	92.37[0.36](0)	92.40[0.41](0.49)	90.07[0.46](4.46)
	(50, 51)	92.43[0.36](0)	90.37[0.48](0.49)	87.70[0.57](4.46)
	(99, 100)	92.58[0.36](0)	89.92[0.54](0.49)	88.06[0.65](4.46)
200	(1, 2)	94.19[0.25](0)	93.31[0.30](0)	93.16[0.33](0.07)
	(100, 101)	93.92[0.25](0)	92.47[0.35](0)	90.18[0.41](0.07)
	(199, 200)	93.71[0.25](0)	91.56[0.39](0)	91.14[0.46](0.07)

*A data example.* We evaluate the use of the proposed estimator on the Lazega’s friendship network data [Lazega et al. (2001); Snijders et al. (2006)]. This dataset contains 71 nodes, and each node represents an attorney. A directed edge from one attorney to another exists if the former considers the later a friend outside of work. In total, there are 575 edges. In-degrees and out-degrees vary from 0 to 22 and 25, respectively. The data set contains 71 nodes with 575 directed edges.

We chose the privacy parameter  $\epsilon$  as 1. Table 2 reports the estimators of  $\alpha_i$  and  $\beta_j$  with their estimated standard errors, and their corresponding in bi-degree sequences. The estimators of parameters for out-degrees range from the minimum  $-9.41$  to maximum  $-0.45$  while those of parameters for in-degrees range from the minimum  $-7.96$  to maximum  $0.97$ .

## 5 Discussion

We have established the asymptotic theory in a class of directed random graph model parameterized by the differentially private bi-sequence and illustrated application to the Probit model. The result shows that statistical inference can be made using the noisy bi-sequence. We assume that the edges are mutually independent in this work. We

Table 2: Lazega’s friendship network dataset: the differentially private estimator of  $\alpha_i$  and  $\beta_j$  with their standard errors and their corresponding bi-degree sequence.

Vertex	$\hat{\alpha}_i$	$\sqrt{\hat{\sigma}_{i,i}}$	$\hat{\beta}_j$	$\sqrt{\hat{\sigma}_{j,j}}$	$d_i^+$	$d_j^-$	Vertex	$\hat{\alpha}_i$	$\sqrt{\hat{\sigma}_{i,i}}$	$\hat{\beta}_j$	$\sqrt{\hat{\sigma}_{j,j}}$	$d_i^+$	$d_j^-$	Vertex	$\hat{\alpha}_i$	$\sqrt{\hat{\sigma}_{i,i}}$	$\hat{\beta}_j$	$\sqrt{\hat{\sigma}_{j,j}}$	$d_i^+$	$d_j^-$
1	-1.61	0.07	0.10	0.06	4	5	25	-1.21	0.04	0.35	0.04	11	10	49	-1.85	0.08	0.00	0.06	4	6
2	-1.84	0.07	0.34	0.04	4	10	26	-1.42	0.04	0.97	0.03	9	22	50	-1.43	0.07	0.19	0.07	8	8
3	-9.41	0.43	-0.24	0.07	0	4	27	-1.08	0.04	0.78	0.03	13	17	51	-1.52	0.05	0.19	0.05	6	7
4	-0.97	0.03	0.61	0.03	15	14	28	-1.09	0.04	0.35	0.04	12	9	52	-1.20	0.04	0.61	0.03	11	14
5	-1.85	0.08	0.00	0.06	3	5	29	-1.28	0.05	0.35	0.05	10	10	53	-2.00	0.08	-7.96	0.44	3	0
6	-9.41	0.43	-0.60	0.11	0	2	30	-1.62	0.05	-0.11	0.06	6	5	54	-1.51	0.05	0.42	0.04	7	11
7	-2.20	0.11	-0.60	0.11	2	2	31	-0.45	0.03	0.74	0.03	25	15	55	-9.41	0.43	-0.39	0.08	0	3
8	-2.19	0.43	0.18	0.05	1	7	32	-1.72	0.07	0.19	0.05	4	7	56	-1.43	0.04	0.35	0.04	8	10
9	-1.71	0.06	0.54	0.03	6	14	33	-1.22	0.04	-0.59	0.08	12	3	57	-1.35	0.04	0.48	0.04	9	12
10	-1.04	0.03	-0.22	0.07	14	4	34	-1.84	0.05	0.27	0.04	6	11	58	-1.03	0.03	0.55	0.04	13	12
11	-1.71	0.05	0.60	0.03	5	14	35	-1.35	0.04	0.35	0.04	9	10	59	-1.85	0.06	-0.39	0.07	5	4
12	-0.73	0.03	0.02	0.04	22	8	36	-1.28	0.04	0.48	0.04	9	11	60	-1.84	0.07	0.19	0.05	4	8
13	-1.08	0.03	0.88	0.03	14	20	37	-2.20	0.03	-0.39	0.05	0	1	61	-1.85	0.08	-0.24	0.08	3	3
14	-1.43	0.04	0.01	0.05	8	6	38	-1.61	0.05	0.41	0.04	8	13	62	-2.00	0.07	-0.24	0.06	4	5
15	-1.73	0.08	-0.39	0.17	4	2	39	-1.43	0.04	0.54	0.03	8	13	63	-2.20	0.17	-7.96	0.08	2	0
16	-1.43	0.04	0.35	0.04	8	10	40	-1.28	0.04	0.19	0.05	10	8	64	-0.71	0.03	0.73	0.03	19	15
17	-0.56	0.03	0.84	0.03	23	18	41	-1.20	0.04	0.72	0.03	12	17	65	-0.54	0.03	0.37	0.05	22	8
18	-1.44	0.04	-0.23	0.05	9	5	42	-0.92	0.03	0.36	0.04	15	9	66	-0.98	0.03	-0.58	0.17	16	3
19	-2.00	0.07	-0.39	0.07	4	4	43	-0.86	0.03	0.67	0.03	15	13	67	-2.20	0.07	-0.92	0.08	4	3
20	-1.09	0.03	0.20	0.05	12	7	44	-9.41	0.04	-7.96	0.04	0	0	68	-1.52	0.05	0.01	0.06	6	5
21	-1.27	0.05	0.77	0.03	8	15	45	-1.62	0.05	-0.23	0.07	6	4	69	-1.72	0.07	-0.23	0.08	5	4
22	-1.44	0.05	0.01	0.06	8	6	46	-1.85	0.08	0.00	0.06	3	5	70	-1.52	0.04	-0.10	0.05	7	5
23	-2.51	0.11	0.10	0.05	1	7	47	-9.40	0.04	-7.96	0.05	0	0	71	-2.51	0.11	0.00	0.05	1	6
24	-0.58	0.03	0.79	0.03	23	17	48	-1.62	0.05	-0.39	0.07	7	4							

should be able to obtain consistent conclusion if the edges are dependent, provided that the conditions stated in Theorem 1 are met. However, the asymptotic normality of the estimator is not clear. To avoid this problem, we need appropriately select a probability distribution for directed random graphs when using the existing method. In the further, we may relax our theoretical conditions to ignore the independence of edges.

## Appendix

We start with some notations. Let  $\mathbb{R}_+ = (0, \infty)$ ,  $\mathbb{R}_0 = [0, \infty)$ ,  $\mathbb{N} = \{1, 2, \dots\}$ ,  $\mathbb{N}_0 = \{0, 1, 2, \dots\}$ . For a vector  $\mathbf{x} = (x_1, \dots, x_n)^\top \in \mathbb{R}^n$ , denote by  $\|\mathbf{x}\|_\infty = \max_{1 \leq i \leq n} |x_i|$  the  $\ell_\infty$ -norm of  $\mathbf{x}$ . For an  $n \times n$  matrix  $J = (J_{i,j})$ ,  $\|J\|_\infty$  denotes the matrix norm induced by the  $\|\cdot\|_\infty$ -norm on vectors in  $\mathbb{R}^n$ :

$$\|J\|_\infty = \max_{\mathbf{x} \neq 0} \frac{\|J\mathbf{x}\|_\infty}{\|\mathbf{x}\|_\infty} = \max_{1 \leq i \leq n} \sum_{j=1}^n |J_{i,j}|.$$

Define the matrix maximum norm  $\|\cdot\|$  for a matrix  $A = (a_{i,j})$  by  $\|A\| := \max_{i,j} |a_{i,j}|$ .

Given  $m, M > 0$ , we say a  $(2n-1) \times (2n-1)$  matrix  $V = (v_{i,j})$  belongs to the matrix

class  $\mathcal{L}_n(m, M)$  if  $V$  satisfies the following conditions:

$$\begin{aligned} m &\leq v_{i,i} - \sum_{j=n+1}^{2n-1} v_{i,j} \leq M, i = 1, \dots, n-1; \quad v_{n,n} = \sum_{j=n+1}^{2n-1} v_{n,j}, \\ v_{i,j} &= 0, i, j = 1, \dots, n, i \neq j, \\ v_{i,j} &= 0, i, j = n+1, \dots, 2n-1, i \neq j, \\ m &\leq v_{i,j} = v_{j,i} \leq M, i = 1, \dots, n, j = n+1, \dots, 2n-1, j \neq n+i, \\ v_{i,n+i} &= v_{n+i,i} = 0, i = 1, \dots, n-1, \\ v_{i,i} &= \sum_{k=1}^n v_{k,i} = \sum_{k=1}^n v_{i,k}, i = n+1, \dots, 2n-1. \end{aligned}$$

Obviously, if  $V \in \mathcal{L}_n(m, M)$ , then  $V$  is a  $(2n-1) \times (2n-1)$  diagonally dominant, symmetric nonnegative matrix and  $V$  has the following structure:

$$V = \begin{pmatrix} V_{11} & V_{12} \\ V_{12}^\top & V_{22} \end{pmatrix},$$

where  $n \times n$  matrix  $V_{11}$  and  $(n-1) \times (n-1)$  matrix  $V_{22}$  are diagonal matrices,  $V_{12}$  is a nonnegative matrix whose nondiagonal elements are positive and diagonal elements equal to zero.

Define  $v_{2n,i} = v_{i,2n} := v_{i,i} - \sum_{j=1; j \neq i}^{2n-1} v_{i,j}$ , for  $i = 1, \dots, 2n-1$  and  $v_{2n,2n} = \sum_{i=1}^{2n-1} v_{2n,i}$ . Then  $m \leq v_{2n,i} \leq M$  for  $i = 1, \dots, n-1$ ,  $v_{2n,i} = 0$  for  $i = n, n+1, \dots, 2n-1$  and  $v_{2n,2n} = \sum_{i=1}^n v_{i,2n} = \sum_{i=1}^n v_{2n,i}$ . Generally, the inverse of  $V$ ,  $V^{-1}$ , does not have a closed form. [Yan et al. \(2016\)](#) proposed a matrix  $S = (s_{i,j})$  to approximate  $V^{-1}$ , where

$$s_{i,j} = \begin{cases} \frac{\delta_{i,j}}{v_{i,i}} + \frac{1}{v_{2n,2n}}, & i, j = 1, \dots, n, \\ -\frac{1}{v_{2n,2n}}, & i = 1, \dots, n, j = n+1, \dots, 2n-1, \\ -\frac{1}{v_{2n,2n}}, & i = n+1, \dots, 2n-1, j = 1, \dots, n, \\ \frac{\delta_{i,j}}{v_{i,i}} + \frac{1}{v_{2n,2n}}, & i, j = n+1, \dots, 2n-1, \end{cases}$$

$\delta_{i,j}$  is the Kronecker delta function. Rewrite  $S$  as

$$S = \begin{pmatrix} S_{11} & S_{12} \\ S_{12}^\top & S_{22} \end{pmatrix},$$

where  $n \times n$  matrix  $S_{11} = 1/v_{2n,2n} + \text{diag}(1/v_{1,1}, 1/v_{2,2}, \dots, 1/v_{n,n})$ ,  $S_{12}$  is an  $n \times (n-1)$  matrix whose elements are all equal to  $-1/v_{2n,2n}$  and  $(n-1) \times (n-1)$  matrix  $S_{22} = 1/v_{2n,2n} + \text{diag}(1/v_{n+1,n+1}, 1/v_{n+2,n+2}, \dots, 1/v_{2n-1,2n-1})$ .

## A.1 Proof of Theorem 1

We apply a Newton iterative theorem of [Yan et al. \(2016\)](#) to prove the existence and consistency of the differentially private estimator  $\hat{\boldsymbol{\theta}}$ .

**Proposition 1.** [[Yan et al. \(2016\)](#)]. *Define a system of equations*

$$\begin{aligned} F_i(\boldsymbol{\theta}) &= d_i - \sum_{k=1; k \neq i}^n \mu(\alpha_i + \beta_k), i = 1, \dots, n, \\ F_j(\boldsymbol{\theta}) &= b_j - \sum_{k=1; k \neq j}^n \mu(\alpha_k + \beta_j), j = 1, \dots, n-1, \\ F(\boldsymbol{\theta}) &= (F_1(\boldsymbol{\theta}), \dots, F_n(\boldsymbol{\theta}), F_{n+1}(\boldsymbol{\theta}), \dots, F_{2n-1}(\boldsymbol{\theta}))^\top, \end{aligned}$$

where  $\mu(\cdot)$  is a continuous function with the third derivative. Let  $\Theta \subset \mathbb{R}^{2n-1}$  be a convex set and assume for any  $\mathbf{x}, \mathbf{y}, \mathbf{v} \in \Theta$ ,

$$\begin{aligned} \| [F'(\mathbf{x}) - F'(\mathbf{y})] \mathbf{v} \|_\infty &\leq K_1 \| \mathbf{x} - \mathbf{y} \|_\infty \| \mathbf{v} \|_\infty, \\ \max_{i=1, \dots, 2n-1} \| F'_i(\mathbf{x}) - F'_i(\mathbf{y}) \|_\infty &\leq K_2 \| \mathbf{x} - \mathbf{y} \|_\infty, \end{aligned}$$

where  $F'(\boldsymbol{\theta})$  is the Jacobin matrix of  $F$  on  $\boldsymbol{\theta}$  and  $F'_i(\boldsymbol{\theta})$  is the gradient function of  $F_i$  on  $\boldsymbol{\theta}$ . Consider  $\boldsymbol{\theta}^{(0)} \in \Theta$  with  $\Omega(\boldsymbol{\theta}^{(0)}, 2r) \subset \Theta$ , where  $r = \|[F(\boldsymbol{\theta}^{(0)})]^{-1}F(\boldsymbol{\theta}^{(0)})\|_\infty$ . For any  $\boldsymbol{\theta} \in \Omega(\boldsymbol{\theta}^{(0)}, 2r)$ , we assume

$$F'(\boldsymbol{\theta}) \in \mathcal{L}_n(m, M) \text{ or } -F'(\boldsymbol{\theta}) \in \mathcal{L}_n(m, M).$$

For  $k = 1, 2, \dots$ , define the Newton iterates  $\boldsymbol{\theta}^{(k+1)} = \boldsymbol{\theta}^{(k)} - [F'(\boldsymbol{\theta}^{(k)})]^{-1}F(\boldsymbol{\theta}^{(k)})$ . let

$$\rho = \frac{c_1(2n-1)M^2K_1}{2m^3n^2} + \frac{K_2}{(n-1)m}.$$

If  $\rho r < 1/2$ , then  $\boldsymbol{\theta}^{(k)} \in \Omega(\boldsymbol{\theta}^{(0)}, 2r)$ ,  $k = 1, 2, \dots$ , are well defined and satisfy

$$\|\boldsymbol{\theta}^{(k+1)} - \boldsymbol{\theta}^{(0)}\|_\infty \leq r/(1 - \rho r).$$

Further,  $\lim_{k \rightarrow \infty} \boldsymbol{\theta}^{(k)}$  exists and the limiting point is precisely the solution of  $F(\boldsymbol{\theta}) = 0$  in the range of  $\boldsymbol{\theta} \in \Omega(\boldsymbol{\theta}^{(0)}, 2r)$ .

Besides, we need lemmas of [Yan et al. \(2016\)](#) about the upper bound of  $\|V^{-1}\mathbf{x}\|_\infty$ , which relies on the upper bound of  $\|V^{-1} - S\|$ .

**Lemma 4.** [Yan et al. (2016)]. If  $V \in \mathcal{L}_n(m, M)$  with  $M/m = o(n)$ , then for large enough  $n$ ,

$$\|V^{-1} - S\| \leq \frac{c_1 M^2}{m^3(n-1)^2},$$

where  $c_1$  is a constant that does not depend on  $M$ ,  $m$  and  $n$ .

**Lemma 5.** [Yan et al. (2016)]. If  $V \in \mathcal{L}_n(m, M)$  with  $M/m = o(n)$ , then for a vector  $\mathbf{x} \in \mathbb{R}^{2n-1}$ ,

$$\begin{aligned} \|V^{-1}\mathbf{x}\|_\infty &\leq \|(V^{-1} - S)\mathbf{x}\|_\infty + \|S\mathbf{x}\|_\infty \\ &\leq \frac{2c_1(2n-1)M^2\|\mathbf{x}\|_\infty}{m^3(n-1)^2} + \frac{|x_{2n}|}{v_{2n,2n}} + \max_{i=1,\dots,2n-1} \frac{|x_i|}{v_{i,i}}, \end{aligned}$$

where  $x_{2n} := \sum_{i=1}^n x_i - \sum_{i=n+1}^{2n-1} x_i$ .

**Proof of Lemma 2.** Note that  $\{e_i^+\}_{i=1}^n$  and  $\{e_j^-\}_{j=1}^n$  are independently discrete Laplace random variables. Let  $[c]$  be the integer part of  $c$  ( $c = \kappa\sqrt{\log n}$ ). Then we have

$$\begin{aligned} P(|e_i| \leq c) &= \frac{1-\lambda}{1+\lambda}(1 + 2\lambda^1 + \dots + 2\lambda^{[c]}) \\ &= \frac{1+\lambda - 2\lambda^{[c]+1}}{1+\lambda} = 1 - \frac{2\lambda^{[c]+1}}{1+\lambda}. \end{aligned}$$

Therefore, we have

$$P(\max_i |e_i| > c) = 1 - \prod_{i=1}^n P(|e_i| \leq c) = 1 - \left(1 - \frac{2\lambda^{[c]+1}}{1+\lambda}\right)^n. \quad (\text{A.1})$$

Note that when  $\epsilon(c+1) > 2\log 2$ ,  $e^{-\epsilon(c+1)/2} < 1/2$ . Since the function  $f(x) = 1 - (1-x)^n$  is an increasing function on  $x$  when  $x \in (0, 1)$ , we have

$$1 - \left(1 - \frac{2e^{-\epsilon(c+1)/2}}{1 + e^{-\epsilon/2}}\right)^n \leq 1 - (1 - 2e^{-\epsilon(c+1)/2})^n,$$

on the other hand,  $(1-x)^n \geq 1 - nx$  when  $x \in (0, 1)$ . So, we have

$$1 - (1 - 2e^{-\epsilon(c+1)/2})^n \leq 1 - (1 - n \times 2e^{-\epsilon(c+1)/2}) = 2ne^{-\epsilon(c+1)/2}.$$

Assume that  $\epsilon(c+1) \geq 4\log n$ , we have

$$P(\max_i |e_i| > c) \leq \frac{2}{n}. \quad (\text{A.2})$$



By Hoeffding's inequality [[Hoeffding \(1963\)](#)], we have

$$P(|d_i^+ - \mathbb{E}(d_i^+)| \geq \sqrt{n \log n}) \leq 2 \exp \left\{ -\frac{2n \log n}{n-1} \right\} = O\left(\frac{1}{n^2}\right).$$

Therefore,

$$P(\max_i |d_i^+ - \mathbb{E}(d_i^+)| \geq \sqrt{n \log n}) \leq n \times \frac{1}{n^2} = \frac{1}{n}.$$

Similarly, we have

$$P(\max_j |d_j^- - \mathbb{E}(d_j^-)| \geq \sqrt{n \log n}) \leq n \times \frac{1}{n^2} = \frac{1}{n}.$$

So, with probability at least  $1 - 3/n$ , we have

$$\max_{i=1,\dots,n} |z_i^+ - \mathbb{E}(z_i^+)| \leq \max_i |d_i^+ - \mathbb{E}(d_i^+)| + \max_i |e_i^+| \leq \sqrt{n \log n} + \kappa \sqrt{\log n}.$$

Similarly, with probability at least  $1 - 3/n$ , we have

$$\max_{j=1,\dots,n} |z_j^- - \mathbb{E}(z_j^-)| \leq \sqrt{n \log n} + \kappa \sqrt{\log n}.$$

Let A and B be the events:

$$A = \left\{ \max_{i=1,\dots,n} |z_i^+ - \mathbb{E}(z_i^+)| \leq \sqrt{n \log n} + \kappa \sqrt{\log n} \right\},$$

$$B = \left\{ \max_{j=1,\dots,n} |z_j^- - \mathbb{E}(z_j^-)| \leq \sqrt{n \log n} + \kappa \sqrt{\log n} \right\}.$$

Consequently, as  $n$  goes to infinity, we have

$$\mathbb{P}(A \cap B) \geq 1 - \mathbb{P}(A^c) - \mathbb{P}(B^c) \geq 1 - 6/n \rightarrow 1.$$

This completes the proof of Lemma [2](#).

**Proof of Theorem [1](#).** In the Newton's iterative step, we set the initial value  $\boldsymbol{\theta}^{(0)} :=$

$\boldsymbol{\theta}^* \in \Theta$ . Let  $F(\boldsymbol{\theta}) = (F_1(\boldsymbol{\theta}), \dots, F_n(\boldsymbol{\theta}), F_{n+1}(\boldsymbol{\theta}), \dots, F_{2n-1}(\boldsymbol{\theta}))^\top$ . By (2.2), for  $i = 1, \dots, n$

$$\begin{aligned} \frac{\partial F_i}{\partial \alpha_l} &= 0, \quad l = 1, \dots, n, l \neq i, \quad \frac{\partial F_i}{\partial \alpha_i} = - \sum_{k=1, k \neq i}^n \frac{\partial \mu(\alpha_i + \beta_k)}{\partial \alpha_i}, \\ \frac{\partial F_i}{\partial \beta_j} &= - \frac{\partial \mu(\alpha_i + \beta_j)}{\partial \beta_j}, \quad j = 1, \dots, n-1, j \neq i, \quad \frac{\partial F_i}{\partial \beta_i} = 0, \end{aligned}$$

and for  $j = 1, \dots, n-1$ ,

$$\begin{aligned} \frac{\partial F_{n+j}}{\partial \alpha_l} &= - \frac{\partial \mu(\alpha_l + \beta_j)}{\partial \alpha_l}, \quad l = 1, \dots, n, l \neq j, \quad \frac{\partial F_{n+j}}{\partial \alpha_j} = 0. \\ \frac{\partial F_{n+j}}{\partial \beta_l} &= 0, \quad l = 1, \dots, n-1, l \neq j, \quad \frac{\partial F_{n+j}}{\partial \beta_j} = - \sum_{k=1, k \neq j}^n \frac{\partial \mu(\alpha_k + \beta_j)}{\partial \beta_j}. \end{aligned}$$

By (2.5), we get  $-F'(\boldsymbol{\theta}) \in \mathcal{L}_n(m, M)$ . To apply Proposition 1, we need to calculate  $r$  and  $\rho r$  in this proposition. By Lemma 5, we have

$$\begin{aligned} r &= \| [F'(\boldsymbol{\theta}^*)]^{-1} F(\boldsymbol{\theta}^*) \|_\infty \\ &\leq \frac{2c_1(2n-1)M^2}{m^3(n-1)^2} \|F(\boldsymbol{\theta}^*)\|_\infty + \frac{|F_{2n}(\boldsymbol{\theta}^*)|}{v_{2n,2n}} + \max_{i=1, \dots, 2n-1} \frac{|F_i(\boldsymbol{\theta}^*)|}{v_{i,i}} \\ &\leq \left( \frac{2c_1(2n-1)M^2}{m^3(n-1)^2} + \frac{1}{(n-1)m} + \frac{1}{m(n-1)} \right) (\sqrt{n \log n} + \kappa \sqrt{\log n}) \\ &= \frac{M^2}{m^3(n-1)} \left( \frac{2c_1(2n-1)}{n-1} + \frac{2m^2}{M^2} \right) (\sqrt{n \log n} + \kappa \sqrt{\log n}) \\ &\leq \frac{c_2 M^2}{nm^3} (\sqrt{n \log n} + \kappa \sqrt{\log n}), \end{aligned}$$

where  $c_2$  is a constant that does not depend on  $M$ ,  $m$  and  $n$ . Let

$$F'_i(\boldsymbol{\theta}) = (F'_{i,1}(\boldsymbol{\theta}), \dots, F'_{i,2n-1}(\boldsymbol{\theta}))^\top := \left( \frac{\partial F_i}{\partial \alpha_1}, \dots, \frac{\partial F_i}{\partial \alpha_n}, \frac{\partial F_i}{\partial \beta_1}, \dots, \frac{\partial F_i}{\partial \beta_{n-1}} \right)^\top.$$

Then, for  $i = 1, \dots, n$ , we have

$$\begin{aligned} \frac{\partial^2 F_i}{\partial \alpha_s \partial \alpha_l} &= 0, \quad s \neq l, \quad \frac{\partial^2 F_i}{\partial \alpha_i^2} = \sum_{k \neq i} \frac{\partial^2 \mu(\alpha_i + \beta_k)}{\partial \alpha_i^2}, \\ \frac{\partial^2 F_i}{\partial \beta_s \partial \alpha_i} &= \frac{\partial^2 \mu(\alpha_i + \beta_s)}{\partial \beta_s \partial \alpha_i}, \quad s = 1, \dots, n-1, s \neq i, \quad \frac{\partial^2 F_i}{\partial \beta_i \partial \alpha_i} = 0, \\ \frac{\partial^2 F_i}{\partial \beta_j^2} &= \frac{\partial^2 \mu(\alpha_i + \beta_j)}{\partial \beta_j^2}, \quad j = 1, \dots, n-1, \quad \frac{\partial^2 F_i}{\partial \beta_s \partial \beta_l} = 0, \quad s \neq l. \end{aligned}$$

By the mean value theorem for vector-valued functions [Lang (1993), page 341], we have

$$F'_i(\mathbf{x}) - F'_i(\mathbf{y}) = J^{(i)}(\mathbf{x} - \mathbf{y}), \quad \mathbf{x}, \mathbf{y} \in \Theta,$$

where  $J_{s,l}^{(i)} = \int_0^1 \frac{\partial F'_{i,s}}{\partial \theta_l}(t\mathbf{x} + (1-t)\mathbf{y})dt$ ,  $s, l = 1, \dots, 2n-1$ . By (2.6), we get

$$\max_s \sum_l |J_{s,l}^{(i)}| \leq 2\eta_1(n-1), \quad \sum_{s,l} |J_{s,l}^{(i)}| \leq 4\eta_1(n-1).$$

Similarly, for  $i = n+1, \dots, 2n-1$ , we also have  $F'_i(\mathbf{x}) - F'_i(\mathbf{y}) = J^{(i)}(\mathbf{x} - \mathbf{y})$ , and the above inequalities. Correspondingly, for any  $i$ , we get

$$\|F'_i(\mathbf{x}) - F'_i(\mathbf{y})\|_\infty \leq \|J^{(i)}\|_\infty \|\mathbf{x} - \mathbf{y}\|_\infty \leq 2\eta_1(n-1) \|\mathbf{x} - \mathbf{y}\|_\infty,$$

and for any vector  $\mathbf{v} \in \mathbb{R}^{2n-1}$ ,

$$\|[F'(\mathbf{x}) - F'(\mathbf{y})]\mathbf{v}\|_\infty \leq \|\mathbf{x} - \mathbf{y}\|_\infty \|\mathbf{v}\|_\infty \sum_{s,l} |J_{s,l}^{(i)}| \leq 4\eta_1(n-1) \|\mathbf{x} - \mathbf{y}\|_\infty \|\mathbf{v}\|_\infty.$$

Thus, we can choose  $K_1 = 4\eta_1(n-1)$  and  $K_2 = 2\eta_1(n-1)$ . Again, we also

$$\begin{aligned} \rho &= \frac{4c_1(2n-1)M^2}{2m^3n^2} \eta_1(n-1) + \frac{1}{(n-1)m} 2\eta_1(n-1) \\ &= \frac{M^2}{m^3} \left( \frac{4c_1(2n-1)(n-1)}{2n^2} + \frac{2m^2}{M^2} \right) \eta_1 \\ &\leq \frac{c_3 M^2 \eta_1}{m^3}, \end{aligned}$$

where  $c_3$  is a constant that does not depend on  $M, m$  and  $\eta_1$ . Combining with (2.9), we get

$$\begin{aligned} \rho r &\leq \frac{c_3 M^2 \eta_1}{m^3} \left( \frac{c_2 M^2}{nm^3} (\sqrt{n \log n} + \kappa \sqrt{\log n}) \right) \\ &\leq O \left( \frac{M^4 \eta_1}{nm^6} (\sqrt{n \log n} + \kappa \sqrt{\log n}) \right) = o(1). \end{aligned}$$

Thus, all conditions in Proposition 1 are satisfied. By Lemma 2, inequality (2.7) holds with probability approaching one such that (2.10) holds.

## A.2 Proof of Theorem 2

Before beginning to prove Lemma 3, we give the following Proposition.

**Proposition 2.** [Fan (2023)]. If  $M_u/m_u = o(n)$ , then for any fixed  $k \geq 1$ , as  $n \rightarrow \infty$ , the vector consisting of the first  $k$  elements of  $S\{\mathbf{g} - \mathbb{E}(\mathbf{g})\}$  is asymptotically multivariate normal with mean zero and covariance matrix  $Z$ .

**Proof of Lemma 3.** There are two cases to consider.

(i) By Yan (2021), we have

$$\left| \sum_{i=1}^n e_i^+ \right| = O_p(\kappa(n \log n)^{1/2}), \quad \left| \sum_{i=1}^n e_i^- \right| = O_p(\kappa(n \log n)^{1/2}). \quad (\text{A.3})$$

Since  $\tilde{g}_i - g_i = e_i^+$  for  $i = 1, \dots, n$ , and  $\tilde{g}_{n+j} - g_{n+j} = e_j^-$  for  $j = 1, \dots, n-1$ , we have

$$\begin{aligned} [S(\tilde{\mathbf{g}} - \mathbb{E}\mathbf{g})]_i &= [S(\mathbf{g} - \mathbb{E}\mathbf{g})]_i + [S(\tilde{\mathbf{g}} - \mathbf{g})]_i \\ &= [S(\mathbf{g} - \mathbb{E}\mathbf{g})]_i + (-1)^{1(i,j>n)} \frac{\sum_{i=1}^n e_i^+ - \sum_{j=1}^{n-1} e_j^-}{u_{2n,2n}} \\ &= [S(\mathbf{g} - \mathbb{E}\mathbf{g})]_i + O_p\left(\frac{\kappa(\log n)^{1/2}}{n^{1/2}m_u}\right). \end{aligned}$$

So if  $\frac{\kappa(\log n)^{1/2}}{m_u} = o(1)$ , then we have

$$[S(\tilde{\mathbf{g}} - \mathbb{E}\mathbf{g})]_i = [S(\mathbf{g} - \mathbb{E}\mathbf{g})]_i + o_p(n^{-1/2}).$$

Consequently, the first part of Lemma 3 immediately follows Proposition 2.

(ii)  $s_n/v_{2n,2n}^{1/2} \rightarrow c$  for some constant  $c$ . Let  $\tilde{e} = \sum_{i=1}^n e_i^+ - \sum_{j=1}^{n-1} e_j^-$  and  $\tilde{a}_{i,j} = a_{i,j} - \mathbb{E}a_{i,j}$ . Denote

$$\begin{aligned} H &:= \begin{pmatrix} \frac{\tilde{g}_1 - \mathbb{E}g_1}{u_{1,1}^{1/2}} \\ \vdots \\ \frac{\tilde{g}_k - \mathbb{E}g_k}{u_{k,k}^{1/2}} \\ \frac{\tilde{g}_{2n} - \mathbb{E}g_{2n}}{u_{2n,2n}^{1/2}} \\ \frac{\tilde{e}}{s_n} \end{pmatrix} = \begin{pmatrix} \frac{\sum_{j=1}^k \tilde{a}_{1,j}}{u_{1,1}^{1/2}} \\ \vdots \\ \frac{\sum_{j=1}^k \tilde{a}_{k,j}}{u_{k,k}^{1/2}} \\ \frac{\sum_{i=1}^k \tilde{a}_{i,n}}{u_{2n,2n}^{1/2}} \\ 0 \end{pmatrix} + \begin{pmatrix} \frac{\sum_{j=k+1}^n \tilde{a}_{1,j}}{u_{1,1}^{1/2}} \\ \vdots \\ \frac{\sum_{j=k+1}^n \tilde{a}_{k,j}}{u_{k,k}^{1/2}} \\ \frac{\sum_{i=k+1}^n \tilde{a}_{i,n}}{u_{2n,2n}^{1/2}} \\ \frac{\tilde{e}}{s_n} \end{pmatrix} \\ &:= I_1 + I_2. \end{aligned}$$

Since  $|a_{i,j}| \leq 1$  and  $u_{i,i} \rightarrow \infty$  as  $n \rightarrow \infty$ ,  $|\sum_{j=1}^k \tilde{a}_{i,j}|/u_{i,i} = o(1)$  for  $i = 1, \dots, k$  with fixed  $k$ . So  $I_1 = o(1)$ .

Next, we will consider  $I_2$ . Recall that  $s_n^2 = \text{Var}(\tilde{e})$ . By the large sample theory,  $(\tilde{e} - \mathbb{E}\tilde{e})/s_n$  converges in distribution to the standard normal distribution if  $s_n \rightarrow \infty$ . By the central limit theorem for the bounded case in Loève (1977) (page 289),  $\sum_{j=k+1}^n \tilde{a}_{i,j}/u_{i,i}^{1/2}$  converges in distribution to the standard normal distribution for any fixed  $i$  if  $M/m_u =$

$o(n)$ . Since  $\tilde{a}_{i,j}$ 's ( $1 \leq i \leq k, j = k+1, \dots, n$ ),  $\tilde{a}_{i,n}$ 's and  $\tilde{e}$  are mutually independent,  $I_2$  converges in distribution to a  $k+2$ -dimensional standardized normal distribution with covariance matrix  $I_{k+2}$ , where  $I_{k+2}$  denotes the  $(k+2) \times (k+2)$  dimensional identity matrix. Let

$$C = \begin{pmatrix} \frac{\sqrt{u_{1,1}}}{v_{1,1}}, & 0, & \dots, & 0, & \frac{\sqrt{u_{2n,2n}}}{v_{2n,2n}}, & \frac{s_n}{v_{2n,2n}} \\ 0, & \frac{\sqrt{u_{2,2}}}{v_{2,2}}, & \dots, & 0, & \frac{\sqrt{u_{2n,2n}}}{v_{2n,2n}}, & \frac{s_n}{v_{2n,2n}} \\ & & \dots & & & \\ 0, & 0, & \dots, & \frac{\sqrt{u_{k,k}}}{v_{k,k}}, & \frac{\sqrt{u_{2n,2n}}}{v_{2n,2n}}, & \frac{s_n}{v_{2n,2n}} \end{pmatrix}.$$

Then

$$[S(\tilde{\mathbf{g}} - \mathbb{E}\mathbf{g})]_{i=1,\dots,k} = CH.$$

Since  $s_n^2/v_{2n,2n} \rightarrow c^2$  for some constant  $c$ , all positive entries of  $C$  are in the same order  $n^{1/2}$ . So  $CM$  converges in distribution to the  $k$ -dimensional multivariate normal distribution with mean  $(\overbrace{0, \dots, 0}^k)$  and covariance matrix

$$\text{diag}\left(\frac{u_{1,1}}{v_{1,1}^2}, \dots, \frac{u_{k,k}}{v_{k,k}^2}\right) + \left(\frac{u_{2n,2n}}{v_{2n,2n}^2} + \frac{s_n^2}{v_{2n,2n}^2}\right) \mathbf{1}_k \mathbf{1}_k^\top,$$

where  $\mathbf{1}_k$  is a  $k$ -dimensional column vector with all entries 1. Before proving Theorem 2, we first establish two lemmas.

**Lemma 6.** [Yan et al. (2016)]. Let  $W = V^{-1} - S$  and  $R = \text{Cov}[W(\mathbf{g} - \mathbb{E}\mathbf{g})]$ , then

$$\|R\| \leq \|V^{-1} - S\| + \frac{3M}{m^2(n-1)^2}. \quad (\text{A.4})$$

**Lemma 7.** If (2.14) holds, then for any  $i$ ,

$$\hat{\boldsymbol{\theta}}_i - \boldsymbol{\theta}_i^* = [V^{-1}(\tilde{\mathbf{g}} - \mathbb{E}(\mathbf{g}))]_i + o_p(n^{-1/2}). \quad (\text{A.5})$$

*Proof.* Let  $\hat{r}_{i,j} = \hat{\alpha}_i + \hat{\beta}_j - \alpha_i^* - \beta_j^*$  and assume

$$\hat{\rho}_n := \max_{i \neq j} |\hat{r}_{i,j}| = O_p\left(\frac{c_2 M^2}{nm^3}(\sqrt{n \log n} + \kappa \sqrt{\log n})\right).$$

For any  $1 \leq i \neq j \leq n$ , by the Taylor's expansion, we get

$$\mu(\hat{\alpha}_i + \hat{\beta}_j) - \mu(\alpha_i^* + \beta_j^*) = \mu'(\alpha_i^* + \beta_j^*) \hat{r}_{i,j} + h_{i,j},$$

where  $h_{i,j} = \frac{1}{2} \mu''(\hat{\theta}_{i,j}) \hat{r}_{i,j}^2$  and  $\hat{\theta}_{i,j} = t_{i,j}(\alpha_i^* + \beta_j^*) + (1 - t_{i,j})(\hat{\alpha}_i + \hat{\beta}_j)$ ,  $t_{i,j} \in (0, 1)$ . By (2.3),

we have

$$\tilde{\mathbf{g}} - \mathbb{E}(\mathbf{g}) = V(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*) + \mathbf{h}.$$

Equivalently,

$$\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^* = V^{-1}(\tilde{\mathbf{g}} - \mathbb{E}(\mathbf{g})) + V^{-1}\mathbf{h}. \quad (\text{A.6})$$

where  $\mathbf{h} = (h_1, \dots, h_{2n-1})^\top$  and

$$h_i = \sum_{k=1, k \neq i}^n h_{i,k}, i = 1, \dots, n,$$

$$h_{n+i} = \sum_{k=1, k \neq i}^n h_{k,i}, i = 1, \dots, n-1.$$

By (2.6), we get

$$|h_{i,j}| = \left| \frac{1}{2} \mu''(\hat{\theta}_{i,j}) \hat{r}_{i,j}^2 \right| \leq \frac{1}{2} \eta_1 \hat{r}_{i,j}^2 \leq \frac{1}{2} \eta_1 \hat{\rho}_n^2,$$

$$|h_i| \leq \sum_{j \neq i} |h_{i,j}| \leq \frac{1}{2} (n-1) \eta_1 \hat{\rho}_n^2.$$

Note that  $(S\mathbf{h})_i = \frac{h_i}{v_{i,i}} + (-1)^{1(i>n)} \frac{h_{2n}}{v_{2n,2n}}$ ,  $h_{2n} := \sum_{i=1}^n h_i - \sum_{i=n+1}^{2n-1} h_i = \sum_{i=1, i \neq k}^n h_{i,n} \leq \frac{n-1}{2} \eta_1 \hat{\rho}_n^2$  and  $(V^{-1}\mathbf{h})_i = (S\mathbf{h})_i + (W\mathbf{h})_i$ . By Lemma 4, we get

$$\begin{aligned} |(V^{-1}\mathbf{h})_i| &\leq |(S\mathbf{h})_i| + |(W\mathbf{h})_i| \\ &\leq \frac{|h_i|}{v_{i,i}} + \frac{|h_{2n}|}{v_{2n,2n}} + \|W\|_{\max} \times \left[ (2n-1) \max_i |h_i| \right] \\ &\leq O\left(\frac{\eta_1 \hat{\rho}_n^2}{m}\right) + O\left(\frac{M^2}{m^3(n-1)^2} \times \frac{1}{2} (2n-1)(n-1) \eta_1 \hat{\rho}_n^2\right) \\ &\leq O\left(\frac{m^2 + M^2}{2m^3} \eta_1 \hat{\rho}_n^2\right) \\ &= O\left(\frac{M^4(m^2 + M^2)}{2m^9 n^2} \eta_1 (n + 2\kappa n^{1/2} + \kappa^2) \log n\right) \\ &\leq O\left((n + 2\kappa n^{1/2} + \kappa^2) \frac{M^6 \eta_1 \log n}{m^9 n^2}\right). \end{aligned}$$

Again, by (2.14), we obtain

$$|(V^{-1}\mathbf{h})_i| \leq |(S\mathbf{h})_i| + |(W\mathbf{h})_i| = o(n^{-1/2}). \quad (\text{A.7})$$

□

**Proof of Theorem 2.** Note that  $V^{-1} = W + S$ , we have

$$(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta}^*)_i = [S\{\tilde{\mathbf{g}} - \mathbb{E}(\mathbf{g})\}]_i + [W\{\tilde{\mathbf{g}} - \mathbb{E}(\mathbf{g})\}]_i + o_p(n^{-1/2}).$$

By (A.2),  $\|\tilde{\mathbf{g}} - \mathbf{g}\|_\infty = O_p(\kappa\sqrt{\log n})$ . So by Lemma 4, we have

$$[W(\tilde{\mathbf{g}} - \mathbf{g})]_i = O_p(n \frac{M^2}{m^3 n^2} \kappa \sqrt{\log n}) = O_p(\frac{\kappa M^2 \sqrt{\log n}}{n m^3}).$$

If  $\frac{\kappa M^2}{m^3} = o((n/\log n)^{1/2})$ , then  $[W\{\tilde{\mathbf{g}} - \mathbf{g}\}]_i = o_p(n^{-1/2})$ . Combing Lemma 6, it yields

$$[W(\tilde{\mathbf{g}} - \mathbb{E}\mathbf{g})]_i = [W(\tilde{\mathbf{g}} - \mathbf{g})]_i + [W(\mathbf{g} - \mathbb{E}\mathbf{g})]_i = o_p(n^{-1/2}).$$

Consequently,

$$(\hat{\boldsymbol{\theta}} - \boldsymbol{\theta})_i = [S(\tilde{\mathbf{g}} - \mathbb{E}\mathbf{g})]_i + o_p(n^{-1/2}).$$

Theorem 2 immediately follows from Lemma 3.

## References

- Blitzstein J, Diaconis P. A sequential importance sampling algorithm for generating random graphs with prescribed degrees. *Internet mathematics*, 2011, 6(4): 489-522.
- Chatterjee S, Diaconis P, Sly A. Random graphs with a given degree sequence. *The Annals of Applied Probability*, 2011: 1400-1435.
- Dwork C, McSherry F, Nissim K, et al. Calibrating noise to sensitivity in private data analysis. *Theory of Cryptography: Third Theory of Cryptography Conference*, Springer Berlin Heidelberg, 2006: 265-284.
- Fan Y. Inference in a class of directed random graph models with an increasing number of parameters. *Communications in Statistics-Theory and Methods*, 2023, 52(13): 4491-4513.
- Hay M, Miklau G, Jensen D, et al. Resisting structural re-identification in anonymized social networks. *Proceedings of the VLDB Endowment*, 2008, 1(1): 102-114.
- Hay M, Li C, Miklau G, et al. Accurate estimation of the degree distribution of private networks. *Ninth IEEE International Conference on Data Mining. IEEE*, 2009: 169-178.
- Helleringer S, Kohler H P. Sexual network structure and the spread of HIV in Africa: evidence from Likoma Island, Malawi. *Aids*, 2007, 21(17): 2323-2332.
- Hoeffding W. Probability inequalities for sums of bounded random variables. *Journal of the American statistical association*. 1963, 58(301): 13-30.
- Holland P W, Leinhardt S. An exponential family of probability distributions for directed

- graphs (with discussion). *Journal of the American Statistical Association*, 1981, 76(373): 33-50.
- Jernigan C, Mistree B F T. Gaydar: Facebook friendships expose sexual orientation. *First Monday*, 2009.
- Karwa V, Slavković A. Inference using noisy degrees: Differentially private  $\beta$ -model and synthetic graphs. *The Annals of Statistics*. 2016, 44: 87-112.
- Kasiviswanathan S P, Nissim K, Raskhodnikova S, et al. Analyzing graphs with node differential privacy. *Theory of Cryptography: 10th Theory of Cryptography Conference*. Springer Berlin Heidelberg, 2013: 457-476.
- Lang S. *Real and Functional Analysis*. Springer, New York. 1993.
- Lazega E. *The collegial phenomenon: The social mechanisms of cooperation among peers in a corporate law partnership*. Oxford University Press, USA, 2001.
- Loève M. *Probability theory I*. 4th ed. Springer, New York. 1977.
- Lu W, Miklau G. Exponential random graph estimation under differential privacy. *Proceedings of the 20th ACM SIGKDD international conference on Knowledge discovery and data mining*. 2014: 921-930.
- Narayanan A, Shmatikov V. De-anonymizing social networks. In *30th IEEE symposium on security and privacy*. IEEE, 2009: 173-187.
- Nissim K, Raskhodnikova S, Smith A. Smooth sensitivity and sampling in private data analysis. *Proceedings of the thirty-ninth annual ACM symposium on Theory of computing*. 2007: 75-84.
- Pan L, Yan T. Asymptotics in the  $\beta$ -model for networks with a differentially private degree sequence. *Communications in Statistics-Theory and Methods*, 2020, 49(18): 4378-4393.
- Snijders T A B, Pattison P E, Robins G L, et al. New specifications for exponential random graph models. *Sociological methodology*, 2006, 36(1): 99-153.
- Wang Q, Wang Q, Luo J. Asymptotics in a probit model for directed networks. *Communications in Statistics-Theory and Methods*, 2022, 51(11): 3463-3479.
- Yan T, Xu J. A central limit theorem in the *beta*-model for undirected random graphs with a diverging number of vertices. *Biometrika*, 2013, 100(2): 519-524.
- Yan T, Leng C, Zhu J. Asymptotics in directed exponential random graph models with an increasing bi-degree sequence. *The Annals of Statistics*, 2016, 44: 31-57.
- Yan T. Directed networks with a differentially private bi-degree sequence. *Statistica Sinica*. 2021, 31: 2031-2050.