

# On a Finite Group Generated by Subnormal Supersoluble Subgroups

Victor S. Monakhov\*

**Abstract.** Supersolubility of a finite group  $G = \langle A, B \rangle$  with the nilpotent derived subgroup  $G'$  is established under the condition that the subgroups  $A$  and  $B$  are both subnormal and supersoluble.

**Mathematics Subject Classification (2010).** 20D10; 20D35.

**Keywords.** Finite group, supersoluble group, subnormal subgroup, derived subgroup.

## 1 Main Result

All groups in this paper are finite. We use the standard notations and terminology of [3].

B. Huppert [4] and R. Baer [1] gave the first examples of nonsupersoluble groups that were a product of normal supersoluble subgroups. R. Baer [1] established supersolubility of a group  $G = AB$  with the nilpotent derived subgroup  $G'$  and normal supersoluble subgroups  $A$  and  $B$ . A. F. Vasil'ev and T. I. Vasil'eva [7] showed that instead of nilpotency of the derived subgroup  $G'$  it is enough to require nilpotency of the  $\mathcal{A}$ -residual, where  $\mathcal{A}$  is the formation of all groups with abelian Sylow subgroups. In these results, normality of subgroups  $A$  and  $B$  can be weakened to subnormality of  $A$  and  $B$  [5]. This theme was developed in many papers, see for example [2].

If a group  $G$  is generated by subnormal subgroups  $X$  and  $Y$ , then it is not always true that  $G = XY$ . The simplest examples are the dihedral group of order 8 and nonabelian group of order  $p^3$  and exponent  $p$ . The normal closure  $X^G$  of a subnormal supersoluble subgroup  $X$  in a group  $G$  can be a nonsupersoluble subgroup. For example,  $G = PSU_3(2) \rtimes C_2$  [6, SmallGroup(144,182)] contains a nonsupersoluble maximal subgroup  $H = S_3 \wr C_2$  [6, SmallGroup(72,40)] of index 2. A subgroup  $X = S_3 \times S_3$  of  $H$  is supersoluble and not normal in  $G$ . Since  $|H : X| = 2$ ,  $X^G = H$  and  $X$  is subnormal in  $G$ .

Next, we write  $G = \langle A, B \rangle$  when a group  $G$  is generated by subgroups  $A$  and  $B$ .

In this paper, we prove the following theorem.

**Theorem 1.1.** *Let  $A$  and  $B$  be subnormal supersoluble subgroups of a group  $G$  and let  $G = \langle A, B \rangle$ . Then  $G$  is metanilpotent and has a Sylow tower of supersoluble type. Furthermore,  $G$  is supersoluble when one of the following conditions holds.*

---

\*Department of Mathematics and Programming Technologies, Francisk Skorina Gomel State University, Belarus, victor.monakhov@gmail.com

- (1)  $G^A$  is nilpotent.
- (2)  $(|A : A'|, |B : B'|) = 1$ .

*Proof.* Let  $p \in \pi(G)$  and let  $A_p$  and  $B_p$  be Sylow  $p$ -subgroups of  $A$  and  $B$ , respectively. Suppose that  $p$  is the largest prime in  $\pi(G)$ . Then  $A_p$  and  $B_p$  are normal in  $A$  and  $B$ , respectively [3, VI.9.1]. Therefore  $A_p$  and  $B_p$  are subnormal in  $G$  and  $\langle A_p, B_p \rangle \leq O_p(G)$ . By induction,  $G/O_p(G)$  has a Sylow tower of supersoluble type. Since  $AO_p(G)/O_p(G)$  and  $BO_p(G)/O_p(G)$  are subnormal  $p'$ -subgroups of  $G/O_p(G)$ ,

$$G/O_p(G) = \langle AO_p(G)/O_p(G), BO_p(G)/O_p(G) \rangle$$

is a  $p'$ -group. Hence  $G$  has a Sylow tower of supersoluble type.

Since  $A'$  is nilpotent [3, VI.9.1] and subnormal in  $G$ , we have  $A' \leq F(A) \leq F(A)^G \leq F(G)$ . Consequently,

$$AF(G)/F(G) \cong A/(A \cap F(G))$$

is abelian and subnormal in  $G/F(G)$ . Similarly,

$$B' \leq F(B) \leq F(B)^G \leq F(G), \quad BF(G)/F(G) \cong B/(B \cap F(G)),$$

therefore  $BF(G)/F(G)$  is abelian and subnormal in  $G/F(G)$ . It is clear that

$$G/F(G) = \langle AF(G)/F(G), BF(G)/F(G) \rangle. \quad (*)$$

So,  $(AF(G)/F(G))^{G/F(G)}$  and  $(BF(G)/F(G))^{G/F(G)}$  are normal in  $G/F(G)$  and nilpotent. Hence

$$G/F(G) = (AF(G)/F(G))^{G/F(G)}(BF(G)/F(G))^{G/F(G)}$$

is nilpotent, and  $G$  is metanilpotent.

(1) Let  $G^A$  be nilpotent. Then  $G^A \leq F(G)$ . Since  $G$  is metanilpotent, we conclude that  $G/F(G)$  is nilpotent, hence  $G/F(G)$  is abelian and  $G' \leq F(G)$ . Therefore  $AG'$  and  $BG'$  is normal in  $G$ . According to [5, Lemma 10],  $AG'$  and  $BG'$  are supersoluble. Hence  $G = (AG')(BG')$  is supersoluble by Baer's Theorem.

(2) Let  $(|A : A'|, |B : B'|) = 1$ . Since  $A'B' \subseteq F(G)$ , we get

$$(|AF(G)/F(G)|, |BF(G)/F(G)|) = 1.$$

From (\*), it follows that

$$G/F(G) = AF(G)/F(G) \times BF(G)/F(G)$$

is abelian and  $G' \leq F(G)$ . Therefore  $G = (AG')(BG')$  is supersoluble.  $\square$

**Corollary 1.1.1.** *Let  $A$  and  $B$  be subnormal supersoluble subgroups of a group  $G$  and let  $G = \langle A, B \rangle$ . If the derived subgroup of  $G$  is nilpotent, then  $G$  is supersoluble.*

## References

- [1] R. Baer, *Classes of finite groups and their properties*. Illinois J. Math., **1** (1957), 115–187. <https://doi.org/10.1215/ijm/1255379396>
- [2] A. Ballester-Bolinches, R. Estaban-Romero, M. Asaad, *Products of finite groups*. Berlin, New York, Walter de Gruyter, 2010. <https://doi.org/10.1515/9783110220612>
- [3] B. Huppert, *Endliche Gruppen. I*. Berlin, Springer-Verlag, 1967.
- [4] B. Huppert, *Monomiale Darstellung endlicher Gruppen*. Nagoya Math. J., **3** (1953), 93–94. <https://doi.org/10.1017/S0027763000017025>
- [5] V.S. Monakhov, I.K. Chirik, *On the supersoluble residual of a product of subnormal supersoluble subgroups*. Siberian Math. J., **58**:2 (2017), 271B–280. <https://doi.org/10.17377/smzh.2017.58.209>
- [6] *The GAP Group: GAP — Groups, Algorithms, and Programming*. Ver. 4.11.0 released on 29 February 2020, <http://www.gap-system.org>.
- [7] A. F. Vasil’ev, T. I. Vasilyeva, *On finite groups whose principal factors are simple groups*. Russian Math. (Iz. VUZ), **41**:11 (1997), 8–12.