

Algebraic Multiscale Method for two-dimensional elliptic problems

Kanghun Cho*, Imbunm Kim†, Raehyun Kim‡, Dongwoo Sheen†

Thursday 27th January, 2022

Abstract

We introduce an algebraic multiscale method for two-dimensional problems. The method uses the generalized multiscale finite element method based on the quadrilateral nonconforming finite element spaces. Differently from the one-dimensional algebraic multiscale method, we apply the dimension reduction techniques to construct multiscale basis functions. Also moment functions are considered to impose continuity between local basis functions. Some representative numerical results are presented.

Keywords. Multiscale, algebraic multiscale method, heterogeneous coefficient.

1 Introduction

In this paper we propose an algebraic multiscale method for two-dimensional elliptic problems. This is an extension of our previous work for the one-dimensional case [3]. We consider the multiscale model problem given by

$$-\nabla \cdot (\kappa(x)\nabla u) = f \text{ in } \Omega, \quad (1.1)$$

where κ is a heterogeneous coefficient and $f \in H^{-1}(\Omega)$.

Assume that we are only given a linear system (3.3), which is obtained from a discretization of micro-scale governing equation, but any detailed information on the coefficient κ and the source term f are not given. In this situation, our target is to construct a macro-scale linear system by using the information and finally to find numerical solutions which possess similar properties of the solutions obtained by multiscale methods.

*Samsung Fire & Marine Insurance Co., Ltd., 14, Seocho-daero 74-gil, Seocho-gu, Seoul 06620, Korea;

†Department of Mathematics, Seoul National University, Seoul 08826, South Korea;

‡Department of Mathematics, University of California, Berkeley, CA 94720-3840, USA; Emails: serein@snu.ac.kr, ikim@snu.ac.kr, rhkim79@math.berkeley.edu, sheen@snu.ac.kr

Multiscale methods have been actively developed in various manners including heterogeneous multiscale methods [1, 2], multiscale hybridizable discontinuous Galerkin methods [4, 8, 9], and multiscale finite element methods [7, 10, 11]. Here we adopt the generalized multiscale finite element method (GMsFEM) [6, 12] based on nonconforming finite element space [5, 13]. However, the method described in this paper applies for the conforming GMsFEM as well.

The GMsFE spaces consist of snapshot function spaces, offline function spaces, and moment function spaces. First, snapshot functions are obtained by solving κ -harmonic problems in each macro element. Then offline functions are constructed by applying suitable dimension reduction techniques to snapshot function space. In one-dimensional case we choose offline functions identical to the snapshot functions since there are only two snapshot functions in each macro element. However in higher dimension, it is necessary to apply such techniques since we have many more snapshot functions which yields huge computational cost. The moment functions are also needed in order to impose continuity between local offline functions, which make a remarkable difference between the 1D and 2D cases.

This paper is organized as follows. In Section 2, we briefly review the nonconforming generalized multiscale finite element method (GMsFEM) based on finite element spaces. Then the algebraic multiscale method for two-dimensional elliptic problem is introduced in Section 3 following the framework of GMsFEM. Section 4 is devoted to energy norm error estimate of the proposed method. In Section 5, representative numerical results are presented. Conclusions are given in Section 6.

2 Preliminaries

In this section we briefly review the framework of the generalized multiscale finite element method (GMsFEM) based on the quadrilateral nonconforming finite element introduced in [5], following [12]. We only consider two-dimensional elliptic boundary problems here, but the framework can be extended to higher dimensional cases and used for other multiscale methods.

Let U be any open subset of \mathbb{R}^2 . Denote the seminorm, norm, and inner product of the Sobolev space $H^k(U)$ by $|\cdot|_{k,U}$, $\|\cdot\|_{k,U}$, and $(\cdot, \cdot)_{k,U}$, respectively. For the space $H^0(U) = L^2(U)$, we abbreviate $(\cdot, \cdot)_{0,U}$ as $(\cdot, \cdot)_U$. Given $f \in H^{-1}(\Omega)$, consider the following elliptic boundary problem:

$$\begin{cases} -\nabla \cdot (\kappa(\mathbf{x}) \nabla u) = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (2.1)$$

where Ω is a simply connected polygonal domain in \mathbb{R}^2 , and κ is a highly heterogeneous coefficient. The weak formulation of (2.1) is to seek $u \in H_0^1(\Omega)$ such that

$$a(u, v) = F(v), \quad v \in H_0^1(\Omega) \quad (2.2)$$

where $a(u, v) = \int_{\Omega} \kappa \nabla u \cdot \nabla v \, d\mathbf{x}$ and $F(v) = \int_{\Omega} f v \, d\mathbf{x}$. Let $\mathcal{T}_h := \bigcup_{j=1}^{N_h} \{T_j\}$ be a family of shape regular triangulations of Ω and V_h be a finite element basis function space based on \mathcal{T}_h . The mesh parameter h is given by

$$h = \max_{j=1, \dots, N_h} \text{diam}(T_j).$$

Let $V_{h,0}$ be the set of all elements in V_h , whose DOFs related to the boundary $\partial\Omega$ vanish. Then the finite element approximation of (2.2) is defined as the solution $u_h \in V_{h,0}$ of the discrete problem

$$a_h(u_h, v_h) = F_h(v_h), \quad v_h \in V_{h,0}, \quad (2.3)$$

where $a_h(u, v) = \sum_{T_j \in \mathcal{T}_h} \int_{T_j} \kappa \nabla u \cdot \nabla v \, d\mathbf{x}$ and $F_h(v) = \sum_{T_j \in \mathcal{T}_h} \int_{T_j} f v \, d\mathbf{x}$. In GMsFEM, we also need to have another shape regular triangulations $\mathcal{T}^H := \bigcup_{J=1}^{N^H} \{T^J\}$ of Ω . We suppose that every $T^J \in \mathcal{T}^H$ consists of a connected union of $T_j \in \mathcal{T}_h$, which makes \mathcal{T}_h be a refinement of \mathcal{T}^H . Here, and in what follows, we refer two triangulations \mathcal{T}_h and \mathcal{T}^H to micro-scale and macro-scale triangulations, respectively. The macro mesh parameter H is given by

$$H = \max_{J=1, \dots, N^H} \text{diam}(T^J).$$

Let V^H be a finite element basis function space associated with \mathcal{T}^H , and $V^{H,0}$ be the set of all elements in V^H , whose DOFs related to $\partial\Omega$ vanish. Then the GMsFE approximation of (2.2) is equivalent to find $u^H \in V^{H,0}$ such that

$$a_h(u^H, v^H) = F_h(v^H), \quad v^H \in V^{H,0}. \quad (2.4)$$

2.1 Framework of nonconforming GMsFEM

The success in using the GMsFEM depends on the construction of corresponding finite element space. V^H must contain the essential properties of V_h as well as the coefficient κ , while the dimension of V^H is significantly reduced compared to that of V_h .

The GMsFE space V^H is composed of two components. The first one is the offline function space which is a spectral decomposition of the snapshot function space, and used to represent the solution in each macro element. The second one is the moment function space which is used to impose continuity between local offline functions. Let a micro-scale basis function space V_h be given. For each macro element $T \in \mathcal{T}^H$, denote the restriction of V_h to T by $V_h(T)$. Also denote the set of all macro edges in \mathcal{T}^H by $\mathcal{E}^H := \bigcup_{J=1}^{N^E} \{E^J\}$, and the set of all interior macro edges by $\mathcal{E}^{H,0}$. Then the process of constructing GMsFE spaces is organized into the following framework:

1. Construct a snapshot function space $V^{\text{snap}} = \bigcup_{T \in \mathcal{T}^H} V^{\text{snap}}(T)$, where $V^{\text{snap}}(T)$ is a subspace of $V_h(T)$ for each macro element $T \in \mathcal{T}^H$. In general, $V^{\text{snap}}(T)$ is chosen to be the span of κ -harmonic functions in T .

2. Construct an offline function space $V^{\text{off}} = \bigcup_{T \in \mathcal{T}^H} V^{\text{off}}(T)$, where $V^{\text{off}}(T)$ is obtained by applying a suitable dimension reduction technique to $V^{\text{snap}}(T)$ for each macro element $T \in \mathcal{T}^H$. We may use generalized eigenvalue decomposition, the singular value decomposition, the proper orthogonal decomposition, and so on.
3. Construct a moment function space $\mathcal{M}_H = \bigcup_{E \in \mathcal{E}^H} \mathcal{M}_H(E)$. $\mathcal{M}_H(E)$ may consist of local κ -harmonic functions in appropriate neighborhood of E . The moment functions are used to glue offline functions through each macro interior edge $E \in \mathcal{E}^{H,0}$.
4. Construct the nonconforming GMsFE spaces V^H and $V^{H,0}$ based on V^{off} and \mathcal{M}_H . They are defined as

$$V^H = \left\{ \psi \in V^{\text{off}} \mid \langle [\psi]_E, \zeta \rangle_E = 0, \forall \zeta \in \mathcal{M}_H(E), \forall E \in \mathcal{E}^{H,0} \right\},$$

$$V^{H,0} = \left\{ \psi \in V^{\text{off}} \mid \langle [\psi]_E, \zeta \rangle_E = 0, \forall \zeta \in \mathcal{M}_H(E), \forall E \in \mathcal{E}^H \right\}.$$

Here $[\psi]_E$ stands for the jump of ψ across macro edge E .

2.2 Notations for the DSSY element and edge-based basis functions

We implement the rectangular DSSY (Douglas-Santos-Sheen-Ye) nonconforming element to construct micro-scale basis function space V_h . Since the DSSY elements are based on the horizontal-type and vertical-type edges, it is more natural to label the edges and basis functions in these two types. For $j = 1, \dots, N_x$ and $k = 1, \dots, N_y$, let Ω_{jk} be the $(j, k)^{\text{th}}$ rectangle with the four vertices $(x_j, y_k), (x_{j-1}, y_k), (x_{j-1}, y_{k-1})$, and (x_j, y_{k-1}) , and vertical edges $e_{jk}, e_{j-1,k}$, and horizontal edges $f_{jk}, f_{j,k-1}$. Edge-based basis functions are given respectively. That is, the two basis functions $\phi_{j,k}$ and $\phi_{j-1,k}$ are associated with the edges e_{jk} and $e_{j-1,k}$, and the two basis functions $\psi_{j,k}$ and $\psi_{j,k-1}$ associated with the edges f_{jk} and $f_{j,k-1}$. See Figure 2.1 for an illustration.




Figure 2.1: The basis functions $\phi_{j'k}$ associated with vertical type edge $e_{j'k}$ for $j' = j-1, j$ and the basis functions $\psi_{jk'}$ associated with horizontal type edge $f_{jk'}$ for $k' = k-1, k$ in element Ω_{jk} with vertices $(x_{j'}, y_{k'})$'s.

3 Algebraic Multiscale Method

In this section, we design an algebraic multiscale method for two-dimensional elliptic problems. We assume that all the components in A and b in the micro-scale linear system $Ax = b$ are known, which is constructed by the finite element method to find $u_h \in V_h$ such that

$$a_h(u_h, v_h) = (f, v_h)_\Omega \quad \forall v_h \in V_h, \quad (3.1)$$

where $a_h(u_h, v_h) = \sum_{j,k} (\kappa \nabla u_h, \nabla v_h)_{\Omega_{j,k}}$. Here we do not assume that any a priori knowledge is given for the coefficient κ and the exterior source term f .

On our procedure, we use the GMsFEM under the following assumptions:

1. The micro-scale mesh is rectangular.
2. The micro-scale linear system is constructed by the DSSY nonconforming finite element method.
3. The coefficient κ is assumed to be constant on each micro element.

Let $V_h = \text{Span}(\{\psi_{jk}\}_{j,k} \cup \{\phi_{jk}\}_{j,k})$ be the DSSY nonconforming finite element space associated with \mathcal{T}_h , where $\psi_{j,k}$ is the DSSY basis functions associated with DOF at midpoint on horizontal micro edge f_{jk} , and $\phi_{j,k}$ is the DSSY basis functions associated with DOF at midpoint on vertical micro edge e_{jk} (see Figure 2.1). Then the micro-scale solution u_h is represented by

$$u_h = \sum_{j'=0}^{n_x} \sum_{k'=1}^{n_y} \alpha_{j'k'} \phi_{j'k'} + \sum_{j'=1}^{n_x} \sum_{k'=0}^{n_y} \beta_{j'k'} \psi_{j'k'}. \quad (3.2)$$

Here, and in what follows, the indices α and β stand for the coefficients for horizontal and vertical edges, respectively. Test (3.1) with u_h represented by (3.2) against $v_h = \phi_{jk}$ and $v_h = \psi_{jk}$ to obtain

$$\begin{aligned} \sum_{j'=0}^{n_x} \sum_{k'=1}^{n_y} \alpha_{j'k'} a_h(\phi_{j'k'}, \phi_{jk}) + \sum_{j'=1}^{n_x} \sum_{k'=0}^{n_y} \beta_{j'k'} a_h(\psi_{j'k'}, \phi_{jk}) \\ = (f, \phi_{jk}), \quad j = 0, \dots, n_x, k = 1, \dots, n_y, \end{aligned} \quad (3.3a)$$

$$\begin{aligned} \sum_{j'=0}^{n_x} \sum_{k'=1}^{n_y} \alpha_{j'k'} a_h(\phi_{j'k'}, \psi_{jk}) + \sum_{j'=1}^{n_x} \sum_{k'=0}^{n_y} \beta_{j'k'} a_h(\psi_{j'k'}, \psi_{jk}) \\ = (f, \psi_{jk}), \quad j = 1, \dots, n_x, k = 0, \dots, n_y. \end{aligned} \quad (3.3b)$$

Taking into account of the supports of basis functions, we get the following linear system for the micro-scale elliptic problem:

$$\begin{aligned} A_{jkjk}^{\alpha,\alpha} \alpha_{jk} + A_{j-1jk}^{\alpha,\alpha} \alpha_{j-1k} + A_{j+1jk}^{\alpha,\alpha} \alpha_{j+1k} \\ + A_{jkjk}^{\beta,\alpha} \beta_{jk} + A_{jk-1k}^{\beta,\alpha} \beta_{jk-1} + A_{j+1kj}^{\beta,\alpha} \beta_{j+1k} + A_{j+1k-1j}^{\beta,\alpha} \beta_{j+1k-1} \end{aligned}$$

$$= f_{jk}^\alpha, \quad j = 0, \dots, n_x, k = 1, \dots, n_y, \quad (3.4a)$$

$$\begin{aligned} & A_{jkjk}^{\beta,\beta} \beta_{jk} + A_{jk-1jk}^{\beta,\beta} \beta_{jk-1} + A_{jk+1jk}^{\beta,\beta} \beta_{jk+1} \\ & + A_{jkjk}^{\alpha,\beta} \alpha_{jk} + A_{j-1kjk}^{\alpha,\beta} \alpha_{j-1k} + A_{jk+1jk}^{\alpha,\beta} \alpha_{jk+1} + A_{j-1k+1jk}^{\alpha,\beta} \alpha_{j-1k+1} \\ & = f_{jk}^\beta, \quad j = 1, \dots, n_x, k = 0, \dots, n_y, \end{aligned} \quad (3.4b)$$

where $A_{j'k'jk}^{\alpha,\alpha} = a_h(\phi_{j'k'}, \phi_{jk})$, $A_{j'k'jk}^{\alpha,\beta} = a_h(\phi_{j'k'}, \psi_{jk})$, $A_{j'k'jk}^{\beta,\alpha} = a_h(\psi_{j'k'}, \phi_{jk})$, $A_{j'k'jk}^{\beta,\beta} = a_h(\psi_{j'k'}, \psi_{jk})$, $f_{jk}^\alpha = (f, \phi_{jk})$ and $f_{jk}^\beta = (f, \psi_{jk})$.

A direct computation of the component of the stiffness matrix on $\Omega_{jk} = (x_{j-1}, x_j) \times (y_{k-1}, y_k)$ gives

$$(\nabla \phi_{jk}, \nabla \phi_{jk})_{\Omega_{jk}} = \frac{37 h_{x_j}}{28 h_{y_k}} + \frac{65 h_{y_k}}{28 h_{x_j}}, \quad (3.5a)$$

$$(\nabla \psi_{jk}, \nabla \phi_{jk})_{\Omega_{jk}} = (\nabla \psi_{jk-1}, \nabla \phi_{jk})_{\Omega_{jk}} = -\frac{37 h_{x_j}^2 + h_{y_k}^2}{28 h_{x_j} h_{y_k}}, \quad (3.5b)$$

$$(\nabla \phi_{j-1k}, \nabla \phi_{jk})_{\Omega_{jk}} = \frac{37 h_{x_j}}{28 h_{y_k}} + \frac{9 h_{y_k}}{28 h_{x_j}}. \quad (3.5c)$$

Analogous components are obtained by replacing Ω_{jk} by Ω_{jk-1} . Furthermore, we have similar results for ψ_{jk} ; just h_{x_j} and h_{y_k} are exchanged in (3.5). Set $\gamma_{jk} = \frac{h_{y_k}}{h_{x_j}}$. By a direct computation, one gets the following expressions:

$$\begin{aligned} A_{jkjk}^{\beta,\beta} &= \left(\frac{65}{28} \frac{1}{\gamma_{jk}} + \frac{37}{28} \gamma_{jk} \right) \kappa_{jk} + \left(\frac{65}{28} \frac{1}{\gamma_{jk+1}} + \frac{37}{28} \gamma_{jk+1} \right) \kappa_{jk+1}, \\ A_{jk-1jk}^{\beta,\beta} &= \left(\frac{9}{28} \frac{1}{\gamma_{jk}} + \frac{37}{28} \gamma_{jk} \right) \kappa_{jk}, \\ A_{jk+1jk}^{\beta,\beta} &= \left(\frac{9}{28} \frac{1}{\gamma_{jk+1}} + \frac{37}{28} \gamma_{jk+1} \right) \kappa_{jk+1}, \\ A_{jkjk}^{\alpha,\beta} &= A_{j-1kjk}^{\alpha,\beta} = -\frac{37}{28} \left(\frac{1}{\gamma_{jk}} + \gamma_{jk} \right) \kappa_{jk}, \\ A_{jk+1jk}^{\alpha,\beta} &= A_{j-1k+1jk}^{\alpha,\beta} = -\frac{37}{28} \left(\frac{1}{\gamma_{jk+1}} + \gamma_{jk+1} \right) \kappa_{jk+1}. \end{aligned}$$

First we need to deduce the coefficient values κ_{jk} and mesh sizes h_{x_j}, h_{y_k} from the micro-scale linear system (3.4). The result is formulated as the following proposition.

Proposition 3.1. κ_{jk} and h_{x_j}, h_{y_k} can be determined from the linear system (3.4):

$$\kappa_{jk} = -(A_{jk-1jk}^{\beta,\beta} + A_{jkjk}^{\alpha,\beta}) \gamma_{jk}, \quad (3.6a)$$

$$h_{x_j} = \frac{1}{\sum_{k=1}^{n_y} \gamma_{jk}}, \quad h_{y_k} = \frac{1}{\sum_{j=1}^{n_x} \frac{1}{\gamma_{jk}}}, \quad (3.6b)$$

$$\gamma_{jk} = \sqrt{\frac{28}{37} \left(\frac{A_{jkjk}^{\alpha,\beta}}{A_{jk-1jk}^{\beta,\beta} + A_{jkjk}^{\alpha,\beta}} \right) - 1}. \quad (3.6c)$$

Proof. At each rectangular elements, except for the 4 corner elements, we can derive at least two information about κ_{jk} from the stiffness matrix. One is $A_{j-1kjk}^{\alpha,\alpha}$ or $A_{jk-1jk}^{\beta,\beta}$ and the other is one of $A_{jkjk}^{\alpha,\beta}$, $A_{jkjk-1}^{\alpha,\beta}$, $A_{j-1kjk}^{\alpha,\beta}$, $A_{j-1kjk-1}^{\alpha,\beta}$. For example, when we have $A_{jk-1jk}^{\beta,\beta}$ and $A_{jkjk}^{\alpha,\beta}$ for left vertical element, we can derive the following equalities:

$$A_{jk-1jk}^{\beta,\beta} + A_{jkjk}^{\alpha,\beta} = -\frac{\kappa_{jk}}{\gamma_{jk}}, \quad \frac{A_{jkjk}^{\alpha,\beta}}{A_{jk-1jk}^{\beta,\beta} + A_{jkjk}^{\alpha,\beta}} = \frac{37}{28}(1 + \gamma_{jk}^2).$$

Hence κ_{jk} and γ_{jk} is derived by

$$\kappa_{jk} = -(A_{jk-1jk}^{\beta,\beta} + A_{jkjk}^{\alpha,\beta}) \gamma_{jk}, \quad \gamma_{jk} = \sqrt{\frac{28}{37} \left(\frac{A_{jkjk}^{\alpha,\beta}}{A_{jk-1jk}^{\beta,\beta} + A_{jkjk}^{\alpha,\beta}} \right) - 1}.$$

Other cases follows in a similar way.

At the corner, we cannot get the value of $A_{j-1kjk}^{\alpha,\alpha}$ or $A_{jk-1jk}^{\beta,\beta}$ from the stiffness matrix. That is, there is only one valid information about κ_{jk} and γ_{jk} . In this case, we need the ratio information from adjacent elements and this is the reason why we adopt the rectangular mesh. First, we can derive γ_{jk} using following relation about ratio, $\gamma_{jk} = \frac{\gamma_{jk+1}\gamma_{j+1k}}{\gamma_{j+1k+1}}$. Since the above formula are valid at every micro element except corners, three of γ_{jk} , γ_{jk+1} , γ_{j+1k} , γ_{j+1k+1} are known and the unknown one would be determined by the ratio information. Then κ_{jk} can be easily derived from the ratio information and one of $A_{jkjk}^{\alpha,\beta}$, $A_{jkjk-1}^{\alpha,\beta}$, $A_{jkjk}^{\beta,\alpha}$, $A_{jkj-1k}^{\beta,\alpha}$. Now we have every κ_{jk} and γ_{jk} value across all elements, and mesh sizes h_{x_j} and h_{y_k} are determined by following equations:

$$\sum_{k=1}^{n_y} \gamma_{jk} = \frac{\sum_{k=1}^{n_y} h_{y_j}}{h_{x_j}} = \frac{1}{h_{x_j}}, \quad \sum_{j=1}^{n_x} \frac{1}{\gamma_{jk}} = \frac{\sum_{j=1}^{n_x} h_{x_j}}{h_{y_k}} = \frac{1}{h_{y_k}}.$$

This completes the proof. \square

3.1 Construction of GMsFE spaces

In this section, we present the detailed procedure for constructing GMsFE spaces using the approximated values κ_{jk} , h_{x_j} , and h_{y_k} , obtained as in (3.6). Denote by ϕ_{jk}^T and ψ_{jk}^T the basis functions on macro element $T \in \mathcal{T}^H$ associated with macro vertical and horizontal edges, respectively.

3.2 Snapshot function space V^{snap}

We first construct local snapshot function space $V^{\text{snap}}(T)$ in each macro element $T \in \mathcal{T}^H$. Since snapshot functions are used to compute multiscale basis functions, we may choose $V^{\text{snap}}(T)$ as all micro-scale basis functions in T . Or smaller space such as the span of κ -harmonic functions in T can be considered to reduce the cost of constructing V^H .

Let $\tilde{u}_l^T \in V_h(T)$ be the solutions of following local κ -harmonic problems:

$$\begin{cases} -\nabla \cdot (\kappa(\mathbf{x})\nabla \tilde{u}_l^T) = 0 & \text{in } T, \\ \tilde{u}_l^T = \delta_l^T & \text{on } \partial T, \end{cases} \quad (3.7)$$

where $\delta_l^T \in V_h(T)$ is the function which equals to one for the j -th micro-scale mesh DOF on ∂T and zeros for the other DOFs on ∂T . Observe that δ_l^T is one of $\phi_{0k}^T, \phi_{n_x^T k}^T, \psi_{j0}^T, \psi_{jn_y^T}^T$ for $j = 1, \dots, n_x^T$ and $k = 1, \dots, n_y^T$. That is, \tilde{u}_l^T is the solution of

$$a_T(\tilde{u}_l^T, v^T) = 0 \quad \forall v^T \in V_{h,0}(T) \quad (3.8)$$

satisfying $\tilde{u}_l^T - \delta_l^T \in V_{h,0}(T)$, where $a_T(u, v) = \sum_{T_j \in T} \int_{T_j} \kappa \nabla u \cdot \nabla v \, d\mathbf{x}$.

Let $\tilde{u}_l^T \in V_h(T)$ is represented by

$$\tilde{u}_l^T = \sum_{j'=0}^{n_x^T} \sum_{k'=1}^{n_y^T} \alpha_{j'k'}^T \phi_{j'k'}^T + \sum_{j'=1}^{n_x^T} \sum_{k'=0}^{n_y^T} \beta_{j'k'}^T \psi_{j'k'}^T.$$

Then (3.8) leads to the following equations by setting $v^T = \phi_{jk}^T$ and $v^T = \psi_{jk}^T$:

$$\begin{aligned} \sum_{j'=0}^{n_x^T} \sum_{k'=1}^{n_y^T} \alpha_{j'k'}^T a_T(\phi_{j'k'}^T, \phi_{jk}^T) + \sum_{j'=1}^{n_x^T} \sum_{k'=0}^{n_y^T} \beta_{j'k'}^T a_T(\psi_{j'k'}^T, \phi_{jk}^T) &= 0, \\ j = 0, \dots, n_x^T, k = 1, \dots, n_y^T, \end{aligned} \quad (3.9a)$$

$$\begin{aligned} \sum_{j'=0}^{n_x^T} \sum_{k'=1}^{n_y^T} \alpha_{j'k'}^T a_T(\phi_{j'k'}^T, \psi_{jk}^T) + \sum_{j'=1}^{n_x^T} \sum_{k'=0}^{n_y^T} \beta_{j'k'}^T a_T(\psi_{j'k'}^T, \psi_{jk}^T) &= 0, \\ j = 1, \dots, n_x^T, k = 0, \dots, n_y^T. \end{aligned} \quad (3.9b)$$

If we take the supports of basis functions into consideration, we have the following linear system for the snapshot function \tilde{u}_l^T :

$$\begin{aligned} &\tilde{A}_{jkjk}^{\alpha,\alpha} \alpha_{jk} + \tilde{A}_{j-1kj}^{\alpha,\alpha} \alpha_{j-1k} + \tilde{A}_{j+1kj}^{\alpha,\alpha} \alpha_{j+1k} \\ &+ \tilde{A}_{jkjk}^{\beta,\alpha} \beta_{jk} + \tilde{A}_{jk-1jk}^{\beta,\alpha} \beta_{jk-1} + \tilde{A}_{j+1kj}^{\beta,\alpha} \beta_{j+1k} + \tilde{A}_{j+1k-1jk}^{\beta,\alpha} \beta_{j+1k-1} \\ &= 0, \quad j = 0, \dots, n_x^T, k = 1, \dots, n_y^T, \quad (3.10a) \\ &\tilde{A}_{jkjk}^{\beta,\beta} \beta_{jk} + \tilde{A}_{jk-1jk}^{\beta,\beta} \beta_{jk-1} + \tilde{A}_{jk+1jk}^{\beta,\beta} \beta_{jk+1} \\ &+ \tilde{A}_{jkjk}^{\alpha,\beta} \alpha_{jk} + \tilde{A}_{j-1kj}^{\alpha,\beta} \alpha_{j-1k} + \tilde{A}_{jk+1jk}^{\alpha,\beta} \alpha_{jk+1} + \tilde{A}_{j-1k+1jk}^{\alpha,\beta} \alpha_{j-1k+1} \end{aligned}$$

$$= 0, \quad j = 1, \dots, n_x^T, \quad k = 0, \dots, n_y^T, \quad (3.10b)$$

where $\tilde{A}_{j'k'jk}^{\alpha,\alpha} = a_T(\phi_{j'k'}^T, \phi_{jk}^T)$, $\tilde{A}_{j'k'jk}^{\alpha,\beta} = a_T(\phi_{j'k'}^T, \psi_{jk}^T)$, $\tilde{A}_{j'k'jk}^{\beta,\alpha} = a_T(\psi_{j'k'}^T, \phi_{jk}^T)$, and $\tilde{A}_{j'k'jk}^{\beta,\beta} = a_T(\psi_{j'k'}^T, \psi_{jk}^T)$. Since each component of the system (3.10) can be computed from the approximate values of κ_{jk} and h_{x_j}, h_{y_k} as given in (3.6), we can compute the snapshot function \tilde{u}_l^T from the matrix components $A_{jklm}^{\alpha,\beta}$'s and $A_{jklm}^{\beta,\beta}$'s in the linear system (3.4).

Denote the number of all snapshot functions in T by $\mathcal{N}^{\text{snap}}(T)$ and zero extension of \tilde{u}_l^T outside T by u_l^T . Then the local snapshot function space $V^{\text{snap}}(T)$ is defined as the space spanned by u_l^T :

$$V^{\text{snap}}(T) = \text{Span} \left\{ u_l^T \in V_h(T) \mid j = 1, \dots, \mathcal{N}^{\text{snap}}(T) \right\}.$$

Then the snapshot function space V^{snap} is defined as the union of such local snapshot function spaces:

$$V^{\text{snap}} = \bigcup_{T \in \mathcal{T}^H} V^{\text{snap}}(T).$$

3.2.1 Oversampling technique

The oversampling technique reduces the resonance error caused by wrong (local) boundary condition δ_l^T . It consist of the restriction of the solution $u_l^{T^+}$ of (3.7) on an extended region T^+ to the original domain T . We denote the local oversampled snapshot function space by $V^{\text{snap},+}(T^+)$, which is defined as

$$V^{\text{snap},+}(T^+) = \text{Span} \left\{ u_l^{T^+} \in V_h(T^+) \mid j = 1, \dots, \mathcal{N}^{\text{snap},+}(T^+) \right\}.$$

Then the oversampled snapshot function space $V^{\text{snap},+}$ is given as follows:

$$V^{\text{snap},+} = \bigcup_{T \in \mathcal{T}^H} V^{\text{snap},+}(T^+).$$

Again we remark that the oversampled snapshot function space is completely characterized by the information on the linear system (3.4).

3.3 Offline function space V^{off}

An offline function space V^{off} is obtained by applying a suitable dimension reduction technique to the snapshot function space V^{snap} . For example, we may use the generalized eigenvalue decomposition. For each macro element $T \in \mathcal{T}^H$, consider the following spectral problem to find $(\lambda_l^T, u_l^T) \in \mathbb{R} \times V^{\text{snap}}(T)$:

$$a_T(u_l^T, v^T) = \lambda_l^T (\kappa u_l^T, v^T)_T, \quad \forall v^T \in V^{\text{snap}}(T). \quad (3.11)$$




Figure 3.1: Multiscale mesh on Ω . T_1, T_2 are macro elements and $\omega(E)^+$ is an oversampled neighborhood of macro edge E .

Since any function $v \in V^{\text{snap}}(T)$ is represented by

$$v^T = \sum_{j'=0}^{n_x^T} \sum_{k'=1}^{n_y^T} \alpha_{j'k'}^T \phi_{j'k'}^T + \sum_{j'=1}^{n_x^T} \sum_{k'=0}^{n_y^T} \beta_{j'k'}^T \psi_{j'k'}^T,$$

we can construct the linear system of (3.11) and calculate the offline function u_l^T from the information on the linear system (3.4).

We suppose that the eigenvalues are sorted in ascending order as

$$0 \leq \lambda_1^T \leq \lambda_2^T \leq \dots \leq \lambda_{\mathcal{N}^{\text{snap}}(T)},$$

and the eigenfunctions are normalized by $(\kappa u_l^T, u_l^T) = 1$. Then the local offline function space $V^{\text{off}}(T)$ is defined as the space spanned by a number of dominant eigenfunctions u_l^T , which is related to l -th smallest eigenvalue λ_l^T . We may choose $\mathcal{L}(T)$ eigenfunctions, where $\mathcal{L}(T)$ is considerably small number compared to $\mathcal{N}^{\text{snap}}(T)$. In short, $V^{\text{off}}(T)$ is given by

$$V^{\text{off}}(T) = \text{Span} \left\{ u_l^T \in V^{\text{snap}}(T) \mid l = 1, \dots, \mathcal{L}(T) \right\},$$

and the offline function space V^{off} is defined as

$$V^{\text{off}} = \bigcup_{T \in \mathcal{T}^H} V^{\text{off}}(T).$$

3.4 Moment function space \mathcal{M}_H

Since the offline functions are defined independently in each macro element $T \in \mathcal{T}^H$, we need to glue those functions through each macro interior edge $E \in \mathcal{E}^{H,0}$. Moment functions play an important role here, as they are used to impose continuity between offline functions in neighboring macro elements. On each macro edge E , let $\omega(E)^+$ be an oversampled neighborhood of E . As we

construct local snapshot space, the moment function $\zeta_l^E \in V_h(\omega(E)^+)$ can be obtained by solving the following local κ -harmonic problem:

$$\begin{cases} -\nabla \cdot (\kappa(\mathbf{x})\nabla \zeta_l^E) = 0 \text{ in } \omega(E)^+, \\ \zeta_l^E = \delta_l^E \text{ on } \partial\omega(E)^+, \end{cases} \quad (3.12)$$

where $\delta_l^E \in V_h(\omega(E)^+)$ is the function which equals to one for the l -th micro-scale mesh DOF on $\partial\omega(E)^+$ and zeros for the other DOFs. We can construct ζ_l^E as we build the snapshot functions, by replacing T to $\omega(E)^+$ in Section 3.2. We collect the traces of ζ_l^E on E and perform a singular value decomposition to them. Denote $m(E)$ linearly independent singular vectors by s_k^E , where s_k^E is arranged in descending order with respect to its norm:

$$\|s_k^E\|_E^2 = \mu_k^E, \quad \mu_1^E \geq \mu_2^E \geq \dots \geq \mu_{m(E)}^E > 0. \quad (3.13)$$

Then the local moment function space $\mathcal{M}_H(E)$ on E is given by

$$\mathcal{M}_H(E) = \text{Span} \left\{ s_k^E \mid 1 \leq k \leq \mathcal{L}(E) \right\},$$

and the moment function space \mathcal{M}_H is defined as

$$\mathcal{M}_H = \bigcup_{E \in \mathcal{E}^H} \mathcal{M}_H(E).$$

It is straightforward to see that the calculation of the moment function space \mathcal{M}_H is completely dependent on the micro-scale linear system (3.4).

3.4.1 Another method for constructing moment function space

We may consider another method for constructing moment function space in order to reduce the computational cost. That is, the moment function space can be made up of the traces of the snapshot functions. For each macro edge $E \in \mathcal{E}^H$, denote the collection of such traces by

$$\mathcal{M}_h(E) := \text{Span} \left\{ u^T|_E \mid u^T \in V^{\text{snap}}(T), E \subset \partial T \right\}.$$

We perform a singular value decomposition to $\mathcal{M}_h(E)$ and choose the first $\mathcal{L}(E)$ dominant modes of $\mathcal{M}_h(E)$, which span the local moment function space. This method makes us avoid to solve local boundary value problems (3.12).

3.5 Nonconforming GMsFE spaces V^H and $V^{H,0}$

The nonconforming GMsFE spaces V^H and $V^{H,0}$ are defined as

$$V^H = \left\{ u \in V^{\text{off}} \mid \langle [u]_E, \zeta \rangle_E = 0, \forall \zeta \in \mathcal{M}_H(E), \forall E \in \mathcal{E}^{H,0} \right\}, \quad (3.14)$$

$$V^{H,0} = \left\{ u \in V^{\text{off}} \mid \langle [u]_E, \zeta \rangle_E = 0, \forall \zeta \in \mathcal{M}_H(E), \forall E \in \mathcal{E}^H \right\}, \quad (3.15)$$

where $[u]_E$ denotes the jump of u across the edge E . Since V^{off} and \mathcal{M}_H are defined as the union of local function spaces, it is possible to construct V^H and $V^{H,0}$ locally. Let $E \in \mathcal{E}^{H,0}$ be a common macro edge for two macro elements T_1 and T_2 (see Figure 3.1.) Suppose that the local moment function space $\mathcal{M}_H(E)$ is constructed from κ -harmonic functions in $\omega(E) := T_1 \cup T_2$. Then the continuity condition for $u \in V^{\text{off}}(T_1) \cup V^{\text{off}}(T_2)$ imposed by $\mathcal{M}_H(E)$ is given as follows:

$$\begin{aligned} \langle [u]_E, \zeta \rangle_E &= 0, \quad \forall \zeta \in \mathcal{M}_H(E), \\ \langle u, \zeta \rangle_{E'} &= 0, \quad \forall \zeta \in \mathcal{M}_H(E'), \quad \forall E' \subset \partial\omega(E). \end{aligned} \quad (3.16)$$

Finally we define the local GMsFE space $V^H(\omega(E))$ as

$$V^H(\omega(E)) = \left\{ u \in V^{\text{off}}(T_1) \cup V^{\text{off}}(T_2) \mid u \text{ satisfies (3.16)} \right\}.$$

Then the GMsFE space $V^{H,0}$ can be obtained by

$$V^{H,0} = \bigcup_{E \in \mathcal{E}^{H,0}} V^H(\omega(E)).$$

V^H also can be constructed similarly by considering $E \in \mathcal{E}^H$ in the above argument.

Remark 3.2. *It is remarkable that there may exist macro bubble functions $u \in V^{\text{off}}(T)$ on T , which satisfy*

$$\langle u, \zeta \rangle_{E'} = 0, \quad \forall \zeta \in \mathcal{M}_H(E'), \quad \forall E' \subset \partial T. \quad (3.17)$$

Denote the space of macro bubble functions on T_j by

$$B_H(T) = \left\{ u \in V^{\text{off}}(T) \mid u \text{ satisfies (3.17)} \right\}.$$

Then the GMsFE space $V^{H,0}$ is obtained by

$$V^{H,0} = \left(\bigcup_{T \in \mathcal{T}^H} B_H(T) \right) \cup \left(\bigcup_{E \in \mathcal{E}^{H,0}} V^H(\omega(E)) \right).$$

Remark 3.3. *The dimension of GMsFE space $V^{H,0}$ may depend on the dimension of local moment function space. For each macro element $T \in \mathcal{T}^H$, we practically take the dimension of local offline function space as*

$$\mathcal{L}(T) = \sum_{E' \subset \partial T} \mathcal{L}(E').$$

Then the dimension of $V^H(\omega(E))$ is given by

$$\dim \left(V^H(\omega(E)) \right) \geq \mathcal{L}(T_1) + \mathcal{L}(T_2) - \mathcal{L}(E) - \sum_{E' \subset \partial\omega(E)} \mathcal{L}(E') = \mathcal{L}(E).$$

If there are no macro bubble functions, it follows that

$$\dim(V^{H,0}) = \sum_{E \in \mathcal{E}^{H,0}} \mathcal{L}(E).$$

3.5.1 Construction of b^M

Now we have the GMsFE space $V^{H,0} = \text{Span}(\{\psi_{JK}^L\}_{J,K,L} \cup \{\phi_{JK}^L\}_{J,K,L})$, where ψ_{JK}^L and ϕ_{JK}^L denote the L -th multiscale basis function associated with the JK -th horizontal macro edge f_{JK} and JK -th vertical macro edge e_{JK} , respectively. Suppose that $\Omega_{JK} \cup \Omega_{J+1K}$ is composed of $n_x^{JK} \times n_y^{JK}$ micro-scale elements. Then ϕ_{JK}^L is represented by

$$\phi_{JK}^L = \sum_{j'=0}^{n_x^{JK}} \sum_{k'=1}^{n_y^{JK}} \alpha_{j'k'}^{JK} \phi_{j'k'}^{JK} + \sum_{j'=1}^{n_x^{JK}} \sum_{k'=0}^{n_y^{JK}} \beta_{j'k'}^{JK} \psi_{j'k'}^{JK},$$

where $\phi_{j'k'}^{JK}$ and $\psi_{j'k'}^{JK}$ are the micro-scale basis functions of vertical and horizontal type in $\Omega_{JK} \cup \Omega_{J+1K}$, respectively. Therefore it is obvious that the components of b^M can be derived from the summation of that of micro-scale right hand side b , which is given by f_{jk}^α and f_{jk}^β in (3.4). The same argument holds for ψ_{JK}^L , which completes the construction of b^M .

4 Numerical results

Example 4.1. Consider the following elliptic problem:

$$\begin{cases} -\nabla \cdot (\kappa(\mathbf{x}) \nabla u) = f & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (4.1)$$

where $\Omega = (0, 1)^2$ and $\kappa(\mathbf{x}) = 1 + (1 + x_1)(1 + x_2) + \epsilon \sin(10\pi x_1) \sin(5\pi x_2)$. The source term f is generated by the exact solution

$$u(x_1, x_2) = \sin(3\pi x_1) x_2 (1 - x_2) + \epsilon \sin(\pi x_1 / \epsilon) \sin(\pi x_2 / \epsilon).$$

We compare numerical results of GMsFEM and AMS(algebraic multiscale method). We use uniform rectangular meshes and the ratio H/h is fixed as 10. The AMS solutions are calculated from the information on the linear system $Ax = b$ obtained by the application of the finite element method based on the DSSY nonconforming element for the elliptic problem (4.1).

The relative energy and L^2 errors for the macro-scale solutions are reported for various ϵ in Tables 4.1–3. We observe almost same error behaviors in both methods.

5 Conclusion

In this paper, we design the algebraic multiscale method for two-dimensional elliptic problems. The generalized multiscale finite element method is used based on the DSSY nonconforming finite element space. The macro-scale linear system is constructed only using the algebraic information on the components of the micro-scale system. The proposed method shows almost identical numerical results with those obtained by the GMsFEM.

$\frac{1}{H}$	$\frac{1}{h}$	$\dim(V^{H,0})$	GMsFEM		AMS	
			Rel. Energy	Rel. L^2	Rel. Energy	Rel. L^2
5	50	400	0.884	0.388	0.884	0.389
10	100	1800	0.871	0.363	0.871	0.363
20	200	7600	0.346	0.676E-01	0.346	0.678E-01
40	400	31200	0.181	0.181E-01	0.181	0.182E-01

Table 4.1: Convergence for $\epsilon = 0.1$.

$\frac{1}{H}$	$\frac{1}{h}$	$\dim(V^{H,0})$	GMsFEM		AMS	
			Rel. Energy	Rel. L^2	Rel. Energy	Rel. L^2
5	50	400	0.885	0.625	0.885	0.625
10	100	1800	0.355	0.118	0.355	0.119
20	200	7600	0.186	0.316E-01	0.186	0.320E-01
40	400	31200	0.940E-01	0.803E-02	0.939E-01	0.823E-02

Table 4.2: Convergence for $\epsilon = 0.2$.

$\frac{1}{H}$	$\frac{1}{h}$	$\dim(V^{H,0})$	GMsFEM		AMS	
			Rel. Energy	Rel. L^2	Rel. Energy	Rel. L^2
5	50	400	0.335	0.130	0.335	0.132
10	100	1800	0.173	0.342E-01	0.173	0.351E-01
20	200	7600	0.884E-01	0.879E-02	0.885E-02	0.928E-02
40	400	31200	0.444E-01	0.221E-02	0.444E-01	0.246E-02

Table 4.3: Convergence for $\epsilon = 0.5$.

Acknowledgment

DS was supported in part by National Research Foundation of Korea (NRF-2017R1A2B3012506 and NRF-2015M3C4A7065662).

References

- [1] A. Abdulle. The finite element heterogeneous multiscale method: a computational strategy for multiscale pdes. *GAKUTO International Series Mathematical Sciences and Applications*, 31(EPFL-ARTICLE-182121):135–184, 2009.
- [2] A. Abdulle, E. Weinan, B. Engquist, and E. Vanden-Eijnden. The heterogeneous multiscale method. *Acta Numerica*, 21:1–87, 2012.
- [3] K. Cho, R. Lim, and D. Sheen. Algebraic multiscale methods for one-dimensional elliptic problems. *this journal*, 2002.




Figure 4.1: Algebraic multiscale solution of $\epsilon = 0.2$ when $1/H = 40, 1/h = 400$.




Figure 4.2: Micro-Scale reference solution of $\epsilon = 0.2$ when $1/h = 400$.

- [4] K. Cho and M. Moon. Multiscale hybridizable discontinuous galerkin method for elliptic problems in perforated domains. *Journal of Computational and Applied Mathematics*, 365:112346, 2020.
- [5] J. Douglas Jr, J. E. Santos, D. Sheen, and X. Ye. Nonconforming galerkin

- methods based on quadrilateral elements for second order elliptic problems. *ESAIM: Mathematical Modelling and Numerical Analysis*, 33(4):747–770, 1999.
- [6] Y. Efendiev, J. Galvis, and T. Y. Hou. Generalized multiscale finite element methods (gmsfem). *Journal of Computational Physics*, 251:116–135, 2013.
- [7] Y. Efendiev and T. Y. Hou. *Multiscale finite element methods: theory and applications*, volume 4. Springer Science & Business Media, 2009.
- [8] Y. Efendiev, R. Lazarov, M. Moon, and K. Shi. A spectral multiscale hybridizable discontinuous galerkin method for second order elliptic problems. *Computer Methods in Applied Mechanics and Engineering*, 292:243–256, 2015.
- [9] Y. Efendiev, R. Lazarov, and K. Shi. A multiscale HDG method for second order elliptic equations. Part I. Polynomial and homogenization-based multiscale spaces. *SIAM Journal on Numerical Analysis*, 53(1):342–369, 2015.
- [10] Y. R. Efendiev, T. Y. Hou, and X.-H. Wu. Convergence of a nonconforming multiscale finite element method. *SIAM Journal on Numerical Analysis*, 37(3):888–910, 2000.
- [11] T. Hou, X.-H. Wu, and Z. Cai. Convergence of a multiscale finite element method for elliptic problems with rapidly oscillating coefficients. *Mathematics of Computation of the American Mathematical Society*, 68(227):913–943, 1999.
- [12] C. S. Lee and D. Sheen. Nonconforming generalized multiscale finite element methods. *Journal of Computational and Applied Mathematics*, 311:215–229, 2017.
- [13] C. Park and D. Sheen. p_1 -nonconforming quadrilateral finite element methods for second-order elliptic problems. *SIAM Journal on Numerical Analysis*, 41(2):624–640, 2003.