

A TOTAL CUNTZ SEMIGROUP FOR C^* -ALGEBRAS OF STABLE RANK ONE

QINGNAN AN AND ZHICHAO LIU

ABSTRACT. In this paper, we show that the unitary Cuntz semigroup is not a complete invariant for AD algebras of real rank zero. Moreover, we prove that for separable, unital C^* -algebras with stable rank one and real rank zero, the unitary Cuntz semigroup can be recovered functorially from the invariant K_* . Then we introduce the refinement of unitary Cuntz semigroup, say the total Cuntz semigroup, which is a new invariant for separable C^* -algebras of stable rank one, is a well-defined continuous functor from C^* -algebras to the category $\underline{\text{Cu}}^\sim$. Furthermore, we show that the total Cuntz semigroup is a complete invariant for a large class of C^* -algebras of real rank zero.

INTRODUCTION

Recently, Cantier introduced the unitary Cuntz semigroup, denote by Cu_1 , as an invariant which captures the K_1 information [7]. As an application, the author obtained the classification theorem for real rank zero AD algebras (which is called AH_d algebras in [7]) in terms of unitary Cuntz semigroup. In this paper, we present an example of two non-isomorphic AD algebras of real rank zero with isomorphic unitary Cuntz semigroup, therefore Corollary 5.21 in [7] is not correct. The author used Theorem 7.3 of [12] to get the corollary. However it was pointed out by Elliott-Gong-Su (See [15]), Theorem 7.3 (and Theorem 7.1) of [12] is only true for simple case (See example 2.19 of [15]). We will also prove the following theorem which is a converse of Theorem 5.20 of [7] for real rank zero case.

Theorem. 2.9. *Let A, B be separable, unital C^* -algebra with stable rank one and real rank zero. Then*

$$(\text{Cu}_1(A), ([1_A, 0])) \cong (\text{Cu}_1(B), ([1_B, 0]))$$

if and only if

$$(K_*(A), K_*^+(A), [1_A]) \cong (K_*(B), K_*^+(B), [1_B]).$$

Since Cu_1 is not a complete invariant for AD algebras of real rank zero, we introduce the refinement of unitary Cuntz semigroup, denoted by $\underline{\text{Cu}}$, and establish the functorial properties, we show that $\underline{\text{Cu}}$ is a continuous functor

2000 *Mathematics Subject Classification.* Primary 46L35, Secondary 46K80 19K35.

Key words and phrases. Total Cuntz semigroup; Total K-theory; Stable rank one; Invariant.

from the category of separable C^* -algebras with stable rank one, denoted by C_{sr1}^* , to a subcategory of Cu^\sim . We have

Theorem. 4.5. *Let $A = \varinjlim (A_i, \phi_{ij})$ be an inductive system in the category C_{sr1}^* . Then*

$$\underline{\text{Cu}}\text{-}\varinjlim \underline{\text{Cu}}(A_i) \cong \underline{\text{Cu}}(A).$$

Then we show that the total K -theory \underline{K} can be recovered functorially from $\underline{\text{Cu}}$. Furthermore, we prove that the total Cuntz semigroup can also be recovered from the total K -theory for a large class of C^* -algebras.

Theorem. 5.9. *Let A, B be AHD algebras with stable rank one and real rank zero. Then*

$$(\underline{\text{Cu}}(A), ([1_A], 0, \oplus_{p=1}^\infty (0, 0))) \cong (\underline{\text{Cu}}(B), ([1_B], 0, \oplus_{p=1}^\infty (0, 0))).$$

if and only if

$$(\underline{K}(A), \underline{K}(A)_+, [1_A]) \cong (\underline{K}(B), \underline{K}(B)_+, [1_B]).$$

Note that the total K -theory \underline{K} is a complete invariant for AHD algebras (including AD algebras and AH algebras with slow dimension growth, see [17]) of real rank zero, then $\underline{\text{Cu}}_u$ is also a complete invariant for this class. The question whether all the separable, unital C^* -algebra with stable rank one and real rank zero can be classified by \underline{K} is still open. Furthermore, there are also many interesting consequences for the structure of C^* -algebras of stable rank one and their Cuntz semigroups ([2, 22]).

1. PRELIMINARIES

1.1. Let A be a unital C^* -algebra. A is said to have stable rank one, written $sr(A) = 1$, if the set of invertible elements of A is dense. A is said to have real rank zero, written $rr(A) = 0$, if the set of invertible self-adjoint elements is dense in the set A_{sa} of self-adjoint elements of A . If A is not unital, let us denote the unitization of A by \tilde{A} . A non-unital C^* -algebra is said to have stable rank one (or real rank zero) if its unitization has stable rank one (or real rank zero). Denote by C_{sr1}^* the category of separable C^* -algebras with stable rank one.

1.2 ([12]). AD algebra is an inductive limit of finite direct sums of the form $M_n(\tilde{I}_p)$ and $M_n(C(X))$, where

$$I_p = \{f \in C([0, 1], M_p(\mathbb{C})) \mid f(0) = 0 \text{ and } f(1) \in \mathbb{C} \cdot 1_p\}$$

is the Elliott-Thomsen dimension-drop interval algebra and X is one of the following finite connected CW complexes: $\{pt\}$, \mathbb{T} , $[0, 1]$. (In [7, 5.4], AD algebra is called AH_d -algebra.)

1.3. (The Cuntz semigroup of a C^* -algebra) Denote the cone of positive elements in A by A_+ . Let $a, b \in A_+$. One says that a is *Cuntz subequivalent* to b , denoted by $a \precsim b$, if there exists a sequence (r_n) in A such that $r_n^* b r_n \rightarrow$

a . One says that a is *Cuntz equivalent* to b , denoted by $a \sim b$, if $a \precsim b$ and $b \precsim a$. The *Cuntz semigroup* of A is defined as $\text{Cu}(A) = (A \otimes \mathcal{K})_+ / \sim$. We will denote the class of $a \in (A \otimes \mathcal{K})_+$ in $\text{Cu}(A)$ by $\langle a \rangle$.

1.4. (The category Cu) Let (S, \leq) be a positively ordered semigroup and let x, y in S . We say that x is way-below y , and we write $x \ll y$ if, for all increasing sequences in S that have a supremum, if $y \leq \sup_{n \in \mathbb{N}} z_n$, then there exists k such that $x \leq z_k$. This is an auxiliary relation on S , called the compact-containment relation and sometimes referred to as the way-below relation. In particular $x \ll y$ implies $x \leq y$ and we say that x is a compact element whenever $x \ll x$.

We say that S is an abstract Cu-semigroup if it satisfies the following order-theoretic axioms:

(O1): Every increasing sequence of elements in S has a supremum.

(O2): For any $x \in S$, there exists a \ll -increasing sequence $(x_n)_{n \in \mathbb{N}}$ in S such that $\sup_{n \in \mathbb{N}} x_n = x$.

(O3): Addition and the compact containment relation are compatible.

(O4): Addition and suprema of increasing sequences are compatible.

A Cu-morphism between two Cu-semigroups is a positively ordered monoid morphism that preserves the compact containment relation and suprema of increasing sequences.

1.5. Let A be a C^* -algebra and $\text{Lat}(A)$ denote the collection of ideals in A , equipped with the partial order given by inclusion of ideals. For any ideal I in A , then $\text{Cu}(I)$ is an ideal of $\text{Cu}(A)$ (See [3, 5.1] and [8, 3.1] for more details). For any $x \in \text{Cu}(A)$, write

$$I_x = \{y \in \text{Cu}(A) \text{ such that } y \leq \infty \cdot x\}$$

the ideal of $\text{Cu}(A)$ generated by x , where

$$\infty \cdot x = \langle \bigoplus_{n=1}^{\infty} \frac{1}{n} a \rangle \in \text{Cu}(A),$$

for any $[a] = x$, $a \in (A \otimes \mathcal{K})_+$.

For any separable C^* -algebra A , $I \rightarrow \text{Cu}(I)$ defines a lattice isomorphism between the lattice $\text{Lat}(A)$ of closed two-sided ideals of A and the lattice $\text{Lat}(\text{Cu}(A))$ of ideals of $\text{Cu}(A)$, denote this isomorphism by Γ , i.e.,

$$\Gamma : \text{Lat}(A) \rightarrow \text{Lat}(\text{Cu}(A)), \quad \Gamma(I) = \text{Cu}(I).$$

1.6. Let $A = \lim(A_n, \phi_{n,m})$ be an inductive limit of C^* -algebras. It is shown in [9, Theorem 2] that the Cuntz semigroup preserving the inductive limits of sequences (this was generalized to arbitrary inductive limits in [3, Corollary 3.2.9], see also [23, Lemma 3.8]), each element $s \in \text{Cu}(A)$ can be represented by an increasing sequence (s_1, s_2, \dots) , with $s_1 \in \text{Cu}(A_1)$, $s_2 \in \text{Cu}(A_2)$, \dots – increasing in the sense that for each i the image of $s_i \in \text{Cu}(A_i)$ in $\text{Cu}(A_{i+1})$ is less than or equal to s_{i+1} . Recall that two sequences $(s_i) \leq (t_i)$ if for any i and any $s \in S_i$ with $s \ll s_i$, eventually $s \ll t_j$ (in S_j). If $(s_i) \leq (t_i)$ and

$(t_i) \leq (s_i)$, then $(s_i) \sim (t_i)$, we denote the class of (s_i) in $\text{Cu}(A)$ by $[(s_i)]$, then

$$\text{Cu}(A) = \text{Cu} - \varinjlim \text{Cu}(A_i).$$

The following lemma combines [9, Theorem 2] and [3, Proposition 5.1.6].

Lemma 1.7. *Let $A = \varinjlim (A_n, \phi_{n,m})$, $s \in \text{Cu}(A)$ and (s_1, s_2, \dots) be an increasing sequence with $[(s_i)] = s$, where $s_i \in \text{Cu}(A_i)$ for any i . Then*

$$I_s \cong \text{Cu} - \varinjlim I_{s_i} \quad \text{and} \quad \Gamma^{-1}(I_s) = \varinjlim (\Gamma^{-1}(I_{s_i}), \phi_{m,n}),$$

where I_{s_i} is the ideal generated by s_i in $\text{Cu}(A_i)$.

Proof. The first part is from the definition of $\varinjlim I_{s_i}$. Every $y \in I_s$ is corresponding to an increasing sequence (y_i) with $[(y_i)] = y$, and $y \leq \infty \cdot s$ if and only if $[(y_i)] \leq \infty \cdot [(s_i)]$.

Now we prove the second part. By [8, Proposition 3.1.3], we know that the natural map Γ between $\text{Lat}(A)$ and $\text{Lat}(\text{Cu}(A))$ is an isomorphism of completed lattices. Since both $\Gamma^{-1}(I_s)$ and $\varinjlim \Gamma^{-1}(I_{s_i})$ are ideals in A , then $\Gamma \circ \Gamma^{-1}(I_s)$ and $\Gamma(\varinjlim \Gamma^{-1}(I_{s_i}))$ are ideals in $\text{Cu}(A)$.

Note that $\Gamma \circ \Gamma^{-1}(I_s) = I_s$ is generated by s in $\text{Cu}(A)$ and $\Gamma(\varinjlim \Gamma^{-1}(I_{s_i}))$ is generated by $[(s_1, s_2, \dots)]$ in $\text{Cu}(A)$, then we have

$$\Gamma \circ \Gamma^{-1}(I_s) = \Gamma(\varinjlim \Gamma^{-1}(I_{s_i})).$$

Hence,

$$\Gamma^{-1}(I_s) = \varinjlim \Gamma^{-1}(I_{s_i}).$$

□

Remark 1.8. In the rest parts, we will use I_s and $\Gamma^{-1}(I_s)$ frequently, in order to save notations, we will also use $I_s \triangleleft A$ to denote $\Gamma^{-1}(I_s) \triangleleft A$. and we may simply write $\Gamma^{-1}(I_s) = \varinjlim \Gamma^{-1}(I_{s_i})$ by $I_s = C^* - \varinjlim I_{s_i}$.

The following theorem is [4, Theorem 2.8].

Theorem 1.9. *If B is a full hereditary C^* -subalgebra of A and if each of A and B has a strictly positive element, then B is stably isomorphic to A .*

Then for any $a \in A_+$, the hereditary algebra $\text{Her } a$ generated by a and the ideal $\text{Ideal}(a)$ generated by a have the same K -theory. We will use this fact frequently.

1.10. (Unitary Cuntz semigroup) Recall the definition of unitary Cuntz semigroup introduced by Cantier in [7, 3.3]. Let A be a C^* -algebra of stable rank one and let

$$H(A) := \{(a, u) \mid a \in (A \otimes \mathcal{K})_+, u \in \mathcal{U}(\widetilde{\text{Her } a})\}.$$

Write $(a, u) \simeq_1 (b, v)$ if $a \precsim b$ and $[\theta_{ab}(u)] = [v]$ in $K_1(\text{Her } b)$, where θ_{ab} is an injection from $\text{Her } a$ to $\text{Her } b$, we say $(a, u) \sim_1 (b, v)$ if $(a, u) \simeq_1 (b, v)$ and $(b, v) \simeq_1 (a, u)$.

Define the unitary Cuntz semigroup of A by

$$\text{Cu}_1(A) := H(A) / \sim_1,$$

Let A be a C^* -algebra of stable rank one. Denote $\text{Lat}_f(A)$ the subset of $\text{Lat}(A)$ consisting of all those ideals of A that containing a full positive element. Hence, any ideal in $\text{Lat}_f(A)$ is singly-generated. Define $\text{Cu}_f(I) := \{\langle a \rangle \in \text{Cu}(A) \mid I_a = I\}$, $\text{Cu}_f(I)$ consists of the elements of $\text{Cu}(A)$ that are full in $\text{Cu}(I)$.

The following theorem illustrates the main structure of $\text{Cu}_1(A)$.

Theorem 1.11. ([7, Theorem 1.6]) *Let A be a C^* -algebra, then*

$$\text{Cu}_1(A) \cong \coprod_{I \in \text{Lat}(A)} \text{Cu}_f(I) \times K_1(I).$$

If there is an $I \in \text{Lat}(A) \setminus \text{Lat}_f(A)$, we have $\text{Cu}_f(I) = \emptyset$. Thus, one can replace the “ $\text{Lat}(A)$ ” in the above theorem by “ $\text{Lat}_f(A)$ ”. We mention that whenever A is separable, then $\text{Lat}(A) = \text{Lat}_f(A)$.

1.12. [3, 5.5] Denote Mon_{\leq} the category of ordered monoid and POM the category consisting of positive ordered monoid. Given an $M \in \text{POM}$, denote its Cu-completion by $\text{Cu}(M)$ (see more details in [3, 3.2]), Thus, we obtain a functor

$$\text{Cu} : \text{POM} \rightarrow \text{Cu}, \quad M \mapsto \text{Cu}(M), \quad \text{for all } M \in \text{POM},$$

mapping the POM-morphism $f : M \rightarrow N$ into the induced Cu-morphism $\text{Cu}(f) : \text{Cu}(M) \rightarrow \text{Cu}(N)$.

Conversely, give a Cu-semigroup S , we let S_c denote the set of compact elements in S . It is easy to see that S_c is a submonoid of S and we equip it with the order induced by S . Then we get another functor

$$\text{Re} : \text{Cu} \rightarrow \text{POM}, \quad S \mapsto S_c, \quad \text{for all } S \in \text{Cu},$$

It is obvious that the two functors Cu and Re establish an equivalence of the category POM and the full subcategory of Cu consisting of algebraic Cu-semigroups.

1.13. The *unitary Cuntz category*, written Cu^\sim , is the subcategory of Mon_{\leq} whose objects are ordered monoids satisfying the Cuntz axioms (O1)-(O4) and such that $0 \ll 0$ and morphisms are Mon_{\leq} -morphisms that respect suprema of increasing sequences and the compact-containment relation.

Let $S, T \in \text{Cu}^\sim$, then S_c is a submonoid of S and we equip it with the order induced by S . Let $f : S_c \rightarrow T_c$ be a Cu^\sim -morphism, denote the induced Grothendieck map by

$$\text{Gr}(f) : \text{Gr}(S_c) \rightarrow \text{Gr}(T_c).$$

2. Cu_1 AND K_*

Cantier has shown that for any unital separable C^* -algebras of stable rank one, the invariant K_* can be recovered from Cu_1 , in this section, under the real rank zero setting, we prove the converse.

Definition 2.1. [14, Definition 1.2.1] Let A be a (unital) C^* -algebra and set $K_*(A) = K_0(A) \oplus K_1(A)$. Define $K_*(A)^+ = \{([p], [u \oplus (1_k - p)])\}$, where $p \in M_k(A)$ is a projection, $u \in pM_k(A)p$ is a unitary and $\mathbf{1}_k \in M_k(A)$ is the unit. Note that we also use $[u]$ to denote the K_1 class $[u + (1_k - p)]_{K_1(A)}$.

Example 2.2. For any C^* -algebra A of stable rank one, $\text{Cu}(A)$ has weak cancellation (see [13],[20]), i.e., $x + z \ll y + z$ implies $x \leq y$ for $x, y, x \in \text{Cu}(A)$, but this is not true for $\text{Cu}_1(A)$, we present an example here.

Set

$$A = \{f \in M_2(C[0, 1]) \mid f(0) = \begin{pmatrix} \lambda & 0 \\ 0 & \mu \end{pmatrix} \text{ and } f(1) = \begin{pmatrix} \delta & 0 \\ 0 & \mu \end{pmatrix}, \lambda, \mu, \delta \in \mathbb{C}\}.$$

Then $K_1(A) = 0$ and $sr(A) = 1$.

Let $q = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \in A$ be a projection, then $K_1(I_{[q]}) = K_1(\text{Her } q) = \mathbb{Z}$.

Choose

$$([q], 0), ([q], 1) \in \text{Cu}_1(A),$$

Since $\delta_{I_{[q]}A} : K_1(I_{[q]}) \rightarrow K_1(A) = 0$, then

$$([q], 0) + ([1_A], 0) = ([q] + [1_A], \delta_{I_{[q]}A}(0) + 0) = ([q] + [1_A], 0),$$

$$([q], 1) + ([1_A], 0) = ([q] + [1_A], \delta_{I_{[q]}A}(1) + 0) = ([q] + [1_A], 0).$$

But $([q], 0) \neq ([q], 1)$.

This means $\text{Cu}_1(A)$ doesn't satisfy the cancellation of compact element in the sense of [7, Proposition 5.16]. The basic reason is that the natural map $\delta_{I_{[q]}A} = K_1(\iota) : K_1(I_{[q]}) \rightarrow K_1(A)$ is not injective, though $\iota : I_{[q]} \rightarrow A$ itself is injective and $K_0(\iota) : K_0(I_{[q]}) \rightarrow K_0(A)$ is injective.

Moreover,

$$\begin{aligned} \alpha : \text{Cu}_1(A)_c &\rightarrow K_*^+(A) \\ ([q], [u]) &\rightarrow ([q], \delta_{I_{[q]}A}[u]) \end{aligned}$$

α is just a surjective map but not an isomorphism. This means the proof of [7, Theorem 5.20] is not correct, but we point out that the conclusion is still true. We will modify it for our refinement version, Theorem 3.13.

Definition 2.3. Give $S \in \text{Cu}$, we say that S is an *algebraic* Cu -semigroup if every element in S is the supremum of an increasing sequence of compact elements.

Theorem 2.4 ([9], Corollary 5). *If A is a C^* -algebra with $rr(A) = 0$, then $\text{Cu}(A)$ is algebraic. For C^* -algebras with $sr(A) = 1$, the converse holds.*

The following lemma is well known, one can see Corollary 4.2.9 and Corollary 4.2.11 in [18].

Lemma 2.5. *Let A be a separable, unital C^* -algebra with $rr(A) = 0$ and $sr(A) = 1$, p be a projection. Then the natural map*

$$K_1(\iota) : K_1(pAp) \rightarrow K_1(A)$$

is injective.

Further, we have

Lemma 2.6. *Let A be a separable, unital C^* -algebra with $rr(A) = 0$ and $sr(A) = 1$. If $I \triangleleft A$, i.e., I is an ideal in A , the natural map*

$$K_1(\iota) : K_1(I) \rightarrow K_1(A)$$

is injective.

Proof. Suppose that a is a strictly positive elements of the ideal I , by assumption, we have $\langle a \rangle$ is the supremum of the following increasing sequence

$$(\langle p_1 \rangle, \langle p_2 \rangle, \dots),$$

where p_i is projection in A , for each $i = 1, 2, \dots$.

Denote $\text{Ideal}(p_1)$ the ideal generated by p_1 in A . Then we have

$$\text{Ideal}(p_1) \triangleleft \text{Ideal}(p_2) \triangleleft \dots \triangleleft I \triangleleft A.$$

This means that $I = C^* - \varinjlim I_{\langle p_i \rangle}$.

By Lemma 2.5 and Theorem 1.9, the natural embedding map $\iota_i : p_i A p_i \rightarrow A$ induces an injective map

$$K_1(\iota_i) : K_1(\text{Ideal}(p_i)) \rightarrow K_1(A).$$

Then the natural embedding map $\iota : I \rightarrow A$ also induces an injective map

$$K_1(\iota) : K_1(I) \rightarrow K_1(A)$$

. □

Corollary 2.7. *Let A be a separable, unital C^* -algebra with $rr(A) = 0$ and $sr(A) = 1$. If we have the following short exact sequence*

$$0 \rightarrow I \rightarrow A \rightarrow B \rightarrow 0,$$

then the exponential map $\sigma : K_0(B) \rightarrow K_1(I)$ is 0.

Remark 2.8. Brown and Pedersen [6] proved that for any real rank zero C^* -algebra, every projection in A/I can be lifted to A , hence, by six-term sequence, a more generalized version for the above conclusion is still true.

The following result is essentially contained in [7], but there is one important deficient part left for the proof, now we present our method.

Theorem 2.9. *Let A, B be separable, unital C^* -algebra with $rr(A) = 0$ and $sr(A) = 1$. Then*

$$(\text{Cu}_1(A), ([1_A, 0])) \cong (\text{Cu}_1(B), ([1_B, 0]))$$

if and only if

$$(K_*(A), K_*^+(A), [1_A]) \cong (K_*(B), K_*^+(B), [1_B]).$$

Proof. The forward direction is essentially [7, Theorem 5.20], even though there are some mistakes as we mentioned in Example 2.2, the proof is almost true.

For the backward direction, it is a combination of [7, Theorem 3.23] and Lemma 2.6. For the convenience of readers, we list a proof in detail for this part. Suppose that

$$(K_*(A), K_*^+(A), [1_A]) \cong (K_*(B), K_*^+(B), [1_B]).$$

Denote the graded isomorphism by $\Delta : K_*(A) \rightarrow K_*(B)$, then

$$\Delta(K_*^+(A)) = K_*^+(B), \quad \Delta([1_A]) = [1_B].$$

As the restriction map of Δ :

$$\Delta_{0,+} : K_0^+(A) \rightarrow K_0^+(B)$$

is a POM-isomorphism, by 1.12, we have

$$\text{Cu}(\Delta_{0,+}) : \text{Cu}(A) \rightarrow \text{Cu}(B)$$

is also a Cu-isomorphism. For convenience we denote $\text{Cu}(\Delta_{0,+})$ by Φ .

Note that Φ induces a completed lattice isomorphism between $\text{Lat}(\text{Cu}(A))$ and $\text{Lat}(\text{Cu}(B))$, which also induces a completed lattice isomorphism between $\text{Lat}(A)$ and $\text{Lat}(B)$. We just denote both the lattice isomorphisms by the same symbol Ψ . Then we have the following commutative diagram

$$\begin{array}{ccc} \text{Lat}(A) & \xrightarrow{\Psi} & \text{Lat}(B) \\ \downarrow \Gamma & & \downarrow \Gamma \\ \text{Lat}(\text{Cu}(A)) & \xrightarrow{\Psi} & \text{Lat}(\text{Cu}(B)). \end{array}$$

Each ideal $I \in \text{Lat}(A)$ is corresponding to an ideal $J \in \text{Lat}(B)$, and $\text{Cu}(I) \cong \text{Cu}(J)$, then $\text{Cu}_f(I) \cong \text{Cu}_f(J)$. Now we only need to check $K_1(I) \cong K_1(J)$.

By Lemma 2.6, both $K_1(I), K_1(J)$ are subgroup of $K_1(A), K_1(B)$, respectively. Denote the restriction map $\Delta|_{K_1(A)}$ by

$$\Delta_1 : K_1(A) \rightarrow K_1(B),$$

Δ_1 is an isomorphism.

Suppose a to be a strictly positive elements of the ideal I , and u is a unitary in \tilde{I} , then

$$(\langle a \rangle, [u]) \in \text{Cu}_1(A) \quad \text{and} \quad J = \Gamma^{-1}(I_{\Phi(\langle a \rangle)}).$$

Since the ideal I can be written as an inductive limit of algebras cutting down by projections, i.e.,

$$I = \lim_{i \rightarrow \infty} p_i A p_i.$$

There exists an i_0 large enough such that $[u] \in K_1(\text{Ideal}(p_{i_0}))$.

Note that

$$(\langle p_{i_0} \rangle, [u]) \in K_*^+(A),$$

then we have

$$\Delta(\langle p_{i_0} \rangle, [u]) \in K_*^+(B).$$

Hence,

$$\Delta_1([u]) \in K_1(\Gamma^{-1}(I_{\Phi(\langle p_{i_0} \rangle)})),$$

where, $I_{\Phi(\langle p_{i_0} \rangle)}$ is the ideal in $\text{Cu}(B)$ generated by $\Phi(\langle p_{i_0} \rangle)$.

Since Φ is a Cu-morphism and

$$\langle p_{i_0} \rangle \leq \langle a \rangle \in \text{Cu}(A),$$

we have

$$\Phi(\langle p_{i_0} \rangle) \leq \Phi(\langle a \rangle) \in \text{Cu}(B).$$

There is a nature inclusion $I_{\Phi(\langle p_{i_0} \rangle)} \subset I_{\Phi(\langle a \rangle)}$, by Lemma 2.6, then

$$K_1(\Gamma^{-1}(I_{\Phi(\langle p_{i_0} \rangle)})) \subset K_1(\Gamma^{-1}(I_{\Phi(\langle a \rangle)})) = K_1(J).$$

Then we obtain $\Delta_1([u]) \in K_1(J)$.

Since Δ_1 is an isomorphism, the restriction map $\Delta_1|_{K_1(I)}$ is automatically injective and the map $\Delta_1^{-1}|_{K_1(J)}$ is the inverse of $\Delta_1|_{K_1(I)}$. Then Δ_1 induces an isomorphism from $K_1(I)$ to $K_1(J)$.

By Theorem 1.11, we define $H : \text{Cu}_1(A) \rightarrow \text{Cu}_1(B)$ as follows:

$$\begin{aligned} H : \text{Cu}_1(A) &\rightarrow \text{Cu}_1(B) \\ (\langle a \rangle, [u]) &\rightarrow (\langle b \rangle, [v]), \end{aligned}$$

where $u \in U(\text{Ideal}(a))$, $v \in U(\text{Ideal}(b))$ and $\langle b \rangle = \Delta(\langle a \rangle)$, it is obvious that $H(\langle 1_A \rangle, 0) = (\langle 1_B \rangle, 0)$ H is additive and preserves the order defined by Cantier, then H is an isomorphism.

Now we have

$$(\text{Cu}_1(A), ([1_A, 0])) \cong (\text{Cu}_1(B), ([1_A, 0])).$$

This ends the proof. \square

Theorem 2.10 (Example 2.19 of [15]). *There exist two AD algebras A and B of real rank zero satisfying that*

$$(K_*(A), K_*^+(A), [1_A]) \cong (K_*(A), K_*^+(A), [1_A]),$$

but

$$A \not\cong B.$$

From the above theorem of Elliott-Gong-Su and Theorem 2.9, we get the following corollary directly.

Corollary 2.11. *There exist two AD algebras A and B of real rank zero satisfying that $(\text{Cu}_1(A), ([1_A, 0])) \cong (\text{Cu}_1(B), ([1_A, 0]))$, but $A \not\cong B$.*

Then Cu_1 is not a complete invariant for AD algebras (AH_d algebras).

Remark 2.12. It is also prove that in [15] the total K-theory is a complete invariant for AD algebras of real rank zero. The total K-theory works quite well in the classification of non-simple C^* -algebras (see [16, 11, 1]). Then it is necessary to give a generalized version of Cuntz semigroup for \underline{K} just like Cu_1 for K_* .

3. THE TOTAL CUNTZ SEMIGROUP

In this section, we introduce the total Cuntz semigroup, which is a refinement of the unitary Cuntz semigroup. We show that $\underline{\text{Cu}}$ is a functor from C^* -algebras of stable rank one to the total Cuntz category.

3.1. (The total K-theory) [21]. For $p \geq 2$, the mod- p K-theory groups are defined by

$$K_*(A; \mathbb{Z}_p) = K_*(A \otimes C_0(W_p)),$$

where W_p denotes the Moore space obtained by attaching the disk to the circle by a degree p -map.

For a C^* -algebra A , the total K-theory of A is defined by

$$\underline{K}(A) = \bigoplus_{p=0}^{\infty} K_*(A; \mathbb{Z}_p),$$

with $K_*(A; \mathbb{Z}_p) = K_*(A)$ for $p = 0$, $K_*(A; \mathbb{Z}_p) = 0$ for $p = 1$.

This is a $\mathbb{Z}_2 \times \mathbb{Z}^+$ graded group. (See more details in [21] and [11, Section 4].) Assume that A is a C^* -algebra of stable rank one, the following is a positive cone for total K-theory of A ([11, Definition 4.6]):

$$\underline{K}(A)_+ = \{(q, u, \oplus_{p=1}^{\infty} (s_p, s^p)) : q \in K_0^+(A), (q, u, \oplus_{p=1}^{\infty} (s_p, s^p)) \in \underline{K}_{I_q}(A)\},$$

where I_q is the ideal of A generated by q and $\underline{K}_{I_q}(A)$ denotes the image of $\underline{K}(I_q)$ in $\underline{K}(A)$.

Proposition 3.2. ([11, 4.10-4.11]) $\underline{K}(\cdot)$ and $\underline{K}(\cdot)_+$ are continuous functors for C^* -algebras of stable rank one.

Definition 3.3. Let A be a C^* -algebra of stable rank one. Define

$$\underline{\text{Cu}}(A) \triangleq \coprod_{I \in \text{Lat}(A)} \text{Cu}_f(I) \times K_1(I) \times \bigoplus_{p=1}^{\infty} K_*(I, \mathbb{Z}_p).$$

We equip $\underline{\text{Cu}}(A)$ with addition and order as follows: For any

$$(x, u, \oplus_{p=1}^{\infty} (s_p, s^p)) \in \text{Cu}_f(I_x) \times K_1(I_x) \times \bigoplus_{p=1}^{\infty} K_*(I_x, \mathbb{Z}_p)$$

and

$$(y, v, \oplus_{p=1}^{\infty}(t_p, t^p)) \in \text{Cu}_f(I_y) \times K_1(I_y) \times \bigoplus_{p=1}^{\infty} K_*(I_y, \mathbb{Z}_p),$$

then

$$\begin{aligned} & (x, u, \oplus_{p=1}^{\infty}(s_p, s^p)) + (y, v, \oplus_{p=1}^{\infty}(t_p, t^p)) \\ &= (x + y, \delta_{I_x I_{x+y}}(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) + \delta_{I_y I_{x+y}}(0, v, \oplus_{p=1}^{\infty}(t_p, t^p))) \end{aligned}$$

and

$$(x, u, \oplus_{p=1}^{\infty}(s_p, s^p)) \prec (y, v, \oplus_{p=1}^{\infty}(t_p, t^p))$$

if,

$$x \leq y \text{ and } \delta_{I_x I_y}(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) = (0, v, \oplus_{p=1}^{\infty}(t_p, t^p)),$$

where $\delta_{I_x I_y}$ is the nature map $\underline{K}(I_x) \rightarrow \underline{K}(I_y)$, we regard I_x and I_y as the corresponding ideals in A .

It is easy to see that $(\underline{\text{Cu}}(A), +, \prec)$ is a partially ordered monoid, and $\underline{\text{Cu}}(A)_+ \triangleq \{x \in \underline{\text{Cu}}(A) : 0 \prec x\} = \text{Cu}(A)$. If A is simple, we will have

$$\underline{\text{Cu}}(A) = (\text{Cu}(A) \setminus \{0\}) \times K_1(A) \times \bigoplus_{p=1}^{\infty} K_*(A, \mathbb{Z}_p) \cup \{0, 0, \bigoplus_{p=1}^{\infty}(0, 0)\}.$$

Proposition 3.4. *Let A be a C^* -algebra of stable rank one, the addition and order \prec for $\underline{\text{Cu}}(A)$ are compatible, i.e., for any*

$$(x_1, u_1, \oplus_{p=1}^{\infty}(s_{1,p}, s^{1,p})) \prec (y_1, v_1, \oplus_{p=1}^{\infty}(t_{1,p}, t^{1,p}))$$

and

$$(x_2, u_2, \oplus_{p=1}^{\infty}(s_{2,p}, s^{2,p})) \prec (y_2, v_2, \oplus_{p=1}^{\infty}(t_{2,p}, t^{2,p})),$$

we have

$$\begin{aligned} & (x_1, u_1, \oplus_{p=1}^{\infty}(s_{1,p}, s^{1,p})) + (x_2, u_2, \oplus_{p=1}^{\infty}(s_{2,p}, s^{2,p})) \\ & \prec (y_1, v_1, \oplus_{p=1}^{\infty}(t_{1,p}, t^{1,p})) + (y_2, v_2, \oplus_{p=1}^{\infty}(t_{2,p}, t^{2,p})). \end{aligned}$$

Proof. From assumption, we have

$$x_1 \leq y_1, \quad x_2 \leq y_2,$$

$$\delta_{I_{x_1} I_{y_1}}(0, u_1, \oplus_{p=1}^{\infty}(s_{1,p}, s^{1,p})) = (0, v_1, \oplus_{p=1}^{\infty}(t_{1,p}, t^{1,p}))$$

and

$$\delta_{I_{x_2} I_{y_2}}(0, u_2, \oplus_{p=1}^{\infty}(s_{2,p}, s^{2,p})) = (0, v_2, \oplus_{p=1}^{\infty}(t_{2,p}, t^{2,p})).$$

Since

$$\begin{aligned} & (x_1, u_1, \oplus_{p=1}^{\infty}(s_{1,p}, s^{1,p})) + (x_2, u_2, \oplus_{p=1}^{\infty}(s_{2,p}, s^{2,p})) \\ &= (x_1 + x_2, \delta_{I_{x_1} I_{x_1+x_2}}(0, u_1, \oplus_{p=1}^{\infty}(s_{1,p}, s^{1,p})) + \delta_{I_{x_2} I_{x_1+x_2}}(0, u_2, \oplus_{p=1}^{\infty}(s_{2,p}, s^{2,p}))) \end{aligned}$$

and

$$\begin{aligned} & (y_1, v_1, \oplus_{p=1}^{\infty}(t_{1,p}, t^{1,p})) + (y_2, v_2, \oplus_{p=1}^{\infty}(t_{2,p}, t^{2,p})) \\ &= (y_1 + y_2, \delta_{I_{y_1} I_{y_1+y_2}}(0, v_1, \oplus_{p=1}^{\infty}(t_{1,p}, t^{1,p})) + \delta_{I_{y_2} I_{y_1+y_2}}(0, v_2, \oplus_{p=1}^{\infty}(t_{2,p}, t^{2,p}))). \end{aligned}$$

Note that the following diagram is commutative naturally

$$\begin{array}{ccccc}
 \underline{K}(I_{x_1}) & \longrightarrow & \underline{K}(I_{y_1}) & & \\
 & \searrow & & \searrow & \\
 & & \underline{K}(I_{x_1+x_2}) & \longrightarrow & \underline{K}(I_{y_1+y_2}) \\
 & \nearrow & & \nearrow & \\
 \underline{K}(I_{x_2}) & \longrightarrow & \underline{K}(I_{y_2}) & &
 \end{array}$$

Then we have

$$\begin{aligned}
 & \delta(I_{x_1+x_2}I_{y_1+y_2})(\delta_{I_{x_1}I_{x_1+x_2}}(0, u_1, \oplus_{p=1}^{\infty}(s_{1,p}, s^{1,p})) + \delta_{I_{x_2}I_{x_1+x_2}}(0, u_2, \oplus_{p=1}^{\infty}(s_{2,p}, s^{2,p}))) \\
 &= \delta_{I_{y_1}I_{y_1+y_2}}(\delta_{I_{x_1}I_{y_1}}(0, v_1, \oplus_{p=1}^{\infty}(t_{1,p}, t^{1,p})) + \delta_{I_{y_2}I_{y_1+y_2}}(\delta_{I_{x_1}I_{y_1}}(0, v_2, \oplus_{p=1}^{\infty}(t_{2,p}, t^{2,p}))) \\
 &= \delta_{I_{y_1}I_{y_1+y_2}}((0, v_1, \oplus_{p=1}^{\infty}(t_{1,p}, t^{1,p}))) + \delta_{I_{y_2}I_{y_1+y_2}}((0, v_2, \oplus_{p=1}^{\infty}(t_{2,p}, t^{2,p}))) \\
 &= (0, v_1, \oplus_{p=1}^{\infty}(t_{1,p}, t^{1,p})) + (0, v_2, \oplus_{p=1}^{\infty}(t_{2,p}, t^{2,p})).
 \end{aligned}$$

Combine this with

$$x_1 + x_2 \leq y_1 + y_2 \in \text{Cu}(A).$$

The conclusion is true. \square

We now show that $(\text{Cu}(A), +, \prec)$ satisfies the Cuntz axioms in 1.4.

Lemma 3.5. *Let A be a C^* -algebra of stable rank one, $x_n \in \text{Cu}(A)$ be an increasing sequence with supremum x . Then for any*

$$u \in K_1(I_x), (s_p, s^p) \in K_*(I_x, \mathbb{Z}_p),$$

there exist $N \in \mathbb{N}$ such that for all $n > N$, there exist

$$u_n \in K_1(I_{x_n}), (s_{n,p}, s^{n,p}) \in K_*(I_{x_n}, \mathbb{Z}_p)$$

such that

$$(x_n, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p})) \prec (x, u, \oplus_{p=1}^{\infty}(s_p, s^p)).$$

Proof. Since $x = \sup_n x_n \in \text{Cu}(A)$, by Lemma 1.7, we have

$$I_x = C^* - \lim_{\longrightarrow} I_{x_n},$$

Then by continuity listed in Proposition 3.2, we have

$$\underline{K}(I_x) = \varinjlim (\underline{K}(I_{x_n}), \delta_{x_n x_m}).$$

Hence, there exists N large enough such that for all $n \geq N$,

$$u_n \in K_1(I_{x_n}), \delta_{I_{x_n}I_x}(0, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p})) = (0, u, \oplus_{p=1}^{\infty}(s_p, s^p)).$$

That is

$$(x_n, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p})) \prec (x, u, \oplus_{p=1}^{\infty}(s_p, s^p)).$$

\square

Corollary 3.6. *Let A be a C^* -algebra of stable rank one. Then any increasing sequence $(x_n, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p}))$ has a supremum $(x, u, \oplus_{p=1}^{\infty}(s_p, s^p))$ in $\underline{\text{Cu}}(A)$. In particular, $x = \sup_n x_n$, and for any $n \in \mathbb{N}$,*

$$\delta_{I_{x_n} I_x}(0, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p})) = (0, u, \oplus_{p=1}^{\infty}(s_p, s^p)).$$

Proof. Since $\text{Cu}(A)$ satisfies (O1), there exists $x \in \text{Cu}(A)$ such that $x = \sup_n x_n$, and

$$I_x = C^* - \lim_{\rightarrow} I_{x_n}.$$

Then

$$\underline{\text{K}}(I_x) = \lim_{\rightarrow} (\underline{\text{K}}(I_{x_n}), \delta_{x_n x_m}).$$

Note that

$$\delta_{x_n x_{n+1}}(0, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p})) = (0, u_{n+1}, \oplus_{p=1}^{\infty}(s_{n+1,p}, s^{n+1,p})).$$

We may choose $(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) \in \underline{\text{K}}(I_x)$ be the limit

$$(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) := \lim_{\rightarrow} \delta_{x_n x}(0, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p}))$$

Then for any $n \in \mathbb{N}$,

$$(x_n, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p})) \prec (x, u, \oplus_{p=1}^{\infty}(s_p, s^p)).$$

Now we check that $(x, u, \oplus_{p=1}^{\infty}(s_p, s^p))$ is in fact the supremum of the sequence, choose any $(y, v, \oplus_{p=1}^{\infty}(t_p, t^p)) \succ (x_n, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p}))$, then $x = \sup_n x_n \leq y$ and

$$\delta_{x_n y}(0, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p})) = (0, v, \oplus_{p=1}^{\infty}(t_p, t^p)).$$

It is obviously that

$$\delta_{xy}(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) = (0, v, \oplus_{p=1}^{\infty}(t_p, t^p)).$$

Hence,

$$(x, u, \oplus_{p=1}^{\infty}(s_p, s^p)) \prec (y, v, \oplus_{p=1}^{\infty}(t_p, t^p)).$$

□

Theorem 3.7. *Let A be a C^* -algebra of stable rank one and let*

$$(x, u, \oplus_{p=1}^{\infty}(s_p, s^p)), (y, v, \oplus_{p=1}^{\infty}(t_p, t^p)) \in \underline{\text{Cu}}(A).$$

Then

$$(x, u, \oplus_{p=1}^{\infty}(s_p, s^p)) \ll (y, v, \oplus_{p=1}^{\infty}(t_p, t^p)).$$

if and only if $x \ll y$ in $\text{Cu}(A)$ and

$$\delta_{I_x I_y}(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) = (0, v, \oplus_{p=1}^{\infty}(t_p, t^p)).$$

Proof. Suppose that $(x, u, \oplus_{p=1}^{\infty}(s_p, s^p)) \ll (y, v, \oplus_{p=1}^{\infty}(t_p, t^p))$, then

$$\delta_{I_x I_y}(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) = (0, v, \oplus_{p=1}^{\infty}(t_p, t^p)),$$

we only need to prove $x \ll y$.

Let $\{y_n\}$ be an increasing sequence in $\text{Cu}(A)$ with supremum y . By Lemma 1.7, we have

$$\underline{\mathbf{K}}(I_y) = \lim_{\rightarrow} (\underline{\mathbf{K}}(I_{y_n}), \delta_{I_{y_n} I_{y_m}}).$$

There exists $n_0 \in \mathbb{N}$ such that for any $n \geq n_0$,

$$(0, v_n, \oplus_{p=1}^{\infty} (t_{n,p}, t^{n,p}) \leq (0, v, \oplus_{p=1}^{\infty} (t_p, t^p)),$$

where $(0, u_n, \oplus_{p=1}^{\infty} (t_{n,p}, t^{n,p}) \in \underline{\mathbf{K}}(I_{y_n})$.

Then $(y_n, v_n, \oplus_{p=1}^{\infty} (t_{n,p}, t^{n,p}))$ is an increasing sequence in $\underline{\mathbf{Cu}}(A)$ with supremum $(y, v, \oplus_{p=1}^{\infty} (t_p, t^p))$. From assumption, there exists $N \in \mathbb{N}$ such that for any $n \geq N$,

$$(x, u, \oplus_{p=1}^{\infty} (s_p, s^p)) \prec (y_n, v_n, \oplus_{p=1}^{\infty} (t_{n,p}, t^{n,p})),$$

Then we have $x \precsim y_n$, that is $x \ll y$.

Conversely, for any increasing sequence $y_n \in \text{Cu}(A)$ with supremum y . Then there exists N large enough such that for any $n > N$,

$$x \leq y_n \leq y, \quad \text{i.e.,} \quad I_x \subset I_{y_n} \subset I_y$$

and

$$\delta_{I_{y_n} I_y} (0, v_n, \oplus_{p=1}^{\infty} (t_{n,p}, t^{n,p})) = (0, v, \oplus_{p=1}^{\infty} (t_p, t^p)).$$

where $(0, v_n, \oplus_{p=1}^{\infty} (t_{n,p}, t^{n,p}) \in \underline{\mathbf{K}}(I_{y_n})$.

Now we have

$$\delta_{I_x I_{y_n}} (0, u, \oplus_{p=1}^{\infty} (s_p, s^p)) = (0, v_n, \oplus_{p=1}^{\infty} (t_{n,p}, t^{n,p})),$$

this means that

$$(x, u, \oplus_{p=1}^{\infty} (s_p, s^p)) \prec (y_n, v_n, \oplus_{p=1}^{\infty} (t_{n,p}, t^{n,p})).$$

Hence,

$$(x, u, \oplus_{p=1}^{\infty} (s_p, s^p)) \ll (y, v, \oplus_{p=1}^{\infty} (t_p, t^p)).$$

□

Corollary 3.8. *Let A be a C^* -algebra of stable rank one, then $(x, u, \oplus_{p=1}^{\infty} (s_p, s^p))$ is compact in $\underline{\mathbf{Cu}}(A)$ if and only if x is compact in $\text{Cu}(A)$.*

Corollary 3.9. *Let A be a C^* -algebra of stable rank one, then A is of real rank zero if and only if $\underline{\mathbf{Cu}}(A)$ is algebraic.*

Then we obtain the following theorem.

Theorem 3.10. *Let A be a C^* -algebra of stable rank one, then $(\underline{\mathbf{Cu}}(A), +, \prec)$ satisfies axioms (O1), (O2), (O3), and (O4).*

Proof. (O1): It is exactly Corollary 3.6.

(O2): $(x, u, \oplus_{p=1}^{\infty} (s_p, s^p)) \in \underline{\mathbf{Cu}}(A)$, from the (O2) of $\text{Cu}(A)$, we have a \ll -increasing sequence $\{x_n\} \in \text{Cu}(A)$. Then by Lemma 3.5 and Theorem 3.7, we achieve the proof for (O2) of $(\underline{\mathbf{Cu}}(A), \prec)$.

(O3): Let

$$(x_1, u_1, \oplus_{p=1}^{\infty} (s_{1,p}, s^{1,p})) \ll (y_1, v_1, \oplus_{p=1}^{\infty} (t_{1,p}, t^{1,p}))$$

and

$$(x_2, u_2, \oplus_{p=1}^{\infty}(s_{2,p}, s^{2,p})) \ll (y_2, v_2, \oplus_{p=1}^{\infty}(t_{2,p}, t^{2,p})).$$

We have

$$\begin{aligned} & (x_1, u_1, \oplus_{p=1}^{\infty}(s_{1,p}, s^{1,p})) + (x_2, u_2, \oplus_{p=1}^{\infty}(s_{2,p}, s^{2,p})) \\ & \prec (y_1, v_1, \oplus_{p=1}^{\infty}(t_{1,p}, t^{1,p})) + (y_2, v_2, \oplus_{p=1}^{\infty}(t_{2,p}, t^{2,p})). \end{aligned}$$

From (O3) of $\text{Cu}(A)$, we also have $x_1 + x_2 \ll y_1 + y_2$, therefore, by Theorem 3.7, we get (O3) for $(\underline{\text{Cu}}(A), \prec)$.

(O4): Let $\{(x_n, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p}))\}$ and $\{(y_n, v_n, \oplus_{p=1}^{\infty}(t_{n,p}, t^{n,p}))\}$ be two increasing sequences in $\underline{\text{Cu}}(A)$. Let

$$(x, u, \oplus_{p=1}^{\infty}(s_p, s^p)) = \sup_n \{(x_n, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p}))\}$$

and

$$(y, v, \oplus_{p=1}^{\infty}(t_p, t^p)) = \sup_{n \in \mathbb{N}} \{(y_n, v_n, \oplus_{p=1}^{\infty}(t_{n,p}, t^{n,p}))\}.$$

From (O4) of $\text{Cu}(A)$, we have $\sup_{n \in \mathbb{N}} \{x_n + y_n\} = x + y$. By the compatibility of order \prec and addition, we know

$$\{(x_n, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p})) + (y_n, v_n, \oplus_{p=1}^{\infty}(t_{n,p}, t^{n,p}))\}$$

is also an increasing sequence. Note that

$$\delta_{I_{x_n} I_x}(0, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p})) = (0, u, \oplus_{p=1}^{\infty}(s_p, s^p))$$

and

$$\delta_{I_{y_n} I_y}(0, v_n, \oplus_{p=1}^{\infty}(t_{n,p}, t^{n,p})) = (0, v, \oplus_{p=1}^{\infty}(t_p, t^p)).$$

This implies

$$\begin{aligned} & \delta_{I_{x_n+y_n} I_{x+y}}(\delta_{I_{x_n} I_{x_n+y_n}}(0, u_n, \oplus_{p=1}^{\infty}(s_{n,p}, s^{n,p})) + \delta_{I_{y_n} I_{x_n+y_n}}(0, v_n, \oplus_{p=1}^{\infty}(t_{n,p}, t^{n,p}))) \\ & = \delta_{I_x I_{x+y}}(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) + \delta_{I_y I_{x+y}}(0, v, \oplus_{p=1}^{\infty}(t_p, t^p)). \end{aligned}$$

Using Corollary 3.6, we get (O4) for $(\underline{\text{Cu}}(A), +, \prec)$. \square

Definition 3.11. Define the total Cuntz category as follows:

$$\underline{\text{Cu}}^{\sim} = \{S \in \text{Cu}^{\sim} \mid \text{Gr}(S_c) \text{ is a } \Lambda - \text{module}\},$$

where Λ denotes the category of Bockstein operations (see [11, 4.1]).

Let $X, Y \in \underline{\text{Cu}}^{\sim}$, the morphism ϕ from X to Y satisfies the following two conditions:

(1) ϕ is a $\underline{\text{Cu}}^{\sim}$ -morphisms, i.e., ϕ is Mon_{\leq} -morphism and preserves suprema of increasing sequences and the compact-containment relation.

(2) The induced Grothendieck map $\text{Gr}(\phi_c) : X_c \rightarrow Y_c$ is Λ -linear, i.e., $\text{Gr}(\phi_c)$ preserve the action of the category Λ .

We call these morphisms $\underline{\text{Cu}}$ -morphisms.

Remark 3.12. Let A and B be C^* -algebras. I_x and J_y are ideals of A and B generated by $x \in \text{Cu}(A)$ and $y \in \text{Cu}(B)$, respectively. $\psi : A \rightarrow B$ is a $*$ -homomorphism satisfying $\psi(I_x) \subset J_y$.

Then ψ induces the two morphisms:

$$\underline{K}(\psi) : \underline{K}(A) \rightarrow \underline{K}(B) \quad \text{and} \quad \underline{K}(\psi_{I_x I_y}) : \underline{K}(I_x) \rightarrow \underline{K}(J_y).$$

In general, $\underline{K}(\psi_{I_x I_y})$ is not the restriction of $\underline{K}(\psi)$, as $\underline{K}(I_x)$ is not a subgroup of $\underline{K}(A)$.

It is easy check that

$$\underline{\text{Cu}}(\psi)(x, u, \oplus_{p=1}^{\infty}(s_p, s^p)) \prec (y, v, \oplus_{p=1}^{\infty}(t_p, t^p))$$

if and only if $\text{Cu}(\psi)(x) \prec y \in \text{Cu}(B)$ and

$$\underline{K}(\psi_{I_x I_y})(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) = (0, v, \oplus_{p=1}^{\infty}(t_p, t^p)).$$

In order to simplify the notations, we will always write

$$\underline{K}(\psi)(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) = (0, v, \oplus_{p=1}^{\infty}(t_p, t^p)) \in \underline{K}(I_y)$$

instead of

$$\underline{K}(\psi_{I_x I_y})(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) = (0, v, \oplus_{p=1}^{\infty}(t_p, t^p)),$$

where $(x, u, \oplus_{p=1}^{\infty}(s_p, s^p)) \in \underline{\text{Cu}}(A)$ and $(y, v, \oplus_{p=1}^{\infty}(t_p, t^p)) \in \underline{\text{Cu}}(B)$.

Theorem 3.13. *The assignment*

$$\begin{aligned} \underline{\text{Cu}} : C_{sr1}^* &\rightarrow \underline{\text{Cu}}^{\sim} \\ A &\rightarrow \underline{\text{Cu}}(A) \\ \phi &\rightarrow \underline{\text{Cu}}(\phi) \end{aligned}$$

from C^* -algebras of stable rank one to the category $\underline{\text{Cu}}^{\sim}$ is a functor.

Proof. The key point is to prove $\text{Gr}(\underline{\text{Cu}}(A)_c) = \underline{K}(A)$, where $\text{Gr}(\cdot)$ is the Grothendick group procedure.

Since A has stable rank one, every compact elements in $\text{Cu}(A)$ can be arised by projections [5, Theorem 5.8], then by Corollary 3.8, every compact elements in $\text{Cu}(A)$ has the form $([e], u, \oplus_{p=1}^{\infty}(s_p, s^p))$, where e is a projection in A .

Define

$$\begin{aligned} \alpha : \underline{\text{Cu}}(A)_c &\rightarrow \underline{K}(A)_+, \\ ([e], u, \oplus_{p=1}^{\infty}(s_p, s^p)) &\rightarrow ([e], \delta_{I_e A}(u), \oplus_{p=1}^{\infty}(\delta_{I_e A}(s_p), \delta_{I_e A}(s^p))). \end{aligned}$$

From the definition we list in 3.1, α is a surjective ordered-monoid-morphism. (α is just surjective, not necessarily injective. Example 2.2 is such an algebra.)

Suppose that

$$\alpha([e], u, \oplus_{p=1}^{\infty}(s_p, s^p)) = \alpha([q], v, \oplus_{p=1}^{\infty}(t_p, t^p)).$$

Then

$$([e], \delta_{I_e A}(u), \oplus_{p=1}^{\infty}(\delta_{I_e A}(s_p), \delta_{I_e A}(s^p))) = ([q], \delta_{I_q A}(v), \oplus_{p=1}^{\infty}(\delta_{I_q A}(t_p), \delta_{I_q A}(t^p))).$$

For $([1_A], 0, \oplus_{p=1}^{\infty}(0, 0)) \in \underline{K}(A)_+$, we have

$$\begin{aligned} & ([1_A] + [e], \delta_{I_e A}(u), \oplus_{p=1}^{\infty}(\delta_{I_e A}(s_p), \delta_{I_e A}(s^p))) \\ &= ([1_A] + [q], \delta_{I_q A}(v), \oplus_{p=1}^{\infty}(\delta_{I_q A}(t_p), \delta_{I_q A}(t^p))). \end{aligned}$$

Now we obtain

$$\begin{aligned} & ([1_A], 0, \oplus_{p=1}^{\infty}(0, 0)) + ([e], u, \oplus_{p=1}^{\infty}(s_p, s^p)) \\ &= ([1_A], 0, \oplus_{p=1}^{\infty}(0, 0)) + ([q], v, \oplus_{p=1}^{\infty}(t_p, t^p)). \end{aligned}$$

This means that for any $x, y \in \underline{Cu}(A)_c$, if $\alpha(x) = \alpha(y)$, under the Grothendieck construction, the difference between x and y vanishes.

Then we have

$$\text{Gr}(\underline{Cu}(A)_c) = \text{Gr}(\alpha(\underline{Cu}(A)_c)) = \text{Gr}(\underline{K}(A)_+) = \underline{K}(A).$$

Since $\underline{K}(A)$ is a Λ -modulo, then $\text{Gr}(\underline{Cu}(A)_c)$ is also a Λ -modulo, now we have $\underline{Cu}(A) \in \underline{Cu}^{\sim}$.

Suppose that $\phi : A \rightarrow B$ is a $*$ -homomorphism, it is obviously that the induced map

$$\underline{Cu}(\phi) : \underline{Cu}(A) \rightarrow \underline{Cu}(B).$$

is a \underline{Cu}^{\sim} -morphism.

Note that

$$\text{Gr}(\underline{Cu}(A)_c) = \underline{K}(A), \quad \text{Gr}(\underline{Cu}(B)_c) = \underline{K}(B).$$

Then the Grothendieck map

$$\text{Gr}(\phi_c) : \underline{K}(A) \rightarrow \underline{K}(B)$$

is exactly the map $\underline{K}(\psi)$, and hence, a Λ -linear map.

Then $\underline{Cu}(\phi)$ is a \underline{Cu}^{\sim} -morphism.

□

4. CONTINUITY OF THE FUNCTOR \underline{Cu}

It is proved in section 3.8 of [7] that Cu_1 is a continuous functor, now we show that \underline{Cu} is also a continuous functor from C^* -algebras of stable rank one to the category \underline{Cu}^{\sim} .

Definition 4.1. Let $S_1 \xrightarrow{\beta_{1,2}} S_2 \xrightarrow{\beta_{2,3}} \dots$ be a sequence in the category \underline{Cu}^{\sim} . We say (s_1, s_2, \dots) is an eventually-increasing sequence if, $s_1 \in S_1$, $s_2 \in S_2, \dots$ and there exists $k \in \mathbb{N}$ such that for any $i, j > k$, $\beta_{i,j}(s_i) \prec s_j$. Denote \mathcal{S} the collection of all the eventually-increasing sequences.

Let $(s_i), (t_i) \in \mathcal{S}$, define the addition operation

$$(s_i) + (t_i) = (s_i + t_i).$$

and the pre-order relation

$$(s_i) \leq (t_i)$$

if there exists $k \in \mathbb{N}$ such that for any $i \geq k$ and $x \in S_i$ with $x \ll s_i$, there is an $m \geq i$ such that $\beta_{i,j}(x) \ll t_j$ (in S_j) for any $j \geq m$. We say $(s_i) \sim (t_i)$,

if $(s_i) \leq (t_i)$ and $(t_i) \leq (s_i)$. We denote the equivalent class of (s_i) in \mathcal{S} by $[(s_i)]$.

Remark 4.2. Every increasing sequence defined in 1.6 is an eventually-increasing sequence. Every eventually-increasing sequence is equivalent to a sequence of the form

$$(0, 0, \dots, 0, s_k, s_{k+1}, \dots).$$

We may denote this sequence by $(s_i)_{i \geq k}$.

The eventually-increasing sequences are defined for inductive systems in the category Cu^\sim , just like the increasing sequences defined for the category Cu [9]. But for particular cases, we may even have $\beta_{k-1,k}(S_{k-1}) \cap s_k^\prec = \emptyset$, where $s_k^\prec \triangleq \{s \in S_k : s \prec s_k\}$. Then the increasing sequence is not applicable, which means the notion of eventually-increasing is necessary.

To build an example, one can pick $S_1 = \mathbb{N} \cup \{\infty\}$ and $S_i = ((\mathbb{N} \setminus \{0\}) \cup \{\infty\}) \times \mathbb{Z} \cup \{(0, 0)\}$, for all $i \geq 2$. (For $(m_1, m_2), (n_1, n_2) \in S_i$, we say $(m_1, m_2) \prec (n_1, n_2)$ if $m_1 \leq n_1$ and $m_2 = n_2$.) Set $\beta_{1,2}(n) = (n, 0)$ and $\beta_{i,j} : S_i \rightarrow S_j$ is identity map for $j > i \geq 2$. Then $(S_i, \beta_{i,j})$ forms an inductive system in the category Cu^\sim .

Let $s_2 = (1, 1)$. Now we have $\beta_{1,2}(S_1) \cap s_2^\prec = \emptyset$. This is the reason we use the “eventually-increasing sequence” to replace the “increasing sequence” for the inductive system in Cu^\sim -category. (One can also pick $S_1 = \text{Cu}_1(\mathbb{C})$ and $S_i = \text{Cu}_1(C(S^1))$, for all $i \geq 2$, which is also such an example.)

It is easy to verify that $(s_i + t_i)$ belongs to \mathcal{S} and the pre-order relation is compatible with addition. With a similar proof of the first part of [9, Theorem 2], we have the following:

Lemma 4.3. *Let $S_1 \rightarrow S_2 \rightarrow \dots$ be a sequence in the category Cu^\sim . Then*

$$\text{Cu}^\sim\text{-}\lim_{\rightarrow} S_i \cong \mathcal{S} / \sim.$$

Remark 4.4. Let $(S_i, \beta_{i,j})$ be an inductive system in Cu^\sim -category, and $S = \mathcal{S} / \sim$. Suppose we have an object $T \in \text{Cu}^\sim$ -category satisfying that for any $j \geq i \in \mathbb{N}$, the following diagram is commutative.

$$\begin{array}{ccc} S_i & \xrightarrow{\psi_i} & T \\ & \searrow \beta_{i,j} & \uparrow \psi_j \\ & & S_j \end{array}$$

Then by universal property of inductive limit, there is a unique Cu^\sim -map ω such that the following is also commutative for any $i \in \mathbb{N}$.

$$\begin{array}{ccc} S_i & \xrightarrow{\psi_i} & T \\ & \searrow \beta_{i,\infty} & \uparrow \exists! \omega \\ & & S \end{array}$$

Given $s = (s_i)_{i \geq k_0} \in \mathcal{S}$, for each $i \geq k_0$, we have

$$\phi_{i,\infty}(s_i) = [(\phi_{i,j}(s_i))_{j \geq i}],$$

and hence from the commutativity,

$$\psi_i(s_i) = \omega([(\phi_{i,j}(s_i))_{j \geq i}]).$$

Then as

$$[s] = \sup_{i \geq k_0} [\phi_{i,\infty}(s_i)]$$

and ω preserves the supreme, we get

$$\omega([s]) = \omega(\sup_{i \geq k_0} [\phi_{i,\infty}(s_i)]) = \sup_{i \geq k_0} \omega([\phi_{i,\infty}(s_i)]) = \sup_{i \geq k_0} \psi_i(s_i).$$

Theorem 4.5. *Let $A = \varinjlim (A_i, \phi_{i,j})$ be an inductive system in the category C_{sr1}^* . Then*

$$\underline{\text{Cu}} - \varinjlim \underline{\text{Cu}}(A_i) \cong \underline{\text{Cu}}(A).$$

Proof. Since $\underline{\text{Cu}}(A_i)$ are objects in the category Cu^\sim . Denote

$$S = \text{Cu}^\sim - \varinjlim \underline{\text{Cu}}(A_i).$$

Then S is the inductive limit in the category Cu^\sim of the sequence

$$\underline{\text{Cu}}(A_1) \rightarrow \underline{\text{Cu}}(A_2) \rightarrow \cdots.$$

For $\underline{\text{Cu}}(A) \in \text{Cu}^\sim$ and the compatible sequence of maps

$$\underline{\text{Cu}}(\phi_{i,\infty}) : \underline{\text{Cu}}(A_i) \rightarrow \underline{\text{Cu}}(A).$$

There exists a unique compatible Cu^\sim -morphism $\omega : S \rightarrow \underline{\text{Cu}}(A)$ such that the following diagram is commutative:

$$\begin{array}{ccc} \underline{\text{Cu}}(A_i) & \xrightarrow{\underline{\text{Cu}}(\phi_{i,\infty})} & \underline{\text{Cu}}(A) \\ & \searrow & \uparrow \exists! \omega \\ & & S \end{array}$$

Now we need only to prove that ω is a Cu^\sim -isomorphism.

First, we show that ω is surjective. For any $(x, u, \oplus_{p=1}^\infty (s_p, s^p)) \in \underline{\text{Cu}}(A)$, by 1.6, there exists an increasing sequence (x_1, x_2, \dots) with $x_1 \in \text{Cu}(A_1)$, $x_2 \in \text{Cu}(A_2), \dots$ such that

$$x = [(x_1, x_2, \dots)],$$

Let I_{x_i} be the ideal of A_i generated by $x_i \in \text{Cu}(A_i)$, I_x be the ideal of A generated by $x \in \text{Cu}(A)$. Then

$$I_{x_1} \rightarrow I_{x_2} \rightarrow \cdots \rightarrow I_x \triangleleft A,$$

with

$$I_x = C^* - \lim I_{x_i}.$$

Note that $(0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) \in \underline{K}(I_x)$, by the continuity of \underline{K} , there exists $k \in \mathbb{N}$ and $(0, u_k, \oplus_{p=1}^{\infty}(s_{k,p}, s^{k,p})) \in \underline{K}(I_{x_k})$ satisfying

$$\underline{K}(\phi_{k,\infty})(0, u_k, \oplus_{p=1}^{\infty}(s_{i,p}, s^{i,p})) = (0, u, \oplus_{p=1}^{\infty}(s_p, s^p)) \in \underline{K}(I_x).$$

and for any $i > k$,

$$(0, u_i, \oplus_{p=1}^{\infty}(s_{i,p}, s^{i,p})) = \underline{K}(\phi_{k,i})(0, u_k, \oplus_{p=1}^{\infty}(s_{k,p}, s^{k,p})) \in \underline{K}(I_{x_i}).$$

Then we obtain an eventually-increasing sequence $(x_j, u_j, \oplus_{p=1}^{\infty}(s_{j,p}, s^{j,p}))$ (if $j < k$, one can choose any $(x_j, u_j, \oplus_{p=1}^{\infty}(s_{j,p}, s^{j,p})) \in \underline{\text{Cu}}^{\sim}(A_j)$). By Lemma 4.3, there exists an $s \in S$ such that

$$s = [(x_j, u_j, \oplus_{p=1}^{\infty}(s_{j,p}, s^{j,p}))].$$

Then by the commutativity of

$$\begin{array}{ccc} \underline{\text{Cu}}(A_i) & \xrightarrow{\underline{\text{Cu}}(\phi_{i,\infty})} & \underline{\text{Cu}}(A) \\ & \searrow \Phi_i & \uparrow \exists! \omega \\ & & S, \end{array}$$

we get

$$\omega([(x_j, u_j, \oplus_{p=1}^{\infty}(s_{j,p}, s^{j,p}))]) = (x, u, \oplus_{p=1}^{\infty}(s_p, s^p)).$$

Second, we show ω is injective. Let $f = (x_i, u_i, \oplus_{p=1}^{\infty}(s_{i,p}, s^{i,p}))_{i \geq k_1}$ and $g = (y_j, v_j, \oplus_{p=1}^{\infty}(t_{j,p}, t^{j,p}))_{j \geq k_2}$ be two eventually-increasing sequences of inductive system $(\underline{\text{Cu}}(A_i), \underline{\text{Cu}}(\phi_{ij}))$ in the category Cu^{\sim} . Suppose $\omega(f) = \omega(g) \in \underline{\text{Cu}}(A)$, we need only to prove $f \sim g$.

From Remark 4.4, we have

$$\sup_{i \geq k_1} \underline{\text{Cu}}(\phi_{i,\infty})(x_i, u_i, \oplus_{p=1}^{\infty}(s_{i,p}, s^{i,p})) = \sup_{j \geq k_2} \underline{\text{Cu}}(\phi_{j,\infty})(y_j, v_j, \oplus_{p=1}^{\infty}(t_{j,p}, t^{j,p})),$$

then

$$[(x_i)] = \sup \text{Cu}(\phi_{i,\infty})(x_i) = \sup \text{Cu}(\phi_{j,\infty})(y_j) = [(y_j)], \text{ i.e., } (x_i) \sim (y_i).$$

Denote $z = [(x_i)] = [(y_i)] \in \text{Cu}(A)$, then

$$I_{x_1} \rightarrow I_{x_2} \rightarrow \cdots \rightarrow I_z \triangleleft A,$$

$$I_{y_1} \rightarrow I_{y_2} \rightarrow \cdots \rightarrow I_z \triangleleft A$$

and

$$I_z = \text{C}^* - \lim I_{x_i} = \text{C}^* - \lim I_{y_i}.$$

Now we have

$$\begin{array}{ccccccc} \underline{K}(I_{x_1}) & \xrightarrow{\varphi_{1,2}} & \underline{K}(I_{x_2}) & \longrightarrow & \cdots & \longrightarrow & \underline{K}(I_z) \\ & & & & & & \downarrow = \\ \underline{K}(I_{y_1}) & \xrightarrow{\psi_{1,2}} & \underline{K}(I_{y_2}) & \longrightarrow & \cdots & \longrightarrow & \underline{K}(I_z), \end{array}$$

where the maps φ_{ij} and ψ_{ij} are in fact $\underline{K}((\phi_{i,j})_{I_{x_i}I_{x_j}})$ and $\underline{K}((\phi_{i,j})_{I_{y_i}I_{y_j}})$, respectively.

As $(x_i) \prec (y_i)$, there exists $k \in \mathbb{N}$ such that for any $i \geq k$ and $c \in \text{Cu}(A_i)$ with $c \ll x_i$, there exists m_c such that $\text{Cu}(\phi_{i,j})(c) \ll y_j$ (in $\text{Cu}(A_j)$), for all $j \geq m_c$. Choose $(c, w, \oplus_{p=1}^{\infty}(r_p^i, r_p^i)) \in \underline{\text{Cu}}(A_i)$ with

$$(c, w, \oplus_{p=1}^{\infty}(r_{i,p}, r_{i,p}^{i,p})) \ll (x_i, u_i, \oplus_{p=1}^{\infty}(s_{i,p}, s_{i,p}^{i,p})),$$

then we have

$$\delta_{I_{c}I_{x_i}}(0, w, \oplus_{p=1}^{\infty}(r_{i,p}, r_{i,p}^{i,p})) = (0, u_i, \oplus_{p=1}^{\infty}(s_{i,p}, s_{i,p}^{i,p})), \quad \forall i \geq k.$$

Now we have the natural commutative diagram:

$$\begin{array}{ccccc} I_c & \longrightarrow & I_{x_i} & \longrightarrow & I_z. \\ \downarrow & & & \nearrow & \\ I_{\text{Cu}(\phi_{ij})(c)} & \longrightarrow & I_{y_j} & & \end{array}$$

Then the following diagram commutes at the level of \underline{K} :

$$\begin{array}{ccccc} \underline{K}(I_c) & \xrightarrow{\delta_{I_c I_{x_i}}} & \underline{K}(I_{x_i}) & \longrightarrow & \underline{K}(I_z). \\ \downarrow & & & \nearrow & \\ \underline{K}(I_{\text{Cu}(\phi_{ij})(c)}) & \xrightarrow{\delta_{I_{\text{Cu}(\phi_{ij})(c)} I_{y_j}}} & \underline{K}(I_{y_j}) & & \end{array}$$

Note that

$$\underline{K}(\phi_{i,\infty})(0, u_i, \oplus_{p=1}^{\infty}(s_{i,p}, s_{i,p}^{i,p})) = \underline{K}(\phi_{j,\infty})(0, v_j, \oplus_{p=1}^{\infty}(t_{j,p}, t_{j,p}^{j,p})) \in \underline{K}(I_z),$$

then there exists $m \geq m_c$ such that

$$\delta_{I_{\text{Cu}(\phi_{i,j})(c)} I_{y_j}} \circ \underline{K}(\phi_{ij})(0, w, \oplus_{p=1}^{\infty}(r_{i,p}, r_{i,p}^{i,p})) = (0, v_j, \oplus_{p=1}^{\infty}(t_{j,p}, t_{j,p}^{j,p})) \in \underline{K}(I_{y_j})$$

holds for any $j \geq m$.

Recall that $c \ll y_j$, by Theorem 3.7, we have

$$(c, w, \oplus_{p=1}^{\infty}(r_{i,p}, r_{i,p}^{i,p})) \ll (y_j, v_j, \oplus_{p=1}^{\infty}(t_{j,p}, t_{j,p}^{j,p}))$$

holds for any $j \geq m$. Then we get $f \leq g$.

Similarly, $(y_i) \leq (x_i)$ will implies $g \leq f$. Now we have ω is injective.

Now we have $\underline{\text{Cu}}(A) \cong S$, by Theorem 3.13, we have S belongs to the category $\underline{\text{Cu}}^\sim$. This completes the proof. \square

5. RECOVER THE TOTAL \underline{K} -THEROY

In this section, we show that $\underline{\text{Cu}}$ contains more information than \underline{K} , hence, $\underline{\text{Cu}}$ is a complete invariant for certain real rank zero algebras.

5.1. Let (S, u) is a $\underline{\text{Cu}}^\sim$ -semigroup with compact order unit. Now a $\underline{\text{Cu}}^\sim$ -morphism between (S, u) and (T, v) is a $\underline{\text{Cu}}^\sim$ -morphism $f : S \rightarrow T$ such that $f(u) \leq v$. Denote the the category of $\underline{\text{Cu}}^\sim$ -semigroups with compact order-unit by $\underline{\text{Cu}}_u^\sim$.

After adding a unit, we transform Theorem 3.13 into the following:

Lemma 5.2. *The assignment*

$$\begin{aligned} \underline{\text{Cu}}_u : C_{sr1}^* &\rightarrow \underline{\text{Cu}}_u^\sim \\ A &\rightarrow (\underline{\text{Cu}}(A), ([1_A], 0, \oplus_{p=1}^\infty(0, 0))) \\ \phi &\rightarrow \underline{\text{Cu}}(\phi) \end{aligned}$$

from the category of unital C^* -algebra of stable rank one to the category $\underline{\text{Cu}}^\sim$ is a covariant functor.

And from the proof of Theorem 3.13, for any C^* -algebras A with stable rank one, we have

$$(\underline{\text{K}}(A), \underline{\text{K}}(A)_+, [1_A]) \cong (\text{Gr}(\underline{\text{Cu}}(A)_c), \alpha(\underline{\text{Cu}}(A)_c), ([1_A], 0, \oplus_{p=1}^\infty(0, 0))).$$

This means we do recover $\underline{\text{K}}$ from $\underline{\text{Cu}}$ as we anticipated.

Proposition 5.3. *Let*

$$\begin{aligned} \underline{H} : \underline{\text{Cu}}_u^\sim &\rightarrow \Lambda - \text{module} \\ (S, u) &\rightarrow (\text{Gr}(S_c), \alpha(S_c), [u]) \\ \phi &\rightarrow \text{Gr}(\phi_c), \end{aligned}$$

where $\alpha(S_c)$ is the image of S_c in $\text{Gr}(S_c)$. Then the functor \underline{H} yields a nature isomorphism $\underline{H} \circ \underline{\text{Cu}}_u \cong \underline{\text{K}}$, that is, for any C^* -algebras A, B with stable rank one, if

$$(\underline{\text{Cu}}(A), [1_A], 0, \oplus_{p=1}^\infty(0, 0)) \cong (\underline{\text{Cu}}(B), [1_B], 0, \oplus_{p=1}^\infty(0, 0)),$$

then

$$(\underline{\text{K}}(A), \underline{\text{K}}(A)_+, [1_A]) \cong (\underline{\text{K}}(B), \underline{\text{K}}(B)_+, [1_B]).$$

To achieve a total version of Theorem 2.9, we list the following.

Definition 5.4. We say that an ideal I in a C^* -algebra A is *K-pure* if both sequences

$$0 \rightarrow K_\delta(I) \rightarrow K_\delta(A) \rightarrow K_\delta(A/I) \rightarrow 0, \quad \delta = 0, 1$$

are pure exact. We say a C^* -algebra A is *K-pure*, if all ideals in A are *K-pure*.

Lemma 5.5. *If an ideal I in a C^* -algebra A is K-pure, then for any $p \geq 2$, the sequences*

$$0 \rightarrow K_*(I; \mathbb{Z}_p) \rightarrow K_*(A; \mathbb{Z}_p) \rightarrow K_*(A/I; \mathbb{Z}_p) \rightarrow 0$$

are exact.

Proof. For any $p \geq 2$, we have the following well known commutative diagram induced by the natural embedding $\iota : I \rightarrow A$.

$$\begin{array}{ccccccc}
 K_\delta(I) & \xrightarrow{\times p} & K_\delta(I) & \xrightarrow{\rho_\delta^I} & K_\delta(I, \mathbb{Z}_p) & \longrightarrow & K_{1-\delta}(I) \\
 K_\delta(\iota) \downarrow & & K_\delta(\iota) \downarrow & & K_\delta(\iota, \mathbb{Z}_p) \downarrow & & K_{1-\delta}(\iota) \downarrow \\
 K_\delta(A) & \xrightarrow{\times p} & K_\delta(A) & \xrightarrow{\rho_\delta^A} & K_\delta(A, \mathbb{Z}_p) & \longrightarrow & K_{1-\delta}(A).
 \end{array}$$

By assumption, we have $K_\delta(\iota)$ are injective for $\delta = 0, 1$. Note that the following diagram is exact, hence, what we need to show is $K_\delta(\iota, \mathbb{Z}_p)$ are injective maps for $\delta = 0, 1$.

$$\begin{array}{ccccc}
 & & K_\delta(\iota, \mathbb{Z}_p) & & \\
 & & \uparrow & & \\
 K_\delta(I, \mathbb{Z}_p) & \xrightarrow{\quad} & K_\delta(A, \mathbb{Z}_p) & \longrightarrow & K_\delta(A/I, \mathbb{Z}_p) \\
 & & & & \downarrow \\
 K_{1-\delta}(A/I, \mathbb{Z}_p) & \longleftarrow & K_{1-\delta}(A, \mathbb{Z}_p) & \xleftarrow{K_{1-\delta}(\iota, \mathbb{Z}_p)} & K_{1-\delta}(I, \mathbb{Z}_p).
 \end{array}$$

Given $a_1 \in K_\delta(I, \mathbb{Z}_p)$ with $K_\delta(\iota, \mathbb{Z}_p)(a_1) = 0$, the standard diagram chasing gives $a_2 \in K_\delta(I) \subset K_\delta(A)$ and $a_3 \in K_\delta(A)$ such that

$$\rho_\delta^I(a_2) = a_1 \quad \text{and} \quad p \times a_3 = a_2.$$

Since

$$0 \rightarrow K_\delta(I) \rightarrow K_\delta(A) \rightarrow K_\delta(A/I) \rightarrow 0, \quad \delta = 0, 1$$

are pure exact, i.e., for $p \geq 2$ we have

$$p \times K_\delta(I) = K_\delta(I) \cap \{p \times K_\delta(A)\},$$

there is an $a_4 \in K_\delta(I)$ such that $p \times a_4 = a_2$. By exactness, we have $a_1 = 0$. \square

5.6. ([17, 2.2-2.3]) An \mathcal{AHD} algebra is an inductive limit of finite direct sums of the form $M_n(I_p)$ and $PM_n(C(X))P$, where I_p is the Elliott-Thomsen dimension-drop interval algebra and X is one of the following finite connected CW complexes: $\{pt\}$, \mathbb{T} , $[0, 1]$, $T_{II,k}$. $P \in M_n(C(X))$ is a projection and $T_{II,k}$ is the 2-dimensional connected simplicial complex with $H^1(T_{II,k}) = 0$ and $H^2(T_{II,k}) = \mathbb{Z}/k\mathbb{Z}$. (In [11], this class is called \mathcal{ASH} algebras.)

Proposition 5.7. ([10, Proposition 4.4]) *Suppose an extension*

$$0 \rightarrow I \rightarrow A \rightarrow A/I \rightarrow 0$$

is given. If A is an \mathcal{AHD} algebra of real rank zero, we have

- (i) *I and A/I are \mathcal{AHD} algebras of real rank zero.*
- (ii) *The ideal I in A is K -pure.*
- (iii) *For any $p \geq 2$ the sequences*

$$0 \rightarrow K_*(I; \mathbb{Z}_p) \rightarrow K_*(A; \mathbb{Z}_p) \rightarrow K_*(A/I; \mathbb{Z}_p) \rightarrow 0$$

are pure exact.

That is, all the $A\mathcal{H}\mathcal{D}$ algebras of real rank zero are K -pure.

Theorem 5.8. ([11, Proposition 4.12]) *Let A, B be separable C^* -algebras of real rank zero and stable rank one. Let $\phi : \underline{K}(A) \rightarrow \underline{K}(B)$ be a morphism of $\mathbb{Z}_2 \times \mathbb{Z}^+$ -graded groups. The following are equivalent.*

- (i) $\phi(\underline{K}(A))_+ \subset \underline{K}(B)_+$.
- (ii) $\phi(K_0^+(A)) \subset K_0^+(B)$ and $\phi(\underline{K}_I(A)) \subset \underline{K}_{I'}(B)$ for all ideals I in A .

Theorem 5.9. *Let A, B be K -pure (or $A\mathcal{H}\mathcal{D}$) algebras with $rr(A) = 0$ and $sr(A) = 1$. Then*

$$(\underline{\text{Cu}}(A), ([1_A], 0, \oplus_{p=1}^{\infty}(0, 0))) \cong (\underline{\text{Cu}}(B), ([1_B], 0, \oplus_{p=1}^{\infty}(0, 0)))$$

if and only if

$$(\underline{K}(A), \underline{K}(A)_+, [1_A]) \cong (\underline{K}(B), \underline{K}(B)_+, [1_B]).$$

Proof. The forward direction is our Proposition 5.3. For the backward direction, it is a combination of Corollary 3.9 and Lemma 5.5.

One can also rewrite a detail version like the proof of Theorem 2.9. As all $A\mathcal{H}\mathcal{D}$ algebras of real rank zero are K -pure, we only need to prove the K -pure case. Since $\phi : \underline{K}(A) \rightarrow \underline{K}(B)$ is a isomorphism of $\mathbb{Z}_2 \times \mathbb{Z}^+$ -graded groups satisfying $\phi(\underline{K}(A))_+ = \underline{K}(B)_+$. Then by Theorem 5.8, we get $\phi(K_0^+(A)) = K_0^+(B)$ and $\phi(\underline{K}_I(A)) = \underline{K}_J(B)$ for all ideals I in A and the relative J in B .

Note that for any ideal I of A , Lemma 5.5 shows that the map $K_*(I; \mathbb{Z}_p)$ is a sub-group of $K_*(A; \mathbb{Z}_p)$, for all $p \geq 2$. So do J and B . Thus we have the mod- p diagrams commutative for all $p \geq 2$ and $\delta = 0, 1$.

$$\begin{array}{ccccc}
 K_{\delta}(I) & \longrightarrow & K_{\delta}(I, \mathbb{Z}_n) & \longrightarrow & K_{1-\delta}(I) \\
 \parallel & \searrow & \parallel & \searrow & \parallel \\
 & & K_{\delta}(A) & \longrightarrow & K_{\delta}(A, \mathbb{Z}_n) & \longrightarrow & K_{1-\delta}(A) \\
 & & \parallel & & \parallel & & \parallel \\
 & & K_{\delta}(B) & \longrightarrow & K_{\delta}(B, \mathbb{Z}_n) & \longrightarrow & K_{1-\delta}(B) \\
 & & \parallel & & \parallel & & \parallel \\
 K_{\delta}(J) & \longrightarrow & K_{\delta}(J, \mathbb{Z}_n) & \longrightarrow & K_{1-\delta}(J)
 \end{array}$$

The rest part is as same as we did for K_* and Cu_1 .

□

Corollary 5.10. *By restricting to the category $\underline{\text{Cu}}^{\sim}$, we can recover \underline{K} from $\underline{\text{Cu}}_u$ through \underline{H} . A fortiori, we have $\underline{\text{Cu}}_u$ is a complete invariant for unital $A\mathcal{H}\mathcal{D}$ algebras of real rank zero with slow dimension growth (see Theorem 9.1 in [11]).*

Proof. Let

$$\Psi : \underline{\text{Cu}}_u(A) \rightarrow \underline{\text{Cu}}_u(B)$$

be a $\underline{\text{Cu}}^\sim$ -isomorphism with

$$\Psi([1_A], 0, \oplus_{p=1}^\infty (0, 0)) \leq ([1_B], 0, \oplus_{p=1}^\infty (0, 0)).$$

By Theorem 5.9 and [11, 9.1], the isomorphism from $(\underline{K}(A), \underline{K}(A)_+, [1_A])$ to $(\underline{K}(B), \underline{K}(B)_+, [1_B])$ can be lifted to a isomorphism ψ from A to B .

Under the assumption, ψ also induce the original $\underline{\text{Cu}}^\sim_u$ -isomorphism Ψ :, since $\underline{\text{Cu}}(A), \underline{\text{Cu}}(B)$ are exactly the completions of $\underline{K}(A)_+, \underline{K}(B)_+$, respectively. \square

Remark 5.11. For any separable, unital C^* -algebras with stable rank one and real rank zero, the unitary Cuntz semigroup can be recovered functorially from the invariant K_* (Theorem 2.9). But we don't know whether the total version Theorem 5.9 is still true or not, if we delete the assumption of “ K -pure (or $A\mathcal{H}\mathcal{D}$ algebras)”.

The key point is that the sequences

$$0 \rightarrow K_i(I) \rightarrow K_i(A) \rightarrow K_i(A/I) \rightarrow 0$$

are always exact; while for any $p \geq 2$, we are not sure whether the sequences

$$0 \rightarrow K_*(I; \mathbb{Z}_p) \rightarrow K_*(A; \mathbb{Z}_p) \rightarrow K_*(A/I; \mathbb{Z}_p) \rightarrow 0$$

are exact or not.

These above exact sequences will lead the commutative Mod- p diagram for $\delta = 0, 1$, which will generalize Theorem 5.9.

One more observation is that if we replaced “ K -pure (or $A\mathcal{H}\mathcal{D}$ algebras)” by “UCT”, then by [19], we have

$$K_*(I; \mathbb{Z}_p) \cong K_*(J; \mathbb{Z}_p).$$

The problem is we don't know if this isomorphism is compatible with that in the following diagram, while all arrows with solid line are natural maps.

$$\begin{array}{ccccc}
 K_\delta(I) & \xrightarrow{\quad} & K_\delta(I, \mathbb{Z}_n) & \xrightarrow{\quad} & K_{1-\delta}(I) \\
 \parallel & \searrow & \parallel & \searrow & \parallel \\
 & & K_\delta(A) & \xrightarrow{\quad} & K_\delta(A, \mathbb{Z}_n) & \xrightarrow{\quad} & K_{1-\delta}(A) \\
 & & \parallel & & \parallel & & \parallel \\
 & & K_\delta(B) & \xrightarrow{\quad} & K_\delta(B, \mathbb{Z}_n) & \xrightarrow{\quad} & K_{1-\delta}(B) \\
 & & \parallel & & \parallel & & \parallel \\
 K_\delta(J) & \xrightarrow{\quad} & K_\delta(J, \mathbb{Z}_n) & \xrightarrow{\quad} & K_{1-\delta}(J)
 \end{array}$$

At last, we have the following question.

Question 5.12. Let A be a C^* -algebra of stable rank one and real rank zero, does $(\underline{K}(A), \underline{K}(A)_+)$ determine $\underline{\text{Cu}}(A)$?

Note that if we replace the definition of $\underline{\text{Cu}}(A)$ by

$$\underline{\text{Cu}}^I(A) \triangleq \coprod_{I \in \text{Lat}(A)} \text{Cu}_f(I) \times K_1^I(A) \times \bigoplus_{p=1}^{\infty} K_*^I(A, \mathbb{Z}_p),$$

where $K_1^I(A)$ and $K_*^I(A, \mathbb{Z}_p)$ are images of $K_1(I)$ and $K_*(I, \mathbb{Z}_p)$ in $K_1(A)$ and $K_*(A, \mathbb{Z}_p)$, respectively.

We point that $\underline{\text{Cu}}^I(A)$ is also a continuous invariant, it is routine to get that $\underline{\text{Cu}}^I(A)$ and $(\underline{K}(A), \underline{K}(A)_+)$ are equivalent invariants for any C^* -algebras of stable rank one and of real rank zero. And for both simple case and *ASH* of real zero case, it is obvious that $\underline{\text{Cu}}^I$ and $\underline{\text{Cu}}$ coincide with each other.

As supplementary, we raised one more question.

Question 5.13. Let A be a C^* -algebra of stable rank one, when do we have $\underline{\text{Cu}}^I(A)$ and $\underline{\text{Cu}}(A)$ determine each other?

Under the real rank zero setting, an affirmative answer to Question 5.13 will give an affirmative to Question 5.12. But in general, these questions remain open for us.

ACKNOWLEDGEMENTS

The research of the first author was supported by the Fundamental Research Funds for the Central Universities (No.:2412021QD001) and NNSF of China(No.:12101113, No.:11920101001). The second author was support by NNSF of China (No.:12101102).

REFERENCES

- [1] Q. An, G. A. Elliott, Z. Li and Z. Liu. The classification of certain *ASH* C^* -algebras of real rank zero. *J. Topol. Anal.*, to appear. [10](#)
- [2] R. Antoine, F. Perera, L. Robert and H. Thiel, C^* -algebras of stable rank one and their Cuntz semigroups. *Duke Math. J.*, 171(2022), 33–99. [2](#)
- [3] R. Antoine, F. Perera and H. Thiel, Tensor Products and Regularity Properties of Cuntz Semigroups, *Mem. Amer. Math. Soc.*, 251 (2018), viii+191. [3](#), [4](#), [5](#)
- [4] L. G. Brown, Stable isomorphism of hereditary subalgebras of C^* -algebras. *Pacific J. Math.*, 71 (2) (1977), 335–348. [4](#)
- [5] N. P. Brown and A. Ciuperca, Isomorphism of Hilbert modules over stably finite C^* -algebras, *J. Funct. Anal.*, 257 (1) (2009), 332–339. [16](#)
- [6] L. G. Brown and G. K. Pedersen, C^* -algebras of real rank zero, *J. Funct. Anal.*, 99 (1991), 131–149. [7](#)
- [7] L. Cantier, A unitary semigroup for C^* -algebras of stable rank one, *J. Funct. Anal.*, 281 (9) (2021), 109175. [1](#), [2](#), [4](#), [5](#), [6](#), [7](#), [8](#), [17](#)
- [8] A. Ciuperca, Some properties of the Cuntz semigroup and an isomorphism theorem for a certain class of non-simple C^* -algebras, ProQuest LLC, Ann Arbor, MI, 2008. Thesis (Ph.D.)-University of Toronto (Canada). [3](#), [4](#)
- [9] K. T. Coward, G. A. Elliott and C. Ivanescu, The Cuntz semigroup as an invariant for C^* -algebras, *J. Reine Angew. Math.*, 623 (2008), 161–193. [3](#), [4](#), [6](#), [18](#)
- [10] M. Dadarlat and S. Eilers. Compressing coefficients while preserving ideals in the K-theory for C^* -algebras, *K-Theory*, 14 (1998), 281–304. [23](#)

- [11] M. Dadarlat and G. Gong, A classification result for approximately homogeneous C^* -algebras of real rank zero, *Geom. Funct. Anal.*, 7 (4) (1997), 646–711. [10](#), [15](#), [23](#), [24](#), [25](#)
- [12] G. A. Elliott, On the classification of C^* -algebras of real rank zero, *J. Reine Angew. Math.*, 443 (1993), 179–219. [1](#), [2](#)
- [13] G. A. Elliott, Hilbert modules over C^* -algebras of stable rank one, *C. R. Math. Acad. Sci. Soc. R. Can.*, 29 (2007), 48–51. [6](#)
- [14] G. A. Elliott and G. Gong, On the classification of C^* -algebras of real rank zero II, *Ann. of Math.*, 144 (2) (1996), 497–610. [6](#)
- [15] G. A. Elliott, G. Gong and H. Su, On the classification of C^* -algebras of real rank zero. IV. Reduction to local spectrum of dimension two, *Operator Algebras and their applications[C]// Fields Inst Commun.* 1998. [1](#), [9](#), [10](#)
- [16] G. Gong, Classification of C^* -Algebras of Real Rank Zero and Unsuspended E-Equivalence Types, *J. Funct. Anal.*, 152 (2) (1998), 281–329. [10](#)
- [17] G. Gong, C. Jiang and L. Li, Hausdorffified algebraic K_1 -group and invariants for C^* -algebras with the ideal property, *Ann. K-Theory*, 5 (1) (2020), 43–78. [2](#), [23](#)
- [18] H. Lin, *An Introduction to the Classification of Amenable C^* -Algebras*, World Scientific Publishing Co. Inc., River Edge, NJ, 2001. [7](#)
- [19] J. Rosenberg and C. Schochet, The Künneth theorem and the universal coefficient theorem for Kasparov’s generalized K-functor, *Duke Math. J.*, 55 (1987), 431–474. [25](#)
- [20] M. Rørdam and W. Winter, The Jiang–Su algebra revisited, *J. Reine Angew. Math.*, 642 (2010), 129–155. [6](#)
- [21] C. Schochet, Topological methods for C^* -algebras IV: Mod p K-theory, *Pacific J. Math.*, 114 (1984), 447–468. [10](#)
- [22] H. Thiel, Ranks of operators in simple C^* -algebras with stable rank one. *Commun. Math. Phys.*, 377 (2020), 37–76. [2](#)
- [23] H. Thiel and E. Vilalta, Covering dimension of Cuntz semigroups. *Adv. Math.*, to appear. [3](#)

SCHOOL OF MATHEMATICS AND STATISTICS, NORTHEAST NORMAL UNIVERSITY, CHANGCHUN, 130024, CHINA

Email address: qingnanan1024@outlook.com

SCHOOL OF MATHEMATICAL SCIENCES, DALIAN UNIVERSITY OF TECHNOLOGY, DALIAN, 116024, CHINA

Email address: lzc.12@outlook.com