A Characterization of Weak Proximal Normal Structure and Best Proximity Pairs

Abhik Digar, Rafael Espínola García and G. Sankara Raju Kosuru

Mathematics Subject Classification (2010). 47H10, 46C20, 54H25.

Abstract. The aim of this paper is to address an open problem given in [Kirk, W. A., Shahzad, Naseer, Normal structure and orbital fixed point conditions, J. Math. Anal. Appl. vol 463(2), (2018) 461–476]. We give a characterization of weak proximal normal structure using best proximity pair property. We also introduce a notion of pointwise cyclic contraction wrt orbits and therein prove the existence of a best proximity pair in the setting of reflexive Banach spaces.

1. Introduction and Preliminaries

Let A, B be two non-empty subsets of a Banach space and T be a cyclic mapping on $A \cup B$ $(T(A) \subseteq B, T(B) \subseteq A)$. A pair $(x, y) \in A \times B$ is said to be a best proximity pair for T if $||x - Tx|| = ||y - Ty|| = d(A, B) = \inf\{||x - y|| : d(A, B) = \inf\{|x - y|| :$ $x \in A, y \in B$. The geometry of Banach spaces plays a crucial role for the existence of best proximity pairs. The analysis of proximal normal structure and weak or semi-normal structure, the property UC, the projectional property due to Eldred et al. [2], Moosa [4], Suzuki et al. [9], G. S. Raju et al. [7] etc., respectively are widely used to prove the existence of a best proximity pair for cyclic maps. We denote $\sup\{||x-y||: y \in B\}$ for $x \in A$ by $\delta(x, B)$. We shall say that the pair (A, B) is proximinal pair if for every x in A (resp. in B), there exists y in B (resp. in A) such that ||x - y|| = d(A, B). Further, if such a y is unique, then (A, B) is said to be a sharp proximinal pair [7]. In this case we denote y by x'. Also, (A, B) is said to be a proximinal parallel pair if (A, B) is sharp proximinal and B = A + h for some $h \in X$ [3]. It is shown in [3] that if X is strictly convex and A, B are weakly compact convex subsets of X, then every (A_0, B_0) is a non-empty proximinal parallel pair. Here $A_0 = \{x \in A : \text{there exists } y \in B \text{ such that } ||x - y|| = d(A, B)\}$

and $B_0 = \{x \in B : \text{ there exists } y \in A \text{ such that } ||x - y|| = d(A, B)\}.$ Also, in [7], the authors have given example(s) of sharp proximinal pair which are not parallel. In [2], the author introduced a geometrical notion called proximal normal structure to prove the existence of a best proximity pair of a relatively nonexpansive mapping $(||Tx - Ty|| \le ||x - y||)$ for all $x \in A, y \in B$.) We say (A, B) has proximal normal structure ([2]) [respectively weak proximal normal structure ([5]) if (A, B) is convex and for any closed bounded [respectively weakly compact] convex proximinal pair (H_1, H_2) of subsets of (A, B) for which $d(H_1, H_2) = d(A, B)$ and $\delta(H_1, H_2) > \delta(H_1, H_2)$ $d(H_1, H_2)$, there exists $(x, y) \in (H_1, H_2)$ such that $\delta(x, H_2) < \delta(H_1, H_2)$ and $\delta(y, H_1) < \delta(H_1, H_2)$. It is well known that every non-empty closed bounded convex pair (A, B) of a uniformly convex Banach space has proximal normal structure. In fact, every non-empty compact convex pair (A, B)of a Banach space has proximal normal structure. It is proved (Proposition 3.2 in [5]) that a bounded convex pair has proximal normal structure if and only if it doesn't contain any proximal diametral sequence. A pair $(\{x_n\}, \{y_n\})$ of sequences in (A, B) with $||x_n - y_n|| = d(A, B), n \ge 1$ is said to be a proximal diametral sequence ([5]) if $d(A, B) < \delta(\{x_n\}, \{y_n\})$ and $\max\{\lim_{n \to \infty} d(x_{n+1}, \operatorname{co}(\{y_1, y_2, ..., y_n\})), \lim_{n \to \infty} d(y_{n+1}, \operatorname{co}(\{x_1, x_2, ..., x_n\}))\} = 0$ $\delta(\{x_n\}, \{y_n\})$. It is easy to see that proximal normal structure coincides with weak proximal normal structure in reflexive Banach spaces [5]. Moreover, therein the author proved the existence of a best proximity pair in the settings of a reflexive Banach space. Recently, in [6], the authors posed an open problem for the existence of a best proximity pair for a more general class of mappings, called relatively orbital nonexpansive mappings. Also therein the authors indicated that an affirmative answer may provide a characterization of proximal normal structure. Motivated by this, we aim to give a partial affirmative answer for the same. We also provide a characterization of weak proximal normal structure by using the existence of a best proximity pair for relatively orbital nonexpansive mappings. Finally, we introduce the notion of pointwise cyclic contraction wrt orbits and prove the minimal invariant subsets of such a map have nondiametral points. This guarantees the existence of a best proximity pair for such a class in the setting of a reflexive Banach space. Finally, we prove the existence of a best proximity pair for the class of pointwise cyclic contraction wrt orbits.

2. Existence of Best Proximity Pairs

Let A, B be two closed convex subsets of a Banach space X. Let $T : A \cup B \to A \cup B$ be a cyclic map. If T admits a best proximity pair, then $A_0 \neq \emptyset \neq B_0$. Also, if T is relatively nonexpansive, then $A_0 \cup B_0$ is cyclically invariant under T ($TA_0 \subseteq B_0, TB_0 \subseteq A_0$). The following theorem is due to Eldred *et al.* [2]

Theorem 2.1. Let (K_1, K_2) be a non-empty weakly compact convex pair in a Banach space and suppose (K_1, K_2) has proximal normal structure. Then every relatively nonexpansive mapping T on $A \cup B$ has a best proximity pair in (K_1, K_2) .

The main tool to prove the same is to use the geometrical notion called "proximal normal structure" on $A_0 \cup B_0$. Later many authors established the existence of a best proximity pair for relatively nonexpansive mappings in different settings using variants of geometry ([3],[4],[8],[9]). In [5], Moosa introduced pointwise relatively nonexpansive mappings involving orbits and therein proved the existence of a best proximity pair for such a class of mappings. Recently, in 2018, Kirk and Shahzad discussed the existence of a best proximity pair for relatively nonexpansive mappings and therein they raised the question "can the assumption that T is relatively nonexpansive in Theorem 2.1 be replaced by the assumption that T is relatively nonexpansive wrt orbits?" T is said to be relatively nonexpansively mappings wrt orbits if $||Tx - Ty|| \leq r_x (\mathcal{O}(y)) = \delta(x, \{y, Ty, T^2y, ...\})$. Using the following example, we can conclude that the answer is negative for the above open problem.

Example 2.2. Let $A = \{x \in \mathbb{R} : -2 \le x \le -1\}, B = \{x \in \mathbb{R} : 1 \le x \le 2\}.$ Define

$$T(x) = \begin{cases} -x, & \text{if } x \in A\\ -1 - \frac{x}{2}, & \text{if } x \in B. \end{cases}$$

Let $y \in B$. For any n, $T^{2n}y = 1 + \frac{2^{n-1}-1}{2^{n-1}} + \frac{y}{2^n} = 2 - \frac{1}{2^{n-1}} + \frac{y}{2^n}$ and $T^{2n+1}y = -1 - \frac{T^{2n}y}{2} = -2 + \frac{1}{2^n} - \frac{y}{2^{n+1}}$. Now, for any $x \in A, y \in B$, $||Tx - Ty|| = |(-x) - (-1 - \frac{y}{2})| \le 2 - x = r_x (\mathcal{O}(y))$.

It is to be observed that a cyclic map T on $A \cup B$ that satisfies $||Tx - Ty|| \leq r_x (\mathcal{O}(y))$ does not guarantee $A_0 \cup B_0$ is cyclically invariant under T. Hence, it is not reasonable to expect the existence of a best proximity pair for such a map T. To overcome this, we redefine the relatively orbital nonexpansive mappings. For $x \in A \cup B$, we denote $\{T^{2n}x : n \in \mathbb{N} \cup \{0\}\}$ by $\mathcal{O}^2(x)$.

Definition 2.3. Let A, B be two non-empty subsets of a Banach space X. A cyclic map $T: A \cup B \to A \cup B$ is said to be a relatively orbital nonexpansive mapping if

- (i) ||Tx Ty|| = d(A, B) if ||x y|| = d(A, B) for $x \in A, y \in B$.
- (ii) for all $x \in A, y \in B$, $||Tx Ty|| \le \min\{r_x (\mathcal{O}^2(y)), r_y (\mathcal{O}^2(x))\}.$

It is worth mentioning that relatively orbital nonexpanive mapping is not necessarily relatively nonexpansive.

Example 2.4. Let $A = \{(0, x) \in \mathbb{R}^2 : 0 \le x \le 1\}, B = \{(1, y) \in \mathbb{R}^2 : 0 \le y \le 1\}$ and $T : A \cup B \to A \cup B$ be defined by

$$x \in A, \ T(x) = \begin{cases} (1, \frac{x}{4}) & \text{if } x \ge \frac{1}{2}; \\ (1, \frac{x}{2}) & \text{if } x < \frac{1}{2}. \end{cases}$$

$$y \in B, \ T(y) = \begin{cases} (0, \frac{y}{4}) & \text{if } y \ge \frac{1}{2}; \\ (0, \frac{y}{2}) & \text{if } y < \frac{1}{2}. \end{cases}$$

We see that T is not relatively nonexpansive but relatively orbital nonexpansive mapping.

Let (A, B) be a non-empty sharp proximinal pair in Banach space and T be a relatively orbital nonexpansive mapping on $A \cup B$. Then it is easy to see that (A_0, B_0) is cyclically invariant under T and Tx' = (Tx)'. We say that (A, B) is said to satisfy the weak best proximity pair property (WBPP) if every relatively orbital nonexpansive mapping on $A \cup B$ has a best proximity pair. The following theorem ensures that every non-empty weakly compact convex pair of subsets of a strictly convex Banach space satisfying the WBPP. The following theorem is in a way different than Theorem 2.6 of [5]. For the sake the completeness, we prove the same here.

Theorem 2.5. Let A, B be two non-empty weakly compact convex substes of a Banach space X. If (A, B) is a sharp proximinal pair having weak proximal normal structure, then (A, B) has WBPP.

Proof. Let \mathscr{F} denote the collection of non-empty closed bounded convex proximinal pair (E_1, E_2) of subsets of (A_0, B_0) with (E_1, E_2) cyclically invariant under T and $d(E_1, E_2) = d(A, B)$. $\mathscr{F} \neq \emptyset$, since $(A_0, B_0) \in \mathscr{F}$. By Zorn's Lemma \mathscr{F} has a minimal element under the set inclusion order " \subseteq ", say, (F_1, F_2) . If (F_1, F_2) is a singleton pair, we have $\delta(F_1, F_2) = d(A, B)$, i.e., T has a best proximity pair. Suppose (F_1, F_2) is not singleton. By weak proximal normal structure, there exist $(x_1, y_1) \in (F_1, F_2)$ such that $m_1 = \delta(x_1, F_2) < \delta(F_1, F_2)$; $m_2 = \delta(y_1, F_1) < \delta(F_1, F_2)$. Set $m = \max\{m_1, m_2\}$. Define

$$L_1 = \{x \in F_1 : \delta(x, F_2) \le m\}$$

$$L_2 = \{y \in F_2 : \delta(y, F_1) \le m\}.$$

 $L_1 \neq \emptyset, L_2 \neq \emptyset$, since $x_1 \in L_1, y_1 \in L_2$. Being closed subset of a weakly compact subset, L_1, L_2 are weakly compact. To see L_1 is convex, let $a, b \in L_1$. For any $\lambda \in [0, 1]$,

 $\delta(\lambda a + (1 - \lambda)b, F_2) \leq \lambda \delta(a, F_2) + (1 - \lambda)\delta(b, F_2) \leq \lambda m + (1 - \lambda)m = m.$ Hence we can conclude that (L_1, L_2) is a convex pair. Let $v \in F_2$. Suppose the unique best approximation of an element $z \in A \cup B$ is denoted by z'. Then

$$\begin{aligned} \left\| \frac{x_1 + y_1'}{2} - v \right\| &\leq \frac{1}{2} \left[\|x_1 - v\| + \|y_1' - v\| \right] \\ &= \frac{1}{2} \left[\|x_1 - v\| + \|y_1 - v'\| \right] \\ &\leq \frac{1}{2} \left[\delta(x_1, F_2) + \delta(y_1, F_1) \right] \\ &\leq m. \end{aligned}$$

Since $v \in F_2$ is arbitrary, $\delta\left(\frac{x_1+y'_1}{2}, F_2\right) \leq m$. Hence, $\frac{x_1+y'_1}{2} \in L_1$ Similarly, $\frac{x'_1+y_1}{2} \in L_2$. Moreover, $\left\|\frac{x_1+y'_1}{2} - \frac{x'_1+y_1}{2}\right\| = d(A, B)$. Hence, $d(L_1, L_2) = d(A, B)$. To see (L_1, L_2) is a proximinal pair, let $x \in L_1$. Then $x \in F_1$ and hence $x' \in F_2$. Therefore $\delta(x', F_1) = \delta(x, F_2) \leq m$. Thus $x' \in L_2$. It infers (L_1, L_2) is a proximinal pair. Thus, $L_2 = \{x' \in F_2 : x \in L_1\}$.

Next, let $x \in L_1, v \in F_2$. Then, $||Tx - Tv|| \leq r_x (\mathcal{O}^2(v)) = \delta(x, \mathcal{O}^2(v)) \leq \delta(x, F_2) \leq m$. It follows that $T(F_2) \subset B(Tx; m) \cap F_1 = F'_1$. Similarly, $T(F_1) \subset B(Tx'; m) \cap F_2 = F'_2$. Clearly, $(F'_1, F'_2) \in \mathscr{F}$. By minimality, $F'_1 = F_1, F'_2 = F_2$. Then $F_1 \subseteq B(Tx; m)$ and $F_2 \subseteq B(Tx'; m)$. For any $u \in F_1, ||u - Tx|| \leq m$, hence, $\delta(Tx, F_1) \leq m$. Therefore, $Tx \in L_2$. Hence, $T(L_1) \subseteq L_2$. Further, if $y \in L_2$, then $y' \in L_1$. This implies $Ty' = (Ty)' \in L_2$. Thus $Ty \in L_1$. As $y \in L_2$ is arbitrary, we have $T(L_2) \subseteq L_1$. Hence, $(L_1, L_2) \in \mathscr{F}$. For $x \in L_1, y \in L_2, ||x - y|| \leq \delta(x, F_2) \leq m < \delta(F_1, F_2)$. This infers that $\delta(L_1, L_2) < \delta(F_1, F_2)$. This contradicts the minimality of (F_1, F_2) .

Let T be a cyclic map on $A \cup B$. We say that the pair (A, B) has a proximinal nondiametral pair if there exists $(x, y) \in A \times B$ such that $\max\{\delta(x, B), \delta(y, A)\} < \delta(A, B)$ whenever $d(A, B) < \delta(A, B)$. A similar technique can be used to obtain the following:

Theorem 2.6. Let (A, B) be a non-empty closed bounded convex proximinal pair of subsets of a Banach space and let T be a relatively orbital nonexpansive mapping on $A \cup B$. If T has a nonempty closed bounded convex minimal cyclically invariant pair (A, B) having a nondiametral pair then T has a best proximity pair.

Example 2.7. Let A, B and T as in the Example 2.4. It is easy to see that ((0,0), (1,0)) is a best proximity pair.

3. Characterization of weak proximal normal structure

Let (A, B) be a bounded convex proximinal pair of a Banach space X. A nonconstant pair of sequences $(\{x_n\}, \{y_n\})$ of (A, B) is said to be a proximinal diametral sequence if $||x_n - y_n|| = d(A, B)$ for every $n \in \mathbb{N}$ and $\delta(\{x_n\}, \{y_n\}) = \lim_{n \to \infty} d(x_{n+1}, \operatorname{co}(\{y_1, y_2, ..., y_n\})) = \lim_{n \to \infty} d(y_{n+1}, \operatorname{co}(\{x_1, x_2, ..., x_n\}))$. It is to be observed that if d(A, B) = 0, then the proximinal diametral sequence turns out to be a diametral sequence in $A \cap B$ in the sense of Brodskii and Milman ([1]). Using a similar argument employed in the proof of Theorem 2.5 ([2]) one can obtain the following:

Theorem 3.1. A bounded convex pair (A, B) of a Banach space X has proximal normal structure if and only if it does not contain a proximinal diametral sequence.

Let (A, B) be a non-empty weakly compact convex sharp proximinal pair of subsets of a Banach space having WBPP. Suppose (A, B) does not have proximal weak normal structure. Then by Theorem 3.1, (A, B) has a proximinal diametral sequence, say, $(\{x_n\}, \{y_n\})$. Consequently, $\lim_{n \to \infty} d(x_{n+1}, \operatorname{co}(\{y_1, y_2, \ldots, \delta(\{x_n\}, \{y_n\}) = \lim_{n \to \infty} d(y_{n+1}, \operatorname{co}(\{x_1, x_2, \ldots, x_n\}))$.

Since, (A, B) is weakly compact, there exists a subsequence $(\{x_{n_k}\}, \{y_{n_k}\})$ of $(\{x_n\}, \{y_n\})$ which is weakly convergent. It is easy to see that the sequence $(\{x_{n_k}\}, \{y_{n_k}\})$ is a proximinal diametral subsequence. Hence, without loss of any generality, we may assume that the sequence $(\{x_n\}, \{y_n\})$ is proximinal diametral and weakly convergent. Now, $H = \overline{\operatorname{co}}(\{x_1, x_2, \ldots\}), K = \overline{\operatorname{co}}(\{y_1, y_2, \ldots\})$ are weakly compact convex subsets of A, B respectively. Define $T: H \cup K \to H \cup K$ by

$$T(x) = \begin{cases} y_1, & \text{if } x \notin \{x_n : n \in \mathbb{N}\} \\ y_{n+1}, & \text{if } x = x_n \text{ for some } n \in \mathbb{N}; \end{cases}$$
$$T(y) = \begin{cases} x_1, & \text{if } y \notin \{y_n : n \in \mathbb{N}\} \\ x_{n+1}, & \text{if } y = y_n \text{ for some } n \in \mathbb{N}. \end{cases}$$

Clearly, $\delta(H, K) = \delta(\{x_n\}, \{y_n\})$ and $\lim_{n \to \infty} ||x_n - z|| = \delta(H, K) = \lim_{n \to \infty} ||y_n - v||$ for any $z \in K, v \in H$. Hence, $r_x(\mathcal{O}^2(y)) = \delta(H, K)$ for each $x \in H, y \in K$. Now,

$$||Tx - Ty|| \le \delta(H, K) = r_x \left(\mathcal{O}^2(y)\right)$$
 for each $x \in H, y \in K$.

Also, if $(x, y) \in H \times K$ with ||x - y|| = d(H, K), then ||Tx - Ty|| = d(H, K). Therefore T is a relatively orbital nonexpansive mapping. As (A, B) is a sharp proximinal pair, then so is (H, K) and T does not have any best proximity pair. Thus we have the following:

Proposition 3.2. Let A, B be two non-empty weakly compact convex substes of a Banach space X. If (A, B) is a sharp proximinal pair and (A, B) has WBPP, then (A, B) has weak proximal normal structure.

By Theorem 2.5 and Proposition 3.2 we have the following characterization:

Theorem 3.3. Let A, B be two non-empty weakly compact convex subsets of a Banach space X. If (A, B) is a sharp proximinal pair, then (A, B) has weak proximal normal structure if and only if every relatively orbital nonexpansive mapping $T : A \cup B \to A \cup B$ has a best proximity pair.

4. Pointwise Cyclic Contraction wrt Orbits

Let (A, B) be a pair of subsets of a normed linear space. A cyclic map T on $A \cup B$ is said to be a proximal pointwise contraction if for any $x \in A$, there exists $\alpha(x) \in [0, 1)$ such that $||Tx - Ty|| \leq \alpha(x)||x - y||$ ([10]). Later many authors obtained the existence of a best proximity pair for certain types of pointwise cyclic contractions ([8], [11], [12]). Now we introduce the notion of pointwise cyclic contraction wrt orbits and prove the existence of a best

proximity pair for such a map. Our result is a generalization of the main results given in the aforementioned articles.

Definition 4.1. A cyclic map T on a non-empty pair (A, B) of subsets of a Banach space is said to be pointwise cyclic contraction wrt orbits if it satisfies

- (i) ||Tx Ty|| = d(A, B) whenever ||x y|| = d(A, B) for $(x, y) \in A \times B$;
- (ii) for each $(x, w) \in (A, B)$ there exists $\alpha(x), \alpha(w) \in (0, 1)$ such that $||Tx - Ty|| \leq \alpha(x)r_x \left(\mathcal{O}^2(y)\right) + (1 - \alpha(x)) d(A, B)$ for all $y \in B$, and $||Tw - Tu|| \leq \alpha(w)r_w \left(\mathcal{O}^2(u)\right) + (1 - \alpha(w)) d(A, B)$ for all $u \in A$.

It is easy to see that every pointwise cyclic contraction mapping wrt orbits is relatively orbital nonexpansive.

Theorem 4.2. Suppose (A, B) is a closed, weakly compact, convex, sharp proximinal pair of a Banach space X and $T : A \cup B \to A \cup B$ is a pointwise cyclic contraction wrt orbits. Then T has a best proximity pair.

Proof. Let \mathscr{F} denote the collection of all non-empty proximal closed convex subsets (H_1, H_2) of (A_0, B_0) such that $TH_1 \subseteq H_2, TH_2 \subseteq H_1$ and $d(H_1, H_2) = d(A, B)$. Since $A_0 \cup B_0 \in \mathscr{F}$, we have $\mathscr{F} \neq \emptyset$. By Zorn's lemma, \mathscr{F} has a minimal, say, (K_1, K_2) . Let $(x, y) \in (K_1, K_2)$ such that $||x - y|| = d(K_1, K_2) = d(A, B)$. If $\delta(x, K_2) = d(A, B)$, then $d(A, B) = d(K_1, K_2) \leq ||x - Tx|| \leq \delta(x, K_2) = d(A, B)$. This infers ||x - Tx|| = d(A, B). Since, T is pointwise cyclic contraction wrt orbits, we have $||Tx - T^2x|| = d(A, B)$. Therefore, (x, Tx) is a best proximity pair. Similarly, if $\delta(y, K_1) = d(A, B)$, then (y, Ty) is a best proximity pair. Hence, we may assume that $\delta(x, K_2) > d(A, B)$ and $\delta(y, K_1) > d(A, B)$. Define

$$K_x = \{ z \in K_1 : ||z - Tx|| \le \alpha(x)\delta(x, K_2) + (1 - \alpha(x)) d(A, B) \}; K_y = \{ w \in K_2 : ||w - Ty|| \le \alpha(y)\delta(y, K_1) + (1 - \alpha(x)) d(A, B) \}.$$

Since

$$||Tx - Ty|| = d(A, B) = \alpha(x)d(A, B) + (1 - \alpha(x))d(A, B) < \alpha(x)\delta(x, K_2) + (1 - \alpha(x))d(A, B).$$

Then $(Ty, Tx) \in (K_x, K_y)$ and hence $K_x \neq \emptyset \neq K_y$. It is easy to see that (K_x, K_y) is convex. If $\{u_n\}_{n=1}^{\infty} \subset K_x$ is a sequence converges to $u \in X$ weakly, then $u \in K_1$. Now, $||u - Tx|| \leq \liminf\{||u_n - Tx|| : n \in \mathbb{N}\} \leq \alpha(x)\delta(x, K_2) + (1 - \alpha(x))d(A, B)$. Then $u \in K_x$ and K_x is closed. Further, for any $u \in K_x$, $||Tu - Ty|| \leq \alpha(y)r_y(\mathcal{O}^2(u)) + (1 - \alpha(y))d(A, B) \leq \alpha(y)\delta(y, K_1) + (1 - \alpha(y))d(A, B)$. This implies that $Tu \in K_y$. Hence, $TK_x \subseteq K_y$. Similarly, $TK_y \subseteq K_x$. Therefore, $(K_x, K_y) \in \mathscr{F}$. By minimality, $K_x = K$, $K_y = K_2$. Now, for any $w \in K_2$, $||w - Ty|| \leq \alpha(y)\delta(y, K_1) + (1 - \alpha(y))d(A, B) < \delta(K_1, K_2)$. Hence, $\delta(Ty, K_2) < \delta(K_1, K_2)$. Similarly, $\delta(Tx, K_1) < \delta(K_1, K_2)$. Thus (K_1, K_2) has a proximinal nondiametral pair. By Theorem 2.6, T has a best proximity pair.

References

- [1] Brodskiĭ, M. S., Mil'man, D. P., On the center of a convex set, Doklady Akad. Nauk SSSR (N.S.), vol 59, (1948) 837–840.
- [2] Eldred, A. Anthony, Kirk, W. A., Veeramani, P., proximal normal structure and relatively nonexpansive mappings, Studia Math., vol 171(3), (2005) 283–293.

[3] Espínola, Rafa, A new approach to relatively nonexpansive mappings, Proc. Amer. Math. Soc., vol 136(6), (2008) 1987–1995.

- [4] Gabeleh, Moosa, Shahzad, Naseer, Seminormal structure and fixed points of cyclic relatively nonexpansive mappings, Abstr. Appl. Anal., (2014) 1085-3375.
- [5] Gabeleh, Moosa, A characterization of proximal normal structure via proximal diametral sequences, J. Fixed Point Theory Appl. vol 19(4), (2017) 2909–2925.
- [6] Kirk, W. A., Shahzad, Naseer, Normal structure and orbital fixed point conditions, J. Math. Anal. Appl. vol 463(2), (2018) 461–476.
- [7] Raju Kosuru, G. Sankara, Veeramani, P., On existence of best proximity pair theorems for relatively nonexpansive mappings, J. Nonlinear Convex Anal., vol 11(1), (2010) 71–77.
- [8] Kosuru, G. Sankara Raju, Veeramani, P., A note on existence and convergence of best proximity points for pointwise cyclic contractions, Numer. Funct. Anal. Optim., vol 32(7), (2011) 821–830.
- [9] Suzuki, Tomonari, Kikkawa, Misako, Vetro, Calogero, The existence of best proximity points in metric spaces with the property UC, Nonlinear Anal., vol **71(7-8)**, (2009) 2918–2926.
- [10] Anuradha, J. and Veeramani, P., Proximal pointwise contraction, Topology Appl., vol 156(18), (2009) 2942–2948.
- [11] Mongkolkeha, Chirasak and Kumam, Poom, Best proximity points for asymptotic proximal pointwise weaker Meir-Keeler-type ψ -contraction mappings, J. Egyptian Math. Soc., vol **21(2)**, (2013) 87–90.
- [12] Gabeleh, Moosa, On generalized pointwise noncyclic contractions without proximal normal structure, Ann. Funct. Anal., vol 9(2), (2018) 220–232.

Abhik Digar Department of Mathemtics IIT Ropar Rupnagar - 140 001 Punjab, India. e-mail: abhikdigar@gmail.com

Rafael Espínola García Departamento de Análisis Matemático, Facultad de Matemáticas, IMUS, Universidad de Sevilla, 41010, Sevilla, Spain. e-mail: espinola@us.es G. Sankara Raju Kosuru Department of Mathemtics IIT Ropar Rupnagar - 140 001 Punjab, India. e-mail: raju@iitrpr.ac.in