arXiv:2201.10793v2 [math.DG] 13 Feb 2022

THE PERTURBATION OF THE DE RHAM HODGE OPERATOR AND THE KASTLER-KALAU-WALZE TYPE THEOREM FOR MANIFOLDS WITH BOUNDARY

SIYAO LIU TONG WU YONG WANG*

Abstract. In this paper, we give Lichnerowicz type formulas for the perturbation of the de Rham Hodge operator. We prove the Kastler-Kalau-Walze type theorems for the perturbation of the de Rham Hodge operator on 4-dimensional and 6-dimensional compact manifolds with or without boundary. Some concrete examples of the perturbation of the de Rham Hodge operator are provided for our main theorems.

1. Introduction

The noncommutative residue was found in [1, 2]. Since noncommutative residues are of great importance to the study of noncommutative geometry, more and more attention has been attached to the study of noncommutative residues. By using the noncommutative residue, Connes derived a conformal 4-dimensional Polyakov action analogy [3]. Connes showed us that the noncommutative residue on a compact manifold M coincided with the Dixmiers trace on pseudodifferential operators of order $-\dim M$ in [4]. Connes put forward that the noncommutative residue of the square of the inverse of the Dirac operator was proportioned to the Einstein-Hilbert action, which is called the Kastler-Kalau-Walze theorem now. Kastler gave a brute-force proof of this theorem [5]. In the same time, Kalau and Walze proved this theorem in the normal coordinates system [6]. Ackermann proved this theorem by using the heat kernel expansion, in [7]. The result of Connes was extended to the higher dimensional case [8]. Fedosov et al. gave the definition about the noncommutative residues on Boutet de Monvel algebra [9].

On the other hand, Wang generalized the Connes' results to the case of manifolds with boundary in [10, 11], and proved the Kastler-Kalau-Walze type theorem for the Dirac operator and the signature operator on lower-dimensional manifolds with boundary. In [12, 13], for the Dirac operator and the signature operator, Wang computed $\widetilde{\mathrm{Wres}}[\pi^+D^{-1}\circ \pi^+D^{-1}]$, in these cases the boundary term vanished. But for $\widetilde{\mathrm{Wres}}[\pi^+D^{-1}\circ \pi^+D^{-3}]$, authors got a nonvanishing boundary term [14], and gave a theoretical explanation for

2010 Mathematics Subject Classification: 53C40; 53C42.

Key words and phrases: Perturbation of the de Rham Hodge operator; Lichnerowicz type formulas; Noncommutative residue; Kastler-Kalau-Walze type theorems.

* *Corresponding author*.

gravitational action on boundary. In other words, Wang provided a kind of method to study the Kastler-Kalau-Walze type theorem for manifolds with boundary. In [15], Wei and Wang proved the Kastler-Kalau-Walze type theorem for modified Novikov operators on compact manifolds. In [16], Wu-Wang-Wang obtained two Lichnerowicz type formulas for the Dirac-witten operators, and gave the proof of Kastler-Kalau-Walze type theorems for the Dirac-witten operators on 4-dimensional and 6-dimensional compact manifolds with (resp.without) boundary. In [17], Wang proved a Kastler-Kalau-Walze type theorem for perturbations of the Dirac operators on compact manifolds with or without boundary.

The motivation of this paper is to prove the Kastler-Kalau-Walze type theorems for the perturbation of the de Rham Hodge operator. Specifically, we calculate $\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+(D_A^*)^{-1}]$, $\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+{D_A}^{-1}]$, $\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+(D_A^*D_AD_A^*)^{-1}]$ and $\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+{D_A}^{-3}]$, for D_A see (2.5).

A brief description of the organization of this paper is as follows. In Section 2, this paper will firstly introduce the basic notions of the perturbation of the de Rham Hodge operator, by means of which we can compute the Lichnerowicz formulas for the perturbation of the de Rham Hodge operator. And then we present the Kastler-Kalau-Walze type theorems for the perturbation of the de Rham Hodge operator on n-dimensional compact manifolds without boundary. In the next section, we calculate $\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+(D_A^*)^{-1}]$ and $\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+{D_A}^{-1}]$ for 4-dimensional compact manifolds with boundary. In Section 4, we prove the Kastler-Kalau-Walze type theorems on 6-dimensional compact manifolds with boundary for the perturbation of the de Rham Hodge operator.

2. The perturbation of the de Rham Hodge operator and its Lichnerowicz formulas

We give some definitions and basic notions which we will use in this paper. Let M be a n-dimensional ($n\geq 3$) oriented compact Riemannian manifold with a Riemannian metric g^M. And let ∇^L be the Levi-Civita connection about g^M. In the local coordinates $\{x_i;1\leq i\leq n\}$ and the fixed orthonormal frame $\{\widetilde{e_1},\cdots,\widetilde{e_n}\}$, the connection matrix $(\omega_{s,t})$ is defined by

$$\nabla^L(\widetilde{e_1},\cdots,\widetilde{e_n})=(\widetilde{e_1},\cdots,\widetilde{e_n})(\omega_{s,t}). \tag{2.1}$$

Let $\epsilon(\widetilde{e_j*})$, $\iota(\widetilde{e_j*})$ be the exterior and interior multiplications respectively, $\widetilde{e_j*}$ be the dual base of $\widetilde{e_j}$ and $c(\widetilde{e_j})$ be the Clifford action. Suppose that ∂_i is a natural local frame on TM and $(g^{ij})_{1\leq i,j\leq n}$ is the inverse matrix associated to the metric matrix $(g_{ij})_{1\leq i,j\leq n}$ on M. Write

$$c(\widetilde{e_j})=\epsilon(\widetilde{e_j*})-\iota(\widetilde{e_j*});\ \overline{c}(\widetilde{e_j})=\epsilon(\widetilde{e_j*})+\iota(\widetilde{e_j*}), \tag{2.2}$$

which satisfies

$$
\begin{aligned}
&c(\widetilde{e_i})c(\widetilde{e_j})+c(\widetilde{e_j})c(\widetilde{e_i})=-2\delta_i^j;\\
&\overline{c}(\widetilde{e_i})c(\widetilde{e_j})+c(\widetilde{e_j})\overline{c}(\widetilde{e_i})=0;\\
&\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})+\overline{c}(\widetilde{e_j})\overline{c}(\widetilde{e_i})=2\delta_i^j.
\end{aligned}
\tag{2.3}
$$

By [12], we have the signature operator

$$
D=d+\delta=\sum_{i=1}^{n}c(\widetilde{e_i})\left[\widetilde{e_i}+\frac{1}{4}\sum_{s,t}\omega_{s,t}(\widetilde{e_i})[\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t})-c(\widetilde{e_s})c(\widetilde{e_t})]\right].
\tag{2.4}
$$

We define the perturbation of the de Rham Hodge operator D_A as follows:

$$
\begin{aligned}
D_A&=d+\delta+A\\
&=\sum_{i=1}^{n}c(\widetilde{e_i})\left[\widetilde{e_i}+\frac{1}{4}\sum_{s,t}\omega_{s,t}(\widetilde{e_i})[\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t})-c(\widetilde{e_s})c(\widetilde{e_t})]\right]+A,
\end{aligned}
\tag{2.5}
$$

then we have

$$
\begin{aligned}
D_A^*&=d+\delta+A^*\\
&=\sum_{i=1}^{n}c(\widetilde{e_i})\left[\widetilde{e_i}+\frac{1}{4}\sum_{s,t}\omega_{s,t}(\widetilde{e_i})[\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t})-c(\widetilde{e_s})c(\widetilde{e_t})]\right]+A^*,
\end{aligned}
\tag{2.6}
$$

where

$$
A=\overline{c}(X_1)\cdots\overline{c}(X_k)c(X_{k+1})\cdots c(X_{k+t});
\tag{2.7}
$$

$$
A^*=(-1)^t c(X_{k+t})\cdots c(X_{k+1})\overline{c}(X_k)\cdots\overline{c}(X_1),
\tag{2.8}
$$

where $X_1,\cdots,X_{k+t}$ are the smooth vector fields on M.

By some simple calculations, we get Lichnerowicz formulas.

Theorem 2.1. *The following equalities hold:*

$$
\begin{aligned}
D_A^2=&-[g^{ij}(\nabla^1_{\partial_i}\nabla^1_{\partial_j}-\nabla^1_{\nabla^L_{\partial_i}\partial_j})]-\frac{1}{8}\sum_{i,j,k,l}R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l})+\frac{1}{4}s+A^2\\
&+\frac{1}{4}\sum_{i=1}^{n}[c(\widetilde{e_i})A+Ac(\widetilde{e_i})]^2-\frac{1}{2}\sum_{j=1}^{n}[\nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A)c(\widetilde{e_j})-c(\widetilde{e_j})\nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A)],
\end{aligned}
\tag{2.9}
$$

$$
\begin{aligned}
D_A^*D_A=&-[g^{ij}(\nabla^2_{\partial_i}\nabla^2_{\partial_j}-\nabla^2_{\nabla^L_{\partial_i}\partial_j})]-\frac{1}{8}\sum_{i,j,k,l}R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l})+\frac{1}{4}s+A^*A\\
&+\frac{1}{4}\sum_{i=1}^{n}[c(\widetilde{e_i})A+A^*c(\widetilde{e_i})]^2-\frac{1}{2}\sum_{j=1}^{n}[\nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A^*)c(\widetilde{e_j})-c(\widetilde{e_j})\nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A)],
\end{aligned}
\tag{2.10}
$$

where s is the scalar curvature.

Now we prove the Theorem 2.1. Let M be a smooth compact oriented Riemannian n-dimensional manifolds without boundary and N be a vector bundle on M. We say that P is a differential operator of Laplace type, if it has locally the form

$$P = -(g^{ij}\partial_i\partial_j + A^i\partial_i + B), \tag{2.11}$$

where ∂_i is a natural local frame on TM and $(g^{ij})_{1\leq i,j\leq n}$ is the inverse matrix associated to the metric matrix $(g_{ij})_{1\leq i,j\leq n}$ on M, and A^i and B are smooth sections of $\mathrm{End}(N)$ on M (endomorphism). If P satisfies the form (2.11), then there is a unique connection ∇ on N and a unique endomorphism E such that

$$P = -[g^{ij}(\nabla_{\partial_i}\nabla_{\partial_j} - \nabla_{\nabla^L_{\partial_i}\partial_j}) + E], \tag{2.12}$$

where ∇^L is the Levi-Civita connection on M. Moreover (with local frames of T^*M and N), $\nabla_{\partial_i} = \partial_i + \omega_i$ and E are related to g^{ij}, A^i and B through

$$\omega_i = \frac{1}{2}g_{ij}\big(A^i + g^{kl}\Gamma^j_{kl}\mathtt{id}\big), \tag{2.13}$$

$$E = B - g^{ij}\big(\partial_i(\omega_j) + \omega_i\omega_j - \omega_k\Gamma^k_{ij}\big), \tag{2.14}$$

where Γ^j_{kl} is the Christoffel coefficient of ∇^L.

Let $g^{ij} = g(dx_i, dx_j)$, $\xi = \sum_k \xi_j dx_j$ and $\nabla^L_{\partial_i}\partial_j = \sum_k \Gamma^k_{ij}\partial_k$, we denote that

$$\sigma_i = -\frac{1}{4}\sum_{s,t}\omega_{s,t}(\widetilde{e_i})c(\widetilde{e_s})c(\widetilde{e_t});\ a_i = \frac{1}{4}\sum_{s,t}\omega_{s,t}(\widetilde{e_i})\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t}); \tag{2.15}$$

$$\xi^j = g^{ij}\xi_i;\ \Gamma^k = g^{ij}\Gamma^k_{ij};\ \sigma^j = g^{ij}\sigma_i;\ a^j = g^{ij}a_i.$$

Then, the perturbation of the de Rham Hodge operator D_A and D^*_A can be written as

$$D_A = \sum_{i=1}^{n} c(\widetilde{e_i})[\widetilde{e_i} + \sigma_i + a_i] + A; \tag{2.16}$$

$$D^*_A = \sum_{i=1}^{n} c(\widetilde{e_i})[\widetilde{e_i} + \sigma_i + a_i] + A^*. \tag{2.17}$$

By [18], the local expression of $(d+\delta)^2$ is

$$(d+\delta)^2 = -\Delta_0 - \frac{1}{8}\sum_{i,j,k,l} R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l}) + \frac{1}{4}s. \tag{2.18}$$

By [7]and[18], we have

$$-\Delta_0 = -g^{ij}(\nabla^L_i\nabla^L_j - \Gamma^k_{ij}\nabla^L_k). \tag{2.19}$$

We note that

$$D^2_A = (d+\delta)^2 + (d+\delta)A + A(d+\delta) + A^2, \tag{2.20}$$

$$(d+\delta)A + A(d+\delta) = \sum_{i,j} g^{ij}[c(\partial_i)A + Ac(\partial_i)]\partial_j + \sum_{i,j} g^{ij}[c(\partial_i)\partial_j(A) + c(\partial_i)a_jA + c(\partial_i)\sigma_jA + Ac(\partial_i)a_j + Ac(\partial_i)\sigma_j], \tag{2.21}$$

then we obtain

$$\begin{aligned} D_A^2 = &-\sum_{i,j} g^{ij}[\partial_i\partial_j + 2\sigma_i\partial_j + 2a_i\partial_j - \Gamma_{ij}^k\partial_k + (\partial_i\sigma_j) + (\partial_i a_j) + \sigma_i\sigma_j + \sigma_i a_j + a_i\sigma_j \\ &+ a_i a_j - \Gamma_{ij}^k\sigma_k - \Gamma_{ij}^k a_k] + \sum_{i,j} g^{ij}[c(\partial_i)\partial_j(A) + c(\partial_i)a_jA + c(\partial_i)\sigma_jA + Ac(\partial_i)a_j \\ &+ Ac(\partial_i)\sigma_j] + \sum_{i,j} g^{ij}[c(\partial_i)A + Ac(\partial_i)]\partial_j - \frac{1}{8}\sum_{i,j,k,l} R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l}) \\ &+ \frac{1}{4}s + A^2. \end{aligned} \tag{2.22}$$

Similarly, we have

$$\begin{aligned} D_A^*D_A = &-\sum_{i,j} g^{ij}[\partial_i\partial_j + 2\sigma_i\partial_j + 2a_i\partial_j - \Gamma_{ij}^k\partial_k + (\partial_i\sigma_j) + (\partial_i a_j) + \sigma_i\sigma_j + \sigma_i a_j + a_i\sigma_j \\ &+ a_i a_j - \Gamma_{ij}^k\sigma_k - \Gamma_{ij}^k a_k] + \sum_{i,j} g^{ij}[c(\partial_i)\partial_j(A) + c(\partial_i)a_jA + c(\partial_i)\sigma_jA + A^*c(\partial_i)a_j \\ &+ A^*c(\partial_i)\sigma_j] + \sum_{i,j} g^{ij}[c(\partial_i)A + A^*c(\partial_i)]\partial_j - \frac{1}{8}\sum_{i,j,k,l} R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l}) \\ &+ \frac{1}{4}s + A^*A. \end{aligned} \tag{2.23}$$

By (2.13), (2.14) and (2.22), we have

$$(\omega_i)_{D_A^2} = \sigma_i + a_i - \frac{1}{2}[c(\partial_i)A + Ac(\partial_i)], \tag{2.24}$$

(2.25)

$$
\begin{aligned}
E_{D_A^2} &= \frac{1}{8}\sum_{i,j,k,l} R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l}) - \sum_{i=1}^{n} c(\partial_i)\partial^i(A) - \sum_{i=1}^{n} c(\partial_i)a^i A - \sum_{i=1}^{n} c(\partial_i)\sigma^i A \\
&+ \frac{1}{2}\sum_{j=1}^{n}\partial^j[c(\partial_j)A + Ac(\partial_j)] - \frac{1}{2}\sum_{k=1}^{n}\Gamma^k[c(\partial_k)A + Ac(\partial_k)] - \frac{1}{4}s - A^2 \\
&+ \sum_{i,j}\frac{g^{ij}}{2}[c(\partial_i)A + Ac(\partial_i)](\sigma_j + a_j) + \sum_{i,j}\frac{g^{ij}}{2}(\sigma_i + a_i)[c(\partial_j)A + Ac(\partial_j)] \\
&- \sum_{i,j}\frac{g^{ij}}{4}[c(\partial_i)A + Ac(\partial_i)][c(\partial_j)A + Ac(\partial_j)] - \sum_{i=1}^{n} Ac(\partial_i)a^i - \sum_{i=1}^{n} Ac(\partial_i)\sigma^i.
\end{aligned}
$$

Since E is globally defined on M, taking normal coordinates at x_0, we have $\sigma^i(x_0) = 0$, $a^i(x_0) = 0$, $\partial^j[c(\partial_j)](x_0) = 0$, $\Gamma^k(x_0) = 0$, $g^{ij}(x_0) = \delta_i^j$, so that

$$
\begin{aligned}
E_{D_A^2}(x_0) &= \frac{1}{8}\sum_{i,j,k,l} R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l}) - \frac{1}{4}\sum_{i=1}^{n}[c(\partial_i)A + Ac(\partial_i)]^2 \\
&+ \frac{1}{2}\sum_{j=1}^{n}[\partial^j(A)c(\partial_j) - c(\partial_j)\partial^j(A)] - \frac{1}{4}s - A^2 \\
&= \frac{1}{8}\sum_{i,j,k,l} R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l}) - \frac{1}{4}\sum_{i=1}^{n}[c(\widetilde{e_i})A + Ac(\widetilde{e_i})]^2 \\
&+ \frac{1}{2}\sum_{j=1}^{n}[\widetilde{e_j}(A)c(\widetilde{e_j}) - c(\widetilde{e_j})\widetilde{e_j}(A)] - \frac{1}{4}s - A^2 \\
&= \frac{1}{8}\sum_{i,j,k,l} R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l}) - \frac{1}{4}\sum_{i=1}^{n}[c(\widetilde{e_i})A + Ac(\widetilde{e_i})]^2 \\
&+ \frac{1}{2}\sum_{j=1}^{n}[\nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A)c(\widetilde{e_j}) - c(\widetilde{e_j})\nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A)] - \frac{1}{4}s - A^2.
\end{aligned} \tag{2.26}
$$

Similarly, we have

$$
\begin{aligned}
E_{D_A^* D_A}(x_0) &= \frac{1}{8}\sum_{i,j,k,l} R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l}) - \frac{1}{4}\sum_{i=1}^{n}[c(\widetilde{e_i})A + A^*c(\widetilde{e_i})]^2 \\
&+ \frac{1}{2}\sum_{j=1}^{n}[\nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A^*)c(\widetilde{e_j}) - c(\widetilde{e_j})\nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A)] - \frac{1}{4}s - A^*A.
\end{aligned} \tag{2.27}
$$

By (2.12), we get Theorem 2.1.

According to the detailed descriptions in [7], we know that the noncommutative residue

of a generalized laplacian $\widetilde{\Delta}$ is expressed as

$$(n-2)\Phi_2(\widetilde{\Delta}) = (4\pi)^{-\frac{n}{2}}\Gamma(\frac{n}{2})\widetilde{res}(\widetilde{\Delta}^{-\frac{n}{2}+1}), \tag{2.28}$$

where $\Phi_2(\widetilde{\Delta})$ denotes the integral over the diagonal part of the second coefficient of the heat kernel expansion of $\widetilde{\Delta}$. Now let $\widetilde{\Delta} = D_A^2$. Since D_A^2 is a generalized laplacian, we can suppose $D_A^2 = \Delta - E$, then, we have

$$\mathrm{Wres}(D_A^2)^{-\frac{n-2}{2}} = \frac{(n-2)(4\pi)^{\frac{n}{2}}}{(\frac{n}{2}-1)!}\int_M \mathrm{tr}(\frac{1}{6}s + E_{D_A^2})d\mathrm{Vol}_{\mathrm{M}}, \tag{2.29}$$

$$\mathrm{Wres}(D_A^* D_A)^{-\frac{n-2}{2}} = \frac{(n-2)(4\pi)^{\frac{n}{2}}}{(\frac{n}{2}-1)!}\int_M \mathrm{tr}(\frac{1}{6}s + E_{D_A^* D_A})d\mathrm{Vol}_{\mathrm{M}}, \tag{2.30}$$

where Wres denote the noncommutative residue.
By applying the formulae shown in (2.26), (2.27), (2.29) and (2.30), we get:

Theorem 2.2. *If M is a n-dimensional compact oriented manifolds without boundary, we have the following:*

(2.31)

$$\mathrm{Wres}(D_A^2)^{-\frac{n-2}{2}} = \frac{(n-2)(4\pi)^{\frac{n}{2}}}{(\frac{n}{2}-1)!}\int_M \mathrm{tr}\Big(-\frac{1}{12}s + (\frac{n}{2}-1)A^2 - \frac{1}{2}\sum_{j=1}^{n} Ac(\widetilde{e_j})Ac(\widetilde{e_j})\Big)d\mathrm{Vol}_{\mathrm{M}},$$

(2.32)

$$\mathrm{Wres}(D_A^* D_A)^{-\frac{n-2}{2}} = \frac{(n-2)(4\pi)^{\frac{n}{2}}}{(\frac{n}{2}-1)!}\int_M \mathrm{tr}\Big(-\frac{1}{12}s + (\frac{n}{2}-1)A^*A - \frac{1}{4}\sum_{j=1}^{n} Ac(\widetilde{e_j})Ac(\widetilde{e_j})$$
$$-\frac{1}{4}\sum_{j=1}^{n} A^*c(\widetilde{e_j})A^*c(\widetilde{e_j}) + \frac{1}{2}\sum_{j=1}^{n}\nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A^*)c(\widetilde{e_j}) - \frac{1}{2}\sum_{j=1}^{n} c(\widetilde{e_j})\nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A)\Big)d\mathrm{Vol}_{\mathrm{M}},$$

where s is the scalar curvature.

3. A Kastler-Kalau-Walze type theorem for 4-dimensional manifolds with boundary

Firstly, we explain the basic notions of Boutet de Monvel's calculus and the definition of the noncommutative residue for manifolds with boundary that will be used throughout the paper. For the details, see Ref.[12].

Let $U \subset M$ be a collar neighborhood of ∂M which is diffeomorphic with $\partial M \times [0,1)$. By the definition of $h(x_n) \in C^\infty([0,1))$ and $h(x_n) > 0$, there exists $\widehat{h} \in C^\infty((-\varepsilon, 1))$ such that $\widehat{h}|_{[0,1)} = h$ and $\widehat{h} > 0$ for some sufficiently small $\varepsilon > 0$. Then there exists a metric g'

on $\widetilde{M} = M \bigcup_{\partial M} \partial M \times (-\varepsilon, 0]$ which has the form on $U \bigcup_{\partial M} \partial M \times (-\varepsilon, 0]$

$$g' = \frac{1}{\widehat{h}(x_n)} g^{\partial M} + dx_n^2, \tag{3.1}$$

such that $g'|_M = g$. We fix a metric g' on the $\widetilde{M}$ such that $g'|_M = g$.

We define the Fourier transformation F' by

$$F' : L^2(\mathbf{R}_t) \to L^2(\mathbf{R}_v);\ F'(u)(v) = \int e^{-ivt} u(t) dt \tag{3.2}$$

and let

$$r^+ : C^\infty(\mathbf{R}) \to C^\infty(\widetilde{\mathbf{R}^+});\ f \to f|\widetilde{\mathbf{R}^+};\ \widetilde{\mathbf{R}^+} = \{x \geq 0; x \in \mathbf{R}\}. \tag{3.3}$$

where $\Phi(\mathbf{R})$ denotes the Schwartz space and $\Phi(\widetilde{\mathbf{R}^+}) = r^+\Phi(\mathbf{R})$, $\Phi(\widetilde{\mathbf{R}^-}) = r^-\Phi(\mathbf{R})$.

We define $H^+ = F'(\Phi(\widetilde{\mathbf{R}^+}))$; $H_0^- = F'(\Phi(\widetilde{\mathbf{R}^-}))$ which satisfies $H^+ \perp H_0^-$. We have the following property: $h \in H^+$ (H_0^-) if and only if $h \in C^\infty(\mathbf{R})$ which has an analytic extension to the lower (upper) complex half-plane $\{\mathrm{Im}\xi < 0\}$ $(\{\mathrm{Im}\xi > 0\})$ such that for all nonnegative integer l,

$$\frac{d^l h}{d\xi^l}(\xi) \sim \sum_{k=1}^{\infty} \frac{d^l}{d\xi^l}(\frac{c_k}{\xi^k}), \tag{3.4}$$

as $|\xi| \to +\infty, \mathrm{Im}\xi \leq 0$ $(\mathrm{Im}\xi \geq 0)$.

Let H' be the space of all polynomials and $H^- = H_0^- \bigoplus H'$; $H = H^+ \bigoplus H^-$. Denote by π^+ (π^-) respectively the projection on H^+ (H^-). For calculations, we take $H = \widetilde{H} =$ {rational functions having no poles on the real axis} ($\tilde{H}$ is a dense set in the topology of H). Then on $\tilde{H}$,

$$\pi^+ h(\xi_0) = \frac{1}{2\pi i} \lim_{u \to 0^-} \int_{\Gamma^+} \frac{h(\xi)}{\xi_0 + iu - \xi} d\xi, \tag{3.5}$$

where Γ^+ is a Jordan close curve included $\mathrm{Im}(\xi) > 0$ surrounding all the singularities of h in the upper half-plane and $\xi_0 \in \mathbf{R}$. Similarly, define π' on $\tilde{H}$,

$$\pi' h = \frac{1}{2\pi} \int_{\Gamma^+} h(\xi) d\xi. \tag{3.6}$$

So, $\pi'(H^-) = 0$. For $h \in H \bigcap L^1(\mathbf{R})$, $\pi' h = \frac{1}{2\pi} \int_{\mathbf{R}} h(v) dv$ and for $h \in H^+ \bigcap L^1(\mathbf{R})$, $\pi' h = 0$.

Let M be a n-dimensional compact oriented manifold with boundary ∂M. Denote by $\mathcal{B}$ Boutet de Monvel's algebra, we recall the main theorem in [9, 12].

Theorem 3.1. [9]**(Fedosov-Golse-Leichtnam-Schrohe)** *Let X and ∂X be connected,* $\dim X = n \geq 3$, $A = \begin{pmatrix} \pi^+ P + G & K \\ T & S \end{pmatrix} \in \mathcal{B}$ *, and denote by p, b and s the local symbols*

of P, G and S respectively. Define:

$$\widetilde{\mathrm{Wres}}(A) = \int_X \int_{\mathbf{S}} \mathrm{tr}_E\left[p_{-n}(x,\xi)\right]\sigma(\xi)dx + 2\pi \int_{\partial X}\int_{\mathbf{S}'} \left\{\mathrm{tr}_E\left[(\mathrm{tr}b_{-n})(x',\xi')\right] + \mathrm{tr}_F\left[s_{1-n}(x',\xi')\right]\right\}\sigma(\xi')dx', \tag{3.7}$$

Then a) $\widetilde{\mathrm{Wres}}([A,B]) = 0$, for any $A, B \in \mathcal{B}$; b) It is a unique continuous trace on $\mathcal{B}/\mathcal{B}^{-\infty}$.

Definition 3.2. [12] *Lower dimensional volumes of spin manifolds with boundary are defined by*

$$\mathrm{Vol}_n^{(p_1,p_2)}M := \widetilde{\mathrm{Wres}}[\pi^+D^{-p_1}\circ\pi^+D^{-p_2}], \tag{3.8}$$

By [12], we get

$$\widetilde{\mathrm{Wres}}[\pi^+D^{-p_1}\circ\pi^+D^{-p_2}] = \int_M\int_{|\xi|=1}\mathrm{trace}_{\wedge^*T^*M}[\sigma_{-n}(D^{-p_1-p_2})]\sigma(\xi)dx + \int_{\partial M}\Phi \tag{3.9}$$

and

(3.10)

$$\Phi = \int_{|\xi'|=1}\int_{-\infty}^{+\infty}\sum_{j,k=0}^{\infty}\sum\frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!}\times\mathrm{trace}_{\wedge^*T^*M}[\partial_{x_n}^j\partial_{\xi'}^\alpha\partial_{\xi_n}^k\sigma_r^+(D^{-p_1})(x',0,\xi',\xi_n) \times\partial_{x'}^\alpha\partial_{\xi_n}^{j+1}\partial_{x_n}^k\sigma_l(D^{-p_2})(x',0,\xi',\xi_n)]d\xi_n\sigma(\xi')dx',$$

where the sum is taken over $r+l-k-|\alpha|-j-1=-n,\ r\le -p_1, l\le -p_2$.

Since $[\sigma_{-n}(D^{-p_1-p_2})]|_M$ has the same expression as $\sigma_{-n}(D^{-p_1-p_2})$ in the case of manifolds without boundary, so locally we can compute the first term by [5], [6], [12], [19].

For any fixed point $x_0\in\partial M$, we choose the normal coordinates U of x_0 in ∂M (not in M) and compute $\Phi(x_0)$ in the coordinates $\widetilde{U} = U\times[0,1)\subset M$ and the metric $\frac{1}{h(x_n)}g^{\partial M} + dx_n^2$. The dual metric of g^M on $\widetilde{U}$ is $h(x_n)g^{\partial M}+dx_n^2$. Write $g_{ij}^M = g^M(\frac{\partial}{\partial x_i},\frac{\partial}{\partial x_j})$; $g_M^{ij} = g^M(dx_i,dx_j)$, then

$$[g_{i,j}^M] = \begin{bmatrix}\frac{1}{h(x_n)}[g_{i,j}^{\partial M}] & 0\\ 0 & 1\end{bmatrix};\ [g_M^{i,j}] = \begin{bmatrix}h(x_n)[g_{\partial M}^{i,j}] & 0\\ 0 & 1\end{bmatrix} \tag{3.11}$$

and

$$\partial_{x_s}g_{ij}^{\partial M}(x_0) = 0, 1\le i,j\le n-1;\ g_{ij}^M(x_0) = \delta_{ij}. \tag{3.12}$$

We review the following three lemmas.

Lemma 3.3. [12] *With the metric g^M on M near the boundary*

$$\partial_{x_j}(|\xi|^2_{g^M})(x_0) = \begin{cases}0,\ \text{if } j<n,\\ h'(0)|\xi'|^2_{g^{\partial M}},\ \text{if } j=n;\end{cases} \tag{3.13}$$

$$\partial_{x_j}[c(\xi)](x_0) = \begin{cases}0,\ \text{if } j<n,\\ \partial x_n(c(\xi'))(x_0),\ \text{if } j=n,\end{cases} \tag{3.14}$$

where $\xi = \xi' + \xi_n dx_n$.

Lemma 3.4. [12] *With the metric* g^M *on* M *near the boundary*

$$\omega_{s,t}(\widetilde{e_i})(x_0) = \begin{cases} \omega_{n,i}(\widetilde{e_i})(x_0) = \frac{1}{2}h'(0), \text{ if } s = n, t = i, i < n; \\ \omega_{i,n}(\widetilde{e_i})(x_0) = -\frac{1}{2}h'(0), \text{ if } s = i, t = n, i < n; \\ \omega_{s,t}(\widetilde{e_i})(x_0) = 0, \text{ other cases}, \end{cases} \tag{3.15}$$

where $(\omega_{s,t})$ *denotes the connection matrix of Levi-Civita connection* ∇^L.

Lemma 3.5. [12]

$$\Gamma^k_{st}(x_0) = \begin{cases} \Gamma^n_{ii}(x_0) = \frac{1}{2}h'(0), \text{ if } s = t = i, k = n, i < n; \\ \Gamma^i_{ni}(x_0) = -\frac{1}{2}h'(0), \text{ if } s = n, t = i, k = i, i < n; \\ \Gamma^i_{in}(x_0) = -\frac{1}{2}h'(0), \text{ if } s = i, t = n, k = i, i < n, \\ \Gamma^i_{st}(x_0) = 0, \text{ other cases}. \end{cases} \tag{3.16}$$

Similar to (3.9) and (3.10), we firstly compute

(3.17)
$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+(D_A^*)^{-1}] = \int_M \int_{|\xi|=1} \mathrm{trace}_{\wedge^* T^*M}[\sigma_{-4}((D_A^* D_A)^{-1})]\sigma(\xi)dx + \int_{\partial M} \Psi,$$

where

$$\Psi = \int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!} \times \mathrm{trace}_{\wedge^* T^*M}[\partial^j_{x_n} \partial^\alpha_{\xi'} \partial^k_{\xi_n} \sigma_r^+(D_A)^{-1} \tag{3.18}$$
$$(x', 0, \xi', \xi_n) \times \partial^\alpha_{x'} \partial^{j+1}_{\xi_n} \partial^k_{x_n} \sigma_l((D_A^*)^{-1})(x', 0, \xi', \xi_n)] d\xi_n \sigma(\xi') dx',$$

the sum is taken over $r + l - k - j - |\alpha| - 1 = -4, r \leq -1, l \leq -1$.

Then we can compute the interior of $\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+(D_A^*)^{-1}]$,

$$\int_M \int_{|\xi|=1} \mathrm{trace}_{\wedge^* T^*M}[\sigma_{-4}((D_A^* D_A)^{-1})]\sigma(\xi)dx = 32\pi^2 \int_M \mathrm{tr}\Big(-\frac{1}{12}s + A^*A \tag{3.19}$$
$$- \frac{1}{4}\sum_{j=1}^n Ac(\widetilde{e_j})Ac(\widetilde{e_j}) - \frac{1}{4}\sum_{j=1}^n A^*c(\widetilde{e_j})A^*c(\widetilde{e_j}) + \frac{1}{2}\sum_{j=1}^n \nabla^{\wedge^* T^*M}_{\widetilde{e_j}}(A^*)c(\widetilde{e_j})$$
$$- \frac{1}{2}\sum_{j=1}^n c(\widetilde{e_j})\nabla^{\wedge^* T^*M}_{\widetilde{e_j}}(A)\Big) d\mathrm{Vol_M}.$$

Now we need to compute $\int_{\partial M} \Psi$. Since, some operators have the following symbols.

Lemma 3.6. *The following identities hold:*

$$\begin{aligned}(3.20)\qquad \sigma_1(D_A) &= \sigma_1(D_A^*) = ic(\xi);\\ \sigma_0(D_A) &= \frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})c(\widetilde{e_i})\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t}) - \frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})c(\widetilde{e_i})c(\widetilde{e_s})c(\widetilde{e_t}) + A;\\ \sigma_0(D_A^*) &= \frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})c(\widetilde{e_i})\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t}) - \frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})c(\widetilde{e_i})c(\widetilde{e_s})c(\widetilde{e_t}) + A^*.\end{aligned}$$

Write

$$(3.21)\qquad D_x^\alpha \ = (-i)^{|\alpha|}\partial_x^\alpha;\ \sigma(D_A) = p_1 + p_0;\ \sigma({D_A}^{-1}) = \textstyle\sum_{j=1}^\infty q_{-j}.$$

By the composition formula of pseudodifferential operators, we have

$$\begin{aligned}(3.22)\qquad 1 = \sigma(D_A \circ {D_A}^{-1}) &= \sum_\alpha \frac{1}{\alpha!}\partial_\xi^\alpha[\sigma(D_A)]D_x^\alpha[\sigma({D_A}^{-1})]\\ &= (p_1+p_0)(q_{-1}+q_{-2}+q_{-3}+\cdots)\\ &\quad + \sum_j(\partial_{\xi_j}p_1 + \partial_{\xi_j}p_0)(D_{x_j}q_{-1} + D_{x_j}q_{-2} + D_{x_j}q_{-3} + \cdots)\\ &= p_1q_{-1} + (p_1q_{-2} + p_0q_{-1} + \sum_j \partial_{\xi_j}p_1 D_{x_j}q_{-1}) + \cdots,\end{aligned}$$

so

$$(3.23)\qquad q_{-1} = p_1^{-1};\ q_{-2} = -p_1^{-1}[p_0p_1^{-1} + \sum_j \partial_{\xi_j}p_1 D_{x_j}(p_1^{-1})].$$

Lemma 3.7. *The following identities hold:*

$$\begin{aligned}(3.24)\qquad \sigma_{-1}({D_A}^{-1}) = \sigma_{-1}((D_A^*)^{-1}) &= \frac{ic(\xi)}{|\xi|^2};\\ \sigma_{-2}({D_A}^{-1}) &= \frac{c(\xi)\sigma_0(D_A)c(\xi)}{|\xi|^4} + \frac{c(\xi)}{|\xi|^6}\sum_j c(dx_j)\Big[\partial_{x_j}(c(\xi))|\xi|^2 - c(\xi)\partial_{x_j}(|\xi|^2)\Big];\\ \sigma_{-2}((D_A^*)^{-1}) &= \frac{c(\xi)\sigma_0(D_A^*)c(\xi)}{|\xi|^4} + \frac{c(\xi)}{|\xi|^6}\sum_j c(dx_j)\Big[\partial_{x_j}(c(\xi))|\xi|^2 - c(\xi)\partial_{x_j}(|\xi|^2)\Big].\end{aligned}$$

We denote tr as shorthand of trace. When $n = 4$, then $\mathrm{tr}_{\wedge^*T^*M}[\mathtt{id}] = \dim(\wedge^*(4)) = 16$, since the sum is taken over $r + l - k - j - |\alpha| - 1 = -4,\ r \le -1, l \le -1$, then we have the following five cases:

case a) I) $r = -1,\ l = -1,\ k = j = 0,\ |\alpha| = 1$

By applying the formula shown in (3.18), we can calculate

(3.25)
$$\Psi_1 = -\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\sum_{|\alpha|=1}\mathrm{trace}[\partial_{\xi'}^{\alpha}\pi_{\xi_n}^{+}\sigma_{-1}({D_A}^{-1})\times\partial_{x'}^{\alpha}\partial_{\xi_n}\sigma_{-1}((D_A^*)^{-1})](x_0)d\xi_n\sigma(\xi')dx'.$$

case a) II) $r=-1,\ l=-1,\ k=|\alpha|=0,\ j=1$

By (3.18), we get

$$\Psi_2 = -\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial_{x_n}\pi_{\xi_n}^{+}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}^{2}\sigma_{-1}((D_A^*)^{-1})](x_0)d\xi_n\sigma(\xi')dx'. \tag{3.26}$$

case a) III) $r=-1,\ l=-1,\ j=|\alpha|=0,\ k=1$

By (3.18), we calculate that

(3.27)
$$\Psi_3 = -\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial_{\xi_n}\pi_{\xi_n}^{+}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}\partial_{x_n}\sigma_{-1}((D_A^*)^{-1})](x_0)d\xi_n\sigma(\xi')dx'.$$

Similar to the formulae (2.17)-(2.31) in [12], we have

$$\Psi_1+\Psi_2+\Psi_3=0. \tag{3.28}$$

case b) $r=-2,\ l=-1,\ k=j=|\alpha|=0$

Similarly, we get

$$\Psi_4 = -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\pi_{\xi_n}^{+}\sigma_{-2}({D_A}^{-1})\times\partial_{\xi_n}\sigma_{-1}((D_A^*)^{-1})](x_0)d\xi_n\sigma(\xi')dx'. \tag{3.29}$$

We first compute

(3.30)
$$\sigma_{-2}({D_A}^{-1})(x_0)=\frac{c(\xi)\sigma_0(D_A)(x_0)c(\xi)}{|\xi|^4}+\frac{c(\xi)}{|\xi|^6}c(dx_n)\left[\partial_{x_n}[c(\xi')](x_0)|\xi|^2-c(\xi)h'(0)|\xi|^2_{\partial M}\right],$$

where

(3.31)
$$\sigma_0(D_A)(x_0)=\frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})(x_0)c(\widetilde{e_i})\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t})-\frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})(x_0)c(\widetilde{e_i})c(\widetilde{e_s})c(\widetilde{e_t})+A.$$

We denote

$$b_0^1(x_0)=\frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})(x_0)c(\widetilde{e_i})\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t}); \tag{3.32}$$

$$b_0^2(x_0)=-\frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})(x_0)c(\widetilde{e_i})c(\widetilde{e_s})c(\widetilde{e_t}). \tag{3.33}$$

Means that

$$\begin{aligned}\pi^+_{\xi_n}\sigma_{-2}({D_A}^{-1})(x_0)|_{|\xi'|=1}&=\pi^+_{\xi_n}\Big(\frac{c(\xi)b_0^1(x_0)c(\xi)}{(1+\xi_n^2)^2}\Big)+\pi^+_{\xi_n}\Big(\frac{c(\xi)Ac(\xi)}{(1+\xi_n^2)^2}\Big)\\&+\pi^+_{\xi_n}\Big(\frac{c(\xi)b_0^2(x_0)c(\xi)+c(\xi)c(dx_n)\partial_{x_n}[c(\xi')](x_0)}{(1+\xi_n^2)^2}-h'(0)\frac{c(\xi)c(dx_n)c(\xi)}{(1+\xi_n^2)^3}\Big).\end{aligned}$$

Since

$$b_0^1(x_0)c(dx_n)=-\frac{1}{4}h'(0)\sum_{i=1}^{n-1}c(\widetilde{e_i})\overline{c}(\widetilde{e_i})c(\widetilde{e_n})\overline{c}(\widetilde{e_n}), \tag{3.34}$$

then by the relation of the Clifford action and $\mathrm{tr}AB=\mathrm{tr}BA$, we have the equalities:

$$\mathrm{tr}[c(\widetilde{e_i})\overline{c}(\widetilde{e_i})c(\widetilde{e_n})\overline{c}(\widetilde{e_n})]=0(i<n);\ \mathrm{tr}[b_0^1(x_0)c(dx_n)]=0.$$

Therefore

$$\partial_{\xi_n}\sigma_{-1}((D_A^*)^{-1})(x_0)|_{|\xi'|=1}=\partial_{\xi_n}q_{-1}(x_0)|_{|\xi'|=1}=i\left(\frac{c(dx_n)}{1+\xi_n^2}-\frac{2\xi_nc(\xi')+2\xi_n^2c(dx_n)}{(1+\xi_n^2)^2}\right). \tag{3.35}$$

Hence, we have

$$\begin{aligned}&\mathrm{tr}\Big[\pi^+_{\xi_n}\left(\frac{c(\xi)b_0^1(x_0)c(\xi)}{(1+\xi_n^2)^2}\right)\times\partial_{\xi_n}\sigma_{-1}((D_A^*)^{-1})(x_0)\Big]|_{|\xi'|=1}\\&=\frac{1}{2(1+\xi_n^2)^2}\mathrm{tr}[b_0^1(x_0)c(\xi')]+\frac{i}{2(1+\xi_n^2)^2}\mathrm{tr}[b_0^1(x_0)c(dx_n)]\\&=\frac{1}{2(1+\xi_n^2)^2}\mathrm{tr}[b_0^1(x_0)c(\xi')].\end{aligned} \tag{3.36}$$

We note that $i<n$, $\int_{|\xi'|=1}\{\xi_{i_1}\cdots\xi_{i_{2d+1}}\}\sigma(\xi')=0$, so $\mathrm{tr}[c(\xi')c(dx_n)]$ has no contribution for computing case b),

$$-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}\Big[\pi^+_{\xi_n}\Big(\frac{c(\xi)b_0^1(x_0)c(\xi)}{(1+\xi_n^2)^2}\Big)\times\partial_{\xi_n}\sigma_{-1}((D_A^*)^{-1})\Big](x_0)d\xi_n\sigma(\xi')dx'=0. \tag{3.37}$$

Since

$$\begin{aligned}(3.38)\qquad &\mathrm{tr}\Big[\pi^+_{\xi_n}\Big(\frac{c(\xi)Ac(\xi)}{(1+\xi_n^2)^2}\Big)\times\partial_{\xi_n}\sigma_{-1}((D_A^*)^{-1})(x_0)\Big]|_{|\xi'|=1}\\ &=\frac{1}{2(1+\xi_n^2)^2}\mathrm{tr}[Ac(\xi')]+\frac{i}{2(1+\xi_n^2)^2}\mathrm{tr}[Ac(dx_n)],\end{aligned}$$

then, we have

$$\begin{aligned}(3.39)\qquad &-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}\Big[\pi^+_{\xi_n}\Big(\frac{c(\xi)Ac(\xi)}{(1+\xi_n^2)^2}\Big)\times\partial_{\xi_n}\sigma_{-1}((D_A^*)^{-1})\Big](x_0)d\xi_n\sigma(\xi')dx'\\ &=-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{1}{2(1+\xi_n^2)^2}\mathrm{tr}[Ac(\xi')]d\xi_n\sigma(\xi')dx'\\ &-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{i}{2(1+\xi_n^2)^2}\mathrm{tr}[Ac(dx_n)]d\xi_n\sigma(\xi')dx'\\ &=\frac{\Omega_3}{2}\mathrm{tr}[Ac(dx_n)]\int_{\Gamma^+}\frac{1}{(\xi_n-i)^2(\xi_n+i)^2}dx'\\ &=\frac{\Omega_3}{2}\mathrm{tr}[Ac(dx_n)]2\pi i[\frac{1}{(\xi_n+i)^2}]^{(1)}|_{\xi_n=i}dx'\\ &=\frac{\pi}{4}\Omega_3\mathrm{tr}[Ac(dx_n)]dx'.\end{aligned}$$

We have

$$(3.40)\qquad \pi^+_{\xi_n}\Big(\frac{c(\xi)b_0^2(x_0)c(\xi)+c(\xi)c(dx_n)\partial_{x_n}[c(\xi')](x_0)}{(1+\xi_n^2)^2}\Big)-h'(0)\pi^+_{\xi_n}\Big(\frac{c(\xi)c(dx_n)c(\xi)}{(1+\xi_n)^3}\Big):=B_1-B_2,$$

where

$$\begin{aligned}(3.41)\qquad B_1=\frac{-1}{4(\xi_n-i)^2}&[(2+i\xi_n)c(\xi')b_0^2(x_0)c(\xi')+i\xi_nc(dx_n)b_0^2(x_0)c(dx_n)-i\partial_{x_n}c(\xi')\\ &+(2+i\xi_n)c(\xi')c(dx_n)\partial_{x_n}c(\xi')+ic(dx_n)b_0^2(x_0)c(\xi')+ic(\xi')b_0^2(x_0)c(dx_n)]\end{aligned}$$

and

$$(3.42)\qquad B_2=\frac{h'(0)}{2}\left(\frac{c(dx_n)}{4i(\xi_n-i)}+\frac{c(dx_n)-ic(\xi')}{8(\xi_n-i)^2}+\frac{3\xi_n-7i}{8(\xi_n-i)^3}(ic(\xi')-c(dx_n))\right).$$

A simple calculation shows that

$$\begin{aligned}(3.43)\qquad \mathrm{tr}[B_2\times\partial_{\xi_n}\sigma_{-1}((D_A^*)^{-1})(x_0)]|_{|\xi'|=1}&=\frac{i}{2}h'(0)\frac{-i\xi_n^2-\xi_n+4i}{4(\xi_n-i)^3(\xi_n+i)^2}\mathrm{tr}[\mathtt{id}]\\ &=8ih'(0)\frac{-i\xi_n^2-\xi_n+4i}{4(\xi_n-i)^3(\xi_n+i)^2}.\end{aligned}$$

Similarly, we have

$$\text{tr}[B_1 \times \partial_{\xi_n}\sigma_{-1}((D_A^*)^{-1})(x_0)]|_{|\xi'|=1} = \frac{-8ic_0}{(1+\xi_n^2)^2} + 2h'(0)\frac{\xi_n^2 - i\xi_n - 2}{(\xi_n - i)(1+\xi_n^2)^2}, \tag{3.44}$$

where $b_0^2 = c_0 c(dx_n)$ and $c_0 = -\frac{3}{4}h'(0)$.
By (3.44) and (3.45), we have

$$\begin{aligned} &- i\int_{|\xi'|=1}\int_{-\infty}^{+\infty} \text{trace}[(B_1 - B_2) \times \partial_{\xi_n}\sigma_{-1}((D_A^*)^{-1})](x_0)d\xi_n\sigma(\xi')dx' \\ &= -\Omega_3 \int_{\Gamma^+} \frac{8c_0(\xi_n - i) + 4ih'(0)}{(\xi_n - i)^3(\xi_n + i)^2} d\xi_n dx' \\ &= \frac{9}{2}\pi h'(0)\Omega_3 dx'. \end{aligned} \tag{3.45}$$

Hence, we have

$$\Psi_4 = \frac{9}{2}\pi h'(0)\Omega_3 dx' + \frac{\pi}{4}\Omega_3 \text{tr}[Ac(dx_n)]dx'. \tag{3.46}$$

case c) $r = -1,\ l = -2,\ k = j = |\alpha| = 0$

Using (3.18), we get

$$\Psi_5 = -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty} \text{trace}[\pi_{\xi_n}^+\sigma_{-1}({D_A}^{-1}) \times \partial_{\xi_n}\sigma_{-2}((D_A^*)^{-1})](x_0)d\xi_n\sigma(\xi')dx'. \tag{3.47}$$

Considering (3.5) and (3.6), we have

$$\pi_{\xi_n}^+\sigma_{-1}({D_A}^{-1})(x_0)|_{|\xi'|=1} = \frac{c(\xi') + ic(dx_n)}{2(\xi_n - i)}. \tag{3.48}$$

Since

$$\sigma_{-2}((D_A^*)^{-1})(x_0) = \frac{c(\xi)\sigma_0(D_A^*)(x_0)c(\xi)}{|\xi|^4} + \frac{c(\xi)}{|\xi|^6}c(dx_n)\left[\partial_{x_n}[c(\xi')](x_0)|\xi|^2 - c(\xi)h'(0)|\xi|_{\partial_M}^2\right], \tag{3.49}$$

where

$$\begin{aligned} \sigma_0(D_A^*)(x_0) &= \frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})(x_0)c(\widetilde{e_i})\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t}) - \frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})(x_0)c(\widetilde{e_i})c(\widetilde{e_s})c(\widetilde{e_t}) + A^* \\ &= b_0^1(x_0) + b_0^2(x_0) + A^*, \end{aligned} \tag{3.50}$$

then

(3.51)

$$\begin{aligned}\partial_{\xi_n}\sigma_{-2}((D_A^*)^{-1})(x_0)|_{|\xi'|=1} &= \partial_{\xi_n}\Bigg(\frac{c(\xi)(b_0^1(x_0)+b_0^2(x_0)+A^*)c(\xi)}{|\xi|^4}\\ &\quad+\frac{c(\xi)}{|\xi|^6}c(dx_n)\left[\partial_{x_n}[c(\xi')](x_0)|\xi|^2-c(\xi)h'(0)|\xi|^2_{\partial_M}\right]\Bigg)\\ &= \partial_{\xi_n}\Bigg(\frac{c(\xi)b_0^2(x_0)c(\xi)}{|\xi|^4}+\frac{c(\xi)}{|\xi|^6}c(dx_n)\left[\partial_{x_n}[c(\xi')](x_0)|\xi|^2-c(\xi)h'(0)\right]\Bigg)\\ &\quad+\partial_{\xi_n}\left(\frac{c(\xi)b_0^1(x_0)c(\xi)}{|\xi|^4}\right)+\partial_{\xi_n}\left(\frac{c(\xi)A^*c(\xi)}{|\xi|^4}\right).\end{aligned}$$

By computation, we have

(3.52)

$$\partial_{\xi_n}\left(\frac{c(\xi)b_0^1(x_0)c(\xi)}{|\xi|^4}\right)=\frac{c(dx_n)b_0^1(x_0)c(\xi)}{|\xi|^4}+\frac{c(\xi)b_0^1(x_0)c(dx_n)}{|\xi|^4}-\frac{4\xi_n c(\xi)b_0^1(x_0)c(\xi)}{|\xi|^6};$$

$$\partial_{\xi_n}\left(\frac{c(\xi)A^*c(\xi)}{|\xi|^4}\right)=\frac{c(dx_n)A^*c(\xi)}{|\xi|^4}+\frac{c(\xi)A^*c(dx_n)}{|\xi|^4}-\frac{4\xi_n c(\xi)A^*c(\xi)}{|\xi|^6}.\tag{3.53}$$

For the sake of convenience in writing, we denote

$$q_{-2}^1=\frac{c(\xi)b_0^2(x_0)c(\xi)}{|\xi|^4}+\frac{c(\xi)}{|\xi|^6}c(dx_n)\left[\partial_{x_n}[c(\xi')](x_0)|\xi|^2-c(\xi)h'(0)\right],\tag{3.54}$$

then

$$\begin{aligned}\partial_{\xi_n}(q_{-2}^1) &= \frac{1}{(1+\xi_n^2)^3}[(2\xi_n-2\xi_n^3)c(dx_n)b_0^2(x_0)c(dx_n)+(1-3\xi_n^2)c(dx_n)b_0^2(x_0)c(\xi')\\ &\quad+(1-3\xi_n^2)c(\xi')b_0^2(x_0)c(dx_n)-4\xi_n c(\xi')b_0^2(x_0)c(\xi')+(3\xi_n^2-1)\partial_{x_n}c(\xi')\\ &\quad-4\xi_n c(\xi')c(dx_n)\partial_{x_n}c(\xi')+2h'(0)c(\xi')+2h'(0)\xi_n c(dx_n)]\\ &\quad+6\xi_n h'(0)\frac{c(\xi)c(dx_n)c(\xi)}{(1+\xi_n^2)^4}.\end{aligned}\tag{3.55}$$

By (3.48) and (3.52), we have

$$\begin{aligned}&\mathrm{tr}\left[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})(x_0)\times\partial_{\xi_n}\left(\frac{c(\xi)b_0^1(x_0)c(\xi)}{|\xi|^4}\right)\right]|_{|\xi'|=1}\\ &=\frac{-1}{2(\xi_n-i)(\xi_n+i)^3}\mathrm{tr}[b_0^1(x_0)c(\xi')]+\frac{i}{2(\xi_n-i)(\xi_n+i)^3}\mathrm{tr}[b_0^1(x_0)c(dx_n)]\\ &=\frac{-1}{2(\xi_n-i)(\xi_n+i)^3}\mathrm{tr}[b_0^1(x_0)c(\xi')],\end{aligned}\tag{3.56}$$

it is shown that

$$
\begin{aligned}
(3.57)\qquad & -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\operatorname{trace}\left[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}\left(\frac{c(\xi)b_0^1c(\xi)}{|\xi|^4}\right)\right](x_0)d\xi_n\sigma(\xi')dx' \\
&= -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{-1}{2(\xi_n-i)(\xi_n+i)^3}\operatorname{tr}[b_0^1(x_0)c(\xi')]d\xi_n\sigma(\xi')dx' \\
&= 0.
\end{aligned}
$$

Similarly to (3.56), we have

$$
\begin{aligned}
(3.58)\qquad & \operatorname{tr}\left[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})(x_0)\times\partial_{\xi_n}\left(\frac{c(\xi)A^*c(\xi)}{|\xi|^4}\right)\right]|_{|\xi'|=1} \\
&= \frac{-1}{2(\xi_n-i)(\xi_n+i)^3}\operatorname{tr}[A^*c(\xi')]+\frac{i}{2(\xi_n-i)(\xi_n+i)^3}\operatorname{tr}[A^*c(dx_n)],
\end{aligned}
$$

then

$$
\begin{aligned}
(3.59)\qquad & -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\operatorname{trace}\left[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}\left(\frac{c(\xi)A^*c(\xi)}{|\xi|^4}\right)\right](x_0)d\xi_n\sigma(\xi')dx' \\
&= -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{-1}{2(\xi_n-i)(\xi_n+i)^3}\operatorname{tr}[A^*c(\xi')]d\xi_n\sigma(\xi')dx' \\
&-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{i}{2(\xi_n-i)(\xi_n+i)^3}\operatorname{tr}[A^*c(dx_n)]d\xi_n\sigma(\xi')dx' \\
&= \frac{\Omega_3}{2}\operatorname{tr}[A^*c(dx_n)]\int_{\Gamma^+}\frac{1}{(\xi_n-i)(\xi_n+i)^3}d\xi_ndx' \\
&= \frac{\Omega_3}{2}\operatorname{tr}[A^*c(dx_n)]2\pi i[\frac{1}{(\xi_n+i)^3}]^{(1)}|_{\xi_n=i}dx' \\
&= -\frac{\pi}{4}\Omega_3\operatorname{tr}[A^*c(dx_n)]dx'.
\end{aligned}
$$

Observing (3.48) and (3.55), we have

(3.60)

$$
\operatorname{tr}[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}(q_{-2}^1)](x_0)|_{|\xi'|=1}=\frac{12h'(0)(i\xi_n^2+\xi_n-2i)}{(\xi-i)^3(\xi+i)^3}+\frac{48h'(0)i\xi_n}{(\xi-i)^3(\xi+i)^4}.
$$

By $\int_{|\xi'|=1}\{\xi_{i_1}\cdots\xi_{i_{2d+1}}\}\sigma(\xi')=0$ and (3.60), we have

$$
\begin{aligned}
(3.61)\qquad & -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{tr}[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}(q^1_{-2})](x_0)d\xi_n\sigma(\xi')dx'\\
&=-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{12h'(0)(i\xi_n^2+\xi_n-2i)}{(\xi-i)^3(\xi+i)^3}+\frac{48h'(0)i\xi_n}{(\xi-i)^3(\xi+i)^4}d\xi_n\sigma(\xi')dx'\\
&=-i\Omega_3\int_{\Gamma^+}\frac{12h'(0)(i\xi_n^2+\xi_n-2i)}{(\xi-i)^3(\xi+i)^3}+\frac{48h'(0)i\xi_n}{(\xi-i)^3(\xi+i)^4}d\xi_n dx'\\
&=-i\Omega_3\frac{2\pi i}{2!}[\frac{12h'(0)(i\xi_n^2+\xi_n-2i)}{(\xi+i)^3}]^{(2)}|_{\xi_n=i}dx'-i\Omega_3\frac{2\pi i}{2!}[\frac{48h'(0)i\xi_n}{(\xi+i)^4}]^{(2)}|_{\xi_n=i}dx'\\
&=-\frac{9}{2}\pi h'(0)\Omega_3dx'.
\end{aligned}
$$

Then,

$$
(3.62)\qquad \Psi_5=-\frac{9}{2}\pi h'(0)\Omega_3dx'-\frac{\pi}{4}\Omega_3\mathrm{tr}[A^*c(dx_n)]dx'.
$$

In summary,

$$
(3.63)\qquad \Psi=\Psi_1+\Psi_2+\Psi_3+\Psi_4+\Psi_5=\frac{\pi}{4}\Omega_3\mathrm{tr}[Ac(dx_n)]dx'-\frac{\pi}{4}\Omega_3\mathrm{tr}[A^*c(dx_n)]dx'.
$$

Applying (3.17), (3.19) and (3.63), we can assert that:

Theorem 3.8. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, D_A be the perturbation of the de Rham Hodge operator on $\widetilde{M}$, then*

(3.64)

$$
\begin{aligned}
&\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+(D_A^*)^{-1}]=32\pi^2\int_M\mathrm{tr}\Big(-\frac{1}{12}s+A^*A-\frac{1}{4}\sum_{j=1}^n Ac(\widetilde{e_j})Ac(\widetilde{e_j})\\
&-\frac{1}{4}\sum_{j=1}^n A^*c(\widetilde{e_j})A^*c(\widetilde{e_j})+\frac{1}{2}\sum_{j=1}^n\nabla^{\wedge^*T^*M}_{\widetilde{e_j}}(A^*)c(\widetilde{e_j})-\frac{1}{2}\sum_{j=1}^n c(\widetilde{e_j})\nabla^{\wedge^*T^*M}_{\widetilde{e_j}}(A)\Big)d\mathrm{Vol_M}\\
&+\int_{\partial M}\frac{\pi}{4}\Omega_3\mathrm{tr}[Ac(dx_n)]d\mathrm{Vol}_{\partial\mathrm{M}}-\int_{\partial M}\frac{\pi}{4}\Omega_3\mathrm{tr}[A^*c(dx_n)]d\mathrm{Vol}_{\partial\mathrm{M}},
\end{aligned}
$$

where s is the scalar curvature.

When $A=c(X)$, then we have

$$
\begin{aligned}
(3.65)\qquad &\int_M\int_{|\xi|=1}\mathrm{trace}_{\wedge^*T^*M}[\sigma_{-4}((D_A^*D_A)^{-1})]\sigma(\xi)dx\\
&=512\pi^2\int_M\Big(-\frac{1}{12}s+2|X|^2+\sum_{j=1}^n g(\nabla^{TM}_{\widetilde{e_j}}X,\widetilde{e_j})\Big)d\mathrm{Vol_M}
\end{aligned}
$$

and

$$\int_{\partial M}\Psi=-\int_{\partial M}8\pi\Omega_3 g(\partial x_n,X)d\mathrm{Vol}_{\partial \mathrm{M}}. \tag{3.66}$$

We can immediately state the following corollary:

Corollary 3.9. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=c(X)$, then*

$$\begin{aligned}\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+(D_A^*)^{-1}]&=512\pi^2\int_M\Big(-\frac{1}{12}s+2|X|^2+\sum_{j=1}^n g(\nabla^{TM}_{\widetilde{e_j}}X,\widetilde{e_j})\Big)d\mathrm{Vol}_{\mathrm{M}}\\&-\int_{\partial M}8\pi\Omega_3 g(\partial x_n,X)d\mathrm{Vol}_{\partial \mathrm{M}},\end{aligned} \tag{3.67}$$

where s is the scalar curvature.

When $A=\overline{c}(X)$, we can get

$$\int_M\int_{|\xi|=1}\mathrm{trace}_{\wedge^*T^*M}[\sigma_{-4}((D_A^*D_A)^{-1})]\sigma(\xi)dx=512\pi^2\int_M\Big(-\frac{1}{12}s-|X|^2\Big)d\mathrm{Vol}_{\mathrm{M}}, \tag{3.68}$$

and

$$\int_{\partial M}\Psi=0. \tag{3.69}$$

Now, we compute $\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+(D_A^*)^{-1}]$.

Corollary 3.10. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=\overline{c}(X)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+(D_A^*)^{-1}]=512\pi^2\int_M\Big(-\frac{1}{12}s-|X|^2\Big)d\mathrm{Vol}_{\mathrm{M}}, \tag{3.70}$$

where s is the scalar curvature.

When $A=c(X)c(Y)$, then $A^*=c(Y)c(X)$.
By computation, we have

$$\begin{aligned}&\mathrm{tr}[A^*A]=\mathrm{tr}[(-|X|^2)(-|Y|^2)]=|X|^2|Y|^2\mathrm{tr}[\mathtt{id}],\\&\mathrm{tr}[c(dx_n)A]=0,\mathrm{tr}[c(dx_n)A^*]=0,\end{aligned} \tag{3.71}$$

$$\sum_{j=1}^{n}\mathrm{tr}[Ac(\widetilde{e_j})Ac(\widetilde{e_j})]=\sum_{j=1}^{n}\mathrm{tr}[c(X)c(Y)c(\widetilde{e_j})c(X)c(Y)c(\widetilde{e_j})] \tag{3.72}$$

$$=-\sum_{j=1}^{n}\mathrm{tr}[c(X)c(Y)c(X)c(\widetilde{e_j})c(Y)c(\widetilde{e_j})]-2\sum_{j=1}^{n}g(\widetilde{e_j},X)\mathrm{tr}[c(X)c(Y)c(Y)c(\widetilde{e_j})]$$

$$=-\sum_{j=1}^{n}\mathrm{tr}[c(X)c(Y)c(X)c(\widetilde{e_j})c(Y)c(\widetilde{e_j})]-2|X|^2|Y|^2\mathrm{tr}[\mathtt{id}]$$

$$=(n-4)|X|^2|Y|^2\mathrm{tr}[\mathtt{id}]+(4-2n)g(X,Y)^2\mathrm{tr}[\mathtt{id}]$$

$$\sum_{j=1}^{n}\mathrm{tr}[A^*c(\widetilde{e_j})A^*c(\widetilde{e_j})]=\sum_{j=1}^{n}\mathrm{tr}[c(Y)c(X)c(\widetilde{e_j})c(Y)c(X)c(\widetilde{e_j})] \tag{3.73}$$

$$=-\sum_{j=1}^{n}\mathrm{tr}[c(Y)c(X)c(Y)c(\widetilde{e_j})c(X)c(\widetilde{e_j})]-2\sum_{j=1}^{n}g(\widetilde{e_j},Y)\mathrm{tr}[c(Y)c(X)c(X)c(\widetilde{e_j})]$$

$$=-\sum_{j=1}^{n}\mathrm{tr}[c(Y)c(X)c(Y)c(\widetilde{e_j})c(X)c(\widetilde{e_j})]-2|X|^2|Y|^2\mathrm{tr}[\mathtt{id}]$$

$$=(n-4)|X|^2|Y|^2\mathrm{tr}[\mathtt{id}]+(4-2n)g(X,Y)^2\mathrm{tr}[\mathtt{id}]$$

$$\sum_{j=1}^{n}\mathrm{tr}[\nabla^{\bigwedge^*T^*M}_{\widetilde{e_j}}(A^*)c(\widetilde{e_j})]=\sum_{j=1}^{n}\mathrm{tr}[\nabla^{\bigwedge^*T^*M}_{\widetilde{e_j}}\big(c(Y)c(X)\big)c(\widetilde{e_j})] \tag{3.74}$$

$$=\sum_{j=1}^{n}\mathrm{tr}[\nabla^{\bigwedge^*T^*M}_{\widetilde{e_j}}\big(c(Y)\big)c(X)c(\widetilde{e_j})+c(Y)\nabla^{\bigwedge^*T^*M}_{\widetilde{e_j}}\big(c(X)\big)c(\widetilde{e_j})]$$

$$=\sum_{j=1}^{n}\mathrm{tr}[c(\nabla^{TM}_{\widetilde{e_j}}Y)c(X)c(\widetilde{e_j})+c(Y)c(\nabla^{TM}_{\widetilde{e_j}}X)c(\widetilde{e_j})]=0,$$

$$\sum_{j=1}^{n}\mathrm{tr}[c(\widetilde{e_j})\nabla^{\bigwedge^*T^*M}_{\widetilde{e_j}}(A)]=\sum_{j=1}^{n}\mathrm{tr}[c(\widetilde{e_j})\nabla^{\bigwedge^*T^*M}_{\widetilde{e_j}}\big(c(X)c(Y)\big)] \tag{3.75}$$

$$=\sum_{j=1}^{n}\mathrm{tr}[c(\widetilde{e_j})\nabla^{\bigwedge^*T^*M}_{\widetilde{e_j}}\big(c(X)\big)c(Y)+c(\widetilde{e_j})c(X)\nabla^{\bigwedge^*T^*M}_{\widetilde{e_j}}\big(c(Y)\big)]$$

$$=\sum_{j=1}^{n}\mathrm{tr}[c(\widetilde{e_j})c(\nabla^{TM}_{\widetilde{e_j}}X)c(Y)+c(\widetilde{e_j})c(X)c(\nabla^{TM}_{\widetilde{e_j}}Y)]=0.$$

By applying the formulae shown in (3.19), we can calculate

$$
\begin{aligned}
(3.76)\quad & \int_M \int_{|\xi|=1} \mathrm{trace}_{\wedge^* T^*M}[\sigma_{-4}((D_A^* D_A)^{-1})]\sigma(\xi)dx \\
&= 32\pi^2 \int_M \mathrm{tr}\Big(-\frac{1}{12}s + A^*A - \frac{1}{4}\sum_{j=1}^{n} Ac(\widetilde{e_j})Ac(\widetilde{e_j}) - \frac{1}{4}\sum_{j=1}^{n} A^*c(\widetilde{e_j})A^*c(\widetilde{e_j}) \\
&+ \frac{1}{2}\sum_{j=1}^{n} \nabla^{\wedge^* T^*M}_{\widetilde{e_j}}(A^*)c(\widetilde{e_j}) - \frac{1}{2}\sum_{j=1}^{n} c(\widetilde{e_j})\nabla^{\wedge^* T^*M}_{\widetilde{e_j}}(A)\Big) d\mathrm{Vol_M} \\
&= 32\pi^2 \int_M \Big(-\frac{1}{12}s + |X|^2|Y|^2 - \frac{1}{4}(-4)g(X,Y)^2 - \frac{1}{4}(-4)g(X,Y)^2\Big)\mathrm{tr}[\mathtt{id}]d\mathrm{Vol_M} \\
&= 512\pi^2 \int_M \Big(-\frac{1}{12}s + |X|^2|Y|^2 + 2g(X,Y)^2\Big) d\mathrm{Vol_M},
\end{aligned}
$$

and

$$
(3.77)\qquad \int_{\partial M} \Psi = 0.
$$

We can claim the following corollary:

Corollary 3.11. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)c(Y)$, then*

$$
(3.78)\quad \widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+ (D_A^*)^{-1}] = 512\pi^2 \int_M \Big(-\frac{1}{12}s + |X|^2|Y|^2 + 2g(X,Y)^2\Big) d\mathrm{Vol_M},
$$

where s is the scalar curvature.

When $A = c(X)\overline{c}(Y)$, similar to (3.78), we can get:

Corollary 3.12. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)\overline{c}(Y)$, then*

$$
(3.79)\qquad \widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+ (D_A^*)^{-1}] = 512\pi^2 \int_M \Big(-\frac{1}{12}s + 2|X|^2|Y|^2\Big) d\mathrm{Vol_M},
$$

where s is the scalar curvature.

When $A = \overline{c}(X)\overline{c}(Y)$, similar to (3.78), we can get the following corollary:

Corollary 3.13. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)\overline{c}(Y)$, then*

$$
(3.80)\quad \widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+ (D_A^*)^{-1}] = 512\pi^2 \int_M \Big(-\frac{1}{12}s - |X|^2|Y|^2 + 4g(X,Y)^2\Big) d\mathrm{Vol_M},
$$

where s is the scalar curvature.

When $A = c(X)c(Y)c(Z)$, then $A^* = -c(Z)c(Y)c(X)$.
By computation, we have

(3.81)

$$\mathrm{tr}[A^*A] = -\mathrm{tr}[(-|X|^2)(-|Y|^2)(-|Z|^2)] = |X|^2|Y|^2|Z|^2\mathrm{tr}[\mathtt{id}],$$

(3.82)

$$\sum_{j=1}^{n}\mathrm{tr}[Ac(\widetilde{e}_j)Ac(\widetilde{e}_j)] = \sum_{j=1}^{n}\mathrm{tr}[c(X)c(Y)c(Z)c(\widetilde{e}_j)c(X)c(Y)c(Z)c(\widetilde{e}_j)]$$
$$= (n-6)|X|^2|Y|^2|Z|^2\mathrm{tr}[\mathtt{id}] + (8-2n)|X|^2g(Y,Z)^2\mathrm{tr}[\mathtt{id}] + (8-2n)|Y|^2g(X,Z)^2\mathrm{tr}[\mathtt{id}]$$
$$+ (8-2n)|Z|^2g(X,Y)^2\mathrm{tr}[\mathtt{id}] + (4n-16)g(X,Y)g(X,Z)g(Y,Z)\mathrm{tr}[\mathtt{id}]$$

(3.83)

$$\sum_{j=1}^{n}\mathrm{tr}[A^*c(\widetilde{e}_j)A^*c(\widetilde{e}_j)] = \sum_{j=1}^{n}\mathrm{tr}[(-c(Z)c(Y)c(X))c(\widetilde{e}_j)(-c(Z)c(Y)c(X))c(\widetilde{e}_j)]$$
$$= (n-6)|X|^2|Y|^2|Z|^2\mathrm{tr}[\mathtt{id}] + (8-2n)|X|^2g(Y,Z)^2\mathrm{tr}[\mathtt{id}] + (8-2n)|Y|^2g(X,Z)^2\mathrm{tr}[\mathtt{id}]$$
$$+ (8-2n)|Z|^2g(X,Y)^2\mathrm{tr}[\mathtt{id}] + (4n-16)g(X,Y)g(X,Z)g(Y,Z)\mathrm{tr}[\mathtt{id}]$$

(3.84)

$$\sum_{j=1}^{n}\mathrm{tr}[\nabla^{\bigwedge^*T^*M}_{\widetilde{e}_j}(A^*)c(\widetilde{e}_j)] = \sum_{j=1}^{n}\mathrm{tr}[\nabla^{\bigwedge^*T^*M}_{\widetilde{e}_j}\big(-c(Z)c(Y)c(X)\big)c(\widetilde{e}_j)]$$
$$= \big[-\sum_{j=1}^{n}g(\nabla^{TM}_{\widetilde{e}_j}Z,\widetilde{e}_j)g(X,Y) + \sum_{j=1}^{n}g(Y,\widetilde{e}_j)g(X,\nabla^{TM}_{\widetilde{e}_j}Z) - \sum_{j=1}^{n}g(X,\widetilde{e}_j)g(Y,\nabla^{TM}_{\widetilde{e}_j}Z)$$
$$- \sum_{j=1}^{n}g(Z,\widetilde{e}_j)g(X,\nabla^{TM}_{\widetilde{e}_j}Y) + \sum_{j=1}^{n}g(\nabla^{TM}_{\widetilde{e}_j}Y,\widetilde{e}_j)g(X,Z) - \sum_{j=1}^{n}g(X,\widetilde{e}_j)g(\nabla^{TM}_{\widetilde{e}_j}Y,Z)$$
$$- \sum_{j=1}^{n}g(Z,\widetilde{e}_j)g(\nabla^{TM}_{\widetilde{e}_j}X,Y) + \sum_{j=1}^{n}g(Y,\widetilde{e}_j)g(\nabla^{TM}_{\widetilde{e}_j}X,Z) - \sum_{j=1}^{n}g(\nabla^{TM}_{\widetilde{e}_j}X,\widetilde{e}_j)g(Y,Z)\big]\mathrm{tr}[\mathtt{id}],$$

(3.85)

$$\sum_{j=1}^{n}\mathrm{tr}[c(\widetilde{e}_j)\nabla^{\bigwedge^*T^*M}_{\widetilde{e}_j}(A)] = \sum_{j=1}^{n}\mathrm{tr}[c(\widetilde{e}_j)\nabla^{\bigwedge^*T^*M}_{\widetilde{e}_j}\big(c(X)c(Y)c(Z)\big)]$$
$$= \big[\sum_{j=1}^{n}g(\widetilde{e}_j,Z)g(\nabla^{TM}_{\widetilde{e}_j}X,Y) - \sum_{j=1}^{n}g(\nabla^{TM}_{\widetilde{e}_j}X,Z)g(\widetilde{e}_j,Y) + \sum_{j=1}^{n}g(Y,Z)g(\widetilde{e}_j,\nabla^{TM}_{\widetilde{e}_j}X)$$
$$+ \sum_{j=1}^{n}g(\widetilde{e}_j,Z)g(X,\nabla^{TM}_{\widetilde{e}_j}Y) - \sum_{j=1}^{n}g(X,Z)g(\widetilde{e}_j,\nabla^{TM}_{\widetilde{e}_j}Y) + \sum_{j=1}^{n}g(\nabla^{TM}_{\widetilde{e}_j}Y,Z)g(\widetilde{e}_j,X)$$
$$+ \sum_{j=1}^{n}g(\widetilde{e}_j,\nabla^{TM}_{\widetilde{e}_j}Z)g(X,Y) - \sum_{j=1}^{n}g(X,\nabla^{TM}_{\widetilde{e}_j}Z)g(\widetilde{e}_j,Y) + \sum_{j=1}^{n}g(Y,\nabla^{TM}_{\widetilde{e}_j}Z)g(\widetilde{e}_j,X)\big]\mathrm{tr}[\mathtt{id}],$$

(3.86)
$$\mathrm{tr}[c(dx_n)A^*] = -g(\partial x_n, X)g(Y,Z)\mathrm{tr}[\mathtt{id}] + g(\partial x_n, Y)g(X,Z)\mathrm{tr}[\mathtt{id}] - g(\partial x_n, Z)g(X,Y)\mathrm{tr}[\mathtt{id}],$$
(3.87)
$$\mathrm{tr}[c(dx_n)A] = g(\partial x_n, X)g(Y,Z)\mathrm{tr}[\mathtt{id}] - g(\partial x_n, Y)g(X,Z)\mathrm{tr}[\mathtt{id}] + g(\partial x_n, Z)g(X,Y)\mathrm{tr}[\mathtt{id}],$$
then we have
$$\int_M \int_{|\xi|=1} \mathrm{trace}_{\wedge^* T^*M}[\sigma_{-4}((D_A^* D_A)^{-1})]\sigma(\xi)dx = 512\pi^2 \int_M \Big(-\frac{1}{12}s + 2|X|^2|Y|^2|Z|^2$$
$$- \sum_{j=1}^4 g(\nabla^{TM}_{\widetilde{e_j}} Z, \widetilde{e_j})g(X,Y) + \sum_{j=1}^4 g(Y, \widetilde{e_j})g(X, \nabla^{TM}_{\widetilde{e_j}} Z) - \sum_{j=1}^4 g(X, \widetilde{e_j})g(Y, \nabla^{TM}_{\widetilde{e_j}} Z)$$
$$- \sum_{j=1}^4 g(Z, \widetilde{e_j})g(X, \nabla^{TM}_{\widetilde{e_j}} Y) + \sum_{j=1}^4 g(\nabla^{TM}_{\widetilde{e_j}} Y, \widetilde{e_j})g(X,Z) - \sum_{j=1}^4 g(X, \widetilde{e_j})g(\nabla^{TM}_{\widetilde{e_j}} Y, Z)$$
$$- \sum_{j=1}^4 g(Z, \widetilde{e_j})g(\nabla^{TM}_{\widetilde{e_j}} X, Y) + \sum_{j=1}^4 g(Y, \widetilde{e_j})g(\nabla^{TM}_{\widetilde{e_j}} X, Z) - \sum_{j=1}^4 g(\nabla^{TM}_{\widetilde{e_j}} X, \widetilde{e_j})g(Y,Z)\Big) d\mathrm{Vol_M}$$
and
$$\int_{\partial M} \Psi = \int_{\partial M} 8\pi\Omega_3 g(\partial x_n, X)g(Y,Z)d\mathrm{Vol}_{\partial \mathrm{M}} - \int_{\partial M} 8\pi\Omega_3 g(\partial x_n, Y)g(X,Z)d\mathrm{Vol}_{\partial \mathrm{M}}$$
$$+ \int_{\partial M} 8\pi\Omega_3 g(\partial x_n, Z)g(X,Y)d\mathrm{Vol}_{\partial \mathrm{M}}. \tag{3.88}$$

Corollary 3.14. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)c(Y)c(Z)$, then*

(3.89)
$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+ (D_A^*)^{-1}] = 512\pi^2 \int_M \Big(-\frac{1}{12}s + 2|X|^2|Y|^2|Z|^2$$
$$- \sum_{j=1}^4 g(\nabla^{TM}_{\widetilde{e_j}} Z, \widetilde{e_j})g(X,Y) + \sum_{j=1}^4 g(Y, \widetilde{e_j})g(X, \nabla^{TM}_{\widetilde{e_j}} Z) - \sum_{j=1}^4 g(X, \widetilde{e_j})g(Y, \nabla^{TM}_{\widetilde{e_j}} Z)$$
$$- \sum_{j=1}^4 g(Z, \widetilde{e_j})g(X, \nabla^{TM}_{\widetilde{e_j}} Y) + \sum_{j=1}^4 g(\nabla^{TM}_{\widetilde{e_j}} Y, \widetilde{e_j})g(X,Z) - \sum_{j=1}^4 g(X, \widetilde{e_j})g(\nabla^{TM}_{\widetilde{e_j}} Y, Z)$$
$$- \sum_{j=1}^4 g(Z, \widetilde{e_j})g(\nabla^{TM}_{\widetilde{e_j}} X, Y) + \sum_{j=1}^4 g(Y, \widetilde{e_j})g(\nabla^{TM}_{\widetilde{e_j}} X, Z) - \sum_{j=1}^4 g(\nabla^{TM}_{\widetilde{e_j}} X, \widetilde{e_j})g(Y,Z)\Big) d\mathrm{Vol_M}$$
$$+ \int_{\partial M} 8\pi\Omega_3 g(\partial x_n, X)g(Y,Z)d\mathrm{Vol}_{\partial \mathrm{M}} - \int_{\partial M} 8\pi\Omega_3 g(\partial x_n, Y)g(X,Z)d\mathrm{Vol}_{\partial \mathrm{M}}$$
$$+ \int_{\partial M} 8\pi\Omega_3 g(\partial x_n, Z)g(X,Y)d\mathrm{Vol}_{\partial \mathrm{M}},$$

where s is the scalar curvature.

When $A=\overline{c}(X)c(Y)c(Z)$, similar to Corollary 3.14, we have:

Corollary 3.15. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=\overline{c}(X)c(Y)c(Z)$, then*

(3.90)
$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+(D_A^*)^{-1}]=512\pi^2\int_M\Big(-\frac{1}{12}s+|X|^2|Y|^2|Z|^2-2|X|^2g(Y,Z)^2\Big)d\mathrm{Vol_M},$$

where s is the scalar curvature.

When $A=\overline{c}(X)\overline{c}(Y)c(Z)$, we can get the following corollary:

Corollary 3.16. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=\overline{c}(X)\overline{c}(Y)c(Z)$, then*

(3.91)
$$\begin{aligned}\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+(D_A^*)^{-1}]&=512\pi^2\int_M\Big(-\frac{1}{12}s+2|Z|^2g(X,Y)^2+\sum_{j=1}^4g(\nabla^{TM}_{\widetilde{e_j}}X,Y)g(Z,\widetilde{e_j})\\&+\sum_{j=1}^4g(X,\nabla^{TM}_{\widetilde{e_j}}Y)g(Z,\widetilde{e_j})+\sum_{j=1}^4g(X,Y)g(\nabla^{TM}_{\widetilde{e_j}}Z,\widetilde{e_j})\Big)d\mathrm{Vol_M}\\&-\int_{\partial M}8\pi\Omega_3g(\partial x_n,Z)g(X,Y)d\mathrm{Vol}_{\partial\mathrm{M}},\end{aligned}$$

where s is the scalar curvature.

When $A=\overline{c}(X)\overline{c}(Y)\overline{c}(Z)$, we get:

Corollary 3.17. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=\overline{c}(X)\overline{c}(Y)\overline{c}(Z)$, then*

(3.92)
$$\begin{aligned}\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+(D_A^*)^{-1}]&=512\pi^2\int_M\Big(-\frac{1}{12}s+3|X|^2|Y|^2|Z|^2-4|X|^2g(Y,Z)^2\\&-4|Y|^2g(X,Z)^2-4|Z|^2g(X,Y)^2+8g(X,Y)g(X,Z)g(Y,Z)\Big)d\mathrm{Vol_M},\end{aligned}$$

where s is the scalar curvature.

Next, we also prove the Kastler-Kalau-Walze type theorem for 4-dimensional manifolds with boundary associated to ${D_A}^2$. By (3.9) and (3.10), we will compute

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+{D_A}^{-1}]=\int_M\int_{|\xi|=1}\mathrm{trace}_{\wedge^*T^*M}[\sigma_{-4}({D_A}^{-2})]\sigma(\xi)dx+\int_{\partial M}\Phi,\tag{3.93}$$

where

(3.94)

$$\Phi=\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\sum_{j,k=0}^{\infty}\sum\frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!}\times \mathrm{trace}_{\wedge^*T^*M}[\partial_{x_n}^j\partial_{\xi'}^{\alpha}\partial_{\xi_n}^k\sigma_r^+({D_A}^{-1})(x',0,\xi',\xi_n)$$
$$\times\partial_{x'}^{\alpha}\partial_{\xi_n}^{j+1}\partial_{x_n}^k\sigma_l({D_A}^{-1})(x',0,\xi',\xi_n)]d\xi_n\sigma(\xi')dx',$$

and the sum is taken over $r+l-k-j-|\alpha|=-3,\ r\leq -1, l\leq -1$.

By Theorem 2.2, we compute the interior of $\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+{D_A}^{-1}]$, then

$$\begin{aligned}(3.95)\qquad &\int_M\int_{|\xi|=1}\mathrm{trace}_{\wedge^*T^*M}[\sigma_{-4}({D_A}^{-2})]\sigma(\xi)dx\\ &=32\pi^2\int_M\mathrm{tr}\bigg(-\frac{1}{12}s+A^2-\frac{1}{2}\sum_{j=1}^{n}Ac(\widetilde{e_j})Ac(\widetilde{e_j})\bigg)d\mathrm{Vol_M}.\end{aligned}$$

When $n=4$, then $\mathrm{tr}_{\wedge^*T^*M}[\mathtt{id}]=\dim(\wedge^*(4))=16$, where tr as shorthand of trace, the sum is taken over $r+l-k-j-|\alpha|=-3,\ r\leq -1, l\leq -1$, then we have the following five cases:

case a) I) $r=-1,\ l=-1,\ k=j=0,\ |\alpha|=1$

By (3.94), we get

$$(3.96)\quad \Phi_1=-\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\sum_{|\alpha|=1}\mathrm{tr}[\partial_{\xi'}^{\alpha}\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{x'}^{\alpha}\partial_{\xi_n}\sigma_{-1}({D_A}^{-1})](x_0)d\xi_n\sigma(\xi')dx'.$$

case a) II) $r=-1,\ l=-1,\ k=|\alpha|=0,\ j=1$

Likewise, we get

$$(3.97)\quad \Phi_2=-\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial_{x_n}\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial^2_{\xi_n}\sigma_{-1}({D_A}^{-1})](x_0)d\xi_n\sigma(\xi')dx'.$$

case a) III) $r=-1,\ l=-1,\ j=|\alpha|=0,\ k=1$

Observing (3.94), we get

$$(3.98)\quad \Phi_3=-\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial_{\xi_n}\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}\partial_{x_n}\sigma_{-1}({D_A}^{-1})](x_0)d\xi_n\sigma(\xi')dx'.$$

By Lemma 3.7, we have $\sigma_{-1}({D_A}^{-1})=\sigma_{-1}((D_A^*)^{-1})$.
In combination with the calculation,

$$(3.99)\qquad \Phi_1+\Phi_2+\Phi_3=0.$$

case b) $r=-2,\ l=-1,\ k=j=|\alpha|=0$

By applying the formulae shown in (3.94), we get

$$\Phi_4=-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\pi^+_{\xi_n}\sigma_{-2}({D_A}^{-1})\times\partial_{\xi_n}\sigma_{-1}({D_A}^{-1})](x_0)d\xi_n\sigma(\xi')dx'. \tag{3.100}$$

By Lemma 3.7, we have $\sigma_{-1}({D_A}^{-1})=\sigma_{-1}((D_A^*)^{-1})$. Then, we have

$$\Phi_4=\frac{9}{2}\pi h'(0)\Omega_3dx'+\frac{\pi}{4}\Omega_3\mathrm{tr}[Ac(dx_n)]dx', \tag{3.101}$$

where Ω_4 is the canonical volume of S^4.

case c) $r=-1,\ l=-2,\ k=j=|\alpha|=0$

By (3.94), we can calculate

$$\Phi_5=-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}\sigma_{-2}({D_A}^{-1})](x_0)d\xi_n\sigma(\xi')dx'. \tag{3.102}$$

By (3.5) and (3.6), we have

$$\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})(x_0)|_{|\xi'|=1}=\frac{c(\xi')+ic(dx_n)}{2(\xi_n-i)}. \tag{3.103}$$

Since

$$\sigma_{-2}({D_A}^{-1})(x_0)=\frac{c(\xi)\sigma_0(D_A)(x_0)c(\xi)}{|\xi|^4}+\frac{c(\xi)}{|\xi|^6}c(dx_n)\left[\partial_{x_n}[c(\xi')](x_0)|\xi|^2-c(\xi)h'(0)|\xi|^2_{\partial_M}\right], \tag{3.104}$$

where

$$\begin{aligned}\sigma_0(D_A)(x_0)&=\frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})(x_0)c(\widetilde{e_i})\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t})-\frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})(x_0)c(\widetilde{e_i})c(\widetilde{e_s})c(\widetilde{e_t})+A\\&=b_0^1(x_0)+b_0^2(x_0)+A,\end{aligned} \tag{3.105}$$

hence

(3.106)

$$\begin{aligned}\partial_{\xi_n}\sigma_{-2}({D_A}^{-1})(x_0)|_{|\xi'|=1} &= \partial_{\xi_n}\bigg(\frac{c(\xi)(b_0^1(x_0)+b_0^2(x_0)+A)c(\xi)}{|\xi|^4}\\ &\quad+\frac{c(\xi)}{|\xi|^6}c(dx_n)\left[\partial_{x_n}[c(\xi')](x_0)|\xi|^2-c(\xi)h'(0)|\xi|^2_{\partial_M}\right]\bigg)\\ &= \partial_{\xi_n}\bigg(\frac{c(\xi)b_0^2(x_0)c(\xi)}{|\xi|^4}+\frac{c(\xi)}{|\xi|^6}c(dx_n)\left[\partial_{x_n}[c(\xi')](x_0)|\xi|^2-c(\xi)h'(0)\right]\bigg)\\ &\quad+\partial_{\xi_n}\bigg(\frac{c(\xi)b_0^1(x_0)c(\xi)}{|\xi|^4}\bigg)+\partial_{\xi_n}\bigg(\frac{c(\xi)Ac(\xi)}{|\xi|^4}\bigg).\end{aligned}$$

By calculation, we have

(3.107)

$$\partial_{\xi_n}\bigg(\frac{c(\xi)b_0^1(x_0)c(\xi)}{|\xi|^4}\bigg)=\frac{c(dx_n)b_0^1(x_0)c(\xi)}{|\xi|^4}+\frac{c(\xi)b_0^1(x_0)c(dx_n)}{|\xi|^4}-\frac{4\xi_n c(\xi)b_0^1(x_0)c(\xi)}{|\xi|^6};$$

(3.108)

$$\partial_{\xi_n}\bigg(\frac{c(\xi)Ac(\xi)}{|\xi|^4}\bigg)=\frac{c(dx_n)Ac(\xi)}{|\xi|^4}+\frac{c(\xi)Ac(dx_n)}{|\xi|^4}-\frac{4\xi_n c(\xi)Ac(\xi)}{|\xi|^6}.$$

For brevity, we denote

(3.109)

$$q_{-2}^1=\frac{c(\xi)b_0^2(x_0)c(\xi)}{|\xi|^4}+\frac{c(\xi)}{|\xi|^6}c(dx_n)\left[\partial_{x_n}[c(\xi')](x_0)|\xi|^2-c(\xi)h'(0)\right],$$

then

(3.110)

$$\begin{aligned}\partial_{\xi_n}(q_{-2}^1) &= \frac{1}{(1+\xi_n^2)^3}[(2\xi_n-2\xi_n^3)c(dx_n)b_0^2(x_0)c(dx_n)+(1-3\xi_n^2)c(dx_n)b_0^2(x_0)c(\xi')\\ &\quad+(1-3\xi_n^2)c(\xi')b_0^2(x_0)c(dx_n)-4\xi_n c(\xi')b_0^2(x_0)c(\xi')+(3\xi_n^2-1)\partial_{x_n}c(\xi')\\ &\quad-4\xi_n c(\xi')c(dx_n)\partial_{x_n}c(\xi')+2h'(0)c(\xi')+2h'(0)\xi_n c(dx_n)]\\ &\quad+6\xi_n h'(0)\frac{c(\xi)c(dx_n)c(\xi)}{(1+\xi_n^2)^4}.\end{aligned}$$

We calculate

(3.111)

$$\begin{aligned}&\mathrm{tr}\bigg[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})(x_0)\times\partial_{\xi_n}\bigg(\frac{c(\xi)b_0^1(x_0)c(\xi)}{|\xi|^4}\bigg)\bigg]|_{|\xi'|=1}\\ &=\frac{-1}{2(\xi_n-i)(\xi_n+i)^3}\mathrm{tr}[b_0^1(x_0)c(\xi')]+\frac{i}{2(\xi_n-i)(\xi_n+i)^3}\mathrm{tr}[b_0^1(x_0)c(dx_n)]\\ &=\frac{-1}{2(\xi_n-i)(\xi_n+i)^3}\mathrm{tr}[b_0^1(x_0)c(\xi')],\end{aligned}$$

then

$$
\begin{aligned}
(3.112)\qquad & -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\operatorname{trace}\left[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}\left(\frac{c(\xi)b_0^1c(\xi)}{|\xi|^4}\right)\right](x_0)d\xi_n\sigma(\xi')dx'\\
&=-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{-1}{2(\xi_n-i)(\xi_n+i)^3}\operatorname{tr}[b_0^1(x_0)c(\xi')]d\xi_n\sigma(\xi')dx'\\
&=0.
\end{aligned}
$$

Likewise, we have

$$
\begin{aligned}
(3.113)\qquad & \operatorname{tr}\left[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})(x_0)\times\partial_{\xi_n}\left(\frac{c(\xi)Ac(\xi)}{|\xi|^4}\right)\right]|_{|\xi'|=1}\\
&=\frac{-1}{2(\xi_n-i)(\xi_n+i)^3}\operatorname{tr}[Ac(\xi')]+\frac{i}{2(\xi_n-i)(\xi_n+i)^3}\operatorname{tr}[Ac(dx_n)],
\end{aligned}
$$

then by $\int_{|\xi'|=1}\{\xi_{i_1}\cdots\xi_{i_{2d+1}}\}\sigma(\xi')=0$, we have

$$
\begin{aligned}
(3.114)\qquad & -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\operatorname{trace}\left[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}\left(\frac{c(\xi)Ac(\xi)}{|\xi|^4}\right)\right](x_0)d\xi_n\sigma(\xi')dx'\\
&=-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{-1}{2(\xi_n-i)(\xi_n+i)^3}\operatorname{tr}[Ac(\xi')]d\xi_n\sigma(\xi')dx'\\
&-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{i}{2(\xi_n-i)(\xi_n+i)^3}\operatorname{tr}[Ac(dx_n)]d\xi_n\sigma(\xi')dx'\\
&=\frac{\Omega_3}{2}\operatorname{tr}[Ac(dx_n)]\int_{\Gamma^+}\frac{1}{(\xi_n-i)(\xi_n+i)^3}|_{\xi_n=i}dx'\\
&=\frac{\Omega_3}{2}\operatorname{tr}[Ac(dx_n)]2\pi i[\frac{1}{(\xi_n+i)^3}]^{(1)}|_{\xi_n=i}dx'\\
&=-\frac{\pi}{4}\Omega_3\operatorname{tr}[Ac(dx_n)]dx'.
\end{aligned}
$$

By (3.103) and (3.109), we have

(3.115)

$$
\operatorname{tr}[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}(q^1_{-2})](x_0)|_{|\xi'|=1}=\frac{12h'(0)(i\xi_n^2+\xi_n-2i)}{(\xi-i)^3(\xi+i)^3}+\frac{48h'(0)i\xi_n}{(\xi-i)^3(\xi+i)^4}.
$$

We compute that

$$
\begin{aligned}
(3.116)\quad & -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{tr}[\pi^{+}_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}(q^{1}_{-2})](x_0)d\xi_n\sigma(\xi')dx' \\
&=-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{12h'(0)(i\xi_n^2+\xi_n-2i)}{(\xi-i)^3(\xi+i)^3}+\frac{48h'(0)i\xi_n}{(\xi-i)^3(\xi+i)^4}d\xi_n\sigma(\xi')dx' \\
&=-i\Omega_3\int_{\Gamma^+}\frac{12h'(0)(i\xi_n^2+\xi_n-2i)}{(\xi-i)^3(\xi+i)^3}+\frac{48h'(0)i\xi_n}{(\xi-i)^3(\xi+i)^4}d\xi_n dx' \\
&=-i\Omega_3\frac{2\pi i}{2!}[\frac{12h'(0)(i\xi_n^2+\xi_n-2i)}{(\xi+i)^3}]^{(2)}|_{\xi_n=i}dx'-i\Omega_3\frac{2\pi i}{2!}[\frac{48h'(0)i\xi_n}{(\xi+i)^4}]^{(2)}|_{\xi_n=i}dx' \\
&=-\frac{9}{2}\pi h'(0)\Omega_3 dx'.
\end{aligned}
$$

Then,

$$
\Phi_5=-\frac{9}{2}\pi h'(0)\Omega_3dx'-\frac{\pi}{4}\Omega_3\mathrm{tr}[Ac(dx_n)]dx'. \tag{3.117}
$$

So

$$
\Phi=\Phi_1+\Phi_2+\Phi_3+\Phi_4+\Phi_5=0. \tag{3.118}
$$

Theorem 3.18. *Let M be a 4-dimensional oriented compact manifold with the boundary ∂M and the metric g^M as above, D_A be the perturbation of the de Rham Hodge operator on $\widetilde{M}$, then*

$$
\begin{aligned}
&\widetilde{\mathrm{Wres}}[\pi^{+}{D_A}^{-1}\circ\pi^{+}{D_A}^{-1}] \\
&=32\pi^2\int_M\mathrm{tr}\Big(-\frac{1}{12}s+A^2-\frac{1}{2}\sum_{j=1}^{n}Ac(\widetilde{e_j})Ac(\widetilde{e_j})\Big)d\mathrm{Vol_M}.
\end{aligned}
$$

where s is the scalar curvature.

We can directly state the following facts as a corollary of Theorem 3.18.

Corollary 3.19. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=c(X)$, then*

$$
\widetilde{\mathrm{Wres}}[\pi^{+}{D_A}^{-1}\circ\pi^{+}{D_A}^{-1}]=512\pi^2\int_M\Big(-\frac{1}{12}s\Big)d\mathrm{Vol_M}, \tag{3.119}
$$

where s is the scalar curvature.

When $A=\overline{c}(X)$, we can get the following corollary:

Corollary 3.20. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=\overline{c}(X)$, then*

$$
\widetilde{\mathrm{Wres}}[\pi^{+}{D_A}^{-1}\circ\pi^{+}{D_A}^{-1}]=512\pi^2\int_M\Big(-\frac{1}{12}s-|X|^2\Big)d\mathrm{Vol_M}, \tag{3.120}
$$

where s is the scalar curvature.

When $A = c(X)c(Y)$, similar to Corollary 3.20, we have :

Corollary 3.21. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)c(Y)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+{D_A}^{-1}] = 512\pi^2\int_M\Big(-\frac{1}{12}s + |X|^2|Y|^2 + 2g(X,Y)^2\Big)d\mathrm{Vol_M}, \tag{3.121}$$

where s is the scalar curvature.

When $A = c(X)\overline{c}(Y)$, we compute $\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+{D_A}^{-1}]$.

Corollary 3.22. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)\overline{c}(Y)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+{D_A}^{-1}] = 512\pi^2\int_M\Big(-\frac{1}{12}s\Big)d\mathrm{Vol_M}, \tag{3.122}$$

where s is the scalar curvature.

A simple calculation shows that:

Corollary 3.23. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)\overline{c}(Y)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+{D_A}^{-1}] = 512\pi^2\int_M\Big(-\frac{1}{12}s - |X|^2|Y|^2 + 4g(X,Y)^2\Big)d\mathrm{Vol_M}, \tag{3.123}$$

where s is the scalar curvature.

When $A = c(X)c(Y)c(Z)$, we have:

Corollary 3.24. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)c(Y)c(Z)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+{D_A}^{-1}] = 512\pi^2\int_M\Big(-\frac{1}{12}s\Big)d\mathrm{Vol_M}, \tag{3.124}$$

where s is the scalar curvature.

When $A = \overline{c}(X)c(Y)c(Z)$, we notice that:

Corollary 3.25. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)c(Y)c(Z)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+{D_A}^{-1}] = 512\pi^2\int_M\Big(-\frac{1}{12}s + |X|^2|Y|^2|Z|^2 - 2|X|^2g(Y,Z)^2\Big)d\mathrm{Vol_M}, \tag{3.125}$$

where s is the scalar curvature.

When $A = \overline{c}(X)\overline{c}(Y)c(Z)$, we have:

Corollary 3.26. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)\overline{c}(Y)c(Z)$, then*

(3.126)
$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+ {D_A}^{-1}] = 512\pi^2 \int_M \Big(-\frac{1}{12}s - 2|X|^2|Y|^2|Z|^2 + 2|Z|^2 g(X,Y)^2 \Big) d\mathrm{Vol_M},$$

where s is the scalar curvature.

When $A = \overline{c}(X)\overline{c}(Y)\overline{c}(Z)$, then $A^* = \overline{c}(Z)\overline{c}(Y)\overline{c}(X)$.
By computation, we have

$$\begin{aligned}
(3.127)\quad & \mathrm{tr}[A^2] = \mathrm{tr}[|X|^2|Y|^2|Z|^2] = |X|^2|Y|^2|Z|^2 \mathrm{tr}[\mathtt{id}], \\
& \mathrm{tr}[c(dx_n)A^*] = 0, \mathrm{tr}[c(dx_n)A] = 0, \\
(3.128)\quad & \sum_{j=1}^{n} \mathrm{tr}[Ac(\widetilde{e_j})Ac(\widetilde{e_j})] = \sum_{j=1}^{n} \mathrm{tr}[\overline{c}(X)\overline{c}(Y)\overline{c}(Z)\overline{c}(X)\overline{c}(Y)\overline{c}(Z)] \\
& = -\sum_{j=1}^{n} \mathrm{tr}[\overline{c}(Y)\overline{c}(X)\overline{c}(Z)\overline{c}(X)\overline{c}(Y)\overline{c}(Z)] + 2\sum_{j=1}^{n} g(X,Y)\mathrm{tr}[\overline{c}(Z)\overline{c}(X)\overline{c}(Y)\overline{c}(Z)] \\
& = -n|X|^2|Y|^2|Z|^2\mathrm{tr}[\mathtt{id}] + 2n|X|^2 g(Y,Z)^2\mathrm{tr}[\mathtt{id}] + 2n|Y|^2 g(X,Z)^2\mathrm{tr}[\mathtt{id}] \\
& + 2n|Z|^2 g(X,Y)^2\mathrm{tr}[\mathtt{id}] - 4n g(X,Y)g(X,Z)(Y,Z)\mathrm{tr}[\mathtt{id}]
\end{aligned}$$

We can calculate

(3.129)
$$\begin{aligned}
& \int_M \int_{|\xi|=1} \mathrm{trace}_{\wedge^* T^*M}[\sigma_{-4}({D_A}^{-2})]\sigma(\xi)dx \\
& = 32\pi^2 \int_M \mathrm{tr}\Big(-\frac{1}{12}s + A^2 - \frac{1}{2}\sum_{j=1}^{n} Ac(\widetilde{e_j})Ac(\widetilde{e_j}) \Big) d\mathrm{Vol_M} \\
& = 32\pi^2 \int_M \Big(-\frac{1}{12}s + |X|^2|Y|^2|Z|^2 + 2|X|^2|Y|^2|Z|^2 - 4|X|^2 g(Y,Z)^2 - 4|Y|^2 g(X,Z)^2 \\
& - 4|Z|^2 g(X,Y)^2 + 8g(X,Y)g(X,Z)(Y,Z) \Big) \mathrm{tr}[\mathtt{id}] d\mathrm{Vol_M} \\
& = 512\pi^2 \int_M \Big(-\frac{1}{12}s + 3|X|^2|Y|^2|Z|^2 - 4|X|^2 g(Y,Z)^2 - 4|Y|^2 g(X,Z)^2 \\
& - 4|Z|^2 g(X,Y)^2 + 8g(X,Y)g(X,Z)(Y,Z) \Big) d\mathrm{Vol_M},
\end{aligned}$$

and

$$(3.130)\qquad \int_{\partial M} \Psi = 0.$$

Corollary 3.27. *Let M be a 4-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)\overline{c}(Y)\overline{c}(Z)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+ {D_A}^{-1}] = 512\pi^2 \int_M \Big(-\frac{1}{12}s + 3|X|^2|Y|^2|Z|^2 - 4|X|^2 g(Y,Z)^2 - 4|Y|^2 g(X,Z)^2 - 4|Z|^2 g(X,Y)^2 + 8g(X,Y)g(X,Z)g(Y,Z)\Big) d\mathrm{Vol_M}, \tag{3.131}$$

where s is the scalar curvature.

4. A Kastler-Kalau-Walze type theorem for 6-dimensional manifolds with boundary

Firstly, we prove the Kastler-Kalau-Walze type theorems for 6-dimensional manifolds with boundary. From [14], we know that

$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+ (D_A^* D_A D_A^*)^{-1}] = \int_M \int_{|\xi|=1} \mathrm{trace}_{\wedge^* T^*M}[\sigma_{-4}((D_A^* D_A)^{-2})]\sigma(\xi)dx + \int_{\partial M} \overline{\Psi}, \tag{4.1}$$

where

$$\begin{aligned}\overline{\Psi} = &\int_{|\xi'|=1} \int_{-\infty}^{+\infty} \sum_{j,k=0}^{\infty} \sum \frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!} \times \mathrm{trace}_{\wedge^* T^*M}[\partial_{x_n}^j \partial_{\xi'}^{\alpha} \partial_{t\xi_n}^k \sigma_r^+ ({D_A}^{-1})(x',0,\xi',\xi_n) \\ &\times \partial_{x'}^{\alpha} \partial_{\xi_n}^{j+1} \partial_{x_n}^k \sigma_l((D_A^* D_A D_A^*)^{-1})(x',0,\xi',\xi_n)]d\xi_n \sigma(\xi')dx',\end{aligned} \tag{4.2}$$

and the sum is taken over $r + \ell - k - j - |\alpha| - 1 = -6,\ r \leq -1, \ell \leq -3$. By Theorem 2.2, we compute the interior term of (4.1), then

$$\begin{aligned}&\int_M \int_{|\xi|=1} \mathrm{trace}_{\wedge^* T^*M}[\sigma_{-4}((D_A^* D_A)^{-2})]\sigma(\xi)dx \\ &= 128\pi^3 \int_M \mathrm{tr}\Big(-\frac{1}{12}s + 2A^*A - \frac{1}{4}\sum_{j=1}^n Ac(\widetilde{e_j})Ac(\widetilde{e_j}) - \frac{1}{4}\sum_{j=1}^n A^* c(\widetilde{e_j}) A^* c(\widetilde{e_j}) \\ &+ \frac{1}{2}\sum_{j=1}^n \nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A^*)c(\widetilde{e_j}) - \frac{1}{2}\sum_{j=1}^n c(\widetilde{e_j})\nabla^{\bigwedge^* T^*M}_{\widetilde{e_j}}(A)\Big) d\mathrm{Vol_M}.\end{aligned} \tag{4.3}$$

Next, we compute $\int_{\partial M}\overline{\Psi}$. By computation, we get

(4.4)

$$
\begin{aligned}
D_A^*D_AD_A^* &= \sum_{i=1}^n c(\widetilde{e_i})\langle\widetilde{e_i},dx_l\rangle(-g^{ij}\partial_l\partial_i\partial_j)+\sum_{i=1}^n c(\widetilde{e_i})\langle\widetilde{e_i},dx_l\rangle\Big\{-(\partial_l g^{ij})\partial_i\partial_j-g^{ij}\Big(4(\sigma_i\\
&+a_i)\partial_j-2\Gamma_{ij}^k\partial_k\Big)\partial_l\Big\}+\sum_{i=1}^n c(\widetilde{e_i})\langle\widetilde{e_i},dx_l\rangle\Big\{-2(\partial_l g^{ij})(\sigma_i+a_i)\partial_j+g^{ij}(\partial_l\Gamma_{ij}^k)\partial_k\\
&-2g^{ij}[(\partial_l\sigma_i)+(\partial_l a_i)]\partial_j+(\partial_l g^{ij})\Gamma_{ij}^k\partial_k+\sum_{j,k}\Big[\partial_l\Big(c(\widetilde{e_j})A+A^*c(\widetilde{e_j})\Big)\Big]\langle\widetilde{e_j},dx_k\rangle\partial_k\\
&+\sum_{j,k}\Big(c(\widetilde{e_j})A+A^*c(\widetilde{e_j})\Big)\Big[\partial_l\langle\widetilde{e_j},dx_k\rangle\Big]\partial_k\Big\}+\sum_{i=1}^n c(\widetilde{e_i})\langle c(\widetilde{e_i}),dx_l\rangle\partial_l\Big\{-g^{ij}\Big[(\partial_i\sigma_j)\\
&+(\partial_i a_j)+\sigma_i\sigma_j+\sigma_i a_j+a_i\sigma_j+a_ia_j-\Gamma_{ij}^k\sigma_k-\Gamma_{ij}^k a_k\Big]+\sum_{i,j}g^{ij}\Big[c(\partial_i)\partial_j(A)\\
&+c(\partial_i)\sigma_jA+c(\partial_i)a_jA+A^*c(\partial_i)\sigma_j+A^*c(\partial_i)a_j\Big]-\frac{1}{8}\sum_{i,j,k,l}R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l})\\
&+\frac{1}{4}s+A^*A\Big\}+\Big[\sum_{i=1}^n c(\widetilde{e_i})(\sigma_i+a_i)+A^*\Big](-g^{ij}\partial_i\partial_j)+\sum_{i=1}^n c(\widetilde{e_i})\langle\widetilde{e_i},dx_l\rangle\Big\{2\sum_{j,k}\\
&\Big[c(\widetilde{e_j})A+A^*c(\widetilde{e_j})\Big]\times\langle\widetilde{e_j},dx_k\rangle\Big\}\partial_l\partial_k+\Big[\sum_{i=1}^n c(\widetilde{e_i})(\sigma_i+a_i)+A^*\Big]\Big\{-\sum_{i,j}g^{ij}\Big[2(\sigma_i\\
&+a_i)\partial_j-\Gamma_{ij}^k\partial_k+(\partial_i\sigma_j)+(\partial_i a_j)+\sigma_i\sigma_j+\sigma_i a_j+a_i\sigma_j+a_ia_j-\Gamma_{ij}^k\sigma_k-\Gamma_{ij}^k a_k\Big]\\
&+\sum_{i,j}g^{ij}\Big(c(\partial_i)A+A^*c(\partial_i)\Big)\partial_j+\sum_{i,j}g^{ij}\Big[c(\partial_i)\partial_j(A)+c(\partial_i)\sigma_jA+c(\partial_i)a_jA\\
&+A^*c(\partial_i)\sigma_j+A^*c(\partial_i)a_j\Big]-\frac{1}{8}\sum_{i,j,k,l}R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l})+\frac{1}{4}s+A^*A\Big\}.
\end{aligned}
$$

Then, we obtain

Lemma 4.1. *The following identities hold:*

(4.5)

$$
\begin{aligned}
\sigma_2(D_A^*D_AD_A^*) &= \sum_{i,j,l}c(dx_l)\partial_l(g^{ij})\xi_i\xi_j+c(\xi)(4\sigma^k+4a^k-2\Gamma^k)\xi_k+2[|\xi|^2A-c(\xi)A^*c(\xi)]\\
&\quad+\frac{1}{4}|\xi|^2\sum_{s,t,l}\omega_{s,t}(\widetilde{e_l})[c(\widetilde{e_l})\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t})-c(\widetilde{e_l})c(\widetilde{e_s})c(\widetilde{e_t})]+|\xi|^2A^*;\\
\sigma_3(D_A^*D_AD_A^*) &= ic(\xi)|\xi|^2.
\end{aligned}
$$

Write

$$\sigma(D_A^*D_AD_A^*)=p_3+p_2+p_1+p_0;\ \sigma((D_A^*D_AD_A^*)^{-1})=\sum_{j=3}^{\infty}q_{-j}. \tag{4.6}$$

By the composition formula of pseudodifferential operators, we have

(4.7)

$$\begin{aligned}1=\sigma((D_A^*D_AD_A^*)\circ(D_A^*D_AD_A^*)^{-1})&=\sum_{\alpha}\frac{1}{\alpha!}\partial_\xi^\alpha[\sigma(D_A^*D_AD_A^*)]D_x^\alpha[(D_A^*D_AD_A^*)^{-1}]\\&=(p_3+p_2+p_1+p_0)(q_{-3}+q_{-4}+q_{-5}+\cdots)\\&+\sum_j(\partial_{\xi_j}p_3+\partial_{\xi_j}p_2+\partial_{\xi_j}p_1+\partial_{\xi_j}p_0)\\&(D_{x_j}q_{-3}+D_{x_j}q_{-4}+D_{x_j}q_{-5}+\cdots)\\&=p_3q_{-3}+(p_3q_{-4}+p_2q_{-3}+\sum_j\partial_{\xi_j}p_3D_{x_j}q_{-3})+\cdots,\end{aligned}$$

by (4.7), we have

$$q_{-3}=p_3^{-1};\ q_{-4}=-p_3^{-1}[p_2p_3^{-1}+\sum_j\partial_{\xi_j}p_3D_{x_j}(p_3^{-1})]. \tag{4.8}$$

By Lemma 4.1, we have some symbols of operators.

Lemma 4.2. *The following identities hold:*

(4.9)

$$\begin{aligned}\sigma_{-3}((D_A^*D_AD_A^*)^{-1})&=\frac{ic(\xi)}{|\xi|^4};\\\sigma_{-4}((D_A^*D_AD_A^*)^{-1})&=\frac{c(\xi)\sigma_2(D_A^*D_AD_A^*)c(\xi)}{|\xi|^8}+\frac{c(\xi)}{|\xi|^8}\Big(|\xi|^2c(dx_n)\partial_{x_n}c(\xi')-2h'(0)c(dx_n)c(\xi)\\&+2\xi_nc(\xi)\partial_{x_n}c(\xi')+4\xi_nh'(0)\Big).\end{aligned}$$

When $n=6$, then $\mathrm{tr}_{\wedge^*T^*M}[\mathtt{id}]=64$, where tr as shorthand of trace. Since the sum is taken over $r+\ell-k-j-|\alpha|-1=-6,\ r\le-1,\ell\le-3$, then we have the $\int_{\partial_M}\overline{\Psi}$ is the sum of the following five cases:

case (a) (I) $r=-1,l=-3,j=k=0,|\alpha|=1.$

By (4.2), we obtain

(4.10)

$$\overline{\Psi}_1=-\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\sum_{|\alpha|=1}\mathrm{trace}[\partial_{\xi'}^\alpha\pi_{\xi_n}^+\sigma_{-1}({D_A}^{-1})\times\partial_{x'}^\alpha\partial_{\xi_n}\sigma_{-3}((D_A^*D_AD_A^*)^{-1})](x_0)d\xi_n\sigma(\xi')dx'.$$

case (a) (II) $r=-1, l=-3, |\alpha|=k=0, j=1.$

By (4.2), we have

(4.11)
$$\overline{\Psi}_2=-\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\operatorname{trace}[\partial_{x_n}\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial^2_{\xi_n}\sigma_{-3}((D^*_AD_AD^*_A)^{-1})](x_0)d\xi_n\sigma(\xi')dx'.$$

case (a) (III) $r=-1, l=-3, |\alpha|=j=0, k=1.$

It is easy to check that

(4.12)
$$\overline{\Psi}_3=-\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\operatorname{trace}[\partial_{\xi_n}\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}\partial_{x_n}\sigma_{-3}((D^*_AD_AD^*_A)^{-1})](x_0)d\xi_n\sigma(\xi')dx'.$$

For [14], we have

$$\overline{\Psi}_1+\overline{\Psi}_2+\overline{\Psi}_3=\frac{10}{2}\pi h'(0)\Omega_4dx'. \tag{4.13}$$

case (b) $r=-1, l=-4, |\alpha|=j=k=0.$

By observing (4.2), we have

(4.14)
$$\begin{aligned}\overline{\Psi}_4&=-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\operatorname{trace}[\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}\sigma_{-4}((D^*_AD_AD^*_A)^{-1})](x_0)d\xi_n\sigma(\xi')dx'\\&=i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\operatorname{trace}[\partial_{\xi_n}\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\sigma_{-4}((D^*_AD_AD^*_A)^{-1})](x_0)d\xi_n\sigma(\xi')dx'.\end{aligned}$$

In the normal coordinate, $g^{ij}(x_0)=\delta^j_i$ and $\partial_{x_j}(g^{\alpha\beta})(x_0)=0$, if $j<n$; $\partial_{x_j}(g^{\alpha\beta})(x_0)=h'(0)\delta^\alpha_\beta$, if $j=n$. So by [12], when $k<n$, we have $\Gamma^n(x_0)=\frac{5}{2}h'(0)$, $\Gamma^k(x_0)=0$, $\delta^n(x_0)=0$ and $\delta^k=\frac{1}{4}h'(0)c(\widetilde{e_k})c(\widetilde{e_n})$. Then, we obtain

(4.15)
$$\begin{aligned}\sigma_{-4}((D^*_AD_AD^*_A)^{-1})(x_0)=\frac{1}{|\xi|^8}c(\xi)\Big(&h'(0)c(\xi)\sum_{k<n}\xi_kc(\widetilde{e_k})c(\widetilde{e_n})-h'(0)c(\xi)\sum_{k<n}\xi_k\overline{c}(\widetilde{e_k})\overline{c}(\widetilde{e_n})\\&-5h'(0)\xi_nc(\xi)+2[|\xi|^2A-c(\xi)A^*c(\xi)]+\frac{5}{4}|\xi|^2h'(0)c(\widetilde{e_i})\overline{c}(dx_n)\overline{c}(\widetilde{e_i})\\&-\frac{1}{4}|\xi|^2h'(0)c(dx_n)+|\xi|^2A^*\Big)c(\xi)+\frac{c(\xi)}{|\xi|^8}\Big(|\xi|^2c(dx_n)\partial_{x_n}c(\xi')\\&-2h'(0)c(dx_n)c(\xi)+2\xi_nc(\xi)\partial_{x_n}c(\xi')+4\xi_nh'(0)\Big),\end{aligned}$$

$$\partial_{\xi_n}\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})(x_0)|_{|\xi'|=1}=-\frac{c(\xi')+ic(dx_n)}{2(\xi_n-i)^2}. \tag{4.16}$$

By (4.15) and (4.16), we have

$$
\begin{aligned}
(4.17)\qquad & \mathrm{tr}[\partial_{\xi_n}\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\sigma_{-4}(D_A^*D_AD_A^*)^{-1}](x_0)|_{|\xi'|=1}\\
&=-\frac{64h'(0)}{2(\xi_n-i)^2(1+\xi_n^2)^4}\big(-\frac{15}{4}i+16\xi_n+\frac{19}{2}i\xi_n^2+8\xi_n^3+\frac{21}{4}i\xi_n^4\big)\\
&+\frac{2+4i\xi_n-2\xi_n^2}{2(\xi_n-i)^2(1+\xi_n^2)^3}\mathrm{tr}\big[Ac(\xi')\big]+\frac{-2i+4\xi_n+2i\xi_n^2}{2(\xi_n-i)^2(1+\xi_n^2)^3}\mathrm{tr}\big[Ac(dx_n)\big]\\
&+\frac{3+2i\xi_n+\xi_n^2}{2(\xi_n-i)^2(1+\xi_n^2)^3}\mathrm{tr}\big[A^*c(\xi')\big]+\frac{i+2\xi_n+3i\xi_n^2}{2(\xi_n-i)^2(1+\xi_n^2)^3}\mathrm{tr}\big[A^*c(dx_n)\big].
\end{aligned}
$$

Consequently,

$$
\begin{aligned}
(4.18)\qquad \overline{\Psi}_4&=i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{-64h'(0)(-\frac{15}{4}i+16\xi_n+\frac{19}{2}i\xi_n^2+8\xi_n^3+\frac{21}{4}i\xi_n^4)}{2(\xi_n-i)^2(1+\xi_n^2)^4}d\xi_n\sigma(\xi')dx'\\
&+i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{-2i+4\xi_n+2i\xi_n^2}{2(\xi_n-i)^2(1+\xi_n^2)^3}\mathrm{tr}\big[Ac(dx_n)\big]d\xi_n\sigma(\xi')dx'\\
&+i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{i+2\xi_n+3i\xi_n^2}{2(\xi_n-i)^2(1+\xi_n^2)^3}\mathrm{tr}\big[A^*c(dx_n)\big]d\xi_n\sigma(\xi')dx'\\
&=\frac{111}{2}\pi h'(0)\Omega_4dx'-\frac{3}{8}\pi\Omega_4\mathrm{tr}\big[Ac(dx_n)\big]dx'+\frac{1}{16}\pi\Omega_4\mathrm{tr}\big[A^*c(dx_n)\big]dx'.
\end{aligned}
$$

case (c) $r=-2,l=-3,|\alpha|=j=k=0.$

By (4.2), we obtain

$$
(4.19)\quad \overline{\Psi}_5=-i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\pi^+_{\xi_n}\sigma_{-2}({D_A}^{-1})\times\partial_{\xi_n}\sigma_{-3}((D_A^*D_AD_A^*)^{-1})](x_0)d\xi_n\sigma(\xi')dx'.
$$

By Lemma 4.1 and Lemma 4.2, we have

(4.20)

$$
\sigma_{-2}({D_A}^{-1})(x_0)=\frac{c(\xi)\sigma_0(D_A)(x_0)c(\xi)}{|\xi|^4}+\frac{c(\xi)}{|\xi|^6}\sum_j c(dx_j)\left[\partial_{x_j}(c(\xi))|\xi|^2-c(\xi)\partial_{x_j}(|\xi|^2)\right](x_0),
$$

where

(4.21)

$$
\sigma_0(D_A)(x_0)=\frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})(x_0)c(\widetilde{e_i})\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t})-\frac{1}{4}\sum_{i,s,t}\omega_{s,t}(\widetilde{e_i})(x_0)c(\widetilde{e_i})c(\widetilde{e_s})c(\widetilde{e_t})+A.
$$

On the other hand,

$$
(4.22)\qquad \partial_{\xi_n}\sigma_{-3}((D_A^*D_AD_A^*)^{-1})(x_0)|_{|\xi'|=1}=\frac{-8i\xi_nc(\xi')}{(1+\xi_n^2)^5}+\frac{i(1-7\xi_n^2)c(dx_n)}{(1+\xi_n^2)^5}.
$$

It is easy to obtain that

$$\pi^+_{\xi_n}\Big(\sigma_{-2}(D_A{}^{-1})\Big)(x_0)|_{|\xi'|=1} = \pi^+_{\xi_n}\Big(\frac{c(\xi)\sigma_0(D_A)(x_0)c(\xi)+c(\xi)c(dx_n)\partial_{x_n}[c(\xi')](x_0)}{(1+\xi_n^2)^2}\Big) - h'(0)\pi^+_{\xi_n}\Big(\frac{c(\xi)c(dx_n)c(\xi)}{(1+\xi_n^2)^3}\Big). \tag{4.23}$$

We denote

$$\sigma_0(D_A)(x_0) = b_0^1(x_0)+b_0^2(x_0)+A. \tag{4.24}$$

Then, we obtain

$$\begin{aligned}\pi^+_{\xi_n}\Big(\sigma_{-2}(D_A{}^{-1})\Big)(x_0)|_{|\xi'|=1} &= \pi^+_{\xi_n}\Big(\frac{c(\xi)b_0^2(x_0)c(\xi)+c(\xi)c(dx_n)\partial_{x_n}[c(\xi')](x_0)}{(1+\xi_n^2)^2}\Big)\\ &\quad - h'(0)\pi^+_{\xi_n}\Big(\frac{c(\xi)c(dx_n)c(\xi)}{(1+\xi_n^2)^3}\Big)+\pi^+_{\xi_n}\Big(\frac{c(\xi)b_0^1(x_0)c(\xi)}{(1+\xi_n^2)^2}\Big)\\ &\quad + \pi^+_{\xi_n}\Big(\frac{c(\xi)Ac(\xi)}{(1+\xi_n^2)^2}\Big).\end{aligned} \tag{4.25}$$

Furthermore,

$$\begin{aligned}\pi^+_{\xi_n}\Big(\frac{c(\xi)b_0^1(x_0)c(\xi)}{(1+\xi_n^2)^2}\Big) &= \pi^+_{\xi_n}\Big(\frac{c(\xi')b_0^1(x_0)c(\xi')}{(1+\xi_n^2)^2}\Big)+\pi^+_{\xi_n}\Big(\frac{\xi_n c(\xi')b_0^1(x_0)c(dx_n)}{(1+\xi_n^2)^2}\Big)\\ &\quad + \pi^+_{\xi_n}\Big(\frac{\xi_n c(dx_n)b_0^1(x_0)c(\xi')}{(1+\xi_n^2)^2}\Big)+\pi^+_{\xi_n}\Big(\frac{\xi_n^2 c(dx_n)b_0^1(x_0)c(dx_n)}{(1+\xi_n^2)^2}\Big)\\ &= -\frac{c(\xi')b_0^1(x_0)c(\xi')(2+i\xi_n)}{4(\xi_n-i)^2}-\frac{ic(\xi')b_0^1(x_0)c(dx_n)}{4(\xi_n-i)^2}\\ &\quad -\frac{ic(dx_n)b_0^1(x_0)c(\xi')}{4(\xi_n-i)^2}-\frac{i\xi_n c(dx_n)b_0^1(x_0)c(dx_n)}{4(\xi_n-i)^2}.\end{aligned} \tag{4.26}$$

For the sake of convenience in writing,

$$\pi^+_{\xi_n}\Big(\frac{c(\xi)b_0^2(x_0)c(\xi)+c(\xi)c(dx_n)\partial_{x_n}(c(\xi'))(x_0)}{(1+\xi_n^2)^2}\Big) - h'(0)\pi^+_{\xi_n}\Big(\frac{c(\xi)c(dx_n)c(\xi)}{(1+\xi_n)^3}\Big) := B_1 - B_2, \tag{4.27}$$

where

$$\begin{aligned}B_1 = \frac{-1}{4(\xi_n-i)^2}\big[&(2+i\xi_n)c(\xi')b_0^2(x_0)c(\xi')+i\xi_n c(dx_n)b_0^2(x_0)c(dx_n)\\ &+(2+i\xi_n)c(\xi')c(dx_n)\partial_{x_n}c(\xi')+ic(dx_n)b_0^2(x_0)c(\xi')+ic(\xi')b_0^2(x_0)c(dx_n)-i\partial_{x_n}c(\xi')\big]\end{aligned} \tag{4.28}$$

and

$$B_2 = \frac{h'(0)}{2}\Big(\frac{c(dx_n)}{4i(\xi_n-i)}+\frac{c(dx_n)-ic(\xi')}{8(\xi_n-i)^2}+\frac{3\xi_n-7i}{8(\xi_n-i)^3}\big(ic(\xi')-c(dx_n)\big)\Big). \tag{4.29}$$

By (4.22) and (4.28), we have

$$\begin{aligned}(4.30)\quad &\mathrm{tr}[B_1\times\partial_{\xi_n}\sigma_{-3}((D_A^*D_AD_A^*)^{-1})(x_0)]|_{|\xi'|=1}=\mathrm{tr}\Big[\frac{-1}{4(\xi_n-i)^2}\big[(2+i\xi_n)c(\xi')b_0^2(x_0)c(\xi')\\&+i\xi_nc(dx_n)b_0^2(x_0)c(dx_n)+ic(dx_n)b_0^2(x_0)c(\xi')+(2+i\xi_n)c(\xi')c(dx_n)\partial_{x_n}c(\xi')\\&+ic(\xi')b_0^2(x_0)c(dx_n)-i\partial_{x_n}c(\xi')\big]\times\frac{-8i\xi_nc(\xi')+(i-7i\xi_n^2)c(dx_n)}{(1+\xi_n^2)^5}\Big]\\&=-32h'(0)\frac{-2i+9\xi_n+14i\xi_n^2-7\xi_n^3}{4(\xi_n-i)^2(1+\xi_n^2)^5}-64h'(0)\frac{-\frac{3}{2}i+12\xi_n+\frac{21}{2}i\xi_n^2}{4(\xi_n-i)^2(1+\xi_n^2)^5}\\&=8h'(0)\frac{-5-28i\xi_n+7\xi_n^2}{(\xi_n-i)^6(\xi_n+i)^5}.\end{aligned}$$

Similarly, we have

$$\begin{aligned}(4.31)\quad &\mathrm{tr}[B_2\times\partial_{\xi_n}\sigma_{-3}((D_A^*D_AD_A^*)^{-1})(x_0)]|_{|\xi'|=1}=\mathrm{tr}\Big\{\frac{h'(0)}{2}\Big[\frac{c(dx_n)}{4i(\xi_n-i)}+\frac{c(dx_n)-ic(\xi')}{8(\xi_n-i)^2}\\&+\frac{3\xi_n-7i}{8(\xi_n-i)^3}[ic(\xi')-c(dx_n)]\Big]\times\frac{-8i\xi_nc(\xi')+(i-7i\xi_n^2)c(dx_n)}{(1+\xi_n^2)^5}\Big\}\\&=-8h'(0)\frac{4i-23\xi_n-14i\xi_n^2+7\xi_n^3}{(\xi_n-i)^7(\xi_n+i)^5},\end{aligned}$$

thus

$$(4.32)\quad \mathrm{tr}[B_1-B_2\times\partial_{\xi_n}\sigma_{-3}((D_A^*D_AD_A^*)^{-1})(x_0)]|_{|\xi'|=1}=8h'(0)\frac{9i-56\xi_n-49i\xi_n^2+14\xi_n^3}{(\xi_n-i)^7(\xi_n+i)^5}.$$

We calculate that

$$\begin{aligned}(4.33)\qquad &\mathrm{tr}\Big[\pi^+_{\xi_n}\Big(\frac{c(\xi)b_0^1(x_0)c(\xi)}{(1+\xi_n^2)^2}\Big)\times\partial_{\xi_n}\sigma_{-3}((D_A^*D_AD_A^*)^{-1})(x_0)\Big]\Big|_{|\xi'|=1}\\&=\frac{-2-16i\xi_n+14\xi_n^2}{4(\xi_n-i)^7(\xi_n+i)^5}\mathrm{tr}[b_0^1(x_0)c(\xi')].\end{aligned}$$

Similar calculations to (4.33), it is shown that

$$\begin{aligned}(4.34)\qquad &\mathrm{tr}\Big[\pi^+_{\xi_n}\Big(\frac{c(\xi)Ac(\xi)}{(1+\xi_n^2)^2}\Big)\times\partial_{\xi_n}\sigma_{-3}((D_A^*D_AD_A^*)^{-1})(x_0)\Big]\Big|_{|\xi'|=1}\\&=\frac{-2-16i\xi_n+14\xi_n^2}{4(\xi_n-i)^7(\xi_n+i)^5}\mathrm{tr}[Ac(\xi')]+\frac{-2i+16\xi_n+14i\xi_n^2}{4(\xi_n-i)^7(\xi_n+i)^5}\mathrm{tr}[Ac(dx_n)].\end{aligned}$$

By $\int_{|\xi'|=1}\xi_1\cdots\xi_{2q+1}\sigma(\xi')=0$, we have

$$
\begin{aligned}
\overline{\Psi}_5 &= -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty} 8h'(0)\frac{9i-56\xi_n-49i\xi_n^2+14\xi_n^3}{(\xi_n-i)^7(\xi_n+i)^5}d\xi_n\sigma(\xi')dx' \\
&\quad - i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{-2-16i\xi_n+14\xi_n^2}{4(\xi_n-i)^7(\xi_n+i)^5}\mathrm{tr}[b_0^1(x_0)c(\xi')]d\xi_n\sigma(\xi')dx' \\
&\quad - i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{-2-16i\xi_n+14\xi_n^2}{4(\xi_n-i)^7(\xi_n+i)^5}\mathrm{tr}[Ac(\xi')]d\xi_n\sigma(\xi')dx' \\
&\quad - i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{-2i+16\xi_n+14i\xi_n^2}{4(\xi_n-i)^7(\xi_n+i)^5}\mathrm{tr}[Ac(dx_n)]d\xi_n\sigma(\xi')dx' \\
&\quad = -i8h'(0)\Omega_4\frac{2\pi i}{6!}\Big[\frac{9i-56\xi_n-49i\xi_n^2+14\xi_n^3}{(\xi_n+i)^5}\Big]^{(5)}|_{\xi_n=i}dx' \\
&\quad - i\Omega_4\mathrm{tr}[Ac(dx_n)]\frac{2\pi i}{6!}\Big[\frac{-2i+16\xi_n+14i\xi_n^2}{4(\xi_n+i)^5}\Big]^{(5)}|_{\xi_n=i}dx' \\
&\quad = -\frac{105}{4}\pi h'(0)\Omega_4dx'+\frac{161}{512}\pi\Omega_4\mathrm{tr}[Ac(dx_n)]dx'.
\end{aligned}
\tag{4.35}
$$

Now $\overline{\Psi}$ is the sum of the cases (a), (b) and (c), then

$$
\overline{\Psi}=\frac{137}{4}\pi h'(0)\Omega_4dx'-\frac{31}{512}\pi\Omega_4\mathrm{tr}\big[Ac(dx_n)\big]dx'+\frac{1}{16}\pi\Omega_4\mathrm{tr}\big[A^*c(dx_n)\big]dx'.
\tag{4.36}
$$

Theorem 4.3. *Let M be a 6-dimensional compact oriented manifold with the boundary ∂M and the metric g^M as above, D_A be the perturbation of the de Rham Hodge operator on $\widetilde{M}$, then*

(4.37)

$$
\begin{aligned}
&\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+(D_A^*D_AD_A^*)^{-1}]=128\pi^3\int_M\mathrm{tr}\Big(-\frac{1}{12}s+2A^*A-\frac{1}{4}\sum_{j=1}^n Ac(\widetilde{e_j})Ac(\widetilde{e_j}) \\
&-\frac{1}{4}\sum_{j=1}^n A^*c(\widetilde{e_j})A^*c(\widetilde{e_j})+\frac{1}{2}\sum_{j=1}^n\nabla^{\bigwedge^*T^*M}_{\widetilde{e_j}}(A^*)c(\widetilde{e_j})-\frac{1}{2}\sum_{j=1}^n c(\widetilde{e_j})\nabla^{\bigwedge^*T^*M}_{\widetilde{e_j}}(A)\Big)d\mathrm{Vol_M} \\
&+\int_{\partial M}\frac{137}{4}\pi h'(0)\Omega_4d\mathrm{Vol}_{\partial\mathrm{M}}-\int_{\partial M}\frac{31}{512}\pi\Omega_4\mathrm{tr}\big[Ac(dx_n)\big]d\mathrm{Vol}_{\partial\mathrm{M}} \\
&+\int_{\partial M}\frac{1}{16}\pi\Omega_4\mathrm{tr}\big[A^*c(dx_n)\big]d\mathrm{Vol}_{\partial\mathrm{M}},
\end{aligned}
$$

where s is the scalar curvature.

We can state the following facts as a corollary of Theorem 4.3.

Corollary 4.4. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)$, then*

(4.38)

$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+(D_A^* D_A D_A^*)^{-1}] = 8192\pi^3 \int_M \Big(-\frac{1}{12}s + 4|X|^2 + \sum_{j=1}^{n} g(\nabla^{TM}_{\widetilde{e_j}} X, \widetilde{e_j}) \Big) d\mathrm{Vol_M}$$
$$+ \int_{\partial M} \frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial \mathrm{M}} + \int_{\partial M} \frac{63}{8}\pi\Omega_4 g(\partial x_n, X) d\mathrm{Vol}_{\partial \mathrm{M}},$$

where s is the scalar curvature.

When $A = \overline{c}(X)$, we can get the following corollary:

Corollary 4.5. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+(D_A^* D_A D_A^*)^{-1}] = 8192\pi^3 \int_M \Big(-\frac{1}{12}s - |X|^2 \Big) d\mathrm{Vol_M} + \int_{\partial M} \frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial \mathrm{M}}, \tag{4.39}$$

where s is the scalar curvature.

When $A = c(X)c(Y)$, we compute $\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+(D_A^* D_A D_A^*)^{-1}]$.

Corollary 4.6. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)c(Y)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+(D_A^* D_A D_A^*)^{-1}] = 8192\pi^3 \int_M \Big(-\frac{1}{12}s + |X|^2|Y|^2 + 4g(X,Y)^2 \Big) d\mathrm{Vol_M}$$
$$+ \int_{\partial M} \frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial \mathrm{M}},$$

where s is the scalar curvature.

When $A = c(X)\overline{c}(Y)$, we obtain:

Corollary 4.7. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)\overline{c}(Y)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+(D_A^* D_A D_A^*)^{-1}] = 8192\pi^3 \int_M \Big(-\frac{1}{12}s + 4|X|^2|Y|^2 \Big) d\mathrm{Vol_M} + \int_{\partial M} \frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial \mathrm{M}}, \tag{4.40}$$

where s is the scalar curvature.

When $A = \overline{c}(X)\overline{c}(Y)$, similar to Corollary 4.7, we can get:

Corollary 4.8. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)\overline{c}(Y)$, then*

(4.41)

$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+ (D_A^* D_A D_A^*)^{-1}] = 8192\pi^3 \int_M \Big(-\frac{1}{12}s - |X|^2|Y|^2 + 6g(X,Y)^2 \Big) d\mathrm{Vol_M}$$
$$+ \int_{\partial M} \frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial \mathrm{M}},$$

where s is the scalar curvature.

When $A = c(X)c(Y)c(Z)$, we compute $\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+ (D_A^* D_A D_A^*)^{-1}]$.

Corollary 4.9. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)c(Y)c(Z)$, then*

(4.42)

$$\begin{aligned}
&\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+ (D_A^* D_A D_A^*)^{-1}] = 8192\pi^3 \int_M \Big(-\frac{1}{12}s + 2|X|^2|Y|^2|Z|^2 \\
&+ 2|X|^2 g(Y,Z)^2 + 2|Y|^2 g(X,Z)^2 + 2|Z|^2 g(X,Y)^2 - 4g(X,Y)g(X,Z)g(Y,Z) \\
&- \sum_{j=1}^{6} g(\nabla^{TM}_{\widetilde{e}_j} Z, \widetilde{e}_j) g(X,Y) + \sum_{j=1}^{6} g(Y,\widetilde{e}_j) g(X, \nabla^{TM}_{\widetilde{e}_j} Z) - \sum_{j=1}^{6} g(X,\widetilde{e}_j) g(Y, \nabla^{TM}_{\widetilde{e}_j} Z) \\
&- \sum_{j=1}^{6} g(Z,\widetilde{e}_j) g(X, \nabla^{TM}_{\widetilde{e}_j} Y) + \sum_{j=1}^{6} g(\nabla^{TM}_{\widetilde{e}_j} Y, \widetilde{e}_j) g(X,Z) - \sum_{j=1}^{6} g(X,\widetilde{e}_j) g(\nabla^{TM}_{\widetilde{e}_j} Y, Z) \\
&- \sum_{j=1}^{6} g(Z,\widetilde{e}_j) g(\nabla^{TM}_{\widetilde{e}_j} X, Y) + \sum_{j=1}^{6} g(Y,\widetilde{e}_j) g(\nabla^{TM}_{\widetilde{e}_j} X, Z) - \sum_{j=1}^{6} g(\nabla^{TM}_{\widetilde{e}_j} X, \widetilde{e}_j) g(Y,Z) \Big) d\mathrm{Vol_M} \\
&+ \int_{\partial M} \frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial \mathrm{M}} + \int_{\partial M} \frac{1}{8}\pi\Omega_4 g(\partial x_n, X) g(Y,Z) d\mathrm{Vol}_{\partial \mathrm{M}} \\
&- \int_{\partial M} \frac{1}{8}\pi\Omega_4 g(\partial x_n, Y) g(X,Z) d\mathrm{Vol}_{\partial \mathrm{M}} + \int_{\partial M} \frac{1}{8}\pi\Omega_4 g(\partial x_n, Z) g(X,Y) d\mathrm{Vol}_{\partial \mathrm{M}},
\end{aligned}$$

where s is the scalar curvature.

When $A = \overline{c}(X)c(Y)c(Z)$, we get:

Corollary 4.10. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)c(Y)c(Z)$, then*

(4.43)

$$\begin{aligned}
&\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+ (D_A^* D_A D_A^*)^{-1}] \\
&= 8192\pi^3 \int_M \Big(-\frac{1}{12}s + 3|X|^2|Y|^2|Z|^2 - 4|X|^2 g(Y,Z)^2 \Big) d\mathrm{Vol_M} + \int_{\partial M} \frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial \mathrm{M}},
\end{aligned}$$

where s is the scalar curvature.

When $A=\overline{c}(X)\overline{c}(Y)c(Z)$, we conclude that:

Corollary 4.11. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=\overline{c}(X)\overline{c}(Y)c(Z)$, then*

(4.44)

$$\widetilde{\mathrm{Wres}}[\pi^+D_A{}^{-1}\circ\pi^+(D_A^*D_AD_A^*)^{-1}]=8192\pi^3\int_M\Big(-\frac{1}{12}s+4|Z|^2g(X,Y)^2$$
$$+\sum_{j=1}^{6}g(\nabla^{TM}_{\widetilde{e}_j}X,Y)g(Z,\widetilde{e}_j)+\sum_{j=1}^{6}g(X,\nabla^{TM}_{\widetilde{e}_j}Y)g(Z,\widetilde{e}_j)+\sum_{j=1}^{6}g(X,Y)g(\nabla^{TM}_{\widetilde{e}_j}Z,\widetilde{e}_j)\Big)d\mathrm{Vol_M}$$
$$+\int_{\partial M}\frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial\mathrm{M}}+\int_{\partial M}\frac{63}{8}\pi\Omega_4 g(\partial x_n,Z)g(X,Y)d\mathrm{Vol}_{\partial\mathrm{M}},$$

where s is the scalar curvature.

When $A=\overline{c}(X)\overline{c}(Y)\overline{c}(Z)$, we find that:

Corollary 4.12. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=\overline{c}(X)\overline{c}(Y)\overline{c}(Z)$, then*

(4.45)

$$\widetilde{\mathrm{Wres}}[\pi^+D_A{}^{-1}\circ\pi^+(D_A^*D_AD_A^*)^{-1}]=8192\pi^3\int_M\Big(-\frac{1}{12}s+5|X|^2|Y|^2|Z|^2-6|X|^2g(Y,Z)^2$$
$$-6|Y|^2g(X,Z)^2-6|Z|^2g(X,Y)^2+12g(X,Y)g(X,Z)g(Y,Z)\Big)d\mathrm{Vol_M}$$
$$+\int_{\partial M}\frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial\mathrm{M}},$$

where s is the scalar curvature.

Next, we prove the Kastler-Kalau-Walze type theorem for 6-dimensional manifold with boundary associated to $D_A{}^3$. From [14], we know that

$$\widetilde{\mathrm{Wres}}[\pi^+D_A{}^{-1}\circ\pi^+D_A{}^{-3}]=\int_M\int_{|\xi|=1}\mathrm{trace}_{\wedge^*T^*M}[\sigma_{-4}(D_A{}^{-4})]\sigma(\xi)dx+\int_{\partial M}\overline{\Phi},\tag{4.46}$$

where $\widetilde{\mathrm{Wres}}$ denote noncommutative residue on minifolds with boundary,

(4.47)

$$\overline{\Phi}=\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\sum_{j,k=0}^{\infty}\sum\frac{(-i)^{|\alpha|+j+k+1}}{\alpha!(j+k+1)!}\times\mathrm{trace}_{\wedge^*T^*M}[\partial_{x_n}^j\partial_{\xi'}^{\alpha}\partial_{\xi_n}^k\sigma_r^+(D_A{}^{-1})(x',0,\xi',\xi_n)$$
$$\times\partial_{x'}^{\alpha}\partial_{\xi_n}^{j+1}\partial_{x_n}^k\sigma_l(D_A{}^{-3})(x',0,\xi',\xi_n)]d\xi_n\sigma(\xi')dx',$$

and the sum is taken over $r+\ell-k-j-|\alpha|-1=-6,\ r\leq-1,\ell\leq-3$.

By Theorem 2.2, we compute the interior term of (4.46)

$$\begin{aligned}(4.48)\qquad &\int_M\int_{|\xi|=1}\mathrm{trace}_{\wedge^*T^*M}[\sigma_{-4}({D_A}^{-4})]\sigma(\xi)dx\\ &=128\pi^3\int_M \mathrm{tr}\Big(-\frac{1}{12}s+2A^2-\frac{1}{2}\sum_{j=1}^n Ac(\widetilde{e_j})Ac(\widetilde{e_j})\Big)d\mathrm{Vol_M}.\end{aligned}$$

So we only need to compute $\int_{\partial M}\overline{\Phi}$. Let us now turn to compute the specification of ${D_A}^3$.

(4.49)

$$\begin{aligned}{D_A}^3 &=\sum_{i=1}^n c(\widetilde{e_i})\langle\widetilde{e_i},dx_l\rangle(-g^{ij}\partial_l\partial_i\partial_j)+\sum_{i=1}^n c(\widetilde{e_i})\langle\widetilde{e_i},dx_l\rangle\Big\{-(\partial_l g^{ij})\partial_i\partial_j-g^{ij}\Big(4(\sigma_i\\ &+a_i)\partial_j-2\Gamma_{ij}^k\partial_k\Big)\partial_l\Big\}+\sum_{i=1}^n c(\widetilde{e_i})\langle\widetilde{e_i},dx_l\rangle\Big\{-2(\partial_l g^{ij})(\sigma_i+a_i)\partial_j+g^{ij}(\partial_l\Gamma_{ij}^k)\partial_k\\ &-2g^{ij}[(\partial_l\sigma_i)+(\partial_l a_i)]\partial_j+(\partial_l g^{ij})\Gamma_{ij}^k\partial_k+\sum_{j,k}\Big[\partial_l\Big(c(\widetilde{e_j})A+Ac(\widetilde{e_j})\Big)\Big]\langle\widetilde{e_j},dx_k\rangle\partial_k\\ &+\sum_{j,k}\Big(c(\widetilde{e_j})A+Ac(\widetilde{e_j})\Big)\Big[\partial_l\langle\widetilde{e_j},dx_k\rangle\Big]\partial_k\Big\}+\sum_{i=1}^n c(\widetilde{e_i})\langle c(\widetilde{e_i}),dx_l\rangle\partial_l\Big\{-g^{ij}\Big[(\partial_i\sigma_j)\\ &+(\partial_i a_j)+\sigma_i\sigma_j+\sigma_i a_j+a_i\sigma_j+a_i a_j-\Gamma_{ij}^k\sigma_k-\Gamma_{ij}^k a_k\Big]+\sum_{i,j}g^{ij}\Big[c(\partial_i)\partial_j(A)\\ &+c(\partial_i)\sigma_j A+c(\partial_i)a_j A+Ac(\partial_i)\sigma_j+Ac(\partial_i)a_j\Big]-\frac{1}{8}\sum_{i,j,k,l}R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l})\\ &+\frac{1}{4}s+A^2\Big\}+\Big[\sum_{i=1}^n c(\widetilde{e_i})(\sigma_i+a_i)+A\Big](-g^{ij}\partial_i\partial_j)+\sum_{i=1}^n c(\widetilde{e_i})\langle\widetilde{e_i},dx_l\rangle\Big\{2\sum_{j,k}\\ &\Big[c(\widetilde{e_j})A+Ac(\widetilde{e_j})\Big]\times\langle\widetilde{e_j},dx_k\rangle\Big\}\partial_l\partial_k+\Big[\sum_{i=1}^n c(\widetilde{e_i})(\sigma_i+a_i)+A\Big]\Big\{-\sum_{i,j}g^{ij}\Big[2(\sigma_i\\ &+a_i)\partial_j-\Gamma_{ij}^k\partial_k+(\partial_i\sigma_j)+(\partial_i a_j)+\sigma_i\sigma_j+\sigma_i a_j+a_i\sigma_j+a_i a_j-\Gamma_{ij}^k\sigma_k-\Gamma_{ij}^k a_k\Big]\\ &+\sum_{i,j}g^{ij}\Big(c(\partial_i)A+Ac(\partial_i)\Big)\partial_j+\sum_{i,j}g^{ij}\Big[c(\partial_i)\partial_j(A)+c(\partial_i)\sigma_j A+c(\partial_i)a_j A\\ &+Ac(\partial_i)\sigma_j+Ac(\partial_i)a_j\Big]-\frac{1}{8}\sum_{i,j,k,l}R_{ijkl}\overline{c}(\widetilde{e_i})\overline{c}(\widetilde{e_j})c(\widetilde{e_k})c(\widetilde{e_l})+\frac{1}{4}s+A^2\Big\}.\end{aligned}$$

Then, we obtain

Lemma 4.13. *The following identities hold:*

$$\sigma_2({D_A}^3) = \sum_{i,j,l} c(dx_l)\partial_l(g^{ij})\xi_i\xi_j + c(\xi)(4\sigma^k + 4a^k - 2\Gamma^k)\xi_k + 2[|\xi|^2 A - c(\xi)Ac(\xi)] + \frac{1}{4}|\xi|^2 \sum_{s,t,l} \omega_{s,t}(\widetilde{e_l})[c(\widetilde{e_l})\overline{c}(\widetilde{e_s})\overline{c}(\widetilde{e_t}) - c(\widetilde{e_l})c(\widetilde{e_s})c(\widetilde{e_t})] + |\xi|^2 A; \tag{4.50}$$

$$\sigma_3({D_A}^3) = ic(\xi)|\xi|^2.$$

Write

$$\sigma({D_A}^3) \;=\; p_3 + p_2 + p_1 + p_0;\ \sigma({D_A}^{-3}) = \sum_{j=3}^{\infty} q_{-j}. \tag{4.51}$$

By the composition formula of pseudodifferential operators, we have

(4.52)

$$\begin{aligned} 1 = \sigma({D_A}^3 \circ {D_A}^{-3}) &= \sum_{\alpha} \frac{1}{\alpha!}\partial_\xi^\alpha[\sigma({D_A}^3)]D_x^\alpha[\sigma({D_A}^{-3})] \\ &= (p_3 + p_2 + p_1 + p_0)(q_{-3} + q_{-4} + q_{-5} + \cdots) \\ &+ \sum_j (\partial_{\xi_j}p_3 + \partial_{\xi_j}p_2 + \partial_{\xi_j}p_1 + \partial_{\xi_j}p_0)(D_{x_j}q_{-3} + D_{x_j}q_{-4} + D_{x_j}q_{-5} + \cdots) \\ &= p_3q_{-3} + (p_3q_{-4} + p_2q_{-3} + \sum_j \partial_{\xi_j}p_3 D_{x_j}q_{-3}) + \cdots, \end{aligned}$$

by (4.52), we have

$$q_{-3} = p_3^{-1};\ q_{-4} = -p_3^{-1}[p_2p_3^{-1} + \sum_j \partial_{\xi_j}p_3 D_{x_j}(p_3^{-1})]. \tag{4.53}$$

By (4.49)-(4.53), we have some symbols of operators.

Lemma 4.14. *The following identities hold:*

$$\sigma_{-3}({D_A}^{-3}) = \frac{ic(\xi)}{|\xi|^4}; \tag{4.54}$$

$$\sigma_{-4}({D_A}^{-3}) = \frac{c(\xi)\sigma_2({D_A}^3)c(\xi)}{|\xi|^8} + \frac{c(\xi)}{|\xi|^8}\Big(|\xi|^2 c(dx_n)\partial_{x_n}c(\xi') - 2h'(0)c(dx_n)c(\xi) + 2\xi_n c(\xi)\partial_{x_n}c(\xi') + 4\xi_n h'(0)\Big).$$

When $n = 6$, then $\mathrm{tr}_{\wedge^*T^*M}[\mathtt{id}] = 64$, where tr as shorthand of trace. Since the sum is taken over $r + \ell - k - j - |\alpha| - 1 = -6,\ r \le -1, \ell \le -3$, then we have the $\int_{\partial M}\overline{\Phi}$ is the sum of the following five cases:

case (a) (I) $r = -1, l = -3, j = k = 0, |\alpha| = 1.$

By (4.47), we get

(4.55)
$$\overline{\Phi}_1 = -\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\sum_{|\alpha|=1}\mathrm{trace}[\partial_{\xi'}^{\alpha}\pi_{\xi_n}^{+}\sigma_{-1}({D_A}^{-1})\times\partial_{x'}^{\alpha}\partial_{\xi_n}\sigma_{-3}({D_A}^{-3})](x_0)d\xi_n\sigma(\xi')dx'.$$

case (a) (II) $r=-1, l=-3, |\alpha|=k=0, j=1.$

We notice that

$$\overline{\Phi}_2 = -\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial_{x_n}\pi_{\xi_n}^{+}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}^{2}\sigma_{-3}({D_A}^{-3})](x_0)d\xi_n\sigma(\xi')dx'. \tag{4.56}$$

case (a) (III) $r=-1, l=-3, |\alpha|=j=0, k=1.$

By (4.47), we compute that

$$\overline{\Phi}_3 = -\frac{1}{2}\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial_{\xi_n}\pi_{\xi_n}^{+}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}\partial_{x_n}\sigma_{-3}({D_A}^{-3})](x_0)d\xi_n\sigma(\xi')dx'. \tag{4.57}$$

By Lemma 4.2 and Lemma 4.5, we have $\sigma_{-3}((D_A^*D_AD_A^*)^{-1}) = \sigma_{-3}({D_A}^{-3})$, then we obtain

$$\overline{\Phi}_1+\overline{\Phi}_2+\overline{\Phi}_3 = \frac{10}{2}\pi h'(0)\Omega_4 dx', \tag{4.58}$$

where Ω_4 is the canonical volume of S^4.

case (b) $r=-1, l=-4, |\alpha|=j=k=0.$

By (4.47), we notice that

$$\begin{aligned}\overline{\Phi}_4 &= -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\pi_{\xi_n}^{+}\sigma_{-1}({D_A}^{-1})\times\partial_{\xi_n}\sigma_{-4}({D_A}^{-3})](x_0)d\xi_n\sigma(\xi')dx' \\ &= i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\partial_{\xi_n}\pi_{\xi_n}^{+}\sigma_{-1}({D_A}^{-1})\times\sigma_{-4}({D_A}^{-3})](x_0)d\xi_n\sigma(\xi')dx'.\end{aligned} \tag{4.59}$$

In the normal coordinate, $g^{ij}(x_0)=\delta_i^j$ and $\partial_{x_j}(g^{\alpha\beta})(x_0)=0$, if $j<n$; $\partial_{x_j}(g^{\alpha\beta})(x_0) = h'(0)\delta_\beta^\alpha$, if $j=n$. So by [12], when $k<n$, we have $\Gamma^n(x_0)=\frac{5}{2}h'(0)$, $\Gamma^k(x_0)=0$, $\delta^n(x_0)=0$ and $\delta^k=\frac{1}{4}h'(0)c(\widetilde{e_k})c(\widetilde{e_n})$. Then, we obtain

$$\begin{aligned}(4.60)\quad \sigma_{-4}({D_A}^{-3})(x_0) &= \frac{1}{|\xi|^8}c(\xi)\Big(h'(0)c(\xi)\sum_{k<n}\xi_k c(\widetilde{e_k})c(\widetilde{e_n}) - h'(0)c(\xi)\sum_{k<n}\xi_k\overline{c}(\widetilde{e_k})\overline{c}(\widetilde{e_n})\\ &\quad - 5h'(0)\xi_n c(\xi) + 2[|\xi|^2 A - c(\xi)Ac(\xi)] + \frac{5}{4}|\xi|^2 h'(0)c(\widetilde{e_i})\overline{c}(dx_n)\overline{c}(\widetilde{e_i})\\ &\quad - \frac{1}{4}|\xi|^2 h'(0)c(dx_n) + |\xi|^2 A\Big)c(\xi) + \frac{c(\xi)}{|\xi|^8}\Big(|\xi|^2 c(dx_n)\partial_{x_n}c(\xi')\\ &\quad - 2h'(0)c(dx_n)c(\xi) + 2\xi_n c(\xi)\partial_{x_n}c(\xi') + 4\xi_n h'(0)\Big).\end{aligned}$$

By (4.16) and (4.60), we have

$$\begin{aligned}(4.61)\qquad &\mathrm{tr}[\partial_{\xi_n}\pi^+_{\xi_n}\sigma_{-1}({D_A}^{-1})\times\sigma_{-4}({D_A}^{-3})](x_0)|_{|\xi'|=1}\\ &= -\frac{64h'(0)}{2(\xi_n-i)^2(1+\xi_n^2)^4}\big(-\frac{15}{4}i + 16\xi_n + \frac{19}{2}i\xi_n^2 + 8\xi_n^3 + \frac{21}{4}i\xi_n^4\big)\\ &+ \frac{5+6i\xi_n-\xi_n^2}{2(\xi_n-i)^2(1+\xi_n^2)^3}\mathrm{tr}\big[Ac(\xi')\big] + \frac{-i+6\xi_n+5i\xi_n^2}{2(\xi_n-i)^2(1+\xi_n^2)^3}\mathrm{tr}\big[Ac(dx_n)\big].\end{aligned}$$

By applying the formula shown in (4.59), we can calculate

$$\begin{aligned}(4.62)\qquad \overline{\Phi}_4 &= i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{-64h'(0)(-\frac{15}{4}i+16\xi_n+\frac{19}{2}i\xi_n^2+8\xi_n^3+\frac{21}{4}i\xi_n^4)}{2(\xi_n-i)^2(1+\xi_n^2)^4}d\xi_n\sigma(\xi')dx'\\ &+ i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{5+6i\xi_n-\xi_n^2}{2(\xi_n-i)^2(1+\xi_n^2)^3}\mathrm{tr}\big[Ac(\xi')\big]d\xi_n\sigma(\xi')dx'\\ &+ i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\frac{-i+6\xi_n+5i\xi_n^2}{2(\xi_n-i)^2(1+\xi_n^2)^3}\mathrm{tr}\big[Ac(dx_n)\big]d\xi_n\sigma(\xi')dx'\\ &= \frac{111}{2}\pi h'(0)\Omega_4 dx' - \frac{5}{16}\pi\Omega_4\mathrm{tr}\big[Ac(dx_n)\big]dx'.\end{aligned}$$

case (c) $r=-2, l=-3, |\alpha|=j=k=0.$

We calculate

$$(4.63)\qquad \overline{\Phi}_5 = -i\int_{|\xi'|=1}\int_{-\infty}^{+\infty}\mathrm{trace}[\pi^+_{\xi_n}\sigma_{-2}({D_A}^{-1})\times\partial_{\xi_n}\sigma_{-3}({D_A}^{-3})](x_0)d\xi_n\sigma(\xi')dx'.$$

By calculation, we have

$$(4.64)\qquad \overline{\Phi}_5 = -\frac{105}{4}\pi h'(0)\Omega_4 dx' + \frac{161}{512}\pi\Omega_4\mathrm{tr}[Ac(dx_n)]dx'.$$

Now $\overline{\Phi}$ is the sum of the cases (a), (b) and (c), hence

$$(4.65)\qquad \overline{\Phi} = \frac{137}{4}\pi h'(0)\Omega_4 dx' + \frac{1}{512}\pi\Omega_4\mathrm{tr}\big[Ac(dx_n)\big]dx'.$$

Theorem 4.15. *Let M be a 6-dimensional compact oriented manifold with the boundary ∂M and the metric g^M as above, D_A be the perturbation of the de Rham Hodge operator on $\widetilde{M}$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+({D_A}^{-3})]=128\pi^3\int_M \mathrm{tr}\Big(-\frac{1}{12}s+2A^2-\frac{1}{2}\sum_{j=1}^{n}Ac(\widetilde{e_j})Ac(\widetilde{e_j})\Big)d\mathrm{Vol_M}$$
$$+\int_{\partial M}\frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial\mathrm{M}}+\int_{\partial M}\frac{1}{512}\pi\Omega_4\mathrm{tr}\big[Ac(dx_n)\big]d\mathrm{Vol}_{\partial\mathrm{M}}.$$

where s is the scalar curvature.

When $A=c(X)$, we can directly state the subsequent corollary:

Corollary 4.16. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=c(X)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+({D_A}^{-3})]=8192\pi^3\int_M\Big(-\frac{1}{12}s\Big)d\mathrm{Vol_M}$$
$$+\int_{\partial M}\frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial\mathrm{M}}-\int_{\partial M}\frac{1}{8}\pi\Omega_4 g(\partial x_n,X)d\mathrm{Vol}_{\partial\mathrm{M}},\tag{4.66}$$

where s is the scalar curvature.

When $A=\overline{c}(X)$, we can get the following corollary:

Corollary 4.17. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=\overline{c}(X)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+({D_A}^{-3})]=8192\pi^3\int_M\Big(-\frac{1}{12}s-|X|^2\Big)d\mathrm{Vol_M}+\int_{\partial M}\frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial\mathrm{M}},\tag{4.67}$$

where s is the scalar curvature.

When $A=c(X)c(Y)$, we compute $\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+({D_A}^{-3})]$.

Corollary 4.18. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A=c(X)c(Y)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+({D_A}^{-3})]=8192\pi^3\int_M\Big(-\frac{1}{12}s+|X|^2|Y|^2+4g(X,Y)^2\Big)d\mathrm{Vol_M}$$
$$+\int_{\partial M}\frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial\mathrm{M}},\tag{4.68}$$

where s is the scalar curvature.

When $A=c(X)\overline{c}(Y)$, we notice that:

Corollary 4.19. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)\overline{c}(Y)$, then*

(4.69)

$$\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+({D_A}^{-3})] = 8192\pi^3\int_M\Big(-\frac{1}{12}s\Big)d\mathrm{Vol_M} + \int_{\partial M}\frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial\mathrm{M}},$$

where s is the scalar curvature.

When $A = \overline{c}(X)\overline{c}(Y)$, similar to (4.68), we can get the following corollary:

Corollary 4.20. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)\overline{c}(Y)$, then*

(4.70)

$$\begin{aligned}\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+({D_A}^{-3})] &= 8192\pi^3\int_M\Big(-\frac{1}{12}s - |X|^2|Y|^2 + 6g(X,Y)^2\Big)d\mathrm{Vol_M}\\ &+\int_{\partial M}\frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial\mathrm{M}},\end{aligned}$$

where s is the scalar curvature.

When $A = c(X)c(Y)c(Z)$, we compute that:

Corollary 4.21. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = c(X)c(Y)c(Z)$, then*

(4.71)

$$\begin{aligned}\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+({D_A}^{-3})] &= 8192\pi^3\int_M\Big(-\frac{1}{12}s - 2|X|^2|Y|^2|Z|^2 + 2|X|^2g(Y,Z)^2\\ &+2|Y|^2g(X,Z)^2 + 2|Z|^2g(X,Y)^2 - 4g(X,Y)g(X,Z)g(Y,Z)\Big)d\mathrm{Vol_M}\\ &+\int_{\partial M}\frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial\mathrm{M}} + \int_{\partial M}\frac{1}{8}\pi\Omega_4 g(\partial x_n, X)g(Y,Z)d\mathrm{Vol}_{\partial\mathrm{M}}\\ &-\int_{\partial M}\frac{1}{8}\pi\Omega_4 g(\partial x_n, Y)g(X,Z)d\mathrm{Vol}_{\partial\mathrm{M}} + \int_{\partial M}\frac{1}{8}\pi\Omega_4 g(\partial x_n, Z)g(X,Y)d\mathrm{Vol}_{\partial\mathrm{M}},\end{aligned}$$

where s is the scalar curvature.

When $A = \overline{c}(X)c(Y)c(Z)$, we can conclude the following facts:

Corollary 4.22. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)c(Y)c(Z)$, then*

(4.72)

$$\begin{aligned}\widetilde{\mathrm{Wres}}[\pi^+{D_A}^{-1}\circ\pi^+({D_A}^{-3})] &= 8192\pi^3\int_M\Big(-\frac{1}{12}s + 3|X|^2|Y|^2|Z|^2 - 4|X|^2g(Y,Z)^2\Big)d\mathrm{Vol_M}\\ &+\int_{\partial M}\frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial\mathrm{M}},\end{aligned}$$

where s is the scalar curvature.

Computations show that:

Corollary 4.23. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)\overline{c}(Y)c(Z)$, then*

$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+({D_A}^{-3})] = 8192\pi^3 \int_M \Big(-\frac{1}{12}s - 4|X|^2|Y|^2|Z|^2 + 4|Z|^2 g(X,Y)^2 \Big) d\mathrm{Vol_M}$$
$$+ \int_{\partial M} \frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial \mathrm{M}} - \int_{\partial M} \frac{1}{8}\pi\Omega_4 g(\partial x_n, Z) g(X,Y) d\mathrm{Vol}_{\partial \mathrm{M}},$$

where s is the scalar curvature.

When $A = \overline{c}(X)\overline{c}(Y)\overline{c}(Z)$, we can get the following corollary:

Corollary 4.24. *Let M be a 6-dimensional oriented compact manifolds with the boundary ∂M and the metric g^M as above, and let $A = \overline{c}(X)\overline{c}(Y)\overline{c}(Z)$, then*

(4.73)

$$\widetilde{\mathrm{Wres}}[\pi^+ {D_A}^{-1} \circ \pi^+({D_A}^{-3})] = 8192\pi^3 \int_M \Big(-\frac{1}{12}s + 5|X|^2|Y|^2|Z|^2 - 6|X|^2 g(Y,Z)^2$$
$$- 6|Y|^2 g(X,Z)^2 - 6|Z|^2 g(X,Y)^2 + 12 g(X,Y)g(X,Z)g(Y,Z) \Big) d\mathrm{Vol_M}$$
$$+ \int_{\partial M} \frac{137}{4}\pi h'(0)\Omega_4 d\mathrm{Vol}_{\partial \mathrm{M}},$$

where s is the scalar curvature.

5. Acknowledgements

The author was supported in part by NSFC No.11771070. The author thanks the referee for his (or her) careful reading and helpful comments.

References

[1] Guillemin V W.: A new proof of Weyl's formula on the asymptotic distribution of eigenvalues. Adv. Math. 55, no. 2, 131-160, (1985).
[2] Wodzicki M.: Local invariants of spectral asymmetry. Invent. Math. 75(1), 143-178, (1995).
[3] Connes A.: Quantized calculus and applications. 11th International Congress of Mathematical Physics(Paris,1994), Internat Press, Cambridge, MA, 15-36, (1995).
[4] Connes A.: The action functinal in noncommutative geometry. Comm. Math. Phys. 117, 673-683, (1998).
[5] Kastler D.: The Dirac operator and gravitation. Comm. Math. Phys. 166, 633-643, (1995).
[6] Kalau W, Walze M.: Gravity, noncommutative geometry and the Wodzicki residue. J. Geom. Phys. 16, 327-344, (1995).

[7] Ackermann T.: A note on the Wodzicki residue. J. Geom. Phys. 20, 404-406, (1996).
[8] Ugalde W J.: Differential forms and the Wodzicki residue. J. Geom. Phys. 58, 1739-1751, (2008).
[9] Fedosov B V, Golse F, Leichtnam E, Schrohe E.: The noncommutative residue for manifolds with boundary. J. Funct. Anal. 142, 1-31, (1996).
[10] Wang Y.: Diffential forms and the Wodzicki residue for manifolds with boundary. J. Geom. Phys. 56, 731-753, (2006).
[11] Wang Y.: Diffential forms the noncommutative residue for manifolds with boundary in the non-product case. Lett. Math. Phys. 77, 41-51, (2006).
[12] Wang Y.: Gravity and the noncommutative residue for manifolds with boundary. Lett. Math. Phys. 80, 37-56, (2007).
[13] Wang Y.: Lower-dimensional volumes and Kastler-Kalau-Walze type theorem for manifolds with boundary. Commun. Theor. Phys. Vol 54, 38-42, (2010).
[14] Wang J, Wang Y.: The Kastler-Kalau-Walze type theorem for six-dimensional manifolds with boundary. J. Math. Phys. 56, 052501, (2015).
[15] Wei S, Wang Y.: Modified Novikov operators and the Kastler-Kalau-Walze type theorem for manifolds with boundary. Adv. Math. Phys. 1-28, (2020).
[16] Wu T, Wang J, Wang Y.: Dirac-witten operators and the Kastler-Kalau-Walze type theorem for manifolds with boundary. J. Nonliear Math. Phys. arXiv:2103.11842.
[17] Wang Y.: A Kastler-Kalau-Walze type theorem and the spectral action for perturbations of Dirac operators on maniflods with boundary. Abstr. Appl. Anal. 17, 286-299, (2014).
[18] Yu Y.: The index theorem and the heat equation method, nankai tracts in mathematics-Vol.2. World Scientific Publishing, (2001).
[19] Ponge R.: Noncommutative geometry and lower dimensional volumes in Riemannian geometry. Lett. Math. Phys. 83, no.1, 19-32, (2008).

S. Liu
School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China
Email: liusy719@nenu.edu.cn
T. Wu
School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China
Email: wut977@nenu.edu.cn
Y. Wang
School of Mathematics and Statistics, Northeast Normal University, Changchun 130024, China
Email: wangy581@nenu.edu.cn